The arc complexes of bicoloured polygons are balls

Pallavi Panda

Abstract. We prove that the arc complexes of a convex polygon and of a once-punctured polygon with a bicolouring are pseudo-manifolds with boundary and we also give a shelling order. As a consequence we get that the arc complex of an ideal decorated hyperbolic (possibly once-punctured) polygon is a closed ball.

1 Introduction

Given a Euclidean polygon \mathcal{P}_m with $m \ge 4$ vertices, its *arc complex* $\mathcal{A}(\mathcal{P}_m)$ is a pure flag simplicial complex constructed using diagonals and their disjointness (See Definition 2.3). It is a classical result of combinatorics that this complex is a piecewise linear sphere of dimension m-4. Penner [17] proved this result in the context of hyperbolic surfaces and Teichmüller theory. He studied the arc complex of an *ideal polygon*, which is the convex hull in the hyperbolic plane \mathbb{H}^2 of finitely many points (called *ideal*) on the boundary $\partial_{\infty}\mathbb{H}^2$. The diagonals in this case are bi-infinite hyperbolic geodesics with endpoints in this finite set. He attributes the original proof to Whitney.

Topologically, a polygon is a closed disk with *n* marked points on its boundary. More generally, for a finite-type orientable surface with finitely many marked points on its boundary (possibly also with punctures), one constructs its arc complex using embedded arcs whose endpoints the marked points. These were first studied by Harer in [7]. He showed that a specific open dense subset of the arc complex, called the *pruned arc complex*, is an open ball of dimension one less than that of the deformation space of the surface. In [4], Fomin–Shapiro–Thurston established an important link between these arc complexes and cluster algebras. They proved that the arc complexes are subcomplexes of the cluster complexes of some cluster algebras. Furthermore, Fomin and Zelevinksy [5] gave a convex polytopal realisation of the cluster complexes in the finite case. The most famous one is the cluster complex of a convex polygon, whose dual is an associahedron.

The associahedron was discovered by Tamari [22], and then rediscovered ten years later by Stasheff [21]. Its first polytopal realisation was given by Lee [8]. Another famous polytopal realisation, which is used in algebraic topology, was given by Loday in [9]. Sleator–Tarjan–Thurston [20] showed that the *d*-dimensional associahedron has diameter at most 2d - 4 when $d \ge 9$ using hyperbolic geometry. Later Pournin [18] proved the equality using combinatorial methods.

The relationship to cluster algebras was motivated by Penner's Decorated Teichmüller Theory [14],[15]. Penner defined the *lambda length* of a *horoball connection* which is a geodesic arc joining two ideal points decorated by horoballs. These lengths act as coordinates for the decorated

Teichmüller space of a decorated *crowned* hyperbolic surface. This is a non-compact hyperbolic surface with polygonal boundary, where the vertices (called *spikes*) are projections of ideal points decorated with horoballs. Furthermore, using the arc complex, he gave a cell-decomposition of the decorated Teichmüller space.

In [17], Penner studied the topology of the quotient of the arc complex under the canonical action of the pure mapping class group of the surface. He conjectured that this quotient space is a sphere of a certain dimension, but this was later proved to be false by Sullivan. There is a complete list of surfaces (see [16]) for which the statement is true, the ideal polygons and the once-punctured ideal polygons being among them.

In the non-orientable setting, Dupont and Palesi [3] found an analogue of a cluster algebra associated to the arc complexes of non-orientable surfaces. Wilson [24] proved that the arc complex of a Möbius strip with marked points on the boundary is spherical. He also reproved the sphericity of the arc complexes of a convex polygon and a once-punctured polygon. The main ingredient he used was the shellability of pure simplicial complexes. This is an ordering of all the maximal simplices of the complex so that the intersection of the *k*-th simplex with the union of the first (k - 1) simplices is always a pure complex of codimension one. He gave shelling orders for the arc complexes of all three surfaces and then concluded using a result by Danaraj and Klee [1] which states that a shellable *d*-pseudo-manifold without boundary is a combinatorial sphere.

The convex core of a crowned hyperbolic surface without punctures is a compact hyperbolic surface with non-empty totally geodesic boundary. An *admissible deformation* of such a hyperbolic surface is an infinitesimal deformation that uniformly lengthens all non-trivial closed geodesics. Goldman–Labourie–Margulis, in [6], proved that the subspace of admissible deformations forms an open convex cone called the *admissible cone*. One can construct the arc complex of this surface generated by the isotopy classes of embedded arcs with endpoints on the boundary. The pruned arc complex of such a surface once again forms an open ball of dimension one less than that of the Te-ichmüller space, which is obtained by reinterpreting Harer's result. Hyperbolic strip deformations were first introduced by Thurston in [23]. See, for example, Section 1.2 in [2] for the definition. Danciger–Guéritaud–Kassel [2] showed that the pruned arc complex parametrises the positively projectivised admissible cone. To a positively weighted arc, the authors associated a unique admissible deformation of the surface by performing hyperbolic strip deformations along the arc, whose strip width is given by the weight.

Motivated by the above works, in [10] we studied the arc complexes of a decorated ideal polygon $\widehat{\Pi_n^{\diamond}}$ $(n \ge 3)$ and a decorated once-punctured ideal polygon $\widehat{\Pi_n^{\times}}$ $(n \ge 2)$. These are generated by finite arcs whose end points lie on the boundary and infinite arcs with one end on the vertex and the finite endpoint on the boundary. In both of these cases the arc complexes are finite but their topologies were unknown. We proved that the pruned arc complex, which is just the interior of the arc complex in these cases, parametrises all the infinitesimal deformations that uniformly lengthened all horoball connections in these polygons. As a result, we found that the interior of the complexes were open balls. In this paper we show that the full arc complexes are *PL*-balls.

The arc complex of a decorated (resp. once-punctured) polygon $\widehat{\Pi_n^{\diamond}}$ (resp. $\widehat{\Pi_n^{\times}}$) is combina-

torially equivalent to a certain subcomplex of the arc complex of a Euclidean (once-punctured) polygon \mathcal{P}_{2n} (resp. $\mathcal{P}_{2n}^{\times}$), endowed with an alternate bicolouring (see Definition 2.2). In this paper, we prove something more general:

Theorem. The subcomplex $\mathcal{Y}(\mathcal{P}_m)$ of a polygon \mathcal{P}_m with any bicolouring is a closed ball of dimension m-4. Similarly, the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times})$ of a once-punctured polygon \mathcal{P}_m^{\times} with any bicolouring is a closed ball of dimension m-2.

Our proofs are heavily motivated by the works of Wilson and Danaraj-Klee. We use purely combinatorial topology tools while in [10] we use methods from hyperbolic geometry. As corollaries to these theorems we get the arc complexes of the two decorated hyperbolic polygons are closed balls.

Corollary 1.1. For $n \ge 3$, the arc complex $\mathcal{A}(\widehat{\Pi_n^{\diamond}})$ of a decorated ideal polygon $\widehat{\Pi_n^{\diamond}}$ is *PL*-homeomorphic to a closed ball of dimension 2n - 4.

Corollary 1.2. For $n \ge 2$, the arc complex $\mathcal{A}(\widehat{\Pi_n^{\times}})$ of a decorated once-punctured ideal polygon $\widehat{\Pi_n^{\times}}$ is *PL*-homeomorphic to a closed ball of dimension 2n - 2.

These results have applications for the study the boundary of the admissible cone of the decorated triangles using the boundary simplices of the arc complex. A facet on the boundary of the admissible cone corresponds to all those infinitesimal deformations that lengthen every horoball connection except one. As we mentioned above, the full arc complex is rarely a *PL*-manifold so this approach is not possible for "bigger" surfaces. Nonetheless it will shed some useful light on how to parametrise the facets using arc complexes of subsurfaces.

The paper is structured into sections in the following way: Section 2 recapitulates the necessary vocabulary and results from simplicial complexes, arc complexes and hyperbolic geometry. Section 3 contains the proofs of our main theorems. Section 4 describes the link between the arc complexes of the Euclidean polygons and the decorated hyperbolic polygons.

Acknowledgements. This work was done at Université de Luxembourg supported by Luxembourg National Research Fund OPEN grant O19/13865598. I would like to thank my supervisor Hugo Parlier for giving me this opportunity. I would also like to thank Lionel Pournin for helpful discussions and encouragement.

2 Setup

2.1 Simplicial complex

In this section we recall relevant definitions and results on finite simplicial complexes.

A simplicial complex is called *pure* if all of its maximal simplices have the same dimension. The *dual graph* of a simplicial complex is the graph whose vertices are the maximal simplices and two vertices are joined by an edge if the corresponding maximal simplices share a codimension 1 face. A pure simplicial complex is said to be *strongly connected* if its dual graph is connected.

A *d-pseudo-manifold with boundary* is a pure strongly connected *d*-simplicial complex in which every (d-1)-simplex is contained in atmost two *d*-simplices.Note that the boundary of such a simplicial complex is formed by all (d-1)-simplices that are contained in exactly one *d*-simplex.

Next we recall the definition of a shelling of pure simplicial complexes.

Let *X* be a pure finite simplicial complex of dimension *d*. A *shelling* of *X* is an enumeration of its maximal simplices $\mathcal{T} : (C_1, \ldots, C_n)$ such that for every $1 \le k \le n$, the intersection $\begin{pmatrix} k-1 \\ \bigcup \\ j=1 \end{pmatrix} \cap C_k$ is a pure simplicial complex of dimension d-1.

The following is a lemma linking shellability and join of two simplicial complexes that we shall use in the proof of our main theorems. See [19] for a proof.

Lemma 2.1. Two complexes X, Y are shellable if and only if $X \bowtie Y$ is shellable.

We will use the following result by Danaraj and Klee [1] to prove our main theorems.

Theorem 2.2. A shellable *d*-pseudo-manifold with boundary is PL-homeomorphic to a closed ball of dimension d.

2.2 Arcs and arc complexes

In this section we recall the relevant definitions and results the arc complex of polygons that will be used in the rest of the paper.

We denote by $S_{g,n}$ a surface with genus $g (\geq 0)$ and $n (\geq 1)$ marked points on its boundary.

Definition 2.3. The arc complex $\mathcal{A}(S_{g,n})$ of a finite-type surface $S_{g,n}$ with marked points on its boundary is a simplicial complex defined in the following way: the 0-skeleton is given by the embedded arcs with their endpoints on the marked points of the polygon, up to homotopy relative to the endpoints. For $k \ge 1$, every k simplex is given by a (k + 1)-tuple of pairwise disjoint and distinct arcs, up to homotopy.

In this section, we are going to consider only two types of surfaces: convex polygons \mathcal{P}_m $(m \ge 4)$ and once-punctured convex polygons \mathcal{P}_m^{\times} , $(m \ge 2)$. In the case of a convex polygon, these homotopy classes are simply the diagonals joining two vertices of the polgon. To avoid confusion, we will refer to the arcs of a punctured polygon as diagonals as well. A *maximal* diagonal of \mathcal{P}_m^{\times} is the diagonal with both its endpoints coinciding on a vertex of the polygon. The blue diagonal in the left panel of Fig. (2) is a maximal diagonal.

The following is a classical fact from combinatorics. See, for instance, [17] for a proof by Penner.



(a) The diagonals of a hexagon \mathcal{P}_6 and $\mathcal{A}(\mathcal{P}_6)$



(b) The flattened perspective of $\mathcal{A}(\mathcal{P}_6)$ and its dual associahedron of dimension 2

Figure 1

Theorem 2.4. The arc complex of a convex polygon \mathcal{P}_m ($m \ge 4$) is PL-homemorphic to a sphere of dimension m - 4.

Fig.(1a) shows the diagonals and the arc complex of a hexagon. The diagonals corresponding to the 0-skeleton of a maximal simplex of $\mathcal{A}(\mathcal{P}_m)$ decomposes the polygon \mathcal{P}_m into triangles. In the case of a punctured polygon \mathcal{P}_m^{\times} , the diagonals decompose it into triangles and a once-punctured disc with one marked point on its boundary. Hence a maximal simplex will be alternatively referred to as a *triangulation* of the polygon. A triangulation, all of whose diagonals are incident on the same vertex, is called a *fan* triangulation.

The dual graph to $\mathcal{A}(S_{g,n})$ is called a *flip graph*. In [11], [12] and [13], Parlier and Pournin study the diameter growth of flip graphs, up to the action of pure mapping class groups, of certain families $S_{g,n}$.

In the case of a convex polygon, dual graph is the 1-skeleton of a convex polytope is the associahedron, which we mentioned in the introduction. See Fig.(1b) for the associahedron of dimension 3.

The following theorem about the arc complex of once-punctured polygons was proved by Penner in [17].

Theorem 2.5. The arc complex $\mathcal{A}(\mathcal{P}_m^{\times})$ of a punctured m-gon, $(m \ge 2)$, is PL-homeomorphic to a sphere of dimension n - 2.

Fig.2 gives an illustration of the arc complex of a once-punctured quadrilateral. The blue diagonal in the left panel is a maximal arc.



Figure 2: The three types of diagonals and the full arc complex of \mathcal{P}_4^{\times}

Bicolourings. Given a convex (possibly punctured) polygon we consider all possible colourings of its vertices with two colours, say red (R) and blue (B), so that there is a vertex of each colour. Such a colouring is called a *bicolouring*. A bicolouring is called *non-trivial* if there is at least one R - R diagonal. A trivial bicolouring can be of two types:

- 1. there is exactly one red vertex,
- 2. there are exactly two red vertices and they are consecutive.



Figure 3

See Fig.(3) for examples of trivial and non-trivial bicolourings of a quadrilateral \mathcal{P}_4 .

One example of a non-trivial bicolouring of \mathcal{P}_m , with *m* even is the *alternate bicolouring*: for every pair of consecutive vertices, exactly one is blue. This will be used in Section 4 to link the Euclidean polygons with their hyperbolic siblings.

We denote by $\mathcal{Y}(\mathcal{P}_m)$ (resp. $\mathcal{Y}(\mathcal{P}_m^{\times})$) the subcomplex of $\mathcal{A}(\mathcal{P}_m)$ (resp. $\mathcal{A}(\mathcal{P}_m^{\times})$) generated by the R - B and B - B diagonals only. We call these diagonals *permitted* and the R - R diagonals are called *rejected*. A simplex of $\mathcal{Y}(\mathcal{P}_m)$ using permitted diagonals is called *permissible*.

Remark 2.1. Any fan triangulation of a bicoloured polygon based at a blue vertex is permissible.

Remark 2.2. In the case of a trivial bicolouring, the subcomplex is the full arc complex because there are no rejected diagonals. This is why the bicolouring is named "trivial".



Figure 4

Figs. (4) and (5) show the subcomplexes $\mathcal{Y}(\mathcal{P}_6)$ and $\mathcal{Y}(\mathcal{P}_4^{\times})$ when the polygons \mathcal{P}_6 , \mathcal{P}_4^{\times} have alternate bicolouring. In both the cases, they are closed 2-balls.

2.3 Shelling of finite arc complexes

Wilson gave an equivalent definition for shelling in [24], in the context of arc complexes.

Property 1 (Wilson). Let *X* be a pure simplicial complex. There exists an enumeration of all the maximal faces of *X*, denoted by $\mathcal{T} : (C_1, \ldots, C_n)$, such that for every two positive integers *j* and *k* satisfying $j < k \le n$, there exists a positive integer i < k such that





- (i) $C_i \cap C_k$ is a codimension one simplex,
- (ii) $C_i \cap C_k \subset C_i \cap C_k$.

In [24] (Proposition 3.5), Wilson proved that in the context of finite arc complexes, there is an equivalence between shellability and Property (1).

Lemma 2.6. Suppose that $S_{g,n}$ is a surface with finite arc complex. Then the complex $\mathcal{A}(S_{g,n})$ is shellable if and only if it satisfies Property (1).

In the following Lemma we show that the same holds for any pure subcomplex of the arc complex.

Lemma 2.7. A pure codimension zero subcomplex X of $\mathcal{A}(S_{g,n})$ is shellable if and only if it satisfies *Property* (1).

Proof. We show that the enumeration $\mathcal{T} = C_1 \dots, C_n$ given by the property works as a shelling order. So we need to show that for every $1 \le k \le n$, the simplicial complex $B_k := \begin{pmatrix} k-1 \\ \bigcup \\ j=1 \end{pmatrix} \cap C_k$ is a pure simplicial complex of dimension d-1, where d is the dimension of the subcomplex X as well as $\mathcal{A}(S_{g,n})$. Let $\sigma \subset B_k$ be any simplex of codimension more than one. Then it is contained in $C_j \cap C_k$ for some $j \in \{1, \dots, k-1\}$. Since X is a subcomplex of the arc complex, the complex $C_j \cap C_k$ is in fact a simplex. From Property (1) we get that there is an i < k such that $C_i \cap C_k$ is of dimension d-1 and $C_j \cap C_k \subset C_i \cap C_k \subset B_k$. Hence we get that the simplex σ is not maximal in B_k and the dimension of any maximal simplex containing σ is d-1.

Conversely, let us suppose that X is shellable with a shelling order $\mathcal{T}_n = (T_1, \dots, T_n)$. Consider two maximal simplices T_j, T_k with $j < k \le n$. If $T_j \cap T_k = \emptyset$, there is nothing to show. So we assume that $T_j \cap T_k \ne \emptyset$. Once again, the intersection $T_j \cap T_k$ is a simplex because X is a subcomplex of the arc complex. Let $B_k := \begin{pmatrix} k-1 \\ \bigcup i=1 \end{pmatrix} \cap T_k$ as before. From the hypothesis, B_k is a pure simplicial complex of dimension d-1. So we get that $T_j \cap T_k$ is contained in a (d-1)-simplex $T_i \cap T_k$, where T_i is a maximal simplex with i < k. This T_i satisfies the two conditions of Property (1).

The orientable surfaces with finite arc complexes are convex polygons \mathcal{P}_m , once-punctured polygons \mathcal{P}_m^{\times} , annulus with one marked point in one boundary component and *m* marked points on the other boundary component. Note that a once-punctured convex polygon is an annulus with marked points only on one boundary component. In [3], Dupont and Palesi introduce the *quasi* arc complex of a non-orientable surface where they include any one-sided curve in the 0-skeleton. The only non-orientable surface with finite (quasi) arc complex is a Möbius strip with marked points on its boundary. Wilson proved the sphericity of this complex in [24], using shellability. In order to prove the shellability, he first gave shelling orders of the arc complexes of a convex polygon and a once-punctured polygon.

Theorem 2.8 (Wilson, Proposition 2.11, Claim 1.4). *The arc complex* $\mathcal{A}(\mathcal{P}_m)$ *of a convex polygon* \mathcal{P}_m ($m \ge 4$) *is a* (m - 4)*-pseudo-manifold with boundary and is shellable.*

Theorem 2.9 (Wilson, Proposition 2.11, 3.13). *The arc complex* $\mathcal{A}(\mathcal{P}_m^{\times})$ *of a punctured polygon* \mathcal{P}_m^{\times} ($m \ge 2$) *is a* (m - 2)*-pseudo-manifold with boundary and is shellable.*

2.4 Hyperbolic geometry

In this section we recall a few definitions from hyperbolic geometry which will be used in 4 to give the application of our main theorems in the context of hyperbolic polygons, as previewed in the introduction.

The hyperbolic plane, denoted by \mathbb{H}^2 , is the unique (up to isometry) complete simply-connected Riemannian 2-manifold of constant curvature equal to -1. Its boundary, denoted by $\partial_{\infty}\mathbb{H}^2$, is homeomorphic to a circle \mathbb{S}^1 . Its orientation preserving isometry group is isomorphic to PSL(2, \mathbb{R}). This group has three types of elements: elliptic (one fixed point in \mathbb{H}^2), parabolic (one fixed point in $\partial_{\infty}\mathbb{H}^2$) and hyperbolic (two fixed points in $\partial_{\infty}\mathbb{H}^2$). A horocycle based at a point in $\partial_{\infty}\mathbb{H}^2$ is the orbit of a parabolic element fixing that point. A horoball is the convex hull of a horocycle.

An *ideal n-gon*, denoted by Π_n^{\diamond} , is defined as the convex hull in \mathbb{H}^2 of $n \geq 3$ distinct points on $\partial_{\infty}\mathbb{H}^2$. The points on the boundary are called *vertices* and the *edges* are infinite geodesics of \mathbb{H}^2 joining two consecutive vertices. The restriction of the hyperbolic metric to an ideal polygon gives it a complete finite-area hyperbolic metric with geodesic boundary.

For $n \ge 2$, an *ideal once-punctured n-gon*, denoted by Π_n^{\times} , is another non-compact complete hyperbolic surface with geodesic boundary, obtained from an ideal (n + 2)-gon, by identifying two consecutive edges using a parabolic element of PSL $(2, \mathbb{R})$ that fixes the common vertex. The edges of the polygon are the connected components of the boundary. The vertices are the quotients of the vertices of Π_{n+2}^{\diamond} .

3 Main theorems

3.1 Bicoloured convex polygon

In this section we prove that the subcomplex $\mathcal{Y}(\mathcal{P}_m) \subset \mathcal{A}(\mathcal{P}_m)$ is PL-homeomorphic to a closed ball. Firstly we show that this subcomplex is a pseudo-manifold with boundary when the bicolouring is non-trivial.

Lemma 3.1. For $m \ge 4$ and any non-trivial bicolouring of \mathcal{P}_m , the subcomplex $\mathcal{Y}(\mathcal{P}_m)$ of $\mathcal{A}(\mathcal{P}_m)$ is pure of dimension m - 4.

Proof. Let σ be a simplex of $\mathcal{Y}(\mathcal{P}_m)$ of dimension $k \ge 1$. The diagonals corresponding to the 0-simplices of σ decompose the polygon \mathcal{P}_m into k + 2 regions, some of which are untriangulated polygons with at least 4 vertices. Since these diagonals are permitted, every smaller untriangulated polygon has a blue vertex. We triangulate each one of them with a fan triangulation based at one such blue vertex. Since the dimension of the maximal simplex in the uncoloured polygon is m - 4, we get that any maximal simplex containing σ is of dimension m - 4.

Next, we show that the subcomplex is strongly connected.





Lemma 3.2. For $m \ge 4$ and any non-trivial bicolouring, the subcomplex $\mathcal{Y}(\mathcal{P}_m)$ is strongly connected.

Proof. We will prove that *T* is connected by flips to the fan triangulation based at any blue vertex, by induction on *m*. For m = 4, there are two non-trivial bicolourings possible: either alternating *R*, *B* vertices or three *R* vertices. In each case the subcomplex $\mathcal{Y}(\mathcal{P}_m)$ is a single 0-simplex which is also the fan triangulation based simultaneously at the two blue vertices. Suppose that for any bicolouring of polygons with $1, \ldots, m - 1$ vertices, any triangulation *T* is connected by flips to a

fan triangulation. Now, consider the polygon \mathcal{P}_m . We name v_1 any blue vertex and we enumerate the rest of the vertices in the clockwise direction. Let T_0 be the fan triangulation based at v_1 .

Suppose that $l_i \in T \cap T_0$ for some i = 3, ..., m - 2. See the left panel of Fig.(3.2). It divides the polygon \mathcal{P}_m into two smaller polygons $\mathcal{P}_i, \mathcal{P}_{m-i+1}$ whose vertices are $v_1, v_2, ..., v_i$ and $v_i, v_{i+1}, ..., v_1$, respectively. The triangulation T triangulates these two polygons. It is possible that that one of these smaller polygons have a trivial coloring. In this case, we use fact that the dual of the full arc complex (the associahedron) is connected. In particular, the restriction of T to this smaller polygons have a non-trivial bicolouring, we use our induction hypothesis inside each of them to connect T by flips to the fan triangulation based at v_1 . So T, as a triangulation of \mathcal{P}_m , is connected by flips to the triangulation T_0 .

Now we suppose that there are no diagonals in T that are incident at v_1 and since T is a permissible triangulation, the vertex v_1 is contained in a unique triangle with vertices at v_2, v_m such that at least one is blue. The diagonal joining v_2, v_m is the edge of exactly one other triangle, whose third vertex is say $v_j \neq v_1$. Let T' be the triangulation obtained by flipping the diagonal joining v_2, v_m with the permissible diagonal l_j joining v_1, v_j . Now we are in the first case. This concludes the induction step.

Next we show that

Lemma 3.3. Every codimension one simplex of the subcomplex $\mathcal{Y}(\mathcal{P}_m)$ is contained in at most two maximal simplices of $\mathcal{Y}(\mathcal{P}_m)$.

Proof. Let σ be a d – 1-simplex, where d is the dimension of $\mathcal{Y}(\mathcal{P}_m)$. Since the complex $\mathcal{A}(\mathcal{P}_m)$ of an uncoloured polygon is a pseudo-manifold, we have that σ is contained in exactly two d-simplices of $\mathcal{A}(\mathcal{P}_m)$. These two simplices represent two triangulations which have all but one diagonal in common. These two diagonals can be either R - B and B - B or both B - B. Indeed, if both were R - R diagonals or both were R - B diagonals, then there would be a R - R diagonal inside σ bounding the quadrilateral region where the two intersecting diagonals lie. This is impossible because $\sigma \subset \mathcal{Y}(\mathcal{P}_m)$. When both the diagonals are of type B - B, then both the maximal simplices lie inside $\mathcal{Y}(\mathcal{P}_m)$. Otherwise only the maximal simplex generated by R - B, B - B diagonals lies in $\mathcal{Y}(\mathcal{P}_m)$. In this case, σ is a boundary simplex of $\mathcal{Y}(\mathcal{P}_m)$.

Using Lemmas (3.1)-(3.3), we get that

Theorem 3.4. The subcomplex $\mathcal{Y}(\mathcal{P}_m)$ is a *d*-pseudo-manifold with boundary.

As the last step, we prove that the subcomplex is shellable.

Theorem 3.5. For $m \ge 4$, the simplicial complex $\mathcal{Y}(\mathcal{P}_m)$ satisfies Property (1).

Proof. We use induction on the number of vertices. For m = 4, there are two possible colourings. In both the cases, the complex is a single 0-simplex which satisfies the property trivially. Let us



now suppose that the statement is true for $4, \ldots, m - 1$. Due to the bicolouring, there is a edge of the polygon joining one red (the right one, say) and one blue vertex (left one). Enumerate the vertices in the anticlockwise direction starting from the blue vertex so that the red vertex is numbered *m*. See Fig.7. For $i = 3, \ldots, m - 2$, let σ_i be the simplex of $\mathcal{Y}(\mathcal{P}_m)$ generated by the two diagonals joining the vertex *i* to the vertices 1 and *m*. In Fig. (7), the diagonals of σ_3 are drawn in grey and those of σ_5 are in greem. If the vertex 2 is a blue vertex, then let σ_2 be the 0-simplices corresponding to the diagonal joining (2, m). Let σ_{m-1} be the 0-simplices corresponding to the diagonals joining (1, m - 1), which is either R - B or B - B because the vertex 1 is a blue vertex. For each *i*, the simplex σ_i decomposes the polygon \mathcal{P}_m into an *i*-gon, an (m - i + 1)-gon and a triangle with vertices at 1, m, i. So the star of the simplex σ_i , $St(\sigma_i)$, in the complex $\mathcal{Y}(\mathcal{P}_m)$ is the join of $\mathcal{Y}(\mathcal{P}_i)$ and $\mathcal{Y}(\mathcal{P}_{m-i+2})$.

From the induction hypothesis, there exist enumerations of the maximal simplices of $\mathcal{Y}(\mathcal{P}_i)$ and $\mathcal{Y}(\mathcal{P}_{m-i+2})$, respectively, satisfying Property (1). From Lemma (2.7), the complexes $\mathcal{Y}(\mathcal{P}_i)$, $\mathcal{Y}(\mathcal{P}_{m-i+1})$ are shellable. From Lemma (2.1), we get that $St(\sigma_i)$ is shellable. Again from Lemma (2.7), we get an enumeration \mathcal{T}^i of the maximal simplices of $St(\sigma_i)$. Let $J \subset \{2, \ldots, m-1\}$ be the blue vertices. Consider the ordering $\mathcal{T} := \mathcal{T}^{m-1}, \mathcal{T}^{i_1}1, \ldots, \mathcal{T}^{i_p}$, where $i_1 > i_2 > \ldots > i_p$ are all the elements of J. We claim that \mathcal{T} is an ordering of all the maximal simplices of $\mathcal{Y}(\mathcal{P}_m)$ satisfying the conditions of Property(1).

First we show that every triangulation appears in \mathcal{T}^i for exactly one $i \in \{3, m-2\}$. Let T be any triangulation of \mathcal{P}_m generated by R - B and R - R diagonals. Then either there is no diagonal of T incident at the vertex 1, or there is at least one, in which case the last diagonal of T (anti clockwise direction) must join 1 to a blue vertex, say $k \in \{2, m - 1\}$, of the polygon. In Fig. (7), we have that the last blue vertex joined to 1 is at 5. Since this diagonal is the last one, there must be a diagonal of T joining the vertices k and m.

Hence $T \in \mathcal{T}^{j}$. Also, for every $j \neq j'$, we have $\mathcal{T}^{j} \cap \mathcal{T}^{j'} = \emptyset$ because the diagonal (j,m) intersects the diagonal (1, j').

Now suppose that S, T are two maximal simplices of $\mathcal{Y}(\mathcal{P}_m)$ such that S precedes T. We need to find another maximal simplex P preceding T such that P is obtained by flipping a diagonal of



Figure 8: Complementary regions to a maximal diagonal

T such that any diagonal common to *S* and *T* must be common to *P* and *T*. If *S*, *T* belong to the same \mathcal{T}_i , then we can conclude immediately by using the induction hypothesis. So we assume that $S \in \mathcal{T}_j$ and $T \in \mathcal{T}_k$ with j > k. Any diagonal common to both the triangulations *S* and *T* must lie outside the quadrilateral 1, *k*, *j*, *m* because they must be disjoint to the diagonals (1, k), (k, j), (j, m). See Fig.7. Now let *P* be the triangulation obtained from *T* by flipping the diagonal (m, k) by (1, j). Since this operation happens inside the quadrilateral, we get that $S \cap T \subset P \cap T$. Also by construction, $P \in St(\sigma_j)$ which precedes $St(\sigma_k)$ and hence *T*.

From Theorem 2.2, Lemma 3.4 and Theorem 3.5, we get that

Theorem 3.6. For $m \ge 4$, the subcomplex $\mathcal{Y}(\mathcal{P}_m)$ of a convex polygon \mathcal{P}_m with a non-trivial bicolouring is a PL-homeomorphic closed ball of dimension m - 4.

3.2 Punctured bicoloured polygons

In this section, we prove that the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times}) \subset \mathcal{A}(\mathcal{P}_m^{\times})$ is a closed ball of dimension m - 2 for any non-trivial bicolouring. First we show that the subcomplex is a pseudo-manifold with boundary.

Lemma 3.7. For $m \ge 2$ and any non-trivial bicolouring, the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times})$ of $\mathcal{A}(\mathcal{P}_m^{\times})$ is pure of dimension m - 2.

Proof. Let σ be any simplex of $\mathcal{Y}(\mathcal{P}_m^{\times})$. If one of its 0-simplices correspond to a maximal diagonal, one of the regions in its complement is a bicoloured unpunctured polygon. See Fig.(8).

Using Lemma 3.1, we have that σ is contained in some permissible maximal simplex of $\mathcal{Y}(\mathcal{P}_m^{\times})$. If none of the 0-simplices represent a maximal diagonal, let $l \in \sigma$ be the diagonal that separates the puncture from the rest of the diagonals of σ . At least one of the endpoints of l is a blue vertex. We add to σ the maximal diagonal based at this vertex. Again use Lemma 3.1 on its complement to get a maximal simplex of $\mathcal{Y}(\mathcal{P}_m^{\times})$ containing σ .

Next we show that the subcomplex is strongly connected.

Lemma 3.8. For $m \ge 2$ and any non-trivial bicolouring, the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times})$ is strongly connected.

Proof. Let *T* be a permissible triangulation of \mathcal{P}_m^{\times} . Then there is a maximal diagonal based at a blue vertex. Cutting along the diagonal we get a bicoloured unpunctured polgygon, triangulated by the rest of the diagonals of *T*. Let T_0 be the fan triangulation based at one of the two blue vertices given by the maximal diagonal. By Lemma 3.2, *T* is connected to T_0 by flips and all the intermediate triangulations remain disjoint to the maximal diagonal.

Now we show that

Lemma 3.9. Every codimension one simplex of the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times})$ is contained in at most two maximal simplices of $\mathcal{Y}(\mathcal{P}_m^{\times})$.

Proof. Let σ be a codimension one simplex of $\mathcal{Y}(\mathcal{P}_m^{\times})$. From Theorem 2.9 we get that σ is contained in exactly two codimension 0 simplices η, η' of $\mathcal{A}(\mathcal{P}_m^{\times})$. From Lemma 3.7, we know that at least one of η, η' is inside $\mathcal{Y}(\mathcal{P}_m^{\times})$. If none of the 0-simplices of η, η' contain a rejected diagonal, then they both are inside $\mathcal{Y}(\mathcal{P}_m^{\times})$, otherwise only one of them is.

Finally we prove that the the subcomplex is shellable.

Theorem 3.10. For $m \ge 2$, the simplicial complex $\mathcal{Y}(\mathcal{P}_m^{\times})$ satisfies Property (1).

Proof. Like in the proof of Theorem 3.5, we enumerate the vertices in the anticlockwise direction so that the first vertex is blue and the *m*-th is red. Let d_{i_1}, \ldots, d_{i_p} be the maximal diagonals based at the blue vertices i_1, \ldots, i_p , respectively. The star $St(d_{i_j})$ is the coloured arc complex $\mathcal{Y}(\mathcal{P}_{m+1})$ of the unpunctured bicoloured polygon \mathcal{P}_{m+1} obtained cutting \mathcal{P}_m^{\times} along d_{i_j} . Let \mathcal{T}^i be the shelling order given by Theorem 3.5. We claim that $\mathcal{T} := \mathcal{T}^{i_1}, \ldots, \mathcal{T}^{i_p}$ is a shelling order of $\mathcal{Y}(\mathcal{P}_m^{\times})$ satisfying Property 1. Since every triangulation of \mathcal{P}_m^{\times} contains exactly one maximal diagonal based at some blue vertex, \mathcal{T} is an enumeration of the maximal simplices of $\mathcal{Y}(\mathcal{P}_m^{\times})$.

Now suppose that *S*, *T* are two maximal simplices of $\mathcal{Y}(\mathcal{P}_m^{\times})$ such that *S* precedes *T*. If they belong to the same star $\operatorname{St}(d_{i_j})$, then we conclude using the proof of Theorem 3.5. Now let $S \in \operatorname{St}(d_{i_j})$ and $T \in \operatorname{St}(d_{i_k})$ with $j < k \leq p$. The diagonals of $S \cap T$, if any, lie outside the region containing $d_{i_k} \cup d_{i_k}$ bounded by the two diagonals joining the vertices i_j, i_k . See Fig.(9). Let *P* be the triangulation obtained from *T* by flipping the diagonal d_{i_k} with d_{i_j} . So $P \in \operatorname{St}(d_{i_j})$, hence preceding *T* in the order. Also by construction of *P*, we have that $S \cap T \subset P \cap T$. This concludes the proof.

See Fig.(10) for a shelling order for the subcomplex $\mathcal{Y}(\mathcal{P}_4^{\times})$ when \mathcal{P}_4^{\times} is endowed with an alternate colouring. In this case, there are only two permissible maximal diagonals: d_1 and d_3 . Cutting along d_1 , we get an unpunctured pentagon with a non-trivial bicolouring. As in the proof of Theorem (3.6), we consider the stars of σ_2 (purple) and σ_4 (orange). The subcomplex $\mathcal{Y}(\mathcal{P}_5)$ is a closed 1-ball as shown in the bottom right panel. A shelling order of this subcomplex gives a shelling order for $St(d_2)$. Similarly, we get a shelling order for $St(d_3)$. Combining the two, we get a shelling order for $\mathcal{Y}(\mathcal{P}_4^{\times})$.

From Lemmas (3.7),(3.8),(3.9) and Theorem (3.11), we get that,



Figure 10: A shelling order for $\mathcal{Y}(\mathcal{P}_4^{\times})$ with alternate bicolouring

Theorem 3.11. For $m \ge 2$, the subcomplex $\mathcal{Y}(\mathcal{P}_m^{\times})$ of a once-punctured polygon \mathcal{P}_m^{\times} with any non-trivial bicolouring is a closed ball of dimension m - 2.



Figure 11: The two types of decorated hyperbolic polygons

4 Applications: decorated hyperbolic polygons

Decorated Polygons. A vertex *v* of an ideal (possibly punctured) polygon is said to be *decorated* if a horoball, based at *v* is added. For $n \ge 3$, a *decorated ideal n-gon*, denoted by $\widehat{\Pi}_n^{\diamond}$, is an ideal polygon, all of whose vertices are decorated with pairwise disjoint horoballs. Similarly, for $n \ge 2$, a *decorated ideal once-punctured n-gon*, denoted by $\widehat{\Pi}_n^{\times}$, is a once-punctured ideal *n*-gon, all of whose vertices are decorated by $\widehat{\Pi}_n^{\times}$, is a once-punctured ideal *n*-gon, all of a decorated with pairwise disjoint horoballs. See Fig.11 for a decorated triangle and a decorated once-punctured bigon.

On these decorated polygons we consider two types of arcs.

An *arc* on a hyperbolic polygon Π , is an embedding α of a closed interval $I \subset \mathbb{R}$ into Π . There are two possibilities depending on the nature of the interval:

- 1. I = [a, b]: In this case, the arc α is finite. We consider those finite arcs that verify: $\alpha(a), \alpha(b) \in \partial \Pi$ and $\alpha(I) \cap S = \{\alpha(a), \alpha(b)\}.$
- 2. $I = [a, \infty)$: These are embeddings of hyperbolic geodesic rays in the interior of the polygon such that $\alpha(a) \in \partial \Pi$. The infinite end converges to a vertex of the polygon.

An arc α of a polygon Π with non-empty boundary is called *non-trivial* if each connected component of $\Pi \setminus \{\alpha\}$ has at least one decorated vertex.

Let \mathscr{A} be the set of all non-trivial arcs of the two types above. The *arc complex* of a decorated hyperbolic is a simplicial complex $\mathscr{A}(\Pi)$ whose 0-simplices are given by the isotopy classes of arcs in \mathscr{A} fixing the boundary and decorated vertices, and for $k \ge 1$, every k-simplex is given by a (k + 1)-tuple of pairwise disjoint and distinct isotopy classes.

Now we establish a link between these decorated polygons and Euclidean polygons with bicolourings. We start with the polygon \mathcal{P}_{2n} with $n \ge 2$ and consider the alternate R - B bicolouring of its vertices. To every decorated polygon $\widehat{\Pi}_n^{\diamond}$, one can associate the polygon \mathcal{P}_{2n} with an alternate B - R colouring of its vertices, in the following way:

- a decorated vertex of $\widehat{\Pi_n^{\diamond}}$ corresponds to a red vertex of \mathcal{P}_{2n} ,
- an edge of $\widehat{\Pi_n^{\diamond}}$ corresponds to a blue vertex of \mathcal{P}_{2n} ,

such that one *R*-vertex and one *B*-vertex are consecutive in \mathcal{P}_{2n} if and only if the corresponding edge and decorated vertex of $\widehat{\Pi_n^{\diamond}}$ are consecutive. See Fig.(12). Again, we have the bijection:

{Isotopy classes of edge-to-edge arcs of
$$\widehat{\Pi_n^\diamond}$$
} \leftrightarrow { $B - B$ diagonals}

 $\left\{ \text{Isotopy classes of edge-to-vertex arcs of } \widehat{\Pi_n^{\diamond}} \right\} \leftrightarrow \{B - R \text{ diagonals} \}$

So the arc complex $\mathcal{A}(\widehat{\Pi_n^{\diamond}})$ is isomorphic to the subcomplex $\mathcal{Y}(\mathcal{P}_{2n})$ of $\mathcal{A}(\mathcal{P}_{2n})$. By starting



Figure 12: From decorated ideal *n*-gon to Euclidean \mathcal{P}_{2n} with alternate bicolouring

with a punctured polygon $\mathcal{P}_{2n}^{\times}$ with an alternate bicolouring of its vertices and by using the same argument as above, we get that the the arc complex $\mathcal{A}(\widehat{\Pi_n^{\times}})$ is isomorphic to the subcomplex $\mathcal{Y}(\mathcal{P}_{2n}^{\times})$ of $\mathcal{A}(\mathcal{P}_{2n}^{\times})$. From Theorem 3.6 we get that

Corollary 4.1. For $n \ge 4$, the arc complex $\mathcal{A}(\widehat{\Pi_n^\diamond})$ of a decorated ideal polygon $\widehat{\Pi_n^\diamond}$ is *PL*-homeomorphic to a closed ball of dimension 2n - 4.

Finally, from Theorem 3.11, we get that

Corollary 4.2. For $n \ge 2$, the arc complex $\mathcal{A}(\widehat{\Pi_n^{\times}})$ of a decorated once-punctured polygon $\widehat{\Pi_n^{\times}}$ is *PL*-homeomorphic to a closed ball of dimension 2n - 2.

References

- [1] Gopal Danaraj and Victor Klee. Shellings of spheres and polytopes. *Duke Mathematical Journal*, 41(2):443 451, 1974.
- [2] Jeffrey Danciger, François Guéritaud, and Fanny Kassel. Margulis spacetimes via the arc complex. *Inventiones mathematicae*, 204(1):133–193, April 2016.
- [3] Grégoire Dupont and Frédéric Palesi. Quasi-cluster algebras from non-orientable surfaces. *Journal of Algebraic Combinatorics*, 42(2):429–472, 2015.
- [4] Sergey Fomin, Michael Shapiro, and Dylan Thurston. Cluster algebras and triangulated surfaces. Part I: Cluster complexes. Acta Mathematica, 201(1):83 – 146, 2008.
- [5] Sergey Fomin and Andrei Zelevinsky. Cluster algebras ii: Finite type classification. *Inventiones mathematicae*, 154(1):63–121, 2003.
- [6] William M. Goldman, François Labourie, and Gregory Margulis. Proper affine actions and geodesic flows of hyperbolic surfaces. *Annals of Mathematics*, 170(3):1051–1083, November 2009.
- [7] John L. Harer. The virtual cohomological dimension of the mapping class group of an orientable surface. *Inventiones Mathematicae*, 84(1):157–176, February 1986.
- [8] Carl W. Lee. The associahedron and triangulations of the n-gon. *European Journal of Com*binatorics, 10(6):551–560, 1989.
- [9] Jean-Louis Loday. Realization of the stasheff polytope. *Archiv der Mathematik*, 83(3):267–278, 2004.
- [10] Pallavi Panda. Topology of arc complexes of decorated hyperbolic polygons, 2023.
- [11] Hugo Parlier and Lionel Pournin. Flip-graph moduli spaces of filling surfaces. J. Eur. Math. Soc., 19(9):2697–2737, 2017.
- [12] Hugo Parlier and Lionel Pournin. Modular flip-graphs of one-holed surfaces. *European Journal of Combinatorics*, 67:158–173, 2018.
- [13] Hugo Parlier and Lionel Pournin. Once punctured disks, non-convex polygons, and pointihedra. *Annals of Combinatorics*, 22(3):619–640, 2018.
- [14] R. C. Penner. The decorated Teichmüller space of punctured surfaces. Communications in Mathematical Physics, 113(2):299 – 339, 1987.
- [15] R. C. Penner. Decorated teichmüller theory of bordered surfaces. Communications in Analysis and Geometry, 12:793–820, 2002.

- [16] R. C. Penner. The structure and singularities of quotient arc complexes. *Journal of Topology*, 1(3):527–550, 05 2008.
- [17] R. C. Penner. Decorated Teichmüller Theory. The QGM master class series. European Mathematical Society, Zürich, Switzerland, 2012. OCLC: ocn773019621.
- [18] Lionel Pournin. The diameter of associahedra. Advances in Mathematics, 259:13–42, 2014.
- [19] J. Scott Provan and Louis J. Billera. Decompositions of simplicial complexes related to diameters of convex polyhedra. *Mathematics of Operations Research*, 5(4):576–594, 1980.
- [20] Daniel D. Sleator, Robert E. Tarjan, and William P. Thurston. Rotation distance, triangulations, and hyperbolic geometry. J. Amer. Math. Soc., 1:647–681, 1988.
- [21] James Dillon Stasheff. Homotopy associativity of h-spaces. i. Transactions of the American Mathematical Society, 108(2):275–292, 1963.
- [22] Dov Tamari. Monoïdes préordonnés et chaînes de malcev. Bulletin de la Société mathématique de France, 82:53–96, 1954.
- [23] William P. Thurston. Minimal stretch maps between hyperbolic surfaces. arXiv:math/9801039, 1998. arXiv: math/9801039.
- [24] Jon Wilson. Shellability and sphericity of finite quasi-arc complexes. Discrete & Computational Geometry, 59(3):680–706, 2018.