Asymptotic properties of stochastic partial differential equations in the sublinear regime

Le Chen[∗] and Panqiu Xia†

Department of Mathematics and Statistics Auburn University

Tuesday 13th June, 2023

Abstract

In this paper, we investigate stochastic heat equation with sublinear diffusion coefficients. By assuming certain concavity of the diffusion coefficient, we establish non-trivial moment upper bounds and almost sure spatial asymptotic properties for the solutions. These results shed light on the *smoothing intermittency effect* under *weak diffusion* (i.e., sublinear growth) previously observed by Zeldovich *et al.* [\[Zel+87\]](#page-49-0). The sample-path spatial asymptotics obtained in this paper partially bridge a gap in earlier works of Conus *et al.* [\[CJK13;](#page-46-0) [Con+13\]](#page-46-1), which focused on two extreme scenarios: a linear diffusion coefficient and a bounded diffusion coefficient. Our approach is highly robust and applicable to a variety of stochastic partial differential equations, including the one-dimensional stochastic wave equation and the stochastic fractional diffusion equations.

Keywords: Stochastic partial differential equation, sublinear growth, asymptotic concavity, moment bounds, intermittency, spatial asymptotics.

AMS 2010 subject classification. Primary. 60H15; Secondary. 35R60.

Contents

[∗]Research is partially supported by *Mathematics and Physical Sciences-Collaboration Grants for Mathematicians* from *Simons Foundation* (Award Number: 959981). Email: [le.chen@auburn.edu,](mailto:le.chen@auburn.edu)

†Email: pqxia@auburn.edu

1 Introduction

In this paper, we investigate the *stochastic partial differential equations* (SPDEs) in the sublinear regime. More precisely, we focus on the asymptotic behavior of the following *stochastic heat equation* (SHE)

$$
\begin{cases} \frac{\partial}{\partial t}u(t,x) = \frac{1}{2}\Delta u(t,x) + \rho(u(t,x))\dot{W}(t,x), \quad t > 0, \ x \in \mathbb{R}^d, \\ u(0,\cdot) = \mu, \end{cases}
$$
\n(1.1)

where the diffusion coefficient $\rho(\cdot)$ is assumed to be locally bounded and exhibits sublinear growth at infinity. Previous studies have extensively examined the case when the diffusion coefficient $\rho(\cdot)$ has linear growth at infinity, which results in an intermittent solution. In particular, a solution is said to be *intermittent* if the *moment Lyapunov exponents* $\overline{\lambda}_p$ and $\underline{\lambda}_p$ of the solution, defined by

$$
\overline{\lambda}_p := \frac{1}{p} \limsup_{t \to \infty} \log \mathbb{E} \left[|u(t, x)|^p \right] \quad \text{and} \quad \underline{\lambda}_p := \frac{1}{p} \liminf_{t \to \infty} \log \mathbb{E} \left[|u(t, x)|^p \right] \quad (p \ge 2),
$$

satisfy the property that $\lambda_2 > 0$. The literature on this topic is extensive, and interested readers may consult [\[CM94;](#page-45-0) [FK09](#page-47-0); [CD15b](#page-45-1); [Che15](#page-46-2); [KKX17](#page-47-1)] and references therein. Zeldovich *et al.* [\[ZRS90,](#page-49-1) Chapter 9] have observed that intermittency is a universal phenomenon that occurs irrespective of the underlying properties of the instability in a random medium, as long as the random field is of multiplicative type. However, they have also noted in [\[Zel+87](#page-49-0)] (see also [\[ZRS90](#page-49-1), Section 8.9]) that "*smoothing intermittency*", where the high maxima of the solution have smaller growth, should be expected in the presence of "*weak diffusion*"—when $\rho(\cdot)$ exhibits sublinear growth at infinity. The authors substantiated their statement by highlighting the power growth of the moments when $\rho(\cdot)$ is bounded, or more specifically, when

$$
\rho(u) = \frac{u}{1+u},\tag{1.2}
$$

and when the noise is white in time and space-independent. Inspired by their previous work, the objective of this paper is to conduct a comprehensive analysis of the smoothing intermittency property for some more general weak diffusion cases and demonstrate how the moment growth rate of the solution is influenced by that of $\rho(\cdot)$.

The examination of SPDEs in the sublinear regime is also motivated by the requirement for more realistic biological population models. As highlighted by König in the Appendix of $[K\ddot{o}n16]$, the *parabolic Anderson model* (PAM) (i.e., $\rho(u) = \lambda u$) utilized in population dynamics, leads to excessive branching and killing rates that are not reflective of real-world scenarios as it lacks any form of birth or death control. In contrast, introducing a sublinear growth for *ρ* provides an avenue for developing models that may address this issue and provide a more accurate depiction of the dynamics of biological populations.

Both the PAM and the SHE with additive noise, or simply the *additive SHE*, (i.e., $\rho \equiv 1$, or more generally, the case when ρ is bounded) have been extensively studied in the literature. They represent two extreme cases where rich properties have been previously derived. For instance, when $d = 1$, the noise is space-time white noise, and the initial condition μ is constant, Conus *et al.* [\[CJK13\]](#page-46-0) showed that

$$
\begin{cases}\n\sup_{|x| \le R} u(t, x) \approx [\log R]^{1/2} & \text{SHE with bounded } \rho \text{ (Theorem 1.2 ibid.)}; \\
\log \sup_{|x| \le R} u(t, x) \approx [\log R]^{2/3} & \text{PAM (Theorem 1.3 ibid.).}\n\end{cases}
$$
\n(1.3)

Here and in this paper, we use the notation $f(x) \leq g(x)$ to denote that there exists a nonrandom constant $C > 0$ such that $\liminf_{x \to \infty} f(x)/g(x) \leq C$; the notation $f(x) \geq g(x)$ is defined analogously. We also write $f(x) \approx g(x)$ if both $f(x) \lesssim g(x)$ and $f(x) \gtrsim g(x)$. The distinct behaviors exhibited in (1.3) naturally raise questions regarding the dynamics when ρ is neither bounded nor linear, but rather demonstrates sublinear growth, thereby providing a potential interpolation between these two extreme scenarios. To address the scenario of sublinear growth, a significantly different method must be developed, which constitutes the primary contribution of this paper.

Sublinear examples of $\rho(u)$ (for $u > 0$) typically include

$$
\rho(u) = \frac{u}{(r+u)^{1-\alpha}}, \quad \alpha \in [0,1) \text{ and } r \ge 0;
$$
\n(1.4)

$$
\rho(u) = u^{\alpha} \left[\log(e + u^2) \right]^{-\beta}, \quad \text{with} \quad \begin{cases} \alpha = 0 \text{ and } \beta < 0 & \text{Case (i)}, \\ \alpha \in (0, 1) \text{ and } \beta \in \mathbb{R} & \text{Case (ii)}, \\ \alpha = 1 \text{ and } \beta > 0 & \text{Case (iii)}; \end{cases}
$$
(1.5)

and

$$
\rho(u) = u \exp\left(-\beta \left(\log\left(\log(e+u^2)\right)\right)^{\kappa}\right) \quad \text{with } \kappa > 0 \text{ and } \beta > 0. \tag{1.6}
$$

Here, *e* denotes the *Euler constant*. In particular, letting $r = 1$ and $\alpha = 0$ in [\(1.4\)](#page-2-1), we reduce to the case of bounded ρ in [\(1.2\)](#page-1-1). The following important example is a special case of [\(1.4\)](#page-2-1) when $r = 0$:

$$
\rho(u) = u^{\alpha}, \quad \alpha \in (0, 1). \tag{1.7}
$$

It should be noted that SHEs with the diffusion coefficient given in [\(1.7\)](#page-2-2) are closely related to superprocesses; see, e.g., $[Draw93; Eth00; Per02]$ $[Draw93; Eth00; Per02]$ $[Draw93; Eth00; Per02]$ $[Draw93; Eth00; Per02]$. Among the three cases in (1.5) , cases (i) and (iii) are more interesting, as they are logarithmic perturbations of the additive SHE and the PAM, respectively. When $\kappa = 1$, ρ in [\(1.6\)](#page-2-4) reduces to case (iii) of [\(1.5\)](#page-2-3). Figure [1.1](#page-3-0) below illustrates the hierarchy of these examples. One would expect properties, such as the moment growth rates and spatial asymptotics, to transition from those of the additive SHE to those of the PAM. For example, for the case [\(1.4\)](#page-2-1), one would expect a polynomial growth in *t* of moments, while the case [\(1.6\)](#page-2-4) should lead to some exponential growth, but with a sublinear dependence on *t* in the exponent. Note that examples in (1.4) – (1.6) are all for the case when $u \ge 0$. If the solution is signed, one needs to replace u by $|u|$.

Figure 1.1: Examples of ρ given in (1.4) – (1.6) .

Let us proceed to set up the problem. The noise \dot{W} in SHE (1.1) is a centered Gaussian noise that is white in time and homogeneously colored in space. Its covariance structure is given by

$$
\mathbb{E}\left[\dot{W}(t,x)\dot{W}(s,y)\right]=\delta_0(t-s)f(x-y).
$$

Here, δ_0 denotes the Dirac delta measure at 0 and f is the correlation (generalized) function on \mathbb{R}^d , which satisfies the following hypothesis:

Hypothesis 1.1. The correlation function $f : \mathbb{R}^d \to \mathbb{R}$ is a nonnegative and nonnegative-definite (generalized) function that is not identically zero, such that the following *Dalang's condition* [\[Dal99](#page-46-3)] is satisfied:

$$
\int_{\mathbb{R}^d} \frac{\widehat{f}(\mathrm{d}\xi)}{1+|\xi|^2} < \infty,\tag{1.8}
$$

where $\hat{f}(\xi) = \int_{\mathbb{R}^d} f(x)e^{-ix\cdot\xi} dx$ is the Fourier transform of *f*.

By using the Fourier transform and the Plancherel theorem, it is easy to verify that Dalang's condition (1.8) is equivalent to the following condition (see, e.g., [\[Dal99](#page-46-3), Formula (20)]):

$$
h(t) := \int_0^t ds \iint_{\mathbb{R}^{2d}} dy dy' \, p_s(y) p_s(y') f(y - y') = \frac{1}{2} \int_0^{2t} ds \int_{\mathbb{R}^d} dz \, p_s(z) f(z) < \infty, \quad \forall t > 0,\tag{1.9}
$$

where $p_t(x) := (2\pi)^{-d/2} \exp(-|x|^2/(2t))$ refers to the heat kernel. The function *h* plays an essential role in our main result—Theorem [1.4](#page-4-0) below, whose asymptotic behaviors both at infinity and around zero will be postponed to Appendix [A.](#page-40-0) Note that the nonnegativity assumption of *f* in Hypothesis [1.1](#page-3-2) ensures that the function $h(\cdot)$ is an increasing function on \mathbb{R}_+ .

To facilitate the analysis, we introduce the following hypothesis on *ρ*, which covers all examples given in (1.4) – (1.6) with all *u* replaced |*u*|:

Hypothesis 1.2. The diffusion coefficient $\rho : \mathbb{R} \to \mathbb{R}$ satisfies the following properties:

(i) ρ is a locally bounded function.

(ii)
$$
\lim_{x \to \pm \infty} \frac{\rho(x)}{x} = 0;
$$

ρ(*x*)

(iii) There exists a constant $M_0 \geq 0$, such that the function $|\rho|$ is concave separately on $(-\infty, -M_0]$ and $[M_0, \infty)$.

The initial condition μ in [\(1.1\)](#page-1-2) also plays an active role in shaping the properties of the solution, for which we make the following assumption:

Hypothesis [1](#page-4-1).3. The initial condition μ for SHE [\(1.1\)](#page-1-2) is a signed Borel measure¹ on \mathbb{R}^d such that

(i) For any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, it holds that, in the sense of Lebesgue integral,

$$
\mathcal{J}_0(t,x) := \int_{\mathbb{R}^d} \mu(\mathrm{d}y) \, p_t(x-y) \in (-\infty, \infty), \quad \text{or equivalently,}
$$
\n
$$
\mathcal{J}_+(t,x) := \int_{\mathbb{R}^d} |\mu|(\mathrm{d}y) \, p_t(x-y) < \infty,
$$
\n
$$
(1.10)
$$

where $\mu = \mu_+ - \mu_-$ is the Hahn decomposition of μ and $|\mu| = \mu_+ + \mu_-$;

(ii) Moreover, if $d \geq 2$, then for any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$,

$$
\mathcal{J}_1(t,x) := \int_0^t \mathrm{d}s \iint_{\mathbb{R}^{2d}} \mathrm{d}y \mathrm{d}y' \, p_{t-s}(x-y) p_{t-s}(x-y') f(y-y') \mathcal{J}_0^2(s,y) < \infty. \tag{1.11}
$$

As usual, the solution to (1.1) is understood as the mild solution to the corresponding stochastic integral equation:

$$
u(t,x) = \mathcal{J}_0(t,x) + \int_0^t \int_{\mathbb{R}^d} p_{t-s}(x-y) \rho(u(s,y)) W(ds,dy), \quad t > 0, x \in \mathbb{R}^d,
$$
 (1.12)

where the stochastic integral is understood in the sense of Walsh [\[Wal86](#page-48-1); [Dal99](#page-46-3)]. Now we are ready to state our main result as follows:

Theorem 1.4. *Under Hypotheses* [1.1,](#page-3-2) [1.2,](#page-4-2) and [1.3,](#page-4-3) *let* $u(t, x)$ *be a solution to SHE* [\(1.1\)](#page-1-2)*. Then, for all* $(t, x, p) \in \mathbb{R}_+ \times \mathbb{R}^d \times [2, \infty)$ *, it holds that*

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 2(2\pi)^d \left(h(t)^{-1} \mathcal{J}_1(t,x) + 4K_M^2 p h(t) + F^{-1}(2p h(t)) \right). \tag{1.13}
$$

¹We follow the convention that when μ is absolutely continuous with the Lebesgue measure, it is identified as its Lebesgue density.

In [\(1.13\)](#page-4-4), $\mathcal{J}_0(\cdot, \circ)$, $\mathcal{J}_1(\cdot, \circ)$ *and* $h(\cdot)$ *are defined in* [\(1.10\)](#page-4-5), [\(1.11\)](#page-4-6) *and* [\(1.9\)](#page-3-3) *above, respectively. The constant K^M is defined as*

$$
K_M := \sup_{x \in (-M,M)} |\rho(x)|,\tag{1.14}
$$

with M given in part (iii) of Lemma [3.3](#page-12-0) below. The function $F(\cdot)$ *and its inverse* $F^{-1}(\cdot)$ *, both determined by ρ, are defined in* [\(3.7\)](#page-13-0) *and* [\(3.8\)](#page-13-1)*, respectively. In particular, the following statements hold:*

- *(i) If* $d = 1$ *, one can replace* $4h(t)^{-1} \mathcal{J}_1(t, x)$ *in* [\(1.13\)](#page-4-4) *by* $2^{7/2} \pi \mathcal{J}_+^2(t/2, x)$ *(see* [\(1.10\)](#page-4-5) *for notation)*;
- *(ii)* If $|\rho(\cdot)|$ *is concave separately on* \mathbb{R}_+ *and* \mathbb{R}_- *, then one can take* $M = K_M = 0$ *in* [\(1.13\)](#page-4-4) *to obtain*

$$
||u(t,x)||_p^2 \le 2 \mathcal{J}_0^2(t,x) + 2(2\pi)^d \left(h(t)^{-1} \mathcal{J}_1(t,x) + F^{-1}(2p \, h(t)) \right); \tag{1.15}
$$

(iii) If $\rho(\cdot)$ *is not identically* 0 *on* $(-\infty, -M_0] \cup [M_0, \infty)$ *, then there exists a constant* $C > 0$ *such that for all* $(t, x, p) \in [1, \infty) \times \mathbb{R}^d \times [2, \infty),^2$ $(t, x, p) \in [1, \infty) \times \mathbb{R}^d \times [2, \infty),^2$ $(t, x, p) \in [1, \infty) \times \mathbb{R}^d \times [2, \infty),^2$

$$
||u(t,x)||_p^2 \le 2 \mathcal{J}_0^2(t,x) + C\left(h(t)^{-1}\mathcal{J}_1(t,x) + F^{-1}(Cp\,h(t))\right). \tag{1.16}
$$

Moreover, if the initial condition is bounded, i.e., $u_0 \in L^{\infty}(\mathbb{R}^d)$ *, the moment bound in* [\(1.16\)](#page-5-1) *can be simplified as follows:*

$$
||u(t,x)||_p^2 \le C_* F^{-1}(C_* p h(t)) \quad \text{for some } C_* > 0. \tag{1.17}
$$

The general moment bounds in the above theorem demonstrate how different components of the SPDE affect the moment growth of the solution (see Figure [1.2](#page-5-2) below), which is a concrete manifestation of the statement in Zeldovich *et al* [\[Zel+87](#page-49-0)] (see also [\[ZRS90,](#page-49-1) Section 8.9]) that *the behavior of nonlinear solutions depends radically on the time behavior of the potential and on the form of the nonlinearity.* When considering specific scenarios, in order to utilize the above theorem, we have to study the corresponding $F^{-1}(\cdot)$ as detailed in Section [4,](#page-24-0) and $h(t)$ as outlined in Section [A.](#page-40-0) Following these analyses, we summarize the results and derive the specific moment bounds in Section [6](#page-36-0) below.

Figure 1.2: Structure of the moment bounds in (1.15) or (1.16) .

²Here *t* can be relaxed to $t > 0$, but the constant *C* in [\(1.16\)](#page-5-1) will depend on *t* when *t* is close to 0

As some applications of the above moment bounds, we obtain the tail probability of the solution, sample-path Hölder regularity, and further establish some estimates for the sample-path asymptotics in the spatial variable, as shown in (1.21) below. The sample-path spatial asymptotics have been previously studied by Conus *et al.* [\[CJK13](#page-46-0); [Con+13\]](#page-46-1) in two extreme cases, namely, the case when ρ is linear and that when ρ is bounded. The result presented in Theorem [1.8](#page-6-2) below serves as an initial attempt to bridge the gap between these two extreme cases by allowing ρ to have sublinear growth. Here, we have to admit that only the asymptotic upper bounds have been obtained, while the more challenging lower bounds will be left for future investigation.

Theorem 1.5 (Tail probability)**.** *Assume that both Hypotheses [1.1](#page-3-2) and [1.2](#page-4-2) hold. Let u*(*t, x*) *be a solution to SHE* [\(1.1\)](#page-1-2) *with the initial condition* $\mu \equiv 1$ *. Then for all* $t \geq 1$, $x \in \mathbb{R}^d$ *and* $z \geq L_t$,

$$
\mathbb{P}(|u(t,x)| \ge z) \le \exp\left(-(C_*h(t))^{-1}F\left(z^2/(C_*e^2)\right)\right),\tag{1.18}
$$

where the constant $C_* > 0$ *is the same as that in* [\(1.17\)](#page-5-4)*,* $F(\cdot)$ *and* $F^{-1}(\cdot)$ *are defined Definition* 3.6*,*

$$
L_t := e\sqrt{2C_*M^2 + C_*F^{-1}(2C_*h(t))},\tag{1.19}
$$

and the constant M in [\(1.19\)](#page-6-4) *is the same as those in Theorem [1.4.](#page-4-0)*

To state the next two results, we need to introduce the following *improved Dalang condition* for the spatial correlation function *f*:

Hypothesis 1.6. The correlation function $f : \mathbb{R}^d \to \mathbb{R}$ is a nonnegative and nonnegative-definite (generalized) function that is not identically zero such that,

$$
\int_{\mathbb{R}^d} \frac{\hat{f}(\mathrm{d}\xi)}{(1+|\xi|^2)^{1-\eta}} < \infty, \quad \text{for some } \eta \in (0,1). \tag{1.20}
$$

Theorem 1.7 (Hölder regularity). *Under parts (i) and (ii) of Hypothesis [1.2,](#page-4-2) part (i) of Hypothesis* [1.3,](#page-4-3) and Hypothesis [1.6,](#page-6-5) the solution $u(t, x)$ to [\(1.1\)](#page-1-2) has a version which is a.s. η_1 –Hölder *continuous in time and* η_2 –*Hölder continuous in space on* $(0, \infty) \times \mathbb{R}^d$ *for all* $\eta_1 \in (0, \eta/2)$ *and* $\eta_2 \in (0, \eta)$ *, where* η *is given in* [\(1.20\)](#page-6-6)*.*

Theorem 1.8 (Spatial asymptotics)**.** *Assume that both Hypotheses [1.2](#page-4-2) and [1.6](#page-6-5) hold. Suppose that* $\rho(\cdot)$ *is not identically zero on* $(-\infty, -M_0] \cup [M_0, \infty)$ *. Let* $u(t, x)$ *be a solution to SHE* [\(1.1\)](#page-1-2) *with the constant initial condition* $\mu \equiv 1$ *. Then, there exists a positive constant C such that for all* $t > 1$ *,*

$$
\sup_{|x| \le R} u(t, x) \lesssim \sqrt{F^{-1}\Big(C\ h(t) \log R\Big)}, \quad a.s., \text{ as } R \to \infty,
$$
\n(1.21)

where $F^{-1}(\cdot)$ *is defined in* [\(3.8\)](#page-13-1)*.*

The paper is organized as follows. We begin by expanding the discussions of the main results in Section [2,](#page-7-0) including a presentation of the proof strategy of Theorem [1.4.](#page-4-0) We then proceed to prove the main results in Section [3.](#page-11-0) In Section [3.1,](#page-11-1) we introduce several technical lemmas, and their proofs can be found in Section [3.6.](#page-20-0) The proofs of Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) [1.7](#page-6-1) and [1.8](#page-6-2) are provided in Sections [3.2,](#page-14-0) [3.3,](#page-16-0) [3.4](#page-17-0) and [3.5,](#page-18-0) respectively. In Section [4,](#page-24-0) we derive $F(\cdot)$ and $F^{-1}(\cdot)$ for examples in (1.4) (1.4) (1.4) – (1.6) . Section [5](#page-27-0) explores potential generalizations of Theorem 1.4 to other SPDEs. In Section [6,](#page-36-0) we present several concrete examples to illustrate, for example, the property transitions from the additive SHE to the PAM. Lastly, in Appendix [A,](#page-40-0) we present asymptotic formulas for the function $h(\cdot)$ in various cases, and in Appendix [B,](#page-44-0) we provide a supplementary proof for results in Section [6.](#page-36-0)

2 Remarks and strategy of the proof

In the following, we make a few comments/discussions on our assumptions and results, and present the strategy of the proof of Theorem [1.4.](#page-4-0)

Assumptions on initial conditions Here are some comments on the assumptions made for the initial conditions in Hypothesis [1.3:](#page-4-3) **(i)** Following [\[CD15b\]](#page-45-1), we call the initial condition satisfying inequality [\(1.10\)](#page-4-5) the *rough initial condition* for SHE [\(1.1\)](#page-1-2). In particular, the Dirac delta measure is a special case; see also $[Con+14]$. Note that the Dirac delta measure plays an important role in studying the asymptotic properties of the PAM on \mathbb{R} ; see [\[ACQ11;](#page-45-2) [Cor12](#page-46-5)]. As an easy exercise, condition (1.10) is equivalent to

$$
-\infty < \int_{\mathbb{R}^d} e^{-a|x|^2} \mu(\mathrm{d}x) < \infty, \quad \text{for all } a > 0.
$$

(ii) Condition [\(1.11\)](#page-4-6) is only a technical assumption. We believe that this assumption can be removed. This will be left for future investigation. If $\mathcal{J}_0(s, y)^2$ in [\(1.11\)](#page-4-6) is replaced by $\mathcal{J}_0(s, y)\mathcal{J}_0(s, y')$, then due to [\[CK19,](#page-46-6) Lemma 2.7], the integral is finite under Dalang's condition [\(1.8\)](#page-3-1) for all rough initial conditions. This extra condition comes from the application of the inequality $\mathcal{J}_0(s, y)\mathcal{J}_0(s, y') \leq$ 1 $\frac{1}{2}$ $(\mathcal{J}_0^2(s, y) + \mathcal{J}_0^2(s, y'))$ in the proof of Theorem [1.4.](#page-4-0) (iii) In case $d = 1$, condition [\(1.11\)](#page-4-6) is auto-matically satisfied by Lemma [3.9](#page-13-3) below. On the other hand, for $d \geq 2$, condition [\(1.11\)](#page-4-6) excludes the Dirac delta initial condition. This can be seen by setting $f(\cdot) \equiv 1$ (the space-independent noise), then the integral in [\(1.11\)](#page-4-6) with delta initial condition reduces to $\int_0^t s^{-d/2} ds = \infty$. Instead, condition [\(1.11\)](#page-4-6) holds when $\mu(dx) = |x|^{-\ell} dx$ for any $\ell \in (0,1)$. This is due to the bound in [\(3.12\)](#page-13-4) and the fact that for such initial data, $\mathcal{J}_0(t,x) \lesssim t^{-\ell/2}$; see [\[CE22b\]](#page-45-3). Roughly speaking, the Dirac delta measure corresponds to the case when $\ell = d$. Removing part (ii) of Hypothesis [1.3](#page-4-3) will be left for future investigation.

Existence and uniqueness: the global Lipschitz case If ρ is globally Lipschitz continuous such as those in (1.4) – (1.6) , the existence and uniqueness of a random field solution to SHE (1.1) under rough initial conditions is guaranteed (see [\[CD15c](#page-45-4); [CK19](#page-46-6)]). Moreover, since $\rho(0) = 0$ in all examples (1.4) – (1.6) , if the initial condition μ is a nonnegative measure and is not vanishing, then the solution $u(t, x)$ is strictly positive almost surely for all $t > 0$ and $x \in \mathbb{R}^d$ thanks to the *sample-path comparison principle* (see [\[Mue91](#page-48-2); [Shi94;](#page-48-3) [CH19](#page-46-7)]) One can further apply the *stochastic (moment) comparison principle* (see [\[JKM17;](#page-47-5) [CK20\]](#page-46-8)) to bound the moments from above by those of the *parabolic Anderson model* (see [\[CM94\]](#page-45-0)), which corresponds to the case when $\rho(u) = \lambda u$. However, moment bounds obtained in this way are generally too rough and do not exhibit the "smoothing intermittency" effect as observed in Zeldovich *et al* [\[ZRS90\]](#page-49-1).

Existence and uniqueness: the non-Lipschitz case In case when *ρ* is not Lipschitz continuous, such as the example in [\(1.7\)](#page-2-2), Theorem [1.4](#page-4-0) can be used to provide some a priori moment estimates. It is known that establishing the uniqueness of solutions to SHE [\(1.1\)](#page-1-2) in this case is a challenging problem. However, it is commonly known that the existence of solutions to (1.1) can usually be established using some common strategy. One first mollifies the non-Lipschitz *ρ* function into a sequence of globally Lipschitz continuous functions, based on which a sequence of random fields are constructed. Under some mild conditions, it is possible to prove that these random fields are jointly Hölder continuous with a uniform constant. By combining the Hölder continuity with a priori estimates for solutions, such as those given in Theorem [1.4,](#page-4-0) one can apply

the *Kolmogorov-Chentsov criterion* (see, e.g., [\[Kal02,](#page-47-6) Corollary 16.9]) to conclude that there is a convergent subsequence of the aforementioned random fields, whose limit gives rise to one solution to [\(1.1\)](#page-1-2). Since the existence and uniqueness of solution is not the focus of the paper. We will not pursue this direction here. Finally, we would like to point out that the strong or even the weak well-posedness problem of SHEs with non-Lipschitz coefficients is highly involved, and only a few results are known. One may consult [\[Myt98;](#page-48-4) [MPS06;](#page-48-5) [MP11\]](#page-48-6) for the sublinear case, and the recent works in [\[DKZ19;](#page-46-9) [Sal22;](#page-48-7) [CH22](#page-46-10)] for the superlinear growth case.

Generality versus sharpness The moment bounds obtained in Theorem [1.4](#page-4-0) strike a balance between their level of generality and their sharpness. Obtaining sharp moment asymptotics in general can be extremely challenging and is typically only possible in some specific settings. For instance, in the case when $d = 1$, \hat{W} is the space-time white noise, and $\rho(u) = \sqrt{u}$ (the super-Brownian motion) with $u(0, x) \equiv 1$, the second author and his collaborators [\[Hu+23\]](#page-47-7) recently find the following exact moment asymptotics:

$$
\mathbb{E}\left[u(t,x)^p\right] \asymp K^p p!\left(1+t^{(p-1)/2}\right), \quad \text{as } t\to\infty,
$$

which is valid for any positive integer p and $x \in \mathbb{R}$. This asymptotic should be compared with the upper bounds obtained by Theorem [1.4:](#page-4-0)

$$
\mathbb{E}\left[u(t,x)^p\right]\leq K^pp!\left(1+t^{p/2}\right).
$$

The difference between these bounds is a factor of \sqrt{t} , showing that the bound in Theorem [1.4](#page-4-0) is not sharp. Although we only have this case showing that our moment bounds are not sharp, we believe that this is generally the case. Nonetheless, the method used in the proof of Theorem [1.4](#page-4-0) are quite robust and can be easily extended to more general settings and a broad class of stochastic partial differential equations (SPDEs), including the *stochastic wave equation* (SWE) (see Section [5.2\)](#page-31-0), and SPDEs with fractional differential operators (see Section [5.1\)](#page-27-1). Moreover, despite being sub-optimal, the moment bounds obtained in Theorem [1.4](#page-4-0) are sufficient to provide a quantitative description of the "smoothing intermittency" phenomenon introduced by Zeldovich *et al.* [\[ZRS90](#page-49-1)].

Hölder regularity The Hölder continuity for solutions to SHEs has been extensively investigated in the literature. Notably, Konno and Shiga [\[KS88\]](#page-47-8) and Reimers [\[Rei89\]](#page-48-8) examined the Hölder continuity for super-Brownian motions and Fleming–Viot processes driven by space-time white noise with function-valued initial conditions. Sanz-Solé and Sarrá [\[SS02](#page-48-9)], on the other hand, examined the scenario where the noise satisfies the improved Dalang condition (Hypothesis [1.6\)](#page-6-5) and *ρ* satisfies the golobal Lipschitz condition. Recently, Chen and Daland [\[CD14](#page-45-5)] and Chen and Huang [\[CH19](#page-46-7)] generalized results in [\[SS02](#page-48-9)] by allowing rough initial conditions (part (i) of Hypothesis [1.3\)](#page-4-3). Theorem [1.7](#page-6-1) makes a slight extension of [\[CH19](#page-46-7), Theorem 1.8] by allowing the non-Lipschitz condition on ρ , where the global Lipschitz condition is replaced by "locally bounded + linear growth at infinity" of *ρ*. As far as we know, the Hölder continuity for SHEs with $\rho(u) = \sqrt{u}$ starting from the Dirac delta measure (one instance of the rough initial condition) has not been studied in the literature. For the classical initial condition, one may consult, e.g., [\[Xio13](#page-49-2), Theorem 1.4.6].

Spatial asymptotics and tail estimates Theorem [1.8](#page-6-2) provides an almost sure asymptotic upper bound for solutions to (1.1) in space. The proof of this theorem closely follows the approach presented in [\[CJK13](#page-46-0)], which relies on tail estimates (Theorem [1.5\)](#page-6-0) and the Borel–Cantelli lemma. While the moment bounds may not be particularly sharp, we find that the tail estimates are indeed sharp, at least in the case of super Brownian motion; see Proposition [6.2](#page-36-2) with $\alpha = 1/2$ and $\overline{H}u+23$, Proposition 1.4]. As a result, we believe Theorem [1.8](#page-6-2) provides a sharp bound for super-Brownian motion, specifically, for any $t > 0$ fixed,

$$
\sup_{|x| \le R} u(t, x) \asymp \sqrt{t} \log(R), \quad \text{almost surely, as } R \to \infty.
$$

In Proposition [6.2,](#page-36-2) by letting $\alpha = 0$, the spatial asymptotic upper bounds coincide with the exact asymptotics of SHEs with the additive space-time white noise as proved in [\[CJK13](#page-46-0), Theorem 1.2], and with additive spatial colored noises as proved in $[Con+13, Theorem 2.3]$. However, Theorem [1.5](#page-6-0) provides a tail estimates obtained by approximating the Legendre-type transform of *H*, as seen in [\(3.27\)](#page-17-1). This approximation is effective only when $\rho(u)$ grows at a "significantly slower" rate than *u* as $u \to \infty$, such as $\rho(u) = u^{\alpha}$ with $\alpha \in [0,1)$. However, if $\rho(u)$ is "very close" to *u* for large *u*, such as $\rho(u) = u [\log(e+u)]^{-\beta}$ with $\beta \in (0, 1/4)$, part (b) of Proposition [6.6](#page-38-0) below suggests that the results in Theorem [1.5](#page-6-0) and consequently in Theorem [1.8](#page-6-2) can be improve.

Interaction with initial conditions Due to the multiplicative nature of the noise, the initial condition interacts with the noise, playing an active role in shaping the solution. This interaction is evident when one writes out the Picard iteration of the mild solution given by (1.12) . For the PAM, this interaction leads to the following moment bound (see [\[CH19](#page-46-7), Theorem 1.7]):

$$
||u(t,x)||_p^2 \le C \mathcal{J}_0^2(t,x) \Upsilon(t), \quad \text{for all } t > 0, x \in \mathbb{R}^d, \text{ and } p \ge 2,
$$
 (2.1)

where Υ(·) is a function that represents the contribution of the driven noise. The *multiplicative interaction* of the initial conditions and the driven noise, as shown in (2.1) , naturally gives rise to the property of propagation of tall peaks in some space-time cone $\{|x| \leq \kappa t\}$, which was earlier observed in physical contexts, see, e.g., Zeldovich *et al.* [\[ZRS90](#page-49-1), Section 8.10] and later rigorously formulated and proved by Conus and Khoshnevisan [\[CK12](#page-46-11)]. Since then, additional researches (see [\[CD15b](#page-45-1); [CK19](#page-46-6); [HLN17](#page-47-9)]) has expanded upon this cone property. In essence, the cone property states that there exists a space-time cone of size κ within which the moments of the PAM grow exponentially fast, while outside of it, they decay exponentially rapidly. The precise size κ of the cone is known as the *intermittency front*.

However, in the present paper, we obtain an *additive interaction* of the initial data and the noise as shown in Theorem [1.4](#page-4-0) and Figure [1.2.](#page-5-2) This additive interaction arises from the way we solve the inequality (2.3) (see Figure [2.1\)](#page-11-2) or from the application of Lemma [3.8](#page-13-5) in general, where the coefficient *b* corresponds to the initial condition. By sending *b* to zero in (2.3) , the linear equation $x = kx^{\alpha} + b$ transitions from having one unique nonnegative solution to having two nonnegative solutions, one of which is zero. Accordingly, assuming $b = 0$, the inequality $x \leq kx^{\alpha}$ can only imply that $x \in [0, k^{1/(1-\alpha)}]$, but additional information is needed to determine the exact value of *x*. In the context of SPDEs, such additional information may be related to the uniqueness and non-uniqueness of the solution. In fact, when $\rho(u) = |u|^{\alpha}$ with $\alpha \in (0, 3/4)$ and W is the one-dimensional space-time white noise, Mueller *et al* [\[MMP14](#page-48-10)] constructed nontrivial solutions starting from zero initial condition. Therefore, the propagation of high peaks (of polynomial order in this case) will be much more subtle and will be left for future research.

If the diffusion coefficient ρ is globally Lipschitz, the moment comparison theorem may be applied, and thus the moment bounds for the "dominant PAM" can be used to determine an upper bound for the propagation to the solution to (1.1) . In case that ρ is not globally Lipschitz, it is known that solutions to [\(1.1\)](#page-1-2), assuming $\rho(u) = u^{\alpha}$ with $\alpha \in (0,1)$, is compactly supported provided the initial condition is a finite measure, as demonstrated in [\[DIP89;](#page-46-12) [MP92](#page-48-11)]. Additionally,

precise analysis on front propagation (a related but distinct property) for Fisher-KPP equations has been presented in [\[MMQ11;](#page-48-12) [MMR21](#page-48-13)]. As a result, it should be possible to obtain results about propagation of tall peaks, even for non-Lipschitz cases with certain necessary restrictions on initial conditions. We hope that this question can be resolved in the future.

Proof strategy of Theorem [1.4](#page-4-0) The proof of Theorem [1.4](#page-4-0) relies on the concavity of *ρ*. To highlight the strategy of the proof, consider the case of space-time white noise $(f = \delta_0)$ and assume that $M_0 = 0$ in Hypothesis [1.2.](#page-4-2) Using standard arguments involving Burkholder-Davis-Gundy's (see, e.g., [\[Kho14](#page-47-10), Theorem B.1]), Minkowski's and Jensen's inequalities, one can derive the following inequality:

$$
X(t,x) \le k\rho_2(X(t,x)) + b, \quad \text{with} \quad X(t,x) = \int_0^t \mathrm{d}s \int_{\mathbb{R}^d} \mathrm{d}y \, p_{t-s}^2(x-y) \, \|u(s,y)\|_p^2, \tag{2.2}
$$

where $\rho_2(\cdot)$ is a sublinear function defined on \mathbb{R}_+ , *k* and *b* are some constants depending on *p* and *t*. Since ρ_2 is sublinear, any nonnegative solution to inequality (2.2) has to lie in a compact interval, e.g., $[0, F]$. In other words, as X satisfies [\(2.2\)](#page-10-1), we get $X \leq F$; see Lemma [3.8.](#page-13-5)

To illustrate the idea of solving inequality [\(2.2\)](#page-10-1), consider the case when $\rho(u) = u^a, u \ge 0$, with $a \in [0, 1)$ fixed. In this case, $\rho_2(u) = \rho(u)$ and the inequality in [\(2.2\)](#page-10-1) becomes

 $x \leq kx^a + b$, for $x \geq 0$ with $a \in (0,1)$, $b > 0$ and $k > 0$ being fixed. (2.3)

By the concavity of the function x^a , the corresponding equation $x = kx^a + b$ has a unique positive solution, which is denoted by x^* . Hence, inequality (2.3) holds provided that $x \in [0, x^*]$, i.e., x^* is an upper bound estimate for *x*. Moreover, one can apply Newton's method for one step, properly started, to obtain an upper bound for x^* (see Figure [2.1](#page-11-2) for an intuitive display of this procedure):

$$
x^* \le k^{1/(1-a)} + b/(1-a). \tag{2.4}
$$

To handle more general cases, including those where $M_0 \neq 0$ (i.e., ρ is only asymptotic concave), the noise is not white in space, and the initial conditions are more general (not a constant), a more meticulous approach is required. This ultimately leads to the establishment of an inequality in the form of [\(2.2\)](#page-10-1), as shown in [\(3.21\)](#page-15-0) below. Lemma [3.8](#page-13-5) outlines the procedure for identifying an upper bound similar to the one depicted in Figure [2.1,](#page-11-2) which in turn yields a bound akin to [\(2.4\)](#page-10-2) for solutions to (3.21) . Overall, this constitutes the general strategy behind the proof of Theorem [1.4.](#page-4-0)

The procedure outlined in this paper involves establishing inequality [\(3.21\)](#page-15-0) first and then deriving its upper bounds using Lemma [3.8.](#page-13-5) This can be seen as a generalization of the one-variable Bihari–LaSalle inequality [\[Bih56](#page-45-6); [LaS49](#page-47-11)] to multivariate or field cases, as evidenced by [\(3.15\)](#page-14-1). However, there are some subtle differences between the two, such as the convolutional form of the time variable in our setting, as seen in [\(3.15\)](#page-14-1). While the Bihari–LaSalle inequality has been applied to moment estimates for nonlinear stochastic differential equations (SDEs), such as in [\[FZ05\]](#page-47-12), its application to nonlinear SPDEs requires careful handling of the multivariate setting. One common approach is taking the supreme norm on the spatial argument and obtaining a one-variable integral inequality. However, this method is not always applicable, especially for unbounded initial conditions or when a better understanding of how the initial condition enters the iterations of the multiplicative type noise is needed. The challenge of applying the Bihari–LaSalle inequality to nonlinear SPDEs motivates us to formulate inequality [\(3.21\)](#page-15-0) and establish Lemma [3.8.](#page-13-5) These two steps constitute some key components of the proof for Theorem [1.4.](#page-4-0)

Figure 2.1: Applying Newton's method for one step starting from the point $(k^{1/(1-a)}, k^{1/(1-a)} + b)$ on the graph to estimate (find an upper bound of) the solution x^* to the equation $x = kx^a + b$.

3 Proof of the main results

3.1 Technical lemmas

In this part, we provide some technical lemmas that will be used in the proof of Theorem [1.4.](#page-4-0) The proofs of these lemmas are postponed to Section [3.6.](#page-20-0)

Lemma 3.1. Let ρ be a function satisfying Hypothesis [1.2.](#page-4-2) Then there exist two functions g^+ : $[M_0, \infty) \to \mathbb{R}$ and $g^- : (-\infty, -M_0] \to \mathbb{R}$ that satisfy the following properties:

 (i) g^+ *is nonnegative, non-increasing, right-continuous, and it satisfies that*

$$
\lim_{x \to \infty} g^+(x) = 0 \quad and \quad |\rho(x)| - |\rho(M_0)| = \int_{M_0}^x g^+(y) dy \quad \text{for all } x \ge M_0;
$$

(ii) Similarly, g − *is nonnegative, non-decreasing, left-continuous, and it satisfies that*

$$
\lim_{x \to -\infty} g^{-}(x) = 0 \quad and \quad |\rho(x)| - |\rho(-M_0)| = \int_{x}^{-M_0} g^{-}(y) dy \quad \text{for all } x \le -M_0.
$$

Let $\theta_p : \mathbb{R} \to \mathbb{R}_+$ denote the power function $\theta_p(x) = |x|^p$ for $p \in \mathbb{R}$. When $\rho(u) = |u|^{\alpha}$ with $u \geq 0$ and $\alpha \in (0, 1]$, we claim that

$$
\|\rho(u)\|_p^2 \le \rho\left(\|u\|_p^2\right) \quad \text{for all } p > 0.
$$

Indeed, it is clear that

$$
\|\rho(u)\|_p^2 = \left(\theta_{\frac{2}{p}} \circ \mathbb{E} \circ \theta_p \circ |\rho|\right)(u) = \left(\theta_{\frac{2}{p}} \circ \mathbb{E} \circ \theta_p \circ \theta_\alpha\right)(u) = \left(\theta_{\frac{2}{p}} \circ \mathbb{E} \circ \theta_\alpha \circ \theta_p\right)(u)
$$

$$
\leq \left(\theta_{\frac{2}{p}} \circ \theta_\alpha \circ \mathbb{E} \circ \theta_p\right)(u) = \left(\theta_\alpha \circ \theta_{\frac{2}{p}} \circ \mathbb{E} \circ \theta_p\right)(u) = \rho\left(\|u\|_p^2\right),
$$

where the inequality is due to the concavity of the function $\rho = \theta_\alpha$ and we used twice the commutative property:

$$
\rho \circ \theta_p = \theta_\alpha \circ \theta_p = \theta_p \circ \theta_\alpha = \theta_p \circ \rho.
$$

However, for a general ρ that satisfies Hypothesis [1.2,](#page-4-2) we need to introduce ρ_p for this purpose.

Definition 3.2. For any function $\rho : \mathbb{R} \to \mathbb{R}$ and any positive number *p*, let the functions ρ_p^+ , $\rho_p^$ and $\rho_p : \mathbb{R}_+ \to \mathbb{R}_+$ be defined as follows: for all $x \in \mathbb{R}_+$,

$$
\rho_p(x) := \rho_p^+(x) + \rho_p^-(x)
$$
 with $\rho_p^{\pm}(x) := |\rho(\pm x^{1/p})|^p$. (3.1)

Or in other words, $\rho_p^{\pm}(\cdot)$ with $p > 0$ are defined so that

$$
\left|\rho\left(\text{sign}(x)|x|^{1/p}\right)\right|^p = \underbrace{\left(\theta_p \circ |\rho| \circ \theta_{1/p}\right)(x)}_{= \rho_p^+(x)} 1_{\{x \ge 0\}} + \underbrace{\left(\theta_p \circ |\rho| \circ r \circ \theta_{1/p}\right)(x)}_{= \rho_p^-(-x)} 1_{\{x < 0\}} \le \rho_p(|x|),
$$

where $r(x) := -x$ is the reflection function.

The next lemma shows that $\rho_p(\cdot)$ and $\rho_p^{\pm}(\cdot)$ inherit the properties from $\rho(\cdot)$.

Lemma 3.3. *Suppose that* $\rho(\cdot)$ *is a function satisfying Hypothesis [1.2.](#page-4-2) For any* $p > 0$ *, let* $\rho_p(\cdot)$ *,* $\rho_p^+(\cdot)$ *and* $\rho_p^-(\cdot)$ *be given in Definition* [3.2.](#page-12-1) Then, the following properties hold:

(i) ρ_p^+ *and* ρ_p^- *admit the following representations: for any* $x \geq M_0^p$,

$$
\rho_p^+(x) - \rho_p^+(M_0^p) = \int_{M_0^p}^x g_p^+(y) dy \quad \text{with } g_p^+(x) := \left| \rho \left(+x^{1/p} \right) \right|^{p-1} g^+ \left(+x^{1/p} \right) x^{-(p-1)/p} \tag{3.2}
$$

and

$$
\rho_p^-(x) - \rho_p^-(M_0^p) = \int_{M_0^p}^x g_p^-(y) dy \quad \text{with } g_p^-(x) := \left| \rho \left(-x^{1/p} \right) \right|^{p-1} g - \left(-x^{1/p} \right) x^{-(p-1)/p}, \tag{3.3}
$$

respectively, where $g^{\pm}(\cdot)$ *are given in Lemma [3.1;](#page-11-3)*

- *(ii) Both* ρ_p^+ *and* ρ_p^- *are non-decreasing on* $[M_0^p, \infty)$ *;*
- *(iii)* There exists $M \geq M_0$, independent of *p*, such that all functions ρ_p , ρ_p^+ , and ρ_p^- are concave *on* $[M^p, \infty)$ *;*
- *(iv)* If $M_0 = 0$ *in part (iii) of Hypothesis* [1.2,](#page-4-2) then all functions ρ_p^+ , ρ_p^- and ρ_p are concave on R+*.*

Lemma 3.4. *For any* $U \in L^p(\Omega)$ *and* $p > 0$ *, it holds that*

$$
\mathbb{E}\left[\rho_p^+(|U|^p)\mathbb{I}_{[+M,+\infty)}(U)\right] \leq \rho_p^+\left(M^p + \|U\|_p^p\right) \tag{3.4}
$$

and

$$
\mathbb{E}\left[\rho_{p}^{-}(|U|^{p})1\!\!1_{(-\infty,-M]}(U)\right] \leq \rho_{p}^{-}\left(M^{p} + \|U\|_{p}^{p}\right),\tag{3.5}
$$

where the constant M is given in part (iii) of Lemma [3.3.](#page-12-0)

Lemma 3.5. *Suppose that ρ satisfies Hypotheses [1.2.](#page-4-2) Let M be the associated constant given in part (iii) of Lemma [3.3.](#page-12-0) Then for all* $p \geq 2$ *and any* $U \in L^p(\Omega)$ *, it holds that*

$$
\|\rho(U)\|_p^2 \le K_M^2 + \rho_2 \left(M^2 + \|U\|_p^2\right),\tag{3.6}
$$

where K_M *is defined in* [\(1.14\)](#page-5-5)*.* In particular, if $M_0 = 0$ in Hypothesis [1.2,](#page-4-2) then one can take $M = K_M = 0$ *in* [\(3.6\)](#page-13-6)*.*

Definition 3.6. Suppose that ρ satisfies Hypothesis [1.2.](#page-4-2) Let $\rho_2(\cdot)$ be the function given by [\(3.1\)](#page-12-2). Define $F : [M^2, \infty) \to \mathbb{R}_+ \cup {\infty}$ as

$$
F(x) := \frac{x}{4\rho_2(x)}, \quad x \ge M^2,
$$
\n(3.7)

with

$$
\frac{x}{0} := \infty \quad \text{if } x > 0; \quad \text{and} \quad F(0) := \lim_{x \downarrow 0} \frac{x}{4\rho_2(x)} \in \mathbb{R}_+ \cup \{\infty\}, \quad \text{if } M = 0 \text{ and } \rho_2(0) = 0,
$$

where $M > M_0$ is the same as in part (iii) of Lemma [3.3.](#page-12-0) We use F^{-1} to denote the (right) inverse of *F* restricted on $[2M^2, \infty)$ as follows,

$$
F^{-1}(x) := \inf \{ y \in [2M^2, +\infty) \colon F(y) \ge x \}.
$$
 (3.8)

Remark 3.7. Here are some remarks on the functions F and F^{-1} :

- (i) As stated in Definition [3.6,](#page-13-2) we allow $F(x) = \infty$ in case $x > 0$ and $\rho_2(x) = 0$, see e.g., $\rho(x) = 1$ for $|x| < 1$ and $\rho(x) = |x - 1|^{\alpha}$ for $|x| \ge 1$. If $\rho \equiv 0$ on $[M_0, \infty)$, then $F \equiv \infty$ on $[M_0, \infty)$ as well. This implies that $F^{-1} \equiv 2M^2$ on \mathbb{R}_+ . As a result, $F^{-1}(2ph(t))$ as in (1.13) is uniformly bounded in *t*, this coincides with the SHE with additive noise.
- (ii) Under part (ii) of Hypothesis [1.2,](#page-4-2) the set in (3.8) is nonempty for any $x > 0$, and thus $F^{-1}(\cdot)$ is a real-valued function.
- (iii) From the definitions, and noticing that *F* is continuous on $[M^2, \infty)$, it is easy to see that

$$
F \circ F^{-1}(x) \ge x, \quad x \ge 0,
$$
\n(3.9)

and

$$
F^{-1} \circ F(x) \le x, \quad x \ge 2M^2. \tag{3.10}
$$

Lemma 3.8. *Suppose that the function* $\rho(\cdot)$ *satisfies Hypothesis* [1.2.](#page-4-2) Let $\rho_2(\cdot)$ *be defined as in* [\(3.1\)](#page-12-2) *with* $p = 2$ *. Then, thanks to part (iii) of Lemma [3.3,](#page-12-0)* ρ_2 *is concave on* $[M^p, \infty)$ *with some* $M > M_0$ *. For any* $k, b > 0$ *, suppose* $x \geq 0$ *such that* $x \leq k\rho_2(x) + b$ *. Then,*

$$
x \le 2F^{-1}(k) + 2b < \infty. \tag{3.11}
$$

Lemma 3.9. *Under Hypothesis* [1.1](#page-3-2) *and part (i) of Hypothesis* [1.3,](#page-4-3) *for all* $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$, $\mathcal{J}_1(t, x)$ *defined in* [\(1.11\)](#page-4-6) *satisfies that*

$$
\mathcal{J}_1(t,x) \le \int_0^t ds \, k(t-s) \int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) \mathcal{J}_0^2(s,y), \tag{3.12}
$$

where

$$
k(t) := \int_{\mathbb{R}^d} dz \, p_t(z) f(z) = h'\left(\frac{t}{2}\right) < \infty. \tag{3.13}
$$

In particular, when d = 1*, it holds that*

$$
\mathcal{J}_1(t,x) \le 2^{3/2} \pi h(t) \mathcal{J}_+(t/2,x)^2 < \infty. \tag{3.14}
$$

3.2 Proof of moment growth formulas—Theorem [1.4](#page-4-0)

The proof of Theorem [1.4](#page-4-0) consists of four steps:

Step 1. In this step, we use the assumption of ρ given in Hypothesis [1.2](#page-4-2) to obtain the following nonlinear integral inequality $||u(t,x)||_p$:

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8K_M^2p h(t) + 8p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy dy' p_{t-s}(x-y) p_{t-s}(x-y')
$$

$$
\times f(y-y')\rho_2\left(M^2 + ||u(s,y)||_p^2\right).
$$
 (3.15)

Indeed, by the Burkholder-Davis-Gundy and Minkowski's inequalities to the mild form [\(1.12\)](#page-4-7),

$$
||u(t,x)||_p^2 = \left(\mathbb{E} \left[\left(\mathcal{J}_0(t,x) + \int_0^t \int_{\mathbb{R}^d} p_{t-s}(x-y) \rho(u(s,y)) W(\mathrm{d}s, \mathrm{d}y) \right)^p \right] \right)^{2/p}
$$

\n
$$
\leq 2 \mathcal{J}_0^2(t,x) + 8p \left\| \int_0^t \mathrm{d}s \iint_{\mathbb{R}^{2d}} \mathrm{d}y \mathrm{d}y' \, p_{t-s}(x-y) p_{t-s}(x-y') f(y-y') \rho(u(s,y)) \rho(u(s,y')) \right\|_{p/2}
$$

\n
$$
\leq 2 \mathcal{J}_0^2(t,x) + 8p \int_0^t \mathrm{d}s \iint_{\mathbb{R}^{2d}} \mathrm{d}y \mathrm{d}y' \, p_{t-s}(x-y) p_{t-s}(x-y') f(y-y') \left\| \rho(u(s,y)) \rho(u(s,y')) \right\|_{p/2}.
$$

Then an application of the Cauchy-Schwarz inequality for the $L^{p/2}(\Omega)$ -norm yields that

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy dy' f(y - y') p_{t-s}(x - y) ||\rho(u(s,y))||_p
$$

$$
\times p_{t-s}(x - y') ||\rho(u(s,y'))||_p.
$$
 (3.16)

Taking account of the fact that $ab \leq \frac{1}{2}$ $\frac{1}{2}(a^2 + b^2)$ for all $a, b \in \mathbb{R}$, we can further deduce that

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8p \int_0^t ds \iint_{\mathbb{R}^{2d}} dy dy' p_{t-s}(x-y) p_{t-s}(x-y') f(y-y') ||\rho(u(s,y))||_p^2.
$$

Next, applying Lemma [3.5,](#page-13-7) we get

$$
\|\rho(u(s,y))\|_p^2 \le K_M^2 + \rho_2\left(M^2 + \|u(s,y)\|_p^2\right). \tag{3.17}
$$

Plugging [\(3.17\)](#page-14-2) into the previous inequality proves [\(3.15\)](#page-14-1).

Step 2. In this step, we will solve the nonlinear integral inequality [\(3.15\)](#page-14-1). Fix $t > 0$ and $x \in \mathbb{R}^d$. Using the function $h(t)$ in [\(1.9\)](#page-3-3) as a normalization constant and thanks to the concavity of ρ_2 , we can apply Jensen's inequality to the triple integrals in [\(3.15\)](#page-14-1) to write that

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8K_M^2p h(t) + 8ph(t)\rho_2(X),
$$
\n(3.18)

where

$$
X := M^2 + h(t)^{-1}Y \quad \text{with} \quad Y := \int_0^t \mathrm{d}s \iint_{\mathbb{R}^{2d}} \mathrm{d}y \mathrm{d}y' \, p_{t-s}(x-y) p_{t-s}(x-y') f(y-y') \, \|u(s,y)\|_p^2 \, .
$$

It reduces to find an upper bound for X . By using (3.15) , we deduce that

$$
Y \le 2\mathcal{J}_1(t,x) + 8K_M^2p\,h(t)^2 + \mathcal{I}
$$

where

$$
\mathcal{I} := 8p \int_0^t dr \iint_{\mathbb{R}^{2d}} dz dz' f(z - z') \rho_2(M^2 + ||u(r, z)||_p^2)
$$

$$
\times \int_r^t ds \iint_{\mathbb{R}^{2d}} dy dy' \ p_{t-s}(x - y) p_{t-s}(x - y') p_{s-r}(y - z) p_{s-r}(y' - z') f(y - y').
$$

Using the following formula,

$$
p_{t-s}(a)p_s(b) = p_{s(t-s)/t} \left(b - \frac{s}{t} \left(a + b \right) \right) p_t(a+b), \quad \text{for all } 0 < s < t \text{ and } a, b \in \mathbb{R}, \tag{3.19}
$$

we see that

$$
\int_{r}^{t} ds \iint_{\mathbb{R}^{2d}} dy dy' p_{t-s}(x-y) p_{t-s}(x-y') p_{s-r}(y-z) p_{s-r}(y'-z') f(y-y')
$$

\n
$$
= p_{t-r}(x-z) p_{t-r}(x-z') \int_{r}^{t} ds \iint_{\mathbb{R}^{2d}} dy dy' f(y-y')
$$

\n
$$
\times p_{(s-r)(t-s)} \left(y-z - \frac{s-r}{t-r}(x-z) \right) p_{(s-r)(t-s)} \left(y'-z' - \frac{s-r}{t-r}(x-z') \right)
$$

\n
$$
\leq (2\pi)^{-d} p_{t-r}(x-z) p_{t-r}(x-z') h(t), \qquad (3.20)
$$

where the last inequality follows from $[CK19, Lemma 2.6 and inequalities (2.21)–(2.23)].$ Thus,

$$
\mathcal{I} \leq 8 (2\pi)^{-d} p h(t) \int_0^t dr \iint_{\mathbb{R}^{2d}} dz dz' p_{t-r}(x-z) p_{t-r}(x-z') f(z-z') \rho_2 \left(M^2 + ||u(r,z)||_p^2 \right)
$$

$$
\leq 8 (2\pi)^{-d} p h^2(t) \rho_2(X),
$$

due to the concavity of $\rho_2(\cdot)$ (see Lemma [3.3\)](#page-12-0), [\(1.9\)](#page-3-3), and Jensen's inequality. Therefore,

$$
Y \le 2\mathcal{J}_1(t,x) + 8K_M^2p\,h(t)^2 + 8(2\pi)^{-d}ph^2(t)\rho_2(X),
$$

or equivalently,

$$
X \leq \underbrace{M^2 + 2h(t)^{-1} \mathcal{J}_1(t, x) + 8K_M^2 p h(t)}_{:= b(p, t, x)} + \underbrace{8 (2\pi)^{-d} p h(t)}_{:= k(p, t)} \rho_2(X). \tag{3.21}
$$

By Lemma [3.8,](#page-13-5) we have that

$$
X \le 2F^{-1}(k(p,t)) + 2b(p,t,x).
$$

Finally, thanks to the monotonicity of $\rho_2(\cdot)$ when $x \geq M^2$; see Lemma [3.3,](#page-12-0) plugging the above moment bounds back to (3.18) proves the moment following bounds

$$
||u(t,x)||_p^2 \le 2 J_0^2(t,x) + 8K_M^2 p h(t) + 8p h(t) \rho_2 \left(2b(p,t,x) + 2F^{-1}(k(p,t))\right). \tag{3.22}
$$

Step 3. In this step, we will simplify the expression in (3.22) and prove inequality (1.13) . Recall the definition of F^{-1} in [\(3.8\)](#page-13-1), one can show that for any $x \ge 2F^{-1}(k)$,

$$
\frac{\rho_2(x)}{x} = \frac{\rho_2(x) - \rho_2(2F^{-1}(k)) + \rho_2(2F^{-1}(k))}{x - 2F^{-1}(k) + 2F^{-1}(k)} \le \frac{1}{2k},
$$

because

$$
\frac{\rho_2(2F^{-1}(k))}{2F^{-1}(k)} \le \frac{1}{4k} \le \frac{1}{2k} \quad \text{and} \quad \frac{\rho_2(x) - \rho_2(2F^{-1}(k))}{x - 2F^{-1}(k)} \le g_2\left(2F^{-1}(k)\right) \le \frac{1}{2k},
$$

where the last inequality is proved in [\(3.32\)](#page-23-0) below. As a result, concerning the fact that $F^{-1}(k) \ge$ $2M^2$ for all $k > 0$, and $8(2\pi)^{-d} \leq 2$ for all $d \geq 1$, we can write

$$
8p h(t) \rho_2 \left(2b(p,t,x) + 2F^{-1}(k(p,t))\right) \leq \frac{4p h(t)}{k(p,t)} \left(b(p,t,x) + F^{-1}(k(p,t))\right)
$$

$$
\leq 2(2\pi)^d \left(h(t)^{-1} \mathcal{J}_1(t,x) + 4K_M^2 p h(t) + F^{-1}(2p h(t))\right).
$$

Plugging the above upper bound back to (3.22) proves (1.13) .

Step 4. The special case when $d = 1$ is an application of Lemma [3.9](#page-13-3) and the case when $\rho_2(\cdot)$ is concave separately on \mathbb{R}_+ and \mathbb{R}_- is due to Lemma [3.5.](#page-13-7) This proves both parts (i) and (ii). As for part (iii), if $M = 0$, inequality [\(1.16\)](#page-5-1) follows from part (ii). Otherwise, if $M > 0$, then (1.16) holds provided that one can verify that there exists $C > 0$ such that for all $t \geq 1$,

$$
4K_M^2p h(t) + F^{-1}(2p h(t)) \le CF^{-1}(Cph(t)).
$$
\n(3.23)

Indeed, the assumption of part (iii) ensures that $\rho_2(x) \geq c > 0$ with some uniform constant *c* > 0 if *x* > *M* is large enough, which implies that $F^{-1}(k)$ ≥ *ck* for large *k* (see [\(3.7\)](#page-13-0)). Hence, inequality [\(3.23\)](#page-16-1) holds by noticing that *h* is a non-decreasing function such that $h(t) > 0$ for all $t > 0$ under Hypothesis [1.1.](#page-3-2) This proves (1.16) .

Finally, if $\mu(\cdot)$ is bounded, then $\mathcal{J}_0(t,x)$ is bounded on $\mathbb{R}_+ \times \mathbb{R}^d$, and the same is true for $h(t)^{-1} \mathcal{J}_1(t,x)$ (see Lemma [3.9\)](#page-13-3). Then, inequality [\(1.17\)](#page-5-4) is a consequence of the fact that $F^{-1}(k)$ is bounded below by a positive constant for all k large enough because ρ is not identically zero on $(-\infty, -M_0] \cup [M_0, \infty)$. This completes the whole proof of Theorem [1.4.](#page-4-0)

3.3 Proof of tail probability—Theorem [1.5](#page-6-0)

We first prove a lemma:

Lemma 3.10. *Let X be a random variable such that for some function* $\alpha : [2, \infty) \to \mathbb{R}$ *,*

$$
\mathbb{E}[|X|^p] \le \exp(\alpha(p)) < \infty, \quad \text{for all } p \ge 2.
$$

Then, for all z > 0*, it holds that*

$$
\mathbb{P}(|X| \ge z) \le \exp\left(-\alpha^*(\log(z))\right),\tag{3.24}
$$

where $\alpha^*(\cdot)$ *is the Legendre-type transform of* $\alpha(\cdot)$ *on* $[2,\infty)$ *, namely,*

$$
\alpha^*(x) \coloneqq \sup \left\{ xp - \alpha(p) \colon p \in [2, \infty) \right\} \in \mathbb{R} \cup \{\infty\}, \quad \text{for all } x \in \mathbb{R}.
$$
 (3.25)

Proof. The lemma is demonstrated using Chebyshev's inequality: for any $z > 0$ and $p \geq 2$,

$$
\mathbb{P}(|X| \geq z) = \mathbb{P}(|X|^p \geq z^p) \leq z^{-p} \exp(\alpha(p)) = \exp(-[p \log(z) - \alpha(p)]).
$$

Then, [\(3.24\)](#page-16-2) follows immediately.

17

 \Box

Proof of Theorem [1.5.](#page-6-0) For all $t \ge 1$ and $p \ge 2$, we apply Lemma [3.10](#page-16-3) with the moment bounds given in (1.17) of Theorem [1.4](#page-4-0) to see that

$$
\mathbb{P}(u(t,x) \ge z) \le \exp\left(-H^*(\log(z))\right),\tag{3.26}
$$

where H^* : $\mathbb{R} \to \mathbb{R}$ is the Legendre-type transform (see [\(3.25\)](#page-16-4)) of $H: [2, \infty) \to \mathbb{R}$ given by

$$
H(p) := \frac{p}{2} \log \left(C_* F^{-1}(C_* ph(t)) \right), \qquad \text{for all } p \ge 2,
$$
 (3.27)

with the constant C_* given in (1.17) . Next, notice that if

$$
y \ge \max\left\{2^{-1}\log\left(2C_*M^2\right) + 1, 2^{-1}\log\left(C_*F^{-1}(2C_*h(t))\right) + 1\right\},\tag{3.28}
$$

then, we have $e^{2(y-1)}/C_* \ge 2M^2$, and (thanks to [\(3.9\)](#page-13-8))

$$
p^*(y) := (C_*h(t))^{-1} F\left(e^{2(y-1)}/C_*\right) \ge 2.
$$

Therefore,

$$
H(p^*(y)) = (2C_*h(t))^{-1} F\left(e^{2(y-1)}/C_*\right) \log\left(C_*F^{-1} \circ F\left(e^{2(y-1)}/C_*\right)\right)
$$

$$
\leq (2C_*h(t))^{-1} F\left(e^{2(y-1)}/C_*\right) \log\left(e^{2(y-1)}\right) = p^*(y)(y-1),
$$

where we have used (3.10) for the inequality. This yields that

$$
H^*(y) \ge yp^*(y) - H(p^*(y)) = yp^*(y) - p^*(y)(y-1) = (C_*h(t))^{-1} F\left(e^{2(y-1)}/C_*\right). \tag{3.29}
$$

Therefore, (1.18) is justified by plugging (3.29) in (3.26) with *y* replaced by log *z*. Similarly, the expression of L_t in [\(1.19\)](#page-6-4) can be obtained by replacing *y* by $\log z$ in [\(3.28\)](#page-17-4). 囗

3.4 Proof of H¨older regularity—Theorem [1.7](#page-6-1)

Proof of Theorem [1.7.](#page-6-1) The proof follows exactly the same arguments as those in [\[CH19](#page-46-7), Theorem 1.8] with the moment bounds obtained in Lemma [3.11](#page-17-5) below. \Box

Lemma 3.11. *Assume Hypothesis [1.1,](#page-3-2) parts (i) and (ii) of Hypothesis [1.2,](#page-4-2) and part (i) of Hypothesis* [1.3.](#page-4-3) Then the solution $u(t, x)$ to SHE [\(1.1\)](#page-1-2) satisfies that

$$
\|u(t,x)\|_{p} \leq \left(\mu' * p_{t}\right)(x) \left[H\left(t; 32pK^{2}\right)\right]^{1/2}, \quad \text{for all } (t,x,p) \in \mathbb{R}_{+} \times \mathbb{R}^{d} \times [2,\infty),\tag{3.30}
$$

where $\mu' := \sqrt{2} + 2|\mu|$ and $H(t; \lambda)$ is a non-decreasing function of *t* with a nonnegative parameter $\lambda > 0$.

Referring to the precise definition of $H(t; \lambda)$, one can consult [\[CH19](#page-46-7), Formula (2.4)]. It should be noted that $H(0; \lambda) > 0$, which means the function $H(t; \lambda)$ does not introduce any singularity at $t = 0$. In general, $H(t; \lambda)$ exhibits exponential growth rate in *t*; see [\[CK19,](#page-46-6) Lemma 2.5]. Certainly, the moment bounds (3.30) are considerably less accurate compared to those as in (1.13) , especially for large *p* or *t*. This is a worthy trade-off for removing part (ii) of Hypothesis [1.3](#page-4-3) in Lemma [3.11,](#page-17-5) which is sufficient for deducing the Hölder continuity of the solutions in Theorem 1.7 . Notably, to achieve more precise tail estimates in Theorem [1.5,](#page-6-0) additional improved estimates for the moment increments, as illustrated in Lemma [3.12](#page-18-1) below, are required.

Proof of Lemma [3.11.](#page-17-5) Parts (i) and (ii) of Hypothesis [1.2,](#page-4-2) namely, *ρ* is locally bounded and asymptotically sublinear $\lim_{x \to \pm \infty} x^{-1} \rho(x) = 0$, imply that $|\rho(x)| \le K_{\rho}(1+|x|)$ with some universal constant $K_{\rho} > 0$ for all $x \in \mathbb{R}$. Hence, from [\(3.16\)](#page-14-4), we see that for all $(t, x, p) \in \mathbb{R}_+ \times \mathbb{R}^d \times [2, \infty)$,

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8pK_\rho^2 \int_0^t ds \iint_{\mathbb{R}^{2d}} dy dy' f(y - y') p_{t-s}(x - y) \left(1 + ||u(s,y)||_p\right) \times p_{t-s}(x - y') \left(1 + ||u(s,y')||_p\right).
$$

By setting $g(t, x) := 1 + ||u(t, x)||_p$ and denoting the above triple integral by $I(t, x)$, the above inequality implies that

$$
g(t,x)^{2} \leq 2 + 2 \|u(t,x)\|_{p}^{2} \leq \left(\sqrt{2} + 2\mathcal{J}_{+}(t,x)\right)^{2} + 16pK_{\rho}^{2}I(t,x).
$$

Therefore, for $\mu' = \sqrt{2}K_{\rho} + |\mu|$, $g(t, x)$ satisfies the following integral inequality:

$$
g(t,x)^{2} \leq \left[\left(\mu' * p_{t} \right) (x) \right]^{2} + 16 p K_{\rho}^{2} \int_{0}^{t} ds \iint_{\mathbb{R}^{2d}} dy dy' f(y - y') p_{t-s}(x - y) g(s, y) \times p_{t-s}(x - y') g(s, y').
$$

It is easy to see that μ' also satisifes part (i) of Hypothesis [1.3.](#page-4-3) Then, an application of [\[CH19](#page-46-7), Lemma 2.2 with the above μ' and $\lambda = 16pK_{\rho}^2$ implies the moment bound [\(3.30\)](#page-17-6). The property of $H(t; \lambda)$ can be found in [\[CK19](#page-46-6), Lemma 2.5]. \Box

3.5 Proof of spatial asymptotics—Theorem [1.8](#page-6-2)

The proof of Theorem [1.8](#page-6-2) is based on a tail estimate for the solution to SHE (1.1) given in Theorem [1.5.](#page-6-0) We also need moment increments in the space variable, in case of the constant initial condition, with a sharper constant than the one implicitly appearing in Theorem [1.7.](#page-6-1)

Lemma 3.12. *Assume Hypotheses* [1.2](#page-4-2) and [1.6.](#page-6-5) Let $u(t, x)$ be a solution to SHE [\(1.1\)](#page-1-2) with initial *condition* $u_0 \equiv 1$ *. Let* $C_* > 0$ *be the constant in* [\(1.17\)](#page-5-4)*. Then, the following statements are satisfied.*

(i) For all $x, y \in \mathbb{R}^d$, $t \geq 1$, and $p \geq 2$, there exists a constant $C > 0$ such that

$$
||u(t,x) - u(t,y)||_p^2 \leq C F^{-1} (C_* ph(t)) |x - y|^{2\eta};
$$

(ii) For any $x \in \mathbb{R}^d$, with B_x denoting the unit ball centered at x , it holds that

$$
\left\| \sup_{y_1, y_2 \in B_x} |u(t, y_1) - u(t, y_2)| \right\|_p^2 \le C' F^{-1} \left(C_* ph(t) \right),
$$

where $C' > 0$ *is another universal constant.*

Proof. Following the same lines as in the proof of [\[CH19](#page-46-7), Theorem 1.8], one can show that

$$
||u(t,x) - u(t,y)||_p^2 \le 4p \int_0^t \iint_{\mathbb{R}^{2d}} |p_{t-s}(x-z) - p_{t-s}(y-z)||p_{t-s}(x-z') - p_{t-s}(y-z')|
$$

$$
\times ||\rho(u(s,z))||_p ||\rho(u(s,z))||_p f(z-z').
$$

Thanks to (3.6) and part (iii) of Theorem [1.4,](#page-4-0) one can write

$$
\sup_{s \in [0,t]} \sup_{z \in \mathbb{R}^d} ||\rho(u(s,z))||_p^2 \le K_M^2 + \rho_2 \left(M^2 + \sup_{s \in [0,t]} \sup_{x \in \mathbb{R}^d} ||u(s,x)||_p^2 \right) \le K_M^2 + \rho_2 \left(M^2 + C_* F^{-1} (C_* p h(t)) \right),
$$

for all $t \geq 1$, where $C_* > 0$ is the same as in [\(1.17\)](#page-5-4). If $F^{-1}(C_* ph(t)) < 2M^2$, one can apply part (ii) of Lemma [3.3](#page-12-0) to see that

$$
I := \rho_2 \left(M^2 + C_* F^{-1}(C_* ph(t)) \right) \le \rho_2 \left((1 + 2C_*) M^2 \right).
$$

Otherwise, with $g_2(\cdot) \coloneqq g_2^+(\cdot) + g_2^-(\cdot)$ defined as in [\(3.2\)](#page-12-3) and [\(3.3\)](#page-12-4), respectively, we have,

$$
I = \rho_2(M^2) + \int_{M^2}^{M^2 + C_* F^{-1}(C_* ph(t))} g_2(x) dx
$$

\n
$$
\leq \rho_2(M^2) + \frac{C_* F^{-1}(C_* ph(t))}{F^{-1}(C_* ph(t)) - M^2} \int_{M^2}^{F^{-1}(C_* ph(t))} g_2(x) dx \leq (2C_* + 1)\rho_2 \left(F^{-1}(C_* ph(t)) \right).
$$

Combining the above two cases, we have that

$$
I \le K_1 + C_1 \rho_2 \left(F^{-1}(C_* ph(t)) \right) = K_1 + C_1 \frac{F^{-1}(C_* ph(t))}{F \circ F^{-1}(Cph(t))}
$$

$$
\le K_1 + C_1 \frac{F^{-1}(C_* ph(t))}{C_* ph(t)}.
$$

with some universal positive constants C_1 and K_1 . Additionally, following the same idea as in the proof of (3.23) , and noting $h(\cdot)$ is a non-decreasing function with $h(t) > 0$ for $t > 0$, we can further simplify above inequality as follows:

$$
I \leq C_2 p^{-1} F^{-1} \left(C_* ph(t) \right),
$$

where $C_2 > 0$ is a universal constant. Hence,

$$
||u(t,x) - u(t,y)||_p^2 \le 4C_2 F^{-1}(C_* ph(t)) \int_0^t ds \iint_{\mathbb{R}^{2d}} dz dz' f(z - z') |p_{t-s}(x - z) - p_{t-s}(y - z)| \times |p_{t-s}(x - z') - p_{t-s}(y - z')|.
$$

The rest proof of part (i) is the same as Step 1 in the proof of [\[CH19](#page-46-7), Theorem 1.8]. The proof of part (ii) also follows from a classical argument in the proof of Kolmogorov's continuity criterion. Thus, we omit it and refer interested readers to, e.g., $[Dal+09,$ Theorem 4.3 on page 10 for more references. \Box

Now we are ready to present the proof of Theorem [1.8.](#page-6-2)

Proof of Theorem [1.8.](#page-6-2) Let C_1 and $C_2 > 0$ be two generic constants that will be fixed later. Denote

$$
Q(R) := C_1 \sqrt{F^{-1}(C_2h(t)\log(R))}, \quad R > 0.
$$

It is clear that for any $R > 0$ fixed, $Q(R)$ is increasing in both C_1 and C_2 . Following the idea as in [\[CJK13](#page-46-0)], to apply the Borel–Cantelli lemma, we need to estimate

$$
\mathcal{T}_1(R) := \mathbb{P}\left\{\max_{x \in \{y \in \mathbb{Z}^d \colon |y| \le R\}} |u(t,x)| \ge Q(R)\right\}
$$

and

$$
\mathcal{T}_2(R) \coloneqq \mathbb{P}\left\{\max_{x \in \left\{y \in \mathbb{Z}^d \colon |y| \le R\right\}} \sup_{y \in B_x} |u(t,y) - u(t,x)| \ge Q(R)\right\},\,
$$

which come from the following inequality:

$$
\mathbb{P}\left\{\sup_{|x|\leq R}|u(t,x)|\geq 2Q(R)\right\}\leq \mathcal{T}_1(R)+\mathcal{T}_2(R),
$$

for all positive integer *R* such that $Q(R) \geq L_t$; see Theorem [1.5.](#page-6-0)

Let C_* be the constant in [\(1.17\)](#page-5-4). Assume $C_1 = \sqrt{C_*}e$ and $C_2 = (2+2d)C_*$. Then, due to Theorem [1.5,](#page-6-0) we have

$$
\mathcal{T}_1(R) \le \sum_{x \in \{y \in \mathbb{Z}^d : \ |y| \le R\}} \mathbb{P}\{|u(t,x)| \ge Q(R)\}
$$

$$
\le (2R)^d \exp\left(-(C_*h(t))^{-1}F\left(Q(R)^2/(C_*e^2)\right)\right) = R^{-2}.
$$

The estimate for \mathcal{T}_2 is quite similar. By using the same argument as in Theorem [1.5](#page-6-0) and taking account of part (ii) of Lemma [3.12,](#page-18-1) one can show that with some $L_t' > 0$ and the same constants C_* and C' as in part (ii) of Lemma [3.12,](#page-18-1) for all $z \geq L'_t$,

$$
\mathbb{P}\left(\sup_{y_1,y_2\in B_x} |u(t,y_1)-u(t,y_2)| \geq z\right) \leq \exp\left(-(C_*h(t))^{-1}F\left(z^2/(C'e^2)\right)\right).
$$

Then, with $C_1 = \sqrt{C'}e$ and $C_2 = (2 + 2d)C_*$, we have $\mathcal{T}_1(R) \leq R^{-2}$ as well. Therefore, with appropriate $C_1, C_2 > 0$ and $L_t > 0$ the same as in (1.19) , the next inequality holds,

$$
\sum_{R=1}^{\infty} \mathbb{P}\left\{\sup_{|x|\leq R} |u(t,x)| \geq 2Q(R)\right\} \leq \sum_{R=1}^{\lfloor L_t \vee L_t'\rfloor} \mathbb{P}\left\{\sup_{|x|\leq R} |u(t,x)| \geq 2Q(R)\right\}
$$

$$
+ \sum_{R=\lfloor L_t \vee L_t'\rfloor+1}^{\infty} (\mathcal{T}_1(R) + \mathcal{T}_2(R)) < \infty.
$$

 \Box

An application of the first Borel–Cantelli lemma completes the proof of Theorem [1.8.](#page-6-2)

3.6 Proofs of technical lemmas

Proof of Lemma [3.1.](#page-11-3) Since $|\rho|$ is concave on $[M_0, \infty)$, we can find a non-increasing and rightcontinuous function g^+ on $[M_0, \infty)$ such that $|\rho(x)| - |\rho(M_0)| = \int_{M_0}^x g^+(y) dy$ holds for all $x \in$ $[M_0, \infty)$; see [\[NP18](#page-48-14), Theorems 1.4.2 and 1.5.2]. We claim that $g^{\dagger} \geq 0$. Because otherwise, due to part (i) of Hypothesis [1.2](#page-4-2) and the monotonicity of g^+ , we have $\lim_{x\to\infty} \int_M^x g^+(y) = -\infty$, and thus $\lim_{x\to\infty} |\rho(x)| = -\infty$, which is impossible. Therefore, $g^+ \geq 0$. Regarding the limit $\lim_{x\to\infty} g^+(x) = 0$, if it is not true, then $\lim_{x\to\infty} g^+(x) > 0$, which implies that $\lim_{x\to\infty} \rho(x) = 0$ ∞ . Thus, by L'Hôpital's rule, $\lim_{x\to\infty} \frac{\rho(x)}{x} = \lim_{x\to\infty} g^+(x) > 0$, which contradicts part (ii) of Hypothesis [1.2.](#page-4-2) Hence, $\lim_{x\to\infty} g^+(x) = 0$. This proves all properties related to g^+ . The case for \Box $g^{-}(\cdot)$ can be proved in the same way.

Proof of Lemma [3.3.](#page-12-0) The representations in both [\(3.2\)](#page-12-3) and [\(3.3\)](#page-12-4) are direct consequences of the definitions of ρ_p^{\pm} in [\(3.1\)](#page-12-2) and Lemma [3.1.](#page-11-3) Part (ii) is an immediate consequence of part (i). Now we prove part (iii). It suffices to show the case for $\rho_p^+(\cdot)$ since the case for $\rho_p^-(\cdot)$ can be proved in the same way and the case for ρ_p follows from those two cases. Let g^+ be given in [\(3.2\)](#page-12-3). We need to show that g_p^+ is non-increasing on $[M^p, \infty)$ with some $M \geq M_0$. We know that g^+ is non-increasing on (M_0, ∞) , it suffices to show that $\varphi(x) = \frac{|\rho(x)|}{x}$ is non-increasing for *x* large enough. To show this property, we write

$$
|\rho(x)| = \int_{M_0}^x g^+(y) dy + |\rho(M_0)|.
$$

Thus for almost every $x \in (M_0, \infty)$,

$$
\varphi'(x) = x^{-2} \left[g^{+}(x) \left(x - M_0 \right) - \int_{M_0}^x g^{+}(y) dy + g^{+}(x) M_0 - |\rho(M_0)| \right] \le g^{+}(x) M_0 - |\rho(M_0)|,
$$

where the last inequality follows from the fact that g^+ is non-increasing on (M_0, ∞) . Since $g^+(x) \to$ 0 as $x \to \infty$ (see Lemma [3.1\)](#page-11-3), we conclude that $g^+(x)M_0 - |\rho(M_0)| < 0$ for *x* large enough. In other words, there exists $M \geq M_0$, such that φ is non-increasing on $[M, \infty)$. Therefore, ρ_p^+ is concave on $[M^p, \infty)$. For the same reason, we can show that ρ_p^- , and thus ρ_p , are also concave on $[M^p, \infty)$ with a possibly different $M \geq M_0$. This proves part (ii). Finally, from the above argument, we see that when $M_0 = 0$, then $g^+(x)M_0 - |\rho(M_0)| = -|\rho(0)| \le 0$ for all $x \ge 0$. Hence, part (iv) follows. This completes the proof of Lemma [3.3.](#page-12-0)

Proof of Lemma [3.4.](#page-12-5) In the following, we will prove [\(3.4\)](#page-12-6) only. The proof of [\(3.5\)](#page-12-7) is similar. Let *p* > 0 and $U \in L^p(\Omega)$. Set $\alpha := \mathbb{P}(U \geq M)$. It is clear that when $\alpha = 0$, the inequality [\(3.4\)](#page-12-6) is trivially true. So, we may assume that $\alpha > 0$. Since $\rho_p^+(\cdot)$ is concave on $[M^p, \infty)$ (see Lemma [3.3\)](#page-12-0), we can apply Jensen's inequality to see that

$$
\mathbb{E}\left[\rho_p^+\left(|U|^p\right)\mathrm{I\!I}_{[M,\infty)}(U)\right]\leq \alpha\rho_p^+\left(\frac{1}{\alpha}\mathbb{E}\left[|U|^p\mathrm{I\!I}_{[M,\infty)}(U)\right]\right)\leq \alpha\rho_p^+\left(\frac{1}{\alpha}\mathbb{E}\left[|U|^p\right]\right),
$$

where the second inequality is due to the monotonicity of $\rho_p^+(\cdot)$; see Lemma [3.3.](#page-12-0) Denote $x = \mathbb{E}[|U|^p]$. Since $x/\alpha \ge M^p$, another application of the monotonicity and the concavity of $\rho_p^+(\cdot)$ shows that

$$
\rho_p^+(\alpha^{-1}x) \le \rho_p^+(M^p + \alpha^{-1}x) \le \rho_p^+(M^p) + \alpha^{-1}(\rho_p^+(M^p + x) - \rho_p^+(M^p)) \le \alpha^{-1}\rho_p^+(M^p + x).
$$

This implies that

$$
\mathbb{E}\big[\rho_p^+(|U|^p)\mathbb{I}_{[M,\infty)}(U)\big] \leq \rho_p^+ (M^p + \mathbb{E}\left[|U|^p\right]) = \rho_p^+ \left(M^p + \|U\|_p^p\right),
$$

 \Box

which completes the proof of Lemma [3.4.](#page-12-5)

Proof of Lemma [3.5.](#page-13-7) Fix an arbitrary $p \geq 2$. By the subadditivity of $\theta_{2/p}(\cdot)$,

$$
\|\rho(U)\|_p^2 \le \left(\mathbb{E}\left[|\rho(U)|^p 1_{\{|U|\le M\}}\right]\right)^{2/p} + \left(\mathbb{E}\left[|\rho(U)|^p 1_{\{u\ge M\}}\right]\right)^{2/p} + \left(\mathbb{E}\left[|\rho(U)|^p 1_{\{u\le -M\}}\right]\right)^{2/p}
$$

\n
$$
\le K_M^2 + \left(\theta_{2/p} \circ \mathbb{E}_{\ge 0} \circ \theta_p \circ |\rho|\right)(U) + \left(\theta_{2/p} \circ \mathbb{E}_{\le 0} \circ \theta_p \circ |\rho|\right)(U)
$$

\n
$$
= K_M^2 + \left(\theta_{2/p} \circ \mathbb{E}_{\ge 0} \circ \theta_p \circ |\rho| \circ \theta_{1/p} \circ \theta_p\right)(U) + \left(\theta_{2/p} \circ \mathbb{E}_{\le 0} \circ \theta_p \circ |\rho| \circ r \circ \theta_{1/p} \circ \theta_p\right)(U)
$$

\n
$$
= K_M^2 + \left(\theta_{2/p} \circ \mathbb{E}_{\ge 0} \circ \rho_p^+ \circ \theta_p\right)(U) + \left(\theta_{2/p} \circ \mathbb{E}_{\le 0} \circ \rho_p^- \circ \theta_p\right)(U),
$$

where \mathbb{E}_{\geq} denote the expectation under $\{u \geq M\}$, namely, $\mathbb{E}_{\geq} (f(U)) = \mathbb{E} (f(U)1_{\{u \geq M\}})$, and \mathbb{E}_{\leq} is defined in the same way. Set $y \coloneqq M^p + ||U||_p^p$ $_p^p$. By Lemma [3.4,](#page-12-5)

$$
\|\rho(U)\|_p^2 \le K_M^2 + \left(\theta_{2/p} \circ \rho_p^+\right)(y) + \left(\theta_{2/p} \circ \rho_p^-\right)(y) \n= K_M^2 + \left(\theta_{2/p} \circ \theta_p \circ |\rho| \circ \theta_{1/p}\right)(y) + \left(\theta_{2/p} \circ \theta_p \circ |\rho| \circ r \circ \theta_{1/p}\right)(y) \n= K_M^2 + \left(\theta_2 \circ |\rho| \circ \theta_{1/2} \circ \theta_{2/p}\right)(y) + \left(\theta_2 \circ |\rho| \circ r \circ \theta_{1/2} \circ \theta_{2/p}\right)(y) \n= K_M^2 + \left(\rho_2^+ \circ \theta_{2/p}\right)(y) + \left(\rho_2^- \circ \theta_{2/p}\right)(y) \n\le K_M^2 + \rho_2^+ \left(M^2 + \|U\|_p^2\right) + \rho_2^- \left(M^2 + \|U\|_p^2\right),
$$

 \Box where the last step is due to the subadditivity of $\theta_{2/p}(\cdot)$ and the monotonicity of $\rho_2^{\pm}(\cdot)$.

Proof of Lemma [3.8.](#page-13-5) The finiteness of $F^{-1}(k) + b$ is a consequence of part (ii) of Hypothesis [1.2.](#page-4-2) Thus it suffices to show the first inequality in (3.11) . To this end, we first prove it by verifying $x \leq \gamma_0(k) + 2b$, with

$$
\gamma_0(k) := \inf \left\{ x \in (M^2, \infty) \colon \frac{\rho_2(x)}{x} \le \frac{1}{k} \text{ and } g_2(x) \le \frac{1}{2k} \right\}.
$$

Here, $g_2(\cdot) := g_2^+(\cdot) + g_2^-(\cdot)$ with $g_2^+(\cdot)$ and $g_2^-(\cdot)$ defined in [\(3.2\)](#page-12-3) and [\(3.3\)](#page-12-4), respectively. Note that $\gamma_0(k) \geq M^2$ and $\rho_2(\cdot)$ is concave on $[M^2,\infty)$. Thus, if $x \geq \gamma_0(k) + 2b$, we have

$$
k\rho_2(x) + b = k\rho_2(\gamma_0(k)) + k \int_{\gamma_0(k)}^x g_2(y) dy + b
$$

\n
$$
\leq \gamma_0(k) + \frac{1}{2}(x - \gamma_0(k)) + b
$$

\n
$$
\leq \gamma_0(k) + \frac{1}{2}(x - \gamma_0(k)) + \frac{1}{2}(x - \gamma_0(k)) = x.
$$

In the next step, we can show that

$$
\gamma_0(k) \le 2F^{-1}(k)
$$
, for all $k > 0$. (3.31)

First if $M = 0$, then

$$
\frac{2}{F(x)} = \frac{x}{2\rho_2(x)} \le \frac{x}{\rho_2(x)} = x \left(\int_0^x dy g_2(y) \right)^{-1} \le \frac{1}{g_2(x)},
$$

by the non-increasing property of $g_2(\cdot)$, and [\(3.31\)](#page-22-0) follows immediately from the definition of F^{-1} ; see [\(3.8\)](#page-13-1). On the other hand, if $M > 0$. Then, for any $x \ge 2F^{-1}(k) \ge 3F^{-1}(k)/2$, it holds that

$$
g_2(x) \le \frac{\rho_2(x) - \rho_2(M^2)}{x - M^2} \le \frac{\rho_2(3F^{-1}(k)/2) - \rho_2(M^2)}{3F^{-1}(k)/2 - M^2}.
$$

Notice that $F^{-1}(k) \ge 2M^2$ implies $3F^{-1}(k)/2 - M^2 \ge F^{-1}(k)$ and

$$
\rho_2(3F^{-1}(k)/2) - \rho_2(M^2) = \int_{M^2}^{3F^{-1}(k)/2} \mathrm{d}y g_2(y) \leq \frac{3F^{-1}(k)/2 - M^2}{F^{-1}(k) - M^2} \int_{M^2}^{F^{-1}(k)} \mathrm{d}y g_2(y)
$$

$$
\leq 2\rho_2 \left(F^{-1}(k)\right).
$$

Combining above inequalities and [\(3.9\)](#page-13-8), we get

$$
g_2(x) \le \frac{2\rho_2\left(F^{-1}(k)\right)}{F^{-1}(k)} = \frac{1}{2F \circ F^{-1}(k)} \le \frac{1}{2k}.\tag{3.32}
$$

This implies that (3.31) , and thus completes the proof of Lemma [3.8.](#page-13-5)

Proof of Lemma [3.9.](#page-13-3) Without loss generality, we assume that μ is nonnegative. Otherwise, one simply replaces μ by $|\mu|$. By the Plancherel theorem and the fact that f is nonnegative-definite, we can write

$$
\sup_{x \in \mathbb{R}^d} \int_{\mathbb{R}^d} dy \, p_t(x - y) f(y) = \sup_{x \in \mathbb{R}^d} (2\pi)^{-d} \int_{\mathbb{R}^d} d\xi \, e^{-ix \cdot \xi - \frac{t|\xi|^2}{2}} \hat{f}(\xi)
$$

$$
= (2\pi)^{-d} \int_{\mathbb{R}^d} d\xi \, e^{-\frac{t|\xi|^2}{2}} \hat{f}(\xi) = \int_{\mathbb{R}^d} dz \, p_t(z) f(z) = k(t),
$$

from which one proves [\(3.12\)](#page-13-4). As for [\(3.14\)](#page-13-11), we proceed with a general $d \geq 1$ and make the restriction to $d = 1$ when some integrability issue comes up. By the Cauchy-Schwarz inequality, we see that

$$
\int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) \mathcal{J}_0^2(s,y) \n= \int_{\mathbb{R}^d} dy \iint_{\mathbb{R}^{2d}} \mu(\mathrm{d}z) \mu(\mathrm{d}z') \, p_{t-s}(x-y) p_s(y-z) p_s(y-z') \n\leq \int_{\mathbb{R}^{2d}} |\mu|(\mathrm{d}z) |\mu|(\mathrm{d}z') \left(\int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) p_s(y-z)^2 \right)^{1/2} \left(\int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) p_s(y-z')^2 \right)^{1/2} \n= \left[\int_{\mathbb{R}^d} |\mu|(\mathrm{d}z) \left(\int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) p_s(y-z)^2 \right)^{1/2} \right]^2.
$$

Because $p_t^2(x) = (4\pi t)^{-d/2} p_{t/2}(x) \le (2\pi t)^{-d/2} p_t(x)$ for all $(t, x) \in \mathbb{R}_+ \times \mathbb{R}^d$, we can write

$$
\int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) p_s(y-z)^2 \le (2\pi s)^{-d/2} \int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) p_s(y-z) = (2\pi s)^{-d/2} p_t(x-z),
$$

which implies that

$$
\mathcal{J}_1(t,x) \leq (2t)^{d/2} \int_0^t ds \, s^{-d/2} k(t-s) \left(\int_{\mathbb{R}^d} |\mu| (\mathrm{d}z) p_{t/2}(x-z) \right)^2
$$

= $(2t)^{d/2} \mathcal{J}_+(t/2,x)^2 \int_0^t ds \, s^{-d/2} k(t-s).$

Notice that the integral against the time argument in the last expression is finite for all $t > 0$, if and only if $d = 1$. Moreover, in case $d = 1$, we can deduce that

$$
\int_0^t ds \, s^{-1/2} k(t-s) \le (2\pi)^{-1/2} \int_0^t ds \, s^{-1/2} (t-s)^{-1/2} \int_{\mathbb{R}} dz \, e^{-\frac{x^2}{2t}} f(z) = \pi t^{1/2} k(t).
$$

Therefore,

$$
\mathcal{J}_1(t,x) \le \sqrt{2\pi} \, t \, k(t) \, \mathcal{J}_+(t/2,x)^2 < \infty. \tag{3.33}
$$

Finally, notice that $k(t)$ is a nonincreasing function of *t* because $k(t) = (2\pi)^{-d} \int_{\mathbb{R}^d} \hat{f}(\mathrm{d}\xi) e^{-t|\xi|^2/2}$. From (1.9) , we see that

$$
2h(t) \ge 2h(t/2) = \int_0^t ds \, k(s) \ge \int_0^t ds \, k(t) = tk(t).
$$

Plugging this inequality back to (3.33) proves (3.14) . This completes the proof of Lemma [3.9.](#page-13-3) \Box

 \Box

4 Examples on various sublinear diffusion coefficients *ρ*

In this section, we derive the explicit expressions for $F(\cdot)$ and its inverse $F^{-1}(\cdot)$ for the sublinear diffusion coefficients ρ given in examples (1.4) – (1.6) . The results are summarized in Figure [4.2](#page-28-0) at the end of this session.

Proposition 4.1. *The following diffusion coefficient*

$$
\rho(u) = \frac{|u|}{(r+|u|)^{1-\alpha}} \qquad \text{for } u \in \mathbb{R}, \text{ with } r \ge 0 \text{ and } \alpha \in [0,1), \tag{4.1}
$$

which reduces to

$$
\rho(u) = \begin{cases} |u|^{\alpha} & \text{with } \alpha \in [0, 1) & \text{if } r = 0, \\ \frac{|u|}{1 + |u|} & \text{if } \alpha = 0, \end{cases}
$$

satisfies the following properties:

- *1.* ρ *is globally Lipschitz provided that* $r > 0$ *, in which case,* $\rho'(0) = r^{-(1-\alpha)} < \infty$;
- *2. ρ satisfies Hypothesis* [1.2](#page-4-2) *with* $M_0 = 0$;
- *3. the corresponding F and F* −1 *take the following explicit forms:*

$$
F(u) = \frac{1}{8} \left(r + u^{1/2} \right)^{2(1-\alpha)}, \quad \text{for all } u \ge 0,
$$

and

$$
F^{-1}(x) = ((8x)^{1/(2(1-\alpha))} - r)^2 1\!\!1_{\{x \ge 8^{-1}r^{2(1-\alpha)}\}}.
$$
\n(4.2)

In particular, $F^{-1}(x) \leq (8x)^{1/(1-\alpha)}$ *for all* $x \geq 0$ *.*

Proof. The diffusion coefficient ρ given in [\(4.1\)](#page-24-1) is clearly locally bounded, and has sublinear growth:

 $\rho(u) \asymp |u|^{\alpha}$ as $|u| \to \infty$.

Notice that for $u \geq 0$,

$$
\rho'(u) = (r + u)^{\alpha - 2} (r + \alpha u) \to r^{-(1 - \alpha)}
$$
 as $u \to 0$.

 $\rho(\cdot)$ is concave separately on \mathbb{R}_+ and \mathbb{R}_- since

$$
\rho''(u) = -(1 - \alpha) (r + |u|)^{\alpha - 3} (2r + \alpha |u|) \le 0, \text{ for all } u \in \mathbb{R}.
$$

Therefore, ρ satisfies Hypothesis [1.2](#page-4-2) with $M_0 = 0$.

The form of ρ (see also [\(1.4\)](#page-2-1)) makes it easy to compute ρ_2 , *F* and *F*⁻¹. Indeed, in this case, for $u \geq 0$,

$$
\rho_2(u) = \frac{2u}{(r+u^{1/2})^{2(1-\alpha)}} \quad \text{and} \quad F(u) = \frac{1}{8}\left(r+u^{1/2}\right)^{2(1-\alpha)}, \quad \text{for all } u \ge 0.
$$

The expression of F^{-1} is readily to be derived from the above formula for *F*.

 \Box

$$
\begin{array}{c}\n\textbf{Case (iii):} \\
\alpha & \alpha = 1, \beta > 0 \\
\textbf{Case (ii):} \\
\alpha \in (0,1), \beta \in \mathbb{R} \\
\rho(u) = |u|^{\alpha} [\log (e+u^2)]^{-\beta} \\
\textbf{Case (i):} \\
\alpha = 0, \beta < 0 \\
\rho(u) = [\log (e+u^2)]^{-\beta} \\
\textbf{Case (i):} \\
\rho(u) = [\log (e+u^2)]^{-\beta}\n\end{array}
$$

Figure 4.1: Three cases for ρ given in (1.5) and Proposition [4.2.](#page-25-0)

The diffusion term ρ in the following examples does not exhibit global concavity, which motivates us to propose the asymptotic concave condition in Hypothesis [1.2.](#page-4-2)

Proposition 4.2. *Suppose that*

$$
\rho(u) = |u|^{\alpha} \left[\log \left(e + u^2 \right) \right]^{-\beta},
$$

with α, β *fulfilling one of the three conditions given as in* [\(1.5\)](#page-2-3)*; see Figure [4.1.](#page-25-1) Then we have*

- *1.* ρ *is globally Lipschitz only when* $\alpha = 1$ *(for all* $\beta \in \mathbb{R}$ *), in which case,* $\rho'(0) = 1$ *;*
- *2.* ρ *satisfies Hypothesis* [1.2](#page-4-2) *with some* $M_0 > 0$;
- *3. the corresponding F and F* [−]¹ *are given below:*

$$
F(u) = \frac{1}{8}u^{1-\alpha} \left[\log(e+u) \right]^{2\beta}, \quad u > 0,
$$
\n(4.3)

and

$$
\begin{cases}\nF^{-1}(x) \asymp (8x)^{1/(1-\alpha)} \left[\frac{1}{1-\alpha} \log(e+x) \right]^{2\beta/(1-\alpha)}, & x \to \infty, \quad \text{Case } (i) \text{ and } (ii), \\
F^{-1}(x) = \left(\exp\left(2^{3/(2\beta)} x^{1/(2\beta)}\right) - e \right) \mathbb{1}_{\{x > 8^{-1}\}}, & \text{Case } (iii).\n\end{cases} (4.4)
$$

In particular, in Cases (i) and (ii),

$$
F^{-1}(x) \lesssim \left[x \left(\log x \right)^{-2\beta} \right]^{1/(1-\alpha)}, \quad \text{as } x \to \infty.
$$

Proof. It is clear that ρ is locally bounded and has sublinear growth at infinity. For the asymptotic concavity, it is elementary, though tedious, to show that for $u \geq 0$,

$$
\rho'(u) = u^{\alpha - 1} \left[\log \left(e + u^2 \right) \right]^{-\beta} \left(\alpha - \frac{2\beta u^2}{\left(e + u^2 \right) \log \left(e + u^2 \right)} \right) \quad \text{and} \quad \rho''(u) = q_1(u)q_2(u)
$$

with

$$
q_1(u) := (e + u^2)^{-2} u^{\alpha - 2} \left[\log (e + u^2) \right]^{-\beta - 2},
$$

\n
$$
q_2(u) := 4\beta(\beta + 1)u^4 - 2\beta u^2((2\alpha - 1)u^2 + 2\alpha e + e) \log(e + u^2)
$$

\n
$$
+ \alpha(\alpha - 1)(e + u^2)^2 \left[\log (e + u^2) \right]^2.
$$

From the expression of $\rho'(u)$, we see that ρ is globally Lipschitz only when $\alpha = 1$ and in that case $\rho'(0) = 1$. From the expression of $\rho''(u)$, we see that $q_1(u) \geq 0$ for $u > 0$, but $q_2(u)$ may take negative values when u is small. By considering the three cases given in (1.5) (see Figure [4.1\)](#page-25-1), we see that

$$
q_2(u) \asymp \overline{q}_2(u) < 0, \quad |u| \to \infty, \quad \text{with } \overline{q}_2(u) := \begin{cases} 2\beta u^4 \log(e + u^2), & \text{Case (i)}, \\ \alpha(\alpha - 1)(e + u^2)^2 \log^2(e + u^2), & \text{Case (ii)}, \\ -2\beta u^4 \log(e + u^2), & \text{Case (iii)}. \end{cases}
$$

Hence, ρ is asymptotically concave separately on $[M_0, \infty)$ and $(-\infty, -M_0]$ for some constant $M_0 >$ 0. This proves that ρ satisfies Hypothesis [1.2.](#page-4-2)

It is straightforward to derive the expression for *F*:

$$
\rho_2(u) = 2u^{\alpha} [\log(e+u)]^{-2\beta}
$$
 and $F(u) = \frac{u}{4\rho_2(u)} = \frac{1}{8} u^{1-\alpha} [\log(e+u)]^{2\beta}$, $u > 0$,

It remains to prove [\(4.4\)](#page-25-2). In Case (iii), namely, $\alpha = 1$ and $\beta > 0$, $F^{-1}(x)$ can be obtained explicitly as given in the statement of the proposition. However, in Cases (i) and (ii), we have $\alpha \in [0,1)$, and it seems impossible to find the explicit formula for $F^{-1}(x)$. Instead, one can find the exact asymptotics. Set

$$
\Theta(x) \coloneqq x^{1/(1-\alpha)} \left[\log(e+x) \right]^{-2\beta/(1-\alpha)}, \quad (x > 0).
$$

From [\(4.3\)](#page-25-3), we see that, for $\lambda > 0$ and $x > 0$,

$$
F(\lambda\Theta(x)) = \frac{\lambda\Theta(x)}{4\rho_2(\lambda\Theta(x))} = \frac{\lambda^{1-\alpha}x}{8} \left[\frac{\log(e+x)}{\log(e+\lambda x^{1/(1-\alpha)}[\log(e+x)]^{-2\beta/(1-\alpha)})} \right]^{-2\beta}
$$

$$
\approx 8^{-1}\lambda^{1-\alpha}(1-\alpha)^{-2\beta}x, \text{ as } x \to \infty.
$$

Hence, by choosing $\lambda_0 := 8^{1/(1-\alpha)}(1-\alpha)^{2\beta/(1-\alpha)}$, we see that

$$
F(\lambda_0 \Theta(x)) \simeq x
$$
, or equivalently $F^{-1}(x) \simeq \lambda_0 \Theta(x)$, as $x \to \infty$.

This completes the proof of Proposition [4.2.](#page-25-0)

Proposition 4.3. *The diffusion coefficient*

$$
\rho(u) = |u| \exp\left(-\beta \left(\log \log\left(e + u^2\right)\right)^{\kappa}\right) \quad \text{with } \kappa > 0 \text{ and } \beta > 0
$$

have the following properties:

1. *ρ is globally Lipschitz continuous with* $\rho'(0) = 1$;

 \Box

- 2. *ρ* satisfies Hypothesis [1.2](#page-4-2) with some $M_0 > 0$;
- *3. the corresponding F and F* −1 *take the following explicit forms:*

$$
F(u) = \frac{1}{8} \exp(2\beta (\log \log(e+u))^{\kappa}), \text{ for all } u > 0,
$$

and

$$
F^{-1}(x) = \left(\exp\left\{\exp\left(\left((2\beta)^{-1}\log(8x)\right)^{1/\kappa}\right)\right\} - e\right)1\!\!1_{\{x > 1/8\}}.\tag{4.5}
$$

.

Proof. It is easy to see that *ρ* is locally bounded. Since $\beta > 0$, we see that *ρ* has sublinear growth at infinity. Thus, ρ satisfies both parts (i) and (ii) of Hypothesis [1.2.](#page-4-2) Notice that

$$
\rho'(u) = e^{-\beta \log^{\kappa}(\log(e+u^2))} \left(1 - \frac{2\kappa \beta u^2 \log^{\kappa-1}(\log(e+u^2))}{(e+u^2)\log(e+u^2)}\right)
$$

Using the fact that $\log(a+u) \approx \log(a) + u/a$ as $u \to 0_+$ with $a > 0$, we see that

$$
\rho'(u) \approx e^{-\beta e^{-\kappa} u^{2\kappa}} \left(1 - 2\kappa \beta e^{-\kappa} u^{2\kappa}\right) \to 1 \quad \text{as } u \to 0_+, \text{ provided that } \kappa > 0.
$$

Hence, ρ is globally Lipschitz with $\rho'(0) = 1$. As for the asymptotic concavity, we have that $\rho''(u) = q_1(u)q_2(u)$ for $u > 0$ with

$$
q_1(u) = 2\kappa \beta u e^{-\beta \log^{\kappa}(\log(e+u^2))} \frac{\log^{\kappa-2}(\log(e+u^2))}{(e+u^2)^2 \log^2(e+u^2)} \ge 0,
$$

and

$$
q_2(u) = 2\kappa \beta u^2 \log^{\kappa} \left(\log \left(e + u^2 \right) \right) - 2(\kappa - 1)u^2
$$

$$
- \left(\left(u^2 + 3e \right) \log \left(e + u^2 \right) - 2u^2 \right) \log \left(\log \left(e + u^2 \right) \right)
$$

$$
\asymp - u^2 \log \left(e + u^2 \right) \log \left(\log \left(e + u^2 \right) \right) \le 0, \quad \text{as } u \to \infty.
$$

Therefore, ρ is asymptotically concave on \mathbb{R}_+ . This proves that ρ satisfies Hypothesis [1.2.](#page-4-2) Finally, it is routine to compute *F* and F^{-1} . This completes the proof of Proposition [4.3.](#page-26-0) \Box

Remark 4.4. When $M_0 \neq 0$, ρ is only asymptotically concave (separately) on \mathbb{R}_+ and \mathbb{R}_- . Hence, when using the formulas in (4.4) and (4.5) , we require the argument of F^{-1} to be sufficiently large.

5 Moment growth for other SPDEs in the sublinear regime

In this section, we will explore the moment upper bounds for other SPDEs in the sublinear regime.

5.1 Nonlinear stochastic space-time fractional diffusion equations

In this part, we explore the moment upper bounds for the following nonlinear fractional stochastic partial differential equations (*fractional SPDE*) driven by the multiplicative space-time white noise:

$$
\left[\left(\frac{\partial}{\partial t} \right)^b + \frac{1}{2} (-\Delta)^{a/2} \right] u(t, x) = I_t^{\gamma} \left[\rho(u(t, x)) \dot{W}(t, x) \right], \tag{5.1}
$$

Figure [4.2](#page-25-0): Summary of the asymptotics of F^{-1} for ρ in Propositions [4.1,](#page-24-2) 4.2 and [4.3.](#page-26-0)

where $\left(\frac{\partial}{\partial t}\right)^b$, $(-\Delta)^{a/2}$, and I_t^{γ} denote the *Caputo fractional differential operator* with $b \in (0,2)$, fractional Laplacian operator with $a \in (0, 2]$, and the *Riemann-Liouville operator* with $\gamma > 0$, respectively. We assume that equation (5.1) starts from the following initial condition(s):

$$
\begin{cases} u(0, \cdot) = \mu_0, & b \in (1/2, 1], \\ u(0, \cdot) = \mu_0, & \frac{\partial}{\partial t} u(t, \cdot) \Big|_{t=0} = \mu_1, & b \in (1, 2), \end{cases}
$$
(5.2)

where both μ_0 and μ_1 are nonnegative measures. For a more detailed account of this equation, we refer interested readers to [\[CHN19](#page-45-7); [CGS22](#page-45-8); [GSS23](#page-47-13)].

Definition 5.1. A random field $u = \{u(t,x): (t,x) \in \mathbb{R}_+ \times \mathbb{R}^d\}$ is a solution to equation [\(5.1\)](#page-27-3), if it satisfies the following mild formulation:

$$
u(t,x) = \mathcal{J}_0(t,x) + \int_0^t \int_{\mathbb{R}^d} Y(t-s,x-y) \rho(u(s,y)) W(ds,dy),
$$

where

$$
\mathcal{J}_0(t,x) = \begin{cases} \int_{\mathbb{R}^d} \mu_0(\mathrm{d}y) Z(t, x - y), & b \in (0, 1], \\ \int_{\mathbb{R}^d} \mu_0(\mathrm{d}y) Z^*(t, x - y) + \int_{\mathbb{R}^d} \mu_1(\mathrm{d}y) Z(t, x - y), & b \in (1, 2), \end{cases}
$$
(5.3)

 $\{(Z(t,x), Z^*(t,x), Y(t,x))\colon (t,x)\in \mathbb{R}_+\times\mathbb{R}^d\}$ denote the fundamental functions of equation [\(5.1\)](#page-27-3), which can be formulated explicitly in terms of some special functions; see [\[CHN19](#page-45-7), Section 4].

Hypothesis 5.2. (i) Dalang's condition holds for (5.1) , namely, $b + \gamma > \frac{1}{2}(1 + \frac{db}{a})$ and $2a > d$.

(ii) Either $\gamma = 0$ or $a > d = 1$.

Definition 5.3. Let $G : \mathbb{R}_+ \times \mathbb{R}^d \to \mathbb{R}$ be a Borel function. We say that $G^2(\cdot, \circ)$ satisfies a *subsemigroup property* if there exist a positive constant C^* , a reference kernel $\mathcal{G} : \mathbb{R}_+ \times \mathbb{R}^d \to \mathbb{R}_+$, and a locally integrable function $\ell : \mathbb{R}_+ \to \mathbb{R}_+$, such that for all $(s, t, x) \in \mathbb{R}_+^2 \times \mathbb{R}^d$,

$$
\int_{\mathbb{R}^d} \mathcal{G}(t, y) dy = 1, \quad G(t, x)^2 \le \ell(t) \mathcal{G}(t, x) \quad \text{and} \quad \int_{\mathbb{R}^d} \mathcal{G}(t, x - y) \mathcal{G}(s, y) dy \le C^* \mathcal{G}(t + s, x).
$$

Lemma 5.4 (Proposition 5.10 of [\[CHN19](#page-45-7)]). Let $Y(\cdot, \circ)$ be as in Definition [5.1.](#page-28-1) Under Hypothe s es [5.2,](#page-29-0) for some constant $C_0 > 0$, $Y^2(\cdot, \circ)$ satisfies the sub-semigroup property (see Definition [5.3\)](#page-29-1) *with*

$$
\ell(t) = C_0 t^{-\sigma} \quad \text{and} \quad \mathcal{G}(t,x) = \begin{cases} k_b t^{-\frac{bd}{2}} \exp\left(-\frac{1}{2} (t^{-\frac{b}{2}} |x|)^{\lfloor b \rfloor + 1}\right), & a = 2, \\ \frac{k_{a,b} t^{b/a}}{\left(t^{\frac{2b}{a}} + |x|^2\right)^{(d+1)/2}}, & a \in (0,2), \end{cases}
$$
(5.4)

where k_b (resp. $k_{a,b}$) is a uniform constant, depending on *b* (resp. (a, b)) and

$$
\sigma := 2(1 - b - \gamma) + (bd)/a < 1. \tag{5.5}
$$

(Note that $\sigma < 1$ *in* [\(5.5\)](#page-29-2) *is due to part (i) of Hypothesis* [5.2.](#page-29-0)

Let $\mathcal{J}_0(t,x)$ be defined as in [\(5.3\)](#page-28-2) and let $\ell(t)$ and $\mathcal{G}(t,x)$ be given as in [\(5.4\)](#page-29-3). We define the next expression

$$
\mathcal{J}_1(t,x) := \int_0^t ds \,\ell(t-s) \int_{\mathbb{R}^d} dy \,\mathcal{G}(t-s,x-y) \mathcal{J}_0^2(s,y), \quad \text{for all } (t,x) \in \mathbb{R}_+ \times \mathbb{R}^d. \tag{5.6}
$$

Then, thanks to [\[CHN19,](#page-45-7) Lemma 5.12], it holds that $\mathcal{J}_1(t,x) < \infty$ for all $t > 0$ and $x \in \mathbb{R}^d$. As a result of Lemma [5.4,](#page-29-4) we can follow the same idea as in Section [3](#page-11-0) to obtain the following result:

Theorem 5.5. Let $u(t, x)$ be a solution to the fractional SPDE (5.1) driven by the space-time white *noise. Assume that*

- *1. the initial conditions given in* [\(5.2\)](#page-28-3) *satisfy* $\mathcal{J}_0(t,x) < \infty$ *for all* $t > 0$ *and* $x \in \mathbb{R}^d$;
- *2. the diffusion coefficient ρ satisfies Hypothesis [1.2;](#page-4-2)*
- *3. the parameters* (*d, a, b, γ*) *satisfy Hypothesis [5.2.](#page-29-0)*

Then for all $(t, x, p) \in \mathbb{R}_+ \times \mathbb{R}^d \times [2, \infty)$ *, it holds that*

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + K\left(t^{-(1-\sigma)}\mathcal{J}_1(t,x) + Kpt^{1-\sigma} + F^{-1}\left(Kpt^{1-\sigma}\right)\right),\tag{5.7}
$$

where F^{-1} , σ , \mathcal{J}_0 , and \mathcal{J}_1 are defined as in [\(3.8\)](#page-13-1), [\(5.5\)](#page-29-2), [\(5.3\)](#page-28-2) and [\(5.6\)](#page-29-5), respectively, and K is a *positive constant not depending on p and t. In particular,*

(i) if $|\rho(\cdot)|$ *is concave separately on* \mathbb{R}_+ *and* \mathbb{R}_- *, then one can take* $K_2 = 0$ *in* [\(5.7\)](#page-29-6) *to obtain*

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + C\left(t^{-(1-\sigma)}\mathcal{J}_1(t,x) + F^{-1}\left(Cp\,t^{1-\sigma}\right)\right);
$$

(ii) if the initial conditions μ_0 *and* μ_1 *are such that*

$$
\sup_{(t,x)\in\mathbb{R}_+\times\mathbb{R}^d} \mathcal{J}_0(t,x) < \infty,
$$

then there exist constant $C_* > 0$ *, such that for all* $(t, x, p) \in [1, \infty) \times \mathbb{R}^d \times [2, \infty)$ *,*

$$
||u(t,x)||_p^2 \le C_* F^{-1} \left(C_* p \, t^{1-\sigma} \right); \tag{5.8}
$$

(iii) if $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, then the solution *u* is a.s. η_1 –Hölder continuous in time and η_2 –Hölder *continuous in space on* $(0, \infty) \times \mathbb{R}^d$ *for all*

$$
\eta_1 \in \left(0, \frac{1-\sigma}{2}\right) \quad \text{and} \quad \eta_2 \in \left(0, 1 \wedge \frac{\theta}{2}\right),
$$

 $where \ \theta := 2a + \frac{a}{b} \min\{2\gamma - 1, 0\}$;

(iv) if $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, then with the same C_* as in part *(ii),* and L_t as in [\(1.19\)](#page-6-4) with $h(t) = t^{1-\sigma}$, for all $(t, x) \in [1, \infty) \times \mathbb{R}^d$ and $z \geq L_t$,

$$
\mathbb{P}\left(|u(t,x)| \geq z\right) \leq \exp\left(-C_*^{-1}t^{-(1-\sigma)}F\left(z^2/(C_*e^2)\right)\right),\,
$$

and with some universal constant $C' > 0$,

$$
\sup_{|x| \le R} u(t,x) \lesssim \sqrt{F^{-1}\Big(C' \, t^{1-\sigma} \log R\Big)} \,, \quad a.s., \; as \; R \to \infty.
$$

Proof. The proof of parts (i) and (ii) is similar to the proof of Theorem [1.4.](#page-4-0) We only need to point out some differences. In this case, the function $h(t)$ should be computed as follows (thanks to Lemma 5.4 :

$$
h(t) := \int_0^t ds \int_{\mathbb{R}^d} dy \, Y^2(s, y) \le \int_0^t ds \, \ell(s) = Ct^{1-\sigma}.
$$

Now, we apply the same argument as in the proof of Theorem [1.4](#page-4-0) to obtain the following inequality analogous to (3.15) :

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8pK_M^2 \int_0^t ds \int_{\mathbb{R}^d} dy Y(t-s,x-y)^2
$$

+8p $\int_0^t ds \int_{\mathbb{R}^d} dy Y(t-s,x-y)^2 \rho_2 \left(M^2 + ||u(s,y)||_p^2\right).$

where K_M is defined in (1.14) . Using Lemma [5.4,](#page-29-4) we can further deduce that

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + \frac{8C_0 p K_M^2 t^{1-\sigma}}{1-\sigma} + 8C_0 p \int_0^t ds \,(t-s)^{-\sigma} \int_{\mathbb{R}^d} dy \,\mathcal{G}(t-s,x-y)\rho_2 \left(M^2 + ||u(s,y)||_p^2\right).
$$
\n(5.9)

Performing the convolution on both sides of [\(5.9\)](#page-30-0) with $s^{-\sigma} \mathcal{G}(s, z)$ on $[0, t] \times \mathbb{R}^d$, and using the sub-semigroup property of \mathcal{G} , we deduce that

$$
\int_0^t ds \, (t-s)^{-\sigma} \int_{\mathbb{R}^d} dz \, \mathcal{G}(t-s, x-z) \, \|u(s,z)\|_p^2
$$

\n
$$
\leq 2 \int_0^t ds \, (t-s)^{-\sigma} \int_{\mathbb{R}^d} dz \, \mathcal{G}(t-s, x-z) \mathcal{J}_0(s,z)^2 + \frac{8C_0 K_M^2 B (1-\sigma, 2-\sigma) p \, t^{2-2\sigma}}{1-\sigma}
$$

\n
$$
+ 8C_0 C_1 p \, t^{1-\sigma} \int_0^t ds \, (t-s)^{-\sigma} \int_{\mathbb{R}^d} dz \, \mathcal{G}(t-s, x-z) \rho_2 \left(M^2 + \|u(v,z)\|_p^2 \right),
$$

where $B(\cdot, \cdot)$ is the beta function. As a result, we can write the next inequality analogous to (3.21) ,

$$
X \le M^2 + 2t^{-(1-\sigma)} \mathcal{J}_1(t,x) + \frac{8C_0 K_M^2 B(1-\sigma, 2-\sigma)}{1-\sigma} p t^{1-\sigma} + 8C_0 C_1 p t^{1-\sigma} \rho_2(X),
$$

with

$$
X := t^{-(1-\sigma)} \int_0^t ds \, (t-s)^{-\sigma} \int_{\mathbb{R}^d} dz \, \mathcal{G}(t-s, x-z) \, ||u(s, z)||_p^2.
$$

The rest proof for parts (i) and (ii) of the theorem employs the same arguments as in Theorem [1.4.](#page-4-0)

Suppose that $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, then for any $x, y \in \mathbb{R}^d$, and $p \ge 2$,

$$
||u(t,x) - u(t,y)||_p^2 \le 8p \int_0^t ds \int_{\mathbb{R}^d} dz |Y(t-s,x-z) - Y(t-s,y-z)|^2 \rho_2 \left(M^2 + ||u(s,z)||_p^2\right)
$$

$$
\le Cp \int_0^t ds \int_{\mathbb{R}^d} dz |Y(t-s,x-z) - Y(t-s,y-z)|^2 \left(1 + F^{-1}(C_*ps^{1-\sigma})\right)
$$

$$
\le C_{p,t} \left(1 + F^{-1}(C_*pt^{1-\sigma})\right) |x-y|^\theta,
$$
 (5.10)

where the last two inequalities are consequence of (5.7) , parts (i) and (ii) of Hypothesis [1.2,](#page-4-2) and [\[CHN19,](#page-45-7) Proposition 5.4], and $C_{t,p} > 0$ depending on p and t. This proves the Hölder continuity of u in space. The Hölder continuity in time can be shown similarly, which is skipped for conciseness. This completes the proof of part (iii).

Finally, with [\(5.8\)](#page-30-1) and [\(5.10\)](#page-31-1) and observation that $1 + F^{-1}(C_*pt^{1-\sigma}) \leq CF^{-1}(C_*pt^{1-\sigma})$ for all $t \geq 1$ with some universal constant $C > 0$, the proof for part (iv) follows the same lines presented as in the proofs of Theorems 1.5 and 1.8. The proof of Theorem 5.5 is complete. as in the proofs of Theorems [1.5](#page-6-0) and [1.8.](#page-6-2) The proof of Theorem [5.5](#page-29-7) is complete.

Remark 5.6. Here are two comments: (1) From the proof of Theorem [5.5,](#page-29-7) it is evident that the key to the conclusion lies in the fact that the fundamental solution *Y* satisfies Assumption [5.3,](#page-29-1) which is ensured by Lemma [5.4.](#page-29-4) There may be other equations whose mild formulation can be written as in Definition [5.1,](#page-28-1) with *Y* fulfilling Assumption [5.3,](#page-29-1) and for which a moment bound can be deduced using the same approach. (2) For spatially colored noise, it is not clear to us how to formulate a similar factorization formula for the reference kernel. Thus, directly applying our proof becomes non-trivial. We conjecture that the moment upper bound can also be established and leave it as an open problem for interested readers to explore.

5.2 One-dimensional stochastic wave equations

In this part, we consider the one-dimensional *stochastic wave equation* (SWE) driven by the spatial homogeneous noise that is white in time:

$$
\begin{cases}\n\frac{\partial^2}{\partial t^2}u(t,x) = \Delta u(t,x) + \rho(u(t,x))\dot{W}(t,x), & (t,x) \in \mathbb{R}_+ \times \mathbb{R}, \\
u(0,\cdot) = \mu_0, & \frac{\partial}{\partial t}u(t,x)\Big|_{t=0} = \mu_1;\n\end{cases}
$$
\n(5.11)

see, e.g., [\[DM09](#page-46-14); [CD15a;](#page-45-9) [BC16](#page-45-10); [HW21](#page-47-14)], for related results about solutions to SWEs in the linear regime. The solution to [\(5.11\)](#page-31-2) is formulated in the mild form:

$$
u(t,x) = \mathcal{J}_0(t,x) + \int_0^t \int_{\mathbb{R}} G(t-s,x-y) \rho(u(s,y)) W(ds,dy),
$$

with

$$
\mathcal{J}_0(t,x) := \frac{1}{2} \left[\mu_0(x+t) + \mu_0(x-t) \right] + \int_{\mathbb{R}} \mu_1(\mathrm{d}y) G(t,x-y),\tag{5.12}
$$

where, throughout the rest part of this section, $G(t, x) = \frac{1}{2} \mathbb{I}_{[-t,t]}(x)$ refers to the wave kernel on R. Our arguments for the main results Theorem [1.4](#page-4-0) can be extended to this case too:

Theorem 5.7. Let $u(t, x)$ be a solution to the SWE [\(5.11\)](#page-31-2) with initial position $\mu_0 \in L^2_{loc}(\mathbb{R})$ and *initial velocity* μ_1 *which is a locally finite Borel measure on* R. Under Hypothesis [1.1](#page-3-2) (with $d = 1$) *and Hypothesis [1.2,](#page-4-2) it holds that*

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + K_1\left(h(t)^{-1}\mathcal{J}_1(t,x) + K_2p\,h(t) + F^{-1}(2p\,h(t))\right),\tag{5.13}
$$

for all $(t, x, p) \in \mathbb{R}_+ \times \mathbb{R} \times [2, \infty)$ *, where* \mathcal{J}_0 *and* F^{-1} *are given in* [\(5.12\)](#page-32-0) *and* [\(3.8\)](#page-13-1)*, respectively,*

$$
\mathcal{J}_1(t,x) := \int_0^t \mathrm{d}s \iint_{\mathbb{R}^2} \mathrm{d}y \mathrm{d}y' \, G\left(t-s, x-y\right) G\left(t-s, x-y'\right) f(y-y') \mathcal{J}_0^2(s, y),\tag{5.14}
$$

and

$$
h(t) := \int_0^t ds \iint_{\mathbb{R}^2} dy dy' G(s, y) G(s, y') f(y - y'). \tag{5.15}
$$

In [\(5.13\)](#page-32-1)*, K*¹ *and K*² *are some positive constants not depending on t and p. In particular,*

- *(i) if* $|\rho(\cdot)|$ *is concave separately on* \mathbb{R}_+ *and* \mathbb{R}_- *, then one can take* $K_2 = 0$ *in* [\(5.13\)](#page-32-1)*;*
- *(ii) there exist some constants* $C > 0$ *such that the moment bound* [\(1.16\)](#page-5-1) *holds true, where* \mathcal{J}_0 *,* \mathcal{J}_1 *, and h(t) are given in* [\(5.12\)](#page-32-0)*,* [\(5.14\)](#page-32-2)*, and* [\(5.15\)](#page-32-3)*, respectively. Moreover, if* μ_0 *is a bounded function and* $|\mu_1|(\mathbb{R}) < \infty$, then the moment bound in [\(1.16\)](#page-5-1) can be simplified to [\(1.17\)](#page-5-4) with *h*(*t*) *given in* [\(5.15\)](#page-32-3);
- *(iii)* assuming that $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, the solution $u(t, x)$ to [\(5.11\)](#page-31-2) has a version which is *a.s.* η_1 –Hölder continuous in time and η_2 –Hölder continuous in space on $(0, \infty) \times \mathbb{R}$ for all $\eta_1, \eta_2 \in (0, \eta)$ *, where* η *is given in* [\(1.20\)](#page-6-6)*;*
- *(iv)* assuming that $\mu_0 \equiv 1$ *and* $|\mu_1| \equiv 0$ *, with the same* C_* *as in part (ii) and* L_t *given in* [\(1.19\)](#page-6-4) *but with* $h(t)$ *replaced by* [\(5.15\)](#page-32-3)*, then for all* $(t, x) \in [T, \infty) \times \mathbb{R}^d$ *and* $z \geq L_t$ *,*

$$
\mathbb{P}\left(|u(t,x)|\geq z\right)\leq \exp\left(-(C_*h(t))^{-1}F\left(z^2/(C_*e^2)\right)\right),
$$

and under Hypothesis [1.6,](#page-6-5) with some universal constant $C' > 0$,

$$
\sup_{|x| \le R} u(t,x) \lesssim \sqrt{F^{-1}\Big(C'\,h(t)\log R\Big)}\,, \quad a.s.,\; as\; R \to \infty.
$$

We first prove two lemmas.

Lemma 5.8. *The wave kernel function* $G(\cdot, \circ)$ *satisfies the following properties:*

(i) For all $t, s \in \mathbb{R}_+$ and $x, y \in \mathbb{R}$,

$$
G(t, x - y)G(s, y) = 2G(t, x - y)G(s, y)G(t + s, x) \le G(s, y)G(t + s, x); \tag{5.16}
$$

(ii) For all $t > 0$ and $x \in \mathbb{R}$,

$$
tG(t,x) \le G(2t,x)(2t-|x|) = 2\left(G(t,\cdot) * G(t,\cdot)\right)(x) \le 2tG(2t,x); \tag{5.17}
$$

(iii) If the correlation function $f : \mathbb{R} \to \mathbb{R}_+$ satisfies Hypothesis [1.1,](#page-3-2) then for all $t > 0$,

$$
\sup_{(s,x)\in[0,t]\times\mathbb{R}}\int_{\mathbb{R}}\mathrm{d}z\,G\left(s,x+z\right)f(z)\leq 2k(2t)<\infty,\tag{5.18}
$$

where $k(t)$ *is defined as in* [\(3.13\)](#page-13-12) *with* $p_t(x)$ *replaced by* $G(t, x)$ *;*

(*iv*) $G(\cdot, \circ)^2$ satisfies the sub-semigroup property (see Definition [5.3\)](#page-29-1) with

$$
\mathcal{G}(t,x) = \frac{1}{2t} \mathbb{I}_{[-t,t]}(x) = \frac{1}{t} G(t,x), \quad \ell(t) = \frac{t}{2}, \quad \text{and} \quad C^* = 2.
$$

Proof. Part (i) is due to the following identity: for any $s, t \in \mathbb{R}_+$ and $x, y \in \mathbb{R}$,

$$
\mathrm{1\!I}_{[-t,t]}(x-y)1\mathrm{I}_{[-s,s]}(y)=\mathrm{1\!I}_{[-t,t]}(x-y)1\mathrm{I}_{[-s,s]}(y)1\mathrm{I}_{[-(s+t),t+s]}(x).
$$

Part (ii) can be obtained by direct computation. As for part (iii), by the nonnegativity of *f* and non-decreasing property of $t \to G(t, x)$ with $x \in \mathbb{R}$ fixed, we see that

$$
\sup_{(s,x)\in[0,t]\times\mathbb{R}}\int_{\mathbb{R}}\mathrm{d}z\,G(s,x+z)\,f(z)\leq \sup_{x\in\mathbb{R}}\int_{\mathbb{R}}\mathrm{d}z\,G(t,x+z)\,f(z)
$$

Next, thanks to the first inequality in [\(5.17\)](#page-32-4), for any $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$, we have that

$$
\int_{\mathbb{R}} dz G(t, x+z) f(z) \leq t^{-1} \int_{\mathbb{R}} dz G(2t, x+z) (2t - |x+z|) f(z)
$$

\n
$$
= \frac{2}{t} \iint_{\mathbb{R}^2} dy dz G(t, x+y) G(t, y-z) f(z),
$$

\n
$$
= \frac{1}{\pi t} \int_{\mathbb{R}} e^{ix} \left| \frac{\sin(t\xi)}{\xi} \right|^2 \hat{f}(d\xi) \leq \frac{1}{\pi t} \int_{\mathbb{R}} \left| \frac{\sin(t\xi)}{\xi} \right|^2 \hat{f}(d\xi)
$$

\n
$$
= \frac{2}{t} \iint_{\mathbb{R}^2} dy dy' G(t, y) G(t, y') f(y - y').
$$

Then apply the second inequality in [\(5.17\)](#page-32-4) to the above upper bound to see that

$$
\int_{\mathbb{R}} dz G(t, x+z) f(z) \leq 2 \int_{\mathbb{R}} dz G(2t, z) f(z) = 2k(2t).
$$

Notice that thanks to Dalang's condition [\(1.8\)](#page-3-1),

$$
\frac{1}{\pi t} \int_{\mathbb{R}} \left| \frac{\sin(t\xi)}{\xi} \right|^2 \hat{f}(\mathrm{d}\xi) \le C_t \int_{\mathbb{R}} \frac{\hat{f}(\mathrm{d}\xi)}{1+|\xi|^2} < \infty, \quad \text{for all } t > 0.
$$

This proves part (iii).

Finally, as for part (iv), it is straightforward to verify that $\int_{\mathbb{R}} \mathcal{G}(t,x) dx = 1$ and $G(t,x)^2 =$ $\ell(t)G(t, x)$. Moreover,

$$
\int_{\mathbb{R}} \mathcal{G}(t, x - y) \mathcal{G}(s, y) dy = \frac{\mathbb{I}_{[-(t+s), t+s]}(x)}{4st} \int_{\mathbb{R}} \mathbb{I}_{[-t,t]}(x - y) \mathbb{I}_{[-s,s]}(y) dy \n= \frac{\mathbb{I}_{[-(t+s), t+s]}(x)}{4st} \times \text{Leb} \left([-t+x, t+x] \cap [-s, s] \right) \n\leq \frac{2(s \wedge t) \mathbb{I}_{[-(t+s), t+s]}(x)}{4st} = \frac{\mathbb{I}_{[-(t+s), t+s]}(x)}{2(s \vee t)} \leq \frac{\mathbb{I}_{[-(t+s), t+s]}(x)}{s+t} = 2\mathcal{G}(t+s, x).
$$

Thus, the sub-semigroup relation in (5.4) holds with $C^* = 2$. This proves part (iii).

 \Box

Thanks to part (iv) of Lemma [5.8,](#page-32-5) we can obtain the moment bounds using the same arguments as those in Theorem [5.5.](#page-29-7) It is important to note, however, that as discussed in Remark [5.6,](#page-31-3) the method in Theorem [5.5](#page-29-7) is limited to the space-time white noise. To get the desired result for the spatial colored noises, we need the following result, which serves as the analog of Lemma [3.9.](#page-13-3)

Lemma 5.9. *Under the same setting as Theorem [5.7,](#page-32-6)* $\mathcal{J}_1(t,x)$ *defined in* [\(5.14\)](#page-32-2) *is finite for all* $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$ *. In particular, for every* $(t, x) \in \mathbb{R}_+ \times \mathbb{R}$ *,*

$$
\mathcal{J}_1(t,x) \leq 4tk(2t)\left(t\int_{\mathbb{R}}G(t,x-y)\mu_0^2(y)\mathrm{d}y + 2\left(t\int_{\mathbb{R}}|\mu_1|(\mathrm{d}y)G(t,x-y)\right)^2\right) < \infty,
$$

where $k(t)$ *is defined in* [\(3.13\)](#page-13-12) *with* $p_t(x)$ *replaced by* $G(t, x)$ *.*

Proof. We decompose $\mathcal{J}_0(t,x)$ in [\(5.12\)](#page-32-0) into two parts:

$$
\mathcal{J}_{0,1}(t,x) := \frac{1}{2} [\mu_0(x+t) + \mu_0(x-t)] \text{ and } \mathcal{J}_{0,2}(t,x) := \int_{\mathbb{R}} \mu_1(\mathrm{d}y) G(t,x-y).
$$

Then, by change of variable,

$$
\mathcal{J}_{1,1}(t,x) := \int_0^t \mathrm{d}s \iint_{\mathbb{R}^2} \mathrm{d}y \mathrm{d}y' \, G\left(t-s, x-y\right) G\left(t-s, x-y'\right) f(y-y') \mathcal{J}_{0,1}(s,y)^2
$$

=
$$
\int_0^t \mathrm{d}s \iint_{\mathbb{R}^2} \mathrm{d}y \mathrm{d}z \, G\left(t-s, x-y\right) G\left(t-s, x-y+z\right) f(z) \mathcal{J}_{0,1}(s,y)^2.
$$

Using inequality (5.18) , we have

$$
\mathcal{J}_{1,1}(t,x) \le 2k(2t) \int_0^t ds \int_{\mathbb{R}} dy \, G\left(t-s, x-y\right) \mathcal{J}_{0,1}(s,y)^2 =: 2k(2t) \Theta_1(t).
$$

Notice that

$$
\Theta_1(t) \le \int_0^t ds \int_{\mathbb{R}} dy G(t-s, x-y) \left(\mu_0^2(y+s) + \mu_0^2(y-s) \right)
$$

=
$$
\int_{\mathbb{R}} dz \mu_0^2(z) \int_0^t ds \left[G(t-s, x-z+s) + G(t-s, x-z-s) \right].
$$

Because $s - |x - z| \le |x - z \pm s| \le t - s$, we see that the above ds-integral is bounded $t1\mathbf{1}_{\{|x - z| \le t\}} =$ $2tG(t, x-z)$. Hence, the condition $\mu_0 \in L^2_{loc}(\mathbb{R})$ implies that

$$
\mathcal{J}_{1,1}(t,x) \le 2tk(2t)\int_{x-t}^{x+t} \mu_0^2(z)dz = 4t\,k(2t)\int_{\mathbb{R}} dz\,G(t,x-z)\mu_0^2(z) < \infty.
$$

It remains to show that

$$
\mathcal{J}_{1,2}(t,x) \coloneqq \int_0^t ds \int_{\mathbb{R}^2} dy dy' \, G\left(t-s, x-y\right) G\left(t-s, x-y'\right) f(y-y') \mathcal{J}_{0,2}(s, y)^2 < \infty.
$$

Thanks to [\(5.18\)](#page-33-0), we see that

$$
\mathcal{J}_{1,2}(t,x) \le 2k(2t) \int_0^t ds \int_{\mathbb{R}} dy G(t-s,x-y) \mathcal{J}_{0,2}(s,y)^2 =: 2k(2t) \Theta_2(t).
$$

By the Minkowski inequality with respect to the d*y*-integral, we can write

$$
\Theta_2(t) = 2 \int_{\mathbb{R}} \left[G(t-s, x-y) \mathcal{J}_{0,2}(s, y) \right]^2 dy \leq 2 \left[\int_{\mathbb{R}} \left(\int_{\mathbb{R}} G(t-s, x-y) G(s, y-z)^2 dy \right)^{1/2} |\mu_1|(dz) \right]^2.
$$

Due to [\(5.16\)](#page-32-7), we see that $\int_{\mathbb{R}} G(t - s, x - y) G(s, y - z)^2 dy \leq \frac{s}{4}$ $\frac{s}{4}$ **I**_[-t,t]($x - z$). Therefore, $\Theta_2(t) \leq$ *s* $rac{s}{2}$ \int_{x-t}^{x+t} $\int_{x-t}^{x+t} |\mu_1|(\mathrm{d}z)\Big)^2$ and

$$
\mathcal{J}_{1,2}(t,x) \le 2t^2 k(2t) \left(\int_{x-t}^{x+t} |\mu_1| (dz) \right)^2 = 8t^2 k(2t) \left(\int_{\mathbb{R}} |\mu_1| (dz) G(t, x-z) \right)^2 < \infty.
$$
\nn
\nthe proof of Lemma 5.9.

This completes the proof of Lemma [5.9.](#page-34-0)

Now we are ready to prove Theorem [5.7.](#page-32-6)

Proof of Theorem [5.7.](#page-32-6) It suffices to provide the proof for part (i), which follows a similar approach to that of Theorem [1.4,](#page-4-0) with one notable difference arising from the absence of the corresponding formula (3.19) for the heat equation case. However, we can leverage (5.16) to overcome this obstacle and conclude

$$
\int_{r}^{t} ds \iint_{\mathbb{R}^{2}} dy dy' G(t-s, x-y)G(t-s, x'-y')G(s-r, y)G(s-r, y')f(y-y')
$$

\n
$$
\leq G(t-r, x)G(t-r, x') \int_{0}^{t-r} ds \iint_{\mathbb{R}^{2}} dy dy' G(s, y)G(s, y')f(y-y')
$$

\n
$$
= G(t-r, x)G(t-r, x')h(t-r) \leq G(t-r, x)G(t-r, x')h(t).
$$

The last sequence of inequalities play the same role as those in [\(3.20\)](#page-15-3). Following the same idea as in the proof of Theorem [1.4,](#page-4-0) we can show that

$$
X \le M^2 + 2h(t)^{-1} \mathcal{J}_1(t, x) + 8K_M^2 p h(t) + 8ph(t)\rho_2(X),
$$

where K_M is given in (1.14) and

$$
X := M^2 + h(t)^{-1} \int_0^t ds \iint_{\mathbb{R}^2} dy dy' G(t-s, x-y) G(t-s, x-y') f(y-y') ||u(s, y)||_p^2.
$$

The remaining proof follows the same line of reasoning as presented in Theorem [1.4,](#page-4-0) [1.5](#page-6-0) and [1.8,](#page-6-2) if one can show the Hölder continuity for solution(s) to (5.11) under the improved Dalang condition. In the following, we only outline the proof for spatial continuity, and the time continuity can be verified similarly. Under the assumption that $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, we can write

$$
||u(t,x) - u(t,y)||_p^2 \le C \int_0^t ds \iint_{\mathbb{R}^2} dz dz' f(z - z')|G(t - s, x - z) - G(t - s, y - z)|
$$

$$
\times |G(t - s, x - z') - G(t - s, y - z')| ||\rho(u(s, z))||_p ||\rho(u(s, z'))||_p
$$

$$
\le C \int_0^t ds \iint_{\mathbb{R}^2} dz dz' f(z - z')|G(t - s, x - z) - G(t - s, y - z)|
$$

$$
\times |G(t - s, x - z') - G(t - s, y - z')| (1 + F^{-1}(C_* ph(s)))^2.
$$

Then, by using the same arguments as in $[DS05, Theorem 5]$, one finds that *u* is Hölder continuous in space with and exponent $\alpha < \eta$ and constant of the form $CF^{-1}(C_*ph(t))$. The proof of Theorem [5.7](#page-32-6) is complete.

Remark 5.10. Parts (iii) and (iv) of Theorem [5.7](#page-32-6) requires that the initial conditions $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, which is not necessary. They are used for simplification of the proof of the Hölder continuity of the solution. It can be relaxed to e.g. $\mu_0 \in H_2^{\eta}$ $\mathcal{P}_2^{\eta}(\mathbb{R})$ and $\mu_1 \in C(\mathbb{R})$ with $|\mu_1|(\mathbb{R}) < \infty$; see [\[DS05](#page-46-15)].

6 Applications

In this session, we will integrate the specific ρ studied in Section [4](#page-24-0) and the noise structure, as detailed in Appendix [A,](#page-40-0) to derive more precise moment bounds, as well as, tail probabilities and spatial asymptotics. In particular, we will demonstrate the transitions of properties from the additive SHE to the PAM in Section [6.1.](#page-36-1)

6.1 Moment growth, tail probability and spatial asymptotics for SHE

In this part, we showcase examples of the diffusion coefficient *ρ* and apply Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) and [1.8](#page-6-2) to derive moment bounds, tail probabilities, and spatial asymptotics for SHE [\(1.1\)](#page-1-2). We will use *C* to denote a generic constant that may change its value at each appearance, but does not depend on either *p* or *t*.

I. We first study the case when ρ is given in Proposition [4.1.](#page-24-2) We start with a simple example:

Proposition 6.1. *Under the setting of Theorem [1.4](#page-4-0) but with the following diffusion coefficient*

$$
\rho(u) = \frac{|u|}{1+|u|},
$$

there exists a unique solution u to SHE [\(1.1\)](#page-1-2) *such that for all* $t > 0$, $x \in \mathbb{R}^d$ *and* $p \ge 2$,

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8ph(t).
$$
\n(6.1)

Note that the uniqueness comes from the fact that ρ satisfies Lipschitz condition and the moment bound in [\(6.1\)](#page-36-3) coincides with the moment bounds for the SHE with additive noise, the proof of which is straightforward by noticing that

$$
||u(t,x)||_p^2 \le 2\mathcal{J}_0^2(t,x) + 8p \int_0^t ds \int_{\mathbb{R}^{2d}} dy dy' p_{t-s}(x-y) p_{t-s}(x-y') f(y-y').
$$

However, in case of the bounded initial condition and $t \geq 1$, one can also obtain the moment bound [\(6.1\)](#page-36-3) in the form of [\(1.17\)](#page-5-4) by using F^{-1} in [\(4.2\)](#page-24-3) with $\alpha = 0$: $F^{-1}(x) \le 8x$.

The next proposition examines the ρ given in Proposition [4.1](#page-24-2) in detail. We particularly focus on the impacts on the initial conditions on the growth of moments.

Proposition 6.2. *Let u be a solution to SHE* [\(1.1\)](#page-1-2) *under the same setting as Theorem [1.4,](#page-4-0) but with the diffusion coefficient* ρ *given in Proposition [4.1.](#page-24-2) Then, for all* $(p, t, x) \in [2, \infty) \times \mathbb{R}_+ \times \mathbb{R}^d$, *it holds that*

$$
||u(t,x)||_p^2 \le C \left[\mathcal{J}_0^2(t,x) + p h(t) + (p h(t))^{1/(1-\alpha)} + h(t)^{-1} \mathcal{J}_1(t,x) \right].
$$

Furthermore, thanks to Proposition [A.1,](#page-41-1) we have the following special cases:

(i) If $d = 1$ *and* $f = \delta$ *, then*

$$
||u(t,x)||_p^2 \le C \left(\mathcal{J}_0^2(t,x) + p\sqrt{t} + \left(p\sqrt{t} \right)^{1/(1-\alpha)} + \mathcal{J}_+^2(t/2,x) \right). \tag{6.2}
$$

Moreover, if μ *is a bounded function, then,*

$$
||u(t,x)||_p^2 \le C \left(p\sqrt{t}\right)^{1/(1-\alpha)}, \quad t \ge 1.
$$

(ii) If $d = 1$ *and* $f(x) = |x|^{-\beta}$ *with* $\beta \in (0, 1)$ *, then*

$$
||u(t,x)||_p^2 \le C\bigg(\mathcal{J}_0^2(t,x) + pt^{1-\beta/2} + \left(pt^{1-\beta/2}\right)^{1/(1-\alpha)} + \mathcal{J}_+^2(t/2,x)\bigg).
$$

Again, if μ *is bounded, then,*

$$
||u(t,x)||_p^2 \le C \left(p t^{1-\beta/2} \right)^{1/(1-\alpha)}, \quad t \ge 1.
$$

(*iii*) If $f = |x|^{-\beta}$ *with* $\beta \in (0, 2 \wedge d)$ *, then for all* $d \ge 1$ *it holds that*

$$
||u(t,x)||_p^2 \le C \left(\mathcal{J}_0^2(t,x) + pt^{1-\beta/2} + \left(pt^{1-\beta/2} \right)^{1/(1-\alpha)} + t^{-1+\beta/2} \mathcal{J}_1(t,x) \right). \tag{6.3}
$$

Moreover,

(a) if $\mu(x) = |x|^{-\ell}$ *with* $\ell \in (0, 2 \land d)$ *, then we have*

$$
||u(t,x)||_p^2 \le C\left(t^{-\ell} + \left(p\,t^{1-\beta/2}\right)^{1/(1-\alpha)} + p\,t^{1-\beta/2}\right);
$$
\n(6.4)

(b) and if $\mu(x) = e^{\ell|x|}$ with $\ell \in \mathbb{R}$, then with some universal constants C_1 and $C_2 > 0$ it holds *that*

$$
||u(t,x)||_p^2 \le C\left(e^{C_1\ell^2 t + C_2\ell|x|} + pt^{1-\beta/2} + \left(pt^{1-\beta/2}\right)^{1/(1-\alpha)}\right). \tag{6.5}
$$

The proof of this proposition is essentially an application of Proposition [4.1](#page-24-2) together with Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) [1.8,](#page-6-2) and Proposition [A.1.](#page-41-1) Some more details regarding the role of initial conditions are given in Appendix [B.](#page-44-0)

Remark 6.3. (1) It is clear that in Case (iii)–(b) of Proposition [6.2,](#page-36-2) if $\ell > 0$, as $t \to \infty$, $||u(t, x)||_p^2$ *p* is bounded by an exponent function of t , which is dominated by the explosion rate of μ at infinity, and the stochastic fluctuation makes fewer contributions in the moment bounds. (2) When $r > 0$, ρ is globally Lipschitz continuous (see part 1 of Proposition [4.1\)](#page-24-2) and the solution is unique.

An application of Proposition [4.1](#page-24-2) together with Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) and [1.8](#page-6-2) yields the following:

Proposition 6.4. *Under the same setting as Proposition [6.2,](#page-36-2) but with bounded the initial condition. Then for all* $p \geq 2$, $t \geq 1$ *and* $x \in \mathbb{R}^d$, *the following statements hold:*

$$
||u(t,x)||_p^2 \lesssim (ph(t))^{1/(1-\alpha)}, \qquad \text{as } p \lor t \to \infty,
$$
 (6.6)

$$
\log \mathbb{P}\left(|u(t,x)| \geq z\right) \lesssim -\frac{z^{2(1-\alpha)}}{h(t)}, \qquad \text{as } z \to \infty,
$$
\n(6.7)

$$
\sup_{|x| \le R} u(t, x) \lesssim [h(t) \log R]^{1/(2(1-\alpha))}, \qquad a.s. \text{ as } R \to \infty.
$$
 (6.8)

Remark 6.5. In comparison to the tail estimates for super Brownian motions found in $[Hu+23,$ Proposition 1.4], the tail estimate given by (6.7) is sharp for the case when $\alpha = 1/2$. Since when $\alpha = 0$, the bound in [\(6.8\)](#page-37-1) coincides with the exact asymptotics demonstrated in [\[CJK13,](#page-46-0) Theorem 1.2] and [\[Con+13](#page-46-1), Theorem 2.3], we believe that the bound in [\(6.8\)](#page-37-1) is sharp for all $\alpha \in [0,1)$.

II. Now we study the case when ρ is given in Proposition [4.2.](#page-25-0) An application of Proposition [4.2](#page-25-0) together with Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) [1.8,](#page-6-2) and Proposition [A.1](#page-41-1) $(h(t) = \sqrt{t/\pi})$ yields part (a) of the following proposition:

Proposition 6.6. *Let u be a solution to SHE* [\(1.1\)](#page-1-2) *under the setting that* $d = 1$, $\mu \equiv 1$, $f = \delta$ *, and the diffusion coefficient* ρ *given in Proposition* [4.2.](#page-25-0) *Then, for all* $p > 2$, $t > 1$ *, and* $x \in \mathbb{R}$ *, the following statements hold:*

(a) In cases (i) and (ii) in [\(1.5\)](#page-2-3), namely, $(\alpha, \beta) \in (0, 1) \times \mathbb{R}$ or $(\alpha, \beta) \in \{0\} \times (-\infty, 0)$,

$$
||u(t,x)||_p^2 \lesssim \left(p\sqrt{t}\right)^{1/(1-\alpha)} \left[\log\left(p\sqrt{t}\right)\right]^{-2\beta/(1-\alpha)}, \qquad \text{as } p \vee t \to \infty,\tag{6.9}
$$

$$
\log \mathbb{P}\left(|u(t,x)| \geq z\right) \lesssim -z^{2(1-\alpha)}t^{-1/2}\left[\log z\right]^{2\beta}, \qquad \text{as } z \to \infty,
$$
\n(6.10)

$$
\sup_{|x| \le R} u(t, x) \lesssim \left[\sqrt{t} \, \log R \right]^{\frac{1}{2(1-\alpha)}} \left[\log \left(\sqrt{t} \, \log R \right) \right]^{-\frac{\beta}{1-\alpha}}, \quad a.s. \text{ as } R \to \infty. \tag{6.11}
$$

(b) In case (iii) in [\(1.5\)](#page-2-3), namely, $\alpha = 1$ and $\beta > 0$, by setting $\beta^* := \beta \wedge (1/4)$,

$$
||u(t,x)||_p^2 \lesssim \exp\left(C\left(p^2t\right)^{1/(4\beta^*)}\right), \qquad \text{as } p \lor t \to \infty,
$$
 (6.12)

$$
\log \mathbb{P}\left(u(t,x) \ge z\right) \lesssim -t^{-1/2} \left[\log z\right]^{1+2\beta^*}, \qquad \text{as } z \to \infty,
$$
\n(6.13)

$$
\sup_{|x| \le R} u(t, x) \lesssim \exp\left(C\left(\sqrt{t} \log R\right)^{1/(1+2\beta^*)}\right), \qquad a.s. \text{ as } R \to \infty. \tag{6.14}
$$

Proof of part (b) of Proposition [6.6.](#page-38-0) For [\(6.12\)](#page-38-1), if we apply $F^{-1}(\cdot)$ given in [\(4.4\)](#page-25-2) to the moment formula (1.17) , we obtain

$$
||u(t,x)||_p^2 \leq C \exp\left(C\left(p^2 t\right)^{1/(4\beta)}\right).
$$

But by invoking the moment comparison principle (see [\[JKM17](#page-47-5); [CK20\]](#page-46-8)), we find that the exponent of t cannot exceed that of the parabolic Anderson model. Therefore, we use β^* in the exponent of *t* in [\(6.12\)](#page-38-1).

The tail estimate in [\(6.13\)](#page-38-2) is better than that obtained by setting $\alpha = 1$ in [\(6.10\)](#page-38-3), the latter of which produces of the power $2\beta^*$ (instead of $2\beta^* + 1$), namely,

$$
\log \mathbb{P}(|u(t,x)| \ge z) \lesssim -t^{-1/2} \left(\log z\right)^{2\beta^*}, \quad \text{for all } z > 0 \text{ large enough.}
$$

Indeed, the tail probability in (6.13) can be obtained by directly computing $H(p)$ in (3.27) (with $CF^{-1}(Cph(t))$ replaced by the sharper moment upper bound in (6.12) and the corresponding Legendre-type transform $H^*(y)$ in (3.29) :

$$
H(p) = Cp^{\frac{1+2\beta^*}{2\beta^*}}t^{1/(4\beta^*)} \quad \Rightarrow \quad H^*(y) = \frac{C}{\sqrt{t}}y^{1+2\beta^*}, \quad \text{for large } y > 0,
$$

and then plugging *H*∗ (*y*) back to [\(3.26\)](#page-17-3). Finally, applying [\(6.13\)](#page-38-2) in the proof of Theorem [1.8](#page-6-2) proves [\(6.14\)](#page-38-4). \Box **III.** Here we study the case when ρ is given in Proposition [4.3.](#page-26-0) An application of Proposition [4.3](#page-26-0) together with Theorems [1.4,](#page-4-0) [1.5,](#page-6-0) [1.8,](#page-6-2) and Proposition [A.1](#page-41-1) $(h(t) = \sqrt{t/\pi})$ yields the following:

Proposition 6.7. *Suppose that* $d = 1$, $\mu \equiv 1$, $f = \delta$, and the diffusion coefficient ρ given in *Proposition* [4.3](#page-26-0) with $\kappa > 1$. Then there exists a unique solution *u* to SHE [\(1.1\)](#page-1-2) such that for all $p \geq 2$, $x \in \mathbb{R}$, and $t > 1$, the following statements hold:

$$
||u(t,x)||_p^2 \lesssim \exp\left(\exp\left\{\left[(2\beta)^{-1}\log\left(Cp\sqrt{t}\right)\right]^{1/\kappa}\right\}\right), \qquad \text{as } p \lor t \to \infty,
$$

$$
\log \mathbb{P}\left(u(t,x) \geq z\right) \lesssim -\frac{1}{\sqrt{t}} \exp\left\{2\beta\left[\log(2) + \log\left(\log z\right)\right]^\kappa\right\}, \qquad \text{as } z \to \infty,
$$

$$
\sup_{|x| \leq R} u(t,x) \lesssim \exp\left(\frac{1}{2}\exp\left\{\left[(2\beta)^{-1}\log\left(C\sqrt{t}\log R\right)\right]^{1/\kappa}\right\}\right), \qquad a.s. \text{ as } R \to \infty.
$$

We summarize the moment growth rates and the spatial asymptotics in Figure [6.1](#page-39-1) below.

Figure 6.1: Summary of results in Propositions [6.4,](#page-37-2) [6.6](#page-38-0) and [6.7](#page-39-2) in case of $d = 1$, space-time white noise $(f = \delta)$, and a constant initial condition $(\mu \equiv 1)$. The moment asymptotics are considered under the condition that $x \in \mathbb{R}$ is arbitrary but fixed and $t \to \infty$, while the spatial asymptotics are considered under the condition $R \to \infty$, with $t \geq 1$.

6.2 Moment bounds for other SPDEs

In the previous Section [6.1,](#page-36-1) we studied the combinations of various ρ with different types of noise for SHE (1.1) . One can readily construct concrete examples for the fractional SPDE (5.1) by considering various ρ , as provided in Propositions [4.1,](#page-24-2) [4.2,](#page-25-0) and [4.3](#page-26-0) (although only for the spacetime white noise case). Similarly, for the SWE (5.11) , thanks to Proposition [A.2,](#page-43-1) one can easily investigate examples with various noise structures as presented in Proposition [A.2](#page-43-1) and various ρ , such as those given in Propositions [4.1,](#page-24-2) [4.2,](#page-25-0) and [4.3.](#page-26-0) This application is straightforward, so we won't carry out the details here. Instead, we will consider one particular example below.

Let us consider the diffusion coefficient ρ given in Proposition [4.1](#page-24-2) (so that $F^{-1}(x) \leq Cx^{1/(1-\alpha)}$) and restrict ourselves to the one-dimensional space-time case (i.e., $f = \delta_0$ and $d = 1$).

1. Fractional SPDE [\(5.1\)](#page-27-3) with $a = 2$ and $\gamma = 0$: In this case, $\sigma = 2 - \frac{3b}{2}$. By Theorem [5.5,](#page-29-7) we can deduce the moment growth of $u(t, x)$ the solution to equation [\(5.1\)](#page-27-3) is as follows:

$$
||u(t,x)||_p^2 \le C \left(\mathcal{J}_0^2(t,x) + t^{1-3b/2} \mathcal{J}_1(t,x) + pt^{3\beta/2 - 1} + \left(pt^{3b/2 - 1} \right)^{1/(1-\alpha)} \right). \tag{6.15}
$$

2. SWE [\(5.11\)](#page-31-2): Let *u* be a solution to equation [\(5.11\)](#page-31-2). It follows from Theorem [5.7](#page-32-6) that

$$
||u(t,x)||_p^2 \le C\left(\mathcal{J}_0^2(t,x) + t^{-2}\mathcal{J}_1(t,x) + pt^2 + \left(pt^2\right)^{1/(1-\alpha)}\right). \tag{6.16}
$$

Taking account of Lemma [3.9,](#page-13-3) we find that inequalities (6.2) and (6.16) are special cases of inequality (6.15) when $b = 1$ and $b = 2$, respectively. This aligns with the interpolation characteristic of the stochastic heat and wave equation; see [\[CE22a](#page-45-11); [CGS22](#page-45-8)] for more instances. Furthermore, with the initial condition $\mu_0 \equiv 1$ and $\mu_1 \equiv 0$, we can write the tail estimate and spatial asymptotics (for $t > 1$ and $x \in \mathbb{R}$):

1. for the solution to (5.1) :

$$
\log \mathbb{P}\left(|u(t,x)| \geq z\right) \lesssim -t^{1-3b/2} z^{2(1-\alpha)}, \quad \text{as } z \to \infty
$$

and

$$
\sup_{|x| \le R} u(t, x) \lesssim \left(t^{3b/2 - 1} \log R \right)^{1/(2(1-\alpha))}, \quad \text{a.s., as } R \to \infty;
$$

2. for the solution to (5.11) :

$$
\log \mathbb{P}\left(|u(t,x)| \geq z\right) \lesssim -t^{-2}z^{2(1-\alpha)}, \quad \text{as } z \to \infty
$$

and

$$
\sup_{|x| \le R} u(t,x) \lesssim \left(t^2 \log R \right)^{1/(2(1-\alpha))}, \quad \text{a.s., as } R \to \infty.
$$

A Asymptotic behaviors of *h*(*t*)

In this section, we present asymptotic behaviors of *h* as in Theorem [1.4](#page-4-0) at infinity with some wellknown correlation functions in Appendix [A.1;](#page-41-0) see [\[CK19](#page-46-6), Examples 1.2-1.5] and [\[CE22b,](#page-45-3) Section 5.4]. Similar asymptotic behaviors of *h* for the SWE (defined as in Theorem [5.7\)](#page-32-6) are presented in Appendix [A.2](#page-43-0)

A.1 Stochastic heat equations

Proposition A.1. *Let h*(*t*) *be defined in* [\(1.9\)](#page-3-3)*. The following statements hold:*

(i) (*Riesz Kernel) If* $f(x) = |x|^{-\alpha}$ *with* $\alpha \in (0, 2 \wedge d)$ *, then*

$$
h(t) = C t^{1-\alpha/2} \quad \text{with } C = \frac{\Gamma((d-\alpha)/2)}{2^{\alpha/2}(2-\alpha)\Gamma(1+d/2)}.
$$

(ii) (Ornstein-Uhlenbeck kernel) If $f(x) = \exp(-|x|^{\alpha})$ *with* $\alpha > 0$ *, then*

$$
h(t) \asymp \begin{cases} \sqrt{t}, & d = 1, \\ \log(t), & d = 2, \text{ as } t \to \infty, \text{ and } h(t) \asymp t \text{ as } t \downarrow 0, \\ 1, & d \ge 3, \end{cases}
$$
 (A.1)

and in particular, when $\alpha = 2$,

$$
h(t) = \begin{cases} \sqrt{2t+1} - 1 & \text{if } d = 1, \\ 4^{-1} \log(1+2t) & \text{if } d = 2, \\ [(d-2)d]^{-1} \left(1 - (1+2t)^{1-d/2}\right) & \text{if } d \ge 3. \end{cases}
$$
(A.2)

- *(iii) (Brownian case:* $W(t, x) = W(t)$ *)* If $f(x) \equiv 1$ *, then* $h(t) = t$ *for all* $t \ge 0$ *.*
- *(iv) (One-dimensional space-time white case)* If $d = 1$ *and* $f(x) = \delta(x)$ *, then* $h(t) = \sqrt{t/\pi}$ *.*
- *(v)* (Bessel potential; see [\[Gra14](#page-47-15), Section 1.2.2], aka. Matérn correlation family; see [\[GG06\]](#page-47-16)) *Suppose that*

$$
f_{\nu}(x) = \int_{\mathbb{R}^d} \frac{e^{-ix\cdot\xi}}{(1+|\xi|^2)^{\nu/2}} d\xi, \quad \nu > 0.
$$
 (A.3)

Then,

$$
h(t) \asymp \begin{cases} \sqrt{t}, & d = 1, \\ \log(t), & d = 2, \\ 1, & d \ge 3, \end{cases} \quad \text{as } t \to \infty, \quad \text{and} \quad h(t) \asymp t \quad \text{as } t \downarrow 0. \tag{A.4}
$$

(Note that the large time asymptotics is the same as the Ornstein-Uhlenbeck kernel because $f_{\nu}(x)$ *behaves like* $e^{-\frac{|x|}{2}}$ *for* $|x| \geq 2$ *; see* [*Gra14, Proposition 1.2.5]*).

(vi) (Bessel potential as the spectral measure) Suppose that $f_{\nu}(x) = (1 + |x|^2)^{-\nu/2}$ with $\nu > 0$. *Then*

$$
h(t) \asymp \begin{cases} t^{1-\frac{\nu \wedge d}{2}}, & \nu \wedge d < 2, \\ \log(t), & \nu \wedge d = 2, \\ 1, & \nu \wedge d > 2, \end{cases} \text{ as } t \to \infty, \text{ and } h(t) \asymp t \text{ as } t \downarrow 0. \tag{A.5}
$$

Proof. Parts (i), (iii), (iv) and [\(A.2\)](#page-41-2) are from Examples 1.2–1.5 in Appendix of [\[CK19](#page-46-6)]. Moreover, the asymptotics at 0 in $(A.1)$, $(A.4)$ and $(A.5)$ can be formulated by evaluating the following limit by L'Hôpital's rule,

$$
\lim_{t\downarrow 0}h(t)/t=2\lim_{t\downarrow 0}\langle p_{2t},f\rangle=2f(0),
$$

for all *f* that is locally continuously bounded and has growth of order $o(e^{\epsilon x^2})$ for all $\epsilon > 0$ at infinity.

Eq. [\(A.1\)](#page-41-3) *in part (ii):* Note that the correlation function is bounded by 1. Thus,

$$
k(t) = \int_{\mathbb{R}^d} dz p_t(z) f(z) \le \int_{\mathbb{R}^d} dz p_t(z) = 1.
$$

On the other hand, using the polar coordinates, we can write

$$
k(t) = Ct^{-\frac{d}{2}} \int_{\mathbb{R}} dx |x|^{d-1} \exp\left(-\frac{x^2}{2t} - |x|^\alpha\right) \le Ct^{-\frac{d}{2}} \int_{\mathbb{R}} dx |x|^{d-1} \exp(-|x|^\alpha) \le Ct^{-\frac{d}{2}}.
$$

Therefore, for $t > 1$

$$
h(t) = \int_0^{2t} ds \, k(s) \le \int_0^1 ds + C \int_1^{2t} ds \, s^{-d/2} \le \begin{cases} C\left(1+\sqrt{t}\right), & d=1, \\ C\left(1+\log t\right), & d=2, \\ C\left(1+t^{1-d/2}\right), & d \ge 3. \end{cases}
$$

For the lower bound, using the polar coordinates again, we can deduce that for $t \geq 1$,

$$
h(t) \ge C \int_1^{2t} ds \, s^{-\frac{d}{2}} \int_{\mathbb{R}} dx \, |x|^{d-1} \exp\left(-\frac{x^2}{2} - |x|^\alpha\right) \ge C \int_1^{2t} ds \, s^{-d/2}.
$$

This completes the proof of the asymptotics of $h(\cdot)$ at infinity.

Part (v): From the proof of $[CE22b,$ Proposition 5.12], we have that

$$
k(2t) = h'(t) \approx U\left(\frac{d}{2}, \frac{2+d-\nu}{2}, t\right), \quad \text{as } t \to \infty,
$$

where *U* denotes the *confluent hypergeometric function*; see $[Olv+10$, Formula 13.4.4 on page 326. Using asymptotic behavior of $U(a, b, t)$ as $t \to \infty$ (see Formula 13.2.6 on page 322, *ibid.*), one obtains the large time asymptotics in [\(A.4\)](#page-41-4).

Part (vi): From the proof of [\[CE22b](#page-45-3), Proposition 5.13] and by the asymptotic analysis for *U* as in the previous case, we deduce that

$$
k(2t) \asymp t^{-\frac{d}{2}}U\left(\frac{d}{2}, \frac{2+d-\nu}{2}, \frac{1}{4t}\right) \asymp t^{-\frac{\nu \wedge d}{2}}, \quad \text{as } t \to \infty,
$$

from which one obtains the large time asymptotics of $h(t)$ in $(A.5)$. This completes the whole proof of Proposition [A.1.](#page-41-1) \Box

A.2 One-dimensional stochastic wave equations

Proposition A.2. *Let h*(*t*) *be defined as in* [\(5.15\)](#page-32-3)*. The following statements hold:*

(i) (Riesz Kernel) If $f(x) = |x|^{-\alpha}$ with $\alpha \in (0,1)$, then for all $t > 0$,

$$
h(t) = Ct^{3-\alpha} \quad \text{with } C = \frac{2^{1-\alpha}}{(1-\alpha)(2-\alpha)(3-\alpha)}.
$$

(ii) (Ornstein-Uhlenbeck kernel) If $f(x) = \exp(-|x|^{\alpha})$ *with* $\alpha > 0$ *, then,*

$$
h(t) \approx t^2 \quad as \ t \to \infty \ and \quad h(t) \approx t^3 \quad as \ t \downarrow 0. \tag{A.6}
$$

- *(iii) (Brownian case:* $W(t, x) = W(t)$ *)* If $f(x) \equiv 1$ *, then* $h(t) = t^3/3$ *for all* $t \ge 0$ *.*
- *(iv) (Space-time white case) If* $f(x) = \delta(x)$ *, then* $h(t) = t^2/2$ *for all* $t \ge 0$ *.*
- *(v)* (Bessel potential) Suppose that $f_\nu(\cdot)$ is the Bessel potential given in [\(A.3\)](#page-41-6) with $d = 1$ and the *parameter* $\nu > 0$ *. Then*

$$
h(t) \asymp t^2 \quad as \ t \to \infty, \quad and \quad h(t) \asymp t^3 \quad as \ t \downarrow 0. \tag{A.7}
$$

(vi) (Bessel potential as the spectral measure) Suppose that $f_{\nu}(x) = (1 + x^2)^{-\nu/2}$ with $\nu > 0$ and $x \in \mathbb{R}$ *. Then,*

$$
h(t) \approx \begin{cases} t^{3-\nu}, & \nu < 1, \\ t^2 \log(t), & \nu = 1, \\ t^2, & \nu > 1, \end{cases} \quad \text{as } t \to \infty, \quad \text{and} \quad h(t) \approx t^3 \quad \text{as } t \downarrow 0. \tag{A.8}
$$

Proof. Parts (i), (iii) and (iv) are straightforward. Similar as in Proposition [A.1,](#page-41-1) the asymptotics for $t \downarrow 0$ in $(A.6)$ – $(A.8)$ are obtained because

$$
\frac{c_1t^3}{3} = c_1 \int_0^t ds \int_{\mathbb{R}^2} dy dy' G(s, y) G(s, y') \le \int_0^t ds \int_{\mathbb{R}^2} dy dy' G(s, y) G(s, y') f(y - y') \le \frac{c_2t^3}{3},
$$

for all $t \in [0, \epsilon)$ and $f(\cdot)$ such that $0 < c_1 \le f(x) \le c_2$ for all $x \in [-2\epsilon, 2\epsilon]$ with some $\epsilon > 0$. For part (ii), we first note that

$$
h(t) \le C \int_0^t \int_{-s}^s \mathrm{d}y \int_{\mathbb{R}} \mathrm{d}y' f(y - y') \le C t^2.
$$

For the low bounds, assume that $t > 1$. Then

$$
h(t) \ge C \int_1^t ds \int_{[-s,s]^2} dy dy' f(y - y') \mathbb{I}_{[-1,1]}(y - y') \ge C 2^{\frac{3}{2}} e^{-2\alpha} \int_1^t ds \, (s - 1) \ge C(t - 1)^2.
$$

This complete the proof of the large time asymptotics in [\(A.6\)](#page-43-2).

As for part (v), by using the Fourier transform of the wave kernel, we see that

$$
h(t) = (2\pi)^{-1} \int_0^t ds \int_{\mathbb{R}} d\xi \frac{\sin^2(s\xi)}{\xi^2 (1 + \xi^2)^{\nu/2}} \n\leq C \int_0^t ds \left(\int_0^{\frac{\pi}{2s}} d\xi \frac{(s\xi)^2}{\xi^2 (1 + \xi^2)^{\nu/2}} + \int_{\frac{\pi}{2s}}^\infty d\xi \frac{1}{\xi^2 (1 + \xi^2)^{\nu/2}} \right) \n\leq C \int_0^t ds \left(\int_0^{\frac{\pi}{2s}} s^2 d\xi + \int_{\frac{\pi}{2s}}^\infty \frac{1}{\xi^2} d\xi \right) \leq C \int_0^t ds (s + s) \leq Ct^2.
$$

On the other hand, it is clear that

$$
h(t) \ge C \int_0^t ds \int_0^{\frac{\pi}{2s}} d\xi \frac{\sin^2(s\xi)}{\xi^2 (1 + \xi^2)^{\nu/2}} \ge C \int_1^t ds \int_0^{\frac{\pi}{2s}} d\xi \frac{(s\xi)^2}{\xi^2 (1 + \xi^2)^{\nu/2}}
$$

$$
\ge C \int_1^t ds \int_0^{\frac{\pi}{2s}} d\xi \frac{s^2}{(1 + (\pi/2)^2)^{\nu/2}} \ge C(t - 1)^2,
$$

for all $t \geq 1$. This proves the large time asymptotics in $(A.7)$.

As for part (vi), by changing of variables $y - y' = z$ and $y + y' = z'$, we have that $dy dy' = z'$ $2^{-1}dzdz'$. Thus, for $t > 2$,

$$
h(t) \le \frac{1}{8} \int_0^t ds \int_{-2s}^{2s} \int_{-2s}^{2s} \frac{dz dz'}{(1+z^2)^{\nu/2}} \le C \left(1 + \int_0^t ds \, s \int_1^{2s} dz \, z^{-\nu} \right)
$$

and

$$
h(t) \ge \frac{1}{8} \int_2^t \mathrm{d}s \int_{-s/2}^{s/2} \int_{-s/2}^{s/2} \frac{\mathrm{d}z \mathrm{d}z'}{(1+z^2)^{\nu/2}} \ge \frac{1}{8} \int_2^t \mathrm{d}s \, s \int_1^{s/2} \frac{\mathrm{d}z}{(1+z^2)^{\nu/2}} \ge \frac{1}{2^{3+\nu/2}} \int_2^t \mathrm{d}s \, s \int_1^{s/2} \mathrm{d}z z^{-\nu}.
$$

By considering three cases $\nu \in (0, 1)$, $\nu = 1$, and $\nu > 1$ in the above two inequalities, we obtain the large time asymptotics in (A.8). This completes the proof of Proposition A.2. large time asymptotics in $(A.8)$. This completes the proof of Proposition [A.2.](#page-43-1)

B Supplementary proof for Proposition [6.2](#page-36-2)

Proof of Proposition [6.2.](#page-36-2) All formulas in the special cases (i)–(iii), except (6.4) and (6.5) , can be deduced by plugging in the specific $h(t)$ given in Appendix [A.1.](#page-41-0) For (6.4) , by [\[CE22b,](#page-45-3) Example 5.8], we find that

$$
\mathcal{J}_0(t,x) \le c_1 t^{-\ell/2} \quad \text{and} \quad \mathcal{J}_1(t,x) \le c_2 t^{1-\ell-\beta/2}.
$$

Then, plugging these inequalities into (6.3) , one gets (6.4) immediately.

The proof of [\(6.5\)](#page-37-4) also relies on the estimates for $\mathcal{J}_0(t,x)$ and $\mathcal{J}_1(t,x)$. By using the equivalence of norms in Euclidean spaces, there exists $c_2 \geq c_1 > 0$ such that

$$
c_1 \sum_{i=1}^d |x_i| \le |x| = \left(\sum_{i=1}^d x_i^2\right)^{1/2} \le c_2 \sum_{i=1}^d |x_i|.
$$

It follows that

$$
e^{\ell|x|} \le \exp\left(c\ell \sum_{i=1}^d |x_i|\right)
$$
, for all $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$,

where $c = c_2$ if $\ell \geq 0$ and otherwise $c = c_1$. As a result, by using [\[CD15c](#page-45-4), Proposition A.11], we can deduce that

$$
\mathcal{J}_0(t,x) \le \prod_{i=1}^d \int_{\mathbb{R}} \mathrm{d}y_i e^{c\ell|y_i|} \frac{1}{\sqrt{2\pi t}} e^{-\frac{(x_i - y_i)^2}{2t}} \le 2e^{c^2 d\ell^2 t + c\ell|x|}.\tag{B.1}
$$

We still need to estimate $\mathcal{J}_1(t,x)$. Using (6.4) , by [\[CE22b](#page-45-3), Example 5.8], we have

$$
\int_{\mathbb{R}^d} \mathrm{d}y p_t(x-y)|y|^{-\beta} \le c_3 t^{-\beta/2},\tag{B.2}
$$

with some universal constant $c_3 > 0$. Hence, plugging [\(B.1\)](#page-44-1) and [\(B.2\)](#page-44-2) into [\(1.11\)](#page-4-6), it follows that

$$
\mathcal{J}_1(t,x) \le 2c_3 \int_0^t ds(t-s)^{-\beta/2} \int_{\mathbb{R}^d} dy \, p_{t-s}(x-y) e^{2c^2 d\ell^2 s + 2c\ell |y|}.
$$

Finally, following the same argument as in the proof of [\(B.1\)](#page-44-1), one can show that

$$
\mathcal{J}_1(t,x) \le 2c_3 \int_0^t ds(t-s)^{-\beta/2} \exp\left(2c^2 d\ell^2 s + 4c^4 d\ell^2 (t-s) + 2c^2 \ell |x|\right)
$$

$$
\le t^{1-\beta/2} \exp\left(C_1 \ell^2 t + C_2 \ell |x|\right).
$$

This completes the proof of (6.5) .

References

- [ACQ11] Gideon Amir, Ivan Corwin, and Jeremy Quastel. "Probability distribution of the free energy of the continuum directed random polymer in 1 + 1 dimensions". In: *Comm. Pure Appl. Math.* 64.4 (2011), pp. 466–537.
- [BC16] Raluca M. Balan and Daniel Conus. "Intermittency for the wave and heat equations with fractional noise in time". In: *Ann. Probab.* 44.2 (2016), pp. 1488–1534.
- [Bih56] I. Bihari. "A generalization of a lemma of Bellman and its application to uniqueness problems of differential equations". In: *Acta Math. Acad. Sci. Hungar.* 7 (1956), pp. 81– 94.
- [CM94] René A. Carmona and S. A. Molchanov. "Parabolic Anderson problem and intermittency". In: *Mem. Amer. Math. Soc.* 108.518 (1994), pp. viii+125.
- [CD14] Le Chen and Robert C. Dalang. "Hölder-continuity for the nonlinear stochastic heat equation with rough initial conditions". In: *Stoch. Partial Differ. Equ. Anal. Comput.* 2.3 (2014), pp. 316–352.
- [CD15a] Le Chen and Robert C. Dalang. "Moment bounds and asymptotics for the stochastic wave equation". In: *Stochastic Process. Appl.* 125.4 (2015), pp. 1605–1628.
- [CD15b] Le Chen and Robert C. Dalang. "Moments and growth indices for the nonlinear stochastic heat equation with rough initial conditions". In: *Ann. Probab.* 43.6 (2015), pp. 3006– 3051.
- [CD15c] Le Chen and Robert C. Dalang. "Moments, intermittency and growth indices for the nonlinear fractional stochastic heat equation". In: *Stoch. Partial Differ. Equ. Anal. Comput.* 3.3 (2015), pp. 360–397.
- [CE22a] Le Chen and Nicholas Eisenberg. "Interpolating the stochastic heat and wave equations with time-independent noise: solvability and exact asymptotics". In: *Stoch. Partial Differ. Equ. Anal. Comput. (in press)* (2022).
- [CE22b] Le Chen and Nicholas Eisenberg. "Invariant measures for the nonlinear stochastic heat equation with no drift term". In: *preprint arXiv:2209.04771* (2022).
- [CGS22] Le Chen, Yuhui Guo, and Jian Song. "Moments and asymptotics for a class of SPDEs with space-time white noise". In: *preprint arXiv:2206.10069* (2022).
- [CHN19] Le Chen, Yaozhong Hu, and David Nualart. "Nonlinear stochastic time-fractional slow and fast diffusion equations on \mathbb{R}^{d} ". In: *Stochastic Process. Appl.* 129.12 (2019), pp. 5073– 5112.

 \Box

- [CH19] Le Chen and Jingyu Huang. "Comparison principle for stochastic heat equation on \mathbb{R}^{d} ". In: *Ann. Probab.* 47.2 (2019), pp. 989–1035.
- [CH22] Le Chen and Jingyu Huang. "Superlinear stochastic heat equation on R *d* ". In: *Proc. Amer. Math. Soc. (in press)* (2022).
- [CK19] Le Chen and Kunwoo Kim. "Nonlinear stochastic heat equation driven by spatially colored noise: moments and intermittency". In: *Acta Math. Sci. Ser. B (Engl. Ed.)* 39.3 (2019), pp. 645–668.
- [CK20] Le Chen and Kunwoo Kim. "Stochastic comparisons for stochastic heat equation". In: *Electron. J. Probab.* 25 (2020), Paper No. 140, 38.
- [Che15] Xia Chen. "Precise intermittency for the parabolic Anderson equation with an $(1 + 1)$ dimensional time-space white noise". In: *Ann. Inst. Henri Poincaré Probab. Stat.* 51.4 (2015), pp. 1486–1499.
- [CJK13] Daniel Conus, Mathew Joseph, and Davar Khoshnevisan. "On the chaotic character of the stochastic heat equation, before the onset of intermitttency". In: *Ann. Probab.* 41.3B (2013), pp. 2225–2260.
- [Con+13] Daniel Conus, Mathew Joseph, Davar Khoshnevisan, and Shang-Yuan Shiu. "On the chaotic character of the stochastic heat equation, II". In: *Probab. Theory Related Fields* 156.3-4 (2013), pp. 483–533.
- [Con+14] Daniel Conus, Mathew Joseph, Davar Khoshnevisan, and Shang-Yuan Shiu. "Initial measures for the stochastic heat equation". In: *Ann. Inst. Henri Poincaré Probab. Stat.* 50.1 (2014), pp. 136–153.
- [CK12] Daniel Conus and Davar Khoshnevisan. "On the existence and position of the farthest peaks of a family of stochastic heat and wave equations". In: *Probab. Theory Related Fields* 152.3-4 (2012), pp. 681–701.
- [Cor12] Ivan Corwin. "The Kardar-Parisi-Zhang equation and universality class". In: *Random Matrices Theory Appl.* 1.1 (2012), pp. 1130001, 76.
- [Dal+09] Robert Dalang, Davar Khoshnevisan, Carl Mueller, David Nualart, and Yimin Xiao. *A minicourse on stochastic partial differential equations*. Vol. 1962. Held at the University of Utah, Salt Lake City, UT, May 8–19, 2006, Edited by Khoshnevisan and Firas Rassoul-Agha. Springer-Verlag, Berlin, 2009, pp. xii+216. isbn: 978-3-540-85993-2.
- [Dal99] Robert C. Dalang. "Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.'s". In: *Electron. J. Probab.* 4 (1999), no. 6, 29.
- [DKZ19] Robert C. Dalang, Davar Khoshnevisan, and Tusheng Zhang. "Global solutions to stochastic reaction-diffusion equations with super-linear drift and multiplicative noise". In: *Ann. Probab.* 47.1 (2019), pp. 519–559.
- [DM09] Robert C. Dalang and Carl Mueller. "Intermittency properties in a hyperbolic Anderson problem". In: *Ann. Inst. Henri Poincar´e Probab. Stat.* 45.4 (2009), pp. 1150–1164.
- [DS05] Robert C. Dalang and Marta Sanz-Solé. "Regularity of the sample paths of a class of second-order spde's". In: *J. Funct. Anal.* 227.2 (2005), pp. 304–337.
- [DIP89] D. A. Dawson, I. Iscoe, and E. A. Perkins. "Super-Brownian motion: path properties and hitting probabilities". In: *Probab. Theory Related Fields* 83.1-2 (1989), pp. 135–205.
- [Daw93] Donald A. Dawson. "Measure-valued Markov processes". In: *École d'Été de Probabilités de Saint-Flour XXI—1991*. Vol. 1541. Springer, Berlin, 1993, pp. 1–260.
- [Eth00] Alison M. Etheridge. *An introduction to superprocesses*. Vol. 20. American Mathematical Society, Providence, RI, 2000, pp. xii+187. isbn: 0-8218-2706-5.
- [FZ05] Shizan Fang and Tusheng Zhang. "A study of a class of stochastic differential equations with non-Lipschitzian coefficients". In: *Probab. Theory Related Fields* 132.3 (2005), pp. 356–390.
- [FK09] Mohammud Foondun and Davar Khoshnevisan. "Intermittence and nonlinear parabolic stochastic partial differential equations". In: *Electron. J. Probab.* 14 (2009), no. 21, 548– 568.
- [Gra14] Loukas Grafakos. *Modern Fourier analysis*. Third. Vol. 250. Springer, New York, 2014, pp. xvi+624. isbn: 978-1-4939-1229-2; 978-1-4939-1230-8.
- [GSS23] Yuhui Guo, Jian Song, and Xiaoming Song. "Stochastic fractional diffusion equations with Gaussian noise rough in space". In: *preprint arXiv:2303.11939* (2023).
- [GG06] Peter Guttorp and Tilmann Gneiting. "Studies in the history of probability and statistics. XLIX. On the Matérn correlation family". In: *Biometrika* 93.4 (2006), pp. 989– 995.
- [HW21] Yaozhong Hu and Xiong Wang. "Intermittency properties for a large class of stochastic PDEs driven by fractional space-time noises". In: *preprint arXiv:2109.03473* (2021).
- [Hu+23] Yaozhong Hu, Xiong Wang, Panqiu Xia, and Jiayu Zheng. "Moment asymptotics for super-Brownian motions". In: *preprint arXiv:2303.12994* (2023).
- [HLN17] Jingyu Huang, Khoa Lê, and David Nualart. "Large time asymptotics for the parabolic Anderson model driven by space and time correlated noise". In: *Stoch. Partial Differ. Equ. Anal. Comput.* 5.4 (2017), pp. 614–651.
- [JKM17] Mathew Joseph, Davar Khoshnevisan, and Carl Mueller. "Strong invariance and noisecomparison principles for some parabolic stochastic PDEs". In: *Ann. Probab.* 45.1 (2017), pp. 377–403.
- [Kal02] Olav Kallenberg. *Foundations of modern probability*. Second. Springer-Verlag, New York, 2002, pp. xx+638. ISBN: 0-387-95313-2.
- [Kho14] Davar Khoshnevisan. *Analysis of stochastic partial differential equations*. Vol. 119. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 2014, pp. viii+116. isbn: 978-1-4704- 1547-1.
- [KKX17] Davar Khoshnevisan, Kunwoo Kim, and Yimin Xiao. "Intermittency and multifractality: a case study via parabolic stochastic PDEs". In: *Ann. Probab.* 45.6A (2017), pp. 3697–3751.
- [Kön16] Wolfgang König. *The parabolic Anderson model*. Random walk in random potential. Birkh¨auser/Springer, [Cham], 2016, pp. xi+192. isbn: 978-3-319-33595-7; 978-3-319- 33596-4.
- [KS88] N. Konno and T. Shiga. "Stochastic partial differential equations for some measurevalued diffusions". In: *Probab. Theory Related Fields* 79.2 (1988), pp. 201–225.
- [LaS49] J. LaSalle. "Uniqueness theorems and successive approximations". In: *Ann. of Math. (2)* 50 (1949), pp. 722–730.
- [Mue91] Carl Mueller. "Long time existence for the heat equation with a noise term". In: *Probab. Theory Related Fields* 90.4 (1991), pp. 505–517.
- [MMP14] Carl Mueller, Leonid Mytnik, and Edwin Perkins. "Nonuniqueness for a parabolic SPDE with $\frac{3}{4}$ – ϵ -Hölder diffusion coefficients". In: *Ann. Probab.* 42.5 (2014), pp. 2032–2112.
- [MMQ11] Carl Mueller, Leonid Mytnik, and Jeremy Quastel. "Effect of noise on front propagation in reaction-diffusion equations of KPP type". In: *Invent. Math.* 184.2 (2011), pp. 405– 453.
- [MMR21] Carl Mueller, Leonid Mytnik, and Lenya Ryzhik. "The speed of a random front for stochastic reaction-diffusion equations with strong noise". In: *Comm. Math. Phys.* 384.2 (2021), pp. 699–732.
- [MP92] Carl Mueller and Edwin A. Perkins. "The compact support property for solutions to the heat equation with noise". In: *Probab. Theory Related Fields* 93.3 (1992), pp. 325–358.
- [Myt98] Leonid Mytnik. "Weak uniqueness for the heat equation with noise". In: *Ann. Probab.* 26.3 (1998), pp. 968–984.
- [MP11] Leonid Mytnik and Edwin Perkins. "Pathwise uniqueness for stochastic heat equations with Hölder continuous coefficients: the white noise case". In: *Probab. Theory Related Fields* 149.1-2 (2011), pp. 1–96.
- [MPS06] Leonid Mytnik, Edwin Perkins, and Anja Sturm. "On pathwise uniqueness for stochastic heat equations with non-Lipschitz coefficients". In: *Ann. Probab.* 34.5 (2006), pp. 1910– 1959.
- [NP18] Constantin P. Niculescu and Lars-Erik Persson. *Convex functions and their applications*. A contemporary approach, Second edition of [MR2178902]. Springer, Cham, 2018, pp. xvii+415. isbn: 978-3-319-78336-9; 978-3-319-78337-6.
- [Olv+10] Frank W. J. Olver, Daniel W. Lozier, Ronald F. Boisvert, and Charles W. Clark. *NIST handbook of mathematical functions*. With 1 CD-ROM (Windows, Macintosh and UNIX). U.S. Department of Commerce, National Institute of Standards and Technology, Washington, DC; Cambridge University Press, Cambridge, 2010, pp. xvi+951. isbn: 978-0-521-14063-8.
- [Per02] Edwin Perkins. "Dawson-Watanabe superprocesses and measure-valued diffusions". In: *Lectures on probability theory and statistics (Saint-Flour, 1999)*. Vol. 1781. Springer, Berlin, 2002, pp. 125–324.
- [Rei89] Mark Reimers. "One-dimensional stochastic partial differential equations and the branching measure diffusion". In: *Probab. Theory Related Fields* 81.3 (1989), pp. 319–340.
- [Sal22] Michael Salins. "Global solutions for the stochastic reaction-diffusion equation with super-linear multiplicative noise and strong dissipativity". In: *Electron. J. Probab.* 27 (2022), Paper No. 12, 17.
- [SS02] Marta Sanz-Solé and Mònica Sarrà. "Hölder continuity for the stochastic heat equation with spatially correlated noise". In: *Seminar on Stochastic Analysis, Random Fields and Applications, III (Ascona, 1999)*. Vol. 52. Birkhäuser, Basel, 2002, pp. 259–268.
- [Shi94] Tokuzo Shiga. "Two contrasting properties of solutions for one-dimensional stochastic partial differential equations". In: *Canad. J. Math.* 46.2 (1994), pp. 415–437.
- [Wal86] John B. Walsh. "An introduction to stochastic partial differential equations". In: *Ecole ´ d'´et´e de probabilit´es de Saint-Flour, XIV—1984*. Vol. 1180. Springer, Berlin, 1986, pp. 265–439.
- [Xio13] Jie Xiong. *Three classes of nonlinear stochastic partial differential equations*. World Scientific Publishing Co. Pte. Ltd., Hackensack, NJ, 2013, pp. xii+164. ISBN: 978-981-4452-35-9.
- [Zel+87] Ya. B. Zel′ dovich, S. A. Molchanov, A. A. Ruzmauikin, and D. D. Sokoloff. "Selfexcitation of a nonlinear scalar field in a random medium". In: *Proc. Nat. Acad. Sci. U.S.A.* 84.18 (1987), pp. 6323–6325.
- [ZRS90] Ya. B. Zel′ dovich, A. A. Ruzmauikin, and D. D. Sokoloff. *The almighty chance*. Vol. 20. Translated from the Russian by Anvar Shukurov. World Scientific Publishing Co., Inc., River Edge, NJ, 1990, pp. xii+316. isbn: 9971-50-916-4; 9971-50-917-2.