ON AN EXTENSION OF NORI AND LOCAL FUNDAMENTAL GROUP SCHEMES

PAVAN ADROJA AND SANJAY AMRUTIYA

ABSTRACT. This article studies an extended Nori and local fundamental group schemes of Abelian varieties. We also discuss the birational invariance of these group schemes and study their behaviour under the Albanese and étale morphisms.

1. INTRODUCTION

Madhav Nori [13] has introduced the fundamental group scheme of a reduced and connected scheme X over a field k. If X is a proper integral scheme over a field k with $x \in X(k) \neq \emptyset$, then the Nori fundamental group scheme, denoted by $\pi^N(X, x)$, is Tannaka dual to a neutral Tannakian category of *essentially finite* vector bundles over X. Recall that a vector bundle E over X is said to be *finite* if there are two distinct polynomials f and g with non-negative integer coefficients such that f(E) is isomorphic to g(E). A vector bundle E over X is said to be *Nori-semistable* if for every non-constant morphism $f : C \longrightarrow X$ with C a smooth projective curve, the pull-back $f^*E \longrightarrow C$ is semistable of degree zero. A vector bundle E over X is said to be *essentially finite* if it is a Nori-semistable subquotient of a finite vector bundle over X. The category $\mathcal{C}^{N}(X)$ of essentially finite vector bundles is a full subcategory of *Nori semistable* bundles.

Adrian Langer [10, 11] studied various properties of so-called the S-fundamental group scheme. The S-fundamental group scheme of (X, x), denoted by $\pi^{S}(X, x)$ is Tannaka dual to the category of numerically flat bundles. Recall that a vector bundle is numerically flat if and only if it is Nori-semistable. He also gives the structure theorem of $\pi^{S}(X, x)$ for abelian varieties.

Otabe [16] considers a slightly bigger category than the category of finite bundles in characteristic zero. He defined the semi-finite bundle and proved that the category of semi-finite bundles over X form the Tannakian category. The corresponding affine group scheme is called the extended Nori fundamental group scheme, and is denoted by $\pi^{\text{EN}}(X, x)$. If C is an elliptic curve over the field of characteristic zero, then $\pi^{\text{EN}}(C, x)$ is the product of $\pi^N(C, x)$ and $\pi^{\text{uni}}(C, x)$. The analoge of a semi-finite bundle over the field of positive characteristic is discussed in [1]. It is proved that for an elliptic curve C over the field of positive characteristic, $\pi^{\text{EN}}(C, x)$ is isomorphic to $\pi^N(C, x)$. In this paper, we

²⁰⁰⁰ Mathematics Subject Classification. Primary: 14L15, 14F05.

Key words and phrases. Fundamental group schemes, Abelian varieties.

The research work of Pavan Adroja is financially supported by CSIR-UGC fellowship under the grant no. 191620092279.

generalize these two results to higher dimensional abelian varieties (see Theorem 3.3 and Proposition 3.6).

If the characteristic of k is zero, then the birational invariance of the S-fundamental group scheme follows from the birational invariance of the topological fundamental group. It is also known that the Nori fundamental group scheme is birational invariant for smooth projective varieties [13]. If char k = p > 0, then the birational invariance of the S-fundamental group scheme is proved in [8]. One can use the techniques of [8] to show the birational invariance of $\pi^{\text{EN}}(X, x)$. In fact, the proof of birational invariance of $\pi^N(X, x)$ and $\pi^{\text{loc}}(X, x)$ is inbuilt in the proof of [8, Theorem 1.2]. In Corollary 4.5, we provide additional details for completeness. We also study the behaviour of the extended Nori fundamental group scheme under the Albanese and étale morphisms (see Section 4). In the last section, we introduce an extension of the local fundamental group scheme and study it in a similar spirit.

2. Preliminaries

Let X be a proper integral scheme defined over a perfect field k endowed with a rational point $x \in X(k)$.

Definition 2.1. A vector bundle V on X is called unipotent bundle if it has a filtration $V = V_0 \supset V_1 \supset V_2 \supset ... \supset V_n = 0$

such that $V_i/V_{i+1} \simeq \mathcal{O}_X$ for all *i*.

The category of unipotent vector bundles $\mathcal{C}^{\text{uni}}(X)$ over X form the Tannakian category and the corresponding affine group scheme $\pi^{\text{uni}}(X, x)$ is called the unipotent group scheme of X [13, Chapter IV]. Let $\pi^{\text{uni}}_{ab}(X, x)$ be the largest abelian quotient of $\pi^{\text{uni}}(X, x)$.

Proposition 2.2. [13, Chapter IV] Let X be a proper connected reduced scheme over the algebraically closed field k with $x \in X(k)$. If k has characteristic zero, then we have

$$\pi_{\rm ab}^{\rm uni}(X, x) \simeq H^1(X, \mathcal{O}_X)^*.$$

If the characteristic of k is positive, then we have

$$\pi_{\rm ab}^{\rm uni}(X,x) \simeq \varprojlim \hat{G},$$

where \hat{G} denotes the Cartier dual of G, and the inverse limit is taken over all finite local group subschemes G of Pic⁰ X.

2.1. An extension of the Nori fundamental group scheme. S. Otabe [16] introduced the notion of a semi-finite bundle when the base field k is of characteristic zero. Note that an essentially finite vector bundle is finite if k has characteristic zero [13, Chapter-I, §3].

Definition 2.3. A vector bundle V on X is called semi–essentially finite bundle if it has a filtration

$$V = V_0 \supset V_1 \supset V_2 \supset \ldots \supset V_n = 0$$

such that V_i/V_{i+1} is an essentially finite indecomposable bundle, for all i.

Let $\mathcal{C}^{\text{EN}}(X)$ be the Tannakian category of semi–essentially finite bundles over X. The corresponding affine group scheme is denoted by $\pi^{\text{EN}}(X, x)$.

Theorem 2.4. [16, Theorem 1.1] Let C be an elliptic curve over an algebraically closed field k of characteristic zero. For k-rational point $x \in C(k)$, we have

$$\pi^{\mathrm{EN}}(C, x) \simeq \pi^{N}(C, x) \times \pi^{\mathrm{uni}}(C, x).$$

Theorem 2.5. [1, Theorem 4.3] Let C be an elliptic curve over an algebraically closed field k of positive characteristic. For k-rational point $x \in C(k)$, we have

$$\pi^{\mathrm{EN}}(C, x) \simeq \pi^N(C, x).$$

2.2. The S-fundamental group scheme. Let X be a smooth projective k-variety. Recall that a line bundle L over X is said to be numerically effective if the degree of the restriction of L to any irreducible curve C in X is non-negative. A vector bundle E over X is called numerically effective if and only if the tautological line bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ is numerically effective. A vector bundle E is called numerically flat if both E and its dual E^{\vee} are numerically effective.

The category $\mathcal{C}^{nf}(X)$ of numerically flat bundles over X with a k-point x form a neutral Tannakian category and the corresponding affine group scheme is called the S-fundamental group scheme, denote by $\pi^{S}(X, x)$ (see [10, 11] for other equivalent descriptions and various properties of this group scheme).

Let G be any abelian group. Then we can define a diagonal group scheme Diag(G) as a functor $\text{Diag}(G): \operatorname{Alg}_k \longrightarrow \operatorname{Ab}$ by $\operatorname{Diag}(G)(R) = \operatorname{Hom}_{gr}(G, R^{\times})$. Clearly, the functor $\operatorname{Diag}(G)$ is represented by the k-group algbera k[G]. So, the group scheme $\operatorname{Diag}(G)$ is the same as taking $\operatorname{Spec} k[G]$ with the natural k-group scheme structure.

Let Pic X denote the Picard group scheme of X. We denote by $\operatorname{Pic}^{0}X$ its connected component of the identity and by $\operatorname{Pic}^{\tau}X$ the torsion component of the identity. Recall that the Neron–Severi group $\operatorname{NS}(X) = (\operatorname{Pic}X)_{\operatorname{red}}/(\operatorname{Pic}^{0}X)_{\operatorname{red}}$ is finitely generated. We have a short exact sequence

$$0 \to (\operatorname{Pic}^{0} X)_{\operatorname{red}} \to (\operatorname{Pic}^{\tau} X)_{\operatorname{red}} \to \operatorname{NS}(X)_{\operatorname{tor}} \to 0,$$

where $NS(X)_{tor}$ is the torsion group of NS(X). Therefore, we have a short exact sequence

$$0 \to \operatorname{Diag}(\operatorname{NS}(X)_{\operatorname{tor}}) \to \operatorname{Diag}((\operatorname{Pic}^{\tau} X)_{\operatorname{red}}) \to \operatorname{Diag}((\operatorname{Pic}^{0} X)_{\operatorname{red}}) \to 0.$$

We refer to [9] for more details about the Picard scheme.

Theorem 2.6. [11, Theorem 5.9] Let X be a smooth projective variety defined over an algebraically closed field k. If char k = 0, then we have an isomorphism

$$\pi^{S}_{\mathrm{ab}}(X, x) \simeq H^{1}(X, \mathcal{O}_{X})^{*} \times \mathrm{Diag}(\mathrm{Pic}^{\tau}X).$$

If char k = p > 0, then we have

$$\pi^{S}_{\mathrm{ab}}(X, x) \simeq \varprojlim \widehat{G} \times \mathrm{Diag}((\mathrm{Pic}^{\tau} X)_{\mathrm{red}}),$$

where \hat{G} denotes the Cartier dual of G and the inverse limit is taken over all finite local group subschemes G of $\operatorname{Pic}^{0} X$.

Abelian varieties: Let A be an abelian variety of dimension g over an algebraically closed field k. Let $n_A : A \longrightarrow A$ be a multiplication by n map and $A_n := \text{Ker } n_A$. If char k = 0, then we have $A_n \simeq \left(\frac{\mathbb{Z}}{n\mathbb{Z}}\right)^{2g}$ as a group scheme. If char k = p > 0, then we have (see [17]) the following:

- (1) $A_n \simeq \left(\frac{\mathbb{Z}}{n\mathbb{Z}}\right)^{2g}$ when n is relatively prime to p;
- (2) $A_{p^m} \simeq A_{p^m}^* \times \hat{A}_{p^m}^* \times A_{p^m}^0$, where $A_{p^m}^*$ denote the reduced part of A_{p^m} , $\hat{A}_{p^m}^*$ denote Cartier dual of $A_{p^m}^*$ and $A_{p^m}^0$ is local-local part of A_{p^m} for all $m \in \mathbb{N}$. Note that $A_{p^m}^* \simeq (\frac{\mathbb{Z}}{p^m \mathbb{Z}})^r$ and $\hat{A}_{p^m}^* \simeq (\mu_{p^m})^r$, where $0 \le r \le g$ is called the *p*-rank of A;
- (3) $A_{tp^m} \simeq A_t \times A_{p^m}$ when t is relatively prime to p.

The inverse limit $\lim_{n \to \infty} A_{p^n}^*$ is called the *p*-adic discrete Tate group $T_p^d(A)$ and, the inverse limit $\lim_{n \to \infty} A_{p^n}^0$ is called *p*-adic local-local Tate group $T_p^0(A)$.

3. Structure theorem

For a variety X defined over a field of positive characteristic, we denote by $F_X \colon X \to X$ the Frobenius morphism. Recall that a vector bundle V on X is called F-trivial if there is a positive integer n such that $(F_X^n)^*V \simeq \mathcal{O}_X^r$.

3.1. The abelian part of the $\pi^{\text{EN}}(X, x)$. We first give the structure theorem for the abelian part of the group scheme $\pi^{\text{EN}}(X, x)$.

Lemma 3.1. Let X be a smooth projective variety over the algebraically closed field k with char k = p > 0. Then, every essentially finite line bundle over X is a torsion (finite) line bundle.

Proof. Let L be an essentially finite line bundle on X. By [3, Theorem 3.5], there is a connected étale Galois covering $\gamma : \hat{X} \longrightarrow X$ such that $\gamma^* L$ is an F-trivial vector bundle over \hat{X} , i.e., there exist $n \in \mathbb{N}$ such that $(F_{\hat{X}}^n)^* \gamma^* L \simeq \mathcal{O}_{\hat{X}}$. Since $\gamma \circ F_{\hat{X}}^n = F_X^n \circ \gamma$, we have

$$(F_{\hat{X}}^n)^*\gamma^*L \simeq \gamma^*(F_X^n)^*L \simeq \gamma^*(L^{\otimes p^n}) \simeq \mathcal{O}_{\hat{X}}$$

Therefore, $L^{\otimes p^n}$ is étale trivializable, and hence it is a finite line bundle.

Let $\pi_{ab}^{EN}(X, x)$ be the largest abelian quotient of $\pi^{EN}(X, x)$ which is called *abelian part* of $\pi^{EN}(X, x)$.

Let \mathbb{F}_{τ} denote the group of torsion (finite) line bundles on X. Clearly, \mathbb{F}_{τ} is a subgroup of Pic^{τ} X.

Theorem 3.2. Let X be a smooth projective variety defined over an algebraically closed field k. If char k = 0 then we have an isomorphism

$$\pi_{\mathrm{ab}}^{\mathrm{EN}}(X, x) \simeq H^1(X, \mathcal{O}_X)^* \times \mathrm{Diag}(\mathbb{F}_{\tau}).$$

If char k = p > 0 then we have

$$\pi_{\mathrm{ab}}^{\mathrm{EN}}(X, x) \simeq \varprojlim \hat{G} \times \mathrm{Diag}(\mathbb{F}_{\tau}),$$

where \hat{G} denotes the Cartier dual of G and the inverse limit is taken over all finite local group subschemes G of $\operatorname{Pic}^{0} X$.

Proof. Let $\pi_{\text{uni}}^{\text{EN}}(X, x)$ be the largest unipotent quotient of $\pi^{\text{EN}}(X, x)$. By the standard Tannakian consideration, the category of finite dimensional representation of $\pi_{\text{uni}}^{\text{EN}}(X, x)$ is equivalent to the full subcategory of $\mathcal{C}^{\text{EN}}(X)$ whose objects have a filtration with all successive quotients are isomorphic to \mathcal{O}_X . Since $\mathcal{C}^{\text{uni}}(X) \subset \mathcal{C}^{\text{EN}}(X)$, we have $\pi_{\text{uni}}^{\text{EN}}(X, x) \simeq \pi^{\text{uni}}(X, x)$. Therefore, the largest unipotent quotient of $\pi_{\text{ab}}^{\text{EN}}(X, x)$ which is called *unipotent part* of $\pi_{\text{ab}}^{\text{EN}}(X, x)$ is isomorphic to the abelian part of $\pi^{\text{uni}}(X, x)$.

If char k = 0, then by [13, Chapter IV, Proposition 2], the abelian part of $\pi^{\text{uni}}(X, x)$ is $H^1(X, \mathcal{O}_X)^*$. If char k = p > 0, then by [13, Ch. IV, Proposition 6], the abelian part of $\pi^{\text{uni}}(X, x)$ is the inverse limit of Cartier duals of finite local group subscheme of Pic⁰ X.

Since the homomorphism $\pi^{\text{EN}}(X, x) \longrightarrow \mathbb{G}_m$ is one dimensional representation of $\pi^{\text{EN}}(X, x)$ and by Tannakian duality, it gives line bundle in $\mathcal{C}^{\text{EN}}(X)$. So, the group of characters of $\pi^{\text{EN}}(X, x)$ is isomorphic to the group of line bundles in $\mathcal{C}^{\text{EN}}(X)$. By the definition of semi–essentially finite bundle and by Lemma 3.1, we conclude that the character group of $\pi^{\text{EN}}(X, x)$ is the group of torsion line bundles \mathbb{F}_{τ} . By [18, 9.5], $\pi^{\text{EN}}_{\text{ab}}(X, x)$ is a product of unipotent and diagonal part, and hence the result follows.

Theorem 3.3. Let A be an abelian variety defined over an algebraically closed field k of characteristic zero. Then, we have

$$\pi^{\mathrm{EN}}(A,0) \simeq \pi^{N}(A,0) \times \pi^{\mathrm{uni}}(A,0).$$

Proof. From the Theorem 3.2 and [11, Theorem 6.1], the unipotent part of $\pi^{\text{EN}}(A, 0)$ is $H^1(X, \mathcal{O}_X)^*$.

Since $\mathbb{F}_{\tau} = \{L \in \operatorname{Pic}^{0} A : L^{\otimes n} \simeq \mathcal{O}_{A} \text{ for some } n \in \mathbb{N}\}$, we can write \mathbb{F}_{τ} as a direct limit of $U_{i} := \{L \in \operatorname{Pic}^{0} A : L^{\otimes i} \simeq \mathcal{O}_{A}\}.$

$$\mathbb{F}_{\tau} = \varinjlim U_i$$
$$k[\mathbb{F}_{\tau}] = \varinjlim k[U_i]$$
$$\text{Diag}(\mathbb{F}_{\tau}) = \text{Spec } k[\mathbb{F}_{\tau}] = \varinjlim \text{Spec } k[U_i]$$

Note that U_i is an *i*-torsion component of $\operatorname{Pic}^0 A$ and by [17, §13], $\operatorname{Pic}^0 A \simeq A^{\vee}$, where A^{\vee} is a dual abelian variety of A, i.e., U_i is an abstract group A_i^{\vee} which is isomorphic to $(\frac{\mathbb{Z}}{i\mathbb{Z}})^{2g}$ as an abstract group. So, we have the following isomorphism as a group scheme.

Spec
$$k[U_i] \simeq (\mu_i)^{2g} \simeq (\frac{\mathbb{Z}}{i\mathbb{Z}})^{2g} \simeq A_i$$

Diag $(\mathbb{F}_{\tau}) = \varprojlim \operatorname{Spec} k[U_i] = \varprojlim A_i.$

Since $\pi^N(A, 0) \simeq \underline{\lim} A_n$ [14, Remark 3], we conclude the theorem.

Remark 3.4. If char k = 0, then by applying the same method of Theorem 3.2 for the Nori fundamental group scheme, and by using [16, Proposition 2.10(1)], we have $\pi_{ab}^{N}(X, x) \simeq \text{Diag}(\mathbb{F}_{\tau})$. Hence, $\pi_{ab}^{\text{EN}}(X, x) \simeq \pi_{ab}^{\text{uni}}(X, x) \times \pi_{ab}^{N}(X, x)$. For an abelian variety A, the group schemes $\pi^{\text{EN}}(A, 0), \pi^{N}(A, 0)$, and $\pi^{\text{uni}}(A, 0)$ being abelian, we can also conclude the Theorem 3.3.

Remark 3.5. If char k = p > 0, then we have $\mathcal{C}^{\text{uni}}(X) \subset \mathcal{C}^N(X)$ [13, Chapter IV, Proposition 3]. Hence, the unipotent part of $\pi_{ab}^N(X, x)$ is isomorphic to $\pi_{ab}^{\text{uni}}(X, x)$. Also, the diagonal part of $\pi_{ab}^N(X, x)$ is isomorphic to $\text{Diag}(\mathbb{F}_{\tau})$. Hence, $\pi_{ab}^{\text{EN}}(X, x) \simeq \pi_{ab}^N(X, x)$. For an abelian variety A, we get $\pi^{\text{EN}}(A, 0) \simeq \pi^N(A, 0)$. We can give more explicit proof of this as follows:

Proposition 3.6. Let A be an abelian variety defined over an algebraically closed field k of positive characteristic, then we have

$$\pi^{\mathrm{EN}}(A,0) \simeq \pi^{N}(A,0).$$

Proof. From the Theorem 3.2 and [11, Theorem 6.1], the unipotent part of $\pi^{\text{EN}}(A, 0)$ is $T_n^0(A) \times T_n^d(A)$.

In this case, we have $U_i \simeq (\frac{\mathbb{Z}}{i\mathbb{Z}})^{2g}$, when *i* is relatively prime to *p*. For any $m \in \mathbb{N}$, we have $U_{p^m} \simeq (\frac{\mathbb{Z}}{p^m\mathbb{Z}})^r$; where $0 \leq r \leq g$, and $U_{tp^m} \simeq U_i \times U_{p^m}$ when *t* is relatively prime to *p*, as a group. Hence, we get

$$\operatorname{Diag}(\mathbb{F}_{\tau}) \simeq \operatorname{\underline{\lim}} \mathcal{H}_i$$

where \mathcal{H}_i is $(\mu_i)^{2g}$ when *i* is relatively prime to *p*, for any $m \in \mathbb{N}$, we have $\mathcal{H}_{p^m} = (\mu_{p^m})^r$; where $0 \leq r \leq g$, and $\mathcal{H}_{tp^m} = \mathcal{H}_t \times \mathcal{H}_{p^m}$ whenever *t* is not divisible by *p*. Therefore,

$$\pi^{\mathrm{EN}}(A,0) \simeq T_p^0(A) \times T_p^d(A) \times \varprojlim \mathcal{H}_i$$
$$\pi^{\mathrm{EN}}(A,0) \simeq \varprojlim A_{p^n}^0 \times \varprojlim A_{p^n}^r \times \varprojlim \mathcal{H}_i \simeq \varprojlim A_n$$

The last isomorphism follows from the decomposition of A_n and some basic property of an inverse limit. Since $\pi^N(A, 0) \simeq \varprojlim A_n$ [14, Remark 3], we conclude the theorem. \Box

Remark 3.7. If k has characteristic zero, then by Theorem 3.3, it follows that for any semi-finite bundle E on A, there is a unipotent bundle U and a finite bundle F on A such that $E \simeq U \otimes F$. If k has a positive characteristic, then by Proposition 3.6, it follows that every semi-essentially finite bundle over an abelian variety A is essentially finite. This leads to raising the following:

Question: Let X be a smooth projective variety defined over a field k of positive characteristic. Does there exist a semi–essentially finite bundle over X which is not an essentially finite?

If
$$\pi^{\acute{e}t}(X, x) = 0$$
, then by [11, Corollary 8.3], we have $\pi^{S}(X, x) \cong \pi^{EN}(X, x) \cong \pi^{N}(X, x)$.

3.2. Local unipotent group scheme. Let $\mathcal{C}^{\text{loc}}(X)$ denote the category of *F*-trivial bundle on *X*. For a rational point $x \in X(k)$, the category $\mathcal{C}^{\text{loc}}(X)$ is a Tannakian category and $\pi^{\text{loc}}(X, x)$ is the local fundamental group scheme corresponding to it (see [12]). Let us denote $\mathcal{C}^{\text{loc}}_U(X)$ the intersection category of $\mathcal{C}^{\text{loc}}(X)$ and $\mathcal{C}^{\text{uni}}(X)$. So, this category contains vector bundles that are unipotent as well as *F*-trivial. Clearly, $\mathcal{C}^{\text{loc}}_U(X)$ is a Tannakian category. Let $\pi^{\text{loc}}_U(X, x)$ denote the corresponding affine group scheme and $\pi^{\text{loc}}_U(X, x)_{ab}$ denote the maximal abelian quotient of it.

Theorem 3.8. Let X be a smooth projective variety defined over an algebraically closed field of positive characteristic. Then,

$$\pi_U^{\mathrm{loc}}(X, x)_{\mathrm{ab}} \simeq \varprojlim G_{\ell\ell}$$

where the inverse limit is taken over all finite local-local group subschemes $G_{\ell\ell}$ of $\operatorname{Pic}^0 X$.

Proof. We have

$$\pi^{\rm uni}_{\rm ab}(X,x) \simeq \varprojlim \hat{G}$$

where \hat{G} denotes the Cartier dual of G, and the inverse limit is taken over all finite local group subschemes G of $\operatorname{Pic}^0 X$. Since G is a finite local group scheme, we can decompose it as a direct sum of local-local $G_{\ell\ell}$ and local-reduced $G_{\ell\nu}$ group scheme.

$$G \simeq G_{\ell\ell} \oplus G_{\ell},$$
$$\hat{G} \simeq \hat{G}_{\ell\ell} \oplus \hat{G}_{\ell},$$

Recall that a local-local group scheme means the group scheme and its Cartier dual are both local group schemes. Therefore, the dual of local-local group scheme $\hat{G}_{\ell\ell}$ is also a local-local group scheme. The local-reduced group scheme means the group scheme is local, and its Cartier dual is a reduced group scheme. Therefore, the dual of local-reduced scheme $\hat{G}_{\ell *}$ is reduced (more specifically reduced-local).

By the universal property of $\pi^{\text{loc}}(X, x)$, we have $\pi^{\text{loc}}(X, x)$ as the inverse limit of the system of a finite local group scheme, which decomposes into the local-local and local-reduced part. Since $\mathcal{C}_U^{\text{loc}}(X) = \mathcal{C}^{\text{loc}}(X) \cap \mathcal{C}^{\text{uni}}(X)$, we need to find the common component of group scheme in $\pi_{\text{ab}}^{\text{uni}}(X, x)$ and $\pi^{\text{loc}}(X, x)$, which is a local-local group scheme as mentioned above. Hence,

$$\pi_U^{\mathrm{loc}}(X, x)_{\mathrm{ab}} \simeq \varprojlim G_{\ell\ell}$$

where the inverse limit is taken over all finite local-local group subschemes $G_{\ell\ell}$ of $\operatorname{Pic}^0 X$.

Remark 3.9. Let C be an ordinary elliptic curve over an algebraically closed field of characteristic p > 0. Let $\mathcal{I}_{(r,0)}$ denote the isomorphism classes of rank r and degree 0 indecomposable vector bundle on C. From Atiyah's classification of vector bundles over an elliptic curve [2], there is rank r vector bundle $E_r \in \mathcal{I}_{(r,0)}$ such that all other vector bundles in $\mathcal{I}_{(r,0)}$ is of the form $E_r \otimes L$ for some $L \in \text{Pic}^0 C$ also there is an exact sequence

$$0 \to E_{r-1} \to E_r \to \mathcal{O}_C \to 0$$

Hence, E_r is an unipotent vector bundle. From [15, Proposition 2.10], we have $(F_C^n)^* E_r \simeq E_r$ for all $n \in \mathbb{N}$ and hence E_r cannot be *F*-trivial bundle.

Theorem 3.10. Let A be an abelian variety of p-rank r defined over an algebraically closed field k of characteristic p > 0 then, we have

$$\pi^{\mathrm{loc}}(A,0) \simeq T_p^0(A) \times \varprojlim(\mu_{p^n})^r$$

Proof. Diagonal part: As we know, the group of characters is the same as the group of one-dimensional representations and by Tannaka duality, which is the same as the group of line bundles in $\mathcal{C}^{\text{loc}}(A)$. Let \mathbb{F}_p denote the group of line bundle in $\mathcal{C}^{\text{loc}}(A)$.

$$\mathbb{F}_p = \{ L \in \mathcal{C}^{\mathrm{loc}}(A) : (F^n)^* L \simeq \mathcal{O}_A \text{ for some } n \in \mathbb{N} \}$$
$$\mathbb{F}_p = \{ L \in \mathcal{C}^{\mathrm{loc}}(A) : L^{\otimes p^n} \simeq \mathcal{O}_A \text{ where } n \in \mathbb{N} \}$$
$$\mathbb{F}_p \simeq \varinjlim \left(\frac{\mathbb{Z}}{p^n \mathbb{Z}} \right)^r$$
$$\operatorname{Spec} k[\mathbb{F}_p] \simeq \varprojlim (\mu_{p^n})^r$$

Unipotent part: Unipotent part is the affine group scheme corresponding to the Tannakian category $\mathcal{C}^{\text{loc}}(A) \cap \mathcal{C}^{\text{uni}}(A)$, that is $\pi_U^{\text{loc}}(A, 0)$. By Theorem 3.8, we have

$$\pi_U^{\mathrm{loc}}(A,0) \simeq \varprojlim G_{\ell\ell} \simeq T_p^0(A)$$

where the inverse limit is taken over all finite local-local group subschemes $G_{\ell\ell}$ of Pic⁰A. Hence, by [18, 9.5], the result follows.

Remark 3.11. From the above Theorem 3.10, one can observe that the maximal abelian quotient of local fundamental group scheme denoted by $\pi_{ab}^{loc}(X, x)$ is isomorphic to the product of $\pi_U^{loc}(X, x)_{ab}$ and $\text{Diag}(\mathbb{F}_p)$, where \mathbb{F}_p denote the abstract group of p^n -torsion line bundles for all $n \in \mathbb{N}$, which is subgroup of $\text{Pic}^{\tau} X$.

4. Properties

4.1. The Albanese morphism. Let X be a smooth projective variety over an algebraically closed field k and $x \in X$ be a fixed point. The variety Alb X is dual to the reduced scheme underlying $\operatorname{Pic}^{0} X$. Note that Alb X is an abelian variety and let $alb_{X} : X \longrightarrow \operatorname{Alb} X$ be the Albanese morphism mapping x to $0_{\operatorname{Alb} X}$. So, there is an induced homomorphism $\psi : \pi_{\operatorname{ab}}^{\operatorname{EN}}(X, x) \longrightarrow \pi^{\operatorname{EN}}(\operatorname{Alb} X, 0_{\operatorname{Alb} X})$. In this section, we aim to find the kernel of the homomorphism ψ . The kernel of multiplication by n map $n_{S} : S \longrightarrow S$ is denoted by ${}_{n}S$ for a commutative group scheme S.

If the characteristic of k is positive, then we have $\pi_{ab}^{EN}(X, x) \simeq \pi_{ab}^{N}(X, x)$ (see Remark 3.5). By [11, Corollary 7.2], we conclude the same result for the fundamental group scheme $\pi^{EN}(X, x)$.

Theorem 4.1. Let X be a smooth projective variety over an algebraically closed field k of the characteristic zero. Then, we have the following short exact sequence:

$$0 \to \operatorname{Diag}(\operatorname{NS}(X)_{\operatorname{tor}}) \to \pi_{\operatorname{ab}}^{\operatorname{EN}}(X, x) \to \pi^{\operatorname{EN}}(\operatorname{Alb} X, 0_{\operatorname{Alb} X}) \to 0$$

Proof. Let $\mathbb{F}_{\tau}(X)$ and $\mathbb{F}_{\tau}(Alb X)$ denote the group of finite line bundles on X and Alb X respectively. By Theorem 3.2, we have the following

$$\pi_{\mathrm{ab}}^{\mathrm{EN}}(X, x) \simeq H^1(X, \mathcal{O}_X)^* \times \mathrm{Diag}(\mathbb{F}_{\tau}(X)),$$

and by replacing X with Alb X, we have

$$\pi^{\mathrm{EN}}(\mathrm{Alb}\,X, 0_{\mathrm{Alb}\,X}) \simeq H^1(\mathrm{Alb}\,X, \mathcal{O}_{\mathrm{Alb}\,X})^* \times \mathrm{Diag}(\mathbb{F}_{\tau}(\mathrm{Alb}\,X))$$

Since AlbX is an abelian variety over the field of characteristic zero, we have $\operatorname{Pic}^{\tau}(\operatorname{Alb} X) \simeq \operatorname{Pic}^{0}(\operatorname{Alb} X)$ and $\operatorname{Pic}^{0}(\operatorname{Alb} X) \simeq (\operatorname{Pic}^{0} X)_{\operatorname{red}} \simeq \operatorname{Pic}^{0} X$.

Let $T_0(\operatorname{Pic}^0 X)$ denote the tangent space at 0. We know that $T_0(\operatorname{Pic}^0 X) \simeq H^1(X, \mathcal{O}_X)$. On the other hand, we have $T_0(\operatorname{Pic}^0(\operatorname{Alb} X)) \simeq H^1(\operatorname{Alb} X, \mathcal{O}_{\operatorname{Alb} X})$. As $\operatorname{Pic}^0(\operatorname{Alb} X) \simeq$ $\operatorname{Pic}^0 X$, we get $H^1(X, \mathcal{O}_X) \simeq H^1(\operatorname{Alb} X, \mathcal{O}_{\operatorname{Alb} X})$. Hence, the unipotent part of Ker ψ is trivial.

As $\mathbb{F}_{\tau}(X)$ denote the torsion part of the group $\operatorname{Pic}^{\tau} X$, so we can write

$$\mathbb{F}_{\tau}(X) = \varinjlim_{n \in \mathbb{N}} {}_{n} \operatorname{Pic}^{\tau} X$$

and

$$\mathbb{F}_{\tau}(\operatorname{Alb} X) = \lim_{n \in \mathbb{N}} {}_{n}\operatorname{Pic}^{\tau}(\operatorname{Alb} X) = \lim_{n \in \mathbb{N}} {}_{n}\operatorname{Pic}^{0} X.$$

Since

$$0 \to \operatorname{Pic}^0 X \to \operatorname{Pic}^\tau X \to \operatorname{NS}(X)_{\operatorname{tor}} \to 0$$

is an exact sequence, we get an exact sequence

$$0 \rightarrow {}_{n}\operatorname{Pic}^{0} X \rightarrow {}_{n}\operatorname{Pic}^{\tau} X \rightarrow {}_{n}\operatorname{NS}(X)_{\operatorname{tor}} \rightarrow 0$$

for all $n \in \mathbb{N}$.

Since $NS(X)_{tor}$ is finite, there is some $N \in \mathbb{N}$ such that $_N NS(X)_{tor} = NS(X)_{tor}$ and by taking direct limit, we get

$$0 \to \mathbb{F}_{\tau}(\operatorname{Alb} X) \to \mathbb{F}_{\tau}(X) \to \operatorname{NS}(X)_{\operatorname{tor}} \to 0$$
.

So, we have an exact sequence

$$0 \to \operatorname{Diag}(\operatorname{NS}(X)_{\operatorname{tor}}) \to \operatorname{Diag}(\mathbb{F}_{\tau}(X)) \to \operatorname{Diag}(\mathbb{F}_{\tau}(\operatorname{Alb} X)) \to 0.$$

Therefore, the diagonal part of Ker ψ is Diag(NS(X)_{tor}).

Proposition 4.2. Let X be a smooth projective variety over an algebraically closed field k of the characteristic p > 0. Then, we have the following short exact sequence:

$$0 \to \varprojlim_{G \subset \operatorname{Pic}^{0} X} G/G_{\operatorname{red}} \times \operatorname{Diag}\left(\varinjlim_{n \in \mathbb{N}} {}_{p^{n}} \operatorname{NS}(X)_{\operatorname{tor}} \right) \to \pi_{\operatorname{ab}}^{\operatorname{loc}}(X, x) \to \pi^{\operatorname{loc}}(\operatorname{Alb} X, 0_{\operatorname{Alb} X}) \to 0$$

where the inverse limit is taken over all the local-local group subscheme G of $\operatorname{Pic}^{0}X$.

Proof. The Albenese morphism $alb_X : X \longrightarrow Alb X$ induced a morphism between the local fundamental group scheme $\varphi : \pi_{ab}^{loc}(X, x) \longrightarrow \pi^{loc}(Alb X, 0_{Alb X})$. By Remark 3.11, we have

$$\pi_{\rm ab}^{\rm loc}(X, x) \simeq \varprojlim_{G_{\ell\ell} \subset \operatorname{Pic}^0 X} G_{\ell\ell} \times \operatorname{Diag}(\mathbb{F}_p)$$

where the inverse limit is taken over all the local-local group subschemes $G_{\ell\ell}$ of $\operatorname{Pic}^0 X$ and \mathbb{F}_p is the subgroup of $(\operatorname{Pic}^{\tau} X)_{\mathrm{red}}$.

Since $\operatorname{Pic}^{\tau}(\operatorname{Alb} X) \simeq \operatorname{Pic}^{0}(\operatorname{Alb} X) \simeq (\operatorname{Pic}^{0} X)_{\operatorname{red}}$, we have

$$\pi^{\mathrm{loc}}(\mathrm{Alb}\,X, 0_{\mathrm{Alb}\,X}) \simeq \varprojlim_{(G_{\ell\ell})_{\mathrm{red}} \subset (\mathrm{Pic}^0 X)_{\mathrm{red}}} (G_{\ell\ell})_{\mathrm{red}} \times \mathrm{Diag}(\mathbb{F}_p(\mathrm{Alb}\,X))$$

where the inverse limit is taken over all the local-local group subschemes $(G_{\ell\ell})_{\rm red}$ of $(\operatorname{Pic}^0 X)_{\rm red}$ and $\mathbb{F}_p(\operatorname{Alb} X)$ is the subgroup of $\operatorname{Pic}^{\tau}(\operatorname{Alb} X) \simeq (\operatorname{Pic}^0 X)_{\rm red}$.

We have the following exact sequence:

$$0 \to (G_{\ell\ell})_{\mathrm{red}} \to G_{\ell\ell} \to G_{\ell\ell}/(G_{\ell\ell})_{\mathrm{red}} \to 0$$
.

By taking dual and passing through the inverse limit, we get the unipotent part of the kernel.

Note that $\mathbb{F}_p(X)$ denote the group of p^n torsion line bundles for all $n \in \mathbb{N}$. We can write

$$\mathbb{F}_p(X) = \lim_{n \in \mathbb{N}} {}_{p^n}(\operatorname{Pic}^{\tau} X)_{\operatorname{red}}$$

Similarly, we have

$$\mathbb{F}_p(\operatorname{Alb} X) = \varinjlim_{n \in \mathbb{N}} {}_{p^n}(\operatorname{Pic}^0 X)_{\operatorname{red}}$$

For all $n \in \mathbb{N}$, we have the following exact sequence:

$$0 \rightarrow {}_{p^n}(\operatorname{Pic}^0 X)_{\operatorname{red}} \rightarrow {}_{p^n}(\operatorname{Pic}^\tau X)_{\operatorname{red}} \rightarrow {}_{p^n}\operatorname{NS}(X)_{\operatorname{tor}} \rightarrow 0$$

By passing through the direct limit, we get the diagonal part of the kernel.

4.2. Birational invariance.

Proposition 4.3. Let Y be a smooth complete variety over an algebraically closed field k and let $f: X \longrightarrow Y$ be the blow-up of Y along a smooth subvariety $Z \subset Y$. Then $\pi^{\text{EN}}(X, x) \longrightarrow \pi^{\text{EN}}(Y, f(x))$ is an isomorphism.

Proof. Let $E \in \mathcal{C}^{\text{EN}}(X)$. Then by [10, Lemma 8.3], we have f_*E is locally free and $E \simeq f^*f_*E$ as $\mathcal{C}^{\text{EN}}(X)$ is the full subcategory of $\mathcal{C}^{nf}(X)$. By [10, Lemma 8.3] and $\pi^{\text{EN}}(X, x)$ is a quotient of $\pi^S(X, x)$, we have the following commutative diagram in which the horizontal maps are surjective and the above vertical map is an isomorphism. So, the

below map $\pi^{\text{EN}}(X, x) \longrightarrow \pi^{\text{EN}}(Y, f(x))$ is faithfully flat.

$$\begin{array}{ccc} \pi^S(X,x) & \xrightarrow{\simeq} & \pi^S(Y,f(x)) \\ & p & & \downarrow^p \\ \pi^{\mathrm{EN}}(X,x) & \longrightarrow & \pi^{\mathrm{EN}}(Y,f(x)) \end{array}$$

To see that $\pi^{\text{EN}}(X, x) \longrightarrow \pi^{\text{EN}}(Y, f(x))$ is a closed immersion, by [5, Proposition 2.21(b)], it is enough to prove that for every object $E \in \mathcal{C}^{\text{EN}}(X)$, there is an object $E' \in \mathcal{C}^{\text{EN}}(Y)$ such that $E \simeq f^*E'$.

Suppose E is an essentially finite bundle on X. By [4, Theorem 1.1] there is a proper k-scheme P with a proper surjective morphism $p: P \longrightarrow X$ such that p^*E is trivial.

Since $(f \circ p)^*(f_*E) \simeq p^*f^*f_*E \simeq p^*E$ is trivial and $f \circ p$ is proper morphism, it follows that f_*E is an essentially finite bundle. If E is semi-essentially finite bundle on X, then by induction on rank, we can conclude that f_*E is semi-essentially finite bundle. To see this, let E be a semi-essentially finite bundle of rank r on X, i.e., there is a filtration

$$E = E_0 \supset E_1 \supset E_2 \supset \ldots \supset E_{n-1} \supset E_n = 0$$

such that E_i/E_{i+1} is an indecomposable essentially finite bundle. From the filtration, E_{n-1} is an essentially finite subbundle of E. Then, the bundle f_*E_{n-1} is an essentially finite. Let $F = E/E_{n-1}$. Note that F is a semi–essentially finite bundle and rank $F < \operatorname{rank} E$ so, by induction, f_*F is semi–essentially finite bundle. We have the following exact sequences:

$$(4.1) 0 \longrightarrow E_{n-1} \longrightarrow E \longrightarrow F \longrightarrow 0$$

$$(4.2) 0 \longrightarrow f_* E_{n-1} \longrightarrow f_* E \longrightarrow f_* F$$

For any numerically flat bundle V, we have $f^*f_*V \simeq V$ and hence rank $f_*V = \operatorname{rank} V$. It follows that the last map of (4.2) is surjective. Hence, f_*E is a semi-essentially finite bundle.

By the weak factorization theorem [19, 12.4] and Proposition 4.3, we have the following:

Corollary 4.4. Let X and Y be smooth projective varieties over an algebraically closed field k of characteristic zero and $\phi : X \dashrightarrow Y$ be a birational map. Let $x_0 \in X(k)$ be a point where ϕ is defined. Then $\pi^{\text{EN}}(X, x_0)$ is isomorphic to $\pi^{\text{EN}}(Y, \phi(x_0))$.

Corollary 4.5. Let X and Y be smooth projective varieties over an algebraically closed field k of positive characteristic and $\phi: X \dashrightarrow Y$ be a birational map. Let $x_0 \in X(k)$ be a point where ϕ is defined. Then, we have $\pi^{EN}(X, x_0) \simeq \pi^{EN}(Y, \phi(x_0))$ and $\pi^{\text{loc}}(X, x_0) \simeq \pi^{\text{loc}}(Y, \phi(x_0))$. *Proof.* If dim $X \leq 2$, then the result follows from [7, Chapter-V, Proposition 5.3] and Proposition 4.3. Assume that dim $X \geq 3$. Consider the diagram



where Z is the normalization of a closure of the graph of ϕ , the morphisms p and q are birational. In view of [8, Theorem 1.2], we need to show that q_*p^*E is a semi–essentially finite whenever E is a semi–essentially finite bundle on X.

Since q_*p^* commutes with the tensor product, it follows that q_*p^*E is a finite bundle whenever E is a finite bundle on X. Let E be an essentially finite bundle on X. Then, there is $V_1, V_2 \in \mathcal{C}^{\mathrm{nf}}(X)$ such that V_2 is subbundle of V_1, V_1 is subbundle of finite bundle and $E \simeq V_1/V_2$. We have the following exact sequence:

$$0 \to V_2 \to V_1 \to V_1/V_2 \to 0$$
$$0 \to q_* p^* V_2 \to q_* p^* V_1 \to q_* p^* (V_1/V_2) \to 0$$

First note that for any $V \in \mathcal{C}^{nf}(X)$, the rank of q_*p^*V is the same as that of V. To see this, let us adopt the setting mentioned in proof of [8, Theorem 1.2]. Let S be a general smooth surface in Y. Since $q: Z \to Y$ is birational, there exists a smooth projective surface T and a proper birational morphism $\pi: T \longrightarrow S$ with the following commutative diagram:

$$(4.3) \qquad \begin{array}{c} T \xrightarrow{h} Z \\ \pi \downarrow \qquad \qquad \downarrow q \\ S \xrightarrow{i} Y \end{array}$$

Let $W = p^*V$. Then, the morphism $i^*q_*W \to \pi_*h^*W$ is an isomorphism. Hence, we get

 $(i \circ \pi)^* q_* W = \pi^* i^* q_* W \simeq \pi^* \pi_* h^* W \simeq h^* W$

in which last isomorphism follows from [10, Lemma 8.3]. By the commutativity of diagram 4.3, we have $h^*W \simeq (i \circ \pi)^*q_*W \simeq (q \circ h)^*q_*W = h^*q^*q_*W$. Therefore, the rank of W is the same as the rank of q_*W .

Hence, $q_*p^*E \simeq q_*p^*(V_1/V_2) \simeq (q_*p^*V_1)/(q_*p^*V_2)$. This proves that q_*p^*E is an essentially finite bundle on Y.

By the similar arguments as in the proof of Proposition 4.3, using induction on rank, we can prove q_*p^*E is a semi-essentially finite bundle whenever E is a semi-essentially finite bundle.

To see that $\pi^{\text{loc}}(X, x_0) \simeq \pi^{\text{loc}}(Y, \phi(x_0))$, it is enough to prove q_*p^*V is *F*-trivial bundle on *Y* whenever *V* is *F*-trivial on *X*. Let *V* be *F*-trivial bundle on *X*. Then, $W := p^*V$ is *F*-trivial on *Z*. i.e. $(F_Z^n)^*(W) \simeq \mathcal{O}_Z^{\oplus r}$ for some $n, r \in \mathbb{N}$. From [8, Lemma 3.1], we have $(F_X^n)^*(q_*W) \simeq q_*((F_Z^n)^*(W)) \simeq q_*(\mathcal{O}_Z^{\oplus r}) \simeq (q_*\mathcal{O}_Z)^{\oplus r} = (\mathcal{O}_Y)^{\oplus r}$. Hence, q_*p^*V is *F*-trivial on *Y*.

4.3. Behaviour under étale morphism. Let $\psi : X \longrightarrow Y$ be an étale Galois cover with a finite Galois group G. Let $x_0 \in X$ and $y_0 \in Y$ be k-points such that $f(x_0) = y_0$. The pullback by ψ induces a map $\psi^* : \mathcal{C}^{\text{EN}}(Y) \longrightarrow \mathcal{C}^{\text{EN}}(X)$ which is a morphism of Tannakian categories. So, this induces a morphism $\tilde{\psi} : \pi^{\text{EN}}(X, x_0) \longrightarrow \pi^{\text{EN}}(Y, y_0)$. Consider the following diagram:



For any $E \in \mathcal{C}^{EN}(X)$, we have the followings:

$$\psi^*\psi_*E \simeq p_*p^*E \simeq p_{1*}\mu^*E \simeq \bigoplus_{g \in G} g^*E$$

Proposition 4.6. The map $\tilde{\psi} : \pi^{\text{EN}}(X, x_0) \longrightarrow \pi^{\text{EN}}(Y, y_0)$ is a closed immersion.

Proof. If E is an essentially finite bundle on X, then there exist a connected étale Galois cover $\gamma : \hat{X} \longrightarrow X$ such that $\gamma^* E$ is F-trivial bundle (if char k = 0, then $\gamma^* E$ is trivial). Note that $\psi \circ \gamma : \hat{X} \longrightarrow Y$ étale covering of Y. Since

$$(\psi \circ \gamma)^* (\psi_* E) \simeq \gamma^* \psi^* \psi_* E \simeq \gamma^* (\bigoplus_{g \in G} E) \simeq \bigoplus_{g \in G} \gamma^* g^* E$$

,

it follows that $(\psi \circ \gamma)^*(\psi_* E)$ is *F*-trivial (or trivial, if char k = 0). Hence, the bundle $\psi_* E$ is an essentially finite.

Let E be a semi–essentially finite bundle of rank r on X, i.e., there is a filtration

 $E = E_0 \supset E_1 \supset E_2 \supset \ldots \supset E_{n-1} \supset E_n = 0$

such that E_i/E_{i+1} is an indecomposable essentially finite bundle. From the filtration, E_{n-1} is essentially finite subbundle of E and E/E_{n-1} is semi–essentially finite bundle on X. Then, ψ_*E_{n-1} is essentially finite and by induction, we have $\psi_*(E/E_{n-1})$ is semi–essentially finite bundle.

Let r(V) denote the rank of vector bundle V. Consider the following exact sequence:

$$(4.4) 0 \longrightarrow \psi_* E_{n-1} \longrightarrow \psi_* E \longrightarrow \psi_* (E/E_{n-1})$$

We have

$$r(\psi_*(E/E_{n-1})) = |G| \cdot r(E/E_{n-1}) = |G| \cdot (r(E) - r(E_{n-1})) = r(\psi_*E) - r(\psi_*E_{n-1}).$$

Hence, the last map of (4.4) is surjective. Since the category of semi–essentially finite bundle is closed under taking extensions, it follows that ψ_*E is a semi–essentially finite bundle.

We need to prove that any $E \in \mathcal{C}^{EN}(X)$ is subquotient of the $\psi^* E'$ for some $E' \in \mathcal{C}^{EN}(Y)$. Let $E' = \psi_* E$ then $\psi^* \psi_* E \simeq \bigoplus_{g \in G} g^* E$. Clearly, E is subquotient of $\bigoplus_{g \in G} g^* E$. Hence, $\tilde{\psi}$ is a closed immersion.

By using the criteria [6, Theorem A.1] and Proposition 4.6, we have:

Corollary 4.7. If $\psi : X \longrightarrow Y$ is an étale Galois cover with a finite Galois group G, then we get the following exact sequence:

$$0 \to \pi^{\mathrm{EN}}(X, x_0) \to \pi^{\mathrm{EN}}(Y, y_0) \to G \to 0.$$

5. An extension of the local fundamental group scheme

Let X be a smooth projective variety over the algebraically closed field of characteristic p > 0. Let $F_X \colon X \longrightarrow X$ be a Frobenius morphism.

Definition 5.1. A vector bundle V on X is called F-semi trivial if there is a filtration

$$V = V_0 \supset V_1 \supset \ldots \supset V_{n-1} \supset V_n = 0$$

such that V_i/V_{i+1} is indecomposable F-trivial bundle on X for all $i \in \{0, ..., n-1\}$.

Let $\mathcal{C}_E^{\text{loc}}(X)$ denote the category of *F*-semi trivial bundles on *X*. Note that simple objects of this category are indecomposable *F*-trivial bundles on *X*, and this category is closed under an extension. By following arguments of [16, Proposition 2.14], one can prove that the category $\mathcal{C}_E^{\text{loc}}(X)$ is *k*-linear abelian rigid tensor category. We give details for completion in the following Proposition 5.2.

Proposition 5.2. (cf. [16, Proposition 2.14]) The category $\mathcal{C}_E^{\text{loc}}(X)$ is k-linear abelian rigid tensor category.

Proof. First, let us prove $\mathcal{C}_E^{\text{loc}}(X)$ is closed under the tensor product and dual. If E is F-trivial and F is F-semi trivial then we have the following filtration

$$F \otimes E = F_0 \otimes E \supset F_1 \otimes E \supset \ldots \supset F_{n-1} \otimes E \supset F_n \otimes E = 0$$

such that $(F_i \otimes E)/(F_{i+1} \otimes E) \simeq (F_i/F_{i+1}) \otimes E$ is *F*-trivial for all *i*. So, $F \otimes E$ is *F*-semi trivial. Now, let *E* and *F* are *F*-semi trivial bundles. We have the filtration

$$E = E_0 \supset E_1 \supset \dots \supset E_{m-1} \supset E_m = 0$$

such that E_i/E_{i+1} is indecomposable F-trivial for all i. After tensoring with F we have,

$$E \otimes F = E_0 \otimes F \supset E_1 \otimes F \supset \dots \supset E_{m-1} \otimes F \supset E_m \otimes F = 0$$

such that $(E_i \otimes F)/(E_{i+1} \otimes F) \simeq E_i/E_{i+1} \otimes F$ is F-semi trivial.

We have the following exact sequence:

$$0 \to E_{i+1} \otimes F \to E_i \otimes F \to E_i / E_{i+1} \otimes F \to 0$$

As the category $\mathcal{C}_E^{\text{loc}}(X)$ closed under extensions and by induction, we have $E \otimes F$ is *F*-semi trivial. To prove E^{\vee} is *F*-semi trivial, let us consider $(E^{\vee})_i = (E/E_{m-i})^{\vee}$.

$$E_{m-i} \subset E_{m-i-1} \Rightarrow E/E_{m-i} \supset E/E_{m-i-1}$$
$$\Rightarrow (E/E_{m-i})^{\vee} \supset (E/E_{m-i-1})^{\vee} \Rightarrow (E^{\vee})_i \supset (E^{\vee})_{i+1}$$

This gives the filtration

$$E^{\vee} = (E^{\vee})_0 \supset (E^{\vee})_1 \supset \dots \supset (E^{\vee})_{m-1} \supset (E^{\vee})_m = 0$$

such that $(E^{\vee})_i/(E^{\vee})_{i+1} \simeq (E_{m-i-1}/E_{m-i})^{\vee}$ which is *F*-trivial for all *i*. Similarly, by following the arguments as in the proof of [16, Proposition 2.14], it can be checked that $\mathcal{C}_E^{\text{loc}}(X)$ is an abelian category.

For a rational point $x \in X(k)$, the category $\mathcal{C}_E^{\text{loc}}(X)$ with the fiber functor x^* : $\mathcal{C}_E^{\text{loc}}(X) \longrightarrow \text{Vect}_k$ is a neutral Tannakian category. Let $\pi_E^{\text{loc}}(X, x)$ denote the corresponding affine group scheme. In general, the group scheme $\pi_E^{\text{loc}}(X, x)$ is bigger than $\pi^{\text{loc}}(X, x)$ (see Remark 3.9).

Theorem 5.3. Let X be a smooth projective variety defined over an algebraically closed field k of characteristic p > 0. Let $x \in X(k)$ be a rational point then, we have

$$\pi_E^{\mathrm{loc}}(X, x)_{\mathrm{ab}} \simeq \pi_{\mathrm{ab}}^{\mathrm{uni}}(X, x) \times \mathrm{Diag}(\mathbb{F}_p)$$

Proof. Note that $\mathcal{C}^{\text{uni}}(X)$ is the full subcategory of $\mathcal{C}_E^{\text{loc}}(X)$. So, unipotent part of $\pi_E^{\text{loc}}(X, x)_{\text{ab}}$ is $\pi_{\text{ab}}^{\text{uni}}(X, x)$.

For a diagonal part, a group of line bundles in $\mathcal{C}_E^{\text{loc}}(X)$ is same as the group \mathbb{F}_p of line bundles in $\mathcal{C}^{\text{loc}}(X)$. Hence, the theorem follows from the decomposition of a commutative group scheme [18, 9.5].

Corollary 5.4. Let A be an abelian variety of p-rank r defined over an algebraically closed field k of characteristic p > 0 then, we have

$$\pi_E^{\mathrm{loc}}(A,0) \simeq \pi^{\mathrm{uni}}(A,0) \times \varprojlim_{n \in \mathbb{N}} (\mu_{p^n})^r$$

Proof. The result follows using Theorem 5.3 and Theorem 3.10.

Corollary 5.5. We have the following exact sequence:

$$0 \to \lim_{G \subset \operatorname{Pic}^{0} X} \widehat{G/G_{red}} \times \operatorname{Diag}\left(\lim_{n \in \mathbb{N}} p^{n} \operatorname{NS}(X)_{\operatorname{tor}}\right) \to \pi_{E}^{\operatorname{loc}}(X, x)_{\operatorname{ab}} \to \pi_{E}^{\operatorname{loc}}(\operatorname{Alb} X, 0_{\operatorname{Alb} X}) \to 0$$

where the inverse limit is taken over all the local group subschemes G of $\operatorname{Pic}^{0}X$.

Proof. We know that the unipotent part of $\pi_E^{\text{loc}}(X, x)_{ab}$ is same as unipotent part of $\pi_{ab}^N(X, x)$ and the diagonal part of $\pi_E^{\text{loc}}(X, x)_{ab}$ is same as diagonal part of $\pi^{\text{loc}}(X, x)_{ab}$. Hence, the lemma follows from [11, Corollary 7.2] and Proposition 4.2.

Remark 5.6. (1) If $\pi^{\text{ét}}(X, x) = 0$, then all horizontal maps in the following diagram



are isomorphisms [11, Proposition 8.2].

(2) The birational invariance property of the extended local fundamental group scheme follows from the birational invariance property of the local fundamental group scheme (Corollary 4.5) and by the same argument mentioned in Proposition 4.3.

References

- S. Amrutiya, A note on certain Tannakian group schemes, Archivum Mathematicum, Vol. 56 (2020), No. 1, 21-29.
- [2] M. F. Atiyah, Vector bundles over an elliptic curve, Proc. London Math. Soc. 3 (1957) 414-452.
- [3] I. Biswas, Y. Holla, Comparison of fundamental group schemes of a projective variety and an ample hypersurface, Jour. Alg. Geom. 16 (2007), 547-597.
- [4] I. Biswas, J. P. P. Dos Santos, Vector bundles trivialized by proper morphisms and the fundamental group scheme. Journal of the Institute of Mathematics of Jussieu, 10 (2011), pp 225-234.
- [5] P. Deligne, J. S. Milne, Tannakian Categories, in: Hodge cycles, motives and Shimura varieties (by P. Deligne, J. S. Milne, A. Ogus and K. Y. Shih), pp. 101-228, Lecture Notes in Mathematics, 900, Springer-Verlag, Berlin-Heidelberg- New York, 1982
- [6] H. Esnault, P. H. Hai, X. Sun, On Nori's fundamental group scheme. In Geometry and dynamics of groups and spaces, Progr. Math. 265, pages 377–398. Birkhäuser, Basel, 2008.
- [7] R. Hartshorne, Algebraic Geometry, Grad. Texts Maths., 52, Springer, New York-Heidelberg-Berlin, 1977.
- [8] A. Hogadi, V. B. Mehta, Birational invariance of the S-fundamental group scheme Pure Appl. Math. Q. 7 (2011), no. 4, Special Issue: In memory of Eckart Viehweg, 1361-1369.
- [9] S. L. Kleiman, Fundamental Algebraic Geometry: Grothendieck's FGA explained; *The Picard Scheme*, American Mathematical Society, Providence, RI, 2005.
- [10] A. Langer, On S-fundamental group scheme, Ann. Inst. Fourier (Grenoble) 61 (2011), no. 5, 2077-2119.
- [11] A. Langer, On S-fundamental group scheme II, J. Inst. Math. Jussieu 11 (2012), 835-854.
- [12] V. Mehta, S. Subramanian, On the fundamental group scheme. Invent. Math. 148, (2002) 143–150.
- [13] M. V. Nori, The fundamental group-scheme, Proc. Indian Acad. Sci. Math. Sci. 91 (1982), no. 2, p. 73-122.
- [14] M. V. Nori, The fundamental group-scheme of an abelian variety, Math. Annalen 263 (1983), 263–266.
- [15] T. Oda, Vector bundles on an elliptic curve, Nagoya Math. J. 43 (1971), 41-72.
- [16] S. Otabe, An extension of Nori fundamental group, Comm. Algebra 45 (2017), 3422-3448.
- [17] D. Mumford, Abelian varieties, Tata Institute of Fundamental Research Studies in Mathematics, Volume 5 (Oxford University Press, 1970).
- [18] W. C. Waterhouse, Introduction to affine group schemes, Graduate Texts in Mathematics, Volume 66 (Springer, 1979).
- [19] Włodarczyk Jarosław; Toroidal varieties and the weak factorization theorem. Invent. Math. 154 (2003), no. 2, 223–331.

Department of Mathematics, IIT Gandhinagar, Near Village Palaj, Gandhinagar - 382355, India

Email address: pavan.a@iitgn.ac.in

Department of Mathematics, IIT Gandhinagar, Near Village Palaj, Gandhinagar - 382355, India

 $Email \ address: \verb"samrutiya@iitgn.ac.in"$