Continuity of a spatial gradient of a weak solution to a very singular parabolic equation involving the one-Laplacian

Shuntaro Tsubouchi [∗]

Abstract

We consider weak solutions to very singular parabolic equations involving a one-Laplace-type operator, which is singular and degenerate, and a p-Laplace-type operator with $\frac{2n}{n+2} < p < \infty$, where $n \geq 2$ denotes the space dimension. This type of equation is used to describe the motion of a Bingham flow. It has been a long-standing open problem of whether a spatial gradient of a weak solution is continuous in space and time. This paper aims to give an affirmative answer for a wide class of such equations. This equation becomes no longer uniformly parabolic near a facet, the place where a spatial gradient vanishes. To achieve our goal, we show local a priori Hölder continuity of gradients suitably truncated near facets. For this purpose, we consider a parabolic approximate problem and appeal to standard methods, including De Giorgi's truncation and comparisons with Dirichlet heat flows. Our method is a parabolic adjustment of our method developed to prove the corresponding statements for stationary problems.

Mathematics Subject Classification (2020) 35K92, 35B65, 35A35

Keywords Continuity of a gradient, De Giorgi's truncation, Comparison arguments

Contents

[∗]Graduate School of Mathematical Sciences, The University of Tokyo, Japan. Email: tsubos@g.ecc.u-tokyo.ac.jp

1 Introduction

This paper is concerned with regularity for weak solutions to

$$
\partial_t u - \Delta_1 u - \Delta_p u = f \quad \text{in} \quad \Omega_T := \Omega \times (0, T). \tag{1.1}
$$

Here $T \in (0, \infty)$ and $p \in (1, \infty)$ are fixed constants, and $\Omega \subset \mathbb{R}^n$ is an *n*-dimensional bounded Lipschitz domain with $n \ge 2$. The real-valued functions $f = f(x, t)$ and $u = u(x, t)$ are respectively given and unknown for $(x, t) = (x_1, \ldots, x_n, t) \in \Omega_T$. The divergence operators Δ_1 and Δ_p are respectively the one-Laplacian and the p-Laplacian, defined as $\Delta_s u := \text{div} \left(|\nabla u|^{s-2} \nabla u \right)$ for $s \in [1, \infty)$. Here $\nabla u := (\partial_{x_i} u, \ldots, \partial_{x_n} u)$ and $\partial_t u$ respectively denote a spatial and time derivative of a solution $u = u(x, t)$, and div $X := \sum_{j=1}^{n} \partial_{x_j} X_j$ is divergence of a vector field $X = (X_1, \ldots, X_n)$.

The parabolic $(1, p)$ -Laplace equation (1.1) appears in describing the motion of Bingham flows with $p = 2$ (see Section [1.3\)](#page-4-0). Mathematical analysis for this parabolic equation at least goes back to the classical textbook [\[21\]](#page-32-0) in 1976. In particular, the unique existence of weak solutions for initial value problems is well-known, which is based on variational inequalities [\[21,](#page-32-0) Chapter VI]. However, the continuity of a spatial gradient has been a long-standing open problem even when $f = 0$. For stationary problems, this regularity problem was answered affirmatively by the author [\[44\]](#page-34-0), [\[45\]](#page-34-1). By adjusting the methods therein to time-evolutional problems (see Sections $1.1-1.2$ for the details), we would like to prove that a spatial gradient is continuous even for the parabolic equation (1.1) .

Throughout this paper, we let f be in a Lebesgue space $L^q(\Omega_T)$, and mainly assume

$$
p_{\rm c} < p < \infty, \quad \text{and} \quad n + 2 < q \leq \infty,\tag{1.2}
$$

where $p_c := \frac{2n}{n+2}$ $\frac{2n}{n+2}$. This assumption is typical when one considers the continuity of a spatial gradient of a p-Poisson flow (see e.g., [\[7\]](#page-32-1), [\[13,](#page-32-2) Chapters VIII–IX], [\[49,](#page-34-2) Chapter 2]). Under a weaker regularity assumption of f, local gradient bounds for a p-Poisson flow are shown in [\[32\]](#page-33-0). For a technical reason, in addition to [\(1.2\)](#page-1-3), we also assume that the continuous inclusion

$$
L^{q}(\Omega_{T}) = L^{q}(0, T; L^{q}(\Omega)) \hookrightarrow L^{p'}(0, T; W^{-1, p'}(\Omega))
$$
\n(1.3)

holds for any fixed $T \in (0, \infty)$. Here $p' := p/(p-1) \in (1, \infty)$ denotes the Hölder conjugate exponent of $p \in (1, \infty)$. Since the continuous embedding $L^q(\Omega) \hookrightarrow W^{-1,p'}(\Omega)$ is automatically satisfied by (1.2) , the assumption [\(1.3\)](#page-1-4) holds true, provided

$$
\frac{1}{p} + \frac{1}{q} \leqslant 1. \tag{1.4}
$$

It is worth mentioning that for $n \geq 4$, [\(1.4\)](#page-1-5) is automatically satisfied by [\(1.2\)](#page-1-3). Also, for the remaining case $n = 2$ or $n = 3$, [\(1.4\)](#page-1-5) is valid, provided $p \ge 2$ and [\(1.2\)](#page-1-3). Hence, (1.4) is not restrictive in applying out regularity results to mathematical models in fluid mechanics (see Section [1.3\)](#page-4-0). Under (1.2) – (1.3) , we would like to prove $\nabla u \in C^0(\Omega; \mathbb{R}^n)$.

More generally, we consider an anisotropic very singular equation

$$
\partial_t u + \mathcal{L}u = f \quad \text{in} \quad \Omega_T \tag{1.5}
$$

with $\mathcal{L}u := -\text{div}(\nabla E_1(\nabla u) + \nabla E_p(\nabla u)).$ The detailed conditions of E_1 and E_p , which are at least C^2 outside the origin, are explained later in Section [1.4.](#page-4-1) Our settings allow us to choose $E_s(z) :=$ $|z|^s/s$ $(z \in \mathbb{R}^n)$ for $s \in \{1, p\}$. In particular, \mathcal{L} generalizes $-\Delta_1 - \Delta_p$.

1.1 Our strategy

To explain the main difficulty, we formally differentiate (1.5) by x_j . Then, we have

$$
\partial_t \partial_{x_j} u - \text{div} \left(\nabla^2 E(\nabla u) \nabla \partial_{x_j} u \right) = \partial_{x_j} f \quad \text{in} \quad \Omega_T,
$$
\n(1.6)

where $E := E_1 + E_p$. Roughly speaking, we face a problem that the coefficient matrix $\nabla^2 E(\nabla u)$ in [\(1.6\)](#page-1-7) becomes no longer uniform parabolic near a facet, the degenerate region of a gradient. Here uniform parabolicity can be measured by the ratio, defined as the maximum eigenvalue of $\nabla^2 E(z)$ divided by the minimum one of $\nabla^2 E(z)$, which is well-defined for $z \in \mathbb{R}^n \setminus \{0\}$. For elliptic problems, this type of ratio is often called the ellipticity ratio (see e.g., [\[38\]](#page-33-1)). Similarly, we would like to call the ratio parabolicity ratio. For the special case $E(z) = |z| + |z|^p/p$, the parabolicity ratio of $\nabla^2 E(\nabla u)$ is behaves like $1 + |\nabla u|^{1-p}$, which blows up as $\nabla u \to 0$. Thus, it appears hard to obtain quantitative continuity estimates for $\partial_{x_i}u$. Such difficulty is substantially caused by the fact that diffusion of one-Laplacian becomes degenerate in the direction of a gradient, while this diffusion may become singular in other directions. In other words, Δ_1 has anisotropic diffusivity, which is substantially different from Δ_p . This property makes it difficult to handle Δ_1 in the existing regularity theory.

However, when one constrains a region $\{|\nabla u| \geq \delta\}$ for fixed $\delta \in (0, 1)$, then over such a place, the parabolicity ratio of $\nabla^2 E(\nabla u)$ becomes bounded by some constant that depends on δ . In this sense, we may say that [\(1.6\)](#page-1-7) is locally uniformly parabolic outside a facet. With this in mind, we would like to introduce a truncation parameter $\delta \in (0, 1)$, and consider a truncated gradient $\mathcal{G}_{\delta}(\nabla u) \coloneqq (|\nabla u| - \delta)_+ \frac{\nabla u}{|\nabla u}$ $\frac{\nabla u}{|\nabla u|}$. We aim to prove that $\mathcal{G}_{\delta}(\nabla u)$ is locally Hölder continuous in Ω_T for each fixed $\delta \in (0, 1)$. Note that this truncated gradient is supported in $\{|\nabla u| \geq \delta\}$, and therefore such local Hölder regularity will be expected. Here it should be noted that our Hölder estimates of $\mathcal{G}_{\delta}(\nabla u)$ substantially depends on $\delta \in (0, 1)$, and most of our continuity estimates will blow up as δ tends to 0. Also, it is worth mentioning that $\mathcal{G}_{\delta}(\nabla u)$ uniformly converges to $\mathcal{G}_{0}(\nabla u) = \nabla u$ in Ω_T . Thus, although some of our regularity estimates might break as δ tends to 0, we conclude that ∇u is continuous.

To give a rigorous proof, we have to appeal to approximation arguments. Here it should be recalled that the very singular operator $-\Delta_1-\Delta_p$, or more general $\mathcal L$, will become no longer uniformly parabolic near a facet of a solution. This prevents us from applying standard regularity methods, including difference quotient methods, directly to (1.1) or (1.5) . In particular, although we have formally deduced [\(1.6\)](#page-1-7), this equation seems difficult to deal with in the sense of $L^2(0, T; W^{-1,2}(\Omega))$. For this reason, we have to consider a parabolic approximate problem, whose solution has improved regularity properties. Precisely speaking, for an approximation parameter $\varepsilon \in (0, 1)$, we relax the principal part $\mathcal{L}u$ by $\mathcal{L}_{\varepsilon}u_{\varepsilon} := -\text{div}(\nabla E_{\varepsilon}(\nabla u_{\varepsilon}))$. Here E_{ε} is defined to be the convolution of $E = E_1 + E_p$ with the Friedrichs standard mollifier $j_{\varepsilon} \in C_{c}^{\infty}(\mathbb{R}^{n})$. We also approximate $f \in L^{q}(\Omega_{T})$ by $f_{\varepsilon} \in C^{\infty}(\Omega_{T})$, which converges to f in a weak sense. The resulting approximate equation is given by $\partial_t u_\varepsilon + \mathcal{L}_\varepsilon u_\varepsilon = f_\varepsilon$ in Ω_T . For this relaxed problem, we may assume $\nabla u_{\varepsilon} \in L^{\infty}_{loc}$ and $\nabla^2 u_{\varepsilon} \in L^2_{loc}$ (see Section [3.1](#page-15-1) for the details), and therefore u_{ε} satisfies

$$
\partial_t \partial_{x_j} u_\varepsilon - \text{div} \left(\nabla^2 E_\varepsilon (\nabla u_\varepsilon) \nabla \partial_{x_j} u_\varepsilon \right) = \partial_{x_j} f_\varepsilon \quad \text{in} \quad \Omega_T \tag{1.7}
$$

in the weak sense. The approximations of E and f are already found in elliptic problems [\[44,](#page-34-0) $\S2$], where a convergence result for an approximate solution is shown. Following the strategy in [\[44,](#page-34-0) §2.6], we verify that an approximate solution u_{ε} , which satisfies a suitable Dirichlet boundary condition, converges to a weak solution to [\(1.5\)](#page-1-6) strongly in $L^p(W^{1,p})$.

With the aid of approximation arguments, our regularity problem is reduced to local a priori Hölder continuity of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}) := \left(\sqrt{\varepsilon^2 + |\nabla u_{\varepsilon}|^2} - 2\delta\right)_{\perp} \frac{\nabla u_{\varepsilon}}{|\nabla u_{\varepsilon}|^2}$ approximation parameter ε is smaller than $\delta/8$. We should remark that the principal part $\mathcal{L}u$ itself is $\frac{\sqrt{u_{\varepsilon}}}{|\nabla u_{\varepsilon}|}$ for each fixed $\delta \in (0, 1)$, where the relaxed by $\mathcal{L}_{\varepsilon}u_{\varepsilon}$. In particular, for [\(1.7\)](#page-2-0), the parabolicity ratio of $\nabla^2 E_{\varepsilon}(\nabla u_{\varepsilon})$ should be measured by $V_{\varepsilon} := \sqrt{\varepsilon^2 + |\nabla u_{\varepsilon}|^2}$, rather than by $|\nabla u_{\varepsilon}|$. For this reason, we have to consider another truncation mapping $\mathcal{G}_{2\delta,\varepsilon}$, instead of $\mathcal{G}_{2\delta}$. In this paper, we mainly prove Hölder estimates of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$, independent of $\varepsilon \in (0, \delta/8)$. Then, the Arzelà–Ascoli theorem enables us to conclude that $\mathcal{G}_{2\delta}(\nabla u)$ is also α -Hölder continuous for each fixed $\delta \in (0, 1)$. We have to mention again that the exponent $\alpha \in (0, 1)$ substantially depends on δ . Indeed, our a priori estimates imply $\alpha \to 0$ as $\delta \to 0$ (see Section [2.3\)](#page-11-0).

Our proof of the Hölder continuity of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$ is based on the modifications of [\[44\]](#page-34-0), which deals the stationary problem of [\(1.1\)](#page-1-2). Here we mainly use standard methods in the parabolic regularity theory, including De Giorgi's truncation and comparisons with Dirichlet heat flows. Our parabolic computations concerning the Hölder continuity of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$ are inspired by [\[7,](#page-32-1) §3–7], where the Hölder gradient continuity of a vector-valued p-harmonic flow is discussed (see also [\[13\]](#page-32-2), [\[17\]](#page-32-3), [\[18\]](#page-32-4) as

fundamental works). Compared with [\[7\]](#page-32-1), our analysis is rather classical, since we do not use intrinsic scaling arguments at all (see Section [1.2\)](#page-3-0). Instead, we always avoid analysis near the degenerate regions of $V_{\varepsilon} = \sqrt{\varepsilon^2 + |\nabla u_{\varepsilon}|^2}$, so that we may regard [\(1.7\)](#page-2-0) as a uniformly parabolic equation in the classical sense, depending on δ . The local Hölder estimate of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$ is shown, as long as local L^{∞} -bounds of V_{ε} , uniformly for ε , are guaranteed. We give the proof of this a priori continuity estimate under the settings $p \in (1, \infty)$ and $V_{\varepsilon} \in L^{\infty}_{loc}$.

In showing the uniform L^{∞} -bounds of V_{ε} , however, the basic approach will differ, depending on whether p is greater than $p_c = \frac{2n}{n+1}$ $rac{2n}{n+2}$ or not. In this paper, where the supercritical case $p_c < p < \infty$ is considered, we prove local $L^{\infty} - L^p$ estimates of V_{ε} by Moser's iteration. For the critical or subcritical case $1 < p \leq p_c$, even the local L^{∞} -bound of a weak solution appears non-trivial, and some higher integrability assumption is required to prove the gradient continuity (see $[10]$). This remaining case is discussed in another paper [\[46\]](#page-34-3), under the assumption that a weak solution is in $L_{\text{loc}}^s(\Omega_T)$ with $s > n(2-p)/p \geq 2$. It should be remarked that the compact embedding $W_0^{1,p}$ $L^{1,p}(\Omega) \hookrightarrow L^{2}(\Omega)$ holds if and only if $p \in (p_c, \infty]$, and that we cannot apply the Aubin–Lions lemma for $p \in (1, p_c]$. In particular, when p is very close to 1, we will not be able to use any compact embedding concerning a parabolic function space. For this reason, the paper [\[46\]](#page-34-3) fails to deal with any external force term (see also Remark [2.5\)](#page-11-1).

1.2 Literature overview and comparisons with the paper

Section [1.2](#page-3-0) briefly introduces previous results on the elliptic or parabolic regularity for the p-Laplace or the $(1, p)$ -Laplace equation. We also compare this paper with [\[44\]](#page-34-0) and [\[45\]](#page-34-1), the author's recent works on the gradient continuity of the stationary $(1, p)$ -Laplace problem.

For the elliptic p-Laplace problems, $C^{1,\alpha}$ -regularity of a weak solution in the class $W^{1,p}(\Omega)$ is wellestablished in the full range $p \in (1, \infty)$ (see e.g., [\[1\]](#page-31-1), [\[12\]](#page-32-6), [\[23\]](#page-32-7), [\[34\]](#page-33-2), [\[37\]](#page-33-3), [\[42\]](#page-33-4), [\[47\]](#page-34-4)). For the parabolic p -Laplace problems, the Hölder gradient continuity was established by DiBenedetto–Friedman for $p \in (p_c, \infty)$ in [\[17\]](#page-32-3) and [\[18\]](#page-32-4) (see also [\[2\]](#page-31-2), [\[16\]](#page-32-8), [\[48\]](#page-34-5) for weaker results on the gradient continuity). In the remaining case $p \in (1, p_c]$, the same regularity was shown by Choe in [\[10\]](#page-32-5), where a solution $u \in$ $L^p(0, T; W^{1,p}(\Omega)) \cap C([0, T]; L^2(\Omega))$ is also assumed to satisfy $u \in L^s_{loc}(\Omega_T)$ with $s > n(2-p)/p \geq 2$. Without this L^s higher integrability assumption, no improved regularity result is generally expected (see [\[20,](#page-32-9) III. 7]).

In the proof of the Hölder gradient continuity for the parabolic p -Laplace equations, found in [\[7\]](#page-32-1) and [\[13,](#page-32-2) Chapter VIII], careful rescaling arguments for local gradient bounds play an important role. This method is often called an intrinsic scaling argument, which is a powerful tool for deducing variousregularity estimates for the parabolic p-Laplace equations ([\[13\]](#page-32-2), [\[19\]](#page-32-10)). In our analysis for (1.1) , however, we do not appeal to intrinsic scaling arguments, since our equation seems no longer uniformly parabolic on the facet of a solution, even if it is carefully rescaled. Instead, we make careful use of the truncation parameter δ , and try to avoid some delicate cases where a gradient may degenerate. In particular, we use no intrinsic parabolic cylinders, since our analysis works on standard parabolic cylinders, thanks to the careful truncation of a gradient.

For the stationary $(1, p)$ -Laplace problem $-\Delta_1 u - \Delta_p u = f$ with $f = f(x)$ in L^q for some $q \in (n, \infty]$ the continuous differentiability of a solution $u = u(x)$ was first shown in [\[29\]](#page-33-5), where a solution is assumed to be both scalar-valued and convex. Although the proof therein is rather easily carried out thanks to some basic tools from convex analysis, it is heavily based on a maximum principle and the convexity of a solution. Recently, the author removed these technical assumptions in [\[44\]](#page-34-0), [\[45\]](#page-34-1), where it is proved that a truncated gradient $\mathcal{G}_{\delta}(\nabla u)$, defined in Section [1.1,](#page-1-1) is locally Hölder continuous. The author's recent research([\[44\]](#page-34-0), [\[45\]](#page-34-1)) is highly inspired by a paper [\[5\]](#page-31-3), which shows the gradient continuity for a very degenerate elliptic problem of the form $-\text{ div}\left(|\mathcal{G}_1(\nabla v)|^{p'-2}\mathcal{G}_1(\nabla v)\right)=f.$ Here the real-valued functions $v = v(x)$ and $f = f(x)$ are respectively unknown and given. For this degenerate problem, the three papers [\[5\]](#page-31-3), [\[11\]](#page-32-11), [\[39\]](#page-33-6) prove that a truncated gradient $\mathcal{G}_1(\nabla v)$, similarly defined in Section [1.1,](#page-1-1) is continuous, provided $f \in L^q$ with $q \in (n, \infty]$. Among them, [\[5\]](#page-31-3) is based on classical methods, including De Giorgi's truncation and a perturbation argument of Campanato-type, and this works even for the vector-valued case. Following the spirit of [\[5\]](#page-31-3), the author proved the

gradient continuity for $(1, p)$ -Laplace-type problems both in the scalar-valued [\[44\]](#page-34-0) and vector-valued [\[45\]](#page-34-1) cases. The continuity of $\mathcal{G}_0(\nabla u) = \nabla u$ and $\mathcal{G}_1(\nabla v)$ is proved in a qualitative way, by showing quantitative continuity estimates of the truncated gradients $\mathcal{G}_{\delta}(\nabla u)$ and $\mathcal{G}_{1+\delta}(\nabla v)$, whose estimates depend on the truncation parameter $\delta \in (0, 1)$. These continuity results are known only for elliptic cases, although some other regularity results are shown for degenerate parabolic problems in recent papers [\[3\]](#page-31-4), [\[25\]](#page-32-12). In this paper, by adjusting the arguments in [\[44\]](#page-34-0) to the parabolic problem, we aim to establish a gradient continuity result for the singular parabolic equation [\(1.1\)](#page-1-2).

We conclude Section [1.2](#page-3-0) by noting a crucial difference between this paper and the author's previous works [\[44\]](#page-34-0), [\[45\]](#page-34-1), especially on the converence of approximate solutions. To prove the strong L^p -convergence of ∇u_{ε} , we have to make use of a weak compactness argument and a compact embedding for parabolic function spaces. There, the technical assumption [\(1.3\)](#page-1-4) is required in showing the boundedness of the time derivatives in $L^{p'}(W^{-1,p'})$. This is mainly because the parabolic equation (1.1) is treated in the classical $L^{p'}(W^{-1,p'})$ -framework, which is typically used to construct a weak solution to a parabolic p-Laplace equation (see [\[36\]](#page-33-7), [\[40\]](#page-33-8)). For this technical reason, in addition to (1.2) , the exponents p and q have to satisfy another good condition such as (1.4) . Also, the condition $p > p_c$ is carefully used to apply the Aubin–Lions lemma, which plays a crucial role in treating the external force terms. Hence, our parabolic approximation arguments need some restrictions on the exponents p and q, while [\[44\]](#page-34-0) and [\[45\]](#page-34-1) require no restrictive assumptions. This kind of restriction strongly appears when $p \leq p_c$, since the compact embedding $W_0^{1,p}$ $L^{1,p}(\Omega) \hookrightarrow L^2(\Omega)$ no longer holds true. In particular, the Aubin–Lions lemma cannot be applied, which prevents us from dealing with any external force term (see also Remark [2.5\)](#page-11-1). Although our approximation argument partly works even for $p \in (1, p_c]$ (see [\[46,](#page-34-3) Proposition 3.3] for the details), no external force term can be treated.

1.3 Mathematical models

The second-order parabolic equation [\(1.1\)](#page-1-2) appears when one models the motion of a Bingham fluid [\[21,](#page-32-0) Chapter VI], a non-Newtonian fluid that has both plasticity and viscosity properties. Its motion is governed by $\partial_t U + (U \cdot \nabla)U - b_1 \Delta_1 U - b_2 \Delta_2 U + \nabla \pi = 0$ and div $U = 0$ if there is no external force. Here $U = U(x_1, x_2, x_3, t)$ denotes the three-dimensional velocity of a Bingham fluid, and a scalar function $\pi = \pi(x_1, x_2, x_3, t)$ denotes the pressure. The positive constants b_1 and b_2 respectively reflect a Bingham fluid's plasticity and viscosity effects. When $b_1 = 0$, this model becomes the incompressible Navier–Stokes equation. We consider a uni-directional laminar flow in a cylinder $\Omega \times \mathbb{R} \subset \mathbb{R}^3$ with velocity $U = (0, 0, u(x_1, x_2, t))$. Since the equalities div $U = 0$ and $(U \cdot \nabla)U = 0$ are automatically fulfilled, the equation is reduced to $\partial_t u - b_1 \Delta_1 u - b_2 \Delta_2 u = -\partial_{x_3} \pi$ and $\partial_{x_1} \pi = \partial_{x_2} \pi = 0$ by carrying out similar arguments in [\[21,](#page-32-0) Chapter VI, §1.3]. Moreover, the right-hand side $\partial_{x_3}\pi \equiv f$ depends at most on t, since the left-hand side is independent of x_3 . From our main Theorem [1.2,](#page-6-1) given in Section [1.4,](#page-4-1) we conclude that a spatial derivative ∇u is continuous, provided $f = f(t)$ admits L^q-integrability with $q > 4$.

It will be worth mentioning that another important model can be found in the evaporation dynamics of crystal surfaces, modeled by Spohn in a paper [\[41,](#page-33-9) §2]. There the evolution of the height function of the crystal, denoted by h, is modeled as $\partial_t h = -\mu \cdot (-\Delta_1 h - \Delta_3 h)$ with $\mu = \mu(\nabla h) > 0$ standing for the mobility. When μ is constant, we can conclude the continuity of ∇h by our main Theorem [1.2.](#page-6-1) However, Spohn's discussion concerning mobility tells us that μ has to be of the form $\mu = \kappa |\nabla h|$ for some constant $\kappa > 0$. In other words, in his model, the speed of each level set is assumed to move by $\Delta_1 h + \Delta_3 h$, and another equation appears.

1.4 Notations, main results and outline of the paper

Before stating our main theorem, we fix some notations.

The set of all non-negative integers and the set of all natural numbers are respectively denoted by $\mathbb{Z}_{\geq 0} := \{0, 1, 2, \ldots\}$, and $\mathbb{N} := \mathbb{Z}_{\geq 0}\setminus\{0\}$. For a given real number $a \in \mathbb{R}$, we write $a_+ := \max\{a, 0\}$. For given vectors $x = (x_1, \ldots, x_n), y = (y_1, \ldots, y_n) \in \mathbb{R}^n$, the canonical inner product and the Euclidean norm are respectively written by $\langle x | y \rangle := x_1 y_1 + \cdots + x_n y_n \in \mathbb{R}$ and $|x| := \sqrt{\langle x | x \rangle} \in [0, \infty)$. For an $n \times n$ real matrix $A = (A_{j,k})_{j,k}$, we define the standard norms as $||A|| := \sup_{|x|=1} |Ax|$, and

 $|A| := \sqrt{\sum_{j,k=1}^n A_{j,k}^2}$, called the operator norm and the Hilbert–Schmidt norm respectively. For $n \times n$ symmetric real matrices A and B, we write $A \leq B$ when $B - A$ is positive semi-definitive. The symbols id and O stand for the identity matrix and the zero matrix respectively.

We introduce a metric d_p in \mathbb{R}^{n+1} by $d_p((x, t), (y, s)) := \max\left\{|x - y|, \sqrt{|t - s|}\right\}$ for $(x, t), (y, s) \in$ $\mathbb{R}^n \times \mathbb{R}$. With respect to this metric, $dist_p(\mathcal{Q}, \partial_p \Omega_T)$ denotes the distance from the given set $\mathcal{Q} \in \Omega_T$ to the parabolic boundary $\partial_p \Omega_T := (\Omega \times \{0\}) \cup (\partial \Omega \times [0, T])$. For given $x_0 \in \mathbb{R}^n$, $t_0 \in \mathbb{R}$ and $r \in (0, \infty)$, we set an open ball $B_r(x_0) := \{x \in \mathbb{R}^n \mid |x - x_0| < r\}$, a half interval $I_r(t_0) := (t_0 - r^2, t_0]$, and a parabolic cylinder $Q_r(x_0, t_0) := B_r(x_0) \times I_r(t_0)$. The closure and the interior of the half interval $I_r(t_0)$ are respectively denoted by $\overline{I_r} := [t_0 - r^2, t_0]$ and $\overline{I_r} := (t_0 - r^2, t_0)$. The center points x_0 and t_0 are often omitted when they are clear.

For $p \in (1, \infty)$, the Hölder conjugate exponent is denoted by $p' := p/(p-1) \in (1, \infty)$. For k, $m \in \mathbb{N}$, we write $W^{k,p}(\Omega; \mathbb{R}^m)$, and $L^p(\Omega; \mathbb{R}^m)$ respectively by the Sobolev space of k-th order, and the Lebesgue space, equipped with the standard norms. When $m = 1$, we often omit the codomain \mathbb{R}^m , and write $L^p(\Omega)$ and $W^{k,p}(\Omega)$. For $p \in (1, \infty)$, the closed subspace $W_0^{1,p}$ $v_0^{1,p}(\Omega) \subset W^{1,p}(\Omega)$ is defined to be the closure of infinitely differentiable functions compactly supported in Ω , equipped with another equivalent norm $\|\nabla u\|_{L^p(\Omega)} := (\int_{\Omega} |\nabla u|^p \,dx)^{1/p}$ for $u \in W_0^{1,p}$ $\int_0^{1,p}(\Omega)$. With respect to this norm, the dual of $W_0^{1,p}$ $V_0^{1,p}(\Omega)$ is denoted by $W^{-1,p'}(\Omega)$. For $F \in W^{-1,s'}(\Omega)$ and $v \in W_0^{1,s}$ $\mathcal{O}_0^{1,s}(\Omega)$ with $s \in (p_c, \infty)$, the dual pairing is often denoted by $\langle F, v \rangle_{W^{-1, s'}(\Omega), W_0^{1, s}(\Omega)} := F(v)$, or more simply by $\langle F, v \rangle$ when s and Ω are clear. We write $C^0([0, T]; L^2(\Omega))$ for the set of all $L^2(\Omega)$ -valued functions that are strongly continuous in [0, T]. For $k \in \mathbb{Z}_{\geqslant 0}$, $l, m \in \mathbb{N}$, and $U \subset \mathbb{R}^l$, the symbol $C^k(U; \mathbb{R}^m)$ denotes the set of all \mathbb{R}^m -valued functions of the C^k -class in U, and we abbreviate $C^k(U) := C^k(U; \mathbb{R})$.

To simplify notation, we often make use of the abbreviations $X = (x, t)$ and $dX = dxdt$. For a kdimensional Lebesgue measurable set $U \subset \mathbb{R}^k$ with $k \in \mathbb{N}$, the symbol $|U|$ stands for the k-dimensional Lebesgue measure of U. For \mathbb{R}^k -valued functions $g = g(x)$, and $h = h(x, t)$ that are respectively integrable in $U \subset \mathbb{R}^n$, and $U \times I \subset \mathbb{R}^{n+1}$ with $0 < |U|, |I| < \infty$, the average integrals are denoted by $f_U g(x) dx := |U|^{-1} \int_U g(x) dx \in \mathbb{R}^k$, $\oiint_{U \times I} h(x, t) dX := |U|^{-1} |I|^{-1} \iint_U h(x, t) dX \in \mathbb{R}^k$. The former average integral is often denoted by f_U . The average integral of h over a parabolic cylinder Q_r is often denoted by $(h)_{Q_r}$ or $(h)_r$.

The function spaces

$$
X^{p}(0, T; \Omega) := \left\{ u \in L^{p}(0, T; W^{1, p}(\Omega)) \mid \partial_t u \in L^{p'}(0, T; W^{-1, p'}(\Omega)) \right\},\,
$$

and

$$
X_0^p(0, T; \Omega) := \left\{ u \in L^p(0, T; W_0^{1, p}(\Omega)) \middle| \partial_t u \in L^{p'}(0, T; W^{-1, p'}(\Omega)) \right\}
$$

are considered for $p \in (\frac{2n}{n+2}, \infty)$. By the embeddings $W_0^{1,p}$ $L^{1,p}(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow W^{-1,p'}(\Omega)$, we can use some inclusions or embeddings concerning X_0^p $_{0}^{p}(0, T; \Omega)$. More precisely, by the Lions–Magenes lemma [\[40,](#page-33-8) Chapter III, Proposition 1.2], the inclusion X_0^p $L_0^p(0, T; \Omega) \subset C^0([0, T]; L^2(\Omega))$ holds. Also by the Aubin–Lions lemma [\[40,](#page-33-8) Chapter III, Proposition 1.3], the compact embedding X_0^p $_{0}^{p}(0, T; \Omega) \hookrightarrow \hookrightarrow$ $L^p(0, T; L^2(\Omega))$ can be applied.

In this paper, we require $E_1 \in C^0(\mathbb{R}^n) \cap C^2(\mathbb{R}^n \setminus \{0\})$, and $E_p \in C^1(\mathbb{R}^n) \cap C^2(\mathbb{R}^n \setminus \{0\})$ for the smoothness of the density functions. We let E_1 be positively one-homogeneous. In other words, E_1 is assumed to satisfy

$$
E_1(kz) = kE_1(z) \quad \text{for all } k \in (0, \infty), z \in \mathbb{R}^n. \tag{1.8}
$$

We also require the existence of a concave, non-decreasing modulus of continuity $\omega_1: [0, \infty) \to [0, \infty)$ such that $\omega_1(0)=0$ and

$$
\left\|\nabla^2 E_1(z_1) - \nabla^2 E_1(z_2)\right\| \leq \omega_1(|z_1 - z_2|)
$$
\n(1.9)

for all $z_1, z_2 \in \mathbb{R}^n$ with $1/32 \leq |z_j| \leq 3$ for $j \in \{1, 2\}$. The other density E_p admits constants $0 < \lambda_0 < \Lambda_0 < \infty$ such that

$$
|\nabla E_p(z)| \le \Lambda_0 |z|^{p-1} \quad \text{for all } z \in \mathbb{R}^n,
$$
\n(1.10)

$$
\lambda_0 |z|^{p-2} \text{id}_n \leqslant \nabla^2 E_p(z) \leqslant \Lambda_0 |z|^{p-2} \text{id}_n \quad \text{for all } z \in \mathbb{R}^n \setminus \{0\}. \tag{1.11}
$$

Moreover, we assume that there exists a concave, non-decreasing, and continuous function ω_p : $[0, \infty) \rightarrow$ $[0, \infty)$, such that $\omega_p(0) = 0$ and

$$
\left\|\nabla^2 E_p(z_1) - \nabla^2 E_p(z_2)\right\| \leq C_{\delta, M} \omega_p(|z_1 - z_2|/\mu)
$$
\n(1.12)

for all $z_1, z_2 \in \mathbb{R}^n$ with $\mu/32 \leq |z_j| \leq 3\mu$ for $j \in \{1, 2\}$, and $\mu \in (\delta, M - \delta)$. Here δ and M are fixed constants with $0 < 2\delta < M < \infty$, and the constant $C_{\delta,M} \in (0, \infty)$ depends on δ and M.

The definition of a weak solution to [\(1.5\)](#page-1-6) is given as follows.

Definition 1.1. Let the exponents p and q satisfy (1.2) – (1.3) , and $f \in L^q(\Omega_T)$. A function $u \in$ $X^p(0, T; \Omega) \cap C^0([0, T]; L^2(\Omega))$ is called a weak solution to [\(1.5\)](#page-1-6) if there exists a vector field $Z \in$ $L^{\infty}(\Omega_T)$ such that

$$
\int_0^T \langle \partial_t u, \varphi \rangle dt + \iint_{\Omega_T} \langle Z + \nabla E_p(\nabla u) | \nabla \varphi \rangle dX = \iint_{\Omega_T} f \varphi dX \tag{1.13}
$$

for all $\varphi \in X_0^p$ $_0^p(0, T; \Omega)$, and

$$
Z(x, t) \in \partial E_1(\nabla u(x, t)) \quad \text{for a.e. } (x, t) \in \Omega_T.
$$
 (1.14)

Our main result is the following Theorem [1.2.](#page-6-1)

Theorem 1.2. Let p, q, E_1 and E_p satisfy (1.2) – (1.3) and (1.8) – (1.12) . Assume that u is a weak solution to [\(1.5\)](#page-1-6) with $f \in L^q(\Omega_T)$. Then, ∇u is continuous in Ω_T .

This paper is organized as follows. Section [2](#page-6-0) aims to give an approximation scheme for the very singular parabolic equation [\(1.5\)](#page-1-6). The basic strategy is to relax an energy density by convoluting this and the Friedrichs standard mollifier. In Section [2,](#page-6-0) we verify that a weak solution to [\(1.5\)](#page-1-6) can be constructed as a limit of the approximate parabolic equation (Proposition [2.4\)](#page-9-0), where [\(1.4\)](#page-1-5) is required. By basic estimates for approximate solutions, including local gradient bounds (Theorem [2.7\)](#page-12-0), De Giorgi-type oscillation estimates (Proposition [2.9\)](#page-13-0), and Campanato-type growth estimates (Proposition [2.10\)](#page-13-1), we prove a priori Hölder bounds for truncated gradients of approximate solutions (Theorem [2.8\)](#page-12-1). From Proposition [2.4](#page-9-0) and Theorem [2.8,](#page-12-1) we complete the proof of Theorem [1.2.](#page-6-1) Sections [3–](#page-15-0)[6](#page-24-0) provides the proofs of Theorem [2.7,](#page-12-0) and Propositions [2.9](#page-13-0)[–2.10.](#page-13-1) In Section [3,](#page-15-0) we deduce some basic energy estimates from weak formulations. Theorem [2.7](#page-12-0) is shown in Section [4.](#page-18-0) Sections [5–](#page-21-0)[6](#page-24-0) provide various oscillation lemmata. There, depending on the size of a superlevel set, we make different analyses for degenerate or non-degenerate cases. Section [5,](#page-21-0) which deals with the degenerate case, is focused on proving Proposition [2.9.](#page-13-0) Section [6,](#page-24-0) where the non-degenerate case is discussed, we appeal to comparisons with Dirichlet heat flows to show Proposition [2.10.](#page-13-1)

Finally, we would like to mention again that the remaining case $p \in (1, p_c]$ is treated in another paper [\[46\]](#page-34-3) by the author. There, the local boundedness of a solution or its gradient is mainly discussed, and Theorem [2.8](#page-12-1) in this paper, which result is valid for any $p \in (1, \infty)$ as long as uniform gradient bounds are guaranteed, is used without proof.

2 Approximation problems

In Section [2,](#page-6-0) we consider a parabolic approximate problem, based on the convolution of $E = E_1 + E_p$ and the Friedrichs mollifiers. More precisely, we choose and fix a spherically symmetric and nonnegative function $j \in C_c^{\infty}(\mathbb{R}^n)$ such that $||j||_{L^1(\mathbb{R}^n)} = 1$ holds, and its support is the closed ball $\overline{B_1(0)}$. For this smooth function j, we define $j_{\varepsilon}(x) := \varepsilon^{-n} j(x/\varepsilon)$ for $\varepsilon \in (0, 1)$, $x \in \mathbb{R}^n$, often called the Friedrichs mollifier. For $\varepsilon \in (0, 1)$, we define

$$
E_{s,\varepsilon}(z) := (j_{\varepsilon} * E_s)(z) = \int_{\mathbb{R}^n} j_{\varepsilon}(z - y) E_s(y) \, dy \tag{2.1}
$$

for $z \in \mathbb{R}^n$, $s \in \{1, p\}$. Section [2](#page-6-0) aims to show convergence results for weak solutions to approximate problems, based on our straightforward relaxation of $E = E_1 + E_p$ by $E_{\varepsilon} := E_{1,\,\varepsilon} + E_{p,\,\varepsilon}$. Also, we would like to prove our main Theorem [1.2](#page-6-1) by making use of a priori estimates for approximate solutions, which are shown in Sections [3](#page-15-0)[–6.](#page-24-0)

2.1 Relaxation of divergence operators and some basic estimates

In Section [2.1,](#page-7-0) we briefly note some basic estimates of E_1 , $E_{\varepsilon} = E_{1,\varepsilon} + E_{p,\varepsilon}$, and $\mathcal{G}_{2\delta,\varepsilon}$.

By [\(1.8\)](#page-5-0), for any $k \in (0, \infty)$, $z \in \mathbb{R}^n \setminus \{0\}$, E_1 satisfies $\nabla E_1(kz) = \nabla E_1(z)$ and $\nabla^2 E_1(kz) =$ $k^{-1}\nabla^2 E_1(z)$ In particular, by $E_1 \in C^2(\mathbb{R}^n \setminus \{0\})$ and [\(2.3\)](#page-7-1), it is easy to find a constant $K_0 \in (0, \infty)$ such that

$$
|\nabla E_1(z)| \leq K_0 \quad \text{and} \quad O \leq \nabla^2 E_1(z) \leq K_0 |z|^{-1} \text{id}_n \quad \text{for all } z \in \mathbb{R}^n \setminus \{0\},\tag{2.2}
$$

$$
\left\|\nabla^2 E_1(z_1) - \nabla^2 E_1(z_2)\right\| \le \mu^{-1} \omega_1(|z_1 - z_2|/\mu)
$$
\n(2.3)

for all $\mu \in (0, \infty)$, $z_1, z_2 \in \mathbb{R}^n \setminus \{0\}$ with $\mu/32 \leqslant |z_j| \leqslant 3\mu$ for $j \in \{1, 2\}$.

By [\(1.10\)](#page-5-1)–[\(1.11\)](#page-5-2) and [\(2.2\)](#page-7-2), the relaxed density functions $E_{1,\varepsilon}$ and $E_{p,\varepsilon}$, given by [\(2.1\)](#page-6-3), satisfies the following estimates for every $\varepsilon \in (0, 1), z \in \mathbb{R}^n$;

$$
|\nabla E_{1,\varepsilon}(z)| \leq K, \quad O \leq \nabla^2 E_{1,\varepsilon}(z) \leq K \left(\varepsilon^2 + |z|^2\right)^{-\frac{1}{2}} \text{id},\tag{2.4}
$$

$$
|\nabla E_{p,\varepsilon}(z)| \leq \Lambda \left(\varepsilon^2 + |z|^2\right)^{\frac{p-1}{2}}, \quad \lambda \left(\varepsilon^2 + |z|^2\right)^{\frac{p}{2}-1} \operatorname{id} \leq \nabla^2 E_{p,\varepsilon}(z) \leq \Lambda \left(\varepsilon^2 + |z|^2\right)^{\frac{p}{2}-1} \operatorname{id}. \tag{2.5}
$$

Here the positive constants $\lambda = \lambda(\lambda_0, n, p), \Lambda = \Lambda(\Lambda_0, n, p), K = K(K_0, n)$ are independent of $\varepsilon \in (0, 1)$. Moreover, by straightforward computations, we have

$$
\langle \nabla E_{\varepsilon}(z_1) - \nabla E_{\varepsilon}(z_2) | z_1 - z_2 \rangle \geq \begin{cases} c(p)\lambda |z_1 - z_2|^p & (p \geq 2), \\ \lambda(\varepsilon^2 + |z_1|^2 + |z_2|^2)^{p/2-1}|z_1 - z_2|^2 & \text{(otherwise)}, \end{cases}
$$
(2.6)

for all $z_1, z_2 \in \mathbb{R}^n$. Also, it is easy to verify that E_p satisfies

$$
\langle \nabla E_p(z_1) - \nabla E_p(z_2) | z_1 - z_2 \rangle > 0 \quad \text{for any} \quad z_1, z_2 \in \mathbb{R}^n \quad \text{with} \quad z_1 \neq z_2. \tag{2.7}
$$

For the proofs of (2.4) – (2.6) , see [\[43,](#page-33-10) §A] and [\[44,](#page-34-0) §2] (see also [\[22\]](#page-32-13)).

Throughout Sections [2–](#page-6-0)[6,](#page-24-0) we deal with truncation mappings \mathcal{G}_{δ} , $\mathcal{G}_{\delta,\varepsilon} : \mathbb{R}^n \to \mathbb{R}^n$, defined as $\mathcal{G}_{\delta}(z) :=$ $(|z|-\delta)_{+\frac{z}{|z}}$ $\frac{z}{|z|}$, and $\mathcal{G}_{\delta,\,\varepsilon}(z) := \left(\sqrt{\varepsilon^2 + |z|^2} - \delta\right)$ $^{+}$ z $\frac{z}{|z|}$ for $z \in \mathbb{R}^n$. The truncation mapping $\mathcal{G}_{\delta, \varepsilon}$ makes sense as long as $0 < \varepsilon < \delta$ holds, and this is Lipschitz continuous, provided ε is much smaller than δ . More precisely, if $0 < \varepsilon < \delta/8$ holds, then by [\[44,](#page-34-0) Lemma 2.4], we have

$$
|\mathcal{G}_{2\delta,\,\varepsilon}(z_1) - \mathcal{G}_{2\delta,\,\varepsilon}(z_2)| \leq c_\dagger |z_1 - z_2| \quad \text{for all } z_1, \, z_2 \in \mathbb{R}^n \tag{2.8}
$$

with $c_{\uparrow} := 1 + 64/\sqrt{255}$. We also use some basic inequalities given in Lemma [2.1.](#page-7-5)

Lemma 2.1. Let $E_{\varepsilon} = E_{1,\varepsilon} + E_{p,\varepsilon}$ be given by [\(2.1\)](#page-6-3) for $\varepsilon \in (0, 1)$. Assume that non-negative constants δ, ε and M satisfy $\varepsilon < \delta/8$ and $2\delta < M$. Then, there exist constants $0 < C_1 < C_2 < \infty$, depending at most on n, p, λ , Λ , K, δ and M, such that the following estimates [\(2.9\)](#page-7-6)–[\(2.10\)](#page-7-7) hold for all z_1 , $z_2 \in \mathbb{R}^n$ satisfying $\delta \leqslant |z_1| \leqslant M$ and $|z_2| \leqslant M$.

$$
\langle \nabla E_{\varepsilon}(z_1) - \nabla E_{\varepsilon}(z_2) | z_1 - z_2 \rangle \geqslant C_1 |z_1 - z_2|^2. \tag{2.9}
$$

$$
|\nabla E_{\varepsilon}(z_1) - \nabla E_{\varepsilon}(z_2)| \leqslant C_2 |z_1 - z_2|.
$$
\n(2.10)

Moreover, there holds

$$
\left|\nabla^2 E_{\varepsilon}(z_1)(z_1 - z_2) - \left(\nabla E_{\varepsilon}(z_1) - \nabla E_{\varepsilon}(z_2)\right)\right| \leq |z_1 - z_2| \omega(|z_1 - z_2|/\mu)
$$
\n(2.11)

for all $z_1, z_2 \in \mathbb{R}^n$ satisfying $\delta + \frac{\mu}{4} \leqslant |z_1| \leqslant \delta + \mu$, $|z_2| \leqslant \delta + \mu$ with $\delta < \mu < M - \delta$. Here the modulus of continuity $\omega = \omega_{\delta,M}$, which is concave, is of the form $\omega(\sigma) = C_3(\omega_1(\sigma) + \omega_p(\sigma) + \sigma)$ with $C_3 \in (0, \infty)$ depending at most on p, Λ , K , δ and M .

In the special case where $\omega_1 = \omega_1(\sigma)$ and $\omega_p = \omega_p(\sigma)$ behave like σ^{β_0} for some $\beta_0 \in (0, 1)$, a similar continuity estimate to (2.11) is shown in [\[44,](#page-34-0) Lemma 2.6]. The proofs of (2.9) – (2.10) are already given there. Modifying [\[44,](#page-34-0) Lemma 2.6], we would like to give a proof of [\(2.11\)](#page-7-8).

Proof. By $\varepsilon < \delta/8$, we can check that $\frac{\mu^2}{16} \leq \varepsilon^2 + |z_1|^2 \leq 5\mu^2$, and $\varepsilon^2 + |z_2|^2 \leq 5\mu^2$. Combining them with $(2.4)-(2.5)$ $(2.4)-(2.5)$ and $\delta < \mu < M - \delta$, we can easily check that the left-hand side of (2.11) is bounded by some $C = C(p, \Lambda, K, \delta, M) \in (1, \infty)$, from which [\(2.11\)](#page-7-8) is easily concluded when $|z_1 - z_2| > \mu/32$. For the remaining case $|z_1 - z_2| \le \mu/32$, we use the triangle inequality to get $\frac{7}{8}\delta + \frac{\mu}{4} \le |z_1 - y| \le \frac{17}{8}\mu$, For the remaining case $|z_1 - z_2| \le \mu/52$, we use the triangle mequality to get $\frac{1}{8} \sigma + \frac{1}{4} \le |z_1 - y| \le \frac{1}{8} \mu$,
and $\frac{7}{8} \delta + \frac{\mu}{32} \le |z_1 - y + t(z_1 - z_2)| \le 3\mu$ for all $y \in B_{\varepsilon}(0)$. By $\nabla^2 E \in C(\mathbb{R}^n \setminus \{0$ $\nabla^2 E_{\varepsilon} = j_{\varepsilon} * \nabla^2 E$ in $\mathbb{R}^n \setminus \overline{B_{\varepsilon}(0)}$ for any fixed $\hat{\varepsilon} \in (\varepsilon, 1)$. Using this identity, [\(1.12\)](#page-6-2), and [\(2.3\)](#page-7-1) yields

$$
\begin{split}\n\left|\nabla^2 E_{\varepsilon}(z_1)(z_1 - z_2) - \left(\nabla E_{\varepsilon}(z_1) - \nabla E_{\varepsilon}(z_2)\right)\right| \\
&= \left| \left(\int_0^1 \left[\nabla^2 E_{\varepsilon}(z_1) - \nabla^2 E_{\varepsilon}(z_1 + t(z_2 - z_1))\right] dt \right) \cdot (z_1 - z_2) \right| \\
&\leq |z_1 - z_2| \sum_{s=1, p} \int_0^1 \int_{B_{\varepsilon}(0)} \left\|\nabla^2 E_s(z_1 - y - t(z_1 - z_2)) - \nabla^2 E_s(z_1 - y)\right\| j_{\varepsilon}(y) dy dt \\
&\leq |z_1 - z_2| \left[\delta^{-1} \omega_1(|z_1 - z_2|/\mu) + C_{\delta, M} \omega_p(|z_1 - z_2|/\mu)\right],\n\end{split}
$$

from which we obtain [\(2.11\)](#page-7-8).

2.2 Convergence of approximate solutions

For an approximation parameter $\varepsilon \in (0, 1)$, we consider the following Dirichlet boundary value problem;

$$
\begin{cases}\n\partial_t u - \operatorname{div} (\nabla E_{\varepsilon} (\nabla u_{\varepsilon})) = f_{\varepsilon} & \text{in} \quad \Omega_T, \\
u_{\varepsilon} = u_{\star} & \text{on} \quad \partial_p \Omega_T.\n\end{cases}
$$
\n(2.12)

Here $u_{\star} \in X^p(0, T; \Omega) \cap C^0([0, T]; L^2(\Omega))$ is a given function. For the approximation of $f \in L^q(\Omega_T)$, we only require the weak^{*} convergence

$$
f_{\varepsilon} \stackrel{*}{\rightharpoonup} f \quad \text{in} \quad L^q(\Omega_T). \tag{2.13}
$$

In particular, we may let $f_{\varepsilon} \in C^{\infty}(\Omega_T)$, and choose a constant $F \in (0, \infty)$ such that

$$
||f_{\varepsilon}||_{L^{q}(\Omega_{T})} \le F \quad \text{for all} \quad \varepsilon \in (0, 1). \tag{2.14}
$$

The weak solution of [\(2.12\)](#page-8-1) is defined to be a function $u_{\varepsilon} \in u_{\star} + X_0^p$ $v_0^p(0, T; \Omega)$ satisfying $u_{\varepsilon}|_{t=0} = u_{\star}|_{t=0}$ in $L^2(\Omega)$, and

$$
\partial_t u_{\varepsilon} - \text{div} \left(\nabla E_{\varepsilon} (\nabla u_{\varepsilon}) \right) = f_{\varepsilon} \quad \text{in} \quad L^{p'}(0, T; L^{p'}(\Omega)). \tag{2.15}
$$

In other words, the function u_{ε} satisfies

$$
\int_0^T \langle \partial_t u_\varepsilon, \, \varphi \rangle \, \mathrm{d}t + \iint_{\Omega_T} \langle \nabla E_\varepsilon(\nabla u_\varepsilon) \mid \nabla \varphi \rangle \, \mathrm{d}X = \iint_{\Omega_T} f_\varepsilon \varphi \, \mathrm{d}X. \tag{2.16}
$$

for all $\varphi \in L^p(0, T; W_0^{1,p})$ $O_0^{(1,p)}(\Omega)$. The unique existence of the weak solution of (2.12) is well-established in found in [\[36,](#page-33-7) Chapitre 2, §1], [\[40,](#page-33-8) §III.4], we can show the unique existence of [\(2.12\)](#page-8-1). There the continuous inclusions $W_0^{1,p}$ $L^{1,p}(\Omega) \hookrightarrow L^2(\Omega) \hookrightarrow W^{-1,p'}(\Omega)$ is used to construct a weak solution by the Faedo–Galerkin method. It appears that the monotonicity methods in [\[36,](#page-33-7) Chapitre 2, §1], [\[40,](#page-33-8) §III.4] cannot be directly applied to (1.5) , since the density E_1 lacks C^1 -regularity at the origin. In particular, the existence of a subgradient vector field Z itself appears non-trivial. To overcome this difficulty, we invoke a convergence lemma on vector fields (Lemma [2.2\)](#page-8-2). This lemma is a special case of [\[44,](#page-34-0) Lemma 2.8], which helps to prove the convergence of weak solutions.

Lemma 2.2. Let $E_1 \in C^0(\mathbb{R}^n) \cap C^1(\mathbb{R}^n \setminus \{0\}), E_p \in C^1(\mathbb{R}^n)$ satisfy $(1.8)-(1.10)$ $(1.8)-(1.10)$. For each $\varepsilon \in (0, 1)$, define $E_{\varepsilon} = E_{1,\varepsilon} + E_{p,\varepsilon}$ by [\(2.1\)](#page-6-3). Then, the following hold.

(1). For each fixed $v \in L^p(\Omega_T)$, we have $\nabla E_{\varepsilon}(v) \to A_0(v)$ in $L^{p'}(\Omega_T; \mathbb{R}^n)$ as $\varepsilon \to 0$. Here $A_0: \mathbb{R}^n \to$ \mathbb{R}^n is defined as $A_0(z) := \nabla E(z)$ for $z \neq 0$, and $A_0(0) := (j * \nabla E_1)(0)$.

 \Box

(2). Assume that a sequence $\{v_{\varepsilon_k}\}_k \subset L^p(\Omega_T; \mathbb{R}^n)$, where $\varepsilon_k \to 0$ as $k \to 0$, satisfies $v_{\varepsilon_k} \to$ v_0 in $L^p(\Omega_T; \mathbb{R}^n)$ as $k \to \infty$ for some $v_0 \in L^p(\Omega_T; \mathbb{R}^n)$. Then, up to a subsequence,

 $\nabla E_{p,\varepsilon_k}(v_{\varepsilon_k}) \to \nabla E_p(v_0)$ in $L^{p'}(\Omega_T; \mathbb{R}^n)$ and $\nabla E_{1,\varepsilon_k}(v_{\varepsilon_k}) \stackrel{*}{\to} Z$ in $L^{\infty}(\Omega_T; \mathbb{R}^n)$

hold as $k \to \infty$, and the weak^{*} limit $Z \in L^{\infty}(\Omega_T; \mathbb{R}^n)$ satisfies $Z(x, t) \in \partial E_1(v_0(x, t))$ for a.e. $(x, t) \in \Omega_T$.

Applying Lemma [2.2,](#page-8-2) we would like to prove that u_{ε} converges to a weak solution of

$$
\begin{cases}\n\partial_t u + \mathcal{L}u = f \quad \text{in} \quad \Omega_{T-\tau}, \\
u = u_\star \quad \text{on} \quad \partial_p \Omega_{T-\tau},\n\end{cases}
$$
\n(2.17)

for each fixed $\tau \in (0, T)$, in the sense of Definition [2.3](#page-9-1) below.

Definition 2.3. Let p, q satisfy (1.2) – (1.3) . Fix $\tau \in (0, T)$ and $u_{\star} \in X^p(0, T - \tau; \Omega) \cap C^0([0, T - \tau])$ τ]; $L^2(\Omega)$). A function $u \in u_\star + X_0^p$ $p_0^p(0, T - \tau; \Omega)$ is called a weak solution of (2.17) if the identity $u|_{t=0} = u_\star|_{t=0}$ holds in $L^2(\Omega)$, and there exists $Z \in L^\infty(\Omega_{T-\tau})$ such that the pair (u, Z) satisfies (1.13) – (1.14) with T replaced by $T - \tau$.

Proposition 2.4. Let p, q satisfy (1.2) – (1.3) , and fix a function $u_{\star} \in X^p(0, T; \Omega) \cap C^0([0, T]; L^2(\Omega))$. Assume that $\{f_{\varepsilon}\}_{0\leq \varepsilon\leq 1}\subset L^q(\Omega_T)$ satisfy [\(2.13\)](#page-8-3). For each $\varepsilon\in (0, 1)$, we consider $u_{\varepsilon}\in u_{\star}+X_0^p$ $_{0}^{p}(0,\,T;\,\Omega),$ a weak solution of [\(2.12\)](#page-8-1). Then, there uniquely exists a function $u_0 \in u_* + X_0^p$ $_{0}^{p}(0, T-\tau; \Omega)$ for each fixed $\tau \in (0, T)$, such that

$$
u_{\varepsilon_j} \to u_0 \quad \text{in} \quad L^p(0, T - \tau; W^{1, p}(\Omega)), \quad \text{and} \quad \nabla u_{\varepsilon_j} \to \nabla u_0 \quad \text{a.e. in} \quad \Omega_{T - \tau}.
$$
 (2.18)

Moreover, this limit function u_0 is the unique weak solution of [\(2.17\)](#page-9-2).

The strong $L^p(W^{1,p})$ -convergence of weak solutions for parabolic approximate p-Laplace-type problems can be found in [\[32,](#page-33-0) Lemma 3.1]. The proof therein cannot be directly applied to our approximate problems, since the density E_1 , unlike E_p , lacks continuous differentiability at the origin. In particular, we have to deduce [\(2.18\)](#page-9-3) only by the weak convergence of u_{ε} or $\partial_t u_{\varepsilon}$. To prove the strong L^p -convergence of a spatial gradient, we have to appeal to the Aubin–Lions lemma and Lemma [2.2.](#page-8-2) As mentioned in Section [1.2,](#page-3-0) we use the technical assumption [\(1.3\)](#page-1-4) to guarantee $L^{p'}(0, T - \tau/2; W^{-1,p'}(\Omega))$ -bounds of time derivatives. In our proof, this bound estimate is required to make a weak compactness argument, and to apply the compact embedding X_0^p $_{0}^{p}(0, T-\tau/2; \Omega) \hookrightarrow$ $L^p(0, T - \tau/2; L^2(\Omega))$. Since our convergence argument is based on a weak convergence method and a parabolic compact embedding, this is substantially different from the proof of [\[32,](#page-33-0) Lemma 3.1].

Proof. We may choose $\varepsilon_0 \in (0, 1)$ and a constant $c_{\star} \in (K, \infty)$ such that $|E_{\varepsilon}(0)| + |\nabla E_{\varepsilon}(0)| \leq c_{\star}$ for all $\varepsilon \in (0, \varepsilon_0)$. We fix arbitrary $\tau \in (0, T)$, and define $0 < T_2 := T - \tau < T_1 := T - \tau/2 < T_0 := T$. For $j \in \{0, 1, 2\}$, we set a piecewise linear function $\theta_j(t) := \frac{(T_j - t)_+}{T_j} \in [0, 1]$ for $t \in [0, T]$. We first claim that there exists $C \in (1, \infty)$, independent of $\varepsilon \in (0, \varepsilon_0)$, such that

$$
\|\nabla u_{\varepsilon}\|_{L^{p}(\Omega_{T_1})} \leq C. \tag{2.19}
$$

To verify this, we test $\varphi = \varphi_0 := (u_{\varepsilon} - u_{\star})\theta_0$ into [\(2.16\)](#page-8-4). As a result, we have

$$
\mathbf{L}_1 + \mathbf{L}_2 := -\frac{1}{2} \iint_{\Omega_T} |u_{\varepsilon} - u_{\star}|^2 \partial_t \theta_0 \, dX + \iint_{\Omega_T} \langle \nabla E_{\varepsilon} (\nabla u_{\varepsilon}) | \nabla u_{\varepsilon} - \nabla u_{\star} \rangle \theta_0 \, dX
$$

=
$$
\iint_{\Omega_T} f_{\varepsilon} (u_{\varepsilon} - u_{\star}) \theta_0 \, dX - \int_0^T \langle \partial_t u_{\star}, \theta_0 (u_{\varepsilon} - u_{\star}) \rangle \, dt =: \mathbf{R}_1 - \mathbf{R}_2.
$$

For L_2 , we use the convexity of E_ε to obtain

$$
\mathbf{L}_2 \geqslant \iint_{\Omega_T} \left[E_{\varepsilon}(\nabla u_{\varepsilon}) - E_{\varepsilon}(\nabla u_{\star}) \right] \theta_0 \, \mathrm{d}X.
$$

By $\theta_0^p \le \theta_0 \le 1$ and Minkowski's inequality, the integral \mathbf{R}_2 is estimated as follows;

$$
|\mathbf{R}_2| \leq \|\partial_t u_{\star}\|_{L^{p'}(0,T;W^{-1,p'}(\Omega))} \left[\left(\iint_{\Omega_T} |\nabla u_{\varepsilon}|^p \theta_0 \, \mathrm{d}X \right)^{1/p} + \left(\iint_{\Omega_T} |\nabla u_{\star}|^p \theta_0 \, \mathrm{d}X \right)^{1/p} \right].
$$

For the remaining term \mathbf{R}_1 , we use Young's inequality and [\(2.14\)](#page-8-5) to get $|\mathbf{R}_1| \leq \mathbf{L}_1 + C F^2$. By [\(2.6\)](#page-7-4) and straightforward computations found in [\[44,](#page-34-0) §2.6], we can find constants $c_1 \in (0, 1), c_2 \in (1, \infty)$, $c_3 \in (1, \infty)$ such that

$$
c_1(p,\,\lambda)\left(|z|^p - c_2(p,\,c_\star)\right) \leqslant E_\varepsilon(z) \leqslant c_3(p,\,\Lambda,\,K,\,c_\star)(|z|^p + 1)
$$

for all $z \in \mathbb{R}^n$. Carrying out a straightforward absorbing argument, we can find the constant $C \in (1, \infty)$ satisfying (2.19) .

By [\(2.19\)](#page-9-4), it is easy to check that $\{u_{\varepsilon} - u_{\star}\}_{0 < \varepsilon < 1} \subset L^p(0, T; W_0^{1,p})$ $\mathcal{O}_0^{(1,p)}(\Omega)$ is bounded. Moreover, by $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ $(1.3), (2.4)–(2.5), \text{ and } (2.14)–(2.15), \text{ the time derivative } \partial_t u_{\varepsilon} \in L^{p'}(0, T_1; W^{-1,p'}(\Omega)) \text{ is also uniformly}$ bounded for $\varepsilon \in (0, \varepsilon_0)$. Therefore, by a weak compactness argument, there exists a sequence $\{\varepsilon_j\}_{j=1}^{\infty} \subset$ $(0, \varepsilon_0)$, which converges to 0, such that

$$
u_{\varepsilon_j} - u_{\star} \rightharpoonup u_0 - u_{\star}
$$
 in $L^p(0, T_1; W_0^{1, p}(\Omega)),$ (2.20)

$$
\partial_t u_{\varepsilon_j} \stackrel{*}{\rightharpoonup} \partial_t u_0 \quad \text{in} \quad L^{p'}(0, T_1; W^{-1, p'}(\Omega)), \tag{2.21}
$$

for some $u_0 \in u_\star + X_0^p$ $_{0}^{p}(0, T_{1}; \Omega)$. By making use of (2.20) – (2.21) , and integration by parts for the time variable, it is easy to check that $u_0|_{t=0} = u_\star|_{t=0}$ in $L^2(\Omega)$. We also invoke the compact embedding X_0^p $_{0}^{p}(0, T_1; \Omega) \hookrightarrow L^p(0, T_1; L^2(\Omega))$ from the Aubin–Lions lemma. In particular, we are allowed to use a strong convergence

$$
u_{\varepsilon_j} \to u_0 \quad \text{in} \quad L^p(0, T_1; L^2(\Omega)) \tag{2.22}
$$

by taking a subsequence if necessary.

From (2.20) – (2.22) , we would like to show

$$
\tilde{\mathbf{L}} := \iint_{\Omega_{T_1}} \left\langle \nabla E_{\varepsilon_j} (\nabla u_{\varepsilon_j}) - \nabla E_{\varepsilon_j} (\nabla u_0) \mid \nabla u_{\varepsilon_j} - \nabla u_0 \right\rangle \theta_1 \, dX \to 0 \tag{2.23}
$$

as $j \to \infty$. To prove [\(2.23\)](#page-10-3), we test $\varphi = \varphi_1 := (u_{\varepsilon} - u_0)\theta_1$ into [\(2.16\)](#page-8-4) with $\varepsilon = \varepsilon_j$. Then, we have

$$
\tilde{\mathbf{L}}_1 + \tilde{\mathbf{L}} := -\frac{1}{2} \iint_{\Omega_{T_1}} |u_{\varepsilon_j} - u_0|^2 \partial_t \theta_1 \, dX + \tilde{\mathbf{L}} \n= \iint_{\Omega_{T_1}} f_{\varepsilon_j} (u_{\varepsilon_j} - u_0) \theta_1 \, dX - \int_0^{T_1} \langle \partial_t u_0, (u_{\varepsilon_j} - u_0) \theta_1 \rangle_{W^{-1, p'}(\Omega), W_0^{1, p}(\Omega)} \, dt \n- \iint_{\Omega_{T_1}} \langle \theta_1 \nabla E_{\varepsilon_j} (\nabla u_0) | \nabla u_{\varepsilon_j} - \nabla u_0 \rangle \, dX \n=: \tilde{\mathbf{R}}_1 - \tilde{\mathbf{R}}_2 - \tilde{\mathbf{R}}_3.
$$

To estimate $\tilde{\mathbf{R}}_1$, we introduce $\kappa \in (0, 1)$, and define $r = r(\kappa, q) \in (0, q')$ as $r(\kappa, q) \coloneqq \frac{2q\kappa}{(1+\kappa)q}$ $\frac{2q\kappa}{(1+\kappa)q-2}$ for finite q, and $r(\kappa, \infty) := \frac{2\kappa}{1+r}$ $\frac{2\kappa}{1+\kappa}$. By Hölder's inequality and Young's inequality, we have

$$
\begin{aligned} |\tilde{\mathbf{R}}_1| \leqslant \left(\iint_{\Omega_{T_1}}|u_{\varepsilon_j}-u_0|^2 \,\mathrm{d}X\right)^{\frac{1-\kappa}{2}}&\left(\iint_{\Omega_{T_1}}|f_{\varepsilon_j}|^{\frac{2}{1+\kappa}}|u_{\varepsilon_j}-u_0|^{\frac{2\kappa}{1+\kappa}} \,\mathrm{d}X\right)^{\frac{1+\kappa}{2}}\\ \leqslant \tilde{\mathbf{L}}_1+C(\kappa)T_1^{\frac{1-\kappa}{1+\kappa}}&\left(\iint_{\Omega_{T_1}}|f_{\varepsilon_j}|^q \,\mathrm{d}X\right)^{\frac{2}{q(1+\kappa)}}&\left(\iint_{\Omega_{T_1}}|u_{\varepsilon_j}-u_0|^r \,\mathrm{d}X\right)^{\frac{2\kappa}{(1+\kappa)r}}=:\tilde{\mathbf{L}}_1+\tilde{\mathbf{R}}_4. \end{aligned}
$$

As a result, we get $0 \leq \tilde{L} \leq -(\tilde{R}_2 + \tilde{R}_3) + \tilde{R}_4$, and therefore it suffices to check $\tilde{R}_k \to 0$ for each $k \in \{2, 3, 4\}$. The integral $\tilde{\mathbf{R}}_2$ is computed as follows

$$
\tilde{\mathbf{R}}_2 = \int_0^{T_1} \langle \theta_1 \partial_t u_0, (u_{\varepsilon_j} - u_\star) - (u_0 - u_\star) \rangle_{W^{-1, p'}(\Omega), W_0^{1, p}(\Omega)} dt.
$$

Noting $\theta_1 \partial_t u_0 \in L^{p'}(0, T_1; W^{-1, p'}(\Omega))$ and (2.21) , we can deduce $\tilde{\mathbf{R}}_2 \to 0$. For $\tilde{\mathbf{R}}_3$, we recall Lemma [2.2,](#page-8-2) so that we are allowed to use $\theta_1 \nabla E_{\varepsilon_j}(\nabla u_0) \to \theta_1 A_0(\nabla u_0)$ in $L^{p'}(\Omega_{T_1})$. We already know ∇u_{ε_j} \to ∇u_0 in $L^p(\Omega_{T_1})$ by [\(2.20\)](#page-10-0). Combining these convergence results, we can easily notice $\tilde{\mathbf{R}}_3 \to 0$. For $\tilde{\mathbf{R}}_4$, we choose and fix $\kappa \in (0, 1)$ satisfying $0 < r(\kappa) < \min\{p, 2\}$. Then, [\(2.14\)](#page-8-5), [\(2.22\)](#page-10-2) and the continuous embedding $L^p(0, T_1; L^2(\Omega)) \hookrightarrow L^r(\Omega_{T_1})$ yield $\mathbf{R}_4 \to 0$. This completes the proof of [\(2.23\)](#page-10-3).

From [\(2.6\)](#page-7-4), [\(2.19\)](#page-9-4), and [\(2.23\)](#page-10-3), it is easy to deduce $\nabla u_{\varepsilon_j} \to \nabla u_0$ in $L^p(\Omega_{T_2})$, whence we may let [\(2.18\)](#page-9-3) hold, by taking a subsequence if necessary. This fact enables us to apply Lemma [2.2](#page-8-2) [\(2\).](#page-9-5) In particular, by choosing a suitable sequence $\{\varepsilon_j\}_j \subset (0, \varepsilon_0)$, and letting $j \to 0$, we conclude that the limit $u_0 \in u_* + X_0^p$ $_{0}^{p}(0, T_2; \Omega)$ admits a vector field $Z_0 \in L^{\infty}(\Omega_{T_2})$ such that the pair (u_0, Z_0) satisfies [\(1.13\)](#page-6-4)–[\(1.14\)](#page-6-5) with T replaced by T_2 . In other words, the limit u_0 is a weak solution of [\(2.17\)](#page-9-2). Our proof is completed by showing that the weak solution of [\(2.17\)](#page-9-2) is unique. Let another pair $(\tilde{u}, \tilde{Z}) \in (u_{\star} + X_0^p)$ $D_0^p(0, T_2; \Omega) \times L^{\infty}(\Omega_{T_2})$ satisfy (1.13) – (1.14) with T replaced by T_2 . Then,

$$
\int_0^{T_2} \langle \partial_t (u_0 - \tilde{u}), \varphi \rangle_{W^{-1, p'}(\Omega), W_0^{1, p}(\Omega)} dt + \iint_{\Omega_{T_2}} \langle Z_0 - \tilde{Z} | \nabla \varphi \rangle dX
$$
\n
$$
+ \iint_{\Omega_{T_2}} \langle \nabla E_p (\nabla u_0) - \nabla E_p (\nabla \tilde{u}) | \nabla \varphi \rangle dX = 0
$$
\n(2.24)

holds for all $\varphi \in X_0^p$ $p_0^p(0, T_2; \Omega)$. Thanks to the Steklov average, we may test $\varphi = \varphi_2 := (u_0 - \tilde{u})\theta_2$ into (2.24) . Here we may discard the first and the second integrals in the left-hand side of (2.24) , since they are positive. In fact, integrating by parts in time, we easily notice that the first integral in [\(2.24\)](#page-11-2) is nonnegative. For the second integral, we use subgradient inequalities to deduce $\langle Z_0 - Z | \nabla u_0 - \nabla \tilde{u} \rangle \geq 0$ a.e. in Ω_{T_2} . Letting $\tilde{\varepsilon} \to 0$ and recalling [\(2.7\)](#page-7-10), we have $\nabla(u_0 - \tilde{u}) = 0$ in $L^p(\Omega_{T_2})$. Recalling $u_0 - \tilde{u} \in L^p(0, T_2; W_0^{1, p})$ $\tilde{u}_0^{1,p}(\Omega)$) yields $u_0 = \tilde{u}$ in $L^p(0, T_2; W_0^{1,p})$ $\binom{1}{0}$ $\binom{1}{0}$.

Remark 2.5. To see the proof of Proposition [2.4,](#page-9-0) we only use the compact embedding X_0^p $_{0}^{p}(0, T; \Omega) \hookrightarrow \hookrightarrow$ $L^p(0, T; L^2(\Omega))$ to deal with the non-trivial external force term f_{ε} . In the simple case $f_{\varepsilon} \equiv 0$, Proposition [2.4](#page-9-0) is shown without this compact embedding.

2.3 Uniform a priori estimates and proof of Theorem [1.2](#page-6-1)

By Proposition [2.4,](#page-9-0) our gradient continuity problem is reduced to the regularity for $u_{\varepsilon} \in X^p(0, T; \Omega) \cap$ $C^0([0, T]; L^2(\Omega))$, a weak solution to

$$
\partial_t u_{\varepsilon} - \text{div} \left(\nabla E_{\varepsilon} (\nabla u_{\varepsilon}) \right) = f_{\varepsilon} \tag{2.25}
$$

in Ω_T . We prove local Hölder estimates of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$.

We consider [\(2.25\)](#page-11-3) in a parabolic cylinder $Q_R = Q_R(x_*, t_*) \coloneqq B_R(x_*) \times (t_* - R^2, t_*] \subseteq \Omega_T$. Then, we are allowed to use the interior regularity

$$
\nabla u_{\varepsilon} \in L^{\infty}(Q_R; \mathbb{R}^n) \quad \text{and} \quad \nabla^2 u_{\varepsilon} \in L^2(Q_R; \mathbb{R}^{n \times n}). \tag{2.26}
$$

Here we should note the fact that the parabolicity ratio of the Hessian matrix $\nabla^2 E_\varepsilon(\nabla u_\varepsilon)$ is everywhere bounded for each fixed $\varepsilon \in (0, 1)$. Therefore, it is possible to make standard parabolic arguments, including difference quotient methods and the De Giorgi–Nash–Moser theory([\[13,](#page-32-2) Chapters VIII], [\[33,](#page-33-11) Chapters III–V]; see also [\[10,](#page-32-5) Appendix]). Differentiating [\(2.25\)](#page-11-3) with respect to x_i for each $j \in \{1, \ldots, n\}$, we have

$$
-\int_{t_*-R^2}^{t_*} \langle \partial_t \partial_{x_j} u_{\varepsilon}, \, \varphi \rangle \, \mathrm{d}t + \iint_{Q_R} \left[\langle \nabla^2 E_{\varepsilon} (\nabla u_{\varepsilon}) \nabla \partial_{x_j} u_{\varepsilon} \, \middle| \, \nabla \varphi \rangle + f_{\varepsilon} \partial_{x_j} \varphi \right] \, \mathrm{d}X = 0 \tag{2.27}
$$

for all $\varphi \in X_0^2(t_* - R^2, t_*; B_R(x_*)).$

Remark 2.6. The coefficient matrix $\nabla^2 E_\varepsilon(\nabla u_\varepsilon)$ is uniformly parabolic in the classical sense for each fixed $\varepsilon \in (0, 1)$. Hence, by $E_{\varepsilon} \in C^{\infty}(\mathbb{R}^n)$ and $f_{\varepsilon} \in C^{\infty}(\Omega_T)$, it is not restrictive to let $u_{\varepsilon} \in C^{\infty}(\Omega_T)$ by bootstrap arguments. Thanks to this improved regularity, all of our computations, concerning

integrals of time derivatives $\partial_t u_{\varepsilon}$, $\partial_t \partial_{x_i} u_{\varepsilon}$, and $\partial_t V_{\varepsilon}$, make sense without making use of Steklov averages. However, it should be remarked that all of the computations in Sections [3](#page-15-0)[–5](#page-21-0) are carried out as long as we can treat the equation $\partial_t \partial_{x_j} u_\varepsilon - \text{div} \left(\nabla^2 E_\varepsilon (\nabla u_\varepsilon) \nabla \partial_{x_j} u_\varepsilon \right) = \partial_{x_j} f_\varepsilon$ locally in $L^2(W^{-1,2})$, which is guaranteed by [\(2.26\)](#page-11-4).

Theorem [2.7](#page-12-0) states local $L^{\infty} - L^p$ estimates of $V_{\varepsilon} := \sqrt{\varepsilon^2 + |\nabla u_{\varepsilon}|^2}$.

Theorem 2.7. Assume that the exponents p and q satisfy [\(1.2\)](#page-1-3). Let u_{ε} be a weak solution to [\(2.25\)](#page-11-3) in Ω_T with $\varepsilon \in (0, 1)$. Define the positive exponents d and d_1 by $(d, d_1) := (p/2, 2)$ when $p \geq 2$, and otherwise $(d, d_1) := (2p/[p(n+2)-2n], p')$. We also set $d_2 := d(n+2)/p$ when $n \geq 3$, and $d_2 := 2d\sigma/p$ when $n = 2$, where $\sigma \in (2, q/2)$ is arbitrarily fixed. Then, there exists a constant $C \in (1, \infty)$, which depends at most on n, p, q, λ , Λ , K , and σ , such that

$$
\sup_{Q_{\theta R}} V_{\varepsilon} \leq \frac{C}{(1-\theta)^{d_2}} \left(1 + \|f_{\varepsilon}\|_{L^q(Q_R)}^{d_1} + \iint_{Q_R} V_{\varepsilon}^p \, \mathrm{d}X \right)^{\frac{d}{p}} \tag{2.28}
$$

for every fixed $Q_R = Q_R(x_*, t_*) \subseteq \Omega_T$ with $R \in (0, 1)$ and $\theta \in (0, 1)$.

The key estimate in this paper is Theorem [2.8,](#page-12-1) where we do not necessarily require $p \in (p_c, \infty)$, but assume local L^{∞} -bounds of V_{ε} , uniformly for sufficiently small ε . In the case [\(1.2\)](#page-1-3), this assumption is already guaranteed by Theorem [2.7.](#page-12-0)

Theorem 2.8. Let $p \in (1, \infty)$ and $q \in (n+2, \infty]$. Fix $\delta \in (0, 1)$, and let $\varepsilon \in (0, \delta/8)$. Assume that u_{ε} is a weak solution to [\(2.25\)](#page-11-3) in $Q_R \nsubseteq \Omega_T$ with $R \in (0, 1)$ and [\(2.26\)](#page-11-4). Also, let the positive numbers F and μ_0 satisfy [\(2.14\)](#page-8-5) and

$$
\underset{Q_R}{\text{ess sup}} V_{\varepsilon} \le \delta + \mu_0 \tag{2.29}
$$

respectively. Then, the estimates

$$
|\mathcal{G}_{2\delta,\,\varepsilon}(\nabla u_{\varepsilon}(X))| \leq \mu_0 \quad \text{for all} \quad X \in Q_{\rho_0}(x_*,\,t_*),\tag{2.30}
$$

$$
|\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}(X_1)) - \mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}(X_2))| \leq C \left(\frac{d_p(X_1, X_2)}{\rho_0}\right)^{\alpha} \mu_0
$$
\n(2.31)

hold for all $X_1, X_2 \in Q_{\rho_0/2}(x_*, t_*)$. Here the radius $\rho_0 \in (0, R/4)$, the exponent $\alpha \in (0, 1)$, and the constant $C \in (1, \infty)$ depend at most on n, p, q, λ , Λ , K , F , ω_1 , ω_p , μ_0 , and δ .

To prove Theorem [2.8,](#page-12-1) we always assume

$$
\sup_{Q_{2\rho}} |\mathcal{G}_{\delta,\varepsilon}(\nabla u_{\varepsilon})| \leq \mu \leq \mu + \delta \leq M,
$$
\n(2.32)

or equivalently

$$
\sup_{Q_{2\rho}} V_{\varepsilon} \leq \mu + \delta \leq M,\tag{2.33}
$$

for $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \subset Q_R$. Here $M := \mu_0 + \delta$ is a fixed constant, and $\mu \in (0, \mu_0]$ denotes a parameter that stands for a local bound of $\mathcal{G}_{\delta,\varepsilon}(\nabla u_{\varepsilon})$. We introduce a sufficiently small ratio $\nu \in (0, 1/4)$, which is determined later in Section [6,](#page-24-0) and define a superlevel set $S_{\rho,\mu,\nu} := \{(x, t) \in Q_\rho \mid V_\varepsilon(x, t) - \delta > (1-\nu)\mu\},\$ often denoted by S_ρ when μ and ν are clear. We may often let

$$
0 < \delta < \mu,\tag{2.34}
$$

since otherwise $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}) = 0$ holds in $Q_{2\rho}$, and therefore some oscillation estimates related to $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$ become trivial. We introduce an exponent $\beta \in (0, 1)$ by $\beta := 1 - (n + 2)/q$ when $q \in$ $(n + 2, \infty)$. For $q = \infty$, we let β be an arbitrarily fixed number in $(0, 1)$. Dividing the possible cases by measuring the size of $S_{\rho,\mu,\nu} \subset Q_{\rho}$, we prove Propositions [2.9](#page-13-0)[–2.10](#page-13-1) as below.

Proposition 2.9. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \in \Omega_T$ satisfy [\(2.32\)](#page-12-2) and [\(2.34\)](#page-12-3). If

$$
|S_{\rho,\,\mu,\,\nu}| \leqslant (1-\nu)|Q_{\rho}|\tag{2.35}
$$

holds for some $\nu \in (0, 1/4)$, then there exist an exponent $\kappa \in ((\sqrt{\nu}/6)^{\beta}, 1)$ and a radius $\tilde{\rho} \in (0, 1)$, depending at most on n, p, q, λ , Λ , K , F , M , δ , and ν , such that

$$
\sup_{Q_{\sqrt{\nu}\rho/3}} |\mathcal{G}_{\delta,\,\varepsilon}(\nabla u_{\varepsilon})| \leq \kappa\mu,\tag{2.36}
$$

holds, provided $\rho \leq \tilde{\rho}$.

Proposition 2.10. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \in Q_R$ satisfy (2.33) – (2.34) . Then, there exist $\hat{\rho} \in (0, 1)$ and $\nu \in (0, 1/4)$, depending at most on n, p, q, λ , Λ , K, F, M, δ , and $\omega = \omega_{\delta,M}$ given by Lemma [2.1,](#page-7-5) such that if there hold

$$
|S_{\rho,\,\mu,\,\nu}| > (1-\nu)|Q_{\rho}| \tag{2.37}
$$

and $\rho \leq \hat{\rho}$, then the limit $\Gamma_{2\delta,\varepsilon} := \lim_{r\to 0} \left(\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})\right)_r \in \mathbb{R}^n$ exists. Moreover, this limit satisfies

$$
\iint_{Q_r} |\mathcal{G}_{2\delta,\,\varepsilon}(\nabla u_\varepsilon) - \Gamma_{2\delta,\,\varepsilon}|^2 \,\mathrm{d}X \leqslant \left(\frac{r}{\rho}\right)^{2\beta} \mu^2 \quad \text{for all } r \in (0,\,\rho].
$$

The desired exponent α in Theorem [2.8](#page-12-1) is given by

$$
\alpha := \frac{\log \kappa}{\log(\sqrt{\nu}/6)} \in (0, \beta), \tag{2.38}
$$

where ν and $\kappa = \kappa(\nu)$ are respectively determined by Propositions [2.10](#page-13-1) and [2.9.](#page-13-0) To see our proofs of Propositions [2.9](#page-13-0)[–2.10,](#page-13-1) given in Sections [5](#page-21-0)[–6,](#page-24-0) we notice that $\nu \to 0$ and $\kappa \to 1$, as $\delta \to 0$. Therefore, the exponent α determined by [\(2.38\)](#page-13-2) will vanish as $\delta \rightarrow 0$.

We make different analyses, depending on whether a gradient may vanish or not. This will be judged by measuring the ratio $|S_{\rho,\mu,\nu}|/|Q_{\rho}|$. It is worth mentioning that (2.37) , the assumption in Proposition [2.10,](#page-13-1) roughly states that the modulus V_{ε} will stay close to its upper bound $\mu + \delta$. This indicates that the average integral of ∇u_{ε} will never vanish, which is rigorously verified by non-trivial energy estimates. Under the other condition [\(2.35\)](#page-13-4) in Proposition [2.9,](#page-13-0) we will probably face the case where a gradient may vanish. For this degenerate case, we should make suitable truncation to avoid analysis over facets. Such divisions by cases, based on the size of the superlevel set of the modulus, can be found in fundamental works in the p-Laplace regularity theory (e.g., [\[13\]](#page-32-2), [\[37\]](#page-33-3)). In the proof of Propositions [2.9–](#page-13-0)[2.10,](#page-13-1) however, we fully appeal to [\(2.34\)](#page-12-3), and do not use the intrinsic scaling methods found in the p-Laplace regularity theory.

Theorem [2.7,](#page-12-0) Propositions [2.9](#page-13-0) and [2.10](#page-13-1) are respectively shown in Sections [4,](#page-18-0) [5,](#page-21-0) and [6.](#page-24-0) We conclude Section [2](#page-6-0) by giving the proofs of Theorems [2.8](#page-12-1) and [1.2.](#page-6-1)

Proof of Theorem [2.8.](#page-12-1) We set $M := \mu_0 + \delta$, and choose $\nu \in (0, 1/4)$ and $\hat{\rho} \in (0, 1)$ as in Proposition [2.10.](#page-13-1) Corresponding to this ν , we set $\sigma := \sqrt{\nu/6}$, and choose $\kappa \in (\sigma^{\beta}, 1)$ and $\tilde{\rho} \in (0, 1)$ as in Proposition [2.9.](#page-13-0) We set $\rho_0 := \min\{\hat{\rho}, \tilde{\rho}, R/4\} \in (0, R/4)$. Then, (2.32) – (2.33) hold with $(\rho, \mu) := (\rho_0, \mu_0)$ and for every center point $(x_0, t_0) \in Q_{\rho_0}(x_*, t_*).$

We claim that the limit $\Gamma_{2\delta,\varepsilon}(x_0, t_0) := \lim_{r \to 0} (\mathcal{G}_{2\delta,\varepsilon}(\nabla u_\varepsilon))_{Q_r(x_0, t_0)} \in \mathbb{R}^n$ exists for each $(x_0, t_0) \in$ $Q_{\rho_0}(x_*, t_*)$. Moreover, this limit satisfies

$$
\iint_{Q_r(x_0,t_0)} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_\varepsilon) - \Gamma_{2\delta,\varepsilon}(x_0,t_0)|^2 \, \mathrm{d}X \le C \left(\frac{r}{\rho_0}\right)^{2\alpha} \mu_0^2 \tag{2.39}
$$

for all $r \in (0, \rho_0]$. Here the exponent $\alpha \in (0, \beta)$ is defined by [\(2.38\)](#page-13-2), and $C \in (1, \infty)$ depends at most on ν and α . To show [\(2.39\)](#page-13-5), we define $\mu_k := \kappa^k \mu_0$, $\rho_k := \sigma^k \rho_0$ for $k \in \mathbb{N}$. For these decreasing sequences $\{\mu_k\}_{k=0}^{\infty}$, $\{\rho_k\}_{k=0}^{\infty}$, we define

$$
\mathcal{N} \mathrel{\mathop:}= \{ k \in \mathbb{Z}_{\geqslant 0} \mid \mu_k > \delta, \text{ and } |S_{\rho_k, \, \mu_k, \, \nu}(x_0, \, t_0)| \leqslant (1-\nu)|Q_{\rho_k}(x_0, \, t_0)| \},
$$

The proper inclusion $\mathcal{N} \subsetneq \mathbb{Z}_{\geqslant 0}$ is clear, since $\mu_k \to 0$ as $k \to \infty$. Thus, we can define $k_{\star} \in \mathbb{Z}_{\geqslant 0}$ by the minimum number satisfying $k_{\star} \notin \mathcal{N}$. By repeatedly applying Proposition [2.9](#page-13-0) with $(\rho, \mu) = (\rho_k, \mu_k)$ for $k \in \{0, \ldots, k_{\star} - 1\}$, we have

$$
\sup_{Q_{2\rho_k}(x_0, t_0)} |\mathcal{G}_{2\delta, \varepsilon}(\nabla u_{\varepsilon})| \leq \sup_{Q_{2\rho_k}(x_0, t_0)} |\mathcal{G}_{\delta, \varepsilon}(\nabla u_{\varepsilon})| \leq \mu_k
$$
\n(2.40)

for every $k \in \{0, \ldots, k_\star\}$. For $k_\star \notin \mathcal{N}$, there are two possible cases.

If $\mu_{k_{\star}} > \delta$, then $|S_{\rho_{k_{\star}},\mu_{k_{\star}},\nu}| > (1-\nu)|Q_{\rho_{k_{\star}}}$ must hold. By Proposition [2.10,](#page-13-1) the limit $\Gamma_{2\delta,\varepsilon}(x_0, t_0) \in$ \mathbb{R}^n exists. Moreover, we have $|\Gamma_{2\delta,\varepsilon}(x_0, t_0)| \leq \mu_{k_\star}$, and

$$
\iint_{Q_r(x_0,t_0)} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_\varepsilon) - \Gamma_{2\delta,\varepsilon}(x_0,t_0)|^2 dX \leqslant \left(\frac{r}{\rho_{k_\star}}\right)^{2\beta} \mu_{k_\star}^2 \leqslant \left(\frac{r}{\rho_0}\right)^{2\alpha} \mu_0^2
$$

for all $r \in (0, \rho_{k_\star}]$, where we have used $\alpha \leq \beta$ and $\kappa = \sigma^\alpha$. For $r \in (\rho_{k_\star}, \rho_0]$, there corresponds a unique $k \in \{0, \ldots, k_{\star} - 1\}$ such that $\rho_{k+1} < r \leq \rho_k$. Then, it is obvious that $\sigma^k < \sigma^{-1} \cdot \frac{r}{\rho_k}$ $\frac{r}{\rho_0} = \frac{6}{\sqrt{2}}$ $rac{r}{\nu} \cdot \frac{r}{\rho_0}$ $\frac{r}{\rho_0}$. Hence, we can compute

$$
\iint_{Q_r(x_0, t_0)} |\mathcal{G}_{2\delta, \varepsilon}(\nabla u_{\varepsilon}) - \Gamma_{2\delta, \varepsilon}(x_0, t_0)|^2 dX \leq 2 \left(|\Gamma_{2\delta, \varepsilon}(x_0, t_0)|^2 + \sup_{Q_{\rho_k}} |\mathcal{G}_{2\delta, \varepsilon}(\nabla u_{\varepsilon})|^2 \right)
$$

$$
\leq 4\mu_k^2 \leq 4 \left(\frac{6}{\sqrt{\nu}} \right)^{2\alpha} \left(\frac{r}{\rho_0} \right)^{2\alpha} \mu_0^2.
$$

This completes the proof of [\(2.39\)](#page-13-5) in the case $\mu_{k_{\star}} > \delta$.

In the remaining case $0 < \mu_{k_{\star}} \leq \delta$, it is clear that $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}) \equiv 0$ in $Q_{\rho_{k_{\star}}}$. In particular, the identity $\Gamma_{2\delta,\varepsilon}(x_0, t_0) = 0$ holds, and it is easy to check that [\(2.40\)](#page-14-0) is satisfied for all $k \in \mathbb{Z}_{\geq 0}$. For each $r \in (0, \rho_0]$, there corresponds a unique number $k \in \mathbb{Z}_{\geq 0}$ such that $\rho_{k+1} < r \leq \rho_k$. By [\(2.40\)](#page-14-0) and $\sigma^k < \frac{6}{\sqrt{2}}$ $\frac{r}{\overline{\nu}} \cdot \frac{r}{\rho_0}$ $\frac{r}{\rho_0}$, we have

$$
\iint_{Q_r(x_0,t_0)} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_\varepsilon) - \Gamma_{2\delta,\varepsilon}(x_0,t_0)|^2 dX \leq \mu_k^2 = \sigma^{2\alpha k} \mu_0^2 \leqslant \left(\frac{6}{\sqrt{\nu}}\right)^{2\alpha} \left(\frac{r}{\rho_0}\right)^{2\alpha} \mu_0^2.
$$

In any possible cases, we conclude [\(2.39\)](#page-13-5) with $C \coloneqq 4 \cdot (6/\sqrt{\nu})^{2\alpha}$.

The mapping $\Gamma_{2\delta,\varepsilon}$ is a Lebesgue representative of $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon})$ by Lebesgue's differentiation theo-rem. Therefore, [\(2.30\)](#page-12-5) is obvious since $|\Gamma_{2\delta,\varepsilon}(x_0, t_0)| \leq \mu_0$ for all $(x_0, t_0) \in Q_{\rho_0}(x_*, t_*)$. The proof of [\(2.31\)](#page-12-6) is completed by showing that $\Gamma_{2\delta,\varepsilon}$ satisfies $|\Gamma_{2\delta,\varepsilon}(X_1) - \Gamma_{2\delta,\varepsilon}(X_2)| \leqslant C \left(\frac{d_p(X_1, X_2)}{\rho_0}\right)^{\alpha} \mu_0$ for all $X_1 = (x_1, t_1), X_2 = (x_2, t_2) \in Q_{\rho_0/2}(x_*, t_*).$ Firstly, we consider the case $r := d_p(X_1, X_2) \leq 2\rho_0/3.$ We fix a point $X_3 = (x_3, t_3) := \left(\frac{x_1 + x_2}{2}, \min\{t_1, t_2\}\right)$. The inclusions $Q_r(X_3) \subset Q_{3r/2}(X_j) \subset$ $Q_{\rho_0}(X_j) \subset Q_{2\rho_0}(X_j) \subset Q_{4\rho_0}(X_*)$ are easily checked for each $j \in \{1, 2\}$. Applying [\(2.39\)](#page-13-5) with $Q_r(x_0, t_0)$ replaced by $Q_{3r/2}(X_j)$ $(j \in \{1, 2\})$, we obtain

$$
\left|\Gamma_{2\delta,\varepsilon}(X_1) - \Gamma_{2\delta,\varepsilon}(X_2)\right|^2 \leq 2\sum_{j=1}^2 \iint_{Q_r(X_3)} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_\varepsilon) - \Gamma_{2\delta,\varepsilon}(X_j)|^2 \,dX \leq C_{n,\nu} \left(\frac{r}{\rho_0}\right)^{2\alpha} \mu_0^2,
$$

from which [\(2.31\)](#page-12-6) is clear. The remaining case $r > 2\rho_0/3$ is easier, since this condition allows us to compute $|\Gamma_{2\delta,\,\varepsilon}(X_1) - \Gamma_{2\delta,\,\varepsilon}(X_2)| \leq 2\mu_0 \leq 2 \cdot \left(\frac{3r}{2\rho_0}\right)$ $\frac{3r}{2\rho_0}$ $\int_0^\alpha \mu_0$, which completes the proof of [\(2.31\)](#page-12-6).

Proof of Theorem [1.2.](#page-6-1) Fix subdomains $\mathcal{Q} \in \mathcal{Q} \in \Omega_{T-\tau}$ with $\tau \in (0, T)$, and let $\delta \in (0, 1)$ and $\varepsilon_k \in (0, \delta/8)$. For each $\varepsilon = \varepsilon_k$, we consider u_{ε_k} , the unique weak solution of [\(2.12\)](#page-8-1). Proposition [2.4](#page-9-0) allows us to assume that $\nabla u_{\varepsilon_k} \to \nabla u$ a.e. in $\Omega_{T-\tau}$, by taking a subsequence if necessary. Moreover, we can choose the positive constants U_{τ} and F , satisfying $\|\nabla u_{\varepsilon_k}\|_{L^p(\Omega_{T-\tau})} \leq U_{\tau}$ and $\|f_{\varepsilon_k}\|_{L^q(\Omega_T)} \leq F$ respectively. By Theorem [2.7,](#page-12-0) we can find a constant $\mu_0 \in (1, \infty)$, which depends at most on n, p, q, $\lambda, \Lambda, K, U_{\tau}, F$, and $dist_p(\tilde{Q}, \partial_p \Omega_{T-\tau})$, such that $V_{\varepsilon} \leq \delta + \mu_0$ holds a.e. in \tilde{Q} . Thanks to this uniform bound and Theorem [2.8,](#page-12-1) we can apply the Arzelà–Ascoli theorem to the sequence $\{\mathcal{G}_{2\delta, \varepsilon_k}(\nabla u_{\varepsilon_k})\}_k \subset$

 $C^0(Q;\mathbb{R}^n)$. In particular, we may let $\mathcal{G}_{2\delta,\varepsilon_k}(\nabla u_{\varepsilon_k})$ uniformly converge to a Hölder continuous mapping $v_{2\delta}$ in Q. Meanwhile, we have $\mathcal{G}_{2\delta,\varepsilon_k}(\nabla u_{\varepsilon_k}) \to \mathcal{G}_{2\delta}(\nabla u)$ a.e. in $\Omega_{T-\tau}$. These convergence results imply that $\mathcal{G}_{2\delta}(\nabla u) = v_{2\delta}$ a.e. in $\mathcal{Q} \in \Omega_{T-\tau}$. Thus, we conclude $\mathcal{G}_{2\delta}(\nabla u) \in C^0(\Omega_T; \mathbb{R}^n)$ for each fixed $\delta \in (0, 1)$.

By the definition of \mathcal{G}_{δ} , it is obvious that $\sup_{\Omega_T} |\mathcal{G}_{\delta_1}(\nabla u) - \mathcal{G}_{\delta_2}(\nabla u)| \leq |\delta_1 - \delta_2|$ for all $\delta_1, \delta_2 \in (0, 1)$. Therefore, $\mathcal{G}_{\delta}(\nabla u) \in C^0(\Omega_T; \mathbb{R}^n)$ uniformly converges as $\delta \to 0$. Since $\mathcal{G}_{\delta}(\nabla u) \to \nabla u$ a.e. in Ω_T as $\delta \to 0$, this implies $\nabla u \in C^0(\Omega_T; \mathbb{R}^n)$.

3 Basic estimates of approximate solutions

In Section [3,](#page-15-0) we aim to give weak formulations and some local estimates of u_{ε} , which will be used in Sections [5–](#page-21-0)[6.](#page-24-0)

3.1 A basic weak formulation

From [\(2.27\)](#page-11-5), we deduce a basic weak formulation concerning $V_{\varepsilon} = \sqrt{\varepsilon^2 + |\nabla u_{\varepsilon}|^2}$ in a systematic approach.

Lemma 3.1. Let u_{ε} be a weak solution to [\(2.25\)](#page-11-3) in $Q = Q_R(x_*, t_*) \in \Omega_T$ with $\varepsilon \in (0, 1)$ and [\(2.26\)](#page-11-4). Assume that ψ is a non-decreasing, non-negative, and locally Lipschitz function that is differentiable except at finitely many points. For this ψ , we define a convex function

$$
\Psi(s) := \int_0^s \sigma \psi(\sigma) d\sigma \quad \text{for} \quad s \in [0, \infty). \tag{3.1}
$$

Let $\zeta \in X_0^2(t_* - R^2_*, t_*; B_R(x_*)) \cap L^{\infty}(\Omega_T)$ be a non-negative function that is compactly supported in the interior of Q. Then, there holds

$$
2J_0 + 2J_1 + J_2 + J_3 \le n\lambda^{-1}J_4 + 2J_5,\tag{3.2}
$$

where J_0, \ldots, J_5 are the integrals defined as

$$
J_0 := \int_{t_*-R^2}^{t_*} \langle \partial_t [\Psi(V_{\varepsilon})] , \zeta \rangle dt, \quad J_1 := \iint_Q \langle \nabla^2 E_{\varepsilon} (\nabla u_{\varepsilon}) \nabla [\Psi(V_{\varepsilon})] | \nabla \zeta \rangle dX,
$$

\n
$$
J_2 := \iint_Q \langle \nabla^2 E_{\varepsilon} (\nabla u_{\varepsilon}) \nabla V_{\varepsilon} | \nabla V_{\varepsilon} \rangle \zeta \psi'(V_{\varepsilon}) V_{\varepsilon} dX,
$$

\n
$$
J_3 := \sum_{j=1}^n \iint_Q \langle \nabla^2 E_{\varepsilon} (\nabla u_{\varepsilon}) \nabla \partial_{x_j} u_{\varepsilon} | \nabla \partial_{x_j} u_{\varepsilon} \rangle \zeta \psi(V_{\varepsilon}) dX,
$$

\n
$$
J_4 := \iint_Q |f_{\varepsilon}|^2 V_{\varepsilon}^{2-p} (\psi(V_{\varepsilon}) + \psi'(V_{\varepsilon}) V_{\varepsilon}) \zeta dX, \quad J_5 := \iint_Q |f_{\varepsilon}| |\nabla \zeta| \psi(V_{\varepsilon}) V_{\varepsilon} dX.
$$

Before the proof, it should be mentioned that the integrals J_2 and J_3 are non-negative. More precisely, the inequalities

$$
J_2 \ge \lambda \iint_Q V_{\varepsilon}^{p-1} |\nabla V_{\varepsilon}|^2 \zeta \psi'(V_{\varepsilon}) \,dX \quad \text{and} \quad J_3 \ge \lambda \iint_Q V_{\varepsilon}^{p-2} |\nabla^2 u_{\varepsilon}|^2 \zeta \psi(V_{\varepsilon}) \,dX \tag{3.3}
$$

follow from (2.4) – (2.5) and ψ , ψ' , $\zeta \ge 0$.

Proof. For each $j \in \{1, \ldots, n\}$, we test $\varphi := \zeta \psi(V_{\varepsilon}) \partial_{x_j} u_{\varepsilon} \in X_0^2(t_* - R^2_*, t_*; B_R(x_*))$ into [\(2.27\)](#page-11-5). Summing over $j \in \{ 1, \ldots, n \}$ and computing similarly to [\[44,](#page-34-0) Lemma 3.3], we have

$$
J_0 + J_1 + J_2 + J_3 = -\iint_Q f_\varepsilon \left(\psi(V_\varepsilon) \langle \nabla u_\varepsilon | \nabla \zeta \rangle + \zeta \psi'(V_\varepsilon) \langle \nabla u_\varepsilon | \nabla V_\varepsilon \rangle + \zeta \psi(V_\varepsilon) \Delta u_\varepsilon \right) dX
$$

$$
\leq J_5 + \frac{1}{2} (J_2 + J_3) + \frac{n}{2\lambda} J_4
$$

by Young's inequality and [\(3.3\)](#page-15-2). From this, [\(3.2\)](#page-15-3) immediately follows.

 \Box

The weak formulation [\(3.2\)](#page-15-3) is viewed in two ways. The first is that the composite function $\Psi(V_{\varepsilon})$ becomes a subsolution to a certain parabolic equation. This is easy to deduce by discarding the nonnegative terms J_2 and J_3 in [\(3.2\)](#page-15-3). In Section [3.2,](#page-16-0) we suitably choose ψ and Ψ to apply De Giorgi's truncation. The second is that we can obtain local L^2 -estimates for the Hessian matrix $\nabla^2 u_{\varepsilon}$ from the integral J_3 . The detailed computations are given in Section [3.3.](#page-17-0)

3.2 De Giorgi's truncation

We prove that the convex composite function $U_{\delta,\varepsilon} := (V_{\varepsilon} - \delta)^2_+$ is a weak subsolution to a uniformly parabolic equation. This fact implies that as in Lemma [3.2](#page-16-1) below, $U_{\delta,\varepsilon}$ belongs to a certain parabolic De Giorgi class (see [\[33,](#page-33-11) Chapter II, §7], [\[35,](#page-33-12) Chapter VI, §13] as related items).

Lemma 3.2. Let the assumptions of Proposition [2.9](#page-13-0) be verified. Fix a subcylinder $Q_0 := B_r \times$ $(T_0, T_1] \subset Q_{2\rho}$. Then, for all $k \in (0, \infty)$, $\eta \in C_c^1(B_r; [0, 1])$, $\phi_c \in C^1([T_0, T_1]; [0, 1])$ satisfying $\phi_{\rm c}(T_0)=0$, we have

$$
\sup_{\tau \in (T_0, T_1)} \int_{B_r \times \{\tau\}} (U_{\delta, \varepsilon} - k)_+^2 \eta^2 \phi_c \, dx + \iint_{Q_0} |\nabla (U_{\delta, \varepsilon} - k)_+|^2 \eta^2 \phi_c \, dX \tag{3.4}
$$
\n
$$
\leq C \left[\int_{Q_0} (U_{\delta, \varepsilon} - k)_+^2 \left(|\nabla \eta|^2 + |\partial_t \phi_c| \right) \, dX + \mu^4 F^2 |A_k|^{1 - 2/q} \right],
$$
\n
$$
\int_{B_r \times \{\tau\}} (U_{\delta, \varepsilon} - k)_+ \eta^2 \, dx - \int_{B_r \times \{\tau_0\}} (U_{\delta, \varepsilon} - k)_+ \eta^2 \, dx \tag{3.5}
$$
\n
$$
\leq C \left[\int_{Q_0} (U_{\delta, \varepsilon} - k)_+^2 |\nabla \eta|^2 \, dX + \mu^4 F^2 |A_k|^{1 - 2/q} \right] \quad \text{for all } \tau \in (T_0, T_1).
$$

Here $A_k := \{(x, t) \in Q_0 \mid U_{\delta, \varepsilon}(x, t) > k\}$, and the constant $C \in (1, \infty)$ depends at most on n, p, q, λ , Λ , K , M , and δ .

Proof. We apply Lemma [3.1](#page-15-4) with $\psi(\sigma) := 2(1 - \delta/\sigma)_+$, so that $\Psi(V_{\varepsilon}) = U_{\delta, \varepsilon}$. Then, it is easy to compute $V_{\varepsilon}\psi(V_{\varepsilon}) = 2U_{\delta,\varepsilon}^{1/2}$, and $V_{\varepsilon}^{2-p}(\psi(V_{\varepsilon}) + V_{\varepsilon}\psi'(V_{\varepsilon})) \leq 2\delta^{1-p}V_{\varepsilon}$. Remarking that all the integrands in [\(3.2\)](#page-15-3) vanish in the place $\{V_{\varepsilon} \leq \delta\}$, we may replace $\nabla^2 E_{\varepsilon}(\nabla u_{\varepsilon})$ by a measurable matrix $\mathcal{A}_{\delta,\varepsilon}(\nabla u_{\varepsilon})$, which coincides with $\nabla^2 E_{\varepsilon}(\nabla u_{\varepsilon})$ on $\{V_{\varepsilon} > \delta\}$, but $\mathcal{A}_{\delta,\varepsilon}(\nabla u_{\varepsilon}) \equiv \mathrm{id}_n$ on $\{V_{\varepsilon} \leq \delta\}$. This matrix satisfies $\lambda_*\mathrm{id}_n \leqslant \mathcal{A}_{\delta,\varepsilon}(\nabla u_\varepsilon) \leqslant \Lambda_*\mathrm{id}_n$ for some constants $\lambda_* \in (0, 1), \Lambda_* \in (1, \infty)$, depending at most on λ, Λ , K, δ and M. Discarding $J_2 \geq 0$ and $J_3 \geq 0$, we obtain

$$
\int_{T_0}^{T_1} \langle \partial_t U_{\delta,\varepsilon}, \zeta \rangle dt + \iint_{Q_0} \langle \mathcal{A}_{\delta,\varepsilon} (\nabla u_\varepsilon) \nabla U_{\delta,\varepsilon} \mid \nabla \zeta \rangle dX \leq C \mu \left(\iint_{Q_0} \left[|f_\varepsilon|^2 \zeta + |f_\varepsilon| |\nabla \zeta| \right) dX \right]
$$

for any non-negative $\zeta \in X_0^2(T_0, T_1; B_r) \cap L^{\infty}(Q_0)$ that is compactly supported in the interior of Q_0 . We test $\zeta := (U_{\delta, \varepsilon} - k)_+ \eta^2 \phi$, where $\phi \colon [T_0, T_1] \to [0, 1]$ is an arbitrary Lipschitz function satisfying $\phi(T_0) = \phi(T_1) = 0$. Then, we have

$$
-\frac{1}{2} \iint_{Q_0} (U_{\delta, \varepsilon} - k)_+^2 \eta^2 \partial_t \phi \, dX + \lambda_* \iint_{Q_0} |\nabla (U_{\delta, \varepsilon} - k)_+|^2 \eta^2 \phi \, dX
$$

\$\leqslant \frac{\lambda_*}{2} \iint_{Q_0} |\nabla (U_{\delta, \varepsilon} - k)_+|^2 \eta^2 \phi \, dX + C \left[\iint_{Q_0} (U_{\delta, \varepsilon} - k)_+^2 |\nabla \eta|^2 \phi \, dX + \mu^4 \iint_{A_k} |f_{\varepsilon}|^2 \eta^2 \phi \, dX \right],

where we have used Young's inequality and (2.34) . By Hölder's inequality, we obtain

$$
-\frac{1}{2} \iint_{Q_0} (U_{\delta,\varepsilon} - k)_+^2 \eta^2 \partial_t \phi \, dX + \frac{\lambda_*}{2} \iint_{Q_0} |\nabla (U_{\delta,\varepsilon} - k)_+|^2 \eta^2 \phi \, dX
$$

\$\leqslant C \left[\iint_{Q_0} (U_{\delta,\varepsilon} - k)_+^2 |\nabla \eta|^2 \phi \, dX + \mu^4 F^2 |A_k|^{1-2/q} \right].

The desired claims (3.4) – (3.5) are easily deduced by suitably choosing ϕ .

 \Box

3.3 L ²-energy estimates

We deduce energy estimates related to $G_{p,\varepsilon}(\nabla u_{\varepsilon}) = V_{\varepsilon}^{p-1} \nabla u_{\varepsilon}$, under the assumptions [\(2.34\)](#page-12-3) and $0 < \nu < 1/4$. Here $G_{p,\varepsilon} : \mathbb{R}^n \to \mathbb{R}^n$ is the bijective mapping defined as $G_{p,\varepsilon}(z) := (\varepsilon^2 + |z|^2)^{(p-1)/2} z$ for $z \in \mathbb{R}^n$.

Lemma 3.3. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \in \Omega_T$ satisfy [\(2.14\)](#page-8-5) and [\(2.33\)](#page-12-4)–[\(2.34\)](#page-12-3). Then, the following estimates hold for any $\sigma \in (0, 1), \nu \in (0, 1/4)$.

$$
\iint_{Q_{\sigma\rho}} |\nabla \left[G_{p,\varepsilon} (\nabla u_{\varepsilon}) \right]|^2 \, \mathrm{d}X \leq \frac{C\mu^{2p}}{\sigma^{n+2}\rho^2} \left[\frac{1}{(1-\sigma)^2} + F^2 \rho^{2\beta} \right]. \tag{3.6}
$$

$$
|Q_{\sigma\rho}|^{-1} \int_{S_{\sigma\rho,\,\mu,\,\nu}} |\nabla \left[G_{p,\,\varepsilon}(\nabla u_{\varepsilon}) \right]|^2 \, \mathrm{d}X \leq \frac{C\mu^{2p}}{\sigma^{n+2}\rho^2} \left[\frac{\nu}{(1-\sigma)^2} + \frac{F^2 \rho^{2\beta}}{\nu} \right]. \tag{3.7}
$$

Here $C \in (0, \infty)$ in [\(3.6\)](#page-17-1)–[\(3.7\)](#page-17-2) depends at most on n, p, λ , Λ , K , δ , and M.

Proof. For each fixed $\sigma \in (0, 1)$, we choose cut-off functions $\eta \in C_c^1(B_\rho; [0, 1])$ and $\phi_c \in C^1(\overline{I_\rho}; [0, 1])$ satisfying $\phi_c(t_0 - \rho^2) = 0$, $\eta|_{B_{\sigma\rho}} \equiv 1$, and $\phi_c|_{I_{\sigma\rho}} \equiv 1$. For sufficiently small $\tilde{\varepsilon} > 0$, which tends to 0 later, we choose $\phi_h: \overline{I_\rho} \to [0, 1]$ satisfying $\phi_h \equiv 1$ on $[t_0 - \rho^2, t_0 - \tilde{\varepsilon}]$, and linearly interpolated on the other domain with $\phi_h(t_0) = 0$. By $|\nabla V_{\varepsilon}|^2 \leq |\nabla^2 u_{\varepsilon}|^2$, it is easy to check that $|\nabla [G_{p,\varepsilon}(\nabla u_{\varepsilon})]|^2 \leq$ $c_p V_{\varepsilon}^{2p-2} |\nabla^2 u_{\varepsilon}|^2$ for some $c_p \in (1, \infty)$. With this in mind, we apply Lemma [3.1](#page-15-4) with $\psi(s) := \tilde{\psi}(s)\sigma^p$, and $\zeta := \eta^2 \phi$ with $\phi := \phi_c \phi_h$. Here $\tilde{\psi} : [0, \infty) \to [0, \infty)$ is a non-decreasing function that is chosen later. The weak formulation [\(3.2\)](#page-15-3) becomes

$$
\mathbf{L}_{1} + \mathbf{L}_{2} + \mathbf{L}_{3} := -2 \iint_{Q_{\rho}} \eta^{2} \Psi(V_{\varepsilon}) \phi_{\varepsilon} \partial_{t} \phi_{h} dX + \iint_{Q_{\rho}} \left\langle \nabla^{2} E_{\varepsilon}(\nabla u_{\varepsilon}) \nabla V_{\varepsilon} \right| \nabla V_{\varepsilon} \right\rangle \eta^{2} \phi V_{\varepsilon} \psi'(V_{\varepsilon}) dX \n+ \sum_{j=1}^{n} \iint_{Q_{\rho}} \left\langle \nabla^{2} E_{\varepsilon}(\nabla u_{\varepsilon}) \nabla \partial_{x_{j}} u_{\varepsilon} \right| \nabla \partial_{x_{j}} u_{\varepsilon} \right\rangle \eta^{2} \phi \psi(V_{\varepsilon}) dX \n\leq 2 \iint_{Q_{\rho}} \Psi(V_{\varepsilon}) \eta^{2} \phi_{h} \partial_{t} \phi_{c} dX + 4 \iint_{Q_{\rho}} \left| \left\langle \nabla^{2} E_{\varepsilon}(\nabla u_{\varepsilon}) \nabla V_{\varepsilon} \right| \nabla \eta \right\rangle \left| \eta \phi \psi(V_{\varepsilon}) V_{\varepsilon} dX \right. \n+ \frac{n}{2\lambda} \iint_{Q_{\rho}} |f_{\varepsilon}|^{2} V_{\varepsilon}^{2-p} \left(\psi(V_{\varepsilon}) + V_{\varepsilon} \psi'(V_{\varepsilon}) \right) \eta^{2} \phi dX + 4 \iint_{Q_{\rho}} |f_{\varepsilon}| |\nabla \eta| \psi(V_{\varepsilon}) V_{\varepsilon} \eta \phi dX \n=: 2\mathbf{R}_{1} + 4\mathbf{R}_{2} + \frac{n}{2\lambda} \mathbf{R}_{3} + 4\mathbf{R}_{4}.
$$

We may discard $L_1 \geq 0$. By the Cauchy–Schwarz inequality and Young's inequality, \mathbb{R}_2 and \mathbb{R}_4 are estimated as follows:

$$
\begin{array}{rcl}\n\mathbf{4R}_2 & \leqslant & \mathbf{L}_2 + 4 \iint_{Q_\rho} \left\langle \nabla^2 E_\varepsilon(\nabla u_\varepsilon) \nabla \eta \mid \nabla \eta \right\rangle V_\varepsilon \phi \frac{\psi(V_\varepsilon)^2}{\psi'(V_\varepsilon)} \, \mathrm{d}X, \\
\mathbf{4R}_4 & \leqslant & 2\mathbf{R}_3 + 2 \iint_{Q_\rho} V_\varepsilon^{p-1} |\nabla \eta|^2 \phi \frac{\psi(V_\varepsilon)^2}{\psi'(V_\varepsilon)} \, \mathrm{d}X.\n\end{array}
$$

Also, we note that our choice of ψ yields

$$
\frac{\psi(V_{\varepsilon})^2}{\psi'(V_{\varepsilon})} = \frac{V_{\varepsilon}^{p+1}\tilde{\psi}(V_{\varepsilon})^2}{p\tilde{\psi}(V_{\varepsilon}) + V_{\varepsilon}\tilde{\psi}'(V_{\varepsilon})}, \quad \text{and} \quad \psi(V_{\varepsilon}) + V_{\varepsilon}\psi'(V_{\varepsilon}) = V_{\varepsilon}^p\left((p+1)\tilde{\psi}(V_{\varepsilon}) + V_{\varepsilon}\tilde{\psi}'(V_{\varepsilon})\right).
$$

Letting $\tilde{\varepsilon} \to 0$, recalling our choice of η and ϕ_c , and using [\(2.33\)](#page-12-4)–[\(2.34\)](#page-12-3) and [\(3.3\)](#page-15-2), we have

$$
\iint_{Q_{\sigma\rho}} |\nabla \left[G_{p,\varepsilon}(\nabla u_{\varepsilon}) \right]|^2 \tilde{\psi}(V_{\varepsilon}) dX \le \frac{C}{[(1-\sigma)\rho]^2} \iint_{Q_{\rho}} \left(\frac{\mu^{2p} \tilde{\psi}(V_{\varepsilon})^2}{p \tilde{\psi}(V_{\varepsilon}) + V_{\varepsilon} \tilde{\psi}'(V_{\varepsilon})} + \Psi(V_{\varepsilon}) \right) dX \qquad (3.8)
$$

$$
+ C \mu^{2p} \iint_{Q_{\rho}} |f_{\varepsilon}|^2 \left(\tilde{\psi}(V_{\varepsilon}) + V_{\varepsilon} \tilde{\psi}'(V_{\varepsilon}) \right) dX.
$$

When $\tilde{\psi}(s) \equiv p + 2$, which yields $\Psi(s) = s^{p+2}$, we simply use (2.34) to get $\Psi(V_{\varepsilon}) \leq C_{p,\delta,M}\mu^{2p}$. By (3.8) and Hölder's inequality, we have

$$
\iint_{Q_{\sigma\rho}} |\nabla \left[G_{p,\varepsilon}(\nabla u_{\varepsilon}) \right]|^2 \, \mathrm{d}X \leqslant C\mu^{2p} \left[\frac{|Q_{\rho}|}{[(1-\sigma)\rho]^2} + F^2 |Q_{\rho}|^{1-2/q} \right].
$$

Dividing each side by $|Q_{\sigma\rho}| = \sigma^{n+2} |Q_{\rho}|$, we obtain [\(3.6\)](#page-17-1). When $\tilde{\psi}(s) = (s - \delta - k)_+^2$ with $k :=$ $(1 - 2\nu)\mu > \mu/2$, we note that for every $s \in (0, \mu + \delta)$

$$
\Psi(s) = \int_0^s ((\sigma - \delta - k) + (\delta + k))^{p+1} (\sigma - \delta - k)_+^2 d\sigma \leq C(p) \nu^3 \mu^{p+4},
$$

which yields $\Psi(V_{\varepsilon}) \leq C_{p,\delta,M} \nu^3 \mu^{2p+2}$ by [\(2.34\)](#page-12-3). We also note that $k \in (\mu/2, \mu)$ yields

$$
\frac{\tilde{\psi}(V_{\varepsilon})^2}{p\tilde{\psi}(V_{\varepsilon})+V_{\varepsilon}\tilde{\psi}'(V_{\varepsilon})} = \frac{(V_{\varepsilon}-\delta-k)^3_+}{p(V_{\varepsilon}-\delta-k)_++2V_{\varepsilon}} \leq \frac{(2\nu\mu)^3}{2(\delta+k)} \leq 8\nu^3\mu^2,
$$

$$
\tilde{\psi}(V_{\varepsilon})+V_{\varepsilon}\tilde{\psi}'(V_{\varepsilon}) = (V_{\varepsilon}-\delta-k)_+(3V_{\varepsilon}-\delta-k) \leq 2\nu\mu \cdot 2(\delta+\mu+\nu\mu).
$$

Combining these inequalities with [\(2.34\)](#page-12-3), and $\tilde{\psi}(V_{\varepsilon}) \geqslant (\nu \mu)^2$ on $S_{\sigma\rho,\,\mu,\,\nu} \subset Q_{\sigma\rho}$, we get

$$
(\nu\mu)^2 \iint_{S_{\sigma\rho,\,\mu,\,\nu}} |\nabla \left[G_{p,\,\varepsilon}(\nabla u_\varepsilon)\right]|^2 \, \mathrm{d}X \leqslant C\mu^{2p+2} \left[\frac{\nu^3 |Q_\rho|}{[(1-\sigma)\rho]^2} + \nu F^2 |Q_\rho|^{1-2/q} \right],
$$

from [\(3.8\)](#page-17-3). This completes the proof of [\(3.7\)](#page-17-2).

4 Local gradient bounds

In Section [4,](#page-18-0) we prove local uniform bounds of ∇u_{ε} .

4.1 Preliminary

For p-harmonic flows with $\frac{n+2}{2n} < p < \infty$, local gradient bounds are proved in [\[16\]](#page-32-8) by De Giorgi's truncation (see also [\[13,](#page-32-2) Chapter VIII]). Another proof that is fully based on Moser's iteration is given by [\[7,](#page-32-1) §4]. However, the method in [\[7,](#page-32-1) §4] will not work in our problem, since the test functions therein may intersect with a facet of a solution. Here we note that our problem can be seen as uniformly parabolic when a spatial gradient does not vanish. With this in mind, we carefully choose test functions supported in the place $D_k := \{(x, t) \in Q_R \mid |\nabla u_{\varepsilon}(x, t)| > k\}$ for some constant $k \geq 1$. This type of strategy can be found in various elliptic problems whose uniform ellipticity may break on some degenerate regions (see e.g., [\[4\]](#page-31-5), [\[8\]](#page-32-14), [\[24\]](#page-32-15), [\[43\]](#page-33-10)). Among these papers, our proof, based on Moser's iteration, is substantially a modification of [\[43\]](#page-33-10).

To clarify our strategy, we introduce another truncation of a partial derivative $\partial_{x_i}u_{\varepsilon}$. The truncation function g_k is given by $g_k(\sigma) := (\sigma - k)_+ - (-\sigma - k)_+$ for $\sigma \in \mathbb{R}$, and the corresponding truncated partial derivative is of the form $v_{j,k} := g_k(\partial_{x_j}u_{\varepsilon})$ for each $j \in \{1, \ldots, n\}$, where the constant $k \geq 1$ will be determined by $||f_{\varepsilon}||_{L^q(Q_R)}$. We do not directly show reversed Hölder estimates for V_{ε} . Instead, we consider another scalar function $W_k := \sqrt{k^2 + \sum_{j=1}^n v_{j,k}^2}$. In Section [4,](#page-18-0) we mainly aim to deduce local L^{∞} -bounds of W_k . Here we note

$$
V_{\varepsilon} \leqslant c_n W_k \quad \text{in} \quad Q = Q_R, \quad \text{and} \quad W_k \leqslant \sqrt{2} V_{\varepsilon} \quad \text{in} \quad D_k \subset Q_R. \tag{4.1}
$$

since we have let $k \geq 1$ (see [\[43,](#page-33-10) §4.1]). In particular, we are allowed to use

$$
\tilde{\lambda} W_k^{p-2} \mathrm{id}_n \leqslant \nabla^2 E_\varepsilon(\nabla u_\varepsilon) \leqslant \tilde{\Lambda} W_k^{p-2} \mathrm{id}_n \quad \text{in } D_k,\tag{4.2}
$$

where the constants $0 < \tilde{\lambda} < \tilde{\Lambda} < \infty$ depend at most on n, p, λ , Λ , K. With this in mind, we can deduce a weak formulation of W_k , similarly to Lemma [3.1.](#page-15-4) We consider the composite functions ψ

 \Box

and Ψ given in Lemma [3.1,](#page-15-4) and fix an arbitrary non-negative Lipschitz function $\zeta = \zeta(x, t)$ that is compactly supported in the interior of Q . Then, W_k satisfies

$$
2\tilde{J}_0 + 2\tilde{J}_1 + \tilde{J}_2 + \tilde{J}_3 \le \tilde{\lambda}^{-1} n \tilde{J}_4 + 2J_5,
$$
\n(4.3)

where for each $l \in \{0, \ldots, 6\}$, \tilde{J}_l is similarly defined as in Lemma [3.1](#page-15-4) with V_{ε} and $\nabla \partial_{x_j} u_{\varepsilon}$ replaced by W_k and $\nabla v_{j,k}$ respectively. The inequality [\(4.3\)](#page-19-0) is deduced by testing $\phi := \zeta \psi(W_k)v_{j,k}$ into [\(2.27\)](#page-11-5). Here it should be mentioned that this test function is supported in $Q \cap \{\left|\partial_x u_{\varepsilon}\right| > k\} \subset D_k$. Therefore we may replace $\nabla \partial_{x_i} u_{\varepsilon}$ and $\partial_t \partial_{x_i} u_{\varepsilon}$ by $\nabla v_{j,k}$ and $\partial_t v_{j,k}$ respectively, and make use of [\(4.2\)](#page-18-2). Summing over $j \in \{1, \ldots, n\}$, we can obtain [\(4.3\)](#page-19-0), similarly to [\(3.2\)](#page-15-3).

Lemma 4.1. Let u_{ε} be a weak solution to [\(2.25\)](#page-11-3) in $Q_R = Q_R(x_*, t_*) \in \Omega_T$ with $\varepsilon \in (0, 1)$ and [\(2.26\)](#page-11-4). For $Q_r = Q_r(x_*, t_*) \subset Q_R$, fix $\eta \in C_c^1(B_r; [0, 1])$, $\phi_c \in C^1(\overline{I_r}; [0, 1])$ satisfying $\phi_c(t_0 - r^2) = 0$. Then, there exists a constant $C \in (0, \infty)$, depending at most on n, p, λ , Λ and K, such that the following estimate holds for all $\alpha \in [0, \infty)$.

$$
(1+\alpha)^{-1} \left[\sup_{\tau \in \mathring{I}_r} \int_{B_r \times \{\tau\}} W_k^{2(1+\alpha)} \eta^2 \phi_c \, dx + \iint_{Q_r} \left| \nabla W_k^{p/2+\alpha} \right|^2 \eta^2 \phi_c \, dx \right] \tag{4.4}
$$
\n
$$
\leq C \left[\iint_{Q_r} \left(W_k^{p+2\alpha} |\nabla \eta|^2 + W_k^{2(1+\alpha)} |\partial_t \phi_c| \right) \, dx + (1+\alpha) \iint_{Q_r} |f_\varepsilon|^2 W_k^{2+2\alpha-p} \eta^2 \phi_c \, dx \right].
$$

Proof. Let $\phi_h: I_r \to [0, 1]$ be an arbitrary Lipschitz function that is non-increasing and $\phi_h(t_0) = 0$, and we set $\phi := \phi_c \phi_h$. We test $\zeta := \eta^2 \phi$ into [\(4.3\)](#page-19-0) with $\psi(s) := s^{2\alpha}$, and therefore $\Psi(s) = [2(1 + \eta^2 s^2)]$ α)]⁻¹s^{2(1+ α}). Then, we note that [\(4.2\)](#page-18-2) yields

$$
\tilde{J}_2 + \tilde{J}_3 \ge (1 + 2\alpha)\tilde{\lambda} \iint_{Q_r} |\nabla W_k|^2 W_k^{p+2\alpha - 2} \eta^2 \phi \, \mathrm{d}X = \frac{4(1 + 2\alpha)}{(p + 2\alpha)^2} \tilde{\lambda} \iint_{Q_r} \left| \nabla W_k^{p/2 + \alpha} \right|^2 \eta^2 \phi \, \mathrm{d}X,
$$

where we have used $|\nabla W_k|^2 \leq \sum_{j=1}^n |\nabla v_{j,k}|^2$. By [\(4.2\)](#page-18-2) and Young's inequality, we have

$$
\begin{aligned} 2|\tilde{J}_1|+n\tilde{\lambda}^{-1}|\tilde{J}_4|+2|\tilde{J}_5|\leqslant&\frac{(1+2\alpha)\tilde{\lambda}}{2}\iint_{Q_r}|\nabla W_k|^2W_k^{p+2\alpha-2}\eta^2\phi\,\mathrm{d} X+C\iint_{Q_r}W_k^{p+2\alpha}|\nabla \eta|^2\phi\,\mathrm{d} X\\ &+C(1+2\alpha)\iint_{Q_r}|f_\varepsilon|^2W_k^{2+2\alpha-p}\eta^2\phi\,\mathrm{d} X. \end{aligned}
$$

Hence, we obtain

$$
\begin{split} &(1+\alpha)^{-1}\left[-\iint_{Q_r}W_k^{2+2\alpha}\eta^2\phi_c\partial_t\phi_\mathrm{h}\,\mathrm{d} X+\iint_{Q_r}\left|\nabla W_k^{p/2+\alpha}\right|^2\eta^2\phi_c\phi_\mathrm{h}\,\mathrm{d} X\right]\\ &\leqslant C\left[\iint_{Q_r}\left(W_k^{p+2\alpha}|\nabla \eta|^2+W_k^{2(1+\alpha)}|\partial_t\phi_c|\right)\,\mathrm{d} X+(1+\alpha)\iint_{Q_r}|f_\varepsilon|^2W_k^{2+2\alpha-p}\eta^2\phi_c\phi_\mathrm{h}\,\mathrm{d} X\right], \end{split}
$$

where $C \in (0, \infty)$ depends at most on p, $\tilde{\lambda}$, and $\tilde{\Lambda}$. From this estimate, we deduce [\(4.4\)](#page-19-1) by suitably choosing ϕ_{h} , similarly to the proof of Lemma 3.2. choosing ϕ_h , similarly to the proof of Lemma [3.2.](#page-16-1)

We infer an iteration lemma, often used implicitly in Moser's iteration.

Lemma 4.2. Fix the constants $A, B, \kappa \in (1, \infty)$ and $\mu \in (0, \infty)$. Let the sequences $\{Y_i\}_{i=0}^{\infty} \subset [0, \infty)$ and $\{p_l\}_{l=0}^{\infty} \subset (0, \infty)$ satisfy $Y_{l+1}^{p_{l+1}} \leq (AB^l Y_l^{p_l})^{\kappa}$, and $p_l \geq \mu$ (κ^l-1) for all $l \in \mathbb{Z}_{\geq 0}$, and $\kappa^l \overline{p_l}^{-1} \to \mu^{-1}$ as $l \to \infty$. Then, we have $\limsup_{l \to \infty} Y_l \leq A^{\frac{\kappa'}{\mu}} B^{\frac{(\kappa')^2}{\mu}}$ $\overline{H} Y$ $\frac{p_0}{\mu}$.

Proof. By induction, it is easy to check that $Y_l^{p_l} \n\t\leq \prod_{k=1}^l A^{\kappa^{l-k+1}} \prod_{k=1}^l B^{k\kappa^{l-k+1}} Y_0^{\kappa^l p_0}$ holds for all $l \in \mathbb{N}$. By [\[6,](#page-32-16) Lemma 2.3] and the assumptions in Lemma [4.2,](#page-19-2) we obtain

$$
Y_l \leqslant \prod_{k=1}^l A^{\frac{\kappa^{l-k+1}}{\mu(\kappa^{l-1})}} \prod_{k=1}^l B^{\frac{k\kappa^{l-k+1}}{\mu(\kappa^{l-1})}} Y_0^{\frac{\kappa^l p_0}{p_l}} \leqslant A^{\frac{\kappa'}{\mu}} B^{\frac{(\kappa')^2}{\mu}} Y_0^{\frac{\kappa^l p_0}{p_l}}.
$$

Letting $l \to \infty$ completes the proof.

 \Box

4.2 Moser's iteration

We give the proof of Theorem [2.7](#page-12-0) by Moser's iteration.

Proof of Theorem [2.7.](#page-12-0) We define $\kappa \in (1, 2)$ to be $\kappa := 1 + 2/n$ for $n \ge 3$, and $\kappa := \sigma'$ for $n \ge 2$. We also set $\tilde{\kappa} := 1 + 2/n - \kappa \in [0, 1)$. By our choice of σ and q, we have $q/(q - 2) < \kappa$. We define $\pi := \max\{p-1, p/2\} > 0$, and $k := 1 + ||f_{\varepsilon}||_{L^q(Q_R)}^{1/\pi} \ge 1$. Then, the non-negative function $\tilde{f}_{\varepsilon} := W_k^{-2\pi} |f_{\varepsilon}|^2 \in L^{q/2}(Q_R)$ satisfies $\|\tilde{f}_{\varepsilon}\|_{L^q(Q_R)} \leq 1$. The proof is completed by showing

$$
\sup_{Q_{\theta R}} W_k \leqslant \frac{C}{(1-\theta)^{d_2}} \left[\iint_{Q_R} W_k^p \, dX \right]^{d/p} \tag{4.5}
$$

for some $C = C(n, p, q, \lambda, \Lambda, K) \in (1, \infty)$. Indeed, [\(4.1\)](#page-18-3), [\(4.5\)](#page-20-1) and $W_k \le k + V_{\varepsilon}$ yield

$$
\sup_{Q_{\theta R}} V_{\varepsilon} \leq C_n \sup_{Q_{\theta R}} W_k \leq \frac{C}{(1-\theta)^{d_2}} \left(k^p + \iint_{Q_R} V_{\varepsilon}^p \, \mathrm{d}X \right)^{d/p}.
$$

Recalling the definition of k , we conclude (2.28) .

To prove [\(4.5\)](#page-20-1), we deduce reversed Hölder inequalities. For $\alpha \in [0, \infty)$, we define $m(\alpha) := \frac{4+4\alpha}{p+2\alpha}$ $\frac{1}{p+2\alpha} \in$ $[2, 4/p] \cup [4/p, 2]$, and $\tilde{m}(\alpha) := \min\{2, m(\alpha)\} \le 2$. Let the functions $\eta = \eta(x)$ and $\phi_c = \phi_c(t)$ satisfy all the assumptions in Lemma [4.1.](#page-19-3) Then, [\(4.4\)](#page-19-1) becomes

$$
\sup_{\tau \in \mathring{I}_r} \int_{B_r \times \{\tau\}} v_\alpha^m \eta^2 \phi_c \, dx + \iint_{Q_r} |\nabla v_\alpha|^2 \eta^2 \phi_c \, dX \tag{4.6}
$$
\n
$$
\leq C(1+\alpha) \left[\iint_{Q_r} (v_\alpha^2 |\nabla \eta|^2 + v_\alpha^m |\partial_t \phi_c|) \, dX + (1+\alpha) \iint_{Q_r} \tilde{f}_\varepsilon v_\alpha^m \eta^2 \phi_c \, dX \right],
$$

where $v_{\alpha} := W_k^{p/2+\alpha}$. From [\(4.6\)](#page-20-2), we would like to find a constant $C \in (1, \infty)$, which depends at most on $n, \kappa, q, \lambda, \Lambda, \Lambda$, and K, such that

$$
\iint_{Q_r} v_\alpha^{2+(\kappa-1)m} \left(\eta^2 \phi_c\right)^\kappa dx \leq C r^{2\tilde{\kappa}} (1+\alpha)^{\kappa \gamma} \left(\iint_{Q_r} \left(v_\alpha^2 (|\nabla \eta|^2 + 1) + v_\alpha^m |\partial_t \phi|\right) dX\right)^\kappa \tag{4.7}
$$

for some constant $\gamma = \gamma(q) \in [2, \infty)$. To obtain [\(4.7\)](#page-20-3), we use the parabolic Poincaré–Sobolev embedding (see also [\[13,](#page-32-2) Chapter I, Proposition 3.1]), given by

$$
\iint_{Q_r} v_\alpha^{2+(\kappa-1)m} \left(\eta^2 \phi_c\right)^\kappa dx \leq C r^{2\tilde{\kappa}} \left(\iint_{Q_r} \left| \nabla \left(v_\alpha \eta \phi_c^{1/2} \right) \right|^2 dx \right) \left(\sup_{\tau \in \hat{I}_r} \int_{B_r \times \{\tau\}} v_\alpha^m \eta^2 \phi_c dx \right)^{\kappa-1}
$$

for some $C = C(\kappa, q) \in (0, \infty)$, independent of m. This is easily deduced by applying Hölder's inequality to the functions $(v_{\alpha}\eta\phi_c^{1/2})^2 \in L^{\frac{1}{2-\kappa}}(Q_r)$, $(v_{\alpha}^m\eta^2\phi_c)^{\kappa-1} \in L^{\frac{1}{\kappa-1}}(Q_r)$, and the Sobolev embedding to $(v_{\alpha}\eta\phi_c^{1/2})(\cdot, t) \in W_0^{1, 2}$ $L_0^{1,2}(B_r) \hookrightarrow L^{\frac{2}{2-\kappa}}(B_r)$. Combining with [\(4.6\)](#page-20-2), we have

$$
\iint_{Q_r} v_{\alpha}^{2+(\kappa-1)m} \left(\eta^2 \phi_c\right)^{\kappa} dx \leq C(1+\alpha)^{\kappa} r^{2\tilde{\kappa}} \left(\iint_{Q_r} \left(v_{\alpha}^2 |\nabla \eta|^2 + v_{\alpha}^m |\partial_t \phi_c|\right) dX \right)^{\kappa} + C(1+\alpha)^{2\kappa} r^{2\tilde{\kappa}} \left(\iint_{Q_r} \left(v_{\alpha}^{\tilde{m}} \eta^2 \phi_c\right)^{\frac{q}{q-2}} dX \right)^{\kappa(1-2/q)}
$$

.

Here we have used $\|\tilde{f}_{\varepsilon}\|_{L^{q/2}(Q_R)} \leq 1$. If $q = \infty$, then the last estimate obviously yields [\(4.7\)](#page-20-3) with $\gamma := 2$. Otherwise, we interpolate the function $v_{\alpha}^{m} \eta^2 \phi_c$ among the Lebesgue spaces $L^1(Q_r) \subset L^{\frac{q}{q-2}}(Q_r)$ $L^{\kappa}(Q_r)$. Combining with Young's inequality, we obtain

$$
\left(\iint_{Q_r} \left(v_\alpha^{\tilde{m}} \eta^2 \phi_c\right)^{\frac{q}{q-2}} dX\right)^{\kappa\left(1-\frac{2}{q}\right)} \leqslant \left(\iint_{Q_r} v_\alpha^{\tilde{m}} \eta^2 \phi_c dX\right)^{\kappa'\cdot \frac{\kappa q-q-2\kappa}{q}} \left(\iint_{Q_r} \left(v_\alpha^{\tilde{m}} \eta^2 \phi_c\right)^{\kappa} dX\right)^{\kappa'\cdot \frac{2}{q}}
$$

$$
\leq \sigma^{-\frac{2\kappa'}{q-2\kappa'}} \left(\iint_{Q_r} v_\alpha^2 \eta^2 \phi_c \,dX \right)^{\kappa} + \sigma \iint_{Q_r} v_\alpha^{2+(\kappa-1)m} \left(\eta^2 \phi_c \right)^{\kappa} dX
$$

 \Box

for all $\sigma \in (0, \infty)$. Here the last inequality follows from $\tilde{m} \leq 2$, $\tilde{m}\kappa \leq 2 + (\kappa - 1)m$, and $v_{\alpha} \geq 1$, which are obvious by the definitions of k and \tilde{m} . We conclude [\(4.7\)](#page-20-3) with $\gamma := \frac{2q}{q-2\kappa'} \in (2, \infty)$.

Before showing [\(4.5\)](#page-20-1) from [\(4.7\)](#page-20-3), we fix some abbreviations. Let $\theta \in (0, 1)$ be given. For every $l \in \mathbb{Z}_{\geqslant 0}$, we set $r_l := \theta R + 2^{-l}(1-\theta)R$, and define $B_l := B_{r_l}(x_0)$, $I_l := I_{r_l}(t_0)$ and $Q_l := B_l \times I_l =$ $Q_{r_l}(x_0, t_0)$. We set the sequences $\{\alpha_l\}_{l=0}^{\infty} \subset [0, \infty)$ and $\{p_l\}_{l=0}^{\infty} \subset [2, \infty)$ as follows. For $p \ge 2$, we define $\alpha_l := \kappa^l - 1 \in [0, \infty)$ and $p_l := p + 2\alpha_l \in [p, \infty)$ for each $l \in \mathbb{Z}_{\geq 0}$. When $2n/(n+2) < p < 2$, we define $\alpha_l \coloneqq \frac{\lambda}{4}$ $\frac{\lambda}{4}$ $(\kappa^l - 1) \in [0, \infty)$ and $p_l := 2 + 2\alpha_l \in [2, \infty)$ for each $l \in \mathbb{Z}_{\geqslant 0}$. Here we choose a constant $\lambda := n(p-2) + 4 \in \left[8/(n+2), 4 \right]$ for $p \in \left(\frac{2n}{n+2}, 2 \right)$. By Moser's iteration, we prove the boundedness of the non-negative sequence ${Y_l}_{l=0}^{\infty}$ defined as $Y_l := (\iint_{Q_l} W_k^{p_l} dX)^{1/p_l}$ for $l \in \mathbb{Z}_{\geqslant 0}$. Moreover, we aim to give bound estimates for $Y_{\infty} := \limsup_{l \to \infty} Y_l = \sup_{Q_{\theta R}} W_k$. By [\(4.7\)](#page-20-3) with $\eta \in C_c^1(B_{r_l}; [0, 1])$ and $\phi_c \in C^1(\overline{I_{r_i}}; [0, 1])$ chosen suitably, we have

$$
\iint_{Q_{l+1}} W_k^{p_{l+1}}\,\mathrm{d} X \leqslant \frac{C (4\kappa^\gamma)^{l\kappa} R^{2\tilde\kappa}}{[(1-\theta)R]^{2\kappa}} \left(\iint_{Q_l} W_k^{p_l}\,\mathrm{d} X\right)^\kappa
$$

for every $l \in \mathbb{Z}_{\geqslant 0}$. In other words, there holds $Y_{l+1}^{p_{l+1}} \leqslant (AB^l Y_l^{p_l})^{\kappa}$ for every $l \in \mathbb{Z}_{\geqslant 0}$, where $A =$ $(1 - \theta)^{-2}C$ and $B := 4\kappa^{\kappa\gamma}$ for some sufficiently large $C = C(n, p, q, \lambda, \Lambda, K) \in (1, \infty)$. For $p \geq 2$, applying Lemma [4.2](#page-19-2) with $\mu := 2$ yields $Y_{\infty} \leq A^{\frac{\kappa'}{2}} B^{\frac{(\kappa')^2}{2}} Y_0^{\frac{p}{2}}$, which implies [\(4.5\)](#page-20-1). For the remaining case $\frac{2n}{n+2} < p < 2$, we apply Lemma [4.2](#page-19-2) with $\mu := \lambda/2$. By the resulting inequality $Y_{\infty} \leq A^{\frac{2\kappa'}{\lambda}} B^{\frac{2(\kappa')^2}{\lambda}} Y_0^{\frac{4}{\lambda}}$ and Young's inequality, we have

$$
\sup_{Q_{\theta R}} W_k \le \left[\frac{C}{(1-\theta)^{2\kappa'}} \iint_{Q_R} W_k^2 \, \mathrm{d}X \right]^{\frac{2}{\lambda}} \le \frac{1}{2} \sup_{Q_R} W_k + \left[\frac{C}{(1-\theta)^{2\kappa'}} \iint_{Q_R} W_k^p \, \mathrm{d}X \right]^{\frac{1}{p}}
$$

for all $\theta \in (0, 1)$. The desired estimate [\(4.5\)](#page-20-1) follows from [\[26,](#page-33-13) Chapter V, Lemma 3.1].

5 Degenerate case

In Section [5,](#page-21-0) we aim to prove Proposition [2.9](#page-13-0) from the estimates (3.4) – (3.5) . The former (3.4) is used to deduce oscillation lemmata (Lemmata [5.1–](#page-21-2)[5.2\)](#page-22-0). There we make use of a result of the expansion of posivity (Lemma [5.3\)](#page-23-1), which is verified by the latter [\(3.5\)](#page-16-3). As related items, see [\[33,](#page-33-11) Chapters II–III], [\[19,](#page-32-10) Chapters 3–4].

5.1 Levelset estimates

Firstly, we prove various estimates for the level sets of $U_{\delta,\epsilon}$. For given $\tau_0 \in (t_0 - 4\rho^2, t_0)$ and $\gamma \in (0, 1)$, we define $I_{\frac{3}{2}\rho}(\gamma; \tau_0) := (\tau_0, \tau_0 + \gamma(\frac{3}{2})$ $(\frac{3}{2}\rho)^2$] $\subset \tilde{I}_{\frac{3}{2}\rho}(\gamma; \tau_0) := (\tau_0 - \gamma(\frac{3}{2})$ $(\frac{3}{2}\rho)^2$, $\tau_0 + \gamma(\frac{3}{2})$ $(\frac{3}{2}\rho)^2$, and $Q_{\frac{3}{2}\rho}(\gamma; \tau_0) :=$ $B_{\frac{3}{2}\rho}(x_0) \times I_{\frac{3}{2}\rho}(\gamma; \tau_0).$

Lemma 5.1. In addition to the assumptions of Proposition [2.9,](#page-13-0) let

$$
I_{\frac{3}{2}\rho}(\gamma;\,\tau_0) \subset \tilde{I}_{\frac{3}{2}\rho}(\gamma;\,\tau_0) \subset (t_0 - 4\rho^2,\,t_0],\tag{5.1}
$$

$$
\left| \left\{ x \in B_{\rho} \mid U_{\delta,\varepsilon}(x,\,t) \leqslant (1-\hat{\nu})\mu^2 \right\} \right| \geqslant \tilde{\nu}|B_{\rho}| \quad \text{for all } t \in I_{\frac{3}{2}\rho}(\gamma;\,\tau_0),\tag{5.2}
$$

and $\rho^{\beta} \leq 2^{-i_{\star}} \hat{\nu}$ hold for some $i_{\star} \in \mathbb{N}$. Then, we have

$$
\left| Q_{\frac{3}{2}\rho}(\gamma; \tau_0) \cap \left\{ U_{\delta, \varepsilon} \geq (1 - 2^{-i_{\star}} \hat{\nu}) \mu^2 \right\} \right| \leq \frac{C_{\dagger}}{\tilde{\nu}\sqrt{\gamma i_{\star}}} \left| Q_{\frac{3}{2}\rho}(\gamma; \tau_0) \right|, \tag{5.3}
$$

for some $C_{\dagger} \in (1, \infty)$ depending at most on n, p, q, λ , Λ , K , F , δ , and M.

We invoke an isoperimetric inequality for the functions in $W^{1,1}(B_\rho)$ (see e.g., [\[15,](#page-32-17) Chapter 10, §5.1]); for any fixed numbers $-\infty < k < l < \infty$ and $v \in W^{1, 1}(B_{\rho})$, there holds

$$
(l-k)|\{x \in B_{\rho} \mid v(x) > l\}| \leq \frac{C_n \rho^{n+1}}{|\{x \in B_{\rho} \mid v(x) < k\}|} \int_{B_{\rho} \cap \{k < v < l\}} v \, \mathrm{d}x. \tag{5.4}
$$

Proof. We set $k_i := (1 - 2^{-i}\hat{\nu})\mu^2$ and $A_i := Q \cap \{U_{\delta, \varepsilon} > k_i\}$ for each $i \in \mathbb{Z}_{\geq 0}$, where $Q := Q_{\frac{3}{2}\rho}(\gamma; \tau_0)$. By the definition of k_i , $k_{i+1} - k_i = 2^{-i-1}\hat{\nu}\mu^2$ is clear for every $i \in \mathbb{Z}_{\geqslant 0}$. Note that (5.2) enables us to apply [\(5.4\)](#page-22-1) to the sliced function $U_{\delta,\varepsilon}(\cdot, t)$. Hence, we obtain

$$
\frac{\hat{\nu}\mu^2}{2^{i+1}} \left| \left\{ x \in B_{\frac{3}{2}\rho} \; \middle| \; U_{\delta,\varepsilon}(x,\,t) > k_{i+1} \right\} \right| \leqslant \frac{C(n)\rho}{\tilde{\nu}} \int_{B_{\frac{3}{2}\rho} \cap \{k_i < U_{\delta,\varepsilon}(\,\cdot\,,t) < k_{i+1}\}} |\nabla (U_{\delta,\varepsilon}(x,\,t) - k_i)_+| \, \mathrm{d}x
$$

for all $t \in I_{\frac{3}{2}\rho}(\gamma; \tau_0)$, and $i \in \mathbb{Z}_{\geq 0}$. Integrating with respect to $t \in I_{\frac{3}{2}\rho}(\gamma; \tau_0)$ and applying the Cauchy–Schwarz inequality, we get

$$
\frac{\hat{\nu}\mu^2}{2^{i+1}}|A_{i+1}| \leq \frac{C(n)\rho}{\tilde{\nu}} \left(\iint_Q |\nabla (U_{\delta,\varepsilon} - k_i)_+|^2 \, dX \right)^{1/2} (|A_i| - |A_{i+1}|)^{1/2}
$$

for every $i \in \mathbb{Z}_{\geqslant 0}$. To estimate the integration on the right-hand side, we make use of [\(3.4\)](#page-16-2) with $Q_0 := B_{2\rho} \times \tilde{I}_{\frac{3}{2}\rho}(\gamma; \tau_0)$ and $k := k_i$. Choosing suitable cut-off functions $\eta \in C_c^1(B_{2\rho}; [0, 1])$ and $\phi_c \in C^1([t_0 - \gamma(3\rho/2)^2, t_0 + \gamma(3\rho/2)^2]; [0, 1]),$ and noting $(U_{\delta, \varepsilon} - k_i)_+ \leq 2^{-i\hat{\nu}}\mu^2$, we compute

$$
\iint_Q |\nabla (U_{\delta, \varepsilon} - k_i)_+|^2 dX \le C \left[\frac{4^{-i} \hat{\nu} \mu^4}{\gamma \rho^2} |Q_0| + \mu^4 F^2 |Q_0|^{1-2/q} \right] \le \frac{C (\hat{\nu} \mu^2)^2}{4^i \gamma \rho^2} \left[1 + \left(\frac{2^i \rho^\beta}{\hat{\nu}} \right)^2 \right] |Q|
$$

for some constant $C \in (1, \infty)$ depending at most on n, p, q, λ , Λ , K , δ and M. The proof is completed by showing [\(5.3\)](#page-21-4), provided $2^{i*}\rho^{\beta} \leq \hat{\nu}$ and [\(5.2\)](#page-21-3). If $2^{i*}\rho^{\beta} \leq \hat{\nu}$ holds for some fixed $i_{\star} \in \mathbb{N}$, then we can find a constant $C_{\dagger} \in (1, \infty)$ satisfying

$$
|A_{i+1}|^2 \leqslant \frac{C_{\dagger}^2}{\tilde{\nu}^2 \gamma} |Q|(|A_i| - |A_{i+1}|)
$$

for each $i \in \{0, 1, \ldots, i_{\star} - 1\}$. From this, we obtain

$$
i_{\star}|A_{i_{\star}}|^{2} \leq \sum_{i=0}^{i_{\star}-1} |A_{i+1}|^{2} \leq \frac{C_{\dagger}^{2}}{\tilde{\nu}^{2}\gamma}|Q| \cdot |A_{0}| \leq \frac{C_{\dagger}^{2}}{\tilde{\nu}^{2}\gamma}|Q|^{2}, \quad \text{whence} \quad |A_{i_{\star}}| \leq \frac{C_{\dagger}}{\tilde{\nu}\sqrt{\gamma i_{\star}}}|Q|.
$$

Lemma 5.2. Under the assumptions of Proposition [2.9,](#page-13-0) there exists $c_* \in (0, 1)$, depending at most on n, p, q, λ , Λ , K , F , δ , and M , such that

$$
\left| Q_{\frac{3}{2}\rho}(\gamma; \tau_0) \cap \left\{ U_{\delta, \varepsilon} \ge (1 - \nu_0) \mu^2 \right\} \right| \le \alpha_\star \left| Q_{\frac{3}{2}\rho}(\gamma; \tau_0) \right| \quad \text{with} \quad \alpha_\star(\gamma) := c_\star \gamma^{\frac{(n+2)}{4\beta}} \tag{5.5}
$$

and $\rho < \nu_0$ imply

$$
\sup_{\tilde{Q}(\gamma;\,\tau_0)} U_{\delta,\,\varepsilon} \leqslant \left(1 - \frac{\nu_0}{2}\right) \mu^2, \quad \text{where} \quad \tilde{Q}(\gamma;\,\tau_0) := B_\rho \times \left(\tau_0 + \frac{5\gamma}{4}\rho^2,\,\tau_0 + \frac{9\gamma}{4}\rho^2\right). \tag{5.6}
$$

Before the proof, we invoke the parabolic Poincaré–Sobolev inequality

$$
\iint_{B\times I} |v|^{2+\frac{4}{n}} \, \mathrm{d}X \le C(n) \left(\underset{\tau \in I}{\mathrm{ess sup}} \int_{B\times\{\tau\}} |v(x,\,\tau)|^2 \, \mathrm{d}x \right)^{2/n} \iint_{B\times I} |\nabla v|^2 \, \mathrm{d}X \tag{5.7}
$$

for all $v \in L^2(I; W_0^{1, 2})$ $L^{1,2}(B)$ \cap $L^{\infty}(I; L^{2}(B))$, where $I \subset \mathbb{R}$ is a bounded open interval, and $B \subset \mathbb{R}^{n}$ is a bounded open ball. (see [\[13,](#page-32-2) Chapter I, Proposition 3.1]).

Proof. We set $\rho_l := (1 + 2^{-l-1})\rho \in (\rho, 2\rho), \tau_l := \tau_0 + \frac{9}{4}$ *Proof.* We set $\rho_l := (1 + 2^{-l-1})\rho \in (\rho, 2\rho), \ \tau_l := \tau_0 + \frac{9}{4}\gamma\rho^2 - \gamma\rho_l^2 \in (-4\rho^2, 0), \ B_l := B_{\rho_l} \subset B_{2\rho}, \ I_l := (\tau_l, \ \tau_0 + \frac{9}{4}\gamma\rho^2] \subset (t_0 - 4\rho^2, \ t_0]$ and $Q_l := B_l \times I_l \subset Q_{2\rho}$ for each $l \in \mathbb{Z}_{\geq 0}$. We also define a s $\left[\frac{9}{4}\gamma\rho^2\right] \subset (t_0 - 4\rho^2, t_0]$ and $Q_l := B_l \times I_l \subset Q_{2\rho}$ for each $l \in \mathbb{Z}_{\geqslant 0}$. We also define a superlevel set $A_l := Q_l \cap \{U_{\delta,\varepsilon} > k_l\}$, where $k_l := (1 - (2^{-1} + 2^{-l-1})\nu_0)\mu^2$ and a ratio $Y_l := |A_l|/|B_l| \in (0, 1].$ Applying Hölder's inequality and [\(5.7\)](#page-22-2), and suitably choosing ϕ_c and η in [\(3.4\)](#page-16-2), we have

$$
\frac{\nu_0 \mu^2}{2^{l+3}} |A_{l+1}| \leqslant \frac{C \nu_0 \mu^2 \cdot 2^l}{\sqrt{\gamma}} \left[\frac{|Q_l|^{1/q}}{\rho} + \frac{1}{\nu_0} \right] |A_l|^{1 + \frac{\beta}{n+2}}
$$

for some $C \in (1, \infty)$ depending at most on n, p, q, λ , Λ , K , F , δ , and M. Noting $|Q_l|/|Q_{l+1}| \leq$ $(\rho_l / \rho_{l+1}) = [1 + (1 + 2^{l+2})^{-1}]^{n+2} \leq (6/5)^{n+2}$, we have

$$
Y_{l+1} \leqslant \left(\frac{6}{5}\right)^{n+2} \frac{|A_{l+1}|}{|Q_{l}|^{1+\frac{\beta}{n+2}}} \cdot |Q_{l}|^{\frac{\beta}{n+2}} \leqslant \frac{\tilde{C}_{*} \cdot 4^{l}}{2\sqrt{\gamma}} \left[1+\frac{\rho}{\nu_{0}}\right] Y_{l}^{1+\frac{2\beta}{n+2}}
$$

for every $l \in \mathbb{Z}_{\geqslant 0}$. Here \tilde{C}_* depends at most on n, p, q, λ , Λ , K , F , δ , and M. We set $c_* :=$ $\tilde{C}_{*}^{-\frac{n+2}{2\beta}} 4^{-\frac{(n+2)^2}{4\beta^2}}$ $\frac{4\beta^2}{\beta^2}$. Then, if $\rho < \nu_0$ and (5.5) hold, then $Y_0 \le \alpha_0$ and $[15, \text{ Chapter 9}, \text{ Lemma 15.1}]$ imply that $Y_l \to 0$ as $l \to \infty$, which completes the proof.

Finally, we prove Lemma [5.3,](#page-23-1) which is often called the expansion of positivity.

Lemma 5.3. In addition to the assumptions of Proposition [2.9,](#page-13-0) let $U_{\delta,\varepsilon}$ satisfy

$$
\left| \left\{ x \in B_{\rho} \mid U_{\delta, \varepsilon}(x, \tau_0) \le (1 - \nu) \mu^2 \right\} \right| \ge \frac{\nu}{2} |B_{\rho}| \tag{5.8}
$$

for some $\tau_0 \in (t_0 - \rho^2, t_0 - \nu \rho^2/2]$. Then, there exist sufficiently small numbers $\gamma_0 \in (0, \nu/9)$, $\theta_0 \in$ $p(0, 1/2)$ and $\rho_{\star} \in (0, 1)$, which depend at most on n, p, q, λ , Λ , K , δ , M and ν , such that (5.1) – (5.2) hold with $(\gamma, \tilde{\nu}, \hat{\nu}) \coloneqq (\gamma_0, \nu/4, \theta_0 \nu)$, provided $\rho \leq \rho_{\star}$.

Proof. We first note that [\(5.1\)](#page-21-5) always holds provided $\gamma_0 \in (0, \nu/9)$. For $\sigma \in (0, 1)$, we choose a suitable $\eta \in C_c^1(B_\rho; [0, 1])$ satisfying $\eta|_{B_{(1-\sigma)\rho}} \equiv 1$, and apply [\(3.5\)](#page-16-3) with $Q_0 = B_\rho \times (\tau_0, \tau_1]$, where $\tau_1 := \tau_0 + \gamma_0(\frac{3}{2})$ $(\frac{3}{2}\rho)^2$. Define $A_{\theta_0 \nu, r}(\tau) := \{x \in B_r \mid U_{\delta, \varepsilon}(x, \tau) > (1 - \theta_0 \nu) \mu^2\}$ for $r \in (0, \rho], \tau \in (\tau_0, \tau_1].$ We choose $k := (1 - \nu)\mu^2$, so that $(U_{\delta, \varepsilon} - k)_+ \leq \nu \mu^2$. Also, $U_{\delta, \varepsilon} - k \geq (1 - \theta_0)\nu\mu^2$ holds in $A_{\theta_0 \nu, r}$. By $(3.5), |Q_0| \le \gamma_0 \rho^2 |B_\rho|$ $(3.5), |Q_0| \le \gamma_0 \rho^2 |B_\rho|$ and $q > n + 2$, we obtain

$$
(1 - \theta_0)^2 (\nu \mu^2)^2 |A_{\theta_0 \nu, (1 - \sigma)\rho}(\tau)|
$$

\n
$$
\leq (\nu \mu^2)^2 | \{ x \in B_\rho \mid U_{\delta, \varepsilon}(x, \tau_0) > (1 - \nu) \mu^2 \} | + C \left[\frac{(\nu \mu^2)^2}{(\sigma \rho)^2} |Q_0| + \mu^4 F^2 |Q_0|^{1 - 2/q} \right]
$$

\n
$$
\leq (\nu \mu^2)^2 |B_\rho| \left(1 - \frac{\nu}{2} + \frac{C\gamma_0}{\sigma^2} + \frac{CF^2 \rho^{2\beta}}{\nu^2} \right).
$$

Noting $|A_{\theta_0\nu,\,\rho}(\tau)| \leqslant |B_{\rho} \setminus B_{(1-\sigma)\rho}| + |A_{\theta_0\nu,(1-\sigma)\rho}(\tau)|$, we have

$$
|A_{\theta_0 \nu, \rho}(\tau)| \leqslant \left(n\sigma + \frac{1-\nu/2}{(1-\theta_0)^2} + \frac{C_0\gamma_0}{\sigma^2(1-\theta_0)^2} + \frac{C_0\rho^{2\beta}}{(1-\theta_0)^2\nu^2}\right)|B_{\rho}|
$$

for some $C_0 = C_0(n, p, q, \lambda, \Lambda, K, F, \delta, M) \in (1, \infty)$. Finally, we choose $\sigma \in (0, 1), \theta_0 \in (0, 1/2)$, $\gamma_0 \in (0, \nu/9)$ and $\rho_{\star} \in (0, 1)$ satisfying $n\sigma \leq \frac{\nu}{24}$, $\frac{1-\nu/2}{(1-\theta_0)^3}$ $\frac{1-\nu/2}{(1-\theta_0)^2} \leq 1 - \frac{\nu}{8}, \frac{C_0\gamma_0}{\sigma^2(1-\theta_0)}$ $\frac{C_0 \gamma_0}{\sigma^2 (1-\theta_0)^2} \leq \frac{\nu}{24}$, and $\frac{C_0 \rho_\star^{2\beta}}{\nu^2 (1-\theta_0)^2} \leq \frac{\nu}{24}$, so that $|A_{\theta_0\hat{\nu},\rho}(\tau)| \leq (1 - \nu/4)|B_\rho|$ holds for all $\tau \in (\tau_0, \tau_1]$, provided $\rho \leq \rho_\star$.

5.2 Proof of the De Giorgi-type oscillation estimates

We conclude Section [5](#page-21-0) by giving the proof of Proposition [2.9.](#page-13-0)

Proof of Proposition [2.9.](#page-13-0) By [\(2.35\)](#page-13-4), there exists $\tau_0 \in (t_0 - \rho^2, t_0 - \nu \rho^2/2]$ such that [\(5.8\)](#page-23-2) holds. In fact, if such τ_0 never exists, we can easily compute $|Q_\rho \backslash S_{\rho,\nu,\mu}| \leq \frac{\nu}{2} + (1 - \frac{\nu}{2}) \cdot \frac{\nu}{2} < \nu |Q_\rho|$, which contradicts with [\(2.35\)](#page-13-4). We fix such τ_0 that satisfies [\(5.8\)](#page-23-2).

Let $\gamma_0 \in (0, \nu/9)$ be as in Lemma [5.3.](#page-23-1) We set $A := (t_0 - \tau_0)\rho^{-2} \in [\nu/2, 1)$, and choose $l_0 \in \mathbb{N}$, depending at most on $A = A(\nu)$ and γ_0 , as the unique number satisfying $9\gamma_0l_0 \geq 4A > 9\gamma_0(l_0-1)$. We also define $\gamma_1 := 4A/(9l_0)$, and $T_k := \tau_0 + (2 + \frac{k}{4})\gamma_1 \rho^2$ for each $k \in \{1, \ldots, 9l_0 - 8\}$. By the definitions, we have $l_0 \leq 1 + \frac{4}{9\gamma_0} =: L_0$, and $T_{9l_0-8} = t_0$. We fix $i_{\star} \in \mathbb{N}$ satisfying $\kappa := \sqrt{1 - 2^{-(i_{\star}+1)(9L_0-8)}} >$ $(\sqrt{\nu}/6)^{\beta}, \frac{C_{\dagger}}{(\nu/4)\sqrt{\pi}}$ $\frac{C_{\dagger}}{(\nu/4)\sqrt{i_{\star}\gamma_1}} \leq \alpha_{\star}(\gamma_1)$, and $\frac{C_{\dagger}}{\sqrt{i_{\star}(\gamma_1)}}$ $\frac{C_{\dagger}}{i_{\star}(\gamma_1/3)} \leq \alpha_{\star}(\gamma_1/3)$, where $C_{\dagger} \in (1, \infty)$ and $\alpha_{\star} \in (0, 1)$ are respectively given by Lemma [5.1](#page-21-2) and Lemma [5.2.](#page-22-0) By $\gamma_0 \ge \gamma_1 \ge 2\nu/(9L_0), i_{\star}$ is determined by $ν$ and $γ_0$, as well as n, p, q, $λ$, $Λ$, K , F , M , and $δ$. We choose the radius $ρ̃ ∈ (0, ρ_*)$ satisfying $\tilde{\rho}^{\beta} < 2^{-[(9l_0-8)i_{\star}+9(l_0-1)]}\theta_0 \nu$. We claim that

$$
U_{\delta,\varepsilon} \leqslant \left(1 - 2^{-(i_{\star}+1)k}\right)\mu^2 \quad \text{in} \quad B_{\frac{3}{2}\rho} \times I_k \tag{5.9}
$$

for every $k \in \{1, ..., 9l_0 - 8\}$, where $I_k := \left(t_0 + \frac{5}{4}\right)$ $\frac{5}{4}\gamma_1\rho^2$, T_k . To prove [\(5.9\)](#page-24-2) for $k = 1$, we are allowed to apply Lemmata [5.1](#page-21-2)[–5.2](#page-22-0) with $(\gamma, \tilde{\nu}, \tilde{\nu}, \nu_0) := (\gamma_1, \nu/4, \theta_0 \nu, 2^{-i*} \theta_0 \nu)$ by our choice of $\tilde{\rho}$. Therefore, the proof of [\(5.9\)](#page-24-2) with $l_0 = 1$ is completed. For $l_0 \ge 2$, since our choice of $\tilde{\rho} \in (0, 1)$ yields $\tilde{\rho} < \tilde{\rho}^{\beta} < 2^{-(i_{\star}+1)k-i_{\star}}$ for every $k \in \{1, \ldots, 9l_0-9\}$, this allows us to repeatedly apply Lemmata [5.1–](#page-21-2)[5.2](#page-22-0) with $(\gamma, \tilde{\nu}, \tilde{\nu}, \nu_0) := (\gamma_1/3, 1, 2^{-(i_{\star}+1)k}\theta_0\nu, 2^{-(i_{\star}+1)k-i_{\star}}\theta_0\nu)$ for each $k \in \{1, ..., 9l_0-9\}$. Thus, $\left(\frac{\overline{\nu}}{3}\rho\right)^2$ $\frac{4}{9}(\tau_0-t_0)\leqslant-\left(\frac{\sqrt{\nu}}{3}\right)$ we conclude [\(5.9\)](#page-24-2) for every $k \in \{1, ..., 9l_0 - 8\}$. Noting $(\tau_0 + \frac{5}{4})$ $(\frac{5}{4}\gamma_1\rho^2)-t_0 \leq \frac{4}{9}$ and using [\(5.9\)](#page-24-2) with $k = 9l_0 - 8$, and recalling $U_{\delta,\varepsilon} = |\mathcal{G}_{\delta,\varepsilon}(\nabla u_{\varepsilon})|^2$, we finally obtain [\(2.36\)](#page-13-6) with $\kappa > (\sqrt{\nu}/6)^{\beta}.$ \Box

6 Non-degenerate case

Section [6](#page-24-0) aims to give the proof of Proposition [2.10.](#page-13-1) Here we mainly deal with an oscillation energy $\Phi(r) := \int_{Q_r} |\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_r|^2 dX$ for $r \in (0, \rho],$ where $Q_{2\rho} = Q_{2\rho}(x_0, t_0)$ satisfies [\(2.33\)](#page-12-4). To make it successful, we carefully use [\(2.34\)](#page-12-3) and [\(2.37\)](#page-13-3) to verify that the average integral of a gradient will not degenerate.

We recall basic inequalities for $G_{p,\varepsilon}$ defined in Section [3.3.](#page-17-0) To be precise, we use

$$
|G_{p,\varepsilon}(z_1) - G_{p,\varepsilon}(z_2)| \ge c(p) \max\left\{ |z_1|^{p-1}, |z_2|^{p-1} \right\} |z_1 - z_2|
$$
\n(6.1)

for all $z_1, z_2 \in \mathbb{R}^n$ (see [\[44,](#page-34-0) Lemma 2.3 & §3.5]). Moreover, by [\(6.1\)](#page-24-3) and $G_{p,\varepsilon}(0) = 0$, we have

$$
\left|G_{p,\varepsilon}^{-1}(w)\right| \leqslant C(p)|w|^{1/p} \quad \text{for all} \quad w \in \mathbb{R}^n. \tag{6.2}
$$

Throughout Section [6,](#page-24-0) we often use a well-known fact that, for $l \in \mathbb{N}$, there holds

$$
\oint_{U} |f - f_{U}|^{2} dy \le \oint_{U} |f - \xi|^{2} dy \quad \text{for all } \xi \in \mathbb{R}^{n}.
$$
\n(6.3)

for any $f \in L^2(U, \mathbb{R}^n)$, where $U \subset \mathbb{R}^l$ is a Lebesgue measurable set with $0 < |U| < \infty$.

6.1 Energy estimates

From Lemma [3.3,](#page-17-4) we aim to prove Lemma [6.1](#page-24-4) below, which plays an important role in making our comparison arguments successful.

Lemma 6.1. For every $\theta \in (0, 1/16)$, there exist sufficiently small $\nu \in (0, 1/4)$ and $\rho_* \in (0, 1)$, depending at most on n, p, q, λ , Λ , K , F , M , δ , and θ , such that the following estimates [\(6.4\)](#page-24-5)–[\(6.5\)](#page-24-6) hold, if the assumptions of Proposition [2.10](#page-13-1) and $\rho \leq \rho_*$ are verified.

$$
|(\nabla u_{\varepsilon})_{\rho}| \geq \delta + \frac{\mu}{2}.\tag{6.4}
$$

$$
\iint_{Q_{\rho}} |\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_{\rho}|^2 dX \leq \theta \mu^2.
$$
\n(6.5)

In addition to Lemma [3.3,](#page-17-4) we use

$$
|\nabla u_{\varepsilon}| \geq \frac{7}{8}\delta + (1 - \nu)\mu \quad \text{a.e. in } S_{\rho, \mu, \nu},\tag{6.6}
$$

which is easy to deduce by $\varepsilon < \delta/8$. In fact, we have $\delta + (1 - \nu)\mu \leq V_{\varepsilon} \leq \varepsilon + |\nabla u_{\varepsilon}| \leq \frac{\delta}{8} + |\nabla u_{\varepsilon}|$ a.e. in $S_{\rho,\mu,\nu}$. The inequality [\(6.6\)](#page-25-0) also allows us to apply [\(6.1\)](#page-24-3), which yields Lemma [6.2.](#page-25-1)

Lemma 6.2. Under the assumptions of Proposition [2.10,](#page-13-1) there exists a constant $C \in (1, \infty)$, depending at most on n, p, q, λ , Λ , K , F , M and δ , such that

$$
\Phi(\sigma \rho) \leqslant \frac{C\mu^2}{\sigma^{n+2}} \left[(1 - \sigma) + \frac{\sqrt{\nu}}{(1 - \sigma)^3} + \frac{F\rho^{\beta}}{(1 - \sigma)^2 \sqrt{\nu}} \right] \tag{6.7}
$$

for all $\sigma \in (0, 1)$.

Proof. We fix $\sigma \in (0, 1)$, and set \mathbb{R}^n -valued functions

$$
H_1(t) := \int_{B_{\sigma\rho}} \nabla u_{\varepsilon}(x, t) dx, \quad H_p(t) := \int_{B_{\sigma\rho}} G_{p,\varepsilon}(\nabla u_{\varepsilon}(x, t)) dx, \quad \text{and} \quad \Xi(t) := G_{p,\varepsilon}^{-1}(H_p(t)),
$$

defined for $t \in I_\rho$. By the definition of $G_{p,\varepsilon}$, [\(2.33\)](#page-12-4)–[\(2.34\)](#page-12-3) and [\(6.2\)](#page-24-7), it is easy to check $H_1(t) \leq 2\mu$ and $\Xi(t) \leq C_p \mu$. We apply [\(6.3\)](#page-24-8) to $\nabla u_{\varepsilon}(\cdot, t) \in L^2(B_{\sigma \rho}; \mathbb{R}^n)$ to deduce

$$
\Phi(\sigma \rho) \leq 2 \iint_{Q_{\sigma \rho}} |\nabla u_{\varepsilon}(x, t) - H_1(t)|^2 \, \mathrm{d}x \mathrm{d}t + 2 \iint_{Q_{\sigma \rho}} |H_1(t) - (\nabla u_{\varepsilon})_{\sigma \rho}|^2 \, \mathrm{d}x \mathrm{d}t
$$
\n
$$
\leq 2 \iint_{Q_{\sigma \rho}} |\nabla u_{\varepsilon} - \Xi(t)|^2 \, \mathrm{d}x \mathrm{d}t + C \mu \iint_{Q_{\sigma \rho}} |H_1(t) - (\nabla u_{\varepsilon})_{\sigma \rho}| \, \mathrm{d}x \mathrm{d}t
$$
\n
$$
\leq C \mu \left[\iint_{Q_{\sigma \rho}} |\nabla u_{\varepsilon} - \Xi(t)| \, \mathrm{d}x \mathrm{d}t + \sup_{t_1, t_2 \in I_{\sigma \rho}} |H_1(t_1) - H_1(t_2)| \right].
$$

Therefore, [\(6.7\)](#page-25-2) is shown by proving [\(6.8\)](#page-25-3)–[\(6.9\)](#page-25-4) to hold for any $t_1, t_2 \in I_{\sigma \rho}$.

$$
\iint_{Q_{\sigma\rho}} |\nabla u_{\varepsilon}(x, t) - \Xi(t)| \, \mathrm{d}x \mathrm{d}t \le \frac{C\mu}{\sigma^{n+2}} \left[\frac{\sqrt{\nu}}{1 - \sigma} + \frac{F\rho^{\beta}}{\sqrt{\nu}} \right]. \tag{6.8}
$$

$$
|H(t_1) - H(t_2)| \leqslant \frac{C\mu}{\sigma^n} \left[(1 - \sigma) + \frac{\sqrt{\nu}}{(1 - \sigma)^3} + \frac{F\rho^\beta}{(1 - \sigma)^2 \sqrt{\nu}} \right]. \tag{6.9}
$$

To prove [\(6.8\)](#page-25-3) from [\(3.6\)](#page-17-1)–[\(3.7\)](#page-17-2), we decompose $Q_{\sigma\rho} = (Q_{\sigma\rho} \setminus S_{\sigma\rho}) \cup S_{\sigma\rho}$, and note

$$
|Q_{\sigma\rho}\backslash S_{\sigma\rho}| = |Q_{\sigma\rho}\backslash S_{\rho}| \leqslant |Q_{\rho}\backslash S_{\rho}| \leqslant \nu |Q_{\rho}| = \sigma^{-(n+2)}\nu |Q_{\sigma\rho}| \tag{6.10}
$$

by [\(2.37\)](#page-13-3). We also recall [\(6.6\)](#page-25-0), so that [\(6.1\)](#page-24-3) can be applied to $|\nabla u_{\varepsilon}(x, t) - \Xi(t)|$ for $(x, t) \in S_{\sigma \rho}$. Combining with (6.10) and the Poincaré–Wirtinger inequality, we get

$$
\iint_{Q_{\sigma\rho}} |\nabla u_{\varepsilon}(x, t) - \Xi_1(t)| \, dX \le \frac{C\nu\mu}{\sigma^{n+2}} + \frac{C}{\mu^{p-1}} \iint_{Q_{\sigma\rho}} |G_{p,\varepsilon}(\nabla u_{\varepsilon}(x, t)) - H_p(t)| \, dX
$$
\n
$$
\le \frac{C\nu\mu}{\sigma^{n+2}} + \frac{C}{\mu^{p-1}}(\sigma\rho) \iint_{Q_{\sigma\rho}} |\nabla [G_{p,\varepsilon}(\nabla u_{\varepsilon})]| \, dX.
$$

To estimate the last integral, we use the decomposition $Q_{\sigma\rho} = (Q_{\sigma\rho} \setminus S_{\sigma\rho}) \cup S_{\sigma\rho}$ and [\(6.10\)](#page-25-5) again. By Hölder's inequality and (3.6) – (3.7) , we have

$$
\iint_{Q_{\sigma\rho}} |\nabla u_{\varepsilon}(x, t) - \Xi_1(t)| \, dx dt \leq \frac{C\nu\mu}{\sigma^{n+2}} + \frac{C\mu}{\sigma^{n+1}} \left[\frac{\left(1 + \sigma^{n/2+1}\right)\sqrt{\nu}}{1 - \sigma} + \left(\sqrt{\nu} + \frac{\sigma^{n/2+1}}{\sqrt{\nu}}\right) F\rho^{\beta}\right].
$$

By $\nu \in (0, 1/4)$ and $\sigma \in (0, 1)$, it is easy to deduce [\(6.8\)](#page-25-3).

To prove [\(6.9\)](#page-25-4), we set $\tilde{\sigma} := (1 + \sigma)/2 \in (1/2, 1)$, and choose $\eta \in C_c^2(B_{\tilde{\sigma} \rho}; [0, 1])$ satisfying $\eta|_{B_{\sigma \rho}} \equiv 1$. Without loss of generality, we may let t_1 , $t_2 \in I_{\sigma\rho}$ satisfy $t_1 < t_2$. We choose sufficiently small $\tilde{\varepsilon} > 0$, and define a piecewise linear function $\chi: \overline{I_{\tilde{\sigma}\rho}} \to [0, 1]$ such that $\chi \equiv 0$ on $[t_0 - (\tilde{\sigma}\rho)^2, t_1] \cup [t_2, t_0]$, $\chi \equiv 1$ on $[t_1 + \tilde{\varepsilon}, t_2 - \tilde{\varepsilon}]$, and χ is linearly interpolated in $(t_1, t_1 + \tilde{\varepsilon}) \cup (t_2 - \tilde{\varepsilon}, t_2)$. We will later let $\tilde{\varepsilon} \to 0$. We test $\varphi(x, t) := \eta(x)\chi(t)$, which is compactly supported in $Q_{\tilde{\sigma}, \rho}$, into the weak formulation

$$
-\iint_{Q_{\tilde{\sigma}_{\rho}}} \partial_{x_j} u_{\varepsilon} \partial_t \varphi \,dX + \iint_{Q_{\tilde{\sigma}_{\rho}}} \left\langle \nabla E_{\varepsilon} (\nabla u_{\varepsilon}) - \nabla E_{\varepsilon} (\Xi(t)) \mid \nabla \partial_{x_j} \varphi \right\rangle \,dX = \iint_{Q_{\tilde{\sigma}_{\rho}}} f_{\varepsilon} \partial_{x_j} \varphi \,dX.
$$

Letting $\tilde{\varepsilon} \to 0$, we have

$$
\left| \int_{B_{\tilde{\sigma}\rho}} \eta \left(\partial_{x_j} u_{\varepsilon}(x, t_1) - \partial_{x_j} u_{\varepsilon}(x, t_2) \right) dx \right| \leq \iint_{Q_{\tilde{\sigma}\rho}} \left[\left| \nabla E_{\varepsilon} (\nabla u_{\varepsilon}) - \nabla E_{\varepsilon} (\Xi(t)) \right| \left| \nabla^2 \eta \right| + |f_{\varepsilon}| |\nabla \eta| \right] dX
$$

To estimate the first integral on the right-hand side, we use $Q_{\tilde{\sigma}\rho} = (Q_{\tilde{\sigma}\rho} \setminus S_{\tilde{\sigma}\rho}) \cup S_{\tilde{\sigma}\rho}$ and [\(6.10\)](#page-25-5) with σ replaced by $\tilde{\sigma}$. By applying [\(2.10\)](#page-7-7) with $(z_1, z_2) = (\nabla u_{\varepsilon}(x, t), \Xi(t))$ for $(x, t) \in S_{\tilde{\sigma}\rho}$, using [\(6.8\)](#page-25-3) with σ replaced by $\tilde{\sigma}$, and summing over $j \in \{1, ..., n\}$, we get

$$
\frac{1}{|B_{\sigma\rho}|} \left| \int_{B_{\tilde{\sigma}\rho}} \eta \left(\nabla u_{\varepsilon}(x, t_1) - \nabla u_{\varepsilon}(x, t_2) \right) dx \right|
$$
\n
$$
\leq \frac{C(n, \delta, M)\mu}{(1 - \sigma)^2 \rho^2} \cdot \frac{\nu |Q_{\rho}|}{|B_{\sigma\rho}|} + \frac{C(n)\tilde{\sigma}^{n+2}}{(1 - \sigma)^2 \sigma^n} \iint_{Q_{\tilde{\sigma}\rho}} |\nabla u_{\varepsilon}(x, t) - \Xi(t)| dx dt + \frac{C(n)F|Q_{\rho}|^{1-1/q}}{(1 - \sigma)\sigma^n \rho^{n+1}}
$$
\n
$$
\leq \frac{C(n, p, q, \delta, M)\mu}{\sigma^n} \left[\frac{\nu}{(1 - \sigma)^2} + \frac{\sqrt{\nu}}{(1 - \sigma)^3} + \frac{F\rho^{\beta}}{(1 - \sigma)^2 \sqrt{\nu}} + \frac{F\rho^{\beta}}{1 - \sigma} \right].
$$

Recalling our choice of $\tilde{\sigma}$ and η , we can deduce [\(6.9\)](#page-25-4) as follows.

$$
\begin{split} &|H_1(t_1) - H_1(t_2)| \\ &\leq \frac{1}{|B_{\sigma\rho}|} \left[\left| \int_{B_{\tilde{\sigma}\rho} \setminus B_{\sigma\rho}} \eta \left(\nabla u_{\varepsilon}(x, t_1) - \nabla u_{\varepsilon}(x, t_2) \right) dx \right| + \left| \int_{B_{\tilde{\sigma}\rho}} \eta \left(\nabla u_{\varepsilon}(x, t_1) - \nabla u_{\varepsilon}(x, t_2) \right) dx \right| \right] \\ &\leq \frac{C\mu}{\sigma^n} \left[(1 - \sigma) + \frac{\sqrt{\nu}}{(1 - \sigma)^3} + \frac{F\rho^{\beta}}{(1 - \sigma)^2 \sqrt{\nu}} \right]. \end{split}
$$

Proof of Lemma [6.1.](#page-24-4) For $\sigma \in (0, 1)$, we apply (6.3) to $\nabla u_{\varepsilon} \in L^2(Q_{\rho}; \mathbb{R}^n)$, and make use of (6.7) . Then, we have

$$
\Phi(\rho) \leqslant \sigma^{n+2} \Phi(\sigma \rho) + \frac{|Q_{\rho} \backslash Q_{\sigma \rho}|}{|Q_{\rho}|} \cdot 4 \mu^2 \leqslant C_\dagger \mu^2 \left[(1-\sigma) + \frac{\sqrt{\nu}}{(1-\sigma)^3} + \frac{F \rho^\beta}{(1-\sigma)^2 \sqrt{\nu}} \right]
$$

with $C_f \in (0, \infty)$ independent of σ , ν , and ρ . Hence, [\(6.5\)](#page-24-6) holds if $\sigma \in (0, 1), \nu \in (0, 1/4)$, and $\rho_* \in (0, 1)$ satisfy $C_{\dagger}(1-\sigma) \leq \frac{\theta}{3}, \nu \leq \min\left\{\frac{3-8\sqrt{\theta}}{23}, \left(\frac{\theta(1-\sigma)^3}{3C_{\dagger}}\right)\right\}$ $3C_{\dagger}$ $\Big)^2$, and $\frac{C_{\dagger}F\rho^{\beta}_{*}}{(1-\sigma)^2\sqrt{\nu}} \leq \frac{\theta}{3}$ $\frac{\theta}{3}$. We use the triangle inequality, the Cauchy–Schwarz inequality, and (6.5) – (6.6) to get

$$
\begin{split} |(\nabla u_{\varepsilon})_{Q_{\rho}}| &\geq \iint_{Q_{\rho}} |\nabla u_{\varepsilon}| \, \mathrm{d}X - \left| \iint_{Q_{\rho}} \left[|\nabla u_{\varepsilon}| - |(\nabla u_{\varepsilon})_{Q_{\rho}}| \right] \, \mathrm{d}X \right| \\ &\geq \frac{|S_{\rho}|}{|Q_{\rho}|} \operatorname{ess} \inf_{S_{\rho}} |\nabla u_{\varepsilon}| - \sqrt{\Phi(\rho)} \geq (1 - \nu) \left(\frac{7}{8} \delta + (1 - \nu) \mu \right) - \sqrt{\theta} \mu. \end{split}
$$

Recalling (2.34) and our choice of ν , we conclude (6.4) .

 \Box

 \Box

6.2 Higher integrability

Before comparing u_{ε} with heat flows, we prove the higher integrability of $|\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_{\rho}|$ (Lemma [6.3\)](#page-27-1), which can be verified when $(\nabla u_{\varepsilon})_\rho$ does not degenerate.

Lemma 6.3. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \subseteq Q_R(x_*, t_*)$ satisfy $(2.33)-(2.34)$ $(2.33)-(2.34)$. If a vector $\xi \in \mathbb{R}^n$ satisfy

$$
\delta + \frac{\mu}{4} \le |\xi| \le \delta + \mu,\tag{6.11}
$$

then, there exist $\vartheta \in (0, 1)$ and $C \in (1, \infty)$, depending at most on n, p, q, λ , Λ , K , δ and M, such that $2(1 + \vartheta) \leqslant q$, and $\left(\iint_{Q_{\rho/2}} |\nabla u_{\varepsilon} - \xi|^{2(1+\vartheta)} dX\right)^{\frac{1}{1+\vartheta}} \leqslant C\left[\iint_{Q_{\rho}} |\nabla u_{\varepsilon} - \xi|^{2} dX + F^{2} \rho^{2\beta}\right].$

Without proof, we infer Lemma [6.4,](#page-27-2) often called Gehring's lemma.

Lemma 6.4. Fix a parabolic cylinder $Q_{2R} \subset \mathbb{R}^{n+1}$, and the exponents $1 < s < \tilde{s} < \infty$. Assume that non-negative functions $g \in L^s(Q_{2R})$ and $h \in L^{\tilde{s}}(Q_{2R})$ satisfy

$$
\oiint_{Q_r(z)} g^s dX \leq B_0 \left[\left(\oiint_{Q_{2r}(z)} g dX \right)^s + \oiint_{Q_{2r}(z)} h^s dX \right] + \theta \iint_{Q_{2r}(z)} g^s dX
$$

for any parabolic cylinder $Q_{2r}(z) \subset Q_{2R}$. If $\theta \leq \theta_0$ holds for some sufficiently small $\theta_0 = \theta_0(n) \in (0, 1)$, then there exists a constant $\vartheta_0 \in (0, \tilde{s}/s - 1)$, depending at most on n, s, \tilde{s} , and B_0 , such that there holds

$$
\iint_{Q_R} g^{s(1+\vartheta)} dX \leq B \left[\left(\iint_{Q_{2R}} g^s dX \right)^{1+\vartheta} + \iint_{Q_{2R}} h^{s(1+\vartheta)} dX \right]
$$

for every $\vartheta \in (0, \vartheta_0)$. Here the constant B depends at most on n, s, \tilde{s} , B_0 , and ϑ .

Remark 6.5. For elliptic cases, Gehring's lemma is shown by the Calderon–Zygmund cube de-composition and Vitali's covering lemma (see e.g., [\[27,](#page-33-14) $\S6.5$], [\[30,](#page-33-15) $\S6.4$], [\[50\]](#page-34-6)). It is often mentioned without proof that the elliptic arguments therein work even for parabolic cubes of the form $C_{r_0}(x_0, t_0) := K_r(x_0) \times (t_0 - r^2, t_0],$ where $K_r(x_0) := (x_{0,1} - r, x_{0,1} + r) \times \dots (x_{0,n} - r, x_{0,n} + r) \subset \mathbb{R}^n$ for $r_0 \in (0, \infty)$, $x_0 = (x_{0,1}, \ldots, x_{0,n}) \in \mathbb{R}^n$. Indeed, the interior of $C_{2r_0}(x_0, t_0)$ coincides with the open set $\{(x, t) \in \mathbb{R}^{n+1} \mid d'((x, t), (x_0, t_0 - 2r_0^2)) < 2r_0\}$, where d' is another metric in \mathbb{R}^{n+1} , defined as $d'((x, t), (y, s)) \coloneqq \max\left\{ |x_j - y_j|, \sqrt{2|t - s|} \right\}$ for $(x, t), (y, s) \in \mathbb{R}^{n+1}$. Thanks to this fact, we can prove Lemma [6.4](#page-27-2) by carrying out the Calder $\acute{\text{on}}$ –Zygmund cube decomposition, and applying Vitali's covering lemma for this metric d' (see also [\[31,](#page-33-16) §5]).

We provide the proof of Lemma [6.3](#page-27-1) (see also [\[28,](#page-33-17) §2] as the classical result).

Proof of Lemma [6.3.](#page-27-1) It suffices to find $C = C(n, p, q, \lambda, \Lambda, K, M, \delta) \in (0, \infty)$, such that

$$
\iint_{Q_r} |\nabla u_{\varepsilon} - \xi|^2 \, \mathrm{d}X \leq C \left[\left(\iint_{Q_{2r}} |\nabla u_{\varepsilon} - \xi|^{\frac{2n}{n+2}} \, \mathrm{d}X \right)^{\frac{n+2}{n}} + \iint_{Q_{2r}} |\rho f_{\varepsilon}|^2 \, \mathrm{d}X \right] + \theta_0 \iint_{Q_{2r}} |\nabla u_{\varepsilon} - \xi|^2 \, \mathrm{d}X
$$

holds for any $Q_{2r} = Q_{2r}(y_0, s_0) \subset Q_\rho = Q_\rho(x_0, t_0)$, where $\theta_0 = \theta_0(n)$ is given in Lemma [6.4.](#page-27-2) Indeed, the assertion enables us to apply Lemma [6.4](#page-27-2) to the functions $g := |\nabla u_{\varepsilon} - \xi|^{2n \over n+2} \in L^{\frac{n+2}{n}}(Q_\rho)$ and $h := |\rho f_{\varepsilon}|^{\frac{2n}{n+2}} \in L^{\frac{q(n+2)}{2n}}(Q_{\rho}).$ By Hölder's inequality and [\(2.14\)](#page-8-5), we conclude the desired estimate.

Fix $Q_{2r} = Q_{2r}(y_0, s_0) \subset Q_{\rho} = Q_{\rho}(x_0, t_0)$ arbitrarily, and define $w_{\varepsilon}(x, t) := u_{\varepsilon}(x, t) - \langle \xi | x - y_0 \rangle$ for $(x, t) \in Q_{2r}$. Set $r_l := 2^{l/2}r \in [r, 2r]$ for $l \in \{0, 1, 2\}$. For each $l \in \{1, 2\}$, we choose $\eta_l \in$ $C_c^1(B_{r_l}(y_0); [0, 1])$ and $\phi_l \in C^1(\overline{I_{r_l}}; [0, 1])$ satisfying $\phi_l(s_0 - r_l^2) = 0, \eta_l|_{B_{r_{l-1}(y_0)}} \equiv 1$, and $\phi_l|_{I_{r_{l-1}(y_0)}} \equiv 1$ 1. For this η_l , we define

$$
\tilde{w}_{\varepsilon,l}(t) := \left[\int_{B_{2r}} \eta_l^2 dx \right]^{-1} \int_{B_{2r}} w_{\varepsilon}(x, t) \eta_l^2 dx
$$

for $t \in I_{2r}(s_0)$. Let $\phi_h : \overline{I_{2r}(s_0)} \to [0, 1]$ be an arbitrary Lipschitz function that is non-increasing and $\phi_h(s_0) = 0$. We test $\varphi := (w_{\varepsilon} - \tilde{w}_{\varepsilon, l}) \eta_l^2 \phi$ with $\phi := \phi_l \phi_h$ into the weak formulation

$$
-\iint_{I_{2r}} \langle \partial_t \varphi, w_\varepsilon \rangle dt + \iint_{Q_{2r}} \langle \nabla E_\varepsilon(\nabla u_\varepsilon) - \nabla E_\varepsilon(\xi) | \nabla \varphi \rangle dX = \iint_{Q_{2r}} f_\varepsilon \varphi dX,
$$

which holds for any $\varphi \in X_0^2(s_0 - 4r^2, s_0; B_{2r}(y_0))$. Noting that the integral of $\varphi(\cdot, t)$ over B_{2r} vanishes for all $t \in I_{2r}(s_0)$, we can compute

$$
I_1 + I_2 := -\iint_{Q_{2r}} (w_{\varepsilon} - \tilde{w}_{\varepsilon,l}) \eta_l^2 \partial_t \phi \, dX + \iint_{Q_{2r}} \langle \nabla E_{\varepsilon} (\nabla u_{\varepsilon}) - \nabla E_{\varepsilon} (\xi) \mid \nabla w_{\varepsilon} \rangle \eta_l^2 \phi \, dX
$$

=
$$
-2 \iint_{Q_{2r}} \langle \nabla E_{\varepsilon} (\nabla u_{\varepsilon}) - \nabla E_{\varepsilon} (\xi) \mid \nabla \eta_l \rangle \eta_l (w_{\varepsilon} - \tilde{w}_{\varepsilon,l}) \phi \, dX + \iint_{Q_{2r}} f_{\varepsilon} (w_{\varepsilon} - \tilde{w}_{\varepsilon,l}) \varphi \, dX
$$

=:
$$
-2I_3 + I_4.
$$

By (6.11) , we can apply (2.9) – (2.10) to estimate I_2 and I_3 by below and by above respectively. Young's inequality and standard absorbing arguments yield

$$
-\frac{1}{2} \iint_{Q_{2r}} \left(w_{\varepsilon} - \tilde{w}_{\varepsilon,l} \right)^2 \eta_l^2 \phi_l \partial_t \phi_h \, dX + \frac{C_1}{2} \iint_{Q_{2r}} |\nabla w_{\varepsilon}|^2 \eta_l^2 \phi_l \phi_h \, dX
$$

\$\leqslant C \left[\iint_{Q_{2r}} |w_{\varepsilon} - \tilde{w}_{\varepsilon,l}|^2 \left(|\nabla \eta_l|^2 + |\partial_t \phi_l| + r_j^{-2} \right) \, dX + \frac{r_j^2}{2} \iint_{Q_{2r}} |f_{\varepsilon}|^2 \eta_l^2 \phi_l \, dX \right].

Choosing ϕ _h suitably, and noting $|Q_{r_j}| = r_j^2 |B_{r_j}|$, we easily deduce

$$
\sup_{\tau \in I_{r_l}^{\delta}} \int_{B_{r_l} \times \{\tau\}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,l}|^2}{r_l^2} \eta_l^2 \phi_l \,dx + \iint_{Q_{r_l}} |\nabla w_{\varepsilon}|^2 \eta_l^2 \phi_l \,dX
$$

$$
\leq C \left[\iint_{Q_{r_l}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,l}|^2}{r_l^2} \,dX + \iint_{Q_{r_l}} |r f_{\varepsilon}|^2 \,dX \right].
$$

Repeatedly using this estimate with $l = 1$ and $l = 2$, we deduce

$$
\iint_{Q_{r_0}} |\nabla w_{\varepsilon}|^2 dX \le \iint_{Q_{r_1}} |\nabla w_{\varepsilon}|^2 \eta_1^2 \phi_1 dX \le C \iint_{Q_{r_1}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,1}|^2}{r_1^2} dX + C \iint_{Q_{r_1}} |r f_{\varepsilon}|^2 dX, \text{ and}
$$
\n
$$
\sup_{\tau \in I_{r_1}^{\delta}} \int_{B_{r_1} \times \{\tau\}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,1}|^2}{r_1^2} dx
$$
\n
$$
\le 2 \sup_{\tau \in I_{r_1}^{\delta}} \left(\int_{B_{r_1} \times \{\tau\}} \frac{|w_{\varepsilon}(x, \tau) - \tilde{w}_{\varepsilon,2}(\tau)|^2}{r_1^2} dx + \frac{|\tilde{w}_{\varepsilon,2}(\tau) - \tilde{w}_{\varepsilon,1}(\tau)|^2}{r_1^2} \right)
$$
\n
$$
\le C_n \sup_{\tau \in I_{r_1}^{\delta}} \int_{B_{r_1} \times \{\tau\}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,2}|^2}{r_1^2} dx \le C_n \sup_{\tau \in I_{r_2}^{\delta}} \int_{B_{r_2} \times \{\tau\}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,2}|^2}{r_2^2} \eta_2^2 \phi_2 dx
$$
\n
$$
\le C \iint_{Q_{r_2}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,2}|^2}{r_2^2} dX + C \iint_{Q_{r_2}} |r f_{\varepsilon}|^2 dX \le C \iint_{Q_{r_2}} |\nabla w_{\varepsilon}|^2 dX + C \iint_{Q_{r_2}} |r f_{\varepsilon}|^2 dX,
$$

where we have used the Poincarè inequality for $(w_{\varepsilon}-\tilde{w}_{\varepsilon,2})(\cdot, t)$. Let the exponent κ be defined as $\kappa :=$ $2n/(n-2)$ for $n \geq 3$, and otherwise satisfy $\kappa \in (2, \infty)$. Interpolating the function $|w_{\varepsilon}(\cdot, t) - \tilde{w}_{1,\varepsilon}(t)|/r_1$ among the Lebesgue spaces $L^{\kappa}(B_{r_1}) \subset L^2(B_{r_1}) \subset L^{\frac{2n}{n+2}}(B_{r_1})$, applying the Poincaré inequality, and using Hölder's inequality for the time variable, we get

$$
\mathop{\rlap{\hskip2.5pt---}\int}\nolimits_{I_{r_{1}}}\left(\mathop{\rlap{\hskip2.5pt---}\int}\nolimits_{B_{r_{1}}}\frac{|w_{\varepsilon}(x,\,t)-\tilde w_{\varepsilon,1}(t)|^{2}}{r_{1}^{2}}\,\mathrm{d}x\right)^{1/2}\,\mathrm{d}t\leqslant C\left(\mathop{\rlap{\hskip2.5pt---}\int}\nolimits_{Q_{r_{1}}}\left|\nabla w_{\varepsilon}\right|^{2}\mathrm{d}X\right)^{\pi_{1}}\left(\mathop{\rlap{\hskip2.5pt---}\int}\nolimits_{Q_{r_{1}}}\left|\nabla w_{\varepsilon}\right|^{\frac{2n}{n+2}}\mathrm{d}X\right)^{\pi_{2}}
$$

with $(\pi_1, \pi_2) := (\frac{1}{4}, \frac{n+2}{4n})$ for $n \ge 3$, and $(\pi_1, \pi_2) = (\frac{\kappa'}{4}, \frac{\kappa-2}{2(\kappa-1)})$ $\frac{\kappa-2}{2(\kappa-1)}$ for $n = 2$. The interpolation among $L^{\infty}(I_{r_1}) \subset L^2(I_{r_1}) \subset L^1(I_{r_1})$ and Young's inequality yield

$$
\iint_{Q_{r_1}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,1}|^2}{r_1^2} dX \le \left(\sup_{\tau \in I_{r_1}^{\tilde{r}}} \int_{B_{r_1} \times \{\tau\}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,1}|^2}{r_1^2} dx \right)^{1/2} \int_{I_{r_1}} \left(\int_{B_{r_1}} \frac{|w_{\varepsilon} - \tilde{w}_{\varepsilon,1}|^2}{r_1^2} dx \right)^{1/2} dt
$$

$$
\le \sigma \iint_{Q_{r_2}} |\nabla w_{\varepsilon}|^2 dX + C(\sigma) \left[\left(\iint_{Q_{r_2}} |\nabla w_{\varepsilon}|^{\frac{2n}{n+2}} dx \right)^{\frac{n+2}{n}} + \iint_{Q_{r_2}} |r f_{\varepsilon}|^2 dX \right]
$$

for every $\sigma \in (0, 1)$. Noting $\nabla w_{\varepsilon} = \nabla u_{\varepsilon} - \xi$ and $2r \leq \rho$, we complete the proof by choosing sufficiently small $\sigma \in (0, 1)$. \Box

6.3 Comparison estimates

We consider a heat flow v_{ε} , and compare u_{ε} with v_{ε} . More precisely, we choose the function v_{ε} $u_{\varepsilon} + X_0^2(t_0 - \rho^2/4, t_0; B_{\rho/2}(x_0))$ satisfying

$$
\int_{t_0 - \rho^2/4}^{t_0} \langle \partial_t v_{\varepsilon}, \, \varphi \rangle \, \mathrm{d}t + \iint_{Q_{\rho/2}} \langle \nabla^2 E_{\varepsilon} ((\nabla u_{\varepsilon})_{\rho}) \nabla v_{\varepsilon} \mid \nabla \varphi \rangle \, \mathrm{d}X = 0 \tag{6.12}
$$

for all $\varphi \in X_0^2(t_0 - \rho^2/4, t_0; B_{\rho/2}(x_0))$, and $v_{\varepsilon}|_{t=t_0 - \rho^2/4} = u_{\varepsilon}|_{t=t_0 - \rho^2/4}$ in $L^2(B_{\rho/2}(x_0))$. In other words, v_{ε} is the weak solution of the Dirichlet boundary problem

$$
\begin{cases}\n\partial_t v_{\varepsilon} - \text{div} \left(\nabla^2 E_{\varepsilon} ((\nabla u_{\varepsilon})_{\rho}) \nabla v_{\varepsilon} \right) & = 0 \quad \text{in} \quad Q_{\rho/2}(x_0, t_0), \\
v_{\varepsilon} & = u_{\varepsilon} \quad \text{on} \quad \partial_p Q_{\rho/2}(x_0, t_0).\n\end{cases} (6.13)
$$

Lemma 6.6. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \in Q_R(x_*, t_*)$ satisfy (2.33) – (2.34) , and

$$
\delta + \frac{\mu}{4} \le |(\nabla u_{\varepsilon})_{\rho}|. \tag{6.14}
$$

Let v_{ε} be the weak solution of [\(6.13\)](#page-29-1). Then, the following [\(6.15\)](#page-29-2)–[\(6.16\)](#page-29-3) hold.

$$
\iint_{Q_{\rho/2}} |\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_{\rho/2}|^2 dX \leq C \left[\omega \left(\sqrt{\Phi(\rho)/\mu^2} \right)^{\frac{\vartheta}{1+\vartheta}} \Phi(\rho) + F^2 \rho^{2\beta} \right]. \tag{6.15}
$$

$$
\iint_{Q_{\sigma\rho}} |\nabla v_{\varepsilon} - (\nabla v_{\varepsilon})_{\sigma\rho}|^2 dX \le C\sigma^2 \iint_{Q_{\rho/2}} |\nabla v_{\varepsilon} - (\nabla v_{\varepsilon})_{\rho/2}|^2 dX \quad \text{for all } \sigma \in (0, 1/2). \tag{6.16}
$$

Here the exponent ϑ is given by Lemma [6.3,](#page-27-1) and the constant $C \in (0, \infty)$ depends at most on n, p, q, λ , Λ , K , δ , M , and ω .

Proof. For notational simplicity, we abbreviate $\xi := (\nabla u_{\varepsilon})_\rho$, which satisfies [\(6.11\)](#page-27-3) by [\(2.33\)](#page-12-4) and [\(6.14\)](#page-29-4). Hence, it is easy to check $\delta \leq (\varepsilon^2 + |\xi|^2)^{1/2} \leq \frac{\delta}{8} + M$. Combining with [\(2.4\)](#page-7-3)–[\(2.5\)](#page-7-9), we can easily find a sufficiently small constant $m \in (0, 1)$, depending at most on p, λ , Λ , K , δ and M, such that $m \mathrm{id}_n \leq \nabla^2 E_{\varepsilon}(\xi) \leq m^{-1} \mathrm{id}_n$ holds. In other words, the problem [\(6.13\)](#page-29-1) is uniformly parabolic in the classical sense, and therefore it is solvable. Moreover, since the coefficient matrix is constant, we can find a constant $C = C(n, m) \in (1, \infty)$ satisfying [\(6.16\)](#page-29-3). For the unique existence and regularity properties of v_{ε} , we refer the reader to [\[33,](#page-33-11) Chapter IV], [\[35,](#page-33-12) Chapter IV] (see also [\[9\]](#page-32-18)).

We are left to prove (6.15) . By (2.25) and (6.12) , we can deduce

$$
\int_{t_0 - \rho^2/4}^{t_0} \langle \partial_t (u_\varepsilon - v_\varepsilon), \, \varphi \rangle \, \mathrm{d}t + \iint_{Q_{\rho/2}} \langle \nabla^2 E_\varepsilon(\xi) (\nabla u_\varepsilon - \nabla v_\varepsilon) \, | \, \nabla \varphi \rangle \, \mathrm{d}X \tag{6.17}
$$

$$
= \iint_{Q_{\rho/2}} \left\langle \nabla^2 E_{\varepsilon}(\xi) (\nabla u_{\varepsilon} - \xi) - (\nabla E_{\varepsilon}(\nabla u_{\varepsilon}) - \nabla E_{\varepsilon}(\xi)) \mid \nabla \varphi \right\rangle dX + \iint_{Q_{\rho/2}} f_{\varepsilon} \varphi dX
$$

for all $\varphi \in X_0^2(t_0 - \rho^2/4, t_0; B_{\rho/2}(x_0))$. For sufficiently small $\tilde{\varepsilon} > 0$, which tends to 0 later, we choose $\phi_h: \overline{I_{\rho/2}} \to [0, 1]$ satisfying $\phi_h \equiv 1$ on $[t_0 - \rho^2/4, t_0 - \tilde{\varepsilon}]$, and linearly interpolated on the other domain with $\phi_h(t_0) = 0$. We test $\varphi := (u_{\varepsilon} - v_{\varepsilon})\phi_h$ into [\(6.17\)](#page-29-6). Note that we may discard the first integral in [\(6.17\)](#page-29-6), since it is positive. Letting $\tilde{\varepsilon} \to 0$, making use of [\(2.11\)](#page-7-8), and applying the embedding $W_0^{1,2}$ $\chi_0^{1,2}(B_{\rho/2}) \hookrightarrow L^2(B_{\rho/2})$ to $(u_{\varepsilon}-v_{\varepsilon})(\cdot,t) \in W_0^{1,2}$ $b_0^{1,2}(B_{\rho/2})$, we get

$$
m \iint_Q |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX \le C \left(\iint_Q \omega \left(\frac{|\nabla u_{\varepsilon} - \xi|}{\mu} \right)^2 |\nabla u_{\varepsilon} - \xi|^2 dX \right)^{\frac{1}{2}} \left(\int_Q |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX \right)^{\frac{1}{2}} + C(n) \rho \left(\iint_Q |f_{\varepsilon}|^2 dX \right)^{\frac{1}{2}} \left(\iint_Q |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX \right)^{\frac{1}{2}}.
$$

with $Q = Q_{\rho/2}(x_0, t_0)$. By making use of Hölder's inequality, and applying Jensen's inequality to the concave function ω , we have

$$
\iint_Q |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX \leq C \left[\left(\iint_Q |\nabla u_{\varepsilon} - \xi|^{2(1+\vartheta)} dX \right)^{\frac{1}{1+\vartheta}} \omega \left(\sqrt{\Phi(\rho)/\mu^2} \right)^{\frac{\vartheta}{1+\vartheta}} + F^2 \rho^{2\beta} \right],
$$

where we note $\omega(|\nabla u_{\varepsilon} - \xi|/\mu) \leq \omega(4)$ by [\(2.33\)](#page-12-4)–[\(2.34\)](#page-12-3). Using Lemma [6.3,](#page-27-1) we conclude [\(6.15\)](#page-29-2). \Box

Lemma 6.7. In addition to the assumptions of Theorem [2.8,](#page-12-1) let the positive numbers μ , M, and a cylinder $Q_{2\rho} = Q_{2\rho}(x_0, t_0) \in Q_R(x_*, t_*)$ satisfy (2.33) – (2.34) . For each $\sigma \in (0, 1/2)$, there exists a sufficiently small $\theta_0 \in (0, 1/16)$, depending on σ , ω , and ϑ , such that if [\(6.5\)](#page-24-6) and [\(6.14\)](#page-29-4) hold with $\theta \le \theta_0$, then there exists a constant $C_* \in (1, \infty)$, depending at most on n, p, q, λ , Λ , K , δ , M and ω , such that

$$
\Phi(\sigma \rho) \le C_* \left[\sigma^2 \Phi(\rho) + \frac{F^2}{\sigma^{n+2}} \rho^{2\beta} \right]. \tag{6.18}
$$

 \Box

Proof. Let v_{ε} be a weak solution of [\(6.13\)](#page-29-1). By applying [\(6.3\)](#page-24-8) to $\nabla u_{\varepsilon} \in L^2(Q_{\sigma\rho}; \mathbb{R}^n)$, $\nabla v_{\varepsilon} \in L^2(Q_{\rho/2}; \mathbb{R}^n)$, and making use of (6.5) and (6.15) – (6.16) , we have

$$
\Phi(\sigma \rho) \leq \iint_{Q_{\sigma \rho}} |\nabla u_{\varepsilon} - (\nabla v_{\varepsilon})_{\sigma \rho}|^2 dX \n\leq \frac{2}{(2\sigma)^{n+2}} \iint_{Q_{\rho/2}} |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX + 2 \iint_{Q_{\sigma \rho}} |\nabla v_{\varepsilon} - (\nabla v_{\varepsilon})_{\sigma \rho}|^2 dX \n\leq C \left[\left(\sigma^2 + \frac{1}{\sigma^{n+2}} \right) \iint_{Q_{\rho/2}} |\nabla u_{\varepsilon} - \nabla v_{\varepsilon}|^2 dX + \sigma^2 \Phi(\rho) \right] \n\leq \frac{C_*}{2} \left[\Phi(\rho) \frac{\omega(\theta^{1/2})^{\frac{\vartheta}{1+\vartheta}}}{\sigma^{n+2}} + \frac{F^2}{\sigma^{n+2}} \rho^{2\beta} + \sigma^2 \Phi(\rho) \right].
$$

Choosing $\theta_0 \in (0, 1/16)$ such that $\omega(\theta_0^{1/2})$ $\frac{\vartheta}{1+\vartheta} \leq \sigma^{n+4}$, we conclude [\(6.18\)](#page-30-1).

6.4 Proof of the Campanato-type growth estimates

Before the proof of Proposition [2.10,](#page-13-1) we infer an elementary lemma without proof.

Lemma 6.8. Provided that $g \in L^2(Q_\rho; \mathbb{R}^n)$ admits a constant $A \in (0, \infty)$ satisfying $\oiint_{Q_r} |g - g_r|$ $(g)_{Q_r}$ ² dX \leq $(\frac{r}{\rho})^{2\beta}$ for all $r \in (0, \rho]$, the limit $G_0 := \lim_{r \to 0} (g)_{Q_r} \in \mathbb{R}^n$ exists, and admits a constant $c_{\dagger \dagger} = c_{\dagger \dagger} (n, \beta) \in (1, \infty)$ such that $\iint_{Q_r} |g - G_0| \, dX \leq c_{\dagger \dagger} A(\frac{r}{\rho})$ $(\frac{r}{\rho})^{2\beta}$ for all $r \in (0, \rho]$.

Proof of Proposition [2.10.](#page-13-1) Let c_t , c_t be the positive constants found in [\(2.8\)](#page-7-11) and Lemma [6.8.](#page-30-2) The constants $\sigma \in (0, 1/2)$ and $\theta \in (0, 1/16)$ are determined to satisfy

$$
C_*\sigma^{2(1-\beta)} \leq \frac{1}{2}
$$
, and $\theta \leq \min\left\{\frac{\sigma^{n+2}}{64}, \theta_0(\sigma, \vartheta, \omega), \frac{\sigma^{n+2+2\beta}}{c_1c_{\dagger\dagger}}\right\}$,

where $\theta_0 \in (0, 1/16)$ and $C_* \in (0, \infty)$ are given by Lemma [6.7.](#page-30-3) Corresponding to this θ , we choose $\nu = \nu(\theta) \in (0, 1/4)$ and $\rho_* = \rho_*(\theta, \nu) \in (0, 1)$ as in Lemma [6.1.](#page-24-4) We choose $\hat{\rho} \in (0, \rho_*]$ satisfying $C_*F^2\hat{\rho}^{2\beta} \leq \theta\sigma^{n+2+2\beta}/2$. Assume $\rho \in (0, \hat{\rho}]$, and set $\rho_k := \sigma^k \rho$ for each $k \in \mathbb{Z}_{\geq 0}$. By iteration, we would like to prove

$$
\Phi(\rho_k) \leq \sigma^{2k\beta} \theta \mu^2, \quad \text{and} \quad |(\nabla u_{\varepsilon})_{\rho_k}| \geq \delta + \left[\frac{1}{2} - \frac{1}{8} \sum_{j=1}^{k-1} 2^{-j}\right] \mu \geq \delta + \frac{\mu}{4},\tag{6.19}
$$

for every $k \in \mathbb{Z}_{\geqslant 0}$. By [\(2.37\)](#page-13-3) and $\rho_0 \leqslant \rho_*,$ we can use Lemma [6.1](#page-24-4) to conclude [\(6.19\)](#page-31-6) for $k = 0$. Let [\(6.19\)](#page-31-6) be valid for $k \in \mathbb{Z}_{\geqslant 0}$. This induction hypothesis allows us to apply Lemma [6.7](#page-30-3) over a smaller cylinder Q_{ρ_k} . In particular, [\(6.18\)](#page-30-1) with ρ replaced by ρ_k implies $\Phi(\rho_{k+1}) \leq C_* \sigma^{2(1-\beta)} \cdot \sigma^{2\beta} \Phi(\rho_k) +$ C_*F^2 $\frac{C_{\ast}F^2}{\sigma^{n+2}}\left(\sigma^k\hat{\rho}\right)^{2\beta}\mu^2 \leq \sigma^{2(k+1)\beta}\theta\mu^2$. Also, the Cauchy–Schwarz inequality and the triangle inequality yield $|(\nabla u_{\varepsilon})_{\rho_{k+1}}| \geqslant |(\nabla u_{\varepsilon})_{\rho_{k+1}} - (\nabla u_{\varepsilon})_{\rho_{k}}| \geqslant |(\nabla u_{\varepsilon})_{\rho_{k}}| - \sigma^{-\frac{n+2}{2}} \sqrt{\Phi(\rho_{k})}$, by which and the induction hypothesis, [\(6.19\)](#page-31-6) holds true for $k + 1 \in \mathbb{Z}_{\geq 0}$.

For every $r \in (0, \rho]$, there corresponds a unique $k \in \mathbb{Z}_{\geq 0}$ such that $\rho_{k+1} < r \leq \rho_k$. Repeatedly applying [\(6.3\)](#page-24-8) to $\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}), \nabla u_{\varepsilon} \in L^2(Q_r; \mathbb{R}^n)$ and using [\(2.8\)](#page-7-11) and [\(6.19\)](#page-31-6), we have

$$
\iint_{Q_r} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}) - (\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}))_r|^2 dX \le \iint_{Q_r} |\mathcal{G}_{2\delta,\varepsilon}(\nabla u_{\varepsilon}) - \mathcal{G}_{2\delta,\varepsilon}((\nabla u_{\varepsilon})_r)|^2 dX
$$

\n
$$
\le c_\dagger^2 \iint_{Q_r} |\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_r|^2 dX \le c_\dagger^2 \sigma^{-(n+2)} \iint_{Q_{\rho_k}} |\nabla u_{\varepsilon} - (\nabla u_{\varepsilon})_{\rho_k}|^2 dX
$$

\n
$$
\le c_\dagger \sigma^{2k\beta - (n+2)} \theta \mu^2 \le \frac{c_\dagger \theta}{\sigma^{n+2+2\beta}} \left(\frac{\rho}{r}\right)^{2\beta} \mu^2.
$$

 \Box

From Lemma [6.8](#page-30-2) and our choice of θ , we complete the proof of Proposition [2.10.](#page-13-1)

Acknowledgments

The author gratefully acknowledges Prof. Yoshikazu Giga for his valuable comments on mathematical models for Bingham fluids and crystal surfaces. The author was supported by the Japan Society for the Promotion of Science through JSPS KAKENHI Grant Number 22KJ0861 during the preparation of the paper.

References

- [1] E. Acerbi and N. Fusco. Regularity for minimizers of nonquadratic functionals: the case $1 < p < 2$. J. Math. Anal. Appl., 140(1):115–135, 1989.
- [2] N. D. Alikakos and L. C. Evans. Continuity of the gradient for weak solutions of a degenerate parabolic equation. J. Math. Pures Appl. (9), 62(3):253–268, 1983.
- [3] P. Ambrosio and A. Passarelli di Napoli. Regularity results for a class of widely degenerate parabolic equations. Adv. Calc. Var., 2023.
- [4] L. Beck and G. Mingione. Lipschitz bounds and nonuniform ellipticity. *Comm. Pure Appl. Math.*, 73(5):944–1034, 2020.
- [5] V. Bögelein, F. Duzaar, R. Giova, and A. Passarelli di Napoli. Higher regularity in congested traffic dynamics. Math. Ann., 385(3-4):1823–1878, 2023.
- [6] V. B¨ogelein, F. Duzaar, N. Liao, and C. Scheven. Boundary regularity for parabolic systems in convex domains. J. Lond. Math. Soc. (2), 105(3):1702–1751, 2022.
- [7] V. Bögelein, F. Duzaar, N. Liao, and C. Scheven. Gradient Hölder regularity for degenerate parabolic systems. Nonlinear Anal., 225:Paper No. 113119, 61, 2022.
- [8] L. Brasco. Global L^{∞} gradient estimates for solutions to a certain degenerate elliptic equation. Nonlinear Anal., 74(2):516–531, 2011.
- [9] S. Campanato. Equazioni paraboliche del secondo ordine e spazi $\mathcal{L}^{2,\theta}(\Omega,\delta)$. Ann. Mat. Pura Appl. (4) , 73:55-102, 1966.
- [10] H. J. Choe. Hölder regularity for the gradient of solutions of certain singular parabolic systems. Comm. Partial Differential Equations, 16(11):1709–1732, 1991.
- [11] M. Colombo and A. Figalli. Regularity results for very degenerate elliptic equations. J. Math. Pures Appl. (9), 101(1):94–117, 2014.
- [12] E. DiBenedetto. $C^{1+\alpha}$ local regularity of weak solutions of degenerate elliptic equations. Nonlinear Anal., 7(8):827–850, 1983.
- [13] E. DiBenedetto. Degenerate Parabolic Equations. Universitext. Springer-Verlag, New York, 1993.
- [14] E. DiBenedetto. Real Analysis. Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkh¨auser Boston, Inc., Boston, MA, 2002.
- [15] E. DiBenedetto. Partial Differential Equations. Cornerstones. Birkhäuser Boston, Ltd., Boston, MA, second edition, 2010.
- [16] E. DiBenedetto and A. Friedman. Regularity of solutions of nonlinear degenerate parabolic systems. J. Reine Angew. Math., 349:83–128, 1984.
- [17] E. DiBenedetto and A. Friedman. Hölder estimates for nonlinear degenerate parabolic systems. J. Reine Angew. Math., 357:1–22, 1985.
- [18] E. DiBenedetto and A. Friedman. Addendum to: "Hölder estimates for nonlinear degenerate parabolic systems". J. Reine Angew. Math., 363:217–220, 1985.
- [19] E. DiBenedetto, U. Gianazza and V. Vespri. Harnack's inequality for degenerate and singular parabolic equations. Springer Monographs in Mathematics. Springer, New York, 2012.
- [20] E. DiBenedetto and M. A. Herrero. Nonnegative solutions of the evolution p-Laplacian equation. Initial traces and Cauchy problem when $1 < p < 2$. Arch. Ration. Mech. Anal., 111(3):225–290, 1990.
- [21] G. Duvaut and J.-L. Lions. Inequalities in Mechanics and Physics, volume 219 of Grundlehren der Mathematischen Wissenschaften. Springer-Verlag, Berlin-New York, 1976.
- [22] L. Esposito and G. Mingione. Some remarks on the regularity of weak solutions of degenerate elliptic systems. Rev. Mat. Complut., 11(1):203–219, 1998.
- [23] L. C. Evans. A new proof of local $C^{1,\alpha}$ regularity for solutions of certain degenerate elliptic p.d.e. J. Differential Equations, 45(3):356–373, 1982.
- [24] I. Fonseca, N. Fusco, and P. Marcellini. An existence result for a nonconvex variational problem via regularity. ESAIM Control Optim. Calc. Var., 7:69–95, 2002.
- [25] A. Gentile and A. Passarelli di Napoli. Higher regularity for weak solutions to degenerate parabolic problems. Calc. Var. Partial Differ. Equ., 2023.
- [26] M. Giaquinta. Multiple integrals in the calculus of variations and nonlinear elliptic systems, volume 105 of Annals of Mathematics Studies. Princeton University Press, Princeton, NJ, 1983.
- [27] M. Giaquinta and L. Martinazzi. An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs, volume 11 of Appunti. Scuola Normale Superiore di Pisa (Nuova Serie) [Lecture Notes. Scuola Normale Superiore di Pisa (New Series)]. Edizioni della Normale, Pisa, second edition, 2012.
- [28] M. Giaquinta and M. Struwe. On the partial regularity of weak solutions of nonlinear parabolic systems. Math. Z., 179(4):437–451, 1982.
- [29] Y. Giga and S. Tsubouchi. Continuity of derivatives of a convex solution to a perturbed one-Laplace equation by p-Laplacian. Arch. Ration. Mech. Anal., 244(2):253–292, 2022.
- [30] E. Giusti. Direct Methods in the Calculus of Variations. World Scientific Publishing Co., Inc., River Edge, NJ, 2003.
- [31] N. V. Krylov. On diffusion processes with drift in a morrey class containing L_{d+2} . J. Dyn. Differ. Equ., 35: 2813–2831 2023.
- [32] T. Kuusi and G. Mingione. Potential estimates and gradient boundedness for nonlinear parabolic systems. Rev. Mat. Iberoam., 28(2):535–576, 2012.
- [33] O. A. Ladyženskaja, V. A. Solonnikov, and N. N. Ural'ceva. *Linear and Quasilinear Equations* of Parabolic Type. Translations of Mathematical Monographs, Vol. 23. American Mathematical Society, Providence, R.I., 1968. Translated from the Russian by S. Smith.
- [34] J. L. Lewis. Regularity of the derivatives of solutions to certain degenerate elliptic equations. Indiana Univ. Math. J., 32(6):849–858, 1983.
- [35] G. M. Lieberman. *Second Order Parabolic Differential Equations*. World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [36] J.-L. Lions. Quelques méthodes de résolution des problèmes aux limites non linéaires. Dunod, Paris; Gauthier-Villars, Paris, 1969.
- [37] J. J. Manfredi. Regularity of the Gradient for a Class of Nonlinear Possibly Degenerate Elliptic Equations. ProQuest LLC, Ann Arbor, MI, 1986. Thesis (Ph.D.)–Washington University in St. Louis.
- [38] G. Mingione. Regularity of minima: an invitation to the dark side of the calculus of variations. Appl. Math., 51(4):355–426, 2006.
- [39] F. Santambrogio and V. Vespri. Continuity in two dimensions for a very degenerate elliptic equation. Nonlinear Anal., 73(12):3832–3841, 2010.
- [40] R. E. Showalter. Monotone operators in Banach space and nonlinear partial differential equations, volume 49 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, 1997.
- [41] H. Spohn. Surface dynamics below the roughening transition. Journal de Physique I, 3(1):69–81, 1993.
- [42] P. Tolksdorf. Regularity for a more general class of quasilinear elliptic equations. J. Differential Equations, 51(1):126–150, 1984.
- [43] S Tsubouchi. Local Lipschitz bounds for solutions to certain singular elliptic equations involving the one-Laplacian. Calc. Var. Partial Differential Equations, 60(1):Paper No. 33, 35, 2021.
- [44] S. Tsubouchi. Continuous differentiability of weak solutions to very singular elliptic equations involving anisotropic diffusivity. Adv. Calc. Var., 2023.
- [45] S. Tsubouchi. A weak solution to a perturbed one-Laplace system by p-Laplacian is continuously differentiable. Math. Ann., 388(2):1261–1322, 2024.
- [46] S. Tsubouchi. Gradient continuity for the parabolic $(1, p)$ -Laplace equation under the subcritical case. arXiv preprint [arXiv:2402.04951v](http://arxiv.org/abs/2402.04951)2, 2024.
- [47] N. N. Ural'ceva. Degenerate quasilinear elliptic systems. Zap. Naučn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI), 7:184–222, 1968.
- [48] M. Wiegner. On C_{α} -regularity of the gradient of solutions of degenerate parabolic systems. Ann. Mat. Pura Appl. (4), 145:385–405, 1986.
- [49] Z. Wu, J. Zhao, J. Yin, and H. Li. Nonlinear diffusion equations. World Scientific Publishing Co., Inc., River Edge, NJ, 2001. Translated from the 1996 Chinese original and revised by the authors.
- [50] A. Zatorska-Goldstein. Very weak solutions of nonlinear subelliptic equations. Ann. Acad. Sci. Fenn. Math., 30(2):407–436, 2005.