# ASYMPTOTIC BEHAVIORS OF MULTISCALE MULTIVALUED STOCHASTIC SYSTEMS WITH SMALL NOISES

## HUIJIE QIAO

School of Mathematics, Southeast University, Nanjing, Jiangsu 211189, P.R.China hjqiaoqean@seu.edu.cn

ABSTRACT. In this paper, we consider asymptotic behaviors of multiscale multivalued stochastic systems with small noises. First of all, for general, fully coupled systems for multivalued stochastic differential equations of slow and fast motions with small noises in the slow components, we prove the strong convergence in the functional law of large numbers by the time discretization scheme. Then for those systems whose slow diffusion coefficients don't depend on the fast components, the large deviation principle is established by the weak convergence approach.

#### 1. Introduction

Fix T > 0 and take a complete filtered probability space  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t \in [0,T]}, \mathbb{P})$ . Let  $(W_t^1), (W_t^2)$  be  $d_1$ - and  $d_2$ -dimensional standard Brownian motions defined on  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t \in [0,T]}, \mathbb{P})$ , respectively. Moreover,  $(W_t^1)$  and  $(W_t^2)$  are mutually independent. Consider the following slow-fast system on  $\mathbb{R}^n \times \mathbb{R}^m$ :

$$\begin{cases}
dX_t^{\varepsilon,\gamma} \in -A_1(X_t^{\varepsilon,\gamma})dt + b_1(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \sqrt{\varepsilon}\sigma_1(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dW_t^1, \\
X_0^{\varepsilon,\gamma} = x_0 \in \overline{\mathcal{D}}(A_1), \quad 0 \leqslant t \leqslant T, \\
dY_t^{\varepsilon,\gamma} \in -A_2(Y_t^{\varepsilon,\gamma})dt + \frac{1}{\gamma}b_2(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \frac{1}{\sqrt{\gamma}}\sigma_2(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dW_t^2, \\
Y_0^{\varepsilon,\gamma} = y_0 \in \overline{\mathcal{D}}(A_2), \quad 0 \leqslant t \leqslant T,
\end{cases} \tag{1}$$

where  $A_1, A_2$  are two multivalued maximal monotone operators (See Subsection 2.2), and these mappings  $b_1: \mathbb{R}^n \times \mathbb{R}^m \to \mathbb{R}^n$ ,  $\sigma_1: \mathbb{R}^n \times \mathbb{R}^m \to \mathbb{R}^{n \times d_1}$ ,  $b_2: \mathbb{R}^n \times \mathbb{R}^m \to \mathbb{R}^m$ ,  $\sigma_2: \mathbb{R}^n \times \mathbb{R}^m \to \mathbb{R}^{m \times d_2}$  are all Borel measurable. Here  $0 < \varepsilon < 1$  is a small parameter and  $\gamma := \gamma(\varepsilon) > 0$  represents the other small parameter (depending on  $\varepsilon$ ) which characterizes the ratio of timescales between processes  $X_{\varepsilon,\gamma}^{\varepsilon,\gamma}$  and  $Y_{\varepsilon,\gamma}^{\varepsilon,\gamma}$ .

We interpret the system (1) as a system of slow and fast components, where  $X^{\varepsilon,\gamma}$ ,  $Y^{\varepsilon,\gamma}$  are slow and fast components, respectively. And these systems like (1) are used for modelling in physics, engineering and economics and for optimization and equilibrium problems involving random data; see [1, 2, 3, 6, 7, 32] and the references therein. Different from common multiscale stochastic systems consisting of two stochastic differential equations (SDEs in short) (cf. e.g. [4, 11, 12, 14, 15, 20, 21, 22, 23, 34]), the system (1) contains two multivalued SDEs. Therefore, we call the system (1) a multiscale multivalued stochastic system. Moreover, we emphasize that two operators  $A_1, A_2$  are multivalued,

AMS Subject Classification(2020): 60H10, 60F10, 60F15.

Keywords: Multiscale multivalued stochastic systems with small noises; the functional law of large numbers; the large deviation principle.

This work was partly supported by NSF of China (No.12071071).

nonlinear and not smooth. The first goal of this paper is to show that as  $\varepsilon, \gamma \to 0$ , the solution  $X^{\varepsilon,\gamma}$  of the slow component in the system (1) will converge to the solution of the following average equation

$$\begin{cases} d\bar{X}_t \in -A_1(\bar{X}_t)dt + \bar{b}_1(\bar{X}_t)dt, \\ \bar{X}_0 = x_0 \in \overline{\mathcal{D}(A_1)}, \quad 0 \leqslant t \leqslant T, \end{cases}$$

where  $\bar{b}_1(x) := \int_{\mathbb{R}^m} b_1(x,y) \nu^x(\mathrm{d}y)$  and  $\nu^x$  is the unique invariant probability measure of Eq.(10) (See Subsection 3.1). Concretely speaking, under some suitable assumptions we prove that

$$\lim_{\varepsilon \to 0} \mathbb{E} \Big( \sup_{0 \le t \le T} |X_t^{\varepsilon, \gamma} - \bar{X}_t|^2 \Big) = 0.$$

Recently, Chen and Wu [8] considered the following multiscale stochastic system:

$$\begin{cases}
dX_t^{\gamma} \in -\partial \varphi(X_t^{\gamma}) dt + b_1(X_t^{\gamma}, Y_t^{\gamma}) dt + \sigma_1(X_t^{\gamma}) dW_t^1, \\
X_0^{\gamma} = x_0 \in \overline{\mathcal{D}(\partial \varphi)}, \quad 0 \leqslant t \leqslant T, \\
dY_t^{\gamma} \in -\partial \psi(Y_t^{\gamma}) dt + \frac{1}{\gamma} b_2(X_t^{\gamma}, Y_t^{\gamma}) dt + \frac{1}{\sqrt{\gamma}} \sigma_2(X_t^{\gamma}, Y_t^{\gamma}) dW_t^2, \\
Y_0^{\gamma} = y_0 \in \overline{\mathcal{D}(\partial \psi)}, \quad 0 \leqslant t \leqslant T,
\end{cases} \tag{2}$$

where  $\varphi, \psi$  are lower semicontinuous convex functions on  $\mathbb{R}^n, \mathbb{R}^m$ , respectively,  $\partial \varphi, \partial \psi$  are corresponding subdifferential operators and  $\sigma_1$  is independent of y. There they also obtained the above result. Note that by Example 2.1,  $\partial \varphi, \partial \psi$  are maximal monotone operators and the system (2) is a multiscale multivalued stochastic system. Thus, comparing Theorem 3.1 (See Subsection 3.1) with Theorem 4.3 in [8], we find that our result is more general. That is, the small noise in the slow component can improve the result, which is one of our motivations.

Next, it would be useful to further study the probability of the deviation  $X^{\varepsilon,\gamma}_{\cdot} - \bar{X}_{\cdot}$ . The study of these rare events in the multiscale framework is a difficult problem due to the presence of the underlying fast component. The first step is to develop the associated large deviations theory.

The large deviation principle (LDP in short) is an important topic in the probability theory and has widespread applications in many fields such as thermodynamics, statistics, information theory and engineering. Moreover, the research on the LDP for the common multiscale stochastic systems with the small noises has fruitful results in the literature (cf. [10, 16, 18, 24, 25, 26, 33, 35, 36]). As to the case in this paper, to the best of our knowledge there are few related results in the literature. Therefore, the second goal of this paper is to show the LDP for the following multiscale multivalued stochastic system:

$$\begin{cases}
dX_t^{\varepsilon,\gamma} \in -A_1(X_t^{\varepsilon,\gamma})dt + b_1(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \sqrt{\varepsilon}\sigma_1(X_t^{\varepsilon,\gamma})dW_t^1, \\
X_0^{\varepsilon,\gamma} = x_0 \in \overline{\mathcal{D}}(A_1), \quad 0 \leqslant t \leqslant T, \\
dY_t^{\varepsilon,\gamma} \in -A_2(Y_t^{\varepsilon,\gamma})dt + \frac{1}{\gamma}b_2(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \frac{1}{\sqrt{\gamma}}\sigma_2(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dW_t^2, \\
Y_0^{\varepsilon,\gamma} = y_0 \in \overline{\mathcal{D}}(A_2), \quad 0 \leqslant t \leqslant T.
\end{cases}$$
(3)

Concretely speaking, if the parameters  $\varepsilon$ ,  $\gamma$  satisfy

$$\lim_{\varepsilon \to 0} \frac{\gamma}{\varepsilon} = 0,$$

we prove that the family  $\{X^{\varepsilon,\gamma}, \varepsilon \in (0,1)\}$  satisfies the LDP in  $\mathbb{S} := C([0,T], \overline{\mathcal{D}(A_1)})$ .

In [25], Kushner considered the following multiscale stochastic system:

$$\begin{cases}
dX_t^{\varepsilon,\gamma} = b_1(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \sqrt{\varepsilon}\sigma_1(X_t^{\varepsilon,\gamma})dW_t^1 + dZ_t^{\varepsilon,\gamma}, \\
X_0^{\varepsilon,\gamma} = x_0, \quad 0 \leqslant t \leqslant T, \\
dY_t^{\varepsilon,\gamma} = \frac{1}{\gamma}b_2(X_t^{\varepsilon,\gamma}, Y_t^{\varepsilon,\gamma})dt + \frac{1}{\sqrt{\gamma}}\sigma_2(X_t^{\varepsilon,\gamma})dW_t^2, \\
Y_0^{\varepsilon,\gamma} = y_0, \quad 0 \leqslant t \leqslant T,
\end{cases} \tag{4}$$

where  $Z^{\varepsilon,\gamma}$  is a boundary reflection process depending on  $X^{\varepsilon,\gamma}$  and  $\sigma_2$  is independent of y. There he mentioned that when

$$\lim_{\varepsilon \to 0} \frac{\gamma}{\varepsilon} = 0,$$

and  $b_1, b_2$  are linear, the family  $\{X^{\varepsilon,\gamma}, \varepsilon \in (0,1)\}$  satisfies the LDP. Note that by Example 2.2, if we take special  $A_1$  and  $A_2 = 0$ , the system (3) can become the system (4). Therefore our result (Theorem 3.3) is more general.

Here we mention that based on the order that  $\varepsilon, \gamma$  converge to 0, there are three different regimes, i.e.

$$\lim_{\varepsilon \to 0} \frac{\gamma}{\varepsilon} = \begin{cases} 0, & \text{Regime 1;} \\ c \in (0, \infty), & \text{Regime 2;} \\ \infty, & \text{Regime 3.} \end{cases}$$

For Regime 2 and Regime 3, the LDP of current setting becomes quite complicated and our current approach does seem to work directly anymore. In [33], Spiliopoulos investigated the LDP for all three regimes by means of occupation measures and the characterization of optimal controls. However, for the system (3), due to containing the multivalued operators  $A_1, A_2$ , it is difficult to prove the tightness for a family of occupation measures through Poisson equations. Therefore, in the forthcoming work we will consider the cases of Regime 2 and Regime 3.

The novelty of this work lies in establishing the strong convergence in the functional law of large numbers and the LDP for systems of slow-fast motion as in (1). Since two operators  $A_1$ ,  $A_2$  are multivalued, nonlinear and not smooth, we need some new ideas and approaches so as to reach two goals.

The rest of this paper is organized as follows. In Section 2, we introduce some notations, maximal monotone operators and multivalued SDEs, and also list all assumptions. In Section 3, we formulate the main results. Before proving main results, we make preparations in Section 4. Two main theorems are showed in Section 5 and 6, respectively.

The following convention will be used throughout the paper: C with or without indices will denote different positive constants whose values may change from one place to another.

## 2. Notations and assumptions

In the section, we will recall some notations and some concepts of maximal monotone operators and multivalued SDEs, and finally list all assumptions.

2.1. **Notations.** In this subsection, we introduce some notations used in the sequel.

Let  $|\cdot|$ ,  $||\cdot||$  be the norms of a vector and a matrix, respectively. Let  $\langle\cdot,\cdot\rangle$  be the inner product of vectors on  $\mathbb{R}^n$ .  $U^*$  denotes the transpose of the matrix U.

Let  $\mathcal{B}_b(\mathbb{R}^n)$  be the set of all bounded Borel measurable functions on  $\mathbb{R}^n$ . Let  $C(\mathbb{R}^n)$  be the set of all functions which are continuous on  $\mathbb{R}^n$ .  $C^2(\mathbb{R}^n)$  represents the collection of all functions in  $C(\mathbb{R}^n)$  with continuous derivatives of order up to 2.

2.2. **Maximal monotone operators.** In the subsection, we introduce maximal monotone operators.

For a multivalued operator  $A: \mathbb{R}^n \mapsto 2^{\mathbb{R}^n}$ , where  $2^{\mathbb{R}^n}$  stands for all the subsets of  $\mathbb{R}^n$ , set

$$\mathcal{D}(A) := \left\{ x \in \mathbb{R}^n : A(x) \neq \emptyset \right\},$$

$$Gr(A) := \left\{ (x, y) \in \mathbb{R}^{2n} : x \in \mathcal{D}(A), \ y \in A(x) \right\}.$$

We say that A is monotone if  $\langle x_1 - x_2, y_1 - y_2 \rangle \ge 0$  for any  $(x_1, y_1), (x_2, y_2) \in Gr(A)$ , and A is maximal monotone if

$$(x_1, y_1) \in Gr(A) \iff \langle x_1 - x_2, y_1 - y_2 \rangle \geqslant 0, \quad \forall (x_2, y_2) \in Gr(A).$$

For readers to understand maximal monotone operators very well, we give two examples.

**Example 2.1.** For a lower semicontinuous convex function  $\varphi : \mathbb{R}^n \mapsto (-\infty, +\infty]$ , we assume  $\operatorname{Int}(Dom(\varphi)) \neq \emptyset$ , where  $Dom(\varphi) \equiv \{x \in \mathbb{R}^n; \varphi(x) < \infty\}$  and  $\operatorname{Int}(Dom(\varphi))$  is the interior of  $Dom(\varphi)$ . Define the subdifferential operator of the function  $\varphi$ :

$$\partial \varphi(x) := \{ y \in \mathbb{R}^n : \langle y, z - x \rangle + \varphi(x) \leqslant \varphi(z), \forall z \in \mathbb{R}^n \}.$$

Then  $\partial \varphi$  is a maximal monotone operator.

**Example 2.2.** For a closed convex subset  $\mathcal{O}$  of  $\mathbb{R}^n$ , we suppose  $\operatorname{Int}(\mathcal{O}) \neq \emptyset$ . Define the indicator function of  $\mathcal{O}$  as follows:

$$I_{\mathcal{O}}(x) := \begin{cases} 0, & \text{if } x \in \mathcal{O}, \\ +\infty, & \text{if } x \notin \mathcal{O}. \end{cases}$$

The subdifferential operator of  $I_{\mathcal{O}}$  is given by

$$\partial I_{\mathcal{O}}(x) := \{ y \in \mathbb{R}^n : \langle y, x - z \rangle \geqslant 0, \forall z \in \mathcal{O} \}$$

$$= \begin{cases} \emptyset, & \text{if } x \notin \mathcal{O}, \\ \{0\}, & \text{if } x \in \text{Int}(\mathcal{O}), \\ \Lambda_x, & \text{if } x \in \partial \mathcal{O}, \end{cases}$$

where  $\Lambda_x$  is the exterior normal cone at x. By simple deduction, we know that  $\partial I_{\mathcal{O}}$  is a maximal monotone operator.

In the following, we recall some properties of a maximal monotone operator A (cf.[6, 27]):

- (i)  $\operatorname{Int}(\mathcal{D}(A))$  and  $\overline{\mathcal{D}(A)}$  are convex subsets of  $\mathbb{R}^n$  with  $\operatorname{Int}\left(\overline{\mathcal{D}(A)}\right) = \operatorname{Int}\left(\mathcal{D}(A)\right)$ , where  $\operatorname{Int}(\mathcal{D}(A))$  denotes the interior of the set  $\mathcal{D}(A)$ .
- (ii) For every  $x \in \mathbb{R}^n$ , A(x) is a closed and convex subset of  $\mathbb{R}^n$ .

Take any T > 0 and fix it. Let  $\mathscr{V}_0$  be the set of all continuous functions  $K : [0, T] \mapsto \mathbb{R}^n$  with finite variations and  $K_0 = 0$ . For  $K \in \mathscr{V}_0$  and  $s \in [0, T]$ , we shall use  $|K|_0^s$  to denote the variation of K on [0,s] and write  $|K|_{TV} := |K|_0^T$ . Set

$$\mathcal{A} := \left\{ (X, K) : X \in C([0, T], \overline{\mathcal{D}(A)}), K \in \mathcal{V}_0, \right.$$
and  $\langle X_t - x, dK_t - ydt \rangle \geqslant 0$  for any  $(x, y) \in Gr(A) \right\}.$ 

And about  $\mathcal{A}$  we have the following two results (cf. [7, 38]).

**Lemma 2.3.** For  $X \in C([0,T], \overline{\mathcal{D}(A)})$  and  $K \in \mathcal{V}_0$ , the following statements are equivalent:

- (i)  $(X,K) \in \mathscr{A}$ .
- (ii) For any  $(x,y) \in C([0,T],\mathbb{R}^d)$  with  $(x_t,y_t) \in Gr(A)$ , it holds that

$$\langle X_t - x_t, dK_t - y_t dt \rangle \geqslant 0.$$

(iii) For any  $(X', K') \in \mathcal{A}$ , it holds that

$$\langle X_t - X_t', dK_t - dK_t' \rangle \geqslant 0.$$

**Lemma 2.4.** Assume that  $Int(\mathcal{D}(A)) \neq \emptyset$ . For any  $a \in Int(\mathcal{D}(A))$ , there exist  $M_1 > 0$ , and  $M_2, M_3 \geqslant 0$  such that for any  $(X, K) \in \mathscr{A}$  and  $0 \leqslant s < t \leqslant T$ ,

$$\int_{s}^{t} \langle X_r - a, dK_r \rangle \geqslant M_1 |K|_s^t - M_2 \int_{s}^{t} |X_r - a| dr - M_3 (t - s).$$

2.3. Multivalued SDEs. In the subsection, we introduce multivalued SDEs.

Fix T > 0 and consider the following multivalued SDE on  $\mathbb{R}^n$ :

$$dX_t \in -A(X_t)dt + b(X_t)dt + \sigma(X_t)dW_t^1, \quad 0 \le t \le T,$$
(5)

where A is a maximal monotone operator with  $\operatorname{Int}(\mathcal{D}(A)) \neq \emptyset$ , the coefficients  $b : \mathbb{R}^n \to \mathbb{R}^n$ ,  $\sigma : \mathbb{R}^n \to \mathbb{R}^n \times \mathbb{R}^{d_1}$  are Borel measurable and  $W^1$  is a  $d_1$ -dimensional Brownian motion on a filtered probability space  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t \in [0,T]}, \mathbb{P})$ .

**Definition 2.5.** We say that Eq.(5) admits a strong solution with the initial value  $X_0 \in \overline{\mathcal{D}(A)}$  if there exists a pair of adapted processes (X, K) on  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t \in [0,T]}, \mathbb{P})$  such that

- (i)  $X_t \in \mathscr{F}_t^{W^1}$ , where  $\{\mathscr{F}_t^{W^1}\}_{t \in [0,T]}$  stands for the  $\sigma$ -field filtration generated by  $W^1$ ,
- (ii)  $(X_{\cdot}(\omega), K_{\cdot}(\omega)) \in \mathscr{A} \mathbb{P} \ a.s.,$
- (iii) it holds that

$$\mathbb{P}\left\{\int_0^T (|b(X_s)| + \|\sigma(X_s)\|^2) \mathrm{d}s < +\infty\right\} = 1,$$

and

$$X_t = X_0 - K_t + \int_0^t b(X_s) ds + \int_0^t \sigma(X_s) dW_s^1, \quad 0 \leqslant t \leqslant T, \quad \mathbb{P} \ a.s..$$

2.4. A general criterion of large deviation principles. In this subsection, we present a general criterion to establish the LDP.

Let  $\mathbb S$  be a Polish space. For each  $\varepsilon > 0$ , let  $X^{\varepsilon}$  be a  $\mathbb S$ -valued random variable given on  $(\Omega, \mathscr F, \{\mathscr F_t\}_{t\in [0,T]}, \mathbb P)$ .

**Definition 2.6.** The function I on  $\mathbb{S}$  is called a rate function if for each  $M < \infty$ ,  $\{\varsigma \in \mathbb{S} : I(\varsigma) \leq M\}$  is a compact subset of  $\mathbb{S}$ .

**Definition 2.7.** We say that  $\{X^{\varepsilon}\}$  satisfies the Laplace principle with the rate function I, if for any real bounded continuous function g on  $\mathbb{S}$ ,

$$\lim_{\varepsilon \to 0} \varepsilon \log \mathbb{E} \left\{ \exp \left[ -\frac{g(X^{\varepsilon})}{\varepsilon} \right] \right\}_{5} = -\inf_{\varsigma \in \mathbb{S}} \left( g(\varsigma) + I(\varsigma) \right).$$

The Laplace principle is equivalent to that the family of  $\{X^{\varepsilon}, \varepsilon \in (0,1)\}$  satisfies the LDP in  $(\mathbb{S}, \mathscr{B}(\mathbb{S}))$  with the rate function I. More precisely, for any closed subset  $B_1 \in \mathscr{B}(\mathbb{S})$ ,

$$\limsup_{\varepsilon \to 0} \varepsilon \log \mathbb{P}(X^{\varepsilon} \in B_1) \leqslant -\inf_{\varsigma \in B_1} I(\varsigma),$$

and for any open subset  $B_2 \in \mathscr{B}(\mathbb{S})$ ,

$$\liminf_{\varepsilon \to 0} \varepsilon \log \mathbb{P}(X^{\varepsilon} \in B_2) \geqslant -\inf_{\varsigma \in B_2} I(\varsigma).$$

Let us state the conditions under which the Laplace principle holds. Set  $\mathbb{H} := L^2([0,T];\mathbb{R}^{d_1+d_2})$  and  $||h||_{\mathbb{H}} := (\int_0^T |h(t)|^2 dt)^{\frac{1}{2}}$  for  $h \in \mathbb{H}$ . Let  $\mathcal{A}$  be the collection of predictable processes  $u(\omega,\cdot)$  belonging to  $\mathbb{H}$  a.s.  $\omega$ . For each  $N \in \mathbb{N}$  we define two following spaces:

$$\mathbf{D}_2^N := \left\{ h \in \mathbb{H} : \|h\|_{\mathbb{H}}^2 \leqslant N \right\}, \quad \mathbf{A}_2^N := \left\{ u \in \mathcal{A} : u(\omega, \cdot) \in \mathbf{D}_2^N, \mathbb{P} \ a.s. \ \omega \right\}.$$

We equip  $\mathbf{D}_2^N$  with the weak convergence topology in  $\mathbb{H}$ . So,  $\mathbf{D}_2^N$  is metrizable as a compact Polish space. In the sequel,  $\mathbf{D}_2^N$  will be always endowed with this topology.

Condition 2.8. Let  $\Psi^{\varepsilon}: C([0,T];\mathbb{R}^{d_1+d_2}) \to \mathbb{S}$  be a family of measurable mappings. There exists a measurable mapping  $\Psi^0: C([0,T];\mathbb{R}^{d_1+d_2}) \to \mathbb{S}$  such that

(i) for  $N \in \mathbb{N}$  and  $\{h_{\varepsilon}, \varepsilon > 0\} \subset \mathbf{D}_2^N$ ,  $h \in \mathbf{D}_2^N$ , if  $h_{\varepsilon} \to h$  as  $\varepsilon \to 0$ , then

$$\Psi^0\left(\int_0^{\cdot} h_{\varepsilon}(s) \mathrm{d}s\right) \longrightarrow \Psi^0\left(\int_0^{\cdot} h(s) \mathrm{d}s\right).$$

(ii) for  $N \in \mathbb{N}$  and  $\{u_{\varepsilon}, \varepsilon > 0\} \subset \mathbf{A}_{2}^{N}$ ,  $u \in \mathbf{A}_{2}^{N}$ , if  $u_{\varepsilon}$  converges in distribution to u as  $\varepsilon \to 0$ , then

$$\Psi^{\varepsilon}\left(\sqrt{\varepsilon}W_{\cdot}+\int_{0}^{\cdot}u_{\varepsilon}(s)\mathrm{d}s\right)\stackrel{d}{\longrightarrow}\Psi^{0}\left(\int_{0}^{\cdot}u(s)\mathrm{d}s\right),$$

where W. is a  $d_1 + d_2$ -dimensional Brownian motion.

Given  $\varsigma \in \mathbb{S}$ , let  $\mathbf{D}_{\varsigma} = \{ h \in \mathbb{H} : \varsigma = \Psi^0(\int_0^{\cdot} h(s) ds) \}$ . Let  $I : \mathbb{S} \mapsto [0, \infty]$  be defined by

$$I(\varsigma) = \frac{1}{2} \inf_{h \in \mathbf{D}_{\varsigma}} ||h||_{\mathbb{H}}^{2}.$$

The following result is due to [5, Theorem 4.2].

**Theorem 2.9.** Set  $X^{\varepsilon} := \Psi^{\varepsilon}(\sqrt{\varepsilon}W)$ . Assume that Condition 2.8 holds. Then  $\{X^{\varepsilon}\}$  satisfies the Laplace principle with the rate function I given above.

2.5. **Assumptions.** In the subsection, we give out all the assumptions used in the sequel:  $(\mathbf{H}_{A_1})$   $0 \in \text{Int}(\mathcal{D}(A_1))$ .

 $(\mathbf{H}_{b_1,\sigma_1}^1)$  There exists a constant  $L_{b_1,\sigma_1} > 0$  such that for  $x_i \in \mathbb{R}^n$ ,  $y_i \in \mathbb{R}^m$ , i = 1, 2,

$$|b_1(x_1, y_1) - b_1(x_2, y_2)|^2 + ||\sigma_1(x_1, y_1) - \sigma_1(x_2, y_2)||^2 \leqslant L_{b_1, \sigma_1} (|x_1 - x_2|^2 + |y_1 - y_2|^2).$$

 $(\mathbf{H}_{A_2}) \ 0 \in \operatorname{Int}(\mathcal{D}(A_2)).$ 

 $(\mathbf{H}_{b_2,\sigma_2}^1)$  There exists a constant  $L_{b_2,\sigma_2} > 0$  such that for  $x_i \in \mathbb{R}^n$ ,  $y_i \in \mathbb{R}^m$ , i = 1, 2,

$$|b_2(x_1, y_1) - b_2(x_2, y_2)|^2 + ||\sigma_2(x_1, y_1) - \sigma_2(x_2, y_2)||^2 \leqslant L_{b_2, \sigma_2} (|x_1 - x_2|^2 + |y_1 - y_2|^2).$$

 $(\mathbf{H}_{b_2,\sigma_2}^2)$  There exists a constant  $\beta > 0$  satisfying  $\beta > 2L_{b_2,\sigma_2}$  such that for  $x \in \mathbb{R}^n$ ,  $y_i \in \mathbb{R}^m$ , i = 1, 2,

$$2\langle y_1 - y_2, b_2(x, y_1) - b_2(x, y_2) \rangle + \|\sigma_2(x, y_1) - \sigma_2(x, y_2)\|^2 \leqslant -\beta |y_1 - y_2|^2.$$

 $(\mathbf{H}_{\sigma_2}^3)$   $\sigma_2(x,y)$  is bounded.

**Remark 2.10.** (i)  $(\mathbf{H}_{A_1})$ ,  $(\mathbf{H}_{A_2})$  can be replaced by  $Int(\mathcal{D}(A_1)) \neq \emptyset$ ,  $Int(\mathcal{D}(A_2)) \neq \emptyset$ , respectively. Indeed, if we require  $Int(\mathcal{D}(A_1)) \neq \emptyset$ , for any  $a \in Int(\mathcal{D}(A_1))$ , by shifting the domain of  $A_1$  and defining  $\tilde{b}_1(x,y) = b_1(x-a,y)$ ,  $\tilde{\sigma}_1(x,y) = \sigma_1(x-a,y)$ , this situation becomes the case of  $0 \in Int(\mathcal{D}(A_1))$ .

(ii)  $(\mathbf{H}_{b_1,\sigma_1}^1)$  yields that there exists a constant  $\bar{L}_{b_1,\sigma_1} > 0$  such that for  $x \in \mathbb{R}^n$ ,  $y \in \mathbb{R}^m$ ,

$$|b_1(x,y)|^2 + ||\sigma_1(x,y)||^2 \leqslant \bar{L}_{b_1,\sigma_1}(1+|x|^2+|y|^2). \tag{6}$$

(iii)  $(\mathbf{H}_{b_2,\sigma_2}^1)$  implies that there exists a constant  $\bar{L}_{b_2,\sigma_2} > 0$  such that for  $x \in \mathbb{R}^n$ ,  $y \in \mathbb{R}^m$ ,

$$|b_2(x,y)|^2 + ||\sigma_2(x,y)||^2 \leqslant \bar{L}_{b_2,\sigma_2}(1+|x|^2+|y|^2). \tag{7}$$

(iv) By  $(\mathbf{H}_{b_2,\sigma_2}^1)$  and  $(\mathbf{H}_{b_2,\sigma_2}^2)$ , it holds that for  $x \in \mathbb{R}^n$ ,  $y \in \mathbb{R}^m$ 

$$2\langle y, b_2(x,y)\rangle + \|\sigma_2(x,y)\|^2 \leqslant -\alpha|y|^2 + C(1+|x|^2),\tag{8}$$

where  $\alpha := \beta - 2L_{b_2,\sigma_2}$  and C > 0 is a constant.

**Remark 2.11.** For these assumptions,  $(\mathbf{H}_{A_1})$ ,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$  assure well-posedness. Then we add  $(\mathbf{H}_{b_2,\sigma_2}^2)$  to obtain the functional law of large numbers, and also assume  $(\mathbf{H}_{\sigma_2}^3)$  to establish the large deviation principle.

## 3. FORMULATION FOR MAIN RESULTS

In this section, we provide main results of this paper.

3.1. **The functional law of large numbers.** In this subsection, we state the functional law of large numbers result.

Recall the following slow-fast system on  $\mathbb{R}^n \times \mathbb{R}^m$ :

$$\begin{cases}
dX_{t}^{\varepsilon,\gamma} \in -A_{1}(X_{t}^{\varepsilon,\gamma})dt + b_{1}(X_{t}^{\varepsilon,\gamma}, Y_{t}^{\varepsilon,\gamma})dt + \sqrt{\varepsilon}\sigma_{1}(X_{t}^{\varepsilon,\gamma}, Y_{t}^{\varepsilon,\gamma})dW_{t}^{1}, \\
X_{0}^{\varepsilon,\gamma} = x_{0} \in \overline{\mathcal{D}}(A_{1}), \quad 0 \leqslant t \leqslant T, \\
dY_{t}^{\varepsilon,\gamma} \in -A_{2}(Y_{t}^{\varepsilon,\gamma})dt + \frac{1}{\gamma}b_{2}(X_{t}^{\varepsilon,\gamma}, Y_{t}^{\varepsilon,\gamma})dt + \frac{1}{\sqrt{\gamma}}\sigma_{2}(X_{t}^{\varepsilon,\gamma}, Y_{t}^{\varepsilon,\gamma})dW_{t}^{2}, \\
Y_{0}^{\varepsilon,\gamma} = y_{0} \in \overline{\mathcal{D}}(A_{2}), \quad 0 \leqslant t \leqslant T.
\end{cases} \tag{9}$$

Under  $(\mathbf{H}_{A_1})$ ,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ , by Lemma 4.1 below, we obtain that the system (9) has a unique strong solution  $(X_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{1,\varepsilon,\gamma}, Y_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{2,\varepsilon,\gamma})$ .

Take any  $x \in \mathbb{R}^n$  and fix it. Consider the following multivalued SDE:

$$\begin{cases}
dY_t^x \in -A_2(Y_t^x)dt + b_2(x, Y_t^x)dt + \sigma_2(x, Y_t^x)dW_t^2, \\
Y_0^x = y_0 \in \overline{\mathcal{D}(A_2)}, \quad 0 \leqslant t \leqslant T.
\end{cases}$$
(10)

Under  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ , we know that the above equation has a unique strong solution  $(Y_{\cdot}^{x,y_0}, K_{\cdot}^{2,x,y_0})$  ([30]). Moreover, under  $(\mathbf{H}_{b_2,\sigma_2}^2)$ , by the same deduction as that of [27,

Theorem 3.2], one could obtain that there exists a unique invariant probability  $\nu^x$  for Eq.(10). So, we construct an average equation on  $(\Omega, \mathscr{F}, \{\mathscr{F}_t\}_{t\in[0,T]}, \mathbb{P})$  as follows:

$$\begin{cases}
d\bar{X}_t \in -A_1(\bar{X}_t)dt + \bar{b}_1(\bar{X}_t)dt, \\
\bar{X}_0 = x_0,
\end{cases} (11)$$

where  $\bar{b}_1(x) := \int_{\mathbb{R}^m} b_1(x, y) \nu^x(\mathrm{d}y)$ .

Now, it is the position to state the first main result.

**Theorem 3.1.** Suppose that  $(\mathbf{H}_{A_1})$ ,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$  hold. If

$$\lim_{\varepsilon \to 0} \frac{\gamma}{\varepsilon} = 0,$$

it holds that

$$\lim_{\varepsilon \to 0} \mathbb{E} \Big( \sup_{0 \le t \le T} |X_t^{\varepsilon, \gamma} - \bar{X}_t|^2 \Big) = 0,$$

where  $(\bar{X}, \bar{K})$  is a solution for Eq.(11).

The proof of Theorem 3.1 is placed in Section 5.

**Remark 3.2.** We mention that the conclusion of the above theorem is still right, if  $(\mathbf{H}_{b_1,\sigma_1}^1)$  is replaced by the following nonlipschitz condition:

 $(\mathbf{H}_{b_1,\sigma_1}^{1'})$  There exist concave nondecreasing continuous functions  $\rho_i : \mathbb{R}_+ \mapsto \mathbb{R}_+, i = 1, 2$ satisfying  $\rho_i(0) = 0$  and

$$\int_{0+} \frac{z}{z + \rho_1^2(z) + \rho_2^2(z)} dz = \infty,$$

such that for  $x_i \in \mathbb{R}^n$ ,  $y_i \in \mathbb{R}^m$ , i = 1, 2,

$$|b_1(x_1, y_1) - b_1(x_2, y_2)| + ||\sigma_1(x_1, y_1) - \sigma_1(x_2, y_2)|| \leq \rho_1(|x_1 - x_2|) + \rho_2(|y_1 - y_2|).$$

See [8] for some detailed proofs.

3.2. The large deviation principle. In this subsection, we require that  $\sigma_1$  is independent of y in the system (9), and state the large deviation principle result.

Theorem 3.3. Assume that 
$$(\mathbf{H}_{A_1})$$
,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$   $(\mathbf{H}_{\sigma_2}^3)$  hold. If 
$$\lim_{\varepsilon \to 0} \frac{\gamma}{\varepsilon} = 0,$$

the family  $\{X^{\varepsilon,\gamma}, \varepsilon \in (0,1)\}$  satisfies the LDP in  $\mathbb{S} := C([0,T], \overline{\mathcal{D}(A_1)})$  with the rate function given by

$$I(\varsigma) = \frac{1}{2} \inf_{h \in \mathbf{D}_{\varsigma}: \varsigma = \bar{X}^h} ||h||_{\mathbb{H}}^2,$$

where  $\bar{X}^h$  solves the following equation

$$\begin{cases}
d\bar{X}_t^h \in -A_1(\bar{X}_t^h)dt + \bar{b}_1(\bar{X}_t^h)dt + \sigma_1(\bar{X}_t^h)\pi_1h(t)dt, \\
\bar{X}_0^h = x_0 \in \overline{\mathcal{D}(A_1)},
\end{cases} (12)$$

and  $\pi_1: \mathbb{R}^{d_1+d_2} \mapsto \mathbb{R}^{d_1}$  is a projection operator.

The proof of Theorem 3.3 is placed in Section 6.

**Example 3.4.** The typical example that Theorem 3.3 is applicable to is a particular system of slow-fast motion, where the fast motion is a fast mean reverting process and the slow motion appears due to the interest in short time asymptotics. Concretely speaking, consider the following system on  $\mathbb{R} \times \mathbb{R}$ :

$$\begin{cases}
dX_t^{\varepsilon,\gamma} \in -\partial \varphi(X_t^{\varepsilon,\gamma}) dt + b(Y_t^{\varepsilon,\gamma}) dt + \sqrt{\varepsilon} \sigma(Y_t^{\varepsilon,\gamma}) dW_t^1, \\
X_0^{\varepsilon,\gamma} = x_0 \in \overline{\mathcal{D}}(\partial \varphi), \quad 0 \leqslant t \leqslant T, \\
dY_t^{\varepsilon,\gamma} \in -\partial \psi(Y_t^{\varepsilon,\gamma}) dt + \frac{1}{\gamma} (m - \frac{1}{2} Y_t^{\varepsilon,\gamma}) dt + \frac{1}{\sqrt{\gamma}} dW_t^2, \\
Y_0^{\varepsilon,\gamma} = y_0 \in \overline{\mathcal{D}}(\partial \psi), \quad 0 \leqslant t \leqslant T,
\end{cases}$$
(13)

where  $\varphi, \psi$  are lower semicontinuous convex functions on  $\mathbb{R}$ ,  $m \in \mathbb{R}$ ,  $W^1, W^2$  are 1-dimensional Brownian motions and  $b : \mathbb{R} \mapsto \mathbb{R}$ ,  $\sigma : \mathbb{R} \mapsto \mathbb{R}$  are measurable functions. It is obvious that

$$b_1(x,y) = b(y), \quad \sigma_1(x,y) = \sigma(y), \quad b_2(x,y) = m - \frac{1}{2}y, \quad \sigma_2(x,y) = 1.$$

If  $0 \in \text{Int}(\mathcal{D}(\partial \varphi))$ ,  $0 \in \text{Int}(\mathcal{D}(\partial \psi))$ , and  $b, \sigma$  are Lipschitz continuous, by Lemma 4.1, we know that the system (13) has a unique strong solution  $(X_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{1,\varepsilon,\gamma}, Y_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{2,\varepsilon,\gamma})$ . Note that for  $y_i \in \mathbb{R}$ , i = 1, 2

$$2\langle y_1 - y_2, b_2(x, y_1) - b_2(x, y_2) \rangle = 2\langle y_1 - y_2, m - \frac{1}{2}y_1 - m + \frac{1}{2}y_2 \rangle = -|y_1 - y_2|^2,$$

where  $\beta = 1, L_{b_2,\sigma_2} = \frac{1}{4}, \beta > 2L_{b_2,\sigma_2}$ . Thus, when  $\sigma(y) = \sigma$  for  $\sigma \in \mathbb{R}$ , by Theorem 3.3 we obtain that the family  $\{X^{\varepsilon,\gamma}, \varepsilon \in (0,1)\}$  satisfies the LDP in  $\mathbb{S} := C([0,T], \overline{\mathcal{D}(\partial\varphi)})$  with the rate function given by

$$I(\varsigma) = \frac{1}{2} \inf_{h \in \mathbf{D}_{\varsigma;\varsigma} = \bar{X}^h} ||h||_{\mathbb{H}}^2,$$

where  $\bar{X}^h$  solves the following equation

$$\begin{cases} d\bar{X}_t^h \in -\partial \varphi(\bar{X}_t^h) dt + \bar{b} dt + \sigma \pi_1 h(t) dt, \\ \bar{X}_0^h = x_0 \in \overline{\mathcal{D}(\partial \varphi)}, \end{cases}$$

 $\bar{b} = \int_{\mathbb{R}} b(y) \nu(\mathrm{d}y)$  and  $\nu$  is the unique invariant probability measure of the following equation

$$\begin{cases} dY_t \in -\partial \psi(Y_t) dt + (m - \frac{1}{2}Y_t) dt + dW_t^2, \\ Y_0 = y_0 \in \overline{\mathcal{D}(\partial \psi)}, \quad 0 \leqslant t \leqslant T. \end{cases}$$

This is consistent with the conclusion in [33, Section 6.2].

#### 4. Preparations

In this section, we make preparations so as to prove the main results.

## 4.1. Well-posedness for the system (9).

**Lemma 4.1.** Assume that  $(\mathbf{H}_{A_1})$ ,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$  hold. Then the system (9) has a unique strong solution  $(X_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{1,\varepsilon,\gamma}, Y_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{2,\varepsilon,\gamma})$ .

Since the proof of the above lemma is similar to that of [8, Theorem 6.1], we omit it.

## 4.2. Some estimates for the frozen equation (10).

**Lemma 4.2.** Under  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ - $(\mathbf{H}_{b_2,\sigma_2}^2)$ , it holds that for any  $t \in [0,T]$ ,  $x, x_1, x_2 \in \mathbb{R}^n$ ,  $y_1, y_2 \in \mathbb{R}^m$ 

$$\mathbb{E}|Y_t^{x,y_0}|^2 \leqslant |y_0|^2 e^{-\frac{\alpha}{2}t} + C(1+|x|^2),\tag{14}$$

$$\mathbb{E}|Y_t^{x_1,y_1} - Y_t^{x_2,y_2}|^2 \le |y_1 - y_2|^2 e^{-\alpha t} + C|x_1 - x_2|^2. \tag{15}$$

*Proof.* Applying the Itô formula to  $|Y_t^{x,y_0}|^2 e^{\lambda_1 t}$  for  $\lambda_1 = \frac{\alpha}{2}$  and taking the expectation, we get that

$$\begin{split} \mathbb{E}|Y_{t}^{x,y_{0}}|^{2}e^{\lambda_{1}t} &= |y_{0}|^{2} + \lambda_{1}\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|Y_{s}^{x,y_{0}}|^{2}\mathrm{d}s + 2\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}\langle Y_{s}^{x,y_{0}},\mathrm{d}K_{s}^{2,x,y_{0}}\rangle \\ &+ 2\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}\langle Y_{s}^{x,y_{0}},b_{2}(x,Y_{s}^{x,y_{0}})\rangle\mathrm{d}s + \mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}\|\sigma_{2}(x,Y_{s}^{x,y_{0}})\|^{2}\mathrm{d}s \\ &\leqslant |y_{0}|^{2} + \lambda_{1}\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|Y_{s}^{x,y_{0}}|^{2}\mathrm{d}s + 2\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|v||Y_{s}^{x,y_{0}}|\mathrm{d}s \\ &+ \mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}\left(-\alpha|Y_{s}^{x,y_{0}}|^{2} + C(1+|x|^{2})\right)\mathrm{d}s \\ &\leqslant |y_{0}|^{2} + \lambda_{1}\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|Y_{s}^{x,y_{0}}|^{2}\mathrm{d}s + \frac{\alpha}{2}\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|Y_{s}^{x,y_{0}}|^{2}\mathrm{d}s + C\int_{0}^{t}e^{\lambda_{1}s}\mathrm{d}s \\ &+ \mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}\left(-\alpha|Y_{s}^{x,y_{0}}|^{2} + C(1+|x|^{2})\right)\mathrm{d}s \\ &\leqslant |y_{0}|^{2} + (\lambda_{1} + \frac{\alpha}{2} - \alpha)\mathbb{E}\int_{0}^{t}e^{\lambda_{1}s}|Y_{s}^{x,y_{0}}|^{2}\mathrm{d}s + C(1+|x|^{2})\int_{0}^{t}e^{\lambda_{1}s}\mathrm{d}s \\ &= |y_{0}|^{2} + C(1+|x|^{2})\frac{e^{\lambda_{1}t} - 1}{\lambda_{1}}, \end{split}$$

where  $v \in A_2(0)$  and Lemma 2.3 is used. By the above deduction, it holds that

$$\mathbb{E}|Y_t^{x,y_0}|^2 \leqslant |y_0|^2 e^{-\frac{\alpha}{2}t} + C(1+|x|^2).$$

Next, we deal with the second estimate. Let  $(Y_{\cdot}^{x_1,y_1}, K_{\cdot}^{2,x_1,y_1})$  and  $(Y_{\cdot}^{x_2,y_2}, K_{\cdot}^{2,x_2,y_2})$  be the solutions to Eq.(10) with initial values  $y_1$  and  $y_2$ , respectively. Then it holds that

$$Y_t^{x_1,y_1} - Y_t^{x_2,y_2} = y_1 - y_2 - (K_t^{2,x_1,y_1} - K_t^{2,x_2,y_2}) + \int_0^t \left( b_2(x_1, Y_s^{x_1,y_1}) - b_2(x_2, Y_s^{x_2,y_2}) \right) ds + \int_0^t \left( \sigma_2(x_1, Y_s^{x_1,y_1}) - \sigma_2(x_2, Y_s^{x_2,y_2}) \right) dW_s^2.$$

Applying the Itô formula to  $|Y_t^{x_1,y_1} - Y_t^{x_2,y_2}|^2 e^{\lambda_2 t}$  for  $\lambda_2 = \alpha$  and taking expectation on two sides, we obtain that

$$\mathbb{E}|Y_t^{x_1,y_1} - Y_t^{x_2,y_2}|^2 e^{\lambda_2 t} = |y_1 - y_2|^2 + \lambda_2 \mathbb{E} \int_0^t e^{\lambda_2 s} |Y_s^{x_1,y_1} - Y_s^{x_2,y_2}|^2 ds$$

$$+2\mathbb{E} \int_0^t e^{\lambda_2 s} \langle Y_s^{x_1,y_1} - Y_s^{x_2,y_2}, d(K_s^{2,x_1,y_1} - K_s^{2,x_2,y_2}) \rangle$$

$$+2\mathbb{E} \int_0^t e^{\lambda_2 s} \langle Y_s^{x_1,y_1} - Y_s^{x_2,y_2}, b_2(x_1, Y_s^{x_1,y_1}) - b_2(x_2, Y_s^{x_2,y_2}) \rangle ds$$

$$+\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \|\sigma_{2}(x_{1}, Y_{s}^{x_{1}, y_{1}}) - \sigma_{2}(x_{2}, Y_{s}^{x_{2}, y_{2}})\|^{2} ds$$

$$\leq |y_{1} - y_{2}|^{2} + \lambda_{2} \mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} |Y_{s}^{x_{1}, y_{1}} - Y_{s}^{x_{2}, y_{2}}|^{2} ds$$

$$+2\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \langle Y_{s}^{x_{1}, y_{1}} - Y_{s}^{x_{2}, y_{2}}, b_{2}(x_{1}, Y_{s}^{x_{1}, y_{1}}) - b_{2}(x_{1}, Y_{s}^{x_{2}, y_{2}}) \rangle ds$$

$$+\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \|\sigma_{2}(x_{1}, Y_{s}^{x_{1}, y_{1}}) - \sigma_{2}(x_{1}, Y_{s}^{x_{2}, y_{2}})\|^{2} ds$$

$$+2\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \langle Y_{s}^{x_{1}, y_{1}} - Y_{s}^{x_{2}, y_{2}}, b_{2}(x_{1}, Y_{s}^{x_{2}, y_{2}}) - b_{2}(x_{2}, Y_{s}^{x_{2}, y_{2}}) \rangle ds$$

$$+\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \|\sigma_{2}(x_{1}, Y_{s}^{x_{1}, y_{1}}) - \sigma_{2}(x_{1}, Y_{s}^{x_{2}, y_{2}})\|^{2} ds$$

$$+2\mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} \|\sigma_{2}(x_{1}, Y_{s}^{x_{2}, y_{2}}) - \sigma_{2}(x_{2}, Y_{s}^{x_{2}, y_{2}})\|^{2} ds$$

$$\leq |y_{1} - y_{2}|^{2} + (\lambda_{2} - \beta + 2L_{b_{2}, \sigma_{2}}) \mathbb{E} \int_{0}^{t} e^{\lambda_{2}s} |Y_{s}^{x_{1}, y_{1}} - Y_{s}^{x_{2}, y_{2}}|^{2} ds$$

$$+C|x_{1} - x_{2}|^{2} \frac{e^{\lambda_{2}t} - 1}{\lambda_{2}},$$

which implies that

$$\mathbb{E}|Y_t^{x_1,y_1} - Y_t^{x_2,y_2}|^2 \leqslant |y_1 - y_2|^2 e^{-\alpha t} + C|x_1 - x_2|^2.$$

The proof is complete.

By the above lemma and the definition of  $\nu^x$ , it holds that

$$\int_{\overline{\mathcal{D}(A_2)}} |y|^2 \nu^x (\mathrm{d}y) = \int_{\overline{\mathcal{D}(A_2)}} \mathbb{E} |Y_t^{x,y}|^2 \nu^x (\mathrm{d}y) \leqslant \int_{\overline{\mathcal{D}(A_2)}} \left( |y|^2 e^{-\frac{\alpha}{2}t} + C(1+|x|^2) \right) \nu^x (\mathrm{d}y) 
= e^{-\frac{\alpha}{2}t} \int_{\overline{\mathcal{D}(A_2)}} |y|^2 \nu^x (\mathrm{d}y) + C(1+|x|^2),$$

and furthermore

$$\int_{\overline{\mathcal{D}(A_2)}} |y|^2 \nu^x (\mathrm{d}y) \leqslant C(1+|x|^2). \tag{16}$$

**Lemma 4.3.** Suppose that  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^2)$  hold. Then, for any  $t \in [0,T]$  there exists a constant C > 0 such that

$$|\mathbb{E}b_1(x, Y_t^{x, y_0}) - \bar{b}_1(x)|^2 \leqslant Ce^{-\alpha t} (1 + |x|^2 + |y_0|^2). \tag{17}$$

*Proof.* Based on (16), one can obtain that

$$|\mathbb{E}b_{1}(x, Y_{t}^{x, y_{0}}) - \bar{b}_{1}(x)|^{2} = \left|\mathbb{E}b_{1}(x, Y_{t}^{x, y_{0}}) - \int_{\overline{\mathcal{D}(A_{2})}} b_{1}(x, y)\nu^{x}(\mathrm{d}y)\right|^{2}$$

$$= \left|\mathbb{E}b_{1}(x, Y_{t}^{x, y_{0}}) - \int_{\overline{\mathcal{D}(A_{2})}} \mathbb{E}b_{1}(x, Y_{t}^{x, y})\nu^{x}(\mathrm{d}y)\right|^{2}$$

$$\leqslant \int_{\overline{\mathcal{D}(A_{2})}} \mathbb{E}|b_{1}(x, Y_{t}^{x,y_{0}}) - b_{1}(x, Y_{t}^{x,y})|^{2} \nu^{x}(\mathrm{d}y) \\
\leqslant L_{b_{1},\sigma_{1}} \int_{\overline{\mathcal{D}(A_{2})}} \mathbb{E}|Y_{t}^{x,y_{0}} - Y_{t}^{x,y}|^{2} \nu^{x}(\mathrm{d}y) \\
\stackrel{(15)}{\leqslant} L_{b_{1},\sigma_{1}} \int_{\overline{\mathcal{D}(A_{2})}} |y_{0} - y|^{2} e^{-\alpha t} \nu^{x}(\mathrm{d}y) \\
\leqslant 2L_{b_{1},\sigma_{1}} e^{-\alpha t} \left(|y_{0}|^{2} + \int_{\overline{\mathcal{D}(A_{2})}} |y|^{2} \nu^{x}(\mathrm{d}y)\right) \\
\leqslant Ce^{-\alpha t} (1 + |x|^{2} + |y_{0}|^{2}),$$

where the second inequality is based on  $(\mathbf{H}_{b_1,\sigma_1}^1)$ . The proof is complete.

**Lemma 4.4.** Under  $(\mathbf{H}_{A_2})$ ,  $(\mathbf{H}_{b_1,\sigma_1}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^1)$ ,  $(\mathbf{H}_{b_2,\sigma_2}^2)$ , it holds that for any  $x_1, x_2 \in \mathbb{R}^n$   $|\bar{b}_1(x_1) - \bar{b}_1(x_2)| \leqslant C|x_1 - x_2|. \tag{18}$ 

*Proof.* From (17) and (15), it follows that for any  $x_1, x_2 \in \mathbb{R}^n$ 

$$|\bar{b}_{1}(x_{1}) - \bar{b}_{1}(x_{2})|^{2} \leqslant 3|\bar{b}_{1}(x_{1}) - \mathbb{E}b_{1}(x_{1}, Y_{t}^{x_{1}, y_{0}})|^{2} + 3|\mathbb{E}b_{1}(x_{1}, Y_{t}^{x_{1}, y_{0}}) - \mathbb{E}b_{1}(x_{2}, Y_{t}^{x_{2}, y_{0}})|^{2} + 3|\mathbb{E}b_{1}(x_{2}, Y_{t}^{x_{2}, y_{0}}) - \bar{b}_{1}(x_{2})|^{2} \leqslant 3Ce^{-\alpha t}(1 + |x_{1}|^{2} + |y_{0}|^{2}) + 3C|x_{1} - x_{2}|^{2} + 3Ce^{-\alpha t}(1 + |x_{2}|^{2} + |y_{0}|^{2}).$$

Letting  $t \to \infty$ , we obtain that

$$|\bar{b}_1(x_1) - \bar{b}_1(x_2)| \leq C|x_1 - x_2|,$$

which completes the proof.

## 5. Proof of Theorem 3.1

In this section, we prove Theorem 3.1. The proof consists of two parts. In the first part (Subsection 5.1), we segment the time interval [0, T] by the size  $\delta$ , where  $\delta$  is a fixed positive number depending on  $\gamma$ , and introduce two auxiliary processes:

$$\begin{cases}
\hat{Y}_{t}^{\varepsilon,\gamma} \in -A_{2}(\hat{Y}_{t}^{\varepsilon,\gamma}) dt + \frac{1}{\gamma} b_{2}(X_{t(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{t}^{\varepsilon,\gamma}) dt + \frac{1}{\sqrt{\gamma}} \sigma_{2}(X_{t(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{t}^{\varepsilon,\gamma}) dW_{t}^{2}, \\
\hat{Y}_{0}^{\varepsilon,\gamma} = y_{0} \in \overline{\mathcal{D}}(A_{2}),
\end{cases} (19)$$

$$\begin{cases}
\hat{X}_{t}^{\varepsilon,\gamma} \in -A_{1}(\hat{X}_{t}^{\varepsilon,\gamma}) dt + b_{1}(X_{t(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{t}^{\varepsilon,\gamma}) dt + \sqrt{\varepsilon} \sigma_{1}(X_{t}^{\varepsilon,\gamma}, Y_{t}^{\varepsilon,\gamma}) dW_{t}^{1}, \\
\hat{X}_{0}^{\varepsilon,\gamma} = x_{0} \in \overline{\mathcal{D}(A_{1})},
\end{cases} (20)$$

where  $t(\delta) = [\frac{t}{\delta}]\delta$ , and  $[\frac{t}{\delta}]$  denotes the integer part of  $\frac{t}{\delta}$ .  $(\hat{Y}^{\varepsilon,\gamma}, \hat{K}^{2,\varepsilon,\gamma}), (\hat{X}^{\varepsilon,\gamma}, \hat{K}^{1,\varepsilon,\gamma})$  denote strong solutions of Eq.(19) and (20), respectively. Then we estimate  $X^{\varepsilon,\gamma}, K^{1,\varepsilon,\gamma}, Y^{\varepsilon,\gamma}, \hat{X}^{\varepsilon,\gamma}, \hat{K}^{1,\varepsilon,\gamma}, \hat{Y}^{\varepsilon,\gamma}, \hat{Y}^{\varepsilon,\gamma}$ . In the second part (Subsection 5.2), we present some estimates for the average equation (11).

#### 5.1. Some necessary estimates.

**Lemma 5.1.** Under the assumptions of Theorem 3.1, there exists a constant C > 0 such that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|X_t^{\varepsilon,\gamma}|^2\right) \leqslant C(1+|x_0|^2+|y_0|^2),\tag{21}$$

$$\sup_{t \in [0,T]} \mathbb{E}|Y_t^{\varepsilon,\gamma}|^2 \leqslant C(1+|x_0|^2+|y_0|^2), \tag{22}$$

$$\mathbb{E}|K^{1,\varepsilon,\gamma}|_0^T \leqslant C(1+|x_0|^2+|y_0|^2). \tag{23}$$

*Proof.* First of all, we estimate  $X_t^{\varepsilon,\gamma}$ . Note that  $X_t^{\varepsilon,\gamma}$  satisfies the following equation:

$$X_t^{\varepsilon,\gamma} = x_0 - K_t^{1,\varepsilon,\gamma} + \int_0^t b_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) ds + \sqrt{\varepsilon} \int_0^t \sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) dW_s^1.$$

By applying the Itô formula to  $|X^{\varepsilon,\gamma}_t|^2$ , it holds that

$$|X_{t}^{\varepsilon,\gamma}|^{2} = |x_{0}|^{2} - 2 \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma}, dK_{s}^{1,\varepsilon,\gamma} \rangle + 2 \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma}, b_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) \rangle ds$$
$$+2\sqrt{\varepsilon} \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma}, \sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) dW_{s}^{1} \rangle + \varepsilon \int_{0}^{t} \|\sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma})\|^{2} ds, \quad (24)$$

and furthermore for any  $u \in A_1(0)$ ,

$$|X_t^{\varepsilon,\gamma}|^2 \leqslant |x_0|^2 + 2|u| \int_0^t |X_s^{\varepsilon,\gamma}| ds + \int_0^t |X_s^{\varepsilon,\gamma}|^2 ds + \int_0^t |b_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})|^2 ds + 2\left|\int_0^t \langle X_s^{\varepsilon,\gamma}, \sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) dW_s^1 \rangle\right| + \int_0^t \|\sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})\|^2 ds,$$

where Lemma 2.3 is used in the above inequality. Then the BDG inequality and (6) imply that

$$\mathbb{E}\left(\sup_{s\in[0,t]}|X_{s}^{\varepsilon,\gamma}|^{2}\right)$$

$$\leqslant \left(|x_{0}|^{2}+2|u|T\right)+\left(2|u|+1\right)\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma}|^{2}\mathrm{d}r+\bar{L}_{b_{1},\sigma_{1}}\mathbb{E}\int_{0}^{t}(1+|X_{r}^{\varepsilon,\gamma}|^{2}+|Y_{r}^{\varepsilon,\gamma}|^{2})\mathrm{d}r\right)$$

$$+2\mathbb{E}\sup_{s\in[0,t]}\left|\int_{0}^{s}\left\langle X_{r}^{\varepsilon,\gamma},\sigma_{1}(X_{r}^{\varepsilon,\gamma},Y_{r}^{\varepsilon,\gamma})\mathrm{d}W_{r}^{1}\right\rangle\right|$$

$$\leqslant C(|x_{0}|^{2}+1)+C\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma}|^{2}\mathrm{d}r+C\mathbb{E}\int_{0}^{t}|Y_{r}^{\varepsilon,\gamma}|^{2}\mathrm{d}r\right)$$

$$+C\mathbb{E}\left(\int_{0}^{t}|X_{r}^{\varepsilon,\gamma}|^{2}|\|\sigma_{1}(X_{r}^{\varepsilon,\gamma},Y_{r}^{\varepsilon,\gamma})\|^{2}\mathrm{d}r\right)^{1/2}$$

$$\leqslant C(|x_{0}|^{2}+1)+C\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma}|^{2}\mathrm{d}r+C\mathbb{E}\int_{0}^{t}|Y_{r}^{\varepsilon,\gamma}|^{2}\mathrm{d}r\right)$$

$$+\frac{1}{2}\mathbb{E}\left(\sup_{s\in[0,t]}|X_{s}^{\varepsilon,\gamma}|^{2}\right)+C\mathbb{E}\int_{0}^{t}||\sigma_{1}(X_{r}^{\varepsilon,\gamma},Y_{r}^{\varepsilon,\gamma})||^{2}\mathrm{d}r,$$

and furthermore

$$\mathbb{E}\left(\sup_{s\in[0,t]}|X_s^{\varepsilon,\gamma}|^2\right) \leqslant C(|x_0|^2+1) + C\int_0^t \mathbb{E}|X_r^{\varepsilon,\gamma}|^2 dr + C\int_0^t \mathbb{E}|Y_r^{\varepsilon,\gamma}|^2 dr.$$
 (25)

For  $Y_t^{\varepsilon,\gamma}$ , fix  $v \in A_2(0)$ . Applying the Itô formula to  $|Y_t^{\varepsilon,\gamma}|^2 e^{\lambda t}$  for  $\lambda = \frac{\alpha}{2\gamma}$  and taking the expectation, one could obtain that

$$\begin{split} & \mathbb{E}|Y_t^{\varepsilon,\gamma}|^2 e^{\lambda t} \\ &= |y_0|^2 + \lambda \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma}|^2 e^{\lambda s} \mathrm{d}s - 2\mathbb{E} \int_0^t e^{\lambda s} \langle Y_s^{\varepsilon,\gamma}, \mathrm{d}K_s^{2,\varepsilon,\gamma} \rangle + \frac{2}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} \langle Y_s^{\varepsilon,\gamma}, b_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) \rangle \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} ||\sigma_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})||^2 \mathrm{d}s \\ \stackrel{(8)}{\leqslant} |y_0|^2 + \lambda \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma}|^2 e^{\lambda s} \mathrm{d}s + 2\mathbb{E} \int_0^t e^{\lambda s} |v| |Y_s^{\varepsilon,\gamma}| \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} \Big( -\alpha |Y_s^{\varepsilon,\gamma}|^2 + C(1 + |X_s^{\varepsilon,\gamma}|^2) \Big) \mathrm{d}s \\ & \leqslant |y_0|^2 + \Big(\lambda + \frac{\alpha}{2\gamma} - \frac{\alpha}{\gamma}\Big) \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma}|^2 e^{\lambda s} \mathrm{d}s + C \int_0^t e^{\lambda s} \mathrm{d}s + \frac{C}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} (1 + |X_s^{\varepsilon,\gamma}|^2) \mathrm{d}s \\ & \leqslant |y_0|^2 + C \frac{e^{\lambda t} - 1}{\lambda} + C \frac{e^{\lambda t} - 1}{\gamma \lambda} \left( \mathbb{E} \left( \sup_{s \in [0,t]} |X_s^{\varepsilon,\gamma}|^2 \right) + 1 \right). \end{split}$$

From this, it follows that

$$\mathbb{E}|Y_t^{\varepsilon,\gamma}|^2 \leqslant C(|y_0|^2 + 1) + C\mathbb{E}\left(\sup_{s \in [0,t]} |X_s^{\varepsilon,\gamma}|^2\right). \tag{26}$$

Inserting (26) in (25), by the Gronwall inequality one can get that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|X_t^{\varepsilon,\gamma}|^2\right)\leqslant C(1+|x_0|^2+|y_0|^2),$$

which together with (26) implies that

$$\sup_{t \in [0,T]} \mathbb{E} |Y_t^{\varepsilon,\gamma}|^2 \leqslant C(1 + |x_0|^2 + |y_0|^2).$$

Finally, for  $K^{1,\varepsilon,\gamma}$ , by (24) and Lemma 2.4, it holds that

$$\begin{split} |X_T^{\varepsilon,\gamma}|^2 &= |x_0|^2 - 2\int_0^T \langle X_s^{\varepsilon,\gamma}, \mathrm{d} K_s^{1,\varepsilon,\gamma} \rangle + 2\int_0^T \langle X_s^{\varepsilon,\gamma}, b_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) \rangle \mathrm{d} s \\ &+ 2\sqrt{\varepsilon} \int_0^T \langle X_s^{\varepsilon,\gamma}, \sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) \mathrm{d} W_s^1 \rangle + \varepsilon \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})\|^2 \mathrm{d} s \\ &\leqslant |x_0|^2 - 2M_1 \left| K^{1,\varepsilon,\gamma} \right|_0^T + 2M_2 \int_0^T |X_s^{\varepsilon,\gamma}| \mathrm{d} s + 2M_3 T + \int_0^T |X_s^{\varepsilon,\gamma}|^2 \mathrm{d} s \\ &+ \int_0^T |b_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})|^2 \mathrm{d} s + 2\sqrt{\varepsilon} \int_0^T \langle X_s^{\varepsilon,\gamma}, \sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) \mathrm{d} W_s^1 \rangle \\ &+ \varepsilon \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma})\|^2 \mathrm{d} s, \end{split}$$

and

$$2M_{1}\mathbb{E} \left| K^{1,\varepsilon,\gamma} \right|_{0}^{T} \leq |x_{0}|^{2} + 2(M_{2} + M_{3})T + (2M_{2} + 1) \int_{0}^{T} \mathbb{E} |X_{s}^{\varepsilon,\gamma}|^{2} ds + C \int_{0}^{T} (1 + \mathbb{E} |X_{s}^{\varepsilon,\gamma}|^{2} + \mathbb{E} |Y_{s}^{\varepsilon,\gamma}|^{2}) ds,$$

which together with (21) and (22) yields that

$$\mathbb{E} \left| K^{1,\varepsilon,\gamma} \right|_0^T \leqslant C(1 + |x_0|^2 + |y_0|^2).$$

The proof is complete.

By the same deduction to that in Lemma 5.1, we obtain the following estimate.

**Lemma 5.2.** Under the assumptions of Theorem 3.1, it holds that

$$\sup_{t \in [0,T]} \mathbb{E}|\hat{Y}_t^{\varepsilon,\gamma}|^2 \leqslant C(1+|x_0|^2+|y_0|^2), \tag{27}$$

$$\mathbb{E}\left(\sup_{t\in[0,T]}|\hat{X}_{t}^{\varepsilon,\gamma}|^{2}\right) \leqslant C(1+|x_{0}|^{2}+|y_{0}|^{2}),\tag{28}$$

$$\mathbb{E}|\hat{K}^{1,\varepsilon,\gamma}|_0^T \leqslant C(1+|x_0|^2+|y_0|^2). \tag{29}$$

Next, we estimate some differences.

Lemma 5.3. Under the assumptions of Theorem 3.1, we have that

$$\lim_{l \to 0} \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 = 0, \tag{30}$$

$$\lim_{l \to 0} \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant t \leqslant s+l} |\hat{X}_t^{\varepsilon,\gamma} - \hat{X}_s^{\varepsilon,\gamma}|^2 = 0. \tag{31}$$

*Proof.* Since the proofs of (30) and (31) are similar, we only prove (30). Note that for  $0 \le s < t \le T$ ,

$$X_t^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma} = -K_t^{1,\varepsilon,\gamma} + K_s^{1,\varepsilon,\gamma} + \int_s^t b_1(X_r^{\varepsilon,\gamma}, Y_r^{\varepsilon,\gamma}) dr + \int_s^t \sigma_1(X_r^{\varepsilon,\gamma}, Y_r^{\varepsilon,\gamma}) dW_r^1.$$

Thus, the Itô formula and (6) imply that

$$|X_{t}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}|^{2}$$

$$= -2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, dK_{r}^{1,\varepsilon,\gamma} \rangle + 2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, b_{1}(X_{r}^{\varepsilon,\gamma}, Y_{r}^{\varepsilon,\gamma}) \rangle dr$$

$$+2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, \sigma_{1}(X_{r}^{\varepsilon,\gamma}, Y_{r}^{\varepsilon,\gamma}) dW_{r}^{1} \rangle + \int_{s}^{t} \|\sigma_{1}(X_{r}^{\varepsilon,\gamma}, Y_{r}^{\varepsilon,\gamma})\|^{2} dr$$

$$\leq -2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, dK_{r}^{1,\varepsilon,\gamma} \rangle + \int_{s}^{t} |X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}|^{2} dr$$

$$+ \bar{L}_{b_{1},\sigma_{1}} \int_{s}^{t} (1 + |X_{r}^{\varepsilon,\gamma}|^{2} + |Y_{r}^{\varepsilon,\gamma}|^{2}) dr + 2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, \sigma_{1}(X_{r}^{\varepsilon,\gamma}, Y_{r}^{\varepsilon,\gamma}) dW_{r}^{1} \rangle. (32)$$

Next, we compute  $-2\int_s^t \langle X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}, \mathrm{d} K_r^{1,\varepsilon,\gamma} \rangle$ . Note that  $0 \in \mathrm{Int}(\mathcal{D}(A_1))$ . Thus, there is a  $\theta_0 > 0$  such that for any R > 0 and  $\theta < \theta_0$ ,

$$\left\{x \in B(0,R) : d(x,(\overline{\mathcal{D}(A_1)})^c) \geqslant \theta\right\} \neq \emptyset,$$

where  $B(0,R) := \{x \in \mathbb{R}^n : |x| \leq R\}, d(\cdot,\cdot)$  is the Euclidean distance in  $\mathbb{R}^n$  and  $(\overline{\mathcal{D}(A_1)})^c$  denotes the complement of  $\overline{\mathcal{D}(A_1)}$ . Set

$$g_R(\theta) := \sup \left\{ |z| : z \in A_1(x) \text{ for all } x \in B(0,R) \text{ with } d\left(x, (\overline{\mathcal{D}(A_1)})^c\right) \geqslant \theta \right\},$$

and by the local boundedness of  $A_1$  on  $Int(\mathcal{D}(A_1))$ , it holds that

$$g_R(\theta) < +\infty$$
.

Again put for any l > 0

$$h_R(l) := \inf \left\{ \theta \in (0, \theta_0) : g_R(\theta) \leqslant l^{-1/2} \right\},\,$$

and we have that

$$g_R(l + h_R(l)) \le l^{-1/2} \text{ and } \lim_{l \downarrow 0} h_R(l) = 0.$$

Take  $l_R > 0$  be such that  $l_R + h_R(l_R) < \theta_0$ . For  $0 < l < l_R \wedge 1$ , let  $X_s^{\varepsilon,\gamma,l,R}$  be the projection of  $X_s^{\varepsilon,\gamma}$  on  $\left\{x \in B(0,R): d\left(x,(\overline{\mathcal{D}(A_1)})^c\right) \geqslant l + h_R(l)\right\}$ . Thus, for  $Z_s^{\varepsilon,\gamma,l,R} \in A_1(X_s^{\varepsilon,\gamma,l,R})$ ,  $\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma}| \leqslant R$  and 0 < t - s < l, it holds that

$$-2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}, dK_{r}^{1,\varepsilon,\gamma} \right\rangle$$

$$= -2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma,l,R}, dK_{r}^{1,\varepsilon,\gamma} \right\rangle - 2\int_{s}^{t} \left\langle X_{s}^{\varepsilon,\gamma,l,R} - X_{s}^{\varepsilon,\gamma}, dK_{r}^{1,\varepsilon,\gamma} \right\rangle$$

$$\leq -2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma,l,R}, Z_{s}^{\varepsilon,\gamma,l,R} \right\rangle dr + 2\left(l + h_{R}(l)\right) \left| K^{1,\varepsilon,\gamma} \right|_{0}^{T}$$

$$\leq 4l^{1/2}R + 2\left(l + h_{R}(l)\right) \left| K^{1,\varepsilon,\gamma} \right|_{0}^{T},$$

and furthermore by (32)

$$\sup_{s\leqslant t\leqslant s+l} |X_t^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 I_{\{\sup_{t\in[0,T]} |X_t^{\varepsilon,\gamma}|\leqslant R\}}$$

$$\leqslant \left(4l^{1/2}R + 2\left(l + h_R(l)\right) \left| K^{1,\varepsilon,\gamma} \right|_0^T \right) + \sup_{s\leqslant t\leqslant s+l} \int_s^t |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \mathrm{d}r$$

$$+ \bar{L}_{b_1,\sigma_1} \sup_{s\leqslant t\leqslant s+l} \int_s^t \left(1 + |X_r^{\varepsilon,\gamma}|^2 + |Y_r^{\varepsilon,\gamma}|^2 \right) \mathrm{d}r$$

$$+ C \sup_{s\leqslant t\leqslant s+l} \left| \int_s^t \left\langle X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}, \sigma\left(X_r^{\varepsilon,\gamma}, Y_r^{\varepsilon,\gamma}\right) \right\rangle \mathrm{d}W_r^1 \right| I_{\{\sup_{t\in[0,T]} |X_t^{\varepsilon,\gamma}|\leqslant R\}}.$$

From the above deduction and the BDG inequality, it follows that

$$\mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 I_{\{\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma}| \leqslant R\}}$$

$$\leqslant \left(4l^{1/2}R + 2\left(l + h_R(l)\right) \mathbb{E}|K^{1,\varepsilon,\gamma}|_0^T\right) + C(1 + |x_0|^2 + |y_0|^2)l$$

$$\begin{split} + C \mathbb{E} \left( \int_{s}^{s+l} |X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}|^{2} \|\sigma(X_{r}^{\varepsilon,\gamma}, Y_{r}^{\varepsilon,\gamma})\|^{2} \mathrm{d}r \right)^{1/2} I_{\{\sup_{t \in [0,T]} |X_{t}^{\varepsilon,\gamma}| \leqslant R\}} \\ \leqslant & \left( 4l^{1/2}R + 2 \left( l + h_{R}(l) \right) \mathbb{E} |K^{1,\varepsilon,\gamma}|_{0}^{T} \right) + C(1 + |x_{0}|^{2} + |y_{0}|^{2}) l \\ & + \frac{1}{2} \mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_{t}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}|^{2} I_{\{\sup_{t \in [0,T]} |X_{t}^{\varepsilon,\gamma}| \leqslant R\}} + C(1 + |x_{0}|^{2} + |y_{0}|^{2}) l, \end{split}$$

which yields that

$$\mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 I_{\{\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma}| \leqslant R\}}$$

$$\leqslant (4l^{1/2}R + 2(l+h_R(l)) \mathbb{E} |K^{1,\varepsilon,\gamma}|_0^T) + C(1+|x_0|^2 + |y_0|^2)l.$$

Based on this, letting  $l \to 0$  and then  $R \to \infty$ , one could conclude (30), which completes the proof.

**Lemma 5.4.** Suppose that the assumptions of Theorem 3.1 hold. Then there exists a constant C > 0 such that

$$\sup_{t \in [0,T]} \mathbb{E}|Y_t^{\varepsilon,\gamma} - \hat{Y}_t^{\varepsilon,\gamma}|^2 \leqslant \frac{C}{\alpha} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s+\delta} |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \right). \tag{33}$$

*Proof.* First of all, by (9) and (19), we have that

$$Y_{t}^{\varepsilon,\gamma} - \hat{Y}_{t}^{\varepsilon,\gamma} = -K_{t}^{2,\varepsilon,\gamma} + \hat{K}_{t}^{2,\varepsilon,\gamma} + \frac{1}{\gamma} \int_{0}^{t} \left( b_{2}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) - b_{2}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) \right) \mathrm{d}s$$

$$+ \frac{1}{\sqrt{\gamma}} \int_{0}^{t} \left( \sigma_{2}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) - \sigma_{2}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) \right) \mathrm{d}W_{s}^{2}.$$

Applying the Itô formula to  $|Y_t^{\varepsilon,\gamma} - \hat{Y}_t^{\varepsilon,\gamma}|^2 e^{\lambda t}$  for  $\lambda = \frac{\alpha}{\gamma}$  and taking the expectation, by  $(\mathbf{H}_{b_2,\sigma_2}^1)$  and  $(\mathbf{H}_{b_2,\sigma_2}^2)$  one could obtain that

$$\begin{split} \mathbb{E}|Y_t^{\varepsilon,\gamma} - \hat{Y}_t^{\varepsilon,\gamma}|^2 e^{\lambda t} &= \lambda \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}|^2 e^{\lambda s} \mathrm{d}s - \mathbb{E} \int_0^t 2e^{\lambda s} \langle Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}, \mathrm{d}(K_s^{2,\varepsilon,\gamma} - \hat{K}_s^{2,\varepsilon,\gamma}) \rangle \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t 2e^{\lambda s} \langle Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}, b_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) - b_2(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \rangle \mathrm{d}s \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} \|\sigma_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) - \sigma_2(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma})\|^2 \mathrm{d}s \\ &\leqslant \lambda \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}|^2 e^{\lambda s} \mathrm{d}s + \frac{1}{\gamma} \mathbb{E} \int_0^t 2e^{\lambda s} \langle Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}, b_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) \\ &- b_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \rangle \mathrm{d}s + \frac{1}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} \|\sigma_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) - \sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \|^2 \mathrm{d}s \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t 2e^{\lambda s} \langle Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}, b_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) - b_2(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \rangle \mathrm{d}s \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} \|\sigma_2(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) - \sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \|^2 \mathrm{d}s \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t 2e^{\lambda s} \|\sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) - \sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \|^2 \mathrm{d}s \\ &+ \frac{1}{\gamma} \mathbb{E} \int_0^t 2e^{\lambda s} \|\sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) - \sigma_2(X_s^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \|^2 \mathrm{d}s \end{split}$$

$$\leqslant (\lambda - \frac{\beta}{\gamma} + \frac{2L_{b_2,\sigma_2}}{\gamma}) \mathbb{E} \int_0^t |Y_s^{\varepsilon,\gamma} - \hat{Y}_s^{\varepsilon,\gamma}|^2 e^{\lambda s} ds 
+ \frac{C}{\gamma} \mathbb{E} \int_0^t e^{\lambda s} |X_s^{\varepsilon,\gamma} - X_{s(\delta)}^{\varepsilon,\gamma}|^2 ds 
\leqslant \frac{C}{\gamma} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \right) \frac{e^{\lambda t} - 1}{\lambda}.$$

Finally, it follows that

$$\mathbb{E}|Y_t^{\varepsilon,\gamma} - \hat{Y}_t^{\varepsilon,\gamma}|^2 \leqslant \frac{C}{\alpha} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \right).$$

The proof is complete.

Finally, we estimate  $\mathbb{E} \sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma} - \hat{X}_t^{\varepsilon,\gamma}|^2$ .

**Lemma 5.5.** Assume that the assumptions of Theorem 3.1 hold. Then we have that

$$\mathbb{E} \sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma} - \hat{X}_t^{\varepsilon,\gamma}|^2 \leqslant C \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \right). \tag{34}$$

*Proof.* Note that

$$X_t^{\varepsilon,\gamma} - \hat{X}_t^{\varepsilon,\gamma} = -K_t^{1,\varepsilon,\gamma} + \hat{K}_t^{1,\varepsilon,\gamma} + \int_0^t \left( b_1(X_s^{\varepsilon,\gamma}, Y_s^{\varepsilon,\gamma}) - b_1(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_s^{\varepsilon,\gamma}) \right) \mathrm{d}s.$$

Thus, by the Taylor formula, it holds that

$$|X_{t}^{\varepsilon,\gamma} - \hat{X}_{t}^{\varepsilon,\gamma}|^{2} = -2 \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma} - \hat{X}_{s}^{\varepsilon,\gamma}, d(K_{s}^{1,\varepsilon,\gamma} - \hat{K}_{s}^{1,\varepsilon,\gamma}) \rangle$$

$$+2 \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma} - \hat{X}_{s}^{\varepsilon,\gamma}, b_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) - b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) \rangle ds$$

$$\leqslant \int_{0}^{t} |X_{s}^{\varepsilon,\gamma} - \hat{X}_{s}^{\varepsilon,\gamma}|^{2} ds + L_{b_{1},\sigma_{1}} \int_{0}^{t} (|X_{s}^{\varepsilon,\gamma} - X_{s(\delta)}^{\varepsilon,\gamma}|^{2} + |Y_{s}^{\varepsilon,\gamma} - \hat{Y}_{s}^{\varepsilon,\gamma}|^{2}) ds,$$

and

$$\mathbb{E}\sup_{s\in[0,t]}|X_s^{\varepsilon,\gamma}-\hat{X}_s^{\varepsilon,\gamma}|^2\leqslant \int_0^t\mathbb{E}\sup_{s\in[0,r]}|X_s^{\varepsilon,\gamma}-\hat{X}_s^{\varepsilon,\gamma}|^2\mathrm{d}r+TC\left(\sup_{s\in[0,T]}\mathbb{E}\sup_{s\leqslant r\leqslant s+\delta}|X_r^{\varepsilon,\gamma}-X_s^{\varepsilon,\gamma}|^2\right).$$

where we use (33). The Gronwall inequality yields the required result. The proof is completed.  $\hfill\Box$ 

### 5.2. Some estimates for the average equation (11).

**Lemma 5.6.** Under the assumptions of Theorem 3.1, Eq.(11) has a unique solution  $(\bar{X}_{\cdot}, \bar{K}_{\cdot})$ . Moreover, it holds that

$$\sup_{t \in [0,T]} |\bar{X}_t|^2 \leqslant C(1+|x_0|^2), \tag{35}$$

$$\lim_{l \to 0} \sup_{s \in [0,T]} \sup_{s \le t \le s+l} |\bar{X}_t - \bar{X}_s|^2 = 0.$$
 (36)

*Proof.* First of all, from (18), it follows that Eq.(11) has a unique solution  $(\bar{X}, \bar{K})$ . Then, since the proofs of the required estimates are similar to that for  $X^{\varepsilon,\gamma}$  in Lemma 5.1 and Lemma 5.3, we omit them. The proof is complete.

**Lemma 5.7.** Suppose that the assumptions of Theorem 3.1 hold. Then there exists a constant C > 0 such that

$$\mathbb{E}\left(\sup_{0 \le t \le T} |\hat{X}_t^{\varepsilon, \gamma} - \bar{X}_t|^2\right) \leqslant C(\sqrt{\varepsilon} + \varepsilon) + \Sigma(\gamma).$$

where

$$\Sigma(\gamma) := C \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s+\delta} |\hat{X}_r^{\varepsilon,\gamma} - \hat{X}_s^{\varepsilon,\gamma}|^2 + \sup_{s \in [0,T]} \sup_{s \leqslant t \leqslant s+\delta} |\bar{X}_t - \bar{X}_s|^2 \right)^{1/2}$$

$$+ C \left( (\frac{\gamma}{\delta})^{1/2} + \delta^{1/2} \right) + C \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s+\delta} |X_r^{\varepsilon,\gamma} - X_s^{\varepsilon,\gamma}|^2 \right)$$

$$+ C \sup_{s \in [0,T]} \sup_{s \leqslant t \leqslant s+\delta} |\bar{X}_t - \bar{X}_s|^2.$$

*Proof.* Note that

$$\hat{X}_{t}^{\varepsilon,\gamma} - \bar{X}_{t} = -\hat{K}_{t}^{1,\varepsilon,\gamma} + \bar{K}_{t} + \int_{0}^{t} \left( b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(\bar{X}_{s}) \right) ds + \sqrt{\varepsilon} \int_{0}^{t} \sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) dW_{s}^{1}.$$

Thus, by the Itô formula, it holds that

$$|\hat{X}_{t}^{\varepsilon,\gamma} - \bar{X}_{t}|^{2}$$

$$= -2 \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}, \operatorname{d}(\hat{K}_{s}^{1,\varepsilon,\gamma} - \bar{K}_{s}) \rangle + 2 \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(\bar{X}_{s}) \rangle \operatorname{d}s$$

$$+2\sqrt{\varepsilon} \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}, \sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) \operatorname{d}W_{s}^{1} \rangle + \varepsilon \int_{0}^{t} \|\sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma})\|^{2} \operatorname{d}s$$

$$\leq 2 \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(\bar{X}_{s}) \rangle \operatorname{d}s$$

$$+2\sqrt{\varepsilon} \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}, \sigma_{1}(X_{s}^{\varepsilon,\gamma}, Y_{s}^{\varepsilon,\gamma}) \operatorname{d}W_{s}^{1} \rangle + \varepsilon L_{b_{1},\sigma_{1}} \int_{0}^{t} (1 + |X_{s}^{\varepsilon,\gamma}|^{2} + |Y_{s}^{\varepsilon,\gamma}|^{2}) \operatorname{d}s.$$

Moreover, based on the BDG inequality, we get that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|\hat{X}_{t}^{\varepsilon,\gamma}-\bar{X}_{t}|^{2}\right)$$

$$\leqslant 2\mathbb{E}\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\langle\hat{X}_{s}^{\varepsilon,\gamma}-\bar{X}_{s},b_{1}(X_{s(\delta)}^{\varepsilon,\gamma},\hat{Y}_{s}^{\varepsilon,\gamma})-\bar{b}_{1}(\bar{X}_{s})\rangle\mathrm{d}s\right|$$

$$+2\sqrt{\varepsilon}\mathbb{E}\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\langle\hat{X}_{s}^{\varepsilon,\gamma}-\bar{X}_{s},\sigma_{1}(X_{s}^{\varepsilon,\gamma},Y_{s}^{\varepsilon,\gamma})\mathrm{d}W_{s}^{1}\rangle\right|+C\varepsilon$$

$$\leqslant 2\mathbb{E}\sup_{0\leqslant t\leqslant T}\left|\int_{0}^{t}\langle\hat{X}_{s}^{\varepsilon,\gamma}-\bar{X}_{s},b_{1}(X_{s(\delta)}^{\varepsilon,\gamma},\hat{Y}_{s}^{\varepsilon,\gamma})-\bar{b}_{1}(\bar{X}_{s})\rangle\mathrm{d}s\right|$$

$$+2\sqrt{\varepsilon}C\mathbb{E}\left(\int_{0}^{T}|\hat{X}_{s}^{\varepsilon,\gamma}-\bar{X}_{s}|^{2}\|\sigma_{1}(X_{s}^{\varepsilon,\gamma},Y_{s}^{\varepsilon,\gamma})\|^{2}\mathrm{d}s\right)^{1/2}+C\varepsilon$$

$$\leq 2\mathbb{E} \sup_{0 \leq t \leq T} \left| \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon, \gamma} - \bar{X}_{s}, b_{1}(X_{s(\delta)}^{\varepsilon, \gamma}, \hat{Y}_{s}^{\varepsilon, \gamma}) - \bar{b}_{1}(\bar{X}_{s}) \rangle ds \right| 
+ \frac{1}{2} \mathbb{E} \left( \sup_{0 \leq t \leq T} |\hat{X}_{t}^{\varepsilon, \gamma} - \bar{X}_{t}|^{2} \right) + C(\sqrt{\varepsilon} + \varepsilon).$$
(37)

Next, set

$$I := \mathbb{E} \sup_{0 \le t \le T} \left| \int_0^t \langle \hat{X}_s^{\varepsilon, \gamma} - \bar{X}_s, b_1(X_{s(\delta)}^{\varepsilon, \gamma}, \hat{Y}_s^{\varepsilon, \gamma}) - \bar{b}_1(\bar{X}_s) \rangle \mathrm{d}s \right|,$$

and we are devoted to estimating I. Note that

$$I \leqslant \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{t} \langle \hat{X}_{s}^{\varepsilon,\gamma} - \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s} + \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(\bar{X}_{s}) \rangle ds \right|$$

$$+ \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{t} \langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \rangle ds \right|$$

$$+ \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{t} \langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) - \bar{b}_{1}(\bar{X}_{s(\delta)}) \rangle ds \right|$$

$$+ \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{t} \langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, \bar{b}_{1}(\bar{X}_{s(\delta)}) - \bar{b}_{1}(\bar{X}_{s}) \rangle ds \right|$$

$$=: I_{1} + I_{2} + I_{3} + I_{4}.$$

$$(38)$$

Thus, we estimate  $I_1, I_2, I_3, I_4$ , respectively.

For  $I_1$ , by the Hölder inequality and the linear growth for  $b_1, \bar{b}_1$ , it holds that

$$I_{1} \leqslant C \left( \int_{0}^{T} \mathbb{E}(|\hat{X}_{s}^{\varepsilon,\gamma} - \hat{X}_{s(\delta)}^{\varepsilon,\gamma}|^{2} + |\bar{X}_{s} - \bar{X}_{s(\delta)}|^{2}) ds \right)^{1/2}$$

$$\times \left( \int_{0}^{T} \mathbb{E}(1 + |X_{s(\delta)}^{\varepsilon,\gamma}|^{2} + |\hat{Y}_{s}^{\varepsilon,\gamma}|^{2} + |\bar{X}_{s}|^{2}) ds \right)^{1/2}$$

$$\leqslant C \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |\hat{X}_{r}^{\varepsilon,\gamma} - \hat{X}_{s}^{\varepsilon,\gamma}|^{2} + \sup_{s \in [0,T]} \sup_{s \leqslant t \leqslant s + \delta} |\bar{X}_{t} - \bar{X}_{s}|^{2} \right)^{1/2} .$$
 (39)

For  $I_2$ , by the deduction in **Step 2.**, we know that

$$I_2 \leqslant C\left(\left(\frac{\gamma}{\delta}\right)^{1/2} + \delta^{1/2}\right). \tag{40}$$

For  $I_3$ , the Lipschitz continuity of  $\bar{b}_1$  implies that

$$I_{3} \leqslant C\mathbb{E} \int_{0}^{T} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|(|X_{s(\delta)}^{\varepsilon,\gamma} - \hat{X}_{s(\delta)}^{\varepsilon,\gamma}| + |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|) ds$$

$$\leqslant C \int_{0}^{T} \mathbb{E} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|^{2} ds + C \int_{0}^{T} \mathbb{E} |X_{s(\delta)}^{\varepsilon,\gamma} - \hat{X}_{s(\delta)}^{\varepsilon,\gamma}|^{2} ds$$

$$\leqslant C \int_{0}^{T} \mathbb{E} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|^{2} ds + C \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |X_{r}^{\varepsilon,\gamma} - X_{s}^{\varepsilon,\gamma}|^{2} \right). \tag{41}$$

For  $I_4$ , by the Lipschitz continuity of  $\bar{b}_1$  and (36), it holds that

$$I_{4} \leqslant C \int_{0}^{T} \mathbb{E} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|^{2} ds + C \int_{0}^{T} |\bar{X}_{s} - \bar{X}_{s(\delta)}|^{2} ds$$

$$\leqslant C \int_{0}^{T} \mathbb{E} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|^{2} ds + C \sup_{s \in [0,T]} \sup_{s \leqslant t \leqslant s + \delta} |\bar{X}_{t} - \bar{X}_{s}|^{2}. \tag{42}$$

Combining (39)-(42) with (38), we have that

$$I \leqslant \Sigma(\gamma) + C \int_0^T \mathbb{E} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}|^2 ds,$$

which together with (37) implies that

$$\mathbb{E}\Big(\sup_{0\leqslant t\leqslant T}|\hat{X}_t^{\varepsilon,\gamma}-\bar{X}_t|^2\Big)\leqslant C(\sqrt{\varepsilon}+\varepsilon)+\Sigma(\gamma)+C\int_0^T\mathbb{E}\Big(\sup_{0\leqslant r\leqslant s}|\hat{X}_r^{\varepsilon,\gamma}-\bar{X}_r|^2\Big)\mathrm{d}s.$$

Finally, the Gronwall inequality yields the required estimate.

Step 2. We prove (40).

For  $I_2$ , it holds that

$$I_{2} = \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{\left[\frac{t}{\delta}\right]\delta} \left\langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \right\rangle ds \right.$$

$$\left. + \int_{\left[\frac{t}{\delta}\right]\delta}^{t} \left\langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \right\rangle ds \right| \right)$$

$$\leqslant \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{\left[\frac{t}{\delta}\right]\delta} \left\langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \right\rangle ds \right| \right)$$

$$+ \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{\left[\frac{t}{\delta}\right]\delta}^{t} \left\langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \right\rangle ds \right| \right)$$

$$=: I_{21} + I_{22}$$

Next, we estimate  $I_{21}$ . Note that

$$\begin{split} I_{21} &= \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \left| \int_{0}^{[\frac{t}{\delta}]\delta} \langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \rangle \mathrm{d}s \right| \right) \\ &= \mathbb{E} \left( \sup_{0 \leqslant t \leqslant T} \left| \sum_{k=0}^{[\frac{t}{\delta}]-1} \int_{k\delta}^{(k+1)\delta} \langle \hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}, b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma}) \rangle \mathrm{d}s \right| \right) \\ &\leqslant \mathbb{E} \left( \sup_{0 \leqslant t \leqslant T} \sum_{k=0}^{[\frac{t}{\delta}]-1} \left| \int_{k\delta}^{(k+1)\delta} \langle \hat{X}_{k\delta}^{\varepsilon,\gamma} - \bar{X}_{k\delta}, b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}) \rangle \mathrm{d}s \right| \right. \\ &\leqslant \sum_{k=0}^{[\frac{T}{\delta}]-1} \mathbb{E} \left( \left| \int_{k\delta}^{(k+1)\delta} \langle \hat{X}_{k\delta}^{\varepsilon,\gamma} - \bar{X}_{k\delta}, b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}) \rangle \mathrm{d}s \right| \right) \\ &\leqslant \left[ \frac{T}{\delta} \right] \sup_{0 \leqslant k \leqslant [\frac{T}{\delta}]-1} \mathbb{E} \left( \left| \int_{k\delta}^{(k+1)\delta} \langle \hat{X}_{k\delta}^{\varepsilon,\gamma} - \bar{X}_{k\delta}, b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}) \rangle \mathrm{d}s \right| \right) \end{aligned}$$

$$\leqslant \gamma(\frac{T}{\delta}) \sup_{0 \leqslant k \leqslant [\frac{T}{\delta}] - 1} \mathbb{E}\left(\left|\langle \hat{X}_{k\delta}^{\varepsilon, \gamma} - \bar{X}_{k\delta}, \int_{0}^{\delta/\gamma} (b_{1}(X_{k\delta}^{\varepsilon, \gamma}, \hat{Y}_{\gamma s + k\delta}^{\varepsilon, \gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon, \gamma})) ds\rangle\right|\right)$$

$$\leqslant \gamma(\frac{T}{\delta}) \sup_{0 \leqslant k \leqslant [\frac{T}{\delta}] - 1} \mathbb{E}|\hat{X}_{k\delta}^{\varepsilon, \gamma} - \bar{X}_{k\delta}| \left|\int_{0}^{\delta/\gamma} (b_{1}(X_{k\delta}^{\varepsilon, \gamma}, \hat{Y}_{\gamma s + k\delta}^{\varepsilon, \gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon, \gamma})) ds\right|$$

$$\leqslant \gamma(\frac{T}{\delta}) \sup_{0 \leqslant k \leqslant [\frac{T}{\delta}] - 1} (\mathbb{E}|\hat{X}_{k\delta}^{\varepsilon, \gamma} - \bar{X}_{k\delta}|^{2})^{1/2}$$

$$\times \left(\mathbb{E}\left|\int_{0}^{\delta/\gamma} (b_{1}(X_{k\delta}^{\varepsilon, \gamma}, \hat{Y}_{\gamma s + k\delta}^{\varepsilon, \gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon, \gamma})) ds\right|^{2}\right)^{1/2}.$$

$$(43)$$

Thus, we compute  $\mathbb{E}\left|\int_0^{\delta/\gamma} (b_1(X_{k\delta}^{\varepsilon,\gamma},\hat{Y}_{\gamma s+k\delta}^{\varepsilon,\gamma}) - \bar{b}_1(X_{k\delta}^{\varepsilon,\gamma})) ds\right|^2$ . It is easy to see that

$$\mathbb{E} \left| \int_{0}^{\delta/\gamma} (b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{\gamma s+k\delta}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma})) ds \right|^{2} \\
= 2\mathbb{E} \int_{0}^{\delta/\gamma} \int_{r}^{\delta/\gamma} \langle b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{\gamma s+k\delta}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}), b_{1}(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{\gamma r+k\delta}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}) \rangle ds dr.$$

Therefore, set for  $0 < r < s \le \delta/\gamma$ 

$$\Phi(s,r) := \mathbb{E}\langle b_1(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{\gamma s + k\delta}^{\varepsilon,\gamma}) - \bar{b}_1(X_{k\delta}^{\varepsilon,\gamma}), b_1(X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{\gamma r + k\delta}^{\varepsilon,\gamma}) - \bar{b}_1(X_{k\delta}^{\varepsilon,\gamma})\rangle,$$

and we only investigate  $\Phi(s, r)$ .

For any s>0 and random variables  $\xi,\zeta\in L^2(\Omega,\mathscr{F}_s,\mathbb{P}),\xi\in\overline{\mathcal{D}(A_1)},\zeta\in\overline{\mathcal{D}(A_2)}$ , we consider the following equation

ler the following equation 
$$\begin{cases} \mathrm{d} \check{Y}_t^{\gamma,s,\xi,\zeta} \in -A_2(\check{Y}_t^{\gamma,s,\xi,\zeta}) \mathrm{d}t + \frac{1}{\gamma} b_2(\xi,\check{Y}_t^{\gamma,s,\xi,\zeta}) \mathrm{d}t + \frac{1}{\sqrt{\gamma}} \sigma_2(\xi,\check{Y}_t^{\gamma,s,\xi,\zeta}) \mathrm{d}W_t^2, \\ \check{Y}_s^{\gamma,s,\xi,\zeta} = \zeta. \end{cases}$$

Then it holds that

$$\hat{Y}^{\varepsilon,\gamma}_t = \check{Y}^{\gamma,k\delta,X^{\varepsilon,\gamma}_{k\delta},\hat{Y}^{\varepsilon,\gamma}_{k\delta}}_t, \quad \hat{K}^{2,\varepsilon,\gamma}_t = \check{K}^{2,\gamma,k\delta,X^{\varepsilon,\gamma}_{k\delta},\hat{Y}^{\varepsilon,\gamma}_{k\delta}}_t, \quad t \in [k\delta,(k+1)\delta],$$

and furthermore

$$\Phi(s,r) = \mathbb{E}\langle b_1(X_{k\delta}^{\varepsilon,\gamma}, \check{Y}_{\gamma s + k\delta}^{\gamma,k\delta, X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{k\delta}^{\varepsilon,\gamma}}) - \bar{b}_1(X_{k\delta}^{\varepsilon,\gamma}), b_1(X_{k\delta}^{\varepsilon,\gamma}, \check{Y}_{\gamma r + k\delta}^{\gamma,k\delta, X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{k\delta}^{\varepsilon,\gamma}}) - \bar{b}_1(X_{k\delta}^{\varepsilon,\gamma}) \rangle.$$

Note that  $X_{k\delta}^{\varepsilon,\gamma}, \hat{Y}_{k\delta}^{\varepsilon,\gamma}$  are  $\mathscr{F}_{k\delta}$ -measurable, and for any  $x \in \overline{\mathcal{D}(A_1)}, y \in \overline{\mathcal{D}(A_2)}, \check{Y}_t^{\gamma,k\delta,x,y}$  is independent of  $\mathscr{F}_{k\delta}$ . Thus, we have that

$$\Phi(s,r) = \mathbb{E}\left[\mathbb{E}\left[\langle b_{1}(X_{k\delta}^{\varepsilon,\gamma},\check{Y}_{\gamma s+k\delta}^{\gamma,k\delta,X_{k\delta}^{\varepsilon,\gamma},\hat{Y}_{k\delta}^{\varepsilon,\gamma}}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma}), \right. \\
\left. b_{1}(X_{k\delta}^{\varepsilon,\gamma},\check{Y}_{\gamma r+k\delta}^{\gamma,k\delta,X_{k\delta}^{\varepsilon,\gamma},\hat{Y}_{k\delta}^{\varepsilon,\gamma}}) - \bar{b}_{1}(X_{k\delta}^{\varepsilon,\gamma})\rangle\right|\mathscr{F}_{k\delta}\right]\right] \\
= \mathbb{E}\left[\mathbb{E}\left[\langle b_{1}(x,\check{Y}_{\gamma s+k\delta}^{\gamma,k\delta,x,y}) - \bar{b}_{1}(x), b_{1}(x,\check{Y}_{\gamma r+k\delta}^{\gamma,k\delta,x,y}) - \bar{b}_{1}(x)\rangle\right]\right|_{(x,y)=(X_{k\delta}^{\varepsilon,\gamma},\hat{Y}_{k\delta}^{\varepsilon,\gamma},\hat{Y}_{k\delta}^{\varepsilon,\gamma})}\right].$$

Here, we investigate  $\check{Y}_{\gamma s+k\delta}^{\gamma,k\delta,x,y}$ . On one hand, it holds that

$$\check{Y}_{\gamma s+k\delta}^{\gamma,k\delta,x,y} = y - \check{K}_{\gamma s+k\delta}^{2,\gamma,k\delta,x,y} + \check{K}_{k\delta}^{2,\gamma,k\delta,x,y} + \frac{1}{\gamma} \int_{k\delta}^{\gamma s+k\delta} b_2(x, \check{Y}_r^{\varepsilon,k\delta,x,y}) dr 
+ \frac{1}{\sqrt{\gamma}} \int_{k\delta}^{\gamma s+k\delta} \sigma_2(x, \check{Y}_r^{\gamma,k\delta,x,y}) dW_r^2 
= y - \check{K}_{\gamma s+k\delta}^{2,\gamma,k\delta,x,y} + \check{K}_{k\delta}^{2,\gamma,k\delta,x,y} + \frac{1}{\gamma} \int_0^{\gamma s} b_2(x, \check{Y}_{u+k\delta}^{\gamma,k\delta,x,y}) du 
+ \frac{1}{\sqrt{\gamma}} \int_0^{\gamma s} \sigma_2(x, \check{Y}_{u+k\delta}^{\gamma,k\delta,x,y}) d\tilde{W}_u^2 
= y - \check{K}_s^{2,\gamma,k\delta,x,y} + \int_0^s b_2(x, \check{Y}_{\gamma v+k\delta}^{\gamma,k\delta,x,y}) dv + \int_0^s \sigma_2(x, \check{Y}_{\gamma v+k\delta}^{\gamma,k\delta,x,y}) d\check{W}_v^2,$$

where  $\tilde{W}_u^2 := W_{u+k\delta}^2 - W_{k\delta}^2$  and  $\tilde{\tilde{W}}_v^2 := \frac{1}{\sqrt{\gamma}} \tilde{W}_{\gamma v}^2$  are two m-dimensional standard Brownian motions, and  $\tilde{\check{K}}_s^{2,\gamma,k\delta,x,y} := \check{K}_{\gamma s+k\delta}^{2,\gamma,k\delta,x,y} - \check{K}_{k\delta}^{2,\gamma,k\delta,x,y}$ . On the other hand, note that the frozen equation (10) is written as

$$Y_s^{x,y} = y - K_s^{2,x,y} + \int_0^s b_2(x, Y_r^{x,y}) dr + \int_0^s \sigma_2(x, Y_r^{x,y}) dW_r^2.$$

Thus, for  $s \in [0, \delta/\gamma]$ ,  $\check{Y}_{\gamma s + k\delta}^{\gamma, k\delta, x, y}$  and  $Y_s^{x, y}$  have the same distribution, which implies that

$$\mathbb{E}\left[\langle b_{1}(x, \check{Y}_{\gamma s+k\delta}^{\gamma,k\delta,x,y}) - \bar{b}_{1}(x), b_{1}(x, \check{Y}_{\gamma r+k\delta}^{\gamma,k\delta,x,y}) - \bar{b}_{1}(x)\rangle\right]$$

$$= \mathbb{E}\langle b_{1}(x, Y_{s}^{x,y}) - \bar{b}_{1}(x), b_{1}(x, Y_{r}^{x,y}) - \bar{b}_{1}(x)\rangle$$

$$= \mathbb{E}\left[\mathbb{E}\left[\langle b_{1}(x, Y_{s}^{x,y}) - \bar{b}_{1}(x), b_{1}(x, Y_{r}^{x,y}) - \bar{b}_{1}(x)\rangle\right] \mathscr{F}_{r}^{W^{2}}\right]$$

$$= \mathbb{E}\left[\langle \mathbb{E}\left[b_{1}(x, Y_{s}^{x,y}) \middle| \mathscr{F}_{r}^{W^{2}}\right] - \bar{b}_{1}(x), b_{1}(x, Y_{r}^{x,y}) - \bar{b}_{1}(x)\rangle\right]$$

$$\leqslant \left(\mathbb{E}\left[\mathbb{E}\left[b_{1}(x, Y_{s}^{x,y}) \middle| \mathscr{F}_{r}^{W^{2}}\right] - \bar{b}_{1}(x)\right]^{2}\right)^{1/2} \left(\mathbb{E}|b_{1}(x, Y_{r}^{x,y}) - \bar{b}_{1}(x)|^{2}\right)^{1/2},$$

where  $\mathscr{F}_r^{W^2} = \sigma\{W_u^2, 0 \leqslant u \leqslant r\} \vee \mathcal{N}$ , and  $\mathcal{N}$  denotes the collection of all  $\mathbb{P}$ -zero sets. Moreover, based on (17), we obtain that

$$\left(\mathbb{E}\left|\mathbb{E}\left[b_{1}(x, Y_{s}^{x,y})\middle|\mathscr{F}_{r}^{W^{2}}\right] - \bar{b}_{1}(x)\middle|^{2}\right)^{1/2} \\
= \left(\mathbb{E}\left|\mathbb{E}\left[b_{1}(x, Y_{s-r}^{x,\hat{y}})\right]\middle|_{\hat{y}=Y_{r}^{x,y}} - \bar{b}_{1}(x)\middle|^{2}\right)^{1/2} \leqslant \left(Ce^{-\alpha(s-r)}(1+|x|^{2}+\mathbb{E}|Y_{r}^{x,y}|^{2})\right)^{\frac{1}{2}} \\
\leqslant \left(Ce^{-\alpha(s-r)}(1+|x|^{2}+|y|^{2})\right)^{\frac{1}{2}} \leqslant Ce^{-\alpha(s-r)/2}(1+|x|+|y|),$$

and

$$\begin{split}
\left(\mathbb{E}|b_{1}(x,Y_{r}^{x,y}) - \bar{b}_{1}(x)|^{2}\right)^{1/2} &= \left(\mathbb{E}|b_{1}(x,Y_{r}^{x,y}) - \int_{\overline{\mathcal{D}(A_{2})}} b_{1}(x,u)\nu^{x}(\mathrm{d}u)|^{2}\right)^{1/2} \\
&\leqslant \left(\mathbb{E}\int_{\overline{\mathcal{D}(A_{2})}} |b_{1}(x,Y_{r}^{x,y}) - b_{1}(x,u)|^{2}\nu^{x}(\mathrm{d}u)\right)^{1/2} \\
&\leqslant C\left(\int_{\overline{\mathcal{D}(A_{2})}} \mathbb{E}|Y_{r}^{x,y} - u|^{2}\nu^{x}(\mathrm{d}u)\right)^{1/2} \\
&\leqslant C\left(|y|^{2}e^{-\frac{\alpha}{2}t} + C(1+|x|^{2})\right)^{1/2} \\
&\leqslant C(1+|x|+|y|).
\end{split}$$

Combining the above deduction, by (21) (27) one can have that

$$\Phi(s,r) \leqslant Ce^{-\alpha(s-r)/2}$$
.

Inserting the above inequality in (43), we get that

$$I_{21} \leqslant C\gamma(\frac{T}{\delta}) \sup_{0 \leqslant k \leqslant \lceil \frac{T}{\delta} \rceil - 1} \left( \int_0^{\frac{\delta}{\gamma}} \int_r^{\frac{\delta}{\gamma}} Ce^{-\alpha(s-r)/2} ds dr \right)^{1/2} \leqslant C(\frac{\gamma}{\delta})^{1/2}. \tag{44}$$

Finally, we estimate  $I_{22}$ . By (6) (21) (27) (28) (35) and the Hölder inequality, one could get that

$$\begin{split} I_{22} &\leqslant 2 \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \int_{[\frac{t}{\delta}]\delta}^{t} |\hat{X}_{s(\delta)}^{\varepsilon,\gamma} - \bar{X}_{s(\delta)}| |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma})| \mathrm{d}s \right) \\ &\leqslant 2\delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |\hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}|^{2} \right)^{1/2} \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \int_{[\frac{t}{\delta}]\delta}^{t} |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma})|^{2} \mathrm{d}s \right)^{1/2} \\ &\leqslant 2\delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |\hat{X}_{s}^{\varepsilon,\gamma} - \bar{X}_{s}|^{2} \right)^{1/2} \left( \mathbb{E} \int_{0}^{T} |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma}, \hat{Y}_{s}^{\varepsilon,\gamma}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma})|^{2} \mathrm{d}s \right)^{1/2} \\ &\leqslant C\delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |\hat{X}_{s}^{\varepsilon,\gamma}|^{2} + \sup_{0 \leqslant s \leqslant T} |\bar{X}_{s}|^{2} \right)^{1/2} \left( \int_{0}^{T} (1 + \mathbb{E}|X_{s(\delta)}^{\varepsilon,\gamma}|^{2} + \mathbb{E}|\hat{Y}_{s}^{\varepsilon,\gamma}|^{2} \mathrm{d}s \right)^{1/2} \\ &\leqslant C\delta^{1/2}, \end{split}$$

which together with (44) implies (40). The proof is complete.

At present, we are ready to prove Theorem 3.1.

**Proof of Theorem 3.1.** By (34) and Lemma 5.7, we get that

$$\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_t^{\varepsilon,\gamma}-\bar{X}_t|^2\right) \leqslant 2\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|X_t^{\varepsilon,\gamma}-\hat{X}_t^{\varepsilon,\gamma}|^2\right) + 2\mathbb{E}\left(\sup_{0\leqslant t\leqslant T}|\hat{X}_t^{\varepsilon,\gamma}-\bar{X}_t|^2\right)$$

$$\leqslant C\left(\sup_{s\in[0,T]}\mathbb{E}\sup_{s\leqslant r\leqslant s+\delta}|X_r^{\varepsilon,\gamma}-X_s^{\varepsilon,\gamma}|^2\right) + C(\sqrt{\varepsilon}+\varepsilon) + \Sigma(\gamma).$$

Let us take  $\delta = \gamma^{\iota}$  for  $0 < \iota < 1$ . Thus, as  $\varepsilon$  tends to  $0, \gamma \to 0, \delta \to 0, \gamma/\delta \to 0$  and (30), (31), (36) imply the required limit, which completes the proof.

## 6. Proof of Theorem 3.3

In this section, we prove Theorem 3.3. Moreover, we remind that  $\sigma_1$  is independent of y in this section.

By Lemma 4.1, we know that the system (9) has a unique strong solution  $(X_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{1,\varepsilon,\gamma}, Y_{\cdot}^{\varepsilon,\gamma}, K_{\cdot}^{2,\varepsilon,\gamma})$ . Thus, there exists a functional  $\Psi^{\varepsilon}: C([0,T]; \mathbb{R}^{d_1+d_2}) \mapsto C([0,T], \overline{\mathcal{D}(A_1)})$  such that

$$X^{\varepsilon,\gamma} = \Psi^{\varepsilon}(\sqrt{\varepsilon}W), \quad W := (W^1, W^2).$$

In order to prove the Laplace principle for  $X^{\varepsilon,\gamma}$ , we will verify Condition 2.8 with  $\mathbb{S} = C([0,T],\overline{\mathcal{D}(A_1)})$ .

First of all, we consider the following controlled processes:

$$\begin{cases}
dX_{t}^{\varepsilon,\gamma,u} \in -A_{1}(X_{t}^{\varepsilon,\gamma,u})dt + b_{1}(X_{t}^{\varepsilon,\gamma,u}, Y_{t}^{\varepsilon,\gamma,u})dt + \sigma_{1}(X_{t}^{\varepsilon,\gamma,u})\pi_{1}u(t)dt \\
+\sqrt{\varepsilon}\sigma_{1}(X_{t}^{\varepsilon,\gamma,u})dW_{t}^{1}, \\
X_{0}^{\varepsilon,\gamma,u} = x_{0} \in \overline{\mathcal{D}}(A_{1}), \quad 0 \leqslant t \leqslant T, \\
dY_{t}^{\varepsilon,\gamma,u} \in -A_{2}(Y_{t}^{\varepsilon,\gamma,u})dt + \frac{1}{\gamma}b_{2}(X_{t}^{\varepsilon,\gamma,u}, Y_{t}^{\varepsilon,\gamma,u})dt + \frac{1}{\sqrt{\gamma\varepsilon}}\sigma_{2}(X_{t}^{\varepsilon,\gamma,u}, Y_{t}^{\varepsilon,\gamma,u})\pi_{2}u(t)dt \\
+\frac{1}{\sqrt{\gamma}}\sigma_{2}(X_{t}^{\varepsilon,\gamma,u}, Y_{t}^{\varepsilon,\gamma,u})dW_{t}^{2}, \\
Y_{0}^{\varepsilon,\gamma,u} = y_{0} \in \overline{\mathcal{D}}(A_{2}), \quad 0 \leqslant t \leqslant T, \quad u \in \mathbf{A}_{2}^{N},
\end{cases} (45)$$

where  $\pi_1: \mathbb{R}^{d_1+d_2} \to \mathbb{R}^{d_1}, \pi_2: \mathbb{R}^{d_1+d_2} \to \mathbb{R}^{d_2}$  are two projection operators. Thus, by the Girsanov theorem, the system (45) have a unique strong solution denoted by  $(X^{\varepsilon,\gamma,u}, K^{1,\varepsilon,\gamma,u}, Y^{\varepsilon,\gamma,u}, K^{2,\varepsilon,\gamma,u})$ . Moreover,  $X^{\varepsilon,\gamma,u} = \Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_0^{\cdot} u(s) ds)$ .

Consider the following multivalued differential equation:

$$\begin{cases}
d\bar{X}_t^u \in A_1(\bar{X}_t^u)dt + \bar{b}_1(\bar{X}_t^u)dt + \sigma_1(\bar{X}_t^u)\pi_1u(t)dt, & u \in \mathbf{A}_2^N, \\
\bar{X}_0^u = x_0 \in \overline{\mathcal{D}(A_1)}.
\end{cases} (46)$$

By (18) and  $(\mathbf{H}_{b_1,\sigma_1}^1)$ , it holds that Eq.(46) has a unique solution  $(\bar{X}^u, \bar{K}^u)$ . Define the measurable map  $\Psi^0: C([0,T]; \mathbb{R}^{d_1+d_2}) \mapsto \mathbb{S}$  by  $\Psi^0(\int_0^{\cdot} u(s) ds) = \bar{X}^u$ , and we verify Condition 2.8 through  $\Psi^{\varepsilon}, \Psi^0$ .

In the following, we divide the concrete deduction into two parts. In the first part (Subsection 6.1), we present some key estimates for  $X^{\varepsilon,\gamma,u}, Y^{\varepsilon,\gamma,u}, \bar{X}^u$ . Then we justify Condition 2.8 in the second part (Subsection 6.2).

## 6.1. Some estimates for $X^{\varepsilon,\gamma,u},Y^{\varepsilon,\gamma,u},\bar{X}^u$ .

**Lemma 6.1.** Under the assumptions of Theorem 3.3, for  $\{u_{\varepsilon}, \varepsilon \in (0,1)\} \subset \mathbf{A}_{2}^{N}$ , there exists a constant C > 0 such that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|X_t^{\varepsilon,\gamma,u_\varepsilon}|^2\right)\leqslant C(1+|x_0|^2+|y_0|^2),\tag{47}$$

$$\int_0^T \mathbb{E}|Y_r^{\varepsilon,\gamma,u_\varepsilon}|^2 \mathrm{d}r \leqslant C(1+|x_0|^2+|y_0|^2),\tag{48}$$

$$\mathbb{E}|K^{1,\varepsilon,\gamma,u_{\varepsilon}}|_{0}^{T} \leqslant C(1+|x_{0}|^{2}+|y_{0}|^{2}). \tag{49}$$

*Proof.* First of all, we estimate  $X_t^{\varepsilon,\gamma,u_{\varepsilon}}$ . Note that  $X_t^{\varepsilon,\gamma,u_{\varepsilon}}$  satisfies the following equation:

$$X_t^{\varepsilon,\gamma,u_{\varepsilon}} = x_0 - K_t^{1,\varepsilon,\gamma,u_{\varepsilon}} + \int_0^t b_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}}) ds + \int_0^t \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_1 u_{\varepsilon}(s) ds$$

$$+\sqrt{\varepsilon}\int_0^t \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\mathrm{d}W_s^1.$$

The Itô formula yields that

$$|X_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} = |x_{0}|^{2} - 2\int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{s}^{1,\varepsilon,\gamma,u_{\varepsilon}} \rangle + 2\int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$+2\int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1}u_{\varepsilon}(s) \rangle ds + 2\sqrt{\varepsilon} \int_{0}^{t} \langle X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{s}^{1} \rangle$$

$$+\varepsilon \int_{0}^{t} \|\sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2} ds.$$

$$(50)$$

Then the BDG inequality, Lemma 2.3 and (6) imply that for any  $u \in A_1(0)$ 

$$\begin{split} & \mathbb{E}\left(\sup_{s\in[0,t]}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\right) \\ \leqslant & \left(|x_{0}|^{2}+2|u|T\right)+(2|u|+1)\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}r+\bar{L}_{b_{1},\sigma_{1}}\mathbb{E}\int_{0}^{t}(1+|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}+|Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2})\mathrm{d}r \\ & +2\mathbb{E}\sup_{s\in[0,t]}\left|\int_{0}^{s}\langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}},\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\pi_{1}u_{\varepsilon}(r)\rangle\mathrm{d}r\right|+2\mathbb{E}\sup_{s\in[0,t]}\left|\int_{0}^{s}\langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}},\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\mathrm{d}W_{r}^{1}\rangle\right| \\ \leqslant & C(|x_{0}|^{2}+1)+C\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}r+C\mathbb{E}\int_{0}^{t}|Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}r+\frac{1}{4}\mathbb{E}\left(\sup_{s\in[0,t]}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\right) \\ & +C\mathbb{E}\left(\int_{0}^{t}\|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}r\right)\left(\int_{0}^{t}|u_{\varepsilon}(r)|^{2}\mathrm{d}r\right)+C\mathbb{E}\left(\int_{0}^{t}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}r\right) \\ \leqslant & C(|x_{0}|^{2}+1)+C\mathbb{E}\int_{0}^{t}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}r+C\mathbb{E}\int_{0}^{t}|Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}r \\ & +\frac{1}{2}\mathbb{E}\left(\sup_{s\in[0,t]}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\right)+C\mathbb{E}\int_{0}^{t}\|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}r, \end{split}$$

and furthermore

$$\mathbb{E}\left(\sup_{s\in[0,t]}|X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2\right) \leqslant C(|x_0|^2+1) + C\int_0^t \mathbb{E}|X_r^{\varepsilon,\gamma,u_{\varepsilon}}|^2 dr + C\int_0^t \mathbb{E}|Y_r^{\varepsilon,\gamma,u_{\varepsilon}}|^2 dr.$$
 (51)

For  $Y_t^{\varepsilon,\gamma,u_{\varepsilon}}$ , fix  $v \in A_2(0)$ . Applying the Itô formula to  $|Y_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 e^{\lambda t}$  for  $\lambda = \frac{\alpha}{3\gamma}$  and taking the expectation, one could obtain that

$$\mathbb{E}|Y_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda t} = |y_{0}|^{2} + \lambda \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda s} ds - 2\mathbb{E} \int_{0}^{t} e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{s}^{2,\varepsilon,\gamma,u_{\varepsilon}} \rangle$$

$$+ \frac{2}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$+ \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} ||\sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}})||^{2} ds$$

$$+ \frac{2}{\sqrt{\gamma \varepsilon}} \mathbb{E} \int_{0}^{t} e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{2} u_{\varepsilon}(s) \rangle ds$$

$$\stackrel{(8)}{\leqslant} |y_{0}|^{2} + \lambda \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} e^{\lambda s} ds + 2\mathbb{E} \int_{0}^{t} e^{\lambda s} |v| |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}| ds 
+ \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} \left( -\alpha |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + C(1 + |X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}) \right) ds 
+ \frac{2}{\sqrt{\gamma \varepsilon}} \mathbb{E} \int_{0}^{t} e^{\lambda s} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}| ||\sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}})|| |u_{\varepsilon}(s)| ds.$$

Note that for the third and last terms of the right side for the above inequality

$$2|v||Y_s^{\varepsilon,\gamma,u_\varepsilon}| \leqslant \frac{\alpha}{3\gamma}|Y_s^{\varepsilon,\gamma,u_\varepsilon}|^2 + C|v|^2,$$

and

$$\frac{2}{\sqrt{\gamma\varepsilon}} |Y_s^{\varepsilon,\gamma,u_{\varepsilon}}| \|\sigma_2(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}}) \| |u_{\varepsilon}(s)| 
\leq \frac{\alpha}{3\gamma} |Y_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \frac{C}{\varepsilon} \|\sigma_2(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}}) \|^2 |u_{\varepsilon}(s)|^2 
\leq \frac{\alpha}{3\gamma} |Y_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \frac{C}{\varepsilon} |u_{\varepsilon}(s)|^2,$$

where  $(\mathbf{H}_{\sigma_2}^3)$  is used. Thus, we have that

$$\mathbb{E}|Y_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda t} \leqslant |y_{0}|^{2} + \left(\lambda + \frac{2\alpha}{3\gamma} - \frac{\alpha}{\gamma}\right)\mathbb{E}\int_{0}^{t}|Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda s}ds + C\int_{0}^{t}e^{\lambda s}ds + C\int_{$$

From this, it follows that

$$\mathbb{E}|Y_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 \leqslant C(|y_0|^2+1) + C\mathbb{E}\left(\sup_{s\in[0,t]}|X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2\right) + \frac{C}{\varepsilon}\mathbb{E}\int_0^t e^{-\lambda(t-s)}|u_{\varepsilon}(s)|^2\mathrm{d}s,$$

and furthermore

$$\int_{0}^{t} \mathbb{E}|Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} dr \leqslant CT(|y_{0}|^{2}+1) + C \int_{0}^{t} \mathbb{E}\left(\sup_{s \in [0,r]} |X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\right) dr$$

$$+ \frac{C}{\varepsilon} \mathbb{E} \int_{0}^{t} \int_{0}^{r} e^{-\lambda(r-s)} |u_{\varepsilon}(s)|^{2} ds dr$$

$$\leqslant CT(|y_{0}|^{2}+1) + C \int_{0}^{t} \mathbb{E}\left(\sup_{s \in [0,r]} |X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\right) dr$$

$$+ C\left(\frac{\gamma}{\varepsilon}\right) \mathbb{E} \int_{0}^{t} |u_{\varepsilon}(s)|^{2} ds$$

$$\leqslant CT(|y_0|^2+1)+C\int_0^t \mathbb{E}\left(\sup_{s\in[0,r]}|X_s^{\varepsilon,\gamma,u_\varepsilon}|^2\right)\mathrm{d}r.$$
 (52)

Inserting (52) in (51), by the Gronwall inequality one can get (47) and (48). Finally, for  $K^{1,\varepsilon,\gamma,u_{\varepsilon}}$ , by (50) and Lemma 2.4, it holds that

$$\begin{split} |X_T^{\varepsilon,\gamma,u_{\varepsilon}}|^2 &= |x_0|^2 - 2\int_0^T \langle X_s^{\varepsilon,\gamma,u_{\varepsilon}}, \mathrm{d} K_s^{1,\varepsilon,\gamma,u_{\varepsilon}} \rangle + 2\int_0^T \langle X_s^{\varepsilon,\gamma,u_{\varepsilon}}, b_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle \mathrm{d}s \\ &+ 2\int_0^T \langle X_s^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_1 u_{\varepsilon}(s) \rangle \mathrm{d}s + 2\sqrt{\varepsilon} \int_0^T \langle X_s^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) \mathrm{d}W_s^1 \rangle \\ &+ \varepsilon \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\|^2 \mathrm{d}s \\ &\leqslant |x_0|^2 - 2M_1 \left| K^{1,\varepsilon,\gamma,u_{\varepsilon}} \right|_0^T + 2M_2 \int_0^T |X_s^{\varepsilon,\gamma,u_{\varepsilon}}| \mathrm{d}s + 2M_3 T + \int_0^T |X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 \mathrm{d}s \\ &+ \int_0^T |b_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}})|^2 \mathrm{d}s + \left(\sup_{s \in [0,T]} |X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2\right) \\ &+ \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\|^2 \mathrm{d}s \int_0^T |u_{\varepsilon}(s)|^2 \mathrm{d}s + 2\sqrt{\varepsilon} \int_0^T \langle X_s^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) \mathrm{d}W_s^1 \rangle \\ &+ \varepsilon \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\|^2 \mathrm{d}s, \end{split}$$

and

$$2M_1 \mathbb{E} \left| K^{1,\varepsilon,\gamma,u_{\varepsilon}} \right|_0^T \leqslant |x_0|^2 + 2(M_2 + M_3)T + (2M_2 + 1) \int_0^T \mathbb{E} |X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 ds$$

$$+ \mathbb{E} \left( \sup_{s \in [0,T]} |X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 \right) + C \int_0^T (1 + \mathbb{E} |X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \mathbb{E} |Y_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2) ds,$$

which together with (47) and (48) yields that

$$\mathbb{E} \left| K^{1,\varepsilon,\gamma,u_{\varepsilon}} \right|_{0}^{T} \leqslant C(1+|x_{0}|^{2}+|y_{0}|^{2}).$$

The proof is complete.

**Lemma 6.2.** Under the assumptions of Theorem 3.3, for  $\{u_{\varepsilon}, \varepsilon \in (0,1)\} \subset \mathbf{A}_{2}^{N}$ , we have that

$$\lim_{l \to 0} \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma,u_{\varepsilon}} - X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 = 0.$$
 (53)

*Proof.* Note that for  $0 \le s < t \le T$ ,

$$X_{t}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}} = -K_{t}^{1,\varepsilon,\gamma,u_{\varepsilon}} + K_{s}^{1,\varepsilon,\gamma,u_{\varepsilon}} + \int_{s}^{t} b_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) dr + \int_{s}^{t} \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1}u_{\varepsilon}(r) dr + \int_{s}^{t} \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{r}^{1}.$$

Thus, the Itô formula and (6) imply that

$$|X_t^{\varepsilon,\gamma,u_{\varepsilon}} - X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2$$

$$= -2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{r}^{1,\varepsilon,\gamma,u_{\varepsilon}} \rangle + 2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle dr$$

$$+2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(r) \rangle dr + 2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{r}^{1} \rangle$$

$$+ \int_{s}^{t} \|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2} dr$$

$$\leq -2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{r}^{1,\varepsilon,\gamma,u_{\varepsilon}} \rangle + \int_{s}^{t} |X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} dr$$

$$+ \bar{L}_{b_{1},\sigma_{1}} \int_{s}^{t} (1 + |X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + |Y_{r}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}) dr + 2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(r) \rangle dr$$

$$+2 \int_{s}^{t} \langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{r}^{1} \rangle. \tag{54}$$

Next, we compute  $-2\int_s^t \langle X_r^{\varepsilon,\gamma,u_\varepsilon} - X_s^{\varepsilon,\gamma,u_\varepsilon}, \mathrm{d} K_r^{1,\varepsilon,\gamma,u_\varepsilon} \rangle$ . Let  $X_s^{\varepsilon,\gamma,u_\varepsilon,l,R}$  be the projection of  $X_s^{\varepsilon,\gamma,u_\varepsilon}$  on  $\{x \in B(0,R): d(x,(\overline{\mathcal{D}(A_1)})^c) \geqslant l+h_R(l)\}$  (See the deduction in Lemma 5.3 for  $l,h_R$ ). Thus, for  $Z_s^{\varepsilon,\gamma,u_\varepsilon,l,R} \in A_1(X_s^{\varepsilon,\gamma,u_\varepsilon,l,R})$ ,  $\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma,u_\varepsilon}| \leqslant R$  and 0 < t-s < l, it

holds that

$$-2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{r}^{1,\varepsilon,\gamma,u_{\varepsilon}} \right\rangle$$

$$= -2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon},l,R}, dK_{r}^{1,\varepsilon,\gamma,u_{\varepsilon}} \right\rangle - 2\int_{s}^{t} \left\langle X_{s}^{\varepsilon,\gamma,u_{\varepsilon},l,R} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, dK_{r}^{1,\varepsilon,\gamma,u_{\varepsilon}} \right\rangle$$

$$\leq -2\int_{s}^{t} \left\langle X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon},l,R}, Z_{s}^{\varepsilon,\gamma,u_{\varepsilon},l,R} \right\rangle dr + 2\left(l + h_{R}(l)\right) \left| K^{1,\varepsilon,\gamma,u_{\varepsilon}} \right|_{0}^{T}$$

$$\leq 4l^{1/2}R + 2\left(l + h_{R}(l)\right) \left| K^{1,\varepsilon,\gamma,u_{\varepsilon}} \right|_{0}^{T},$$

and furthermore by (54)

$$\begin{split} \sup_{s\leqslant t\leqslant s+l} &|X^{\varepsilon,\gamma,u_{\varepsilon}}_{t}-X^{\varepsilon,\gamma,u_{\varepsilon}}_{s}|^{2} I_{\{\sup_{t\in[0,T]}|X^{\varepsilon,\gamma,u_{\varepsilon}}_{t}|\leqslant R\}} \\ \leqslant & \left(4l^{1/2}R+2\left(l+h_{R}(l)\right)\left|K^{1,\varepsilon,\gamma,u_{\varepsilon}}\right|^{T}_{0}\right) + \sup_{s\leqslant t\leqslant s+l} \int_{s}^{t} |X^{\varepsilon,\gamma,u_{\varepsilon}}_{r}-X^{\varepsilon,\gamma,u_{\varepsilon}}_{s}|^{2} \mathrm{d}r \\ & + \bar{L}_{b_{1},\sigma_{1}} \sup_{s\leqslant t\leqslant s+l} \int_{s}^{t} \left(1+|X^{\varepsilon,\gamma,u_{\varepsilon}}_{r}|^{2}+|Y^{\varepsilon,\gamma,u_{\varepsilon}}_{r}|^{2}\right) \mathrm{d}r \\ & + 2\sup_{s\leqslant t\leqslant s+l} \left|\int_{s}^{t} \left\langle X^{\varepsilon,\gamma,u_{\varepsilon}}_{r}-X^{\varepsilon,\gamma,u_{\varepsilon}}_{s},\sigma_{1}(X^{\varepsilon,\gamma,u_{\varepsilon}}_{r})\pi_{1}u_{\varepsilon}(r)\right\rangle \mathrm{d}r \right| I_{\{\sup_{t\in[0,T]}|X^{\varepsilon,\gamma,u_{\varepsilon}}_{t}|\leqslant R\}} \\ & + C\sup_{s\leqslant t\leqslant s+l} \left|\int_{s}^{t} \left\langle X^{\varepsilon,\gamma,u_{\varepsilon}}_{r}-X^{\varepsilon,\gamma,u_{\varepsilon}}_{s},\sigma_{1}(X^{\varepsilon,\gamma,u_{\varepsilon}}_{r})\right\rangle \mathrm{d}W^{1}_{r} \right| I_{\{\sup_{t\in[0,T]}|X^{\varepsilon,\gamma,u_{\varepsilon}}_{t}|\leqslant R\}}. \end{split}$$

From the above deduction and the BDG inequality, it follows that

$$\mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma,u_{\varepsilon}} - X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 I_{\{\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma,u_{\varepsilon}}| \leqslant R\}}$$

$$\leqslant \left(4l^{1/2}R + 2\left(l + h_R(l)\right) \mathbb{E} |K^{1,\varepsilon,\gamma,u_{\varepsilon}}|_0^T\right) + C(1 + |x_0|^2 + |y_0|^2)l$$

$$+2\mathbb{E}\int_{s}^{s+l}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}-X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|\|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\||u_{\varepsilon}(r)|\mathrm{d}rI_{\{\sup_{t\in[0,T]}|X_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|\leqslant R\}}$$

$$+C\mathbb{E}\left(\int_{s}^{s+l}|X_{r}^{\varepsilon,\gamma,u_{\varepsilon}}-X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\|\sigma_{1}(X_{r}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}r\right)^{1/2}I_{\{\sup_{t\in[0,T]}|X_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|\leqslant R\}}$$

$$\leqslant \left(4l^{1/2}R+2\left(l+h_{R}(l)\right)\mathbb{E}|K^{1,\varepsilon,\gamma,u_{\varepsilon}}|_{0}^{T}\right)+C\left(1+|x_{0}|^{2}+|y_{0}|^{2}\right)l$$

$$+\frac{1}{2}\mathbb{E}\sup_{s\leqslant t\leqslant s+l}|X_{t}^{\varepsilon,\gamma,u_{\varepsilon}}-X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}I_{\{\sup_{t\in[0,T]}|X_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|\leqslant R\}}+C\left(1+|x_{0}|^{2}+|y_{0}|^{2}\right)l,$$

which yields that

$$\mathbb{E} \sup_{s \leqslant t \leqslant s+l} |X_t^{\varepsilon,\gamma,u_{\varepsilon}} - X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 I_{\{\sup_{t \in [0,T]} |X_t^{\varepsilon,\gamma,u_{\varepsilon}}| \leqslant R\}}$$

$$\leqslant \left(4l^{1/2}R + 2\left(l + h_R(l)\right) \mathbb{E}|K^{1,\varepsilon,\gamma,u_{\varepsilon}}|_0^T\right) + C\left(1 + |x_0|^2 + |y_0|^2\right)l.$$

Based on this, letting  $l \to 0$  and then  $R \to \infty$ , one could conclude (53).

**Lemma 6.3.** Under the assumptions of Theorem 3.3, for any  $u \in \mathbf{A}_2^N$ , it holds that

$$\sup_{t \in [0,T]} |\bar{X}_t^u|^2 \leqslant C(1+|x_0|^2), \tag{55}$$

$$|\bar{K}^u|_0^T \leqslant C(1+|x_0|^2),$$
 (56)

$$\lim_{l \to 0} \sup_{s \in [0,T]} \sup_{s \le t \le s+l} |\bar{X}_t^u - \bar{X}_s^u|^2 = 0.$$
 (57)

Since the proofs of (55), (56) and (57) are similar to that of (47), (49) and (53), respectively, we omit it.

Next, set for  $u_{\varepsilon}, u \in \mathbf{A}_2^N$ 

$$g_{\varepsilon}(t) := \int_0^t \sigma_1(\bar{X}_r^u) \pi_1(u_{\varepsilon}(r) - u(r)) dr,$$

and we have the following result.

**Lemma 6.4.** suppose that the assumptions of Theorem 3.3 hold, and  $u_{\varepsilon}$  converges to u almost surely as  $\varepsilon \to 0$ . Then  $g_{\varepsilon}(\cdot)$  tends to 0 almost surely in  $C([0,T],\mathbb{R}^n)$  as  $\varepsilon \to 0$ .

*Proof.* On one hand,  $\{g_{\varepsilon}; \varepsilon \in (0,1)\}$  is relatively compact in  $C([0,T],\mathbb{R}^n)$ . In fact, by the Ascoli-Arzelá lemma, we only need to justify that

- (i)  $\sup_{\varepsilon \in (0,1)} \sup_{t \in [0,T]} |g_{\varepsilon}(t)| < \infty$ ,
- (ii)  $\{[0,T]\ni t\mapsto g_{\varepsilon}(t); \varepsilon\in(0,1)\}$  is equi-continuous.

For  $0 \le s < t \le T$ , it holds that

$$|g_{\varepsilon}(s) - g_{\varepsilon}(t)| \leqslant \left| \int_{s}^{t} \sigma_{1}(\bar{X}_{r}^{u}) \pi_{1}(u_{\varepsilon}(r) - u(r)) dr \right| \leqslant 2N^{1/2} \left( \int_{s}^{t} \|\sigma_{1}(\bar{X}_{r}^{u})\|^{2} dr \right)^{1/2}$$
$$\leqslant 2N^{1/2} C \left( \int_{s}^{t} (1 + |\bar{X}_{r}^{u}|^{2}) dr \right)^{1/2},$$

where (6) is used. Letting s = 0, by (55) we have that

$$|g_{\varepsilon}(t)| \leq 2N^{1/2}C(1+|x_0|),$$

where C is independent of  $\varepsilon$ . So, (i) holds.

For (ii), noticing that

$$|g_{\varepsilon}(s) - g_{\varepsilon}(t)| \leq 2N^{1/2}C(1+|x_0|)(t-s)^{1/2},$$

we know that (ii) holds.

On the other hand, note that

$$\int_0^t \|\sigma_1(\bar{X}_r^u)\|^2 dr \le C \int_0^t (1 + |\bar{X}_r^u|^2) dr < \infty.$$

Since  $u_{\varepsilon} \to u$  almost surely in  $\mathbf{A}_2^N$  as  $\varepsilon \to 0$ , one get that for any  $t \in [0,T]$ 

$$\lim_{\varepsilon \to 0} g_{\varepsilon}(t) = 0,$$

which implies that

$$\lim_{\varepsilon \to 0} \sup_{t \in [0,T]} |g_{\varepsilon}(t)| = 0.$$

The proof is complete.

Finally we introduce the following auxiliary process:

$$\begin{cases}
\hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}} \in -A_{2}(\hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}}) dt + \frac{1}{\gamma} b_{2}(X_{t(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}}) dt + \frac{1}{\sqrt{\gamma}} \sigma_{2}(X_{t(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{t}^{2}, \\
\hat{Y}_{0}^{\varepsilon,\gamma,u_{\varepsilon}} = y_{0}.
\end{cases} (58)$$

**Lemma 6.5.** Under the assumptions of Theorem 3.3, for  $\{u_{\varepsilon}, \varepsilon \in (0,1)\} \subset \mathbf{A}_{2}^{N}$ , there exists a constant C > 0 such that

$$\sup_{t \in [0,T]} \mathbb{E} |\hat{Y}_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 \leqslant C(1+|x_0|^2+|y_0|^2), \tag{59}$$

$$\int_0^T \mathbb{E}|Y_t^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 dt \leqslant \frac{C}{\alpha} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s+\delta} |X_r^{\varepsilon,\gamma,u_{\varepsilon}} - X_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 \right) + C \frac{\gamma}{\varepsilon}.$$
 (60)

*Proof.* Since the proof of (59) is similar to that for (22), we only prove (60). First of all, by (9) and (58), we have that

$$Y_{t}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}} = -K_{t}^{2,\varepsilon,\gamma,u_{\varepsilon}} + \hat{K}_{t}^{2,\varepsilon,\gamma,u_{\varepsilon}} + \frac{1}{\gamma} \int_{0}^{t} \left( b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - b_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \right) ds$$

$$+ \frac{1}{\sqrt{\gamma}} \int_{0}^{t} \left( \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \right) dW_{s}^{2}$$

$$+ \frac{1}{\sqrt{\gamma\varepsilon}} \int_{0}^{t} \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{2} u_{\varepsilon}(s) ds.$$

Applying the Itô formula to  $|Y_t^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 e^{\lambda t}$  for  $\lambda = \frac{\alpha}{2\gamma}$  and taking the expectation, by  $(\mathbf{H}_{b_2,\sigma_2}^1)$  and  $(\mathbf{H}_{b_2,\sigma_2}^2)$  one could obtain that

$$\mathbb{E}|Y_{t}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{t}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda t}$$

$$= \lambda \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}e^{\lambda s} ds - \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, d(K_{s}^{2,\varepsilon,\gamma,u_{\varepsilon}} - \hat{K}_{s}^{2,\varepsilon,\gamma,u_{\varepsilon}}) \rangle$$

$$+ \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - b_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$\begin{split} & + \frac{1}{\sqrt{\gamma \varepsilon}} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{2} u_{\varepsilon}(s) \rangle \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} \| \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \|^{2} \mathrm{d}s \\ & \leqslant \lambda \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} e^{\lambda s} \mathrm{d}s + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle \\ & - b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle \mathrm{d}s + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} \| \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \|^{2} \mathrm{d}s \\ & + \frac{\alpha}{2\gamma} \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} e^{\lambda s} \mathrm{d}s + \frac{C}{\varepsilon} \mathbb{E} \int_{0}^{t} e^{\lambda s} |u_{\varepsilon}(s)|^{2} \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \langle Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, b_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - b_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} \| \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \|^{2} \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \| \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \|^{2} \mathrm{d}s \\ & + \frac{1}{\gamma} \mathbb{E} \int_{0}^{t} 2e^{\lambda s} \| \sigma_{2}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \sigma_{2}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \|^{2} \mathrm{d}s \\ & \leq (\lambda - \frac{\beta}{\gamma} + \frac{2L_{b_{2},\sigma_{2}}}{\gamma} + \frac{\alpha}{2\gamma}) \mathbb{E} \int_{0}^{t} |Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} e^{\lambda s} \mathrm{d}s + \frac{C}{\varepsilon} \mathbb{E} \int_{0}^{t} e^{\lambda s} |u_{\varepsilon}(s)|^{2} \mathrm{d}s \\ & + \frac{C}{\gamma} \mathbb{E} \int_{0}^{t} e^{\lambda s} |u_{\varepsilon}(s)|^{2} \mathrm{d}s + \frac{C}{\gamma} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s + \delta} |X_{r}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} \right) \frac{e^{\lambda t} - 1}{\lambda}. \end{split}$$

Finally, it follows that

$$\int_0^T \mathbb{E}|Y_t^{\varepsilon,\gamma,u_\varepsilon} - \hat{Y}_t^{\varepsilon,\gamma,u_\varepsilon}|^2 dt \leqslant \frac{C}{\alpha} \left( \sup_{s \in [0,T]} \mathbb{E} \sup_{s \leqslant r \leqslant s+\delta} |X_r^{\varepsilon,\gamma,u_\varepsilon} - X_s^{\varepsilon,\gamma,u_\varepsilon}|^2 \right) + C\left(\frac{\gamma}{\varepsilon}\right).$$

The proof is complete.

## 6.2. Verification for Condition 2.8.

**Lemma 6.6.** Suppose that the assumptions of Theorem 3.3 hold. Assume that  $u, u_{\varepsilon} \in \mathbf{A}_{2}^{N}$ ,  $\varepsilon \in (0,1)$ , and  $u_{\varepsilon}$  converges to u almost surely as  $\varepsilon \to 0$ . Then  $\Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_{0}^{\infty} u_{\varepsilon}(s) ds) \to \Psi^{0}(\int_{0}^{\infty} u(s) ds)$  in probability.

*Proof.* We divide the proof into four steps.

**Step 1.** We estimate the difference of  $\Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_0^{\cdot} u_{\varepsilon}(s)ds), \Psi^{0}(\int_0^{\cdot} u(s)ds)$  in  $\mathbb{S}$ . Note that

$$X^{\varepsilon,\gamma,u_{\varepsilon}} = \Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_{0}^{\cdot} u_{\varepsilon}(s)\mathrm{d}s), \quad \bar{X}^{u} = \Psi^{0}(\int_{0}^{\cdot} u(s)\mathrm{d}s).$$

Thus, set  $Z^{\varepsilon,u_{\varepsilon}}(t) = X_t^{\varepsilon,\gamma,u_{\varepsilon}} - \bar{X}_t^u$ , and it holds that

$$Z^{\varepsilon,u_{\varepsilon}}(t) = -(K_t^{1,\varepsilon,\gamma,u_{\varepsilon}} - \bar{K}_t^u) + \int_0^t \left[ b_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}, Y_s^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_1(\bar{X}_s^u) \right] ds$$

$$+ \int_0^t \left[ \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_1 u_{\varepsilon}(s) - \sigma_1(\bar{X}_s^u) \pi_1 u(s) \right] ds + \sqrt{\varepsilon} \int_0^t \sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}}) dW_s^1.$$

By Itô's formula and Lemma 2.3, we get that

$$|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}$$

$$= -2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), d(K_{s}^{1,\varepsilon,\gamma,u_{\varepsilon}} - \bar{K}_{s}^{u}) \rangle + 2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), b_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{s}^{u}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(s) - \sigma_{1}(\bar{X}_{s}^{u}) \pi_{1} u(s) \rangle ds$$

$$+2\sqrt{\varepsilon} \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{s}^{1} \rangle + \varepsilon \int_{0}^{t} \|\sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2} ds$$

$$\leq 2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), b_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{s}^{u}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(s) - \sigma_{1}(\bar{X}_{s}^{u}) \pi_{1} u(s) \rangle ds$$

$$+2\sqrt{\varepsilon} \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{s}^{1} \rangle + \varepsilon \int_{0}^{t} \|\sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2} ds$$

$$=: J_{1}(t) + J_{2}(t) + J_{3}(t) + J_{4}(t).$$
(61)

For  $J_1(t)$ , note that

$$J_{1}(t) = 2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), b_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), -\bar{b}_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) + \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s), \bar{b}_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{s}^{u}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s) - Z^{\varepsilon,u_{\varepsilon}}(s(\delta)), b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$+2 \int_{0}^{t} \langle Z^{\varepsilon,u_{\varepsilon}}(s(\delta)), b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) \rangle ds$$

$$=: J_{11}(t) + J_{12}(t) + J_{13}(t) + J_{14}(t) + J_{15}(t).$$

So, by the Hölder inequality and the Lipschitz continuity of  $b_1, \bar{b}_1$ , we get that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|J_{11}(t)|\right) + \mathbb{E}\left(\sup_{t\in[0,T]}|J_{12}(t)|\right) + \mathbb{E}\left(\sup_{t\in[0,T]}|J_{13}(t)|\right) 
\leqslant C \int_{0}^{T} \mathbb{E}|Z^{\varepsilon,u_{\varepsilon}}(s)|^{2} ds + C \int_{0}^{T} \mathbb{E}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} ds 
+ C \int_{0}^{T} \mathbb{E}|Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} ds.$$
(62)

And the Hölder inequality and the linear growth of  $b_1, \bar{b}_1$  imply that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|J_{14}(t)|\right) \leqslant C\left(\int_{0}^{T}(\mathbb{E}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}-X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}+\mathbb{E}|\bar{X}_{s}^{u}-\bar{X}_{s(\delta)}^{u}|^{2})\mathrm{d}s\right)^{1/2} \times \left(\int_{0}^{T}(1+\mathbb{E}|X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}+\mathbb{E}|\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2})\mathrm{d}s\right)^{1/2}.$$
(63)

For  $J_2$ , we rewrite it as

$$J_{2}(t) = 2 \int_{0}^{t} \langle Z^{\varepsilon, u_{\varepsilon}}(s), \sigma_{1}(X_{s}^{\varepsilon, \gamma, u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(s) - \sigma_{1}(\bar{X}_{s}^{u}) \pi_{1} u_{\varepsilon}(s) \rangle ds$$

$$+ 2 \int_{0}^{t} \langle Z^{\varepsilon, u_{\varepsilon}}(s), \sigma_{1}(\bar{X}_{s}^{u}) \pi_{1}(u_{\varepsilon}(s) - u(s)) \rangle ds.$$

$$=: J_{21}(t) + J_{22}(t).$$

For  $J_{21}(t)$ , by  $(\mathbf{H}^1_{b_1,\sigma_1})$  and the Hölder inequality, it holds that

$$\mathbb{E} \sup_{t \in [0,T]} |J_{21}(t)| \leqslant 2\sqrt{L_{b_1,\sigma_1}} \mathbb{E} \int_0^T |Z^{\varepsilon,u_{\varepsilon}}(s)| |Z^{\varepsilon,u_{\varepsilon}}(s)| |u_{\varepsilon}(s)| ds$$

$$\leqslant \frac{1}{4} \mathbb{E} \left[ \sup_{t \in [0,T]} |Z^{\varepsilon,u_{\varepsilon}}(t)|^2 \right] + C \mathbb{E} \left( \int_0^T |Z^{\varepsilon,u_{\varepsilon}}(s)| |u_{\varepsilon}(s)| ds \right)^2$$

$$\leqslant \frac{1}{4} \mathbb{E} \left[ \sup_{t \in [0,T]} |Z^{\varepsilon,u_{\varepsilon}}(t)|^2 \right] + C \mathbb{E} \left( \int_0^T |Z^{\varepsilon,u_{\varepsilon}}(s)|^2 ds \right) \left( \int_0^T |u_{\varepsilon}(s)|^2 ds \right)$$

$$\leqslant \frac{1}{4} \mathbb{E} \left[ \sup_{t \in [0,T]} |Z^{\varepsilon,u_{\varepsilon}}(t)|^2 \right] + C \mathbb{E} \left[ \int_0^T |Z^{\varepsilon,u_{\varepsilon}}(s)|^2 ds \right]. \tag{64}$$

For  $J_3(t)$ , from the Burkholder-Davis-Gundy inequality and the linear growth of  $\sigma_1$ , it follows that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|J_{3}(t)|\right) \leq 2\sqrt{\varepsilon}C\mathbb{E}\left(\int_{0}^{T}|Z^{\varepsilon,u_{\varepsilon}}(s)|^{2}\|\sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}s\right)^{1/2}$$

$$\leq \frac{1}{4}\mathbb{E}\left[\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}\right] + \sqrt{\varepsilon}C\mathbb{E}\int_{0}^{T}\|\sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}})\|^{2}\mathrm{d}s$$

$$\leq \frac{1}{4}\mathbb{E}\left[\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}\right] + \sqrt{\varepsilon}C\int_{0}^{T}(1+\mathbb{E}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2})\mathrm{d}s$$

$$\leq \frac{1}{4}\mathbb{E}\left[\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}\right] + \sqrt{\varepsilon}C. \tag{65}$$

For  $J_4(t)$ , by the linear growth of  $\sigma_1$ , we know

$$\mathbb{E}\left(\sup_{t\in[0,T]}|J_4(t)|\right) \leqslant \varepsilon C \int_0^T (1+\mathbb{E}|X_s^{\varepsilon,\gamma,u_\varepsilon}|^2) \mathrm{d}s \leqslant \varepsilon C. \tag{66}$$

Combining (62)-(66) with (61), we can get

$$\mathbb{E}\left(\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}\right) \\
\leqslant C\int_{0}^{T}\mathbb{E}\sup_{r\in[0,s]}|Z^{\varepsilon,u_{\varepsilon}}(r)|^{2}\mathrm{d}s + C\int_{0}^{T}\mathbb{E}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}s + C\int_{0}^{T}\mathbb{E}|Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}\mathrm{d}s \\
+C\left(\int_{0}^{T}(\mathbb{E}|X_{s}^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + \mathbb{E}|\bar{X}_{s}^{u} - \bar{X}_{s(\delta)}^{u}|^{2})\mathrm{d}s\right)^{1/2} \\
\times \left(\int_{0}^{T}(1 + \mathbb{E}|X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + \mathbb{E}|\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2})\mathrm{d}s\right)^{1/2} \\
+4\mathbb{E}\left(\sup_{t\in[0,T]}|J_{15}(t)|\right) + 4\mathbb{E}\left(\sup_{t\in[0,T]}|J_{22}(t)|\right) + C(\sqrt{\varepsilon} + \varepsilon),$$

which together with the Gronwall inequality implies that

$$\mathbb{E}\left(\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|^{2}\right)\leqslant C\left[\Gamma(\varepsilon)+\mathbb{E}\left(\sup_{t\in[0,T]}|J_{15}(t)|\right)+\mathbb{E}\left(\sup_{t\in[0,T]}|J_{22}(t)|\right)+C(\sqrt{\varepsilon}+\varepsilon)\right],(67)$$

where

$$\Gamma(\varepsilon) := C \int_0^T \mathbb{E}|X_s^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^2 ds + C \int_0^T \mathbb{E}|Y_s^{\varepsilon,\gamma,u_{\varepsilon}} - \hat{Y}_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2 ds$$

$$+ C \left( \int_0^T (\mathbb{E}|X_s^{\varepsilon,\gamma,u_{\varepsilon}} - X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \mathbb{E}|\bar{X}_s^u - \bar{X}_{s(\delta)}^u|^2) ds \right)^{1/2}$$

$$\times \left( \int_0^T (1 + \mathbb{E}|X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \mathbb{E}|\hat{Y}_s^{\varepsilon,\gamma,u_{\varepsilon}}|^2) ds \right)^{1/2}.$$

Step 2. We estimate  $\mathbb{E} \sup_{t \in [0,T]} |J_{15}(t)|$ .

For  $J_{15}(t)$ , it holds that

$$\mathbb{E}\sup_{t\in[0,T]}|J_{15}(t)| = 2\left(\mathbb{E}\sup_{0\leqslant t\leqslant T}\Big|\int_{0}^{[\frac{t}{\delta}]\delta}\langle Z^{\varepsilon,u_{\varepsilon}}(s(\delta)),b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}},\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}})\rangle \mathrm{d}s + \int_{[\frac{t}{\delta}]\delta}^{t}\langle Z^{\varepsilon,u_{\varepsilon}}(s(\delta)),b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}},\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}})\rangle \mathrm{d}s\Big|\right)$$

$$\leqslant 2\left(\mathbb{E}\sup_{0\leqslant t\leqslant T}\Big|\int_{0}^{[\frac{t}{\delta}]\delta}\langle Z^{\varepsilon,u_{\varepsilon}}(s(\delta)),b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}},\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}})\rangle \mathrm{d}s\Big|\right)$$

$$+2\left(\mathbb{E}\sup_{0\leqslant t\leqslant T}\Big|\int_{[\frac{t}{\delta}]\delta}^{t}\langle Z^{\varepsilon,u_{\varepsilon}}(s(\delta)),b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}},\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}})\rangle \mathrm{d}s\Big|\right)$$

$$=: J_{151} + J_{152}.$$

Next, we estimate  $I_{151}$ . By the similar deduction to that of  $I_{21}$  in Lemma 5.7, we get that

$$I_{151} \leqslant C(\frac{\gamma}{\delta})^{1/2}.\tag{68}$$

Finally, we estimate  $I_{152}$ . By (6) (47) (59) and the Hölder inequality, one could get that

$$\begin{split} I_{152} &\leqslant 2 \Biggl( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \int_{[\frac{t}{\delta}]\delta}^{t} |Z^{\varepsilon,u_{\varepsilon}}(s(\delta))| |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) | \mathrm{d}s \Biggr) \\ &\leqslant 2 \delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |Z^{\varepsilon,u_{\varepsilon}}(s)|^{2} \right)^{1/2} \left( \mathbb{E} \sup_{0 \leqslant t \leqslant T} \int_{[\frac{t}{\delta}]\delta}^{t} |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) |^{2} \mathrm{d}s \Biggr)^{1/2} \\ &\leqslant 2 \delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |Z^{\varepsilon,u_{\varepsilon}}(s)|^{2} \right)^{1/2} \left( \mathbb{E} \int_{0}^{T} |b_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}, \hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}) |^{2} \mathrm{d}s \Biggr)^{1/2} \\ &\leqslant 2 \delta^{1/2} \left( \mathbb{E} \sup_{0 \leqslant s \leqslant T} |X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + \mathbb{E} \sup_{0 \leqslant s \leqslant T} |\bar{X}_{s}^{u}|^{2} \right)^{1/2} \left( \int_{0}^{T} (1 + \mathbb{E}|X_{s(\delta)}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2} + \mathbb{E}|\hat{Y}_{s}^{\varepsilon,\gamma,u_{\varepsilon}}|^{2}) \mathrm{d}s \right)^{1/2} \\ &\leqslant C \delta^{1/2}, \end{split}$$

which together with (68) implies

$$\mathbb{E} \sup_{t \in [0,T]} |J_{15}(t)| \leqslant C\left(\left(\frac{\gamma}{\delta}\right)^{1/2} + \delta^{1/2}\right). \tag{69}$$

Step 3. We compute  $\mathbb{E} \sup_{t \in [0,T]} |J_{22}(t)|$ .

For  $J_{22}(t)$ , applying the Itô formula to  $\langle Z^{\varepsilon,u_{\varepsilon}}(t),g_{\varepsilon}(t)\rangle$ , we have that

$$\frac{1}{2}J_{22}(t) = \langle Z^{\varepsilon,u_{\varepsilon}}(t), g_{\varepsilon}(t) \rangle + \int_{0}^{t} \langle g_{\varepsilon}(s), d(K_{s}^{1,\varepsilon,\gamma,u_{\varepsilon}} - \bar{K}_{s}^{u}) \rangle 
- \int_{0}^{t} \langle g_{\varepsilon}(s), b_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}, Y_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) - \bar{b}_{1}(\bar{X}_{s}^{u}) \rangle ds 
- \int_{0}^{t} \langle g_{\varepsilon}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) \pi_{1} u_{\varepsilon}(s) - \sigma_{1}(\bar{X}_{s}^{u}) \pi_{1} u(s) \rangle ds 
- \sqrt{\varepsilon} \int_{0}^{t} \langle g_{\varepsilon}(s), \sigma_{1}(X_{s}^{\varepsilon,\gamma,u_{\varepsilon}}) dW_{s}^{1} \rangle 
=: J_{221}(t) + J_{222}(t) + J_{223}(t) + J_{224}(t) + J_{225}(t).$$

For  $J_{221}(t)$ , note that

$$\sup_{t \in [0,T]} |J_{221}(t)| \leqslant \sup_{t \in [0,T]} |Z^{\varepsilon,u_{\varepsilon}}(t)| \sup_{t \in [0,T]} |g_{\varepsilon}(t)|.$$

Thus, by Lemma 6.4 it holds that

$$\sup_{t \in [0,T]} |J_{221}(t)| \stackrel{a.s.}{\to} 0, \quad \varepsilon \to 0.$$

Besides, the fact that  $\sup_{t \in [0,T]} |g_{\varepsilon}(t)| \leq C(1+|x_0|)$  implies that

$$\mathbb{E}\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|\sup_{t\in[0,T]}|g_{\varepsilon}(t)| \leqslant C(1+|x_{0}|)\mathbb{E}\sup_{t\in[0,T]}|Z^{\varepsilon,u_{\varepsilon}}(t)|$$

$$\leq C(1+|x_0|) \left( \mathbb{E} \sup_{t \in [0,T]} |Z^{\varepsilon,u_{\varepsilon}}(t)|^2 \right)^{1/2} \\
\leq C(1+|x_0|) \left( \mathbb{E} \sup_{0 \leq t \leq T} |X_t^{\varepsilon,\gamma,u_{\varepsilon}}|^2 + \mathbb{E} \sup_{0 \leq t \leq T} |\bar{X}_t^u|^2 \right)^{1/2} \\
\leq C(1+|x_0|^2+|y_0|^2),$$

where the last inequality is based on (47), (55). Then the dominated convergence theorem yields that

$$\lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{221}(t)| = 0.$$

For  $J_{222}(t)$ , noticing that

$$\sup_{t \in [0,T]} |J_{222}(t)| \leqslant \sup_{t \in [0,T]} |g_{\varepsilon}(t)| (|K^{1,\varepsilon,\gamma,u_{\varepsilon}}|_0^T + |\bar{K}^u|_0^T),$$

by Lemma 6.4 we obtain that  $\sup_{t\in[0,T]}|J_{222}(t)|\stackrel{a.s.}{\to} 0$ . Then (49), (56) and the dominated convergence theorem imply that

$$\lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{222}(t)| = 0. \tag{70}$$

By the same deduction to that for (70), one can get that

$$\lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{223}(t)| = 0, \quad \lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{224}(t)| = 0.$$

For  $J_{225}(t)$ , the BDG inequality yields that

$$\mathbb{E} \sup_{t \in [0,T]} |J_{225}(t)| \leq C\sqrt{\varepsilon} \mathbb{E} \left( \int_0^T |g_{\varepsilon}(s)|^2 \|\sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\|^2 ds \right)^{1/2}$$

$$\leq C\sqrt{\varepsilon} \left( \mathbb{E} \sup_{t \in [0,T]} |g_{\varepsilon}(t)|^2 + \mathbb{E} \int_0^T \|\sigma_1(X_s^{\varepsilon,\gamma,u_{\varepsilon}})\|^2 ds \right)$$

$$\leq C\sqrt{\varepsilon} (1 + |x_0|^2 + |y_0|^2).$$

Therefore, it holds that

$$\lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{225}(t)| = 0.$$

Combining the above deduction, we get that

$$\lim_{\varepsilon \to 0} \mathbb{E} \sup_{t \in [0,T]} |J_{22}(t)| = 0. \tag{71}$$

Step 4. We prove that  $\Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_{0}^{\cdot} u_{\varepsilon}(s)ds) \to \Psi^{0}(\int_{0}^{\cdot} u(s)ds)$  in probability. Let  $\delta = \gamma^{\iota}$  for  $0 < \iota < 1$  and  $\gamma \to 0, \delta \to 0, \gamma/\delta \to 0$  as  $\varepsilon \to 0$ . Thus, (53) (57) (60) (67) (69) (71) imply that  $X^{\varepsilon,\gamma,u_{\varepsilon}} - \bar{X}^{u} \to 0$  in the mean square and  $\Psi^{\varepsilon}(\sqrt{\varepsilon}W + \int_{0}^{\cdot} u_{\varepsilon}(s)ds) \to \Psi^{0}(\int_{0}^{\cdot} u(s)ds)$  in probability. The proof is complete.

By the similar or even simpler deduction to that of Lemma 6.6, we obtain the following result.

**Lemma 6.7.** Suppose that the assumptions of Theorem 3.3 hold, and  $h_{\varepsilon} \to h$  in  $\mathbf{D}_{2}^{N}$  as  $\varepsilon \to 0$ . Then  $\Psi^{0}(\int_{0}^{\cdot} h_{\varepsilon}(s) ds)$  converges to  $\Psi^{0}(\int_{0}^{\cdot} h(s) ds)$ .

Now, it is the position to prove Theorem 3.3.

## Proof of Theorem 3.3.

By Theorem 2.9, to establish LDP, it is sufficient to verify the two conditions in Condition 2.8. In Lemma 6.7, we have already proved Condition 2.8 (i). Thus we only need to verify Condition 2.8 (ii).

For  $\varepsilon \in (0,1)$  and  $\{u_{\varepsilon}, \varepsilon > 0\} \subset \mathbf{A}_2^N$ ,  $u \in \mathbf{A}_2^N$ , let  $u_{\varepsilon}$  converge to u in distribution. By the Skorohod theorem, there exists a probability space  $(\tilde{\Omega}, \tilde{\mathscr{F}}, \tilde{\mathbb{P}})$ , and  $\mathbf{A}_2^N$ -valued random variables  $\{\tilde{u}_{\varepsilon}\}$ ,  $\tilde{u}$  and a  $d_1 + d_2$ -dimensional Brownian motion  $\tilde{W}$  such that

- (i)  $(\tilde{u}_{\varepsilon}, \tilde{W}) \stackrel{d}{=} (u_{\varepsilon}, W)$  and  $\tilde{u} \stackrel{d}{=} u$ ;
- (ii)  $\tilde{u}_{\varepsilon}$  converges to  $\tilde{u}$  almost surely.

In the following, we construct multivalued SDEs:

In the following, we construct multivalued SDEs:
$$\begin{cases}
dX_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}} \in -A_1(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})dt + b_1(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, Y_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})dt + \sigma_1(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})\pi_1\tilde{u}_{\varepsilon}(t)dt \\
+\sqrt{\varepsilon}\sigma_1(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})d\tilde{W}_t^1, \\
X_0^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}} = x_0 \in \overline{\mathcal{D}}(A_1), \quad 0 \leqslant t \leqslant T, \\
dY_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}} \in -A_2(Y_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})dt + \frac{1}{\gamma}b_2(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, Y_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})dt + \frac{1}{\sqrt{\gamma\varepsilon}}\sigma_2(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, Y_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})\pi_2\tilde{u}_{\varepsilon}(t)dt
\end{cases} (72)$$

$$+\frac{1}{\sqrt{\gamma}}\sigma_2(X_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, Y_t^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}})d\tilde{W}_t^2, \quad \tilde{u}_{\varepsilon} \in \mathbf{A}_2^N, \\
Y_0^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}} = y_0 \in \overline{\mathcal{D}}(A_2), \quad 0 \leqslant t \leqslant T,$$

$$\int d\bar{X}_t^{\tilde{u}} \in A_1(\bar{X}_t^{\tilde{u}})dt + \bar{b}_1(\bar{X}_t^{\tilde{u}})dt + \sigma_1(\bar{X}_t^{\tilde{u}})\pi_1u(t)dt, \quad \tilde{u} \in \mathbf{A}_2^N, \tag{72}$$

$$\begin{cases}
d\bar{X}_t^{\tilde{u}} \in A_1(\bar{X}_t^{\tilde{u}}) dt + \bar{b}_1(\bar{X}_t^{\tilde{u}}) dt + \sigma_1(\bar{X}_t^{\tilde{u}}) \pi_1 u(t) dt, & \tilde{u} \in \mathbf{A}_2^N, \\
\bar{X}_0^{\tilde{u}} = x_0.
\end{cases}$$
(73)

For the system (72) and Eq.(73), each has a unique strong solution denoted by  $(X^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}},$  $K^{1,\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, Y^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}}, K^{2,\varepsilon,\gamma,\tilde{u}_{\varepsilon}})$  and  $(\tilde{X}^{\tilde{u}}, \bar{K}^{\tilde{u}})$  respectively. Moreover, it holds that

$$X^{\varepsilon,\gamma,\tilde{u}_{\varepsilon}} = \Psi^{\varepsilon} \left( \sqrt{\varepsilon} \tilde{W} + \int_{0}^{\cdot} \tilde{u}_{\varepsilon}(s) \mathrm{d}s \right), \quad \bar{X}^{\tilde{u}} = \Psi^{0} \left( \int_{0}^{\cdot} \tilde{u}(s) \mathrm{d}s \right).$$

By Lemma 6.6, we have that  $\Psi^{\varepsilon}(\sqrt{\varepsilon}\tilde{W} + \int_{0}^{\cdot} \tilde{u}_{\varepsilon}(s)ds) \to \Psi^{0}(\int_{0}^{\cdot} \tilde{u}(s)ds)$  in probability, which yields that  $\Psi^{\varepsilon}(\sqrt{\varepsilon}\tilde{W} + \int_{0}^{\cdot} \tilde{u}_{\varepsilon}(s)ds) \to \Psi^{0}(\int_{0}^{\cdot} \tilde{u}(s)ds)$  in distribution. Note that

$$\Psi^{\varepsilon} \left( \sqrt{\varepsilon} \tilde{W} + \int_{0}^{\cdot} \tilde{u}_{\varepsilon}(s) ds \right) \stackrel{d}{=} \Psi^{\varepsilon} \left( \sqrt{\varepsilon} W + \int_{0}^{\cdot} u_{\varepsilon}(s) ds \right),$$

$$\Psi^{0} \left( \int_{0}^{\cdot} \tilde{u}(s) ds \right) \stackrel{d}{=} \Psi^{0} \left( \int_{0}^{\cdot} u(s) ds \right).$$

So,  $\Psi^{\varepsilon}\left(\sqrt{\varepsilon}W + \int_{0}^{\cdot} u_{\varepsilon}(s)ds\right) \to \Psi^{0}\left(\int_{0}^{\cdot} u(s)ds\right)$  in distribution, which is Condition 2.8 (ii).

#### References

- [1] A. Bensoussan, L. Mertz and S. Yam: Long cycle behavior of the plastic deformation of an elastoperfectly-plastic oscillator with noise, C. R. Math. Acad. Sci. Paris, 350(17-18)(2012)853-859.
- [2] A. Bensoussan and J. Turi: Stochastic variational inequalities for elasto-plastic oscillators, C. R. Math. Acad. Sci. Paris, 343(2006)399-406.
- [3] A. Bensoussan and J. Turi: Degenerate Dirichlet problems related to the invariant measure of elasto-plastic oscillators, Appl. Math. Optim., 58(1)(2008)1-27.
- [4] R. Bertram and J. E. Rubin: Multi-timescale systems and fast-slow analysis, Math. Biosci., 287(2017)105-121.

- [5] A. Budhiraja, P. Dupuis, and V. Maroulas: Variational representations for continuous time processes. *Ann. Inst. Henri Poincaré*, *Probab. Stat.*, 47(2011)725-747.
- [6] E. Cépa: Équations différentielles stochastiques multivoques, in: Sém. Prob. XXIX, in: Lecture Notes in Math., 1995, pp. 86-107.
- [7] E. Cépa: Probleme de Skorohod Multivoque, Ann. Prob., 26(1998)500-532.
- [8] Z.-Q. Chen and J. Wu: Averaging principle for stochastic variational inequalities with application to PDEs with nonlinear Neumann conditions, *Journal of Differential Equations*, 328(2022)157-201.
- [9] P. Dupuis and R. Ellis: A Weak Convergence Approach to the Theory of Large Deviations, Wiley, New York, 1997.
- [10] P. Dupuis and K. Spiliopoulos: Large deviations for multiscale diffusion via weak convergence methods, Stoch. Process. Appl., 122(2012)1947-1987.
- [11] W. E and B. Engquist: Multiscale modeling and computations, Not. Am. Math. Soc., 50(2003)1062-1070.
- [12] W. E, D. Liu and E. Vanden-Eijnden: Analysis of multiscale methods for stochastic differential equations, *Commun. Pure Appl. Math.*, 58(2005)1544-1585.
- [13] K. Fang, W. Liu, H. Qiao and F. Zhu: Asymptotic behaviors of small perturbation for multivalued McKean-Vlasov stochastic differential equations, to appear in Applied Mathematics and Optimization, 2023.
- [14] M. Freidlin and A. D. Wentzell: Random Perturbations of Dynamical Systems, translated from the 1979 Russian original by Joseph Szücs, third edition, Springer, Heidelberg, 2012, xxviii+458 pp.
- [15] M. Freidlin and A. D. Wentzell: Long-time behavior of weakly coupled oscillators, J. Stat. Phys., 123(2006)1311-1337.
- [16] J. Gao, W. Hong and W. Liu: Small noise asymptotics of multi-scale McKean-Vlasov stochastic dynamical systems, *Journal of Differential Equations*, 364(2023)521-575.
- [17] J. Gong and H. Qiao: The stability for multivalued McKean-Vlasov SDEs with non-Lipschitz coefficients, https://arxiv.org/abs/2106.12080.
- [18] R. Liptser: Large deviations for two scaled diffusions, Probab. Theory Relat. Fields, 106(1996)71-104.
- [19] D. Liu: Strong convergence of principle of averaging for multiscale stochastic dynamical systems, *Commun. Math. Sci.*, 8(2010)999-1020.
- [20] W. Liu, M. Rökner, X. Sun and Y. Xie: Averaging principle for slow-fast stochastic differential equations with time dependent locally Lipschitz coefficients, J. Differ. Equ., 268(2020)2910-2948.
- [21] R. Z. Khas'minskii, On the principle of averaging the Itô's stochastic differential equations, Kybernetika (Prague), 4(1968) 260-279.
- [22] Y. Kifer, Some recent advances in averaging, in: Modern Dynamical Systems and Applications, Cambridge Univ. Press, Cambridge, 2004, pp. 385-403.
- [23] C. Kuehn: Multiple Time Scale Dynamics, Applied Mathematical Sciences, vol. 191, Springer, Cham, 2015.
- [24] R. Kumar and L. Popovic: Large deviations for multi-scale jump-diffusion processes, Stoch. Process. Appl., 127(2017)1297-1320.
- [25] H. J. Kushner: Large deviations for two-time-scale diffusions with delays, Applied Mathematics and Optimization, 62(2010)295-322.
- [26] A. A. Puhalskii: On large deviations of coupled diffusions with time scale separation, Ann. Probab., 44(2016)3111-3186.
- [27] H. Qiao: Limit theorems of invariant measures for multivalued McKean-Vlasov stochastic differential equations, to appear in *Journal of Mathematical Analysis and Applications*, 2023.
- [28] H. Qiao and J. Gong: Backward multivalued McKean-Vlasov SDEs and associated variational inequalities, Discrete and Continuous Dynamical Systems-S, 16(2023)819-845.
- [29] J. Ren, J. Wu and H. Zhang: General large deviations and functional iterated logarithm law for multivalued stochastic differential equations, J. Theor. Probab., 28(2015)550-586.
- [30] J. Ren, J. Wu and X. Zhang: Exponential ergodicity of non-Lipschitz multivalued stochastic differential equations, Bull. Sci. Math, 134(2010)391-404.
- [31] J. Ren, S. Xu and X. Zhang: Large deviations for multivalued stochastic differential equations, J. Theor. Probab 23 (2010), 1142-1156.

- [32] R. T. Rockafellar and R. J-B Wets: Stochastic variational inequalities: single-stage to multistage, *Math. Program. Ser. B*, 165(1)(2017)331-360.
- [33] K. Spiliopoulos: Large deviations and importance sampling for systems of slow-fast motion, Appl. Math. Optim., 67(2013)123-161.
- [34] A. Y. Verernnikov: On an averaging principle for systems of stochastic differential equations, *Mat. Sb.*, 181(2)(1990)256-268 (in Russian); translation in: *Math. USSR Sb.*, 69(1)(1991)271-284.
- [35] A.Yu. Veretennikov, On large deviations in the averaging principle for SDEs with a full dependence, correction, arXiv: math/0502098v1 [math.PR] (2005). initial article in Annals of Probability, 27(1999)284-296.
- [36] A. Y. Verernnikov: On large deviations for SDEs with small diffusion and averaging, *Stoch. Process.* Appl., 89(2000)69-79.
- [37] H. Zhang: Moderate deviation principle for multivalued stochastic differential equations. Stochastic and Dynamics, 20(2020)1-30.
- [38] X. Zhang: Skorohod problem and multivalued stochastic evolution equations in Banach spaces, Bull. Sci. Math., 131(2007)175-217.