CONTINUUM LIMIT OF SPIN DYNAMICS ON TWO-DIMENSIONAL HEXAGONAL LATTICE

MENGJIA BAI, JINGRUN CHEN, ZHIWEI SUN, YUN WANG

Abstract

This study investigates the atomistic spin system in $CrCl_3$, which exhibits topologically nontrivial meron structures within its layered hexagonal lattice framework. We analyze the complete model of discrete spin dynamics on a two-dimensional hexagonal lattice and demonstrate its convergence to the continuum Landau-Lifshitz-Gilbert equation in the weak sense. The primary challenge lies in defining appropriate difference quotient and interpolation operators for the hexagonal lattice since the loss of symmetry. To address these, we utilized a one-step difference quotient for the 2nd nearest neighbors and introduced novel multi-step difference quotients for the 1st and 3rd nearest neighbors, enabling the integration by parts formula. Additionally, we generalized Ladysenskaya's interpolation operator for hexagonal lattices and provided an alternative strategy for the convergence procedure by applying an isometric mapping property. This work provides necessary tools for analyzing weak convergence in other atomistic nonlinear problems on hexagonal lattices towards the continuum limit.

Key words : Spin dynamics, hexagonal lattice, Landau-Lifshitz-Gilbert equation, interpolation operator, difference method.

1. INTRODUCTION

Nature favors the hexagonal lattice, seen in honeycombs created by insects and graphene, a cutting-edge technology. Nanotechnology has led to the discovery of (quasi) two-dimensional materials with intriguing electronic and magnetic properties, such as superconducting states in twisted graphene and meron spin textures in $CrCl_3$, both possessing a hexagonal lattice. To understand the dynamics of physical quantities in these materials, two models have been developed: (1) an atomistic model that uses ordinary differential equations (ODEs) with degrees of freedom defined on the hexagonal lattice; (2) a phenomenological model that employs partial differential equations (PDEs), treating time and space variables continuously. While it is convenient to solve the phenomenological model, the atomistic model provides an accurate description of the underlying material. Therefore, it is natural to ask about the connection between atomistic and continuum models.

There is an extensive literature addressing material properties from various perspectives, including mechanical [3,12], electronic [9-11], and magnetic properties [6,15], among others. For mechanical property, the consistency between atomistic and continuum models is derived in [9] with stability proven in [12]. The electronic property aims to understand solids at the

quantum or molecular level with rigorous derivation of continuum models established in [9]. Magnetic property involves deriving the continuum limit of a discrete spin system on a cubic lattice as well as identifying a physically relevant thin-film limit starting from a three-dimensional micromagnetic model [6, 15].

Additionally, mathematical works have been done to pass the limit from ODE systems on hexagonal lattices to continuum PDE models [5,8]. However, these studies assume sufficient smoothness of the continuum solutions. This raises the question: if the solution of the continuum model exists only in the weak sense, is it reliable for describing the behavior of an atomistic system on a hexagonal lattice from a macroscopic scale? In this study, we address this question by focusing on the dynamics of discrete spins in $CrCl_3$. The discrete spin model is known for its excitation of various topologically nontrivial structures like domain walls, skyrmions, and merons, Recently, merons, along with antimerons, have been demonstrated to exist in the magnetic structure of hexagonal lattice $CrCl_3$ [2]. In this work, we analyze the complete model of discrete spin dynamics on a two-dimensional hexagonal lattice, and establish a rigorous passage from the discrete spin system to the well-known Landau-Lifshitz-Gilbert (LLG) equation in the weak sense.

To present the main finding, we begin by introducing the spin model for CrCl_3 . The two-dimensional hexagonal lattice point is denoted by \boldsymbol{x}_i , where $i \in \mathbb{N}$, and each atomic spin is represented by $\mathbf{m}(\boldsymbol{x}_i)$ with a unit length. The equilibrium state of spin $\mathbf{m}(\boldsymbol{x}_i)$ is determined by minimizing its energy. In the case of CrCl_3 , this can be expressed as per [2]:

(1)
$$\mathcal{H}[\mathbf{m}] = \mathcal{H}_{e}[\mathbf{m}] + \mathcal{H}_{b}[\mathbf{m}] + \mathcal{H}_{a}[\mathbf{m}] + \mathcal{H}_{z}[\mathbf{m}].$$

The different contributions to the energy in (1) are:

1. **Bilinear Exchange Energy**. The fundamental property of ferromagnetic materials is that the spins experience the presence of an exchange field, decribed by

$$\mathcal{H}_{\mathrm{e}}[\mathbf{m}] = -\frac{1}{2} \sum_{i,j \in \mathbb{N}} J_{ij} \mathbf{m}(\boldsymbol{x}_i) \cdot \mathbf{m}(\boldsymbol{x}_j).$$

Here, $\mathbf{m}(\mathbf{x}_i)$ and $\mathbf{m}(\mathbf{x}_j)$ are localized magnetic moments on Cr atomic sites *i* and *j* coupled by pair-wise exchange interactions. It is usually considered up to the third nearest neighbour for J_{ij} $(J_1 - J_2 - J_3)$, where J_k is constant denote the magnetic quantities of the *k*th nearest neighbour; see Figure 1. Here, we use the notation J_{ij} $(J_1 - J_2 - J_3)$ to represent the following: suppose \mathbf{x}_i is the *k*th nearest neighbour of \mathbf{x}_j , then

$$J_{ij} = \begin{cases} J_k, & \text{for } k = 1, 2, 3; \\ 0, & \text{for } k > 3. \end{cases}$$

2. Biquadratic (BQ) Exchange Energy. BQ exchange interactions involving the hopping of two electrons are critical in the elucidation of the magnetic features of layered magnetic materials [13]. This is described by

$$\mathcal{H}_{\mathrm{b}}[\mathbf{m}] = -\frac{1}{2} \sum_{i,j \in \mathbb{N}} K_{ij} (\mathbf{m}(\boldsymbol{x}_i) \cdot \mathbf{m}(\boldsymbol{x}_j))^2.$$

Its strength is given by the constant $K_{ij}(K)$ where only the nearest neighbours taken into account, i.e., suppose \boldsymbol{x}_i is the *k*th nearest neighbour of \boldsymbol{x}_j , then

$$K_{ij} = \begin{cases} K, & \text{for } k = 1; \\ 0, & \text{for } k > 1. \end{cases}$$

This is the simplest and most natural form of non-Heisenberg coupling.

3. Anisotropy Energy. The anisotropy energy can be described by the terms of the form

$$egin{aligned} \mathcal{H}_{\mathbf{a}}[\mathbf{m}] &= -\sum_{i\in\mathbb{N}}\lambda_{i}ig(\mathbf{m}(oldsymbol{x}_{i})\cdot\mathbf{e}^{x}ig)^{2} \ &- rac{1}{2}\sum_{i,j\in\mathbb{N}}L_{ij}ig(\mathbf{m}(oldsymbol{x}_{i})\cdot\mathbf{e}^{z}ig)ig(\mathbf{m}(oldsymbol{x}_{j})\cdot\mathbf{e}^{z}ig). \end{aligned}$$

Here, the first term is induced by electronic structure of the underlying crystalline lattice with a preferred orientation \mathbf{e}^x . We assume λ_i is the single-ion magnetic anisotropy coefficient. The second term is the symmetric anisotropy exchange energy that can be deduced when considering relativistic effect. It is considered up to the third nearest neighbour for $L_{ij} (L_1^{(ij)} - L_2^{(ij)} - L_3^{(ij)})$, where $L_k^{(ij)} (k = 1, 2, 3)$ denote the magnetic quantities of the *k*th nearest neighbour. More precisely, suppose \mathbf{x}_i is the *k*th nearest neighbour of \mathbf{x}_j , then we have:

$$L_{ij} = \begin{cases} L_k^{(ij)}, & \text{for } k = 1, 2, 3; \\ 0, & \text{for } k > 3. \end{cases}$$

4. Zeeman Energy. In the presence of an external magnetic field \mathbf{B}_i , the magnetization tends to align with it. This translates into an energy term of the form

$$\mathcal{H}_{\mathrm{z}}[\mathbf{m}] = -\sum_{i\in\mathbb{N}} \mu \mathbf{m}(\boldsymbol{x}_i) \cdot \mathbf{B}_i,$$

where μ is the magnetic permeability of vacuum.

Furthermore, note that we consider the hexagonal lattice over \mathbb{R}^2 without the compactness, the following assumption on the coefficients are needed such that the anisotropy energy and Zeeman energy are finite.

Assumption 1. We assume that there exists continuous functions $\lambda(\mathbf{x})$, $L_k(\mathbf{x})$, k = 1, 2, 3 and $\mathbf{B}(\mathbf{x})$ which are $L^1(\mathbb{R}^2)$ bounded. Furthermore, it

holds

$$\lambda_i = \lambda(\boldsymbol{x}_i), \qquad L_k^{(ij)} = \frac{1}{2} \Big(L_k(\boldsymbol{x}_i) + L_k(\boldsymbol{x}_j) \Big) \qquad \mathbf{B}_i = \mathbf{B}(\boldsymbol{x}_i).$$

The time evolution of each atomic spin from any possible initial structure to its equilibrium state follows a discrete spin model that described by the Landau-Lifshitz-Gilbert (LLG) equation:

(2)
$$\frac{\mathrm{d}\mathbf{m}}{\mathrm{d}t} = -\frac{\gamma}{1+\alpha^2} \Big\{ \mathbf{m} \times \mathbf{B}_{\mathrm{eff}} + \alpha \mathbf{m} \times \big(\mathbf{m} \times \mathbf{B}_{\mathrm{eff}} \big) \Big\}.$$

Here, $\gamma > 0$ represents the gyromagnetic constant, and $\alpha > 0$ denotes the damping constant. The effective field, denoted as \mathbf{B}_{eff} , is computed by

(3)
$$\mathbf{B}_{\rm eff} = -\frac{\delta \mathcal{H}}{\delta \mathbf{m}}.$$




FIGURE 1. Illustration of the hexagonal lattice. Red squares, green triangles, and purple crosses, represent the 1st, 2nd, and 3rd-neighbor nodes of the reference node x_i with $i \in \mathbb{N}$.

In an early work [15], Sulem, Sulem, and Bardos analyzed the system (1)-(3) on a square grid. They considered only the exchange energy in (1) and interactions up to the 1st nearest neighbor, without including the damping term ($\alpha = 0$). By utilizing Ladysenskaya's interpolation operator, they demonstrated that the solution to the discrete spin model can be regarded as an approximation of the weak solution of the continuum LLG equation. In this case, it was found that the exchange term corresponds to Laplacian. A similar outcome is obtained in 1 when accounting for damping. Furthermore, the results of 1D using the difference method are presented in [16]. We refer to [15] which shows that the solution to the discrete spin model converges to the smooth solution of the continuum LLG equation if the initial data for (1)-(3) is sufficiently smooth. However, while only a simple model on a square lattice was considered in [15], we study a more "physical" model with the full system (1)-(3) on a two-dimensional hexagonal lattice. Our main objective is to consider the Cauchy problem for LLG system (1)-(3), and investigate its weak convergence as the rescaled lattice constant of the hexagonal lattice approaches 0 towards degenerated nonlinear equations given by (7)-(8).

The main concept is based on [14,15], but our focus is on a hexagonal lattice, which differs from the classical square lattice discussed in those works. Hence we need to introduce new difference quotients and interpolation operators specific to the hexagonal lattice and study their corresponding properties. The primary challenge lies in defining appropriate difference quotient and interpolation operators for the hexagonal lattice since the loss of symmetry, especially when considering the 1st and 3rd nearest neighborhoods. Indeed, each node in square lattice is a 4-fold rotation center, which possesses the rotation symmetry operation as a cyclic group C_4 . While each node in hexagonal lattice is only a 3-fold rotation center. This loss of symmetry is caused by the fact that the hexagonal lattice can be viewed as a union of two triangular lattices staggered from each other, denoted by $\mathcal{G} = \mathcal{G}_1 \cup \mathcal{G}_2$ (see definition in (10)) where \mathcal{G} represents the nodes of hexagonal lattice. As a consequence, we find a one-step difference quotient can be used to conduct the convergence to the continuum operator for 2nd nearest neighborhoods, since the nodes all appear in only one of the two sets, \mathcal{G}_1 and \mathcal{G}_2 . However, when considering the 1st and 3rd nearest neighborhoods, the one-step difference quotient fails in integration by parts. In order to overcome it, we construct several new multi-step difference quotients. Overall, we introduce the procedure as follows:

- Step 1. Defining the one-step difference quotients for 2st nearest neighborhoods.
- Step 2. Constructing new multi-step difference quotients for the 1st and 3rd nearest neighborhoods, which yield the formula of integration by parts.
- Step 3. Introducing some auxiliary difference quotients to establish the connection between the discrete Laplace operator and our new multi-step difference quotient in Step 2.

The above argument of construction can be applicable to other parabolic problems on a hexagonal lattice. On the other hand, interpolation operators play an important role in connecting solutions of discrete and continuum models. As shown in [14], in order to conduct the weak convergence argument, one needs three different interpolation operators which map the discrete function into step function, piece-wise linear function, and hybrid function that is stepped in one dimension and piece-wise linear in other dimensions. However, constructing the last one on hexagonal lattice is difficult. To address this problem, we provide an alternative strategy to prove the convergence by using only two interpolation operators. Parallel to the work in [14], by utilizing the difference quotients and interpolation operators constructed in this work, we finally derive the convergence from the discrete spin model on hexagonal lattice to the continuum LLG equation. As a by-product, our result provides a precise contribution on physical parameters in the atomistic model (1) to the continuum LLG equation, as listed in Table 1. Of particular interest, the bilinear exchange constant J_k (for k = 1, 2, 3) and the biquadratic exchange constant K collectively form a continuum exchange constant given by $\frac{3}{4}J_1 + \frac{9}{2}J_2 + 3J_3 + \frac{3}{2}K$.

1.1. **Parameterized Cauchy problem.** To begin, we will establish the Cauchy problem for system (1)-(3), with the hexagonal lattice constant being parameterized. We denote h^* as the dimensionless ratio between the lattice constant and characteristic length of the system, and h as a small quantity satisfying $0 \le h \le h^*$. Let \mathbf{m}_h be a discrete function defined on a hexagonal lattice with grid size h. Using these parameters, we can express the energy in a parameterized form:

(4)
$$\mathcal{H}_{h}[\mathbf{m}_{h}] = \frac{(h^{*})^{2}}{h^{2}} \left(\mathcal{H}_{e}[\mathbf{m}] + \mathcal{H}_{b}[\mathbf{m}] \right) + \mathcal{H}_{a}[\mathbf{m}] + \mathcal{H}_{z}[\mathbf{m}].$$

It follows that $\mathcal{H}_h[\mathbf{m}_h] = \mathcal{H}[\mathbf{m}_h]$ when $h = h^*$. The corresponding effective field $\mathbf{B}_{\text{eff},h}(\boldsymbol{x}_h)$ can be calculated by

(5)
$$\mathbf{B}_{\text{eff,h}} = -\frac{\delta \mathcal{H}_h}{\delta \mathbf{m}_h}.$$

Moreover, rescaling time as

$$t^* = \frac{\gamma(h^*)^2}{1+\alpha^2}t,$$

and prescribing the initial data, then we set the Cauchy problem for LLG system (2) with parameter h by

(6)
$$\begin{cases} \frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t^{*}}(\boldsymbol{x}_{h}) = -\frac{1}{(h^{*})^{2}} \Big\{ \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}(\boldsymbol{x}_{h}) \\ + \alpha \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}(\boldsymbol{x}_{h})\big) \Big\}, \\ \mathbf{m}_{h}(\boldsymbol{x}_{h},0) = \mathbf{m}_{\mathrm{init},\mathrm{h}}(\boldsymbol{x}_{h}), \quad |\mathbf{m}_{\mathrm{init},\mathrm{h}}(\boldsymbol{x}_{h})| = 1. \end{cases}$$

As we are interested in phenomena occurring at large scales, compared to the lattice mesh size, we take the limit of (6) as h approaches zero.

1.2. **Main result.** We denote H^s (and L^p) for the Sobolev space $H^s(\mathbb{R}^2; \mathbb{R}^3)$ with $s \ge 1$ (and the Lebesgue space $L^p(\mathbb{R}^2; \mathbb{R}^3)$ with $p \in [1, \infty]$, respectively). Additionally, we denote \dot{H}^s as the homogeneous Sobolev space whose seminorm is determined through Fourier transform:

$$\|u\|_{\dot{H}^s}^2 := \frac{1}{2\pi} \int_{\mathbb{R}^2} |\hat{u}(\boldsymbol{\xi})|^2 |\boldsymbol{\xi}|^{2s} \mathrm{d}\boldsymbol{\xi}, \quad \text{where} \quad \hat{u}(\boldsymbol{\xi}) = \int_{\mathbb{R}^2} e^{-i\boldsymbol{x}\cdot\boldsymbol{\xi}} u(\boldsymbol{x}) \mathrm{d}\boldsymbol{x}.$$

In particular, $||u||_{\dot{H}^1} = ||\nabla u||_{L^2}$.

Formally, taking the limit as $h \to 0$ for the discrete system given by equations (4)-(6) yields the following continuum LLG equation:

(7)
$$\begin{cases} \frac{\partial \mathbf{m}}{\partial t} = -\mathbf{m} \times \mathbf{B}_{\text{eff}} - \alpha \mathbf{m} \times (\mathbf{m} \times \mathbf{B}_{\text{eff}}), \\ \mathbf{m}(\boldsymbol{x}, 0) = \mathbf{m}_{\text{init}}(\boldsymbol{x}), \quad |\mathbf{m}_{\text{init}}(\boldsymbol{x})| = 1. \end{cases}$$

Here the continuum effective field \mathbf{B}_{eff} is given by

(8)
$$\mathbf{B}_{\text{eff}} = J^* \Delta \mathbf{m} + L^* (\mathbf{m} \cdot \mathbf{e}^z) \mathbf{e}^z + \lambda^* (\mathbf{m} \cdot \mathbf{e}^x) \mathbf{e}^x + \mu^* \mathbf{B}.$$

The coefficients J^* , L^* , λ^* , and μ^* are precisely linked to the parameters in the discrete model (4) as listed in Table 1.

TABLE 1. Relationship between parameters in the atomistic and continuum Landau-Lifshitz-Gilbert equations. h^* is the dimensionless lattice constant.

Parameter	Relationship
Exchange	$J^* = \frac{3}{4}J_1 + \frac{9}{2}J_2 + 3J_3 + \frac{3}{2}K$
Anisotropy I	$L^* = (h^*)^{-2}(L_1 + L_2 + L_3)$
Anisotropy II	$\lambda^* = (h^*)^{-2}\lambda$
External	$\mu^* = (h^*)^{-2}\mu$

The weak solution to (7)-(8) is defined in the following sense:

Definition 1.1. Suppose $|\boldsymbol{m}(\boldsymbol{x},t)| = 1$ a.e. in $\mathbb{R}^2 \times [0,T]$. The function \boldsymbol{m} is said to be a weak solution of (7)-(8), if

$$\nabla \boldsymbol{m} \in L^{\infty}(0,T;L^2), \quad \frac{\partial \boldsymbol{m}}{\partial t} \in L^2(0,T;L^2), \quad \nabla \cdot (\boldsymbol{m} \times \nabla \boldsymbol{m}) \in L^2(0,T;L^2),$$

and there holds for any $\boldsymbol{\varphi} \in C^\infty_c(\mathbb{R}^2 \times (0,T))$ that

(9)
$$\int_0^T \int_{\mathbb{R}^2} \partial_t \boldsymbol{m} \cdot \boldsymbol{\varphi} \, \mathrm{d}\boldsymbol{x} \, \mathrm{d}t = -\int_0^T \int_{\mathbb{R}^2} \boldsymbol{m} \times \mathbf{B}_{\mathrm{eff}} \cdot \boldsymbol{\varphi} \, \mathrm{d}\boldsymbol{x} \, \mathrm{d}t \\ -\alpha \int_0^T \int_{\mathbb{R}^2} \boldsymbol{m} \times (\boldsymbol{m} \times \mathbf{B}_{\mathrm{eff}}) \cdot \boldsymbol{\varphi} \, \mathrm{d}\boldsymbol{x} \, \mathrm{d}t$$

with initial value $\mathbf{m}(x, 0) = \mathbf{m}_{init}(x)$.

Note that the definition above is reasonable because $\mathbf{m} \times \Delta \mathbf{m} = \nabla \cdot (\mathbf{m} \times \nabla \mathbf{m}) \in L^2(0,T;L^2).$

Now let us state the main results.

Theorem 1.1. (Convergence). Suppose the Assumption (1) holds. Assume the initial data satisfies $\nabla \boldsymbol{m}_{init}(\boldsymbol{x}) \in L^2(\mathbb{R}^2)$ in equation (7). Given $\boldsymbol{m}_{init,h}(\boldsymbol{x}) \in \widetilde{H}^1_{loc,h}(\mathcal{G})$ and $p_h \boldsymbol{m}_{init,h} \to \boldsymbol{m}_{init}$ strongly in $\dot{H}^1(\mathbb{R}^2)$, the discrete LLG equation (4)-(6) admits a unique solution \boldsymbol{m}_h . The solution \boldsymbol{m}_h

converges to a weak solution \mathbf{m}^* to the continuum LLG equation (7)-(8) in the following way,

 $p_h \boldsymbol{m}_h, \ q_h \boldsymbol{m}_h \to \boldsymbol{m}^* \text{ strongly in } C([0,T]; L^2_{loc}(\mathbb{R}^2)),$ $p_h \boldsymbol{m}_h \rightharpoonup \boldsymbol{m}^* \text{ weakly in } L^{\infty}(0,T; \dot{H}^1(\mathbb{R}^2)).$

Here we note that q_h , p_h are linear interpolation operators on a hexagonal lattice, whose definitions and properties will be introduced in Section 4 and Propositions 3.3 and 3.4. For a vector-valued function \mathbf{m}_h , $q_h \mathbf{m}_h$, $p_h \mathbf{m}_h$ are defined to be $(q_h m_h^1, q_h m_h^2, q_h m_h^3)$ and $(p_h m_h^1, p_h m_h^2, p_h m_h^3)$ respectively.

Remark 1.1. The uniqueness of the weak solution obtained in Theorem 1.1 is unclear, as discussed in [1]. However, if a unique strong solution $m \in C^3(\mathbb{R}^2 \times [0,T])$ exists, then it can be shown that $m^* = m$, which is referred to as "weak-strong uniqueness" [7].

This article is organized as follows. Section 2 introduces the notation of difference quotients on a hexagonal lattice and derives the discrete formula for integration by parts. In Section 3, we deduce the weak convergence procedure and prove Theorem 1.1 using a priori energy estimate, leaving the calculation for later. Section 4 presents the construction and proof of properties of proper interpolation operators on a hexagonal lattice. Finally, in the Appendix, we transform the LLG system (6) into a discrete PDE type equation and derive uniform estimates for discrete solutions.

2. NOTATIONS ON HEXAGONAL LATTICE

Let \mathcal{G} denote the collection of all the nodes of hexagonal lattice, with the parameterized side length h > 0. Note that it takes account up to three layers of nearest neighbors of point-to-point interactions in the energy (1). For any point $\boldsymbol{x}_h \in \mathcal{G}$, we use N_k , k = 1, 2, 3, to represent the number of nearest neighbors of \boldsymbol{x}_h in the *k*th layer, with

$$N_1 = 3, N_2 = 6, N_3 = 3,$$

and write the *j*th neighbor in the *k*th layer of \boldsymbol{x}_h by $\boldsymbol{x}_h^{k,j}$ where $j = 1, \ldots, N_k$. Figure 2 shows the arrangement mode of $\boldsymbol{x}_h^{k,j}$ in relation to \boldsymbol{x}_h . It is worth noting that the points in \mathcal{G} can be classified into two categories based on the distribution of their 1st-layer nearest neighbors: regular triangle type (Figure 2(a)) and inverted triangle type (Figure 2(b)). Let us write $\mathcal{G} = \mathcal{G}_1 \cup \mathcal{G}_2$, where

(10)

 $\mathcal{G}_1 = \{ \boldsymbol{x}_h \in \mathcal{G} \mid ext{the nearest neighbors of } \boldsymbol{x}_h ext{ form a regular triangle } \}, \ \mathcal{G}_2 = \{ \boldsymbol{x}_h \in \mathcal{G} \mid ext{the nearest neighbors of } \boldsymbol{x}_h ext{ form an inverted triangle } \}.$




FIGURE 2. The 1st, 2nd, and 3rd nearest neighbors of $\boldsymbol{x}_h \in \mathcal{G}$ with two categories: (a), $\boldsymbol{x}_h \in \mathcal{G}_1$; (b), $\boldsymbol{x}_h \in \mathcal{G}_2$

2.1. One-step difference quotients and discrete space. Let us introduce the difference quotients and Laplace operator on a hexagonal lattice. We can define the one-step difference quotient for the kth nearest neighbors and jth direction as follows:

(11)
$$\widetilde{D}^{k,j}u_h(\boldsymbol{x}_h) = \frac{u_h(\boldsymbol{x}_h^{k,j}) - u_h(\boldsymbol{x}_h)}{d_k h} \quad j = 1, \dots, N_k, \quad k = 1, 2, 3.$$

Here the normalized constants $d_1 = 1$, $d_2 = \sqrt{3}$, and $d_3 = 2$. The notation of backward difference quotient is necessary for the formula of integration by parts. Define the backward difference quotient

(12)
$$\widetilde{D}_B^{k,j} = \begin{cases} -\widetilde{D}^{k,j}, & \text{if } k = 1, 3\\ -\widetilde{D}^{k,j^*}, & \text{if } k = 2 \end{cases}$$

with $j^* = (j+3) \mod 6$. It follows by the definition that

(13)
$$\widetilde{D}^{k,j}u_h(\boldsymbol{x}_h) = \widetilde{D}_B^{k,j}u_h(\boldsymbol{x}_h^{k,j}).$$

The discrete Laplace operator in the kth nearest neighbors is given by

(14)
$$\widetilde{\Delta}_k u_h(\boldsymbol{x}_h) = \sum_{j=1}^{N_k} \frac{u_h(\boldsymbol{x}_h^{k,j}) - u_h(\boldsymbol{x}_h)}{l_k h^2}$$

with $l_1 = 3/4$, $l_2 = 9/2$, and $l_3 = 3$. The gradient operator in the kth nearest neighbors on hexagonal lattice is denoted by

(15)
$$\widetilde{\nabla}_k = g_k(\widetilde{D}^{k,1}, \widetilde{D}^{k,2}, \cdots, \widetilde{D}^{k,N_k})$$

with $g_1 = g_3 = \sqrt{6}/3, g_2 = \sqrt{3}/3.$

We introduce some function spaces on a hexagonal lattice. The Hilbert space with an inner product given by

(16)
$$(u_h, v_h)_h = \frac{3\sqrt{3}}{4} h^2 \sum_{\boldsymbol{x}_h \in \mathcal{G}} u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h).$$

And \widetilde{H}_h^1 denotes the Hilbert space with inner product given by

$$(u_h, v_h)_{\widetilde{H}_h^1} = (u_h, v_h)_h + (\widetilde{\nabla}_1 u_h, \widetilde{\nabla}_1 v_h)_h.$$

2.2. Integration by parts for one-step difference quotients. The integration by parts formula is crucial in energy estimation. The following theorems indicate the one-step difference quotients satisfy the formula of integration by parts and Green's formula on hexagonal lattice.

Lemma 2.1. For every function $u_h(\boldsymbol{x}_h)$, $v_h(\boldsymbol{x}_h)$ defined on \mathcal{G} , it holds that (17)

$$u_h(\boldsymbol{x}_h) \cdot \widetilde{D}^{k,j} v_h(\boldsymbol{x}_h) + \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h^{k,j}) \cdot v_h(\boldsymbol{x}_h^{k,j}) = \widetilde{D}^{k,j} \{ u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h) \}$$

and

(18)
$$\widetilde{\Delta}_k u_h(\boldsymbol{x}_h) = \frac{d_k^2}{2l_k} \sum_{j=1}^{N_k} \widetilde{D}_B^{k,j} \{ \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h) \},$$

where k = 1, 2, 3 and $j = 1, ..., N_k$.

Proof. The formula (17) follows directly from the definition (11) and the equation (13). Now let us prove the formula (18). It follows from (11) and (13) that

$$\begin{split} \widetilde{D}^{k,j} \big\{ D_B^{k,j} u_h(\boldsymbol{x}_h) \big\} = & \frac{1}{d_k h} \big\{ \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h^{k,j}) - \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h) \big\} \\ = & \frac{1}{d_k h} \big\{ \widetilde{D}^{k,j} u_h(\boldsymbol{x}_h) - \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h) \big\}. \end{split}$$

Summing up for $j = 1, \ldots, N_k$ yields

(19)
$$\sum_{j=1}^{N_k} \widetilde{D}_B^{k,j} \left\{ \widetilde{D}_B^{k,j} u_h(\boldsymbol{x}_h) \right\} = \frac{2}{d_k h} \sum_{j=1}^{N_k} \widetilde{D}_{k,j}^{k,j} u_h(\boldsymbol{x}_h).$$

On the other hand, it follows from (14)

(20)
$$\widetilde{\Delta}_k u_h(\boldsymbol{x}_h) = \frac{d_k}{l_k h} \sum_{j=1}^{N_k} \widetilde{D}^{k,j} u_h(\boldsymbol{x}_h).$$

Hence the formula (18) is proved.

Since $\sum_{\boldsymbol{x}_h \in \mathcal{G}} \widetilde{D}^{k,j} f_h(\boldsymbol{x}_h) = 0$, the equation (17) indicates the formula of integration by parts

$$\left(u_h, \, \widetilde{D}^{k,j} v_h\right)_h + \left(\widetilde{D}_B^{k,j} u_h, \, v_h\right)_h = 0.$$

However, it is important to point out that the one-step difference quotient $D^{k,j}$ has opposite directions on adjacent lattice nodes, which makes the weak convergence process fail. To fix this problem, we introduce multiplier

$$\kappa(oldsymbol{x}_h) = egin{cases} 1, & ext{if} oldsymbol{x}_h \in \mathcal{G}_1, \ -1, & ext{if} oldsymbol{x}_h \in \mathcal{G}_2, \end{cases}$$

such that $\kappa \widetilde{D}^{k,j} u_h$ retains the uniform direction on the lattice. When k = 2, note that $\kappa(\boldsymbol{x}_h) = \kappa(\boldsymbol{x}_h^{k,j})$, we can deduce directly the following formula of integrating by parts.

Proposition 2.1. For every function $u_h(\boldsymbol{x}_h)$, $v_h(\boldsymbol{x}_h)$ defined on \mathcal{G} , it holds that

(21)
$$(u_h, \kappa \widetilde{D}^{2,j} v_h)_h + (\kappa \widetilde{D}_B^{2,j} u_h, v_h)_h = 0.$$

2.3. Multi-step difference quotients. For the case k = 1 or 3, the operator $\kappa D^{k,j}$ does not satisfy the integration by parts property (21). To overcome this problem, we introduce multi-step difference quotients for the 1st and 3rd nearest neighbors. We denote the two-step difference quotient \mathbb{D}_x^k for k = 1, 3, which is given by

(22)
$$\widetilde{\mathbb{D}}_{x}^{k}u_{h}(\boldsymbol{x}_{h}) = \frac{\sqrt{3}}{3} \big\{ \kappa \widetilde{D}^{k,3} - \kappa \widetilde{D}^{k,2} \big\} u_{h}(\boldsymbol{x}_{h}),$$

and $\widetilde{\mathbb{D}}_y^k$ denotes the three-step difference quotient in the kth nearest neighbors that is constructed as

(23)
$$\widetilde{\mathbb{D}}_{y}^{k}u_{h}(\boldsymbol{x}_{h}) = \frac{1}{3} \left\{ 2\kappa \widetilde{D}^{k,1} - \kappa \widetilde{D}^{k,2} - \kappa \widetilde{D}^{k,3} \right\} u_{h}(\boldsymbol{x}_{h}).$$

We will use the operators $\widetilde{\mathbb{D}}^k_x$ and $\widetilde{\mathbb{D}}^k_y$ to conduct a weak convergence procedure when k equals 1 or 3. In the following we prove that operators $\widetilde{\mathbb{D}}_x^k$ and $\widetilde{\mathbb{D}}_{y}^{k}$ yield the formula of integration by parts.

Proposition 2.2. When k = 1, 3, for every $u_h(\boldsymbol{x}_h)$, $v_h(\boldsymbol{x}_h)$ defined on \mathcal{G} , it holds that

(24)
$$(u_h, \widetilde{\mathbb{D}}_x^k v_h)_h + (\widetilde{\mathbb{D}}_x^k u_h, v_h)_h = 0.$$

and

(25)
$$(u_h, \widetilde{\mathbb{D}}_y^k v_h)_h + (\widetilde{\mathbb{D}}_y^k u_h, v_h)_h = 0.$$

Proof. Multiplying the equation (17) by $\kappa(\boldsymbol{x}_h)$ yields

$$u_h(\boldsymbol{x}_h) \cdot \kappa(\boldsymbol{x}_h) \widetilde{D}^{k,j} v_h(\boldsymbol{x}_h) + \kappa(\boldsymbol{x}_h^{k,j}) \widetilde{D}^{k,j} u_h(\boldsymbol{x}_h^{k,j}) \cdot v_h(\boldsymbol{x}_h^{k,j}) \\ = \kappa(\boldsymbol{x}_h) \widetilde{D}^{k,j} \{ u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h) \}.$$

Applying the definition in (22), one can deduce from above equation

$$u_h(\boldsymbol{x}_h) \cdot \widetilde{\mathbb{D}}_x^k v_h(\boldsymbol{x}_h) + \widetilde{\mathbb{D}}_x^k u_h(\boldsymbol{x}_h^{k,j}) \cdot v_h(\boldsymbol{x}_h^{k,j}) = \widetilde{\mathbb{D}}_x^k \{ u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h) \},\$$

By summing (2.3) over all $\boldsymbol{x}_h \in \mathcal{G}$, we obtain the equation (24), which follows from the fact that $\sum_{\boldsymbol{x}_h \in \mathcal{G}} \widetilde{\mathbb{D}}_x^k f_h(\boldsymbol{x}_h) = 0$. The equation (25) can be proved in the same way, for which we omit the details.

2.4. Auxiliary difference quotients. Finally, it remains to establish the connection between discrete Laplace operator (18) and the multi-step difference quotients (22)-(23). To achieve this, we need some auxiliary difference quotients. For k = 1, 3 and r = 0, 1, define

(26)
$$\widetilde{\mathfrak{D}}_{r}^{k}u_{h}(\boldsymbol{x}_{h}) = \begin{cases} \left(\frac{2}{3}\right)^{r}\widetilde{D}^{k,1}u_{h}(\boldsymbol{x}_{h}), & \text{for } \boldsymbol{x}_{h} \in \mathcal{G}_{1} \\ \left(\frac{1}{3}\right)^{r}\left\{\widetilde{D}^{k,2} + \widetilde{D}^{k,3}\right\}u_{h}(\boldsymbol{x}_{h}), & \text{for } \boldsymbol{x}_{h} \in \mathcal{G}_{2}, \end{cases}$$

and its backward difference quotient is given by

(27)
$$\widetilde{\mathfrak{D}}_{r,B}^{k}u_{h}(\boldsymbol{x}_{h}) = \begin{cases} -\left(\frac{1}{3}\right)^{r} \{\widetilde{D}^{k,2} + \widetilde{D}^{k,3}\}u_{h}(\boldsymbol{x}_{h}), & \text{for } \boldsymbol{x}_{h} \in \mathcal{G}_{1}, \\ -\left(\frac{2}{3}\right)^{r} \widetilde{D}^{k,1}u_{h}(\boldsymbol{x}_{h}), & \text{for } \boldsymbol{x}_{h} \in \mathcal{G}_{2}. \end{cases}$$

By the definitions of (23)-(27), one can directly deduce that

(28)
$$\widetilde{\mathbb{D}}_{y}^{k}u_{h}(\boldsymbol{x}_{h}) = \{\widetilde{\mathfrak{D}}_{1}^{k} + \widetilde{\mathfrak{D}}_{1,B}^{k}\}u_{h}(\boldsymbol{x}_{h}),$$

Moreover, we have the following decomposition of the quadratic form of one-step operator $\widetilde{D}^{k,j}$:

Lemma 2.2. For any u_h , v_h defined on \mathcal{G} and k = 1, 3, the following decomposition holds:

(29)
$$\sum_{j=1}^{3} \widetilde{D}^{k,j} u_h \cdot \widetilde{D}^{k,j} v_h$$
$$= \frac{3}{2} \{ \widetilde{\mathfrak{D}}_1^k u_h \cdot \widetilde{\mathfrak{D}}_0^k v_h + \widetilde{\mathfrak{D}}_{1,B}^k u_h \cdot \widetilde{\mathfrak{D}}_{0,B}^k v_h + \widetilde{\mathbb{D}}_x^k u_h \cdot \widetilde{\mathbb{D}}_x^k v_h \}.$$

Proof. Note that for any u_h , v_h defined on \mathcal{G} , one has

$$\begin{split} \sum_{j=1}^{3} \widetilde{D}^{k,j} u_h \cdot \widetilde{D}^{k,j} v_h = \widetilde{D}^{k,1} u_h \cdot \widetilde{D}^{k,1} v_h + \frac{1}{2} \{ \widetilde{D}^{k,2} + \widetilde{D}^{k,3} \} u_h \cdot \{ \widetilde{D}^{k,2} + \widetilde{D}^{k,3} \} v_h \\ + \frac{1}{2} \{ \widetilde{D}^{k,2} - \widetilde{D}^{k,3} \} u_h \cdot \{ \widetilde{D}^{k,2} - \widetilde{D}^{k,3} \} v_h, \end{split}$$

then (29) follows by substituting the definitions (22)-(27) into the above equation.

The formula (29) gives us the connection between the discrete Laplace operator and the multi-step difference quotients $\widetilde{\mathbb{D}}_x^k$, $\widetilde{\mathbb{D}}_y^k$, by the help of (18) and (28). We remark that, the auxiliary operators $\widetilde{\mathfrak{D}}_r^k$ and $\widetilde{\mathfrak{D}}_{r,B}^k$ cannot be used to conduct weak convergence directly, since they neither have uniform direction on the lattice when r = 0, nor yield the integration by parts when r = 1.

With these notations, we conclude the following convergence result:

Lemma 2.3. Assume that q_h is the interpolation operator given by Definition 4.1. For $\varphi \in C_c^2(\mathbb{R}^2)$, let $\varphi_h(\boldsymbol{x}_h) = \varphi(\boldsymbol{x}_h)$. When k = 1, 3, it holds that

(30)
$$q_h\{\widetilde{\mathbb{D}}_x^k\varphi_h\} \to (-1)^i \frac{\partial}{\partial x}\varphi, \quad \text{in } L^2$$
$$q_h\{\widetilde{\mathfrak{D}}_0^k\varphi_h\}, \ q_h\{\widetilde{\mathfrak{D}}_{0,B}^k\varphi_h\}, \ q_h\{\widetilde{\mathbb{D}}_y^k\varphi_h\} \to (-1)^i \frac{\partial}{\partial y}\varphi \quad \text{in } L^2$$

with i = 0 for k = 1, and i = 1 for k = 3. When k = 2, it holds that

(31)
$$q_h \{\kappa \widetilde{D}^{2,j} \varphi_h\}, \ q_h \{\kappa \widetilde{D}_B^{2,j} \varphi_h\} \to \frac{\partial}{\partial \nu^j} \varphi \quad in \ L^2, \quad j = 1, \cdots, 6,$$

here ν^{j} denotes the fixed unit vector independent of \mathbf{x} , given by $\nu^{1} = -\nu^{4} = (-1/2, \sqrt{3}/2), \ \nu^{2} = -\nu^{5} = (-1, 0), \ \nu^{3} = -\nu^{6} = (-1/2, -\sqrt{3}/2).$

Lemma 2.3 follows directly from Taylor expansion and the property (iii) of q_h , so we omit the details here.

3. Proof of Theorem 1.1

In this section, we will provide an overview of the proof for Theorem 1.1. We recall that the discrete LLG equation reads:

(32)
$$\begin{cases} \frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t^{*}}(\boldsymbol{x}_{h}) = -\frac{1}{(h^{*})^{2}} \Big\{ \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}(\boldsymbol{x}_{h}) \\ + \alpha \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}(\boldsymbol{x}_{h})\big) \Big\}, \\ \mathbf{m}_{h}(\boldsymbol{x}_{h},0) = \mathbf{m}_{\mathrm{init},\mathrm{h}}(\boldsymbol{x}_{h}), \quad |\mathbf{m}_{\mathrm{init},\mathrm{h}}(\boldsymbol{x}_{h})| = 1. \end{cases}$$

By virtue of Cauchy-Lipshitz-Picard Theorem [4], it is guaranteed that there exists a unique solution $\mathbf{m}_h \in C^1([0, +\infty))$ for the ODE system (32). Let us revisit the discrete energy (4) and provide the expression for the total effective field in (5). We write $\mathbf{B}_{\text{eff},h}(\boldsymbol{x}_h) = \mathbf{B}_1(\boldsymbol{x}_h) + \mathbf{B}_2(\boldsymbol{x}_h) + \mathbf{B}_3(\boldsymbol{x}_h) + \mathbf{B}_4(\boldsymbol{x}_h)$, where

$$\begin{split} \mathbf{B_1}(\boldsymbol{x_h}) &= -\frac{(h^*)^2}{h^2} \frac{\delta \mathcal{H}_e}{\delta \mathbf{m}_h}, \qquad \mathbf{B_2}(\boldsymbol{x_h}) = -\frac{(h^*)^2}{h^2} \frac{\delta \mathcal{H}_b}{\delta \mathbf{m}_h}, \\ \mathbf{B_3}(\boldsymbol{x_h}) &= -\frac{\delta \mathcal{H}_a}{\delta \mathbf{m}_h}, \qquad \mathbf{B_4}(\boldsymbol{x_h}) = -\frac{\delta \mathcal{H}_z}{\delta \mathbf{m}_h}. \end{split}$$

Then one can calculate from (4) that

(33)

$$\begin{cases}
\mathbf{B}_{1}(\boldsymbol{x}_{h}) = (h^{*})^{2} \sum_{k=1}^{3} \sum_{j=1}^{N_{k}} \frac{J_{k}}{h^{2}} \mathbf{m}_{h}(\boldsymbol{x}_{h}^{k,j}) \\
\mathbf{B}_{2}(\boldsymbol{x}_{h}) = 2(h^{*})^{2} \sum_{j=1}^{3} \frac{K}{h^{2}} \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{m}_{h}(\boldsymbol{x}_{h}^{1,j})\right) \mathbf{m}_{h}(\boldsymbol{x}_{h}^{1,j}), \\
\mathbf{B}_{3}(\boldsymbol{x}_{h}) = \sum_{k=1}^{3} \sum_{j=1}^{N_{k}} \left(L_{k}(\boldsymbol{x}_{h}) + L_{k}(\boldsymbol{x}_{h}^{k,j})\right) \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}^{k,j}) \cdot \mathbf{e}^{z}\right) \mathbf{e}^{z} \\
+ 2\lambda(\boldsymbol{x}_{h}) \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{x}\right) \mathbf{e}^{x}, \\
\mathbf{B}_{4}(\boldsymbol{x}_{h}) = \mu \mathbf{B}(\boldsymbol{x}_{h}).
\end{cases}$$

3.1. Energy estimate. In the following, we introduce the discrete PDE type equation:

(34)
$$\frac{\mathrm{d}\mathbf{m}_h}{\mathrm{d}t^*} = -\mathbf{m}_h \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}} - \alpha \mathbf{m}_h \times (\mathbf{m}_h \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}}),$$

where the new effective field, denoted as $\mathbf{B}_{\text{eff},h}^*$, is expressed using the difference quotient notation. It is given by the equation $\mathbf{B}_{\text{eff},h}^*(\boldsymbol{x}_h) = \mathbf{B}_1^*(\boldsymbol{x}_h) + \mathbf{B}_2^*(\boldsymbol{x}_h) + \mathbf{B}_3^*(\boldsymbol{x}_h) + \mathbf{B}_4^*(\boldsymbol{x}_h)$ with $\mathbf{B}_3^* = \mathbf{B}_3^{(1)} + \mathbf{B}_3^{(2)}$, which can computed by: (35) $\begin{cases} \mathbf{B}_1^*(\boldsymbol{x}_h) = \sum_{k=1}^3 l_k J_k \widetilde{\Delta}_k \mathbf{m}_h(\boldsymbol{x}_h) \\ \mathbf{B}_2^*(\boldsymbol{x}_h) = 2l_1 K \widetilde{\Delta}_1 \mathbf{m}_h(\boldsymbol{x}_h) + 2K \sum_{j=1}^3 \left(\mathbf{m}_h(\boldsymbol{x}_h) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h)\right) \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h), \\ \mathbf{B}_3^{(1)}(\boldsymbol{x}_h) = \frac{2}{(h^*)^2} \lambda(\boldsymbol{x}_h) \left(\mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^x\right) \mathbf{e}^x + \frac{2}{(h^*)^2} \sum_{k=1}^3 N_k L_k(\boldsymbol{x}_h) \left(\mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z\right) \mathbf{e}^z, \\ \mathbf{B}_3^{(2)}(\boldsymbol{x}_h) = \frac{h^2}{(h^*)^2} \sum_{k=1}^3 l_k \left\{ L_k(\boldsymbol{x}_h) \left(\widetilde{\Delta}_k \mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \right) \mathbf{e}^z + \left(\widetilde{\Delta}_k(L_k \mathbf{m}_h)(\boldsymbol{x}_h) \cdot \mathbf{e}^z \right) \mathbf{e}^z \right\} \\ \mathbf{B}_4^*(\boldsymbol{x}_h) = \frac{\mu}{(h^*)^2} \mathbf{B}(\boldsymbol{x}_h). \end{cases}$

The relation between the equation above and the descrete LLG equation (32)-(33) is concluded in the following proposition.

Proposition 3.1. Let m_h be the solution of equation (32) with effective field $\mathbf{B}_{\text{eff},h}$ given by (33). Then m_h also satisfies the discrete equation (34), with a PDE type effective field $\mathbf{B}_{\text{eff},h}^*(\boldsymbol{x}_h)$ given by (35).

Moreover, the solution \mathbf{m}_h has the following energy estimate.

Proposition 3.2. Suppose the Assumption (1) holds. If m_h is the solution of the equations (34)-(35), it yields

(36)
$$|\boldsymbol{m}_h(\boldsymbol{x}_h, t)| = 1, \quad \sup_{t \ge 0} \|\widetilde{\nabla}_k \boldsymbol{m}_h(\cdot, t)\|_{\widetilde{L}^2_h}^2 \le C,$$

Furthermore, given T > 0, it satisfies the following estimates

(37)
$$\int_0^T \|\boldsymbol{m}_h \times \mathbf{B}^*_{\text{eff},h}\|_{\tilde{L}^2_h}^2 \mathrm{d}t \le C, \quad \int_0^T \|\frac{\partial \boldsymbol{m}_h}{\partial t}\|_{\tilde{L}^2_h}^2 \mathrm{d}t \le C,$$

where C is a constant that depends on the initial data, $\|\mathbf{B}\|_{L^1}$, $\|L\|_{L^1}$, and $\|\lambda\|_{L^1}$, but is independent of h.

The Proposition 3.1 and 3.2 will be verified in the Appendix.

3.2. Generalization of Ladysenskaya's interpolation operators. We now introduce interpolation operators, denoted as q_h and p_h , which are defined in Definitions 4.1 and 4.2. The construction of these operators is explained in Section 4. In order to prove Theorem 1.1, it is necessary for the operators q_h and p_h to satisfy certain properties.

Proposition 3.3. For any u_h , $v_h \in \widetilde{L}_h^2$, the linear interpolation q_h defined in Definition 4.1 satisfies

- (i) (isometric mapping) $\int_{\mathbb{R}^2} q_h u_h \cdot q_h v_h d\mathbf{x} = (u_h, v_h)_h$,
- (ii) $q_h\{u_h \cdot v_h\} = q_h u_h \cdot q_h v_h,$ (iii) $\inf_{\boldsymbol{x}_h \in V_E} u_h(\boldsymbol{x}_h) \leq q_h u_h(\boldsymbol{x}) \leq \sup_{\boldsymbol{x}_h \in V_E} u_h(\boldsymbol{x}_h), \text{ for any } \boldsymbol{x} \in E.$

Here E represents a unit cell in the hexagonal lattice, and V_E is its vertex point.

Proposition 3.4. For any $u_h \in \widetilde{L}_h^2$, $v_h \in \widetilde{H}_h^1$, the linear interpolation p_h defined in Definition 4.2 satisfies

- (*iv*) $||p_h u_h||_{L^2} \le C ||u_h||_{\widetilde{L}^2_h}$, $||\nabla p_h v_h||_{L^2} \le C ||\widetilde{\nabla}_1 v_h||_{\widetilde{L}^2_h}$,
- (v) $||p_h v_h q_h v_h||_{L^2} \le h ||\widetilde{\nabla}_1 v_h||_{\widetilde{L}^2_{+}}$

where the constant C is independent of u_h , v_h and h.

For brevity, the proofs of Propositions 3.3 and 3.4 are left in Section 4.3.

Remark 3.1. Here, the required properties are provided to support the weak convergence argument.

Remark 3.2. Similar interpolation operators p_h and q_h were introduced in [14] for the square lattice case. For any discrete function u_h , the function $q_h u_h$ is constructed as a step function in \mathbb{R}^2 , and $p_h u_h$ is constructed as a piece-wise linear function in \mathbb{R}^2 .

Besides q_h and p_h , one more interpolation operator r_h was suggested in [14] to conduct the weak convergence of derivatives. More precisely, denote a differential operator D and its discrete form D, and given a convergent sequence

 $q_h u_h, p_h u_h \to u \text{ strongly}, \qquad q_h \widetilde{D} u_h \to v \text{ weakly}.$

in L^2 . r_h was introduced to prove v = Du.

Defining an interpolation operator r_h on a hexagonal lattice poses difficulties. Instead, we propose an alternative strategy to prove v = Du in this work. This approach will be applied in the upcoming proof of (43) and (44).

3.3. Convergence of \mathbf{m}_h and its derivatives. In this subsection, we will prove the convergence of the discrete solution \mathbf{m}_h and its derivatives, with the aid of the uniform estimates in Proposition 3.2 and interpolation operator q_h and p_h which satisfy Propositions 3.3 and 3.4.

Assume that q_h and p_h are interpolation operators satisfying Propositions 3.3 and 3.4. Applying properties (i) and (iv), one can deduce from (36)-(37)

(38) $p_{h}\mathbf{m}_{h} \text{ remains in a bounded set of } L^{\infty}(0,T;\dot{H}^{1}) \cap L^{\infty}(0,T;L^{\infty}),$ $\frac{d}{dt}p_{h}\mathbf{m}_{h}, \frac{d}{dt}q_{h}\mathbf{m}_{h} \text{ remain in a bounded set of } L^{2}(0,T;L^{2}).$

As a consequence of Aubin-Lions Lemma, there exists a subsequence of $p_h \mathbf{m}_h$ converging strongly to some function \mathbf{m}^* in $C([0, T]; L^2_{\text{loc}})$. Due to the property (v) and (36), it as well holds the convergence

(39)
$$q_h \mathbf{m}_h \to \mathbf{m}^* \text{ strongly in } C([0,T]; L^2_{\text{loc}}),$$

and thus converges almost everywhere. Note that $|q_h \mathbf{m}_h(\boldsymbol{x})| = 1$ for all $\boldsymbol{x} \in \mathbb{R}^2$, it follows from the properties (ii)-(iii) that

$$|\mathbf{m}^*| = 1 \quad \text{a.e in } \mathbb{R}^2.$$

Moreover, the statement (38) tells that a subsequence of $q_h \mathbf{m}_h$ (which we also denote by $q_h \mathbf{m}_h$) satisfies

(41)
$$\frac{\partial}{\partial t}q_h\mathbf{m}_h \to \frac{\partial \mathbf{m}^*}{\partial t}$$
 weakly in $L^2(0,T;L^2)$.

Now we consider the convergence of spatial derivatives. When k = 2, according to the estimate (36), $q_h \{\kappa \tilde{D}^{k,j} \mathbf{m}_h\}$ remains in a bounded set of $L^{\infty}(0,T,L^2)$. Thus it has a subsequence which converges to some limit function weakly * in $L^{\infty}(0,T,L^2)$. We assert that, this limit function is $\frac{\partial \mathbf{m}^*}{\partial \nu^j}$. In fact, due to the formula of integration by parts and the property (i) for q_h , it holds that

$$\int_{\mathbb{R}^2} q_h \{\kappa \widetilde{D}^{2,j} \mathbf{m}_h\} \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} = -\int_{\mathbb{R}^2} q_h \mathbf{m}_h \cdot q_h \{\kappa \widetilde{D}_B^{2,j} \boldsymbol{\varphi}_h\} \mathrm{d}\boldsymbol{x},$$

here, φ_h is the discrete function satisfying $\varphi_h(\boldsymbol{x}_h) = \varphi(\boldsymbol{x}_h)$ for every $\varphi(x) \in C_c^{\infty}(\mathbb{R}^2)$. By virtue of Lemma 2.3 and (39), one has

(42)
$$\int_{\mathbb{R}^2} q_h \mathbf{m}_h \cdot q_h \left\{ \kappa \widetilde{D}_B^{2,j} \boldsymbol{\varphi}_h \right\} \mathrm{d}\boldsymbol{x} \to \int_{\mathbb{R}^2} \mathbf{m}^* \cdot \frac{\partial \boldsymbol{\varphi}}{\partial \nu_j} \mathrm{d}\boldsymbol{x}$$

16

On the other hand

$$\begin{split} \left| \int_{\mathbb{R}^2} q_h \{ \kappa \widetilde{D}^{2,j} \mathbf{m}_h \} \cdot \boldsymbol{\varphi} \mathrm{d}\boldsymbol{x} - \int_{\mathbb{R}^2} q_h \{ \kappa \widetilde{D}^{2,j} \mathbf{m}_h \} \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} \right| \\ = \left| \int_{\mathbb{R}^2} q_h \{ \kappa \widetilde{D}^{2,j} \mathbf{m}_h \} \cdot (\boldsymbol{\varphi} - q_h \boldsymbol{\varphi}_h) \mathrm{d}\boldsymbol{x} \right| \le \left\| q_h \{ \kappa \widetilde{D}^{2,j} \mathbf{m}_h \} \right\|_{L^2} \| \boldsymbol{\varphi} - q_h \boldsymbol{\varphi}_h \|_{L^2}, \end{split}$$

which together with (42) implies that

(43)
$$q_h \{ \kappa \widetilde{D}^{2,j} \mathbf{m}_h \} \to \frac{\partial \mathbf{m}^*}{\partial \nu^j} \quad \text{weakly in } L^\infty(0,T,L^2),$$

for all $j = 1, \cdots, 6$.

As for the case k = 1, 3, note that the sequence $\widetilde{\mathbb{D}}_x^k \mathbf{m}_h$ and $\widetilde{\mathbb{D}}_y^k \mathbf{m}_h$ are also bounded in $L^{\infty}(0, T, \widetilde{L}_h^2)$ by (36), one can repeat the above step and use the result of integration by parts in Lemma 2.2 and strong convergence in Lemma 2.3, to derive that

(44)
$$q_h\{\widetilde{\mathbb{D}}_x^k\mathbf{m}_h\} \to \frac{\partial \mathbf{m}^*}{\partial x}, \quad q_h\{\widetilde{\mathbb{D}}_y^k\mathbf{m}_h\} \to \frac{\partial \mathbf{m}^*}{\partial y},$$

weakly in $L^{\infty}(0, T, L^2)$.

3.4. Convergence of degenerated Laplace term. Let us consider the limit for the integral of degenerated Laplace term

$$T_h := \int_{\mathbb{R}^2} q_h \{ \mathbf{m}_h \times \widetilde{\Delta}_k \mathbf{m}_h \} \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x}.$$

According to Lemma 2.1 and the property (i) of q_h ,

(45)
$$T_h = -\frac{d_k^2}{2l_k} \sum_{j=1}^{N_k} \int_{\mathbb{R}^2} q_h \{ \mathbf{m}_h \times \widetilde{D}^{k,j} \mathbf{m}_h \} \cdot q_h \{ \widetilde{D}^{k,j} \boldsymbol{\varphi}_h \} \mathrm{d}\boldsymbol{x}.$$

Now we assert that

(46)

$$\lim_{h\to 0} T_h = -\int_{\mathbb{R}^2} \mathbf{m}^* \times \nabla \mathbf{m}^* \cdot \nabla \varphi d\boldsymbol{x} = \int_{\mathbb{R}^2} \left\{ \nabla \cdot (\mathbf{m}^* \times \nabla \mathbf{m}^*) \right\} \cdot \varphi d\boldsymbol{x}.$$

Proof of (46). The proof is divided into two cases: k = 2 and k = 1, 3. For the case k = 2, due to the fact $\kappa^2(\boldsymbol{x}_h) = 1$, one has

$$T_{h} = -\frac{d_{k}^{2}}{2l_{k}} \sum_{j=1}^{N_{k}} \int_{\mathbb{R}^{2}} q_{h} \{\mathbf{m}_{h} \times \kappa \widetilde{D}^{k,j} \mathbf{m}_{h}\} \cdot q_{h} \{\kappa \widetilde{D}^{k,j} \boldsymbol{\varphi}_{h}\} \mathrm{d}\boldsymbol{x}$$
$$= -\frac{1}{3} \sum_{j=1}^{6} \int_{\mathbb{R}^{2}} q_{h} \mathbf{m}_{h} \times q_{h} \{\kappa \widetilde{D}^{2,j} \mathbf{m}_{h}\} \cdot q_{h} \{\kappa \widetilde{D}^{2,j} \boldsymbol{\varphi}_{h}\} \mathrm{d}\boldsymbol{x}.$$

Let $h \to 0$, it follows from (39), (43) and Lemma 2.3 that

$$\lim_{h\to 0} T_h = -\frac{1}{3} \sum_{j=1}^6 \int_{\mathbb{R}^2} \left(\mathbf{m}^* \times \frac{\partial \mathbf{m}^*}{\partial \nu^j} \right) \cdot \frac{\partial \varphi}{\partial \nu^j} \mathrm{d}\boldsymbol{x} = -\int_{\mathbb{R}^2} \mathbf{m} \times \nabla \mathbf{m} \cdot \nabla \varphi \mathrm{d}\boldsymbol{x},$$

with $\nu^j, 1 \le j \le 6$, given in Lemma 2.3.

For the case k = 1, 3, applying the formula (29) to the right-hand side of (45), one gets

$$\begin{split} T_{h} &= -\int_{\mathbb{R}^{2}} \left(q_{h} \{ \mathbf{m}_{h} \times \widetilde{\mathfrak{D}}_{1}^{k} \mathbf{m}_{h} \} \cdot q_{h} \{ \widetilde{\mathfrak{D}}_{0}^{k} \boldsymbol{\varphi}_{h} \} \right) \\ &+ \left(q_{h} \{ \mathbf{m}_{h} \times \widetilde{\mathfrak{D}}_{1,B}^{k} \mathbf{m}_{h} \} \cdot q_{h} \{ \widetilde{\mathfrak{D}}_{0,B}^{k} \boldsymbol{\varphi}_{h} \} \right) + \left(q_{h} \{ \mathbf{m}_{h} \times \widetilde{\mathbb{D}}_{x}^{k} \mathbf{m}_{h} \} \cdot q_{h} \{ \widetilde{\mathbb{D}}_{x}^{k} \boldsymbol{\varphi}_{h} \} \right) \mathrm{d}\boldsymbol{x}. \end{split}$$

Thus using the strong convergence for $\widetilde{\mathfrak{D}}_{0}^{k} \widetilde{\mathfrak{D}}_{0,B}^{k}$ by (30), together with formula (28), one can write

$$\lim_{h \to 0} T_h = -\lim_{h \to 0} \int_{\mathbb{R}^2} \left(q_h \{ \mathbf{m}_h \times \widetilde{\mathbb{D}}_x^k \mathbf{m}_h \} \cdot \frac{\partial \varphi}{\partial x} \right) + \left(q_h \{ \mathbf{m}_h \times \widetilde{\mathbb{D}}_y^k \mathbf{m}_h \} \cdot \frac{\partial \varphi}{\partial y} \right) \mathrm{d}x.$$

Using convergence result (39), (44), and property (ii) of q_h , it finally leads to (46), for k = 1, 3.

3.5. Convergence of equation. For every $\varphi \in C_c^2$, define $\varphi_h(\boldsymbol{x}_h) = \varphi(\boldsymbol{x}_h)$. Applying interpolation operator q_h to the equation (34), and taking $q_h \varphi_h$ as a test function, one gets

(47)
$$\int_{\mathbb{R}^2} \frac{\partial}{\partial t} q_h \mathbf{m}_h \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} = -\int_{\mathbb{R}^2} q_h \{\mathbf{m}_h \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}}\} \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} - \alpha \int_{\mathbb{R}^2} q_h \{\mathbf{m}_h \times (\mathbf{m}_h \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}})\} \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x}$$

First, we consider the limit of the first term on right-hand side and prove that

(48)
$$\int_{\mathbb{R}^2} q_h(\mathbf{m}_h \times \mathbf{B}^*_{\text{eff},h}) \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} \to \int_{\mathbb{R}^2} \mathbf{m}^* \times \mathbf{B}_{\text{eff}} \cdot \boldsymbol{\varphi} \mathrm{d}\boldsymbol{x}.$$

Substitute the expression of $\mathbf{B}_{\text{eff},h}^* = \mathbf{B}_1^* + \mathbf{B}_2^* + \mathbf{B}_3^* + \mathbf{B}_4^*$ in (35) to the left-hand side of equation (48). Then one can easily pass to the limit in (48) by utilizing (39) and (46) with two exemptions: the second term in \mathbf{B}_2^* , and the term $\mathbf{B}_3^{(2)}$. For the second term in \mathbf{B}_2^* , let us denote it by

$$\widetilde{\mathbf{B}}_{2}^{*} = 2K \sum_{j=1}^{3} \int_{\mathbb{R}^{2}} q_{h} \{ (\mathbf{m}_{h} \cdot \widetilde{D}^{1,j} \mathbf{m}_{h}) \mathbf{m}_{h} \times \widetilde{D}^{1,j} \mathbf{m}_{h} \} \cdot q_{h} \boldsymbol{\varphi}_{h} \mathrm{d}\boldsymbol{x},$$

Since $|\mathbf{m}_h(\boldsymbol{x}_h)| = 1$ for all \boldsymbol{x}_h , it follows from the definition of $\widetilde{D}^{1,j}$ that

$$\mathbf{m}_h(\boldsymbol{x}_h) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h) = \mathbf{m}_h(\boldsymbol{x}_h^{1,j}) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h^{1,j}),$$

$$\mathbf{m}_h(\boldsymbol{x}_h) \times \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h) = -\mathbf{m}_h(\boldsymbol{x}_h^{1,j}) \times \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h^{1,j}).$$

Consequently,

(49)
$$(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h})) \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h})$$
$$= \frac{h}{2} \widetilde{D}^{1,j} \Big\{ \big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h}) \big) \mathbf{m}_{h}(\boldsymbol{x}_{h}) \times \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h}) \Big\}.$$

Substituting (49) into F_1 , and applying integration by parts in Lemma 2.2 and Hölder's inequality implies

$$|\widetilde{\mathbf{B}}_{2}^{*}| \leq hK_{1} \sum_{j=1}^{3} \|\widetilde{D}^{1,j}\boldsymbol{\varphi}_{h}\|_{\widetilde{L}_{h}^{\infty}} \|\widetilde{D}^{1,j}\mathbf{m}_{h}\|_{\widetilde{L}_{h}^{2}}^{2},$$

which tends to 0 as $h \to 0$. Finally, for the term $\mathbf{B}_3^{(2)}$, utilizing integration by parts yields

$$\begin{split} &\int_{\mathbb{R}^2} q_h(\mathbf{m}_h \times \mathbf{B}_3^{(2)}) \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} \\ = & \frac{h^2}{(h^*)^2} \sum_{k=1}^3 \sum_{j=1}^{N_k} \frac{d_k^2}{2} \int_{\mathbb{R}^2} q_h \big\{ \widetilde{D}^{k,j}(L_k \mathbf{m}_h) \cdot \mathbf{e}^z \big\} \cdot q_h \big\{ \widetilde{D}^{k,j}(\mathbf{m}_h \times \mathbf{e}^z \cdot \boldsymbol{\varphi}_h) \big\} \mathrm{d}\boldsymbol{x} \\ &+ \frac{h^2}{(h^*)^2} \sum_{k=1}^3 \sum_{j=1}^{N_k} \frac{d_k^2}{2} \int_{\mathbb{R}^2} q_h \big\{ \widetilde{D}^{k,j} \mathbf{m}_h \cdot \mathbf{e}^z \big\} \cdot q_h \big\{ \widetilde{D}^{k,j}(L_k \mathbf{m}_h \times \mathbf{e}^z \cdot \boldsymbol{\varphi}_h) \big\} \mathrm{d}\boldsymbol{x}. \end{split}$$

Therefore by Hölder's inequality one can derive

$$\begin{split} & \left| \int_{\mathbb{R}^2} q_h(\mathbf{m}_h \times \mathbf{B}_3^{(2)}) \cdot q_h \boldsymbol{\varphi}_h \mathrm{d}\boldsymbol{x} \right| \\ \leq & \frac{h^2}{(h^*)^2} \sum_{k=1}^3 \sum_{j=1}^{N_k} d_k^2 \Big(\|\widetilde{D}^{k,j} \mathbf{m}_h\|_{\widetilde{L}_h^2} \|\widetilde{D}^{k,j} \boldsymbol{\varphi}_h\|_{\widetilde{L}_h^2} + \|\boldsymbol{\varphi}_h\|_{\widetilde{L}_h^\infty} \|\widetilde{D}^{1,j} \mathbf{m}_h\|_{\widetilde{L}_h^2}^2 \Big), \end{split}$$

thus it tends to 0 as $h \to 0$. Hence the assertion (48) is proved.

Furthermore, inequality (36) demonstrates that the term $q_h \{\mathbf{m}_h \times \mathbf{B}_{\text{eff},h}^*\}$ is bounded in $L^2(0, T, L^2)$. Therefore there is a subsequence which converges weakly in $L^2(0, T, L^2)$. Considering (48), it can be inferred that

(50)
$$q_h \{ \mathbf{m}_h \times \mathbf{B}^*_{\text{eff},h} \} \to \mathbf{m}^* \times \mathbf{B}_{\text{eff}}$$
 weakly in $L^2(0,T,L^2)$.

Let us return to the equation (47). One can take the limit as $h \to 0$ for the first term in (47) by using (48). Moreover, the last term in (47) can pass to the limit by utilizing (39) and (50). Consequently, by taking the limit in (47), we can conclude that limiting function \mathbf{m}^* satisfies the continuity equation (9).

4. INTERPOLATION TO HEXAGONAL LATTICE

In this section, we provide the construction of interpolation operators q_h and p_h , which fulfill Propositions 3.3 and 3.4. Following the approach used in [14] for the square lattice case, we define the interpolation function $q_h u_h$ as a step function and $p_h u_h$ as a piece-wise linear function for any discrete function u_h . 4.1. Step interpolation function. To introduce the definition of q_h , we first need to establish a mapping from lattice points in \mathcal{G} to open sets in \mathbb{R}^2 . We use open set $\Omega_{\boldsymbol{x}_h} \subset \mathbb{R}^2$ to denote an adjoint area of $\boldsymbol{x}_h \in \mathcal{G}$, such that

if $\boldsymbol{x}_h \in \mathcal{G}_1$, then $\Omega_{\boldsymbol{x}_h}$ is the trapezoid on the upper right side of \boldsymbol{x}_h ,

if $\boldsymbol{x}_h \in \mathcal{G}_2$, then $\Omega_{\boldsymbol{x}_h}$ is the trapezoid on the lower left side of \boldsymbol{x}_h ,

as shown in Figure 3. It is easy to verify that the sets $\Omega_{\boldsymbol{x}_h}$ are mutually disjoint for all $\boldsymbol{x}_h \in \mathcal{G}$. Furthermore, the entire space of \mathbb{R}^2 is covered by $\bigcup_{\boldsymbol{x}_h \in \mathcal{G}} \overline{\Omega}_{\boldsymbol{x}_h}$.




FIGURE 3. Lattice division 1.

Definition 4.1. Given any function $u_h(\boldsymbol{x}_h)$ defined on \mathcal{G} , define a step function $q_h u_h(\boldsymbol{x})$ as

 $q_h u_h(\boldsymbol{x}) = u_h(\boldsymbol{x}_h), \text{ for any } \boldsymbol{x} \in \Omega_{\boldsymbol{x}_h}, \ \boldsymbol{x}_h \in \mathcal{G}.$

4.2. **Piece-wise linear interpolation function.** The step function $q_h u_h$ is simple to evaluate, but it lacks continuity. Therefore, we require a piecewise linear function $p_h u_h$ that maintains higher regularity from u_h . Before defining p_h , let us first discuss the construction strategy.

Let $\boldsymbol{x}_h \in \mathcal{G}$ be fixed. We denote the lattice point \boldsymbol{x}_h by the letter A, and use B, C, and D to represent other vertex points of $\Omega_{\boldsymbol{x}_h}$ as shown in Figure 4. For any function u_h defined on \mathcal{G} , the value of $p_h u_h(\boldsymbol{x}^*)$ for any point $\boldsymbol{x}^* \in \Omega_{\boldsymbol{x}_h}$ is given by:

- To begin with, we can determine the value of u_h at lattice points A, B, C, and D to evaluate $p_h u_h$. Additionally, we can use the linear Lagrangian interpolation formula to calculate $p_h u_h$ on lines AB and CD. Next, we need to draw a plumb line through x^* that intersects with AB and CD at points E and F respectively.
- Finally, we note that the values of $p_h u_h$ on E and F are given in the first step. Using linear Lagrangian interpolation formula again, we can obtain the value of $p_h u_h$ on line EF. Therefore, we can determine the value of $p_h u_h(\boldsymbol{x}^*)$.




FIGURE 4. Lattice division 2.

Following the above steps, we can introduce the definition of operator p_h .

Definition 4.2. For any function u_h defined on hexagonal lattice \mathcal{G} , define the piece-wise linear function $p_h u_h(\boldsymbol{x}^*)$ as

$$p_h u_h(\boldsymbol{x}^*) = u_h(\boldsymbol{x}_h) + \frac{y_1}{h} \left(u_h(\boldsymbol{x}_h^{1,3}) - u_h(\boldsymbol{x}_h) \right) + \frac{y_2}{y_1 + h} \left(u_h(\boldsymbol{x}_h^{1,1}) - u_h(\boldsymbol{x}_h) \right) \\ + \frac{y_1 y_2}{h(y_1 + h)} \left(u_h(\boldsymbol{x}_h) - u_h(\boldsymbol{x}_h^{1,3}) - u_h(\boldsymbol{x}_h^{1,1}) + u_h(\boldsymbol{x}_h^{2,6}) \right),$$

for any $\mathbf{x}^* \in \Omega_{\mathbf{x}_h}$, $\mathbf{x}_h \in \mathcal{G}$. As shown in Figure 4, the pair (y_1, y_2) is determined by \mathbf{x}^* ,

$$y_1 = \frac{2}{\sqrt{3}} |(\boldsymbol{x}^* - \boldsymbol{x}_h) \cdot \boldsymbol{e}_1|, \quad y_2 = \frac{1}{\sqrt{3}} |(\boldsymbol{x}^* - \boldsymbol{x}_h) \cdot \boldsymbol{e}_1| + |(\boldsymbol{x}^* - \boldsymbol{x}_h) \cdot \boldsymbol{e}_2|.$$

4.3. Proof of Propositions 3.3 and 3.4. Now let us prove the above interpolation operators q_h , p_h constructed on hexagonal lattice satisfy Propositions 3.3 and 3.4.

For any function $u_h, v_h \in \tilde{L}_h^2$, it follows from Definition 4.1 that

(51)
$$\int_{R^2} q_h^* u_h \cdot q_h^* v_h d\boldsymbol{x} = \sum_{\boldsymbol{x}_h \in \mathcal{G}} \int_{\Omega_{\boldsymbol{x}_h}} u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h) d\boldsymbol{x}$$
$$= \frac{3\sqrt{3}}{4} h^2 \sum_{\boldsymbol{x}_h \in \mathcal{G}} u_h(\boldsymbol{x}_h) \cdot v_h(\boldsymbol{x}_h) = (u_h, v_h)_h.$$

Hence the map q_h satisfies the isometry property (i) stated in Proposition 3.3, and it is a linear transformation from \tilde{L}_h^2 to L^2 . Additionally, properties (ii)-(iii) can be directly derived for q_h based on its definition.

As for the operator p_h , straightforward calculations show that

(52)
$$|p_h u_h(\boldsymbol{x})| \le |u_h(\boldsymbol{x}_h)| + |u_h(\boldsymbol{x}_h^{1,1})| + |u_h(\boldsymbol{x}_h^{1,3})| + |u_h(\boldsymbol{x}_h^{2,6})|.$$

Moreover, since $\nabla y_1 = (2/\sqrt{3}, 0)$ and $\nabla y_2 = (1/\sqrt{3}, 1)$, one can obtain

(53)
$$|\nabla(p_h^*u_h(\boldsymbol{x}))| \le |\widetilde{D}^{1,1}u_h(\boldsymbol{x}_h)| + |\widetilde{D}^{1,3}u_h(\boldsymbol{x}_h)| + |\widetilde{D}^{1,2}u_h(\boldsymbol{x}_h^{1,1})|.$$

Therefore by substitution one can deduce

$$\|p_h v_h\|_{L^2}^2 = \sum_{\boldsymbol{x}_h \in \mathcal{G}} \int_{\Omega_{\boldsymbol{x}_h}} |p_h v_h|^2 \mathrm{d}\boldsymbol{x} \le \sum_{\boldsymbol{x}_h \in \mathcal{G}} |\Omega_{\boldsymbol{x}_h}| \sup_{\boldsymbol{x} \in \Omega_{\boldsymbol{x}_h}} |p_h u_h(\boldsymbol{x})|^2 \le 16 \|u_h\|_{\tilde{L}^2_h}^2.$$

and

$$\|\nabla(p_h v_h)\|_{L^2}^2 \leq \sum_{\boldsymbol{x}_h \in \mathcal{G}} |\Omega_{\boldsymbol{x}_h}| \sup_{\boldsymbol{x} \in \Omega_{\boldsymbol{x}_h}} |\nabla(p_h u_h(\boldsymbol{x}))| \leq 6 \sum_{j=1}^3 \|\widetilde{D}^{1,j} v_h\|_{\widetilde{L}^2_h}^2.$$

Hence p_h satisfies the property (iv) in Proposition 3.4.

Regarding property (v), note that for any given $\boldsymbol{x} \in \Omega_{\boldsymbol{x}_h}$ where $\boldsymbol{x}_h \subset \mathcal{G}$, we can use the definitions of p_h and q_h to derive the following:

$$|p_h u_h(\boldsymbol{x}) - q_h u_h(\boldsymbol{x})| \le h \sum_{j=1}^3 |\widetilde{D}^{1,j} u_h|.$$

Substituting it into the right-hand side of equation

$$\int_{\mathbb{R}^2} (p_h u_h - q_h u_h)^2 \mathrm{d}\boldsymbol{x} = \sum_{\boldsymbol{x}_h \in \mathcal{G}} \int_{\Omega_{\boldsymbol{x}_h}} (p_h u_h - q_h u_h)^2 \mathrm{d}\boldsymbol{x},$$

then one finally derives for $u_h \in \widetilde{H}_h^1$ that

$$\int_{\mathbb{R}^2} (p_h u_h - q_h u_h)^2 \mathrm{d}\boldsymbol{x} \le h^2 \sum_{j=1}^3 \|\widetilde{D}^{1,j} u_h\|_{\widetilde{L}^2_h}^2.$$

The proof of Propositon 3.4 is completed.

Appendix A. Proof of Proposition 3.1

Firstly, it is easy to verify that $\mathbf{m}_h(\mathbf{x}_h, t)$ remains of unit length by multiplying equation (34) with \mathbf{m}_h and then deriving

(54)
$$\frac{\mathrm{d}}{\mathrm{d}t}|\mathbf{m}_h(\boldsymbol{x}_h,t)|^2 = 0.$$

To prove the proposition, it suffices to prove that

(55)
$$\frac{1}{(h^*)^2}\mathbf{m}_h(\boldsymbol{x}_h) \times \mathbf{B}_l(\boldsymbol{x}_h) = \mathbf{m}_h(\boldsymbol{x}_h) \times \mathbf{B}_l^*(\boldsymbol{x}_h), \quad l = 1, 2, 3, 4$$

with \mathbf{B}_l and \mathbf{B}_l^* given in (33) and (35), respectively. For the convenience of reading, we rewrite the equation (33) in the following:

(56)
$$\begin{cases} \mathbf{B}_{1}(\boldsymbol{x}_{h}) = (h^{*})^{2} \sum_{k=1}^{3} \sum_{j=1}^{N_{k}} \frac{J_{k}}{h^{2}} \mathbf{m}_{h}(\boldsymbol{x}_{h}^{k,j}) \\ \mathbf{B}_{2}(\boldsymbol{x}_{h}) = 2(h^{*})^{2} \sum_{j=1}^{3} \frac{K}{h^{2}} \Big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{m}_{h}(\boldsymbol{x}_{h}^{1,j}) \Big) \mathbf{m}_{h}(\boldsymbol{x}_{h}^{1,j}), \\ \mathbf{B}_{3}(\boldsymbol{x}_{h}) = \sum_{k=1}^{3} \sum_{j=1}^{N_{k}} \Big(L_{k}(\boldsymbol{x}_{h}) + L_{k}(\boldsymbol{x}_{h}^{k,j}) \Big) \Big(\mathbf{m}_{h}(\boldsymbol{x}_{h}^{k,j}) \cdot \mathbf{e}^{z} \Big) \mathbf{e}^{z} \\ + 2\lambda(\boldsymbol{x}_{h}) \Big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{x} \Big) \mathbf{e}^{x}, \\ \mathbf{B}_{4}(\boldsymbol{x}_{h}) = \mu \mathbf{B}(\boldsymbol{x}_{h}). \end{cases}$$

And the equation (35) is rewritten in (62). Now let us calculate the left-hand side of (55) by using (56). Note that

(57)
$$\mathbf{m}_h(\boldsymbol{x}_h^{k,j}) = d_k h \widetilde{D}^{k,j} \mathbf{m}_h(\boldsymbol{x}_h) + \mathbf{m}_h(\boldsymbol{x}_h).$$

Applying (57), together with unit-length property of \mathbf{m}_h , one has

$$\frac{1}{(h^*)^2}\mathbf{m}_h(\boldsymbol{x}_h) \times \mathbf{B}_1(\boldsymbol{x}_h) = \sum_{k=1}^3 \sum_{j=1}^{N_k} \frac{J_k d_k}{h} \mathbf{m}_h(\boldsymbol{x}_h) \times \widetilde{D}^{k,j} \mathbf{m}_h(\boldsymbol{x}_h),$$

and,

$$\frac{1}{(h^*)^2} \mathbf{m}_h(\boldsymbol{x}_h) \times \mathbf{B}_2(\boldsymbol{x}_h)$$

=2 $\sum_{j=1}^3 \frac{K d_1}{h} \Big(1 + d_1 h \mathbf{m}_h(\boldsymbol{x}_h) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h) \Big) \mathbf{m}_h(\boldsymbol{x}_h) \times \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h).$

Therefore combining them with (20) results in the equation (55) when l = 1, 2. The second term in \mathbf{B}_3 and the term \mathbf{B}_4 is trivial. Now we consider the first term in \mathbf{B}_3 and denote it by

$$\mathbf{B}_3^* = \sum_{k=1}^3 \sum_{j=1}^{N_k} \left(L_k(\boldsymbol{x}_h) + L_k(\boldsymbol{x}_h^{k,j}) \right) \left(\mathbf{m}_h(\boldsymbol{x}_h^{k,j}) \cdot \mathbf{e}^z \right) \mathbf{e}^z.$$

One can make the use of following transformation:

$$\sum_{k=1}^{3} \sum_{j=1}^{N_k} L_k(\boldsymbol{x}_h) \Big(\mathbf{m}_h(\boldsymbol{x}_h^{k,j}) \cdot \mathbf{e}^z \Big) \mathbf{e}^z$$
$$= \sum_{k=1}^{3} N_k L_k(\boldsymbol{x}_h) \Big(\mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \Big) \mathbf{e}^z + \sum_{k=1}^{3} \sum_{j=1}^{N_k} d_k h L_k(\boldsymbol{x}_h) \Big(\widetilde{D}^{k,j} \mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \Big) \mathbf{e}^z,$$

and

$$\sum_{k=1}^{3} \sum_{j=1}^{N_k} L_k(\boldsymbol{x}_h^{k,j}) \Big(\mathbf{m}_h(\boldsymbol{x}_h^{k,j}) \cdot \mathbf{e}^z \Big) \mathbf{e}^z$$

$$= \sum_{k=1}^{3} N_k L_k(\boldsymbol{x}_h) \Big(\mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \Big) \mathbf{e}^z + \sum_{k=1}^{3} \sum_{j=1}^{N_k} d_k h \Big(\widetilde{D}^{k,j} (L_k \mathbf{m}_h) (\boldsymbol{x}_h) \cdot \mathbf{e}^z \Big) \mathbf{e}^z.$$

Using the transformation above, together with (20), it follows that

$$\mathbf{B}_{3}^{*} = 2\sum_{k=1}^{3} N_{k}L_{k}(\boldsymbol{x}_{h}) \Big(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z}\Big) \mathbf{e}^{z} + \sum_{k=1}^{3} l_{k}h^{2}L_{k}(\boldsymbol{x}_{h}) \Big(\widetilde{\Delta}^{k}\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z}\Big) \mathbf{e}^{z} + \sum_{k=1}^{3} l_{k}h^{2} \Big(\widetilde{\Delta}^{k}(L_{k}\mathbf{m}_{h})(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z}\Big) \mathbf{e}^{z}.$$

Substitution of the expression above leads to equation (55) for the case when l = 3. Hence the proposition is proved.

APPENDIX B. DISCRETE ENERGY DISSIPATION

Before we prove Proposition 3.2, an energy dissipation property for the discrete equations (34)-(35) is needed. Instead of using the energy $\mathcal{H}_h[\mathbf{m}_h]$ as given in (4), we introduce another form of energy, denoted by $\mathcal{H}_h^*[\mathbf{m}_h]$, which is obtained through a difference quotient (58)

$$\begin{aligned} &\mathcal{H}_{h}^{(30)} \mathbf{H}_{h}^{*}[\mathbf{m}_{h}] \\ &= \sum_{k=1}^{3} J_{k} \frac{d_{k}^{2}}{4g_{k}^{2}} \|\widetilde{\nabla}_{k}\mathbf{m}_{h}\|_{\widetilde{L}_{h}^{2}}^{2} + K \frac{d_{1}^{2}}{2g_{1}^{2}} \|\widetilde{\nabla}_{1}\mathbf{m}_{h}\|_{\widetilde{L}_{h}^{2}}^{2} - K \frac{d_{1}^{2}}{2g_{1}^{2}} \|\mathbf{m}_{h} \cdot \widetilde{\nabla}_{1}\mathbf{m}_{h}\|_{\widetilde{L}_{h}^{2}}^{2} \\ &+ \frac{h^{2}}{(h^{*})^{2}} \sum_{k=1}^{3} \frac{d_{k}^{2}}{2g_{k}^{2}} \|L_{k}^{1/2} \widetilde{\nabla}^{k}\mathbf{m}_{h} \cdot \mathbf{e}^{z}\|_{\widetilde{L}_{h}^{2}}^{2} \\ &+ \frac{h^{2}}{(h^{*})^{2}} \sum_{k=1}^{3} \sum_{j=1}^{N_{k}} \frac{d_{k}^{2}}{2} \Big((\widetilde{D}^{k,j}L_{k}) \big(\mathbf{m}_{h} \cdot \mathbf{e}^{z} \big) \mathbf{e}^{z}, \ \widetilde{D}^{k,j}\mathbf{m}_{h} \Big)_{h} \\ &- \frac{1}{(h^{*})^{2}} \sum_{k=1}^{3} N_{k} \|L_{k}^{1/2}\mathbf{m}_{h} \cdot \mathbf{e}^{z}\|_{\widetilde{L}_{h}^{2}}^{2} - \frac{1}{(h^{*})^{2}} \|\lambda^{1/2}\mathbf{m}_{h} \cdot \mathbf{e}^{x}\|_{\widetilde{L}_{h}^{2}}^{2} - \frac{\mu}{(h^{*})^{2}} \big(\mathbf{m}_{h}, \mathbf{B} \big)_{h} \end{aligned}$$

Then we have:

Proposition B.1. Let m_h be the solution to the equations (34)-(35). The following energy dissipation holds:

(59)
$$\mathcal{H}_{h}^{*}[\boldsymbol{m}_{h}(t)] + \alpha \int_{0}^{t} \|\boldsymbol{m}_{h}(s) \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}^{*}(s)\|_{\tilde{L}_{h}^{2}}^{2} \mathrm{d}s \leq \mathcal{H}_{h}^{*}[\boldsymbol{m}_{h}(0)],$$

holds for every $t \in [0, +\infty)$, where the energy \mathcal{H}_h^* is given by (58).

24

Proof. Since

$$\left(\mathbf{m}_h \times (\mathbf{m}_h \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}^*)\right) \cdot \mathbf{B}_{\mathrm{eff},\mathrm{h}}^* = -\left|\mathbf{m}_h \times \mathbf{B}_{\mathrm{eff},\mathrm{h}}^*\right|^2,$$

taking the \widetilde{L}_h^2 inner-product of (34) and ${\bf B}_{\rm eff,h}^*,$ one has

(60)
$$-\left(\frac{\mathrm{d}\mathbf{m}_h}{\mathrm{d}t}, \ \mathbf{B}^*_{\mathrm{eff},\mathrm{h}}\right)_h + \alpha \|\mathbf{m}_h(t) \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}}(t)\|^2_{\tilde{L}^2_h} = 0$$

We assert that

(61)
$$-\left(\frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t}, \ \mathbf{B}_{\mathrm{eff},\mathrm{h}}^{*}\right)_{h} = \frac{\mathrm{d}}{\mathrm{d}t}\mathcal{H}_{h}^{*}[\mathbf{m}_{h}].$$

Then the inequality (59) follows by substituting (61) into (60) and integration over [0, t].

Now let us compute left-hand side of (61) term by term by applying (35). For the convenience of reading, we rewrite the equation (35) here: (62)

$$\begin{cases} \mathbf{B}_{1}^{*}(\boldsymbol{x}_{h}) = \sum_{k=1}^{3} l_{k} J_{k} \widetilde{\Delta}_{k} \mathbf{m}_{h}(\boldsymbol{x}_{h}) \\ \mathbf{B}_{2}^{*}(\boldsymbol{x}_{h}) = 2 l_{1} K \widetilde{\Delta}_{1} \mathbf{m}_{h}(\boldsymbol{x}_{h}) + 2 K \sum_{j=1}^{3} \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h}) \right) \widetilde{D}^{1,j} \mathbf{m}_{h}(\boldsymbol{x}_{h}), \\ \mathbf{B}_{3}^{(1)}(\boldsymbol{x}_{h}) = \frac{2}{(h^{*})^{2}} \lambda(\boldsymbol{x}_{h}) \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{x} \right) \mathbf{e}^{x} + \frac{2}{(h^{*})^{2}} \sum_{k=1}^{3} N_{k} L_{k}(\boldsymbol{x}_{h}) \left(\mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z} \right) \mathbf{e}^{z}, \\ \mathbf{B}_{3}^{(2)}(\boldsymbol{x}_{h}) = \frac{h^{2}}{(h^{*})^{2}} \sum_{k=1}^{3} l_{k} \left\{ L_{k}(\boldsymbol{x}_{h}) \left(\widetilde{\Delta}_{k} \mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z} \right) \mathbf{e}^{z} + \left(\widetilde{\Delta}_{k}(L_{k} \mathbf{m}_{h})(\boldsymbol{x}_{h}) \cdot \mathbf{e}^{z} \right) \mathbf{e}^{z} \right\} \\ \mathbf{B}_{4}^{*}(\boldsymbol{x}_{h}) = \frac{\mu}{(h^{*})^{2}} \mathbf{B}(\boldsymbol{x}_{h}). \end{cases}$$

For the term \mathbf{B}_1^* in (62), we apply (18) and deduce that

$$-\left(\frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t}, \ \mathbf{B}_{1}^{*}\right)_{h} = -\sum_{k=1}^{3} J_{k} \frac{d_{k}^{2}}{2} \left(\frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t}, \ \widetilde{D}^{k,j} \{\widetilde{D}_{B}^{k,j}\mathbf{m}_{h}\}\right)_{h}$$
$$= \sum_{k=1}^{3} J_{k} \frac{d_{k}^{2}}{2} \left(\widetilde{D}^{k,j} \frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t}, \ \widetilde{D}^{k,j}\mathbf{m}_{h}\right)_{h} = \frac{1}{2} \frac{\mathrm{d}}{\mathrm{d}t} \sum_{k=1}^{3} J_{k} \frac{d_{k}^{2}}{2g_{k}^{2}} \|\widetilde{\nabla}_{k}\mathbf{m}_{h}\|_{\widetilde{L}_{h}^{2}}^{2},$$

where we used (15) and Lemma 2.1. The first term in \mathbf{B}_2^* can be checked in a similar way. For the second term in \mathbf{B}_2^* , we assert that

(63)
$$-\sum_{j=1}^{3} \left(\frac{\mathrm{d}\mathbf{m}_{h}}{\mathrm{d}t}, \ \left(\mathbf{m}_{h} \cdot \widetilde{D}^{1,j}\mathbf{m}_{h}\right) \widetilde{D}^{1,j}\mathbf{m}_{h} \right)_{h}$$
$$= -\frac{d_{1}^{2}}{4g_{1}^{2}} \frac{\mathrm{d}}{\mathrm{d}t} \left\| \mathbf{m}_{h}(\boldsymbol{x}_{h}) \cdot \widetilde{\nabla}_{1}\mathbf{m}_{h}(\boldsymbol{x}_{h}) \right\|_{\widetilde{L}^{2}_{h}}^{2}.$$

25

To clarify, we can calculate the inner product on the left-hand side of equation (63) for the interior and gather all terms that involve points \boldsymbol{x}_h and its neighbor $\boldsymbol{x}_h^{1,j}$. This yields two resulting terms:

(64)
$$-\left(\mathbf{m}_h(\boldsymbol{x}_h)\cdot\widetilde{D}^{1,j}\mathbf{m}_h(\boldsymbol{x}_h)\right)\cdot\left(\frac{\mathrm{d}\mathbf{m}_h}{\mathrm{d}t}(\boldsymbol{x}_h)\cdot\widetilde{D}^{1,j}\mathbf{m}_h(\boldsymbol{x}_h)\right),$$

and

(65)
$$-\left(\mathbf{m}_h(\boldsymbol{x}_h^{1,j}) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h^{1,j})\right) \cdot \left(\frac{\mathrm{d}\mathbf{m}_h}{\mathrm{d}t}(\boldsymbol{x}_h^{1,j}) \cdot \widetilde{D}^{1,j} \mathbf{m}_h(\boldsymbol{x}_h^{1,j})\right).$$

By using exchange of notation in (13) and unit-length property, adding (64) and (65) together equals

$$-\frac{1}{2}\frac{\mathrm{d}}{\mathrm{d}t}\big|\mathbf{m}_h(\boldsymbol{x}_h)\cdot\widetilde{D}^{1,j}\mathbf{m}_h(\boldsymbol{x}_h)\big|^2.$$

Then equation (63) follows by the notation in (15).

Finally, one can check that the computation of terms $\mathbf{B}_3^{(1)}$ and \mathbf{B}_4^* is trivial. As for the term $\mathbf{B}_3^{(2)}$, we can deduce from (15) and Lemma 2.1:

$$l_k \Big(L_k(\boldsymbol{x}_h) \big(\widetilde{\Delta}_k \mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \big) \mathbf{e}^z, \, \frac{\mathrm{d}}{\mathrm{d}t} \mathbf{m}_h(\boldsymbol{x}_h) \Big)_h \\ = -\sum_{j=1}^{N_k} \frac{d_k^2}{2} \Big(\big(\widetilde{D}^{k,j} \mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \big) \mathbf{e}^z, \, \widetilde{D}^{k,j} \big\{ L_k(\boldsymbol{x}_h) \frac{\mathrm{d}}{\mathrm{d}t} \mathbf{m}_h(\boldsymbol{x}_h) \big\} \Big)_h,$$

and,

$$l_k \Big(\big(\widetilde{\Delta}_k (L_k \mathbf{m}_h) (\boldsymbol{x}_h) \cdot \mathbf{e}^z \big) \mathbf{e}^z, \, \frac{\mathrm{d}}{\mathrm{d}t} \mathbf{m}_h (\boldsymbol{x}_h) \Big)_h \\ = -\sum_{j=1}^{N_k} \frac{d_k^2}{2} \Big(\big(\widetilde{D}^{k,j} (L_k \mathbf{m}_h) (\boldsymbol{x}_h) \cdot \mathbf{e}^z \big) \mathbf{e}^z, \, \widetilde{D}^{k,j} \frac{\mathrm{d}}{\mathrm{d}t} \mathbf{m}_h (\boldsymbol{x}_h) \Big)_h.$$

Therefore summing up one can derive

$$-\left(\frac{\mathrm{d}\mathbf{m}_h}{\mathrm{d}t}, \ \mathbf{B}_3^{(2)}\right)_h = \frac{h^2}{(h^*)^2} \frac{d_k^2}{2} \frac{\mathrm{d}}{\mathrm{d}t} \sum_{j=1}^{N_k} \left(\left(\widetilde{D}^{k,j} (L_k \mathbf{m}_h) (\boldsymbol{x}_h) \cdot \mathbf{e}^z \right) \mathbf{e}^z, \ \widetilde{D}^{k,j} \mathbf{m}_h(\boldsymbol{x}_h) \right)_h,$$

where the right-hand side can be written as

$$\frac{h^2}{(h^*)^2} \frac{d_k^2}{2} \left\{ \frac{\mathrm{d}}{\mathrm{d}t} \frac{1}{g_k^2} \| L_k^{1/2} \widetilde{\nabla}^k \mathbf{m}_h(\boldsymbol{x}_h) \cdot \mathbf{e}^z \|_{\widetilde{L}_h^2}^2 + \frac{\mathrm{d}}{\mathrm{d}t} \sum_{j=1}^{N_k} \left((\widetilde{D}^{k,j} L_k) \left(\mathbf{m}_h \cdot \mathbf{e}^z \right) \mathbf{e}^z, \ \widetilde{D}^{k,j} \mathbf{m}_h \right)_h \right\}$$

This completes the proof.

APPENDIX C. PROOF OF PROPOSITION 3.2

The equation (54) implies that $|\mathbf{m}_h| = 1$, and the energy inequality (59) deduce

(66)
$$\mathcal{H}_{h}^{*}[\mathbf{m}_{h}(t)] \leq \mathcal{H}_{h}^{*}[\mathbf{m}_{h}(0)]$$

Notice that using the assumption 1 and $|\mathbf{m}_h| = 1$, we have the estimate: (67)

$$\frac{1}{(h^*)^2} \sum_{k=1}^3 N_k \|L_k^{1/2} \mathbf{m}_h \cdot \mathbf{e}^z\|_{\widetilde{L}_h^2}^2 + \frac{1}{(h^*)^2} \|\lambda^{1/2} \mathbf{m}_h \cdot \mathbf{e}^x\|_{\widetilde{L}_h^2}^2 + \frac{\mu}{(h^*)^2} \big(\mathbf{m}_h, \mathbf{B}\big)_h \le C,$$

where constant C depends only on $\|\mathbf{B}\|_{L^1}$, $\|L\|_{L^1}$ and $\|\lambda\|_{L^1}$. Thus applying (58)-(59), it leads to

$$\begin{split} \sum_{k=1}^{3} J_k \frac{d_k^2}{4g_k^2} \| \widetilde{\nabla}_k \mathbf{m}_h \|_{\widetilde{L}_h^2}^2 + K \frac{d_1^2}{2g_1^2} \Big(\| \widetilde{\nabla}_1 \mathbf{m}_h \|_{\widetilde{L}_h^2}^2 - \| \mathbf{m}_h \cdot \widetilde{\nabla}_1 \mathbf{m}_h \|_{\widetilde{L}_h^2}^2 \Big) \\ + \frac{h^2}{(h^*)^2} \sum_{k=1}^{3} \frac{d_k^2}{2g_k^2} \| L_k^{1/2} \, \widetilde{\nabla}^k \mathbf{m}_h \cdot \mathbf{e}^z \|_{\widetilde{L}_h^2}^2 \le \mathcal{H}_h^* [\mathbf{m}_h(0)] + C. \end{split}$$

Here, the second term on left-hand side is non-negative. Hence,

$$\sum_{k=1}^{3} J_k \frac{d_k^2}{4g_k^2} \|\widetilde{\nabla}_k \mathbf{m}_h\|_{\widetilde{L}^2_h}^2 \leq \mathcal{H}_h^*[\mathbf{m}_h(0)] + C,$$

which leads to the second inequality in (36).

On the other hand, the equation (34) implies that

$$\int_0^T \left\| \frac{\partial \mathbf{m}_h}{\partial t} \right\|_{\tilde{L}^2_h}^2 \mathrm{d}t = (1 + \alpha^2) \int_0^T \|\mathbf{m}_h \times \mathbf{B}^*_{\mathrm{eff},\mathrm{h}} \|_{\tilde{L}^2_h}^2 \mathrm{d}t.$$

Hence, the inequality (3.2) is also proved, by virtue of Proposition B.1. The proof of Proposition 3.2 is completed.

Acknowledgments

This work was supported by NSFC grant 11971021 (J. Chen). The work of Zhiwei Sun is supported by China Scholarship Council grant 202006920083.

References

- François Alouges and Alain Soyeur, On global weak solutions for Landau-Lifshitz equations: existence and nonuniqueness, Nonlinear Analysis: Theory, Methods & Applications 18 (1992), no. 11, 1071–1084.
- 2. Mathias Augustin, Sarah Jenkins, Richard FL Evans, Kostya S Novoselov, and Elton JG Santos, *Properties and dynamics of meron topological spin textures in the two-dimensional magnet* CrCl₃, Nature Communications **12** (2021), no. 1, 185.
- 3. Xavier Blanc, Claude Le Bris, and P-L Lions, From molecular models to continuum mechanics, Archive for Rational Mechanics and Analysis 164 (2002), 341–381.
- 4. Haim Brezis, Functional analysis, Sobolev spaces and partial differential equations, Springer, 2011.

- Suzan C Buranay and Nouman Arshad, Hexagonal grid approximation of the solution of the heat equation on special polygons, Advances in Difference Equations 2020 (2020), no. 1, 1–24.
- Antonio DeSimone, Robert V Kohn, Stefan Müller, and Felix Otto, A reduced theory for thin-film micromagnetics, Communications on Pure and Applied Mathematics 55 (2002), no. 11, 1408–1460.
- Giovanni Di Fratta, Michael Innerberger, and Dirk Praetorius, Weak-strong uniqueness for the Landau-Lifshitz-Gilbert equation in micromagnetics, Nonlinear Analysis: Real World Applications 55 (2020), 103–122.
- Adiguzel A Dosiyev and Emine Celiker, Approximation on the hexagonal grid of the Dirichlet problem for Laplace's equation, Boundary Value Problems 2014 (2014), no. 1, 1–19.
- Weinan E and Jianfeng Lu, The continuum limit and QM-continuum approximation of quantum mechanical models of solids, Communications in Mathematical Sciences 5 (2007), no. 3, 679–696.
- _____, The electronic structure of smoothly deformed crystals: Wannier functions and the Cauchy-Born rule, Archive for Rational Mechanics and Analysis 199 (2011), no. 2, 407–433.
- 11. _____, The Kohn-Sham equation for deformed crystals, vol. 221, Memoirs of the American Mathematical Society, 2013.
- Weinan E and Pingbing Ming, Cauchy-Born rule and the stability of crystalline solids: static problems, Archive for Rational Mechanics and Analysis 183 (2007), no. 2, 241– 297.
- Alexey Kartsev, Mathias Augustin, Richard FL Evans, Kostya S Novoselov, and Elton JG Santos, *Biquadratic exchange interactions in two-dimensional magnets*, npj Computational Materials 6 (2020), no. 1, 150.
- 14. Olga A Ladyzhenskaya, *The boundary value problems of mathematical physics*, Springer Science & Business Media, 2013.
- Pierre-Louis Sulem, Catherine Sulem, and Claude Bardos, On the continuous limit for a system of classical spins, Communications in Mathematical Physics 107 (1986), no. 3, 431–454.
- 16. Yulin Zhou, Boling Guo, and Shaobin Tan, Existence and uniqueness of smooth solution for system of ferromagnetic chain, Science in China (series A) **34** (1991), 257–266.