Resolution of Non-Singularities and the Absolute Anabelian Conjecture

Emmanuel Lepage

June 13, 2023

Introduction

Grothendieck anabelian conjecture is concerned with the recovery, for smooth curves over number fields, of a curve from its étale fundamental group. More precisely, it conjectures that if X and Y are two hyperbolic curves over a number field K with absolute Galois group G_K , then $\operatorname{Isom}_K(X,Y) \to \operatorname{Isom}_{G_K}(\pi_1^{\operatorname{alg}}(X), \pi_1^{\operatorname{alg}}(Y))/\sim$ is bijective, where \sim is given by composing with inner automorphisms. This conjecture was proved by S. Mochizuki in [4], where it is proved much more generally for sub-*p*adic fields.

In this article we will only be interested in the case where K is a p-adic field. In [7], Mochizuki then asks if one can remove the assumption of lying over an isomorphism of absolute Galois groups. This gives us the following conjecture:

Conjecture 0.1. Let K, L be finite extensions of \mathbb{Q}_p , and X/L, Y/K be two hyperbolic curves. Then the map $\mathrm{Isom}(X, Y) \to \mathrm{Isom}(\pi_1^{\mathrm{alg}}(X), \pi_1^{\mathrm{alg}}(Y))/\sim$ is bijective.

The conjecture was proved by Mochizuki in [7] in the case where X and Y are of quasi-Belyi type. This quasi-Belyi type condition doesn't contain any proper curve. The goal of this paper is to extend this result to a wider family of curves, that in particular contains every Mumford curve :

Theorem 0.2. Let K, L be finite extensions of \mathbb{Q}_p , and X/L, Y/K be two hyperbolic curves satisfying resolution of non singularities (see definition 1.1), for example two hyperbolic Mumford curves. Then the map $\mathrm{Isom}(X,Y) \to \mathrm{Isom}(\pi_1^{\mathrm{alg}}(X), \pi_1^{\mathrm{alg}}(Y))/\sim$ is bijective.

A theorem of Mochizuki ([7, Thm 2.9]) claims that to prove that it is sufficient to characterize group-theoretically among the partial Galois sections $G_{K'} \to \pi_1^{\text{alg}}(X)$ those which come from a closed point $x \in X(K')$ for any finite extension K' of K. The idea of the proof then relies on the reconstruction of the Berkovich space |X| (as a topological space) of the curve from the étale fundamental group, as was done in [3, Thm. 3.10] for the geometric tempered fundamental group. From a result of Mochizuki ([5, Thm 2.7]), one can recover from $\pi_1^{\text{alg}}(X)$ the dual graph $\mathbb{G}(X)$ of the special fiber of the stable model of $X_{\overline{K}}$. By applying this reconstruction to all the finite étale coverings, the recovering of the Berkovich space follows from a homeomorphism $|\overline{X}| = (\lim_{Z} |\mathbb{G}(Z)|) / \pi_1^{\text{alg}}(X)$, where Z goes through connected pointed finite étale covers of X and \overline{X} is the smooth compactification of X. It is then sufficient to characterized rigid points inside the Berkovich space |X|.

The Berkovich points on curves are traditionally classified into four classes, where the type 1 points correspond to \mathbb{C}_p -points. The type 2 points are type 3 points are easy to distinguish from the others by the local topological behavior. The main issue is then to distinguish between type 4 and type 1 points. In this paper, we present two different ways of doing it. First through a metric characterization, using Mochizuki's recovery of the log-structure of the special fiber of the stable model [5, Thm 2.7]). Secondly, by showing that the geometric decomposition group of type 4 points is non trivial. More precisely, we will see that a \mathbb{Z}_p -torsor of $X_{\mathbb{C}_p}$ does not decompose at a type 4 points if it is not a topological torsor. Locally in a small disk, this \mathbb{Z}_p -torsor comes from Kummer classes of invertible functions, and by applying the log-differential, whose image is bounded, one gets that these invertible functions cannot have p^n th root for big enough n. Then the rigid points are characterized among type 1 points by the fact that their image in the absolute Galois group G_K is open.

1 Resolution of Non-Singularities

Let K be a complete nonarchimedean field of mixed characteristics (0, p). Let O_K be its ring of integers and \widetilde{K} be its residue field.

Let $X = \overline{X} \setminus D$ be a hyperbolic curve over K. Let X^{an} be the associated Berkovich space. As a set, X^{an} can be seen as the union of the set of closed points of X and the set of \mathbb{R} -valued valuations v on the fraction field K(X) extending the valuation on K. The completion of K(X) for this valuation will be denoted by $\mathcal{H}(v)$. Since K(X)/K is an extension of transcendance degree one, for every such valuation v, $r(v) + t(v) \leq 1$ where $r(v) = \operatorname{rk}(v(K(X)^{\times})/v(K^{\times}))$ and t(v) is the transcendance degree of the residue field $\widehat{\mathcal{H}(v)}$ of $\mathcal{H}(v)$ as an extension of \widetilde{K} .

A type 2 point of X^{an} then corresponds to a valuation v such that t(v) = 1, and a type 3 point corresponds to a valuation v such that r(v) = 1. If a point x of X^{an} is a rigid point or comes from a valuation such that K(X) embeds isometrically into the completion of an algebraic closure of K, then x is said to be of type 1. Otherwise it is of type 4.

If \overline{X} has a stable model \mathfrak{X} over O_K , there is a natural specialization map sp : $X^{\mathrm{an}} \to \mathfrak{X}_s$ where $\mathfrak{X}_s = \mathfrak{X}_{\widetilde{K}}$ is the special fiber of \mathfrak{X} . Then the generic point z of any irreducible component of \mathfrak{X}_s has a unique preimage v_z and this preimage is a type 2 point, and the residue field $\mathcal{H}(v_z)$ is isomorphic to k(z). Such a type 2 point will be called a skeletal point of X^{an} , and the set of skeletal points will be denoted by V(X). Skeletal points are the vertices of the minimal triangulation of X^{an} in the sense of [1, §5.1.13]. The connected components of $X^{\mathrm{an}} \setminus V(X)$ are given by the preimages by sp of the closed points in \mathfrak{X}_s . Moreover, if K is algebraically closed and x is a closed point of \mathfrak{X}_s , then sp⁻¹(x) is analytically isomorphic to :

• a punctured open disk if x is a cusp;

- an open disk if x is a smooth point;
- an open annulus if x is a node.

In particular if v is a type two point of X^{an} such that $\mathcal{H}(v)$ is the fraction field of a curve of genus > 0, then v is skeletal.

Definition 1.1. ([3, Def. 2.1])

Let C be an algebraically closed complete nonarchimedean field of mixed characteristics (0, p). Let X be a hyperbolic curve over C. X is said to satisfy *resolution of non-singularities* (or RNS(X) for short) if for every point x of type 2 of X^{an}, there exists a finite étale cover $f : Y \to X$ such that $f^{-1}(x) \cap V(Y) \neq \emptyset$.

More generally, if X is a hyperbolic curve over a complete field K of mixed characteristics (0, p), then we will write RNS(X) for $\text{RNS}(X_{\text{C}})$, where C is the completion of an algebraic closure of K.

Proposition 1.1. Let $f : X \to Y$ be a map of hyperbolic curves over K.

- 1. If $f: X \to Y$ is a dominant map, then RNS(Y) implies RNS(X).
- 2. If $f: X \to Y$ is finite étale, then RNS(X) and RNS(Y) are equivalent.

Proof. One can assume K to be algebraically closed.

- 1. Let $x \in X^{an}$ be a point of type 2, and let y = f(x). Then y is also of type 2, and by RNS(Y), there exists a finite étale cover $g : Z \to Y$ and $z \in g^{-1}(y) \cap V(Z)$. Let $f_Z : X \times_Y Z \to Z$ be the map obtained by pulling back f and $g_X : X \times_Y Z \to X$ be the finite étale cover obtained by pulling back g. Then there exists $z' \in g_X^{-1}(x) \cap f_Z^{-1}(z) \subset g_X^{-1}(x) \cap f_Z^{-1}(x) \cap V(X \times_Y Z)$.
- 2. By (i), RNS(Y) implies RNS(X). Let us assume RNS(X) and let $y \in Y^{an}$ be a point of type 2, and let $x \in X^{an}$ be a preimage of y by f. Then there exists a finite étale cover $g : Z \to X$ and $z \in g^{-1}(x) \cap V(Z)$. Then $fg : Z \to Y$ is a finite étale cover and $z \in (fg)^{-1}(y) \cap V(Z)$, so that RNS(Y) is true.

Proposition 1.2. ([3, Th. 0.1])

1. If \overline{X} is a Mumford curve over $\overline{\mathbb{Q}}_p$ and $X \subset \overline{X}_{\mathbb{C}_p}$ is a Zariski open subset of $X_{\mathbb{C}_p}$ which is a hyperbolic curve over \mathbb{C}_p , then $\text{RNS}(X_{\mathbb{C}_p})$ is true.

2. If X is a \mathbb{C}_p -curve of quasi-Belyi type, then RNS(X) is true.

The case of Mumford curves over $\overline{\mathbb{Q}}_p$ follows from a more general criterion in terms of the following Hodge-Tate map. Let X be a smooth adic space over K, and let $\nu : X_{\text{pro-\acute{e}t}} \to X_{\acute{e}t}$ be the natural map from its pro-étale site to its étale site. Then the short exact sequence on $X_{\text{pro-\acute{e}t}}$:

$$1 \to \mathbb{Z}_p(1) \to \lim_{\times p} \mathcal{O}_X^{\times} \to \mathcal{O}_X^{\times} \to 1$$

induces a boundary map $\kappa : \mathcal{O}_{X_{\acute{e}t}}^{\times} = \nu_* \mathcal{O}_X^{\times} \to R^1 \nu_* \mathbb{Z}_p(1)$. There exists a unique $\mathcal{O}_{X_{\acute{e}t}}$ -linear map $\phi : \Omega^1_{X_{\acute{e}t}} \to R^1 \nu_* \widehat{\mathcal{O}}_{X_{\acute{e}t}}^{\times}$ such that the following diagram commutes (see [8, Lem. 3.24]):

$$\begin{array}{c} \mathcal{O}_{X_{\mathrm{\acute{e}t}}}^{\times} \xrightarrow{\kappa} R^{1} \nu_{*} \mathbb{Z}_{p}(1) \\ \downarrow^{dlog} & \downarrow^{i} \\ \Omega_{X_{\mathrm{\acute{e}t}}}^{1} \xrightarrow{\phi} R^{1} \nu_{*} \widehat{\mathcal{O}}_{X_{\mathrm{\acute{e}t}}}^{\times} \end{array}$$

and ϕ is an isomorphism. By taking global sections of $\phi^{-1}i$, one gets a map:

$$\operatorname{HT}_X : H^1(X, \mathbb{Z}_p(1)) \to H^0(X, \Omega^1_X).$$

- **Definition 1.2.** 1. Let $HTvLoc_1(X) \subset X(K)$ (for Hodge-Tate vanishing locus of X) be the set of points such that there exists $c \in H^1(Y, \mathbb{Z}_p(1))$ such that $HT_Y(c)$ has vanishing order $e < +\infty$ at y such that e + 1 is not a power of p.
 - 2. Let $\operatorname{HTvLoc}_{1}^{\text{f-\acute{e}t}}(X) = \bigcup_{f:Y \to X} f(\operatorname{HTvLoc}_{1}(Y)) \subset X(K)$, where the union is indexed through all finite étale covers of X.
 - 3. Let $HTv_1(X)$ be the following property: $HTvLoc_1^{f-\acute{e}t}(X)$ is dense in X(K).

Proposition 1.3. Let X be a hyperbolic curve over K. Then $HTv_1(X)$ implies RNS(X).

Proof. By the argument of [3, Prop. 2.2] it is enough to show that for every disk D in X, there exists a finite étale cover $g: Z \to X$ such that $g^{-1}(D) \cap V(Z) \neq \emptyset$.

By definition of $\operatorname{HTv}_1(X)$, there exists a finite étale cover $f: Y \to X, y \in f^{-1}(D)$ and $c \in H^1(Y, \mathbb{Z}_p(1))$ such that the vanishing order e of $HT_Y(c)$ at y is not of the form $p^n - 1$ for any $n \in \mathbb{N}$.

For $n \in \mathbb{N}$, let $c_n \in H^1(Y, \mu_{p^n})$ be the image of c by the map induced from the surjection $\mathbb{Z}_p(1) \to \mu_{p^n}$ and let $g_n : Y_n \to Y$ be the corresponding μ_{p^n} -torsor. Let D' be a discal neighborhood of y in $f^{-1}(D)$. Since $H^1(D', \mathbf{G}_m) = 0$, the restriction $c_{n,D'}$ of c_n to D' is in the image of the Kummer map $\kappa_n : \mathcal{O}^{\times}(D') \to H^1(D', \mu_{p^n})$. Let $a_n \in \mathcal{O}^{\times}(D')$ such that $\kappa_n(a_n) = c_{n,D'}$ and $a_n(y) = 1$. There exists $b_n \in \mathcal{O}^{\times}(D')$ such that $a_n = b_n^{p^n} a_{n+1}$ and $b_n(y) = 1$. After replacing D' by a smaller disk D'', there exists r < 1 such that for every $n \in \mathbb{N}$, $|b_n - 1|_{D'} \leq r$, and therefore $(b_n^{p^n})_n$ converges uniformly to 1 on D'. Therefore $(a_{n,D'})_n$ converges to $a \in \mathcal{O}^{\times}(D'')$ and $c_{D''}$ is the image of a by the boundary map $\kappa : \mathcal{O}_{Y_{\text{ét}}}^{\times} = \nu_* \mathcal{O}_Y^{\times} \to R^1 \nu_* \mathbb{Z}_p(1)$. Therefore $HT_Y(c)_{D''} = dlog(a)$ and [3, Prop. 2.4] shows that, for n big enough, there exists $y_n \in V(Y_n) \cap g_n^{-1}(D'') \subset V(Y_n) \cap (fg_n)^{-1}(D)$.

2 Reconstruction of the Berkovich topology

If X is a hyperbolic curve over K, let \overline{X} be its smooth compactification, so that $X = \overline{X} \setminus D_X$ for some divisor D_X on \overline{X} . Let |X| be the Berkovich space of X, considered as a topological space.

If \widetilde{X} is a universal pro-object in the category of finite étale cover of X, we will use the following notations : $|\widetilde{X}| = \varprojlim_{\widetilde{X} \to Y} |Y|$ and $|\widetilde{X}|_c = \varprojlim_{\widetilde{X} \to Y} |\overline{Y}|$, where Y goes through \widetilde{X} -pointed finite étale cover of X.

Proposition 2.1. Let X_1/K and X_2/K_2 be two p-adic hyperbolic curves satisfying rns and let $\widetilde{X}_i/\widetilde{K}_i$ be a universal pro-cover of X_i/K_i . Let $\alpha : \pi_1(X_1, \widetilde{X}_1) \to \pi_1(X_2, \widetilde{X}_2)$ be an isomorphism of fundamental groups. Then there is a unique homeomorphism of pro-Berkovich spaces $\tilde{\alpha} : |\widetilde{X}_1| \to |\widetilde{X}_2|$ and $\tilde{\alpha}_c : |\widetilde{X}_1|_c \to |\widetilde{X}_2|_c$ which are equivariant in the sense that the following diagram commutes:

$$\pi_1(X_1, \widetilde{X}_1) \times |\widetilde{X}_1|_c \longrightarrow |\widetilde{X}_1|_c$$

$$\downarrow^{\alpha \times \widetilde{\alpha}} \qquad \qquad \qquad \downarrow^{\widetilde{\alpha}}$$

$$\pi_1(X_2, \widetilde{X}_2) \times |\widetilde{X}_2|_c \longrightarrow |\widetilde{X}_2|_c$$

In particular, by quotienting by the étale fundmental group and the geometric étale fundamental groups, one gets homeomorphisms:

$$\begin{split} \bar{\alpha}: \overline{X}_1^{\mathrm{an}} \to \overline{X}_2^{\mathrm{an}}, \\ \bar{\alpha}_{\mathbb{C}_p}: \overline{X}_{1,\mathbb{C}_p}^{\mathrm{an}} \to \overline{X}_{2,\mathbb{C}_p}^{\mathrm{an}} \end{split}$$

Proof. Let $\Pi_i = \pi_1(X_i, \widetilde{X}_i)$, $\Delta_i = \pi_1(X_{i, \overline{K}_i}, \widetilde{X}_i) \subset \pi_1(X_i, \widetilde{X}_i)$ and $G_i = \Pi_i / \Delta_i \simeq \operatorname{Gal}(\overline{K_i}/K_i)$. Then $\alpha(\Delta_1) = \Delta_2$ by [5, Lem. 1.1.4] and therefore α induces an isomorphism $\alpha_G : G_1 \to G_2$.

Let $\Delta_i^{(p')}$ be the maximal pro-prime-to-p quotient of Δ_i and $\Pi_i^{(p')} = \Pi_i / \operatorname{Ker}(\Delta_i \to \Delta_i^{(p')})$ be the maximal geometrically pro-prime-to-p quotient of Π_i . Let $\alpha_{geom}^{(p')} : \Delta_1^{(p')} \to \Delta_2^{(p')}$ and $\alpha^{(p')} : \Pi_1^{(p')} \to \Pi_2^{(p')}$ be the morphisms induced by α .

Let H_1 be an open subgroup of Π_1 , and let $H_2 = \alpha(H_1)$. Let Y_1 be the connected finite étale cover of X_1 corresponding to H_1 and let Y_2 be the cover of X_2 corresponding to H_2 . Denote by L_i the field of constants of Y_i . Then cofinally on Y_1 , both Y_1/L_1 and Y_2/L_2 have stable reduction. Denote by \mathfrak{Y}_1 and \mathfrak{Y}_2 the corresponding stable models endowed with the natural log-structure. According to [5, Thm 2.7], α induces natural isomorphisms of log-special fibers $\alpha_{H_1,\log}: \mathfrak{Y}_{1,s} \to \mathfrak{Y}_{2,s}$. More precisely, when one identifies the prime-to-p geometric log fundamental group $\pi_1^{\log-\text{geom}}(\mathfrak{Y}_{1,s})$ of $\mathfrak{Y}_{i,s}$ with $(H_i \cap \Delta_i)^{(p')}$ along the specialisation map, the morphism $\pi_1^{\log-\text{geom}}(\mathfrak{Y}_{1,s}) \to \pi_1^{\log-\text{geom}}(\mathfrak{Y}_{1,s})$ induced by $\alpha_{H_1,\log}$ coincides with $\alpha_{H_1,geom}^{(p')}: (H_1 \cap \Delta_1)^{(p')} \to (H_2 \cap \Delta_2)^{(p')}$. In particular, one gets an isomorphism of the geometric dual graphs $\mathbb{G}_{Y_1} \to \mathbb{G}_{Y_2}$ such that if z_1 is an edge or a vertex in \mathbb{G}_{Y_1} with image z_2 in \mathbb{G}_{Y_2} , then $\alpha_{H_1,geom}^{(p')}(D_{z_1}) = D_{z_2}$ where D_{z_i} is the decomposition group of z_i in $(H_i \cap \Delta_i)^{(p')}$. The construction (and unicity) of $\overline{\alpha}$ is then perfectly similar to the one in [3, Thm. 3.10], that we now recall.

If K_i is an open subgroup of Δ_i , let $\tilde{V}(K_i)$ (resp. $\tilde{E}(K_i)$) be the set of decomposition groups of vertices of \mathbb{G}_{Y_i} in $K_i^{(p')}$ where $Y_i \to X_i$ is the finite étale cover

corresponding to H_i , where H_i is any open subgroup of Π_i such that $K_i = H_i \cap \Delta_i$ and Y_i has stable reduction ($\tilde{V}(K_i)$ and $\tilde{E}(K_i)$ do not depend on the choice of H_i). Let $V(K_i) = \tilde{V}(K_i)/K_i$ and $E(K_i) = \tilde{E}(K_i)/K_i$ be the set of conjugation classes.

Let $K''_i \subset K'_i \subset K_i$ be three open subsets of Δ_i such that K'_i and K''_i are normal in K_i . let $j_{K''_i,K'_i} : (K''_i)^{(p')} \to (K'_i)^{(p')}$ be the induced map. Then for every $D \in \tilde{V}(K'_i)$ there exists a unique $D' \in \tilde{V}(K''_i)$ up to conjugation in K_i such that $j_{K''_i,K'_i}(D')$ is open in D. This induces an injection $i_{K''_i,K'_i} : V(K'_i)/K_i \to V(K''_i)/K_i$.

Let $D \in \tilde{E}(K_i)$. Let K' be a normal open subgroup of K_i and let $j : K'^{(p')} \to K^{(p')}$ be the induced map. One denotes by

 $A_{D,K'} := \{D' \in \tilde{V}(K') | j(D') \text{ is conjugate to an open subgroup of } D\}/K_i \subset V(K')/K_i.$

If $K'' \subset K'$, then $i_{K'',K'}$ maps $A_{D,K''}$ to $A_{D,K'}$, and let

$$A_D = \lim_{K'} A_{D,K'}$$

Let $D_1 \in \tilde{V}(K_i)$ such that $D \subset D_1$ and let D_2 be the only other subgroup in $\tilde{V}(K_i)$ such that $D \subset D_2$. The full subgraph $\mathbb{G}(A_{D,K'})$ of $\mathbb{G}_{K'}/K_i$ with vertices

$$A_{D,K'} \cup \{i_{K',K_i}(D_1), i_{K',K_i}(D_2)\}$$

is a line, so that $A_{D,K'}$ is naturally an ordered set for which D_1 is the minimal element and D_2 is the maximal. If one exchanges the role of D_1 and D_2 , the order is reversed.

The injective map $A_{D,K''} \to A_{D,K'}$ is increasing, so that one obtains an order on A_D , which is then, non-canonically, isomorphic to $\mathbf{Q} \cap (0,1)$ as an ordered set. Let \widehat{A}_D be the Dedekind completion of A_D : \widehat{A}_D is an ordered topological space noncanonically isomorphic to [0,1] as an ordered set. One denotes by $e(D_1)$ the minimal element of A_D . As a set \widehat{A}_D does not depend on D_1 but the order is reversed when one replace D_1 by D_2 . One then defines the topological space

$$S(K_i) := \left(V(K_i) \coprod_{D \in E(K_i)} \widehat{A}_D \right) / \sim$$

where \sim is generated by

 $\forall D \subset D_1, \quad D_1 \sim e(D_1).$

Since \widehat{A}_D is (a priori non canonically) homeomorphic to [0, 1], $S(K_i)$ is (a priori non canonically) homeomorphic to the geometric realization of $\mathbb{G}(K_i)$.

Let $\widetilde{B}_i = \varinjlim_{K_i} S(K_i)$. Then one has a natural homeomorphism $|\widetilde{X}_1|_c \to \widetilde{B}_i$ (see [3, Lem. 3.9]). Then the family of isomorphisms of graphs $\mathbb{G}(K_1) \to \mathbb{G}(K_2)$ induce a homeomorphism $\widetilde{B}_1 \to \widetilde{B}_2$, and therefore one gets a homeomorphism $\widetilde{\alpha}_c : |\widetilde{X}_1|_c \to |\widetilde{X}_2|_c$. It preserves cusps, for example by [6], so that it also induces a homeomorphism $\widetilde{\alpha} : |\widetilde{X}_1| \to |\widetilde{X}_2|$. The action of Π_i on itself by conjugacy induce an action of Π_i on \widetilde{B}_i , so that the homeomorphism $\varinjlim_{K_i} S(K_i) \to |\widetilde{X}_1|_c$ is Π_i -equivariant, and $\widetilde{B}_1 \to \widetilde{B}_2$ is also equivariant.

The uniqueness is identical to [3, Prop. 3.11].

The map $\bar{\alpha}_{C_n}$ is deduced from the fact that $\alpha(\Delta_1) = \Delta_2$.

3 The canonical metric

Let \mathbb{G} be a finite connected graph. A metric on \mathbb{G} is just a map $l: E(\mathbb{G}) \to \mathbb{R}_{>0}$ where $E(\mathbb{G})$ is the set of edges of \mathbb{G} . If x, y are two vertices of \mathbb{G} , a path from x to y is a sequence $p = (x = x_0, e_1, x_1, \dots, e_n, x_n = y)$ where for all i, x_i is a vertex of \mathbb{G} and e_i is an edge of \mathbb{G} abutting at x_{i-1} and x_i . We then set $d(x, y) = \inf_p l(p)$ where the infimum goes through all paths from x to y and $l(p) = \sum_{i=1}^n l(e_i)$ if $p = (x = x_0, e_1, x_1, \dots, e_n, x_n = y)$ is a path.

Let X^{\log}/k^{\log} be a semistable log curve over a log point, and let \mathbb{G}_X be the dual graph of X. Let us fix $\pi \in M_{k^{\log}} \setminus \{0\}$. Let e be a node of X^{\log} , then $\overline{M}_{X,e}$ is isomorphic to $\overline{M}_{k^{\log}} \oplus \mathbb{N}u \oplus \mathbb{N}v)/(u+v=a)$ for some $a \in \overline{M}_{k^{\log}} \setminus \{0\}$. Note that u and v are characterized as the generators of the faces of $\overline{M}_{X,e}$ isomorphic to N, so that, in particular, a is preserved by automorphisms of $\overline{M}_{X,e}$. Then $a = \pi^r$ for a unique $r \in \mathbb{Q}_{>0}$, and one can set $l_{X^{\log},\pi}(e) = r$ to get a metric $d_{X^{\log},\pi}$ on \mathbb{G}_X .

If X is an analytic \mathbb{C}_p -curve in the sense of Berkovich, the set $X_{[2]}$ of points of type 2 is endowed with a canonical metric (see [1, Prop. 4.5.7]). If $x, y \in X_{[2]}$, let an annular path from x to y be a sequence $p = (x = x_0, A_1, x_1, A_2, \dots, A_n, x_n = y)$, where for all $i, x_i \in X_2$ and A_i is a annulus embedded in X with endpoints x_{i-1} and x_i . The length of the path $p = (x = x_0, A_1, x_1, A_2, \dots, A_n, x_n = y)$ is then defined to be $l(p) = \sum_{i=1}^n r_{norm}(A_i)$ and the distance between x and y is $d(x, y) = \inf_p l(p)$ where the infimum goes through all annular paths from x to y. If X is topologically simply connected, d(x, y) = l(p) for all annular path $p = (x = x_0, A_1, x_1, A_2, \dots, A_n, x_n = y)$ such that, for all $i \neq j, A_i \cap A_j = \emptyset$.

Let \mathfrak{X} be a proper semistable log curve over \mathbb{C}_p with smooth generic fiber. Let $\pi : \mathfrak{X}_{\eta}^{\mathrm{an}} \to \mathfrak{X}_s$ be the specialisation map from the analytic generic fiber to the special fiber. Let $V(\mathfrak{X}_s)$ be the set of generic points of \mathfrak{X}_s , $N(\mathfrak{X}_s)$ be the set of nodal points of \mathfrak{X}_s and $S(\mathfrak{X}) = \pi^{-1}(V(\mathfrak{X}_s))$. Then π induces a bijection $S(\mathfrak{X}) \to V(\mathfrak{X}_s)$ and $S(\mathfrak{X})$ is a triangulation of $\mathfrak{X}_{\eta}^{\mathrm{an}}$ in the sense of [1, §5.1.13]. If $z \in N(\mathfrak{X}_s)$, then $\pi^{-1}(z)$ is an open annulus and $r_{\mathrm{norm}}(\pi^{-1}(z)) = l_{\mathfrak{X}_s^{\mathrm{log}},p}(z)$. Moreover, the distance between two points of $S(\mathfrak{X})$ is reached by an annular path $p = (x = x_0, A_1, x_1, A_2, \cdots, A_n, x_n = y)$ such that for all *i* there exists $z_i \in N(\mathfrak{X}_s)$ such that $A_i = \pi^{-1}(z_i)$. When one identifies $S(\mathfrak{X})$ and $V(\mathfrak{X}_s)$ along π , $d_{\mathfrak{X}_s^{\mathrm{log}},p}$ and the restriction of the canonical distance $d_{\mathfrak{X}_{\eta}^{\mathrm{an}}}$ to $S(\mathfrak{X})$ therefore coincide.

Proposition 3.1. *Keep the assumptions and the notations of proposition (2.1). Then* $\bar{\alpha}$ *preserves the points of type 2 and 3 and preserves the canonical distance on points of type 2: if* $x, y \in X_{1,[2]}$ *, then* $d(\bar{\alpha}(x), \bar{\alpha}(y)) = d(x, y)$ *.*

Proof. In \overline{X}_1^{an} , a point x is of type 2 if and only if $\overline{X}_1^{an} \setminus \{x\}$ has more than two connected components and is of type 3 if and only if $\overline{X}_1^{an} \setminus \{x\}$ has exactly two connected component. In particular points of type 2 (resp. 3) are preserved by arbitrary automorphisms, and therefore by $\overline{\alpha}$.

Let $x, y \in X_{1,[2]}^{an}$ and let $p = (x = x_0, A_1, x_1, A_2, \dots, A_n, x_n = y)$ be an annular path. By resolution of non singularities, there exists a finite étale cover $f : Y_1 \to X_1$ such that for every $i \leq n$, $f^{-1}(x_i) \cap V(Y_1) \neq \emptyset$. One can moreover assume that $f : Y_1 \to X_1$ is a Galois cover, and let $G = \text{Gal}(Y_1/X_1)$. Then, since $V(Y_1)$ is Gequivariant, for every $i, f^{-1}(x_i) \subset V(Y_1)$. Up to refining p, one may assume that one can lift p to a path $(y_0, B_1, y_1, B_2, \dots, B_n, y_n)$ of Y_1 such that $B_i = \operatorname{sp}_{Y_1}^{-1}(e_i)$ where e_i is a node of the special fiber $\mathfrak{Y}_{1,s}$ of the stable model of Y_1 . Then $r_{\operatorname{norm}}(A_i) = l_{\mathfrak{Y}_{1,s}^{\log}}(e_i)/|\operatorname{Stab}_G(\tilde{e}_i)|$, and $l(p) = \sum_i l_{\mathfrak{Y}_{1,s}^{\log}}(e_i)/|\operatorname{Stab}_G(\tilde{e}_i)|$. Let Y_2 be the corresponding cover of X_2 . Since the isomorphism of graphs $\mathbb{G}_{Y_1} \to \mathbb{G}_{Y_2}$ is *G*-equivariant and actually comes from an isomorphism of log-schemes $\mathfrak{Y}_{1,s}^{\log} \to \mathfrak{Y}_{2,s}^{\log}$, one deduces that $\bar{\alpha}$ preserves the length of p, and therefore the metric.

Proposition 3.2. *Keep the assumptions and the notations of proposition* (2.1)*. Then* $\bar{\alpha}$ *preserves the types of the points.*

Proof. In \overline{X}_{1}^{an} , a point x is of type 2 if and only if $\overline{X}_{1}^{an} \setminus \{x\}$ has more than two connected components and is of type 3 if and only if $\overline{X}_{1}^{an} \setminus \{x\}$ has exactly two connected component. In particular points of type 2 (resp. 3) are preserved by arbitrary automorphisms, and therefore by $\overline{\alpha}$. We still need to distinguish points of type 1 from points of type 4. $(\overline{X}_{i}^{an})_{(2)}$ is endowed with a natural metric d_{i} and a point x of \overline{X}_{i}^{an} is of type 1 if and only if for every sequence $(x_{n}) \in (\overline{X}_{i}^{an})_{(2)}^{\mathbb{N}}$ that converges to x, $d(x_{0}, x_{n})$ goes to infinity. But the restriction of d_{i} to $V(X_{i})$ can be recovered (up to a multiplicative constant, which is determined by the absolute ramification index of K_{i} , which can be recovered from $\pi_{1}(X_{i})$) by the log-structure on $\mathfrak{X}_{i,s}$. By doing so for every $\mathfrak{Y}_{i,s}$, one gets that for every $x, x' \in (\overline{X}_{1}^{an})_{(2)}, d_{1}(x, x') = d_{2}(\overline{\alpha}(x), \overline{\alpha}(x'))$. But a point x of \overline{X}_{i}^{an} is of type 1 if and only if for every sequence $(x_{n}) \in (\overline{X}_{i}^{an})_{(2)}^{\mathbb{N}}$ that converges to $x, d(x_{0}, x_{n})$ goes to $x, d(x_{0}, x_{n})$ goes to infinity. Therefore, $\overline{\alpha}$ also preserves points of type 1.

Theorem 3.3. Let X_1/K_1 and X_2/K_2 be two p-adic hyperbolic curves satisfying rns and let $\widetilde{X}_i/\widetilde{K}_i$ be a universal cover of X_1/K_1 . Let $\alpha : \pi_1(X_1, \widetilde{X}_1) \to \pi_1(X_2, \widetilde{X}_2)$ be an isomorphism of fundamental groups. Then α is induced by a unique isomorphism $\widetilde{X}_1 \to \widetilde{X}_2$.

Proof. Let \tilde{x}_1 be a closed point of \tilde{X}_1 . Then the decomposition group $D_{\tilde{x}_1}$ is the stabilizer of \tilde{x}_1 for the action of $\pi_1(X_1)$ on $\tilde{X}_1|_c$. Therefore, the equivariance of $\bar{\alpha}$ tells us that $\alpha(D_{\tilde{x}_1}) = D_{\bar{\alpha}(\tilde{x}_1)}$. According to Prop. 2.1, $\bar{\alpha}(\tilde{x}_1)$ is of type 1, so that $\mathcal{H}(x_1)$ is a closed subfield of \mathbb{C}_p . Since $k(x_1)$ is a finite extension of \mathbb{Q}_p , the image of $D_{\tilde{x}_1}$ inside G_{K_1} is open, therefore $\alpha(D_{\tilde{x}_1}) = D_{\bar{\alpha}(\tilde{x}_1)}$ maps to an open subgroup of G_{K_2} , so that $\mathcal{H}(x_2)$ is a finite extension of \mathbb{Q}_p . Therefore $\alpha(D_{\tilde{x}_1})$ is the decomposition group of a closed point of \tilde{X}_2 . The result then follows from [7, Thm 2.9].

4 Geometric characterization of the type of points

Let C be a algebraically closed complete non archimedean field of mixed characteristics (0, p).

Proposition 4.1. Let $D = \mathcal{M}(C\{T\})$ be a disk; let η be its Gauss point, and let $c \in H^1(D, \mathbb{Z}_p(1)) = \operatorname{Hom}(\pi_1^{\text{\'et}}(D), \mathbb{Z}_p(1))$. For $n \ge 1$, let $f_n : Y_n \to D$ be the corresponding μ_{p^n} -torsor, so that $\pi_1^{\text{\'et}}(Y_n) = c^{-1}(p^n \mathbb{Z}_p(1)) \subset \pi_1^{\text{\'et}}(D)$.

 $1. \ |\frac{HT_D(c)}{dT}|_{\eta} \le 1;$

2. For every $n \ge 1$, if f_n splits over η , then $|\frac{HT_D(c)}{dT}|_{\eta} \le \frac{1}{p^n}$.

Proof. 1. If $c = \kappa(f)$ with $f \in O^{\times}(D)$, then $HT_D(c) = \frac{df}{f}$. Up to multiplying f by a constant in C, one can assume that $|f|_{\eta} = 1$. Then $|\frac{HT_D(c)}{dT}|_{\eta} = |\frac{df}{dT}|_{\eta} \leq 1$. In general, the restriction of c to any proper closed disk is in the image of the Kummer map. In particular, if z is the Gauss point of a closed disk of radius $r \in (0,1) \cap |\mathbb{C}^{\times}|$, one deduce from the previous argument applied to the parameter $a^{-1}T$ where |a| = r that $|\frac{HT_D(c)}{dT}|_{\eta_r} \leq \frac{1}{r}$. When r converges to 1, one gets that $|\frac{HT_D(c)}{dT}|_{\eta} \leq 1$.

2. Assume f_n splits over η . Then it splits over a neighborhood of η , which contains a small annulus $\{r \leq |T| \leq 1\}$. By gluing with the trivial μ_{p^n} -torsor over the disk $\{0 \leq |\frac{1}{T}| \leq \frac{1}{r}\}$ centered at infinity, one obtains a μ_{p^n} -torsor over \mathbb{P}^1 , which is necessarily trivial. Therefore f_n is a trivial μ_{p^n} -torsor.

The pullback $f_n^* c \in \operatorname{Hom}(\pi_1^{\operatorname{alg}}(Y_n), \mathbb{Z}_p(1))$ has its image in $\operatorname{Hom}(\pi_1^{\operatorname{alg}}(Y_n), p^n \mathbb{Z}_p(1))$, so there exists $c' \in \operatorname{Hom}(\pi_1^{\operatorname{alg}}(Y_n), \mathbb{Z}_p(1))$ such that $f_n^* c = p^n c'$, and so $f_n^* HT_{Y_n}(c) = p^n HT_{Y_n}(c')$. If f_n is a trivial cover, let $s: D \to Y_n$ be a section of f_n . Then $HT_D(c) = p^n HT_D(s^*c')$ and therefore $|\frac{HT_D(c)}{dT}|_{\eta} = \frac{1}{p^n}|\frac{HT_D(s^*c')}{dT}|_{\eta} \leq \frac{1}{p^n}$.

Proposition 4.2. Let X be a hyperbolic curve over C. Let $x \in X^{\text{an}}$ be a type 4 point. Let $c \in Hom(\pi_1^{\text{ét}}(X), \mathbb{Z}_p(1))$ such that $\operatorname{HT}_X(c) \neq 0$. Then c is not split over x: its restriction $c_{D_x} \in \operatorname{Hom}(D_x, \mathbb{Z}_p(1))$ to the decomposition group D_x is non-zero.

Proof. Let $f_n: Y_n \to X$ be the μ_{p^n} -torsor induced by c.

The point x is contained in a closed disk. Since $\operatorname{HT}(c)$ vanishes in D only at finitely many points of type 1, one can assume after replacing D by a smaller disk that $\operatorname{HT}(c)$ does not vanish on D. Let T be a parameter on D. Then $\frac{HT(c)}{dT}$ has constant norm λ on D. Let us normalize the radii of subdisks of D such that $r(\{|T-a| \leq \alpha\}) = \alpha$. Then if $f_n : Y_n \to X$ is split over a strict closed disk D then $r(D) \leq \frac{\lambda}{p^n}$. Then there exists a decreasing sequence of strict closed disks $(D_k)_{k\in\mathbb{N}}$ in D such that $x = \lim_k \eta_{D_k}$ where η_{D_k} is the Gauss point of D_k . Since x if of type 4, $r := \lim_k r(D_k) > 0$. In particular, there exists $n \in \mathbb{N}$ such that $\frac{\lambda}{p^n} < r$. Therefore, for every $k \in \mathbb{N}, \frac{\lambda}{p^n} < r(D_k)$, so that $Y_n \to X$ does not split over x_k . Therefore it does not split over x either. In particular $c_{D_x} \neq 0$.

This criterion gives another proof of proposition 3.2 that does not use the recovery of the log structure on the special fiber. This allows to deduce the following purely geometric analogue of 3.2.

Definition 4.1. Let Π be a topological group.

One denotes by RNS(Π) the following property: there exists a hyperbolic curve X over an algebraically closed complete non-archimedean field C of mixed characteristics (0, p) which satisfies resolution of non-singularity and an isomorphism $a : \Pi \to \pi_1^{\text{temp}}(X, \widetilde{X})$.

Such an isomorphism $a: \Pi \to \pi_1^{\text{temp}}(X, \widetilde{X})$ will be called a geometric setting of Π .

Let Π be a topological group such that $\text{RNS}(\Pi)$ is true. By [2, Th. 3.39], one has a group-theoretical graph that recovers the graph of the stable reduction in the same way as in [6] over discretely valued fields. More precisely, one sets $\mathbb{G}(\Pi)$ to be the graph whose vertices are conjugacy classes of maximal compact subgroups of $\Pi^{\text{prime}-\text{to}-p}$ and whose edges are conjugacy classes of non-trivial intersections of two distinct maximal compact subgroups of $\Pi^{\text{prime}-\text{to}-p}$. Then any geometric setting a : $\Pi \to \pi_1^{\text{temp}}(X)$ induces an isomorphism $\mathbb{G}(\Pi) \to \mathbb{G}(X)$.

Then, the construction of [3, §3.3] (or the construction in the proof of 2.1) extends without modification to this context (the only slight difference is that in the notations of [3, §3.3], A_e is no longer isomorphic to $\mathbb{Q} \cap (0, 1)$ in general, but to $v(\mathbb{C}^{\times}) \cap (0, 1)$ where $v(\mathbb{C}^{\times})$ is the value group of C): one obtains a topological space $\widetilde{X}_c(\Pi)$ endowed with a continuous action of Π , and for every geometric setting $a : \Pi \to \pi_1^{\text{temp}}(X, \widetilde{X})$ a unique equivariant homeomorphism $\widetilde{a}_c : \widetilde{X}_c(\Pi) \to |\widetilde{X}^{\text{an}}|_c$.

Let $\overline{X}(\Pi) = \widetilde{X}_c(\Pi)/\Pi$. If $\tilde{x} \in \widetilde{X}_c(\Pi)$, let $D_{\tilde{x}} = \operatorname{Stab}_{\Pi}(\tilde{x})$. If $x \in \overline{X}(\Pi)$, let $D_x = D_{\tilde{x}}$, where \tilde{x} is a preimage of x in $\widetilde{X}(\Pi)$, so that it is only well-defined up to conjugation. If $\Pi = \pi_1^{\operatorname{temp}}(X)$, then D_x coincides with the image of the outer morphism $G_{\mathcal{H}(x)} \to \Pi_X$ induced by the morphism of analytic spaces $\mathcal{M}(\mathcal{H}(x)) \to X^{\operatorname{an}}$, where $G_{\mathcal{H}(x)}$ is the absolute Galois group of $\mathcal{H}(x)$.

Cusps of $\overline{X}(\Pi)$ are end-points x of $\overline{X}(\Pi)$ (in the sense that the set of open neighborhoods U of x such that $U \setminus \{x\}$ is connected are cofinal among open neighborhoods of x) such that D_x is isomorphic to $\widehat{\mathbb{Z}}$, and we denote by $X(\Pi) = \overline{X}(\Pi) \setminus \{\text{cusps}\}$. Then for every geometric setting $a : \Pi \to \pi_1^{\text{temp}}(X, \widetilde{X})$, \tilde{a}_c restricts to a homeomorphism $X(\Pi) \to |X^{\text{an}}|$ (it follows for example from the following proposition 4.3).

Definition 4.2. Let $x \in X(\Pi)$. Let l be a prime number different from p. Then x is said to be of type:

- 1 if $D_x = \{1\};$
- 2 if D_x has a non abelian pro-*l* subquotient;
- 3 if D_x has a non trivial pro-*l* subquotient but every pro-*l* subquotient of D_x is abelian.
- 4 if D_x is a non-trivial pro-p group.

Proposition 4.3. For any geometric setting $a : \Pi \to \pi_1^{\text{temp}}(X)$, the isomorphism $\bar{a} : X(\Pi) \simeq |X^{\text{an}}|$ preserves the types of points. In particular, the type of a point in definition 4.2 is independent from l.

Proof. From the definition 4.2, it is clear that a point in $X(\Pi_Z)$ cannot be of two different types. It is therefore sufficient to show that if $x \in Z^{\text{an}}$ is of a given type, $D_x \subset \Pi_Z$ satisfies the corresponding property in definition 4.2.

1. Assume x is of type 1. Then $D_x = \{1\}$ because $\mathcal{H}(x) = C$ is algebraically closed.

- 2. Assume x is of type 2. By resolution of non-singularities, there exists a finite cover f : Y → Z and y ∈ f⁻¹({x}) ∩ V(Y). Then D_y = D_x ∩ Π_Y (for some choice of D_x inside its conjugacy class), and the image of D_y in Π_Y^{prime-to-p} is isomorphic to π₁^{ét}(U)^{prime-to-p} where U is the smooth locus in the irreducible component of the special fiber of the stable model of Y corresponding to the specialisation of y. Since U is an hyperbolic curve over the residue field C of C, D_y has a non abelian pro-l quotient.
- 3. Assume x is of type 3. By resolution of non-singularities, there exists a finite Galois cover f : Y → Z and y ∈ f⁻¹({x}) such that y belongs to the skeleton of Y^{an}. Then the image of D_y in Π^{prime-to-p}_Y is isomorphic to Z(1)^{prime-to-p}, and therefore has a non trivial pro-l quotient.

Since $\mathcal{H}(x)$ has an algebraically closed residue field, the absolute galois group $G_{\mathcal{H}(x)}$ of $\mathcal{H}(x)$ is equal to its inertia subgroup $I_{\mathcal{H}(x)}$. Since the wild inertia subgroup $W_{\mathcal{H}(x)}$ is a pro-*p* group, every subquotient of $G_{\mathcal{H}(x)}$ is isomorphic to a subquotient of $I_{\mathcal{H}(x)}/W_{\mathcal{H}(x)}$ and is therefore abelian (cf. [1, §2.3.19]).

4. Assume x is of type 4. Then according to proposition 4.2, D_x is non trivial. Since D_x is a quotient of the absolute Galois group of H(x), it is enough to show that its absolute galois group G_{H(x)} is a p-group. Since the residue field of H(x) is algebraically closed and its value group is divisible, G_{H(x)} is equal to its wild inertia subgroup W_{H(x)}, which is a pro-p group.

References

- [1] Antoine Ducros, La structure des courbes analytique.
- [2] Sylvain Gaulhiac, Reconstruction anabélienne du squelette des courbes analytiques, Annales de l'Institut Fourier 73 (2023), no. 3, 999–1084 (fr).
- [3] Emmanuel Lepage, *Resolution of nonsingularities for Mumford curves*, Publ. Res. Inst. Math. Sci. **49** (2013), no. 4, 861–891.
- [4] Shinichi Mochizuki, *The local pro-p anabelian geometry of curves*, Inventiones Mathematicae **138** (1999), 319–423.
- [5] _____, *The absolute anabelian geometry of hyperbolic curves*, Galois theory and modular forms, Kluwer Academic publishers, 2003, pp. 77–122.
- [6] _____, Semi-graphs of anabelioids, Publications of the Research Institute of Mathematical Sciences 42 (2006), no. 1, 221–322.
- [7] _____, Topics in anabelian geometry II: Decomposition groups and endomorphisms, J. Math. Sci. Univ Tokyo **20** (2013), 171–269.
- [8] Peter Scholze, *Perfectoid spaces: A survey*, Current Developments in Mathematics 2012 (2013).