ON THE TOPOLOGY OF FIBER-TYPE CURVES: A ZARISKI PAIR OF AFFINE NODAL CURVES

J.I. COGOLLUDO-AGUSTÍN AND EVA ELDUQUE

ABSTRACT. In this paper we explore conditions for a curve in a smooth projective surface to have a free product of cyclic groups as the fundamental group of its complement. It is known that if the surface is \mathbb{P}^2 , then such curves must be of fiber type, i.e. a finite union of fibers of an admissible map onto a complex curve. In this setting, we exhibit an infinite family of Zariski pairs of fiber-type curves, that is, pairs of plane projective fiber-type curves whose tubular neighborhoods are homeomorphic, but whose embeddings in \mathbb{P}^2 are not. This includes a Zariski pair of curves in \mathbb{C}^2 with only nodes as singularities (and the same singularities at infinity) whose complements have non-isomorphic fundamental groups, one of them being free. Our examples show that the position of nodes also affects the topology of the embedding of fiber-type curves.

1. INTRODUCTION

In a series of recent papers by different authors ([8, 7, 2, 10]) there is a growing interest in studying the connection between the geometric properties of a smooth connected complex quasi-projective surface U and its fundamental group. This paper continues [10], which deals with the case where $\pi_1(U)$ is a free product of cyclic groups.

One of the main results by the authors in this direction is [10, Theorem 1.3], which gives geometric necessary conditions for a curve inside of a smooth projective surface to have a free product of cyclic groups as the fundamental group of its complement. The flavor of this result can be summarized as follows: if X is a smooth connected complex projective surface and $D \subset X$ is a curve such that $\pi_1(X \setminus D)$ is a free product of cyclic groups, then there exists a smooth projective curve S and an admissible (see Definition 2.1) map $F : X \setminus \mathcal{B} \to S$ inducing an isomorphism $\pi_1(X \setminus D) \to \pi_1^{\text{orb}}(S)$, where \mathcal{B} is a finite set of points in D and $\pi_1^{\text{orb}}(S)$ is the orbifold fundamental group for some orbifold structure on S induced by the multiplicity of the fibers of $F_{|X \setminus D}$. Moreover, D is the closure of a finite union of fibers of F and of special null-homology horizontal components. See Theorem 2.14 in section 2.3 for the precise statement when $X = \mathbb{P}^2$.

However, the existence of the admissible map F (sometimes also called a pencil map) such that D is (essentially) a finite union of fibers is not a sufficient condition for the fundamental group of $U := X \setminus D$ to be a free product of cyclic groups. For instance, if $X = \mathbb{P}^2$ and D contains only generic fibers of an admissible map of a certain kind (satisfying Condition 2.3 in section 2.1), then $\pi_1(U)$ is a free product of cyclic groups (see [10, Theorem 1.2]). On the other extreme, if D contains all atypical fibers of an admissible map, then $\pi_1(U)$ tends to not be a free product of cyclic groups, but rather a semidirect product of the fundamental groups of two

²⁰²⁰ Mathematics Subject Classification. 32S25, 32S55, 32S05, 32S20, 57K31.

The authors are partially supported by the Spanish Government PID2020-114750GB-C31. The first author is partially supported by the Departamento de Ciencia, Universidad y Sociedad del Conocimiento del Gobierno de Aragón (Grupo de referencia E22_20R "Álgebra y Geometría"). The second author is partially supported by the Ramón y Cajal Grant RYC2021-031526-I funded by MCIN/AEI /10.13039/501100011033 and by the European Union NextGenerationEU/PRTR.

smooth quasi-projective curves [10, Lemma 2.18, Corollary 2.19]. Thus, it is natural to ask if it is possible to formulate sufficient conditions for the fundamental group of a curve complement to be a free product of cyclic groups in terms of the topology of the fibers contained in D in the case when some of these fibers contain only mild singularities (for example nodes).

In section 3, we describe this problem in precise terms by defining a partial order in the subsets of the set of atypical values B_F of the admissible map $F: X \setminus \mathcal{B} \to S$. The set of atypical values B_F is the minimal set of values $B_F \subset S$ for which $F: (X \setminus \mathcal{B}) \setminus F^{-1}(B_F) \to S \setminus B_F$ is a locally trivial fibration, which is well known to be a finite set. The set of all minimal elements of this partial order in the set of subsets of B_F is called the threshold of atypical values and, if X is \mathbb{P}^2 , it completely describes the remarkable phenomenon of $\pi_1(X \setminus D)$ being a free product of cyclic groups for curves D which are the closure of two or more fibers of F.

In section 4 we explore whether or not the "free product of cyclic groups" condition for $\pi_1(X \setminus D)$ could be determined by any topological information provided by the atypical fibers of F which lie in D. To answer this question in the negative, we give in section 4.1 a particular example of a Zariski pair of curves, both of which consist of two homeomorphic fibers in \mathbb{C}^2 of the same polynomial map, plus the line at infinity. A Zariski pair (see [3]) refers to two algebraic curves $D_1, D_2 \subset \mathbb{P}^2$ with homeomorphic regular neighborhoods, but non-homeomorphic embeddings, that is, $(\mathbb{P}^2, D_1) \not\cong (\mathbb{P}^2, D_2)$. In particular, both curves have the same *combinatorial type* (see the discussion after Definition 2 in [5]) which includes the same degree and the same local types of singularities.

The previously mentioned example discussed in section 4.1 is a Zariski pair of two projective curves of degree 13, each one of them being the union of two nodal sextics and a line. In fact, both curves share the line and a sextic. Considering the line as the line at infinity, their complement can be thought of as the complement in \mathbb{C}^2 of two affine nodal sextics whose fundamental groups are not isomorphic, namely, one of them is free and the other one is not. Moreover, since they are constructed as the union of two fibers of a polynomial map, their singularities at infinity coincide, hence the difference in fundamental groups seems to be determined by the position of the nodes on the affine plane alone. Note that other examples of Zariski pairs where there is a connection between the topology of their complement and the position of singular points (including nodes) have been exhibited before (see [18, 6]). However, the remarkable aspect of the present family of examples stands for the fact that the only special position of singularities is given by the nodes. As far as we know, this phenomenon has never been described before -it is known that projective nodal curves have abelian fundamental groups (cf. [13, 12]) and hence there cannot be a Zariski pair of projective nodal curves.

In section 4.2 it is shown that both curve complements have the same Alexander invariants and that twisted Alexander polynomials can tell these fundamental groups apart, which is the first example in the literature to our knowledge. In section 4.3 we prove that the union of the two sextics for both of these curves yields two projective curves of degree 12 which also form a Zariski pair. Finally, in section 4.4 we compute the threshold of atypical values corresponding to the morphism $\mathbb{P}^2 \dashrightarrow \mathbb{P}^1$ which gives rise to the Zariski pairs in sections 4.1 and 4.3. This completely characterizes all the (infinitely many) Zariski pairs whose irreducible components are fibers of this morphism, and such that the fundamental groups of their complements satisfy that one of them is a free product of cyclic groups (see Theorem 4.12).

2. Preliminaries

2.1. Admissible maps. The following definition is found in Arapura [1].

Definition 2.1. A surjective morphism $F: U \to S$ from a smooth quasi-projective surface U to a smooth projective curve S is admissible if it admits a surjective holomorphic enlargement $\hat{F}: \hat{U} \to S$ with connected fibers, where \hat{U} is a smooth compactification of U.

Remark 2.2. Let $F : U \to S$ be a surjective morphism from a smooth quasiprojective surface to a smooth projective curve. Then F is admissible if and only if F has connected generic fibers. Indeed, if $F : U \to S$ is admissible and $\hat{F} : \hat{U} \to S$ is a holomorphic enlargement with connected fibres, the generic fiber of \hat{F} is smooth by generic smoothness on the target. Hence, the generic fiber of F is a Zariski open subset of a smooth connected compact curve, which is connected. For the other implication, suppose that $\hat{F} : \hat{U} \to S$ is a holomorphic enlargement of F, where \hat{U} is a smooth compactification of U. Then, Stein factorization implies that \hat{F} factors as a map with connected fibers $G : X \to S'$ and a surjective finite morphism $h : S' \to S$, where S' can be taken to be a a smooth projective curve. Since F has connected generic fibers, and F also factors through h, h must be a branched cover of degree d = 1, so h is a homeomorphism and thus \hat{F} has connected fibers.

Most of the maps that will appear in this paper will be dominant rational morphisms $F : \mathbb{P}^2 \longrightarrow \mathbb{P}^1$ such that the restriction to its maximal domain of definition is an admissible map. Any dominant rational morphism F can be expressed by a surjective morphism $F = [f : g] : U \to \mathbb{P}^1$ when restricted to its maximal domain of definition U, where $f, g \in \mathbb{C}[x, y, z]$ are homogeneous polynomials of the same degree with no non-constant common factors, and $U = \mathbb{P}^2 \setminus \mathcal{B}$, where \mathcal{B} is the finite set $V(f) \cap V(g)$. In particular, most of the maps that will appear in this paper will be of the form described below:

Condition 2.3. $F = [f_q^p : f_p^q] : U \to \mathbb{P}^1$ is such that

- $p, q \in \mathbb{Z}_{>1}$, with gcd(p,q) = 1,
- f_p, f_q ∈ C[x, y, z] are homogeneous polynomials of degrees p and q respectively, with no non-constant common factors,
- neither f_p nor f_q is a k-th power of another polynomial in $\mathbb{C}[x, y, z]$ for any $k \geq 2$,
- $U = \mathbb{P}^2 \setminus \mathcal{B}$, where \mathcal{B} is the finite set $V(f_p) \cap V(f_q) \in \mathbb{P}^2$,
- The map F has no multiple fibers outside $\{[0:1], [1:0]\}$.

The last point in Condition 2.3 is necessary (up to change of coordinates in \mathbb{P}^1) for F to be admissible, as shown in the following remark.

Remark 2.4. Let $F: U \to \mathbb{P}^1$ be the restriction to its maximal domain of definition of a dominant rational map $F: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$. If $F: U \to \mathbb{P}^1$ is admissible, [11, Proposition 2.8] implies that the number of multiple fibers of F does not exceed 2, so, after a change of coordinates in \mathbb{P}^1 , we may assume that the multiple fibers of F lie over a subset of $\{[0:1], [1:0]\}$.

Remark 2.5. Let $f : \mathbb{C}^2 \to \mathbb{C}$ be a non-constant polynomial map with connected generic fibers. Then, f extends to an admissible map $F : \mathbb{P}^2 \setminus \mathcal{B} = [\overline{f}(x, y, z) : z^d] \to \mathbb{P}^1$, where \overline{f} is the homogenization of f with respect to the variable z, d is the degree of f, and $\mathcal{B} = V(\overline{f}(x, y, z)) \cap V(z)$.

It is easy to construct admissible maps satisfying Condition 2.3, as exemplified by the following result. **Lemma 2.6.** Let $F = [f_{kq}^p : f_{kp}^q] : U = \mathbb{P}^2 \setminus (V(f_{kq}) \cap V(f_{kp})) \to \mathbb{P}^1$ such that $k \in \mathbb{Z}_{\geq 1}$, gcd(p,q) = 1, and $f_{kq}, f_{kp} \in \mathbb{C}[x, y, z]$ are irreducible homogeneous polynomials of degree kq and kp respectively that are not constant multiples of one another. Then, F is an admissible map. Moreover, if k = 1, then F also satisfies Condition 2.3 (maybe after a change of coordinates in \mathbb{P}^1 if either p or q equal 1).

Proof. Let $\hat{F}: X \to \mathbb{P}^1$ be a resolution of indeterminacies of the pencil $F: \mathbb{P}^2 \dashrightarrow \mathbb{P}^1$. In particular, X is a smooth simply connected projective surface. Using Stein factorization and the fact that X is simply connected, \hat{F} factors as $H \circ \hat{G}$, where $\hat{G}: X \to \mathbb{P}^1$ has connected fibers and $H: \mathbb{P}^1 \to \mathbb{P}^1$ is finite, generically d: 1. Our goal is to show that d = 1. Let $G := \hat{G}_{|U}: U \to \mathbb{P}^1$, which is an admissible map by construction.

The map H (resp. G) is of the form $[h_1(x, y) : h_2(x, y)]$ (resp. [f(x, y, z) : g(x, y, z)], where $h_1, h_2 \in \mathbb{C}[x, y]$ (resp. $f, g \in \mathbb{C}[x, y, z]$) are homogeneous polynomials of the same degree with no non-constant common factors. Hence,

$$f_{kq}^p = h_1(f,g), \quad f_{kp}^q = h_2(f,g),$$

Since f_{kq} and f_{kp} are irreducible and every homogeneous polynomial in $\mathbb{C}[x, y]$ decomposes as a product of linear forms, we have that there exist $[a:b], [c:d] \in \mathbb{P}^1$ and $m, l \in \mathbb{Z}_{\geq 1}$ such that

$$h_1(x,y) = (bx - ay)^m, \quad m \mid p, \quad h_2(x,y) = (dx - cy)^l, \quad l \mid q,$$

and since h_1 and h_2 have the same degree, m = l. The condition that gcd(p,q) = 1 implies that m = l = 1. Hence, H is a change of coordinates in \mathbb{P}^1 , which concludes the proof of the fact that F is admissible. By Remark 2.4, if k = 1, F also satisfies Condition 2.3, possibly after a change of coordinates in \mathbb{P}^1 .

Example 2.7. The result of Lemma 2.6 is no longer true if the irreducibility assumption is dropped. For example, if $f_p = x^p - y^p$ and $f_q = x^q + y^q$ and p is odd, then $[f_p^q : f_q^p] : \mathbb{P}^2 \setminus \{[0:0:1]\} \to \mathbb{P}^1$ is not an admissible map, since the closure of any of its fibers is a union of lines in \mathbb{P}^2 going through the point [0:0:1].

2.2. Orbifold fundamental groups and morphisms. Let S be a smooth projective curve of genus g. We may endow it with an orbifold structure by choosing $\varphi : S \to \mathbb{Z}_{\geq 0}$ such that $\varphi(P) \neq 1$ only for a finite number of points. Let $\Sigma = \Sigma_0 \cup \Sigma_+ \subset \overline{S}$ be the points for which $\varphi(P) = 0$ if $P \in \Sigma_0$ and $\varphi(Q) = m_Q > 1$ if $Q \in \Sigma_+$. We will denote this structure by $S_{(n+1,\bar{m})}$, where $n+1 = \#\Sigma_0$, and \bar{m} is a $(\#\Sigma_+)$ -tuple whose entries are the corresponding m_Q 's.

As a word of caution for the following definitions, the term orbifold might be misleading: we do not need to develop any theory of orbifolds or V-manifolds in this context, but rather use the orbifold structures to highlight the existence of multiple fibers of an admissible map. This will become clear throughout the section.

Definition 2.8 (Orbifold fundamental group of a smooth projective curve). Let $S_{(n+1,\bar{m})}$ be a smooth projective curve endowed with an orbifold structure. The orbifold fundamental group of $S_{(n+1,\bar{m})}$ is defined by

$$\pi_1^{\operatorname{orb}}(S_{(n+1,\bar{m})}) := \pi_1(S \setminus \Sigma) / \langle \mu_P^{\varphi(P)}, P \in \Sigma \rangle,$$

where μ_P is a meridian in $S \setminus \Sigma$ around $P \in \Sigma$, and $\langle \mu_P^{\varphi(P)}, P \in \Sigma \rangle$ is the normal subgroup generated by the $\mu_P^{\varphi(P)}$'s.

Note that if S has genus g, $\pi_1^{\text{orb}}(S_{(n+1,\bar{m})})$ is hence generated by $\{a_i, b_i\}_{i=1,\ldots,g} \cup \{\mu_P\}_{P \in \Sigma}$ and presented by the relations

(1)
$$\mu_P^{m_P} = 1$$
, for $P \in \Sigma_+$, and $\prod_{P \in \Sigma} \mu_P = \prod_{i=1,\dots,g} [a_i, b_i].$

In particular, if $\Sigma_0 \neq \emptyset$, (1) shows that $\pi_1^{\text{orb}}(S_{(n+1,\bar{m})})$ is a free product of cyclic groups as follows

$$\pi_1^{\mathrm{orb}}(S_{(n+1,\bar{m})}) \cong \pi_1(S \setminus \Sigma_0) \ast \left(\ast_{P \in \Sigma_+} \left(\frac{\mathbb{Z}}{m_P \mathbb{Z}} \right) \right) \cong \mathbb{F}_r \ast \mathbb{Z}_{m_1} \ast \cdots \ast \mathbb{Z}_{m_s},$$

where $r = 2g - 1 + \#\Sigma_0 = 2g + n$, $s = \#\Sigma_+$, and $\bar{m} = (m_1, \ldots, m_s)$. Note that any finitely generated free product of cyclic groups can be realized as an orbifold fundamental group of a smooth projective curve in this way.

Definition 2.9 (Orbifold morphism). Let U be a smooth algebraic variety. A dominant algebraic morphism $F: U \to S_{(n+1,\bar{m})}$ defines an orbifold morphism if for all $P \in S$ such that $\varphi(P) > 0$, the divisor $F^*(P)$ is a $\varphi(P)$ -multiple.

Definition 2.10 (Maximal orbifold structures). Let $F : U \to S_{(n+1,\bar{m})}$ be an orbifold morphism. The orbifold $S_{(n+1,\bar{m})}$ is said to be maximal (with respect to F) if $F(U) = S \setminus \Sigma_0$ and for all $P \in F(U)$ the divisor $F^*(P)$ is not an *n*-multiple for any $n > \varphi(P)$.

The following result is well known (see [4, Prop. 1.4], for example)

Remark 2.11. Let $F: U \to S_{(n+1,\bar{m})}$ be an orbifold morphism. Then, F induces a morphism $F_*: \pi_1(U) \to \pi_1^{\text{orb}}(S_{(n+1,\bar{m})})$. Moreover, if the generic fiber of F is connected, then F_* is surjective.

We end this section with important notational conventions that will be used throughout the rest of the paper:

- Whenever we refer to a dominant algebraic morphism $F: U \to S$ as an orbifold morphism, we will assume that S is endowed with the orbifold structure that turns F into an orbifold morphism in such a way that the orbifold structure is maximal.
- The only orbifold structures that will appear are the maximal ones.
- In particular, if $F: U \to S$ is surjective and $B \subset S$, the notation $F_*: \pi_1\left(U \setminus F^{-1}(B)\right) \to \pi_1^{\text{orb}}(S \setminus B)$ will refer to the morphism endowed by $F_{|U \setminus F^{-1}(B)}: U \setminus F^{-1}(B) \to S$ on (orbifold) fundamental groups, where S is endowed with the orbifold structure with is maximal with respect to $F_{|U \setminus F^{-1}(B)}$ (which has $\Sigma_0 = B$).

2.3. Fiber-type curves.

Definition 2.12 (Fiber-type curve). Let X be a smooth projective surface, and let C be a curve in X. We say that a C is a fiber-type curve if, for some complement of a finite number of points in X called U, there exists an admissible map $F: U \to S$ to a smooth projective curve S such that C is the closure in X of a finite number of fibers of F.

Remark 2.13. Let $f : \mathbb{C}^2 \to \mathbb{C}$ be a non-constant polynomial map with connected generic fibers. We will also refer to the union C of a finite number of fibers of fas an *affine fiber-type curve*. Indeed, both points of view agree if we want to study the topology of their complements: f extends to an admissible map $F : \mathbb{P}^2 \setminus \mathcal{B} =$ $[\overline{f}(x, y, z) : z^d] \to \mathbb{P}^1$ as in Remark 2.5. The complement in \mathbb{P}^2 of the union of \overline{C} and the line at infinity (which is the closure in \mathbb{P}^2 of the fiber of F over [1:0]) coincides with $\mathbb{C}^2 \setminus C$.

As mentioned in the introduction, the curves in a smooth projective surface X whose complements have a fundamental group which is a free product of cyclic groups are (essentially) fiber-type curves (with perhaps some extra null-homotopic irreducible components). In fact, if $X = \mathbb{P}^2$, the curves must actually be fiber-type curves, as exemplified by the following result:

Theorem 2.14 (Main Theorem for curves in \mathbb{P}^2 [10, Cor. 3.14]). Let D be a curve in \mathbb{P}^2 . Suppose that $\pi_1(\mathbb{P}^2 \setminus D)$ is a free product of cyclic groups. Then, there exists $r \geq 0$ and $m_1 \geq m_2 \geq 1$ with $gcd(m_1, m_2) = 1$ such that $\pi_1(\mathbb{P}^2 \setminus D) \cong \mathbb{F}_r * \mathbb{Z}_{m_1} * \mathbb{Z}_{m_2}$. Moreover, there exists $F = [f_q^p : f_p^q] : U = \mathbb{P}^2 \setminus \mathcal{B} \to \mathbb{P}^1$ an admissible map as in Condition 2.3 such that:

(i) F induces an orbifold morphism

$$\mathcal{F}_{|}: \mathbb{P}^2 \setminus D \to \mathbb{P}^1_{(r+1,\bar{m})},$$

where $\mathbb{P}^1_{(r+1,\bar{m})}$ is maximal with respect to $F_{|}$, and \bar{m} takes the following value: (m_1, m_2) if $m_2 \ge 2$, (m_1) if $m_1 \ge 2$ and $m_2 = 1$, and it is empty if $m_1 = m_2 = 1.$

- (ii) $F_*: \pi_1(\mathbb{P}^2 \setminus D) \to \pi_1^{\operatorname{orb}}(\mathbb{P}^1_{(r+1,\bar{m})})$ is an isomorphism.
- (iii) $D = \overline{F^{-1}(\Sigma_0)} \subset \mathbb{P}^2$ is a fiber-type curve which is the closure of the union of r + 1 irreducible fibers of F.

More concretely,

- (1) If $m_1 > m_2 > 1$, then $p = m_1$, $q = m_2$ and the pencil $F = [f_{m_2}^{m_1} : f_{m_1}^{m_2}]$ has exactly two multiple fibers corresponding to $[0:1], [1:0] \notin \Sigma_0$.
- (2) If $m_1 > m_2 = 1$, then $p = m_1$, $[1:0] \in \Sigma_0$, and the pencil $F = [f_q^{m_1}: f_{m_1}^q]$ has at least one multiple fiber corresponding to $[0:1] \notin \Sigma_0$.
- (3) If $m_1 = m_2 = 1$, then $F = [f_q^p : f_p^q]$ has at most two multiple fibers corresponding to $[0:1], [1:0] \in \Sigma_0$.

Remark 2.15. This result implies that, if an affine curve satisfies that the fundamental group of its complement in \mathbb{C}^2 is a free product of cyclic groups, then it is a free group, and moreover, the affine curve must consist of a finite union of irreducible fibers of a polynomial map $f : \mathbb{C}^2 \to \mathbb{C}$.

The goal of Section 4 is to find Zariski pairs in this setting, where the irreducible components of both curves will be closures of fibers of the same admissible map.

We introduce the following notation.

Notation 2.16. Let U be a smooth quasi-projective surface and let S be a smooth projective curve. If $F: U \to S$ is an admissible map and $B \subset S$ is a finite set, we let U_B be

$$U_B = U \setminus F^{-1}(B).$$

Remark 2.17. Let $F = [f : g] : U = \mathbb{P}^2 \setminus \mathcal{B} \to \mathbb{P}^1$ be an admissible map, where $f, g \in \mathbb{C}[x, y, z]$ are homogeneous polynomials of degree $d \geq 1$ with no non-constant common factors, and $\mathcal{B} = V(f) \cap V(g)$. Let B be a non-empty finite subset of \mathbb{P}^1 , and let $D = \overline{F^{-1}(B)}$. Then $U_B = \mathbb{P}^2 \setminus D$.

The following result can be found in [10, Thm. 1.4, Cor. 4.9].

Theorem 2.18 (Addition-Deletion Lemma). Let U be a smooth quasi-projective surface and let $F: U \to S$ be an admissible map to a smooth projective curve S. Assume $B \subset S$, where $\#B = n \geq 1$, and let $P \in S \setminus (B_F \cup B)$, where B_F is the (finite) set of atypical values of F.

Then the following are equivalent:

(1) $F_*: \pi_1(U_B) \to \pi_1^{\operatorname{orb}}(S \setminus B)$ is an isomorphism, (2) $F_*: \pi_1(U_{B \cup \{P\}}) \to \pi_1^{\operatorname{orb}}(S \setminus (B \cup \{P\}))$ is an isomorphism,

and in that case,

$$\pi_1(U_{B\cup\{P\}}) \cong \mathbb{Z} * \pi_1(U_B).$$

Moreover, the implication (2) \Rightarrow (1) also holds if $P \in B_F \setminus B$.

Remark 2.19. We will refer to $(1) \Rightarrow (2)$ as the Addition Lemma (which is true for generic fibers) and to $(2) \Rightarrow (1)$ as the Deletion Lemma (which is true for all fibers).

3. Threshold of atypical values

Let U be a smooth quasi-projective surface and let S be a smooth projective curve. Let $F: U \to S$ be an admissible map and let $B \subset S$ be a finite set of points. The goal of this section is to study how the presence of atypical values of F in Brelates to the induced morphism $F_*: \pi_1(U_B) \to \pi_1^{\mathrm{orb}}(S \setminus B)$ being an isomorphism.

To study this problem, we start by defining the set

$$\mathcal{T}_F := \left\{ T \subset B_F \left| \begin{array}{c} F_* : \pi_1(U_{\{P\} \cup (B_F \setminus T)}) \to \pi_1^{\mathrm{orb}}\left(S \setminus (\{P\} \cup (B_F \setminus T))\right) \\ \text{is an isomorphism} \end{array} \right\},$$

where $P \in S \setminus B_F$. The following result summarizes some of the properties of \mathcal{T}_F .

Lemma 3.1. Let U be a smooth quasi-projective surface and let S be a smooth projective curve. Let $F: U \to S$ be an admissible map, let $B_F \subset S$ be its set of atypical values, and let $P \in S \setminus B_F$. The following hold:

- (1) The set \mathcal{T}_F is independent of the choice of $P \in S \setminus B_F$.
- (2) If $T \in \mathcal{T}_F$ and T' is such that $T \subset T' \subset B_F$, then $T' \in \mathcal{T}_F$. (3) $\mathcal{T}_F \neq \emptyset$ if and only if $F_* : \pi_1(U_P) \to \pi_1^{\text{orb}}(S \setminus \{P\})$ is an isomorphism.

Proof. Let $P, Q \in S \setminus B_F$, let $T \subset B_F$ and suppose that

$$F_*: \pi_1(U_{\{P\}\cup (B_F\setminus T)}) \to \pi_1^{\operatorname{orb}}(S\setminus (\{P\}\cup (B_F\setminus T))))$$
 is an isomorphism.

By the Addition Lemma (Theorem 2.18), one can add the generic fiber of F at Q, that is, $F_* : \pi_1(U_{\{P,Q\}\cup(B_F\setminus T)}) \to \pi_1^{\operatorname{orb}}(S \setminus \{P,Q\} \cup (B_F \setminus T))$ is an isomorphism. Using now the Deletion Lemma, the fiber at P can be removed and $F_*: \pi_1(U_{\{Q\}\cup (B_F\setminus T)}) \to \pi_1^{\operatorname{orb}}(S\setminus \{Q\}\cup (B_F\setminus T))$ is an isomorphism. This proves part (1). Part (2) follows directly from the Deletion Lemma. Part (2) implies that $\mathcal{T}_F \neq \emptyset$ if and only if $B_F \in \mathcal{T}_F$, which implies part (3). \square

The previous result suggests the following definition.

Definition 3.2. Let U be a smooth quasi-projective surface, let S be a smooth projective curve. Let $F: U \to S$ be an admissible map and let $B_F \subset S$ be its set of atypical values. We say that a set $T \subset B_F$ is a threshold set of (atypical) values for F if it is a minimal set of \mathcal{T}_F (with respect to the inclusion of sets).

Remark 3.3 (Non-empty \mathcal{T}_F , projective case). Let $F = [f_q^p : f_p^q] : U = \mathbb{P}^2 \setminus \mathcal{B} \to \mathbb{P}^1$ be an admissible map satisfying Condition 2.3. If $P \in \mathbb{P}^1 \setminus B_F$, then $F_*: \pi_1(U_P) \to \pi_1^{\text{orb}}(S \setminus \{P\})$ is an isomorphism by [10, Theorem 1.2], hence, Lemma 3.1(3) implies $\mathcal{T}_F \neq \emptyset$.

Example 3.4 (Empty \mathcal{T}_F). Consider $\mathcal{B} = V(h_2) \cap V(h_4)$, where h_d is an irreducible homogeneous polynomial in $\mathbb{C}[x, y, z]$ of degree d. Define $U = \mathbb{P}^2 \setminus \mathcal{B}, S = \mathbb{P}^1$, and $F: U \to S$ is described as $F = [h_2^2: h_4]$. By Lemma 2.6, F is an admissible map, however $F_*: G_1 := \pi_1(U_P) \to \pi_1^{\text{orb}}(S \setminus \{P\}) =: G_2$ is not an isomorphism for any $P \notin B_F$, since the abelianizations $G_1/G_1' = \mathbb{Z}_4$, $G_2/G_2' = \mathbb{Z}_2$ are not isomorphic. Hence, Lemma 3.1(3) implies $\mathcal{T}_F = \emptyset$.

The purpose of \mathcal{T}_F is to characterize the property that $F_*: \pi_1(U_B) \to \pi_1^{\mathrm{orb}}(S \setminus B)$ defines an isomorphism, regardless of whether or not B contains a generic fiber, as the following result shows.

Proposition 3.5. Let U be a smooth quasi-projective surface, S a smooth projective curve and $F: U \to S$ be an admissible map. Consider $B \subset S$ a finite non-empty set. Then the following are equivalent:

- (1) $F_*: \pi_1(U_B) \to \pi_1^{\operatorname{orb}}(S \setminus B)$ is an isomorphism.
- (2) There exists $T \in \mathcal{T}_F$ such that $B \cap T = \emptyset$.

(3) $B_F \setminus B \in \mathcal{T}_F$.

Proof. Note that $(2) \Leftrightarrow (3)$ follows immediately from Lemma 3.1(2).

To show (1) \Leftrightarrow (3) note that the Addition-Deletion Lemma (Theorem 2.18) implies that

$$F_*: \pi_1(U_B) \to \pi_1^{\operatorname{orb}}(S \setminus B)$$

is an isomorphism if and only if

$$F_*: \pi_1(U_{(B\cap B_F)\cup\{P\}}) \to \pi_1^{\operatorname{orb}}(S \setminus ((B\cap B_F)\cup\{P\}))$$

is an isomorphism for any $P \in S \setminus (B_F \cup B)$, which is true iff $B_F \setminus B \in \mathcal{T}_F$. \Box

In this paper, we will compute the threshold sets of atypical values in cases which fall under the realm of Remark 3.3. In these cases, the threshold sets of atypical values contain more information about the fundamental groups of the complements of the fiber-type curves that arise from F than the definition of a threshold set of atypical values may suggest, as exemplified by the following result, which may be used in combination with Proposition 3.5.

Proposition 3.6. Let $F = [f : g] : U = \mathbb{P}^2 \setminus \mathcal{B} \to \mathbb{P}^1$ be an admissible map, where $f, g \in \mathbb{C}[x, y, z]$ are homogeneous polynomials of degree $d \ge 1$ with no non-constant common factors, and $\mathcal{B} = V(f) \cap V(g)$. Let $B \subset \mathbb{P}^1$ be such that $\#B \ge 2$. Then, the following are equivalent:

- (1) $\pi_1(U_B)$ is a free product of cyclic groups.
- (2) F satisfies (i)–(iii) of Theorem 2.14, so in particular,

$$F_*: \pi_1(U_B) = \pi_1\left(\mathbb{P}^2 \setminus \overline{F^{-1}(B)}\right) \to \pi_1^{\operatorname{orb}}(\mathbb{P}^1 \setminus B) \quad is \ an \ isomorphism.$$

Moreover, possibly after a change of coordinates in \mathbb{P}^1 , F also satisfies Condition 2.3.

Proof. The implication $(2) \Rightarrow (1)$ is trivial. Let us now see the implication $(1) \Rightarrow (2)$ if $\#B \ge 2$. Assume that $\pi_1(U_B)$ is a free product of cyclic groups. By Theorem 2.14 there exists a pencil $G = [g_u^v : g_v^u]$ satisfying Condition 2.3 and (i)–(iii) of Theorem 2.14. In particular, every irreducible component of $\overline{F^{-1}(B)} \subset \mathbb{P}^2$ is the union of the closure of finitely many irreducible fibers of G. Let $[\alpha_1 : \beta_1], [\alpha_2 : \beta_2]$ be distinct points in B. Then,

$$\beta_1 f - \alpha_1 g = g_u^{k_1} g_v^{k_2} \prod_{i=1}^{\kappa_3} (b_i g_u^v - a_i g_v^u)^{l_i},$$

$$\beta_2 f - \alpha_2 g = g_u^{k_4} g_v^{k_5} \prod_{j=1}^{k_6} (d_j g_u^v - c_j g_v^u)^{m_j},$$

for some $k_j \in \mathbb{Z}_{\geq 0}$, j = 1, ..., 6 with $k_1 + k_2 + k_3 \geq 1$, $k_4 + k_5 + k_6 \geq 1$, $l_i \in \mathbb{Z}_{\geq 1}$ for all $i = 1, ..., k_3$, $m_j \in \mathbb{Z}_{\geq 1}$ for all $j = 1, ..., k_6$, and $[a_i : b_i], [c_j : d_j]$ distinct points in $\mathbb{P}^1 \setminus \{[1:0], [0:1]\}$ for $i = 1, ..., k_3$ and $j = 1, ..., k_6$.

Note that, since G satisfies (iii) of Theorem 2.14 and Condition 2.3, $b_i g_u^v - a_i g_v^u$ and $d_j g_u^v - c_j g_v^u$ are irreducible polynomials for all $i = 1, \ldots, k_3$ and for all $j = 1, \ldots, k_6$. Looking at the degrees of the homogeneous polynomials in the previous two equations, we obtain that

$$d = k_1 u + k_2 v + uv \sum_{i=1}^{k_3} l_i = k_4 u + k_5 v + uv \sum_{j=1}^{k_6} m_j.$$

Since u and v are coprime, the last equality implies that $v \mid (k_1-k_4)$ and $u \mid (k_2-k_5)$. Note that, since different fibers of F do not have any irreducible components in common, at least one element of the sets $\{k_1, k_4\}$ and $\{k_2, k_5\}$ is 0, so $v \mid k_1, k_4$ and $v \mid k_2, k_5$. Hence, there exist homogeneous polynomials $h_1, h_2 \in \mathbb{C}[x, y]$ of the same degree d' such that $f = h_1(g_u^v, g_v^u)$ and $g = h_2(g_u^v, g_v^u)$. In other words, Ffactors as the composition of G and $[h_1 : h_2] : \mathbb{P}^1 \to \mathbb{P}^1$. Since F is admissible, then d' = 1, and hence $[h_1 : h_2]$ is a change of coordinates in \mathbb{P}^1 . In particular, F satisfies Condition 2.3 (after possibly a change of coordinates in \mathbb{P}^1), and also (i)–(iii) of Theorem 2.14, including the fact that

$$F_*: \pi_1(U_B) \to \pi_1^{\operatorname{orb}}(\mathbb{P}^1 \setminus B)$$

is an isomorphism.

In other words, Propositions 3.5 and 3.6 show that finding all the threshold sets of atypical values for an admissible map F amounts to completely characterizing when the fundamental group of $U_B = \mathbb{P}^2 \setminus \overline{F^{-1}(B)}$ is a free product of cyclic groups for all finite $B \subset \mathbb{P}^1$ such that $\#B \ge 2$.

Remark 3.7. If #B = 1 and $B \subset \mathbb{P}^2 \setminus B_F$, the result of Proposition 3.6 holds by [10, Theorem 1.2].

However, if #B = 1 and $B \subset B_F$, the result is no longer true:

Example 3.8. Let $f_2, f_3 \in \mathbb{C}[x, y, z]$ be irreducible homogeneous polynomials of degrees 2 and 3 respectively. By Lemma 2.6, $[f_2^3 : f_3^2]$ restricted to its maximal domain of definition is an admissible map. By [10, Theorem 1.2], $\pi_1(\mathbb{P}^2 \setminus V(f_2^3 + \lambda f_3^2)) \cong \mathbb{Z}_2 * \mathbb{Z}_3$ for $\lambda \in \mathbb{C}$ generic. In particular, $f_2^3 + \lambda f_3^2$ is an irreducible degree 6 homogeneous polynomial. Let $f_5 \in \mathbb{C}[x, y, z]$ be an irreducible degree 5 homogeneous polynomial. By Lemma 2.6, the restriction of the pencil $F = [(f_2^3 + \lambda f_3^2)^5 : f_5^6]$ to its maximal domain of definition is an admissible map. For this F, if $B = \{[0:1]\}$ we have that $\pi_1(U_B) = \mathbb{Z}_2 * \mathbb{Z}_3$ is a free product of cyclic groups, but

$$F_*: \pi_1(U_B) \to \pi_1^{\operatorname{orb}}(\mathbb{P}^1 \setminus B) = \mathbb{Z}_6$$

is not an isomorphism.

4. ZARISKI PAIRS OF FIBER-TYPE CURVES

Consider $F: \mathbb{P}^2 \to \mathbb{P}^1$ the rational morphism defined as $[\bar{f}(x, y, z) : z^6]$, where $f(x, y) = x^6 + y^6 + 6xy$ and $\bar{f}(x, y, z) = x^6 + y^6 + 6xyz^4$. This rational map is not well defined on the base points $\mathcal{B} = \{[x : y : 0] \in \mathbb{P}^2 \mid x^6 + y^6 = 0\}$. The generic fiber of the resulting admissible map $F: U \to \mathbb{P}^1$, where $U = \mathbb{P}^2 \setminus \mathcal{B}$, is a smooth sextic (with six points removed). The set of atypical values of F is $B_F = \{0, \infty, \pm 4, \pm 4i\} \subset \mathbb{P}^1 \equiv \mathbb{C} \cup \{\infty\}$. The fiber $F^{-1}(\infty)$ is the only multiple fiber and it has multiplicity six. The fiber $F^{-1}(0)$ is a singular sextic with one node. The remaining atypical fibers are irreducible sextics with six nodes. In particular, F is an admissible map satisfying Condition 2.3. For any finite set $B \subset \mathbb{P}^1$ we will denote $C_B := \overline{F^{-1}(B)}$. The purpose of this section is to study $\pi_1(U_B)$ for $U_B := \mathbb{P}^2 \setminus C_B$ for different choices of B. In particular, we will exhibit a Zariski pair of affine nodal curves and another Zariski pair of projective sextics of fiber-type.

4.1. A Zariski pair of affine nodal curves. Using the notation presented above, we consider the curve C_{B_1} (resp. C_{B_2}) for the set $B_1 = \{4, -4, \infty\}$ (resp. $B_2 = \{4, 4i, \infty\}$). Using the identification $\mathbb{P}^2 \setminus \{z = 0\} = \mathbb{P}^2 \setminus \overline{C_{\infty}} \equiv \mathbb{C}^2$ we can consider U_{B_j} as the complement in \mathbb{C}^2 of the affine curves D_j defined as a union of the two fibers over $B_j \setminus \{\infty\}$ of the polynomial map $(x, y) \mapsto f(x, y) = x^6 + y^6 + 6xy$, for j = 1, 2. In this context one has the following.

Theorem 4.1. The curves (D_1, D_2) described above form a Zariski pair of affine nodal curves of degree 12. Moreover, $\pi_1(U_{B_2}) = \mathbb{F}_2$ whereas

(2)
$$\pi_1(U_{B_1}) = \langle x, y, u, v : [u, x] = [v, y] = 1, u = [v^{-1}, x], v = [u, y] \rangle$$

fits in the following free-by-free extension

(3)
$$1 \to \mathbb{Z}u * \mathbb{Z}v \to \pi_1(U_{B_1}) \to \mathbb{Z}x * \mathbb{Z}y \to \mathbb{I}$$

and is not free.

Proof. Since the coefficients of the equations of the curves can be given in a number field such as $\mathbb{Q}(i)$, one can obtain presentations of the fundamental groups accurately using the package *Sirocco* in the computer software SageMath¹ [17].

In order to give a comprehensive proof of this result we will perform a series of transformations to D_j in order to simplify it and be able to calculate its braid monodromy and apply Zariski-Van Kampen's method. Consider the double cover $\mathbb{C}^2 \xrightarrow{\sigma} \mathbb{C}^2$ of symmetries defined as $\sigma(x, y) = (xy, x + y) = (u, v)$. The discriminant of σ is given by $\Delta_{\sigma}(u, v) = v^2 - 4u$ and σ_{\parallel} defines an unramified double cover outside $\sigma^{-1}(\Delta_{\sigma}) = x - y = 0$. Note that $f(x, y) = g(\sigma(x, y))$, where g(u, v) = $v^6 - 6v^4u + 9v^2u^2 - 2u^3 + 6u$. Consider $\mathbb{C}^2 \xrightarrow{\sigma} \mathbb{C}^2$ the cover $\delta(u, v) = (u, v^2)$ associated with the reflection over $\ell_1 = \{v = 0\}$, which is an unramified double cover outside of the preimage of ℓ_1 . One obtains that $f(x, y) = (h \circ \delta \circ \sigma)(x, y)$, where $h(u, v) = v^3 - 6v^2u + 9vu^2 - 2u^3 + 6u$. In order to compute $G_j = \pi_1(U_{B_j})$, we will first calculate $\tilde{G}_j = \pi_1(\mathbb{C}^2 \setminus (H_{B_j \setminus \{\infty\}} \cup \ell_1 \cup \ell_2))$, where $H_B = h^{-1}(B)$ and $\ell_2 = \{v = 4u\}$. Once \tilde{G}_j is computed one can obtain G_j after a sequence of index 2 subgroups associated with the double covers and quotients by meridians in order to eliminate the ramification locus.

Note that the polynomial map h produces an elliptic fibration whose generic fiber is a smooth cubic. This fibration is ramified at $B_h = \{\pm 4, \pm 4i\}$ whose fibers are of type I_1 (nodal cubics). In addition, the line ℓ_1 (resp. ℓ_2) is tangent to the fibers H_B , for $B = \{0, \pm 4\}$ (resp. $B = \{\pm 4i\}$). This implies that $f^{-1}(\lambda), \lambda \in B_h$ contains 4 nodes coming from the preimages of the nodes at H_λ plus two nodes coming from the tangency at $\ell_1 \cap H_\lambda$ (resp. $\ell_2 \cap H_\lambda$). Finally, the fiber $f^{-1}(0)$ contains only one node since the tangency $\ell_1 \cap H_\lambda$ produces a node after the preimage of δ , which is on the ramification locus of σ and hence produces only one node after the preimage by σ .



FIGURE 1. Real curve $H_{B_1 \setminus \{\infty\}} \cup \ell_1 \cup \ell_2$

¹A program to do so is available in the public repository

https://riemann.unizar.es/~jicogo/software/Zariski-Pair.sage using Sage with the beta package from https://github.com/enriqueartal/sage/tree/puiseux

Using the real picture (see Figure 1) one can obtain a braid monodromy for $H_{B_1 \setminus \{\infty\}} \cup \ell_1 \cup \ell_2$ and use Zariski-Van Kampen's method to obtain a presentation of the fundamental group $\tilde{G}_1 = \pi_1(\mathbb{C}^2 \setminus (H_{B_1 \setminus \{\infty\}} \cup \ell_1 \cup \ell_2))$ as

(4)
$$\tilde{G}_{1} = \left\langle \begin{array}{c} [\gamma_{1}, \gamma_{2}] = [\gamma_{2}, \gamma_{\pm}] = [\gamma_{1}^{2}, \gamma_{\pm}] = 1 \\ (\gamma_{1}, \gamma_{2}, \gamma_{+}, \gamma_{-}: (\gamma_{1}\gamma_{+})^{2} = (\gamma_{+}\gamma_{1})^{2}, (\gamma_{1}\gamma_{-})^{2} = (\gamma_{-}\gamma_{1})^{2} \\ [\gamma_{-}, \gamma_{+}\gamma_{1}\gamma_{+}^{-1}] = [\gamma_{+}, \gamma_{-}^{-1}\gamma_{1}\gamma_{-}] = 1 \end{array} \right\rangle,$$

where γ_j is a meridian of the line ℓ_j (i = 1, 2) and γ_{\pm} is a meridian of $H_{\pm 4}$. The fundamental group of the double cover δ is obtained as the kernel \tilde{K}_1 of the morphism $\tilde{G}_1 \to \mathbb{Z}_2$ defined as $\gamma_1 \mapsto 1$, $\gamma_2 \mapsto 0$, $\gamma_{\pm} \mapsto 0$.

The group K_1 is generated by $\gamma_1^2, \gamma_2, \tilde{\gamma}_2 := \gamma_1 \gamma_2 \gamma_1, \gamma_{\pm}, \tilde{\gamma}_{\pm} := \gamma_1 \gamma_{\pm} \gamma_1$ and has relations

$$\begin{split} \tilde{\gamma}_2 &= \gamma_2 \gamma_1^2 = \gamma_1^2 \gamma_2, \qquad [\gamma_\pm, \gamma_2] = [\tilde{\gamma}_\pm, \gamma_2] = 1, \\ [\gamma_\pm, \gamma_1^2] &= [\tilde{\gamma}_\pm, \gamma_1^2] = 1, \qquad [\gamma_\pm, \tilde{\gamma}_\pm] = 1, \\ \gamma_1^2 \tilde{\gamma}_+ \gamma_- \gamma_+ &= \gamma_+ \tilde{\gamma}_- \tilde{\gamma}_+ \qquad \gamma_1^2 \gamma_- \gamma_+ \tilde{\gamma}_- = \tilde{\gamma}_- \tilde{\gamma}_+ \gamma_-. \end{split}$$

Since γ_1^2 is the meridian of the conic $\delta^*(\ell_1)$, the quotient

$$\tilde{K}_1/\langle \gamma_1^2 \rangle = \mathbb{Z}\gamma_2 \times \left\langle \gamma_{\pm}, \tilde{\gamma}_{\pm} : \begin{array}{c} [\gamma_{\pm}, \tilde{\gamma}_{\pm}] = 1, \\ \tilde{\gamma}_{+}\gamma_{-}\gamma_{+} = \gamma_{+}\tilde{\gamma}_{-}\tilde{\gamma}_{+}, \gamma_{-}\gamma_{+}\tilde{\gamma}_{-} = \tilde{\gamma}_{-}\tilde{\gamma}_{+}\gamma_{-} \end{array} \right\rangle.$$

is the fundamental group of the complement Z_1 of $g^{-1}(B_1 \setminus \{\infty\}) \cup \ell_2$. Now, the fundamental group K_1 of $\sigma^{-1}(Z_1)$ is isomorphic to the kernel of the homomorphism $\tilde{K}_1/\langle \gamma_1^2 \rangle \to \mathbb{Z}_2$ defined by $\gamma_2 \mapsto 1$, $\gamma_{\pm} \mapsto 0$, $\tilde{\gamma}_{\pm} \mapsto 0$, that is,

$$K_1 = \mathbb{Z}\gamma_2^2 \times \left\langle \gamma_{\pm}, \tilde{\gamma}_{\pm} : \begin{array}{c} [\gamma_{\pm}, \tilde{\gamma}_{\pm}] = 1, \\ \tilde{\gamma}_+ \gamma_- \gamma_+ = \gamma_+ \tilde{\gamma}_- \tilde{\gamma}_+, \gamma_- \gamma_+ \tilde{\gamma}_- = \tilde{\gamma}_- \tilde{\gamma}_+ \gamma_- \end{array} \right\rangle.$$

After factoring out the normal subgroup generated by γ_2^2 , the meridian of $\sigma^*(\delta^*(\ell_2))$, one obtains the desired presentation in (2) where $x := \gamma_+, y := \gamma_-, u := \tilde{\gamma}_+ \gamma_+^{-1}$, and $v := \tilde{\gamma}_- \gamma_-^{-1}$.

Now consider the case $B_2 = \{4, 4i, \infty\}$. The strategy to follow is similar to that of the previous case. However, note that the curve $H_{B_2 \setminus \{\infty\}}$ is not real anymore. A way around this is to consider $h(ui, vi) - 4i = -i(\tilde{h}(u, v) - 4)$, where $\tilde{h}(u, v) = v^3 - 6v^2u + 9vu^2 - 2u^3 - 6u$. This allows us to consider the braid monodromy along real loops (for H_4, ℓ_1, ℓ_2) and purely imaginary loops for H_{4i} as a braid monodromy over a real path for $\tilde{H}_4 = \tilde{h}^{-1}(4)$. This produces the following presentation,

(5)
$$\tilde{G}_2 = \pi_1(\mathbb{C}^2 \setminus (H_{B_2 \setminus \{\infty\}} \cup \ell_1 \cup \ell_2)) = \mathbb{Z}\gamma_1 \times \mathbb{Z}\gamma_2 \times \langle \gamma_+, \gamma_- \rangle \cong \mathbb{Z}^2 \times \mathbb{F}_2,$$

where γ_j is a meridian of the line ℓ_j (j = 1, 2) and γ_+ (resp. γ_-) is a meridian of H_4 (resp. H_{4i}). Following the same strategy as above $\tilde{K}_2 \cong \mathbb{Z}\gamma_1^2 \times \mathbb{Z}\gamma_2 \times \mathbb{F}_2$, so $\tilde{K}_2/\langle \gamma_1^2 \rangle \cong \mathbb{Z}\gamma_2 \times \mathbb{F}_2$. Consequently $K_2 \cong \mathbb{Z}\gamma_2 \times \mathbb{F}_2$, and $G_2 \cong \mathbb{F}_2$.

To prove the short exact sequence (3), note that the kernel of the projection map $x \mapsto x, y \mapsto y, u \mapsto 1, v \mapsto 1$ is generated by $u_w := wuw^{-1}$ and $v_w := wvw^{-1}$ for any $w \in \mathbb{Z}x * \mathbb{Z}y$. From the presentation (2) of G_1 , one obtains the following recursive set of relations

$$\begin{array}{ll} u_{wx} = u_{wx^{-1}} = u_w, & v_{wy} = v_{wy^{-1}} = v_w, \\ u_{wy} = v_w^{-1} u_w, & v_{wx} = v_w u_w, \\ u_{wy^{-1}} = v_w u_w, & v_{wx^{-1}} = v_w u_w^{-1}. \end{array}$$

This allows one to reduce the set of generators of the kernel to $u_1 = u$ and $v_1 = v$ and no further relations. Since the abelianization of G_1 has rank 2 and finitely generated free groups are Hopfian, the short exact sequence (3) shows that G_1 cannot be free. Remark 4.2. Note that both spaces U_{B_1} and U_{B_2} are associated with affine pencils without multiple fibers and hence the epimorphisms of their fundamental groups onto the free group \mathbb{F}_2 must have finitely generated kernels as shown by Catanese in [7, Thm. 5.4].

Let $\omega := e^{\frac{1}{12}\pi i}$. Note that the isomorphism

(6)
$$\begin{array}{ccc} \underline{\omega}: & \mathbb{P}^2 & \to & \mathbb{P}^2 \\ & & [x:y:z] & \mapsto & [\omega x: \omega^5 y:z], \end{array}$$

takes $\overline{F^{-1}([a:b])}$ to $\overline{F^{-1}([ai:b])}$ for all $[a:b] \in \mathbb{P}^1$. Using this transformation, Theorem 4.1 immediately implies the following.

Corollary 4.3. $\pi_1(U_{\{4,-4,\infty\}}) \cong \pi_1(U_{\{4i,-4i,\infty\}})$ is the non-free group with presentation (2).

We end this section by finding more affine fiber-type curves $D \subset \mathbb{C}^2$ whose irreducible components are atypical fibers of f and such $\pi_1(\mathbb{C}^2 \setminus D)$ is also free.

Proposition 4.4. Let
$$\underline{\omega} : \mathbb{P}^2 \to \mathbb{P}^2$$
 be the isomorphism from equation (6). Then,
 $\mathbb{F}_3 \cong \pi_1(U_{\{0,4,4i,\infty\}}) \stackrel{\underline{\omega}}{\cong} \pi_1(U_{\{0,-4,4i,\infty\}}) \stackrel{\underline{\omega}}{\cong} \pi_1(U_{\{0,-4,-4i,\infty\}}) \stackrel{\underline{\omega}}{\cong} \pi_1(U_{\{0,4,-4i,\infty\}}).$

Proof. Similar, but more involved calculations as those detailed in the proof of Theorem 4.1, show that $G_0 = \pi_1(U_{B_0}) = \mathbb{F}_3$ for $B_0 = \{0, 4, 4i, \infty\}$. In fact, using the notation from the proof of Theorem 4.1, we have that

$$\tilde{G}_0 = \pi_1(\mathbb{C}^2 \setminus (H_{B_0 \setminus \{\infty\}} \cup \ell_1 \cup \ell_2)) = \mathbb{Z}\gamma_1 \times \mathbb{Z}\gamma_2 \times \langle \gamma_0, \gamma_+, \gamma_- \rangle \cong \mathbb{Z}^2 \times \mathbb{F}_3,$$

where γ_1 (resp. γ_2) are meridians of the lines ℓ_1 (resp. ℓ_2), γ_0 is a meridian around H_0 and γ_+ (resp. γ_-) is a meridian of H_4 (resp. H_{4i}).

4.2. Twisted Alexander polynomials. The purpose of this section is to show that twisted Alexander polynomials can also distinguish the fundamental groups $G_1 = \pi_1(U_{B_1})$ and $G_2 = \pi_1(U_{B_2})$. To our knowledge, this is the first example in the literature of a Zariski pair whose groups have the same Alexander invariant but different twisted Alexander polynomials. Moreover, Lemma 4.6 shows that the position of the nodes affects the fundamental groups and twisted Alexander polynomials are sensitive to this special position. Classical Alexander polynomials and characteristic varieties are known not to be sensitive to the position of nodes (see for instance [14] and [15, 2.5]). We refer the reader to [9, Section 2] for a quick introduction on twisted Alexander polynomials and how to compute them.

We first compute the Alexander invariant of the complements U_{B_1} and U_{B_2} .

Proposition 4.5. The Alexander invariant of both $\mathbb{C}^2 \setminus D_1$ and $\mathbb{C}^2 \setminus D_2$ is $\mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$.

Proof. Let us denote by M_j the Alexander invariant of $\mathbb{C}^2 \setminus D_j$, that is, $M_j := G'_j/G''_j$ as an *R*-module where $R = \mathbb{C}[G_j/G'_j] = \mathbb{C}[t_1^{\pm 1}, t_2^{\pm 1}]$. It is immediate that $M_2 = [x, y]R$. As for M_1 , one can use the presentation (2), where the abelian classes of x and y generate G_1/G'_1 and $u, v \in G'_1$. Using [5, §2.5], this implies that M_1 is generated by [x, y], u, and v and whose relations are given by rewriting the relations in (3) as elements of the *R*-module M_1 . Since [x, q] = 0 in M_1 as long as $q \in G'_1$, the first two relations in (3) become trivial and the last ones become u = v = 0. This shows $M_1 \cong M_2 \cong R$.

By Proposition 4.5, the characteristic varieties of U_{B_1} and U_{B_2} are both $(\mathbb{C}^*)^2$. Analogously, the Alexander polynomials associated with any homomorphism ε : $G_j \to \mathbb{Z}^2$ are trivial. The following result shows that the nodes of the curve D_1 are in special position, unlike those in D_2 . **Lemma 4.6.** The 12 nodes in the curve D_1 are contained in three lines containing 4 nodes each. There are no three nodes aligned in D_2 .

Proof. We use the notation introduced in the proof of Theorem 4.1. It is easy to check that the node of $H_{\pm 4}$ (resp. $H_{\pm 4i}$) has coordinates $P_{\pm 4} = (u, v) = (\pm 1, \pm 3)$ (resp. $P_{\pm 4i} = (u, v) = (\pm \frac{i}{4}, \pm \frac{i}{4})$). Also, $H_{\pm 4}$ (resp. $H_{\pm 4i}$) is tangent to the line $\ell_1 = \{v = 0\}$ (resp. $\ell_2 = \{v = 4u\}$) at the point $Q_{\pm 4} = (\pm 1, 0)$ (resp. $Q_{\pm 4i} = (\pm i, \pm 4i)$). The line $\ell_+ = \{v = 3u\}$ joining P_4 and P_{-4} is transformed through $\delta \circ \sigma$ into the singular conic $Q_+ = \{3x^2 + 5xy + 3y^2 = 0\}$ containing the 8 nodes which are in the preimage of $P_{\pm 4}$. Each of the lines in the conic Q_+ contains 4 of those nodes. In addition, the line ℓ_1 is transformed through σ into the interval the remaining 4 nodes which are the preimage of the tangencies $Q_{\pm 4}$.

On the other hand, let $\ell_{-} = \{(4-i)v - (12-i)u + 2i = 0\}$ be the line joining P_4 and P_{4i} . The preimage of ℓ_{-} through $\delta \circ \sigma$ is a smooth conic Q_{-} containing the 8 nodes preimage of P_4 and P_{4i} . This proves that no three of those preimages can be aligned. An analogous argument rules out the existence of a line joining three points in the preimage of any two of the points of $S = \{P_4, P_{4i}, Q_4, Q_{4i}\}$.

Assume that there is a line joining a preimage of P_4 , P_{4i} , and Q_4 , and consider the quartic consisting on the four lines obtained from it by applying the deck transformations of the global 4:1 covering induced by the restriction of $\delta \circ \sigma$. This invariant quartic is the preimage by $\delta \circ \sigma$ is a conic which is tangent to ℓ_1 at Q_4 , since the preimages of Q_4 are double points in the quartic. However, this complete reducible quartic has four extra double points. Since they cannot be on the ramification locus, they have to be on a fiber. Hence, the conic has to be singular. This contradicts the condition about the tangency with ℓ_1 . An analogous argument rules out the existence of a line joining any three points in the preimage of three of the points of S.

This is the announced result showing that twisted Alexander polynomials can be sensitive to the position of nodes.

Proposition 4.7. The twisted Alexander polynomial for unitary representations in U(2) distinguishes $\pi_1(U_{B_1})$ and $\pi_1(U_{B_2})$.

Proof. By definition, note that if $G = \mathbb{F}_r$ is a free group, then $\Delta_{\rho,\varepsilon}(t) = 1$ for any representation ρ of G and any homomorphism $\varepsilon : G/G' \to \mathbb{Z}$. In particular, $\Delta_{\rho_2,\varepsilon_2}(t) = 1$ for any twisted Alexander polynomial of the free group G_2 . Hence, it is enough to find a representation, say $\rho_1 : G_1 \to U(2)$ and a homomorphism $\varepsilon_1 : G_1/G'_1 \to \mathbb{Z}$ such that $\Delta_{\rho_1,\varepsilon_1}(t) \neq 1$. Consider $\varepsilon_1 : G_1/G'_1 \to \mathbb{Z}$ defined as $\varepsilon_1(x) = \varepsilon_1(y) = 1$, $\varepsilon_1(u) = \varepsilon_1(v) = 0$ and the irreducible representation $\rho_1 : G_1 \to U(2)$ defined as:

$$\rho_1(x) := \begin{pmatrix} \xi & 0 \\ 0 & \bar{\xi} \end{pmatrix}, \qquad \rho_1(y) := \frac{\sqrt{2}}{2} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}, \\
\rho_1(u) := i \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \qquad \rho_1(v) := \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix},$$

where $\xi = \frac{\sqrt{2}}{2}(1+i)$ is a primitive 8th root of unity. One can check that A_1 below is equivalent to the Fox matrix for the relations in (2) with respect to ρ_1 and ε_1 .

$$A_{1} = \begin{pmatrix} * & 0 & 0 & (t - \bar{\xi}) & 0 \\ 0 & 0 & 0 & (t - \xi) \\ \hline * & (t - \sqrt{2}) & it & \sqrt{2} \alpha(t) & 0 \\ \hline -it & (t - \sqrt{2}) & 0 & \sqrt{2} \alpha(t) \\ \hline * & (t - \xi) & -(t - \xi) & \xi(t - i\sqrt{2})(t - \bar{\xi}) & \xi(t - \sqrt{2})(t - \xi) \\ \hline & (t - \bar{\xi}) & (t - \bar{\xi}) & \bar{\xi}(t - \sqrt{2})(t - \bar{\xi}) & \bar{\xi}(t + \sqrt{2})(t - \xi) \end{pmatrix}$$

We omit the first two columns since they will not be necessary for the calculation of the twisted Alexander polynomial. Following Wada [19, Lemma 1] one needs to compute the Fitting ideal generated by the (4×4) minors of the submatrix of A_1 obtained after deleting the first two columns. This ideal is generated by $\Delta_1(t) = \alpha(t)^2$, where $\alpha(t) = t^2 - \sqrt{2}t + 1 = (t - \xi)(t - \overline{\xi})$. One also needs to compute $\Delta_0(t) = \det(\rho_1(x)t - I_2) = \alpha(t)$. Hence, the twisted Alexander polynomial $\Delta_{\rho_1,\varepsilon_1}(t)$ can be obtained as

$$\Delta_{\rho_1,\varepsilon_1}(t) = \frac{\Delta_1(t)}{\Delta_0(t)} = \alpha(t) = (t-\xi)(t-\bar{\xi}).$$

Remark 4.8. The roots of $\Delta_{\rho_1,\varepsilon_1}(t)$ are in fact eigenvalues of the unitary representation ρ_1 (see [16, Thm. 5.4]). It is also worth noting that the representation ρ_1 factors though one of the degree-two representations of the *binary octahedral group* BO₄₈. The group BO₄₈ is the finite group of symmetries of the octahedron and it is a non-abelian solvable subgroup of order 48 of U(2).

4.3. A projective Zariski pair. In this section we continue using the notation presented at the beginning of §4. One can obtain a Zariski pair of projective curves in \mathbb{P}^2 as follows.

Theorem 4.9. The projective curves $(C_{\{4,-4\}}, C_{\{4,4i\}})$ form a Zariski pair of projective curves of degree 12 whose complements in \mathbb{P}^2 have non-isomorphic fundamental groups. More concretely, $\pi_1(U_{\{4,4i\}}) = \mathbb{Z} * \mathbb{Z}_6$, whereas

(7)
$$\pi_1(U_{\{4,-4\}}) \cong \pi_1(U_{\{4,-4,\infty\}}) / \langle xy^2 x^2 y u x (vy)^2 (ux)^2 vy \rangle,$$

where $\langle \gamma \rangle$ denotes the normal subgroup of $\pi_1(U_{\{4,-4,\infty\}})$ generated by γ .

Proof. Let us denote $B'_1 = \{4, -4\}$ and $B'_2 = \{4, 4i\}$. The proof of $\pi_1(U_{B'_2}) = \mathbb{Z} * \mathbb{Z}_6$ is a consequence of Theorem 4.1 and the Deletion Lemma in Theorem 2.18. The presentation of $\pi_1(U_{B'_1})$ can be obtained from the presentation of $\pi_1(U_{B_1})$ by quotienting by the normal subgroup generated by the meridian $\tilde{\gamma}_{\infty}$ around the line at infinity ℓ_{∞} . This meridian can be obtained by lifting the square γ^2_{∞} of the meridian around the line at infinity by the coverings σ and δ from the proof of Theorem 4.1. The real picture given in Figure 1 shows that $\gamma^{-1}_{\infty} = \gamma_2 \gamma_+ \gamma_-^2 \gamma_+^2 \gamma_- \gamma_1$. Following the transformations given by the covering, one obtains

$$\tilde{\gamma}_{\infty}^{-1} = \gamma_+ \gamma_-^2 \gamma_+^2 \gamma_- \tilde{\gamma}_+ \tilde{\gamma}_-^2 \tilde{\gamma}_+^2 \tilde{\gamma}_- = xy^2 x^2 y u x (vy)^2 (ux)^2 vy.$$

The fact that the group (7) is not isomorphic to $\mathbb{Z}*\mathbb{Z}_6$ can be obtained by calculating the twisted Alexander polynomial of $\bar{G}_1 := \pi_1(U_{B'_1})$ associated with the morphism $\bar{\varepsilon}: \bar{G}_1/\bar{G}'_1 \to \mathbb{Z}$ defined as $\bar{\varepsilon}(x) = 1$, $\bar{\varepsilon}(y) = -1$, $\bar{\varepsilon}(u) = \bar{\varepsilon}(v) = 0$ resulting from the composition of the abelianization morphism and its projection onto the free part. Note that $\Delta_{\bar{\rho}_2,\bar{\varepsilon}}(\bar{G}_2)$ is trivial for any representation $\bar{\rho}_2$ and any epimorphism $\bar{\varepsilon}: \bar{G}_2/\bar{G}'_2 \cong \mathbb{Z} \times \mathbb{Z}_6 \to \mathbb{Z}$. However, consider the character $\bar{\rho}_1: \bar{G}_1 \to \mathbb{C}^*$ defined as $\bar{\rho}_1(x) = -\frac{\sqrt{2}}{2}i, \ \bar{\rho}_1(y) = \sqrt{2}i, \ \bar{\rho}_1(u) = \bar{\rho}_1(v) = 1$. Since $\Delta_{\bar{\rho}_1,\bar{\varepsilon}}(\bar{G}_1) = t + \frac{\sqrt{2}}{2}i$ is not trivial, then $\bar{G}_1 \ncong \mathbb{Z} * \mathbb{Z}_6$.

Remark 4.10. As seen in the proof of Lemma 4.6, the three lines containing the 12 nodes in $C_{\{4,-4\}}$ are the preimages of $\ell_1 = \{v = 0\}$ and $\ell_+ = \{v = 3u\}$. The latter passes through the nodes $(\pm 1, \pm 3)$ and it intersects the two cubics $H_{\pm 4}$ transversally at two other smooth points, namely $(\mp 2, \mp 6)$, and hence the preimage of ℓ_+ by $\delta \circ \sigma$ (which splits in two lines $6x + (5 + \sqrt{11}i)y = 0$ and $6x + (5 - \sqrt{11}i)y = 0$) contains 8 nodes (4 each line) and it intersects $C_{\{4,-4\}}$ transversally at the remaining 8 points of intersection (again, 4 each line). On the other hand, ℓ_1 is tangent to the two

cubics at $(\pm 1, 0)$ and it intersects them transversally at two other points $(\mp 2, 0)$. This implies that its preimage (a ramification line x + y = 0) passes through 4 nodes (the preimage of a simple tangency to a ramification locus of order two becomes two nodes) and through 4 additional transversal intersections (the preimages of $(\pm 2, 0)$). In particular, the three lines passing through the 12 nodes are concurrent and the remaining intersection points of any of these three lines with the curve $C_{\{4,-4\}}$ are transversal intersections (4 transversal intersections per line). This stresses the fact that the only difference between the position of the singularities of the curves $(C_{\{4,-4\}}, C_{\{4,4i\}})$ is the contribution of the special position of the nodes, and the same goes for $(C_{B_1} = C_{\{4,-4,\infty\}}, C_{B_2} = C_{\{4,4i,\infty\}})$ (that is, the projective curves whose complements in \mathbb{P}^2 are $\mathbb{C}^2 \setminus D_1$ and $\mathbb{C}^2 \setminus D_2$ respectively).

4.4. Threshold sets of atypical values for F.

The following result shows that all the threshold sets of atypical values for F have two elements from $\{4, -4, 4i, -4i\}$, but not all choices are allowed.

Proposition 4.11. There are four threshold sets of atypical values for F, namely $T \in \mathcal{T}_F$ is minimal if and only if $T = \{\varepsilon_1 4, \varepsilon_2 4i\}$, where $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$.

Proof. By Theorem 4.9, $F_*: \pi_1(U_{\{4,-4\}}) \to \pi_1^{\operatorname{orb}}(\mathbb{P}^1 \setminus \{4,-4\}) \cong \mathbb{Z} * \mathbb{Z}_6$ is not an isomorphism. By Propositions 3.5 and 3.6 one has $B_F \setminus \{4,-4\} = \{0,4i,-4i,\infty\} \notin \mathcal{T}_F$. Using the transformation $\underline{\omega}$ from (6) we also obtain that $\{0,4,-4,\infty\} \notin \mathcal{T}_F$. Hence, by Lemma 3.1(2), no subset of $\{0,4,-4,\infty\}$ or of $\{0,4i,-4i,\infty\}$ can be in \mathcal{T}_F . In particular, no singleton can be a threshold set of atypical values for F. Also, $T \in \mathcal{T}_F$ implies $T \supset \{\varepsilon_1 4, \varepsilon_2 4i\}$ for some choice $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$. The converse follows from Propositions 3.5, 3.6 and 4.4

Finally, we explicitly state the consequence of the computation of threshold sets.

Theorem 4.12. Let $F: U \to \mathbb{P}^1$ be given by $F = [x^6 + y^6 + 6xyz^4 : z^6]$, where $U = \mathbb{P}^2 \setminus \{ [x:y:0] \mid x^6 + y^6 = 0 \}$. Let $B_F \subset \mathbb{P}^1$ be the set of atypical values of F. For every finite set $A \subset \mathbb{P}^1$, let $C_A = \overline{F^{-1}(A)} \subset \mathbb{P}^2$.

Then, the following is a complete list of all Zariski pairs of curves $(C_A, C_{A'})$ such that the fundamental group of the complement of C_A in \mathbb{P}^2 is a free product of cyclic groups:

$\pi_1(\mathbb{P}^2 \setminus C_A)$	A	A'
$\mathbb{F}_{r+1} * \mathbb{Z}_6$	$\{\varepsilon_1 4, \varepsilon_2 4i\} \cup B$	$\{4\alpha, -4\alpha\} \cup B'$
\mathbb{F}_{r+2}	$\{\varepsilon_1 4, \varepsilon_2 4i, \infty\} \cup B$	$\{4\alpha, -4\alpha, \infty\} \cup B'$
$\mathbb{F}_{r+2} * \mathbb{Z}_6$	$\{0,\varepsilon_14,\varepsilon_24i\}\cup B$	$\{0, 4\alpha, -4\alpha\} \cup B'$
\mathbb{F}_{r+3}	$\{0,\varepsilon_14,\varepsilon_24i,\infty\}\cup B$	$\{0, 4\alpha, -4\alpha, \infty\} \cup B'$

where $B, B' \subset \mathbb{P}^1 \setminus B_F$ are finite, r := #B = #B', $\varepsilon_1, \varepsilon_2 \in \{\pm 1\}$, and $\alpha \in \{1, i\}$.

Proof. Recall from the beginning of the section, that $B_F = \{0, \infty, \pm 4, \pm 4i\}$. Let's write $A = A_1 \cup B$, where $A_1 = A \cap B_F$ and $B = A \setminus B_F$ (and analogously we define A'_1 and B' from A'). Since the closure of every fiber of F is an irreducible curve in \mathbb{P}^2 , Zariski pairs of the form $(C_A, C_{A'})$ must satisfy that $\#A_1 = \#A'_1, \#B = \#B'$, and the combinatorial type of the fibers at A_1 are in one-to-one correspondence with those at A'_1 . Also note that, if $A_1 = A'_1$ one can use an isotopy of the base that fixes B_F and takes B to B' to show that (\mathbb{P}^2, C_A) is isotopic to $(\mathbb{P}^2, C_{A'})$ and hence $(\mathbb{P}^2, C_A) \cong (\mathbb{P}^2, C_{A'})$.

Let $(C_A, C_{A'})$ be a Zariski pair, such that $\pi_1(\mathbb{P}^2 \setminus C_A)$ is a free product of cyclic groups. We will prove that A (resp. A') must be as in the middle (right-most) column in the table above for one row.

Assume $\#(A_1 \cap \{\pm 4, \pm 4i\}) = \#(A'_1 \cap \{\pm 4, \pm 4i\}) \leq 1$. Since the transformation $\underline{\omega} : \mathbb{P}^2 \to \mathbb{P}^2$ from equation (6) acts transitively on the fibers of F at $\{\pm 4, \pm 4i\}$ and fixes the fibers at $\{0, \infty\}$, up to a power of $\underline{\omega}$ one can assume $A_1 = A'_1$. By the first paragraph, $(\mathbb{P}^2, C_A) \cong (\mathbb{P}^2, C_{A'})$, so $(C_A, C_{A'})$ cannot be a Zariski pair.

Therefore, $\#(A_1 \cap \{\pm 4, \pm 4i\}) = \#(A'_1 \cap \{\pm 4, \pm 4i\}) \geq 2$. This means in particular $\#A_1 = \#A'_1 \geq 2$ and hence Propositions 3.5 and 3.6 imply $\pi_1(\mathbb{P}^2 \setminus C_A)$ is a free product of cyclic groups if and only if $B_F \setminus A \in \mathcal{T}_F$. By Proposition 4.11, the only possible A satisfying the hypothesis are shown on the middle column of the table above. Note that A' could be either a set from the middle column on the table above (shown as A) or a set from the right-most column. Since $\#A'_1 \geq 2$, Proposition 3.6 implies that $\pi_1(\mathbb{P}^2 \setminus C_{A'_1})$ is a free product of cyclic groups in the former case, whereas it is not in the latter. This shows that all cases on the table are Zariski pairs that can be distinguished by the groups of their complements.

Finally, if $\pi_1(\mathbb{P}^2 \setminus C_{A'})$ and $\pi_1(\mathbb{P}^2 \setminus C_A)$ are isomorphic to a free product of cyclic groups (that is, both A_1 and A'_1 are shown in the middle column on the table above), then one can use $\underline{\omega}$ and assume $A_1 = A'_1$. Again, as above, $(\mathbb{P}^2, C_A) \cong (\mathbb{P}^2, C_{A'})$. \Box

References

- D. Arapura, Geometry of cohomology support loci for local systems. I, J. Algebraic Geom. 6 (1997), no. 3, 563–597. MR 1487227
- D. Arapura, Toward the structure of fibered fundamental groups of projective varieties, J. Éc. polytech. Math. 4 (2017), 595–611. MR 3665609
- 3. E. Artal, Sur les couples de Zariski, J. Algebraic Geom. 3 (1994), no. 2, 223–247. MR 1257321
- E. Artal and J.I. Cogolludo, On the connection between fundamental groups and pencils with multiple fibers, J. Singul. 2 (2010), 1–18. MR 2763015
- E. Artal, J.I. Cogolludo, and H.o. Tokunaga, A survey on Zariski pairs, Algebraic geometry in East Asia—Hanoi 2005, Adv. Stud. Pure Math., vol. 50, Math. Soc. Japan, Tokyo, 2008, pp. 1–100. MR MR2409555
- S. Bannai and T. Shirane, Nodal curves with a contact-conic and Zariski pairs, Adv. Geom. 19 (2019), no. 4, 555–572. MR 4015190
- F. Catanese, Fibred Kähler and quasi-projective groups, no. suppl., 2003, Special issue dedicated to Adriano Barlotti, pp. S13–S27. MR 2028385
- J.I. Cogolludo and A. Libgober, Free quotients of fundamental groups of smooth quasiprojective varieties, Proc. Edinb. Math. Soc. (2) 64 (2021), no. 4, 924–946. MR 4349417
- J. I. Cogolludo Agustín and V. Florens, Twisted Alexander polynomials of plane algebraic curves, J. Lond. Math. Soc. (2) 76 (2007), no. 1, 105–121. MR 2351611
- J.I. Cogolludo-Agustín and E. Elduque, Quasi-projective varieties whose fundamental group is a free product of cyclic groups, Preprint available at arXiv: 2210.17282.
- J.I. Cogolludo-Agustín and A. Libgober, Mordell-Weil groups of elliptic threefolds and the Alexander module of plane curves, J. Reine Angew. Math. 697 (2014), 15–55. MR 3281651
- P. Deligne, Le groupe fondamental du complément d'une courbe plane n'ayant que des points doubles ordinaires est abélien (d'après W. Fulton)., Bourbaki Seminar, Vol. 1979/80,, Lecture Notes in Math., 842,, ,, 1981, pp. 1–10,. MR 636513
- W. Fulton, On the fundamental group of the complement of a node curve, Ann. of Math. (2) 111 (1980), no. 2, 407–409. MR 569076
- A. Libgober, Alexander invariants of plane algebraic curves, Singularities, Part 2 (Arcata, Calif., 1981), Proc. Sympos. Pure Math., vol. 40, Amer. Math. Soc., Providence, RI, 1983, pp. 135–143. MR 713242
- <u>—</u>, Characteristic varieties of algebraic curves, Applications of algebraic geometry to coding theory, physics and computation (Eilat, 2001), Kluwer Acad. Publ., Dordrecht, 2001, pp. 215–254. MR 1 866 902
- Mon vanishing loci of Hodge numbers of local systems, Manuscripta Math. 128 (2009), no. 1, 1–31. MR 2470184
- M.A. Marco-Buzunáriz and M. Rodríguez, Sirocco: A library for certified polynomial root continuation, Mathematical Software–ICMS 2016: 5th International Conference, Berlin, Germany, July 11-14, Proceedings 5 (2016), 191–197.
- H. Tokunaga, Geometry of irreducible plane quartics and their quadratic residue conics, J. Singul. 2 (2010), 170–190. MR 2763025

 M. Wada, Twisted Alexander polynomial for finitely presentable groups, Topology 33 (1994), no. 2, 241–256. MR 1273784

Departamento de Matemáticas, IUMA, Universidad de Zaragoza, C. Pedro Cerbuna 12, 50009 Zaragoza, Spain.

Email address: jicogo@unizar.es

Departamento de Matemáticas, Universidad Autónoma de Madrid, ICMAT, 28049 Madrid, Spain.

 $Email \ address:$ eva.elduque@uam.es