A BINARY-MATROID ANALOGUE OF A GRAPH CONNECTIVITY THEOREM OF JAMISON AND MULDER

CAMERON CRENSHAW AND JAMES OXLEY

ABSTRACT. Let G be a graph such that, whenever two vertices x and y of G are joined by three internally disjoint paths, x and y are adjacent. Jamison and Mulder determined that the set of such graphs coincides with the set of graphs that can be built from cycles and complete graphs via 1-sums and parallel connections. This paper proves an analogous result for binary matroids.

1. INTRODUCTION

Jamison and Mulder [3] defined a graph G to be Θ_3 -closed if, whenever distinct vertices x and y of G are joined by three internally disjoint paths, x and y are adjacent. For disjoint graphs G_1 and G_2 , a 1-sum of G_1 and G_2 is a graph that is obtained by identifying a vertex of G_1 with a vertex of G_2 . Following Jamison and Mulder, we define a 2-sum of G_1 and G_2 to be a graph that is obtained by identifying an edge of G_1 with an edge of G_2 . Note that, in contrast to some other definitions of this operation, we retain the identified edge as an edge of the resulting graph. The main result of Jamison and Mulder's paper is the following.

Theorem 1.1. A connected graph G is Θ_3 -closed if and only if G can be built via 1-sums and 2-sums from cycles and complete graphs.

This paper generalizes Theorem 1.1 to binary matroids; all matroids considered here are binary unless stated otherwise. The terminology and notation follow [5] with the following additions. We will use P_r to denote the rank-r binary projective geometry, PG(r-1,2). A theta-graph is a graph that consists of two distinct vertices and three internally disjoint paths between them. A theta-graph in a matroid M is a restriction of M that is isomorphic to the cycle matroid of a theta-graph. Equivalently, it is a restriction of M that is isomorphic to a matroid that is obtained from $U_{1,3}$ by a sequence of series extensions. The series classes of a theta-graph are its arcs. Let T be a theta-graph of M with arcs A_1 , A_2 , and A_3 . If M has an element e such that, for every i, either $A_i \cup e$ is a circuit of M, or $A_i = \{e\}$, then e completes T in M, and T is said to be complete. A matroid M is

Date: June 14, 2023.

²⁰¹⁰ Mathematics Subject Classification. 05B35, 05C40.

Key words and phrases. theta-graph, binary matroid, parallel connection.

matroid Θ_3 -closed if every theta-graph of M is complete. The next theorem is the main result of this paper.

Theorem 1.2. A matroid M is matroid Θ_3 -closed if and only if M can be built via direct sums and parallel connections from circuits, cycle matroids of complete graphs, and projective geometries.

Suppose M is isomorphic to the cycle matroid of a graph G. Two vertices in G that are joined by three internally disjoint paths are adjacent via an edge e exactly when the corresponding theta-graph of M is completed by e. In other words, G is Θ_3 -closed if and only if M is matroid Θ_3 -closed. This allows us to refer to M as Θ_3 -closed without ambiguity. We will denote the class of Θ_3 -closed matroids by Θ_3 .

Section 2 introduces supporting results. The 3-connected matroids that are Θ_3 -closed are characterized in Section 3, and the proof of Theorem 1.2 appears in Section 4.

2. Preliminaries

Our first proposition collects some essential properties of Θ_3 -closed matroids. These properties will be used frequently and often implicitly.

Proposition 2.1. If $M \in \Theta_3$, then

- (i) $\operatorname{si}(M) \in \Theta_3$;
- (ii) $M|F \in \Theta_3$ for every flat F of M; and
- (iii) $M/e \in \Theta_3$ for every $e \in E(M)$.

Proof. Parts (i) and (ii) are straightforward. For part (iii), let T be a theta-graph of M/e. Then $[(M/e)|T]^*$ is obtained from $U_{2,3}$ by adding elements in parallel to the existing elements. Since M is binary, it follows that $M^*/(E(M) - (T \cup e))$ is obtained from $[(M/e)|T]^*$, that is, from $M^*/(E(M) - (T \cup e)) \setminus e$, by adding e as a coloop or by adding e in parallel to one of the existing elements. Thus, e is either a loop in $M|(T \cup e)$, or is in series with another element. Hence, since T is complete in M, it is complete in M/e.

Evidently, a matroid is in Θ_3 if and only if its connected components are in Θ_3 . This will also be used implicitly throughout the paper. The following is an immediate consequence of Proposition 2.1.

Corollary 2.2. If $M \in \Theta_3$ and N is a parallel minor of M, then $N \in \Theta_3$.

From, for example, [5, Exercise 8.3.3], if $M = M_1 \oplus_2 M_2$, then M_1 and M_2 are parallel minors of M. The next result now follows from Corollary 2.2.

Corollary 2.3. If $M \oplus_2 N$ is in Θ_3 , then M and N are in Θ_3 .

To see that the converse of the last corollary fails, observe that $M(K_{2,4})$ is not in Θ_3 although it is the 2-sum of two copies of a matroid in Θ_3 .

We conclude this section with a result about constructing larger matroids in Θ_3 from smaller ones. Recall that, for sets X and Y in a matroid M, the *local connectivity* between X and Y, denoted $\sqcap(X,Y)$, is defined by $\sqcap(X,Y) = r(X) + r(Y) - r(X \cup Y)$. We will use the following result about local connectivity from, for example, [5, Lemma 8.2.3].

Lemma 2.4. Let X_1 , X_2 , Y_1 , and Y_2 be subsets of the ground set of a matroid M. If $X_1 \supseteq Y_1$ and $X_2 \supseteq Y_2$, then $\sqcap(X_1, X_2) \ge \sqcap(Y_1, Y_2)$.

Proposition 2.5. For matroids M and N, the parallel connection P(M, N) is in Θ_3 if and only if $M \in \Theta_3$ and $N \in \Theta_3$.

Proof. Let p be the basepoint of the parallel connection. When p is a loop or a coloop of M, the matroid P(M, N) is $M \oplus (N/p)$ or $(M/p) \oplus N$, respectively. In these cases, it follows using Proposition 2.1 that the result holds. Thus we may assume that p is neither a loop nor a coloop of M or N. Suppose $P(M, N) \in \Theta_3$. Let B_M be a basis for M containing p. Extend B_M to a basis B for P(M, N). After contracting both the elements of $B - B_M$ in P(M, N) as well as all of the resulting loops, the remaining elements of E(N) - p are parallel to p. We deduce that M, and similarly N, is a parallel minor of P(M, N). Hence, by Corollary 2.2, M and N are in Θ_3 .

Conversely, suppose that $M, N \in \Theta_3$ and let T be a theta-graph of P(M, N) with arcs A_1, A_2 , and A_3 . Then we may assume that $|A_i| \ge 2$ for each i, otherwise T is complete. Suppose $p \in A_1$. Then $A_1 \cup A_2$ is a circuit containing p, so it is contained in E(M) or E(N) depending on which of these sets contains $A_1 - p$. It follows that the same set contains A_2 and, likewise A_3 , so T is complete. Hence we may assume that $p \notin T$.

Suppose that each of the arcs of T meets both $E(M \setminus p)$ and $E(N \setminus p)$. Let $T_M = E(T) \cap E(M)$, and similarly for T_N . Note that T_M and T_N are independent, and $T_M \cup T_N = E(T)$, so

$$\sqcap (T_M, T_N) = r(T_M) + r(T_N) - r(T_M \cup T_N)$$

= $|T_M| + |T_N| - (|T_M| + |T_N| - 2)$
= 2.

However, $\sqcap (E(M), E(N)) = 1$, contradicting Lemma 2.4.

Next, suppose that each of A_1 and A_2 meets both $E(M \setminus p)$ and $E(N \setminus p)$. Then, from above, we may assume that $A_3 \subseteq E(M \setminus p)$. The circuits $A_1 \cup A_3$ and $A_2 \cup A_3$ have the form $(C_1 - p) \cup (D_1 - p)$ and $(C_2 - p) \cup (D_2 - p)$, respectively, for circuits C_1 and C_2 of M containing p, and circuits D_1 and D_2 of N containing p. Because $A_3 \subseteq E(M)$ and $A_1 \cap A_2 = \emptyset$, it follows that $D_1 - p$ and $D_2 - p$ are disjoint. However, since M is binary, $D_1 \triangle D_2$ contains a circuit of P(M, N) that is properly contained in the circuit $A_1 \cup A_2$, a contradiction.

Now, suppose that A_1 meets both $E(M \setminus p)$ and $E(N \setminus p)$. Then, from above, each of the remaining arcs of T lies in $E(M \setminus p)$ or $E(N \setminus p)$. We may assume that $A_2 \subseteq E(M \setminus p)$. Suppose $A_3 \subseteq E(N \setminus p)$. Then the circuits $A_1 \cup A_2$ and $A_3 \cup A_2$ have the form $(C_1 - p) \cup (D_1 - p)$ and $(C_3 - p) \cup (D_3 - p)$, respectively, for circuits C_1 and C_3 of M containing p, and circuits D_1 and D_3 of N containing p. Now, since $A_2 \subseteq E(M \setminus p)$ and A_1 meets $E(M \setminus p)$, the set $C_1 - p$ properly contains A_2 . Further, as A_3 does not meet $E(M \setminus p)$, we have that $A_2 = C_3 - p$. This means $A_2 \cup p$ is the circuit C_3 , but $A_2 \cup p$ is properly contained in C_1 , a contradiction.

We conclude that $A_3 \subseteq E(M \setminus p)$. Form T' from T by replacing the portion of A_1 in $E(N \setminus p)$ by p. Observe that T' is isomorphic to a series minor of T, so T' is a theta-graph. Moreover, T' is a theta-graph of M, so it is completed in M by an element f. Now, since T and T' share an arc, f also completes T in P(M, N).

We are left to consider the case when each arc of T is contained in either E(M) or E(N). If all three arcs belong to E(M), say, then T is complete in M, and so is complete in P(M, N). Otherwise, p completes T.

3. The 3-Connected Θ_3 -closed Matroids

The proof of Theorem 1.2 will use the canonical tree decomposition of Cunningham and Edmonds [2] and, in support of that approach, this section proves the following 3-connected form of Theorem 1.2.

Theorem 3.1. Let M be a simple 3-connected Θ_3 -closed matroid. Then M is a projective geometry or the cycle matroid of a complete graph.

The proof of this theorem relies on the next two propositions.

Proposition 3.2. If M is a simple matroid in Θ_3 and M has a spanning $M(K_{r+1})$ -restriction, then $M \cong M(K_{r+1})$ or $M \cong P_r$.

Proof. Take a standard binary representation for P_r , and view M as the restriction of P_r to the set X of vectors. Recall that the number of nonzero entries of a vector is its *weight*, and that the *distance* between two vectors is the number of coordinates upon which they disagree. Because M has an $M(K_{r+1})$ -restriction, we may assume that X contains the set Z of vectors of weight one or two. We may assume that $Z \neq X$. Then M has an element e of weight at least three. We shall establish that $M \cong P_r$ by proving the following three assertions.

- (i) M has an element of weight three;
- (ii) if the matroid M has every element of weight k-1 and an element of weight k, for some k exceeding two, then M has every element of weight k; and
- (iii) if M has every element of weight k, where $3 \le k < r$, then M has an element of weight k + 1.

Let e_i denote the weight-1 element whose nonzero entry is in the *i*th position. To show (i), we may assume e has weight $k \ge 4$. Say $e = e_1 + e_2 + \cdots + e_k$. Let $Y = \{e, e_1, e_2, e_4, e_5, \ldots, e_k, e_1 + e_3, e_2 + e_3\}$. Then M|Y is a theta-graph having arcs $\{e, e_4, e_5, \ldots, e_k\}$, $\{e_1, e_2 + e_3\}$, and $\{e_2, e_1 + e_3\}$. This theta-graph forces $e_1 + e_2 + e_3$ to be an element of M, so (i) holds.

To prove (ii), we may assume k < r. Suppose g is an element of weight k not in M, and let f be an element of weight k in M with minimum distance from g. Let s label a row where f is 1 and g is 0, and let t label a row where g is 1 and f is 0. Next, as $k \ge 3$, there are two additional rows, u and v, distinct from s where f is 1. Now, the set $\{f, e_u, e_v, e_s + e_t\}$ is independent, so the arcs $\{f, e_s + e_t\}$, $\{e_u, f + e_u + e_s + e_t\}$, and $\{e_v, f + e_v + e_s + e_t\}$ form a theta-graph in M. This theta-graph implies that $f + e_s + e_t$ belongs to M. However, $f + e_s + e_t$ has weight k and is a smaller distance from g than f, a contradiction. Thus (ii) holds.

Finally, let f be an element of weight k + 1 for some k with $3 \le k < r$. By symmetry, we may assume that the set of rows in which f is nonzero contains $\{1, 2, 3\}$. Then $\{f, e_1, e_2, e_3\}$ is independent in M, and the sets $\{e_1, f + e_1\}$, $\{e_2, f + e_2\}$, and $\{e_3, f + e_3\}$ are the arcs of a theta-graph in M. This theta-graph shows that f belongs to M. Thus (iii) holds. Hence the proposition holds as well.

The second proposition that we use to prove Theorem 3.1 will follow from the following three results.

Lemma 3.3. Let M be a simple rank-r matroid in Θ_3 . Suppose that

(i) $r \ge 4;$

(ii) E(M) has a subset P such that $M|P \cong P_{r-1}$; and

(iii) E(M) - P contains at least three elements.

Then $M \cong P_r$.

Proof. View M as a restriction of P_r , and let $\{e, f, g\}$ be a subset of E(M) - P. Let p be a point in $E(P_r) - P$ that is not in $\{e, f, g\}$. Observe that, for each x in $\{e, f, g\}$, the third point on the line in P_r containing $\{x, p\}$ is in P. Thus there are three lines of M that meet at p. Provided p is not coplanar with $\{e, f, g\}$, these lines define a theta-graph in M that is completed by p, so p is in M. It remains to show that the point q of $E(P_r) - (P \cup \{e, f, g\})$ that is coplanar with $\{e, f, g\}$ belongs to M. But one easily checks that $P_r \setminus q$ is not in Θ_3 when $r \ge 4$. Thus $M \cong P_r$.

Corollary 3.4. Let M be a simple rank-r matroid in Θ_3 with $r \ge 3$. If M has a basis B and an element b in B so that, for each $\{x, y\} \subseteq B - b$, the set $\{b, x, y\}$ spans an F_7 -restriction of M, then $M \cong P_r$.

Proof. Let $B = \{b_1, b_2, \ldots, b_r\}$ with $b = b_1$. If r = 3, then the result is immediate, so suppose $r \ge 4$. By induction, $M | \operatorname{cl}(B - b_r)$ is isomorphic to P_{r-1} . Since $M | \operatorname{cl}(\{b_1, b_2, b_r\}) \cong F_7$, we see that this restriction contains an independent set of three elements that avoids $\operatorname{cl}(B - b_r)$. Lemma 3.3 now implies that $M \cong P_r$.

The next result was proved by McNulty and Wu [4, Lemma 2.10].

Lemma 3.5. Let M be a 3-connected binary matroid with at least four elements. Then, for any two distinct elements e and f of M, there is a connected hyperplane containing e and avoiding f.

For a simple binary matroid M, we now define the smallest Θ_3 -closed matroid whose ground set contains E(M). Let $M_0 = M$ and r(M) = r. Suppose M_0, M_1, \ldots, M_k have been defined. The simple binary matroid M_{k+1} is obtained from M_k by ensuring that, whenever T is an incomplete theta-graph of M_k , the element x that completes T is in $E(M_{k+1})$. Since each M_i is a restriction of P_r , there is a j for which $M_{j+1} = M_j$. When this first occurs, we call M_j the Θ_3 -closure of M. Evidently this is well defined. By associating M with its ground set, the Θ_3 -closure is a closure operator (but not necessarily a matroid closure operator) on the set of subsets of the ground set of any projective geometry containing M.

Proposition 3.6. Let M be a simple 3-connected matroid in Θ_3 , and let k be an integer exceeding two. If M has a simple minor N whose Θ_3 -closure is P_k , then M is a projective geometry.

Proof. Take subsets X and Y of E(M) such that $M/X \setminus Y = N$ with X independent and Y coindependent in M. The matroid M/X is in Θ_3 and has N as a spanning restriction. Therefore M/X has P_k as a restriction, so P_k is a minor of M. From here, the proof is by induction on the rank, r, of M.

If r = k, the result is immediate, so assume r > k. By Seymour's Splitter Theorem, P_k can be obtained from M by a sequence of single-element contractions and deletions, all while staying 3-connected. Let e be the first element that is contracted in this sequence. Note that $\operatorname{si}(M/e)$ is a 3-connected member of Θ_3 that has P_k as a minor. By induction, $\operatorname{si}(M/e) \cong P_{r-1}$. Fix an embedding of M in P_r . We may assume that $M \ncong P_r$. Then some line ℓ of P_r through e is not contained in E(M). For each subset Z of $E(P_r)$, let $\operatorname{cl}_P(Z)$ be its closure in P_r . Since $\operatorname{si}(M/e) \cong P_{r-1}$, there is an element s of E(M) that is in $\ell - \{e\}$. Let t be the point of P_r in $\ell - \{e\}$ that is not in E(M).

3.6.1. Let F be a rank-4 flat of M containing ℓ -t. Then M|F is isomorphic to one of $P(F_7, U_{2,3})$ or $F_7 \oplus U_{1,1}$ where the F_7 -restriction of M|F contains s but avoids e.

To see this, first note that, by Proposition 2.1(ii), M|F is in Θ_3 . Recall that each line of $cl_P(F)$ through e contains another point of F. Then there are three planes, π_1 , π_2 , and π_3 , of M|F containing $\ell - t$. Let \mathcal{P} be this set of planes. Therefore, each plane in \mathcal{P} has at least one pair of points that are not on ℓ such that these two points are collinear with either s or t. Call such a pair of points a *target pair*. The rest of the proof frequently relies of finding theta-graphs, particularly in rank 4. It maybe helpful to note that such a theta-graph will be isomorphic to $M(K_{2,3})$. Geometrically, this is three non-coplanar lines, each full except for shared point. This common point completes the theta-graph.

Suppose π_1 has a target pair collinear with t. Note that if two distinct planes in \mathcal{P} each have a target pair collinear with t, then these pairs, along




FIGURE 1. In the proof of 3.6.1, each of $M|\pi_2$ and $M|\pi_3$ has this form.




FIGURE 2. The matroid $M|(\pi_2 \cup \pi_3)$ in the proof of 3.6.1.

with $\{e, s\}$, are the arcs of an incomplete theta-graph in M|F, an impossibility. Consequently, neither π_2 nor π_3 has a target pair collinear with t, so both have the form in Figure 1. The restriction of M to $\pi_2 \cup \pi_3$ is given in Figure 2.

The plane π_1 adds a pair of points collinear with t to $M|(\pi_2 \cup \pi_3)$, and one readily checks that the addition of this pair gives a restriction of M|F that is isomorphic to $F_7 \oplus_2 U_{2,3}$. A theta-graph of M|F now gives that M|F has a restriction isomorphic to $P(F_7, U_{2,3})$. It follows that $M|F \cong P(F_7, U_{2,3})$ otherwise, by Lemma 3.3, $M|F \cong P_4$ and we obtain the contradiction that $t \in F$.

We may now suppose that no π_i has a target pair collinear with t. It follows that each target pair in each π_i is collinear with s, and that e is the only element of F outside of the target pairs. Observe that the target pairs must be coplanar as, otherwise, we can find an incomplete theta-graph in M|F. Thus $M|F \cong F_7 \oplus U_{1,1}$. Noting that e is not in the F_7 -restriction of M|F, we conclude that 3.6.1 holds. Since M is 3-connected, it follows from 3.6.1 that $r \ge 5$. By Lemma 3.5, there is a connected hyperplane H of M containing s and avoiding e. Let $s = b_1$ and let $\{b_1, b_2, \ldots, b_{r-1}\}$ be a basis B of M|H. For distinct elements iand j of $\{2, 3, \ldots, r-1\}$, let $F_{i,j}$ denote the rank-4 flat $M|\operatorname{cl}(\{e, s, b_i, b_j\})$ of M. By 3.6.1, $M|F_{i,j}$ is isomorphic to $F_7 \oplus U_{1,1}$ or $P(F_7, U_{2,3})$. Let X be the subset of $F_{i,j}$ such that $M|X \cong F_7$, and recall that $e \notin X$ and $s \in X$. The hyperplane H either contains X or meets X along one of the lines $\operatorname{cl}(\{s, b_i\})$ or $\operatorname{cl}(\{s, b_i\})$. We deduce the following.

3.6.2. For each pair $\{i, j\} \subseteq \{2, 3, ..., r-1\}$, at least one of $s + b_i$ or $s + b_j$ is in E(M).

Suppose $s + b_2$ is not in E(M). By 3.6.2, the element $s + b_i$ belongs to E(M) for every i in $\{3, 4, \ldots, r-1\}$. Consequently, for each pair $\{i, j\}$ in $\{3, 4, \ldots, r-1\}$, the hyperplane H contains the copy of F_7 in $M|F_{i,j}$, and this F_7 is spanned by $\{s, b_i, b_j\}$. Corollary 3.4 now implies that $M|\operatorname{cl}(B-b_2) \cong P_{r-2}$.

Now, since M|H is connected, H contains an element f that is not in $\operatorname{cl}(B-b_2) \cup b_2$. The line $\operatorname{cl}(\{b_2, f\})$ meets the projective geometry $\operatorname{cl}(B-b_2)$, so $b_2 + f$ is also in M. Now consider $Y = \operatorname{cl}(\{e, s, b_2, b_2 + f\})$. The intersection $H \cap Y$ contains the line $\{b_2, f, b_2 + f\}$ and also the element s. By applying 3.6.1 to M|Y, we see that $H \cap Y$ is an F_7 -restriction containing s and b_2 . Since $s + b_2$ is not in E(M), we have a contradiction.

We conclude that $s + b_i$ is in E(M) for every i in $\{2, 3, \ldots, r-1\}$. The flat $M|F_{i,j}$ now meets H in an F_7 -restriction for every pair $\{i, j\} \subseteq \{2, 3, \ldots, r-1\}$ so, by Corollary 3.4, M|H is a projective geometry. Finally, as M is 3-connected, there is an independent set of three elements in E(M) avoiding H. Hence, by Lemma 3.3, $M \cong P_r$.

We are now ready to prove the main result of this section.

Proof of Theorem 3.1. Let r be the rank of M. If M is graphic, then Theorem 1.1 gives that $M \cong M(K_{r+1})$, so we may assume that M is not graphic. Thus M has a minor N isomorphic to F_7 , F_7^* , $M^*(K_{3,3})$, or $M^*(K_5)$. By Proposition 3.6, it now suffices to show that the Θ_3 -closure, $\Theta(N)$, of N is a projective geometry.

This is immediate when $N \cong F_7$, so suppose N is isomorphic to F_7^* , labelled as in Figure 3. The theta-graphs of N imply that, in $\Theta(N)$, the plane containing $\{1, 2, 5, 6\}$ is isomorphic to F_7 . Proposition 3.6 now implies that M is isomorphic to P_r .

Next, suppose $N \cong M^*(K_{3,3})$. The complement of N in P_4 is $U_{2,3} \oplus U_{2,3}$; let x be an element of this complement. The elementary quotient of Nobtained by extending N by x and then contracting x is shown in Figure 4. The three pairwise-skew 2-element circuits of this quotient correspond to three lines in the extension of N by x where the union of these lines has rank four. Thus N contains a theta-graph that is completed by x. It follows




FIGURE 3. The matroid F_7^* .




FIGURE 4. A quotient of $M^*(K_{3,3})$.




FIGURE 5. The graph K_5 .

that x and, symmetrically, every point in the complement of N in P_4 , belongs to $\Theta(N)$. Lemma 3.3 now implies that $\Theta(N) \cong P_4$.

Now suppose $N \cong M^*(K_5)$, where K_5 is labelled as in Figure 5. Figure 6 gives a corresponding binary representation of N. The dual of a theta-graph with arcs of size at least two is a triangle with no trivial parallel classes.

				4						
Γ	1	0	0	0	0	0	1	0	0	$ \begin{bmatrix} 1 \\ 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} $
	0	1	0	0	0	0	1	0	1	0
	0	0	1	0	0	0	1	1	0	0
	0	0	0	1	0	0	0	1	0	1
	0	0	0	0	1	0	0	0	1	1
L	0	0	0	0	0	1	0	1	1	0

FIGURE 6. A binary representation of $M^*(K_5)$.

Therefore, we can detect theta-graph restrictions of N by contracting elements of N^* to produce such a triangle. For example, $N^*/5, 8$ is the dual of a theta-graph in N with arcs $\{1, 2\}, \{3, 7\},$ and $\{0, 4, 6, 9\}$. This theta-graph is completed by the element $[1 \ 1 \ 0 \ 0 \ 0]^T$, so this element belongs to $\Theta(N)$. The following is a list of duals of theta-graphs of N and the corresponding elements of $\Theta(N)$ that they produce using this reasoning.

- $N^*/5, 8$ gives $[1 \ 1 \ 0 \ 0 \ 0]^T \in \Theta(N)$.
- $N^*/4, 9$ gives $\begin{bmatrix} 1 & 0 & 1 & 0 & 0 \end{bmatrix}^T \in \Theta(N).$
- $N^*/3, 9$ gives $\begin{bmatrix} 1 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}^T \in \Theta(N).$
- $N^*/2, 8$ gives $[1 \ 0 \ 0 \ 1 \ 0]^T \in \Theta(N).$
- $N^*/0, 6$ gives $\begin{bmatrix} 0 & 1 & 1 & 0 & 0 \end{bmatrix}^T \in \Theta(N)$.
- $N^*/1, 8$ gives $[0\ 1\ 0\ 0\ 1\ 0]^T \in \Theta(N)$.
- $N^*/0, 3$ gives $\begin{bmatrix} 0 & 1 & 0 & 0 & 0 \end{bmatrix}_{T}^{T} \in \Theta(N).$
- $N^*/1,9$ gives $[0\ 0\ 1\ 1\ 0\ 0]^T \in \Theta(N)$.
- $N^*/0, 2$ gives $[0 \ 0 \ 1 \ 0 \ 0 \ 1]^T \in \Theta(N).$
- $N^*/6, 7$ gives $[0 \ 0 \ 0 \ 1 \ 1 \ 0]^T \in \Theta(N).$
- $N^*/5, 7$ gives $[0 \ 0 \ 0 \ 1 \ 0 \ 1]^T \in \Theta(N)$.
- $N^*/4, 7$ gives $[0 \ 0 \ 0 \ 0 \ 1 \ 1]^T \in \Theta(N).$

It is now straightforward to find theta-graphs in $\Theta(N)$ that are completed by the elements $[1 \ 0 \ 0 \ 0 \ 1]^T$, $[0 \ 1 \ 0 \ 1 \ 0 \ 0]^T$, and $[0 \ 0 \ 1 \ 0 \ 1 \ 0]^T$, so $\Theta(N)$ contains every vector of weight 1 or 2. Thus $\Theta(N)$ properly contains $M(K_7)$, so, by Proposition 3.2, $\Theta(N) \cong P_6$.

4. The Main Result

After a review of canonical tree decompositions, this section proves Theorem 1.2. For a set $\{M_1, M_2, \ldots, M_n\}$ of matroids, a *matroid-labelled tree* with vertex set $\{M_1, M_2, \ldots, M_n\}$ is a tree T such that

- (i) if e is an edge of T with endpoints M_i and M_j , then $E(M_i) \cap E(M_j) = \{e\}$, and $\{e\}$ is not a separator of M_i or M_j ; and
- (ii) $E(M_i) \cap E(M_j)$ is empty if M_i and M_j are non-adjacent.

The matroids M_1, M_2, \ldots, M_n are called the *vertex labels* of T. Now suppose e is an edge of T with endpoints M_i and M_j . We obtain a new matroidlabelled tree T/e by contracting e and relabelling the resulting vertex with $M_i \oplus_2 M_j$. As the matroid operation of 2-sum is associative, T/X is well defined for all subsets X of E(T).

Let T be a matroid-labelled tree for which $V(T) = \{M_1, M_2, \ldots, M_n\}$ and $E(T) = \{e_1, e_2, \ldots, e_{n-1}\}$. Then T is a *tree decomposition* of a connected matroid M if

- (i) $E(M) = (E(M_1) \cup E(M_2) \cup \cdots \cup E(M_n)) \{e_1, e_2, \dots, e_{n-1}\};$
- (ii) $|E(M_i)| \ge 3$ for all *i* unless |E(M)| < 3, in which case n = 1 and $M = M_1$; and
- (iii) M labels the single vertex of T/E(T).

In this case, the elements $\{e_1, e_2, \ldots, e_{n-1}\}$ are the *edge labels* of T. Cunningham and Edmonds [2] (see also [5, Theorem 8.3.10]) proved the next theorem that says that M has a *canonical tree decomposition*, unique to within relabelling of the edges.

Theorem 4.1. Let M be a 2-connected matroid. Then M has a tree decomposition T in which every vertex label is 3-connected, a circuit, or a cocircuit, and there are no two adjacent vertices that are both labelled by circuits or are both labelled by cocircuits. Moreover, T is unique to within relabelling of its edges.

We now complete the proof of our main result.

Proof of Theorem 1.2. Since circuits, cycle matroids of complete graphs, and projective geometries are in Θ_3 , by Proposition 2.5, every matroid that can be built from such matroids by a sequence of parallel connections is in Θ_3 .

To prove the converse, we begin by noting that loops can be added via direct sums and that parallel elements can be added via parallel connections of circuits, so we may assume M is simple. Let T be the canonical tree decomposition of M. The proof is by induction on |V(T)|.

If |V(T)| = 1, then M is 3-connected and the result holds by Theorem 3.1. Now assume T has at least two vertices, and let N be a matroid labelling a leaf of T. Since M is simple, N is not a cocircuit. We may now write $M = N \oplus_2 M_1$, where, by Corollary 2.3, N and M_1 are in Θ_3 . Thus, by Theorem 3.1, N is a circuit, the cycle matroid of a complete graph of rank at least three, or a projective geometry of rank at least three. Moreover, by induction, M_1 is a parallel connection of circuits, cycle matroids of complete graphs, and projective geometries.

Let N_1 be the label of the neighbor of N in T, and suppose N_1 is not a cocircuit. In this case, each of N and N_1 is a circuit, the cycle matroid of a complete graph of rank at least three, or a projective geometry of rank at least three, and they are not both circuits. Therefore, if p is the basepoint of

the 2-sum $N \oplus_2 N_1$, there are circuits in N and N_1 that form a theta-graph that is completed by p, a contradiction. Thus N_1 is a cocircuit.

Now let k be the degree of N_1 in T. Evidently N_1 has at least k elements, but, since M is simple, N_1 has at most k + 1 elements. If k = 2, then N_1 has three elements as N_1 labels a vertex of T. Otherwise $k \ge 3$, so there are circuits in M that form a theta-graph that is completed by an element of N_1 . Hence N_1 has k + 1 elements, and therefore corresponds to a parallel connection of its neighbors. It now follows that M is the parallel connection of circuits, cycle matroids of complete graphs, and projective geometries. \Box

Acknowledgements

The authors thank the referee for suggesting a number of improvements to the paper.

References

- Bixby, R. E., Kuratowski's and Wagner's Theorems for matroids, J. Combin. Theory Ser. B 22 (1977), 31–53.
- [2] Cunningham, W. H., A combinatorial decomposition theory, Ph.D. thesis, University of Waterloo, 1973.
- [3] Jamison, R. E. and Mulder, H. M., Graphs constructible from cycles and complete graphs, *Discrete Math.* 340 (2017), 2301–2306.
- [4] McNulty, J. and Wu, H., Connected hyperplanes in binary matroids, J. Combin. Theory Ser. B 79 (2000), 87–97.
- [5] Oxley, J. G., *Matroid Theory*, Second edition, Oxford University Press, New York, 2011.
- [6] Seymour, P. D., Decomposition of regular matroids, J. Combin. Theory Ser. B 28 (1980), 305–359.

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA *Email address*: ccrens5@lsu.edu

MATHEMATICS DEPARTMENT, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LOUISIANA *Email address*: oxley@math.lsu.edu