Nonlinear Generalized Ridge Regression

Robert L. Obenchain Risk Benefit Statistics

October 2023

Abstract

A Two-Stage approach is described that literally "straightens out" any potentially nonlinear relationship between a y-outcome variable and each of $p \ge 2$ potential x_i -predictor variables. The y-outcome is then predicted from all p of these "linearized" $s(x_i)$ -predictors using the form of Generalized Ridge Regression that is most likely to yield minimal MSE risk under Normal distribution-theory. These estimates are then compared and contrasted with those from the Generalized Additive Model that uses the same x_i variables.

Keywords: Efficient Generalized Ridge Regression, Normal-theory Maximum Likelihood, Generalized Additive Model, Mean-Squared-Error Risk, R-functions.

1 Introduction

We discuss a simple "Two-Stage" approach that combines use of nonlinear gam() smooth "spline" functions from the mgcv R-package, Wood (2001,2023), Wood and Augustin (2002), Wood, Pya and Saefken (2016) in stage one, with optimally biased estimation of linear models using the eff.ridge() function from the RXshrink R-package, Obenchain (2022,2023), in stage two.

This new multiple regression strategy is applicable when $p \ge 2$ different individual *x*-vectors are initially used alone (one-at-a-time) to make *p* (different) nonlinear predictions of a single observed *y*-outcome vector. Especially when the number, *n*, of observations available is large [e.g. n =2,793 in the example analyses], using the gam() function from the *mgcv R*-package, Wood (2023), provides superior speed and efficiency.

Our numerical example will use p = 6 x-predictors of y-Outcome values of "Age Adjusted Circulatory-Respiratory Mortality" rates for 2, 793 US Counties in 2016. A CSV file containing the data can be downloaded from *Dryad*, Young and Obenchain (2022).

Upon completing this "Initial Phase" of analysis, the analyst forms an R-data.frame containing the following 2*p+1 variables: (a) the observed y-outcome vector, (b) the p x-predictor variables,

TIDDE I TRaines and Descriptions of Variables								
Var	Name	Description	Min	Max				
у	AACRmort	Age Adjusted Circulatory-Respiratory Mortality	95.5	46.00				
x1	Avoc	Anthropogenic Volatile Organic Compounds	0.22282	2.8906				
$\mathbf{x}2$	Bvoc	Biogenic Volatile Organic Compounds	0.2611	3.3092				
x3	PREMdeath	Premature Death Index (before Age 75)	2853	36469				
x4	ASmoke	Adult Smoking Percentage	0.06735	0.41202				
x5	ChildPOV	Children Living in Poverty Index	0.0290	0.6630				
x6	IncomIEQ	Income Inequality Index	2.932	8.929				

TABLE 1 – Names and Descriptions of Variables

and (c) the p y-predictor vectors, each of which is a semi-parametric, nonlinear function of only one of the given x-predictor vectors. Note that gam() fitting uses a form of Generalized Ridge Regression that minimizes a Generalized Cross Validation (GCV) statistic to provide "automatic smoothing".

Finally, the analyst uses the eff.ridge() and MLcalc() functions from the "RXshrink" R-package, Obenchain (2023), to first generate "TRACE" graphics such as those displayed in Figure 4 and then to calculate the relative MSE risk statistics reported in Table 3. Again, we find that the spline estimates with MaximumLikelihood of being "optimally biased" under Normal distribution-theory, Obenchain (2022), generally tend to achieve both lower "residual error" and lower "relative" MSE risk compared to *unbiased* (BLUE) least squares estimates.

2 Linear Rank Deficiency

Regression models typically assume that (1) the n by p "centered" X-matrix, (I - 11'/n)X, is of full (column) rank, p, and that (2) an "intercept" term, μ , is included to assure that the "fit" (a line, plane or hyper-plane) passes through $y = \bar{y}$ in the p + 1 dimensional Euclidean space where all x-coordinates equal their mean vector. Specifically,

$$\hat{\mu} = \bar{y} - \bar{x}'\hat{\beta} \ . \tag{1}$$

Note that this *intercept estimate* automatically changes when the $\hat{\beta}$ coefficient vector is shrunken via generalized ridge regression, allowing the implied "fit" to pivot about \bar{x} in *p*-space.

3 Visual Validation using Spline *x*-Coordinates

The syxi() and plot.syxi() functions in Version 2.3 of the "RXshrink" CRAN *R*-Package, Obenchain (2023), compute and display three types of predictions of a single *y*-outcome variable. The two

scatter plots in each row of Figures 1, 2 and 3 display x_i vs. y in the left-hand plot and the gam() spline $s(x_i)$ fitted values vs. y in the right-hand plot. Note that each left-hand scatter plot contains both a purely "linear" $lm(y \sim x_i)$ fit (dashed RED) and a clearly "nonlinear" fit (BLUE).

The right-hand plots in these Figures provide the statistical information that an analyst needs to perform "Visual Validation." Specifically, the fitted spline *abscissa* in each right-hand scatter provides a nonlinear transformation that, in my experience, frequently tends to make the $lm(y \sim s(x_i))$ fit (solid BLUE) "LOOK" rather reasonable. On the other hand, this transformed scatter may lack homogeneity of variance.

4 Linear GRR Shrinkage

Early proposals for choosing the "k-factor" defining the 1-parameter shrinkage path of Hoerl and Kennard (1970) were based mostly upon "heuristics." In sharp contrast, shrinkage based upon Normal-theory Maximum Likelihood estimation has a firm theoretical foundation, Obenchain (1975, 1977, 1978). The Efficient p-parameter shrinkage-path is a two-piece linear spline with a single interior Knot at the overall "best" (minimum MSE risk) vector of point-estimates, Obenchain (2022).

Ridge computations and *TRACE* displays for the Efficient Path (plus other visualizations) are implemented in version 2.3 of the *RXshrink R*-package, Obenchain (2023). Each *TRACE* typically displays estimates of $p \ge 2$ quantities that change as *shrinkage* occurs. The "coef" *TRACE* displays the *p* fitted linear-model β -coefficients, while the "rmse" *TRACE* plots *relative* mean-squared-error estimates given by the diagonal elements of the MSE-matrix divided by the OLS-estimate of σ^2 .

5 Overall Extent of Shrinkage

The *multicollinearity allowance*, *m*, measures the "extent" of shrinkage applied.

$$m = p - \delta_1 - \dots - \delta_p = rank(X) - trace(\Delta).$$
⁽²⁾

Note that $0 \le m \le p$, Obenchain (1977). Besides being the rank of X, p is also the *trace* of the OLS Hat-matrix, XX^+ . Similarly, $trace(\Delta)$ is like a measure of "rank" for the *diagonal* $(p \times p) \Delta$ shrinkage-factor matrix.

6 Correlations within Groups of Variables

Note that the 2,793 observed y-Outcomes are positively correlated with the six given x-variables and also tend to be slightly more positively correlated with the six gam() spline prediction vectors,



Figure 1: The initial pair of gam() predictions of $AACRmort \ y$ -outcomes are the x1 = Avoc and x2 = Bvoc Volatile Organic Components of PM2.5 air pollution. Note that the relationship between Avoc and AACRmort is distinctly non-monotone, while Bvoc has nearly monotone (possibly causal) effects at its higher levels. While Avoc and x6 = IncomIEQ are the two least important predictors of mortality, Bvoc is slightly more important.



Figure 2: The third and fourth pairs of gam() predictions of $AACRmort \ y$ -outcomes are shown here. x3 = PREMdeath is quite curved and has the highest Adjusted R-square, confirmed by s3. x4 = ASmoke has nearly monotone effects on mortality and the second highest Adjusted R-square, confirmed by s4.



Figure 3: The fifth and sixth pairs of gam() predictions of y-outcomes are (x5 = ChildPOV, s5) and (x6 = IncomIEQ, s6). While x5 = ChildPOV is the third most important predictor of mortality, x6 = IncomIEQ is not very important.

as shown in Table 2.

TABL	E 2 - Pea	arson Cor	relations	s within	Two Se	ts of Se	ven Varia	ables
variable	У	x1	$\mathbf{x}2$	x3	x4	$\mathbf{x5}$	x6	
у	1.0000							
x1	0.2489	1.00000						
x2	0.4589	0.58472	1.0000					
x3	0.6421	0.08896	0.4217	1.00000				
x4	0.6047	0.32707	0.4622	0.67611	1.0000			
x5	0.5524	0.11336	0.4884	0.69932	0.6605	1.0000		
x6	0.3040	0.13933	0.4163	0.41804	0.3800	0.5708	1.0000	
variable	У	s1	s2	s3	s4	$\mathbf{s5}$	$\mathbf{s6}$	
у	1.0000							
s1	0.3888	1.0000						
s2	0.4809	0.5991	1.0000					
s3	0.6769	0.3077	0.5117	1.0000				
s4	0.6156	0.5002	0.5485	0.6621	1.0000			
s5	0.5656	0.3029	0.5192	0.7348	0.6607	1.0000		
s6	0.3379	0.1884	0.4378	0.4389	0.3775	0.5775	1.0000	

...

MSE Risk Comparisons 7

Applied researchers should consider transforming the individual x-variables available to them using nonlinear spline methods to potentially improve prediction of their target y-outcome variable. The one-variable-at-a-time first stage approach illustrated here provides a viable way to start. Once a researcher identifies $p \ge 2$ predictor variables, two different sets of Maximum Likelihood estimates under Normal distribution-theory (Unbiased or Optimally Shrunken) can be computed and compared in several ways.

c a

Note that Table 3 (on page 9) shows that the shrunken gam() model achieves not only lower residual error than the shrunken Linear model but also lower MSE risks "relative to" that lower residual error for 5 of the 6 spline predictor variables!

Note that the optimally shrunken $x_3 = PREM death$ coefficient has an estimated MSE risk of 0.5064 * 0.000756 = 0.000383, while that of its s3 counterpart is estimated to be larger: 0.4791 *0.000911 = 0.000436. All differences in MSE risks and relative risks again tend to be small in this large scale EPA Particulate Matter example, but achieving numerical risk reductions for 5



Figure 4: The two Traces of β -coefficients in the Top Row show that relative magnitudes of coefficient estimates are slightly unstable, with less shrinkage (to m = 0.775 rather than to m = 1.07) being appropriate when all six nonlinear transformations of the given x-variables are used in the model. The corresponding pair of *Relative Risk* (MSE/σ^2) Traces in the bottom row confirm that the six nonlinear predictors of mortality (AACRmort) rates provide better predictions. See Table 3 for more detail on this finding.

TABLE 3 -	Kev	Statistical	Comparis	sons for	the L	inear a	and NI	Models
		Statistical	Comparin			inour (1 III GUOI

Linear Model Formula	$y \sim x1$	+x2	+x3	+x4	+x5	+x6
Residual Mean Square	0.506395					
Residual Std. Error	0.711615					
OLS Beta Coefficients	0.060975	0.136652	0.389470	0.216750	0.099290	-0.063270
ML Optimally Biased Betas	0.064848	0.134431	0.384555	0.218586	0.101132	-0.062424
OLS Relative MSE Risks	0.000613	0.000745	0.000821	0.000822	0.000968	0.000519
ML Minimum Relative Risks	0.000292	0.000428	0.000756	0.000380	0.000524	0.000488
dMSE Estimates	0.9996	0.9413	0.9964	0.9429	0.9728	0.003282
Transformed Model Formula	$y \sim s1$	+s2	+s3	+s4	+s5	+s6
Residual Mean Square	0.479143					
Residual Std. Error	0.692202					
OLS Beta Coefficients	0.098704	0.056765	0.451541	0.214582	0.035727	-0.005387
ML Optimally Biased Betas	0.103079	0.056651	0.443230	0.221146	0.034597	-0.006164
OLS Relative MSE Risks	0.000590	0.000715	0.000852	0.000815	0.000996	0.000533
ML Minimum Relative Risks	0.000380	0.000559	0.000911	0.000790	0.000939	0.000453
dMSE Estimates	0.9997	0.4332	0.9961	0.8721	0.9359	0.9882

out of 6 relevant predictor variables certainly suggests that this simple "Two-Stage" approach deserves further study of its ability to estimate the effects of "x" and "s" variables and to predict "y-outcomes".

8 Summary

Here, we have provided a fairly large example (p = 6 x-predictors and n = 2,793 observations) of our "two-stage" approach. This appears to be a straight-forward "common sense" way to transform predictor variables and, ultimately, to end up with potentially "better" unbiased and/or shrunken predictions of the y-outcome.

In early arXiv papers on this topic, we called this two-stage approach "non-parametric". Others might prefer to call it "semi-parametric smoothing". Any transformation applied to an x-variable in stage-one must be nonlinear!

Software implementing accurate computations and providing clear "visual insights" into the strengths and weaknesses of alternative methods are indispensable components of an adequate *Tool Bag* for today's applied researchers.

9 References

Hastie, T. and Tibshirani, R. (1990). "Generalized Additive Models." Chapman and Hall.

Obenchain, R. L. (1975). "Ridge analysis following a preliminary test of the shrunken hypothesis." *Technometrics* 17, 431-441. http://doi.org/10.1080/00401706.1975.10489369

Obenchain, R. L. (1977). "Classical F-tests and confidence regions for ridge regression." *Techno*metrics 19, 429-439. http://doi.org/10.1080/00401706.1977.10489582

Obenchain, R. L. (1978). "Good and Optimal Ridge Estimators." Annals of Statistics 6, 1111–1121. http://doi.org/10.1214/aos/1176344314

Obenchain, R. L. (2022). "Efficient Generalized Ridge Regression", Open Statistics 3, 1–18. https://doi.org/10.1515/stat-2022-0108

Obenchain, R. L. (2023). "*RXshrink*: Maximum Likelihood Shrinkage using Generalized Ridge or Least Angle Regression Methods", ver 2.3, https://CRAN.R-project.org/package=RXshrink

Wood, S. N. (2001), "mgcv: GAMs and Generalized Ridge Regression for R." R News 1(2):20-25.

Wood, S. N. and Augustin, N. H. (2002), "GAMs with integrated model selection using penalized regression splines and applications to environmental modelling." *Ecological Modelling* 157: 157–177.

Wood, S. N., Pya N. and Saefken B. (2016), "Smoothing parameter and model selection for general smooth models (with discussion). *Journal of the American Statistical Association* 111: 1548–1575. http://doi:10.1080/01621459.2016.1180986

Wood, S. N. (2023), "mgcv: Mixed GAM Computation Vehicle with Automatic Smoothness Estimation", ver. 1.8-42, https://CRAN.R-project.org/package=mgcv

Young, S. S. and Obenchain, R. L. (2022). "EPA Particulate Matter Data", Dryad [Data Archive]. https://doi.org/10.5061/dryad.63xsj3v58