LOG-CONCAVITY FOR PARTITIONS WITHOUT SEQUENCES

LUKAS MAUTH

Abstract. We prove log-concavity for the function counting partitions without sequences. We use an exact formula for a mixed-mock modular form of weight zero, explicit estimates on modified Kloosterman sums and the saddle point method.

1. Introduction and statement of results

A *partition* of a non-negative integer n is a finite non-increasing sequence of positive integers which sum to n. We denote the number of partitions of n by $p(n)$. A classical tool for studying partitions is its generating function [\[1\]](#page-13-0)

$$
P(q) := \sum_{n=0}^{\infty} p(n)q^n = \prod_{n=1}^{\infty} \frac{1}{1 - q^n} = \frac{1}{(q;q)_{\infty}},
$$
\n(1.1)

where for $a \in \mathbb{C}$ and $n \in \mathbb{N} \cup \{\infty\}$ we define the q-Pochhammer symbol $(a)_n = (a;q)_n := \prod_{k=1}^{n-1}(1-aq^j)$. This form of the generating function was used by G. H. Hardy and S. Ramanujan to develop the Circle Method and led them to prove the following asymptotics [\[14\]](#page-14-0)

$$
p(n) \sim \frac{1}{4n\sqrt{3}} e^{\pi\sqrt{\frac{2n}{3}}}, \quad n \to \infty.
$$
 (1.2)

Later, H. Rademacher improved the Circle Method to obtain an exact formula for $p(n)$ [\[16\]](#page-14-1). To state this formula, define the *Kloostermann sums*

$$
A_k(n) := \sum_{h \pmod{k}} \omega_{h,k} e^{\frac{-2\pi i n h}{k}},
$$

where $*$ indicates that h only runs over those residue classes that are coprime to k and $\omega_{h,k}$ is a 24k-th root of unity defined by

$$
\omega_{h,k} := \begin{cases}\n\left(\frac{-k}{h}\right) e^{-\pi i \left(\frac{1}{4}(2-hk-h) + \frac{1}{12}\left(k-\frac{1}{k}\right)\left(2h-h'+h^2h'\right)\right)} & \text{if } h \text{ is odd,} \\
\left(\frac{-h}{k}\right) e^{-\pi i \left(\frac{1}{4}(k-1) + \frac{1}{12}\left(k-\frac{1}{k}\right)\left(2h-h'+h^2h'\right)\right)} & \text{if } k \text{ is odd,}\n\end{cases}
$$

where h' is a solution to $hh' \equiv -1 \pmod{k}$, and $\left(\frac{1}{k}\right)$ denotes the Kronecker symbol. Furthermore, let I_{κ} denote the modified Bessel function of order κ . Rademacher's exact formula for $p(n)$ is then given by

$$
p(n) = \frac{2\pi}{(24n-1)^{3/4}} \sum_{k=1}^{\infty} \frac{A_k(n)}{k} I_{\frac{3}{2}}\left(\frac{\pi\sqrt{24n-1}}{6k}\right).
$$

²⁰²⁰ Mathematics Subject Classification. 11B57, 11F03, 11F20, 11F30, 11F37, 11P82.

Key words and phrases. Circle Method, η -function, partitions.

2 LUKAS MAUTH

The first term in the sum recovers the asymptotics found by Hardy and Ramanujan. Rademacher's proof made extensive use of the fact that the generating function $P(q)$ is essentially a modular form. Continuing their study, Rademacher and H. Zuckerman showed exact formulas for the Fourier coefficients (at any cusp) of all modular forms of *negative* weight of any finite index subgroup of $SL_2(\mathbb{Z})$ [\[17,](#page-14-2) [21\]](#page-14-3).

From a combinatorial point of view it is a classical question to ask whether $p(n)$ is log-concave, where we call a sequence $\{a_n\}$ *log-concave* if it satisfies for all $n \geq 1$ the inequality

$$
a_n^2 \ge a_{n-1}a_{n+1}.
$$

Many important sequences that arise naturally in combinatorics are known to be log-concave, among them are binomial coefficients, Stirling numbers and Bessel numbers [\[19\]](#page-14-4). The first proof that $p(n)$ is log-concave for $n \geq 26$ and all even $n < 26$ was given by J.-L. Nicolas [\[12\]](#page-14-5) and was later reproved independently by S. Desalvo and I. Pak [\[8\]](#page-14-6). K. Ono, S. Pujahari and L. Rolen showed in [\[15\]](#page-14-7) that the function counting plane partitions is log-concave for all $n \geq 12$. All these proofs follow the same principle. Using the Circle Method (or an exact formula obtained by the Circle Method) one obtains strong asymptotics with explicit error bounds. The main term is easily seen to be log-concave and the difficulty lies in finding analytically a small bound after which the main term dominates the error term in the log-concavity condition. The remaining case are then checked directly.

In this paper we intend to carry out this program for the function $p_2(n)$ which is defined as the number of partitions of n that do not contain any consecutive integers as parts. These types of partitions were studied by P. MacMahon [\[11\]](#page-14-8) and arise in certain probability models and the study of threshold growth in cellular automata [\[3,](#page-14-9) [9\]](#page-14-10). G. Andrews showed [\[2\]](#page-13-1) the following formula for the generating function:

$$
G_2(q) := \sum_{n=0}^{\infty} p_2(n)q^n = \frac{(-q^3;q^3)_{\infty}}{(q^2;q^2)_{\infty}}\chi(q)
$$

where $\chi(q)$ denotes the third order mock theta function

$$
\chi(q) := \sum_{n=0}^{\infty} \frac{(-q;q)_n}{(-q^3;q^3)_n} q^{n^2}.
$$

In contrast to $p(n)$ the generating function of $p_2(n)$ is not anymore a modular form but the product of a mock theta function and a modular form with an overall weight of 0. This increases difficulty of proving asymptotics for $p_2(n)$. Bringmann and K. Mahlburg first succeeded in proving an asymptotic formula for $p_2(n)$ [\[7\]](#page-14-11) and recent work [\[5\]](#page-14-12) by W. Bridges and K. Bringmann improves their result and gives an exact formula for $p_2(n)$, and is the first example of an exact formula for the coefficients of a mixed-mock modular form. To state their result we need to introduce some notation. For $b \in \mathbb{R}, k \in \mathbb{N}$, and $\nu \in \mathbb{Z}$ define

$$
\mathcal{I}_{b,k,\nu}(n) := \int_{-1}^1 \frac{\sqrt{1-x^2} I_1\left(\frac{2\pi}{k}\sqrt{2bn(1-x^2)}\right)}{\cosh\left(\frac{\pi i}{k}\left(\nu-\frac{1}{6}\right)-\frac{\pi}{k}\sqrt{\frac{b}{3}}x\right)} dx.
$$

Moreover, for $\nu, n \in \mathbb{Z}$ and for all $k \in \mathbb{N}$ we define the following Kloosterman sums

$$
K_{k}^{[4]}(\nu; n) := \sum_{\substack{0 \le h < k \\ \gcd(h,k)=1}} \frac{\omega_{h,k}\omega_{h,\frac{k}{2}}\omega_{3h,k}}{\omega_{3h,\frac{k}{2}}} e^{\frac{\pi i}{k}(-3\nu^{2}+v)h'} e^{\frac{2\pi i n h}{k}}, \quad \gcd(k, 6) = 2,
$$
\n
$$
K_{k}^{[6]}(\nu; n) := \sum_{\substack{0 \le h < k \\ \gcd(h,k)=1}} \frac{\omega_{h,k}\omega_{2h,k}\omega_{h,\frac{k}{3}}}{\omega_{2h,\frac{k}{3}}} e^{\frac{\pi i}{k}(-3\nu^{2}+v)h'} e^{\frac{2\pi i n h}{k}}, \quad \gcd(k, 6) = 3,
$$
\n
$$
\gcd(k, 6) = 3,
$$
\n
$$
K_{k}^{[8]}(\nu; n) := \sum_{\substack{0 \le h < k \\ \gcd(h,k)=1}} \frac{\omega_{h,k}\omega_{2h,k}\omega_{3h,k}}{\omega_{6h,k}} e^{\frac{\pi i}{k}(-3\nu^{2}-v)h'} e^{\frac{2\pi i n h}{k}}, \quad \gcd(k, 6) = 1,
$$
\n
$$
K_{k}(n) := \sum_{\substack{1 \le h \le k \\ \gcd(h,k) = 1}} \frac{\omega_{h,k}\omega_{2h,k}\omega_{6h,k}}{\omega_{3h,k}^{3}} e^{-\frac{2\pi i n h}{k}}, \quad \gcd(k, 6) = 1.
$$

The exact formula for $p_2(n)$ then reads as follows,

Theorem. For $n \geq 2$ we have

$$
p_2(n) = \frac{\pi}{6\sqrt{n}} \sum_{\substack{k \geq 1 \\ \gcd(k,6)=1}} \frac{K_k(n)}{k^2} I_1\left(\frac{2\pi\sqrt{n}}{3k}\right)
$$

+
$$
\frac{\pi}{18\sqrt{6n}} \sum_{\substack{k \geq 1 \\ \gcd(k,6)=1}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \nu \pmod{k}}} (-1)^{\nu} K_k^{[8]}(\nu; n) \mathcal{I}_{\frac{1}{18},k,\nu}(n)
$$

+
$$
\frac{5\pi}{36\sqrt{6n}} \sum_{\substack{k \geq 1 \\ \gcd(k,6)=2}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \nu \pmod{k}}} (-1)^{\nu} K_k^{[4]}(\nu; n) \mathcal{I}_{\frac{5}{36},k,\nu}(n)
$$

+
$$
\frac{\pi}{6\sqrt{6n}} \sum_{\substack{k \geq 1 \\ \gcd(k,6)=3}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \nu \pmod{k}}} (-1)^{\nu} K_k^{[6]}(\nu; n) \mathcal{I}_{\frac{1}{6},k,\nu}(n).
$$
 (1.3)

We are going to use this formula to obtain strong asymptotics for $p_2(n)$ in terms of elementary functions with an explicit error term. To carry out this program we need explicit estimates for the Kloostermans sums and the integrals $\mathcal{I}_{b,k,v}$. We will derive these estimates in Section 2. It turns out that if we need a really strong estimate for $\mathcal{I}_{\frac{1}{18},1,0}(n)$. This will be done with the saddle point method in Section 3. In Section 4 we are going to prove our main theorem

Theorem. We have for $n \geq 482$ and all even $2 \leq n < 482$ that

$$
p_2^2(n) - p_2(n-1)p_2(n+1) \ge 0.
$$

ACKNOWLEDGEMENTS

The author wishes to thank Kathrin Bringmann and Walter Bridges for suggesting and supervising this project, William Craig, Johann Franke and Badri Vishal Pandey for helpful discussions concerning log-concavity problems. Furthermore, the author wishes to thank Ben Kane, Andreas Mono, Joshua Males and Caner Nazaroglu for helping with the verification of the remaining 450000 cases for log-concavity. The author recieved funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No. 101001179).

2. Estimates for Kloosterman sums and Bessel Integrals

First we are going to estimates the terms that come from $k \geq 2$ in [\(1.3\)](#page-2-0) as they are exponentially smaller than the first term. We consider two different ranges, depending on whether k is small or large. We say that k is small if $k < 2\sqrt{n}$ and we say that k is large if $k \ge 2\sqrt{n}$. We state the following premliminary results. First we need explicit upper bounds for the Bessel function $I_1(x)$ for small and large x.

Lemma 2.1. *The following are true:*

$$
I_1(x) \le \sqrt{\frac{2}{\pi x}}e^x, \quad x > 1,
$$

$$
I_1(x) \le x, \quad 0 \le x \le 1.
$$

We want to write the modified Kloosterman sums in terms of classical ones. To that end, recall the definition of the *classical Kloosterman* sum for $a, b \in \mathbb{Z}$ and $k \in \mathbb{N}$ we define

$$
K(a, b, k) = \sum_{h \pmod{k}} e^{\frac{2\pi i}{k}(ah + b[h]_k)},
$$

where $[h]_k$ denotes the inverse of h mod k. For classical Kloosterman sums we have the Weil bound, see [\[10\]](#page-14-13) for a proof. Let $\tau(n)$ denote the number of divisors of n.

Theorem 2.2 (Weil bound for Kloosterman sums). For $a, b \in \mathbb{Z}$ and $k \in \mathbb{N}$ we have

$$
|K(a,b,k)| \le \tau(n)\sqrt{\gcd(a,b,k)}k^{\frac{1}{2}}.
$$

Furthermore, it is well known that $\tau(n) \ll_{\varepsilon} n^{\varepsilon}$. The constant, depending on ε can be worked out explicitly. We take the following result from [\[13\]](#page-14-14).

Lemma 2.3. We have for all $n \geq 1$ that

$$
\tau(n) \le 576 \left(\frac{n}{21621600} \right)^{\frac{1}{4}} \le 9n^{\frac{1}{4}}.
$$

We also need the following result from [\[5\]](#page-14-12) on the multipliers of the Kloosterman sums $K_k^{[4]}$ $k^{[4]}$, $K_k^{[6]}$ and $K_k^{[8]}$ $\kappa_k^{\text{[o]}}$ to write them in terms of classical Kloosterman sums.

Lemma 2.4. *We have*

$$
\frac{\omega_{h,k}\omega_{h,\frac{k}{2}}\omega_{3h,k}}{\omega_{3h,\frac{k}{2}}} = e^{\frac{2\pi i}{k}\left(\frac{k(k+2)}{8}h - \frac{k^2+2}{18}h'\right)},
$$

$$
\frac{\omega_{h,k}\omega_{2h,k}\omega_{h,\frac{k}{3}}}{\omega_{2h,\frac{k}{3}}} = (-1)^{\frac{k+1}{2}}e^{\frac{4\pi ikh}{9} - \frac{2\pi i}{24k}(k^2-3)h'}
$$

$$
\frac{\omega_{h,k}\omega_{2h,k}\omega_{3h,k}}{\omega_{6h,k}} = (-1)^{\frac{k+1}{2}}e^{\frac{10\pi i(k^2-1)h'}{72k}}.
$$

,

We are now ready to estimate the modified Kloosterman sums appearing in (1.3) . This is an explicit version of the estimates given in [\[5\]](#page-14-12) and follows their proof closely.

Lemma 2.5. *For all* $k \geq 1$ *and* $0 \leq \nu \leq k$ *, we have*

$$
\left|K_k^{[4]}(\nu; n)\right| \le 26\sqrt{n}k^{\frac{3}{4}}, \quad \gcd(k, 6) = 2,
$$

$$
\left|K_k^{[6]}(\nu; n)\right| \le 27\sqrt{n}k^{\frac{3}{4}}, \quad \gcd(k, 6) = 3,
$$

$$
\left|K_k^{[8]}(\nu; n)\right| \le 9\sqrt{n}k^{\frac{3}{4}}, \quad \gcd(k, 6) = 1,
$$

$$
|\mathcal{K}_k(n)| \le k.
$$

Proof. Suppose $gcd(k, 6) = 2$. We then write using Lemma [2.4](#page-3-0)

$$
K_k^{[4]}(\nu;n) = \sum_{\substack{0 \le h < k \\ \gcd(h,k)=1}} e^{\frac{2\pi i}{k}} \left(\left(\frac{k(k+2)}{8} - n \right) h + \left(\frac{-k^2 + 2}{18} - \frac{3\nu^2 + \nu}{2} \right) h' \right).
$$

We now change variables $h' \mapsto -3h'$ and $h \mapsto [-1]_k[3]_k h$ to obtain

$$
K_k^{[4]}(\nu; n) = K\left([-1]_k[3]_k\left(\frac{k(k+2)}{8} - n\right), \frac{k^2 + 2}{6} + \frac{9\nu^2 + 3\nu}{2}, k\right)
$$

Using Weil's bound, Theorem [2.2,](#page-3-1) we estimate this by

$$
K_k^{[4]}(\nu; n) \le \tau(k) \sqrt{\gcd\left([-1]_k[3]_k\left(\frac{k(k+2)}{8}-n\right), \frac{k^2+2}{6}+\frac{9\nu^2+3\nu}{2}, k\right)} k^{\frac{1}{2}}.
$$

We continue to estimate

$$
\gcd\left([-1]_k[3]_k\left(\frac{k(k+2)}{8}-n\right), \frac{k^2+2}{6}+\frac{9\nu^2+3\nu}{2}, k\right) \le \gcd\left([-1]_k[3]_k\left(\frac{k(k+2)}{8}-n\right), k\right)
$$

$$
\le \gcd\left([-1]_k[3]_k 8n, k\right) = \gcd(8n, k) \le 8n.
$$

Combined with the Lemma [2.3](#page-3-2) this yields as claimed

$$
|K_k^{[4]}(\nu;n)| \leq 9k^{\frac{1}{4}}\sqrt{8n}k^{\frac{1}{2}} \leq 26\sqrt{n}k^{\frac{3}{4}}.
$$

The estimates for $K_k^{[6]}$ $k^{[6]}(\nu; n)$ and $K_k^{[8]}$ $\mathcal{K}_k(\nu; n)$ follow in the same way. The estimate for $\mathcal{K}_k(n)$ is the trivial bound. \Box

In the next step we rewrite the integrals $\mathcal{I}_{b,k,\nu}(n)$. We start with the following elementary result, whose proof is a straightforward calculation.

Lemma 2.6. *Let* $a, b \in \mathbb{R}$ *and* $x \in [0, 1]$ *. Then,*

$$
f_{a,b}(x) := \frac{1}{\cosh(a+ib)} + \frac{1}{\cosh(a-ib)} = 4 \frac{\cos(a)\cosh(bx)}{\cos(2a) + \cosh(2bx)}.
$$

In particular, $f_{a,b}$ *is a real valued function with constant sign on* [0, 1]. *Furthermore*, $|f_{a,b}(x)|$ *is monotonically decreasing on* [0, 1].

Corollary 2.7. *Let* $b \in \mathbb{R}, k \in \mathbb{N}$ *and* $\nu \in \mathbb{Z}$ *. We set* $a := \frac{\pi}{k} \left(\nu - \frac{1}{6} \right)$ $\frac{1}{6}$) and $b := \frac{\pi}{k} \sqrt{\frac{b}{3}}$ 3 . *Then, we have the estimate*

$$
|\mathcal{I}_{b,k,\nu}(n)| \leq |2\sec(a)|I_1\left(\frac{2\pi}{k}\sqrt{2bn}\right).
$$

Proof. We use the previous lemma to write

$$
\mathcal{I}_{b,k,\nu}(n) = \int_0^1 f_{a,b}(x) \sqrt{1 - x^2} I_1\left(\frac{2\pi}{k} \sqrt{2bn(1 - x^2)}\right).
$$

Recall that $|f_{a,b}(x)|$ is decreasing. Now, the result follows immediately as the maximum of the inte-
grand in absolute value is at $x = 0$ and by noting that $f_{a,b}(0) = 2 \sec(a)$. grand in absolute value is at $x = 0$ and by noting that $f_{a,b}(0) = 2 \sec(a)$.

2.1. **Estimates for large k.** In this section we always assume that $k \geq 2\sqrt{n}$.

Lemma 2.8. *Let* $n \geq 1$ *. Furthermore, let* $b \in \mathbb{R}$ *and* $\nu \in \mathbb{N}$ *. Then,*

$$
\sum_{\nu \pmod{k}} |\mathcal{I}_{b,k,\nu}(n)| \leq 32\pi \log(k)\sqrt{2bn}.
$$

Proof. By Corollary [2.7](#page-5-0) and Lemma [2.1](#page-3-3) we find that

$$
|\mathcal{I}_{b,k,\nu}(n)| \leq 2 \left| \sec \left(\frac{\pi}{k} \left(\nu - \frac{1}{6} \right) \right) \right| \frac{2\pi}{k} \sqrt{2bn}.
$$

For $x \in [0,1], x \neq \frac{\pi}{2}$ $\frac{\pi}{2}$ one has the inequality

$$
\left|\frac{1}{\cos(x)}\right| \le \frac{\pi}{2} \frac{1}{|x - \frac{\pi}{2}|}
$$

.

Using this one inequality, one sees easily that

$$
\sum_{\nu \pmod{k}} \left| \frac{1}{\cos\left(\frac{\pi}{k}\left(\nu - \frac{1}{6}\right)\right)} \right| \leq 8k \log(k),
$$

which finishes the proof immediately. \Box

This allows us to estimate the contribution from large k to the sum in [\(1.3\)](#page-2-0).

Lemma 2.9. *Let* $n \geq 1$ *. Then, we have*

$$
\frac{\pi}{6\sqrt{n}} \left| \sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=1}} \frac{K_k(n)}{k^2} I_1\left(\frac{2\pi\sqrt{n}}{3k}\right) \right|
$$
\n
$$
+ \frac{\pi}{18\sqrt{6n}} \left| \sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=1}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \gcd(k,6)=1}} (-1)^{\nu} K_k^{[8]}(\nu; n) \mathcal{I}_{\frac{1}{18},k,\nu}(n) \right|
$$
\n
$$
+ \frac{5\pi}{36\sqrt{6n}} \left| \sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=2}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \nu \pmod{k} }} (-1)^{\nu} K_k^{[4]}(\nu; n) \mathcal{I}_{\frac{5}{36},k,\nu}(n) \right|
$$
\n
$$
+ \frac{\pi}{6\sqrt{6n}} \left| \sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=3}} \frac{1}{k^2} \sum_{\substack{\nu \pmod{k} \\ \nu \pmod{k} }} (-1)^{\nu} K_k^{[6]}(\nu; n) \mathcal{I}_{\frac{1}{6},k,\nu}(n) \right| \le 25265n^{\frac{7}{8}}.
$$
\n(2.1)

Proof. We start with the first sum. By Lemma [2.1](#page-3-3) and Lemma [2.5](#page-4-0) we see

$$
\sum_{\substack{k\geq 2\sqrt{n}\\ \gcd(k,6)=1}}\frac{|\mathcal{K}_k(n)|}{k^2}\left|I_1\left(\frac{2\pi\sqrt{n}}{3k}\right)\right|\leq \frac{2\pi}{3}\sum_{k\geq \sqrt{n}}\frac{1}{k^2}\leq 2\pi\int_{\sqrt{n}}^{\infty}\frac{1}{x^2}dx=\frac{2\pi}{\sqrt{n}}\leq 2\pi n^{\frac{7}{8}}.
$$

For the second sum we find using Lemma [2.5](#page-4-0) and Lemma [2.8](#page-5-1) that

$$
\sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=1}} \frac{1}{k^2} \sum_{\nu \pmod{k}} \left| K_k^{[8]}(\nu; n) \right| \left| \mathcal{I}_{\frac{1}{18}, k, \nu}(n) \right| \le 9\sqrt{n} \sum_{k \ge 2\sqrt{n}} \frac{1}{k^{\frac{5}{4}}} \sum_{\nu \pmod{k}} \left| \mathcal{I}_{\frac{1}{18}, k, \nu}(n) \right|
$$

$$
\le 96\pi n \sum_{k \ge \sqrt{n}} \frac{\log(k)}{k^{\frac{5}{4}}} \le 300\pi n \sum_{k \ge \sqrt{n}} \frac{1}{k^{\frac{9}{8}}} \le 600\pi n \int_{\sqrt{n}}^{\infty} \frac{1}{t^{\frac{9}{8}}} dt \le 2400\pi n^{\frac{7}{8}},
$$

where we used $log(n) \leq 3n^{\frac{1}{8}}$ for all $n \geq 1$. The other sums can be estimated similarly and we obtain the bounds

$$
\sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=2}} \frac{1}{k^2} \sum_{\nu \pmod{k}} \left| K_k^{[4]}(v;n) \right| \left| \mathcal{I}_{\frac{5}{36},k,\nu}(n) \right| \le 2640 \pi n^{\frac{7}{8}},
$$

$$
\sum_{\substack{k \ge 2\sqrt{n} \\ \gcd(k,6)=3}} \frac{1}{k^2} \sum_{\nu \pmod{k}} \left| K_k^{[6]}(\nu;n) \right| \left| \mathcal{I}_{1/6,k,\nu}(n) \right| \le 3000 \pi n^{\frac{7}{8}}.
$$

Furthermore, we have that

$$
\max\left(\frac{5\pi}{36\sqrt{6n}}, \frac{\pi}{6\sqrt{6n}}, \frac{\pi}{6\sqrt{n}}, \frac{\pi}{18\sqrt{6n}}\right) = \frac{\pi}{\sqrt{6n}} \le \frac{1}{\sqrt{n}}.
$$

Combining all estimates proves the claim.

2.2. **Estimates for small k.** We assume throughout this section that $k \leq 2\sqrt{n}$. We start with the corresponding result to Lemma [2.8](#page-5-1) for small k .

Lemma 2.10. *Let* $n \geq 1$ *be fixed. Then we have*

$$
\frac{\pi}{6\sqrt{n}} \left| \sum_{\substack{2 \le k \le 2\sqrt{n} \\ \gcd(k,6)=1}} \frac{K_k(n)}{k^2} I_1\left(\frac{2\pi\sqrt{n}}{3k}\right) \right|
$$
\n
$$
+ \frac{\pi}{18\sqrt{6n}} \left| \sum_{\substack{2 \le k \le 2\sqrt{n} \\ \gcd(k,6)=1}} \frac{1}{k^2} \sum_{\nu \pmod{k}} (-1)^{\nu} K_k^{[8]}(\nu; n) \mathcal{I}_{\frac{1}{18},k,\nu}(n) \right|
$$
\n
$$
+ \frac{5\pi}{36\sqrt{6n}} \left| \sum_{\substack{2 \le k \le 2\sqrt{n} \\ \gcd(k,6)=2}} \frac{1}{k^2} \sum_{\nu \pmod{k}} (-1)^{\nu} K_k^{[4]}(\nu; n) \mathcal{I}_{\frac{5}{36},k,\nu}(n) \right|
$$
\n
$$
+ \frac{\pi}{6\sqrt{6n}} \left| \sum_{\substack{2 \le k \le 2\sqrt{n} \\ \gcd(k,6)=3}} \frac{1}{k^2} \sum_{\nu \pmod{k}} (-1)^{\nu} K_k^{[6]}(\nu; n) \mathcal{I}_{\frac{1}{6},k,\nu}(n) \right| \le 4e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}}. \tag{2.2}
$$

Proof. First of all we start by identifying the dominating term in this sum. We set $a_{\nu} = \frac{\pi}{2}$ $rac{\pi}{2}$ $\left(\nu-\frac{1}{6}\right)$ $\frac{1}{6}$. By Corollary [2.7](#page-5-0) for $k = 2$ we have

$$
\left| \mathcal{I}_{\frac{1}{18},2,\nu}(n) \right| \leq 2 |\sec(a_{\nu})| I_1 \left(\frac{\pi}{3} \sqrt{n} \right),
$$

$$
\left| \mathcal{I}_{\frac{5}{36},2,\nu}(n) \right| \leq 2 |\sec(a_{\nu})| I_1 \left(\frac{\pi \sqrt{10}}{6} \sqrt{n} \right),
$$

$$
\left| \mathcal{I}_{\frac{1}{6},2,\nu}(n) \right| \leq 2 |\sec(a_{\nu})| I_1 \left(\frac{\sqrt{3}\pi}{3} \sqrt{n} \right).
$$

By monotonicity of the Bessel function we see that the last term grows the fastest. Lemma [2.1](#page-3-3) gives

$$
I_1\left(\frac{\sqrt{3}\pi}{3}\sqrt{n}\right) \le \frac{\sqrt{2}3^{1/4}}{\pi n^{\frac{1}{4}}}e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}} \le e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}}.
$$

Combining this with the trivial estimate on the Kloosterman shows that the expression from the statement is bounded by

$$
\frac{2}{\sqrt{n}} \sum_{k=2}^{2\sqrt{n}} e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}} \le 4e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}}.
$$

 \Box

For two functions f, g we introduce the notation

$$
f = O_{\leq}(g),
$$

if $f = O(g)$ and the implicit constant can be chosen to be 1. Combing the results we obtain the following Lemma.

Lemma 2.11. *Let* $n \geq 1$ *. Then*

$$
p_2(n) = \frac{\pi}{6\sqrt{n}} I_1\left(\frac{2\pi\sqrt{n}}{3}\right) + \frac{\pi}{18\sqrt{6n}} \mathcal{I}_{\frac{1}{18},1,0}(n) + O_{\leq}\left(4e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}} + 2565n^{\frac{7}{8}}\right).
$$

3. ASYMPTOTIC FORMULA FOR $p_2(n)$ AND LOG-CONCAVITY

For proving log-concavity strong asymptotics are required. Thus, we will have to use a strong asymptotic formula for the Bessel function and $\mathcal{I}_{\frac{1}{18},1,0}(n)$. The latter is the more difficult problem. We will find such a formula with the saddle point method. We define the following list of real numbers:

$$
a_1 := \frac{1}{4\sqrt{3}}, \quad a_2 := \frac{1}{18\sqrt{2}}, \quad a_3 := -\frac{3\sqrt{3}}{64\pi}, \quad a_4 := -\frac{324 + 5\pi^2}{3888\sqrt{2}\pi}, \quad a_5 := -\frac{45\sqrt{3}}{2048\pi^2} \quad a_6 := \frac{1080 + 17\pi^2}{186624\sqrt{2}},
$$

$$
a_7 := -\frac{945\sqrt{3}}{32768\pi^3}, \quad a_8 := -\frac{349920 + 33048\pi^2 + 455\pi^4}{40310784\sqrt{2}\pi}, \quad a_9 := -\frac{127575\sqrt{3}}{2097152\pi^4}.
$$

Theorem 3.1. *Let* $n \geq 1$ *. Then,*

$$
p_2(n) = \left(\frac{a_1}{n^{\frac{3}{4}}} + \frac{a_2}{n} + \frac{a_3}{n^{\frac{5}{4}}} + \frac{a_4}{n^{\frac{3}{2}}} + \frac{a_5}{n^{\frac{7}{4}}} + \frac{a_6}{n^{\frac{7}{4}}} + \frac{a_7}{n^{\frac{9}{4}}} + \frac{a_8}{n^{\frac{5}{4}}} + \frac{a_9}{n^{\frac{11}{2}}} + \frac{a_9}{n^{\frac{11}{4}}} + O_{\leq}\left(16\frac{e^{\frac{2\pi}{3}\sqrt{n}}}{n^3}\right). \tag{3.1}
$$

Proof. Define

$$
a_k(\nu) := (-1)^k \frac{\left(\frac{1}{2} - \nu\right)_n \left(\frac{1}{2} + \nu\right)_n}{2^k k!}, \quad k = 0, 1, 2, \dots
$$

$$
(\lambda)_n := \frac{\Gamma(\lambda + n)}{\Gamma(\lambda)}.
$$

We have for $\nu \in \mathbb{N}$ the well-known asymptotic formulas (see [\[20\]](#page-14-15)),

$$
I_{\nu}(z) \sim \frac{e^z}{\sqrt{2\pi z}} \sum_{k=0}^{\infty} (-1)^k \frac{a_k(\nu)}{z^k}.
$$

In [\[4\]](#page-14-16) K. Banerjee worked out explicit error bounds in the asymptotic formulas for the Bessel functions of non-negative order. More explicitly, we have in our cases

$$
\frac{\pi}{6\sqrt{n}}I_1\left(\frac{2\pi\sqrt{n}}{3}\right) = \left(\frac{1}{4\sqrt{3}n^{\frac{3}{4}}}-\frac{3\sqrt{3}}{64\pi n^{\frac{5}{4}}}-\frac{45\sqrt{3}}{2048\pi^2 n^{\frac{7}{4}}}-\frac{945\sqrt{3}}{32768\pi^3 n^{\frac{9}{4}}}-\frac{127575\sqrt{3}}{2097152\pi^4 n^{\frac{11}{4}}}\right)e^{\frac{2\pi}{3}\sqrt{n}} + O_{\leq}\left(\frac{0.0795351}{n^{\frac{13}{4}}}e^{\frac{2\pi}{3}\sqrt{n}}\right).
$$
\n(3.2)

By Lemma [2.6](#page-4-1) and since the integrand is an even function, we can write with $a = -\frac{\pi}{6}$ $\frac{\pi}{6}$ and $b=0$ that

$$
\mathcal{I}_{\frac{1}{18},1,0}(n) = \frac{1}{2} \int_{-1}^{1} f_{a,b}(x) \sqrt{1 - x^2} e^{\frac{2\pi}{3} \sqrt{n(1-x^2)}} dx.
$$

We now cut off the integral at $\frac{1}{8}$ and insert the expansion of I_1 up to order 3 and obtain

$$
\frac{\pi}{18\sqrt{6n}} \mathcal{I}_{\frac{1}{18},1,0}(n) = \frac{1}{72\sqrt{2n^{\frac{3}{4}}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{\frac{1}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx \n- \frac{1}{128\sqrt{2\pi n^{\frac{5}{4}}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{-\frac{1}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx \n- \frac{15}{4096\sqrt{2\pi^2 n^{\frac{7}{4}}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{-\frac{3}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx \n- \frac{315}{65536\sqrt{2\pi^3 n^{\frac{9}{4}}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{-\frac{5}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx \n+ O \leq \left(\frac{27}{160\sqrt{2\pi^4 n^{\frac{11}{4}}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{-\frac{7}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx \right) \n+ O \leq \left(\frac{1}{2} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x) (1-x^2)^{\frac{1}{4}} e^{\frac{\sqrt{7}\pi}{4}\sqrt{n}} dx \right).
$$
\n(3.3)

We will now use the saddle point method to expand each of these terms into a power series. We illustrate this process for the first term. We introduce the notation $A := \frac{2\pi}{3}\sqrt{n}$, $f(x) := f_{a,b}(x)(1-x^2)^{\frac{1}{4}}$
and $g(x) = \sqrt{1-x^2}$. Then, we can write

$$
\int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^2)^{\frac{1}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}} dx = \int_{-\frac{1}{8}}^{\frac{1}{8}} f(x) e^{Ag(x)} dx = \frac{1}{\sqrt{A}} \int_{-\frac{1}{8}\sqrt{A}}^{\frac{1}{8}\sqrt{A}} f\left(\frac{y}{\sqrt{A}}\right) e^{Ag\left(\frac{y}{\sqrt{A}}\right)}. (3.4)
$$

The functions $g(x)$ and $f(x)$ are analytic on the interval $\left[-\frac{1}{8}\right]$ $\frac{1}{8}, \frac{1}{8}$ $\frac{1}{8}$ and we can expand them into power series

$$
Ag\left(\frac{y}{\sqrt{A}}\right) = A - \frac{y^2}{2} - \frac{y^4}{8A} - \frac{y^6}{16A^2} - \frac{5y^8}{128A^3} - \frac{7y^{10}}{256A^4} - \frac{21y^{12}}{1024A^5} + O \le \left(j\frac{y^{14}}{A^6}\right)
$$

\n
$$
e^{Ag\left(\frac{y}{\sqrt{A}}\right)} = e^A e^{-\frac{y^2}{2}} \left(1 - \frac{y^4}{8A} - \frac{y^6}{16A^2} - \frac{5y^8}{128A^3} - \frac{7y^{10}}{256A^4} - \frac{21y^{12}}{1024A^5} + \frac{jy^{14}}{A6}\right)
$$

\n
$$
+ \frac{1}{2} \left(-\frac{y^4}{8A} - \frac{y^6}{16A^2} - \frac{5y^8}{128A^3} - \frac{7y^{10}}{256A^4} - \frac{21y^{12}}{1024A^5} + \frac{jy^{14}}{A^6}\right)^2
$$

\n
$$
+ \frac{1}{6} \left(-\frac{y^4}{8A} - \frac{y^6}{16A^2} - \frac{5y^8}{128A^3} - \frac{7y^{10}}{256A^4} - \frac{21y^{12}}{1024A^5} + \frac{jy^{14}}{A^6}\right)^3
$$

\n
$$
+ O \le \left(k \left(-\frac{y^4}{8A} - \frac{y^6}{16A^2} - \frac{5y^8}{128A^3} - \frac{7y^{10}}{256A^4} - \frac{21y^{12}}{1024A^5} + \frac{jy^{14}}{A^6}\right)^4\right),
$$

$$
f\left(\frac{y}{\sqrt{A}}\right) = f(0)\left(1 - \frac{y^2}{4A} - \frac{5\pi^2y^2}{324A} - \frac{3y^4}{32A^2} + \frac{5\pi^2y^4}{1296A^2} + \frac{17\pi^4y^4}{69984A^2} - \frac{7y^6}{128A^3} + \frac{5\pi^2y^6}{3456A^3} - \frac{17\pi^4y^6}{279936A^3} - \frac{91\pi^6y^6}{22674816A^3} - \frac{77y^8}{2048A^4} + \frac{35\pi^2y^8}{41472A^4} - \frac{17\pi^4y^8}{746496A^4} + \frac{91\pi^6y^8}{90699264A^4} + \frac{207913\pi^8y^8}{3085588961280A^4} + \frac{\sqrt{3}ly^9}{4A^{\frac{9}{2}}}\right),
$$

where j, k and ℓ are positive constants given by

$$
j = \max_{x \in \left[-\frac{1}{8}, \frac{1}{8}\right]} \left| \frac{d^{14}}{dx^{14}} \frac{g(x)}{14!} \right| \le 0.0504,
$$

\n
$$
k = \max_{x \in \left[-\frac{1}{8}, \frac{1}{8}\right]} \left| \frac{d^2}{dx^2} \frac{e^x}{2} \right| = \frac{e^{\frac{1}{8}}}{2},
$$

\n
$$
\ell = \max_{x \in \left[-\frac{1}{8}, \frac{1}{8}\right]} \left| \frac{d^{10}}{dx^{10}} \frac{f(x)}{10!} \right| \le 0.109.
$$

We now multiply out the series expansions in the integrand and obtain

$$
f\left(\frac{y}{\sqrt{A}}\right)e^{Ag\left(\frac{y}{\sqrt{A}}\right)} = T(y, A) + O_{\leq}(R(y, A)),
$$

where we have the main term

$$
T(y, A) = 1 + \frac{-162y^2 - 10\pi^2y^2 - 81y^4}{648A} + \frac{-26244y^4 + 1080\pi^2y^4 + 68\pi^4y^4 - 8748y^6 + 540\pi^2y^6 + 2187y^8}{279936A^2} + \frac{-9920232y^6 + 262440\pi^2y^6 - 11016\pi^4y^6 - 728\pi^6y^6 - 2125764y^8 + 87480\pi^2y^8}{181398528A^3} + \frac{-5508\pi^4y^8 + 1062882y^{10} - 21870\pi^2y^{10} - 59049y^{12}}{181398528A^3}
$$

and the error term

$$
R(y, A) = \sum_{n=4}^{N} \frac{P_n(y)}{A^n},
$$

for some $N > 0$ and even polynomials $P_n(y)$. We now extend the range of integration in [\(3.4\)](#page-9-0) to the whole real line and obtain

$$
\frac{1}{\sqrt{A}} \int_{-\frac{1}{8}\sqrt{A}}^{\frac{1}{8}\sqrt{A}} f\left(\frac{y}{\sqrt{A}}\right) e^{Ag\left(\frac{y}{\sqrt{A}}\right)} dy = \frac{f(0)e^A}{\sqrt{A}} \int_{-\infty}^{\infty} e^{-\frac{y^2}{2}} T(y, A) dy
$$

+ $O \le \left(\frac{f(0)e^A}{\sqrt{A}} \int_{-\frac{1}{8}\sqrt{A}}^{\frac{1}{8}\sqrt{A}} e^{-\frac{y^2}{2}} (T(y, A) + E(y, A)) dy + \frac{f(0)e^A}{\sqrt{A}} \int_{-\infty}^{\infty} e^{-\frac{y^2}{2}} E(y, A) dy\right).$ (3.5)

We now reduced the integrals to Gaussian integrals which we can evaluate. Performing these integrations we find that for $n \geq 435000$

$$
\frac{1}{72\sqrt{2}n^{\frac{3}{4}}}\int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^2)^{\frac{1}{4}}e^{\frac{2\pi}{3}\sqrt{n(1-x^2)}}dx
$$

$$
= \left(\frac{1}{18\sqrt{2}} \cdot \frac{1}{n} - \frac{5(81+2\pi^2)}{7776\sqrt{2}\pi} \cdot \frac{1}{n^{\frac{3}{2}}} + \frac{6561+3780\pi^2+68\pi^4}{746496\sqrt{2}\pi^2} \cdot \frac{1}{n^2} - \frac{5(-1240029+503010\pi^2+49572\pi^4+728\pi^6)}{322486272\sqrt{2}\pi^3} \cdot \frac{1}{n^{\frac{5}{2}}}\right)e^{\frac{2\pi}{3}\sqrt{n}} + O \le \left(\frac{90\sqrt{\frac{2}{\pi}}}{n^3}e^{\frac{2\pi}{3}\sqrt{n}}\right).
$$

We apply the exact same method to other integrals and obtain for $n \geq 435000$

$$
\frac{1}{128\sqrt{2}\pi n^{\frac{5}{4}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^{2})^{-\frac{1}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^{2})}} dx
$$
\n
$$
= \left(\frac{1}{32\sqrt{2}\pi} \cdot \frac{1}{n^{3/2}} - \frac{81+10\pi^{2}}{13824\sqrt{2}\pi^{2}} \cdot \frac{1}{n^{2}} + \frac{-10935+1620\pi^{2}+68\pi^{4}}{1327104\sqrt{2}\pi^{3}} \cdot \frac{1}{n^{5/2}}\right) e^{\frac{2\pi}{3}\sqrt{n}}
$$
\n
$$
+ O_{\leq}\left(\frac{5}{64\pi^{3/2}n^{3}} e^{\frac{2\pi}{3}\sqrt{n}}\right),
$$
\n
$$
\frac{15}{4096\sqrt{2}\pi^{2}n^{\frac{7}{4}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^{2})^{-\frac{3}{4}} e^{\frac{2\pi}{3}\sqrt{n(1-x^{2})}} dx
$$
\n
$$
= \left(\frac{15}{1024\sqrt{2}\pi^{2}} \cdot \frac{1}{n^{2}} + \frac{5\left(243-10\pi^{2}\right)}{147456\sqrt{2}\pi^{3}}\right) \cdot \frac{1}{n^{5/2}} e^{\frac{2\pi}{3}\sqrt{n}} + O_{\leq}\left(\frac{75}{2048\pi^{5/2}n^{3}} e^{\frac{2\pi}{3}\sqrt{n}}\right),
$$
\n
$$
\frac{315}{65536\sqrt{2}\pi^{3}n^{\frac{9}{4}}} \int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^{2})^{-\frac{5}{4}} e^{\frac{2\pi}{3}\sqrt{n}(1-x^{2})} dx
$$
\n
$$
= \frac{315}{16384\sqrt{2}\pi^{3}} \cdot \frac{1}{n^{5/2}} e^{\frac{2\pi}{3}\sqrt{n}} + O_{\leq}\left(\frac{1575}{32768n^{3}\pi^{7/2}n^{3}} e^{\frac{2\pi}{3}\sqrt{n}}\right),
$$
\n
$$
\frac
$$

Adding up all these terms we obtain for $n \geq 45000$ the asymptotic formula

$$
p_2(n) = \left(\frac{a_1}{n^{\frac{3}{4}}} + \frac{a_2}{n} + \frac{a_3}{n^{\frac{5}{4}}} + \frac{a_4}{n^{\frac{3}{2}}} + \frac{a_5}{n^{\frac{7}{4}}} + \frac{a_6}{n^2} + \frac{a_7}{n^{\frac{9}{4}}} + \frac{a_8}{n^{\frac{5}{2}}} + \frac{a_9}{n^{\frac{11}{4}}}\right)e^{\frac{2\pi}{3}\sqrt{n}}
$$

+
$$
O \le \left(\frac{55296\sqrt{2} + 7875\sqrt{\pi} + 6000\pi^{3/2} + 12800\pi^{5/2} + 204800\pi^{7/2}}{163840\pi^4 n^3} + 0.00795351\left(\frac{1}{n}\right)^{13/4} + \frac{27\left(\frac{1}{n}\right)^{7/2}}{16\pi^{9/2}}\right)e^{\frac{2\pi}{3}\sqrt{n}} + O \le \left(\frac{1}{2}\int_{-\frac{1}{8}}^{\frac{1}{8}} f_{a,b}(x)(1-x^2)^{\frac{1}{4}}e^{\frac{\sqrt{7}\pi}{4}\sqrt{n}} dx\right) + O \le \left(4e^{\frac{\sqrt{3}\pi}{3}\sqrt{n}} + 2565n^{\frac{7}{8}}\right).
$$

An easy calculation shows that the error term can be bounded by

$$
O_{\leq}\left(\frac{1}{n^3}e^{\frac{2\pi}{3}\sqrt{n}}\right)
$$

for all $n \geq 435000$. If we call the main term of our asymptotic formula $G(n)$, then we find

$$
\max_{1 \le n \le 435000} \frac{n^3 \left| p_2(n) - G(n) \right|}{e^{\frac{2\pi}{3}\sqrt{n}}} \le 16. \tag{3.6}
$$

This concludes the proof. \Box

Theorem 3.2. We have for $n \geq 482$ and all even $2 \leq n < 482$, that

$$
p_2^2(n) - p_2(n-1)p_2(n+1) \ge 0.
$$

Proof. We write that the asymptotic formula from Theorem [3.1](#page-8-0) in the form

$$
p_2(n) = \mathcal{P}(n) + O_{\leq}(\mathcal{E}(n)).
$$

Then, we estimate

$$
p_2^2(n) - p_2(n-1)p_2(n+1) \geq (\mathcal{P}(n) - \mathcal{E}(n))^2 - (\mathcal{P}(n-1) + \mathcal{E}(n-1))(\mathcal{P}(n+1) + \mathcal{E}(n+1)).
$$

Using Taylor expansion we write this as

$$
\frac{\pi}{216\sqrt{6}} \cdot \frac{1}{n^{\frac{13}{4}}} + \frac{\pi}{288n^3} - E(n),
$$

where

$$
\begin{split} |E(n)| &\leq \frac{6.055140198378155\times 10^{-6}}{n^{\frac{35}{2}}}+\frac{0.00535139}{n^{\frac{69}{4}}}+\frac{1.18221}{n^{17}}+\frac{0.0141338}{n^{\frac{67}{4}}}+\frac{0.000646658}{n^{\frac{33}{2}}} \\ &+\frac{0.000254429}{n^{\frac{65}{4}}}+\frac{0.000742507}{n^{16}}+\frac{0.000588099}{n^{\frac{63}{4}}}+\frac{0.00321441}{n^{\frac{31}{2}}}+\frac{0.0232749}{n^{\frac{61}{4}}}+\frac{4.30165}{n^{15}} \\ &+\frac{0.0726311}{n^{\frac{59}{4}}}+\frac{0.00242495}{n^{\frac{59}{2}}}+\frac{0.000930963}{n^{\frac{57}{4}}}+\frac{0.002270314}{n^{14}}+\frac{0.00214691}{n^{\frac{55}{4}}}+\frac{0.0116994}{n^{\frac{27}{2}}} \\ &+\frac{0.0938434}{n^{\frac{53}{4}}}+\frac{17.6703}{n^{13}}+\frac{0.288354}{n^{\frac{54}{4}}}+\frac{0.0100125}{n^{\frac{25}{2}}}+\frac{0.00381031}{n^{\frac{49}{4}}}+\frac{0.0111939}{n^{\frac{47}{4}}}+\frac{0.113621}{n^{\frac{47}{4}}} \\ &+\frac{23.2169}{n^{\frac{23}{4}}}+\frac{0.447083}{n^{\frac{45}{4}}}+\frac{25.0755}{n^{11}}+\frac{0.62135}{n^{\frac{43}{4}}}+\frac{0.0296742}{n^{\frac{21}{4}}}+\frac{0.0169767}{n^{\frac{41}{4}}}+\frac{0.0785117}{n^{10}} \\ &+\frac{0.278106}{n^{\frac{39}{4}}}+\frac{42.2773}{n^{\frac{19}{
$$

We conclude that

$$
\frac{\pi}{216\sqrt{6}} \cdot \frac{1}{n^{\frac{13}{4}}} + \frac{\pi}{288n^3} - E(n) > 0
$$

for $n \geq 5092$. The finitely many remaining cases are verified directly.

Remark. (i) In [\[7\]](#page-14-11) Bringmann and Mahlburg obtained the asymptotic formula

$$
p_2(n) \sim \left(\frac{1}{4\sqrt{3n^{\frac{3}{4}}}} + \frac{1}{18\sqrt{2}n}\right) e^{\frac{2\pi}{3}\sqrt{n}}.
$$

However, this formula is not strong enough to prove log-concavity. Indeed, if one expands the logconcavity condition with abstract coefficients one sees that an asymptotic formula with precision to at least order $n^{-\frac{9}{4}}$ is necessary. We chose to expand up to order $n^{-\frac{11}{4}}$ to reduce the number of finitely many cases we have to check directly.

(ii) By looking at the quotient (3.6) for large values of n it seems that the optimal implicit constant can be chosen as $C \approx 0.69$, while we can optimally show for large n

$$
C \approx \frac{55296\sqrt{2} + 7875\sqrt{\pi} + 6000\pi^{3/2} + 12800\pi^{5/2} + 204800\pi^{7/2}}{163840\pi^4} \approx 0.727135.
$$

Furthermore, the quotient (3.6) is always larger for odd values of n, then for even values of n, which is in line with the fact that log-concavity fails for small odd n .

REFERENCES

- [1] G. Andrews, The Theory of Partitions, Encycl. Math. and Its Appl., vol. 2, G.-C. Rota ed., Addison-Wesley, Reading, 1976 (Reissued: Cambridge University Press, Cambridge 1998).
- [2] G. Andrews, Partitions with short sequences and mock theta functions, Proc. Nat. Acad. Sci. 102, 2005

- [3] G. Andrews, H. Eriksson, F. Petrov and D. Romik, *Integrals, partitions and MacMahon's theorem*, J. Comb. Theory (A) 114, 2007.
- [4] K. Banerjee, Inequalities for the modified Bessel function of first kind of non-negative order, pre-print, 2022.
- [5] W. Bridges and K. Bringmann, A Rademacher-type exact formula for partitions without sequences, [arXiv:2304.01377,](http://arxiv.org/abs/2304.01377) 2023.
- [6] K. Bringmann, B. Kane, L. Rolen and Z. Tripp, Fractional partitions and conjectures of Chern–Fu–Tang and Heim–Neuhauser, Transactions of the American Mathematical Society Series B, 2021.
- [7] K. Bringmann and K. Mahlburg, An extension of the Hardy-Ramanujan Circle Method and applications to partitions without sequences, American Journal of Mathematics 133, 2011.
- [8] S. Desalvo and I. Pak, *Log-concavity of the partition function*, Ramanujan J. **38**, 2015.
- [9] A. Holroyd, T. Liggett and D. Romik, Integrals, partitions, and cellular automata, Trans. Amer. Math. Soc. 356, 2004.
- [10] H. Iwaniec and E. Kowalski, Analytic Number Theory, American Mathematical Society, 2004.
- [11] P. MacMahon, Combinatory Analysis, Cambridge University Press, 1916.
- [12] J.-L. Nicolas, Sur les entiers N pour lesquels il y a beaucoup de groupes abéliens d'ordre N, Ann. Inst. Fourier 28, 1978.
- [13] J.-L. Nicolas, On highly composite numbers, In Ramanujan Revisited, Proceedings of the Centenary Conference, University of Illinois at Urbana-Champaign, 1987.
- [14] G. Hardy and S. Ramanujan, Asymptotic formulae in combinatory analysis, Proc. London Math. Soc. Ser. 2 17, 1918, 75–115.
- [15] K. Ono, S. Pujahari and L. Rolen *Turán inequalities for the plane partition function*, Advances in Mathematics Volume 409, Part B, 2022
- [16] H. Rademacher, A convergent series for the partition function $p(n)$, PNAS February 1, 1937 23 (2), 78–84.
- [17] H. Rademacher and H. Zuckermann, On the Fourier Coefficients of Certain Modular Forms of Positive Dimension, Annals of Mathematics, Second Series, Vol. 39, No. 2, 1938.
- [18] S. Ramanujan, *Collected Papers*, Cambridge University Press, Cambridge, 1927; reprinted by Chelsea, New York, 1962; reprinted by American Mathematical Society, Providence, 2000.
- [19] R. Stanley, Log-concave and unimodal sequences in algebra, combinatorics, and geometry, Ann. New York Acad. Sci. 576, 500–535, 1989
- [20] N. Temme, Asymptotic Methods for Integrals, World Scientific, 2014.
- [21] H. Zuckerman, On the coefficients of certain modular forms belonging to subgroups of the modular group, Trans. Amer. Math. Soc. 45, 1939.

Department of Mathematics and Computer Science, Division of Mathematics, University of Cologne, Weyertal 86-90, 50931 Cologne, Germany

Email address: lmauth@uni-koeln.de