Quasi-stationary distributions for time-changed symmetric α -stable processes killed upon hitting zero

Zhe-Kang Fang * Yong-Hua Mao * Tao Wang *

Abstract

For a time-changed symmetric α -stable process killed upon hitting zero, under the condition of entrance from infinity, we prove the existence and uniqueness of quasi-stationary distribution (QSD). The exponential convergence to the QSD from any initial distribution is proved under conditions on transition densities.

Keywords and phrases: Quasi-stationary distribution; stable process; time-change; entrance at infinity; ground state.

Mathematics Subject classification(2020): 60G52 60F99

1 Introduction and main results

Quasi-stationary distribution (QSD in short) is a good measurement to describe the long-time behavior of the absorbing Markov process when the process is conditioned to survive. Many efforts were made to study the existence, uniqueness, the domains of attraction of QSDs and the convergence rate to a QSD for various Markov processes, cf. [6, 10, 11] for Markov chains, [3, 6, 15] for diffusion processes, and [12, 13, 16, 22] for general Markov processes under some additional conditions.

In this paper, we will study QSDs for time-changed symmetric stable processes killed upon hitting zero. Let $X := (X_t)_{t \ge 0}$ be a symmetric α -stable process on \mathbb{R} with generator $\Delta^{\alpha/2} := -(-\Delta)^{\alpha/2}$, $\alpha \in (1, 2)$, where $-(-\Delta)^{\alpha/2}$ is the fractional Laplacian. Consider the following stochastic differential equation:

$$dY_t = \sigma(Y_{t-}) \, dX_t, \tag{1.1}$$

where σ is a strictly positive continuous function on \mathbb{R} . By [7, Proposition 2.1], there is a unique weak solution $Y = (Y_t)_{t \ge 0}$ to the SDE (1.1), and Y can also be expressed as a time-changed process $Y_t := X_{\zeta_t}$, where

$$\zeta_t := \inf\left\{s > 0 : \int_0^s \sigma\left(X_u\right)^{-\alpha} \, \mathrm{d}u > t\right\}.$$

^{*}Laboratory of Mathematics and Complex Systems(Ministry of Education), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, P.R. China.

By [18, Remark 43.12], the process is pointwise recurrent. Suppose $\mu(dx) := \sigma(x)^{-\alpha} dx$ is a probability measure.

Let $T_0 = \inf\{t > 0 : Y_t = 0\}$, $T_{\infty} = \lim_{R \to +\infty} T_{(-R,R)^c}$, and $T_A = \inf\{t > 0, Y_t \in A\}$, for any Borel subset $A \subseteq \mathbb{R}$. According to [14, Theorem 2.3] and the strong Markov property, a direct calculation (see Appendix for more details) leads to the conclusion that for any $x \neq 0$, $\mathbb{P}_x[T_0 < T_{\infty}] = 1$, which means the processes are almost-surely absorbed by 0. Let Y^0 be the (sub-)process of Y killed upon 0, with transition function

 $P_t^0(x, A) = \mathbb{P}_x[Y_t \in A, t < T_0], \text{ for any } x \in \mathbb{R}^0 := \mathbb{R} \setminus \{0\} \text{ and } A \in \mathscr{B}(\mathbb{R}^0).$

We call a probability measure ν is a QSD for Y^0 if for any $t \ge 0$ and any $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\mathbb{P}_{\nu}[Y_t \in A | t < T_0] = \frac{\mathbb{P}_{\nu}[Y_t \in A, t < T_0]}{\mathbb{P}_{\nu}[t < T_0]} = \nu(A),$$

where $\mathbb{P}_{\nu}(\cdot) = \int_{\mathbb{R}^0} \mathbb{P}_x(\cdot)\nu(\mathrm{d}x).$

Döring and Kyprianou [7] studied the entrance and exit from infinity for this process and Wang [25] studied the exponential and strong ergodicity. The main purpose of this paper is to study the QSD for time-changed α -stable processes killed upon zero. We will consider the existence and uniqueness of QSD, Yaglom limit, domain of attraction and the speed of convergence to the QSD.

In this paper, we will assume the following condition always holds:

$$I^{\sigma,\alpha} := \int_{\mathbb{R}} \sigma(x)^{-\alpha} |x|^{\alpha-1} \mathrm{d}x < \infty.$$
(1.2) I

By [25, Theorem 1.4], (1.2) is equivalent to strong ergodicity for Y; meanwhile, according to [7, Table 2], (1.2) holds if and only if $\pm \infty$ are entrance from infinity.

Denote by $L^2(\mathbb{R}^0, \mu)$ the space of square integrable measurable functions on \mathbb{R}^0 with respect to μ . We first present the result on compactness for killed transition semigroup $(P_t^0)_{t>0}$ under the above condition.

compact Theorem 1.1. If $I^{\sigma,\alpha} < \infty$, then $(P_t^0)_{t>0}$ is compact on $L^2(\mathbb{R}^0;\mu)$.

Under entrance from infinity (1.2), by Theorem 1.1, there exists a complete orthonormal set of eigenfunctions $\{\psi_n\}_{n\geq 0}$ with $\|\psi_n\|_{L^2(\mathbb{R}^0;\mu)} = 1$, such that $P_t^0\psi_n = e^{-\lambda_n t}\psi_n$ for any $n\geq 0$ and $t\geq 0$, where $\{\lambda_n\}_{n\geq 0}$ are eigenvalues of generator of $(P_t^0)_{t\geq 0}$, such that $0<\lambda_0<\lambda_1\leq\cdots\to+\infty$, where the positivity and simplicity of λ_0 will be proved in Appendix, Proposition A.2. The principal eigenfunction ψ_0 is called the **ground state**. By using the ground state ψ_0 , we can state our results on QSDs.

do<->ent Theorem 1.2. If
$$I^{\sigma,\alpha} < \infty$$
, then Y^0 has a unique QSD:

$$\nu(A) = \frac{\int_A \psi_0 \, \mathrm{d}\mu}{\int_{\mathbb{R}^0} \psi_0 \, \mathrm{d}\mu}, \ A \in \mathscr{B}(\mathbb{R}^0).$$
(1.3) Inu

Furthermore, ν is a Yaglom limit of Y^0 , that is for any $x \in \mathbb{R}^0$ and any subset $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\lim_{t \to \infty} \mathbb{P}_x[Y_t \in A | t < T_0] = \nu(A).$$

Moreover, there exists $0 < C(x) < \infty$, such that

$$\|\mathbb{P}_x[Y_t \in \cdot | t < T_0] - \nu\|_{TV} \le C(x)e^{-(\lambda_1 - \lambda_0)t}, \qquad (1.4) \quad \text{Yaglom-exp}$$

where $\|\eta\|_{TV} := \sup_{|f| \leq 1} |\eta(f)|$ denotes the total variation of a signed measure η .

Next, under some additional assumptions of transition density functions, we can prove ν attracts all probability measures η on \mathbb{R}^0 , and the exponential convergence in total variation.

do_con

eg

Theorem 1.3. Assume $I^{\sigma,\alpha} < \infty$. Let

$$p_t^0(x,y) := \frac{\mathrm{d}P_t^0(x,\cdot)}{\mathrm{d}\mu}(y)$$

If for any $r_0 > 0$, $p_t^0(x, y)$ satisfies that $\sup_{x \in [-r_0, r_0] \setminus \{0\}} p_2^0(x, x) < +\infty$, then ν attracts all probability measures η on \mathbb{R}^0 , that is, for any subset $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\lim_{t \to \infty} \mathbb{P}_{\eta}[Y_t \in A | t < T_0] = \nu(A).$$
(1.5) [con_ordin]

Furthermore, if $\sup_x p_2^0(x,x) < +\infty$, then for any probability measure η on \mathbb{R}^0 ,

$$\|\mathbb{P}_{\eta}[Y_t \in \cdot | t < T_0] - \nu(\cdot)\|_{TV} \le \frac{Ce^{-(\lambda_1 - \lambda_0)t} \|\eta - \nu\|_{TV}}{\eta(\psi_0)}, \tag{1.6}$$

where C is a constant independent of η .

Remark 1.4. The Lyapunov function condition is also an important method for proving the existence and uniqueness of QSD, and we refer the reader to [12, Theorem 2.2] for the recent result. Note that the Lyapunov function condition [12, C3] implies

$$\lim_{n \to \infty} \sup_{x \in \mathbb{R}} \mathbb{E}_x[T_{[-n,n]}] = 0,$$

which is equivalent to entrance from infinity by Lemma 3.2 in this paper.

Note that when σ is a polynomial, the conditions in above theorem can be explicitly characterized, and we obtain the following conclusion.

Example 1.5. Consider the polynomial case: $\sigma(x) = \left(\frac{2}{\alpha\gamma-1}\right)^{1/\alpha} (1+|x|)^{\gamma}$ (where $\left(\frac{2}{\alpha\gamma-1}\right)^{1/\alpha}$ is the normalizing constant so that the μ is a probability measure). When $\gamma > 1$, Y^0 has a unique QSD ν given by (1.3), ν is the Yaglom limit of Y^0 and (1.4) holds. Furthermore, 1.6 holds if and only if $\gamma > 1$.

An outline of this paper is as follows. In Section 2, we recall some basic notions and properties about killed processes and Green functions. In section 3, we prove the compactness of the semigroup of killed processes (Theorem 1.1), and the strict positivity, continuity and boundedness of ground state, which plays an important role in subsequent proofs. In section 4, we prove our main results on QSDs (Theorem 1.2, Theorem 1.3 and Example 1.5), and give some corollaries.

2 Killed processes and Green functions

Recall that X is a one-dimensional symmetric α -stable process with $\alpha \in (1, 2)$, and Y is its time-changed process. According to [9, Theorem 6.2.1], it is known that Y is μ -symmetric. From [5, Section 1], the Dirichlet form $(\mathscr{E}, \mathscr{F})$ associated with Y is given by

$$\mathscr{E}(f,g) = \frac{1}{2} \int_{\mathbb{R}} \int_{\mathbb{R}} (f(x) - f(y))(g(x) - g(y)) \frac{C_{\alpha} \mathrm{d}x \mathrm{d}y}{|x - y|^{1 + \alpha}},$$

and

$$\mathscr{F} = \{ f \in L^2(\mu) : \mathscr{E}(f, f) < \infty \},\$$

where $C_{\alpha} = \frac{\alpha 2^{\alpha-1} \Gamma((\alpha+1)/2)}{\sqrt{\pi} \Gamma(1-\alpha/2)}$.

Given an open subset $B \subseteq \mathbb{R}$, let Y^B be the killed (sub-)process of Y killed upon exiting B with transition function

$$P_t^B(x, A) = \mathbb{P}_x[Y_t \in A, t < T_{B^c}], \text{ for any } x \in B \text{ and } A \in \mathscr{B}(B).$$

Define

$$\mathscr{F}^B = \{ f \in \mathscr{F}, \tilde{f} = 0, \text{ q.e. on } B^c \}, \quad \mathscr{E}^B = \mathscr{E} \text{ on } \mathscr{F}^B \times \mathscr{F}^B,$$

where q.e. stands for quasi-everywhere, and \tilde{f} is a quasi-continuous modification of f (cf. [17, Section 2.2]). By [17, Theorem 3.5.7], $(\mathscr{E}^B, \mathscr{F}^B)$ is a regular Dirichlet form on $L^2(B; \mu)$ associated with Y^B ; Y^B is symmetric with respect to the measure $\mu|_B(dx)$ (where $L^2(B, \mu)$ be the space of square integrable measurable functions on B with respect to μ). We write $(\mathcal{L}^B, \mathcal{D}(\mathcal{L}^B))$ for the infinitesimal generator of (P_t^B) in $L^2(B; \mu)$, with

$$\mathcal{D}\left(\mathcal{L}^B\right) := \left\{ u \in L^2(B;\mu) : \lim_{t \to 0} \frac{P_t^B u - u}{t} \text{ exists in } L^2(B;\mu) \right\},\,$$

and

$$\mathcal{L}^B f = \lim_{t \to 0} \frac{P_t^B f - f}{t}$$
 in $L^2(B; \mu)$, for any $f \in \mathcal{D}(\mathcal{L}^B)$.

Let

$$G^B(x, \mathrm{d}y) = \int_0^{+\infty} P_t^B(x, \mathrm{d}y) dt$$

be the Green potential measure and denote the Green operator by

$$G^B f(x) = \int_B f(y) G^B(x, \mathrm{d}y).$$

Simultaneously, the killed process X^B and its Green potential measure $G_X^B(x, dy)$ are defined similarly. Let $G_X^B(x, y)$ be the Green function of X^B , that is, for any $x, y \in B$, $G_X^B(x, dy) = G_X^B(x, y) dy$. According to [25, Section 2], it should be pointed out here that the Green operator of Y^B has a strong relationship with the Green operator of X^B :

$$G^{B}f(x) = \int_{B} f(y)G^{B}_{X}(x,y)\sigma(y)^{-\alpha}\mathrm{d}y, \ \forall f \in \mathscr{B}(\mathbb{R}).$$

$$(2.1) \quad \underline{\mathsf{Y}}_{X}$$

On some occasions, the Green function of X^B can be expressed explicitly, for example:

(1) [2, Lemma 4] $B = \mathbb{R}^0$: for any $x, y \neq 0$,

$$G_X^0(x,y) := G_X^{\{0\}^c}(x,y) = \frac{\omega_\alpha}{2} \left(|y|^{\alpha-1} + |x|^{\alpha-1} - |y-x|^{\alpha-1} \right), \qquad (2.2) \quad \boxed{\mathbf{G}_{\mathbf{X}}^{\mathbf{X}} \mathbf{0}}$$

where $\omega_{\alpha} = -\frac{1}{\cos(\pi \alpha/2)\Gamma(\alpha)}$. By direct calculation, we have (see [25, Page 9])

$$G_X^0(x,y) \le \omega_\alpha(|x|^{\alpha-1} \land |y|^{\alpha-1}).$$
(2.3) [esti for GO]

(2) [8, (11)] $B = \mathbb{R} \setminus [-1, 1]^c$: for any $x, y \in [-1, 1]^c$,

$$G_X^{[-1,1]^c}(x,y) = c_\alpha \left(|x-y|^{\alpha-1} h\left(\frac{|xy-1|}{|x-y|}\right) - (\alpha-1)h(x)h(y) \right),$$
(2.4) **G_X^1c**

where $c_{\alpha} = 2^{1-\alpha} / (\Gamma(\alpha/2)^2)$, and

$$h(x) = \int_{1}^{|x|} \left(z^2 - 1\right)^{\frac{\alpha}{2} - 1} dz$$

Furthermore, according to [8, Lemma 3.3],

$$\lim_{x \to \infty} G_X^{[-1,1]^c}(x,y) = K_{\alpha}h(y), \qquad (2.5) \quad \boxed{\mathbf{G}_X^{1}\mathrm{clim}}$$

where K_{α} is a constant and defined by

$$K_{\alpha} = \frac{2c_{\alpha}(1-\frac{\alpha}{2})\Gamma(\frac{\alpha}{2})}{\Gamma(1-\frac{\alpha}{2})} \int_{1}^{\infty} \frac{h'(v)}{1+v} \mathrm{d}v < \infty.$$

Besides, thanks to the self-similarity of X, for any R > 0,

$$G_X^{[-R,R]^c}(x,y) = R^{\alpha-1} G_X^{[-1,1]^c}(\frac{x}{R},\frac{y}{R}).$$
(2.6) sel_1

3 Compactness of killed semigroups and properties of the ground states

Let $P_t^0 := P_t^{\{0\}^c}$ be the semigroup of Y killed upon hitting 0. Denote by $Y^0 := Y^{\{0\}^c}$, $\mathcal{L}^0 := \mathcal{L}^{\{0\}^c}$ and $G^0 := G^{\{0\}^c}$ the killed process, the generator of P_t^0 , and the Green operator on \mathbb{R}^0 respectively. In this section, we first prove that under the condition $I^{\sigma,\alpha} < \infty$, $(P_t^0)_{t>0}$ is compact; therefore, we can study the properties of ground state, which is crucial to our proofs of main results on QSDs.

Firstly, we prove Theorem 1.1 and G^0 is a Hilbert-Schmidt operator on $L^2(\mathbb{R}^0;\mu)$ under the condition $I^{\sigma,\alpha} < \infty$.

Proof of Theorem 1.1. By [23, Theorem 0.3.9], to demonstrate $(P_t^0)_{t>0}$ is compact, we just need to prove 0 is belong to resolvent set $\rho(\mathcal{L}^0)$ of \mathcal{L}^0 and G^0 is compact.

Firstly, we prove $0 \in \rho(\mathcal{L}^0)$, that is, the inverse of $-\mathcal{L}^0$, $G^0 = (-\mathcal{L}^0)^{-1}$ is bounded in the operator norm from $L^2(\mathbb{R}^0;\mu)$ to $L^2(\mathbb{R}^0;\mu)$:

$$||G^0||_{L^2(\mathbb{R}^0;\mu)\to L^2(\mathbb{R}^0;\mu)} < \infty.$$

Note that by (2.3),

$$\begin{aligned} \|G^0 f\|_{L^1(\mathbb{R}^0;\mu)} &= \int_{\mathbb{R}^0} \int_{\mathbb{R}^0} G^0(x,y) f(y) \mu(\mathrm{d}y) \mu(\mathrm{d}x) \\ &\leq \int_{\mathbb{R}^0} \int_{\mathbb{R}^0} \omega_\alpha |x|^{\alpha-1} f(y) \mu(\mathrm{d}y) \mu(\mathrm{d}x) \\ &= \omega_\alpha I^{\sigma,\alpha} \|f\|_{L^1(\mathbb{R}^0;\mu)} .xa \end{aligned}$$

Therefore, $||G^0||_{L^1(\mathbb{R}^0;\mu)\to L^1(\mathbb{R}^0;\mu)} \leq \omega_{\alpha} I^{\sigma,\alpha} < \infty$, and

$$\|G^{0}\|_{L^{\infty}(\mathbb{R}^{0};\mu)\to L^{\infty}(\mathbb{R}^{0};\mu)} = \|G^{0}\|_{L^{1}(\mathbb{R}^{0};\mu)\to L^{1}(\mathbb{R}^{0};\mu)} < \infty.$$

From the Riesz-Thorin theorem, it follows that $\|G^0\|_{L^2(\mathbb{R}^0;\mu)\to L^2(\mathbb{R}^0;\mu)} < \infty$. By the definition of resolvent, $0 \in \rho(\mathcal{L}^0)$.

Next, we show that G^0 is a Hilbert-Schmidt operator, which implies that G^0 is compact.

Note that

$$\int_{\mathbb{R}^{0}} \int_{\mathbb{R}^{0}} (G^{0}(x,y))^{2} \mu(\mathrm{d}x)\mu(\mathrm{d}y) \leq \int_{\mathbb{R}^{0}} \int_{\mathbb{R}^{0}} \omega_{\alpha}^{2} (|x|^{\alpha-1} \wedge |y|^{\alpha-1})^{2} \mu(\mathrm{d}x)\mu(\mathrm{d}y) \\
\leq \omega_{\alpha}^{2} \int_{\mathbb{R}^{0}} |x|^{\alpha-1} \mu(\mathrm{d}x)\omega_{\alpha} \int_{\mathbb{R}^{0}} |y|^{\alpha-1} \mu(\mathrm{d}y) = (\omega_{\alpha} I^{\sigma,\alpha})^{2} < \infty,$$
(3.1)

therefore $G^0(x,y) \in L^2(\mathbb{R}^0 \times \mathbb{R}^0; \mu \times \mu)$, and hence G^0 is a Hilbert-Schmidt operator.

Theorem 1.1 indicates that there exists a complete orthonormal set of eigenfunctions $\{\psi_n\}_{n\geq 0}$ with $\|\psi_n\|_{L^2(\mathbb{R}^0;\mu)} = 1$, such that $P_t^0\psi_n = e^{-\lambda_n t}\psi_n$ for any $n\geq 0$ and $t\geq 0$, where $\{\lambda_n\}_{n\geq 0}$ are eigenvalues of generator of $(P_t^0)_{t\geq 0}$, satisfying $0 < \lambda_0 < \lambda_1 \leq \cdots \rightarrow +\infty$, where the positivity and simplicity of λ_0 will be proved in Appendix, Proposition A.2. The principal eigenfunction ψ_0 is called the **ground state**. Next, we are going to prove some basic properties about ψ_0 . We prove the positivity and continuity of ψ_0 on \mathbb{R}^0 in the following theorem.

pos_con Theorem 3.1. If $I^{\sigma,\alpha} < \infty$, then ψ_0 can be chosen to be strictly positive and continuous on \mathbb{R}^0 .

Proof. Firstly, we prove ψ_0 can be chosen to be nonnegative.

Actually, by using the proof by contradiction, we have $\mu(\{x : \psi_0(x) \ge 0\}) = 1$ or $\mu(\{x : \psi_0(x) \le 0\}) = 1$. Assume that $\mu(\{x : \psi_0(x) > 0\}) > 0$ and $\mu(\{x : \psi_0(x) < 0\}) > 0$. Then $\text{Leb}(\{(x, y) : \psi_0(x)\psi_0(y) < 0\}) > 0$, where $\text{Leb}(\cdot)$ is the Lebesgue measure. Note that on $\{(x, y) : \psi_0(x)\psi_0(y) \ge 0\}$,

$$(\psi_0(x) - \psi_0(y))^2 = (|\psi_0(x)| - |\psi_0(y)|)^2,$$

while on $\{(x, y) : \psi_0(x)\psi_0(y) < 0\},\$

$$(\psi_0(x) - \psi_0(y))^2 > (|\psi_0(x)| - |\psi_0(y)|)^2.$$

Therefore,

$$\begin{aligned} \mathscr{E}(\psi_0,\psi_0) &= \frac{1}{2} \left(\int_{\{(x,y):\psi_0(x)\psi_0(y)\ge 0\}} + \int_{\{(x,y):\psi_0(x)\psi_0(y)< 0\}} \right) (\psi_0(x) - \psi_0(y))^2 \frac{C_\alpha \mathrm{d}x\mathrm{d}y}{|x-y|^{1+\alpha}} \\ &> \frac{1}{2} \left(\int_{\{(x,y):\psi_0(x)\psi_0(y)\ge 0\}} + \int_{\{(x,y):\psi_0(x)\psi_0(y)< 0\}} \right) (|\psi_0(x)| - |\psi_0(y)|)^2 \frac{C_\alpha \mathrm{d}x\mathrm{d}y}{|x-y|^{1+\alpha}} \\ &= \mathscr{E}(|\psi_0|, |\psi_0|). \end{aligned}$$

(3.2) inf contradio

However, by the definition of ground state and the first Dirichlet eigenvalue,

$$\mathscr{E}(\psi_0, \psi_0) = \lambda_0 = \inf\{\mathscr{E}(f, f) : \mu(f^2) = 1, f(0) = 0, f \in \mathscr{F}\} \le \mathscr{E}(|\psi_0|, |\psi_0|),$$

which contradicts (3.2). Thus $\mu\{x : \psi_0(x) > 0\} = 0$ or $\mu\{x : \psi_0(x) < 0\} = 0$. If $\mu\{x : \psi_0(x) > 0\} = 0$, then $-\psi_0$ satisfies $P_t^0(-\psi_0) = e^{-\lambda_0 t}(-\psi_0)$ and $\mu\{x : -\psi_0(x) < 0\} = 0$. So we can always choose eigenfunction ψ_0 satisfying

$$\mu(\{x:\psi_0(x)\ge 0\})=1. \tag{3.3} \ |\texttt{psi_0<0=0}|$$

Note that by the definition of ground state, it is easy to see that

$$G^{0}\psi_{0}(x) = \int_{0}^{\infty} P_{t}^{0}\psi_{0}(x)dt = \int_{0}^{\infty} e^{-\lambda_{0}t}\psi_{0}(x)dt = \frac{\psi_{0}(x)}{\lambda_{0}}.$$

Thus by combining the above equality, (2.1), and (3.3), we get that for any $x \neq 0$,

$$\frac{\psi_0(x)}{\lambda_0} = \int_{\mathbb{R}^0} G_X^0(x, y) \psi_0(y) \mu(\mathrm{d}y) \ge 0, \qquad (3.4) \quad \text{dom con thm}$$

which proves ψ_0 is nonnegative.

Next, we prove ψ_0 is strictly positive.

Actually,

$$G_X^0(x,y) > 0$$
 for any $x, y \neq 0$. (3.5) positive Gree

According to (2.2), by noting that $G_X^0(x,y) = G_X^0(-x,-y)$, we just need to prove that for any x > 0, $y \neq 0$, $G_X^0(x,y) > 0$. Indeed, $G_X^0(x,\cdot)$ is strictly decreasing in $(-\infty,0)$, $(x,+\infty)$ and strictly increasing in (0,x). Since for any y > x, by [25, Page 592],

$$G_X^0(x,y) > \frac{\omega_\alpha}{2} (x \wedge y)^{\alpha - 1} = \frac{\omega_\alpha}{2} x^{\alpha - 1} > 0,$$

and $G_X^0(x,0) = 0$, then $G_X^0(x,y) > 0$ for any $y \neq 0$, x > 0.

Now, if there would exist $x_0 \neq 0$ such that $\psi_0(x_0) = 0$, then by (3.4),

$$0 = \frac{\psi_0(x_0)}{\lambda_0} = \int_{\mathbb{R}^0} G_X^0(x_0, y) \psi_0(y) \mu(\mathrm{d}y).$$

By (3.5), and the non-negativity of ψ_0 , we would have $\psi_0 = 0$, μ -a.e., which is contradictory to $\|\psi_0\|_{L^2(\mathbb{R}^0;\mu)} = 1$. So ψ_0 is strictly positive on \mathbb{R}^0 .

Finally, to prove the continuity of ψ_0 on \mathbb{R}^0 , w.l.o.g., we assume $x_0 > 0$. According to (2.3), for any $|x - x_0| < 1 \wedge \frac{x_0}{2}$,

$$\int_{\mathbb{R}^0} G_X^0(x,y)\psi_0(y)\mu(\mathrm{d}y) \le \omega_\alpha \int_{\mathbb{R}^0} (|x| \wedge |y|)^{\alpha-1}\psi_0(y)\mu(\mathrm{d}y) \le \omega_\alpha (x_0+1)^{\alpha-1}\mu(\psi_0) < \infty.$$

By using (3.4), the continuity of $G(\cdot, y)$ on \mathbb{R}^0 for any $y \neq 0$ and dominated convergence theorem, we have thus proved Theorem 3.1.

Next, to prove the boundedness of ψ_0 , we need some lemmas.

equ_con Lemma 3.2. $I^{\sigma,\alpha} := \int_{\mathbb{R}} \sigma(x)^{-\alpha} |x|^{\alpha-1} dx < \infty$ if and only if

$$\lim_{R \to \infty} \sup_{x \in \mathbb{R}} \mathbb{E}_x[T_{[-R,R]}] = 0.$$

Proof. If $I^{\sigma,\alpha} < +\infty$, from (2.1), (2.3),

$$\sup_{x} \mathbb{E}_{x}[T_{[-R,R]}] = \sup_{x \in [-R,R]^{c}} \int_{[-R,R]^{c}} G^{[-R,R]^{c}}(x, \mathrm{d}y)$$

$$= \sup_{x \in [-R,R]^{c}} \int_{[-R,R]^{c}} G^{[-R,R]^{c}}_{X}(x,y)\mu(\mathrm{d}y)$$

$$\leq \sup_{x \in [-R,R]^{c}} \int_{[-R,R]^{c}} G^{0}_{X}(x,y)\mu(\mathrm{d}y)$$

$$\leq \omega_{\alpha} \int_{[-R,R]^{c}} |y|^{\alpha-1}\mu(\mathrm{d}y),$$

By letting $R \to +\infty$, we obtain that $\sup_x \mathbb{E}_x[T_{[-R,R]}] \to 0$.

On the contrary, if $I^{\sigma,\alpha} = \infty$, then for any R > 0, by using (2.4), (2.5) and (2.6),

$$\sup_{x} \mathbb{E}_{x}[T_{[-R,R]}] = \sup_{x \in [-R,R]^{c}} \int_{[-R,R]^{c}} R^{\alpha-1} G_{X}^{[-1,1]^{c}} \left(\frac{x}{R}, \frac{y}{R}\right) \mu(\mathrm{d}y)$$

$$\geq \int_{[-R,R]^{c}} \liminf_{x \to +\infty} R^{\alpha-1} G_{X}^{[-1,1]^{c}} \left(\frac{x}{R}, \frac{y}{R}\right) \mu(\mathrm{d}y)$$

$$= \int_{[-R,R]^{c}} R^{\alpha-1} K_{\alpha} h\left(\frac{y}{R}\right) \mu(\mathrm{d}y)$$

$$\geq \frac{K_{\alpha}}{\alpha} \int_{[-R,R]^{c}} R^{\alpha-1} \left(\frac{|y|^{\alpha-1}}{R^{\alpha-1}} - 1\right) \mu(\mathrm{d}y) = +\infty.$$

equ_rem Remark 3.3. According to the proof of [25, Theorem 1.4], it is easy to verify that $I^{\sigma,\alpha} < +\infty$ is equivalent to $\sup_{x \in \mathbb{R}} \mathbb{E}_x[T_0] < +\infty$; now by Proposition 3.2, we know $I^{\sigma,\alpha} < +\infty$ is also equivalent to $\sup_{x \in \mathbb{R}} \mathbb{E}_x[T_{[-1,1]}] < +\infty$.

bou_lem Lemma 3.4. If $I^{\sigma,\alpha} < \infty$, then for any $\lambda > 0$, there exists a constant $R = R(\lambda)$, such that

$$\sup_{x\in\mathbb{R}^0}\mathbb{E}_x[e^{\lambda T_{[-R,R]}}]<+\infty.$$

Proof. It follows from Lemma 3.2 that, for any $\epsilon > 0$, there exists $R = R(\epsilon)$, such that

$$\sup_{x} \mathbb{E}_x[T_{[-R,R]}] \le \epsilon.$$

Using Markov's inequality, we have

$$\sup_{x} \mathbb{P}_x[T_{[-R,R]} > 1] \le \epsilon.$$

By Markov property, for any $n \ge 2$,

$$\mathbb{P}_{x}\left(T_{[-R,R]} > n\right) = \mathbb{E}_{x}\left[\mathbf{1}_{\{T_{[-R,R]} > 1\}}\mathbf{1}_{\{T_{[-R,R]} \circ \theta_{1} > n-1\}}\right]$$
$$= \mathbb{E}_{x}\left[\mathbf{1}_{\{T_{[-R,R]} > 1\}}\mathbb{E}_{X_{1}}\left[\mathbf{1}_{\{T_{[-R,R]} > n-1\}}\right]\right]$$
$$\leq \mathbb{P}_{x}\left(T_{[-R,R]} > 1\right)\sup_{x}\mathbb{P}_{x}\left(T_{[-R,R]} > n-1\right).$$

Then by induction,

$$\sup_{x} \mathbb{P}_x[T_{[-R,R]} > n] \le \epsilon^n.$$

By Fubini Theorem, it comes to the fact that for any $\lambda > 0$, we can take $\epsilon < e^{-\lambda}$ and $R = R(\epsilon)$ such that for any $x \in \mathbb{R}^0$,

$$\mathbb{E}_{x}[e^{\lambda T_{[-R,R]}}] = \int_{0}^{+\infty} \lambda e^{\lambda s} \mathbb{P}_{x}[T_{[-R,R]} > s] dt \leq \sum_{i=0}^{+\infty} \lambda e^{\lambda(i+1)} \mathbb{P}_{x}[T_{[-R,R]} > i] + 1$$
$$\leq \sum_{i=0}^{+\infty} \lambda e^{\lambda(i+1)} \epsilon^{i} + 1 \leq \frac{\lambda e^{\lambda}}{1 - e^{\lambda} \epsilon} + 1 < +\infty.$$

Using Lemma 3.4 and following the proof of [20, Lemma 5.3 and Theorem 5.4], we obtain the following corollary.

bou Corollary 3.5. If $I^{\sigma,\alpha} < \infty$, then ψ_0 is bounded.

4 Proof of the main results on QSD

We verify the main results on QSD in this section. Firstly, we prove Theorem 1.2, which shows the existence and uniqueness of the QSD for Y^0 , the existence of Yaglom limit and the exponential convergence to Yaglom limit when starting at a single point $x \neq 0$. Secondly, we provide a sufficient condition for exponential convergence to the QSD for any initial distribution (Theorem 1.3). Finally, we focus on Example 1.5, which indicates the condition $I^{\sigma,\alpha} < +\infty$ is a sufficient and necessary condition for uniform exponential convergence on some occasions.

First of all, we prove Theorem 1.2.

Proof of Theorem 1.2. (1) Firstly, we prove

$$\nu(\mathrm{d}x) = \frac{\psi_0(x)\mu(\mathrm{d}x)}{\int_{\mathbb{R}^0}\psi_0(x)\mu(\mathrm{d}x)}.$$

is a QSD for Y^0 .

Since $\psi_0 \in L^2(\mathbb{R}^0, \mu)$ and μ is a finite measure, $\psi_0 \in L^1(\mathbb{R}^0, \mu)$. According to [9, Lemma 4.1.3] and the μ -symmetry of Y, we know that Y^0 is μ -symmetric. By using Theorem 1.1 and Theorem 3.1, we get that for all $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\begin{aligned} \mathbb{P}_{\nu}[Y_t \in A | t < T_0] &= \frac{\int_{\mathbb{R}^0} \mathbb{P}_x[Y_t \in A, t < T_0]\nu(\mathrm{d}x)}{\int_{\mathbb{R}^0} \mathbb{P}_x[t < T_0]\nu(\mathrm{d}x)} \\ &= \frac{\int_{\mathbb{R}^0} P_t^0 \mathbf{1}_A(x)\psi_0(x)\mu(\mathrm{d}x)}{\int_{\mathbb{R}^0} P_t^0 \mathbf{1}_(x)\psi_0(x)\mu(\mathrm{d}x)} \\ &= \frac{\int_{\mathbb{R}^0} P_t^0 \psi_0(x)\mathbf{1}_A(x)\mu(\mathrm{d}x)}{\int_{\mathbb{R}^0} P_t^0 \psi_0(x)\mathbf{1}_A(x)\mu(\mathrm{d}x)} \\ &= \frac{\int_{\mathbb{R}^0} e^{-\lambda_0 t}\psi_0(x)\mathbf{1}_A(x)\mu(\mathrm{d}x)}{\int_{\mathbb{R}^0} e^{-\lambda_0 t}\psi_0(x)\mathbf{1}_A(x)\mu(\mathrm{d}x)} \\ &= \nu(A). \end{aligned}$$

It comes to the conclusion that ν is a QSD for Y^0 .

(2) Secondly, we turn to prove the uniqueness of the QSD. Assume η is also a QSD for Y^0 , and there exists $A \in \mathscr{B}(\mathbb{R}^0)$ such that $\eta(A) > \nu(A)$.

From the proof of [24, Proposition 3.5], for any $x \in \mathbb{R}$ and t > 0, $P_t(x, dy)$ has a density function $p_t(x, y)$ with respect to μ . Using [4, (3)], we have for any $x \in \mathbb{R}^0$ and t > 0, $P_t^0(x, dy)$ has a density function $p_t^0(x, y)$ with respect to μ satisfying

$$P_t^0(x, \mathrm{d}y) = p_t^0(x, y)\mu(\mathrm{d}y), \qquad p_t^0(x, y) = p_t^0(y, x), \text{ for any } x, y \neq 0.$$

From [6, Theorem 2.2], there exists $\lambda > 0$, such that for any $A \in \mathbb{R}^0$,

$$\mathbb{P}_{\eta}[Y_t \in A, t < T_0] = e^{-\lambda t} \eta(A).$$

It follows that

$$\int_{\mathbb{R}^0} p_t^0(x, y) \mathbf{1}_A(y) \mu(\mathrm{d}y) \eta(\mathrm{d}x) = e^{-\lambda t} \eta(A).$$

Then η is absolutely continuous with respect to the Lebesgue measure. Let $\eta(dx) = \eta(x)dx$.

By using Doob-*h* transform, we define Y^{ψ_0} with the transition semigroup as follows:

$$P_t^{\psi_0} f = e^{\lambda_0 t} \frac{P_t^0(\psi_0 f)}{\psi_0}, \quad \text{for any } t > 0.$$

By definition of symmetry and conservativity, we know that Y^{ψ_0} is $\psi_0^2 \mu$ -symmetric and conservative. Then we prove Y^{ψ_0} is irreducible. Indeed, let $A \subseteq \mathbb{R}^0$ be a $P_t^{\psi_0}$ -invariant measurable set satisfying $\psi_0^2 \mu(A) > 0$, according to [9, Lemma 1.6.1], $P_t^{\psi_0} \mathbf{1}_A(x) = 0 \ \psi_0^2 \mu$ a.e. on $\mathbb{R}^0 \setminus A$ for any t > 0. Therefore, $P_t^0(\psi_0 \mathbf{1}_A)(x) = 0 \ \psi_0^2 \mu$ -a.e. on $\mathbb{R}^0 \setminus A$ for any t > 0, which yields that $G^0(\psi_0 \mathbf{1}_A)(x) = 0 \ \psi_0^2 \mu$ -a.e. on $\mathbb{R}^0 \setminus A$. Since $G^0(x, y) > 0$ for any $x \neq 0$ and $y \neq 0$, $G^0(\psi_0 \mathbf{1}_A)(x) > 0$ for any $x \neq 0$. So $\psi_0^2 \mu(\mathbb{R}^0 \setminus A) = 0$, which means Y^{ψ_0} is irreducible. Note that for any compact subsets $K, F \subseteq \mathbb{R}^0$ and any t > 0,

$$\eta(K) = \frac{\mathbb{P}_{\eta}[Y_t \in K, t < T_0]}{\mathbb{P}_{\eta}[t < T_0]} \le \frac{\mathbb{P}_{\eta}[Y_t \in K, t < T_0]}{\mathbb{P}_{\eta}[Y_t \in F, t < T_0]} = \frac{\int_{\mathbb{R}^0} P_t^0 \mathbf{1}_K(x)\eta(\mathrm{d}x)}{\int_{\mathbb{R}^0} P_t^0 \mathbf{1}_F(x)\eta(\mathrm{d}x)}$$

$$= \frac{\int_{\mathbb{R}^0} \psi_0(x) P_t^{\psi_0}(\frac{\mathbf{1}_K}{\psi_0})(x)\eta(\mathrm{d}x)}{\int_{\mathbb{R}^0} \psi_0(x) P_t^{\psi_0}(\frac{\mathbf{1}_F}{\psi_0})(x)\eta(\mathrm{d}x)}.$$
(4.1) [etaK]

Note that by the continuity and positivity of ψ_0 ,

$$\left|\frac{\mathbf{1}_K}{\psi_0}\right| \le \frac{1}{\inf_{x \in K} \psi_0(x)} < \infty.$$

Combining this with the irreducibility and conservativity of Y^{ψ_0} , [21, Theorem 2.2] is valid, so

$$P_t^{\psi_0}\left(\frac{\mathbf{1}_K}{\psi_0}\right) \to \frac{1}{\mu(\mathbb{R}^0)} \int_K \psi_0 \mathrm{d}\mu, \text{ a.e., as } t \to +\infty,$$

and

$$P_t^{\psi_0}\left(\frac{\mathbf{1}_F}{\psi_0}\right) \to \frac{1}{\mu(\mathbb{R}^0)} \int_F \psi_0 \mathrm{d}\mu, \text{ a.e., as } t \to +\infty.$$

Therefore, by dominated convergence theorem, let $t \to \infty$ in (4.1), we get that

$$\eta(K) \le \frac{\int_K \psi_0(x)\mu(\mathrm{d}x)}{\int_F \psi_0(x)\mu(\mathrm{d}x)}.$$

Then for any $x \neq 0$, by taking $K = [x, x + \Delta x]$ and $F \uparrow \mathbb{R}^0$, we arrive at

$$\eta(x) = \lim_{\Delta x \to 0} \frac{\eta([x, x + \Delta x])}{\Delta x} \le \lim_{\Delta x \to 0} \frac{\int_{[x, x + \Delta x]} \psi_0(x) \mu(\mathrm{d}x)}{\Delta x \mu(\psi_0)} = \frac{\psi_0(x) \sigma(x)^{-\alpha}}{\mu(\psi_0)}$$

It follows that for any $A \in \mathscr{B}(\mathbb{R}^0)$, $\eta(A) \leq \nu(A)$, which is a contradiction. Thus ν is the unique QSD.

(3) Thirdly, we turn to the proof of the existence of Yaglom limit of Y^0 . Using Theorem 1.1 and following the proof of [16, Corollary 24], we have for any $A \in \mathscr{B}(\mathbb{R}^0)$ and any t > 2,

$$\begin{split} \|e^{\lambda_{0}t}P_{t}^{0}\mathbf{1}_{A}-\langle\psi_{0},\mathbf{1}_{A}\rangle\psi_{0}\|_{L^{2}(\mathbb{R}^{0};\mu)}^{2} \\ &=\|\sum_{n=1}^{+\infty}e^{(\lambda_{0}-\lambda_{n})t}\langle\psi_{n},\mathbf{1}_{A}\rangle\psi_{n}\|_{L^{2}(\mathbb{R}^{0};\mu)}^{2} =\sum_{n=1}^{+\infty}e^{2(\lambda_{0}-\lambda_{n})t}\langle\psi_{n},\mathbf{1}_{A}\rangle^{2} \\ &\leq e^{2(\lambda_{0}-\lambda_{1})(t-1)}e^{2\lambda_{0}}\sum_{n=0}^{+\infty}e^{-2\lambda_{n}}\langle\psi_{n},\mathbf{1}_{A}\rangle^{2} \\ &=e^{-2(\lambda_{1}-\lambda_{0})(t-1)}e^{2\lambda_{0}}\|P_{1}^{0}\mathbf{1}_{A}\|_{L^{2}(\mathbb{R}^{0};\mu)}^{2}. \end{split}$$

By using Hölder inequality, we obtain

$$\begin{aligned} |e^{\lambda_0(t-1)} \mathbb{P}_x[Y_t \in A, t < T_0] - \langle \psi_0, \mathbf{1}_A \rangle e^{-\lambda_0} \psi_0(x)| \\ &= |e^{\lambda_0(t-1)} \langle P_{t-1}^0 \mathbf{1}_A, p_1^0(x, \cdot) \rangle - \langle \psi_0, \mathbf{1}_A \rangle \langle \psi_0, p_1^0(x, \cdot) \rangle| \\ &\leq e^{-(\lambda_1 - \lambda_0)(t-2)} e^{\lambda_0} (p_2^0(x, x))^{\frac{1}{2}} \to 0 \text{ as } t \to +\infty, \end{aligned}$$

$$(4.2) \quad \texttt{lim_t3}$$

which yields that

$$\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_x[Y_t \in A, t < T_0] = \psi_0(x) \langle \psi_0, \mathbf{1}_A \rangle, \qquad (4.3) \quad \texttt{lim_t1}$$

and

$$\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_x[t < T_0] = \psi_0(x) \langle \psi_0, \mathbf{1} \rangle.$$
(4.4)
$$[\texttt{lim_t2}]$$

Therefore,

$$\lim_{t \to \infty} \mathbb{P}_x[Y_t \in A | t < T_0] = \frac{\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_x[Y_t \in A, t < T_0]}{\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_x[t < T_0]} = \nu(A).$$

Hence ν is the Yaglom limit of Y^0 .

(4) Finally, we prove the exponential convergence to the Yaglom limit when starting at a single point $x \neq 0$. Fix $x \neq 0$. According to the positivity (Theorem 3.1), boundedness (Corollary 3.5) of ψ_0 , and (4.4),

$$0 < m_x := \inf_{t>1} e^{\lambda_0(t-1)} \mathbb{P}_x[t < T_0] \le \sup_{t>1} e^{\lambda_0(t-1)} \mathbb{P}_x[t < T_0] < \infty.$$

It follows from (4.2) that

$$\begin{split} &|\mathbb{P}_{x}[Y_{t} \in A|t < T_{0}] - \nu(A)| \\ &= \left| \frac{e^{\lambda_{0}(t-1)} \mathbb{P}_{x}[Y_{t} \in A, t < T_{0}]}{e^{\lambda_{0}(t-1)} \mathbb{P}_{x}[t < T_{0}]} - \frac{e^{-\lambda_{0}} \psi_{0}(x) \langle \psi_{0}, \mathbf{1}_{A} \rangle}{e^{-\lambda_{0}} \psi_{0}(x) \langle \psi_{0}, \mathbf{1}_{A} \rangle} \right| \\ &\leq \frac{\psi_{0}(x) \langle \psi_{0}, \mathbf{1} \rangle |e^{\lambda_{0}(t-1)} \mathbb{P}_{x}[Y_{t} \in A, t < T_{0}] - e^{-\lambda_{0}} \psi_{0}(x) \langle \psi_{0}, \mathbf{1}_{A} \rangle |}{m_{x} \psi_{0}(x) \langle \psi_{0}, \mathbf{1} \rangle} \\ &+ \frac{\psi_{0}(x) \langle \psi_{0}, \mathbf{1}_{A} \rangle |e^{-\lambda_{0}} \psi_{0}(x) \langle \psi_{0}, \mathbf{1} \rangle - e^{\lambda_{0}(t-1)} \mathbb{P}_{x}[t < T_{0}]|}{m_{x} \psi_{0}(x) \langle \psi_{0}, \mathbf{1} \rangle} \\ &\leq \frac{2e^{-(\lambda_{1}-\lambda_{0})(t-2)} e^{\lambda_{0}} (p_{2}^{0}(x, x))^{\frac{1}{2}}}{m_{x}}. \end{split}$$
(4.5)

Since

$$\|\mathbb{P}_{x}[Y_{t} \in \cdot | t < T_{0}] - \nu\|_{TV} = 2 \sup_{A \in \mathscr{B}(\mathbb{R}^{0})} |\mathbb{P}_{x}[Y_{t} \in A | t < T_{0}] - \nu(A)|,$$

we arrive at (1.4) by choosing

$$C(x) = \frac{4e^{2\lambda_1}(p_2^0(x,x))^{\frac{1}{2}}}{m_x}$$

in (4.5). Hence we finish the proof of Theorem 1.2.

Next, we consider the problem about the domain of attraction of QSD and the speed of convergence. We prove Theorem 1.3 as follows. The idea of the proof benefits from [26, Theorem 4.3] and [22, Proof of Corollary 2.2.4].

Proof of Theorem 1.3. Firstly, we prove the result about the domain of attraction. We assume that $I^{\sigma,\alpha} < +\infty$ and for any $r_0 > 0$, $\sup_{x \in [-r_0, r_0] \setminus \{0\}} p_2^0(x, x) < +\infty$.

By taking $\lambda = \lambda_0$ in Lemma 3.4, there exists $R_0 > 0$ such that

$$B_1 := \sup_{x \in \mathbb{R}^0} \mathbb{E}_x[e^{\lambda_0 T_{[-R_0, R_0]}}] < +\infty.$$

Combining Corollary 3.5 and (4.2), we have

$$B_2 := \sup_{t>2} \sup_{x \in [-R_0, R_0]} e^{\lambda_0 t} \mathbb{P}_x[T_0 > t] < +\infty.$$

Then for any $x > R_0$, using strong Markov property, we get that

$$\begin{split} \mathbb{P}_{x}[T_{0} > t] &= \mathbb{P}_{x}[T_{[-R_{0},R_{0}]} > t] + \mathbb{P}_{x}[T_{[-R_{0},R_{0}]} \leq t, T_{0} > t] \\ &\leq e^{-\lambda_{0}t} \mathbb{E}_{x}[e^{\lambda_{0}T_{[-R_{0},R_{0}]}}] + \mathbb{E}_{x} \left[\mathbb{P}_{x}[T_{[-R_{0},R_{0}]} \leq t, T_{0} > t | \mathscr{F}_{T_{[-R_{0},R_{0}]}}] \right] \\ &\leq e^{-\lambda_{0}t}B_{1} + \mathbb{E}_{x} \left[\mathbf{1}_{\{T_{[-R_{0},R_{0}]} \leq t\}} \mathbb{P}_{x}[T_{0} > t | \mathscr{F}_{T_{[-R_{0},R_{0}]}}] \right] \\ &\leq e^{-\lambda_{0}t}B_{1} + \int_{0}^{t} \int_{-R_{0}}^{R_{0}} \mathbb{P}_{y}[T_{0} > t - u] \mathbb{P}_{x}[T_{[-R_{0},R_{0}]} \in du, X_{T_{[-R_{0},R_{0}]}} \in dy] \\ &\leq e^{-\lambda_{0}t}B_{1} + \int_{0}^{t} \int_{-R_{0}}^{R_{0}} B_{2}e^{-\lambda_{0}(t-u)}\mathbb{P}_{x}[T_{[-R_{0},R_{0}]} \in du, X_{T_{[-R_{0},R_{0}]}} \in dy] \\ &= e^{-\lambda_{0}t}B_{1} + e^{-\lambda_{0}t}B_{2} \int_{0}^{t} e^{\lambda_{0}u}\mathbb{P}_{x}[T_{[-R_{0},R_{0}]} \in du] \\ &\leq e^{-\lambda_{0}t}B_{1} + e^{-\lambda_{0}t}B_{2}\mathbb{E}_{x}[e^{\lambda_{0}T_{[-R_{0},R_{0}]}] \\ &\leq e^{-\lambda_{0}t}B_{1}(B_{2} + 1) < +\infty. \end{split}$$

Then by the above analysis,

$$\sup_{x \in \mathbb{R}^0} \sup_{t>2} e^{\lambda_0(t-1)} \mathbb{P}_x[Y_t \in A, t < T_0] \le \sup_{x \in \mathbb{R}^0} \sup_{t>2} e^{\lambda_0(t-1)} \mathbb{P}_x[t < T_0] < +\infty.$$

Therefore, by using dominated convergence theorem and 4.3, we have

$$\begin{split} \lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_{\eta}[Y_t \in A, t < T_0] &= \lim_{t \to \infty} \int_{\mathbb{R}^0} e^{\lambda_0 t} \mathbb{P}_x[Y_t \in A, t < T_0] \eta(\mathrm{d}x) \\ &= \int_{\mathbb{R}^0} \lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_x[Y_t \in A, t < T_0] \eta(\mathrm{d}x) \\ &= \int_{\mathbb{R}^0} \psi_0(x) \langle \psi_0, \mathbf{1}_A \rangle \eta(\mathrm{d}x), \end{split}$$

where we use (4.3) in the last equality. Similarly,

$$\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_{\eta}[t < T_0] = \int_{\mathbb{R}^0} \psi_0(x) \langle \psi_0, 1 \rangle \eta(\mathrm{d}x).$$

Thus we arrive at the conclusion that

$$\lim_{t \to \infty} \mathbb{P}_{\eta}[Y_t \in A | t < T_0] = \frac{\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_{\eta}[Y_t \in A, t < T_0]}{\lim_{t \to \infty} e^{\lambda_0 t} \mathbb{P}_{\eta}[t < T_0]} = \nu(A).$$

Secondly, we prove the result about the speed of convergence.

Assume that $I^{\sigma,\alpha} < +\infty$ and $\sup_x p_2^0(x,x) < +\infty$. Let $h_t(x) = e^{\lambda_0 t} \mathbb{P}_x[T_0 > t]$ and $h(x) = \psi_0(x)\mu(\psi_0)$. Then $\eta(h_t) = e^{\lambda_0 t} \mathbb{P}_\eta[T_0 > t]$ and $\eta(h) = \eta(\psi_0)\mu(\psi_0)$. We observe that

$$\begin{aligned} \|\mathbb{P}_{\eta}[Y_{t} \in \cdot | t < T_{0}] - \nu \|_{TV} &= \left\| \frac{\mathbb{P}_{\eta}[Y_{t} \in \cdot, t < T_{0}]}{\mathbb{P}_{\eta}[t < T_{0}]} - \nu \right\|_{TV} \\ &= \left\| \frac{e^{\lambda_{0}t}\eta P_{t}^{0}}{\eta(h_{t})} - \nu \right\|_{TV} \\ &= \left\| e^{\lambda_{0}t}\eta P_{t}^{0}(\frac{1}{\eta(h_{t})} - \frac{1}{\eta(h)} + \frac{1}{\eta(h)}) - \nu \right\|_{TV} \\ &\leq \left\| \frac{e^{\lambda_{0}t}\eta(h - h_{t})\eta P_{t}^{0}}{\eta(h_{t})\eta(h)} \right\|_{TV} + \left\| \frac{e^{\lambda_{0}t}\eta P_{t}^{0} - \eta(h)\nu}{\eta(h)} \right\|_{TV}, \quad (4.6) \quad \boxed{\operatorname{con}_{-4}} \end{aligned}$$

where $\eta P_t^0(\cdot) := \int_{\mathbb{R}^0} P_t^0(x, \cdot) \eta(\mathrm{d}x)$. Next, we prove the result by estimating the last two items of (4.6).

According to (4.2), if $\sup_x p_2^0(x,x) < +\infty$, then there exists a constant C_1 such that for any $A \in \mathscr{B}(\mathbb{R}^0)$,

$$|e^{\lambda_0 t} P_t^0 \mathbf{1}_A(x) - \psi_0(x) \int_A \psi_0(y) \mu(\mathrm{d}y)| \le C_1 e^{-(\lambda_1 - \lambda_0)t}.$$
(4.7) [con_1]

Since $\nu(dx) = \frac{\psi_0(x)\mu(dx)}{\mu(\psi_0)}$, then for any probability measure η on \mathbb{R}^0 , we have

$$\begin{aligned} \left\| e^{\lambda_0 t} \eta P_t^0 - \eta(\psi_0) \mu(\psi_0) \nu \right\|_{TV} \\ &= 2 \sup_{A \in \mathscr{B}(\mathbb{R}^0)} \left| e^{\lambda_0 t} \eta P_t^0(A) - \eta(\psi_0) \int_A \psi_0(y) \mu(\mathrm{d}y) \right| \\ &= 2 \sup_{A \in \mathscr{B}(\mathbb{R}^0)} \left| e^{\lambda_0 t} \int_{\mathbb{R}^0} P_t^0 \mathbf{1}_A(x) \eta(\mathrm{d}x) - \int_{\mathbb{R}^0} \psi_0(x) \int_A \psi_0(y) \mu(\mathrm{d}y) \eta(\mathrm{d}x) \right| \\ &\leq 2 \sup_{A \in \mathscr{B}(\mathbb{R}^0)} \int_{\mathbb{R}^0} \left| e^{\lambda_0 t} P_t^0 \mathbf{1}_A(x) - \psi_0(x) \int_A \psi_0(y) \mu(\mathrm{d}y) \right| \eta(\mathrm{d}x) \\ &\leq 2 C_1 e^{-(\lambda_1 - \lambda_0) t}. \end{aligned}$$
(4.8) con_2

Let $\eta - \nu = \tilde{\eta}$. $\tilde{\eta}$ is a signed measure and its Hahn decomposition is denoted by $\tilde{\eta} = \tilde{\eta}_{+} - \tilde{\eta}_{-}$. By calculation, it is easy to prove that $\nu(h) = \nu(h_t) = 1$ and $e^{\lambda_0 t} \nu P_t^0 = \nu$, so

$$\tilde{\eta}(h-h_t) = \eta(h-h_t), \qquad e^{\lambda_0 t} \eta P_t^0 - \eta(h)\nu = e^{\lambda_0 t} \tilde{\eta} P_t^0 - \tilde{\eta}(h)\nu.$$

Using (4.8), it follows that

$$\begin{aligned} \|e^{\lambda_0 t} \eta P_t^0 - \eta(h)\nu\|_{TV} &= \|e^{\lambda_0 t} \tilde{\eta} P_t^0 - \tilde{\eta}(h)\nu\|_{TV} \\ &\leq \|e^{\lambda_0 t} \tilde{\eta}_+ P_t^0 - \tilde{\eta}_+(h)\nu\|_{TV} + \|e^{\lambda_0 t} \tilde{\eta}_- P_t^0 - \tilde{\eta}_-(h)\nu\|_{TV} \\ &\leq 2C_1 e^{-(\lambda_1 - \lambda_0)t} \|\tilde{\eta}\|_{TV}. \end{aligned}$$

Besides, from (4.7), we obtain that

$$\begin{aligned} \left\| \frac{e^{\lambda_0 t} \eta(h-h_t) \eta P_t^0}{\eta(h_t)} \right\|_{TV} &= \left\| \mathbb{P}_{\eta} [Y_t \in \cdot | t < T_0] \tilde{\eta}(h-h_t) \right\|_{TV} \\ &= 2 \sup_{A \in \mathscr{B}(\mathbb{R}^0)} \mathbb{P}_{\eta} [Y_t \in A | t < T_0] \left| \int_{\mathbb{R}^0} (h(x) - h_t(x)) \tilde{\eta}(\mathrm{d}x) \right| \\ &\leq 2 C_1 e^{-(\lambda_1 - \lambda_0) t} \| \tilde{\eta} \|_{TV}. \end{aligned}$$

Combining with (4.6), it comes to the conclusion by taking $C = 4C_1/\mu(\psi_0)$.

In the following, we point out $I^{\sigma,\alpha} < +\infty$ is necessary for the QSD attracting all probability measures on \mathbb{R}^0 .

do->ent Theorem 4.1. If there exists a QSD π such that for any probability measure η on \mathbb{R}^0 and any subset $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\lim_{t \to \infty} \mathbb{P}_{\eta}[Y_t \in A | t < T_0] = \pi(A),$$

then $I^{\sigma,\alpha} < \infty$.

Proof. According to [6, Theorem 2.2], there exists a constant $\beta > 0$, such that

$$\mathbb{P}_{\pi}[T_0 > t] = e^{-\beta t}, \ \forall \ t \ge 0.$$

$$(4.9) \quad \text{[ex]}$$

By using (4.9), and a similar argument to the proof of [3, Proposition 7.5], we show that for any $\lambda \in (0, \beta)$ and any probability measure η on \mathbb{R}^0 , $\mathbb{E}_{\eta}[e^{\lambda T_0}] < +\infty$. For any $x \neq 0$, by taking $\eta = \delta_x$ (where δ_x is the Dirac measure), we obtain that $\mathbb{E}_x[e^{\lambda T_0}] < +\infty$. Let $g(x) = \mathbb{E}_x[e^{\lambda T_0}] < +\infty$.

We claim that g is bounded. If not, there would exist sequences $\{x_n\}$ such that $g(x_n) \geq 2^n, \forall n \geq 1$. However, if we take $\eta = \sum_{n=1}^{+\infty} \frac{1}{2^n} \delta_{x_n}$, then it is easy to verify that $\mathbb{E}_{\eta}[e^{\lambda T_0}] = +\infty$, which is a contradiction.

Therefore, we arrive at

$$\lambda \sup_{x \in \mathbb{R}^0} \mathbb{E}_x[T_0] + 1 \le \sup_{x \in \mathbb{R}^0} \mathbb{E}_x[e^{\lambda T_0}] < +\infty.$$

Combining it with Remark 3.3, $I^{\sigma,\alpha} < \infty$.

Next, we will obtain a corollary about the exponential moments of the hitting time T_0 , and prove the first Dirichlet eigenvalue λ_0 equals to the uniform decay rate

$$\lambda'_0 := \lim_{t \to +\infty} -\frac{1}{t} \log \sup_{x \in \mathbb{R}^0} \mathbb{P}_x[T_0 > t].$$

Corollary 4.2. If $I^{\sigma,\alpha} < \infty$, and for any $r_0 > 0$, $\sup_{x \in [-r_0,r_0] \setminus \{0\}} p_2^0(x,x) < +\infty$, then (1) we have

$$\sup_{x \in \mathbb{R}^0} \mathbb{E}_x[e^{\lambda T_0}] < +\infty \text{ if and only if } \lambda < \lambda_0, \tag{4.10} \text{ exp_lam}$$

(2) $\lambda_0 = \lambda'_0$.

Proof. (1) The "only if" implication follows from [15, Claim 2.4] and the monotonicity of exponential function. Hence we only prove "if" part. By the assumptions, Theorem 1.3 holds. Note that

$$\mathbb{P}_{\nu}[T_0 > t] = \int_{\mathbb{R}^0} P_t^0 1 \mathrm{d}\nu = \frac{\langle P_t^0 1, \psi_0 \rangle}{\mu(\psi_0)} = \mathrm{e}^{-\lambda_0 t},$$

so by proof of Theorem 4.1, we know for any $\lambda < \lambda_0$, $\sup_x \mathbb{E}_x[e^{\lambda T_0}] < +\infty$.

(2) Using (4.4), for any $x \neq 0$,

$$\lambda_0 = \lim_{t \to \infty} -\frac{1}{t} \log \mathbb{P}_x[T_0 > t].$$

It follows that

$$\limsup_{t \to +\infty} -\frac{1}{t} \log \sup_{x \in \mathbb{R}^0} \mathbb{P}_x[T_0 > t] \le \lim_{t \to \infty} -\frac{1}{t} \log \mathbb{P}_x[T_0 > t] = \lambda_0.$$

For any $\lambda < \lambda_0$, using (4.10) and Chebyshev inequality,

$$\sup_{x \in \mathbb{R}^0} \mathbb{P}_x[T_0 > t] \le e^{-\lambda t} \sup_{x \in \mathbb{R}^0} \mathbb{E}_x[e^{\lambda T_0}],$$

which yields that

$$\liminf_{t \to +\infty} -\frac{1}{t} \log \sup_{x \in \mathbb{R}^0} \mathbb{P}_x[T_0 > t] \ge \lambda,$$

which completes the proof.

) |exp distri

It should be mentioned here that for some classical cases, such as Example 1.5, the exponential convergence to QSD (1.6) is equivalent to the entrance from infinity. Now we prove Example 1.5 as follows.

Proof of Example 1.5. According to [25, Corollary 6], $I^{\sigma,\alpha} < +\infty$ holds if and only if $\gamma > 1$. Therefore, when $\gamma > 1$, by Theorem 1.2, Y^0 has unique QSD ν given by (1.3), ν is the Yaglom limit of Y^0 and (1.4) holds.

Furthermore, by [24, (3.11)], when $\gamma > 1$, for the transition density $p_t(x, y)$ with respect to μ , by choosing t = 1, we have there exists a constant $C_2 > 0$ such that

$$p_1(x,y) \le C_2, \quad x,y \in \mathbb{R}.$$

Note that

$$p_2(x,x) = \int_{\mathbb{R}} p_1(x,y)^2 \mu(\mathrm{d}y) \le C_2^2,$$

thus we know that

$$\sup_{x} p_2^0(x,x) \le \sup_{x} p_2(x,x) < +\infty.$$

Therefore, (1.6) holds in this case. Noting that the exponential convergence (1.6) implies that (1.5), so by Theorem 4.1, (1.6) means that $\gamma > 1$. Thus the exponential convergence to QSD (1.6) is equivalent to the entrance from infinity in this example.

Finally, let's discuss the quasi-ergodic distribution for Y^0 , which is a related topic.

def_QED Definition 4.3. Let ρ be a probability measure on \mathbb{R}^0 . We say ρ is a quasi-ergodic distribution(QED) for Y^0 , if for any $x \neq 0$ and any $A \in \mathscr{B}(\mathbb{R}^0)$,

$$\lim_{t \to \infty} \mathbb{E}_x \left(\frac{1}{t} \int_0^t \mathbf{1}_A(X_s) \mathrm{d}s | T_0 > t \right) = \rho(A).$$

Using (4.3), (4.4) and Corollary 3.5, it is easy to prove the following statement in the same way as [13, Theorem 3.1].

Corollary 4.4. Assume $I^{\sigma,\alpha} < \infty$, then for any bounded and measurable functions f, g on \mathbb{R}^0 , $x \neq 0$ and 0 ,

(1)
$$\lim_{t \to \infty} \mathbb{E}_x[f(X_{pt})g(X_t)|T_0 > t] = \int_{\mathbb{R}} f(y)m(\mathrm{d}y) \int_{\mathbb{R}} g(y)\nu(\mathrm{d}y),$$

(2)
$$\lim_{t \to \infty} \mathbb{E}_x[f(X_{pt})g(X_{qt})|T_0 > t] = \int_{\mathbb{R}} f(y)m(\mathrm{d}y) \int_{\mathbb{R}} g(y)m(\mathrm{d}y) dy$$

(3)
$$\lim_{t\to\infty} \mathbb{E}_x\left(\frac{1}{t}\int_0^t f(X_s)\mathrm{d}s|T_0>t\right) = \int_{\mathbb{R}} f(y)m(\mathrm{d}y),$$

where

$$m(\mathrm{d}y) = \psi_0(y)^2 \mu(\mathrm{d}y)$$
 and ν is the QSD for Y^0

Appendix

Proposition A.1 If $\alpha \in (1, 2)$, then for all $x \neq 0$, $\mathbb{P}_x[T_0 < T_\infty] = 1$.

Proof. According to [1, Chapter V] and Blumenthal-Getoor-McKean theorem, it is easy to prove the following formula:

$$\mathbb{P}_{x}[T_{0} < T_{[R,+\infty)} \land T_{(-\infty,-R]}] = \frac{G_{X}^{(-R,R)}(x,0)}{G_{X}^{(-R,R)}(0,0)}.$$
(4.11) [non_exp]

Using [14, Theorem 2.2.3], it follows that

$$G_X^{(-R,R)}(x,0) = \frac{2^{1-\alpha} |x|^{\alpha-1}}{\Gamma(\frac{\alpha}{2})^2} \int_1^{\frac{R}{|x|}} (s+1)^{\alpha/2-1} (s-1)^{\alpha/2-1} ds$$
$$G_X^{(-R,R)}(0,0) = \frac{2^{1-\alpha} R^{\alpha-1}}{\Gamma(\frac{\alpha}{2})^2 (\alpha-1)}.$$

It comes to the conclusion by letting $R \to +\infty$ and using L'Hôspital's rule in (4.11).

Proposition A.2 If $I^{\sigma,\alpha} < \infty$, then $\lambda_0 > 0$. Furthermore, λ_0 is simple.

Proof. If $I^{\sigma,\alpha} < \infty$, then according to [25, Theorem 1.4], the process Y is strongly ergodic, thus Y is exponentially ergodic. Therefore, by [25, Theorem 1.1],

$$\delta := \sup_{x} |x|^{\alpha - 1} \int_{\mathbb{R} \setminus (-|x|, |x|)} \sigma(y)^{-\alpha} \mathrm{d}y < \infty.$$

So by [25, Theorem 1.3], $\lambda_0 \ge (4\omega_\alpha \delta)^{-1} > 0$.

According to [19, Theorem 6.6] and spectral representation theorem, the spectral radius of G^0 is a simple eigenvalue and equals to λ_0^{-1} , which implies λ_0 is a simple eigenvalue.

Acknowledgements This work was supported by the National Nature Science Foundation of China (Grant No. 12171038), National Key Research and Development Program of China (2020YFA0712901).

References

- **BG68** [1] R. M. Blumenthal and R. K. Getoor. *Markov processes and potential theory*. Pure and Applied Mathematics, Vol. 29 Academic Press, 1968.
- bz06 [2] K. Bogdan and T. Zak. On Kelvin transformation. J. Theoret. Probab., 19(1):89–120, 2006.
- [3] P. Cattiaux, P. Collet, A. Lambert, Servet Martínez, S. Méléard, and J. San Martín. Quasi-stationary distributions and diffusion models in population dynamics. Ann. Probab., 37(5):1926–1929, 2009.
 - **CXW14** [4] X. Chen and J. Wang. Intrinsic ultracontractivity for general lévy processes on bounded open sets. *Illinois J. Math.*, 58:1117–1144, 2014.

- CW14[5] Z.-Q. Chen and J. Wang. Ergodicity for time-changed symmetric stable processes.Stoch. Proc. Appl., 124(9):2799–2823, 2014.
- CMS13 [6] P. Collet, S. Martínez, and J. San Martin. Quasi-stationary distributions. Markov chains, diffusions and dynamical systems. Probab. Appl. (N.Y.), Springer, Heidelberg, 2013.
- DK20 [7] L. Doring and A. E. Kyprianou. Entrance and exit at infinity for stable jump diffusions. Ann. Probab., 48(3):1220–1265, 2020.
- DKW20 [8] L. Doring, A. E. Kyprianou, and P. Weissmann. Stable processes conditioned to avoid an interval. *Stoch. Proc. Appl.*, 130:471–487, 2020.
- FOT11 [9] M. Fukushima, Y. Oshima, and M. Takeda. Dirichlet Forms and Symmetric Markov Processes, extended ed. De Gruyter Studies in Mathematics 19. de Gruyter, Berlin., 2011.
- [I0] W.-J. Gao and Y.-H. Mao. Quasi-stationary distribution for the birth-death process with exit boundary. J. Math. Anal. Appl., 427:114–125, 2015.
- <u>GMZ22</u> [11] W.-J. Gao, Y.-H. Mao, and C. Zhang. The birth-death processes with regular boundary: stationarity and quasi-stationarity. *Acta Math. Sin. (Engl. Ser.)*, 38(5).
- [GNW23] [12] A. Guillin, B. Nectoux, and L. Wu. Quasi-stationary distribution for strongly Feller Markov processes by Lyapunov functions and applications to hypoelliptic Hamiltonian systems. *Preprint hal-03068461*, 2023.
- [HYZ19] [13] G. He, G. Yang, and Y. Zhu. Some conditional limiting theorems for symmetric Markov processes with tightness property. *Electron. Commun. Probab.*, 24(60):1–11, 2019.
- **KA18** [14] A. E. Kyprianou. Stable processes, self-similarity and the unit ball. *ALEA Lat. Am. J. Probab. Math. Stat.*, 15(1):617–690, 2018.
- LJ12 [15] J. Littin C. Uniqueness of quasistationary distributions and discrete spectra when ∞ is an entrance boundary and 0 is singular. J. Appl. Probab., 49(3):719–730, 2012.
- MV12 [16] S. Méléard and D. Villemonais. Quasi-stationary distributions and population processes. *Probab. Surv.*, 9:340–410, 2012.
- [013] [17] Y. Oshima. Semi-Dirichlet forms and Markov processes. Berlin: De Gruyter Studies in Mathematics., 2013.
- **Sa99** [18] K. Sato. Lévy processes and infinitely divisible distributions. Cambridge Studies in Advanced Mathematics **68**. Cambridge: Cambridge Univ. Press., 1999.
- <u>SH74</u> [19] H. H. Schaefer. *Banach Lattices and Positive Operators*. Die Grundlehren der mathematischen Wissenschaften, vol. 215, Springer, 1974.
- T18 [20] M. Takeda. Compactness of symmetric Markov semi-groups and boundedness of eigenfunctions. *Trans. Amer. Math. Soc.*, 372(6):3905–3920, 2019.

- [T19] [21] M. Takeda. Existence and uniqueness of quasi-stationary distributions for symmetric markov processes with tightness property. J. Theoret. Probab., 32:2006–2019, 2019.
- **V19** [22] A. Velleret. Exponential quasi-ergodicity for processes with discontinuous trajectories. *arXiv.1902.01441*, 2019.
- wfy05 [23] F.-Y. Wang. Functional Inequalities, Markov Semigroups, and Spectral Theory. Beijing: Science press, 2005.
- WZ21 [24] J. Wang and L.T. Zhang. Functional inequalities for time-changed symmetric α -stable processes. *Front. Math. China*, 16(2):595–622, 2021.
- W21+ [25] T. Wang. Exponential and strong ergodicity for one-dimensional time-changed symmetric stable processes. *Bernoulli*, 29(1):580–596, 2023.
- [ZH16] [26] H. Zhang and G. He. Domain of attraction of quasi-stationary distribution for onedimensional diffusions. *Front. Math. China*, 11:411–421, 2016.