FREE BOUNDARY FLOW WITH SURGERY

ROBERT HASLHOFER

ABSTRACT. In this paper, we prove the existence of mean curvature flow with surgery for mean-convex surfaces with free boundary. To do so, we implement our recent new approach for constructing flows with surgery without a prior estimates in the free boundary setting. The flow either becomes extinct in finite time or for $t \to \infty$ converges smoothly in the one or two sheeted sense to a finite collection of stable connected minimal surfaces with empty or free boundary (in particular, there are no surgeries for t sufficiently large). Our free boundary flow with surgery will be applied in forthcoming work with Ketover, where we will address the existence problem for 3 free boundary minimal disks in convex balls.

Contents

1. Introduction	1
2. Definitions and basic properties	3
3. Distance to free boundary level set flow	5
4. Hybrid compactness theorem for free boundary flows	6
5. Excluding microscopic surgeries	9
6. Multiplicity-one for free boundary flows	10
6.1. Large blowups with entropy at most two	10
6.2. Sheeting theorem for Λ -blowup sequences	12
6.3. Ruling out generalized limit flows with density two	13
7. Partial regularity and convexity	16
8. Canonical neighborhoods and existence theorem	17
References	19

1. INTRODUCTION

Geometric flows with surgery provide a controlled way of flowing through singularities by cutting along a suitable collection of necks and gluing in standard caps. Mean curvature flow with surgery was first constructed in the setting of two-convex hypersurfaces in \mathbb{R}^N for $N \geq 4$ in pioneering work by Huisken-Sinestrari [HS09], which in particular yielded a topological classification of two-convex hypersurfaces. The case N = 3 has then been solved independently by Brendle-Huisken [BH16] and Kleiner and the author [HK17]. The construction has been later generalized to other ambient manifolds [BH18, HK19], low entropy flows [MW21, DH22], and higher codimensions [Ngu20, LN21]. Applications include the

Date: June 14, 2023.

work on moduli-spaces in [BHH21, BHH19], the construction of foliations in [HK19, LM23], and the proof of the low entropy Schönflies conjecture in [DH22].

However, a problem that until recently seemed inaccessible is the construction of a flow with surgery in the setting of mean-convex free boundary surfaces. The reason for this is that both the approach by Brendle-Huisken [BH16] and the approach by Kleiner and the author [HK17] crucially rely on the noncollapsing result of Andrews [And12], which is only available in the setting without boundary. In this paper, we solve this problem for mean-convex surfaces with free boundary in any smooth convex domain D. More precisely, throughout this paper, D denotes a smooth connected compact domain in a Riemannian three-manifold (N, g), such that ∂D is nonempty and has positive second fundamental form. Our main result gives existence, as well as long-time behaviour, of a flow with surgery with free boundary starting at any mean-convex free boundary initial surface $M_0 = \partial K_0 \subset D$:

Theorem 1.1. Given any smooth compact strictly mean-convex free boundary domain $K_0 \subset D$, there exists a free boundary flow with surgery starting at K_0 . Moreover, the flow either becomes extinct in finite time or for $t \to \infty$ converges smoothly in the one or two-sheeted sense to a finite collection of stable connected minimal surfaces with empty or free boundary (in particular, there are no surgeries for t sufficiently large).

For the purpose of the present paper a free boundary flow with surgery is a free boundary (δ, \mathcal{H}) -flow in the domain D as introduced in Definition 2.4. In particular, $\delta > 0$ is a small parameter that captures the quality of the surgery necks and half necks, and \mathcal{H} is a triple of curvature scales $H_{\text{trigger}} \gg H_{\text{neck}} \gg H_{\text{thick}} \gg 1$, which is used to specify more precisely when and how surgeries are performed. Moreover, there is a finite set of surgery times t_i , where (i) some necks or half necks in the presurgery domain $K_{t_i}^-$ are replaced by caps or half caps yielding a domain $K_{t_i}^{\sharp} \subseteq K_{t_i}^-$, and/or (ii) some connected components covered entirely by high curvature regions are discarded yielding the postsurgery domain $K_{t_i}^+ \subseteq K_{t_i}^{\sharp}$.

As a byproduct of our existence proof we also obtain a canonical neighborhood result:

Theorem 1.2. Given any $\varepsilon > 0$, regions of sufficiently large curvature are ε -close (after rescaling to unit curvature) for any sufficiently good surgery parameters to either (i) the evolution of a standard cap preceded by a round shrinking cylinder, or a round shrinking sphere, translating bowl or ancient oval, or (ii) the evolution of a standard half cap preceded by a round shrinking half cylinder, or a round shrinking half cylinder, round shrinking half sphere, translating half bowl or ancient half oval.

Remark 1.3. The existence and canonical neighborhood theorem will be applied in forthcoming work with Ketover [HK], based on the methodology from our earlier paper [HK19], to address the existence problem for 3 free boundary minimal disks in convex balls.

To prove our results we implement our recent new approach from [Has23], which is based on the theory of weak solutions rather than a priori estimates for smooth solutions, in the free boundary setting. Specifically, to begin with, after introducing the necessary definitions, we first prove a quantitative estimates for the space-time Hausdorff distance between the free boundary flow with surgery and the free boundary level set flow. We then study sequences \mathcal{K}^j of free boundary (δ, \mathcal{H}^j) -flows, with the same mean-convex initial condition $K_0 \subset D$, where $\delta \leq \bar{\delta}$ and where the curvature scales \mathcal{H}^j improve along the sequence. Given any sequence of rescaling factors $\lambda_j \to \infty$, we consider the blowup sequence $\tilde{\mathcal{K}}^j$, which is obtained from \mathcal{K}^j by translating X_j to the origin and parabolically rescaling by λ_j . To keep track of multiplicities we also consider the associated family of Radon measures $\tilde{\mathcal{M}}^j$.

We then establish a hybrid compactness theorem, which allows us to pass to a limits of such generalized blowup sequences, which are smooth near the surgery regions but potentially singular in all other regions. Moreover, using Edelen's monotonicty formula for free boundary flows from [Ede20] we rule out microscopic surgeries, namely the potential scenario that one has convergence to a plane or half plane with surgeries satisfying $\lambda_j/H_{\text{neck}}^j \to 0$.

Next, we generalize the theory of mean-convex Brakke flows with free boundary by Edelen, Ivaki, Zhu and the author [EHIZ22] to our setting of hybrid limits of free boundary flows with surgery, and in particular establish multiplicity-one, regularity and convexity. As a consequence of these results, taking also into account the recent classification of ancient solutions from [BC19, ADS20], we then establish the canonical neighborhood theorem.

Using the canonical neighborhood theorem, via a continuity argument similarly as in [HK17], we can establish the existence of free boundary flow with surgery on arbitrarily large time intervals. Finally, borrowing an argument of Brendle-Huisken [BH18], in combination with the long-time limit results from [EHIZ22] and our distance estimate to the free boundary level set flow, we can upgrade this to existence for all times and convergence.

Acknowledgments. I thank Jonathan Zhu for helpful discussions. This research has been supported by an NSERC Discovery Grant and a Sloan Research Fellowship.

2. Definitions and basic properties

In this section, we define free boundary flows with surgery and establish some basic properties. Let us begin with the following flexible notion of free boundary δ -flows:

Definition 2.1 (free boundary δ -flow). A free boundary δ -flow \mathcal{K} in D is a collection of finitely many smooth free boundary strictly mean-convex mean curvature flows $\{K_t^i \subset D\}_{t \in [t_{i-1}, t_i]}$ $(i = 1, \ldots, k; t_0 < \ldots < t_k)$ such that:

- (a) for each i = 1, ..., k 1, the final time slices of some collection of disjoint strong δ -necks or strong half δ -necks (see Definition 2.2) are replaced by pairs of standard caps or standard half caps (see Definition 2.3), giving a domain $K_{t_i}^{\sharp} \subseteq K_{t_i}^i =: K_{t_i}^-$.
- or standard half caps (see Definition 2.3), giving a domain $K_{t_i}^{\sharp} \subseteq K_{t_i}^i =: K_{t_i}^-$. (b) the initial time slice of the next flow, $K_{t_i}^{i+1} =: K_{t_i}^+$, is obtained from $K_{t_i}^{\sharp}$ by discarding some connected components.
- (c) there exists some $r_{\sharp} = r_{\sharp}(\mathcal{K}) > 0$, such that all necks and half necks in item (a) have radius $r \in [\frac{1}{2}r_{\sharp}, 2r_{\sharp}]$.

The above definition relies on the following two further definitions:

Definition 2.2 (strong δ -neck and strong half δ -neck). We say that a free boundary δ -flow $\mathcal{K} = \{K_t \subset D\}_{t \in I}$ has a

- strong δ -neck with center $p \in \text{Int}(D)$ and radius r at time $t_0 \in I$, if the rescaled flow $\{r^{-1}(\exp_p|_{B_{1/\delta}(0)})^{-1}(K_{t_0+r^2t})\}_{t\in(-1,0]}$ is δ -close in $C^{\lfloor 1/\delta \rfloor}$ in $B_{1/\delta}(0) \times (-1,0]$ to the evolution of a solid round cylinder $\bar{B}^2 \times \mathbb{R}$ with radius 1 at t = 0.
- strong half δ -neck with center $p \in \partial D$ and radius r at time $t_0 \in I$, if the rescaled flow $\{r^{-1}(\Phi_p^{-1}|_{B_{1/\delta}^+(0)})^{-1}(K_{t_0+r^2t})\}_{t\in(-1,0]}$ is δ -close in $C^{\lfloor 1/\delta \rfloor}$ in $B_{1/\delta}^+(0) \times (-1,0]$ to the evolution of a solid round half cylinder $\bar{B}^2_+ \times \mathbb{R}$ with radius 1 at t = 0.

Here, we recall that for any $p \in \partial D$ the normal exponential map defines a diffeomorphism Φ_p from an open neighborhood of p in D to an open half ball $B_{\rho}^+(0) \subset \mathbb{R}^3_+ := \mathbb{R}_{\geq 0} \times \mathbb{R}^2$.

Definition 2.3 (replacement by standard caps or standard half caps). Given a cap separation parameter $\Gamma < \infty$ and any $\delta \leq \frac{1}{100\Gamma}$, we say that the final time slice of a strong δ -neck with center $p \in \text{Int}(D)$ and radius r is replaced by a pair of standard caps or the final time slice of a strong half δ -neck with center $p \in \partial D$ and radius r is replaced by a pair of standard half caps, respectively, if the presurgery domain K^- is replaced by a postsurgery domain $K^{\sharp} \subseteq K^-$ with free boundary in D, such that:

- (a) the modification takes places inside $B = B(p, 5\Gamma r)$.
- (b) there are bounds for the second fundamental form and its derivatives:

$$\sup_{\partial K^{\sharp} \cap B} |\nabla^{\ell} A| \le C_{\ell} r^{-1-\ell} \qquad (\ell = 0, 1, 2, \ldots).$$

(c) for every point $p_{\sharp} \in \partial K^{\sharp} \cap B$ with $\lambda_1(p_{\sharp}) < 0$, there is a point $p_- \in \partial K^- \cap B$ with

$$\frac{\lambda_1}{H}(p_-) \le \frac{\lambda_1}{H}(p_{\sharp}).$$

(d) the domain $r^{-1}(\exp_p|_{B_{10\Gamma_r}(0)})^{-1}(K^{\sharp})$ is $\delta'(\delta)$ -close in $B(0, 10\Gamma)$ to a pair of standard caps, or the domain $r^{-1}(\Phi_p|_{B_{10\Gamma_r}(0)})^{-1}(K^{\sharp})$ is $\delta'(\delta)$ -close in $B^+(0, 10\Gamma)$ to a pair of standard half caps, respectively, that are at distance Γ from the origin. Here, $\delta'(\delta) \to 0$ as $\delta \to 0$.

Here, a standard cap $K^{\text{st}} \subset \mathbb{R}^3$ is a smooth convex domain that coincides with a solid round half-cylinder of radius 1 outside a ball of radius 2. Similarly, we call $K^{\text{st}} \cap \mathbb{R}^3_+$ a standard half cap.

A free boundary flow with surgery is a free boundary (δ, \mathcal{H}) -flow defined as follows:

Definition 2.4 (free boundary flow with surgery). A free boundary (δ, \mathcal{H}) -flow in D, where $\mathcal{H} = (H_{\text{thick}}, H_{\text{neck}}, H_{\text{trigger}})$, is a free boundary δ -flow $\{K_t \subset D\}_{t \ge 0}$ with smooth free boundary strictly mean-convex initial condition $K_0 \subset D$ such that:

- (a) $H \leq H_{\text{trigger}}$ everywhere, and surgery and/or discarding occurs precisely at times t when $H = H_{\text{trigger}}$ somewhere.
- (b) The collection of necks and half necks in Definition 2.1(a) is a minimal collection of solid δ -necks and solid half δ -necks of curvature H_{neck} which separate the set $\{H = H_{\text{trigger}}\}$ from $\{H \leq H_{\text{thick}}\}$ in K_t^- .

- (c) K_t^+ is obtained from K_t^{\sharp} by discarding precisely those connected components with $H > \frac{1}{10}H_{\text{neck}}$ everywhere. For each pair of facing surgery caps or surgery half caps, precisely one is discarded.
- (d) If a strong δ -neck or strong half δ -neck from item (b) also is a strong $\hat{\delta}$ -neck or strong half $\hat{\delta}$ -neck for some $\hat{\delta} < \delta$, then property (d) of Definition 2.3 also holds with $\hat{\delta}$ instead of δ .

As a consequence of the definitions we have the following basic properties:

Proposition 2.5 (basic properties). There exist $\overline{\delta} > 0$ and $\Gamma_0 < \infty$, such that any free boundary δ -flow \mathcal{K} in D with surgery quality $\delta \leq \overline{\delta}$ and cap separation parameter $\Gamma \geq \Gamma_0$ satisfies the following:

- (a) If p is the center of a surgery neck or surgery half neck of radius r, then there are no other surgeries in $B(p, \frac{1}{10}\delta^{-1}r)$.
- (b) For every ball B we have $|\partial K_{t_1} \cap B| \leq |\partial K' \cap B|$ for every K' that agrees with K_{t_1} outside B and satisfies $K_{t_1} \subseteq K' \subseteq K_{t_0}$ for some $t_0 < t_1$.

Moreover, any free boundary (δ, \mathcal{H}) -flow in D for $t \leq T$ satisfies $H > C^{-1}$ and $|A| \leq CH$, where $C = C(D, K_0, T) < \infty$.

Proof. The spatial separation of surgeries follows directly from the definitions. Next, the onesided minimization follows using the geometric measure theory argument in [Whi00, Section 3]. Finally, the claimed bounds for H and |A|/H follow from [Ede16] (which simplifies a lot in our setting thanks to the convexity of D), and the definition of surgeries.

Convention 2.6. We now fix a suitable standard cap K^{st} and cap separation parameter $\Gamma < \infty$. Moreover, throughout this paper $\bar{\delta} > 0$ and $\varepsilon > 0$ denote sufficiently small constants (by convention, these constants can be decreased finitely many times as needed or convenient).

3. DISTANCE TO FREE BOUNDARY LEVEL SET FLOW

In this section, we prove a quantitative estimate for the distance to the free boundary level set flow from [GS93]. We will identify free boundary δ -flows with their spacetime track

(3.1)
$$\mathcal{K} = \bigcup_{t} K_t \times \{t\} \subset D \times \mathbb{R},$$

where we set $K_{t_i} := K_{t_i}^-$ at surgery times to ensure that \mathcal{K} is a closed subset of space-time.

Proposition 3.1 (distance to free boundary level set flow). Given any $T < \infty$, there exist constants $\bar{\delta} = \bar{\delta}(D,T) > 0$ and $C = C(D, K_0, T) < \infty$ such that if \mathcal{K} is a free boundary (δ, \mathcal{H}) -flow $(\delta \leq \bar{\delta})$ with initial condition K_0 , where $H_{\text{neck}} \geq C$, and \mathcal{L} is the free boundary level set flow with the same initial condition K_0 , then

(3.2)
$$d_{\mathrm{H}}\left(\mathcal{K} \cap \{t \leq T\}, \mathcal{L} \cap \{t \leq T\}\right) \leq CH_{\mathrm{neck}}^{-1},$$

where $d_{\rm H}$ denotes the Hausdorff distance of the spacetime tracks.

Proof. Recall from [EHIZ22] that the free boundary level set flow is the maximal family of closed sets starting at K_0 that does not bump into any smooth free boundary subsolution of the mean curvature flow. Observing that free boundary (δ, \mathcal{H})-flows do not bump into any smooth free boundary subsolution of the mean curvature flow, we thus get $\mathcal{K} \subseteq \mathcal{L}$.

To estimate the distance from the other direction, observe that similarly as in [Has23, Claim 3.2], there exists an $\eta < \infty$, such that if t is a surgery time of \mathcal{K} and $B_r \subseteq K_t^- \cap \operatorname{Int}(D)$ is a geodesic ball of radius $r > \eta H_{\operatorname{neck}}^{-1}$, then $B_r \subseteq K_t^+$. Also, note that there is a $C_0 = C_0(K_0) < \infty$, such that for all $\gamma > 0$ small enough there is some $\tau \in [C_0^{-1}\gamma, C_0\gamma]$, such that

(3.3)
$$d(K_{\tau}, D \setminus K_0) = \gamma.$$

Denoting by $-\kappa$ a lower bound for the Ricci curvature, we choose $\gamma = \eta e^{\kappa T} / H_{\text{neck}}$, which is allowed provided H_{neck} is large enough. With the corresponding τ , we let \mathcal{L}^{τ} be the level set flow with initial condition $L_0^{\tau} = K_{\tau}$.

Claim 3.2 (evolution of distance). We have $L_t^{\tau} \subseteq K_t$ for all $t \leq T$ with the estimate

(3.4)
$$d(L_t^{\tau}, D \setminus K_t) \ge \gamma e^{-\kappa t}.$$

Proof. Consider the function $f(t) := e^{\kappa t} d(L_t^{\tau}, D \setminus K_t)$, and note that $f(0) = \gamma$. Now, denoting by $\frac{d}{dt}$ the lim inf of difference quotients and by H the mean curvature in the viscosity sense, away from the surgery times we can estimate

(3.5)
$$\frac{d}{dt}f(t) \ge \kappa e^{\kappa t} d(L_t^{\tau}, D \setminus K_t) + e^{\kappa t} (H(p) - H(q)) \ge 0,$$

where we observed that thanks to the strict convexity of D the distance at time t is realized by points $p \in \partial L_t^{\tau} \cap \operatorname{Int}(D)$ and $q \in \partial K_t \cap \operatorname{Int}(D)$, and these points satisfy $H(p) - H(q) \geq -\kappa d(p,q)$ by mean curvature comparison. Finally, if t_i is a surgery time, then $d(L_{t_i}^{\tau}, D \setminus K_{t_i}^{-}) \geq \gamma e^{-\kappa t}$ together with the above observation implies $d(L_{t_i}^{\tau}, D \setminus K_{t_i}^{+}) \geq \gamma e^{-\kappa t}$. \Box

Summarizing, we have shown that $\mathcal{L}^{\tau} \subseteq \mathcal{K} \subseteq \mathcal{L}$, where τ is comparable to H_{neck}^{-1} . Since \mathcal{L}^{τ} and \mathcal{L} just differ by a shift by τ in time direction, this implies the assertion.

4. Hybrid compactness theorem for free boundary flows

Let \mathcal{K}^j be a sequence of free boundary (δ, \mathcal{H}^j) -flows, where $\delta \leq \overline{\delta}$, with the same strictly mean-convex initial condition K_0 , and suppose that $\mathcal{H}^j \to \infty$, where we use the abbreviation

(4.1)
$$\mathcal{H}^{j} \to \infty \quad :\Leftrightarrow \quad \min\left(H^{j}_{\text{thick}}, \frac{H^{j}_{\text{neck}}}{H^{j}_{\text{thick}}}, \frac{H^{j}_{\text{trigger}}}{H^{j}_{\text{neck}}}\right) \to \infty.$$

Let $\lambda_j \to \infty$ be a sequence of rescaling factors, and let $X_j = (p_j, t_j) \in \partial \mathcal{K}^j$ be a sequence of space-time points satisfying $t_j \leq T$ for some $T < \infty$. Consider the blowup sequence

(4.2)
$$\widetilde{\mathcal{K}}^j := \mathcal{D}_{\lambda_j}(\mathcal{K}^j - X_j),$$

which is obtained from \mathcal{K}^{j} by translating X_{j} to the origin and parabolically rescaling by λ_{j} , and where for ease of notation we pretend that D is a subset of three-dimensional Euclidean space. Moreover, we also consider the associated family of Radon measures

(4.3)
$$\widetilde{\mathcal{M}}^{j} = \left\{ \widetilde{\mu}^{j}_{t} = \mathcal{H}^{2} \lfloor \partial \widetilde{K}^{j}_{t} \right\},$$

where we set $K_t := K_t^-$ at surgery times. Furthermore, we can assume that

(4.4)
$$\Lambda := \lim_{j \to \infty} \frac{\lambda_j}{H_{\text{neck}}^j} \in [0, \infty]$$

exists. We then say that $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ is a Λ -blowup sequence. For $\Lambda = 0$ we have:

Proposition 4.1 (small blowups). If $\Lambda = 0$, then a subsequence of $\widetilde{\mathcal{K}}^{j}$ Hausdorff converges to a limit \mathcal{K} , which is a blowup limit of the free boundary level set flow of K_0 .

Proof. Let \mathcal{L} be the free boundary level set flow of K_0 . By Proposition 3.1 (distance from free boundary level set flow) we have $d_{\rm H}(\mathcal{K}^j \cap \{t \leq 2T\}, \mathcal{L} \cap \{t \leq 2T\}) \leq C/H^j_{\rm neck}$, hence

(4.5)
$$d_{\mathrm{H}}(\mathcal{\tilde{K}}^{j} \cap \{t \leq \lambda_{j}^{2}T\}, \mathcal{D}_{\lambda_{j}}(\mathcal{L} - X_{j}) \cap \{t \leq \lambda_{j}^{2}T\}) \leq C\lambda_{j}/H_{\mathrm{neck}}^{j}.$$

Since $\lim_{j\to\infty} \lambda_j / H^j_{\text{neck}} = 0$ by assumption, it follows that $\widetilde{\mathcal{K}}^j$ and $\mathcal{D}_{\lambda_j}(\mathcal{L} - X_j)$ converge subsequentially in the Hausdorff sense to the same limit \mathcal{K} . This proves the proposition. \Box

Next, for $0 < \Lambda < \infty$ the limiting objects will be either ancient Brakke δ -flows in \mathbb{R}^3 as in [Has23, Definition 4.2] or ancient free boundary Brakke δ -flows in \mathbb{R}^3_+ defined as follows:

Definition 4.2 (ancient free boundary Brakke δ -flow). An ancient free boundary Brakke δ -flow in \mathbb{R}^3_+ is a pair $(\mathcal{K}, \mathcal{M})$ consisting of a nested family of closed sets $\mathcal{K} = \{K_t \subset \mathbb{R}^3_+\}_{t \in (-\infty,T)}$ and a family of Radon measures $\mathcal{M} = \{\mu_t\}_{t \in (-\infty,T)}$ in \mathbb{R}^3_+ , for which there exists a constant $r_{\sharp} = r_{\sharp}(\mathcal{K}, \mathcal{M}) \in (0, \infty)$ and a disjoint collection of two-sided parabolic balls $P_i = B(p_i, 50\Gamma r_{\sharp}) \times (t_i - \frac{1}{2}r_{\sharp}^2, t_i + \varepsilon r_{\sharp}^2)$ such that:

- (a) For $t \in (t_i \frac{1}{2}r_{\sharp}^2, t_i + \varepsilon r_{\sharp}^2)$, we have $\mu_t \lfloor B(p_i, 50\Gamma r_{\sharp}) = \mathcal{H}^2 \lfloor \partial K_t \cap B(p_i, 50\Gamma r_{\sharp})$, and for $t \neq t_i$ the sets $K_t \cap B(p_i, 50\Gamma r_{\sharp})$ are smooth and evolve by free boundary mean curvature flow. At time t_i a strong δ -neck or strong half δ -neck (see Definition 2.2) of radius $r_i \in [r_{\sharp}/2, 2r_{\sharp}]$ centered at (p_i, t_i) is replaced by a pair of standard caps or standard half caps (see Definition 2.3), and possibly some connected components of K_{t_i} are discarded.
- (b) Considering the somewhat smaller $P'_i = B(p_i, 25\Gamma r_{\sharp}) \times (t_i \frac{1}{4}r_{\sharp}^2, t_i + \frac{1}{2}\varepsilon r_{\sharp}^2)$, we have that \mathcal{M} is a free boundary integral Brakke flow away from $\bigcup_i P'_i$, and $\partial \mathcal{K} \setminus \bigcup_i P'_i = \operatorname{spt} \mathcal{M} \setminus \bigcup_i P'_i$.

Here, we use the notion of free boundary integral Brakke flows from Edelen [Ede20], and in particular recall that the support consists of all space-time points where the (reflected) Gaussian density is at least one.

Remark 4.3 (reflection). In light of [Ede20, Proposition 4.6] any ancient free boundary Brakke δ -flow in \mathbb{R}^3_+ can be reflected to a flow without boundary in \mathbb{R}^3 , which satisfies all the axioms of an ancient Brakke δ -flow in \mathbb{R}^3 with the caveat that at surgery times the closeness to a standard cap from item (d) of Definition 2.3 now only holds in the C^1 sense.

Next, convergence of Λ -blowup sequences to Brakke δ -flows is defined similarly as in [Has23, Definition 4.2], and convergence to free boundary Brakke δ -flows is defined as follows:

Definition 4.4 (convergence to free boundary Brakke δ -flow). Given $0 < \Lambda < \infty$, a Λ blowup sequence $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ converges to a free boundary Brakke δ -flow $(\mathcal{K}, \mathcal{M})$ if:

- (a) The space-time tracks $\widetilde{\mathcal{K}}^{j}$ Hausdorff converge to the space-time track \mathcal{K} .
- (b) If $P_i = B(p_i, 50\Gamma r_{\sharp}) \times (t_i \frac{1}{2}r_{\sharp}^2, t_i + \varepsilon r_{\sharp}^2)$ is a surgery region of $(\mathcal{K}, \mathcal{M})$ as in Definition 4.2 (ancient free boundary Brakke δ -flow), then for some $(p_i^j, t_i^j) \in \widetilde{\mathcal{K}}^j$ converging to (p_i, t_i) the forwards and backwards portion $\{\widetilde{K}_{t+t_i}^j - p_i^j\}_{t \in (0,\varepsilon r_{\sharp}^2)}$ and $\{\widetilde{K}_{t+t_i}^j - p_i^j\}_{t \in (-\frac{1}{2}r_{\sharp}^2,0]}$ converge smoothly to $\{K_{t+t_i} - p_i\}_{t \in (0,\varepsilon r_{\sharp}^2)}$ and $\{K_{t+t_i} - p_i\}_{t \in (-\frac{1}{2}r_{\sharp}^2,0]}$, respectively, in $B(0, 50\Gamma r_{\sharp})$.
- (c) $\widetilde{\mathcal{M}}^{j}$ converges to \mathcal{M} in the sense of free boundary Brakke flows away from $\cup_{i} P'_{i}$.

Theorem 4.5 (hybrid compactness). Any Λ -blowup sequence $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ with $\Lambda \in (0, \infty)$ has a subsequence that converges to a limit $(\mathcal{K}, \mathcal{M})$ that is either an ancient Brakke δ -flow in \mathbb{R}^3 or an ancient free boundary Brakke δ -flow in \mathbb{R}^3_+ .

Proof. If $\lambda_j d(p_j, \partial D) \to \infty$, then the proof of [Has23, Theorem 4.4] applies, yielding convergence to an ancient Brakke δ -flow in \mathbb{R}^3 . Hence, after shifting the base-points by controlled rescaled distance we can assume from now on that $p_j \in \partial D$. We can also assume that $\widetilde{\mathcal{K}}^j$ Hausdorff converges to a limit $\mathcal{K} = \{K_t \subset \mathbb{R}^3_+\}_{t \in (-\infty,T)}$, which is a nested family of closed sets.

Since $\Lambda > 0$, for any $R < \infty$ the number N_R^j of surgery centers of $\widetilde{\mathcal{K}}^j$ in the two-sided parabolic ball $P(0, R) = B(0, R) \times (-R^2, R^2)$ is uniformly bounded. After passing to a subsequence, we can assume that $N_R^j = N_R$ is independent of j. Moreover, denoting by $\{(p_i^j, t_i^j)\}_{i=1}^{N_R}$ the surgery centers of $\widetilde{\mathcal{K}}^j$ in P(0, R), and denoting by $r_i^j \in [\Lambda_j/2, 2\Lambda_j]$ the neck radii, where $\Lambda_j := \lambda_j/H_{\text{neck}}^j$, after passing to a further subsequence we can assume that

(4.6)
$$(p_i^j, t_i^j) \to (p_i, t_i)$$
 and $r_i^j \to r_i \in [\Lambda/2, 2\Lambda].$

Set $r_{\sharp} := \Lambda$, and recall that $\varepsilon > 0$ is a small fixed constant by Convention 2.6. Arguing similarly as in the proof of [Has23, Claim 4.5], where we now use the local regularity theorem for free boundary flows from [Ede20], for each $P_i = B(p_i, 50\Gamma r_{\sharp}) \times (t_i - \frac{1}{2}r_{\sharp}^2, t_i + \varepsilon r_{\sharp}^2)$ we get

(4.7)
$$\limsup_{j \to \infty} \sup_{\partial \widetilde{\mathcal{K}}^j \cap P_i} |\nabla^{\ell} A| < \infty \qquad (\ell = 0, 1, 2, \ldots).$$

Hence, the convergence in the two-sided parabolic balls P_i is smooth in the sense of Definition 4.4(b). Moreover, away from $\bigcup_i P'_i$ we can pass to a subsequential weak limit using the compactness theorem for free boundary integral Brakke flows from [Ede20]. In the surgery regions we define our measure by declaring that $\mu_t(A) := \mathcal{H}^2(\partial K_t \cap A)$ for $A \subseteq B(p_i, 50\Gamma r_{\sharp})$ and $t \in (t_i - \frac{1}{2}r_{\sharp}^2, t_i + \varepsilon r_{\sharp}^2)$, and observe that this is consistent in the overlap regions. We now consider a sequence $R_j \to \infty$, and pass to a diagonal subsequence of the above

We now consider a sequence $R_j \to \infty$, and pass to a diagonal subsequence of the above to obtain a global limit $(\mathcal{K}, \mathcal{M})$. Observe that our limit satisfies all the properties listed in Definition 4.2(a), and that \mathcal{M} is a free boundary integral Brakke flow away from $\cup_i P'_i$. Finally, arguing similarly as in the last paragraph of the proof [Has23, Theorem 4.4], where we now use the (reflected) Gaussian density from [Ede20], we see that the support of \mathcal{M} agrees with $\partial \mathcal{K}$ away from $\cup_i P'_i$. This finishes the proof of the theorem.

Finally, let us deal with the case $\Lambda = \infty$:

Proposition 4.6 (large blowups). If $\Lambda = \infty$, then $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ subsequentially converges either (a) in the Hausdorff and Brakke sense to a mean-convex flow in \mathbb{R}^3 or a mean-convex free boundary flow in \mathbb{R}^3_+ or (b) smoothly to a static or backwards/forwards quasistatic multiplicity-one plane in \mathbb{R}^3 or multiplicity-one free boundary half plane in \mathbb{R}^3_+ .

Proof. If for every $R < \infty$ the two-sided parabolic ball P(0, R) does not contain points modified by surgeries for infinitely many j, then the conclusion (a) holds. Assume now that there is some $R < \infty$, such that P(0, R) contains points modified by by surgeries for all j. Since $\Lambda = \infty$, arguing similarly as in the proof of (4.7) we see that for each $\rho < \infty$ we have

(4.8)
$$\limsup_{j \to \infty} \sup_{\partial \widetilde{\mathcal{K}}^j \cap P(0,\rho)} |\nabla^{\ell} A| = 0 \qquad (\ell = 0, 1, 2, \ldots).$$

It follows that conclusion (b) holds. This finishes the proof of the proposition.

A large portion of this paper will deal with analyzing the limits constructed above:

Definition 4.7 (generalized limit flow). A Λ -generalized limit flow is any limit (\mathcal{K}, \mathcal{M}) provided by Theorem 4.5 (hybrid compactness) or Proposition 4.6 (large blowups).

To conclude this section, let us observe that generalized limit flows inherit the one-sided minimization property in the following sense:

Corollary 4.8 (one-sided minimization for generalized limit flows). Let $(\mathcal{K}, \mathcal{M})$ be a Λ generalized limit flow (see Definition 4.7). Suppose γ is a 1-cycle that at some time t bounds
a 2-chain in ∂K_t , and $\mathcal{H}^2(\operatorname{sing} \partial K_t) = 0$. Then, there exists a 2-chain Σ supported in K_t with $\partial \Sigma = \gamma$, such that $\mathcal{H}^2(\Sigma) \leq \mathcal{H}^2(\Sigma')$ for any 2-chain Σ' bounded by γ .

Proof. This follows from the one-sided minimization for smooth flows (see Proposition 2.5), via the same argument as in the the proof of [Whi00, Theorem 6.1]. \Box

5. Excluding microscopic surgeries

The goal of this section is to prove the following theorem.

Theorem 5.1 (no microscopic surgeries). Suppose $\widetilde{\mathcal{K}}^j$ is a 0-blowup sequence (with $\delta \leq \overline{\delta}$ small enough) that Hausdorff converges in space-time to a (quasi-)static multiplicity-one plane or half plane. Then, for any $R < \infty$ there exists a $j_0 = j_0(R) < \infty$, such that for all $j \geq j_0$ the two-sided parabolic ball P(0, R) contains no points modified by surgeries.

Proof. Observing that the argument from the proof of [Has23, Theorem 4.4] already rules out microscopic surgeries in the interior, it suffices to rule out microscopic surgeries at the boundary. Specifically, pretending for ease of notation that $D \subset \mathbb{R}^3_+$ with $0 \in \partial D$, suppose towards a contradiction that there is a 0-blowup sequence $\widetilde{\mathcal{K}}^j$ that Hausdorff converges to

 $\{x_1 \geq 0\} \cap \{x_3 \geq 0\} \times (-\infty, 0]$, but such that there are surgeries at half necks of radius $r_j \to 0$ centered at (0, 0). Set $t_0^j = r_j^2/2$, and instead of Huisken's quantity Θ from [Hui90] consider Edelen's quantity Θ_R from [Ede20, Definition 5.1.1], centered at $X_0^j = (0, t_0^j)$, namely

(5.1)
$$\Theta_R\left(\widetilde{\mathcal{M}}^j, X_0^j, \tau\right) = \int_{\partial \widetilde{K}^j_{t_0^j - \tau}} \left(\theta(x, \tau) + \theta(R_j x, \tau)\right) \, dA(x),$$

where θ denotes the truncated Gaussian kernel defined by

(5.2)
$$\theta(x,\tau) = \frac{1}{4\pi\tau} \exp\left(-\frac{|x|^2}{4\tau}\right) \left(1 - \left(\frac{\rho^2}{\tau}\right)^{3/4} \frac{|x|^2 - \alpha\tau}{\rho^2}\right)_+^4,$$

for suitable choice of $\rho > 0$ and $\alpha > 0$, and $R_j x$ denotes the reflection of x across $\partial(\lambda_j D)$. For small backwards time, say $\tau_j = r_j^2$, we are δ -close to a half neck, and thus get

(5.3)
$$\liminf_{j \to \infty} \Theta_R\left(\widetilde{\mathcal{M}}^j, X_0^j, r_j^2\right) > 3/2.$$

Now, by [Ede20, Theorem 5.5] the function $\tau \mapsto \Theta_R(\widetilde{\mathcal{M}}^j, X_0^j, \tau)$ is almost monotone if there are no surgeries. Note also that discarding connected components has the good sign. Finally, arguing similarly as in the proof of [HK17, Claim 2.17] we see that the cumulative error in Edelen's monotonicity inequality due to surgeries between $\tau = r_j^2$ and $\tau = \varepsilon \rho^2$ is less than 1/100, provided j is sufficiently large. Hence, closeness to a multiplicity-one half plane at scale $\tau = \varepsilon \rho^2$ for j large enough gives the desired contradiction with (5.3).

6. Multiplicity-one for free boundary flows

In this section, we prove that every generalized limit flow has multiplicity-one. To this end, we will adapt the arguments from [EHIZ22] to our setting of flows with surgeries.

6.1. Large blowups with entropy at most two. In this subsection, fixing the initial domain $K_0 \subset D$, we consider the class C of all blowup limits $(\mathcal{K}, \mathcal{M})$ given by case (a) of Proposition 4.6 (large blowups), such that

- (1) $\lim_{\tau\to\infty} \Theta(\mathcal{M}, 0, \tau) \leq 2$ respectively $\lim_{\tau\to\infty} \Theta(\widetilde{\mathcal{M}}, 0, \tau) \leq 2$,
- (2) $(\mathcal{K}, \mathcal{M})$ respectively $(\widetilde{\mathcal{K}}, \widetilde{\mathcal{M}})$ is not a static or quasistatic multiplicity-two plane.

Here, in case $(\mathcal{K}, \mathcal{M})$ is defined in \mathbb{R}^3_+ we denote by $(\widetilde{\mathcal{K}}, \widetilde{\mathcal{M}})$ the reflected flow in \mathbb{R}^3 .

Proposition 6.1 (partial regularity). For $(\mathcal{K}, \mathcal{M}) \in \mathcal{C}$ tangent flows at singular points cannot be static or quasi-static. In particular, the dimension of the singular set is at most 1.

Here, the singular set is defined as the collection of all space-time points that do not have a backwards parabolic neighborhood in which the flow is smooth with multiplicity-one. *Proof.* In case $(\mathcal{K}, \mathcal{M})$ is defined in \mathbb{R}^3_+ we consider the reflected flow $(\widetilde{\mathcal{K}}, \widetilde{\mathcal{M}})$. By the equality case of Huisken's monotonicity formula [Hui90] and the definition of the class \mathcal{C} no tangent flow can be a static or quasistatic plane of higher multiplicity. Together with standard stratification [Whi97], remembering that we are working in \mathbb{R}^3 , the assertion follows. \Box

Corollary 6.2 (static or quasistatic limits). If $(\mathcal{K}, \mathcal{M}) \in \mathcal{C}$ is static or quasi-static, then one of the following five cases occurs:

- (1) \mathcal{K} is a (quasi-)static half space in \mathbb{R}^3 , and \mathcal{M} is the (quasi-)static plane $\partial \mathcal{K}$.
- (2) \mathcal{M} is a pair of two (quasi-)static parallel multiplicity-one planes in \mathbb{R}^3 and \mathcal{K} is the region in between.
- (3) \mathcal{K} is a (quasi-)static quarter space in \mathbb{R}^3_+ , and \mathcal{M} is the (quasi-)static half plane $\partial \mathcal{K}$ with multiplicity-one.
- (4) \mathcal{M} is a pair of(quasi-)static multiplicity-one half planes in \mathbb{R}^3_+ with free boundary and \mathcal{K} is the region in between.
- (5) \mathcal{M} is a (quasi-)static multiplicity one plane in \mathbb{R}^3_+ parallel to $\partial \mathbb{R}^3_+$, and \mathcal{K} is the region in between.

Proof. Since it is (quasi-)static, $\partial \mathcal{K}$ must be smooth and flat by Proposition 6.1 (partial regularity), and hence a union of one or two disjoint (quasi-)static multiplicity-one planes or half planes. The result then follows from one-sided minimization (Corollary 4.8).

Theorem 6.3 (separation theorem). Let $(\mathcal{K}, \mathcal{M}) \in \mathcal{C}$. In case the flow is defined in \mathbb{R}^3_+ , suppose that there is a half plane H perpendicular to $\partial \mathbb{R}^3_+$, such that $H \subseteq \bigcap_t K_t$, and suppose the complement of $\cap_t K_t$ contains points on each side of H. Then, $(\mathcal{K}, \mathcal{M})$ is static, and K_t is the region between two parallel half planes perpendicular to $\partial \mathbb{R}^3_+$. Moreover, a similar statement holds in case the flow is defined in entire space and contains a plane.

Proof. Using Proposition 6.1 (partial regularity) and Corollary 4.8 (one-sided minimization for generalized limit flows) we can follow the proof of [EHIZ22, Theorem 6.4] to show that the reflected flow $\widetilde{\mathcal{M}}$ splits into two components $\widetilde{\mathcal{M}}_{\pm}$, each contained in the respective halfspace defined by \widetilde{H} , which is obtained from H by reflection. Since each Brakke flow $\widetilde{\mathcal{M}}_{\pm}$ has density at least 1, but the sum of densities is at most 2, each $\widetilde{\mathcal{M}}_{\pm}$ must have density exactly 1 and hence be a multiplicity-one plane. Finally, since $\widetilde{K}_t \supseteq \widetilde{H}$ is in particular nonempty for all t, we conclude that each plane $\widetilde{\mathcal{M}}_{\pm}$ is static. This implies the assertion.

Now, as in [Whi00, Section 4], for a set $S \subseteq D$, a point $x \in D$, and a radius r > 0, the relative thickness of S in B(x, r) is defined by

(6.1)
$$\operatorname{Th}(S, x, r) = \frac{1}{r} \inf_{|v|=1} \sup_{y \in S \cap B(x, r)} |\langle v, y - x \rangle|.$$

Theorem 6.4 (Bernstein-type theorem). There exists an $\varepsilon > 0$ with the following significance. If $(\mathcal{K}, \mathcal{M}) \in \mathcal{C}$ is defined in \mathbb{R}^3_+ and there is a point x such that

(6.2)
$$\limsup_{r \to \infty} \operatorname{Th}(K_{-r^2}, x, r) < \varepsilon \quad \text{and} \quad \liminf_{r \to \infty} \frac{\operatorname{dist}(K_{r^2}, x)}{r} < 1,$$

then \mathcal{M} is either a pair of static parallel multiplicity-one half planes with free boundary or a static multiplicity-one plane parallel to $\partial \mathbb{R}^3_+$. In either case \mathcal{K} is the region in between the planes of \mathcal{M} and $\partial \mathbb{R}^3_+$. Similarly, if $(\mathcal{K}, \mathcal{M})$ is defined in \mathbb{R}^3 , then under the same assumptions \mathcal{M} is a pair of static parallel multiplicity-one planes, with \mathcal{K} the region in between.

Proof. The statement for flows in \mathbb{R}^3 follows from the proof of [Has23, Theorem 6.4], so we focus on the case of free boundary flows in \mathbb{R}^3_+ .

Since \mathcal{K} is nested and does not bump into any smooth subsolution of the free boundary mean curvature flow, choosing ε small enough we can apply the expanding hole result from [EHIZ22, Corollary 6.7] to infer that $\Sigma := \bigcap_t K_t$ is nonempty. Now, consider the flows obtained by translating $(\mathcal{K}, \mathcal{M})$ by (0, -T) and let $(\mathcal{K}', \mathcal{M}')$ be a limit as $T \to \infty$. Then $(\mathcal{K}', \mathcal{M}')$ is a static flow with $K'_t = \Sigma$ at any time t. Note that $(\mathcal{K}', \mathcal{M}')$ either is a multiplicitytwo half plane, or belongs to the class \mathcal{C} and hence by Corollary 6.2 (static or quasistatic limits) and assumption (6.2) is either the region in between a pair of multiplicity-one free boundary half planes or the region bounded by $\partial \mathbb{R}^3_+$ and a parallel multiplicity-one plane. Hence, Theorem 6.3 (separation theorem), applied directly in the former cases and applied to the reflected flow in the final case, respectively, implies the result.

6.2. Sheeting theorem for Λ -blowup sequences. In this subsection, we establish a sheeting theorem for Λ -blowup sequences with $\Lambda > 0$. We start with the following lemma:

Lemma 6.5 (slab rescaling). Let $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ be a Λ -blowup sequence with $\Lambda > 0$, and suppose that

(6.3)
$$d_{\mathrm{H}}(\widetilde{\mathcal{K}}^{j} \cap P(0,2), (V \times \mathbb{R}) \cap P(0,2)) \to 0,$$

where V is either a plane or a half plane. Then, there exists a sequence $\mu_j \to \infty$, such that $\mathcal{D}_{\mu_j} \widetilde{\mathcal{M}}^j$ converges smoothly to either (a) a pair of parallel planes in \mathbb{R}^3 or (b) a pair of parallel half planes with free boundary in \mathbb{R}^3_+ or (c) a multiplicity-one plane parallel to $\partial \mathbb{R}^3_+$. Moreover, in all cases $\mathcal{D}_{\mu_j} \widetilde{\mathcal{K}}^j$ converges to the enclosed region.

Proof. Fixing $\varepsilon > 0$ small enough, let $\mu_j < \infty$ be the largest number such that for all $r \in [\mu_j^{-1}, 1]$ we have

(6.4)
$$\operatorname{Th}(\tilde{K}^{j}_{-r^{2}}, 0, r) \leq \varepsilon \quad \text{and} \quad d(\tilde{K}^{j}_{r^{2}}, 0) \leq r.$$

Assumption (6.3) implies that $\mu_j \to \infty$, so remembering also that $\Lambda > 0$ we in particular get $\mu_j \lambda_j / H^j_{\text{neck}} \to \infty$, and thus by Proposition 4.6 (large blowups) we can take a subsequential limit of $\mathcal{D}_{\mu_j}(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$. By construction, any such limit $(\mathcal{K}, \mathcal{M})$ satisfies

(6.5)
$$\operatorname{Th}(K_{-r^2}, 0, r) \le \varepsilon \quad \text{and} \quad d(K_{r^2}, 0) \le r \quad \text{for all } r \ge 1,$$

with at least one inequality being non-strict for r = 1. In particular, we must be in case (a) of Proposition 4.6. Taking also into account Corollary 4.8 (one-sided minimization for generalized limit flows) we thus infer that $(\mathcal{K}, \mathcal{M}) \in \mathcal{C}$. Hence, Theorem 6.4 (Bernstein-type theorem) and the local regularity theorem [Whi05, Ede20] imply the assertion. Denote by $D^j = \lambda_j D$ the domain of the rescaled flow $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$. To construct a separating surface in case (a) and (b) of the above lemma, we let S_t^j be the set of centers of open balls B, such that $B \cap D^j \subseteq K_t^j$ and $\overline{B} \cap D^j$ touches ∂K_t^j at two or more points. Set

(6.6)
$$\mathcal{S}^j = \bigcup_t S^j_t \cap D^j.$$

Theorem 6.6 (sheeting theorem). Let $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$ be a Λ -blowup sequence with $\Lambda > 0$, and suppose that

(6.7)
$$d_{\mathrm{H}}(\mathcal{K}^{j} \cap P(0,4), (V \times \mathbb{R}) \cap P(0,4)) \to 0,$$

where either $D^j \to \mathbb{R}^3$ and V is a plane, or $D^j \to \mathbb{R}^3_+$ and V is a free boundary half plane. Then, for all j large, $S^j \cap P(0,1)$ is a C^1 -hypersurface that divides $\partial \mathcal{K}^j$ into two nonempty components.

Proof. Observe that thanks to (6.7) and $\Lambda > 0$, for *j* large enough there are no points modified by surgery in P(0,3). Hence, following the proof of [EHIZ22, Theorem 6.10], where we now use Lemma 6.5 (slab rescaling) in lieu of [EHIZ22, Lemma 6.9], yields the assertion.

6.3. Ruling out generalized limit flows with density two. As above, fixing the initial condition $K_0 \subset D$, we consider all generalized limit flows $(\mathcal{K}, \mathcal{M})$ as in Definition 4.7. We recall that they arise as limits of Λ -blowup sequences, where $\Lambda \in (0, \infty]$.

Given any closed subset $\mathcal{K}' \subset \mathbb{R}^3 \times \mathbb{R}$, similarly as in [Whi00, Section 9] we denote by $\phi(\mathcal{K}')$ the infimum of s > 0 such that

(6.8)
$$d_{\mathrm{H}}(\mathcal{K}' \cap P_{-}(0, 1/s), (V \times \mathbb{R}) \cap P_{-}(0, 1/s)) < s$$

for some plane $V \subset \mathbb{R}^3$ through the origin, and similarly as in [EHIZ22, Section 6] we denote by $\phi_+(\mathcal{K}')$ the infimum of s > 0 such that

(6.9)
$$d_{\mathrm{H}}(\mathcal{K}' \cap P_{-}(0, 1/s), (H \times \mathbb{R}) \cap P_{-}(0, 1/s)) < s$$

for some half plane $H \subset \mathbb{R}^3_+$ that meets $\partial \mathbb{R}^3_+$ orthogonally at the origin.

Lemma 6.7 (isolation). There exists an $\varepsilon > 0$, such that if $(\mathcal{K}', \mathcal{M}')$ is a tangent flow to a generalized limit flow at X = (0, t), then we have:

- (1) If 0 is a boundary point and $\phi_+(\mathcal{K}') < \varepsilon$, then $\phi_+(\mathcal{K}') = 0$.
- (2) If 0 is a boundary point and $\phi(\mathcal{K}') < \varepsilon$, then $\phi(\mathcal{K}') = 0$.
- (3) If 0 is an interior point and $\phi(\mathcal{K}') < \varepsilon$, then $\phi(\mathcal{K}') = 0$.

Proof. Observing that the argument from the proof of [Has23, Lemma 6.8] already shows (3), it suffices to show (1) and (2). Specifically, given any sequence of tangent flows $(\mathcal{K}^j, \mathcal{M}^j)$ at $X_j = (0, t_j)$ to Λ_j -generalized limit flows in \mathbb{R}^+_3 , where $\Lambda_j > 0$, with $\phi_+(\mathcal{K}^j) \to 0$ or $\phi(\mathcal{K}^j) \to 0$, we must show that for large j we have $\phi_+(\mathcal{K}^j) = 0$ or $\phi(\mathcal{K}^j) = 0$, respectively.

Note that in particular each $(\mathcal{K}^j, \mathcal{M}^j)$ is an ∞ -generalised limit flow. Hence, in case (1) as a consequence of Theorem 6.6 (sheeting theorem), similarly as in [EHIZ22, Lemma 6.16], for large j there are functions f_j and g_j , defined on an exhaustion of $\mathbb{R}^2_+ \times (-\infty, 0)$, such that:

- (1) Either $f_j < g_j$ everywhere, or $f_j \equiv g_j$.
- (2) For any $U \subset \mathbb{R}^3_+$ and $[a,b] \subset (-\infty,0)$, for j large enough the region K^j_t coincides in U with the region between graph (f_i) and graph (g_i) for all $t \in [a,b]$.
- (3) f_j and g_j converge smoothly on compact subsets to 0.
- (4) f_j and g_j solve the graphical (free boundary) mean curvature flow equation.
- (5) f_i and g_i are nondecreasing and nonincreasing in time, respectively.

A similar statement holds in case (2), where the functions f_j and g_j are now defined on an exhaustion of $\partial \mathbb{R}^3_+ \times (-\infty, 0)$ and we can simply take $f_j \equiv 0$.

Moreover, since $\Lambda_j > 0$, by monotonicity each tangent flow is backwardly self-similar, so

(6.10)
$$f_j(rx, r^2t) = rf_j(x, t), \qquad g_j(rx, r^2t) = rg_j(x, t).$$

Now, if $f_j < g_j$ for infinitely many j, then using the Harnack inequality similarly as in [Whi00, Case 1 in the proof of Theorem 9.1], where in case (1) we consider the reflected functions on $\mathbb{R}^2 \times (-\infty, 0)$, we can find $c_j > 0$, such that a subsequence of $c_j(g_j - f_j)$ converges smoothly on compact subsets to the constant function $u \equiv 1$. However, using (6.10) we infer that $u(rx, r^2t) = ru(x, t)$, which is absurd.

Thus, $f_j \equiv g_j$ for large j. But then the functions are constant in t, and together with the self-similarity we infer that $f_j \equiv g_j$ is 1-homogenous. Remembering smoothness this implies linearity. Taking also into account the free boundary condition in case (1), we conclude that $\phi_+(\mathcal{K}^j) = 0$ or $\phi(\mathcal{K}^j) = 0$, respectively, for j large.

Lemma 6.8 (minimal surface). Let $(\mathcal{K}, \mathcal{M})$ be a generalized limit flow. Suppose $(\mathcal{K}', \mathcal{M}')$ is a tangent flow of $(\mathcal{K}, \mathcal{M})$ taken taken at a density-two point X = (x, t). Then:

- (1) If x is a boundary point, and \mathcal{K}' is a static or quasistatic half plane, then there exist an open neighborhood U of x in \mathbb{R}^3_+ , an open interval (a, b), and a properly embedded smooth free boundary minimal surface Σ in U, such that $K_{\tau} \cap U = \Sigma$ for all $\tau \in (a, b)$.
- (2) If x is a boundary point, and \mathcal{K}' is a static or quasistatic plane, then there exists a neighborhood U of 0 in \mathbb{R}^3_+ , and an open interval (a, b), such that $K_{\tau} \cap U = \partial \mathbb{R}^3_+ \cap U$ for all $\tau \in (a, b)$.
- (3) If x is an interior point, and \mathcal{K}' is a static or quasistatic plane, then there is an open neighborhood U of x in \mathbb{R}^3 , an open interval (a, b), and a properly embedded smooth minimal surface Σ in U, such that $K_{\tau} \cap U = \Sigma$ for all $\tau \in (a, b)$.

Furthermore, in the cases (1) and (2) we have $\Theta(\widetilde{\mathcal{M}}, \cdot) \geq 2$ on all of $\widetilde{\Sigma} \times (-\infty, b]$ or $(\partial \mathbb{R}^3_+ \cap U) \times (-\infty, b]$, respectively, and in case (3) we have $\Theta(\mathcal{M}, \cdot) \geq 2$ on all of $\Sigma \times (-\infty, b]$.

Proof. Note that there are no surgeries in a spacetime neighborhood of X. Hence, the assertions (1), (2) and (3) follow from Lemma 6.7 (isolation) and Theorem 6.6 (sheeting theorem) exactly as in the proof of [EHIZ22, Lemma 6.22].

For the density, in case (1) let \mathcal{Z} be the set of points of \mathcal{M} at which none of the tangent flows is planar (which is necessarily in the complement of the surgery region). By general stratification results [Whi97], the parabolic Hausdorff dimension of \mathcal{Z} is at most 1. In particular, the spatial projection $\pi(\mathcal{Z})$ has Hausdorff dimension at most 1. So by upper semicontinuity of the density, it is enough to show that $\Theta(\widetilde{\mathcal{M}}, (y, t)) = 2$ for all $y \in \widetilde{\Sigma} \setminus \pi(\mathcal{Z})$

14

and t < b. To this end, fix $y \in \widetilde{\Sigma} \setminus \pi(\mathcal{Z})$, and let $T^* = \sup\{\tau < b \mid \Theta(\widetilde{\mathcal{M}}, (y, \tau)) \neq 2\}$. Clearly $T^* \leq a$. Suppose towards a contradiction that $T^* > -\infty$. Note that a neighborhood of (y, T^*) is unmodified by surgeries, since otherwise the density near (y, T^*) would be less than 2. Now, consider a tangent flow $(\mathcal{K}', \mathcal{M}')$ at (y, T^*) . If it is a static multiplicity 2 plane, then applying the first part of the theorem shows that the density is 2 for times close to T^* , which contradicts the definition of T^* . If $(\mathcal{K}', \mathcal{M}')$ was a quasistatic plane or a static plane of multiplicity 1, then we would obtain a contradiction with the fact that $\Theta(\widetilde{\mathcal{M}}, (y, \tau)) = 2$ for $\tau \in [T^*, b)$. Thus, $(y, T^*) \in \mathcal{Z}$, contradicting the choice of y. Observing that a similar argument applies in case (2) and (3) as well, this concludes the proof of the lemma.

Theorem 6.9 (multiplicity-one for generalized limit flows). Given any mean-convex initial data $K_0 \subset D$, static or quasistatic density-two planes cannot occur as generalized limit flows.

Proof. Given a mean-convex initial condition $K_0 \subset D$, denote by $\varepsilon > 0$ the smaller one of the two constants from Lemma 6.7 (isolation) and [EHIZ22, Theorem 6.20]. Consider any Λ -blowup sequence $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j) = (\mathcal{D}_{\lambda_j} \mathcal{K}^j, \mathcal{D}_{\lambda_j} \mathcal{M}^j)$, where for ease of notation we pretend that $D \subset \mathbb{R}^3_+$ and $0 \in \partial \mathcal{K}^j$. Thanks to Proposition 4.1 (small blowups) and the multiplicity-one theorem for blowup limits of free boundary mean-convex level set flow from [EHIZ22] we may assume that $\Lambda > 0$. Now, suppose towards a contradiction that $(\widetilde{\mathcal{K}}^j, \widetilde{\mathcal{M}}^j)$, converges to a limit $(\widetilde{\mathcal{K}}^{\infty}, \widetilde{\mathcal{M}}^{\infty})$, which is a static or quasistatic density-two plane or half plane.

We first analyze the case where $(\widetilde{\mathcal{K}}^{\infty}, \widetilde{\mathcal{M}}^{\infty})$ is a density-two half plane. Let $\mu_j > 0$ be the largest number such that

(6.11)
$$\phi_+(\mathcal{D}_{\mu_j}\mathcal{K}^j) \ge \varepsilon/2.$$

Note that $\mu_j \to 0$. After passing to a subsequence, we can also assume that $\mathcal{D}_{\mu_i} \widetilde{\mathcal{K}}^j$ Hausdorff converges to a limit \mathcal{K} . By construction, we have

(6.12)
$$\phi_+(\mathcal{K}) \ge \varepsilon/2 \quad \text{but} \quad \phi_+(\mathcal{D}_\lambda \mathcal{K}) \le \varepsilon/2 \text{ for } \lambda \ge 1.$$

If we had $\mu_j \lambda_j / H^j_{\text{neck}} \to 0$, then by Proposition 4.1 (small blowups) we would see that \mathcal{K} is a blowup limit or homothetic copy of the level set flow. But then, using (6.12) and arguing similarly as in the prof of [EHIZ22, Theorem 6.23], we could construct a blowup limit of the free boundary level set flow that is a non-planar minimal cone. Thus, after passing to a subsequence we can assume that $\mu_j \lambda_j / H^j_{\text{neck}} \to \Lambda' > 0$. In particular, $\mu_j \lambda_j \to \infty$ and \mathcal{K} is a generalized limit flow that comes with a family of Radon measures \mathcal{M} , which is provided by Theorem 4.5 (hybrid compactness) or Proposition 4.6 (large blowups), respectively.

Now, by Lemma 6.7 (isolation) and (6.12), any tangent flow to $(\mathcal{K}, \mathcal{M})$ at the space-time origin must be a static or quasistatic multiplicity-two half plane. So we can apply Lemma 6.8 (minimal surface) to obtain a free boundary minimal surface $\Sigma \subset \mathbb{R}^3_+$ containing the origin and a real number b, such that the associated reflected quantities satisfy

(6.13)
$$\Theta(\widetilde{\mathcal{M}}, \cdot) \ge 2 \quad \text{on} \quad \widetilde{\Sigma} \times (-\infty, b].$$

In particular, it follows that $\Lambda' < \infty$. Now translate $(\mathcal{K}, \mathcal{M})$ by $(x, t) \mapsto (x, t+j)$ and using Theorem 4.5 (hybrid compactness) along $j \to \infty$ pass to a subsequential limit to get a static

Brakke δ -flow ($\mathcal{K}', \mathcal{M}'$) satisfying

(6.14)
$$K'_t = \bigcup_s K_s \quad \forall t, \quad \text{and} \quad \Theta(\widetilde{\mathcal{M}}', \cdot) \ge 2 \text{ on } \widetilde{\Sigma} \times (-\infty, \infty).$$

Observe that since the flow is static, it does not contain any surgeries. Moreover, by (6.12) it is not planar. Hence, taking a tangent flow at $-\infty$ to $(\widetilde{\mathcal{K}}', \widetilde{\mathcal{M}}')$ we obtain a non-planar static minimal cone, which gives the desired contradiction.

Finally, if $(\widetilde{\mathcal{K}}^{\infty}, \widetilde{\mathcal{M}}^{\infty})$ is a density-two plane, then we can obtain a similar contradiction as above, provided we now work with the more general quantity $\psi = \min(\varphi, \varphi_h)$, similarly as in the proof of [EHIZ22, Theorem 6.23]. This concludes the proof of the theorem. \Box

7. PARTIAL REGULARITY AND CONVEXITY

In this section, we show that generalized limit flows have small singular set and nonnegative second fundamental form.

Theorem 7.1 (partial regularity). The singular set of any generalized limit flow has parabolic Hausdorff dimension at most 1.

Proof. By Corollary 4.8 (one-sided minimization for generalized limit flows) and Theorem 6.9 (multiplicity-one for generalized limit flows) nontrivial cones and higher-multiplicity planes cannot occur as tangent flows of generalized limit flows. Using this, the assertion follows from standard dimension reduction and the local regularity theorem [Whi97, Whi05, Ede20]. \Box

Proposition 7.2 (rigidity). Let $(\mathcal{K}, \mathcal{M})$ be a generalized limit flow. If $0 \in \partial K_0$ is a regular point, and H(0,0) = 0, then $(\mathcal{K}, \mathcal{M}) \cap \{t \leq 0\}$ is a flat density-one plane or half plane.

Proof. Suppose that (0,0) is a regular point and H(0,0) = 0. Then by the strict maximum principle, possibly applied to the reflected flow, we have $H \equiv 0$ in some backwards parabolic ball $P_{-}(0,\rho)$. By Theorem 7.1 (partial regularity) we can choose a time $t_0 \in (-\rho^2, 0)$ at which the solution is completely smooth. Then, again by the strict maximum principle, there is an entire connected component $\Sigma \subset M_{t_0}$ that contains the origin and on which the mean curvature vanishes identically. Note that Σ must be noncompact, since there are no compact minimal surfaces in Euclidean space and no compact free boundary minimal surfaces in Euclidean halfspace. Next, again by the smallness of the singular set any $X \in$ $\Sigma \times (-\infty, t_0] \setminus \text{sing} \mathcal{M}$ can be connected to $(0, t_0)$ by a time-like space-time curve that entirely avoids the singular set. Together with the strict maximum principle this yields

(7.1)
$$\Sigma \times (-\infty, t_0] \subseteq \partial \mathcal{K}.$$

In particular, there are no surgeries near Σ . Now, by Corollary 4.8 (one-sided minimization for generalized limit flows) and Theorem 6.9 (multiplicity-one for generalized limit flows) in the case $\Lambda > 0$, and by Proposition 4.1 (convergence to level-set flow) and the one-sided minimization and multiplicity-one theorem for blowups of the level-set flow from [EHIZ22] in the case $\Lambda = 0$, the tangent cone at infinity of Σ or $\widetilde{\Sigma}$, respectively, must be a multiplicityone plane. Hence, by monotonicity, Σ or $\widetilde{\Sigma}$, respectively, is flat. Finally, by White's strong half space result [Whi03, Theorem 7] there cannot be any other connected components. \Box

Theorem 7.3 (nonnegative second fundamental form). Let $(\mathcal{K}, \mathcal{M})$ be a generalized limit flow. Then at every regular point all principal curvatures are nonnegative.

Proof. Fixing $K_0 \subset D$, suppose towards a contradiction that there is a sequence of generalized limit flows $(\mathcal{K}^j, \mathcal{M}^j)$ and a sequence of regular points X_j such that $\frac{\lambda_1}{H}(X_j)$ converges to an infimal value $\gamma < 0$. Note that $\gamma > -\infty$ thanks to the bound $|A| \leq CH$ from Proposition 2.5 (basic properties). Moreover, by translating and scaling we may assume that $X_j = 0$ and

(7.2)
$$\sup_{P(0,1)} |A_{\partial \widetilde{\mathcal{K}}^j}| \le 1 \le \sup_{\overline{P(0,1)}} |A_{\partial \widetilde{\mathcal{K}}^j}|.$$

If there is no r > 0 such that the flow is unmodified by surgeries in P(0, r), then after adjusting our sequence, using in particular item (c) of Definition 2.3 (replacement by standard caps or standard half caps), we may assume that (0, 0) lies in the presurgery domain.

If $(\mathcal{K}^j, \mathcal{M}^j)$ is a 0-blowup sequence, then using Proposition 4.1 (small blowups) and Theorem 5.1 (no microscopic surgeries) we obtain contradiction with the convexity theorem for blowup limits of the free boundary level-set flow from [EHIZ22]. Hence, by Theorem 4.5 (hybrid compactness) and Proposition 4.6 (large blowups) we may assume that $(\mathcal{K}^j, \mathcal{M}^j)$ converges to a generalized limit flow $(\mathcal{K}, \mathcal{M})$. By (7.2) and Proposition 7.2 (rigidity) the limit $(\mathcal{K}, \mathcal{M})$ must have strictly positive mean curvature. Hence λ_1/H attains a strictly negative minimum at the space-time origin, which, possibly after considering the reflected flow, contradicts the strict maximum principle. This proves the theorem.

8. CANONICAL NEIGHBORHOODS AND EXISTENCE THEOREM

In this final section, we prove the canonical neighborhood theorem and the existence theorem for free boundary flows with surgery.

Theorem 8.1 (canonical neighborhoods). Suppose \mathcal{K}^j is a sequence of (δ, \mathcal{H}^j) -flows starting at a smooth compact strictly mean-convex domain $K_0 \subset D$, such that $\delta \leq \overline{\delta}$ and $\mathcal{H}^j \to \infty$. Then, for any sequence of space-time points $X_j = (x_j, t_j) \in \partial \mathcal{K}^j$ with $\sup_j t_j < \infty$ and $H(X_j) \to \infty$, the rescaled flows $\mathcal{D}_{H(X_j)}(\mathcal{K}^j - X_j)$ subsequentially converge to either

- the evolution of a standard cap preceded by a round shrinking cylinder, or a round shrinking cylinder, round shrinking sphere, translating bowl or ancient oval, or
- the evolution of a standard half cap preceded by a round shrinking half cylinder, or a round shrinking half cylinder, round shrinking half sphere, translating half bowl or ancient half oval.

Proof. Consider the blowup sequence $\widetilde{\mathcal{K}}^j := \mathcal{D}_{\lambda_j}(\mathcal{K}^j - X_j)$, where λ_j is chosen such that (8.1) $\sup_{P(0,1)} |A_{\partial \widetilde{\mathcal{K}}^j}| \le 1 \le \sup_{\overline{P(0,1)}} |A_{\partial \widetilde{\mathcal{K}}^j}|.$ In light of Proposition 4.1 (small blowups) and Proposition 4.6 (large blowups) we may assume that $\lambda_j/H_{\text{neck}}^j \to \Lambda \in (0, \infty)$, since [EHIZ22, Corollary 1.3] or the argument from its proof, respectively, already yields the conclusion in the other cases. Hence, by Theorem 4.5 (hybrid compactness), considering the associated family of Radon measures $\widetilde{\mathcal{M}}^j$ defined as in (4.3), we can pass to a subsequential limit $(\mathcal{K}, \mathcal{M})$. Define $\mathcal{K}^0 = \{K_t^0\}_{t\leq 0}$ by for each $t \leq 0$ setting K_t^0 to be the connected component of K_t that contains the origin. Note that all these sets in fact contain a closed ball B of positive radius thanks to (8.1). By Theorem 7.1 (partial regularity) together with Theorem 7.3 (nonnegative second fundamental form) and connectedness, the sets K_t^0 are smooth and convex for almost every t. Remembering in particular the way surgeries are performed, we see that convexity in fact holds at all $t \leq 0$. Now given any $p \in \partial K_t^0$ the convex hull of p and B is contained in K_t^0 , and consequently, remembering Corollary 4.8 (one-sided minimization for generalized limit flows) and Theorem 6.9 (multiplicity-one for generalized limit flows), any tangent flow at (p, t) must be a density-one plane or half plane. Hence, K_t^0 is smooth and convex for all $t \leq 0$.

If there are no surgeries, then by the classification from [BC19, ADS20], the flow \mathcal{K}^0 or its reflection $\widetilde{\mathcal{K}^0}$, respectively, must be a round shrinking cylinder, round shrinking sphere, translating bowl or ancient oval. Assume now \mathcal{K}^0 does contain a surgery, let $T \leq 0$ be a surgery time and let $N \subset K_T^0$ be a surgery neck or half neck of quality $\overline{\delta}$ sitting in the backward time slice. Note that N is the limit of some \widetilde{N}^j in the approximators $\widetilde{\mathcal{K}}^j$. By part (b) of Definition 2.4 (free boundary flow with surgery) we can find a curve γ_j in the approximator connecting $\{H = H_{\text{trig}}^j\}$ and $\{H \leq H_{\text{th}}^j\}$, such that it passes through \widetilde{N}^j but avoids all other $\overline{\delta}$ -necks and half $\overline{\delta}$ -necks of the disjoint collection. We can assume that the curve γ_j enters and leaves \widetilde{N}^j exactly once. Let p_j be the center of \widetilde{N}^j . Since \mathcal{K}^0 is smooth with strictly positive mean curvature, and since $\mathcal{H}^j \to \infty$, given any $R < \infty$, for j large enough the curve γ_j must start and end outside $B(p_j, R)$. Thus, $K_T^{0,-} \setminus N$ has at least two unbounded components. Since $K_T^{0,-}$ is connected, $K_T^{0,-} \setminus N$ must have exactly two components. We have thus shown that $K_T^{0,-}$ has two ends, and consequently it contains a line, and all prior time slices contain this line as well. Hence, at each fixed time the convex set splits off an \mathbb{R} -factor, and thus there cannot be any other surgeries. It follows that \mathcal{K}^0 or $\widetilde{\mathcal{K}^0}$, respectively, is a round cylindrical flow for t < T followed by the unique evolution of the standard cap for t > T.

Finally, since all the ancient solutions from the above sweep out the entire space for $t \to -\infty$ it follows that there are in fact no other connected components, i.e. $\mathcal{K}^0 = \mathcal{K}$, and since $H(X_j) \to H(0) \in (0, \infty)$, remembering also Theorem 5.1 (no microscopic surgeries), we can rescale by $H(X_j)$ instead of λ_j to conclude the proof of the theorem.

We can now prove our main theorem, which we restate here for convenience of the reader:

Theorem 8.2 (free boundary flow with surgery). Given any smooth compact strictly meanconvex free boundary domain $K_0 \subset D$, for suitable choice of the surgery parameters δ and \mathcal{H} , there exists a free boundary (δ, \mathcal{H}) -flow $\{K_t\}_{t \in [0,\infty)}$ starting at K_0 . Moreover, the flow either becomes extinct in finite time or for $t \to \infty$ converges smoothly in the one or two-sheeted sense to a finite collection of stable connected minimal surfaces with empty or free boundary (in particular, there are no surgeries for t sufficiently large).

Proof. Given any $T < \infty$, we will first prove the existence of a free boundary (δ, \mathcal{H}) -flow $\{K_t\}$ starting at K_0 and defined on the finite interval [0, T] via a continuity argument similarly as in [HK17, Section 4.2]. To this end, fixing $\bar{\delta} > 0$ small enough, suppose towards a contradiction that there is a sequence \mathcal{K}^j of free boundary (δ, \mathcal{H}_j) -flows with $\delta \leq \bar{\delta}$ and $\mathcal{H}^j \to \infty$, that can only be defined on a maximal time interval $[0, T_j]$ for some $T_j < T$. Then, it must be the case that we cannot find a minimal collection of strong δ -necks and strong half δ -necks in $K_{T_j}^j$ as required in Definition 2.4 (free boundary flow with surgery), since otherwise we could perform surgeries along an 'innermost' such collection of centers p, i.e. one for which $\sum_p \text{dist}(p, \{H = H_{\text{trig}}^j\})$ is minimal, and run smooth free boundary mean curvature flow for a short time, contradicting the maximality of T_j . So our goal is to produce a minimal separating collection of strong δ -necks for large j.

Let \mathcal{I}_j be the set of points $p \in \partial K_{T_j}^j$ with $H(p) > H_{\text{neck}}^j$, and let \mathcal{J}_j be the set of points $p \in \partial K_{T_j}^j$ with $H(p) = H_{\text{neck}}^j$. Then, similarly as in [HK17, Claim 4.6] there is a large constant $C < \infty$, such that the union $V_j = \bigcup_{p \in \mathcal{J}_j} B(p, CH^{-1}(p))$, for j large enough, separates $\{H = H_{\text{trig}}^j\}$ from $\{H \leq H_{\text{th}}^j\}$ in the domain $K_{T_j}^j$. Let $\hat{\mathcal{J}}_j \subseteq \mathcal{J}_j$ be a minimal subset such that the union of balls $\bigcup_{p \in \hat{\mathcal{J}}_j} B(p, CH^{-1}(p))$ has the separation property. Then, using Theorem 8.1 (canonical neighborhoods) and arguing similarly as in the proof of [HK17, Claim 4.7] we see that for large j, every $p \in \hat{\mathcal{J}}_j$ is a strong δ -neck point or strong half δ -neck point. This collection is disjoint for large j, since otherwise two intersecting δ -necks or half δ -necks with the separation property; this gives the desired contradiction and thus proves the existence on the interval [0, T].

Finally, it is known by [EHIZ22, Theorem 1.5] that the free boundary level set flow of K_0 either becomes extinct in finite time or or for $t \to \infty$ converges smoothly in the one or two-sheeted sense to a finite collection of stable minimal surfaces or stable free boundary minimal surfaces. Hence, applying the above result for $T < \infty$ sufficiently large and taking also into account Proposition 3.1 (distance to free boundary level set flow), by mimicking the argument from Brendle-Huisken [BH18], we can get long-time existence and convergence.

References

- [ADS20] S. Angenent, P. Daskalopoulos, and N. Sesum. Uniqueness of two-convex closed ancient solutions to the mean curvature flow. Ann. of Math. (2), 192(2):353–436, 2020.
- [And12] B. Andrews. Noncollapsing in mean-convex mean curvature flow. *Geom. Topol.*, 16(3):1413–1418, 2012.
- [BC19] S. Brendle and K. Choi. Uniqueness of convex ancient solutions to mean curvature flow in \mathbb{R}^3 . Invent. Math., 217(1):35–76, 2019.

- [BH16] S. Brendle and G. Huisken. Mean curvature flow with surgery of mean convex surfaces in \mathbb{R}^3 . Invent. Math., 203(2):615–654, 2016.
- [BH18] S. Brendle and G. Huisken. Mean curvature flow with surgery of mean convex surfaces in threemanifolds. J. Eur. Math. Soc. (JEMS), 20(9):2239–2257, 2018.
- [BHH19] R. Buzano, R. Haslhofer, and O. Hershkovits. The moduli space of two-convex embedded tori. Int. Math. Res. Not. IMRN, (2):392–406, 2019.
- [BHH21] R. Buzano, R. Haslhofer, and O. Hershkovits. The moduli space of two-convex embedded spheres. J. Differential Geom., 118(2):189–221, 2021.
- [DH22] J. Daniels-Holgate. Approximation of mean curvature flow with generic singularities by smooth flows with surgery. *Adv. Math.*, 410(part A):Paper No. 108715, 42, 2022.
- [Ede16] N. Edelen. Convexity estimates for mean curvature flow with free boundary. Adv. Math., 294:1–36, 2016.
- [Ede20] N. Edelen. The free-boundary Brakke flow. J. Reine Angew. Math., 758:95–137, 2020.
- [EHIZ22] N. Edelen, R. Haslhofer, M. Ivaki, and J. Zhu. Mean convex mean curvature flow with free boundary. Comm. Pure Appl. Math., 75(4):767–817, 2022.
- [GS93] Y. Giga and M. Sato. Neumann problem for singular degenerate parabolic equations. Differential Integral Equations, 6(6):1217–1230, 1993.
- [Has23] R. Haslhofer. Flows with surgery revisited. arXiv:2305.16267, 2023.
- [HK] R. Hashofer and D. Ketover. Free boundary minimal disks in convex balls. *in preparation*.
- [HK17] R. Haslhofer and B. Kleiner. Mean curvature flow with surgery. *Duke Math. J.*, 166(9):1591–1626, 2017.
- [HK19] R. Haslhofer and D. Ketover. Minimal 2-spheres in 3-spheres. Duke Math. J., 168(10):1929–1975, 2019.
- [HS09] G. Huisken and C. Sinestrari. Mean curvature flow with surgeries of two-convex hypersurfaces. Invent. Math., 175(1):137–221, 2009.
- [Hui90] G. Huisken. Asymptotic behavior for singularities of the mean curvature flow. J. Differential Geom., 31(1):285-299, 1990.
- [LM23] Y. Liokumovich and D. Maximo. Waist inequality for 3-manifolds with positive scalar curvature. In *Perspectives in scalar curvature. Vol. 2*, pages 799–831. World Sci. Publ., Hackensack, NJ, 2023.
- [LN21] M. Langford and H. Nguyen. Quadratically pinched hypersurfaces of the sphere via mean curvature flow with surgery. *Calc. Var. Partial Differential Equations*, 60(6):Paper No. 216, 33, 2021.
- [MW21] A. Mramor and S. Wang. Low entropy and the mean curvature flow with surgery. *Calc. Var. Partial Differential Equations*, 60(3):Paper No. 96, 28, 2021.
- [Ngu20] H. Nguyen. High codimension mean curvature flow with surgery. arXiv:2004.07163, 2020.
- [Whi97] B. White. Stratification of minimal surfaces, mean curvature flows, and harmonic maps. J. Reine Angew. Math., 488:1–35, 1997.
- [Whi00] B. White. The size of the singular set in mean curvature flow of mean convex sets. J. Amer. Math. Soc., 13(3):665–695, 2000.
- [Whi03] B. White. The nature of singularities in mean curvature flow of mean-convex sets. J. Amer. Math. Soc., 16(1):123–138, 2003.
- [Whi05] B. White. A local regularity theorem for mean curvature flow. Ann. of Math. (2), 161(3):1487–1519, 2005.

ROBERT HASLHOFER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, 40 ST GEORGE STREET, TORONTO, ON M5S 2E4, CANADA

E-mail: roberth@math.toronto.edu

20