

# On the $\alpha$ -index of minimally $k$ -(edge-)connected graphs for small $k$ \*

Jiayu Lou<sup>a,b</sup>, Ligong Wang<sup>a,b,†</sup>, Ming Yuan<sup>a</sup>

<sup>a</sup>School of Mathematics and Statistics, Northwestern Polytechnical University,  
Xi'an, Shaanxi 710129, P.R. China.

<sup>b</sup> Xi'an-Budapest Joint Research Center for Combinatorics, Northwestern Polytechnical University,  
Xi'an, Shaanxi 710129, P.R. China.

## Abstract

Let  $G$  be a graph with adjacency matrix  $A(G)$  and let  $D(G)$  be the diagonal matrix of vertex degrees of  $G$ . For any real  $\alpha \in [0, 1]$ , Nikiforov defined the  $A_\alpha$ -matrix of a graph  $G$  as  $A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$ . The largest eigenvalue of  $A_\alpha(G)$  is called the  $\alpha$ -index or the  $A_\alpha$ -spectral radius of  $G$ . A graph is minimally  $k$ -(edge-)connected if it is  $k$ -(edge-)connected and deleting any arbitrary chosen edge always leaves a graph which is not  $k$ -(edge-)connected. In this paper, we characterize the minimally 2-edge-connected graphs and minimally 3-connected graph with given order having the maximum  $\alpha$ -index for  $\alpha \in [\frac{1}{2}, 1)$ , respectively.

**Key Words:** minimally 2-edge-connected graph, minimally 3-connected graph,  $\alpha$ -index, extremal graph

**AMS Subject Classification (2020):** 05C50, 05C40, 05C35.

## 1 Introduction

Throughout this paper, we only consider simple, finite and undirected graphs. Let  $G = (V(G), E(G))$  be a graph with vertex set  $V(G) = \{v_1, v_2, \dots, v_n\}$  and edge set  $E(G)$ . The order of  $G$  is the number  $n = |V(G)|$  of its vertices and the size of  $G$  is the number  $m = |E(G)|$  of its edges. The neighborhood  $N_G(v)$  (or,  $N(v)$  for short) of a vertex  $v$  is  $\{u \in V(G) : uv \in E(G)\}$  and the number  $d_G(v) = |N_G(v)|$  (or,  $d(v)$  for short) is the degree of  $v$ . The closed neighbor set is defined as  $N_G[v] = N_G(v) \cup \{v\}$  (or,  $N[v]$  for short). Let  $\Delta(G)$  and  $\delta(G)$  be the maximum degree and minimum degree of  $G$ , respectively. The average degree of the neighbors of a vertex  $u$  of  $G$  is  $m(u) = \frac{\sum_{uv \in E(G)} d(v)}{d(u)}$ . For graphs  $F$  and  $G$ , we use  $H \subseteq G$  to denote that  $G$  contains  $H$  as a subgraph. For a vertex set  $S \subseteq V(G)$ , let  $G[S]$  be the subgraph of  $G$  induced by  $S$ . For  $A, B \subset V(G)$ , we denote by  $e(A)$  the number of edges in  $G[A]$  and by  $e(A, B)$  the number of edges with one endpoint in  $A$  and one endpoint in  $B$ . A

\*Supported by the National Natural Science Foundation of China (No. 12271439).

†Corresponding author.

E-mail addresses: lgwangmath@163.com, jyloumath@163.com, ym19980508@mail.nwpu.edu.cn.

cut vertex (resp., cut edge) of a graph is a vertex (resp., edge) whose deletion increases the number of components of a graph. A cycle  $C$  of  $G$  is said to have a chord if there is an edge of  $G$  that joins a pair of non-adjacent vertices from  $C$ .

Denote by  $P_n$ ,  $C_n$  and  $K_{a,n-a}$  the path, the cycle and the complete bipartite graph on  $n$  vertices, respectively. For two graphs  $G_1$  and  $G_2$ , we define  $G_1 \cup G_2$  to be their disjoint union. The join  $G_1 \vee G_2$  is obtained from  $G_1 \cup G_2$  by joining every vertex of  $G_1$  with every vertex of  $G_2$  by an edge. Let  $W_n = K_1 \vee C_{n-1}$  be the wheel graph on  $n$  vertices. For  $k \geq 1$ , denote by  $F_k$  the graph on  $2k + 1$  vertices consisting of  $k$  triangles which intersect in exactly one common vertex.

The adjacency matrix of  $G$  is defined as the matrix  $A(G) = (a_{u,v})_{u,v \in V(G)}$  with  $a_{u,v} = 1$  if  $u, v$  are adjacent, and  $a_{u,v} = 0$  otherwise. Let  $D(G)$  be the diagonal matrix of vertex degrees of  $G$ . The Laplacian matrix and signless Laplacian matrix of  $G$  are defined as  $L(G) = D(G) - A(G)$  and  $Q(G) = D(G) + A(G)$ , respectively. The largest eigenvalue of  $A(G)$  is called the index or the spectral radius of  $G$ , and the largest eigenvalue of  $Q(G)$  is called the  $Q$ -index or the signless Laplacian spectral radius of  $G$ . The largest eigenvalue of  $L(G)$  is called the  $L$ -index or the Laplacian spectral radius of  $G$ . To track the gradual change of  $A(G)$  into  $Q(G)$ , Nikiforov [19] proposed to study the convex linear combination  $A_\alpha(G)$  of  $A(G)$  and  $D(G)$  defined by

$$A_\alpha(G) = \alpha D(G) + (1 - \alpha)A(G)$$

for any real  $\alpha \in [0, 1]$ . Note that  $A(G) = A_0(G)$ ,  $D(G) = A_1(G)$  and  $Q(G) = 2A_{\frac{1}{2}}(G)$ . The largest eigenvalue of  $A_\alpha(G)$ , denoted by  $\rho_\alpha(G)$ , is called the  $\alpha$ -index or the  $A_\alpha$ -spectral radius of  $G$ . Denote by  $\lambda(M)$  the largest eigenvalue of a matrix  $M$ . For a connected graph  $G$ ,  $A_\alpha(G)$  is irreducible. By the Perron-Frobenius Theorem,  $\rho_\alpha(G)$  is positive, and there exists a unique positive unit eigenvector corresponding to  $\rho_\alpha(G)$ , which is called the  $\alpha$ -Perron vector of  $G$ .

Let  $\mathcal{G}$  be a specified class of graphs. Brualdi and Solheid [3] proposed the problem of characterizing the graphs for which the maximal or the minimal spectral invariants are attained in  $\mathcal{G}$ . For many different  $\mathcal{G}$  and different spectral invariants (such as index,  $L$ -index and  $Q$ -index), this problem has been extensively studied (see [7, 11, 16, 28]). For  $A_\alpha$ -index, there are some relative results in [5, 13, 14]. We refer the interested reader to the surveys [4, 12, 18] for more results.

A graph is  $k$ -connected (resp.,  $k$ -edge-connected) if removing fewer than  $k$  vertices (resp., edges) always leaves the remaining graph connected, and is minimally  $k$ -(edge-)connected if it is  $k$ -connected (resp.,  $k$ -edge-connected) and deleting any arbitrary chosen edge always leaves a graph which is not  $k$ -connected (resp.,  $k$ -edge-connected). In recent works, some researchers restrict  $\mathcal{G}$  to (minimally)  $k$ -(edge-)connected graphs of order  $n$  or size  $m$ . A graph is minimally 1-(edge-)connected if and only if it is a tree. It is natural to consider the problem of finding the graph with the maximum ( $Q$ -)index among all minimally  $k$ -(edge-)connected graphs for  $k \geq 2$ . Fan, Goryainov and Lin [8] asked the following question for  $k \geq 2$ .

**Problem 1.1.** *What is the maximum ( $Q$ -)index and what are the corresponding extremal graphs among minimally  $k$ -(edge-)connected graph for  $k \geq 2$ ?*

Chen and Guo [6] characterized the minimally 2-(edge-)connected graphs with given order having the maximum index. Lou, Min and Huang [17] characterized the minimally 2-(edge-)connected graphs with given size having the maximum index. Fan, Goryainov and Lin [8] not only determined the minimally 2-(edge-)connected graphs with given order having the maximum  $Q$ -index, but characterized the minimally 3-connected graphs with given order having the maximum ( $Q$ -)index. Guo and Zhang [9] characterized the (minimally) 2-connected

graphs with given size having the maximum  $Q$ -index. Zhang and Guo [27] determined the (minimally) 2-connected graphs with given size having the maximum  $L$ -index. Lou, Wang and Yuan [15] determined the minimally 2-connected graphs with given order or size having the maximum  $\alpha$ -index, respectively. They also asked the following question for  $\alpha$ -index.

**Problem 1.2.** *What is the maximum  $\alpha$ -index and what are the corresponding extremal graphs among minimally  $k$ -(edge)-connected graph for  $k \geq 2$ ?*

In this paper, we characterize the minimally 2-edge-connected graphs and minimally 3-connected graph with given order having the maximum  $\alpha$ -index for  $\alpha \in [\frac{1}{2}, 1)$ , respectively.

**Theorem 1.1.** *Let  $G$  be a minimally 2-edge-connected graph with order  $n$  and  $\alpha \in [\frac{1}{2}, 1)$ .*

- (i) *If  $n \geq 7$  is an odd number, then  $\rho_\alpha(G) \leq \rho_\alpha(F_{\frac{n-1}{2}})$ , with equality if and only if  $G \cong F_{\frac{n-1}{2}}$ .*
- (ii) *If  $n \geq 8$  is an even number, then  $\rho_\alpha(G) \leq \rho_\alpha(K_{2,n-2})$ , with equality if and only if  $G \cong K_{2,n-2}$ .*

**Theorem 1.2.** *Let  $G$  be a minimally 3-connected graph with order  $n \geq 7$ . If  $\alpha \in [\frac{1}{2}, 1)$ , then  $\rho_\alpha(G) \leq \rho_\alpha(W_n)$ , with equality if and only if  $G \cong W_n$ .*

The remainder of the paper is organized as follows. In Section 2, we recall some useful lemmas that will be used later. In Sections 3 and 4, we give proofs of Theorems 1.1 and 1.2, respectively.

## 2 Preliminaries

In this section, we introduce some useful lemmas that are used in the proof of our main results.

**Lemma 2.1.** ([19]) *If  $G$  is a graph with no isolated vertices, then*

$$\rho_\alpha(G) \leq \max_{u \in V(G)} \left\{ \alpha d(u) + \frac{1-\alpha}{d(u)} \sum_{uv \in E(G)} d(v) \right\}.$$

*If  $\alpha \in (\frac{1}{2}, 1)$  and  $G$  is connected, with equality if and only if  $G$  is regular.*

**Lemma 2.2.** ([24]) *For any graph  $G$  and  $\alpha \in [0, 1)$ . Then*

$$\rho_\alpha(G) \leq \max_{uv \in E(G)} \left\{ \frac{\alpha (d(u) + d(v)) + \sqrt{\alpha^2 (d(u) - d(v))^2 + 4(1-\alpha)^2 m(u)m(v)}}{2} \right\}.$$

*If  $G$  is connected, the equality holds if and only if  $G$  is regular or bipartite semi-regular.*

**Lemma 2.3.** ([19]) *Let  $G$  be a graph with  $\Delta(G) = \Delta$ . If  $\alpha \in [0, \frac{1}{2}]$ , then*

$$\rho_\alpha(G) \geq \alpha(\Delta + 1).$$

*If  $\alpha \in [\frac{1}{2}, 1)$ , then*

$$\rho_\alpha(G) \geq \alpha\Delta + \frac{(1-\alpha)^2}{\alpha}.$$

**Lemma 2.4.** ([2]) *If  $G$  is a minimally  $k$ -(edge-)connected graph, then  $\delta(G) = k$ .*

**Lemma 2.5.** ([2]) *If  $G$  is a minimally 2-edge-connected graph with order  $n$ , then  $m \leq 2n - 2$ .*

**Lemma 2.6.** ([17]) *If  $G$  is a minimally 2-edge-connected graph, then no cycle of  $G$  has a chord.*

**Lemma 2.7.** ([1]) *If  $G$  is a minimally 3-connected graph with order  $n$ , then each cycle contains at least two vertices of degree 3.*

**Lemma 2.8.** ([20, 25]) *Let  $G$  be a connected graph with  $\alpha \in [0, 1]$ . For  $u, v \in V(G)$ , suppose  $N \subseteq N(v) \setminus (N(u) \cup \{u\})$ . Let  $G' = G - \{vw : w \in N\} + \{uw : w \in N\}$ . Let  $X = (x_1, x_2, \dots, x_n)^T$  be the  $\alpha$ -Perron vector of  $G$  corresponding to  $\rho_\alpha(G)$ . If  $N \neq \emptyset$  and  $x_u \geq x_v$ , then  $\rho_\alpha(G') > \rho_\alpha(G)$ .*

**Lemma 2.9.** ([19]) *Let  $G_1$  be a  $r_1$ -regular graph of order  $n_1$ , and  $G_2$  be a  $r_2$ -regular graph of order  $n_2$ . Then the largest eigenvalue of  $A_\alpha(G_1 \vee G_2)$  is*

$$\rho_\alpha(G_1 \vee G_2) = \lambda \begin{pmatrix} r_1 + \alpha n_2 & (1 - \alpha)^2 n_1 n_2 \\ 1 & r_2 + \alpha n_1 \end{pmatrix}.$$

**Lemma 2.10.** ([19]) *Let  $a \geq b \geq 1$ . If  $\alpha \in [0, 1]$ , then the largest eigenvalue of  $A_\alpha(K_{a,b})$  is*

$$\rho_\alpha(K_{a,b}) = \frac{1}{2}(\alpha(a+b) + \sqrt{\alpha^2(a+b)^2 + 4ab(1-2\alpha)}).$$

It is obvious that  $\rho_\alpha(K_{2,n-2}) = \frac{1}{2}(\alpha n + \sqrt{\alpha^2 n^2 + 8(1-2\alpha)(n-2)})$ .

**Lemma 2.11.** *Let  $n$  be an odd integer. If  $\alpha \in [0, 1]$ , then the largest eigenvalue of  $A_\alpha(F_{\frac{n-1}{2}})$  is*

$$\rho_\alpha(F_{\frac{n-1}{2}}) = \frac{1}{2}(\alpha n + 1 + \sqrt{\alpha^2 n^2 - 10\alpha n + 12\alpha + 4n - 3}).$$

**Proof.** Since  $F_{\frac{n-1}{2}} = K_1 \vee \frac{n-1}{2}K_2$ , then by Lemma 2.9, we have

$$\rho_\alpha(F_{\frac{n-1}{2}}) = \lambda \begin{pmatrix} \alpha(n-1) & (1-\alpha)^2(n-1) \\ 1 & 1+\alpha \end{pmatrix}.$$

By computation, we have

$$\rho_\alpha(F_{\frac{n-1}{2}}) = \frac{1}{2}(\alpha n + 1 + \sqrt{\alpha^2 n^2 - 10\alpha n + 12\alpha + 4n - 3}). \quad \square$$

### 3 Proof of Theorem 1.1

In this section, we give the proof of Theorem 1.1.

**Proof.** (i) Let  $G$  be the graph having the maximum  $\alpha$ -index among all minimally 2-edge-connected graphs with odd order  $n \geq 7$ . Then  $\Delta(G) \leq n-1$ . Notice that  $F_{\frac{n-1}{2}}$  is a minimally 2-edge-connected graph. It follows that  $\rho_\alpha(G) \geq \rho_\alpha(F_{\frac{n-1}{2}})$ . By Lemma 2.3, we have

$$\rho_\alpha(F_{\frac{n-1}{2}}) > \alpha\Delta(F_{\frac{n-1}{2}}) + \frac{(1-\alpha)^2}{\alpha} = \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} \quad (1)$$

for  $\alpha \in [\frac{1}{2}, 1)$ .

Let  $w$  be a vertex of  $G$  such that

$$\alpha d(w) + \frac{1-\alpha}{d(w)} \sum_{uv \in E(G)} d(v) = \max_{u \in V(G)} \left\{ \alpha d(u) + \frac{1-\alpha}{d(u)} \sum_{uv \in E(G)} d(v) \right\}.$$

By Lemma 2.1, we have

$$\rho_\alpha(G) \leq \alpha d(w) + \frac{1-\alpha}{d(w)} \sum_{uv \in E(G)} d(v). \quad (2)$$

Let  $A_1$  be the isolated vertex set of  $G[N(w)]$ . Denote by  $B = V(G) \setminus N[w]$  and  $A_2 = N(w) \setminus A_1$ . Now we give Claims 1-3 to complete the proof of Theorem 1.1 (i).

**Claim 1.**  $G[N(w)] = sK_2 \cup tK_1$ , where  $s, t \geq 0$ .

**Proof.** Firstly, we assert that  $G[N(w)]$  is  $P_3$ -free. Otherwise, there exists a  $P_3 = v_1v_2v_3$  in  $G[N(w)]$  such that  $wv_2$  is a chord of cycle  $wv_1v_2v_3w$ , which contradicts with Lemma 2.6. Secondly, we assert that  $G[N(w)]$  is  $C_l$ -free, where  $l \geq 3$ . Otherwise, there exists a wheel graph  $W_{l+1}$  in  $G$  such that there is a cycle containing a chord, a contradiction. Hence, we have  $G[N(w)] = sK_2 \cup tK_1$ , where  $s, t \geq 0$ .  $\square$

**Claim 2.**  $N_B(u) = \emptyset$  for any  $u \in A_2$ .

**Proof.** Otherwise, there exists a vertex  $v \in B$  that is adjacent to a vertex  $u_2 \in A_2$ . We assume that  $u'_2 \in N_{A_2}(u_2)$ . If  $N_B(v) = \emptyset$ , then there exists a vertex  $u \in A$  adjacent to  $v$  by Lemma 2.4. We consider three cases regarding vertex  $u$ . If  $u \in A_1$ , then a chord  $wu_2$  exists in cycle  $wu'_2u_2vuw$ , a contradiction. If  $u \in A_2$  and  $u = u'_2$ , then  $u'_2u_2$  is a chord of the cycle  $wu'_2vu_2w$ , a contradiction. If  $u \in A_2$  and  $u \neq u'_2$ , then there exists a chord  $wu_2$  in cycle  $wu'_2u_2vuw$ , a contradiction. Hence we have  $N_B(v) \neq \emptyset$ .

Denote by  $P(u \rightarrow v)$  the path from  $u$  to  $v$ , then there exists a path  $P(v \rightarrow v_t)$  in  $G[B]$  such that  $v_t$  is adjacent to a vertex  $u' \in A$ . Otherwise  $u_2v$  will be a cut edge. If  $u' \in A_1$ , then a chord  $wu_2$  exists in cycle  $wu'_2u_2P(v \rightarrow v_t)u'w$ , a contradiction. If  $u' \in A_2$  and  $u' = u'_2$ , then  $u'_2u_2$  is a chord of cycle  $wu_2P(v \rightarrow v_t)u'w$ , a contradiction. If  $u' \in A_2$  and  $u' \neq u'_2$ , then  $G' = G - u_2v - u_2v_t + uv + uv_t$  is a minimally 2-edge-connected graph. By Lemma 2.8, we have  $\rho(G') > \rho(G)$ , which contradicts the maximality of  $G$ .  $\square$

**Claim 3.**  $\Delta(G) = n-1$ .

**Proof.** If  $\Delta(G) \leq n-2$ , then  $|B| \geq 1$ . By Lemma 2.4, we have  $2 \leq d(w) \leq \Delta(G) \leq n-2$ . We consider the following two cases.

**Case 1.**  $e(N(w)) = 0$ . In this case, we have  $e(N(w), B) \leq d(w)|B|$ . It follows that

$$\sum_{uv \in E(G)} d(v) = d(w) + 2e(N(w)) + e(N(w), B) \leq d(w)(|B| + 1).$$

Combining this with (2), we have

$$\begin{aligned}\rho_\alpha(G) &\leq \alpha d(w) + (1 - \alpha)(|B| + 1) = \alpha d(w) + (1 - \alpha)(n - d(w)) \\ &= \alpha d(w) + (1 - \alpha)n = (2\alpha - 1)d(w) + (1 - \alpha)n \\ &\leq (2\alpha - 1)(n - 2) + (1 - \alpha)n.\end{aligned}$$

Noting that

$$\alpha(n - 1) + \frac{(1 - \alpha)^2}{\alpha} - ((2\alpha - 1)(n - 2) + (1 - \alpha)n) = \frac{(2\alpha - 1)^2}{\alpha} \geq 0 \quad (3)$$

for  $n \geq 7$  and  $\alpha \in [\frac{1}{2}, 1)$ , we have  $\rho_\alpha(G) \leq \alpha(n - 1) + \frac{(1 - \alpha)^2}{\alpha}$ . Combining this with (1), we have  $\rho_\alpha(G) < \rho_\alpha(F_{\frac{n-1}{2}})$ , a contradiction.

**Case 2.**  $e(N(w)) \geq 1$ .

By Claims 1 and 2, we have  $e(N(w), B) \leq (d(w) - 2e(N(w)))|B|$ . It follows that  $\sum_{wv \in E(G)} d(v) \leq d(w) + 2e(N(w)) + (d(w) - 2e(N(w)))|B|$ . Combining this with (2) and  $|B| \geq 1$ , we have

$$\begin{aligned}\rho_\alpha(G) &\leq \alpha d(w) + (1 - \alpha)(d(w) + 2e(N(w)) + (d(w) - 2e(N(w)))|B|) \\ &= \alpha d(w) + (1 - \alpha)(|B| + 1) + \frac{2(1 - \alpha)e(N(w))}{d(w)}(1 - |B|) \\ &\leq \alpha d(w) + (1 - \alpha)(n - d(w)) \\ &\leq (2\alpha - 1)(n - 2) + (1 - \alpha)n.\end{aligned}$$

Combining with (1) and (3), we have  $\rho_\alpha(G) < \rho_\alpha(F_{\frac{n-1}{2}})$ , a contradiction.

Combining the above arguments, we have  $\Delta(G) = n - 1$ .  $\square$

By Claim 3, we have  $\Delta(G) = 1$ . Notice that  $G[N(w)]$  consists of some edges and isolated vertices. By the maximality of  $\rho_\alpha(G)$ , we have  $G \cong F_{\frac{n-1}{2}}$ .

(ii) Let  $G$  be a minimally 2-edge-connected graph with even order  $n \geq 8$ . Then  $\Delta(G) \leq n - 2$  by Lemmas 2.4 and 2.6. Notice that  $K_{2, n-2}$  is a minimally 2-edge-connected graph. By Lemma 2.3, we have

$$\rho_\alpha(K_{2, n-2}) > \alpha \Delta(K_{2, n-2}) + \frac{(1 - \alpha)^2}{\alpha} = \alpha(n - 2) + \frac{(1 - \alpha)^2}{\alpha} \quad (4)$$

for  $\alpha \in [\frac{1}{2}, 1)$ .

If  $\Delta(G) = n - 2$ , then  $|B| = 1$ . If  $e(N(w)) = 0$ , then  $G \cong K_{2, n-2}$  by Lemma 2.4. If  $e(N(w)) \geq 1$ , we assume that  $B = \{v\}$  and  $u_i \in A_i$  for  $i = 1, 2$ . By analysis, when  $e(N(w)) = 1$ , the vertices  $w$  and  $u_2$  make the right-hand side of the inequality in Lemma 2.2 obtain the maximum value. It follows that

$$\begin{aligned}\rho_\alpha(G) &\leq \max_{uv \in E(G)} \left\{ \frac{\alpha(d(u) + d(v)) + \sqrt{\alpha^2(d(u) - d(v))^2 + 4(1 - \alpha)^2 m(u)m(v)}}{2} \right\} \\ &= \frac{\alpha(d(w) + d(u_2)) + \sqrt{\alpha^2(d(w) - d(u_2))^2 + 4(1 - \alpha)^2 m(w)m(u_2)}}{2} \\ &= \frac{\alpha n + \sqrt{\alpha^2(n - 4)^2 + 8(1 - \alpha)^2(n - 3)}}{2}.\end{aligned}$$

for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ .

Noting that  $\alpha^2(n-4)^2 + 8(1-\alpha)^2(n-3) - (\alpha^2n^2 + 8(1-2\alpha)(n-2)) = -8(\alpha-1)^2 < 0$ , we have

$$\frac{\alpha n + \sqrt{\alpha^2(n-4)^2 + 8(1-\alpha)^2(n-3)}}{2} < \frac{\alpha n + \sqrt{\alpha^2n^2 + 8(1-2\alpha)(n-2)}}{2}$$

for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ . By Lemma 2.10, we have  $\rho_\alpha(G) < \rho_\alpha(K_{2,n-2})$ .

Next, we consider  $\Delta(G) \leq n-3$ . By Lemma 2.4, we have  $2 \leq d(w) \leq \Delta(G) \leq n-3$ . By Lemma 2.1, we have

$$\rho_\alpha(G) \leq \alpha d(w) + \frac{1-\alpha}{d(w)} \sum_{wv \in E(G)} d(v). \quad (5)$$

Now, we prove that  $\rho_\alpha(G) < \rho_\alpha(K_{2,n-2})$  for  $2 \leq d(w) \leq n-3$ . We consider the following two cases.

**Case 1.**  $d(w) = 2$ .

**Subcase 1.1.**  $e(N(w)) = 0$ .

If  $e(B) = 0$ , then  $G \cong K_{2,n-2}$  by Lemma 2.4. If  $e(B) \geq 1$ , then there exists at least one path  $P_n$  ( $n \geq 2$ ) in  $B$ . Since every cycle in  $G$  has no chord, only the endpoints in  $P_n$  are connected to each of the two vertices in  $N(w)$ , and the remaining vertices are not connected to any vertex of  $N(w)$ . It follows that  $e(N(w), B) \leq 2 + d(w)(|B| - 2) = 2n - 8$ . Thus,  $\sum_{wv \in E(G)} d(v) = d(w) + 2e(N(w)) + e(N(w), B) \leq 2n - 6$ . Combining this with (5), we have

$$\rho_\alpha(G) \leq 2\alpha + \frac{1-\alpha}{2} \sum_{wv \in E(G)} d(v) \leq 2\alpha + (1-\alpha)(n-3).$$

Noting that

$$\alpha(n-2) + \frac{(1-\alpha)^2}{\alpha} - (2\alpha + (1-\alpha)(n-3)) = \frac{(2\alpha^2 - \alpha)n - 6\alpha^2 + \alpha + 1}{\alpha} \geq 0$$

for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ , then we have  $\rho_\alpha(G) \leq \alpha(n-2) + \frac{(1-\alpha)^2}{\alpha}$ . Combining this with (4), we have  $\rho_\alpha(G) < \rho_\alpha(K_{2,n-2})$  for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ .

**Subcase 1.2.**  $e(N(w)) = 1$ .

By Lemma 2.6, we know that each vertex in  $B$  exactly has one neighbor in  $N(w)$ . It follows that  $e(N(w), B) < n-3$ . Thus,  $\sum_{wv \in E(G)} d(v) = d(w) + 2e(N(w)) + e(N(w), B) < n+1$ . Combining this with (5), we have

$$\rho_\alpha(G) \leq 2\alpha + \frac{1-\alpha}{2} \sum_{wv \in E(G)} d(v) < 2\alpha + \frac{(1-\alpha)(n+1)}{2}.$$

Noting that

$$\alpha(n-2) + \frac{(1-\alpha)^2}{\alpha} - (2\alpha + \frac{(1-\alpha)(n+1)}{2}) = \frac{(3\alpha^2 - \alpha)n - 5\alpha^2 - 5\alpha + 2}{\alpha} \geq 0$$

for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ , then we have  $\rho_\alpha(G) < \alpha(n-2) + \frac{(1-\alpha)^2}{\alpha}$ . Combining this with (4), we have  $\rho_\alpha(G) < \rho_\alpha(K_{2,n-2})$  for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ .

**Case 2.**  $3 \leq d(w) \leq n - 3$ .

In order to prove  $\rho_\alpha(G) < \rho_\alpha(K_{2,n-2}) = \frac{1}{2}(\alpha n + \sqrt{\alpha^2 n^2 + 8(n-2)(1-2\alpha)})$ , it is enough to prove  $\rho_\alpha(G)^2 - \alpha n \rho_\alpha(G) + 2(2\alpha - 1)(n - 2) < 0$ . For convenience, we denote  $A_\alpha(G) = A_\alpha$ ,  $A(G) = A$  and  $D(G) = D$ . Let

$$B = (b_{ij})_{n \times n} = A_\alpha^2 - \alpha n A_\alpha + 2(2\alpha - 1)(n - 2)I_n,$$

where  $I_n$  is the  $n \times n$  identity matrix. Let  $c_u(B)$  be the sum of all elements in the  $u$ -th column of  $B$ . Then we have the following claim.

**Claim 2.1.**  $c_u(B) < 0$  for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ .

**Proof.** Since  $A_\alpha = \alpha D + (1 - \alpha)A$ , then

$$\begin{aligned} B &= (\alpha D + (1 - \alpha)A)^2 - \alpha n(\alpha D + (1 - \alpha)A) + 2(2\alpha - 1)(n - 2)I_n \\ &= \alpha^2 D^2 + (1 - \alpha)^2 A^2 + \alpha(1 - \alpha)DA + \alpha(1 - \alpha)AD - \alpha^2 nD - (\alpha n - \alpha^2 n)A \\ &\quad + 2(2\alpha - 1)(n - 2)I_n. \end{aligned}$$

It is easy to see that  $c_u(A) = c_u(D) = d(u)$ ,  $c_u(A^2) = c_u(DA) = \sum_{uv \in E(G)} d(v)$  and  $c_u(AD) = d^2(u)$ . By Lemma 2.5, we have  $\sum_{uv \in E(G)} d(v) < |E(G)| \leq 2n - 2$ . It follows that

$$\begin{aligned} c_u(B) &= \alpha^2 d^2(u) + (1 - \alpha)^2 \sum_{uv \in E(G)} d(v) + \alpha(1 - \alpha) \sum_{uv \in E(G)} d(v) + \alpha(1 - \alpha)d^2(u) - \alpha^2 nd(u) \\ &\quad - (\alpha n - \alpha^2 n)d(u) + 2(2\alpha - 1)(n - 2) \\ &= \alpha d^2(u) + (1 - \alpha) \sum_{uv \in E(G)} d(v) - \alpha nd(u) + 2(2\alpha - 1)(n - 2) \\ &< \alpha d^2(u) + (1 - \alpha)(2n - 2) - \alpha nd(u) + 2(2\alpha - 1)(n - 2) \\ &= \alpha(d^2(u) - nd(u) + 2n - 6) + 2 \\ &\leq \max \{ \alpha(9 - 3n + 2n - 6), \alpha((n - 3)^2 - n(n - 3) + 2n - 6) \} + 2 \\ &= \alpha(-n + 3) + 2 < 0 \end{aligned}$$

for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ .

This completes the proof of the claim.  $\square$

Let  $X = (x_1, x_2, \dots, x_n)^T$  be the  $\alpha$ -Perron vector of  $G$  corresponding to  $\rho_\alpha(G)$  satisfying  $\sum_{i=1}^n x_i = 1$ . Then

$$BX = (\rho_\alpha(G)^2 - \alpha n \rho_\alpha(G) + 2(2\alpha - 1)(n - 2))X.$$

Hence we have

$$\begin{aligned} &\rho_\alpha(G)^2 - \alpha n \rho_\alpha(G) + 2(2\alpha - 1)(n - 2) \\ &= \sum_{i=1}^n (\rho_\alpha(G)^2 - \alpha n \rho_\alpha(G) + 2(2\alpha - 1)(n - 2))x_i \\ &= \sum_{i=1}^n (BX)_i = \sum_{i=1}^n \left( \sum_{j=1}^n b_{ij} x_j \right) = \sum_{j=1}^n \left( \sum_{i=1}^n b_{ij} \right) x_j = \sum_{j=1}^n c_j(B) x_j < 0. \end{aligned}$$

Combining the above arguments, we have  $\rho_\alpha(G) \leq \rho_\alpha(K_{2,n-2})$  for  $n \geq 8$  and  $\alpha \in [\frac{1}{2}, 1)$ , with equality if and only if  $G \cong K_{2,n-2}$ .

These complete the proof.  $\square$



## 4 Proof of Theorem 1.2

In this section, we give the proof of Theorem 1.2.

**Proof.** Let  $G$  be a minimally 3-connected graph with order  $n$ . Notice that  $W_n$  is a minimally 3-connected graph. By Lemma 2.3, we have

$$\rho_\alpha(W_n) > \alpha\Delta(W_n) + \frac{(1-\alpha)^2}{\alpha} = \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} \quad (6)$$

for  $\alpha \in [\frac{1}{2}, 1)$ . Let  $w$  be a vertex of  $G$  such that

$$\alpha d(w) + \frac{1-\alpha}{d(w)} \sum_{uv \in E(G)} d(v) = \max_{u \in V(G)} \left\{ \alpha d(u) + \frac{1-\alpha}{d(u)} \sum_{uv \in E(G)} d(v) \right\}.$$

By Lemma 2.1, we have

$$\rho_\alpha(G) \leq \alpha d(w) + \frac{1-\alpha}{d(w)} \sum_{uv \in E(G)} d(v). \quad (7)$$

Denote by  $B = V(G) \setminus N[w]$ . We assert that  $d_{N(w)}(u) \leq 2$  for  $u \in N(w)$ . Otherwise, we assume that  $d_{N(w)}(u_1) \geq 3$  for  $u_1 \in N(w)$  and  $u'_1 \in N_{N(w)}(u_1)$ . Then there exists a cycle  $wu_1u'_1w$  with  $d(w) \geq 4$  and  $d(u_1) \geq 4$ , which contradicts Lemma 2.7. Hence, we have  $d_{N(w)}(u) \leq 2$  for  $u \in N(w)$ . If  $\Delta(G) = n-1$ , then  $G[N(w)]$  is the union of some cycles by Lemma 2.4. It follows that  $G \cong W_n$ . Otherwise  $w$  is a cut vertex. If  $\Delta(G) \leq n-2$ , then  $3 \leq d(w) \leq n-2$  by Lemma 2.4. We have the following claim.

**Claim 1.**  $G[N(w)]$  contains no cycles for  $3 \leq d(w) \leq n-2$ .

**Proof.** If there exists a cycle  $C_q$  in  $G[N(w)]$ , then  $W_{q+1} \subseteq G$ . Since  $3 \leq d(w) \leq n-2$ , we have  $3 \leq q \leq n-2$ , then  $|V(W_{q+1})| < n$ . Notice that every vertex of  $W_{q+1}$  is contained in some  $K_3$ . By Lemma 2.7, there exists exactly one vertex  $u \in V(W_{q+1})$  such that  $d(u) \geq 4$ . It follows that  $u$  is a cut vertex, which contradicts the fact that  $G$  is a minimally 3-connected graph.  $\square$

By Claim 1, we know that  $G[N(w)]$  is the union of some paths and isolated vertices. We consider the following two cases.

**Case 1.**  $e(N(w)) = 0$ .

In this case, we have  $e(N(w), B) \leq d(w)|B| = d(w)(n-d(w)-1)$ . It follows that

$$\sum_{uv \in E(G)} d(v) = 2e(N(w)) + d(w) + e(N(w), B) \leq d(w)(n-d(w)).$$

Combining with (7), we have

$$\begin{aligned} \rho_\alpha(G) &\leq \alpha d(w) + (1-\alpha)(n-d(w)) \\ &= (2\alpha-1)d(w) + (1-\alpha)n \leq (n-4)\alpha + 2. \end{aligned}$$

Noting that

$$\alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} - ((n-4)\alpha + 2) = \frac{(2\alpha-1)^2}{\alpha} \geq 0$$

for  $\alpha \in [\frac{1}{2}, 1)$ , we have  $\rho_\alpha(G) \leq \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha}$ . Combining this with (6), we have

$$\rho_\alpha(G) \leq \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} < \rho_\alpha(W_n)$$

for  $\alpha \in [\frac{1}{2}, 1)$ .

**Case 2.**  $e(N(w)) \geq 1$ .

We consider the following two subcases.

**Subcase 2.1.**  $d(w) = 3$ .

In this case,  $|B| = n - 4$ . If  $e(N(w)) = 1$ , by Lemma 2.7, we have  $e(N(w), B) \leq (d(w) - 1)|B| + 1$ . If  $e(N(w)) = 2$ , we assume  $P_3 = u_1u_2u_3 \in G[N(w)]$ , then  $N_B(u_2) \neq \emptyset$ . Otherwise,  $G - \{u_1, u_3\}$  is disconnected, which contradicts the fact that  $G$  is a minimally 3-connected graph. By Lemma 2.7, we have  $e(N(w), B) \leq 2 + (d(w) - 2)|B|$ . Hence we have  $e(N(w), B) \leq (3 - e(N(w)))|B| + e(N(w))$  for  $e(N(w)) = 1, 2$ . It follows that

$$\begin{aligned} \sum_{wv \in E(G)} d(v) &= 2e(N(w)) + d(w) + e(N(w), B) \\ &\leq 2e(N(w)) + 3 + (3 - e(N(w)))(n - 4) + e(N(w)) \\ &= (3 - e(N(w)))n + 7e(N(w)) - 9. \end{aligned}$$

Combining with (7), we have

$$\begin{aligned} \rho_\alpha(G) &\leq 3\alpha + \frac{1-\alpha}{3}((3 - e(N(w)))n + 7e(N(w)) - 9) \\ &\leq \max \left\{ 3\alpha + \frac{1-\alpha}{3}(2n - 2), 3\alpha + \frac{1-\alpha}{3}(n + 5) \right\} \\ &= 3\alpha + \frac{2}{3}(1-\alpha)(n - 1). \end{aligned}$$

Noting that

$$\alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} - (3\alpha + \frac{2}{3}(1-\alpha)(n-1)) = \frac{(5\alpha^2 - 2\alpha)n - 11\alpha^2 - 4\alpha + 3}{3\alpha} \geq 0$$

for  $n \geq 7$  and  $\alpha \in [\frac{1}{2}, 1)$ , we have  $\rho_\alpha(G) \leq \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha}$ . Combining this with (6), we have

$$\rho_\alpha(G) \leq \alpha(n-1) + \frac{(1-\alpha)^2}{\alpha} < \rho_\alpha(W_n)$$

for  $n \geq 7$  and  $\alpha \in [\frac{1}{2}, 1)$ .

**Subcase 2.2.**  $4 \leq d(w) \leq n - 2$ .

Let  $P_{t_i}$  ( $1 \leq i \leq s$ ,  $t_i \geq 2$ ) be the  $s$  paths in  $G[N(w)]$ . Then  $e(N(w)) = \sum_{i=1}^s (t_i - 1)$ . We assert that  $|B| \geq 2$ . Otherwise, if  $|B| = 1$ , we assume that  $B = \{v\}$ . Denote by  $c$  the number of components in  $G[N(w)]$ . If  $c = 1$ , then  $P_{d(w)}$  is the only one path in  $G[N(w)]$ . By Lemma 2.7, we have  $d(v) = 2$ , which contradicts Lemma 2.4. If  $c \geq 2$ , then  $G - \{w, v\}$  is disconnected, which contradicts the fact that  $G$  is a minimally 3-connected graph. Hence, we have  $|B| \geq 2$ . By Lemma 2.7, we have  $e(N(w), B) \leq 2s + (d(w) - \sum_{i=1}^s t_i)|B|$ . Hence we

have

$$\begin{aligned}
\sum_{wv \in E(G)} d(v) &= 2e(N(w)) + d(w) + e(N(w), B) \\
&\leq 2 \sum_{i=1}^s (t_i - 1) + d(w) + 2s + (d(w) - \sum_{i=1}^s t_i) |B| \\
&= d(w)(|B| + 1) + (2 - |B|) \sum_{i=1}^s t_i.
\end{aligned}$$

Combining with (7) and  $|B| \geq 2$ , we have

$$\begin{aligned}
\rho_\alpha(G) &\leq \alpha d(w) + \frac{1-\alpha}{d(w)} (d(w)(|B| + 1) + (2 - |B|) \sum_{i=1}^s t_i) \\
&\leq \alpha d(w) + (1 - \alpha)(|B| + 1) \\
&= \alpha d(w) + (1 - \alpha)(n - d(w)) \\
&\leq (n - 4)\alpha + 2.
\end{aligned}$$

Noting that

$$\alpha(n - 1) + \frac{(1 - \alpha)^2}{\alpha} - ((n - 4)\alpha + 2) = \frac{(2\alpha - 1)^2}{\alpha} \geq 0$$

for  $\alpha \in [\frac{1}{2}, 1)$ , we have  $\rho_\alpha(G) \leq \alpha(n - 1) + \frac{(1-\alpha)^2}{\alpha}$ . Combining this with (6), we have

$$\rho_\alpha(G) \leq \alpha(n - 1) + \frac{(1 - \alpha)^2}{\alpha} < \rho_\alpha(W_n)$$

for  $\alpha \in [\frac{1}{2}, 1)$ .

Combining the above arguments, we have  $\rho_\alpha(G) \leq \rho_\alpha(W_n)$  for  $n \geq 7$  and  $\alpha \in [\frac{1}{2}, 1)$ , with equality if and only if  $G \cong W_n$ .

These complete the proof.  $\square$

## Data availability

No data was used for the research described in the article.

## Conflict of interest

The authors declare that they have no conflict of interest.

## References

- [1] B. Bollobás, *Extremal Graph Theory*, Academic Press, London, 1978.
- [2] J.A. Bondy, U.S.R. Murty, *Graph Theory*, Springer, New York, 2008.
- [3] R.A. Brualdi, E.S. Solheid, On the spectral radius of complementary acyclic matrices of zeros and ones, *SIAM J. Algebra. Discrete Methods* 7 (1986) 265–272.

- [4] M.Z. Chen, X.-D. Zhang, Some new results and problems in spectral extremal graph theory (in Chinese) *J. Anhui Univ. Nat. Sci.* 42 (2018) 12–25.
- [5] Q.Q. Chen, Q.X. Huang, The maximal  $A_\alpha$ -spectral radius of graphs with given matching number, *Linear Multilinear Algebra* 20 (2022) 5193–5206.
- [6] X.D. Chen, L.T. Guo, On minimally 2-(edge)-connected graphs with extremal spectral radius, *Discrete Math.* 342 (2019) 2092–2099.
- [7] X. Du, L. Shi, Graphs with small independence number minimizing the spectral radius, *Discrete Math. Algorithms Appl.* 5 (3) (2013) 1350017.
- [8] D.D. Fan, S. Goryainov, H.Q. Lin, On the (signless Laplacian) spectral radius of minimally  $k$ -(edge)-connected graphs for small  $k$ , *Discrete Appl. Math.* 305 (2021) 154–163.
- [9] S.-G. Guo, R. Zhang, Sharp upper bounds on the  $Q$ -index of (minimally) 2-connected graphs with given size, *Discrete Appl. Math.* 320 (2022) 408–415.
- [10] S.-G. Guo, R. Zhang, The sharp upper bounds on the  $A_\alpha$ -spectral radius of  $C_4$ -free graphs and Halin graphs, *Graphs Combin.* 38 (2022) 19.
- [11] W. Li, A. Chang, The minimal Laplacian spectral radius of trees with given matching number, *Linear Multilinear Algebra.* 62 (2014) 218–228.
- [12] Y.T. Li, W.J. Liu, L.H. Feng, A survey on spectral conditions for some extremal graph problems, *Adv. Math.* 51 (2) (2022) 193–258.
- [13] C. Liu, Z.M. Yan, J.P. Li, The maximum  $A_\alpha$ -spectral radius of  $t$ -connected graphs with bounded matching number, *Discrete Math.* 346 (2023) 113447.
- [14] M.H. Liu, C.H. Chen, S.-G. Guo, et al., The  $A_\alpha$ -spectral radius of dense graphs, *Linear Multilinear Algebra* 71 (2023) 1044–1053.
- [15] J.Y. Lou, L.G. Wang, M. Yuan, On the  $\alpha$ -index of minimally 2-connected graphs with given order or size, <https://arxiv.org/abs/2301.03389>.
- [16] Z.Z. Lou, J.M. Guo, Z.W. Wang, Maxima of L-index and Q-index: graphs with given size and diameter, *Discrete Math.* 344 (2021) 112533.
- [17] Z.Z. Lou, G. Min, Q.X. Huang, On the spectral radius of minimally 2-(edge)-connected graphs with given size, *Electron. J. Combin.* 30 (2023) P2.23.
- [18] V. Nikiforov, Some new results in extremal graph theory, in: *Surveys in Combinatorics 2011*, London Math. Soc. Lecture Note Ser. 392 (2011) 141–181.
- [19] V. Nikiforov, Merging the  $A$ - and  $Q$ -spectral theories, *Appl. Anal. Discrete Math.* 11 (2017) 81–107.
- [20] V. Nikiforov, O. Rojo, On the  $\alpha$ -index of graphs with pendent paths, *Linear Algebra Appl.* 550 (2018) 87–104.
- [21] M. Plummer, On minimal blocks, *Trans. Amer. Math. Soc.* 134 (1968) 85–94.

- [22] Z. Stanić, *Inequalities for Graph Eigenvalues*, Cambridge University Press, New York, 2015.
- [23] D. Stevanović, *Spectral Radius of Graphs*, Academic Press, New York, 2015.
- [24] S. Wang, D. Wong, F. Tian, Bounds for the largest and the smallest  $A_\alpha$ -eigenvalues of a graph in terms of vertex degrees, *Linear Algebra Appl.* 590 (2020) 210–223.
- [25] J. Xue, H.Q. Lin, S.T. Liu, J.L. Shu, On the  $A_\alpha$ -spectral radius of a graph, *Linear Algebra Appl.* 550 (2018) 105–120.
- [26] J. Xue, R.F. Liu, J.X. Guo, et al., The maximum spectral radius of irregular bipartite graphs, *Adv. in Appl. Math.* 142 (2023) 102433.
- [27] R. Zhang, S.-G. Guo, Maxima of the Laplacian spectral radius of (minimally) 2-connected graphs with fixed size, *Linear Algebra Appl.* 651 (2022) 390–406.
- [28] W. Q. Zhang, The maximum spectral radius of  $t$ -connected graphs with bounded matching number, *Discrete Math.* 345 (2022) 112775.