THE RANK OF THE CARTIER OPERATOR ON PICARD CURVES

VAHID NOUROZI AND FARHAD RAHMATI*

ABSTRACT. For an algebraic curve \mathcal{X} defined over an algebraically closed field of characteristic p > 0, *a*-number $a(\mathcal{X})$ is the dimension of the space of exact holomorphic differentials on \mathcal{X} . We computed the *a*-number for a family of certain Picard curves using the action of the Cartier operator on $H^0(\mathcal{X}, \Omega^1)$.

1. INTRODUCTION

Let \mathcal{X} be a geometrically irreducible, projective, and non-singular algebraic curve defined over the finite field \mathbb{F}_{ℓ} of order ℓ . Let $\mathcal{X}(\mathbb{F}_{\ell})$ denote the set of \mathbb{F}_{ℓ} rational points for \mathcal{X} . In studying curves over finite fields, determining the size of $\mathcal{X}(\mathbb{F}_{\ell})$ is a fundamental problem. The exact primary result here is the Hasse-Weil bound, which asserts that

$$| #\mathcal{X}(\mathbb{F}_{\ell}) - (\ell+1) | \leq 2g\sqrt{\ell},$$

where $g = g(\mathcal{X})$ is the genus of \mathcal{X} .

In this paper, we consider Picard curves of genus 3 over \mathbb{F}_{q^2} of characteristic p > 3. Let \mathcal{X} be a Picard curve. Then \mathcal{X} can be defined by an affine equation of the form

$$y^3 = f(x), \tag{1.1}$$

where f(x) is a polynomial of degree 4, without multiple roots. It is easy to see that the maximal curve \mathcal{X} is super-singular because all the slopes of its Newton polygon are equal to 1/2. This implies that the Jacobian $X := \text{Jac}(\mathcal{X})$ has no *p*-torsion points over $\overline{\mathbb{F}}_p$. A relevant invariant of the *p*-torsion group scheme of the Jacobian of the curve is the *a*-number.

Consider multiplication by p-morphism $[p]: X \to X$, a finite flat morphism of degree p^{2g} . These factors were $[p] = V \circ F$. Here, $F: X \to X^{(p)}$ is the relative Frobenius morphism originating from the p-power map on the sheaf structure, and the Verschiebung morphism $V: X^{(p)} \to X$ is the dual of F. The multiplicationby-p kernel on X is defined as X[p]. An important invariant is the *a*-number $a(\mathcal{X})$ of curve \mathcal{X} defined by

$$a(\mathcal{X}) = \dim_{\bar{\mathbb{F}}_p} \operatorname{Hom}(\alpha_p, X[p]),$$

where α_p is the kernel of the Frobenius endomorphism in the group scheme $\operatorname{Spec}(k[X]/(X^p))$. Another definition for the *a*-number is

$$a(\mathcal{X}) = \dim_{\mathbb{F}_p}(\operatorname{Ker}(F) \cap \operatorname{Ker}(V)).$$

Key words and phrases. a-number; Cartier operator; Super-singular Curves; Picard Curves.

^{*} Corresponding author.

A few results on the rank of the Cartier operator (especially *a*-number) of curves are introduced by Kodama and Washio [9], González [6], Pries and Weir [18], Yui [21]. Besides that, Vahid talked about the rank of the Cartier of the maximal curves in [16], the hyperelliptic curve in [13], and the maximal function fields of \mathbb{F}_{q^2} in [12, 17]. Vahid also wrote about these topics in his [15] dissertation.

In Section 3, we prove that the *a*-number of the Picard curves with Equation (1.1) is 0 for $p \equiv 1 \mod 3$, and is 1 for $p \equiv 2 \mod 3$, see Theorem 3.1. The proofs directly use the Cartier operator action on $H^0(\mathcal{X}, \Omega^1)$. Finally, we provide an example of the *a*-number of Picard curves obtained from the Hasse-Witt matrix.

2. The Cartier operator

Let k be an algebraically closed field of characteristic p > 0. Let \mathcal{X} be a curve defined over k. The Cartier operator is a 1/p-linear operator acting on the sheaf $\Omega^1 := \Omega^1_{\mathcal{X}}$ of differential forms on \mathcal{X} with positive characteristics, p > 0.

Let $K = k(\mathcal{X})$ be the function field of curve \mathcal{X} of genus g defined over k. The separating variable for K is an element $x \in K \setminus K^p$. We can write any function $f \in K$ uniquely in form

$$f = f_0^p + f_1^p x + \dots + f_{p-1}^p x^{p-1},$$

where $f_i \in K$, for $i = 0, \cdots, p - 1$.

Definition 2.1. (Cartier Operator). Let $\omega \in \Omega_{K/K_p}$. There exist f_0, \dots, f_{p-1} such that $\omega = (f_0^p + f_1^p x + \dots + f_{p-1}^p x^{p-1}) dx$. The Cartier operator \mathfrak{C} is defined by

$$\mathfrak{C}(\omega) := f_{p-1} dx.$$

This definition does not depend on the choice of x (see [19, Proposition 1]).

We refer the reader to [19, 2, 3, 20] for proofs of the following statements.

Proposition 2.2. (Global Properties of \mathfrak{C}). For all $\omega \in \Omega_{K/K_q}$ and all $f \in K$,

•
$$\mathfrak{C}(f^p\omega) = f\mathfrak{C}(\omega);$$

- $\mathfrak{C}(\omega) = 0 \Leftrightarrow \exists h \in K, \omega = dh;$
- $\mathfrak{C}(\omega) = \omega \Leftrightarrow \exists h \in K, \omega = dh/h.$

Remark 2.3. Moreover, one can easily show that

$$\mathfrak{C}(x^j dx) = \begin{cases} 0 & \text{if} \quad p \nmid j+1\\ x^{s-1} dx & \text{if} \quad j+1 = ps. \end{cases}$$

If div(ω) is effective, then differential ω is holomorphic. The set $H^0(\mathcal{X}, \Omega^1)$ of holomorphic differentials is a *g*-dimensional *k*-vector subspace of Ω^1 , such that $\mathfrak{C}(H^0(\mathcal{X}, \Omega^1)) \subseteq H^0(\mathcal{X}, \Omega^1)$. If \mathcal{X} is a curve, then the *a*-number of \mathcal{X} equals the dimension of the kernel of the Cartier operator $H^0(\mathcal{X}, \Omega^1)$ (or equivalently, the dimension of the space of the exact holomorphic differentials on \mathcal{X}) (see [10, 5.2.8]).

The following theorem is based on Gorenstein; see [7, Theorem 12].

Theorem 2.4. A differential $\omega \in \Omega^1$ is holomorphic if and only if it has the form $(h(x, y)/F_y)dx$, where H : h(X, Y) = 0 is a canonical adjoint.

Theorem 2.5. [11] With the above assumptions,

$$\mathfrak{C}(h\frac{dx}{F_y}) = \left(\frac{\partial^{2p-2}}{\partial x^{p-1}\partial y^{p-1}}(F^{p-1}h)\right)^{\frac{1}{p}}\frac{dx}{F_y}$$

for any $h \in K(\mathcal{X})$.

The differential operator ∇ is defined by

$$\nabla = \frac{\partial^{2p-2}}{\partial x^{p-1} \partial y^{p-1}}$$

has the property

$$\nabla(\sum_{i,j} c_{i,j} X^i Y^j) = \sum_{i,j} c_{ip+p-1,jp+p-1} X^{ip} Y^{jp}.$$
(2.1)

3. The *a*-number of Picard Curves

This section considers the Picard curves over a finite field with q^2 elements of characteristic p > 3. Let \mathcal{X} be the Picard curve of genus 3 over \mathbb{F}_{q^2} . Then \mathcal{X} can be defined by an affine equation of the form

$$y^3 = f(x),$$
 (3.1)

where f(x) is a polynomial of degree 4 without multiple roots. From Theorem 2.4, one can find a basis for the space $H^0(\mathcal{X}, \Omega^1)$ of holomorphic differentials on \mathcal{X} , namely

$$\mathcal{B} = \left\{ z_i = \frac{h_i dx}{f_y} \right\},\,$$

where $h_1 = 3, h_2 = 3x$ and $h_3 = 3y$ for $1 \le i \le 3$. Therefore, $\mathcal{B} = \{\frac{dx}{y^2}, \frac{xdx}{y^2}, \frac{dx}{y}\}.$

Theorem 3.1. Suppose that $p \not\equiv 0 \mod 3$ and f(x) have a non-zero constantcoefficient, then

- 1. If $p \equiv 1 \mod 3$, then $a(\mathcal{X}) = 0$.
- 2. If $p \equiv 2 \mod 3$, then $a(\mathcal{X}) = 1$.

Proof. 1. Let us compute the image of $\mathfrak{C}(\omega)$ for any $\omega \in \mathcal{B}$. Taking p = 3r+1 and $r \in \mathbb{Z}$. Since p-1 = 3r. We get 2p-2 = 6r. So

$$\begin{aligned} \mathfrak{C}(xdx/y^2) &= \mathfrak{C}(y^{2p-2}y^{-2p}xdx) = y^{-2}\mathfrak{C}(xf(x)^{2r}dx) \\ &= y^{-2}\sum_{i=0}^{2r}c_i\mathfrak{C}(x^{i+1}dx) = c_rdx \neq 0. \\ \mathfrak{C}(dx/y^2) &= \mathfrak{C}(y^{2p-2}y^{-2p}dx) = y^{-2}\mathfrak{C}(f(x)^{2r}dx) \end{aligned}$$

$$= y^{-2} \sum_{i=0}^{2r} c_i \mathfrak{C}(x^i dx) = c_r dx \neq 0.$$

Now we let p - 1 = 3r,

$$\begin{aligned} \mathfrak{C}(dx/y) &= \mathfrak{C}(y^{p-1}y^{-p}dx) = y^{-1}\mathfrak{C}(f(x)^r dx) \\ &= y^{-1}\sum_{i=0}^r c_i \mathfrak{C}(x^i dx) = c_r dx \neq 0. \end{aligned}$$

Thus, $a(\mathcal{X}) = 0$.

2. We assume that p = 3r + 2 and $r \in \mathbb{Z}$. We get p - 2 = 3r. Hence

$$\begin{split} \mathfrak{C}(xdx/y^2) &= \mathfrak{C}(y^{p-2}y^{-p}xdx) = y^{-1}\mathfrak{C}(xf(x)^rdx) \\ &= y^{-1}\sum_{i=0}^r c_i\mathfrak{C}(x^{i+1}dx) = c_rdx \neq 0. \\ \mathfrak{C}(dx/y^2) &= \mathfrak{C}(y^{p-2}y^{-p}dx) = y^{-1}\mathfrak{C}(f(x)^rdx) \\ &= y^{-1}\sum_{i=0}^r c_i\mathfrak{C}(x^idx) = c_rdx \neq 0. \end{split}$$

Finally, from Remark 2.3 we have

$$\begin{aligned} \mathfrak{C}(dx/y) &= \mathfrak{C}(y^{p-1}y^{-p}dx) = y^{-1}\mathfrak{C}(f(x)^{\frac{3r-1}{3}}dx) \\ &= y^{-1}\sum_{i=0}^{\frac{3r-1}{3}}c_i\mathfrak{C}(x^i dx) = 0. \end{aligned}$$

Therefore, $a(\mathcal{X}) = 1$.

Where the coefficients $c_i \in k$ are obtained from the expansion

$$y^{p-1} = f(x)^{(p-1)/2} = \sum_{i=0}^{N} c_i x^i$$
 with $N = \frac{p-1}{2}$.

In [4], Estrada Sarlabous showed that if $F^{p-1} = \sum_{i,j} c_{i,j} x^i y^j$, then the Hasss-Wit matrix of \mathcal{B} is

$$H = \begin{pmatrix} c_{p-1,p-1} & c_{2p-1,p-1} & c_{p-1,2p-1} \\ c_{p-2,p-1} & c_{2p-2,p-1} & c_{p-2,2p-1} \\ c_{p-1,p-2} & c_{2p-1,p-2} & c_{p-1,2p-2} \end{pmatrix}.$$

Since $F = y^3 - f(x)$, $c_{i,j} = 0$ for $j \not\equiv 0 \mod 3$. Thus, we have

1. $p \equiv 1 \mod 3$. Then

$$H = \begin{pmatrix} c_{p-1,p-1} & c_{2p-1,p-1} & 0\\ c_{p-2,p-1} & c_{2p-2,p-1} & 0\\ 0 & 0 & c_{p-1,2p-2} \end{pmatrix}.$$
 (3.2)

where $c_{p-1,p-1}$, $c_{2p-1,p-1}$, $c_{p-2,p-1}$, $c_{2p-2,p-1}$ are the respective coefficients of $x^{p-1}, x^{2p-1}, x^{p-2}$ and x^{2p-2} in $f(x)^{(2p-2)/3} \mod p$. $c_{p-1,2p-2}$ is the coefficient of x^{p-1} in $f(x)^{(p-1)/3} \mod p$.

2. $p \equiv 2 \mod 3$. Then

$$H = \begin{pmatrix} 0 & 0 & c_{p-1,2p-1} \\ 0 & 0 & c_{p-2,2p-1} \\ c_{p-1,p-2} & c_{2p-1,p-2} & 0 \end{pmatrix}.$$
 (3.3)

where $c_{p-1,2p-1}$ and $c_{p-2,2p-1}$ are the coefficients of x^{p-1} and x^{p-2} in $f(x)^{(p-2)/3}$ mod p, and $c_{p-1,p-2}$, and $c_{2p-1,p-2}$ are the coefficients of x^{p-1} and x^{2p-1} in $f(x)^{(2p-1)/3}$ mod p.

Example 3.2. Consider the Picard curves with the following equation:

$$y^3 = x^4 + 1.$$

We calculate the a-number with the Hasse-Wit matrix of this curve with p = 5and p = 13. Firstly, suppose that p = 5 in this case from equation (3.3), we have

$$H = \left(\begin{array}{rrr} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 3 & 0 & 0 \end{array}\right).$$

So rank(H) = 2. From the definition of *a*-number, it is clear that $a(\mathcal{X}) = g(\mathcal{X}) - \operatorname{rank}(H)$. Therefore, $a(\mathcal{X}) = 1$. Finally, assuming that p = 13 in this case from equation (3.2), we have

$$H = \left(\begin{array}{rrr} 4 & 0 & 0\\ 0 & 2 & 0\\ 0 & 0 & 4 \end{array}\right).$$

So rank(H) = 3, hence $a(\mathcal{X}) = 0$. In addition, we achieve this result from MAGMA computation [1].

Acknowledgements. This paper was written while Vahid Nourozi was visiting Unicamp (Universidade Estadual de Campinas) supported by TWAS/Cnpq (Brazil) with fellowship number 314966/2018 - 8. We are extremely grateful to the referee for their valuable comments and suggestions, which led us to find correct references for many assertions and improve the exposition.

References

- W. Bosma, J. Cannon, and C. Playoust. The Magma algebra system. I. The user language. J. Symbolic Comput., 24(3-4):235–265, 1997. Computational algebra and number theory (London, 1993).
- P. Cartier. Une nouvelle opération sur les formes différentielles. C. R. Acad. Sci. Paris, 244 (1957), 426-428.
- P. Cartier. Questions de rationalité des diviseurs en géométrie algébrique. Bull. Soc. Math. France, 86 (1958), 177-251.
- J. Estrada Sarlabous. On the Jacobian varieties of Picard curves defined over fields of characteristic p > 0. Math. Nachr., 152:392-340, 1991.
- R. Fuhrmann and F. Torres, The genus of curves over finite fields with many rational points, Manuscripta Math. 89 (1996), 103-106.
- J. González, Hasse-Witt matrices for the Fermat curves of prime degree, *Tohoku Math. J.* 49 (1997) 149-163.
- D. Gorenstein, An arithmetic theory of adjoint plane curves, Trans. Am. Math. Soc. 72 (1952) 414-436.
- J.W.P. Hirschfeld, G. Korchm´aros and F. Torres, Algebraic Curves Over a Finite Field, Princeton Series in Applied Mathematics. Princeton Univ. Press, Princeton, NJ, 2008. xx+ 696 pp.
- T. Kodama, T. Washio, Hasse-Witt matrices of Fermat curves, Manuscr. Math. 60 (1988) 185-195.
- K.-Z. Li, F. Oort, Moduli of Supersingular Abelian Varieties, *Lecture Notes in Mathematics*, vol.1680, Springer-Verlag, Berlin, 1998, iv+116pp.
- M. Montanucci, P. Speziali, The a-numbers of Fermat and Hurwitz curves. J. Pure Appl. Algebra 222 (2018) 477-488.
- 12. Nourozi, Vahid, and Farhad Rahmati. The rank of the Cartier operator of the certain F_q^2 -Maximal function fields. *Missouri Journal of Mathematical Sciences*, 34, no. 2 (2022): 184–190.

- V. Nourozi, F. Rahmati, and S. Tafazolian. The a-number of certain hyperelliptic curves. Iranian Journal of Science and Technology, Transactions A: Science, 46, no. 4 (2022): 1235–1239.
- 14. V. Nourozi, and S. Tafazolian. The a-number of maximal curves of the third largest genus. *AUT Journal of Mathematics and Computing*, 3, no. 1 (2022): 11–16.
- 15. V. Nourozi. The rank Cartier operator and linear system on curves= Classificação do operador Cartier e sistemas lineares na curva. Doctoral dissertation., 2021.
- V. Nourozi, S. Tafazolian, and F. Rahamti. The *a*-number of jacobians of certain maximal curves. *Transactions on Combinatorics*, 10, no. 2 (2021): 121–128.
- 17. V. Nourozi, and B. Mosallaei. The *a*-number of $y^{q^2+q+1} = x^{q^2+1} + x^q$ over finite field. arXiv preprint arXiv:2401.13707 (2024).
- R. Pries, C. Weir, The Ekedahl-Oort type of Jacobians of Hermitian curves, Asian J. Math. 19 (2015) 845-869.
- C. S. Seshadri. L'opération de Cartier. Applications. In Variétés de Picard, volume 4 of Séminaire Claude Chevalley. Secrétariat Mathématiques, Paris, 1958-1959.
- M. Tsfasman, S. Vladut, and D. Nogin. Algebraic geometric codes: basic notions, volume 139 of Mathematical Surveys and Monographs. American Mathematical Society, Providence, RI, (2007(.
- N. Yui, On the Jacobian Varieties of Hyperelliptic Curves over Fields of Characteristic p. J. Algebra, 52, (1978), 378-410.

FACULTY OF MATHEMATICS AND COMPUTER SCIENCE, AMIRKABIR UNIVERSITY OF TECH-NOLOGY, (TEHRAN POLYTECHNIC), 424 HAFEZ AVE., TEHRAN 15914, IRAN

Email address: nourozi@aut.ac.ir; nourozi.v@gmail.com *Email address*: frahmati@aut.ac.ir