## Primal-Proximity Spaces

Ahmad Al-Omari<sup>1</sup>, Murad Özkoç<sup>2</sup>, Santanu Acharjee<sup>3</sup> <sup>1</sup>Al al-Bayt University Faculty of Sciences, Department of Mathematics P.O. Box 130095, Mafraq 25113, Jordan <sup>2</sup>Muğla Sıtkı Koçman University, Faculty of Science Department of Mathematics 48000, Menteşe-Muğla, Turkey <sup>3</sup>Department of Mathematics, Guwahati-781014, Assam, India e-mail: <sup>1</sup>omarimutah1@yahoo.com, <sup>2</sup>murad.ozkoc@mu.edu.tr <sup>3</sup>sacharjee326@gmail.com

#### Abstract

The main purpose of this paper is to introduce and study the primalproximity spaces. Also, we define two new operators via primal proximity spaces and investigate some of their fundamental properties. In addition, we obtain a new topology, which is weaker than old one, via these new operators. Moreover, we not only discuss some of their properties but also enrich with some examples.

2010 AMS Classifications: 54E05, 54A05, 54A10, 54B99.

**Keywords:** Primal, primal-proximity, Kuratowski closure operator, primal topological space

#### 1 Introduction

In many areas of mathematics, topology plays very crucial roles. Applications of many topological ideas, to solve various problems of nature, have attracted researchers of different branches of science and social sciences. Many new notions have been introduced in topology, which have enriched topology with several new areas of research. Some of the most important classical structures of topology are filters [31], ideals [15], and grills [3]. The definition of ideal was first introduced by Kuratowski [15]. On the hand, the notion of grill was introduced in [3]. It is important to observe that the notion of ideal is the dual of filter, but ideal has helped researchers to introduce many new areas of topology viz. ideal topological space [7], *I*-proximity [10], etc. But to the best of our knowledge, no literature was available on dual structure of grill prior to [1].

Recently, Acharjee et al. [1] introduced a new structure called 'primal'. They obtain not only some fundamental properties related to primal but also some relationships between topological spaces and primal topological spaces. Primals [1] come across as the dual of the notion of grills while the dual of filters are ideals. Later, Al-Omari et al. [2] introduced several new operators in primal topological spaces using primal. On the other hand, the notion of proximity [4] is also an important notion in the area of topology as well. Several forms of this notion such as *I*-proximity [10],  $\mu$ -proximity [20, 24], quasi proximity [21], and multiset proximity [12] have been studied by several researchers. Moreover, applications of proximity can be found in pattern recognition [32], region based theory of space [33, 34], artificial intelligence [35], spatial analysis [25], etc. One may refer to [5, 6, 8, 9, 11, 13, 14, 16, 17, 18, 19, 23, 26, 27, 28, 29, 30] and many others for proximity.

In section 3 of this paper, we introduce a new type of proximity called primalproximity. Also, we define point-primal proximity operator and investigate some of its fundamental properties in section 4. In addition, we prove that this operator is a Kuratowski closure operator under special condition. Moreover, we define one more operator via point-primal proximity operator. This operator come across as a Kuratowski closure operator without any condition. Furthermore, we give not only some relationships but also several examples.

#### 2 Preliminaries

In this section, we discuss some preliminary definitions which will be used in next sections.

**Definition 2.1.** [1] Let X be a non-emptyset. A collection  $\mathcal{P} \subseteq 2^X$  is called a primal on X if it satisfies the following conditions:

- (i)  $X \notin \mathcal{P}$ ,
- (ii) if  $A \in \mathcal{P}$  and  $B \subseteq A$ , then  $B \in \mathcal{P}$ ,
- (iii) if  $A \cap B \in \mathcal{P}$ , then  $A \in \mathcal{P}$  or  $B \in \mathcal{P}$ .

**Corollary 2.2.** [1] Let X be a non-emptyset. A collection  $\mathcal{P} \subseteq 2^X$  is a primal on X if and only if it satisfies the following conditions:

(i) 
$$X \notin \mathcal{P}_{i}$$

(ii) if  $B \notin \mathcal{P}$  and  $B \subseteq A$ , then  $A \notin \mathcal{P}$ ,

(iii) if  $A \notin \mathcal{P}$  and  $B \notin \mathcal{P}$ , then  $A \cap B \notin \mathcal{P}$ .

**Example 2.3.** [1] Let X be a non-emptyset. Then,  $\mathcal{P} = \{A \subseteq X : |A^c| \ge \aleph_0\}$  is a primal on X, where  $\aleph_0$  is the lowest infinite cardinal number.

**Example 2.4.** [1] Let X be a non-emptyset. Then,  $\mathcal{P} = \{A \subseteq X : |A^c| > \aleph_0\}$  is a primal on X, where  $\aleph_0$  is the lowest infinite cardinal number.

**Definition 2.5.** [4] A binary relation  $\delta$  on  $2^X$  is called an (Efremovič) proximity on X if  $\delta$  satisfies the following conditions:

- 1.  $A\delta B \Rightarrow B\delta A$ ,
- 2.  $A\delta(B \cup C) \Leftrightarrow A\delta B \text{ or } A\delta C$ ,
- 3.  $A\delta B \Rightarrow A \neq \emptyset$  and  $B \neq \emptyset$ ,
- 4.  $A \cap B \neq \emptyset \Rightarrow A\delta B$ ,
- 5. if  $A \not \otimes B$ , then there exists  $C, D \subseteq X$  such that  $A \not \otimes C^c, D^c \not \otimes B$  and  $C \cap D = \emptyset$ .

A proximity space is a pair  $(X, \delta)$  consisting of a set X and a proximity relation on X. We shall write  $A\delta B$  if the sets  $A, B \subseteq X$  are  $\delta$ -related, otherwise we shall write  $A \not \delta B$ . Throughout this paper, the space  $(X, \delta, \mathcal{P})$  means an Ef-proximity space  $(X, \delta)$  with a primal  $\mathcal{P}$  on X. Now, we define the following definition which will be used in section 5.

**Definition 2.6.** In a space  $(X, \delta, \mathcal{P})$ , we say that a subset A of X is locally in  $\mathcal{P}$ at  $x \in X$  if there exists a  $\delta$ -neighborhood U of x such that  $U^c \cup A^c \notin \mathcal{P}$ . Also for a subset A of X, the primal local function of A with respect to  $\delta$  and  $\mathcal{P}$ , denoted by  $A^{\diamond}(\delta, \mathcal{P})$ , simply  $A^{\diamond}(\mathcal{P})$  or  $A^{\diamond}$ , is the set  $\bigcup \{x \in X : A \text{ is not primal locally in } \mathcal{P}$ at  $x\}$  i.e.,  $A^{\diamond}(\delta, \mathcal{P}) = \bigcup \{x \in X : U^c \cup A^c \notin \mathcal{P}, \text{ for every } \delta\text{-neighborhood } U \text{ of } x\}$ .

#### **3** Primal-Proximity Spaces

In this section, we introduce the notion of primal-proximity on X and investigate some of its fundamental properties. **Definition 3.1.** A binary relation  $\hookrightarrow$  on  $2^X$  with a primal  $\mathcal{P}$  on a non-emptyset X is called a primal-proximity on X if  $\hookrightarrow$  satisfies the following conditions:

- (1) if  $A \hookrightarrow B$ , then  $B \hookrightarrow A$ ;
- (2)  $A \hookrightarrow (B \cup C)$  if and only if  $A \hookrightarrow B$  or  $A \hookrightarrow C$ ;
- (3) if  $A^c \notin \mathcal{P}$ , then  $A \nleftrightarrow B$  for all  $B \subseteq X$ ;
- (4) if  $(A \cap B)^c \in \mathcal{P}$ , then  $A \hookrightarrow B$ ;
- (5) if  $A \nleftrightarrow B$ , then there exist  $C, D \subseteq X$  such that  $A \nleftrightarrow C^c$ ,  $D^c \nleftrightarrow B$  and  $(C \cap D)^c \notin \mathcal{P}$ .

**Definition 3.2.** A primal-proximity space is a pair  $(X, \hookrightarrow)$  consisting of a set X and primal-proximity relation on a non-emptyset X. We write  $A \hookrightarrow B$  if the sets  $A, B \subseteq X$  are  $\hookrightarrow$ -related, otherwise we write  $A \nleftrightarrow B$ .

**Remark 3.3.** Let X be a non-emptyset and  $A \subseteq X$  such that  $\mathcal{P} = 2^X \setminus \{X\}$ .

- 1. If  $x \in A$ , then  $\{x\} \hookrightarrow A$ .
- 2. If  $A \nleftrightarrow B$ , then  $A \cap B = \emptyset$ .

Suppose  $A \cap B \neq \emptyset$ . Then, there exists at least one point in X such that  $a \in A \cap B$ . Therefore,  $(A \cap B)^c \neq X$ . Hence,  $(A \cap B)^c \in \mathcal{P}$  since  $\mathcal{P} = 2^X \setminus \{X\}$ . It follows that  $A \hookrightarrow B$ , which is impossible. Therefore,  $A \cap B = \emptyset$ .

**Corollary 3.4.** Let  $\hookrightarrow$  be a primal-proximity on a non-emptyset X. Then, the followings hold:

- (1) if  $B \nleftrightarrow A$ , then  $A \nleftrightarrow B$ ,
- (2)  $A \nleftrightarrow (B \cup C)$  if and only if  $A \nleftrightarrow B$  and  $A \nleftrightarrow C$ ,
- (3) if there exists  $B \subseteq X$  such that  $A \hookrightarrow B$ , then  $A^c \in \mathcal{P}$ ,
- (4) if  $A \not\hookrightarrow B$ , then  $(A \cap B)^c \notin \mathcal{P}$ ,
- (5) if  $A \nleftrightarrow B$ , then there exist  $C, D \subseteq X$  such that  $A \nleftrightarrow C^c$ ,  $D^c \nleftrightarrow B$  and  $(C \cap D)^c \notin \mathcal{P}$ .

**Example 3.5.** Let  $\mathcal{P}$  be a primal on a non-emptyset X and  $A, B \subseteq X$ . We define a binary relation  $\hookrightarrow$  on  $2^X$  as:

$$A \hookrightarrow B \Leftrightarrow A^c, B^c \in \mathcal{P}.$$

Then,  $\hookrightarrow$  is a primal-proximity relation. Indeed, one easily finds that  $\hookrightarrow$  satisfies conditions, (1) to (4). We are to check that  $\hookrightarrow$  also satisfies condition (5). Let  $A \nleftrightarrow B$ . It follows that  $A^c \notin \mathcal{P}$  or  $B^c \notin \mathcal{P}$ . If  $A^c \notin \mathcal{P}$ , by taking  $C = A^c$  and D = A have the required properties. If  $B^c \notin \mathcal{P}$ , by taking C = B and  $D = B^c$ . **Example 3.6.** Let  $\mathcal{P}$  be a primal on a non-emptyset X and  $A, B \subseteq X$ . We define a binary relation  $\hookrightarrow$  on  $2^X$  as:

$$A \hookrightarrow B \Leftrightarrow (A \cap B)^c \in \mathcal{P}.$$

Then,  $\hookrightarrow$  is a primal-proximity on X. It follows directly from the definition that  $\hookrightarrow$  satisfies conditions (1) to (4). To prove that  $\hookrightarrow$  satisfies condition (5), let  $A \nleftrightarrow B$ . It follows that  $(A \cap B)^c \notin \mathcal{P}$ . If we take  $C := B^c$  and D := B, the results proves.

**Example 3.7.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space such that  $\mathcal{P} = 2^X \setminus \{X\}$ . Let  $(X, \tau)$  be a normal space and  $A, B \subseteq X$ . Define a binary relation  $\hookrightarrow$  on  $2^X$  as:

$$A \hookrightarrow B \Leftrightarrow (cl(A) \cap cl(B))^c \in \mathcal{P},$$

where the closure is taken with respect to  $\tau$ . Then, the binary relation  $\hookrightarrow$  is a primal-proximity on X.

*Proof.* (1)  $A \hookrightarrow B \Leftrightarrow (cl(A) \cap cl(B))^c \in \mathcal{P} \Leftrightarrow (cl(B) \cap cl(A))^c \in \mathcal{P} \Leftrightarrow B \hookrightarrow A.$ 

(2) Let  $A, B, C \subseteq X$ .  $A \hookrightarrow (B \cup C) \Leftrightarrow (cl(A) \cap cl(B \cup C))^c \in \mathcal{P}$   $\Leftrightarrow (cl(A) \cap (cl(B) \cup cl(C)))^c \in \mathcal{P}$   $\Leftrightarrow ((cl(A) \cap cl(B)) \cup (cl(A) \cap cl(C)))^c \in \mathcal{P}$   $\Leftrightarrow (cl(A) \cap cl(B))^c \cap (cl(A) \cap cl(C))^c \in \mathcal{P}$   $\Leftrightarrow (cl(A) \cap cl(B))^c \in \mathcal{P} \text{ or } (cl(A) \cap cl(C))^c \in \mathcal{P}$  $\Leftrightarrow A \hookrightarrow B \text{ or } A \hookrightarrow C.$ 

(3) Let 
$$A \hookrightarrow B$$
.  
 $A \hookrightarrow B \Leftrightarrow (cl(A) \cap cl(B))^c \in \mathcal{P} \Rightarrow (cl(A))^c \in \mathcal{P}$   
 $\mathcal{P} = 2^X \setminus \{X\} \} \Rightarrow (cl(A))^c \neq X$   
 $\Rightarrow cl(A) \neq \emptyset \Rightarrow A \neq \emptyset \Rightarrow A^c \neq X$   
 $\mathcal{P} = 2^X \setminus \{X\} \} \Rightarrow A^c \in \mathcal{P}.$ 

(4) Let 
$$A \not\hookrightarrow B$$
.  
 $A \not\hookrightarrow B \Rightarrow (cl(A) \cap cl(B))^c \notin \mathcal{P}$   
 $\mathcal{P} = 2^X \setminus \{X\}$   $\} \Rightarrow (cl(A) \cap cl(B))^c = X$   
 $\Rightarrow cl(A) \cap cl(B) = \emptyset \Rightarrow A \cap B = \emptyset \Rightarrow (A \cap B)^c = X$   
 $\mathcal{P} = 2^X \setminus \{X\}$   $\} \Rightarrow (A \cap B)^c \notin \mathcal{P}.$ 

(5) Let  $A \nleftrightarrow B$ . Then,  $(cl(A) \cap cl(B))^c \notin \mathcal{P}$ . So  $cl(A) \cap cl(B) = \emptyset$ . So that, since  $(X, \tau)$  is normal space there exist two disjoint open sets in  $\tau$ , C and Dsuch that  $cl(A) \subseteq C$  and  $cl(B) \subseteq D$ . Hence,  $C^c$  is closed and  $cl(A) \cap C^c =$   $\emptyset$ . This implies  $cl(A) \nleftrightarrow C^c$ . Since  $C \cap D = \emptyset$ , we have  $C \subseteq D^c$ . It follows that  $cl(C) \subseteq D^c$  since  $D^c$  is closed. Therefore,  $cl(C) \cap cl(B) = \emptyset$ and  $(cl(C) \cap cl(B))^c \notin \mathcal{P}$ . Hence,  $C \nleftrightarrow B$ . Let  $E = C^c$ . Then,  $A \nleftrightarrow B$ implies that there exists a subset E such that  $A \nleftrightarrow E$  and  $E^c \nleftrightarrow B$  and  $(E \cap E^c)^c \notin \mathcal{P}$ .

#### 4 Point-Primal Proximity Operator

This section introduces point-primal proximity operator. Here, we study several properties of a primal-proximity space using this operator.

**Definition 4.1.** Let  $(X, \hookrightarrow)$  be a primal-proximity space. Then, the operator  $(\stackrel{\rightarrow}{\cdot}): 2^X \to 2^X$  defined by  $\stackrel{\rightarrow}{A}:= \{x \in X | \{x\} \hookrightarrow A\}$  is said to be point-primal proximity operator. Moreover,  $\stackrel{\rightarrow}{A}$  is said to be point-primal proximity of A.

We now provide the following lemma without the proof.

**Lemma 4.2.** Let  $\mathcal{P}$  be a primal on a non-emptyset X. If  $A \hookrightarrow B$ ,  $A \subseteq C$ , and  $B \subseteq D$ , then  $C \hookrightarrow D$ .

**Lemma 4.3.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . If  $B \nleftrightarrow A$ , then  $\stackrel{\hookrightarrow}{A} \subseteq B^c$ .

*Proof.* Suppose  $\overrightarrow{A} \cap B \neq \emptyset$ . Then there exists at least a point  $x \in \overrightarrow{A} \cap B$ . So  $x \in \overrightarrow{A}$  and  $x \in B$ , i.e.  $\{x\} \hookrightarrow A$  and  $\{x\} \subseteq B$  by Lemma 4.2 implies that  $A \hookrightarrow B$ , which is a contradiction. Hence,  $\overrightarrow{A} \subseteq B^c$ .

**Theorem 4.4.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . If  $B \nleftrightarrow A$ , then  $B \nleftrightarrow \overset{\rightarrow}{A}$ .

Proof. Let  $B \nleftrightarrow A$ . Then by (5) of Definition 3.1 there exist  $C, D \subseteq X$  such that  $B \nleftrightarrow C^c$ ,  $D^c \nleftrightarrow A$  and  $(C \cap D)^c \notin \mathcal{P}$ . This result, combined with Lemma 4.3, implies that  $\overrightarrow{A} \subseteq D$ . Now, we want to prove that  $\overrightarrow{A} \subseteq C^c$ . Let  $x \in \overrightarrow{A}$ , then  $\{x\} \hookrightarrow A$ . Suppose  $x \in C$  then implies that  $x \in C \cap D$  and  $(C \cap D)^c \subseteq X \setminus \{x\}$ , so  $X \setminus \{x\} \notin \mathcal{P}$ . Then, by Definition 3.1 (3), we have  $\{x\} \nleftrightarrow A$ , which a contradiction. Hence,  $x \in C^c$ . So  $\overrightarrow{A} \subseteq C^c$ . Now, we have by Lemma 4.2  $B \nleftrightarrow \overrightarrow{A}$ . Hence, the theorem is proven.

Due to Theorem 4.4 and (1) of Definition 3.1, we have the following corollary.

**Corollary 4.5.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . If  $B \nleftrightarrow A$ , then  $\stackrel{\leftrightarrow}{B} \nleftrightarrow \stackrel{\leftrightarrow}{A}$ .

**Theorem 4.6.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . Then, the following properties hold:

1. if  $A \subseteq B$ , then  $\overrightarrow{A} \subseteq \overrightarrow{B}$ ; 2.  $(A \cap B) \subseteq \overrightarrow{A} \cap \overrightarrow{B}$ ; 3.  $\overrightarrow{A} \cup \overrightarrow{B} = (A \cup B)$ ; 4.  $\overrightarrow{A} \subseteq \overrightarrow{A}$ ; 5. if  $A^c \notin \mathcal{P}$ , then  $\overrightarrow{A} = \emptyset$ ; 6.  $\overrightarrow{\emptyset} = \emptyset$ ; 7.  $\overrightarrow{A} \setminus \overrightarrow{B} \subseteq (A \setminus B)$ ; 8. if  $B^c \notin \mathcal{P}$ , then  $(A \cup B) = \overrightarrow{A} = (A \setminus B)$ ; 9. if  $[(A \setminus B) \cup (B \setminus A)]^c \notin \mathcal{P}$ , then  $\overrightarrow{A} = \overrightarrow{B}$ .

*Proof.* 1) Let  $A \subseteq B$  and  $x \in \overset{\leftrightarrow}{A}$ . Then  $\{x\} \hookrightarrow A$ . Since  $A \subseteq B$ , by Lemma 4.2, we have  $\{x\} \hookrightarrow B$ . Hence,  $x \in \overset{\leftrightarrow}{B}$ .

2) Let  $A, B \subseteq X$ . From (1), it is not difficult to see that  $(A \cap B) \subseteq A$  and  $(A \cap B) \subseteq B$ . Thus, we get  $(A \cap B) \subseteq A \cap B$ .

3) Let  $A, B \subseteq X$ . From (1), we can easily find that  $\overrightarrow{A} \subseteq (\overrightarrow{A \cup B})$  and  $\overrightarrow{B} \subseteq (\overrightarrow{A \cup B})$ . Thus, obviously  $\overrightarrow{A} \cup \overrightarrow{B} \subseteq (\overrightarrow{A \cup B})$ .

Conversely, let  $y \in (A \cup B)$ . Then,  $\{y\} \hookrightarrow A \cup B$ . Due to Definition 3.1, either  $\{y\} \hookrightarrow A$  or  $\{y\} \hookrightarrow B$ . It indicates that either  $y \in \overset{\leftrightarrow}{A}$  or  $y \in \overset{\leftrightarrow}{B}$ . So, we can conclude that  $(A \cup B) \subseteq \overset{\leftrightarrow}{A} \cup \overset{\leftrightarrow}{B}$ .

4) Let  $A \subseteq X$  and let  $x \notin A$ . Then,  $\{x\} \not\hookrightarrow A$ . So, by Theorem 4.4 we have  $\{x\} \nleftrightarrow \overset{\leftrightarrow}{A}$ . Hence,  $x \notin \overset{\leftrightarrow}{A}$ . Thus, we get  $\overset{\leftrightarrow}{A} \subseteq \overset{\leftrightarrow}{A}$ .

5) Let  $A^c \notin \mathcal{P}$ . Then, by (3) of Definition 3.1,  $A \nleftrightarrow B$  for all subsets B of X. Therefore, we have  $A \nleftrightarrow \{x\}$  for all  $x \in X$ . Again, by (1) of Definition 3.1,  $\{x\} \nleftrightarrow A$  for all  $x \in X$ . This means  $x \notin \overset{\rightarrow}{A}$  for all  $x \in X$ . Hence,  $\overset{\rightarrow}{A} = \emptyset$ .

6) Let  $\overleftrightarrow{\emptyset} \neq \emptyset$ . Thus, we assume that  $x \in \overleftrightarrow{\emptyset}$ . Hence,  $\{x\} \hookrightarrow \emptyset$ . By (1) of Definition 3.1,  $\emptyset \hookrightarrow \{x\}$ . Again, by (3) of Definition 3.1,  $\emptyset^c = X \in \mathcal{P}$ ; which is a contradiction to the definition of primal  $\mathcal{P}$ . Hence,  $\overset{\leftrightarrow}{\emptyset} = \emptyset$ .

7) For all  $A, B \subseteq X, A = (A \setminus B) \cup (A \cap B)$ , by (3) we have  $\overrightarrow{A} = (\overrightarrow{A \setminus B}) \cup (A \cap B) \subseteq (\overrightarrow{A \setminus B}) \subseteq (\overrightarrow{A \setminus B}) \cup \overrightarrow{B}$ . Hence,  $\overrightarrow{A} \setminus \overrightarrow{B} \subseteq (\overrightarrow{A \setminus B})$ . 8) If  $B^c \notin \mathcal{P}$ , then  $(\overrightarrow{A \cup B}) = \overrightarrow{A} \cup \overrightarrow{B} = \overrightarrow{A} \cup \emptyset = \overrightarrow{A}$ . Also,  $\overrightarrow{A} \setminus \overrightarrow{B} \subseteq (\overrightarrow{A \setminus B})$ , then  $\overrightarrow{A} \subseteq (\overrightarrow{A \setminus B})$ . And  $(\overrightarrow{A \setminus B}) = (\overrightarrow{A \cap B^c}) \subseteq \overrightarrow{A} \cap \overrightarrow{B^c} \subseteq \overrightarrow{A}$ . Hence,  $(\overrightarrow{A \cup B}) = \overrightarrow{A \cup B} = \overrightarrow{A \cup B}$ .  $\stackrel{\leftrightarrow}{A} = (A \stackrel{\leftrightarrow}{\setminus} B).$ 

9) If  $[(A \setminus B) \cup (B \setminus A)]^c \notin \mathcal{P}$ , then  $(A \setminus B)^c \notin \mathcal{P}$  and  $(B \setminus A)^c \notin \mathcal{P}$ . Since  $\overrightarrow{A} = [(A \setminus B) \cup (A \cap B)]$  and  $(A \setminus B)^c \notin \mathcal{P}$ , by using (8)  $\overrightarrow{A} = (A \cap B) \subseteq \overrightarrow{B}$ . It follows that  $\overrightarrow{A} \subseteq \overrightarrow{B}$ . Similarly, since  $\overrightarrow{B} = [(B \setminus A) \cup (B \cap A)]$  and  $(B \setminus A)^c \notin \mathcal{P}$ , by using (8)  $\overrightarrow{B} = (B \cap A) \subseteq \overrightarrow{A}$ . It follows that  $\overrightarrow{B} \subseteq \overrightarrow{A}$ . Hence,  $\overrightarrow{A} = \overrightarrow{B}$ .

**Remark 4.7.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A \subseteq X$ . The inclusion  $A \subseteq A$  need not be true in general as shown by following example.

**Example 4.8.** Let  $X = \{a, b, c\}, \mathcal{P} = \{\emptyset, \{b\}, \{c\}, \{b, c\}\}$  and the binary relation  $\hookrightarrow$  on  $2^X$  defined as Example 3.6. For the subset  $A = \{b\}$ , we have  $A = \{b\} \notin$  $\emptyset = \overset{\hookrightarrow}{A}.$ 

**Theorem 4.9.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . Then, the following statements hold:

- 1.  $A \cap \overleftrightarrow{B} = \emptyset$ , for all  $A^c \notin \mathcal{P}$  and  $B \subseteq X$ ,
- 2.  $\{x\} \hookrightarrow X \text{ for all } x \in X \text{ if and only if } \mathcal{P} = 2^X \setminus \{X\}.$
- 3. if  $\mathcal{P} = 2^X \setminus \{X\}$ , then  $\overleftrightarrow{X} = X$ .

*Proof.* (1) Let  $A^c \notin \mathcal{P}$  and suppose  $A \cap \overset{\smile}{B} \neq \emptyset$ . It follows that  $A \nleftrightarrow B$  since  $A^c \notin \mathcal{P}$  and also  $\overset{\smile}{B} \notin A^c$ . Hence, by Lemma 4.3 we have  $A \hookrightarrow B$ , which is a contradiction. Thus,  $A \cap \overset{\smile}{B} = \emptyset$ .

(2) If  $\{x\} \hookrightarrow X$  for all  $x \in X$ , then by (3) of Corollary 3.4, we have  $\{x\}^c \in \mathcal{P}$  for all  $x \in X$ . Hence,  $\mathcal{P} = 2^X \setminus \{X\}$ . Conversely, if  $\mathcal{P} = 2^X \setminus \{X\}$ , then  $(\{x\} \cap X)^c = (\{x\})^c \in \mathcal{P}$  and by (4) of Definition 3.1, we have  $\{x\} \hookrightarrow X$  for all  $x \in X$ .

(3) Let  $x \in X$ . Since  $\mathcal{P} = 2^X \setminus \{X\}$ , then  $(\{x\})^c = (\{x\} \cap X)^c \in \mathcal{P}$  and by (4) of Definition 3.1, we get  $\{x\} \hookrightarrow X$  for all  $x \in X$ . Hence,  $\overset{\leftrightarrow}{X} = X$ .

**Theorem 4.10.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space. If  $A, B, C \subseteq X$  and  $B \subsetneq C$  such that  $A \nleftrightarrow B$  but  $A \hookrightarrow C$ , then  $A \hookrightarrow (C \setminus B)$ .

*Proof.* Let  $A, B, C \subseteq X$  and  $B \subsetneq C$ . We consider  $D = C \setminus B$ . Since  $A \hookrightarrow C$ , then  $A \hookrightarrow B \cup (C \setminus B) = B \cup D$ . Then by Definition 3.1,  $A \hookrightarrow B$  or  $A \hookrightarrow D$ . Now,  $A \hookrightarrow B$  is not possible since we consider  $A \nleftrightarrow B$ . Then obviously,  $A \hookrightarrow (C \setminus B)$ .  $\Box$ 

**Theorem 4.11.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . If  $A \nleftrightarrow B$ , then there exists  $C \subseteq X$  such that  $A \nleftrightarrow C$  and  $B \nleftrightarrow C^c$ .

*Proof.* Since  $A \nleftrightarrow B$ , thus by (5) of Definition 3.1, there exist  $M, N \subseteq X$  such that  $A \nleftrightarrow M^c$ ,  $N^c \nleftrightarrow B$  and  $(M \cap N)^c \notin \mathcal{P}$ . Let  $M = X \setminus C$  and N = C. Then,  $(M \cap N)^c = X \notin \mathcal{P}$ . Also,  $A \nleftrightarrow (X \setminus C)^c$ ,  $C^c \nleftrightarrow B$ . It yields  $A \nleftrightarrow C$ ,  $B \nleftrightarrow C^c$ . Hence, the proof is completed.

**Corollary 4.12.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B, C \subseteq X$ . If  $A \nleftrightarrow B$  and  $B \hookrightarrow C$ , then  $A \nleftrightarrow C$ .

# 5 Proximal Closed Sets and $\stackrel{\leftrightarrow}{\tau}$ Topology

In this section, proximal closed sets are defined. Moreover, various results between a primal-proximity space and a primal topological space are obtained using proximal closed sets and related notions.

**Definition 5.1.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space. Then, a subset F of X is called proximity-closed if and only if  $\{x\} \hookrightarrow F$  implies  $x \in F$ .

**Lemma 5.2.** If there is a point  $x \in X$  such that  $A \hookrightarrow \{x\}$  and  $\{x\} \hookrightarrow B$ , then  $A \hookrightarrow B$ .

*Proof.* Suppose  $A \nleftrightarrow B$ , by Theorem 4.11, there exists a subset C such that  $A \nleftrightarrow C$  and  $C^c \nleftrightarrow B$ . Now, either  $x \in C$  or  $x \in C^c$ .

Case (1): If  $x \in C$ , then  $A \nleftrightarrow \{x\}$ . For if  $A \hookrightarrow \{x\}$ , then  $A \hookrightarrow C$ , by Lemma 4.2 which is a contradiction.

Case (2): If  $x \in C^c$ , then  $\{x\} \nleftrightarrow B$ . Therefore, if  $A \hookrightarrow \{x\}$  and  $\{x\} \hookrightarrow B$ , then  $A \hookrightarrow B$ .

**Theorem 5.3.** The collection of complements of all proximity-closed sets of  $(X, \hookrightarrow, \mathcal{P})$  forms a topology on X. This topology is denoted by  $\overleftrightarrow{\tau}$ .

*Proof.* Since X and  $\emptyset$  are proximity-closed in  $(X, \hookrightarrow, \mathcal{P})$ , their complements  $\emptyset$  and X are in  $\overleftrightarrow{\tau}$ .

Let  $\{F_i : i \in I\}$  be a collection of proximity-closed sets. If  $\{x\} \hookrightarrow \bigcap \{F_i : i \in I\}$ , then  $\{x\} \hookrightarrow F_i$  for every  $i \in I$  by Lemma 4.2. Since  $F_i$  is proximity-closed,  $x \in F_i$ for every  $i \in I$ . Hence,  $x \in \bigcap \{F_i : i \in I\}$  and  $\bigcap \{F_i : i \in I\}$  is proximity-closed. Therefore, if  $(X \setminus F_i) \in \vec{\tau}$  for every  $i \in I$ , then  $\bigcup \{X \setminus F_i : i \in I\}$  is the complement of  $\bigcap \{F_i : i \in I\}$  which belongs to  $\vec{\tau}$ .

Finally, let  $F_1$  and  $F_2$  be two proximity-closed sets. If  $\{x\} \hookrightarrow F_1 \cup F_2$ , then  $\{x\} \hookrightarrow F_1$  or  $\{x\} \hookrightarrow F_2$ . Thus,  $x \in F_1$  or  $x \in F_2$  since  $F_1$  and  $F_2$  are proximityclosed. This implies  $x \in F_1 \cup F_2$ . Thus,  $F_1 \cup F_2$  is proximity-closed. Therefore, if  $X \setminus F_1 \in \vec{\tau}$  and  $X \setminus F_2 \in \vec{\tau}$ , then  $(X \setminus F_1) \cap (X \setminus F_2) = X \setminus (F_1 \cup F_2) \in \vec{\tau}$ . Hence,  $\vec{\tau}$  is a topology on X.

**Theorem 5.4.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space. The set  $\overrightarrow{A}$  is the closure of A where the closure is taken with respect to the topology  $\overrightarrow{\tau}$  and denoted by  $cl_{\overrightarrow{\tau}}(A)$ .

*Proof.* Let  $x \in \overset{\smile}{A}$ . Then  $\{x\} \hookrightarrow A$ . By Lemma 4.2,  $\{x\} \hookrightarrow cl_{\overrightarrow{\tau}}(A)$  since  $A \subseteq cl_{\overrightarrow{\tau}}(A)$  and  $cl_{\overrightarrow{\tau}}(A)$  is proximity-closed in  $\overset{\smile}{\tau}$ . Thus,  $x \in cl_{\overrightarrow{\tau}}(A)$ . Hence,  $\overset{\smile}{A} \subseteq cl_{\overrightarrow{\tau}}(A)$ .

Conversely, let  $x \notin \vec{A}$ . Then  $\{x\} \nleftrightarrow A$ . By Theorem 4.11, there exists a subset C such that  $\{x\} \nleftrightarrow C$  and  $C^c \nleftrightarrow A$ . Since there is no point of  $C^c$  which is related to A, then  $\vec{A} \subseteq C$ . By Lemma 4.2,  $\{x\} \nleftrightarrow cl_{\overrightarrow{\tau}}(A)$ . Thus,  $\vec{A}$  is proximity-closed in  $\overrightarrow{\tau}$ . Therefore,  $cl_{\overrightarrow{\tau}}(A) \subseteq \vec{A}$ . Hence,  $cl_{\overrightarrow{\tau}}(A) = \vec{A}$ .

**Definition 5.5.** [15] The the operator  $\Phi : 2^X \to 2^X$  is a Kuratowski closure operator provided:

(1)  $\Phi(\emptyset) = \emptyset;$ 

- (2)  $A \subseteq \Phi(A)$  for every  $A \in 2^X$ ;
- (3)  $\Phi(A \cup B) = \Phi(A) \cup \Phi(B)$  for any  $A, B \in 2^X$ ;
- (4)  $\Phi(\Phi(A)) = \Phi(A)$  for every  $A \in 2^X$ .

**Theorem 5.6.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space such that  $\mathcal{P} = 2^X \setminus \{X\}$ . Then, the operator  $A := \{x \in X | \{x\} \hookrightarrow A\}$  on a primal-proximity space  $(X, \hookrightarrow, \mathcal{P})$  is a Kuratowski closure operator.

Proof. (1) By (6) of Theorem 4.6,  $\overrightarrow{\emptyset} = \emptyset$ . (2) If  $x \in A$ , then  $\{x\} \hookrightarrow A$ . Hence,  $x \in \overrightarrow{A}$ . This shows that  $A \subseteq \overrightarrow{A}$ . (3) By (3) of Theorem 4.6,  $(A \cup B) = \overrightarrow{A} \cup \overrightarrow{B}$ . (4) By (4) of Theorem 4.6, we have always  $\overrightarrow{A} \subseteq \overrightarrow{A}$ . Now, let  $x \notin \overrightarrow{A}$ . Then  $\{x\} \nleftrightarrow \overrightarrow{A}$ . By (4) of Corollary 3.4, we have  $(\{x\} \cap \overrightarrow{A})^c \notin \mathcal{P}$ . Since  $\mathcal{P} = 2^X \setminus \{X\}$ ,

we get  $\left(\{x\} \cap \stackrel{\leftrightarrow}{A}\right)^c = X$  which means that  $\{x\} \cap \stackrel{\leftrightarrow}{A} = \emptyset$ . Thus, we have  $x \notin \stackrel{\leftrightarrow}{A}$ .

Hence,  $\overrightarrow{A} \subseteq \overrightarrow{A}$  and  $\overrightarrow{A} = \overrightarrow{A}$  which completes the proof and this topology is denoted by  $\overrightarrow{\tau}$ .

**Theorem 5.7.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space. Then, the operator  $cl^* : 2^X \to 2^X$  defined by  $cl^*(A) := A \cup \overset{\smile}{A}$  satisfies Kuratowski closure axioms and induces a topology on X called  $\tau^*$  is given by  $\tau^* = \{A \subseteq X | cl^*(A^c) = A^c\}.$ 

*Proof.* (1) By (6) of Theorem 4.6, we have  $cl^*(\emptyset) = \emptyset \cup \overleftrightarrow{\emptyset} = \emptyset$ .

- (2) Let  $A \subseteq X$ . Since  $cl^*(A) := A \cup A$ , we have  $A \subseteq cl^*(A)$ .
- (3) Let  $A, B \subseteq X$ . By (3) of Theorem 4.6, we have

$$cl^{*}(A \cup B) = (A \cup B) \cup (\overrightarrow{A \cup B})$$
$$= (A \cup B) \cup \left(\overrightarrow{A \cup B}\right)$$
$$= \left(A \cup A\right) \cup \left(\overrightarrow{A \cup B}\right)$$
$$= cl^{*}(A) \cup cl^{*}(B).$$

(4) Let  $A \subseteq X$ . By (4) of Theorem 4.6, we have

$$cl^{*}(cl^{*}(A)) = cl^{*}(A) \cup cl^{*}(A)$$

$$= \left(A \cup \overrightarrow{A}\right) \cup \left(A \cup \overrightarrow{A}\right)$$

$$= \left(A \cup \overrightarrow{A}\right) \cup \left(\overrightarrow{A} \cup \overrightarrow{A}\right)$$

$$= \left(A \cup \overrightarrow{A}\right) \cup \left(\overrightarrow{A} \cup \overrightarrow{A}\right)$$

$$= A \cup \overrightarrow{A}$$

$$= cl^{*}(A). \square$$

**Theorem 5.8.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space. Then the following properties hold:

1.  $B \nleftrightarrow A$  if and only if  $B \nleftrightarrow cl^*(A)$ . 2.  $cl^*\left(\stackrel{\leftrightarrow}{A}\right) = \stackrel{\leftrightarrow}{A}$ . 3.  $cl^*\left(\stackrel{\leftrightarrow}{A}\right) = cl^*\stackrel{\leftrightarrow}{(A)}$ .

*Proof.* (1) Let  $B \nleftrightarrow A$ . Then, by Theorem 4.4, we have  $B \nleftrightarrow \overrightarrow{A}$ . Hence, by (2) of Definition 3.1,  $B \nleftrightarrow (A \cup \overrightarrow{A}) = cl^*(A)$  if and only if  $B \nleftrightarrow A$  and  $B \nleftrightarrow \overrightarrow{A}$ .

(2) Let  $A \subseteq X$ . By (4) of Theorem 4.6, we have

$$cl^*\left(\stackrel{\leftrightarrow}{A}\right) = \stackrel{\leftrightarrow}{A} \cup \stackrel{\leftrightarrow}{A} = \stackrel{\leftrightarrow}{A}.$$

(3) Let  $A \subseteq X$ . By (3) of Theorem 4.6, we have

$$cl^*\left(\stackrel{\leftrightarrow}{A}\right) = \stackrel{\leftrightarrow}{A} \cup \stackrel{\leftrightarrow}{A} = \left(\stackrel{\leftrightarrow}{A} \cup \stackrel{\leftrightarrow}{A}\right) = cl^*(A).$$

**Theorem 5.9.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B, H \subseteq X$  such that  $A \subseteq B$ . If  $A \hookrightarrow B$  and  $\{b\} \hookrightarrow H$  for all  $b \in B$ , then  $A \hookrightarrow H$ .

*Proof.* Suppose  $A \nleftrightarrow H$ , then there exist  $C, D \subseteq X$  such that  $A \nleftrightarrow C^c$ ,  $D^c \nleftrightarrow B$ and  $(C \cap D)^c \notin \mathcal{P}$ . This result, combined with  $A \hookrightarrow B$  and (2) of Definition 3.1, implies that  $B \nsubseteq C^c$ , that is  $B \cap C \neq \emptyset$ . It follows that there is a point  $x \in X$ such that  $\{x\} \hookrightarrow H$  and  $x \in C$ . Then, there are two cases either  $x \in D$  or  $x \notin D$ .

Case 1: If  $x \in D$ . Hence  $X \setminus \{x\} \notin \mathcal{P}$ , by (3) of Definition 3.1, implies  $\{x\} \nleftrightarrow H$  for any subset H of X, which is contradiction.

Case 2:  $x \in D^c$ , then  $\{x\} \nleftrightarrow B$ . This result, combined with (3) and (4) of Definition 3.1, implies  $\{x\} \nleftrightarrow H$ , which is contradiction. Hence,  $A \hookrightarrow H$ .

**Example 5.10.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space and  $\hookrightarrow$  be a binary relation on  $2^X$  defined as  $A \hookrightarrow B$  if and only if  $(A \cap cl(B))^c \in \mathcal{P}$ . Then " $\hookrightarrow$ " is not a primal-proximity relation on  $2^X$  but satisfes (2),(3),(4) and (5) of Definition 3.1. Hence, in this case  $\tau \subseteq \tau^*$ .

Proof. We want to show that  $cl^*(A) \subseteq cl(A)$  for all  $A \subseteq X$ . Let  $x \in cl^*(A) = A \cup A$ . Then,  $x \in A$  or  $x \in A$ . If  $x \in A$ , then  $x \in cl(A)$ . Now if  $x \in A$ , then  $\{x\} \hookrightarrow A$ . Hence,  $(\{x\} \cap cl(A))^c \in \mathcal{P}$  and so  $(\{x\} \cap cl(A))^c \neq X$ . Thus,  $\{x\} \cap cl(A) \neq \emptyset$  which means  $x \in cl(A)$ . Therefore,  $\tau \subseteq \tau^*$ .

**Example 5.11.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space and  $\hookrightarrow$  be a binary relation on  $2^X$  defined as  $A \hookrightarrow B$  if and only if  $(A \cap cl^{\diamond}(B))^c \in \mathcal{P}$ . Then " $\hookrightarrow$ " is not a primal-proximity relation on  $2^X$  but satisfy (2), (3), (4) and (5) of Definition 3.1. Hence, in this case  $\tau^{\diamond} \subseteq \tau^*$ .

*Proof.* We want to show that  $cl^*(A) \subseteq cl^{\diamond}(A)$  for all  $A \subseteq X$ . Let  $x \in cl^*(A) = A \cup \overrightarrow{A}$ . Then  $x \in A$  or  $x \in \overrightarrow{A}$ . If  $x \in A$ , then  $x \in A \subseteq A \cup A^{\diamond} = cl^{\diamond}(A)$ . Now if  $x \in \overrightarrow{A}$ , then  $\{x\} \hookrightarrow A$ . Hence,  $(\{x\} \cap cl^{\diamond}(A))^c \in \mathcal{P}$  and so  $(\{x\} \cap cl^{\diamond}(A))^c \neq X$ . Thus,  $\{x\} \cap cl^{\diamond}(A) \neq \emptyset$  which means  $x \in cl^{\diamond}(A)$ . Therefore,  $\tau^{\diamond} \subseteq \tau^*$ .

**Definition 5.12.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space. Then, X is said to be a primal-regular space if for all  $x \in X$  and  $\tau^{\diamond}$ -closed set F such that  $(\{x\} \cap F)^c \notin \mathcal{P}$ there exist two open sets H, G such that  $x \in H$  and  $F \subseteq G$  and  $(H \cap G)^c \notin \mathcal{P}$ .

**Theorem 5.13.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space. Let X be a primalregular space and  $\hookrightarrow$  be a binary relation on  $2^X$  as defined in Example 5.11, then  $\tau^{\diamond} = \tau^*$ .

*Proof.* In order to prove the theorem, it suffices to show  $cl^{\diamond}(A) = cl^{*}(A)$  for all subsets A of X.

Let  $x \in cl^*(A)$ . Then,  $x \in A$  or  $x \in A$ . If  $x \in A$ , then  $x \in cl^\diamond(A)$ . Now if  $x \in A$ , then  $\{x\} \hookrightarrow A$ . Hence,  $(\{x\} \cap cl^\diamond(A))^c \in \mathcal{P}$  which means  $\{x\} \cap cl^\diamond(A) \neq \emptyset$ . Consequently, we have  $x \in cl^\diamond(A)$ . Thus,  $cl^*(A) \subseteq cl^\diamond(A)$ . Now, let  $x \notin cl^*(A)$ . Then,  $x \notin A$  and  $x \notin A$ . It follows that  $\{x\} \nleftrightarrow A$ and hence by Example 5.11 implies that  $(\{x\} \cap cl^{\diamond}(A))^c \notin \mathcal{P}$ . Since X is primalregular space and  $\tau^c \subseteq \tau^{\diamond c}$ , there exist two open sets H and G such that  $x \in H$ and  $A \subseteq cl^{\diamond}(A) \subseteq G$  and  $(H \cap G)^c \notin \mathcal{P}$ . Hence,  $(H \cap A)^c \notin \mathcal{P}$  and since  $x \in H \in \tau$ and  $(H \cap A)^c \notin \mathcal{P}$ , then  $x \notin A^{\diamond}$ . So,  $x \notin cl^{\diamond}(A)$ . It follows that  $cl^{\diamond}(A) \subseteq cl^*(A)$ . Hence,  $cl^{\diamond}(A) = cl^*(A)$ .

**Example 5.14.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space and  $\hookrightarrow$  be a binary relation on  $2^X$  defined as  $A \hookrightarrow B$  if and only if  $(cl^{\diamond}(A) \cap cl^{\diamond}(B))^c \in \mathcal{P}$ . Then, " $\hookrightarrow$ " is not a primal-proximity relation on  $2^X$  but satisfies (1)-(4) of Definition 3.1.

**Definition 5.15.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space. Then, X is said to be a primal-normal space if for two  $\tau^{\diamond}$ -closed sets  $F_1, F_2$  such that  $(F_1 \cap F_2)^c \notin \mathcal{P}$ , there exist two open sets H and G such that  $F_1 \subseteq H$  and  $F_2 \subseteq G$  and  $(H \cap G)^c \notin \mathcal{P}$ .

**Theorem 5.16.** Let  $(X, \tau, \mathcal{P})$  be a primal topological space. If X is a primalnormal space and a binary relation defined as in Example 5.14 and  $(X, \tau)$  is  $T_1$ space, then  $\tau^{\diamond} = \tau^*$ .

*Proof.* In order to prove the theorem, it suffices to show that  $cl^{\diamond}(A) = cl^{*}(A)$  for all subsets A of X.

Let  $x \in cl^*(A)$ . Then,  $x \in A$  or  $x \in \vec{A}$ . If  $x \in A$ , then  $x \in cl^{\diamond}(A)$ . Now, if  $x \in \vec{A}$ , then  $\{x\} \hookrightarrow A$ . Hence,  $(cl^{\diamond}(\{x\}) \cap cl^{\diamond}(A))^c \in \mathcal{P}$ . Since  $(X, \tau)$  is  $T_1$ -space and  $\tau^c \subseteq \tau^{\diamond c}$ , then  $[\{x\} \cap cl^{\diamond}(A)]^c \in \mathcal{P}$  and so  $\{x\} \cap cl^{\diamond}(A) \neq \emptyset$ . Consequently, we have  $x \in cl^{\diamond}(A)$ . Hence,  $cl^*(A) \subseteq cl^{\diamond}(A)$ .

Now, let  $x \notin cl^*(A)$ . Then,  $x \notin A$  and  $x \notin A$ . It follows that  $\{x\} \nleftrightarrow A$ and hence by Example 5.14 implies that  $(cl^{\diamond}(\{x\}) \cap cl^{\diamond}(A))^c \notin \mathcal{P}$ . Since  $(X, \tau)$  is primal-normal space,  $T_1$ -space and  $\tau^c \subseteq \tau^{\diamond c}$ , there exist two open sets H and Gsuch that  $\{x\} \subseteq H$ ,  $A \subseteq cl^{\diamond}(A) \subseteq G$  and  $(H \cap G)^c \notin \mathcal{P}$ . Hence,  $(H \cap A)^c \notin \mathcal{P}$ . Since  $x \in H \in \tau$  and  $(H \cap A)^c \notin \mathcal{P}$ , thus  $x \notin A^{\diamond}$ . So,  $x \notin cl^{\diamond}(A)$ . It follows that  $cl^{\diamond}(A) \subseteq cl^*(A)$  and hence,  $cl^{\diamond}(A) = cl^*(A)$ .

**Theorem 5.17.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A \subseteq X$ . Then,  $A \in \overleftrightarrow{\tau}$  if and only if  $\{x\} \nleftrightarrow A^c$  for every  $x \in A$ .

*Proof.* Let  $A \in \overleftarrow{\tau}$  and  $x \in A$ . Then,  $A^c$  is proximity-closed in  $\overleftarrow{\tau}$  and  $x \notin A^c$ . Hence, we get  $\{x\} \nleftrightarrow A^c$ .

Conversely, if for every  $x \in A$ , we have  $\{x\} \nleftrightarrow A^c$ , then  $\{x\} \hookrightarrow A^c$  implies that  $x \notin A$ . This means that  $\{x\} \hookrightarrow A^c$  implies  $x \in A^c$ . Hence,  $A^c$  is proximity-closed in  $\overrightarrow{\tau}$ . Thus,  $A \in \overrightarrow{\tau}$ .

**Theorem 5.18.** Let  $(X, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$  such that  $A \nleftrightarrow B$ . Then the following conditions hold:

- 1.  $cl_{\overrightarrow{\tau}}(B) \subseteq A^c$ , where  $cl_{\overrightarrow{\tau}}(B)$  means the closure of B with respect to  $\overleftarrow{\tau}$ .
- 2. if  $\mathcal{P} = 2^X \setminus \{X\}$ , then  $B \subseteq int_{\overrightarrow{\tau}}(A^c)$  where  $int_{\overrightarrow{\tau}}(A^c)$  means the interior of  $A^c$  with respect to  $\overrightarrow{\tau}$ .

*Proof.* (1) Since the closure is taken with respect to  $\overleftarrow{\tau}$  and  $A \nleftrightarrow B$ , we have  $\overrightarrow{B} = cl_{\overrightarrow{\tau}}(B) \subseteq A^c$ .

(2) If  $x \in B$ , then  $\{x\} \hookrightarrow B$ . This implies that  $\{x\} \nleftrightarrow A$ . Because if  $\{x\} \hookrightarrow A$ , then  $A \hookrightarrow B$  by Lemma 5.2. Hence,  $x \notin cl_{\overrightarrow{\tau}}(A)$  which means  $x \in (cl_{\overrightarrow{\tau}}(A))^c = int_{\overrightarrow{\tau}}(A^c)$ . Hence, we have  $B \subseteq int_{\overrightarrow{\tau}}(A^c)$ .

**Theorem 5.19.** Let  $(X, \, \hookrightarrow, \mathcal{P})$  be a primal-proximity space and  $A, B \subseteq X$ . Then,  $A \hookrightarrow B$  if and only if  $cl_{\overrightarrow{\tau}}(A) \hookrightarrow cl_{\overrightarrow{\tau}}(B)$ , where  $cl_{\overrightarrow{\tau}}(A)$  means the closure of A with respect to  $\overrightarrow{\tau}$ .

*Proof.* If  $A \hookrightarrow B$ , then by Lemma 4.2,  $cl_{\overrightarrow{\tau}}(A) \hookrightarrow cl_{\overrightarrow{\tau}}(B)$  since  $A \subseteq cl_{\overrightarrow{\tau}}(A)$  and  $B \subseteq cl_{\overrightarrow{\tau}}(B)$ .

If  $A \nleftrightarrow B$ , then there exists a subset E of X such that  $A \nleftrightarrow E$  and  $E^c \nleftrightarrow B$ and  $(E \cap E^c)^c \notin \mathcal{P}$ . Hence,  $cl_{\overrightarrow{\tau}}(B) \subseteq E$  by (1) of Theorem 5.18. This implies that  $A \nleftrightarrow cl_{\overrightarrow{\tau}}(B)$ . Because if  $A \hookrightarrow cl_{\overrightarrow{\tau}}(B)$  then by Lemma 4.2, then  $A \hookrightarrow E$ since  $cl_{\overrightarrow{\tau}}(B) \subseteq E$ . Now, if  $A \nleftrightarrow B$  then  $A \nleftrightarrow cl_{\overrightarrow{\tau}}(B)$ . Also,  $cl_{\overrightarrow{\tau}}(B) \nleftrightarrow A$  by similar prove again it follows that  $cl_{\overrightarrow{\tau}}(B) \nleftrightarrow cl_{\overrightarrow{\tau}}(A)$ . Hence,  $A \hookrightarrow B$  if and only if  $cl_{\overrightarrow{\tau}}(A) \hookrightarrow cl_{\overrightarrow{\tau}}(B)$ .

### 6 Conclusion

In this paper, we introduced a new type of proximity space called primal-proximity space. Later, we defined point-primal proximity operator and investigated some of its fundamental properties. We also proved that this operator is a Kuratowski closure operator under special condition. Moreover, one more operator via pointprimal proximity operator was defined. Furthermore, we gave not only some relationships but also several examples.

Conflict of interest: The authors declare that there is no conflict of interest.

#### References

- S. Acharjee, M. Ozkoç and F.Y. Issaka, *Primal topological spaces*, arXiv:2209.12676v1 [math.GM]
- [2] A. Al-Omari, S. Acharjee and M. Ozkoç, A new operator of primal topological spaces, arXiv:2210.17278v [math.GN]
- [3] G. Choquet, Sur les notions de filter et grille, Comptes Rendus Acad.Sci. Paris, 224 (1947), 171-173.
- [4] V.A. Efremovič, The geometry of proximity, Mat. Sbornik 31(73) (1952), 189-200.
- [5] R.A. Hosny and O.A. Tantawy, New proximities from old via ideals, Acta Mathematica Hungarica, 110(1-2) (2006), 37–50.
- [6] R.A. Hosny, Relations and applications on proximity structures, Gen, 11(1) (2012), 24–40.
- [7] D. Jankovic and T. R. Hamlett, New topologies from old via ideals, American math. monthly, 97(4) (1990), 295-310.
- [8] W.J. Thron, Proximity structures and grills, Math. Ann., 206 (1973), 35-62.
- [9] B. Roy and M.N. Mukherjee, On a typical topology induced by a grill, Soochow Jour. Math., 33(4) (2007), 771-786.
- [10] A. Kandil, O.A. Tantawy, S.A. El-Sheikh and A. Zakaria, *I-proximity spaces*, Jökull Journal, 63(5) (2013), 237-245.
- [11] A. Kandil, O.A. Tantawy, S.A. El-Sheikh and A. Zakaria, New structures of proximity spaces, Information Sciences Letters, 3(3) (2014), 85-89.
- [12] A. Kandil, O.A. Tantawy, S.A. El-Sheikh and A. Zakaria, *Multiset proximity spaces*, Journal of Egyptian Mathematical Society, 24 (2016), 562-567.
- [13] A. Kandil, O.A. Tantawy, S.A. El-Sheikh and A. Zakaria, *Generalized I-proximity spaces*, Journal of Egyptian Mathematical Society, 24 (2016), 562-567.
- [14] A. Kandil, S.A. El-Sheikh, M.M. Yakout and S.A. Hazza, Proximity structures and ideals, Matematicki Vesnik, 67(2) (2015), 130–142.
- [15] K. Kuratowski, Topologie I, Warszawa, 1933.

- [16] S. Leader, On clusters in proximity spaces, Fundamenta Mathematicae, 47(2) (1959), 205–213.
- [17] M. Lodato, On topologically induced generalized proximity relations. II, Pacific Journal of Mathematics, 17(1) (1966), 131–135.
- [18] S. Modak, Topology on grill-filter space and continuity, Bol. Soc. Paran. Mat., 31(2) (2013), 219-230.
- [19] S. Modak, Grill-filter space, Jour. Indian Math. Soc., 80(3-4) (2013), 313-320.
- [20] M.N. Mukherjee, D. Mandal and D. Dipankar, Proximity structure on generalized topological spaces, Afrika Matematika, 30 (2019), 91–100.
- [21] E.F. Steiner, The relation between quasi-proximities and topological spaces, Math. Ann. 155 (1964), 194-195.
- [22] W.J. Thron, Proximity structures and grills, Mathematische Annalen, 206 (1973), 35–62.
- [23] S. Tiwari and P.K. Singh, *Čech rough proximity spaces*, Matematicki Vesnik, 72(1) (2020), 6–16.
- [24] E.D. Yıldırım, μ-proximity structure via hereditary classes, Maejo International Journal of Science and Technology, 15(2) (2021), 129–136.
- [25] J. Brennan and E. Martin (2012). Spatial proximity is more than just a distance measure, Inter. Jour. Human-comp. stud., 70(1) (2012), 88-106.
- [26] A.A. Nasef and A.A. Azzam, Some topological operators via grills, Jour. Linear Top. Alg., 5(3) (2016), 199-204.
- [27] A.A. Azzam, S.S. Hussein and H. Saber Osman, Compactness of topological spaces with grills, Italian. Jour. Pure. Appl. Math., 44 (2020), 198–207.
- [28] A. Talabeigi, On the Tychonoff's type theorem via grills, Bull. Iranian Math. Soc., 42(1) (2016), 37–41.
- [29] N. Boroojerdian and A. Talabeigi, One-point λ-compactification via grills, Iran. Jour. Sci. Tech. Trans.A: Sci., 41 (2017), 909–912.
- [30] M.N. Mukherjee and A. Debray, On H-closed spaces and grills, An. Stiint. Univ. AL. I. Cuza Iasi. Mat.(N. S.), 44 (1998), 1-25.
- [31] S. Willard, *General topology*, Courier Corporation, 2012

- [32] J. F. Peters Local near sets: Pattern discovery in proximity spaces, Mathematics in Comput. Sci., 7 (2013), 87-106.
- [33] G. Dimov and D. Vakarelov, Contact algebras and region-based theory of space: a proximity approach-I, Fund. Info., 74(2-3) (2006), 209-249.
- [34] G. Dimov and D. Vakarelov, Contact algebras and region-based theory of space: proximity approach-II, Fund. Info., 74(2-3) (2006), 251-282.
- [35] I. Düntsch and D. Vakarelov, Region-based theory of discrete spaces: A proximity approach, Annals of Math. Art. Int., 49(1-4) (2007), 5-14.