

On coloring parameters of triangle-free planar (n, m) -graphs

SOUMEN NANDI^a, SAGNIK SEN^b, S TARUNI^b

(a) Netaji Subhas Open University, India.

(b) Indian Institute of Technology Dharwad, India.

Abstract

An (n, m) -graph is a graph with n types of arcs and m types of edges. A homomorphism of an (n, m) -graph G to another (n, m) -graph H is a vertex mapping that preserves the adjacencies along with their types and directions. The order of a smallest (with respect to the number of vertices) such H is the (n, m) -chromatic number of G . Moreover, an (n, m) -relative clique R of an (n, m) -graph G is a vertex subset of G for which no two distinct vertices of R get identified under any homomorphism of G . The (n, m) -relative clique number of G , denoted by $\omega_{r(n,m)}(G)$, is the maximum $|R|$ such that R is an (n, m) -relative clique of G . In practice, (n, m) -relative cliques are often used for establishing lower bounds of (n, m) -chromatic number of graph families.

Generalizing an open problem posed by Sopena [Discrete Mathematics 2016] in his latest survey on oriented coloring, Chakroborty, Das, Nandi, Roy and Sen [Discrete Applied Mathematics 2022] conjectured that $\omega_{r(n,m)}(G) \leq 2(2n+m)^2+2$ for any triangle-free planar (n, m) -graph G and that this bound is tight for all $(n, m) \neq (0, 1)$. In this article, we positively settle this conjecture by improving the previous upper bound of $\omega_{r(n,m)}(G) \leq 14(2n+m)^2+2$ to $\omega_{r(n,m)}(G) \leq 2(2n+m)^2+2$, and by finding examples of triangle-free planar graphs that achieve this bound. As a consequence of the tightness proof, we also establish a new lower bound of $2(2n+m)^2+2$ for the (n, m) -chromatic number of the family of triangle-free planar graphs.

Keywords: graph homomorphisms, (n, m) -graphs, (n, m) -chromatic number, (n, m) -relative clique number, triangle-free planar graphs.

1 Introduction

The focus of this work is on the resolution of a recent conjecture proposed Chakroborty, Das, Nandi, Roy and Sen [9] which is in the context of homomorphisms of (n, m) -graphs, introduced in 2000 by Nešetřil and Raspaud [33]. The conjecture generalizes an open question posed by Sopena [37] in his survey on oriented graphs.

1.1 Colored mixed graphs

An (n, m) -colored mixed graph, or simply, an (n, m) -graph G is a graph with n different types of arcs and m different types of edges. We denote the set of vertices, arcs, and edges of G by $V(G)$, $A(G)$, and $E(G)$ respectively. Also, we denote the underlying graph of G by $und(G)$.

In this article, we restrict ourselves to studying only those (n, m) -graphs whose underlying graphs are simple graphs. We can capture the notions of simple graphs when $(n, m) = (0, 1)$ [38], oriented graphs when $(n, m) = (1, 0)$ [22, 35, 36, 37], 2-edge-colored graphs or signed graphs when $(n, m) = (0, 2)$ [27, 31, 32, 34, 39], and m -edge colored graphs when $(n, m) = (0, m)$ [1]. Therefore, (n, m) -graphs help us investigate well-studied families of graphs in a generalized set-up. Moreover, the upcoming definitions of graph homomorphisms, chromatic numbers, and clique numbers for (n, m) -graphs truly capture the existing concepts of the same for all values of (n, m) , including $(0, 1)$. In this work though, whenever we use the term (n, m) -graphs, we mean it for all values of $(n, m) \neq (0, 1)$ unless otherwise stated, for convenience.

1.2 Homomorphisms, chromatic numbers, and clique numbers

A *homomorphism* of an (n, m) -graph G to another (n, m) -graph H is a vertex mapping $f : V(G) \rightarrow V(H)$ such that for any arc (resp., edge) xy in G , their images induces an arc (resp., edge) $f(x)f(y)$ of the same type in H . If there exists a homomorphism of G to H , then we denote it by $G \rightarrow H$.

Using the concept of homomorphisms of such graphs, Nešetřil and Raspaud [33] further presented a generalization to the notion of chromatic number by introducing the (n, m) -chromatic number of an (n, m) -graph as

$$\chi_{n,m}(G) := \min\{|V(H)| : G \rightarrow H\}.$$

For a family \mathcal{F} of simple graphs, the (n, m) -chromatic number can be defined as

$$\chi_{n,m}(\mathcal{F}) := \max\{\chi_{n,m}(G) : und(G) \in \mathcal{F}\}.$$

This parameter is studied for the family of graphs having bounded acyclic chromatic numbers, bounded arboricity and acyclic chromatic number, bounded maximum degree, sparse graphs, planar graphs and planar graphs with girth restrictions, partial k -trees and partial k -trees with girth restrictions, outerplanar graphs and outerplanar graphs with girth restrictions, etc. across several papers [17, 24, 28, 33].

Bensmail, Duffy, and Sen [4] contributed to this line of work by introducing and studying generalizations of the concept of clique number. The generalization ramifies into two parameters in the context of (n, m) -graphs. An (n, m) -*relative clique* $R \subseteq V(G)$ is a vertex subset satisfying $|f(R)| = |R|$ for all homomorphisms f of G to any H . The (n, m) -relative clique number of a graph G denoted by $\omega_{r(n,m)}(G)$, is,

$$\omega_{r(n,m)}(G) = \max\{|R| : R \text{ is an } (n, m)\text{-relative clique of } G\}.$$

An (n, m) -*absolute clique* $A \subseteq G$ is a subgraph of G satisfying $\chi_{n,m}(A) = |V(A)|$. The (n, m) -absolute clique number of G denoted by $\omega_{a(n,m)}(G)$, is,

$$\omega_{a(n,m)}(G) = \max\{|V(A)| : A \text{ is an } (n, m)\text{-absolute clique of } G\}.$$

For family \mathcal{F} of simple graphs, both the parameters are defined similarly like (n, m) -chromatic number. That is,

$$p(\mathcal{F}) := \max\{p(G) : und(G) \in \mathcal{F}\}$$

where $p \in \{\omega_{r(n,m)}, \omega_{a(n,m)}\}$. The two parameters are primarily being studied for graphs with bounded degrees, planar, partial 2-trees and outerplanar graphs, and their subfamilies due to girth restrictions [4, 9].

1.3 Context and motivation

Observe that the above defined three parameters trivially satisfy the following relation [4]

$$\omega_{a(n,m)}(G) \leq \omega_{r(n,m)}(G) \leq \chi_{n,m}(G).$$

Therefore, the study of one impacts the other. The best known lower and upper bounds for the (n, m) -chromatic number of planar graphs is given below. The lower bounds for (n, m) -chromatic number of planar graphs was given by Fabila-Monroy, Flores, Huemer and Montejano[17] while the upper bound was due to Nešetřil and Raspaud [33]. Let the family of planar graphs with girth at least g is denoted by \mathcal{P}_g .

Theorem 1.1 ([17, 33]). *For the family of planar graphs \mathcal{P}_3 we have,*

$$\begin{aligned} (2n + m)^3 + 2(2n + m)^2 + (2n + m) + 1 &\leq \chi_{n,m}(\mathcal{P}_3) \leq 5(2n + m)^4, \text{ for } m > 0 \text{ even} \\ (2n + m)^3 + (2n + m)^2 + (2n + m) + 1 &\leq \chi_{n,m}(\mathcal{P}_3) \leq 5(2n + m)^4, \text{ otherwise.} \end{aligned}$$

The above theorem can be treated as an approximate analogue of the Four-Color Theorem in the context of (n, m) -graphs. For triangle-free planar graphs, no such dedicated studies have been made, that is, an approximate analogue of the Grötzsch's Theorem is open till date. On the other hand, it turns out, even though finding clique numbers for these two families, that is, the families of planar and triangle-free planar graphs, is trivial for $(n, m) = (0, 1)$, and it is quite a challenging problem for the other values of (n, m) . There have been dedicated studies to explore these parameters even for $(n, m) = (1, 0)$ [12, 13], and later for general values [4, 9].

We recall the existing bounds for the two clique numbers for the families of planar and triangle-free graphs to place our work in context. For convenience, we present all these results under one theorem. The bounds for the (n, m) -absolute clique number of planar graphs \mathcal{P}_3 is proved by Bensmail, Duffy and Sen [4] while the rest are due to Chakraborty, Das, Nandi, Roy and Sen [9].

Theorem 1.2. *For the families \mathcal{P}_3 of planar and \mathcal{P}_4 triangle-free planar graphs, we have*

- (i) $3(2n + m)^2 + (2n + m) + 1 \leq \omega_{a(n,m)}(\mathcal{P}_3) \leq 9(2n + m)^2 + 2(2n + m) + 2$ [4],
- (ii) $3(2n + m)^2 + (2n + m) + 1 \leq \omega_{r(n,m)}(\mathcal{P}_3) \leq 42(2n + m)^2 - 11$ [9],
- (iii) $\omega_{a(n,m)}(\mathcal{P}_4) = (2n + m)^2 + 2$ [9],
- (iv) $(2n + m)^2 + 2 \leq \omega_{r(n,m)}(\mathcal{P}_4) \leq 14(2n + m)^2 + 1$ [9].

If one notices, except the (n, m) -absolute clique number of triangle-free planar graphs, exact general values are not known for the other parameters from Theorem 1.2.

The inception of the analogue to clique number in this set up was due to Klostermeyer and MacGillivray [21] where they defined the notion of absolute clique number for $(1, 0)$ -graphs. In the same work, they proved

$$\omega_{a(1,0)}(\mathcal{P}_4) \leq 14 \text{ and } \omega_{a(1,0)}(\mathcal{P}_3) \leq 36,$$

and conjectured $\omega_{a(1,0)}(\mathcal{P}_3) = 15$. Nandy, Sen and Sopena [30] proved the tight bounds

$$\omega_{a(1,0)}(\mathcal{P}_4) = 6 \text{ and } \omega_{a(1,0)}(\mathcal{P}_3) = 15$$

and thus, positively settled the above mentioned conjecture.

While proving the conjecture, Nandy, Sen and Sopena [30] introduced the notion of relative cliques for $(1, 0)$ -graphs which was a useful tool in their proofs. Then relative clique number of $(1, 0)$ -graphs were studied by Das, Prabhu and Sen [12] in a focused and systematic manner. In particular, they proved

$$10 \leq \omega_{r(1,0)}(\mathcal{P}_4) \leq 14 \text{ and } 15 \leq \omega_{r(1,0)}(\mathcal{P}_3) \leq 32.$$

In the meantime, Sopena [37] enquired the exact values of $\omega_{r(1,0)}(\mathcal{P}_3)$ and $\omega_{r(1,0)}(\mathcal{P}_4)$ in his survey in 2016 (latest to date) on oriented coloring as open problems. Das, Nandi and Sen [13] solved the second problem by showing $\omega_{r(1,0)}(\mathcal{P}_4) = 10$ while the first question remains open still.

Later, Bensmail, Duffy and Sen [4] generalized the notions of relative and absolute clique numbers for all (n, m) -graphs and proved Theorem 1.2(i). They [4] also conjectured that lower bound of Theorem 1.2(i) is tight, generalizing the result $\omega_{a(1,0)}(\mathcal{P}_3) = 15$ due to Nandy, Sen and Sopena [30].

Conjecture 1.3 ([4]). *For the family \mathcal{P}_4 of triangle-free planar graphs we have*

$$\omega_{a(n,m)}(\mathcal{P}_3) = 3(2n + m)^2 + (2n + m) + 1.$$

Extending this direction of research, Chakraborty, Das, Nandi, Roy and Sen [9] proved a tight bound for the (n, m) -absolute clique number of triangle-free planar graphs via Theorem 1.2(iii). They also provided lower and upper bounds for the relative clique numbers of planar and triangle-free planar graphs in Theorem 1.2(ii) and (iv). Moreover, in the same work, they made two conjectures about the exact values of $\omega_{r(n,m)}(\mathcal{P}_3)$ and $\omega_{r(n,m)}(\mathcal{P}_4)$, generalizing the open problems proposed by Sopena [37] in the context of $(1, 0)$ -graphs.

Conjecture 1.4 ([9]). *For the family \mathcal{P}_3 of planar graphs we have*

$$\omega_{r(n,m)}(\mathcal{P}_3) = 3(2n + m)^2 + (2n + m) + 1.$$

Conjecture 1.5 ([9]). *For the family \mathcal{P}_4 of triangle-free planar graphs we have*

$$\omega_{r(n,m)}(\mathcal{P}_4) = 2(2n + m)^2 + 2.$$

1.4 Our contributions

As mentioned before, Das, Nandi and Sen [13] solved Conjecture 1.5 for $(n, m) = (1, 0)$. Moreover, they noted that the solution for $(n, m) = (0, 2)$ is similar. Thus, we may consider the conjecture to be solved when $(2n + m) = 2$. In this article, we positively settle this conjecture when $(2n + m) \geq 3$ by proving the following.

Theorem 1.6. *For the family \mathcal{P}_4 of triangle-free planar graphs we have*

$$\omega_{r(n,m)}(\mathcal{P}_4) = 2(2n + m)^2 + 2$$

for $(2n + m) \geq 3$.

As a direct consequence, we find a new lower bound for the (n, m) -chromatic number of triangle-free planar graphs.

Theorem 1.7. *For the family \mathcal{P}_4 of triangle-free planar graphs we have*

$$\chi_{n,m}(\mathcal{P}_4) \geq 2(2n + m)^2 + 2$$

for $(2n + m) \geq 3$.

1.5 The significance of relative cliques

The focus parameter of this article, (n, m) -relative cliques, is not such a popularly studied parameter. However, a closer look may imply otherwise, hence the need of this subsection. As (n, m) -absolute cliques are special cases of (n, m) -relative cliques, we will try to point out works done on both areas to highlight the significance of (n, m) -relative cliques. Every point written below is to the best of our knowledge.

- For (n, m) -graphs, there have been two works dedicated on (n, m) -absolute and (n, m) -relative cliques: [4] and [9].
- For oriented graphs, that is, for $(n, m) = (1, 0)$, a number of papers dealt with the notions of absolute and relative cliques: [5, 12, 13, 15, 21, 30]. A similar variant of homomorphisms of oriented graphs, called pushable homomorphisms of oriented graphs, also studies the notions of absolute and relative cliques [7].
- For 2-edge-colored graphs, that is, for $(n, m) = (0, 2)$, a some of papers dealt with the notions of absolute and relative cliques. However, it is usually for a particular variant of homomorphisms of 2-edge colored graphs, called homomorphisms of signed graphs, which recently gained popularity due to its underlying connections with the theory of graph coloring. The notions of absolute and relative cliques exists, and are relevant, for signed graphs also. The following are some works on them: [6, 11, 10, 31].

The (n, m) -relative (resp., absolute) clique numbers are natural lower bounds for (n, m) -chromatic numbers. In practice, one of the most popularly used methods to establish a lower bound for (n, m) -chromatic number of a family of graphs is to provide an

example of a graph from that family having high (n, m) -relative clique number. This trick has been used even before the notions of absolute and relative cliques came into existence. This is possibly the most important significance of the notions of absolute and relative cliques. In the following we list such instances.

- For (n, m) -graphs, lower bounds for (n, m) -chromatic number of several graph families were established using examples having high (n, m) -relative clique number across the following works: [17, 28, 33].
- For oriented graphs, that is, for $(n, m) = (1, 0)$, lower bounds for $(1, 0)$ -chromatic number (popularly known as oriented chromatic number) of several graph families were established using examples having high (n, m) -relative clique number across a number of works. Most of them can be found in the latest survey by Sopena [37]. We would like to make a special mention to [25, 26], where relative cliques and the analysis of their interactions was used to improve the lower bound for the oriented chromatic number of the family of planar graphs. Similar phenomenon can be observed for pushable chromatic number of oriented graphs in the following works: [2, 22].
- For 2-edge colored graphs, that is, for $(n, m) = (0, 2)$, $(0, 2)$ -chromatic number¹ and of chromatic number of signed graphs, the use of absolute and relative cliques to establish lower bounds can be observed in the following works: [3, 31, 34].

In particular for oriented graphs, that is, for $(n, m) = (1, 0)$, which is also historically the oldest and most studied specific case of homomorphisms of (n, m) -graphs, absolute and relative cliques appear in two other significant areas.

- Borodin, Fon-Der-Flaass, Kostochka, Raspaud, and Sopena [8] introduced and studied the notion of deeply critical oriented graphs and asked the existence of “deeply critical $(1, 0)$ -absolute cliques” (even before the term absolute clique came into existence) on prescribed number, of vertices. They themselves proved such existence when the number of vertices are of the form $2 \cdot 3^n - 1$. Later, answering to an open question asked in the above mentioned paper, Duffy, Pavan, Sandeep, and Sen [14] found examples of such $(1, 0)$ -absolute cliques for all odd values of number of vertices, except 1, 3, and 7.
- In relation to the famous degree-diameter problem, Erdős, Rényi and Sós [16] defined the function

$$h_d(n) = \min\{|E(G)| : \text{diam}(G) \leq d \text{ and } |V(G)| = n\}$$

as the minimum number of edges among graphs having diameter at most d . Its directed (oriented) analogue turns out to be very interesting even for $d = 2$. In fact, the oriented graphs having (weak) diameter 2 are exactly the $(1, 0)$ -absolute cliques. A number of works are dedicated in quest of finding the exact description of this function [18, 20, 23].

¹This parameter has been studied under several different names across the literature, such as: 2-edge-colored chromatic number, signified chromatic number, and signed-preserving chromatic number.

Thus, (n, m) -relative cliques are important objects of study in the context of homomorphisms of (n, m) -graphs.

1.6 Organization of the article

This article is structured as follows. In Section 2 we give the necessary definitions and notations used in this article. In Section 3 we prove Theorem 1.6. In Section 4 we share our concluding remarks.

Note: A preliminary version of this article has been published in IWOCA 2022 [29].

2 Preliminaries

In this section, we introduce some notation to help us write the proofs. Also, we follow West [38] for standard definitions, notation and terminology.

Given an (n, m) -graph G the different types of arcs in G are distinguished by n different labels $2, 4, \dots, 2n$. To be precise, an arc xy with label $2i$ is an arc of type i from x to y . In fact, in such a situation, y is called a $2i$ -neighbor of x , or equivalently, x is called a $(2i - 1)$ -neighbor of y . The different types of edges in G are distinguished by m different types of labels $2n + 1, 2n + 2, \dots, 2n + m$. Here, an edge xy with label $2n + j$ is an edge of type $2n + j$ between x and y . In this case, x and y are called $(2n + j)$ -neighbors of each other.

Usually, throughout the article, we use the Greek alphabets such as α, β, γ and their variants (like α', β', γ' , etc.) to denote these labels. Therefore, whenever we use such a symbol, say α , one may assume it to be an integer between 1 and $(2n + m)$. Let us denote the set of all α -neighbors of x as $N^\alpha(x)$. Two vertices x, y agree on a third vertex z if $z \in N^\alpha(x) \cap N^\alpha(y)$ for some α , and disagree on z otherwise.

A *special 2-path* is a 2-path uvw such that u, v disagrees with each other on w . In an (n, m) -graph G , a vertex u sees a vertex v if they are either adjacent, or are connected by a special 2-path. If u and v are connected by a special 2-path with w as the *internal vertex*, then it is said that u sees v via w or equivalently, v sees u via w . We recall a useful characterization of an (n, m) -relative clique due to Bensmail, Duffy and Sen [4].

Lemma 2.1 ([4]). *Two distinct vertices of an (n, m) -graph G are part of a relative clique if and only if they are either adjacent or connected by a special 2-path in G , that is, they see each other.*

3 Proof of Theorem 1.6

The proof is lengthy and involved. So for the reader's convenience, we have presented it under separate indicative headers and have distributed it into different propositions, lemmas, and observations.

3.1 Lower bound

The lower bound of Theorem 1.6 is given by the following proposition.

Proposition 3.1. *There exists a triangle-free planar (n, m) -graph G such that*

$$\omega_{r(n,m)}(G) = 2(2n + m)^2 + 2.$$

Proof. We construct an example to prove this claim. We start with a $K_{2,2(2n+m)^2}$. Let the vertices in the partite set consisting of only two vertices be x and y . Let the vertices from the other partite set be

$$\{g_{\alpha\beta}, g'_{\alpha\beta} : \text{for all } \alpha, \beta \in \{1, 2, \dots, 2n + m\}\}.$$

Next, assign colors and directions to the edges in such a way that $g_{\alpha\beta}$ and $g'_{\alpha\beta}$ become α -neighbors of x and β -neighbors of y . Finally, add special 2-paths of the form $g_{\alpha\beta}h_{\alpha\beta}g'_{\alpha\beta}$ connecting $g_{\alpha\beta}$ and $g'_{\alpha\beta}$.

Notice that, $g_{\alpha\beta}$ (or $g'_{\alpha\beta}$) sees $g_{\alpha'\beta'}$ (or $g'_{\alpha'\beta'}$) via x if $\alpha \neq \alpha'$ or via y if $\beta \neq \beta'$. If $\alpha = \alpha'$ and $\beta = \beta'$, then they are the two vertices $g_{\alpha\beta}$ and $g'_{\alpha\beta}$ and they see each other via $h_{\alpha\beta}$. On the other hand, x and y are adjacent to each $g_{\alpha\beta}$, while they see each other via $g_{\alpha'\beta'}$ for some $\alpha' \neq \beta'$. Thus, the $g_{\alpha\beta}$'s together with x and y , form an (n, m) -relative clique of cardinality $2(2n + m)^2 + 2$. \square

3.2 Strategy and set up for the upper bound

Next, we concentrate on the upper bound. We prove the upper bound by the method of contradiction. Therefore, let us assume that $\omega_{r(n,m)}(\mathcal{P}_4) > 2(2n + m)^2 + 2$.

Let $n_3(G)$ be the number of vertices with a degree at least 3. A *critical (n, m) -relative clique H* for the family \mathcal{P}_4 of triangle-free planar graphs is an (n, m) -graph H satisfying the following properties:

- (i) $und(H) \in \mathcal{P}_4$,
- (ii) $\omega_{r(n,m)}(H) = \omega_{r(n,m)}(\mathcal{P}_4)$,
- (iii) $\omega_{r(n,m)}(H^*) < \omega_{r(n,m)}(H)$, if

$$(n_3(H^*), |V(H^*)|, |E(und(H^*))|) < (n_3(H), |V(H)|, |E(und(H))|)$$

in the dictionary ordering, where $H^* \in \mathcal{P}_4$.

We consider a particular critical (n, m) -relative clique H for the rest of this proof. As H is a triangle-free planar graph, we fix a particular planar embedding of it. Whenever we deal with something dependent on the embedding of H or a portion of it, we refer to this particular fixed embedding. We also fix a particular (n, m) -relative clique R of cardinality $\omega_{r(n,m)}(H)$. Furthermore, the vertices of R are called *good vertices* and those of $S = V(H) \setminus R$ are called the *helper vertices*. Notice that,

$$\omega_{r(n,m)}(H) = |R| = \omega_{r(n,m)}(\mathcal{P}_4) > 2(2n + m)^2 + 2$$

due to our basic assumption. Thus, the goal of the proof is to contradict this strict inequality.




Figure 1: The exceptional configuration of Lemma 3.4. The symbols α and β are adjacency types with respect to x, y , respectively and h_1, h_2, h_3 denote the helper vertices.

3.3 Key lemmas

We recall some useful results due to Chakraborty, Das, Nandi, Roy and Sen [9].

Lemma 3.2 ([9]). *The (n, m) -graph H is connected and the set S of helper vertices in H is an independent set.*

We recall a modified version of another lemma due to Chakraborty, Das, Nandi, Roy and Sen [9] which we use for our proofs.

Lemma 3.3 ([9]). *A vertex $x \in V(H)$ can have at most $2(2n + m)$ good α -neighbors in H for any α .*

We prove a similar result for common good neighbors of two distinct vertices as well, but for an exception. Notice that, it is possible for two vertices x, y of H to agree on three good vertices as depicted in Fig. 1 where the good vertices are also able to see each other via distinct helpers. Turns out this is the only exception, otherwise, two distinct vertices of H can agree on at most 2 good vertices.

Lemma 3.4. *Two distinct vertices x, y of H can agree on at most 2 good vertices, except when x, y agree on exactly three good vertices which are mutually connected by internally vertex disjoint special two paths as in Fig. 1.*

Proof. Suppose x and y have three good neighbors g_1, g_2, g_3 from $N^\alpha(x) \cap N^\beta(y)$, and at least another common neighbor g_4 . Also assume that g_1, g_2, g_3, g_4 are arranged in an anti-clockwise order around x in the fixed planar embedding of H . Notice that, as H is triangle-free, g_1 must see g_3 via some $h \notin \{x, y, g_1, g_2, g_3, g_4\}$. This is not possible to achieve keeping the graph planar. Hence, if x and y have four common good neighbors, then it is not possible for three of their neighbors to agree on x and y .

If x and y have exactly three common neighbors g_1, g_2, g_3 , and all three agree on x and y , then as H is triangle-free, g_i must see g_j via some h_{ij} for all $i < j$. These h_{ij} 's must be distinct, otherwise a $K_{3,3}$ will be created. Hence Fig. 1 is forced. \square




Figure 2: The configuration \mathbb{F}_k .

In general, two vertices in H can have at most $2(2n + m)^2$ many common good neighbors.

Lemma 3.5. *Two distinct vertices x, y of H can have at most $2(2n + m)^2$ good vertices in their common neighborhood.*

Proof. As $(2n + m) \geq 2$, and thus, $2(2n + m)^2 \geq 8$, the exceptional case mentioned in Lemma 3.4 satisfies the condition of the statement of this lemma.

In other case, Lemma 3.4 implies that

$$|N^\alpha(x) \cap N^\beta(y) \cap R| \leq 2.$$

As α, β can be chosen from a set of $(2n + m)$ integers, we can have a total of $(2n + m)^2$ distinct (α, β) pairs. Therefore, there can be at most $2(2n + m)^2$ good vertices in $N(x) \cap N(y)$. \square

3.4 The forbidden configuration \mathbb{F}_k

The configuration \mathbb{F}_k consists of two vertices x, y of H which may not necessarily be good vertices and k common good neighbors of x, y , namely, g_1, g_2, \dots, g_k . Also, we assume a default embedding of \mathbb{F}_k where g_1, g_2, \dots, g_k are arranged in an anti-clockwise order around x . Without loss of generality, we may assume the embedding of \mathbb{F}_k , induced by the fixed embedding of H , to be such that the boundary of the unbounded region is the 4-cycle xg_1yg_kx . Thus the 4-cycle xg_1yg_kx divides the plane into two regions: a bounded region containing other g_i 's called R , and an unbounded region called R_0 . Moreover, the 2-paths of the type xg_iy further subdivides the region R into $(k - 1)$ regions. These regions are called R_1, R_2, \dots, R_{k-1} , where R_i is bounded by $xg_iyg_{i+1}x$, for $i \in \{1, 2, \dots, k - 1\}$. See Fig. 2 for a pictorial representation of this planar embedding.

If it is not possible for H to contain a \mathbb{F}_k for some k , then we say that \mathbb{F}_k is *forbidden* in H . Therefore, Lemma 3.5 essentially says that \mathbb{F}_k , for all $k \geq 2(2n + m)^2 + 1$ is forbidden. Next, we show that \mathbb{F}_k , for all $k \geq 3$ is forbidden as a key step of our proof.

Lemma 3.6. *The configuration \mathbb{F}_k , for all $k \geq 3$ is forbidden in H .*




Figure 3: The private good neighbors of x and y .

Proof. We prove this by strong backward induction. Suppose that the statement is true for all $k \geq t+1$ and we prove that it is true for $k = t$, where $t \geq 3$. The base case is taken care of by Lemma 3.5, where it is proved that \mathbb{F}_k is forbidden for all $k \geq 2(2n+m)^2 + 1$ in H .

Now, notice that, if a good vertex g which is not part of \mathbb{F}_k , belongs to region R_0 (resp., R_1, R_2), then it must see g_2 (resp., g_3, g_1) via x or y . Observe that, it is not possible for g to be adjacent to both x and y , as otherwise, it will create a \mathbb{F}_{t+1} , which is forbidden due to the induction hypothesis.

Therefore, every good vertex of H , other than x, y , is either adjacent to both x and y or exactly one of them. If in case, all of them are adjacent to x , then we can say that there are maximum $2(2n+m)^2$ good vertices in the neighborhood of x due to Lemma 3.3. This implies that H has at most $2(2n+m)^2 + 2$ good vertices, the additional two vertices coming from counting x and y . Thus it is not possible for all good neighbors other than x, y to be adjacent to x . Similarly, it is not possible for all good neighbors other than x, y to be adjacent to y .

Therefore, there must be at least one good vertex which is adjacent to x but not to y and at least one good vertex that is adjacent to y but not to x . Notice that, these vertices must belong to the same region R_i for some $i \in \{0, 1, \dots, R_{k-1}\}$, as otherwise they will not be able to see each other. Hence, without loss of generality, we may assume that all good vertices other than the ones contained in \mathbb{F}_k belong to R_1 . These good vertices are adjacent to exactly one of x and y . Let us assume that x_1, x_2, \dots, x_p are the good vertices adjacent to x and not to y while y_1, y_2, \dots, y_q are the good vertices adjacent to y and not to x . Also suppose that x_1, x_2, \dots, x_p are arranged in an anti-clockwise order around x and y_1, y_2, \dots, y_q are arranged in a clockwise order around y . Also for convenience, x_i 's are called private good neighbors of x and y_j 's are called private good neighbors of y . Similarly, g_i 's are called common good neighbors of x and y . See Fig. 3 for a pictorial reference. Furthermore, due to symmetry, we may assume that $p \geq q \geq 1$.

Observe that, if x has a private good α -neighbor, then it cannot have a common good α -neighbor other than, possibly, g_1, g_2 . A similar statement holds for y and its private neighbors as well. Thus, if x (resp., y) has a (resp., b) different types of adjacencies with its private neighbors, then x, y can have at most $(2n+m-a)(2n+m-b)$ types (with respect to adjacencies with x and y) of common neighbors (other than, possibly,

g_1, g_2). Thus, due to Lemma 3.4, assuming \mathbb{F}_k is not the exceptional case, there are at most $2(2n + m - a)(2n + m - b)$ common good neighbors of x, y (other than, possibly, g_1, g_2). On the other hand, due to Lemma 3.3, there are at most $2(2n + m)a$ private good neighbors of x and $2(2n + m)b$ private good neighbors of y . Notice that, it is possible to count g_1 (resp., g_2) among the number of common good neighbors of x, y if it is neither adjacent to x with any of the a types of adjacencies, nor to y with any of the b types of adjacencies. Otherwise, it is possible to count them along with the private neighbors. Thus the total number of good neighbors in H is at most

$$2(2n + m - a)(2n + m - b) + 2(2n + m)a + 2(2n + m)b + 2 = 2(2n + m)^2 + 2ab + 2.$$

Even though the above equation does not provide the bound we want, we will consider three different cases and in each case a modified version of the above equation will provide us the required bound.

The above equation gets modified when $p \geq q \geq 2$. In that case note that if x_1 is adjacent to y_q , then it is not possible for x_p to see y_1 . Hence, x_1 must see y_q via some h . This will force every x_i to see y_j via h , except for, maybe when $i = j = 1$ or when $i = p, j = q$. If x_1 agrees with y_1 on h , then it is not possible for any other x_i (resp., y_j), for $i, j \neq 1$, to agree with x_1 on h , as otherwise, they cannot see y_1 (resp., x_1). A similar statement holds if x_p agrees with y_q on h . Suppose if there are c types of adjacencies of h among private neighbors of x and d types of adjacencies among private neighbors of y , the value of $p + q$ can be at most $2ac + 2bd$. Therefore, without loss of generality assuming $a \geq b$, the revised upper bound of $p + q$ is

$$p + q \leq 2ac + 2bd \leq 2a(c + d) \leq 2(2n + m)a.$$

Hence, the revised upper bound for the total number of good neighbors in H is

$$2(2n + m - a)(2n + m - b) + 2(2n + m)a + 2 \leq 2(2n + m)^2 + 2.$$

Thus, we have a contradiction in the case when $p \geq q \geq 2$.

Next, let us concentrate on the case when $q = 1$. In this case, each x_i must see y via y_1 or via some h_i . However, if x_i sees y via some h_i and is non-adjacent to y_1 , then it must see y_1 via some h'_i . Hence, in the worst case scenario, x_1 and x_p will see y and y_1 via h_1, h_p and h'_1, h'_p , respectively while the other x_i 's will see y via y_1 . In this case, unless $p \leq 2$, it is not possible for x_1 to agree with x_p on x . Also, x_i 's (for $i \neq 1, p$) disagree with y on y_1 . Therefore, $p + q \leq 2(2n + m)a + 1$ in this scenario. Hence, when $q = 1$ and $p \geq 3$, the total number of good vertices in H is at most

$$2(2n + m - a)(2n + m - 1) + 2(2n + m)a + 1 + 2 \leq 2(2n + m)^2 + 3 - 2(2n + m - a).$$

Observe that $a < 2n + m$, as none of the private neighbors of x can agree with g_3 on x . Hence, $(2n + m - a) \geq 1$, which implies that the total number of good vertices in H is at most $2(2n + m)^2 + 2$.

Thus, we have a contradiction in the case when $p \geq 3$ and $q = 1$.

On the other hand, if $p \leq 2$, then the total number of private neighbors is at most 3. In this scenario, the total number of good vertices in H is at most

$$2(2n + m - a)(2n + m - 1) + 3 + 2 \leq 2(2n + m)^2 - 2(2n + m)(a + 1) + a + 3 + 2.$$

Notice that the above bound is at most $2(2n + m)^2 + 2$ as $2n + m \geq 2$ and $a = 1$ or 2 . Thus, we have a contradiction in the case when $p \leq 2$.

This completes the proof, except when \mathbb{F}_k is the exceptional case. If \mathbb{F}_k is the exceptional case, that is, the one described in Fig. 1, and assume that the common neighbors are α -neighbors of x and β -neighbors of y . Then x (resp., y) cannot have any private α -neighbor (resp. β -neighbor) as they cannot see all the three common neighbors. Therefore, $a, b \leq 2n + m - 1$. As there are exactly three common neighbors, the total number of good vertices in H is at most

$$2(2n + m - 1)(2n + m - 1) + 3 + 2 \leq 2(2n + m)^2 - 4(2n + m) + 7.$$

As $(2n + m) \geq 3$, the above bound is at most $2(2n + m)^2 + 2$. This is a contradiction, and this completes the proof. \square

3.5 Proving the existence of a good vertex with low degree

We now restrict the number of good neighbors for any vertex of H . To do so, we need some supporting results.

Lemma 3.7. *It is not possible for a vertex $x \in V(H)$ to have four or more good neighbors, with at least three of them being α -neighbors.*

Proof. Suppose x has at least four good neighbors g_1, g_2, g_3, g_4 arranged in an anti-clockwise order around x in the fixed embedding of H , where g_1, g_2, g_3 are α -neighbors.

Observe that, as H is a triangle-free graph, the only way for g_1 to see g_3 is via some vertex $h \notin \{x, g_1, g_2, g_3, g_4\}$. Similarly, g_2 must see g_i via some h_i , for all $i \in \{1, 3\}$. Notice that, h, h_1, h_3 are all distinct vertices, as otherwise, a \mathbb{F}_3 will be created which is forbidden due to Lemma 3.6. Also as there is no way for h_1 and h_3 to see g_4 , thus h_1, h_3 cannot be good vertices.

On the other hand, the 4-cycle xg_1hg_3x divides the plane into two connected regions: the bounded region containing g_2 and the unbounded region containing g_4 . Let us call the unbounded region as R . Notice that, the bounded region is further subdivided into three regions, say R_1, R_2 , and R_3 bounded by the cycles $xg_1h_1g_2x$, $xg_2h_3g_3x$, and $g_1h_1g_2h_3g_3hg_1$, respectively.

Notice that, any good vertex g belonging to R_3 must see g_4 via h . However, then x and h will have three common good neighbors g_1, g_3, g_4 , creating a \mathbb{F}_3 . Therefore, R_3 cannot contain any good vertices.

Furthermore, any good vertex belonging to R_1 or R_2 must see g_4 via x , and any vertex belonging to R must see g_2 via x . Thus, every good vertex, maybe except h and x itself, is adjacent to x . As x can have at most $2(2n + m)^2$ good neighbors due to Lemma 3.3, there can be at most $2(2n + m)^2 + 2$ good vertices in H , counting x and h , a contradiction. \square

With the above lemma, we can further restrict the number of good neighbors of any vertex in H .

Lemma 3.8. *A vertex $x \in V(H)$ can have at most $2(2n + m)$ good neighbors in H .*

Proof. Suppose x has at least $2(2n + m) + 1$ good neighbors. Note that $2(2n + m) + 1 \geq 5$ as $2n + m \geq 2$. By the Pigeonhole principle, x must have at least three α -neighbors for some adjacency type α . According to Lemma 3.7, this is not possible, a contradiction. \square

Lemma 3.9. *The (n, m) -graph H has no helper vertex of degree three.*

Proof. Suppose u is a helper vertex in H of degree three having neighbors u_1, u_2 and u_3 . We delete the vertex u and add three degree two helper vertices h_{12}, h_{23}, h_{31} such that h_{ij} is adjacent to u_i and u_j . Moreover, if u_i (resp., u_j) is an α -neighbor of u then it is also an α -neighbor of h_{ij} . Let the so obtained graph be H^* .

Note that H^* is a triangle-free planar graph satisfying $n_3(H^*) < n_3(H)$. Observe that, if two good vertices of H see each other directly or via a helper other than u , then they see each other in the same way in H^* . On the other hand, the only vertices that see each other via u in H are u_i and u_j for $i, j \in \{1, 2, 3\}$ while in H^* they see each other via h_{ij} . Hence, $\omega_{r(n,m)}(H^*) \geq \omega_{r(n,m)}(H)$ as the maximum (n, m) -relative clique R in H remains an (n, m) -relative clique in H^* as well. This contradicts the criticality of H . \square

We recall a useful result for planar graphs.

Theorem 3.10 ([19]). *Every planar graph G with $\delta(G) \geq 4$, contains an edge xy with $\deg(x) = 4$ and $4 \leq \deg(y) \leq 7$, or contains an edge xy , with $\deg(x) = 5$ and $5 \leq \deg(y) \leq 6$. The bounds are the best possible.*

Lemma 3.11. *There exists a good vertex in H of degree at most seven. Moreover, if all good vertices have a degree of at least seven, then there exists a good vertex of degree seven in H with a neighbor which is a helper vertex of degree four.*

Proof. We know that there exists no helper vertex of degree zero or one in H due to its criticality and degree three in H by Lemma 3.9. Let us delete each helper vertex of degree two in H and put an edge between its neighbors. Let the so obtained graph be H' . Note that, two distinct helper vertices of degree two in H do not have the same common neighborhood as H is critical. Thus H' is a simple planar graph with helper vertices of minimum degree four.

Therefore, either H contains a good vertex of degree at most three or $\delta(H) \geq 4$. If the former happens, then we are done. If the latter happens, then by Theorem 3.10, H contains an edge xy such that $\deg(x) + \deg(y) \leq 11$. Without loss of generality, let $\deg(x) \leq \deg(y)$. If x is a good vertex, we are done. If x is a helper vertex, then y has to be a good vertex as helper vertices are independent by Lemma 3.2.

Note that, in case every good vertex in H' has a degree at least seven, by the above argument, there must exist a good vertex y of degree seven with a neighbor x which is a helper vertex of degree four. \square

3.6 Analysis of the structure of H

Let v be the good vertex in H having a degree at most seven. Such a vertex exists due to Lemma 3.11. Let $v_1, v_2, v_3, \dots, v_k$ be its neighbors where $k \leq 7$. Let A_i be the set of good neighbors of v_i other than v such that none of the good vertices in A_i is adjacent to v_j for any $j < i$. Let us assume that the indexing of the good neighbors of v is such that the k -tuple $(|A_1|, |A_2|, |A_3|, \dots, |A_k|)$ attains a maximum in the lexicographic ordering. By Lemma 3.8 $|A_i| \leq 2(2n + m) - 1$ for all $i \in \{1, 2, \dots, k\}$. We would like to improve it by showing that, for $1 \leq p < q < r \leq 7$, we have $|A_p| \cup |A_q| \cup |A_r| \leq 2(2n + m)$.

Lemma 3.12. *For $1 \leq p < q < r \leq 7$, we have $|A_p| \cup |A_q| \cup |A_r| \leq 2(2n + m)$.*

Proof. Suppose on the contrary, for some $1 \leq p < q < r \leq 7$, $|A_p| + |A_q| + |A_r| \geq 2(2n + m) + 1$. Without loss of generality, let us assume that the neighbors v_p, v_q and v_r be arranged in an anti-clockwise manner around v . And let the associated set of neighbors of each v_p, v_q and v_r be $A_p = \{p_1, p_2, \dots, p_a\}$, $A_q = \{q_1, q_2, \dots, q_b\}$ and $A_r = \{r_1, r_2, \dots, r_c\}$, respectively. Furthermore, let the vertices in each of these sets A_i to be arranged in an anti-clockwise manner around v_i in increasing order of indices under the fixed embedding of H for all $i \in \{p, q, r\}$. Following the above construction, by our lexicographic ordering and by Lemma 3.8, we have, $2(2n + m) - 1 \geq |A_p| \geq |A_q| \geq |A_r|$. Also $|A_p| + |A_q| + |A_r| \geq 2(2n + m) + 1 \geq 7$ as $(2n + m) \geq 3$. Hence, $|A_p| \geq 3$ and $|A_q| + |A_r| \geq 2$. As $|A_q| + |A_r| \geq 2$, either $|A_q| \geq |A_r| \geq 1$ or $|A_q| \geq 2$ and $|A_r| \geq 0$. We analyse the cases based on how p in A_p sees q_1 in A_q , where $p = p_1$ if $|A_p| = a = 3$, and $p = p_{a-3}$, if $|A_p| = a \geq 4$. See Fig. 4b for a pictorial representations of the two scenarios.

Case 1: If p sees q_1 by being adjacent to it. Let the region bounded by the cycle $pv_pvv_qq_1p$ containing p_{a-1} be D_1 . Now, q_2 (resp., r_1) can see p_i only via the vertices of D_1 , where $i = \{a - 1, a\}$. However, q_2 (resp., r_1) cannot see p_i via p or v as otherwise a triangle will be formed. Also, q_2 (resp., r_1) cannot see p_i via v_p as v_p cannot be adjacent to q_2 (resp., r_1) due to the definition of A_p . Moreover, q_2 (if it exists) cannot see p_i via q_1 as otherwise a triangle will be formed. Thus, the only possibility for q_2 to see p_{a-1} and p_a is via v_q . If q_2 see both p_{a-1}, p_a via v_q , then v_p, v_q along with their three common good neighbors v, p_{a-1}, p_a will form an \mathbb{F}_3 , which is not possible. Furthermore, r_1 (if it exists) cannot see p_i via v_q due to the definition of A_q . Thus, the only possibility for r_1 to see p_{a-1} and p_a is via q_1 . If r_1 sees both p_{a-1}, p_a via q_1 , then v_p, q_1 along with their three common good neighbors p, p_{a-1}, p_a will form an \mathbb{F}_3 , which is not possible. Therefore, it is not possible for p to see q_1 by being adjacent to it.

Case 2: If p sees q_1 via v_q . Let the region bounded by the cycle pv_pvv_qp containing p_{a-1} be D_2 . Note that q_1 can see p_{a-1} only via the vertices of D_2 . However, q_1 cannot see p_{a-1} via p or v as otherwise a triangle will be formed. Also, q_1 cannot see p_{a-1} via v_p by the definition of A_p . Thus, the only possibility for q_1 to see p_{a-1} is via v_q . If q_1 sees p_{a-1} via v_q , then v_p, v_q along with their three common good neighbors v, p, p_{a-1} will form an \mathbb{F}_3 , which is not possible. Therefore, it is not possible for p to see q_1 via v_q .

Case 3: If p sees q_1 via v_r . Let the region bounded by the cycle $pv_pvv_qq_1v_r p$ containing p_a be D_3 . Note that, q_2 (if it exists) can see p_i only via the vertices of D_3 where $i \in \{a - 1, a\}$. However, q_2 cannot see p_i via p, v or q_1 as otherwise a triangle will be formed. Also, q_2




Figure 4: A pictorial representations of the two scenarios dealt with in the proof of Lemma 3.12.

cannot see p_i via v_p by the definition of A_p . Notice that q_2 cannot see p_i via v_r , as otherwise v_p, v_r along with their three common good neighbors p, p_i, v will form an \mathbb{F}_3 . Thus, the only possibility for q_2 to see p_i is via v_q . If both p_{a-1}, p_a , sees q_2 via v_q , then v_p, v_q along with their three common good neighbors p, p_{a-1}, p_a will form an \mathbb{F}_3 , which is not possible. Thus, q_2 does not exist. Hence, $|A_q| = 1$ and r_1 must exist. Observe that as v_r is adjacent to p, q_1 and r_1 , it contradicts the lexicographic maximality of $(|A_1|, |A_2|, \dots, |A_k|)$. Therefore, it is not possible for p to see q_1 via v_r .

Case 4: If p sees q_1 via a vertex $h \notin \{v, v_p, v_q, v_r\} \cup A_p \cup A_q \cup A_r$. Let the region bounded by the cycle $pv_pvv_qq_1hvp$ containing p_a be D_4 . Now q_2 (resp., r_1) can see p_i only via the vertices of D_4 , where $i \in \{a-1, a\}$ if $a = 3$ and $i \in \{a-2, a-1, a\}$ if $a \geq 4$. However, note that q_2 (resp., r_1) cannot see p_i via p or v as otherwise a triangle will be formed. Also, q_2 (resp., r_1) cannot see p_i via v_p by the definition of A_p . Moreover, q_2 cannot see p_i via q_1 to avoid creating a triangle and r_1 cannot see p_i via v_q by the definition of A_q . Thus, q_2 (if it exists), must see exactly one of p_{a-1}, p_a via v_q and the other via h in order to avoid creating an \mathbb{F}_3 . Similarly, r_1 (if it exists), must see exactly one of p_{a-1}, p_a via q_1 and the other via h in order to avoid creating an \mathbb{F}_3 . Hence, if both q_2 and r_1 existed, a triangle will be created among the vertices v_q, q_1 and one of the p_i s. So, at most one of q_2 or r_1 can exist, which implies that $a \geq 4$. Thus, q_2 (resp., r_1) will see p_{a-2} via v_q (resp., q_1) or h , creating an \mathbb{F}_3 which is not possible. Hence, it is not possible for p to see q_1 via h .

Case 5: If p sees q_1 via r_j for some $j \in \{1, 2, 3, \dots, c\}$. Let the region bounded by the cycle $pv_pvv_r r_j p$ containing p_{a-1} be D_5 . Notice that this bounded region is subdivided

into two regions: (i) D_{51} bounded by the cycle $pv_pvv_qq_1r_jp$ containing p_{a-1} and (ii) D_{52} bounded by the cycle $q_1v_qvv_r r_jq_1$ not containing p_{a-1} .

Suppose there exists some $r \in A_r$ distinct from r_j . If r does not belong to the region D_{52} , and hence does not belong to D_5 , then r can see p_i only via the vertices of D_5 , where $i \in \{a-1, a\}$. However, r cannot see p_i via p, v or r_j , as otherwise a triangle will be formed. Also, r cannot see p_i via v_p by the definition of A_p , and v_r due to the planarity of H . Thus r must be contained in the region D_{52} . Now, r can see p_i only via the vertices of D_{52} . However, we have seen that r cannot see p_i via v, r_j or v_r . Notice that r cannot see both p_{a-1} and p_a via v_q as otherwise, a \mathbb{F}_3 will be formed. Also, r cannot see one of p_{a-1}, p_a via q_1 and the other one via v_q , as otherwise v_q, q_1, r will form a triangle. So the only possibility for r to see p_i s is via q_1 . However, if r sees p_i via q_1 , then q_2 (it exists as $|A_q| \geq |A_r| \geq 2$ in this case) which is in D_{52} can see p_i only via q_1 , which is not possible, as otherwise, a triangle will be formed. Thus, $|A_r| = 1$ and $r_j = r_1$.

If $|A_q| \geq 3$, then q_2 and q_3 belongs to the region D_{52} . Note that, q_j can see p_i only via v_q or r_1 as otherwise, either a triangle will be formed or planarity of H will be violated, where $j = \{2, 3\}$ and $i = \{a-1, a\}$. Also, if both p_{a-1}, p_a are adjacent to v_q (resp., r_1) or if both q_2, q_3 are adjacent to r_1 , an \mathbb{F}_3 will be formed. Thus, there is no possible way for the all the q_j s to see the all the p_i s. Hence, $|A_q| \leq 2$. If $|A_q| = 2$, then $|A_p| \geq 4$. Thus, q_2 will see at least two of p_{a-2}, p_{a-1} and p_a via v_q or r_1 , forming an \mathbb{F}_3 . Hence, $|A_q| = 1$.

If $|A_q| = 1$, then $|A_p| \geq 2(2n+m) - 1 \geq 5$. Also, we will have $p = p_{a-3}$. In this case, p_s must see q_1 via r_1 , as otherwise either a triangle will be created or the definition of A_p will be violated, where $s \in \{1, 2, \dots, a-4\}$. Observe that, this will create an \mathbb{F}_3 unless $(2n+m) = 3$ and $a = 5$. Therefore, for the remainder of the proof let us assume that $(2n+m) = 3$ and $a = 5$.

Observe that q_1 cannot see p_i for $i = \{a-2, a-1, a\}$ via p or v , as otherwise a triangle will be formed. Also, q_1 cannot see p_i via v_p by the definition of A_p or via r_1 as it will create an \mathbb{F}_3 . If p_{a-2} see q_1 via v_q or by being adjacent, then p_{a-1} and p_a cannot see r_1 without creating an \mathbb{F}_3 . Therefore, p_{a-2} must see q_1 via some $h \notin \{v, v_p, v_q, v_r\} \cup A_p \cup A_q \cup A_r$. This will force p_{a-1} and p_a to see r_1 via q_1 .

Now we know that, apart from the good vertices among $\{v, v_p, v_q, v_r\} \cup A_p \cup A_q \cup A_r$, we still have at least ten more good vertices. If such a good vertex belongs to the region D_5 , then it has to be adjacent to q_1 or r_1 for seeing p, p_1 and r_1 . If such a good vertex does not belong to the region D_5 , then also it has to be adjacent to r_1 for seeing q_1 . Thus, by the pigeonhole principle, q_1 or r_1 is adjacent to at least eight good vertices (recall, they are already having three good neighbors in form of p_{a-1}, p_a, r_1 or p_1, p, q_1 , respectively). However, we know due to Lemma 3.8 that a vertex can have at most $2(2n+m)$ good neighbors, which in our case is six as $(2n+m) = 3$. Hence it is not possible for p to see q_1 via r_j for some $j \in \{1, 2, 3, \dots, c\}$.

From all the cases above, we have that p cannot see q_1 , which is a contradiction to our basic assumption. \square

3.7 Concluding the proof

We are now ready to prove the Theorem 1.6.

Proof of Theorem 1.6. Recall that v is a good vertex in H satisfying $k = \deg(v) \leq 7$.

We count the number of possible good neighbors of v . If $k \leq 6$, then by Lemma 3.12 the total number of good vertices in the second neighborhood of v is at most $4(2n + m)$. Moreover, by Lemma 3.11, the total number of good vertices in the closed neighborhood of v is at most seven. Hence, the total number of good vertices in H is,

$$|R| \leq 4(2n + m) + 7 < 2(2n + m)^2 + 2$$

for all $(2n + m) \geq 3$, a contradiction.

If $k = 7$, then by Lemma 3.12, v has at most six good neighbors and at least one neighbor which is a helper of degree four. Thus the total number of good vertices in the second neighborhood of v is at most $4(2n + m) + 3$. Hence, the total number of good vertices in H is,

$$|R| \leq 4(2n + m) + 3 + 7 < 2(2n + m)^2 + 2$$

for all $(2n + m) \geq 4$, a contradiction.

Now, for the case when $(2n + m) = 3$ and when $k = 7$, we give a detailed counting argument as follows. Since $k = 7$, the set of good vertices in the second neighborhood of v is $A_1, A_2, A_3, \dots, A_7$. As $(2n + m) = 3$, we have $|A_i| \leq 2(2n + m) - 1 = 5$ for all $i \in \{1, 2, \dots, 7\}$. Note that, $|A_1| \geq 3$ as by Lemma 3.12, there exists a neighbor of v which is a helper vertex of degree four. Moreover as $|A_1| \geq |A_2| \geq \dots \geq |A_7|$ and $|A_1| + |A_2| + |A_3| \leq 6$, we are left with the following possible cases to deal with.

(i) If $|A_1| = 5$, $|A_2| = 1$, $|A_i| = 0$ for all $i \geq 3$, then the total number of good vertices are at most $|R \cap N[v]| + \sum_{i=1}^7 |A_i| \leq 7 + 6 = 13$.

(ii) If $|A_1| = 4$, $|A_2| = 2$, $|A_i| = 0$ for all $i \geq 3$, then the total number of good vertices are at most $|R \cap N[v]| + \sum_{i=1}^7 |A_i| \leq 7 + 6 = 13$.

(iii) If $|A_1| = 4$ and $|A_i| \leq 1$, for all $i \geq 2$, then the total number of good vertices are at most $|R \cap N[v]| + \sum_{i=1}^7 |A_i| \leq 7 + 10 = 17$.

(iv) If $|A_1| = 3$, $|A_2| = 3$, $|A_i| = 0$ for all $i \geq 3$, then the total number of good vertices are at most $|R \cap N[v]| + \sum_{i=1}^7 |A_i| \leq 7 + 6 = 13$.

(v) If $|A_1| = 3$, $|A_2| \leq 2$, $|A_i| \leq 1$ for all $i \geq 3$, then the total number of good vertices are at most $|R \cap N[v]| + \sum_{i=1}^7 |A_i| \leq 7 + 10 = 17$.

In all the cases above, we see that the number of good vertices is strictly less than $2(2n + m)^2 + 2 = 20$ since $(2n + m) = 3$, a contradiction. \square

4 Conclusions

In the following, we present a table which describes what is known regarding the (n, m) -relative and (n, m) -absolute clique number of the families \mathcal{P}_g of planar graphs having girth at least g . The previous works had successfully found the exact values of $\omega_a(n, m)(\mathcal{P}_g)$ for all $g \geq 4$ and $\omega_r(n, m)(\mathcal{P}_g)$ for all $g \geq 5$. Therefore, finding the exact values were left open only for $\omega_a(n, m)(\mathcal{P}_3)$, $\omega_r(n, m)(\mathcal{P}_3)$ and $\omega_r(n, m)(\mathcal{P}_4)$. In this article, we find the exact value for $\omega_r(n, m)(\mathcal{P}_4)$.

\mathcal{P}_g , for g	$\omega_{a(n,m)}(\mathcal{P}_g)$	$\omega_r(n, m)(\mathcal{P}_g)$
$= 3$	$[3p^2 + p + 1, 9p^2 + 2p + 2]$ [4]	$[3p^2 + p + 1, 42p^2 - 11]$ [9]
$= 4$	$p^2 + 2$ [9]	$2p^2 + 2$ (Theorem 1.6)
$= 5$	$\max(p + 1, 5)$ [9]	$\max(p + 1, 6)$ [9]
$= 6$	$p + 1$ [9]	$\max(p + 1, 4)$ [9]
≥ 7	$p + 1$ [9]	$p + 1$ [9]

Table 1: This is the list of all known lower and upper bounds for $\omega_{a(n,m)}(\mathcal{P}_g)$, $\omega_r(n, m)(\mathcal{P}_g)$ where \mathcal{P}_g denotes the family of planar graphs having girth at least g . Moreover, the list captures the bounds for all $(n, m) \neq (0, 1)$ where $p = 2n + m$.

We know that

$$2(2n + m)^2 \leq \chi_{n,m}(\mathcal{P}_4) \leq 5(2n + m)^4$$

where the lower bound is by Theorem 1.7 and the upper bound follows from the upper bound of the (n, m) -chromatic number of planar graphs due to Nešetřil and Raspaud [33]. As the (n, m) -relative clique number of the triangle-free planar graphs is quadratic in $(2n + m)$, the following may be an interesting question to ask:

Question 4.1. *Does there exist a constant c such that $\chi_{n,m}(\mathcal{P}_4) \leq c(2n + m)^2$ for all $2n + m \geq 3$?*

Acknowledgements: This work is partially supported by IFCAM project “Applications of graph homomorphisms” (MA/IFCAM/18/39), and SERB-MATRICES “Oriented chromatic and clique number of planar graphs” (MTR/2021/000858).

Note: The authors have no relevant financial or non-financial interests to disclose. The manuscript has no associated data. All the authors have contributed equally and their names are listed in the alphabetical order of their last names.

References

- [1] Noga Alon and Timothy H Marshall. Homomorphisms of edge-colored graphs and coxeter groups. *Journal of Algebraic Combinatorics*, 8(1):5–13, 1998.
- [2] Julien Bensmail, Sandip Das, Soumen Nandi, Soumyajit Paul, Théo Pierron, Sagnik Sen, and Eric Sopena. Pushable chromatic number of graphs with degree constraints. *Discrete Mathematics*, 344(1):112151, 2021.

- [3] Julien Bensmail, Sandip Das, Soumen Nandi, Théo Pierron, Sagnik Sen, and Eric Sopena. On the signed chromatic number of some classes of graphs. *Discrete Mathematics*, 345(2):112664, 2022.
- [4] Julien Bensmail, Christopher Duffy, and Sagnik Sen. Analogues of cliques for (m, n) -colored mixed graphs. *Graphs and Combinatorics*, 33(4):735–750, 2017.
- [5] Julien Bensmail, Romaric Duvignau, and Sergey Kirgizov. The complexity of deciding whether a graph admits an orientation with fixed weak diameter. *Discrete Mathematics and Theoretical Computer Science*, 17(3):31–42, 2016.
- [6] Julien Bensmail, Soumen Nandi, Mithun Roy, and Sagnik Sen. Classification of edge-critical underlying absolute planar cliques for signed graphs. *Australasian Journal of Combinatorics*, 77(1):117–135, 2020.
- [7] Julien Bensmail, Soumen Nandi, and Sagnik Sen. On oriented cliques with respect to push operation. *Discrete Applied Mathematics*, 232:50–63, 2017.
- [8] Oleg V Borodin, Dmitry Fon-Der-Flaass, Alexandr V Kostochka, André Raspaud, and Éric Sopena. On deeply critical oriented graphs. *Journal of Combinatorial Theory, Series B*, 81(1):150–155, 2001.
- [9] Dipayan Chakraborty, Sandip Das, Soumen Nandi, Debdeep Roy, and Sagnik Sen. On clique numbers of colored mixed graphs (in press). *Discrete Applied Mathematics*, 2021.
- [10] Sandip Das, Prantar Ghosh, Swathy Prabhu, and Sagnik Sen. Relative clique number of planar signed graphs. *Discrete Applied Mathematics*, 280:86–92, 2020.
- [11] Sandip Das, Soumen Nandi, Sagnik Sen, and Ritesh Seth. The relative signed clique number of planar graphs is 8. In *Algorithms and Discrete Applied Mathematics: 5th International Conference, CALDAM 2019, Kharagpur, India, February 14-16, 2019, Proceedings*, pages 245–253. Springer, 2019.
- [12] Sandip Das, Swathy Prabhu, and Sagnik Sen. A study on oriented relative clique number. *Discrete Mathematics*, 341(7):2049–2057, 2018.
- [13] Soura Sena Das, Soumen Nandi, and Sagnik Sen. The relative oriented clique number of triangle-free planar graphs is 10. In *Conference on Algorithms and Discrete Applied Mathematics*, pages 260–266. Springer, 2020.
- [14] Christopher Duffy, P. D. Pavan, R. B. Sandeep, and Sagnik Sen. On deeply critical oriented cliques. *Journal of Graph Theory*.
- [15] Janusz Dybizbański, Pascal Ochem, Alexandre Pinlou, and Andrzej Szepietowski. Oriented cliques and colorings of graphs with low maximum degree. *Discrete Mathematics*, 343(5):111829, 2020.
- [16] Paul Erdos, Alfréd Rényi, and VT Sós. On a problem of graphs theory. *Studia Sci. Math Hungr. 1*, 1:215–235, 1966.

- [17] R Fabila-Monroy, D Flores, Clemens Huemer, and A Montejano. Lower bounds for the colored mixed chromatic number of some classes of graphs. *Commentationes Mathematicae Universitatis Carolinae*, 49(4):637–645, 2008.
- [18] Zoltán Füredi, Peter Horak, Chandra M Pareek, and Xuding Zhu. Minimal oriented graphs of diameter 2. *Graphs and Combinatorics*, 14(4):345–350, 1998.
- [19] Stanislav Jendrol and H-J Voss. Light subgraphs of graphs embedded in the plane—a survey. *Discrete Mathematics*, 313(4):406–421, 2013.
- [20] Gyula Katona and Endre Szemerédi. On a problem of graph theory. *Studia Scientiarum Mathematicarum Hungarica*, 2:23–28, 1967.
- [21] William F Klostermeyer and Gary MacGillivray. Analogues of cliques for oriented coloring. *Discussiones Mathematicae Graph Theory*, 24(3):373–387, 2004.
- [22] William F Klostermeyer and Gary MacGillivray. Homomorphisms and oriented colorings of equivalence classes of oriented graphs. *Discrete Mathematics*, 274(1-3):161–172, 2004.
- [23] Alexandr V Kostochka, Tomasz Luczak, Gábor Simonyi, and Eric Sopena. On the minimum number of edges giving maximum oriented chromatic number. *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, 49:179–182, 1999.
- [24] Abhiruk Lahiri, Soumen Nandi, Sagnik Sen, and S. Taruni. On chromatic number of (n, m) -graphs. *Extended Abstracts EuroComb 2021*, pages 745–751, 2021.
- [25] TH Marshall. Homomorphism bounds for oriented planar graphs. *Journal of Graph Theory*, 55(3):175–190, 2007.
- [26] TH Marshall. On oriented graphs with certain extension properties. *Ars Combinatoria*, 120:223–236, 2015.
- [27] Amanda Montejano, Pascal Ochem, Alexandre Pinlou, André Raspaud, and Éric Sopena. Homomorphisms of 2-edge-colored graphs. *Discrete Applied Mathematics*, 158(12):1365–1379, 2010.
- [28] Amanda Montejano, Alexandre Pinlou, André Raspaud, and Éric Sopena. Chromatic number of sparse colored mixed planar graphs. *Electronic Notes in Discrete Mathematics*, 34:363–367, 2009.
- [29] Soumen Nandi, Sagnik Sen, and S Taruni. On relative clique number of triangle-free planar colored mixed graphs. In *International Workshop on Combinatorial Algorithms*, pages 439–450. Springer, 2022.
- [30] Ayan Nandy, Sagnik Sen, and Éric Sopena. Outerplanar and planar oriented cliques. *Journal of Graph Theory*, 82(2):165–193, 2016.
- [31] Reza Naserasr, Edita Rollová, and Éric Sopena. Homomorphisms of signed graphs. *Journal of Graph Theory*, 79(3):178–212, 2015.

- [32] Reza Naserasr, Éric Sopena, and Thomas Zaslavsky. Homomorphisms of signed graphs: An update. *European Journal of Combinatorics*, 91:103222, 2021.
- [33] Jaroslav Nešetřil and André Raspaud. Colored homomorphisms of colored mixed graphs. *Journal of Combinatorial Theory, Series B*, 80(1):147–155, 2000.
- [34] Pascal Ochem, Alexandre Pinlou, and Sagnik Sen. Homomorphisms of 2-edge-colored triangle-free planar graphs. *Journal of Graph Theory*, 85(1):258–277, 2017.
- [35] Eric Sopena. The chromatic number of oriented graphs. *Journal of Graph Theory*, 25(3):191–205, 1997.
- [36] Éric Sopena. Oriented graph coloring. *Discrete Mathematics*, 229(1-3):359–369, 2001.
- [37] Éric Sopena. Homomorphisms and colourings of oriented graphs: An updated survey. *Discrete Mathematics*, 339(7):1993–2005, 2016.
- [38] Douglas Brent West. *Introduction to graph theory*, volume 2. Prentice hall Upper Saddle River, 2001.
- [39] Thomas Zaslavsky. Signed graphs. *Discrete Applied Mathematics*, 4(1):47–74, 1982.