DESINGULARIZATION OF SEMIALGEBRAIC SETS. APPLICATIONS

ANTONIO CARBONE AND JOSÉ F. FERNANDO

ABSTRACT. Hironaka's resolution of singularities of algebraic varieties (over a field of characteristic 0) is a widespread celebrated discipline that has many applications in many areas of Mathematics. In this work we approach the desingularization of semialgebraic sets $S \subset \mathbb{R}^n$. To that end, we prove the counterpart to 'Hironaka's desingularization of real algebraic sets' in the semialgebraic setting. In particular, it involves Nash manifolds with corners instead of non-singular real algebraic sets. The obtained results are optimal when S is a closed semialgebraic set. The main tools involved are: Hironaka's desingularization of real algebraic sets, Hironaka's embedded desingularization of real algebraic subsets of non-singular real algebraic sets sets and drilling blow-ups of Nash manifolds with centers closed Nash submanifolds (which were developed in a former work of the second author).

We also show how to 'build' a Nash manifold with corners $\mathcal{Q} \subset \mathbb{R}^n$ from a suitable Nash manifold $M \subset \mathbb{R}^n$ (of its same dimension), which contains \mathcal{Q} as a closed subset, by folding M along the irreducible components of a normal-crossings divisor of M, which is the smallest Nash subset of M that contains the boundary $\partial \mathcal{Q}$ of \mathcal{Q} . In addition, one can choose as the Nash manifold M the Nash 'double' $D(\mathcal{Q})$ of \mathcal{Q} , which is analogous to the Nash double of a Nash manifold with (smooth) boundary, but $D(\mathcal{Q})$ takes into account the peculiarities of the boundary of a Nash manifold with corners.

We include several applications of the previous two results: (1) Weak desingularization of closed semialgebraic sets using Nash manifolds with (smooth) boundary, (2) Representation of compact semialgebraic sets connected by analytic paths as images under Nash maps of closed unit balls, (3) Explicit construction of Nash models for compact orientable smooth surfaces of genus $g \ge 0$, and (4) Nash approximation of continuous semialgebraic maps whose target spaces are Nash manifolds with corners.

1. INTRODUCTION

Hironaka's resolution of singularities [Hi1] on an algebraic variety (over a field of characteristic 0) is a widespread celebrated discipline that has many applications in many areas of Mathematics [Ko, L]. It has also been developed by many other authors in the analytic and subanalytic cases (we refer to Abhyankar [A1, A2, A3], Bierstone-Milman [BM1, BM3, BM2], Villamayor [Vi], Zariski [Z1, Z2, Z3] as an example). For simplicity we restrict to the case when the ground field is \mathbb{R} . The general approach consists of the following: Given a real algebraic set $X \subset \mathbb{R}^n$, one finds a non-singular real algebraic set $X' \subset \mathbb{R}^m$ together with a proper polynomial map $f: X' \to X$ that is a biregular diffeomorphism outside the set of singular points of X. We next recall the precise statement:

Theorem 1.1 (Hironaka's desingularization). Let $X \subset \mathbb{R}^n$ be a real algebraic set. Then there exist a non-singular real algebraic set $X' \subset \mathbb{R}^m$ and a proper polynomial map $f: X' \to X$ such that the restriction $f|_{X'\setminus f^{-1}(\operatorname{Sing}(X))} : X' \setminus f^{-1}(\operatorname{Sing}(X)) \to X \setminus \operatorname{Sing}(X)$ is a diffeomorphism whose inverse map is a regular map.

Date: 13/07/2023.

²⁰¹⁰ Mathematics Subject Classification. Primary: 14P10, 14P20, 14E15; Secondary: 14B05, 57R12, 57Q55.

Key words and phrases. Desingularization, semialgebraic sets, Nash manifolds with corners, folding Nash manifolds, Nash manifolds with boundary, Nash maps and images, semialgebraic sets connected by analytic paths, approximation.

The first author is supported by GNSAGA of INDAM.

The second author is supported by Spanish STRANO PID2021-122752NB-I00 and Grupos UCM 910444. This article has been developed during several one month research stays of the second author in the Dipartimento di Matematica of the Università di Trento. The second author would like to thank the department for the invitations and the very pleasant working conditions.

Remark 1.2. If X is pure dimensional, $X \setminus \text{Sing}(X)$ is dense in X and, as f is proper, it is surjective.

A related important construction (which will also be very useful for our purposes) is Hironaka's embedded resolution of singularities [Hi1], which involves the concept of normal-crossings divisor. Let $X \subset Y \subset \mathbb{R}^n$ be real algebraic sets such that Y is non-singular and has dimension d. Recall that X is a normal-crossings divisor of Y if the irreducible components of X are non-singular, have codimension 1 in Y and constitute a transversal family. More precisely, for each point $x \in X$ there exists a regular system of parameters $\mathbf{x}_1, \ldots, \mathbf{x}_d$ for Y at x such that X is given on an open Zariski neighborhood of x in Y by the equation $\mathbf{x}_1 \cdots \mathbf{x}_k = 0$ for some $k \leq d$.

Theorem 1.3 (Hironaka's embedded desingularization). Let $X \subset Y \subset \mathbb{R}^n$ be real algebraic sets such that Y is non-singular. Then there exists a non-singular real algebraic set $Y' \subset \mathbb{R}^m$ and a proper surjective polynomial map $g: Y' \to Y$ such that $g^{-1}(X)$ is a normal-crossings divisor of Y' and the restriction $g|_{Y'\setminus g^{-1}(X)}: Y'\setminus g^{-1}(X) \to Y\setminus X$ is a diffeomorphism whose inverse map is a regular map.

1.1. Semialgebraic setting. In Real Algebraic Geometry not only algebraic equalities, but also algebraic inequalities appear, and it seems reasonable to explore until what extent it is possible to prove similar results to Hironaka's desingularization (Theorem 1.1) in the context of semialgebraic sets. Recall that a subset $S \subset \mathbb{R}^n$ is *semialgebraic* when it admits a description in terms of a finite boolean combination of polynomial equalities and inequalities, which we will call a semialgebraic description. The category of semialgebraic sets is closed under basic boolean operations but also under usual topological operations: taking closures (denoted by $Cl(\cdot))$, interiors (denoted by $Int(\cdot))$, connected components, etc. If $S \subset \mathbb{R}^m$ and $\mathfrak{T} \subset \mathbb{R}^n$ are semialgebraic sets, a map $f: S \to T$ is *semialgebraic* if its graph is a semialgebraic set. A Nash map on an open semialgebraic set $U \subset \mathbb{R}^n$ is a semialgebraic smooth map $f: U \to \mathbb{R}^m$. Given a semialgebraic set $S \subset \mathbb{R}^n$, a Nash map on S is the restriction to S of a Nash map $F: U \to \mathbb{R}^m$ on an open semialgebraic neighborhood $U \subset \mathbb{R}^n$ of S. A Nash subset $X \subset M$ of a Nash manifold $M \subset \mathbb{R}^n$ (that is, a semialgebraic set that is a smooth submanifold of \mathbb{R}^n) is the zero set of a Nash function $f: M \to \mathbb{R}$, whereas the Nash closure in M of a semialgebraic set $\mathcal{S} \subset M$ is the smallest Nash subset X of M that contains S. In this setting, a Nash normal-crossings divisor of a Nash manifold M is a Nash set $X \subset M$ whose irreducible components are Nash manifolds of codimension 1 in M and constitute a transversal family.

1.2. Desingularization of semialgebraic sets by Nash manifolds with corners. A first step to approach desingularization of semialgebraic sets concerns the choice of a suitable class of 'non-singular semialgebraic domains' to represent any semialgebraic set as the image under a proper polynomial map of one of such domains. As semialgebraic sets $S \subset \mathbb{R}^n$ may have boundaries, which are semialgebraic sets of smaller dimension, we must be careful with the choice of the class of 'non-singular semialgebraic domains'. A lighter approach of the desingularization of semialgebraic sets can be found in [BP] (see Remark 1.5 for a more detailed explanation).

As we would like that the involved maps are polynomial and proper, we find additional rigidity. To lighten the initial approach to the problem, we begin with the resolution of closed semialgebraic sets and we use as 'non-singular semialgebraic domains' Nash manifolds with corners, that is, semialgebraic sets that are smooth submanifolds with corners of \mathbb{R}^n . This choice takes into account not only the fact that a semialgebraic set may have boundary, but also that the Zariski closure of such boundary may have different irreducible components.

Recall that a Nash manifold with corners $\mathcal{Q} \subset \mathbb{R}^n$ is contained as a closed (semialgebraic) subset of a Nash manifold $M \subset \mathbb{R}^n$ of its same dimension [FGR, Prop.1.2]. In fact, we restrict our scope to Nash manifolds $\mathcal{Q} \subset \mathbb{R}^n$ with corners such that the Nash closure in M of the boundary $\partial \mathcal{Q}$ is a (Nash) normal-crossings divisor of M (maybe after shrinking M). Connected Nash manifolds with corners are connected by analytic paths [Fe2, Main Thm.1.4, Lem.8.2, Lem.8.3], so their images under Nash maps (in particular polynomial or regular maps) are connected by analytic paths as a consequence of [Fe2, Main Thm.1.4]. As in Hironaka's results (Theorems 1.1 and 1.3), we would like that each connected component of the 'non-singular semialgebraic domains' covers the largest possible part of the (closed) semialgebraic set $S \subset \mathbb{R}^n$ we want to desingularize. Using the analytic path-connected components of a semialgebraic set $S \subset \mathbb{R}^n$ (Theorem 2.2), we reduce the problem of desingularization to the case of semialgebraic sets connected by analytic paths (Corollary 2.3).

Theorem 1.4 (Desingularization of closed semialgebraic sets). Let $S \subset \mathbb{R}^m$ be a d-dimensional closed semialgebraic set connected by analytic paths and let \overline{S}^{zar} be its Zariski closure. Then there exist:

- (i) A d-dimensional irreducible non-singular real algebraic set $X \subset \mathbb{R}^n$ and a normalcrossings divisor $Y \subset X$.
- (ii) A connected Nash manifold with corners $Q \subset X$ (which is a closed subset of X) whose boundary ∂Q has Y as its Zariski closure.
- (iii) A polynomial map $f : \mathbb{R}^n \to \mathbb{R}^m$ such that the restrictions $f|_X : X \to \overline{S}^{\operatorname{zar}}$ and $f|_{\mathbb{Q}} : \mathbb{Q} \to S$ are proper and $f(\mathbb{Q}) = S$.
- (iv) A closed semialgebraic set $\mathcal{R} \subset S$ of dimension strictly smaller than d such that $S \setminus \mathcal{R}$ and $\Omega \setminus f^{-1}(\mathcal{R})$ are Nash manifolds and the polynomial map $f|_{\Omega \setminus f^{-1}(\mathcal{R})} : \Omega \setminus f^{-1}(\mathcal{R}) \to S \setminus \mathcal{R}$ is a Nash diffeomorphism.

In particular, if \overline{S}^{zar} is compact, also X is compact.

Remark 1.5. Theorem 1.4 is related to a previous result of Bierstone-Parusiński [BP, Rmk.2.6] concerning desingularization of semialgebraic sets (and also of subanalytic sets). However, there are two main differences between their result and ours:

- (1) The 'non-singular (closed) semialgebraic domain' $\operatorname{Cl}(f^{-1}(S \setminus \mathcal{R})) \subset X$ that maps onto the semialgebraic set $S \subset \mathbb{R}^m$ under the polynomial map f does not need to be a Nash manifold with corners in [BP, Rmk.2.6], but only the closure of a union of connected components of $X \setminus Y$.
- (2) If S is connected by analytic paths, $\operatorname{Cl}(f^{-1}(S \setminus \mathcal{R})) \subset X$ does not need to be connected in [BP, Rmk.2.6].

As Nash manifolds with corners are locally compact, their images under proper maps are also locally compact, so Theorem 1.4 cannot be extended directly to general semialgebraic sets. In order to solve this, we will use *Nash quasi-manifolds with corners*, which are essentially Nash manifolds with corners with some 'faces' erased. More precisely, we construct a natural semialgebraic partition $\mathfrak{S}(\mathfrak{Q})$ of any Nash manifold with corners $\mathfrak{Q} \subset \mathbb{R}^n$ that takes into account the structure of its boundary (Definition 4.11). A semialgebraic set $\mathfrak{T} \subset \mathbb{R}^n$ is a *Nash quasimanifold with corners* if $\mathfrak{Q} := \operatorname{Cl}(\mathfrak{T})$ is a Nash manifold with corners and $\mathfrak{Q} \setminus \mathfrak{T}$ is a union of elements of $\mathfrak{S}(\mathfrak{Q})$. In this case, it is necessary to substitute (proper) polynomial maps by (proper) Nash maps. We explain in Example 4.14 why this additional change is mandatory.

Theorem 1.6 (Desingularization of semialgebraic sets). Let $S \subset \mathbb{R}^m$ be a d-dimensional semialgebraic set connected by analytic paths. Then there exist:

- (i) A d-dimensional connected compact (irreducible) non-singular real algebraic set $X \subset \mathbb{R}^n$ and a normal-crossings divisor $Y \subset X$.
- (ii) A connected Nash quasi-manifold with corners S[•] ⊂ X whose closure in X is a compact connected Nash manifold with corners Q[•] ⊂ X and whose boundary ∂Q[•] has Y as its Zariski closure.
- (iii) A Nash map $f : \mathbb{R}^n \to \mathbb{R}^m$ such that the restriction $f|_{\mathbb{S}^{\bullet}} : \mathbb{S}^{\bullet} \to \mathbb{S}$ is proper and $f(\mathbb{S}^{\bullet}) = \mathbb{S}$.
- (iv) A closed semialgebraic set $\mathcal{R} \subset S$ of dimension strictly smaller than d such that $S \setminus \mathcal{R}$ and $S^{\bullet} \setminus f^{-1}(\mathcal{R})$ are Nash manifolds and the Nash map $f|_{S^{\bullet} \setminus f^{-1}(\mathcal{R})} : S^{\bullet} \setminus f^{-1}(\mathcal{R}) \to S \setminus \mathcal{R}$ is a Nash diffeomorphism.

1.2.1. Drilling blow-up. Apart from Hironaka's desingularization (Theorems 1.1 and 1.3), a main tool to prove Theorems 1.4 and 1.6 is the drilling blow-up of a Nash manifold M with center a closed Nash submanifold N, see [Fe2]. We refer the reader to [S, Hi2] for the oriented blow-up of

a real analytic space with center a closed subspace, which is the counterpart of the construction in [Fe2] for the real analytic setting. In [HPV, §5] appears a presentation of the oriented blow-up in the analytic case closer to the drilling blow-up described in [Fe2]. There the authors consider the case of the oriented blow-up of a real analytic manifold M with center a closed real analytic submanifold N whose vanishing ideal inside M is finitely generated (this happens for instance if N is compact). In [Fe1, §3] a similar construction is presented in the semialgebraic setting, which is used to 'appropriately embed' semialgebraic sets in affine spaces. In [FFQU, §3] oriented blow-ups appear to compare regoluous images of \mathbb{R}^2 with regular images of \mathbb{R}^2 . In Section 3 we recall the main properties of drilling blow-up developed in [Fe2] and we improve some of them in order to prove Theorems 1.4 and 1.6.

1.3. Folding Nash manifolds to build Nash manifolds with corners. Recall that a Nash manifold $\Omega \subset \mathbb{R}^n$ with corners is contained as a closed (semialgebraic) subset of a Nash manifold $M \subset \mathbb{R}^n$ of its same dimension and the Nash closure of its boundary $\partial \Omega$ is a Nash normal-crossings divisor of M. Our next result shows that the Nash manifold M can be 'folded' to reconstruct the Nash manifold with corners Ω . Namely,

Theorem 1.7 (Folding Nash manifolds). Let $Q \subset \mathbb{R}^n$ be a d-dimensional Nash manifold with corners. Then there exist

- (i) A d-dimensional Nash manifold $M \subset \mathbb{R}^n$ that contains \mathfrak{Q} as a closed subset.
- (ii) A Nash normal-crossings divisor $Y \subset M$ that is the Nash closure of $\partial \Omega$ in M and satisfies $\Omega \cap Y = \partial \Omega$.
- (iii) A proper Nash map f: Cl(M) → Q such that f|_Q: Q → Q is a semialgebraic homeomorphism close to the identity map and f|_{Int(Q)}: Int(Q) → Int(Q) is a Nash diffeomorphism.

In addition, for each $x \in \partial \Omega$ there exist open semialgebraic neighborhoods $U, V \subset M$ of x equipped with Nash diffeomorphisms $\varphi: U \to \mathbb{R}^d$ and $\psi: V \to \mathbb{R}^d$ and $1 \leq s \leq d$ such that

 $\psi \circ f \circ \varphi^{-1} : \mathbb{R}^d \to \mathbb{R}^d, \ (x_1, \dots, x_d) \mapsto (x_1^2, \dots, x_s^2, x_{s+1}, \dots, x_d).$

In addition, we construct for Nash manifolds with corners $\mathcal{Q} \subset \mathbb{R}^n$ the counterpart of the Nash double of a Nash manifold with (smooth) boundary and such Nash double $D(\mathcal{Q})$ is Nash diffeomorphic around \mathcal{Q} to the Nash manifold M provided by Theorem 1.7 (Theorem 5.13). As a straightforward consequence of Theorems 1.4 and 5.13 and Lemma 2.8, the reader can prove the following:

Corollary 1.8 (Nash ramified covering map). Let $S \subset \mathbb{R}^m$ be a d-dimensional closed semialgebraic set connected by analytic paths. Then there exist:

- (i) A d-dimensional irreducible non-singular real algebraic set $X \subset \mathbb{R}^n$.
- (ii) A polynomial map $f : \mathbb{R}^n \to \mathbb{R}^m$ such that the restriction $f|_C : C \to S$ is proper and f(C) = S for each connected component C of X.
- (iii) A closed semialgebraic set $\mathcal{R} \subset S$ of dimension strictly smaller than d such that $S \setminus \mathcal{R}$ and $X \setminus f^{-1}(\mathcal{R})$ are Nash manifolds and the polynomial map $f|_{X \setminus f^{-1}(\mathcal{R})} : X \setminus f^{-1}(\mathcal{R}) \to S \setminus \mathcal{R}$ is a Nash covering map whose fibers have constant cardinality.

1.4. Some further applications. We next propose some further applications of both techniques proposed above (desingularization of closed semialgebraic sets (connected by analytic paths) and folding of Nash manifolds to construct Nash manifolds with corners).

1.4.1. Weak desingularization of closed semialgebraic sets by Nash manifolds with boundary. A similar result to Theorem 1.4 changing the Nash manifold with corners Ω by a Nash manifold with boundary \mathcal{H} (that is, a semialgebraic set that is a smooth submanifold with (smooth) boundary of \mathbb{R}^n) seems difficult to be obtained (even in the compact case) if we want to keep that the map f is polynomial and that the semialgebraic set \mathcal{R} has dimension strictly smaller than the dimension of \mathcal{S} . We propose the following statement, whose proof is based on the folding of Nash manifolds (Theorem 1.7):

Theorem 1.9 (Weak desingularization of closed semialgebraic sets). Let $S \subset \mathbb{R}^n$ be a ddimensional closed semialgebraic set connected by analytic paths and let $\varepsilon > 0$ be a strictly positive continuous semialgebraic function on S. Then there exist:

- (i) A d-dimensional irreducible non-singular real algebraic set $X \subset \mathbb{R}^m$.
- (ii) A Nash manifold with boundary $\mathfrak{H}_{\varepsilon} \subset \mathbb{R}^m$ such that the Zariski closure Z_{ε} of $\partial \mathfrak{H}_{\varepsilon}$ is a non-singular real algebraic set $Z_{\varepsilon} \subset X$ of dimension d-1 and $\operatorname{Int}(\mathfrak{H}_{\varepsilon})$ is a connected component of $X \setminus Z_{\varepsilon}$.
- (iii) A proper Nash map $f : \mathcal{H}_{\varepsilon} \to S$ such that $f(\mathcal{H}_{\varepsilon}) = S$.
- (iv) The restriction $f|_{\mathcal{H}_{\varepsilon} \setminus f^{-1}(\mathcal{T}_{\varepsilon})} : \mathcal{H}_{\varepsilon} \setminus f^{-1}(\mathcal{T}_{\varepsilon}) \to \mathcal{S} \setminus \mathcal{T}_{\varepsilon}$ is a Nash diffeomorphism, where $\mathcal{T}_{\varepsilon} := \{x \in \mathcal{S} : \operatorname{dist}(x, \mathcal{R}) \leq \varepsilon\}$ for a certain closed semialgebraic set $\mathcal{R} \subset \mathcal{S}$ of dimension strictly smaller than d.

If S is compact, we may assume in addition: ε is constant, X and $\mathcal{H}_{\varepsilon}$ are also compact and X is connected.

Related to the previous result in [Fe2, Proof of Thm.1.4, §8.C.12] it is shown that if $S \subset \mathbb{R}^n$ is a semialgebraic set connected by analytic paths, then there exists a Nash manifold \mathcal{H} with smooth boundary and a surjective Nash map $f : \mathcal{H} \to S$. However, the Nash map provided there is far from being proper even if S is a closed semialgebraic set.

1.4.2. Nash images of the closed unit ball. In Section 6.2 we provide an alternative proof of the following result proposed in [CF].

Theorem 1.10 (Compact Nash images). Let $S \subset \mathbb{R}^n$ be a d-dimensional compact semialgebraic set. The following assertions are equivalent:

- (i) There exists a Nash map $f : \mathbb{R}^d \to \mathbb{R}^n$ such that $f(\overline{\mathcal{B}}_d) = S$.
- (ii) S is connected by analytic paths.

A related (and much more previous in time) question is to determine until what extent we can represent semialgebraic sets connected by analytic paths using polynomial maps in the vein of Gamboa's problem proposed in the 1990 *Oberwolfach reelle algebraische Geometrie* week [G] (see also [E, §3.IV, p.69] and [FGU] for a survey concerning this problem):

Problem 1.11 (Gamboa's problem). To characterize the (semialgebraic) subsets of \mathbb{R}^n that are either polynomial or regular images of \mathbb{R}^m .

The reader can confront Theorem 1.10 with partial results [FU, Thm.1.4 & Thm.1.6] concerning polynomial images of the closed unit balls. Polynomial images of models connected by polynomial paths (affine spaces, closed balls etc.) are connected by polynomial paths, which is a stronger condition than being connected by analytic paths. In general, semialgebraic sets do not contain rational paths and consequently not even polynomial paths. By [C, V] a generic complex hypersurface Z of \mathbb{CP}^m of degree $d \ge 2m - 2$ for $m \ge 4$ and of degree $d \ge 2m - 1$ for m = 2, 3 does not contain rational curves. If $S \subset \mathbb{R}^m$ is a semialgebraic set whose Zariski closure in \mathbb{RP}^m is a generic hypersurface of high enough degree, then its Zariski closure Z in \mathbb{CP}^m does not contain rational curves, so S cannot contain rational paths. This means in particular that general semialgebraic sets (even if they are connected by analytic paths) do not contain polynomial paths, so they are not polynomial images of either affine spaces or closed balls. In view of Theorem 1.4 and Corollary 1.8 the obstruction to solve Problem 1.11 mainly concentrates on non-singular real algebraic sets, which in general do not contain polynomial paths.

1.4.3. *Explicit Nash models for orientable compact smooth surfaces.* Theorem 1.7 provides an explicit procedure to construct Nash models for orientable compact smooth surfaces, see §6.3.

1.4.4. Approximation of continuous semialgebraic maps by Nash maps. Approximation of maps of certain type by maps of a better subtype is an important tool in Mathematics. A celebrated example is Whitney's approximation theorem [Wh]: a continuous map (defined on a locally compact subset of an affine space) whose target space is a C^r submanifold M of \mathbb{R}^n for $r \in$

 $\mathbb{N} \cup \{\infty, \omega\}$ can be approximated by a \mathcal{C}^r map whose target space is M. Whitney approximation theorem has been extended in many directions, like the case of manifolds with boundary (using partitions of unit and collars, [ORR, Ch. III, Thm. 6.1]). New results on approximation in Whitney's style have been proved in [FGh2] when the target space has 'singularities' and admits 'nice' triangulations. As a consequence: every continuous map between a locally compact subset $X \subset \mathbb{R}^m$ and a smooth manifold with corners $\mathbb{Q} \subset \mathbb{R}^n$ can be approximated by a smooth map with respecto to the \mathcal{C}^0 topology.

There are also relevant results concerning approximation in the semialgebraic setting. Efroymson [Ef] showed that every continuous semialgebraic function on a Nash manifold can be approximated by Nash functions. Shiota [Sh] improved this result in many directions. He proved relative and absolute versions with (strong) control on finitely many derivatives of the approximation map. Approximation results also appear when the target space has singularities. In [BFR, Thm.1.7] the target space is a Nash set with monomial singularities and approximation of continuous semialgebraic maps by Nash maps is proved under certain restrictions. In [FGh1] it is presented differentiable approximation of continuous semialgebraic maps when the target space admits 'nice' triangulations. As a consequence: every continuous semialgebraic map between a compact semialgebraic set $X \subset \mathbb{R}^m$ and a Nash manifold with corners $\Omega \subset \mathbb{R}^n$, can be approximated by a \mathcal{C}^r semialgebraic map with respect to the \mathcal{C}^0 semialgebraic topology.

The techniques developed in [FGh1] make an essential use of C^r semialgebraic partitions of unity, so the proofs proposed there do not extend to Nash approximation. However, as an additional application of Theorem 1.7, we prove approximation of continuous semialgebraic maps by Nash maps when the target space is a Nash manifold with corners.

Theorem 1.12 (Nash approximation). Let $S \subset \mathbb{R}^m$ be a locally compact semialgebraic set, $\Omega \subset \mathbb{R}^n$ a Nash manifold with corners and $h : S \to \Omega$ a continuous semialgebraic map. Then there exist Nash maps $g : S \to \Omega$ arbitrarily close to h with respect to the C^0 semialgebraic topology.

1.5. Structure of the article. The article is organized as follows. In Section 2 we recall the concepts and main properties of regular and smooth points of a semialgebraic set. We also analyze the main properties of Nash manifolds with corners and we refer the reader to [FGR] for further details. In this section also the concept of analytic path-connected components of a semialgebraic set appears and we recall the existence and uniqueness of such components. In addition, we show that each connected Nash manifold can be understood as a connected component of an irreducible non-singular algebraic set whose connected components are pairwise Nash diffeomorphic. We also prove in this section Theorems 1.4 and 1.6 for the 1-dimensional case. In Section 3 we recall the main properties of the drilling blow-up of a Nash manifold along a closed Nash submanifold proposed in [Fe2, §3]. We also present some additional new properties of the drilling blow-up that we need to prove Theorems 1.4 and 1.6. In Section 4 we prove Theorem 1.4 and 1.6 (for semialgebraic sets of dimension $d \ge 2$), whereas in Section 5 we prove Theorem 1.7 and we construct the Nash double of a Nash manifold with corners (Theorem 5.13). Section 6 is devoted to prove Theorems 1.9, 1.10 and 1.12. We also present an explicit procedure to construct Nash models for orientable compact smooth surfaces.

1.6. Acknowledgements. The authors are very grateful to E. Bierstone for useful comments and to S. Schramm for a careful reading of the final version and for the suggestions to refine its redaction.

2. Preliminary results

In this section we collect some preliminary concepts and results that will be used freely along this article. We include them for the sake of completeness and to ease the reading of the article. We refer the reader to [Fe2, FGR] for further details.

2.1. Regular points versus smooth points. The set $\operatorname{Reg}(S)$ of regular points of a semialgebraic set $S \subset \mathbb{R}^n$ is defined as follows. Let X be the Zariski closure of S in \mathbb{R}^n and \widetilde{X} the complexification of X, that is, the smallest complex algebraic subset of \mathbb{C}^n that contains X. Define $\operatorname{Reg}(X) := X \setminus \operatorname{Sing}(\widetilde{X})$ and let $\operatorname{Reg}(S)$ be the interior of $S \setminus \operatorname{Sing}(\widetilde{X})$ in $\operatorname{Reg}(X)$. Observe that $\operatorname{Reg}(S)$ is a finite union of disjoint Nash manifolds maybe of different dimensions. We refer the reader to [Fe2, §2.A] for further details concerning the set of regular points of a semialgebraic set.

A point $x \in S$ is *smooth* if there exists an open neighborhood $U \subset \mathbb{R}^n$ of x such that $U \cap S$ is a Nash manifold. It holds that each regular point is a smooth point, but the converse is not always true even if S = X is a real algebraic set [Fe2, Ex.2.1]. The set Sth(S) of smooth points of a semialgebraic set $S \subset \mathbb{R}^n$ is by [St] a semialgebraic subset of \mathbb{R}^n (and consequently a union of Nash submanifolds of \mathbb{R}^n possibly of different dimension), which contains Reg(S) (maybe as a proper subset as it happens in [Fe2, Ex.2.1]), and it is open in S. The set of points of dimension k of Sth(S) is either the empty-set or a Nash manifold of dimension k for each $k = 0, 1, \ldots, d$. In particular, if S is pure dimensional, Sth(S) is a Nash submanifold of \mathbb{R}^n . If X is a real algebraic set, Sing(X) is always an algebraic subset of X whereas the set $X \setminus Sth(X)$ of non-smooth points is in general only a semialgebraic subset of X (see [Fe2, Ex.2.1]).

If X is a non-singular real algebraic set of dimension d, then $\operatorname{Reg}(X) = \operatorname{Sth}(X)$. Thus, if $S \subset X$ is a pure dimensional semialgebraic set of dimension d, we have $\operatorname{Reg}(S) = \operatorname{Sth}(S)$, because the Zariski closure $\overline{S}^{\operatorname{zar}}$ of S is a union of irreducible components of X, so it is a non-singular real algebraic set of dimension d.

2.2. Analytic path-connected components of a semialgebraic set. We recall next the concept of analytic path-connected components of a semialgebraic set introduced in [Fe2, §9].

Definition 2.1. A semialgebraic set $S \subset \mathbb{R}^n$ admits a decomposition into *analytic path-connected* components if there exist semialgebraic sets $S_1, \ldots, S_r \subset S$ such that:

- (i) Each S_i is connected by analytic paths.
- (ii) If $\mathcal{T} \subset S$ is a semialgebraic set connected by analytic paths that contains S_i , then $S_i = \mathcal{T}$.
- (iii) $S_i \not\subset \bigcup_{j \neq i} S_j$.
- (iv) $S = \bigcup_{i=1}^{r} S_i$.

In [Fe2, Thm.9.2] the existence and uniqueness of the analytic path-connected components of a semialgebraic set is shown.

Theorem 2.2 ([Fe2, Thm.9.2]). Let $S \subset \mathbb{R}^n$ be a semialgebraic set. Then S admits a decomposition into analytic path-connected components and this decomposition is unique. In addition, the analytic path-connected components of a semialgebraic set are closed in S.

Corollary 2.3. Let $S \subset \mathbb{R}^n$ be a semialgebraic set and let $\{S_i\}_{i=1}^r$ be the analytic path-connected components of S. Consider the semialgebraic set $S^* := \bigcup_{i=1}^r S_i \times \{i\} \subset \mathbb{R}^{n+1}$ and the projection $\pi : \mathbb{R}^{n+1} \to \mathbb{R}^n (x, x_n) \mapsto x$. Then the restriction $\pi|_{S^*} : S^* \to S$ is a surjective proper projection such that $\pi(S_i \times \{i\}) = S_i$ for i = 1, ..., r.

2.3. Nash manifolds with corners. Let $\Omega \subset \mathbb{R}^n$ be a Nash manifold with corners. The set of *internal points* of a Nash manifold with corners Ω is $\operatorname{Int}(\Omega) := \operatorname{Sth}(\Omega)$. The *boundary* $\partial \Omega$ of Ω is $\partial \Omega := \Omega \setminus \operatorname{Int}(\Omega) = \Omega \setminus \operatorname{Sth}(\Omega)$. If $\overline{\Omega}^{\operatorname{zar}}$ is a non-singular real algebraic set, then $\operatorname{Reg}(\Omega) = \operatorname{Sth}(\Omega) = \operatorname{Int}(\Omega)$ and $\partial \Omega = \Omega \setminus \operatorname{Reg}(\Omega)$. Otherwise, $\partial \Omega \subset \Omega \setminus \operatorname{Reg}(\Omega)$ and the inclusion could be strict.

A Nash manifold with corners $\mathcal{Q} \subset \mathbb{R}^n$ is locally closed. Consequently, \mathcal{Q} is a closed Nash submanifold with corners of the Nash manifold $\mathbb{R}^n \setminus (\operatorname{Cl}(\mathcal{Q}) \setminus \mathcal{Q})$. In [FGR, Thm.1.11] it is shown that \mathcal{Q} is a closed subset of an affine Nash manifold of its same dimension. Recall that a Nash subset Y of a Nash manifold $M \subset \mathbb{R}^n$ has only Nash normal-crossings in M if for each point $y \in Y$ there exists an open semialgebraic neighborhood $U \subset M$ such that $Y \cap U$ is a Nash normal-crossings divisor of U.

Theorem 2.4 ([FGR, Thm.1.11]). Let $Q \subset \mathbb{R}^n$ be a d-dimensional Nash manifold with corners. There exists a d-dimensional Nash manifold $M \subset \mathbb{R}^n$ that contains Q as a closed subset and satisfies:

- (i) The Nash closure Y of ∂Q in M has only Nash normal-crossings in M and $Q \cap Y = \partial Q$.
- (ii) For every $x \in \partial \Omega$ the smallest analytic germ that contains the germ $\partial \Omega_x$ is Y_x .
- (iii) *M* can be covered by finitely many open semialgebraic subsets U_i (for i = 1, ..., r) equipped with Nash diffeomorphisms $u_i := (u_{i1}, ..., u_{id}) : U_i \to \mathbb{R}^d$ such that:

 $\begin{cases} U_i \subset \operatorname{Int}(\mathcal{Q}) \text{ or } U_i \cap \mathcal{Q} = \emptyset, & \text{ if } U_i \text{ does not meet } \partial \mathcal{Q}, \\ U_i \cap \mathcal{Q} = \{u_{i1} \ge 0, \dots, u_{ik_i} \ge 0\}, & \text{ if } U_i \text{ meets } \partial \mathcal{Q} \text{ (for a suitable } k_i \ge 1). \end{cases}$

The Nash manifold M is called a *Nash envelope of* Q. In general, it is not guaranteed that the Nash closure Y of ∂Q in M is a Nash normal-crossings divisor of M as we show next.




FIGURE 2.1. The teardrop.

Example 2.5. The teardrop $\Omega := \{\mathbf{x} \ge 0, \mathbf{y}^2 \le \mathbf{x}^2 - \mathbf{x}^4\} \subset \mathbb{R}^2$ is a Nash manifold with corners (Figure 2.1). Given any open semialgebraic neighborhood M of Ω in \mathbb{R}^2 the Nash closure of $\partial \Omega$ in M is not a Nash normal-crossings divisor.

We define now Nash manifolds with divisorial corners.

Definition 2.6. A Nash manifold with corners $\Omega \subset \mathbb{R}^n$ is a Nash manifold with *divisorial* corners if there exists a Nash envelope $M \subset \mathbb{R}^n$ such that the Nash closure of $\partial \Omega$ in M is a Nash normal-crossings divisor.

A facet of a Nash manifold with corners $\mathcal{Q} \subset \mathbb{R}^n$ is the (topological) closure in \mathcal{Q} of a connected component of $\mathrm{Sth}(\partial \mathcal{Q})$. As $\partial \mathcal{Q} = \mathcal{Q} \setminus \mathrm{Sth}(\mathcal{Q})$ is semialgebraic, the facets are semialgebraic and finitely many. The non-empty intersections of facets of \mathcal{Q} are the faces of \mathcal{Q} . In [FGR] the following characterization for Nash manifolds with divisorial corners is shown:

Theorem 2.7 ([FGR, Thm.1.12, Cor.6.5]). Let $Q \subset \mathbb{R}^n$ be a d-dimensional Nash manifold with corners. The following assertions are equivalent:

- (i) There exists a Nash envelope $M \subset \mathbb{R}^n$ where the Nash closure of $\partial \Omega$ is a Nash normalcrossings divisor.
- (ii) Every facet \mathfrak{F} of \mathfrak{Q} is contained in a Nash manifold $X \subset \mathbb{R}^n$ of dimension d-1.
- (iii) The number of facets of Ω that contain every given point $x \in \partial \Omega$ coincides with the number of connected components of the germ $Sth(\partial \Omega)_x$.
- (iv) All the facets of Q are Nash manifold with divisorial corners.

If that is the case, the Nash manifold M in (i) can be chosen such that the Nash closure in M of every facet \mathfrak{F} of \mathfrak{Q} meets \mathfrak{Q} exactly along \mathfrak{F} .


Note that properties (iii) and (iv) are intrinsic properties of Ω and do not depend on the Nash envelope M. The faces of a Nash manifold with divisorial corners are again Nash manifolds with divisorial corners. If a Nash envelope $M \subset \mathbb{R}^n$ of Ω satisfies (one of the equivalent) conditions of Theorem 2.7, every open semialgebraic neighborhood $M' \subset M$ of Ω satisfies such conditions. For the rest of this article: A Nash manifold with corners means a Nash manifold with divisorial corners.

2.4. Connected components of a non-singular real algebraic set. The following result allows the reader to understand connected Nash manifolds as connected components of irreducible non-singular real algebraic sets whose connected components are pairwise diffeomorphic.

Lemma 2.8. Let $M \subset \mathbb{R}^n$ be a d-dimensional connected Nash manifold. Then there exists a non-singular irreducible real algebraic set $X \subset \mathbb{R}^{n+k}$ whose connected components are Nash diffeomorphic to M. In addition,

- (i) Each connected component of X projects onto M.
- (ii) Given any two of the connected components of X, there exists an automorphism of X that swaps them.
- (iii) If M is compact, we may assume that X has at most two connected components.

Proof. Consider the Nash map $f : M \hookrightarrow \mathbb{R}^n$, $x \mapsto x$. By Artin-Mazur's description [BCR, Thm.8.4.4] of Nash maps there exist $s \ge 1$ and a non-singular irreducible real algebraic set $Z \subset \mathbb{R}^{2n+s}$ of dimension d, a connected component N of Z and a Nash diffeomorphism $g : M \to N$ such that the following diagram is commutative:



We denote the projection of $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^s$ onto the first space \mathbb{R}^n with π_1 and the projection of $\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^s$ onto the second space \mathbb{R}^n with π_2 . Write m := 2n + s. By [M] applied to N and the union of the remaining connected components of Z there exist finitely many polynomials $P_1, \ldots, P_\ell, Q_1, \ldots, Q_\ell \in \mathbb{R}[\mathbf{x}] := \mathbb{R}[\mathbf{x}_1, \ldots, \mathbf{x}_m]$ such that each Q_j is strictly positive on \mathbb{R}^m and

$$N = Z \cap \Big\{ \sum_{j=1}^{\ell} P_j \sqrt{Q_j} > 0 \Big\}.$$

If M is compact, also N is compact and by [CF, Lem.6.1] there exists a polynomial $P \in \mathbb{R}[x]$ such that $N = Z \cap \{P > 0\}$.

A general N is the projection of the real algebraic set

$$Y := \left\{ (z, y, t) \in Z \times \mathbb{R}^{\ell} \times \mathbb{R} : \left(\sum_{j=1}^{\ell} P_j y_j^2 \right) t^2 - 1 = 0, \ y_j^4 - Q_j = 0 \text{ for } j = 1, \dots, \ell \right\}$$

under $\pi : \mathbb{R}^m \times \mathbb{R}^\ell \times \mathbb{R} \to \mathbb{R}^m$, $(z, y, t) \mapsto z$. If M is compact, N is the projection of the real algebraic set $Y := \{(z, t) \in Z \times \mathbb{R} : P(z)t^2 - 1 = 0\}$ under $\pi : \mathbb{R}^m \times \mathbb{R} \to \mathbb{R}^m$, $(z, t) \mapsto z$. Denote

$$h := \begin{cases} P & \text{if } M \text{ is compact,} \\ \sqrt{\sum_{j=1}^{\ell} P_j \sqrt{Q_j}} & \text{if } M \text{ is non-compact.} \end{cases}$$

Fix $\epsilon := (\epsilon_1, \ldots, \epsilon_\ell, \epsilon_{\ell+1}) \in \{-1, 1\}^{\ell+1}$ (where $\ell = 0$ if N is compact) and let $N_{\epsilon} := Y \cap \{\epsilon_1 y_1 > 0, \ldots, \epsilon_\ell y_\ell > 0, \epsilon_{\ell+1} t > 0\}$. Consider the Nash diffeomorphism

$$\varphi_{\epsilon}: N \to N_{\epsilon}, \ z \mapsto \left(z, \epsilon_1 \sqrt[4]{Q_1(z)}, \dots, \epsilon_{\ell} \sqrt[4]{Q_{\ell}(z)}, \epsilon_{\ell+1} \frac{1}{h(z)}\right)$$

whose inverse map is the restriction of the projection π to N_{ϵ} .

Observe that $\{N_{\epsilon}\}_{\epsilon \in \{-1,1\}^{\ell+1}}$ is the collection of the connected components of Y. As $\pi(N_{\epsilon}) = N$ and using the diagram above, we deduce

$$(\pi_2 \circ \pi)(N_{\epsilon}) = \pi_2(N) = (f \circ \pi_1)(N) = f(M) = M.$$

In addition, each N_{ϵ} is Nash diffeomorphic to M and for $\epsilon \neq \epsilon'$ the polynomial map

$$\phi_{\epsilon,\epsilon'}: \mathbb{R}^m \times \mathbb{R}^\ell \times \mathbb{R} \to \mathbb{R}^m \times \mathbb{R}^\ell \times \mathbb{R}, \ (x, y, t) \mapsto (x, (\epsilon \cdot \epsilon') \cdot (y, t))$$

induces an involution of Y such that $\phi_{\epsilon,\epsilon'}(N_{\epsilon}) = N_{\epsilon'}$. We have denoted

$$(\epsilon \cdot \epsilon') \cdot (y, t) := (\epsilon_1 \epsilon'_1 y_1, \dots, \epsilon_\ell \epsilon'_\ell y_\ell, \epsilon_{\ell+1} \epsilon'_{\ell+1} t)$$

As Z is non-singular, also Y is non-singular. Let X be the irreducible component of Y that contains $N_{(1,...,1)}$, which has at most two connected components if M is compact. Then $k := m + \ell + 1 - n$ and the non-singular real algebraic set X satisfies the requirements in the statement. \Box

2.5. The 1-dimensional case. Let us prove here Theorems 1.4 and 1.6 when $S \subset \mathbb{R}^n$ is a 1-dimensional semialgebraic set connected by analytic paths. As 1-dimensional semialgebraic sets are locally compact, we can merge both results in one by choosing the hypothesis of Theorem 1.6 and obtaining the thesis of Theorem 1.4.

Proof of Theorems 1.4 and 1.6 for the 1-dimensional case. Following the proof of [Fe2, Prop.1.6] we find a 1-dimensional non-singular real algebraic set $X \subset \mathbb{R}^m$, a proper polynomial map $f: X \to \overline{S}^{\text{zar}}$, and a connected 1-dimensional semialgebraic subset $\mathfrak{T} \subset X$ such that:

- (i) The restriction $f|_{X \setminus f^{-1}(\operatorname{Sing}(\overline{S}^{\operatorname{zar}}))} : X \setminus f^{-1}(\operatorname{Sing}(\overline{S}^{\operatorname{zar}})) \to \overline{S}^{\operatorname{zar}} \setminus \operatorname{Sing}(\overline{S}^{\operatorname{zar}})$ is a biregular diffeomorphism.
- (ii) The restriction $f|_{\mathfrak{T}}: \mathfrak{T} \to \mathfrak{S}$ is proper and surjective.

We have proved in [Fe2, Prop.1.6] that the connected components of X are Nash diffeomorphic either to \mathbb{R} or to the sphere \mathbb{S}^1 . Thus, \mathcal{T} is Nash diffeomorphic to either (0,1), [0,1), [0,1] or \mathbb{S}^1 . In the latter two cases ([0,1] and \mathbb{S}^1), the properties stated in Theorem 1.4 follow readily. In the first two cases, we consider $\mathcal{A} := \operatorname{Cl}(\mathcal{T}) \setminus \mathcal{T}$, which is a finite set with at most two elements. Observe that \mathcal{T} is closed in $X \setminus \mathcal{A}$, which is a Zariski open subset of X. Thus, the 1-dimensional non-singular real algebraic set $X' := \{(x, y) \in X \times \mathbb{R} : yh(x) = 1\}$, where $h \in \mathbb{R}[\mathbf{x}]$ is any polynomial whose zero set is \mathcal{A} , together with the projection $\pi : X \times \mathbb{R} \to X$ (onto the first factor) satisfy that $\pi|_{X'} : X' \to X \setminus \mathcal{A}$ is a biregular diffeomorphism. Substituting f by $f' := f \circ \pi|_{X'}$ the properties stated in Theorem 1.4 are proved straightforwardly, as required. \Box

In Section 4 we prove Theorems 1.4 and 1.6 for semialgebraic sets of dimension $d \ge 2$.

3. Drilling blow-up

In this section we recall the main properties of the drilling blow-up of a Nash manifold with center a closed Nash submanifold [Fe2, §5]. We prove some additional results (Remark 3.4, Fact 3.5, Remark 3.10 an §3.4) that we need in the subsequent sections.

3.1. Local structure of the drilling blow-up. Let $M \subset \mathbb{R}^m$ be a *d*-dimensional Nash manifold and $N \subset M$ a closed *e*-dimensional Nash submanifold. As we are interested in the local structure, assume that there exists a Nash diffeomorphism $u := (u_1, \ldots, u_d) : M \to \mathbb{R}^d$ such that $N = \{u_{e+1} = 0, \ldots, u_d = 0\}$. Denote $\psi := u^{-1} : \mathbb{R}^d \equiv \mathbb{R}^e \times \mathbb{R}^{d-e} \to M$. Let $\zeta_{e+1}, \ldots, \zeta_d : \mathbb{R}^d \to \mathbb{R}^k$ be Nash maps such that the vectors $\zeta_{e+1}(y, 0), \ldots, \zeta_d(y, 0)$ are linearly independent for each $y \in \mathbb{R}^e$. Write $z \in \mathbb{R}^{d-e}$ as $z := (z_{e+1}, \ldots, z_d)$. Consider the Nash maps

$$\varphi : \mathbb{R}^d \equiv \mathbb{R}^e \times \mathbb{R}^{d-e} \to \mathbb{R}^k, \ (y,z) \mapsto \zeta_{e+1}(y,z)z_{e+1} + \dots + \zeta_d(y,z)z_d,$$
$$\varphi : \mathbb{R}^e \times \mathbb{R} \times \mathbb{S}^{d-e-1} \to \mathbb{R}^k, \ (y,\rho,w) \mapsto \zeta_{e+1}(y,\rho w)w_{e+1} + \dots + \zeta_d(y,\rho w)w_d$$

and assume that $\varphi(y, z) = 0$ if and only if z = 0. Consider the (well-defined) Nash map:

$$\Phi: \mathbb{R}^e \times \mathbb{R} \times \mathbb{S}^{d-e-1} \to M \times \mathbb{S}^{k-1}, \ (y, \rho, w) \mapsto \Big(\psi(y, \rho w), \frac{\phi(y, \rho, w)}{\|\phi(y, \rho, w)\|}\Big).$$

Fact 3.1. Fix $\epsilon = \pm$ and denote

$$I_{\epsilon} := \begin{cases} [0, +\infty) & \text{ if } \epsilon = +, \\ (-\infty, 0] & \text{ if } \epsilon = -. \end{cases}$$

The closure \widetilde{M}_{ϵ} in $M \times \mathbb{S}^{k-1}$ of the set

$$\Gamma_{\epsilon} := \left\{ \left(\psi(y, z), \epsilon \frac{\varphi(y, z)}{\|\varphi(y, z)\|} \right) \in M \times \mathbb{S}^{k-1} : z \neq 0 \right\}$$

is a Nash manifold with boundary such that:

- (i) $\widetilde{M}_{\epsilon} \subset \operatorname{im}(\Phi)$.
- (ii) The restriction of Φ to $\mathbb{R}^e \times I_\epsilon \times \mathbb{S}^{d-e-1}$ induces a Nash diffeomorphism between $\mathbb{R}^e \times I_\epsilon \times \mathbb{S}^{d-e-1}$ and \widetilde{M}_ϵ . Consequently, $\partial \widetilde{M}_\epsilon = \Phi(\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1})$ and $\Gamma_\epsilon = \operatorname{Int}(\widetilde{M}_\epsilon) = \operatorname{Int}(\widetilde{M}_\epsilon)$ $\Phi(\mathbb{R}^e \times (I_{\epsilon} \setminus \{0\}) \times \mathbb{S}^{d-e-1}).$

Fact 3.2. Denote $\mathcal{R} := \partial \widetilde{M}_+ = \partial \widetilde{M}_-$ and $\widehat{M} := \widetilde{M}_+ \cup \widetilde{M}_- = \Gamma_+ \sqcup \mathcal{R} \sqcup \Gamma_-$. Then Φ induces a Nash diffeomorphism between $\mathbb{R}^e \times \mathbb{R} \times \mathbb{S}^{d-e-1}$ and \widehat{M} , which is the Nash closure of \widetilde{M}_+ and \widetilde{M}_- in $M \times \mathbb{S}^{k-1}$. In addition, the Nash map $\sigma : M \times \mathbb{S}^{k-1} \to M \times \mathbb{S}^{k-1}$, $(a, b) \to (a, -b)$ induces a Nash involution on \widehat{M} without fixed points such that $\sigma(\widetilde{M}_+) = \widetilde{M}_-$ and $\Phi(y, -\rho, -w) = (\sigma \circ \Phi)(y, \rho, w)$ for each $(y, \rho, w) \in \mathbb{R}^e \times \mathbb{R} \times \mathbb{S}^{d-e-1}$.

Fact 3.3. Consider the projection $\pi: M \times \mathbb{S}^{k-1} \to M$ onto the first factor and denote $\pi_{\epsilon} := \pi|_{\widetilde{M_{\epsilon}}}$ for $\epsilon = \pm$. Then

- (i) π_{ϵ} is proper, $\pi_{\epsilon}(\widetilde{M}_{\epsilon}) = M$ and $\Re = \pi_{\epsilon}^{-1}(N)$.
- (ii) The restriction $\pi_{\epsilon}|_{\Gamma_{\epsilon}} : \Gamma_{\epsilon} \to M \setminus N$ is a Nash diffeomorphism. (iii) For each $q \in N$ it holds $\pi_{\epsilon}^{-1}(q) = \{q\} \times \mathbb{S}_{q}^{d-e-1}$ where \mathbb{S}_{q}^{d-e-1} is the sphere of dimension d-e-1 obtained when intersecting the sphere \mathbb{S}^{k-1} with the linear subspace L_q generated by $(\zeta_{e+1} \circ u)(q), \ldots, (\zeta_d \circ u)(q).$




FIGURE 3.1. Local structure of the drilling blow-up \widetilde{M}_+ of M of center N (figure borrowed from [Fe2, Fig.3]).

Denote $\widehat{\pi} := \pi|_{\widehat{M}}$ and consider the commutative diagram.

As a consequence, we have: The Nash maps π_{ϵ} and $\hat{\pi}$ have local representations

 $(x_1, \ldots, x_d) \mapsto (x_1, \ldots, x_e, x_{e+1}, x_{e+1}x_{e+2}, \ldots, x_{e+1}x_d)$

in an open neighborhood of each point $p \in \mathbb{R}$. In addition, $d\pi_p(T_p\widehat{M}) \not\subset T_{\pi(p)}N$.

Remarks 3.4. Denote $g := u \circ \hat{\pi} \circ \Phi$ and $g_+ := g|_{\mathbb{R}^e \times [0, +\infty) \times \mathbb{S}^{d-e-1}} = u \circ \pi_+ \circ \Phi|_{\mathbb{R}^e \times [0, +\infty) \times \mathbb{S}^{d-e-1}}$. Consider the Nash normal-crossings divisor $Z := \{\mathbf{y}_{e+1} \cdots \mathbf{y}_d = 0\} \subset \mathbb{R}^d$. Consider coordinates (w_{e+1}, \ldots, w_d) in \mathbb{R}^{d-e} and the sphere $\mathbb{S}^{d-e-1} = \{\mathbf{w}_{e+1}^2 + \cdots + \mathbf{w}_d^2 = 1\}$.

(i) Write $Z_k := \{y_k = 0\}$ for $k = e + 1, \dots, d$ and observe that

$$g^{-1}(Z_k) = (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times \mathbb{R} \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_k = 0\})),$$

$$g^{-1}_+(Z_k) = (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_k = 0\}))$$

for $k = e + 1, \ldots, d$. Thus,

$$g^{-1}(Z) = (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup \bigcup_{k=e+1}^d (\mathbb{R}^e \times \mathbb{R} \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_k = 0\}))$$

is a Nash normal-crossings divisor and

$$g_{+}^{-1}(Z) = (\mathbb{R}^{e} \times \{0\} \times \mathbb{S}^{d-e-1}) \cup \bigcup_{k=e+1}^{d} (\mathbb{R}^{e} \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_{k} = 0\})).$$

The Nash closure of $g_{+}^{-1}(Z)$ is $g^{-1}(Z)$.

(ii) Let $\epsilon := (\epsilon_{e+1}, \ldots, \epsilon_d)$ where $\epsilon_k = \pm 1$ and denote $\mathfrak{Q}_{\epsilon} := \{\epsilon_{e+1}\mathfrak{y}_{e+1} \ge 0, \ldots, \epsilon_d\mathfrak{y}_d \ge 0\}$. Write $-\epsilon := (-\epsilon_{e+1}, \ldots, -\epsilon_d)$. We have:

$$\operatorname{Cl}(g_+^{-1}(\mathfrak{Q}_{\epsilon} \setminus Z)) = \mathbb{R}^e \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\epsilon_{e+1}\mathfrak{w}_{e+1} \ge 0, \dots, \epsilon_d\mathfrak{w}_d \ge 0\}).$$

Consequently, $\operatorname{Cl}(g_+^{-1}(\mathfrak{Q}_{\epsilon} \setminus Z)) \cap \operatorname{Cl}(g_+^{-1}(\mathfrak{Q}_{-\epsilon} \setminus Z)) = \emptyset$. More generally, if

 $\epsilon := (\epsilon_{e+1}, \dots, \epsilon_m, \epsilon_{m+1}, \dots, \epsilon_d)$ and $\epsilon' := (\epsilon_{e+1}, \dots, \epsilon_m, -\epsilon_{m+1}, \dots, -\epsilon_d)$

where e < m < d, then

$$\mathfrak{Q}_{\epsilon} \cap \mathfrak{Q}_{\epsilon'} = \{\epsilon_{e+1}\mathfrak{y}_{e+1} \ge 0, \dots, \epsilon_m\mathfrak{y}_m \ge 0, \mathfrak{y}_{m+1} = 0, \dots, \mathfrak{y}_d = 0\},\$$

which has dimension e + (d - e) - (d - m) = m. In addition,

$$\begin{aligned} \operatorname{Cl}(g_{+}^{-1}(\mathfrak{Q}_{\epsilon} \setminus Z)) &\cap \operatorname{Cl}(g_{+}^{-1}(\mathfrak{Q}_{\epsilon'} \setminus Z)) \\ &= \mathbb{R}^{e} \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\epsilon_{e+1} \mathtt{w}_{e+1} \ge 0, \dots, \epsilon_{m} \mathtt{w}_{m} \ge 0, \mathtt{w}_{m+1} = 0, \dots, \mathtt{w}_{d} = 0\}), \end{aligned}$$

which has dimension e + 1 + (d - e - 1 - (d - m)) = m.

(iii) Let Y_1, Y_2 be intersections of dimension e+1 of irreducible components of Z that contain N. We may assume $Y_1 = \{y_{e+1} = 0, \dots, y_{d-1} = 0\}$ and $Y_2 = \{y_{e+1} = 0, \dots, y_{d-2} = 0, y_d = 0\}$, so $Y_1 \cap Y_2 = \{y_{e+1} = 0, \dots, y_d = 0\} = N$. Thus,

$$\begin{split} g^{-1}(Y_1) &= (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times \mathbb{R} \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-1} = 0\})), \\ g^{-1}_+(Y_1) &= (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-1} = 0\})), \\ g^{-1}_-(Y_2) &= (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times \mathbb{R} \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-2} = 0, \mathbf{w}_d = 0\})), \\ g^{-1}_+(Y_2) &= (\mathbb{R}^e \times \{0\} \times \mathbb{S}^{d-e-1}) \cup (\mathbb{R}^e \times [0, +\infty) \times (\mathbb{S}^{d-e-1} \cap \{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-2} = 0, \mathbf{w}_d = 0\})), \\ As the intersection \end{split}$$

$$\mathbb{S}^{d-e-1} \cap {\{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-1} = 0\}} \cap {\{\mathbf{w}_{e+1} = 0, \dots, \mathbf{w}_{d-2} = 0, \mathbf{w}_d = 0\}}$$

is empty, we conclude that the intersection

$$Cl(g^{-1}(Y_1 \setminus N)) \cap Cl(g^{-1}(Y_2 \setminus N)) = (g^{-1}(Y_1) \cap Cl(g^{-1}(Y_1 \setminus N))) \cap (g^{-1}(Y_2) \cap Cl(g^{-1}(Y_2 \setminus N)))$$

of the strict transforms of Y_1, Y_2 under g is also empty. Analogously, the intersection

$$Cl(g_{+}^{-1}(Y_{1} \setminus N)) \cap Cl(g_{+}^{-1}(Y_{2} \setminus N)) = (g_{+}^{-1}(Y_{1}) \cap Cl(g_{+}^{-1}(Y_{1} \setminus N))) \cap (g_{+}^{-1}(Y_{2}) \cap Cl(g_{+}^{-1}(Y_{2} \setminus N)))$$

of the strict transforms of Y_1, Y_2 under g_+ is empty.

Next, we analyze the properties of the local structure of drilling blow-up when N has dimension d-1.

Fact 3.5. Assume N has dimension e = d - 1. The Nash diffeomorphism $u := (u_1, \ldots, u_d) : M \to \mathbb{R}^d$ satisfies $N = \{u_d = 0\}$. Recall that $\psi := u^{-1} : \mathbb{R}^d \equiv \mathbb{R}^{d-1} \times \mathbb{R} \to M, \zeta_d : \mathbb{R}^d \to \mathbb{R}^k$ is a Nash map that does not take the value $0 \in \mathbb{R}^k$ and define the Nash map

$$\Phi: \mathbb{R}^{d-1} \times \mathbb{R} \times \{\pm 1\} \to M \times \mathbb{S}^{k-1}, \ (y, \rho, \pm 1) \mapsto \Big(\psi(y, \pm \rho), \pm \frac{\zeta_d(y, \pm \rho)}{\|\zeta_d(y, \pm \rho)\|}\Big).$$

Fix $\epsilon = \pm$ and denote

$$I_{\epsilon} := \begin{cases} [0, +\infty) & \text{if } \epsilon = +, \\ (-\infty, 0] & \text{if } \epsilon = -. \end{cases}$$

The closure \widetilde{M}_{ϵ} in $M \times \mathbb{S}^{k-1}$ of the set

$$\Gamma_{\epsilon} := \left\{ \left(\psi(y, z), \epsilon \frac{z}{|z|} \frac{\zeta_d(y, z)}{\|\zeta_d(y, z)\|} \right) \in M \times \mathbb{S}^{k-1} : z \neq 0 \right\}$$

is a Nash manifold with boundary such that:

- (i) $\widetilde{M}_{\epsilon} \subset \operatorname{im}(\Phi)$.
- (ii) The restriction of Φ to $\mathbb{R}^{d-1} \times I_{\epsilon} \times \{\pm 1\}$ induces a Nash diffeomorphism between $\mathbb{R}^{d-1} \times I_{\epsilon} \times \{\pm 1\}$ and \widetilde{M}_{ϵ} . Consequently, $\partial \widetilde{M}_{\epsilon} = \Phi(\mathbb{R}^{d-1} \times \{0\} \times \{\pm 1\})$ and $\Gamma_{\epsilon} = \operatorname{Int}(\widetilde{M}_{\epsilon}) = \Phi(\mathbb{R}^{d-1} \times (I_{\epsilon} \setminus \{0\}) \times \{\pm 1\})$.

Denote $\Re := \partial \widetilde{M}_+ = \partial \widetilde{M}_-$ and $\widehat{M} := \widetilde{M}_+ \cup \widetilde{M}_- = \Gamma_+ \sqcup \Re \sqcup \Gamma_-$. Then Φ induces a Nash diffeomorphism between $\mathbb{R}^{d-1} \times \mathbb{R} \times \{\pm 1\}$ and \widehat{M} , which is the Nash closure of \widetilde{M}_+ and \widetilde{M}_- in $M \times \mathbb{S}^{k-1}$. In addition, the Nash map $\sigma : M \times \mathbb{S}^{k-1} \to M \times \mathbb{S}^{k-1}, (a,b) \to (a,-b)$ induces a Nash involution on \widehat{M} without fixed points such that $\sigma(\widetilde{M}_+) = \widetilde{M}_-$ and $\Phi(y,-\rho,\pm 1) = (\sigma \circ \Phi)(y,\rho,\mp 1)$ for each $(y,\rho,\pm 1) \in \mathbb{R}^{d-1} \times \mathbb{R} \times \{\pm 1\}.$

Consider the projection $\pi: M \times \mathbb{S}^{k-1} \to M$ onto the first factor and denote $\pi_{\epsilon} := \pi|_{\widetilde{M}_{\epsilon}}$. Then

- (i) π_{ϵ} is proper, $\pi_{\epsilon}(\widetilde{M}_{\epsilon}) = M$ and $\mathcal{R} = \pi_{\epsilon}^{-1}(N)$.
- (ii) The restriction $\pi_{\epsilon}|_{\Gamma_{\epsilon}}: \Gamma_{\epsilon} \to M \setminus N$ is a Nash diffeomorphism.
- (iii) For each $q \in N$ it holds $\pi_{\epsilon}^{-1}(q) = \{q\} \times \{\pm \frac{\{(\zeta_d \circ u)(q)\}}{\|\{(\zeta_d \circ u)(q)\}\|}\}$, that is, each point $q \in N$ has exactly two preimages under π_{ϵ} .

Denote $\widehat{\pi} := \pi|_{\widehat{M}}$ and consider the commutative diagram.

Consequently: The Nash maps π_{ϵ} and $\widehat{\pi}$ have local representations $(x_1, \ldots, x_d) \mapsto (x_1, \ldots, x_d)$ in an open neighborhood of each point $p \in \mathbb{R}$. In addition, $d\pi_p(T_p\widehat{M}) = T_{\pi(p)}M$.

3.2. Global definition. Let $M \subset \mathbb{R}^m$ be a *d*-dimensional Nash manifold and $N \subset M$ a closed *e*-dimensional Nash submanifold. Let $f_1, \ldots, f_k \in \mathcal{N}(M)$ be a finite system of generators of the ideal $\mathcal{I}(N)$ of Nash functions on M vanishing identically on N. Consider the Nash map

$$F: M \setminus N \to \mathbb{S}^{k-1}, x \mapsto \frac{(f_1(x), \dots, f_k(x))}{\|(f_1(x), \dots, f_k(x))\|}.$$

We have:

Fact 3.6. Fix
$$\epsilon = \pm$$
. The closure M_{ϵ} in $M \times \mathbb{S}^{k-1}$ of the graph
 $\Gamma_{\epsilon} := \{(x, \epsilon F(x)) \in M \times \mathbb{S}^{k-1} : x \in M \setminus N\}$

is a Nash manifold with boundary. Denote $\mathcal{R} := \partial \widetilde{M}_+ = \partial \widetilde{M}_-$ and $\widehat{M} := \widetilde{M}_+ \cup \widetilde{M}_- = \Gamma_+ \sqcup \mathcal{R} \sqcup \Gamma_-$, which is the Nash closure of \widetilde{M}_+ and \widetilde{M}_- in $M \times \mathbb{S}^{k-1}$ if M is connected. In addition, \widehat{M} is a Nash manifold and the Nash map $\sigma: M \times \mathbb{S}^{k-1} \to M \times \mathbb{S}^{k-1}$, $(a, b) \to (a, -b)$ induces a Nash involution on \widehat{M} without fixed points that maps \widehat{M}_+ onto \widehat{M}_- .

Fact 3.7. Consider the projection $\pi: M \times \mathbb{S}^{k-1} \to M$ onto the first factor. Denote $\pi_{\epsilon} := \pi|_{\widetilde{M}}$ for $\epsilon = \pm$ and $\widehat{\pi} := \pi|_{\widehat{M}}$. We have:

- (i) π_{ϵ} is proper, $\pi_{\epsilon}(M_{\epsilon}) = M$ and $\Re = \pi_{\epsilon}^{-1}(N)$.
- (ii) The restriction $\pi_{\epsilon}|_{\Gamma_{\epsilon}} : \Gamma_{\epsilon} \to M \setminus N$ is a Nash diffeomorphism.
- (iii) Consider the Nash map $f := (f_1, \ldots, f_k) : M \to \mathbb{R}^k$ (whose components generate $\mathcal{I}(N)$). Fix $q \in N$ and let E_q be any complementary linear subspace of T_qN in T_qM . Then $\pi_{\epsilon}^{-1}(q) = \{q\} \times \mathbb{S}_q^{d-e-1}$, where \mathbb{S}_q^{d-e-1} denotes the sphere of dimension d-e-1 obtained when intersecting \mathbb{S}^{k-1} with the (d-e)-dimensional linear subspace $d_q f(E_q)$. In case e = d - 1, each $q \in N$ has exactly two preimages under π_{ϵ} .
- (iv) The Nash maps π_{ϵ} and $\hat{\pi}$ have local representations of the type

 $(x_1, \ldots, x_d) \mapsto (x_1, \ldots, x_e, x_{e+1}, x_{e+1}x_{e+2}, \ldots, x_{e+1}x_d)$

in an open neighborhood of each point $p \in \mathcal{R}$. In case e = d - 1 the previous local representations are $(x_1, \ldots, x_d) \mapsto (x_1, \ldots, x_d)$. In addition, $d\pi_p(T_p \widehat{M}) \not\subset T_{\pi(p)} N$ and if e = d - 1, we have $d\pi_p(T_pM) = T_{\pi(p)}M$.

Fact 3.8. Up to Nash diffeomorphisms compatible with the respective projections, the pairs $(M_{\epsilon}, \pi_{\epsilon})$ and $(M, \hat{\pi})$ do not depend on the generators f_1, \ldots, f_k of $\mathcal{I}(N)$. Moreover, such Nash diffeomorphisms are unique.

Definition 3.9. The pair (M_+, π_+) is the *drilling blow-up* of the Nash manifold M with center the closed Nash submanifold $N \subset M$ and $(M, \hat{\pi})$ is the twisted Nash double of (M_+, π_+) .

Remark 3.10. Let $N \subset M' \subset M$ be Nash manifolds such that N, M' are closed in M. Let (\widetilde{M}_+, π_+) be the drilling blow-up of M with center N and let $(\widehat{M}, \widehat{\pi})$ be the twisted Nash double of (\widetilde{M}_+, π_+) . Denote $M'^* := \operatorname{Cl}(\pi_+^{-1}(M' \setminus N))$ and $M'^{\bullet} := \operatorname{Cl}(\widehat{\pi}(M' \setminus N))$. Then $(M'^*, \pi_+|_{M'^*})$ is the drilling blow-up of M' with center N and $(M'^{\bullet}, \widehat{\pi}|_{M'^{\bullet}})$ is the twisted Nash double of $(M'^*, \pi_+|_{M'^*}).$

One only has to observe that a finite system of generators $f_1, \ldots, f_k \in \mathcal{N}(M')$ of the ideal $\mathcal{I}(N)$ can be obtained considering a finite system of generators $g_1, \ldots, g_k \in \mathcal{N}(M)$ of the ideal $\mathcal{I}(N)$ and defining $f_j := g_j|_{M'}$ for $j = 1, \ldots, k$.

3.3. Relationship between drilling blow-up and classical blow-up. Let $M \subset \mathbb{R}^m$ be a d-dimensional Nash manifold and $N \subset M$ a closed e-dimensional Nash submanifold. Let $f_1, \ldots, f_k \in \mathcal{N}(M)$ be a system of generators of the ideal $\mathcal{I}(N)$. Define

$$\Gamma' := \{ (x, (f_1(x) : \ldots : f_k(x))) \in M \times \mathbb{RP}^{k-1} : x \in M \setminus N \}.$$

The closure B(M,N) of Γ' in $M \times \mathbb{RP}^{k-1}$ together with the restriction π' to B(M,N) of the projection $M \times \mathbb{RP}^{k-1} \to M$ is the classical *blow-up* of M with center N (in the Nash setting).

Corollary 3.11. Let $(\widehat{M}, \widehat{\pi})$ be the twisted Nash double of the drilling blow-up (\widetilde{M}_+, π_+) . Let $\sigma: \widehat{M} \to \widehat{M}, (a, b) \mapsto (a, -b)$ be the involution of \widehat{M} without fixed points. Consider the Nash $map \ \Theta: M \times \mathbb{S}^{k-1} \to M \times \mathbb{RP}^{k-1}, (p,q) \to (p,[q]) \ and \ its \ restriction \ \theta := \Theta|_{\widehat{M}} : \widehat{M} \to B(M,N).$ We have:

- (i) $\theta(\widehat{M}) = B(M,N), \ \theta \circ \sigma = \theta, \ \pi' \circ \theta = \widehat{\pi} \ and \ \theta^{-1}(a,[b]) = \{(a,b),(a,-b)\} \ for \ each$ $(a, [b]) \in B(M, N).$
- (ii) θ is an unramified two to one Nash covering of B(M, N).

15

3.4. Algebraic description of drilling blow-up. We analyze a general enough situation for which we can guarantee that the twisted (Nash) double of the drilling blow-up is a non-singular real algebraic set. Let $X \subset \mathbb{R}^n$ be a *d*-dimensional non-singular real algebraic set and $Y \subset X$ a *e*-dimensional non-singular real algebraic subset. Let $f_1, \ldots, f_r \in \mathbb{R}[\mathbf{x}]$ be a system of generators of the ideal $\mathcal{I}(Y)$ of polynomials vanishing identically on Y and denote $u := (u_1, \ldots, u_r) \in \mathbb{S}^{r-1}$. Fix $\epsilon = \pm$ and let \widetilde{X}_{ϵ} be the (topological) closure of

$$\Gamma_{\epsilon} := \left\{ (x, u) \in (X \setminus Y) \times \mathbb{S}^{r-1} : \operatorname{rk} \begin{pmatrix} u_1 & \cdots & u_r \\ f_1(x) & \cdots & f_r(x) \end{pmatrix} = 1, \, \epsilon(u_1 f_1(x) + \cdots + u_r f_r(x)) > 0 \right\}$$

in $X \times \mathbb{S}^{r-1}$. Let $\pi : X \times \mathbb{S}^{r-1} \to X$ be the projection (onto the first factor) and \widehat{X} the (topological) closure of

$$\Gamma := \Big\{ (x, u) \in (X \setminus Y) \times \mathbb{S}^{r-1} : \operatorname{rk} \begin{pmatrix} u_1 & \cdots & u_r \\ f_1(x) & \cdots & f_r(x) \end{pmatrix} = 1 \Big\}.$$

Observe that $\widehat{X} = \widetilde{X}_+ \cup \widetilde{X}_-$ is the union of the irreducible components of the real algebraic set

$$\left\{ (x,u) \in X \times \mathbb{S}^{r-1} : \operatorname{rk} \begin{pmatrix} u_1 & \cdots & u_r \\ f_1(x) & \cdots & f_r(x) \end{pmatrix} = 1 \right\}$$

different from $Y \times \mathbb{S}^{r-1}$. If we denote $\widehat{\pi} := \pi|_{\widehat{X}}$, it holds that $(\widehat{X}, \widehat{\pi})$ is the twisted Nash double of the drilling blow-up (\widetilde{X}_+, π_+) of X with center Y (where $\pi_+ := \pi|_{\widetilde{X}_+}$). Proceeding similarly to the proof of [BCR, Prop.3.5.11(i)], one shows that \widehat{X} is a non-singular real algebraic set. In addition, proceeding similarly to the proof of [BCR, Prop.3.5.11(ii)], one shows that there exists a finite open cover of Y by Zariski open subsets U_i such that $\pi^{-1}(U_i)$ is Nash diffeomorphic to $U_i \times \mathbb{S}^{d-e-1}$ via π (use also Fact 3.7(iii)). It holds $\widetilde{X}_{\epsilon} = \widehat{X} \cap \{\epsilon(u_1f_1 + \cdots + u_rf_r) \ge 0\}$ and the Zariski closure of \widetilde{X}_{ϵ} is contained in \widehat{X} .

If $S \subset X$ is a pure dimensional semialgebraic set of dimension d, then $\operatorname{Reg}(S) = \operatorname{Sth}(S)$ and $\mathcal{T} := \operatorname{Cl}(\pi_+^{-1}(S \setminus Y)) \cap \pi_+^{-1}(S)$ is a pure dimensional semialgebraic set of dimension d such that $\operatorname{Reg}(\mathcal{T}) = \operatorname{Sth}(\mathcal{T})$. This is because both X and \hat{X} are non-singular real algebraic sets of dimension d and $\pi_+^{-1}|_{\tilde{X}_+} : \tilde{X}_+ \setminus \pi_+^{-1}(Y) \to X \setminus Y$ is a Nash diffeomorphism.

Remark 3.12. If X is irreducible, then \widehat{X} does not need to be irreducible (even if X is connected).

Consider for instance the plane $X := \mathbb{R}^2$ and the line $Y := \{\mathbf{x} = 0\}$. Then $\mathcal{I}(Y) = (\mathbf{x})$ and $\widehat{X} = \{\mathbf{z} = -1\} \cup \{\mathbf{z} = 1\} \subset \mathbb{R}^3$, which is disconnected. However, \widehat{X} is the Zariski closure of $\widetilde{X}_+ = \{\mathbf{x} \ge 0, \mathbf{z} = 1\} \cup \{\mathbf{x} \le 0, \mathbf{z} = -1\}$ (resp. $\widetilde{X}_- = \{\mathbf{x} \le 0, \mathbf{z} = 1\} \cup \{\mathbf{x} \ge 0, \mathbf{z} = -1\}$).

However, if X is irreducible, at least we have the following result that we prove below:

Lemma 3.13. If X is irreducible, then \widehat{X} is the Zariski closure of \widetilde{X}_{ϵ} for $\epsilon = \pm 1$.

3.4.1. Relationship with algebraic blow-up. Denote $[z] := [z_1 : \cdots : z_r] \in \mathbb{RP}^{r-1}$ and define

$$\Gamma' := \{ (x, [z]) \in (X \setminus Y) \times \mathbb{RP}^{r-1} : \operatorname{rk} \begin{pmatrix} z_1 & \cdots & z_r \\ f_1(x) & \cdots & f_r(x) \end{pmatrix} = 1 \}.$$

The closure B(X,Y) of Γ' in $X \times \mathbb{RP}^{r-1}$ together with the restriction π' to B(X,Y) of the projection $X \times \mathbb{RP}^{r-1} \to Y$ (onto the first factor) is the (algebraic) *blow-up* of X with center Y. Consider the polynomial map $\Theta : X \times \mathbb{S}^{r-1} \to X \times \mathbb{RP}^{r-1}, (x, z) \to (x, [z])$ and its restriction $\theta := \Theta|_{\widehat{X}} : \widehat{X} \to B(X,Y)$. We have:

(i) $\theta(\widehat{X}) = B(X, Y), \pi' \circ \theta = \widehat{\pi} \text{ and } \theta^{-1}(x, [z]) = \{(x, z), (z, -z)\} \text{ for each } (x, [z]) \in B(X, Y).$ (ii) θ is an unramified two to one polynomial covering of B(X, Y).

Proof of Lemma 3.13. As X is irreducible, T := B(X, Y) is irreducible (because $X \setminus Y$ and $T \setminus \pi'^{-1}(Y)$ are biregularly diffeomorphic via $\pi'|_{T \setminus \pi'^{-1}(Y)}$ [BCR, Prop.3.5.8]).

The polynomial map $\theta: \widehat{X} \to T$ is an unramified two to one polynomial covering. If \widehat{X} is irreducible, \widehat{X} is the Zariski closure of \widetilde{X}_{ϵ} for $\epsilon = \pm 1$, because $\dim(\widehat{X}) = \dim(\widetilde{X}_{\epsilon})$. Thus, let us assume \widehat{X} is reducible. Let Z_1, \ldots, Z_ℓ be irreducible components of $\widehat{X} = \widetilde{X}_+ \cup \widetilde{X}_-$.

As \widehat{X} is pure dimensional and non-singular, we deduce Z_i is pure dimensional and nonsingular, so it is a union of connected components of \widehat{X} . Thus, Z_i is an open and closed subset of \widehat{X} . As θ is an open and closed map and each Z_i is an open and closed subset of \widehat{X} , we deduce $\theta(Z_i)$ is a union of connected components of T.

Let \widehat{X} be the complexification of \widehat{X} , \widetilde{Z}_i the complexification of Z_i and \widetilde{T} the complexification of T. Observe that $\widehat{\widehat{X}} = \bigcup_{i=1}^{\ell} \widetilde{Z}_i$.

Consider the polynomial extension $\tilde{\theta}: \hat{X} \to \tilde{T}$ of θ to \hat{X} . Its image is contained in \tilde{T} , because $\theta(\hat{X}) = T$. Observe that $\tilde{\theta}|_{\tilde{Z}_i}$ is a dominant polynomial map, because θ is a local diffeomorphism and Z_i is a union of connected components of \hat{X} . In particular $\tilde{\theta}(\tilde{Z}_i)$ is a Zariski open subset of the irreducible algebraic set \tilde{T} .

Denote with $\mathcal{R}(\widetilde{Z}_i)$ the field of rational functions on \widetilde{Z}_i and with $\mathcal{R}(\widetilde{T})$ the field of rational functions on \widetilde{T} . The map $\widetilde{\theta}|_{\widetilde{Z}_i}^* : \mathcal{R}(\widetilde{T}) \to \mathcal{R}(\widetilde{Z}_i), f \mapsto f \circ \widetilde{\theta}|_{\widetilde{Z}_i}$ is a homomorphism of fields of the same transcendence degree $d = \dim(X)$ over \mathbb{C} . Consequently, $\mathcal{R}(\widetilde{Z}_i)$ is an algebraic extension of $\mathcal{R}(\widetilde{T})$ of finite degree m_i . By [Ha, Prop.7.16] the number of points in a general fiber of $\widetilde{\theta}|_{\widetilde{Z}_i}$ is equal to m_i . As $(\widetilde{\theta}|_{\widetilde{Z}_i})^{-1}(q) \setminus (\theta|_{Z_i})^{-1}(q)$ is even for each $q \in \theta(Z_i)$, we deduce that the fibers of $\theta|_{Z_i} : Z_i \to \theta(Z_i) \subset T$ all have the same cardinality, which is either one or two. Thus, $\theta|_{Z_i} : Z_i \to \theta(Z_i) \subset T$ is either a polynomial diffeomorphism or an unramified two to one polynomial covering for $i = 1, \ldots, \ell$. We distinguish two cases:

CASE 1. If $\theta|_{Z_i} : Z_i \to \theta(Z_i) \subset T$ is an unramified two to one polynomial covering, then $Z_i \cap \widetilde{X}_{\epsilon} \neq \emptyset$ and Z_i is contained in the Zariski closure of \widetilde{X}_{ϵ} .

CASE 2. If $\theta|_{Z_i} : Z_i \to \theta(Z_i) \subset T$ is a polynomial diffeomorphism, let us prove: $\theta(Z_i) = T$, $\ell = 2$ and $\theta|_{Z_i} : Z_i \to T$ is a polynomial diffeomorphism for i = 1, 2.

Suppose that $\theta(Z_i) \neq T$. Then there exists a connected component M of T such that $\theta(Z_i) \cap M = \emptyset$ (recall that $\theta(Z_i)$ is a union of connected components of T). As T is pure dimensional of dimension d, we deduce that M has dimension d. As $\theta|_{Z_i} : Z_i \to \theta(Z_i)$ is a polynomial diffeomorphism, we deduce that m_i is odd. Pick a point $q \in M$ such that $(\theta|_{Z_i})^{-1}(q)$ has exactly m_i points. As the cardinality of $(\tilde{\theta}|_{\tilde{Z}_i})^{-1}(q) \setminus (\theta|_{Z_i})^{-1}(q)$ is even and m_i is odd, the cardinality of $(\theta|_{Z_i})^{-1}(q) = \emptyset$ is also odd, which is a contradiction. Consequently, $\theta(Z_i) = T$.

As $\theta: \hat{X} \to T$ is an unramified two to one polynomial covering, we deduce that it is trivial, that is, $\ell = 2$ and $\theta|_{Z_i}: Z_i \to T$ is a polynomial diffeomorphism for i = 1, 2.

We deduce either:

- Each restriction $\theta|_{Z_i}: Z_i \to \theta(Z_i) \subset T$ is an unramified two to one polynomial covering for $i = 1, \ldots, \ell$ and the Zariski closure of \widetilde{X}_{ϵ} is $\bigcup_{i=1}^{\ell} Z_i = \widehat{X}$, or
- $\ell = 2$ and $\theta|_{Z_i} : Z_i \to T$ is a polynomial diffeomorphism for i = 1, 2.

Suppose that we are under the hypothesis of the second case and pick a point $y \in (\pi')^{-1}(Y) \subset T$. Observe that $\theta^{-1}(y) \subset \widetilde{X}_+ \cap \widetilde{X}_-$, so $\widetilde{X}_\epsilon \cap Z_i \neq \emptyset$ for $\epsilon = \pm$ and i = 1, 2. As \widetilde{X}_ϵ is pure dimensional of dimension d and each Z_i is pure dimensional of dimension d and open and closed in $\widehat{X} = Z_1 \sqcup Z_2$, we conclude that each Z_i is contained in the Zariski closure of \widetilde{X}_ϵ , so \widehat{X} is the Zariski closure of \widetilde{X}_ϵ for $\epsilon = \pm$, as required.

4. Desingularization of semialgebraic sets

In this section we prove Theorems 1.4 and 1.6. We begin recalling the concept of checkerboard set [Fe2, Thm.8.4].

4.1. Checkerboard sets. If $S \subset \mathbb{R}^m$ is a semialgebraic set, we denote $\partial S := \operatorname{Cl}(S) \setminus \operatorname{Reg}(S)$. As S is closed, we have $\partial S = Sing(S)$. If Q is a Nash manifold with corners such that \overline{Q}^{zar} is a nonsingular real algebraic set, then $Int(\Omega) = Reg(\Omega)$ and the differences $\Omega \setminus Int(\Omega)$ and $\Omega \setminus Reg(\Omega)$ define the same semialgebraic set ∂Q . In other situations, one should be careful with the possible ambiguity in the definition of the set $\partial \Omega$. In this section such ambiguity does not appear because all the Zariski closures of all involved Nash manifolds with corners are non-singular.

Definition 4.1. A pure dimensional semialgebraic set $\mathcal{T} \subset \mathbb{R}^n$ is a *checkerboard set* if it satisfies the following properties:

- T^{zar} is a non-singular real algebraic set.
 ∂T^{zar} is a normal-crossings divisor of T^{zar}.
- $\operatorname{Reg}(\mathfrak{T}) = \operatorname{Sth}(\mathfrak{T})$ is connected.

Recall that any checkerboard set is connected by analytic paths [Fe2, Main Thm.1.4, Lem.8.2]. In [Fe2, Thm.8.4] the following crucial result is proved.

Theorem 4.2 ([Fe2, Thm.8.4]). Let $S \subset \mathbb{R}^m$ be a semialgebraic set of dimension $d \geq 2$ connected by analytic paths and denote $X := \overline{S}^{\text{zar}}$. Then there exists a checkerboard set $\mathfrak{T} \subset \mathbb{R}^n$ of dimension d and a proper regular map $f: Y := \overline{\mathfrak{T}}^{\text{zar}} \to X$ such that $f(\mathfrak{T}) = \mathfrak{S}$.

As the map f is proper, if the semialgebraic set S is compact, we may assume that also the checkerboard set \mathcal{T} is compact (see the proof of [Fe2, Thm.8.4]). Even if it is not explicitly quoted in the statement of [Fe2, Thm.8.4], there is actually proved much more. Looking at this proof, we can reformulate the statement of Theorem 4.2 in the following way:

Theorem 4.3. Let $S \subset \mathbb{R}^m$ be a semialgebraic set connected by analytic paths of dimension Theorem 4.5. Let $\mathfrak{S} \subset \mathbb{R}^n$ be a semiangeonale set connected by analytic pairs of annexity $d \geq 2$ and denote $X := \overline{\mathfrak{S}}^{\operatorname{zar}}$. Then there exists a checkerboard set $\mathfrak{T} \subset \mathbb{R}^n$ of dimension d and a proper polynomial map $f : Y := \overline{\mathfrak{T}}^{\operatorname{zar}} \to X$ such that $f(\mathfrak{T}) = \mathfrak{S}$ and the restriction $f|_{\mathfrak{T}}: \mathfrak{T} \to \mathbb{S}$ is also proper. Moreover, there exists a semialgebraic set $\mathfrak{R} \subset \mathbb{S}$ of dimension strictly smaller than d such that $f^{-1}(\mathfrak{R}) \subset \overline{\partial \mathfrak{T}}^{\operatorname{zar}}$, $\mathbb{S} \setminus \mathfrak{R}$ and $\mathfrak{T} \setminus f^{-1}(\mathfrak{R})$ are Nash manifolds and $f|_{\mathfrak{T}\setminus f^{-1}(\mathfrak{R})}:\mathfrak{T}\setminus f^{-1}(\mathfrak{R})\to \mathfrak{S}\setminus\mathfrak{R}$ is a Nash diffeomorphism. In particular, if X is compact, also Y is compact.

4.2. Desingularization of closed checkerboard sets. In order to prove Theorem 1.4, in view of Theorem 4.3 we are reduced to assume that $\mathcal{T} \subset \mathbb{R}^n$ is a closed checkerboard set. We need some preliminary notations and results on checkerboard sets.

4.2.1. Definition and properties of the function e of a checkerboard set. Given a non-singular real algebraic set $X \subset \mathbb{R}^n$ of dimension d and a normal-crossings divisor $Z \subset X$, we denote

$$\begin{aligned} \operatorname{Sing}_0(Z) &:= Z, \\ \operatorname{Sing}_\ell(Z) &:= \operatorname{Sing}(\operatorname{Sing}_{\ell-1}(Z)) \quad \text{for } 1 \leq \ell \leq d. \end{aligned}$$

Observe that if $\operatorname{Sing}_{\ell}(Z) \neq \emptyset$, then $\dim(\operatorname{Sing}_{\ell}(Z)) = d - \ell - 1$. In addition, if $\operatorname{Sing}_{\ell}(Z) = \emptyset$, then $\operatorname{Sing}_k(Z) = \emptyset$ for $k \ge \ell$. In particular, $\operatorname{Sing}_d(Z) = \emptyset$. Recall that if $A_x \subset \mathbb{R}^n_x$ is a set germ at x, its analytic closure $\overline{A}^{\operatorname{an}}_x$ is the smallest analytic set germ of \mathbb{R}^n_x that contains A_x .

Let $\mathcal{T} \subset \mathbb{R}^n$ be a closed checkerboard set and denote $X := \overline{\mathcal{T}}^{\operatorname{zar}}$ and $\partial \mathcal{T} := \mathcal{T} \setminus \operatorname{Reg}(\mathcal{T})$. For each point $x \in \mathcal{T}$ there exists a coordinate system $(\mathbf{u}_1, \ldots, \mathbf{u}_d)$ of the Nash manifold X at x and an integer $0 \le r_x \le d$ such that either $\overline{\partial \mathfrak{T}_x^{\mathrm{an}}} = {\mathbf{u}_1 \cdots \mathbf{u}_{r_x}} = 0}_x$ if $r_x \ge 1$ or $x \in \operatorname{Reg}(\mathfrak{T}) = \mathfrak{T} \setminus \partial \mathfrak{T}$ if $r_x = 0$. We denote with $e_x := e_x(\mathfrak{T}) \le r_x$ the number of indices $1 \le i \le r_x$ such that the germ $\mathfrak{T}_x \setminus {\mathbf{u}_i = 0}_x$ is disconnected. If $r_x \le 1$, then $e_x = 0$. We have the following:

Lemma 4.4. The value $e_x = 0$ if and only if T_x is the germ at x of a Nash manifold with corners.

Proof. The if implication is clear because after changing the sign of some of the variables if necessary, we may assume either $x \in \text{Reg}(\mathcal{T})$ or $\mathcal{T}_x = \{u_1 \ge 0, \dots, u_{r_x} \ge 0\}_x$ for some $1 \le r_x \le d$, so $e_x = 0$.

Conversely, suppose $e_x = 0$. Then, either $x \in \operatorname{Reg}(\mathfrak{T})$ (so $\mathfrak{T}_x = \operatorname{Reg}(\mathfrak{T})_x$ is the germ at x of a Nash manifold) or $\overline{\partial \mathfrak{T}_x}^{\operatorname{an}} = \{\mathfrak{u}_1 \cdots \mathfrak{u}_{r_x} = 0\}_x$ for some $1 \leq r_x \leq d$ and, after changing the sign of some of the variables, we may assume $\mathfrak{T}_x \subset \{\mathfrak{u}_1 \geq 0, \ldots, \mathfrak{u}_{r_x} \geq 0\}_x$, because $e_x = 0$. As $\mathfrak{T}_x \setminus \overline{\partial \mathfrak{T}_x}^{\operatorname{an}} = \operatorname{Reg}(\mathfrak{T})_x \setminus \overline{\partial \mathfrak{T}_x}^{\operatorname{an}}$ is an open and closed germ, $\mathfrak{T}_x \setminus \overline{\partial \mathfrak{T}_x}^{\operatorname{an}}$ is a union of connected components of $\{\mathfrak{u}_1 \cdots \mathfrak{u}_{r_x} \neq 0\}_x$ contained in $\{\mathfrak{u}_1 > 0, \ldots, \mathfrak{u}_{r_x} > 0\}_x$, so $\mathfrak{T}_x \setminus \overline{\partial \mathfrak{T}_x}^{\operatorname{an}} = \{\mathfrak{u}_1 > 0, \ldots, \mathfrak{u}_{r_x} > 0\}_x$. As \mathfrak{T}_x is closed and pure dimensional and $\dim(\overline{\partial \mathfrak{T}_x}^{\operatorname{an}}) < \dim(\mathfrak{T}_x)$, we conclude $\mathfrak{T}_x = \{\mathfrak{u}_1 \geq 0, \ldots, \mathfrak{u}_{r_x} \geq 0\}_x$ is the germ at x of a Nash manifold with corners, as required. \Box

It follows from the previous statement: A closed checkerboard set is a Nash manifold with corners if and only if $e_x(\mathfrak{T}) = 0$ for each $x \in \mathfrak{T}$.

Lemma 4.5. Let $\mathfrak{T} \subset \mathbb{R}^n$ be a closed checkerboard set. Then $e_x \neq 1$, for each $x \in \partial \mathfrak{T}$.

Proof. Let $X := \overline{\mathcal{T}}^{\text{zar}}$. For each $x \in \partial \mathcal{T}$ there exist an open semialgebraic set $U \subset X$ equipped with a Nash diffeomorphism $u := (u_1, \ldots, u_d) : U \to \mathbb{R}^d$ and an integer $1 \leq r_x \leq d$ such that u(x) = 0 and $\overline{\partial \mathcal{T}}_x^{\text{an}} = \{\mathbf{u}_1 \cdots \mathbf{u}_{r_x} = 0\}_x$.

Suppose that $e_x \ge 1$ for some $x \in \partial \mathcal{T}$. As $e_x \ne 0$, then $r_x \ge 2$, because otherwise \mathcal{T}_x is the germ of a Nash manifold with boundary and $e_x = 0$. Up to rename the variables if necessary, we may assume $\mathcal{T}_x \setminus \{\mathbf{u}_1 = 0\}_x$ is disconnected. Suppose that for each $2 \le i \le r_x$ the germ $\mathcal{T}_x \setminus \{\mathbf{u}_i = 0\}_x$ is connected. After changing the signs of some of the variables if necessary, we may assume $\mathcal{T}_x \subset \{\mathbf{u}_2 \ge 0, \ldots, \mathbf{u}_{r_x} \ge 0\}_x$. Proceeding as in the proof of Lemma 4.4, as $\mathcal{T}_x \setminus \overline{\partial \mathcal{T}_x^{\mathrm{an}}} = \operatorname{Reg}(\mathcal{T})_x \setminus \overline{\partial \mathcal{T}_x^{\mathrm{an}}}$ is an open and closed germ, $\mathcal{T}_x \setminus \overline{\partial \mathcal{T}_x^{\mathrm{an}}}$ is a union of connected components of $\{\mathbf{u}_1 \cdots \mathbf{u}_{r_x} \ne 0\}_x$ contained in $\{\mathbf{u}_2 > 0, \ldots, \mathbf{u}_{r_x} > 0\}_x$. As \mathcal{T}_x is closed and pure dimensional and dim $(\overline{\partial \mathcal{T}_x^{\mathrm{an}}) < \operatorname{dim}(\mathcal{T}_x)$, we have only two possibilities:

• $\mathfrak{T}_x = {\mathfrak{u}_1 \ge 0, \dots, \mathfrak{u}_{r_x} \ge 0}_x$ (up to changing the sign of the germ \mathfrak{u}_1 if necessary),

•
$$\mathfrak{T}_x = {\mathfrak{u}_2 \ge 0, \dots, \mathfrak{u}_{r_x} \ge 0}_x.$$

In the first case $e_x = 0$, which contradicts the fact that $\mathfrak{T}_x \setminus \{\mathfrak{u}_1 = 0\}_x$ is disconnected, whereas in the second case $\{\mathfrak{u}_1 = 0\}_x \not\subset \overline{\partial \mathfrak{T}}^{\mathrm{an}}$, which contradicts the fact that $\overline{\partial \mathfrak{T}}^{\mathrm{an}} = \{\mathfrak{u}_1 \cdots \mathfrak{u}_{r_x} = 0\}_x$. Thus, there exists $2 \leq i \leq r_x$ such that $\mathfrak{T}_x \setminus \{\mathfrak{u}_i = 0\}_x$ is disconnected, so $e_x \geq 2$ as required. \Box

We show next that the function $e(\mathcal{T})$ is a semialgebraic function.

Lemma 4.6 (Semialgebricity of $e(\mathfrak{T})$). Let $\mathfrak{T} \subset \mathbb{R}^n$ be a d-dimensional closed checkerboard set and let $0 \leq e \leq d$. The set $\mathfrak{T}_e := \{x \in \mathfrak{T} : e_x = e\}$ is a semialgebraic set and \mathfrak{T}_0 is an open subset of \mathfrak{T} . In addition, if Z is the Zariski closure of $\partial \mathfrak{T}$ and C is a connected component of $\operatorname{Sing}_{\ell}(Z) \setminus \operatorname{Sing}_{\ell+1}(Z)$ for some $0 \leq \ell \leq d-1$, then either $C \cap \mathfrak{T} = \emptyset$ or $C \subset \mathfrak{T}$ and $e(\mathfrak{T})$ is constant on C.

Proof. The boundary $\partial \mathfrak{T}$ is a closed semialgebraic subset of the Nash manifold $X := \overline{\mathfrak{T}}^{\operatorname{zar}}$. For each $x \in \partial \mathfrak{T}$ there exists a coordinate system $(\mathfrak{u}_1, \ldots, \mathfrak{u}_d)$ of X at x and an integer $1 \leq r_x \leq d$ such that $\overline{\partial \mathfrak{T}_x^{\operatorname{an}}} = {\mathfrak{u}_1 \cdots \mathfrak{u}_{r_x}} = 0_x$. By [FGR, Prop.4.4, Prop.4.6] there exist finitely many open semialgebraic sets ${U_i}_{i=1}^s$ equipped with Nash diffeomorphisms $\mathfrak{u}_i := (\mathfrak{u}_{i1}, \ldots, \mathfrak{u}_{id}) : U_i \to \mathbb{R}^d$ and integers $r_i \geq 1$ such that $\overline{\partial \mathfrak{T}_x^{\operatorname{an}}} = {\mathfrak{u}_{i1} \cdots \mathfrak{u}_{ir_i}} = 0_x$ for all $x \in \mathfrak{T} \cap U_i$.

Fix $i \in \{1, \ldots, s\}$ and $J \subset \{1, \ldots, r_i\}$. Reordering the variables if necessary, we may assume $J = \{1, \ldots, m\}$ for some $1 \le m \le r_i$. Let V be a connected component of $U_i \setminus \{\mathbf{u}_{i,m+1} \cdots \mathbf{u}_{i,r_i} = 0\}$. After changing the signs of some of the variables if necessary, we may assume $V := \{\mathbf{u}_{i,m+1} > 0\}$. Consider the semialgebraic set $\mathcal{T}' := \mathcal{T} \cap U_i \cap V$ and the projection

$$\pi_i: \mathbb{R}^d \equiv \mathbb{R}^m \times \mathbb{R}^{d-m} \to \mathbb{R}^m$$

onto the first *m* coordinates. We take coordinates (x_1, \ldots, x_m) on \mathbb{R}^m and (x_{m+1}, \ldots, x_d) on \mathbb{R}^{d-m} . Denote $\Lambda_i := \{\mathbf{x}_{i,m+1} > 0, \ldots, \mathbf{x}_{i,r_i} > 0\} \subset \mathbb{R}^{d-m}$. As $u_i(\mathfrak{T} \cap U_i)$ is the union of connected components of $\mathbb{R}^d \setminus \{\mathbf{x}_1 \ldots \mathbf{x}_{r_i} = 0\}$, there exist $\varepsilon_{i,1}, \ldots, \varepsilon_{i,k} \in \{-1, 1\}^{r_i}$ such that

$$u_i(\mathfrak{T}\cap U_i) = \bigcup_{p=1}^k \{\varepsilon_{1p_1} \mathbf{x}_1 \ge 0, \dots, \varepsilon_{ip_{r_i}} \mathbf{x}_{r_i} \ge 0\}$$

where $\varepsilon_{ip} := (\varepsilon_{ip_1}, \ldots, \varepsilon_{ip_{r_i}})$. Consequently,

$$u_i(\mathcal{T}') = u_i(\mathcal{T} \cap U_i \cap V) = \bigcup_{p=1}^k \{\varepsilon_{1p_1} \mathbf{x}_1 \ge 0, \dots, \varepsilon_{ip_{r_i}} \mathbf{x}_{r_i} \ge 0, \mathbf{x}_{m+1} > 0, \dots, \mathbf{x}_{r_i} > 0\}$$

Observe that

$$\begin{split} \{\varepsilon_{1p_1}\mathbf{x}_1 \ge 0, \dots, \varepsilon_{ip_{r_i}}\mathbf{x}_{r_i} \ge 0, \mathbf{x}_{m+1} > 0, \dots, \mathbf{x}_{r_i} > 0\} \\ &= \begin{cases} \{\varepsilon_{1p_1}\mathbf{x}_1 \ge 0, \dots, \varepsilon_{ip_m}\mathbf{x}_m \ge 0, \mathbf{x}_{m+1} > 0, \dots, \mathbf{x}_{r_i} > 0\} & \text{if } \varepsilon_{ip_{m+1}} = \dots = \varepsilon_{ip_{r_i}} = 1, \\ \varnothing & \text{otherwise.} \end{cases} \end{split}$$

Thus,

$$u_i(\mathfrak{T}') = \bigcup_{\substack{p \in \{1,\dots,k\}\\ (\varepsilon_{ip_{m+1}},\dots,\varepsilon_{ip_r}) = (1,\dots,1)}} \{\varepsilon_{1p_1} \mathbf{x}_1 \ge 0,\dots,\varepsilon_{ip_m} \mathbf{x}_m \ge 0, \mathbf{x}_{m+1} > 0,\dots,\mathbf{x}_{r_i} > 0\} = \pi_i(u_i(\mathfrak{T}')) \times \Lambda_i.$$

Observe that $W := {\mathbf{u}_{i1} = 0, \dots, \mathbf{u}_{im} = 0, \mathbf{u}_{i,m+1} > 0, \dots, \mathbf{u}_{ir_i} > 0} \subset \mathcal{T}$ and for each $x \in W$, we have $e_x(\mathcal{T}) = e_0(\pi_i(u_i(\mathcal{T}')))$, so $e_x(\mathcal{T})$ is constant on W. As each $x \in \mathcal{T} \cap U_i \cap {\mathbf{u}_{i1} \dots \mathbf{u}_{ir_i} = 0}$ belongs to a set of the type $W_{J,\varepsilon} := {\mathbf{u}_{ij} = 0, j \in J} \cap {\varepsilon_j \mathbf{u}_{ij} > 0, j \notin J}$ where $J = {1, \dots, r_i}$ and $\varepsilon_j \in {-1, 1}$, the function $e(\mathcal{T})$ provides a semialgebraic partition of $\partial \mathcal{T} \cap U_i$ for each $i = 1, \dots, s$. In particular, each set \mathcal{T}_e is semialgebraic. As the condition 'to be a Nash manifold with corners' is a local open condition, we deduce \mathcal{T}_0 is an open semialgebraic subset of \mathcal{T} .

We have proved above that if $Z'_i := \{u_{i1} \cdots u_{ir_i} = 0\}$ and C'_i is a connected component of $\operatorname{Sing}_{\ell}(Z'_i) \setminus \operatorname{Sing}_{\ell+1}(Z'_i)$ for some $0 \leq \ell \leq d-1$, then either $C'_i \cap \mathfrak{T} = \emptyset$ or $C'_i \subset \mathfrak{T}$ and $e(\mathfrak{T})$ is constant on C'_i . Let C be a connected component of $\operatorname{Sing}_{\ell}(Z) \setminus \operatorname{Sing}_{\ell+1}(Z)$. As \mathfrak{T} is a checkerboard set, either $C \cap \mathfrak{T} = \emptyset$ or $C \subset \mathfrak{T}$. Assume we are in the second case. As we have proved above, there exists a finite semialgebraic open covering $\mathcal{W}_C := \{W_i\}_{i=1}^p$ of C such that $e(\mathfrak{T})$ is constant on W_i . If $x, y \in C$, there exists $W_{i_1}, \ldots, W_{i_q} \in \mathcal{W}_C$ such that $x \in W_{i_1}, y \in W_{i_q}$ and $W_{i_j} \cap W_{i_{j+1}} \neq \emptyset$ for $j = 1, \ldots, q-1$. As $e(\mathfrak{T})|_{W_{i_j}}$ is constant, we deduce recursively that $e_x(\mathfrak{T}) = e_y(\mathfrak{T})$, as required. \square

We show next that the function $e(\mathcal{T})$ is upper semi-continuous.

Lemma 4.7 (Upper semi-continuity of $e(\mathfrak{T})$). Let $\mathfrak{T} \subset \mathbb{R}^d$ be a d-dimensional closed checkerboard set and let $x \in \partial \mathfrak{T}$. Then $e_x \geq e_y$ for each $y \in \mathfrak{T}$ close enough to x.

Proof. Let k be the maximum of the values $e \ge 0$ such that $x \in \operatorname{Cl}(\mathfrak{T}_e)$. It is enough to check that $e_x \ge k$. Consider the Nash manifold $X := \overline{\mathfrak{T}}^{\operatorname{zar}}$ and let $U \subset X$ be an open semialgebraic neighborhood of x equipped with a Nash diffeomorphism $u := (u_1, \ldots, u_d) : U \to \mathbb{R}^d$ such that u(x) = 0 and $\overline{\partial \mathfrak{T}_x}^{\operatorname{an}} = \{\mathfrak{u}_1 \cdots \mathfrak{u}_r = 0\}_x$. If $e_x < k$, we may assume $\mathfrak{T}_x \subset \{\mathfrak{u}_k \ge 0, \ldots, \mathfrak{u}_r \ge 0\}_x$ and $\partial \mathfrak{T}_x \subset \{\mathfrak{u}_1 \cdots \mathfrak{u}_r = 0\}_x$. Shrinking U if necessary, we have $\mathfrak{T} \cap U \subset \{\mathfrak{u}_k \ge 0, \ldots, \mathfrak{u}_r \ge 0\}$ and $\partial \mathfrak{T} \cap U \subset \{\mathfrak{u}_1 \cdots \mathfrak{u}_r = 0\}$. Thus, $e_y < k$ for each $y \in U$, which is a contradiction because $x \in \operatorname{Cl}(\mathfrak{T}_k)$.

Remark 4.8. As $e(\mathcal{T})$ is upper semi-continuous, the set $\bigcup_{k\geq e} \mathcal{T}_k$ is a closed subset of $\partial \mathcal{T}$ for each $1\leq e\leq d$.

4.2.2. Closed checkerboard sets and drilling blow-up. We want to study now how the value $e_x(\mathcal{T})$ changes after performing a drilling blow-up. We show the following:

Lemma 4.9. Let $S \subset \mathbb{R}^n$ be a d-dimensional closed checkerboard set. Denote $X := \overline{S}^{\text{zar}}$ and $Z := \overline{\partial S}^{\text{zar}}$. Let Z_1, \ldots, Z_r be the irreducible components of Z and Y an irreducible component of $Z_1 \cap \cdots \cap Z_\ell$ for some $2 \leq \ell \leq r$. Let $(\widehat{X}, \widehat{\pi})$ be the twisted Nash double of the drilling blow-up (\widetilde{X}, π_+) of X with center Y. Let $\mathfrak{T} := \operatorname{Cl}(\pi_+^{-1}(\mathbb{S} \setminus Y)) = \pi_+^{-1}(\mathbb{S}) \cap \operatorname{Cl}(\pi_+^{-1}(\mathbb{S} \setminus Y))$ be the strict transform of \mathbb{S} under π_+ . Then \mathfrak{T} is a d-dimensional closed checkerboard set, $e_y(\mathfrak{T}) \leq e_{\pi_+(y)}(\mathbb{S})$ for each $y \in \partial \mathfrak{T} \cap \pi_+^{-1}(Y)$ such that $\mathbb{S}_{\pi_+(y)} \setminus Z_{i,\pi_+(y)}$ is not connected for $i = 1, \ldots, \ell$.

Proof. As $\ell \geq 2$, we have dim $(Y) \leq d - 2$, so Reg $(S) \setminus Y$ is connected, because Reg(S) is a connected *d*-dimensional Nash manifold. As S and T are both pure dimensional, we have

$$\pi_+^{-1}(\operatorname{Reg}(\mathfrak{S})\setminus Y)\subset\operatorname{Sth}(\mathfrak{I})=\operatorname{Reg}(\mathfrak{I})\subset\operatorname{Cl}(\pi_+^{-1}(\mathfrak{S}\setminus Y))=\operatorname{Cl}(\pi_+^{-1}(\operatorname{Reg}(\mathfrak{S})\setminus Y)).$$

Thus, $\operatorname{Reg}(\mathfrak{T})$ is connected and \mathfrak{T} is a checkerboard set, because (see §3.4): it is closed, $\overline{\mathfrak{T}}^{\operatorname{zar}} = \widehat{X}$ is a non-singular real algebraic set and the Zariski closure of $\partial \mathfrak{T}$ is a union of irreducible components of $\widehat{\pi}^{-1}(\overline{\partial S}^{\operatorname{zar}})$, which is a normal-crossings divisor by Remarks 3.4 and 3.10.

As $\pi_+|_{\widetilde{X}\setminus\pi_+^{-1}(Y)}: \widetilde{X}\setminus\pi_+^{-1}(Y)\to X\setminus Y$ is a Nash diffeomorphism, it holds $e_y(\mathfrak{T})=e_{\pi_+(y)}(\mathfrak{S})$ for each $y\in\partial\mathfrak{T}\setminus\pi_+^{-1}(Y)$. Let us see what happens at the points of $\partial\mathfrak{T}\cap\pi_+^{-1}(Y)$. Fix a point $y\in\partial\mathfrak{T}\cap\pi_+^{-1}(Y)$ and denote $x:=\pi_+(y)\in Y$.

Assume that the irreducible components of Z that contain x are $Z_1, \ldots, Z_{r'}$ for some $2 \leq \ell \leq r' \leq r$. Let $U \subset X$ be an open semialgebraic neighborhood of x equipped with a Nash diffeomorphism $u := (u_1, \ldots, u_d) : U \to \mathbb{R}^d$ such that $u(Z \cap U) = \{\mathbf{u}_1 \cdots \mathbf{u}_{r'} = 0\}$ and $u(Y \cap U) = \{\mathbf{u}_1 = 0, \ldots, \mathbf{u}_\ell = 0\}$. Write $e := \dim(Y) = d - \ell$ and assume that $e_x(S) = k \leq r$. Reordering the variables and changing their signs if necessary, we may assume

$$\mathcal{S} \cap U \subset \{\mathbf{u}_1 \ge 0, \dots, \mathbf{u}_m \ge 0, \mathbf{u}_{\ell+1} \ge 0, \dots, \mathbf{u}_s \ge 0\}$$

$$(4.1)$$

for some $0 \le m \le \ell$ and $\ell \le s \le r'$ and both m and s are maximal satisfying (4.1). If m = 0, then $S \cap U \subset \{u_{\ell+1} \ge 0, \dots, u_s \ge 0\}$, whereas if $s = \ell$, then $S \cap U \subset \{u_1 \ge 0, \dots, u_m \ge 0\}$. As $e_x(S) = k$, we have $k = (\ell - m) + (r' - s)$. By Remarks 3.4 we can choose (local) coordinates in \widehat{X} such that π_+ behaves (with the already taken (local) coordinates in X) as the Nash map

$$g_+: [0, +\infty) \times \mathbb{S}^{d-e-1} \times \mathbb{R}^e \to \mathbb{R}^d, \ (\rho, w, z) \mapsto (\rho w, z),$$

where $w := (w_1, ..., w_\ell)$ and $z := (z_{\ell+1}, ..., z_d) \in \mathbb{R}^e = \mathbb{R}^{d-\ell}$. We have

$$g_{+}^{-1}(u(Z \cap U)) = \{\rho^{\mathfrak{c}} \mathfrak{w}_{1} \cdots \mathfrak{w}_{\ell} \mathbf{z}_{\ell+1} \cdots \mathbf{z}_{r'} = 0, \mathfrak{w}_{1}^{2} + \dots + \mathfrak{w}_{\ell}^{2} = 1\},\$$

$$g_{+}^{-1}(u(Y \cap U)) = \{\rho \mathfrak{w}_{1} = 0, \dots, \rho \mathfrak{w}_{\ell} = 0, \mathfrak{w}_{1}^{2} + \dots + \mathfrak{w}_{\ell}^{2} = 1\} = \{\rho = 0, \mathfrak{w}_{1}^{2} + \dots + \mathfrak{w}_{\ell}^{2} = 1\},\$$

$$g_{+}^{-1}(u(S \cap U \setminus Y)) \subset \{\rho \ge 0, \mathfrak{w}_{1} \ge 0, \dots, \mathfrak{w}_{m} \ge 0, \mathbf{z}_{\ell+1} \ge 0, \dots, \mathbf{z}_{s} \ge 0, \mathfrak{w}_{1}^{2} + \dots + \mathfrak{w}_{\ell}^{2} = 1\}.$$

If m = 0, then $S \cap U \subset {u_{\ell+1} \ge 0, \dots, u_s \ge 0}$ and

$$g_+^{-1}(u(\mathcal{S} \cap U \setminus Y)) \subset \{\rho \ge 0, \mathbf{z}_{\ell+1} \ge 0, \dots, \mathbf{z}_s \ge 0, \mathbf{w}_1^2 + \dots + \mathbf{w}_\ell^2 = 1\}.$$

Thus, $e_y(\mathfrak{T}) \leq \ell - 1 + r' - s = k - 1 < k = e_x(\mathfrak{S})$ for each $y \in g^{-1}(x)$. The condition m = 0 means that $\mathfrak{S}_x \setminus Z_{i,x}$ is not connected for $i = 1, \ldots, \ell$.

We assume in the following $m \ge 1$. Observe that $S_x \setminus Z_{i,x}$ is not connected for $i = m + 1, \ldots, \ell$. Let us show $e_y(\mathfrak{T}) \le e_x(\mathfrak{S})$ for each $y \in g^{-1}(x)$. It may happen that for some $y \in g^{-1}(x)$ the previous inequality is strict even if $S_x \setminus Z_{i,x}$ is connected for the indices $i = 1, \ldots, m$. If some $\mathbf{w}_i(y) \ne 0$, this variable has no relevance in the description of \mathfrak{T} locally around y and \mathbf{w}_i behaves as $\pm \sqrt{1 - \sum_{j \ne i} \mathbf{w}_j^2}$. Analogously, if some $\mathbf{z}_j(y) \ne 0$, this variable has no relevance in the description of \mathfrak{T} locally around y. As $\mathbf{w}_1^2 + \cdots + \mathbf{w}_\ell^2 = 1$, there exists an index $1 \le i \le \ell$ such that $\mathbf{w}_i(y) \ne 0$: CASE 1. If $1 \le i \le m$, we may assume i = m. Thus, $\mathbf{w}_m(y) > 0$ and $\mathbf{w}_m = +\sqrt{1 - \sum_{j \ne m} \mathbf{w}_j^2}$. We consider (local) coordinates $(\rho, \mathbf{w}_1, \ldots, \mathbf{w}_{m-1}, \mathbf{w}_{m+1}, \ldots, \mathbf{w}_\ell, \mathbf{z}_{\ell+1}, \ldots, \mathbf{z}_s, \mathbf{z}_{s+1}, \ldots, \mathbf{z}_{r'}, \mathbf{z}_{r'+1}, \ldots, \mathbf{z}_d)$ and

$$e_y(\mathfrak{T}) \le \ell - (m-1+1) + (r'-s) = (\ell - m) + (r'-s) = k = e_x(\mathfrak{S}).$$

CASE 2. If $m+1 \leq i \leq \ell$, we may assume $i = \ell$. Thus, $\mathbf{w}_{\ell}(y) \neq 0$ and $\mathbf{w}_{\ell} = \pm \sqrt{1 - \sum_{j=1}^{\ell-1} \mathbf{w}_i^2}$. We consider (local) coordinates $(\rho, \mathbf{w}_1, \dots, \mathbf{w}_m, \mathbf{w}_{m+1}, \dots, \mathbf{w}_{\ell-1}, \mathbf{z}_{\ell+1}, \dots, \mathbf{z}_s, \mathbf{z}_{s+1}, \dots, \mathbf{z}_{r'}, \mathbf{z}_{r'+1}, \dots, \mathbf{z}_d)$ and

$$e_y(\mathfrak{T}) \le 1 + (\ell - 1) - (m + 1) + (r' - s) = (\ell - m) + (r' - s) - 1 = k - 1 < k = e_x(\mathfrak{S}),$$

as required.

4.2.3. Closed checkerboard sets and Nash manifolds with corners. We are ready to prove Theorem 1.4. After the previous preparation (in particular Theorem 4.3) we are reduced to the case when S is a d-dimensional closed checkerboard set.

Proof of Theorem 1.4. Let us construct the Nash manifold with corners Ω first. Let $S \subset \mathbb{R}^m$ be a *d*-dimensional closed checkerboard set and denote $X_0 := \overline{S}^{\text{zar}}$. Let $Z := \overline{\partial S}^{\text{zar}}$ and Z_1, \ldots, Z_r be its irreducible components. Define $e := \max\{e_x(S) : x \in \partial S\}$.

If e = 0, we conclude by Lemma 4.4 that S is already a Nash manifold with corners. Otherwise, by Lemma 4.5 $e \ge 2$. By Remark 4.8 S_e is a closed subset of ∂S . By Lemma 4.6 S_e is a union of connected components of the semialgebraic sets $\operatorname{Sing}_{\ell}(Z) \setminus \operatorname{Sing}_{\ell+1}(Z)$ for $1 \le \ell \le d-1$ (recall that $e \ge 2$).

Pick a point $x \in S_e$ and assume that Z_1, \ldots, Z_e are the irreducible components of Z such that the germ $S_x \setminus Z_{i,x}$ is not connected for $i = 1, \ldots, e$. Then there exists an open semialgebraic neighborhood $U \subset X$ of x equipped with a Nash diffeomorphism $u := (u_1, \ldots, u_d) : U \to \mathbb{R}^d$ such that $u(x) = 0, Z \cap U = \{\mathbf{u}_1 \cdots \mathbf{u}_r = 0\}, Z_i \cap U = \{\mathbf{u}_i = 0\}$ and $S \cap U \subset \{\mathbf{u}_{e+1} \ge 0, \ldots, \mathbf{u}_r \ge 0\}$. Thus, $\{\mathbf{u}_1 = 0, \ldots, \mathbf{u}_e = 0, \mathbf{u}_{e+1} \ge 0, \ldots, \mathbf{u}_r \ge 0\} \subset S_e$. By Lemma 4.7 the connected component \mathbb{C} of $(Z_1 \cap \cdots \cap Z_e) \setminus \bigcup_{i=e+1}^r Z_i$ that contains $\{\mathbf{u}_1 = 0, \ldots, \mathbf{u}_e = 0, \mathbf{u}_{e+1} \ge 0, \ldots, \mathbf{u}_r \ge 0\}$ is contained in S_e . Thus, the Zariski closure of \mathbb{C} is the irreducible component of $Z_1 \cap \cdots \cap Z_e$ that contains x. As we can repeat the previous argument for each $x \in S_e$, we conclude that the Zariski closure of S_e is a union of irreducible components of the real algebraic set $\bigcup_{\{i_1,\ldots,i_e\} \subset \{1,\ldots,r\}} \bigcap_{j=1}^e Z_{i_j}$. As $e \ge 2$, we have $\dim(\overline{S_e}^{\operatorname{zar}}) \le d-2$.

In addition, for each $x \in S_e$ there exist irreducible components Z_{i_1}, \ldots, Z_{i_e} of Z such that the germ $S_x \setminus Z_{i_j,x}$ is not connected for $j = 1, \ldots, e$. We proceed by double induction on e and the number m of irreducible components of the Zariski closure of $\overline{S_e}^{\text{zar}}$ of S_e .

Let W be an irreducible component of $\overline{\mathcal{S}_e}^{\text{zar}}$, which has dimension $\leq d-2$. Let $(\widehat{X}_0, \widehat{\pi})$ be the twisted Nash double of the drilling blow-up (\widetilde{X}_0, π_+) of X_0 with center W, which is by Subsection 3.4 a non-singular real algebraic set. Let

$$\mathfrak{T} := \operatorname{Cl}(\pi_+^{-1}(\mathfrak{S} \setminus W)) = \pi_+^{-1}(\mathfrak{S}) \cap \operatorname{Cl}(\pi_+^{-1}(\mathfrak{S} \setminus W))$$

be the strict transform of S under π_+ (recall that S is closed). As S is pure dimensional and $S_e \subset \overline{\partial S}^{zar}$ has dimension strictly smaller, $S \setminus W$ is dense in S, so

$$\pi_{+}(\mathfrak{T}) = \pi_{+}(\operatorname{Cl}(\pi_{+}^{-1}(\mathbb{S} \setminus W))) = \operatorname{Cl}(\pi_{+}(\pi_{+}^{-1}(\mathbb{S} \setminus W))) = \operatorname{Cl}(\mathbb{S} \setminus W) = \mathbb{S},$$

because $\pi_+ : \widetilde{X}_0 \to X_0$ is proper and surjective. As $e \ge 2$, we can apply Lemma 4.9 and we deduce: \mathfrak{T} is a checkerboard set, $e_y(\mathfrak{T}) \le e_{\pi_+(y)}(\mathfrak{S})$ for each $y \in \partial \mathfrak{T}$ and $e_y(\mathfrak{T}) < e_{\pi_+(y)}(\mathfrak{S})$ for each $y \in \partial \mathfrak{T} \cap \pi_+^{-1}(W)$ such that $\mathfrak{S}_{\pi_+(y)} \setminus Z_{i_j,\pi_+(y)}$ is not connected for $j = 1, \ldots, e$.

If $\max\{e(\mathfrak{T})_y: y \in \partial \mathfrak{T}\} < e$, by induction hypothesis the statement holds for \mathfrak{T} , so it also holds for \mathfrak{S} . If $\max\{e(\mathfrak{T})_y: y \in \partial \mathfrak{T}\} = e$, the Zariski closure of \mathfrak{T}_e is contained in $\operatorname{Cl}(\widehat{\pi}^{-1}(\overline{\mathfrak{S}_e}^{\operatorname{zar}} \setminus W))$ and it has m-1 irreducible components. As by Lemma 4.4 e = 0 if and only if \mathfrak{T} is a Nash manifold with corners, our inductive argument is consistent. Thus, by induction hypothesis the statement holds for \mathfrak{T} , so it also holds for \mathfrak{S} .

Let $\mathcal{Q} \subset \mathbb{R}^n$ be the Nash manifold with corners obtained by our inductive process. As \mathcal{Q} is a checkerboard set, $\operatorname{Reg}(\mathcal{Q})$ is connected, so also $\mathcal{Q} = \operatorname{Cl}(\operatorname{Reg}(\mathcal{Q}))$ is connected. We have constructed \mathcal{Q} , starting from \mathcal{S} , with a finite number of drilling blow-ups. Namely, we have constructed a finite number of tuples $\{\mathcal{T}_i, (\widetilde{X}_{\mathcal{T}_i}, \pi_{+,i}), (\widehat{X}_{\mathcal{T}_i}, \widehat{\pi}_i)\}_{i=0}^s$ where:

- $\mathcal{T}_{0} := \mathfrak{S}, \ \widetilde{X}_{\mathcal{T}_{0}} := X_{0} = \overline{\mathfrak{S}}^{\operatorname{zar}}, \ \pi_{+,0} := \operatorname{id}_{X_{0}}, \ \widehat{X}_{\mathcal{T}_{0}} := X_{0} \ \text{and} \ \widehat{\pi}_{0} := \operatorname{id}_{X_{0}}.$
- $(\widetilde{X}_{\mathfrak{T}_{i+1}}, \pi_{+,i+1})$ is the drilling blow-up of the irreducible non-singular real algebraic set $\overline{\mathcal{T}}_i^{\operatorname{zar}}$ with center a (suitable) irreducible non-singular real algebraic subset W_i of $\overline{\mathcal{T}}_i^{\operatorname{zar}}$ of dimension $\leq d-2$.
- $(\widehat{X}_{\mathfrak{T}_{i+1}}, \widehat{\pi}_{i+1})$ is the twisted Nash double of $(\widetilde{X}_{\mathfrak{T}_{i+1}}, \pi_{+,i+1})$ and it is a non-singular real algebraic set of dimension d. By Lemma 3.13 $\widehat{X}_{\mathfrak{T}_{i+1}}$ is the Zariski closure of $\widetilde{X}_{\mathfrak{T}_{i+1}}$.

- $\mathcal{T}_{i+1} := \operatorname{Cl}(\pi_{+,i+1}^{-1}(\mathcal{T}_i \setminus W_i))$ is the strict transform of \mathcal{T}_i under $\pi_{+,i+1}$ and $\pi_{+,i+1}(\mathcal{T}_{i+1}) = \mathcal{T}_i$. By Lemma 4.9 \mathcal{T}_{i+1} is a checkerboard set.
- $\pi_{+,i+1}^{-1}(W_i) \cap \mathfrak{I}_{i+1} \subset \partial \mathfrak{I}_{i+1}, \ \pi_{+,i+1}^{-1}(\partial \mathfrak{I}_i \setminus W_i) \subset \partial \mathfrak{I}_{i+1} \text{ and } \pi_{+,i+1}^{-1}(\operatorname{Reg}(\mathfrak{I}_i) \setminus W_i) \subset \operatorname{Reg}(\mathfrak{I}_{i+1}) \text{ (use Fact 3.7(ii) and §3.4). Consequently, } \operatorname{Reg}(\mathfrak{I}_{i+1}) = \pi_{+,i+1}^{-1}(\operatorname{Reg}(\mathfrak{I}_i) \setminus W_i)$ and $\partial \mathfrak{I}_{i+1} = \pi_{+,i+1}^{-1}(\partial \mathfrak{I}_i \setminus W_i) \cup (\pi_{+,i+1}^{-1}(W_i) \cap \mathfrak{I}_{i+1}).$
- (*) $\pi_{+,i+1}(\partial \mathfrak{T}_{i+1}) = \partial \mathfrak{T}_i \cup (W_i \cap \mathfrak{T}_i).$
- (**) $\pi_{+,i+1}|_{\operatorname{Reg}(\mathfrak{T}_{i+1})} : \operatorname{Reg}(\mathfrak{T}_{i+1}) \to \operatorname{Reg}(\mathfrak{T}_i) \setminus (W_i \cap \mathfrak{T}_i)$ is a Nash diffeomorphism (Fact 3.7(ii)). • $\mathfrak{T}_s = \mathfrak{Q}.$

As \mathfrak{T}_i is connected by analytic paths, $\overline{\mathfrak{T}}_i^{\operatorname{zar}}$ is by [Fe2, Lem.7.3] an irreducible component of the pure dimensional non-singular real algebraic set $\widehat{X}_{\mathfrak{T}_i}$ of dimension d. Thus, $\overline{\mathfrak{T}}_i^{\operatorname{zar}}$ is a pure dimensional irreducible non-singular real algebraic set of dimension d. Consequently, the Nash manifold with corners $\mathfrak{Q} = \mathfrak{T}_s$ is (by Lemma 4.9) a checkerboard set, $X := \overline{\mathfrak{Q}}^{\operatorname{zar}}$ is a d-dimensional irreducible non-singular real algebraic set and $Y := \overline{\partial \mathfrak{Q}}^{\operatorname{zar}}$ is a normal-crossings divisor of X. Thus, (i) and (ii) hold. As all the involved polynomials maps $\widehat{\pi}_i$ are proper, if X_0 is compact, then X is also compact.

Consider the polynomial map $f := \hat{\pi}_1 \circ \ldots \circ \hat{\pi}_s : \hat{X}_{\mathfrak{T}_s} \to X_0$. By Fact 3.7(i) $f : \hat{X}_{\mathfrak{T}_s} \to X_0$ is composition of proper maps, so it is proper. Moreover, as S is closed and Ω is obtained from S after a finite number of drilling blow-ups taking strict transforms in each step, Ω is a closed subset of X. Thus, also the restriction $f|_{\Omega} : \Omega \to X_0$ is proper. In addition, as $\hat{\pi}_{i+1}(\mathfrak{T}_{i+1}) = \mathfrak{T}_i$ for $i = 0, \ldots, s - 1$, we conclude that $f(\Omega) = S$.

Let us show (iv). By property (**) applied inductively we deduce

$$f(\operatorname{Reg}(\mathfrak{Q})) \subset \operatorname{Reg}(\mathfrak{T}_0) \setminus ((W_0 \cap \mathfrak{T}_0) \cup \bigcup_{k=1}^{s-1} (\pi_{+,1} \circ \cdots \circ \pi_{+,k}) (W_k \cap \mathfrak{T}_k)),$$

whereas by property (*) applied recursively we have

$$f(\partial \mathfrak{Q}) = \partial \mathfrak{T}_0 \cup (W_0 \cap \mathfrak{T}_0) \cup \bigcup_{k=1}^{s-1} (\pi_{+,1} \circ \cdots \circ \pi_{+,k}) (W_k \cap \mathfrak{T}_k),$$
(4.2)

so $f(\operatorname{Reg}(\mathfrak{Q})) \cap f(\partial \mathfrak{Q}) = \emptyset$. Thus, as $\mathfrak{S} = f(\mathfrak{Q}) = f(\operatorname{Reg}(\mathfrak{Q})) \cup f(\partial \mathfrak{Q})$, we deduce $f(\operatorname{Reg}(\mathfrak{Q})) = \mathfrak{S} \setminus f(\partial \mathfrak{Q})$ and $f^{-1}(f(\partial \mathfrak{Q})) = \partial \mathfrak{Q}$. Property (**) implies $f|_{\operatorname{Reg}(\mathfrak{Q})} = (\pi_{+,1} \circ \cdots \circ \pi_{+,s})|_{\operatorname{Reg}(\mathfrak{Q})}$: $\operatorname{Reg}(\mathfrak{Q}) \to \mathfrak{S} \setminus f(\partial \mathfrak{Q})$ is a Nash diffeomorphism (because it is a composition of finitely many Nash diffeomorphisms). By (4.2) we have $\partial \mathfrak{S} \subset f(\partial \mathfrak{Q})$. The semialgebraic set $\mathfrak{R} := f(\partial \mathfrak{Q}) \subset \mathfrak{S}$ is closed, because f is proper and $\partial \mathfrak{Q}$ is closed. As $\partial \mathfrak{Q}$ has dimension not greater than d-1, we have $\dim(\mathfrak{R}) \leq d-1 < d$. The semialgebraic sets $\mathfrak{S} \setminus \mathfrak{R}$ and $\mathfrak{Q} \setminus f^{-1}(\mathfrak{R})$ are Nash manifolds, because $\mathfrak{S} \setminus \mathfrak{R}$ is an open semialgebraic subset of the Nash manifold $\operatorname{Reg}(\mathfrak{S}) = \mathfrak{S} \setminus \partial \mathfrak{S}$ and $\mathfrak{Q} \setminus f^{-1}(\mathfrak{R}) = \mathfrak{Q} \setminus \partial \mathfrak{Q} = \operatorname{Reg}(\mathfrak{Q})$. Consequently, $f|_{\operatorname{Reg}(\mathfrak{Q})} : \operatorname{Reg}(\mathfrak{Q}) = \mathfrak{Q} \setminus f^{-1}(\mathfrak{R}) \to \mathfrak{S} \setminus \mathfrak{R}$ is a Nash diffeomorphism, as required.




FIGURE 4.1. Desingularization of the closed checkerboard set S (right) by the Nash manifold with corners Q (left).

23

4.3. Desingularization of general checkerboard sets. Let $S \subset \mathbb{R}^m$ be a semialgebraic set connected by analytic paths. To prove Theorem 1.6 we introduce Nash quasi-manifolds with corners, that is, semialgebraic sets $\mathcal{T} \subset \mathbb{R}^n$ whose closure is a Nash manifold with corners $\mathcal{Q} \subset \mathbb{R}^n$ and $\mathcal{Q} \setminus \mathcal{T}$ is a union of some of the strata of certain (Nash) stratification of $\partial \mathcal{Q}$, that we introduce below. Let us recall the definition of (Nash) stratification of a semialgebraic set.

Definition 4.10. Let $S \subset \mathbb{R}^m$ be a semialgebraic set. A (Nash) *stratification* of S is a finite semialgebraic partition $\{S_{\alpha}\}_{\alpha \in A}$ of S, where each S_{α} is a connected Nash submanifold of \mathbb{R}^m and the following property is satisfied: if $S_{\alpha} \cap \operatorname{Cl}(S_{\beta}) \neq \emptyset$ and $\alpha \neq \beta$, then $S_{\alpha} \subset \operatorname{Cl}(S_{\beta})$ and $\dim(S_{\alpha}) < \dim(S_{\beta})$. The S_{α} are called the *strata* of the (Nash) stratification and if $d := \dim(S_{\alpha})$, then S_{α} is a *d-stratum*.

The condition dim(\mathfrak{S}_{α}) < dim(\mathfrak{S}_{β}) follows from [BCR, Prop.2.8.13], because $\mathfrak{S}_{\alpha} \subset \mathrm{Cl}(\mathfrak{S}_{\beta}) \setminus \mathfrak{S}_{\beta}$.

Definition 4.11. Given a *d*-dimensional semialgebraic set $S \subset \mathbb{R}^m$, we consider the following semialgebraic partition of S. Recall that Sth(S) is the set of points $x \in S$ at which the germ S_x is the germ of a Nash manifold (Section 2.1). Define $\Gamma_1 := Sth(S)$ and $\Gamma_k := Sth(S \setminus \bigcup_{j=1}^{k-1} \Gamma_j)$ for $k \geq 2$. Let $s \geq 1$ be the largest index k such that $\Gamma_k \neq \emptyset$. For each $k \geq 1$ let $\Gamma_{k\ell}$ (for $\ell = 1, \ldots, r_k$) be the connected components of Γ_k . The collection $\mathfrak{G}(S) := \{\Gamma_{k\ell} : 1 \leq k \leq s, 1 \leq \ell \leq r_k\}$ is a partition of S. We say that $\mathfrak{G}(S)$ is *compatible with* a semialgebraic set $\mathcal{T} \subset S$ if \mathcal{T} is the union of some of the $\Gamma_{k\ell}$.

Examples 4.12. (i) The semialgebraic partition $\mathfrak{G}(S)$ of a semialgebraic set $S \subset \mathbb{R}^n$ is not in general a stratification of S. Consider for instance the semialgebraic set $S := \{y^2 - x^3 = 0\} \cap (\{z > 0\} \cup \{z \le 0, y \ge 0\}) \subset \mathbb{R}^3$. Then

$$\Gamma_{11} := \{ y^2 - x^3 = 0, y > 0 \}, \ \Gamma_{12} := \{ y^2 - x^3 = 0, y < 0, z > 0 \}, \ \Gamma_{21} = \{ x = 0, y = 0 \}$$

and $\mathfrak{G}(\mathfrak{S}) = {\Gamma_{11}, \Gamma_{12}, \Gamma_{21}}$. Observe that $\Gamma_{21} \cap \operatorname{Cl}(\Gamma_{12}) \neq \emptyset$, but $\Gamma_{21} \not\subset \operatorname{Cl}(\Gamma_{12})$. Thus, $\mathfrak{G}(\mathfrak{S})$ is not a stratification of \mathfrak{S} .

(ii) If $X \subset \mathbb{R}^n$ is a *d*-dimensional non-singular real algebraic set and $Y \subset X$ is a normalcrossings divisor, then $\mathfrak{G}(Y)$ is a stratification of Y.

It is enough to consider local models, that is, $Y := \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\} \subset \mathbb{R}^d$. In fact, we may assume r = d, because $Y = Z \times \mathbb{R}^{d-r}$ where $Z := \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\} \subset \mathbb{R}^r$. The semialgebraic partition $\mathfrak{G}(Y)$ of Y is the collection of semialgebraic sets $\Lambda_{\ell} := \{\mathbf{x}_1 *_1 0, \dots, \mathbf{x}_d *_d 0\}$ where $*_i \in \{<, =, >\}$ and at least one of the symbols $*_i$ is equal to =. The closure of each Λ_{ℓ} is a union of finitely many Λ_k and consequently $\mathfrak{G}(Y)$ is a stratification of Y.

In this case $\operatorname{Sth}(Y) = \operatorname{Reg}(Y)$ and $\operatorname{Sth}(\operatorname{Sing}_{\ell}(Y)) = \operatorname{Reg}(\operatorname{Sing}_{\ell}(Y))$ for each $\ell \ge 1$.

(iii) Let $X \subset \mathbb{R}^n$ be a non-singular algebraic set, $Y \subset X$ is a normal-crossings divisor and S the closure of a union of connected components of $X \setminus Y$. Then $\mathfrak{G}(S)$ is a stratification of S and $\mathfrak{G}(S)$ is compatible with $\partial S = S \setminus \operatorname{Reg}(S)$ and $S \cap Y$.

It is enough to consider local models, so we may assume $Y := \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\} \subset \mathbb{R}^d$. We suppose r = d, because $Y = Z \times \mathbb{R}^{d-r}$ where $Z := \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\} \subset \mathbb{R}^r$. Thus, $S := \bigcup_{\varepsilon \in \mathfrak{F}} \mathcal{Q}_{\varepsilon}$, where $\mathcal{Q}_{\varepsilon} := \{\varepsilon_1 \mathbf{x}_1 \ge 0, \ldots, \varepsilon_d \mathbf{x}_d \ge 0\}$, $\varepsilon := (\varepsilon_1, \ldots, \varepsilon_d)$ and $\mathfrak{F} \subset \{-1, 1\}^d$. The semialgebraic partition $\mathfrak{G}(S)$ of S is a collection of the type $\Gamma_\ell := \{\mathbf{x}_{i_1} *_{i_1} 0, \ldots, \mathbf{x}_{i_\ell} *_{i_\ell} 0\}$ where $0 \le \ell \le d$, $1 \le i_1 < \cdots < i_\ell \le d$ and $*_{i_k} \in \{<, =, >\}$ for $k = 1, \ldots, \ell$. The closure of each Γ_ℓ is a union of finitely many Γ_k , so $\mathfrak{G}(S)$ is a stratification of S. Observe that $\partial S = S \setminus \operatorname{Reg}(S)$ and $S \cap Y$ are unions of finitely many of the sets $\{\mathbf{x}_{i_1} *_{i_1} 0, \ldots, \mathbf{x}_{i_\ell} *_{i_\ell} 0\}$ with the condition that some of the $*_{i_k}$ are equal to =, that is, all of them belong to $\mathfrak{G}(S)$ and $\mathfrak{G}(S)$ is compatible with $\partial S = S \setminus \operatorname{Reg}(S)$.

In this case $\operatorname{Sth}(\mathfrak{S}) = \operatorname{Reg}(\mathfrak{S})$ and $\operatorname{Sth}(\partial^{\ell}\mathfrak{S}) = \operatorname{Reg}(\partial^{\ell}\mathfrak{S})$, where $\partial^{\ell}\mathfrak{S} := \partial(\partial^{\ell-1}\mathfrak{S})$ for each $\ell \geq 2$. This is because $\partial\mathfrak{S} \subset \operatorname{Sing}(Y)$ and $\partial^{\ell}\mathfrak{S} \subset \operatorname{Sing}_{\ell}(Y)$ for each $\ell \geq 2$.

(iv) If $\Omega \subset \mathbb{R}^n$ is a *d*-dimensional Nash manifold with corners, $\mathfrak{G}(\Omega) = \mathfrak{G}(\partial \Omega) \sqcup \mathfrak{G}(\operatorname{Int}(\Omega))$ is a stratification of Ω .

It is enough to apply Theorem 2.7 and (iii).

Definition 4.13. A subset $\mathcal{T} \subset \mathbb{R}^n$ is a Nash quasi-manifold with corners if $\mathcal{Q} := \operatorname{Cl}(\mathcal{T})$ is a Nash manifold with corners and $Q \setminus \mathcal{T}$ is a union of elements of the stratification $\mathfrak{G}(\partial Q)$.

We are ready to prove Theorem 1.6.

Proof of Theorem 1.6. The proof is conducted in several steps and subsequent reductions:

STEP 1. INITIAL PREPARATION. We embed \mathbb{R}^m in \mathbb{RP}^m and the latter in \mathbb{R}^p for p large enough, so we can suppose S is in addition bounded. The previous embedding is a regular map, which is in particular a Nash map. Thus, Cl(S) is compact and the Zariski closure of S is also compact. By Theorem 4.3 we may assume that $S \subset \mathbb{R}^m$ is a checkerboard set, whose Zariski closure X is a d-dimensional non-singular compact algebraic subset of \mathbb{R}^m . In particular, Reg(S) is connected and the Zariski closure Z_1 of $\partial S := \operatorname{Cl}(S) \setminus \operatorname{Reg}(S)$ is a normal-crossings divisor of X. We construct next a semialgebraic partition of Cl(S) as a finite union of Nash manifolds of different dimensions compatible with $\mathcal{T}_1 := \operatorname{Cl}(\mathcal{S}) \setminus \mathcal{S}$ and $\mathcal{T}_2 := \mathcal{S} \setminus \operatorname{Reg}(\mathcal{S})$. Observe that \mathcal{T}_1 and \mathcal{T}_2 are disjoint semialgebraic sets, they have dimensions $\leq d-1$ and $\partial S = T_1 \sqcup T_2$.




FIGURE 4.2. A bounded checkerboard set S (left) and S with the sets T_1 (blue) and \mathcal{T}_2 (red) coloured (right).

Let N_1 be the union of the connected components of $\operatorname{Reg}(\mathcal{T}_1) \sqcup \operatorname{Reg}(\mathcal{T}_2)$ of dimension d-1. Note that the connected components of dimension d-1 of $\operatorname{Reg}(\mathcal{T}_1) \sqcup \operatorname{Reg}(\mathcal{T}_2)$ are in general different from the connected components of dimension d-1 of $\operatorname{Reg}(\partial S)$. As $\dim(\mathfrak{T}_i \setminus \operatorname{Reg}(\mathfrak{T}_i)) \leq d-2$ and $\partial S = \mathcal{T}_1 \sqcup \mathcal{T}_2$, the semialgebraic set $\partial S \setminus N_1$ has dimension $\leq d-2$. Let Z_2 be the Zariski closure of $\partial S \setminus N_1$. Each connected component of the Nash manifold $N'_1 := \partial S \setminus Z_2 = N_1 \setminus Z_2$ has dimension d-1 and it is contained either in \mathcal{T}_1 or \mathcal{T}_2 . In addition, $\partial S \setminus N'_1 \subset Z_2$ has dimension $\leq d-2, Z_2$ is the Zariski closure of $\partial S \setminus N'_1$ (because $\partial S \setminus N_1 \subset \partial S \setminus N'_1 \subset Z_2$ and Z_2 is the Zariski closure of $\partial S \setminus N_1$ and $\operatorname{Cl}(N'_1) \subset \partial S$.

SUBSTEP 1.1. Let us construct recursively:

- Pairwise disjoint semialgebraic sets $N'_k \subset \partial S$ that are either Nash manifolds of dimension d-k, whose connected components are contained in either \mathfrak{T}_1 or \mathfrak{T}_2 , or the empty set,
- Real algebraic sets Z_k of dimension $\leq d-k$,

such that :

- (1) $N'_k \subset Z_k$ is an open semialgebraic subset of Z_k , (2) Z_k is the Zariski closure of $\partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1})$,

(3) $N'_k \cap Z_{k+1} = \emptyset$ and $Z_{k+1} \subset Z_k$, (4) $\partial S \setminus (N'_1 \cup \cdots \cup N'_k) \subset Z_{k+1}$ and $\operatorname{Cl}(N'_k) \subset \partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1})$.

In addition, $\partial S = N'_1 \sqcup \cdots \sqcup N'_d$ and Z_k is the Zariski closure of $N'_k \sqcup \cdots \sqcup N'_d$ for $k = 1, \ldots, d$.

Suppose that we have constructed the Nash manifolds N'_1, \ldots, N'_{k-1} and the real algebraic sets Z_1, \ldots, Z_k satisfying the required conditions and let us construct N'_k and Z_{k+1} . Let N_k be the union of the connected components of dimension d - k of $\operatorname{Reg}(\mathfrak{T}_1 \cap Z_k) \sqcup \operatorname{Reg}(\mathfrak{T}_2 \cap Z_k)$ or the empty set if dim $((Z_k \cap Cl(S)) \setminus S) < d - k$. Observe that N_k is an open semialgebraic subset of Z_k . As dim $((\mathcal{T}_i \cap Z_k) \setminus \text{Reg}(\mathcal{T}_i \cap Z_k)) \leq d - (k+1)$, the semialgebraic set $\partial S \setminus (N'_1 \cup \cdots \cup \mathcal{T}_k)$ $N'_{k-1} \cup N_k$) $\subset (\partial S \cap Z_k) \setminus N_k$ has dimension $\leq d - (k+1)$. Let Z_{k+1} be the Zariski closure of $\partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1} \cup N_k)$, which has dimension $\leq d - (k+1)$.

In case $N_k = \emptyset$, then dim $(Z_k) < d - k$ and $Z_{k+1} = Z_k$. Suppose $N_k \neq \emptyset$. Each connected component of the Nash manifold $N'_k := \partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1} \cup Z_{k+1}) = N_k \setminus Z_{k+1}$ has dimension d - k and it is contained in either \mathcal{T}_1 or \mathcal{T}_2 . In addition, N'_k is an open semialgebraic subset of $Z_k, \partial S \setminus (N'_1 \cup \cdots \cup N'_k) \subset Z_{k+1}$ (so it has dimension $\leq d - (k+1)$) and Z_{k+1} is the Zariksi closure of $\partial S \setminus (N'_1 \cup \cdots \cup N'_k)$ (because $\partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1} \cup N_k) \subset \partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1} \cup N_k) \subset Z_{k+1}$ and Z_{k+1} is the Zariski closure of $\partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1} \cup N_k)$).

As $Z_k \subset Z_\ell$ if $\ell < k$ and $N'_\ell \cap Z_{\ell+1} = \emptyset$, we deduce $N'_\ell \cap Z_k = \emptyset$ if $\ell < k$ and $\operatorname{Cl}(N'_k) \subset Z_k$ does not meet $N'_1 \cup \cdots \cup N'_{k-1}$, so $\operatorname{Cl}(N'_k) \subset \partial S \setminus (N'_1 \cup \cdots \cup N'_{k-1})$. Thus, we have constructed the Nash manifolds N'_k of dimension d - k (or the empty set) and the real algebraic sets Z_k of dimension $\leq d - k$ satisfying the required conditions for $k = 1, \ldots, d - 1$. In particular,

$$\operatorname{Cl}(N'_k) \setminus N'_k \subset \partial S \setminus (N'_1 \cup \cdots \cup N'_k) \subset Z_{k+1}$$

for k = 1, ..., d (and $Z_{d+1} := \emptyset$).

SUBSTEP 1.2. For technical reasons we reorder the indices of the real algebraic sets Z_k and the Nash manifolds N'_k .

Define the real algebraic set $T_k := Z_{d-k}$, which has dimension $\leq k$, and the Nash manifold $M_k := N'_{d-k}$, which is either empty or has dimension k, for $k = 0, \ldots, d-1$. The real algebraic set T_k is the Zariski closure of $M_1 \sqcup \cdots \sqcup M_k$ and each connected component of M_k is either contained in \mathfrak{T}_1 or \mathfrak{T}_2 .

If $M_0 \neq \emptyset$, then M_0 is a finite set and $M_0 = T_0$. Otherwise, if m is the least k such that $M_k \neq \emptyset$, then $\operatorname{Cl}(M_m) \setminus M_m = \emptyset$ (so M_m is closed in \mathbb{R}^n) and T_m is the Zariski closure of M_m . As M_m is open and closed in T_m , we deduce that M_m is a (finite) union of connected components of T_m of dimension m and $M_k = \emptyset$ for $0 \leq k < m$.

Observe that $\dim(T_k) \leq d-k$, $T_k \subset T_{k+1}$, $M_k \subset T_k$, $M_k \cap T_{k-1} = \emptyset$, $\operatorname{Cl}(M_k) \setminus M_k \subset M_0 \sqcup \cdots \sqcup M_{k-1} \subset T_{k-1}$. In addition,

$$\partial S \cap T_k = N'_{d-k} \sqcup \cdots \sqcup N'_d = M_0 \sqcup \cdots \sqcup M_k,$$

$$\mathfrak{T}_1 \cap T_k = (\operatorname{Cl}(S) \setminus S) \cap T_k = (M_0 \cap \mathfrak{T}_1) \sqcup \cdots \sqcup (M_k \cap \mathfrak{T}_1),$$

$$\mathfrak{T}_2 \cap T_k = (S \setminus \operatorname{Reg}(S)) \cap T_k = (M_0 \cap \mathfrak{T}_2) \sqcup \cdots \sqcup (M_k \cap \mathfrak{T}_2),$$

and each intersection $M_{\ell} \cap \mathfrak{T}_i$ is the union of the connected components of the Nash manifold M_{ℓ} contained in \mathfrak{T}_i for i = 1, 2. Observe that $M_{\ell} = (M_{\ell} \cap \mathfrak{T}_1) \sqcup (M_{\ell} \cap \mathfrak{T}_2)$ for $\ell = 0, \ldots, d-1$. STEP 2. INITIAL ALGEBRAIC RESOLUTION PROCEDURE. Denote $E_{-1} := \emptyset, X_{-1} := X, T_{-1,i} := T_i$ and $Y_0 := T_0$. By Theorem 1.3 there exists a non-singular (compact) real algebraic set X_0 and a proper surjective polynomial map $g_0 : X_0 \to X_{-1}$ such that $E_0 := g_1^{-1}(E_{-1} \cup Y_0)$ is a normal-crossings divisor of X_0 and the restriction $g_0|_{X_0 \setminus E_0} : X_0 \setminus E_0 \to X \setminus (E_{-1} \cup Y_0)$ is a Nash diffeomorphism whose inverse map is a regular map. In fact, g_0 is a composition of finitely many blow-ups whose non-singular centers have dimension $\leq \min\{\dim(Y_0), d-2\}$. Denote $T_{0i} := g_0^{-1}(T_{-1,i}) \cap \operatorname{Cl}(g_0^{-1}(T_{-1,i} \setminus (E_{-1} \cup Y_0)))$ the strict transform of $T_{-1,i}$ under g_0 , which is a real algebraic set of the same dimension as T_i and denote $Y_1 := T_{01}$. Observe that no T_{0i} is contained in the real algebraic set E_0 for $i \geq 1$. In particular, dim $(T_{0i} \cap E_0) < \dim(T_{0i})$ for each $i \geq 1$ and no irreducible component of Y_1 is contained in E_0 . We desingularize next $E_0 \cup Y_1$.

We proceed recursively and in the step $k \leq d-1$ we find by Theorem 1.3 a non-singular (compact) real algebraic set X_k and a proper surjective polynomial map $g_k : X_k \to X_{k-1}$ such that $E_k := g_k^{-1}(E_{k-1} \cup Y_k)$ is a normal-crossings divisor of X_k and the restriction $g_k|_{X_k \setminus E_k} :$ $X_k \setminus E_k \to X_{k-1} \setminus (E_{k-1} \cup Y_k)$ is a Nash diffeomorphism whose inverse map is a regular map. In fact, g_k is a composition of finitely many blow-ups whose non-singular centers have dimension $\leq \min\{\dim(Y_k), d-2\}$. Denote

$$T_{ki} := g_k^{-1}(T_{k-1,i}) \cap \operatorname{Cl}(g_k^{-1}(T_{k-1,i} \setminus (E_{k-1} \cup Y_k)))$$

the strict transform of $T_{k-1,i}$ under g_k , which is a real algebraic set of the same dimension as T_i and let $Y_{k+1} := T_{k,k+1}$. Observe that no T_{ki} is contained in the real algebraic set E_k for $i \ge k+1$. In particular, dim $(T_{ki} \cap E_k) < \dim(T_{ki})$ for each $i \ge k+1$ and no irreducible component of Y_{k+1} is contained in E_k . Observe that $Y_d = \emptyset$ (because $T_d = \emptyset$, so $T_{d-1,d} = \emptyset$) and $E_{d-1} = (g_0 \circ \cdots \circ g_{d-1})^{-1}(T_{d-1})$ is a normal-crossings divisor.

STEP 3. PROPERTIES OF THE STRICT TRANSFORM. We begin studying some properties of the composition $g := g_0 \circ \cdots \circ g_{d-1} : X_{d-1} \to X$. We claim: There exists a real algebraic subset $R \subset X$ of dimension $\leq d-2$ such that the restriction $g|_{X_{d-1}\setminus g^{-1}(R)} : X_{d-1}\setminus g^{-1}(R) \to X\setminus R$ is a Nash diffeomorphism.

For each $k = 0, \ldots, d-1$ the polynomial map $g_k : X_k \to X_{k-1}$ (where $X_{-1} := X$) is the composition of finitely many blow-ups whose centers have dimensions $\leq d-2$. Recall that the blow-up $b : \hat{V} \to V$ of a *d*-dimensional non-singular real algebraic set V with center a non-singular real algebraic subset W of dimension $\leq d-2$ provides a Nash diffeomorphism $b|_{\hat{V}\setminus b^{-1}(W)}: \hat{V}\setminus b^{-1}(W) \to V\setminus W$. As the image of a semialgebraic set of dimension $\leq d-2$ is a semialgebraic set of dimension $\leq d-2$, we conclude recursively that there exists a semialgebraic set $R_{k-1} \subset X_{k-1}$ of dimension $\leq d-2$ such that $g_k|_{X_k\setminus g_k^{-1}(R_{k-1})}: X_k\setminus g_k^{-1}(R_{k-1}) \to X_{k-1}\setminus R_{k-1}$ is a Nash diffeomorphism and $(g_0 \circ \cdots \circ g_{k-1})(R_{k-1})$ has dimension $\leq d-2$.

Let R be the Zariski closure of $R_{-1} \cup \bigcup_{k=0}^{d-2} (g_0 \circ \cdots \circ g_k)(R_k)$, which is a real algebraic set of dimension $\leq d-2$. The restriction $g|_{X_{d-1}\setminus g^{-1}(R)} : X_{d-1} \setminus g^{-1}(R) \to X \setminus R$ is a Nash diffeomorphism, as claimed.

SUBSTEP 3.1. Define $S^* := g^{-1}(S) \cap \operatorname{Cl}(g^{-1}(S) \setminus E_{d-1})$ the strict transform of S under g and $S_k := g_k^{-1}(S_{k-1}) \cap \operatorname{Cl}(g_k^{-1}(S_{k-1}) \setminus E_k)$ the strict transform of S_{k-1} under g_k for $k = 0, \ldots, d-1$, where $S_{-1} := S$. We claim: $S^* = S_{d-1}$. Let us prove by induction on ℓ that: $S_{\ell}^* := (g_0 \circ \cdots \circ g_{\ell})^{-1}(S) \cap \operatorname{Cl}((g_0 \circ \cdots \circ g_{\ell})^{-1}(S) \setminus E_{\ell})$ equals S_{ℓ} for each $\ell = 0, \ldots, d-1$.

If $\ell = 0$, then $S_0^* = S_0$. Suppose the result true for $\ell - 1$, that is, $S_{\ell-1}^* = S_{\ell-1}$ and let us check $S_{\ell}^* = S_{\ell}$. Denote $h_{\ell-1} := g_0 \circ \cdots \circ g_{\ell-1}$. We have

$$S_{\ell}^{*} = (g_{0} \circ \dots \circ g_{\ell})^{-1}(S) \cap \operatorname{Cl}((g_{0} \circ \dots \circ g_{\ell})^{-1}(S) \setminus E_{\ell}) = g_{\ell}^{-1}(h_{\ell-1}^{-1}(S)) \cap \operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(S)) \setminus E_{\ell})$$

is the strict transform of $h_{\ell-1}^{-1}(S)$ under g_{ℓ} . It holds $S_{\ell-1}^* = h_{\ell-1}^{-1}(S) \cap \operatorname{Cl}(h_{\ell-1}^{-1}(S) \setminus E_{\ell-1})$ and recall that $E_{\ell} = g_{\ell}^{-1}(E_{\ell-1} \cup Y_{\ell})$.

The strict transform of $S^*_{\ell-1}$ under g_{ℓ} is

$$g_{\ell}^{-1}(\mathbb{S}_{\ell-1}^{*}) \cap \operatorname{Cl}(g_{\ell}^{-1}(\mathbb{S}_{\ell-1}^{*}) \setminus E_{\ell}) = g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \cap g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})) \cap \operatorname{Cl}((g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}) \cap (g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1}) \setminus E_{\ell})).$$

As
$$h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1} \subset \operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})$$
 and $g_{\ell}^{-1}(E_{\ell-1}) \subset E_{\ell}$, we deduce
 $g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell} \subset g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})).$

Consequently,

$$\operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}) \subset g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})),$$
$$g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell} \subset g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})) \setminus E_{\ell}.$$

Thus,

$$\operatorname{Cl}((g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}) \cap (g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1}) \setminus E_{\ell})) = \operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}),$$

$$\operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}) \cap g_{\ell}^{-1}(\operatorname{Cl}(h_{\ell-1}^{-1}(\mathbb{S}) \setminus E_{\ell-1})) = \operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}).$$

We conclude

$$g_{\ell}^{-1}(\mathbb{S}_{\ell-1}^*) \cap \operatorname{Cl}(g_{\ell}^{-1}(\mathbb{S}_{\ell-1}^*) \setminus E_{\ell}) = g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \cap \operatorname{Cl}(g_{\ell}^{-1}(h_{\ell-1}^{-1}(\mathbb{S})) \setminus E_{\ell}) = \mathbb{S}_{\ell}^*.$$

As by induction hypothesis $S_{\ell-1}^* = S_{\ell-1}$, we have $S_{\ell}^* = g_{\ell}^{-1}(S_{\ell-1}) \cap \operatorname{Cl}(g_{\ell}^{-1}(S_{\ell-1}) \setminus E_{\ell}) = S_{\ell}$, as claimed.

SUBSTEP 3.2. We prove next: $\operatorname{Reg}(S^*)$ is a connected d-dimensional Nash manifold and it contains the inverse image $g^{-1}(\operatorname{Reg}(S) \setminus R)$ as a connected dense open semialgebraic subset.

As Reg(S) is a connected Nash manifold and dim $(R) \leq d-2$, also Reg(S) $\setminus R$ is a connected Nash manifold. As the restriction $g|_{X_{d-1}\setminus g^{-1}(R)} : X_{d-1}\setminus g^{-1}(R) \to X\setminus R$ is a Nash diffeomorphism, $g^{-1}(\operatorname{Reg}(S)\setminus R)$ is a connected Nash manifold. As dim $(R) \leq d-2$, S is pure dimensional of dimension d and Reg(S) is dense in S, we deduce Reg(S) $\setminus R$ is a dense open semialgebraic subset of S, so $g^{-1}(\operatorname{Reg}(S)\setminus R)$ is a dense open semialgebraic subset of $g^{-1}(S\setminus R)$.

As dim $(E_{d-1}) = d-1$, $g^{-1}(R) \subset E_{d-1}$ and $g^{-1}(\mathbb{S} \setminus R)$ is pure dimensional of dimension d, we deduce that $g^{-1}(\operatorname{Reg}(\mathbb{S})) \setminus E_{d-1} = g^{-1}(\operatorname{Reg}(\mathbb{S}) \setminus R) \setminus E_{d-1}$ is a dense open semialgebraic subset of $g^{-1}(\mathbb{S}) \setminus E_{d-1}$, so

$$\operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S}) \setminus R) \setminus E_{d-1}) = \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})) \setminus E_{d-1}) = \operatorname{Cl}(g^{-1}(\mathfrak{S}) \setminus E_{d-1}).$$
(4.3)

As $g^{-1}(\operatorname{Reg}(S) \setminus R)$ is a *d*-dimensional Nash manifold and $\dim(E_{d-1}) = d - 1$, we deduce that $g^{-1}(\operatorname{Reg}(S) \setminus R) \setminus E_{d-1}$ is dense in $g^{-1}(\operatorname{Reg}(S) \setminus R)$, so

$$\operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})\setminus R)) = \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})\setminus R)\setminus E_{d-1}).$$

Consequently,

$$g^{-1}(\operatorname{Reg}(\mathfrak{S}) \setminus R) = g^{-1}(\operatorname{Reg}(\mathfrak{S}) \setminus R) \cap \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S}) \setminus R) \setminus E_{d-1})$$
$$\subset g^{-1}(\operatorname{Reg}(\mathfrak{S})) \cap \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})) \setminus E_{d-1}) = \mathfrak{S}^*.$$

As S is connected by analytic paths, we deduce by [Fe2, Lem.7.16] that also S^{*} is connected by analytic paths, so in particular S^{*} is pure dimensional (of dimension d). Thus,

$$S^* \setminus E_{d-1} = g^{-1}(S) \setminus E_{d-1}$$
 (4.4)

is a dense open semialgebraic subset of S^* . As $g^{-1}(\operatorname{Reg}(S) \setminus R)$ is a dense open semialgebraic subset of $g^{-1}(S \setminus R)$ and $g^{-1}(R) \subset E_{d-1}$, we deduce that $g^{-1}(\operatorname{Reg}(S) \setminus R)$ is a dense open semialgebraic subset of S^* . As $g^{-1}(\operatorname{Reg}(S) \setminus R)$ is a *d*-dimensional connected Nash manifold, $g^{-1}(\operatorname{Reg}(S) \setminus R) \subset \operatorname{Reg}(S^*)$, so $\operatorname{Reg}(S^*)$ is connected because it contains a dense connected subset.

SUBSTEP 3.3. Let us prove: $\partial S^* \subset E_{d-1}$.

It holds $\partial S = \operatorname{Cl}(S) \setminus \operatorname{Reg}(S) \subset \bigcup_{k=0}^{d-1} T_k$ and

$$g^{-1}(\partial \mathbb{S}) = g^{-1}(\operatorname{Cl}(\mathbb{S}) \setminus \operatorname{Reg}(\mathbb{S})) \subset \bigcup_{k=0}^{d-1} g^{-1}(T_k)$$

Recall that T_{ki} is the strict transform of $T_{k-1,i}$ under g_k for $i \ge k$, $Y_k := T_{k-1,k}$, $E_k = g_k^{-1}(E_{k-1} \cup Y_k)$ for $k \ge 1$ and $E_0 = g_0^{-1}(T_0)$. Thus,

$$(g_0 \circ \cdots \circ g_{k-1})^{-1}(T_k) \subset T_{k-1,k} \cup E_{k-1} \subset Y_k \cup E_{k-1},$$

so $(g_0 \circ \cdots \circ g_k)^{-1}(T_k) \subset g_k^{-1}(Y_k \cup E_{k-1}) = E_k$ and $g^{-1}(T_k) = (g_0 \circ \cdots \circ g_{d-1})^{-1}(T_k) \subset E_{d-1}$ for each $k = 0, \ldots, d-1$. Thus,

$$g^{-1}(\partial \mathbb{S}) = g^{-1}(\operatorname{Cl}(\mathbb{S}) \setminus \operatorname{Reg}(\mathbb{S})) \subset \bigcup_{k=0}^{d-1} g^{-1}(T_k) \subset E_{d-1}.$$
(4.5)

We deduce using that $g^{-1}(R) \subset E_{d-1}$ and $g^{-1}(\operatorname{Reg}(S) \setminus R) \subset \operatorname{Reg}(S^*)$

$$\operatorname{Cl}(\mathbb{S}^*) \setminus E_{d-1} \subset \operatorname{Cl}(g^{-1}(\mathbb{S})) \cap \operatorname{Cl}(g^{-1}(\mathbb{S}) \setminus E_{d-1}) \setminus E_{d-1}$$
$$= \operatorname{Cl}(g^{-1}(\mathbb{S})) \setminus E_{d-1} \subset g^{-1}(\operatorname{Cl}(\mathbb{S})) \setminus E_{d-1}$$
$$= (g^{-1}(\operatorname{Reg}(\mathbb{S})) \setminus E_{d-1}) \cup (g^{-1}(\partial \mathbb{S}) \setminus E_{d-1}))$$
$$= g^{-1}(\operatorname{Reg}(\mathbb{S})) \setminus E_{d-1} \subset g^{-1}(\operatorname{Reg}(\mathbb{S}) \setminus R) \subset \operatorname{Reg}(\mathbb{S}^*),$$

so $\partial S^* = \operatorname{Cl}(S^*) \setminus \operatorname{Reg}(S^*) \subset E_{d-1}$.

SUBSTEP 3.4. Let $\mathfrak{T}_1^* := \operatorname{Cl}(\mathfrak{S}^*) \setminus \mathfrak{S}^*$ and $\mathfrak{T}_2^* := \mathfrak{S}^* \setminus \operatorname{Reg}(\mathfrak{S}^*)$. Let us check: $g(\mathfrak{T}_i^*) \subset \mathfrak{T}_i$ for i = 1, 2.

Recall that by (4.3) and (4.5) we have

$$\operatorname{Cl}(g^{-1}(\mathfrak{S}) \setminus E_{d-1}) = \operatorname{Cl}(g^{-1}(\operatorname{Cl}(\mathfrak{S})) \setminus E_{d-1}) = \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})) \setminus E_{d-1}).$$
(4.6)

As S is connected by analytic paths, Cl(S) is also connected by analytic paths [Fe2, Lem.7.4]. Thus, the strict transform $Cl(S)^*$ of Cl(S) under g is connected by analytic paths [Fe2, Lem.7.16], so it is pure dimensional of dimension d. Thus,

$$\operatorname{Cl}(\mathfrak{S})^* \setminus E_{d-1} = g^{-1}(\operatorname{Cl}(\mathfrak{S})) \setminus E_{d-1} = g^{-1}(\mathfrak{S}) \setminus E_{d-1} \subset \mathfrak{S}^* \subset \operatorname{Cl}(\mathfrak{S}^*)$$

is a dense subset of $\operatorname{Cl}(S)^*$. As $\operatorname{Cl}(S)^*$ is a closed set that contains S^* , we conclude that $\operatorname{Cl}(S^*) = \operatorname{Cl}(S)^*$ is the strict transform under g of $\operatorname{Cl}(S)$. Consequently, $\mathfrak{T}_1^* = \operatorname{Cl}(S^*) \setminus S^* = g^{-1}(\operatorname{Cl}(S) \setminus S) \cap \operatorname{Cl}(g^{-1}(S) \setminus E_{d-1}) \subset g^{-1}(\mathfrak{T}_1)$, so $g(\mathfrak{T}_1^*) \subset \mathfrak{T}_1$.

In addition, the strict transform $\operatorname{Reg}(S)^*$ of $\operatorname{Reg}(S)$ under g is

$$g^{-1}(\operatorname{Reg}(\mathfrak{S})) \cap \operatorname{Cl}(g^{-1}(\operatorname{Reg}(\mathfrak{S})) \setminus E_{d-1}).$$

As $g^{-1}(\operatorname{Reg}(S))$ is pure dimensional of dimension d, because it is an open semialgebraic subset of X_{d-1} , and E_{d-1} has dimension d-1, we deduce that $g^{-1}(\operatorname{Reg}(S)) \setminus E_{d-1}$ is dense in $g^{-1}(\operatorname{Reg}(S))$, so $\operatorname{Reg}(S)^* = g^{-1}(\operatorname{Reg}(S))$ is an open semialgebraic subset of X_{d-1} . Thus, $\operatorname{Reg}(S)^* \subset \operatorname{Reg}(S^*)$ and by (4.6)

$$\mathfrak{T}_2^* = \mathfrak{S}^* \setminus \operatorname{Reg}(\mathfrak{S}^*) \subset \mathfrak{S}^* \setminus \operatorname{Reg}(\mathfrak{S})^* = g^{-1}(\mathfrak{S} \setminus \operatorname{Reg}(\mathfrak{S})) \cap \operatorname{Cl}(g^{-1}(\mathfrak{S}) \setminus E_{d-1}) \subset g^{-1}(\mathfrak{T}_2),$$

so
$$g(\mathfrak{I}_2^*) \subset \mathfrak{I}_2$$
.
Substant 3.5 Let 7

SUBSTEP 3.5. Let Z be the Zariski closure of $\partial S^* := \operatorname{Cl}(S^*) \setminus \operatorname{Reg}(S^*)$ in the non-singular (compact) real algebraic set $\overline{S^*}^{\operatorname{zar}}$. We claim: Z is a normal-crossings divisor of $X_{d-1} = \overline{S^*}^{\operatorname{zar}}$, S^* is a checkerboard set, $\mathcal{T}_1^* := \operatorname{Cl}(S^*) \setminus S^*$, $\mathcal{T}_2^* := S^* \setminus \operatorname{Reg}(S^*)$, $\mathcal{T}_3^* := \operatorname{Cl}(S^*) \setminus \operatorname{Reg}(\operatorname{Cl}(S^*))$ and $\mathcal{T}_4^* := \operatorname{Reg}(\operatorname{Cl}(S^*)) \setminus S^*$ are unions of elements of the stratification $\mathfrak{G}(Z)$.

We prove first: \mathfrak{T}_1^* and \mathfrak{T}_2^* are unions of elements of $\mathfrak{G}(E_{d-1})$.

Define $T_{-1} := \emptyset$. Let E_{d-1}^k be the Zariski closure of $g^{-1}(T_k \setminus T_{k-1})$ for $k = 0, \ldots, d-1$, which is the union of the irreducible components of $g^{-1}(T_k)$ that are not contained in $g^{-1}(T_{k-1})$. We have $E_{d-1} = \bigcup_{k=0}^{d-1} E_{d-1}^k$ and each irreducible component of E_{d-1} is an irreducible component of E_{d-1}^k for exactly one $k = 0, \ldots, d-1$. Conversely, each irreducible component of E_{d-1}^k is an irreducible component of E_{d-1} , because if H is an irreducible component of E_{d-1}^k , then $g(H) \subset T_k$ and $g(H) \not\subset T_\ell$ if $\ell < k$.

Let $W \in \mathfrak{G}(\partial S^*)$ and L be the Zariski closure of W. As W is a connected Nash manifold, L is an irreducible real algebraic set. Observe that L is an irreducible component of $\operatorname{Sing}_j(E_{d-1})$ for some $j \geq 1$ and W is a connected component of $\operatorname{Reg}(\operatorname{Sing}_j(E_{d-1}))$, so $W \in \mathfrak{G}(E_{d-1})$. Let $k \geq 0$ be such that $g(L) \subset T_k$, but $g(L) \not\subset T_{k-1}$. Then $L \subset E_{d-1}^{\ell}$ for $\ell \geq k$, but $L \not\subset \bigcup_{\ell=0}^{k-1} E_{d-1}^{\ell} =$ $g^{-1}(T_{k-1})$. Observe that W is a connected component of $L \setminus g^{-1}(T_{k-1})$, because

$$L \cap \operatorname{Sing}(\operatorname{Sing}_{i}(E_{d-1})) = L \cap g^{-1}(T_{k-1})$$

(recall that $L \subset E_{d-1}^{\ell}$ for $\ell \geq k$). Then $g(W) \subset \partial S \cap T_k \setminus T_{k-1} = M_k$ is connected. As each connected component of M_k is contained in either \mathfrak{T}_1 or \mathfrak{T}_2 , we deduce either $g(W) \subset \mathfrak{T}_1$ or $g(W) \subset \mathfrak{T}_2$. If $W \cap \mathfrak{T}_i^* \neq \emptyset$, then $W \subset \mathfrak{T}_i^*$, because otherwise also $W \cap \mathfrak{T}_j^* \neq \emptyset$ for $j \in \{1, 2\} \setminus \{i\}$ and g(W) meets \mathfrak{T}_1 and \mathfrak{T}_2 , which is a contradiction because g(W) is contained either in \mathfrak{T}_1 or \mathfrak{T}_2 . Thus, W is contained either in \mathfrak{T}_1^* or \mathfrak{T}_2^* . As $\partial S^* = \mathfrak{T}_1^* \sqcup \mathfrak{T}_2^*$ is covered by the elements of $\mathfrak{G}(\partial S^*) \subset \mathfrak{G}(E_{d-1})$, we conclude that both \mathfrak{T}_1^* and \mathfrak{T}_2^* are unions of elements of $\mathfrak{G}(E_{d-1})$.

We show next: \mathfrak{T}_3^* and \mathfrak{T}_4^* are unions of elements of $\mathfrak{G}(E_{d-1})$.

By (4.4) and (4.5)

$$S^* \setminus E_{d-1} = g^{-1}(S) \setminus E_{d-1} = g^{-1}(\operatorname{Reg}(S)) \setminus E_{d-1} = g^{-1}(\operatorname{Cl}(S)) \setminus E_{d-1} = \operatorname{Cl}(S^*) \setminus E_{d-1}$$

is an open and closed subset of the Nash manifold $X_{d-1} \setminus E_{d-1}$, so it is a union of connected components of the Nash manifold $X_{d-1} \setminus E_{d-1}$. Thus, $\operatorname{Cl}(\mathfrak{S}^*) \setminus E_{d-1} \subset \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))$, so $\mathfrak{T}_3^* :=$

 $\operatorname{Cl}(\mathbb{S}^*) \setminus \operatorname{Reg}(\operatorname{Cl}(\mathbb{S}^*))$ is a (d-1)-dimensional semialgebraic subset contained in E_{d-1} (because $\operatorname{Cl}(\mathbb{S}^*)$ is the closure of a finite union of connected components of $X_{d-1} \setminus E_{d-1}$). In fact, using (local) coordinates, one realizes that both \mathcal{T}_3^* and $\operatorname{Cl}(\mathbb{S}^*) \cap E_{d-1}$ are unions of elements of $\mathfrak{G}(E_{d-1})$ (see Examples 4.12(iii)). Thus,

$$\begin{aligned} \mathfrak{T}_1^* \cap \mathfrak{T}_3^* &= (\operatorname{Cl}(\mathfrak{S}^*) \setminus \mathfrak{S}^*) \cap (\operatorname{Cl}(\mathfrak{S}^*) \setminus \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))) = \operatorname{Cl}(\mathfrak{S}^*) \setminus (\mathfrak{S}^* \cup \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))) \\ &= (\operatorname{Cl}(\mathfrak{S}^*) \cap E_{d-1}) \setminus ((\mathfrak{S}^* \cup \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))) \cap E_{d-1}) \end{aligned}$$

is a union of elements of $\mathfrak{G}(E_{d-1})$, so $(\mathfrak{S}^* \cup \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))) \cap E_{d-1}$ is also a union of elements of $\mathfrak{G}(E_{d-1})$. As

$$\mathfrak{S}^* \cap E_{d-1} = (\operatorname{Cl}(\mathfrak{S}^*) \cap E_{d-1}) \setminus (\operatorname{Cl}(\mathfrak{S}^*) \setminus \mathfrak{S}^*) = (\operatorname{Cl}(\mathfrak{S}^*) \cap E_{d-1}) \setminus \mathfrak{T}_1^*$$

is a union of elements of $\mathfrak{G}(E_{d-1})$, we conclude

$$\begin{aligned} \mathfrak{T}_4^* &:= (\operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*)) \setminus \mathfrak{S}^*) = (\operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*)) \setminus \mathfrak{S}^*) \cap E_{d-1} \\ &= ((\mathfrak{S}^* \cup \operatorname{Reg}(\operatorname{Cl}(\mathfrak{S}^*))) \cap E_{d-1}) \setminus (\mathfrak{S}^* \cap E_{d-1}) \end{aligned}$$

is a union of elements of $\mathfrak{G}(E_{d-1})$.

CURRENT SITUATION (1). The Zariski closure of S^* is X_{d-1} , which is a non-singular (compact) real algebraic set, and the Zariski closure of ∂S^* is a union of irreducible components of E_{d-1} , which is a normal-crossings divisor of X_{d-1} . As S^* is the strict transform of S under g and S is pure dimensional of dimension d, the restriction $g|_{S^*}: S^* \to S$ is a proper surjective map. Take $\mathcal{R} := g(E_{d-1}) \cap S$, which has dimension $\leq d-1$ and observe that $g|_{S^* \setminus g^{-1}(\mathcal{R})} : S^* \setminus g^{-1}(\mathcal{R}) \to S \setminus \mathcal{R}$ is a Nash diffeomorphism, because $g|_{X_{d-1} \setminus E_{d-1}} : X_{d-1} \setminus E_{d-1} \to X \setminus T_{d-1}$ is a Nash diffeomorphism. As $S^* \setminus g^{-1}(\mathcal{R}) = S^* \setminus E_{d-1} \subset S^* \setminus \partial S^*$ is a Nash manifold, its image $S \setminus \mathcal{R}$ under $g|_{X_{d-1} \setminus E_{d-1}}$ is also a Nash manifold. It still remains to improve the construction to achieve that $\operatorname{Cl}(S^*)$ is a Nash manifold with corners.

At this point we assume that the initial situation is the one quoted concerning S^* . For the sake of simplicity we reset all the previous notations above to continue the proof.

STEP 4. FIRST DRILLING DESINGULARIZATION PROCEDURE. We assume: S is a checkerboard set (and Reg(S) is a connected Nash manifold), Cl(S) is compact, the Zariski closure X of S is a nonsingular (compact) real algebraic set, the Zariski closure Z of Cl(S) \Reg(S) is a normal-crossings divisor of X, the semialgebraic sets $\mathcal{T}_1 := Cl(S) \setminus S$, $\mathcal{T}_2 := S \setminus Reg(S)$, $\mathcal{T}_3 := Cl(S) \setminus Reg(Cl(S))$ and $\mathcal{T}_4 := Reg(Cl(S)) \setminus S$ are unions of elements of the stratification $\mathfrak{G}(Z)$.

By Theorem 1.4 applied to Cl(S) there exist:

- (i) A d-dimensional (compact) irreducible non-singular real algebraic set X' and a normalcrossings divisor $Z' \subset X'$.
- (ii) A connected Nash manifold with corners $Q \subset X'$ (which is a closed subset of X') whose boundary ∂Q has Z' as its Zariski closure.
- (iii) A polynomial map $g : \mathbb{R}^n \to \mathbb{R}^m$ such that the restriction $g|_{\mathbb{Q}} : \mathbb{Q} \to \mathrm{Cl}(\mathbb{S})$ is proper and $g(\mathbb{Q}) = \mathrm{Cl}(\mathbb{S})$.
- (iv) A closed semialgebraic set $\mathcal{R} \subset \operatorname{Cl}(\mathcal{S})$ of dimension strictly smaller than d such that $\operatorname{Cl}(\mathcal{S}) \setminus \mathcal{R}$ and $\mathcal{Q} \setminus g^{-1}(\mathcal{R})$ are Nash manifolds and the polynomial map $g|_{\mathcal{Q} \setminus g^{-1}(\mathcal{R})} : \mathcal{Q} \setminus g^{-1}(\mathcal{R}) \to \operatorname{Cl}(\mathcal{S}) \setminus \mathcal{R}$ is a Nash diffeomorphism.

In addition, as X is compact, also X' is compact.

Let $S^* := g^{-1}(S) \cap \operatorname{Cl}(g^{-1}(S) \setminus \mathcal{R})$ be the strict transform of S under g. We claim: The semialgebraic sets $\mathcal{T}_1^* := \operatorname{Cl}(S^*) \setminus S^*$, $\mathcal{T}_2^* := S^* \setminus \operatorname{Reg}(S^*)$, $\mathcal{T}_3^* := \operatorname{Cl}(S^*) \setminus \operatorname{Reg}(\operatorname{Cl}(S^*))$ and $\mathcal{T}_4^* := \operatorname{Reg}(\operatorname{Cl}(S^*)) \setminus S^*$ are unions of elements of the stratification $\mathfrak{G}(Z')$.

Using the properties of the drilling blow-up and especially Remarks 3.4, one deduces straightforwardly that the semialgebraic sets T_1^* , T_2^* , T_3^* and T_4^* are unions of elements of the stratification $\mathfrak{G}(Z')$. To that end, one can almost reproduce the procedure developed in STEP 3 taking into account the particularities of the proof of Theorem 1.4, when one applies it to Cl(\mathcal{S}). CURRENT SITUATION (2). If $\operatorname{Reg}(\operatorname{Cl}(S^*)) = \operatorname{Reg}(S^*)$, then $\mathcal{T}_3^* = \operatorname{Cl}(S^*) \setminus \operatorname{Reg}(S^*) = \partial S^*$ and $\mathcal{T}_4^* = \emptyset$. Thus, S^* is a quasi Nash manifold with corners and the proof is finished for this case. Consequently, to continue we suppose $\operatorname{Reg}(\operatorname{Cl}(S^*)) \neq \operatorname{Reg}(S^*)$. This means that $\mathcal{T}_4^* \neq \emptyset$, because otherwise $\operatorname{Reg}(\operatorname{Cl}(S^*)) \subset S^*$ and consequently $\operatorname{Reg}(\operatorname{Cl}(S^*)) = \operatorname{Reg}(S^*)$ (because $\operatorname{Reg}(\operatorname{Cl}(S^*))$ is an open semialgebraic subset of X' contained in S^* that contains $\operatorname{Reg}(S^*)$).

At this point we assume that the initial situation is the one quoted concerning S^* . For the sake of simplicity we reset all the previous notations above to continue the proof.

STEP 5. SECOND DRILLING DESINGULARIZATION PROCEDURE. We assume in the following: S is a checkerboard set (and in particular Reg(S) is a connected Nash manifold), $\Omega := Cl(S)$ is a compact Nash manifold with corners, the Zariski closure X of S is a non-singular (compact) real algebraic set, the Zariski closure Z of $\Omega \setminus Reg(S)$ is a normal-crossings divisor of X, the semialgebraic sets $\mathcal{T}_1 := \Omega \setminus S$, $\mathcal{T}_2 := S \setminus Reg(S)$, $\mathcal{T}_3 := \Omega \setminus Reg(\Omega)$ and $\mathcal{T}_4 := Reg(\Omega) \setminus S \neq \emptyset$ are unions of elements of the stratification $\mathfrak{G}(Z)$. Let us prove: We may assume in addition $Reg(\mathcal{T}_4)$ is a pure dimensional semialgebraic set of dimension d - 1, $\partial \mathcal{T}_4 = Cl(\mathcal{T}_4) \setminus Reg(\mathcal{T}_4) \subset \partial \Omega$ and $\mathcal{T}_4 = Reg(\mathcal{T}_4)$.

As \mathfrak{T}_4 is a union of elements of the stratification $\mathfrak{G}(Z)$, the semialgebraic set $\operatorname{Cl}(\mathfrak{T}_4)$ is a union of elements of the stratification $\mathfrak{G}(Z)$. If \mathfrak{T}_4 has dimension $\leq d-2$, it is contained in $\operatorname{Sing}(Z)$ and $\emptyset \neq \mathfrak{T}_4 \subset \operatorname{Cl}(\mathfrak{T}_4) \cap \operatorname{Reg}(\mathfrak{Q})$. If \mathfrak{T}_4 has dimension d-1, define \mathcal{M}_4 as the union of the connected components of $\operatorname{Reg}(\mathfrak{T}_4)$ of dimension d-1. Observe that both \mathcal{M}_4 and $\operatorname{Cl}(\mathfrak{T}_4) \setminus \mathcal{M}_4$ are unions of elements of the stratification $\mathfrak{G}(Z)$. Define

$$\mathcal{A}(\mathcal{S}) := \begin{cases} \operatorname{Cl}(\mathcal{T}_4) \cap \operatorname{Reg}(\mathcal{Q}) & \text{if } \dim(\mathcal{T}_4) \leq d-2, \\ (\operatorname{Cl}(\mathcal{T}_4) \setminus \mathcal{M}_4) \cap \operatorname{Reg}(\mathcal{Q}) & \text{if } \dim(\mathcal{T}_4) = d-1. \end{cases}$$

Observe that $\dim(\mathcal{A}(S)) \leq d-2$ and $\mathcal{A}(S) = \emptyset$ if and only if $\dim(\mathfrak{T}_4) = d-1$ and $\operatorname{Cl}(\mathfrak{T}_4) \setminus \mathfrak{M}_4 \subset \partial \mathfrak{Q}$. As $\mathfrak{T}_4 = \operatorname{Reg}(\mathfrak{Q}) \setminus S$, this means that $\operatorname{Reg}(\mathfrak{T}_4) = \mathfrak{M}_4$ is a pure dimensional semialgebraic set of dimension d-1. We next develop an inductive procedure to reduce to the latter case.

Let Y be the Zariski closure of $\mathcal{A}(S)$, which is a union of irreducible components of $\operatorname{Sing}_{\ell}(Z)$ for $\ell = 1, \ldots, d-2$, maybe of different dimensions and denote $\operatorname{Sing}_0(Y) := Y$. Let e be the dimension of Y and Y_k the union of $\operatorname{Sing}_{e-k}(Y)$, which is either the empty set or a real algebraic set of dimension k, and the irreducible components of Y of dimension k for $k = 0, \ldots, e$.

If $\mathcal{A}(S) = \emptyset$, define $\ell(S) = d-1$ and m(S) = 0. If $\mathcal{A}(S) \neq \emptyset$, let $\ell := \ell(S) \leq e$ be the minimum value k such that $Y_k \neq \emptyset$ and m := m(S) the number of irreducible components of Y_ℓ . Observe that Y_ℓ is a pure dimensional non-singular (compact) real algebraic set of dimension $\leq d-2$. We proceed by double induction on ℓ and m.

Let W be an irreducible component of Y_{ℓ} . Let $(\widehat{X}, \widehat{\pi})$ be the twisted Nash double of the drilling blow-up (\widetilde{X}, π_+) of X with center W, which is by Section 3.4 a real algebraic set. Let

$$Q^* := \pi_+^{-1}(Q) \cap \operatorname{Cl}(\pi_+^{-1}(Q \setminus W))$$

be the strict transform of Ω under π_+ . As Ω is pure dimensional and $Y_{\ell} \subset Z$ has dimension strictly smaller, $\Omega \setminus W$ is dense in Ω , so $\pi_+(\Omega^*) = \Omega$, because $\pi_+ : \widetilde{X} \to X$ is proper and surjective. By Lemma 4.9 Ω^* is a checkerboard set and a Nash manifold with corners such that $\pi_+^{-1}(W) \cap \Omega^* \subset \partial \Omega^*$. Let $S^* := \pi_+^{-1}(S) \cap \operatorname{Cl}(\pi_+^{-1}(S \setminus W))$ be the strict transform of S^* under π_+ , which keeps the same properties required to S (to check this fact one proceeds similarly as we have done in STEPS 3 and 4). As $\pi_+^{-1}(W) \cap \Omega^* \subset \partial \Omega^*$, we deduce $\mathcal{A}(S^*) = \pi_+^{-1}(\mathcal{A}(S) \setminus W)$, so $m(S^*) = m(S) - 1$ and

$$\ell(\mathbb{S}^*) \begin{cases} > \ell(\mathbb{S}) & \text{if } m(\mathbb{S}) = 1, \\ = \ell(\mathbb{S}) & \text{if } m(\mathbb{S}) > 1. \end{cases}$$

The restriction $\pi_+|_{\mathbb{S}^*} : \mathbb{S}^* \to \mathbb{S}$ is a surjective proper polynomial map and if $\mathcal{R} := \partial \mathbb{S} \cup W$, the restriction $\pi_+|_{\mathbb{S}^*\setminus\pi_+^{-1}(\mathcal{R})} : \mathbb{S}^*\setminus\pi_+^{-1}(\mathcal{R}) \to \mathbb{S}\setminus\mathcal{R}$ is a Nash diffeomorphism.

We proceed inductively and after finitely many steps we may assume $\mathcal{A}(S) = \emptyset$. Under such assumption, we have: $\operatorname{Reg}(\mathcal{T}_4) = \mathcal{T}_4$.

As $\mathfrak{T}_4 = \operatorname{Reg}(\mathfrak{Q}) \setminus \mathfrak{S} \subset \operatorname{Reg}(\mathfrak{Q})$ and $\partial \mathfrak{T}_4 = \operatorname{Cl}(\mathfrak{T}_4) \setminus \operatorname{Reg}(\mathfrak{T}_4) \subset \partial \mathfrak{Q}$, we deduce $\mathfrak{T}_4 \cap \partial \mathfrak{T}_4 \subset \operatorname{Reg}(\mathfrak{Q}) \cap \partial \mathfrak{Q} = \emptyset$, so $\mathfrak{T}_4 = \operatorname{Reg}(\mathfrak{T}_4)$.

STEP 6. FINAL DRILLING DESINGULARIZATION PROCEDURE. After resetting notations, we assume in the following: S is a checkerboard set (and in particular Reg(S) is a connected Nash manifold), Q := Cl(S) is a compact Nash manifold with corners, the Zariski closure X of S is a non-singular (compact) real algebraic set, the Zariski closure Z of $Q \setminus Reg(S)$ is a normal-crossings divisor of X, the semialgebraic sets $\mathcal{T}_1 := Q \setminus S$, $\mathcal{T}_2 := S \setminus Reg(S)$, $\mathcal{T}_3 := Q \setminus Reg(Q)$, and $\mathcal{T}_4 := Reg(Q) \setminus S \neq \emptyset$ are unions of elements of the stratification $\mathfrak{G}(Z)$. In addition, $\mathcal{T}_4 = Reg(\mathcal{T}_4)$ is a pure dimensional semialgebraic set of dimension d - 1 and $\partial \mathcal{T}_4 = Cl(\mathcal{T}_4) \setminus Reg(\mathcal{T}_4) \subset \partial Q$.

In order to finish the proof, we will take advantage of Fact 3.5. Except for the initial embedding of \mathbb{RP}^m in \mathbb{R}^p , which is a regular map, until this step all the involved maps are polynomial. As we will perform the drilling blow-up of a Nash submanifold of dimension d-1, we have to proceed carefully in order to not disconnect the regular locus of S (Example 4.14). Thus, in the following the involved maps are a priori only Nash maps.

Let C_1, \ldots, C_m be the connected components of \mathcal{T}_4 . Each intersection $\operatorname{Cl}(C_i) \cap C_j = \emptyset$ for $i \neq j$, because $\operatorname{Cl}(C_i) \cap C_j = \operatorname{Cl}(C_i) \cap \mathcal{T}_4 \cap C_j = C_i \cap C_j = \emptyset$ (we have used that the connected components of \mathcal{T}_4 are pairwise disjoint closed subsets of \mathcal{T}_4). As $\operatorname{Reg}(C_i) = C_i$, we have

$$\partial \mathfrak{T}_4 = \operatorname{Cl}(\mathfrak{T}_4) \setminus \operatorname{Reg}(\mathfrak{T}_4) = \operatorname{Cl}(\mathfrak{T}_4) \setminus \mathfrak{T}_4 = \operatorname{Cl}\left(\bigcup_{i=1}^m C_i\right) \setminus \bigcup_{i=1}^m C_i = \bigcup_{i=1}^m \operatorname{Cl}(C_i) \setminus C_i = \bigcup_{i=1}^m \partial C_i.$$
(4.7)

Consequently, $\partial C_i \subset \partial \mathfrak{T}_4 \subset \partial \mathfrak{Q}$ for $i = 1, \ldots, m$. As $\operatorname{Reg}(\mathfrak{T}_4) = \mathfrak{T}_4$ is a union of elements of the stratification $\mathfrak{G}(Z)$, each connected component C_i of $\operatorname{Reg}(\mathfrak{T}_4)$ is a union of elements of the stratification $\mathfrak{G}(Z)$.

SUBSTEP 6.1. We claim: $\operatorname{Cl}(C_i) \cap \operatorname{Cl}(C_j) = \emptyset$ if $i \neq j$.

Assume $\operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2) \neq \emptyset$. As $\operatorname{Cl}(C_i) \cap C_j = \emptyset$ if $i \neq j$, we deduce $\operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2) = (\operatorname{Cl}(C_1) \setminus C_1) \cap (\operatorname{Cl}(C_2) \setminus C_2) = \partial C_1 \cap \partial C_2 \subset \partial \Omega.$ (4.8)

Pick $x \in \operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2)$ and let $U \subset X$ be an open semialgebraic neighborhood of x such that $Z \cap U = \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\}$ in (local) coordinates. As $\mathfrak{G}(Z)$ is compatible with C_1, C_2 , which are connected components of the Nash manifold $\mathfrak{T}_4 = \operatorname{Reg}(\mathfrak{T}_4)$, there exist indices $1 \leq i, j \leq r$ such that $C_1 \cap U \subset \{\mathbf{x}_i = 0\}$ and $C_2 \cap U \subset \{\mathbf{x}_j = 0\}$. If i = j, we suppose $Q \cap U = \{\mathbf{x}_1 \geq 0, \dots, \mathbf{x}_s \geq 0\}$ for some $1 \leq s \leq r-1, C_1 \cap U \subset \{\mathbf{x}_r = 0\}$ and $C_2 \cap U \subset \{\mathbf{x}_r = 0\}$. As $\mathfrak{G}(Z)$ is compatible with C_1 and C_2 , we may assume

$$\{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_{s+1} \ge 0, \dots, \mathbf{x}_{r-2} \ge 0, \mathbf{x}_{r-1} \ge 0, \mathbf{x}_r = 0\} \subset \operatorname{Cl}(C_1) \cap U, \\ \{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_{s+1} *_{s+1} 0, \dots, \mathbf{x}_{r-2} *_{r-2} 0, \mathbf{x}_{r-1} *_{r-1} 0, \mathbf{x}_r = 0\} \subset \operatorname{Cl}(C_2) \cap U,$$

where $*_j \in \{\geq, \leq\}$ for j = s + 1, ..., r - 1. By (4.8)

 $\{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_{s+1} = 0, \dots, \mathbf{x}_r = 0\} \subset \operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2) \cap U$

$$\subset \partial \Omega \cap U = \bigcup_{i=1}^{\circ} \{ \mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_i = 0 \},$$

which is a contradiction. If $i \neq j$, we suppose $\Omega \cap U = \{\mathbf{x}_1 \geq 0, \dots, \mathbf{x}_s \geq 0\}$ for some $1 \leq s \leq r-2$, $C_1 \cap U \subset \{\mathbf{x}_{r-1} = 0\}$ and $C_2 \cap U \subset \{\mathbf{x}_r = 0\}$. As $\mathfrak{G}(Z)$ is compatible with C_1 and C_2 , we may assume

$$\{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_{s+1} \ge 0, \dots, \mathbf{x}_{r-2} \ge 0, \mathbf{x}_{r-1} = 0, \mathbf{x}_r \ge 0\} \subset \operatorname{Cl}(C_1) \cap U, \\ \{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_s \ge 0, \mathbf{x}_{s+1} *_{s+1} 0, \dots, \mathbf{x}_{r-2} *_{r-2} 0, \mathbf{x}_{r-1} \ge 0, \mathbf{x}_r = 0\} \subset \operatorname{Cl}(C_2) \cap U,$$

where $*_i \in \{\geq, \leq\}$ for $j = s + 1, \dots, r - 2$. By (4.8)

$$\begin{aligned} \{\mathtt{x}_1 \ge 0, \dots, \mathtt{x}_s \ge 0, \mathtt{x}_{s+1} = 0, \dots, \mathtt{x}_r = 0\} \subset \operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2) \cap U \\ \subset \partial \mathfrak{Q} \cap U = \bigcup_{i=1}^s \{\mathtt{x}_1 \ge 0, \dots, \mathtt{x}_s \ge 0, \mathtt{x}_i = 0\}, \end{aligned}$$

which is a contradiction. Consequently, $\operatorname{Cl}(C_1) \cap \operatorname{Cl}(C_2) = \emptyset$, as claimed.

SUBSTEP 6.2. Let Γ be a stratum of $\mathfrak{G}(Z)$ contained in \mathfrak{Q} such that Γ is not contained in $\operatorname{Cl}(C_i)$. We prove next: If the Zariski closure of Γ is contained in the Zariski closure of C_i , then $\operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) = \emptyset$.

As Γ is not contained in $\operatorname{Cl}(C_i)$ and the stratification $\mathfrak{G}(Z)$ is compatible with C_i , we have $\operatorname{Cl}(\Gamma) \cap C_i = \emptyset$, so

$$\operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) = \operatorname{Cl}(\Gamma) \cap (\operatorname{Cl}(C_i) \setminus C_i) = \operatorname{Cl}(\Gamma) \cap \partial C_i.$$

Suppose $\operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) \neq \emptyset$, pick $x \in \operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i)$ and let $U \subset X$ be an open semialgebraic neighborhood of x such that $Z \cap U = \{\mathbf{x}_1 \cdots \mathbf{x}_r = 0\}$ in (local) coordinates. We may assume $\mathcal{Q} \cap U = \{\mathbf{x}_1 \geq 0, \dots, \mathbf{x}_s \geq 0\}$ for some $1 \leq s \leq r-1$ and $C_i \cap U \subset \{\mathbf{x}_r = 0\}$. As the Zariski closure of Γ is contained in the Zariski closure of C_i and $x \in \operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) \cap U$, we deduce $\operatorname{Cl}(\Gamma) \cap U \subset \mathcal{Q} \cap U \cap \{\mathbf{x}_r = 0\}$. In addition, $\partial C_i \cap U \subset \partial \mathcal{Q} \cap U = \bigcup_{i=1}^s \{\mathbf{x}_1 \geq 0, \dots, \mathbf{x}_s \geq 0, \mathbf{x}_i = 0\}$. As $\mathfrak{G}(Z)$ is compatible with C_i and $\partial C_i \subset \partial \mathcal{Q}$ (see (4.7)), we conclude

$$C_{i} \cap U = \{\mathbf{x}_{1} > 0, \dots, \mathbf{x}_{s} > 0, \mathbf{x}_{r} = 0\} = \operatorname{Reg}(\mathfrak{Q}) \cap U \cap \{\mathbf{x}_{r} = 0\},\$$

$$\operatorname{Cl}(C_{i}) \cap U = \{\mathbf{x}_{1} \ge 0, \dots, \mathbf{x}_{s} \ge 0, \mathbf{x}_{r} = 0\} = \mathfrak{Q} \cap U \cap \{\mathbf{x}_{r} = 0\},\$$

so $\operatorname{Cl}(\Gamma) \cap U \subset \mathcal{Q} \cap U \cap \{\mathbf{x}_r = 0\} = \operatorname{Cl}(C_i) \cap U$, which is a contradiction. Thus, $\operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) = \emptyset$.

SUBSTEP 6.3. Recall that $\dim(C_i) = d - 1$ for each $i = 1, \ldots, m$. The Zariski closure of C_i is the irreducible component Z_i of Z that contains C_i . The semialgebraic set $Z_i \cap Q \setminus \operatorname{Cl}(C_i)$ is a union of elements of $\mathfrak{G}(Z)$ and it is closed, because otherwise there exists a stratum Γ of $\mathfrak{G}(Z)$ contained in Q such that $\Gamma \not\subset \operatorname{Cl}(C_i)$ but $\operatorname{Cl}(\Gamma) \cap \operatorname{Cl}(C_i) \neq \emptyset$, which is a contradiction by SUBSTEP 6.2. Consider the closed semialgebraic set $K_i := (Z_i \cap Q \setminus \operatorname{Cl}(C_i)) \cup \bigcup_{j \neq i} \operatorname{Cl}(C_j)$ and observe that $K_i \cap \operatorname{Cl}(C_i) = \emptyset$. As both semialgebraic sets are compact (recall that the Zariski closure of S is compact) and disjoint,

$$\varepsilon := \frac{1}{2} \min\{\operatorname{dist}(K_i, \operatorname{Cl}(C_i)): i = 1, \dots, m\} > 0.$$

Define $U_i := \{x \in Z_i : \text{dist}(x, \text{Cl}(C_i)) < \varepsilon\}$, which is an open semialgebraic neighborhood of $\text{Cl}(C_i)$ in Z_i . We claim:

- (1) The union $N := \bigcup_{i=1}^{m} U_i$ is a closed Nash submanifold of the Nash manifold $M_0 := X \setminus \bigcup_{i=1} (\operatorname{Cl}(U_i) \setminus U_i),$
- (2) $Q \subset M_0$ and
- (3) $\mathcal{Q} \cap N = \bigcup_{i=1}^{m} \operatorname{Cl}(C_i) = \operatorname{Cl}(\mathfrak{T}_4).$

We prove first (1). It is clear that N is a closed subset of M_0 . As each U_i is an open semialgebraic subset of the Nash manifold Z_i , to prove that $N \subset M_0$ is a Nash manifold, it is enough to show that $\operatorname{Cl}(U_i) \cap \operatorname{Cl}(U_j) = \emptyset$ if $i \neq j$. If there exists $x \in \operatorname{Cl}(U_i) \cap \operatorname{Cl}(U_j)$, then

$$\operatorname{dist}(\operatorname{Cl}(C_i), \operatorname{Cl}(C_j)) \leq \operatorname{dist}(x, \operatorname{Cl}(C_i)) + \operatorname{dist}(x, \operatorname{Cl}(C_j)) \\ < 2\varepsilon \leq \operatorname{dist}(K_i, \operatorname{Cl}(C_i)) \leq \operatorname{dist}(\operatorname{Cl}(C_i), \operatorname{Cl}(C_j)),$$

which is a contradiction. Consequently, the semialgebraic sets $Cl(U_i)$ for i = 1, ..., m are pairwise disjoint and N is a closed Nash submanifold of M_0 .

We check next (2): $Q \subset M_0$.

Suppose there exists $x \in Q \cap (Cl(U_i) \setminus U_i)$, so

 $x \in Z_i \cap \mathcal{Q} \cap (\mathrm{Cl}(U_i) \setminus U_i) \subset ((Z_i \cap \mathcal{Q}) \setminus \mathrm{Cl}(C_i)) \cap \mathrm{Cl}(U_i),$

because $\operatorname{Cl}(C_i) \subset U_i$. As $x \in \operatorname{Cl}(U_i)$ and $x \in (Z_i \cap \mathfrak{Q}) \setminus \operatorname{Cl}(C_i)$, we have

$$\operatorname{dist}(x,\operatorname{Cl}(C_i)) \le \varepsilon \le \frac{1}{2}\operatorname{dist}(K_i,\operatorname{Cl}(C_i)) \le \frac{1}{2}\operatorname{dist}(Z_i \cap \mathfrak{Q} \setminus \operatorname{Cl}(C_i),\operatorname{Cl}(C_i)) \le \frac{1}{2}\operatorname{dist}(x,\operatorname{Cl}(C_i)),$$

which is a contradiction. Consequently, $\Omega \cap (\operatorname{Cl}(U_i) \setminus U_i) = \emptyset$ for each $i = 1, \ldots, m$. Thus, $\Omega \subset X \setminus \bigcup_{i=1} (\operatorname{Cl}(U_i) \setminus U_i) = M_0$.

As $\operatorname{Cl}(\mathfrak{T}_4) = \bigcup_{i=1}^m \operatorname{Cl}(C_i) \subset \mathfrak{Q} \cap \bigcup_{i=1}^m U_i = \mathfrak{Q} \cap N$, to prove (3): $\mathfrak{Q} \cap N = \operatorname{Cl}(\mathfrak{T}_4)$, it is enough to check: $\mathfrak{Q} \cap U_i \subset \operatorname{Cl}(C_i)$ or, equivalently, $\mathfrak{Q} \cap (U_i \setminus \operatorname{Cl}(C_i)) = \emptyset$ for $i = 1, \ldots, m$.

If $x \in \Omega \cap (U_i \setminus \operatorname{Cl}(C_i)) \subset Z_i$, then $x \in U_i$ and $x \in Z_i \cap \Omega \setminus \operatorname{Cl}(C_i)$, which is a contradiction as we have seen when proving (2). Thus, $\Omega \cap U_i \subset \operatorname{Cl}(C_i)$ for $i = 1, \ldots, m$.

SUBSTEP 6.4. As $\partial \mathfrak{T}_4 \subset \partial \mathfrak{Q}$, we have $\operatorname{Cl}(\mathfrak{T}_4) \cap \operatorname{Reg}(\mathfrak{Q}) = \operatorname{Reg}(\mathfrak{T}_4) \cap \operatorname{Reg}(\mathfrak{Q})$, so $\operatorname{Cl}(\mathfrak{T}_4) \cap \operatorname{Reg}(\mathfrak{Q}) = \mathfrak{T}_4 \cap \operatorname{Reg}(\mathfrak{Q})$. As $\mathfrak{T}_4 = \operatorname{Reg}(\mathfrak{Q}) \setminus \mathfrak{S}$, the difference

$$\operatorname{Reg}(\mathfrak{Q}) \setminus \operatorname{Cl}(\mathfrak{T}_4) = \operatorname{Reg}(\mathfrak{Q}) \setminus \mathfrak{T}_4 = \mathfrak{S} \cap \operatorname{Reg}(\mathfrak{Q}) \subset \mathfrak{S}$$

is an open semialgebraic subset of X contained in S, so $S \cap \operatorname{Reg}(Q) \subset \operatorname{Reg}(S)$. As $\operatorname{Reg}(S) \subset S \cap \operatorname{Reg}(Q)$, we conclude $\operatorname{Reg}(S) = \operatorname{Reg}(Q) \setminus \operatorname{Cl}(\mathcal{T}_4) = \operatorname{Reg}(Q) \setminus (Q \cap N) = \operatorname{Reg}(Q) \setminus N$. In general, N is not a real algebraic set and its Zariski closure Y is not an option because $\operatorname{Reg}(Q) \setminus Y$ might be disconnected (Example 4.14). Thus, the following drilling blow-up is in general only a Nash map. Let $(\widehat{M}, \widehat{\pi})$ be the twisted Nash double of the drilling blow-up (\widetilde{M}, π_+) of M := X with center the closed Nash submanifold N of M, which is by Section 3.2 a Nash manifold. As $\widehat{\pi} : \widehat{M} \to M$ is proper and surjective and M is compact, also \widehat{M} is compact. We have denoted X by M in order to stress that \widehat{M} is a compact Nash manifold, which is not in general a non-singular real algebraic set (but only one of its compact connected components). Let $Q^{\bullet} := \pi_+^{-1}(Q) \cap \operatorname{Cl}(\pi_+^{-1}(Q \setminus N))$ be the strict transform of Q under π_+ . As Q is pure dimensional and $N \subset Z$ has dimension strictly smaller, $Q \setminus N$ is dense in Q, so $\pi_+(Q^{\bullet}) = Q$, because $\pi_+ : \widetilde{M} \to M$ is proper and surjective. By Remarks 3.4 and Fact 3.5 Q^{\bullet} is a Nash manifold with corners such that $\pi_+^{-1}(N) \cap Q^{\bullet} \subset \partial Q^{\bullet}$. Observe that $Q^{\bullet} \setminus \pi_+^{-1}(N) = \pi_+^{-1}(Q \setminus N)$ is Nash diffeomorphic to $Q \setminus N$. Thus,

$$\operatorname{Sth}(\mathfrak{Q}^{\bullet}) = \operatorname{Sth}(\pi_{+}^{-1}(\mathfrak{Q} \setminus N)) = \pi_{+}^{-1}(\operatorname{Reg}(\mathfrak{Q} \setminus N)) = \pi_{+}^{-1}(\operatorname{Reg}(\mathfrak{Q}) \setminus N) = \pi_{+}^{-1}(\operatorname{Reg}(\mathfrak{S})),$$

so $\operatorname{Sth}(\mathbb{Q}^{\bullet})$ is connected, because $\pi_+|_{\widetilde{M}\setminus\pi_+^{-1}(N)}: \widetilde{M}\setminus\pi_+^{-1}(N) \to M\setminus N$ is a Nash diffeomorphism. Let $\mathcal{S}^{\bullet} := \pi_+^{-1}(\mathcal{S}) \cap \operatorname{Cl}(\pi_+^{-1}(\mathcal{S}\setminus N))$ be the strict transform of \mathcal{S} under π_+ , which keeps the same properties required to \mathcal{S} if one changes the operator $\operatorname{Reg}(\cdot)$ by the operator $\operatorname{Sth}(\cdot)$ in each case. To check this fact one proceeds similarly as we have done in STEPS 3, 4 and 5. In addition,

$$\operatorname{Sth}(\mathfrak{S}^{\bullet}) \subset \operatorname{Sth}(\mathfrak{Q}^{\bullet}) = \pi_{+}^{-1}(\operatorname{Reg}(\mathfrak{S})) \subset \operatorname{Sth}(\mathfrak{S}^{\bullet})$$

(because $\operatorname{Reg}(S) \subset M \setminus N$), so $\operatorname{Sth}(S^{\bullet}) = \operatorname{Sth}(Q^{\bullet})$. Observe that $\pi_+|_{S^{\bullet}} : S^{\bullet} \to S$ is a surjective proper Nash map and if $\mathcal{R} := S \setminus \operatorname{Reg}(S)$, the restriction $\pi_+|_{S^{\bullet} \setminus \pi_+^{-1}(\mathcal{R})} : S^{\bullet} \setminus \pi_+^{-1}(\mathcal{R}) \to S \setminus \mathcal{R}$ is a Nash diffeomorphism.

SUBSTEP 6.5. To finish we shall 'algebrize' our construction as much as possible. Recall that by [AK, Thm.1.1] the pair constituted by a compact Nash manifold and a Nash normal-crossings divisor is diffeomorphic to a pair constituted by a non-singular (compact) real algebraic set and a normal-crossings divisor and the previous diffeomorphism preserves Nash irreducible components of the corresponding Nash normal-crossings divisor. By the proof of the approximation results [BFR, Thm.1.7 & Prop.8.2] modified to fit our situation ,we may assume that the previous diffeomorphism is in addition a Nash diffeomorphism. To that end, we have to substitute Efroymson's approximation result [Sh, Thm.II.4.1] for differentiable semialgebraic functions on a Nash manifold by Nash functions by Stone-Weierstrass approximation for differentiable functions on differentiable manifolds by polynomial functions (see also [GS, §6]).

Using the previous fact and [Fe2, Lem.8.3 & Lem.C.1], we may assume in addition (using a suitable Nash embedding of \widetilde{M} in some affine space) that the quasi Nash manifold with corners S^{\bullet} is a checkerboard set, the Nash manifold with corners $Q^{\bullet} = \operatorname{Cl}(S^{\bullet})$ is a checkerboard set, the Zariski closure X^{\bullet} of S^{\bullet} is a connected compact irreducible non-singular real algebraic set, the

Zariski closure Z^{\bullet} of $\partial S^{\bullet} = \mathcal{Q}^{\bullet} \setminus \operatorname{Reg}(S^{\bullet}) = \mathcal{Q}^{\bullet} \setminus \operatorname{Reg}(\mathcal{Q}^{\bullet}) = \partial \mathcal{Q}^{\bullet}$ is a normal-crossings divisor of X^{\bullet} and the stratification $\mathfrak{G}(Z^{\bullet})$ is compatible with $S^{\bullet} \setminus \operatorname{Reg}(S^{\bullet})$.




FIGURE 4.3. Desingularization of the checkerboard set S (right) by the quasi Nash manifold with corners S^{\bullet} (left).

Example 4.14. Let $X := \{\mathbf{x}_1^2 + \dots + \mathbf{x}_n^2 = 1\} \subset \mathbb{R}^n$ and $S := X \cap \{\mathbf{x}_n^2 \leq \frac{1}{4}\} \setminus \{\mathbf{x}_{n-2} \leq 0, \mathbf{x}_{n-1} = 0\}$, which is a checkerboard set whose Zariski closure is X. The real algebraic set X is the (n-1)-dimensional unit sphere, so it is compact and non-singular. The closure $\operatorname{Cl}(S) = X \cap \{\mathbf{x}_n^2 \leq \frac{1}{4}\}$ is a compact Nash manifold with corners. Observe that $\operatorname{Reg}(S) = S \cap \{\mathbf{x}_n^2 < \frac{1}{4}\}$, so

$$\operatorname{Cl}(\mathfrak{S}) \setminus \operatorname{Reg}(\mathfrak{S}) = (X \cap \{\mathfrak{x}_n^2 = \frac{1}{4}\}) \cup (X \cap \{\mathfrak{x}_{n-2} \le 0, \mathfrak{x}_{n-1} = 0\} \cap \{\mathfrak{x}_n^2 \le \frac{1}{4}\}).$$

The Zariski closure of $\operatorname{Cl}(\mathcal{S}) \setminus \operatorname{Reg}(\mathcal{S})$ is

$$Z := (X \cap \{\mathbf{x}_n = \frac{1}{2}\}) \cup (X \cap \{\mathbf{x}_n = -\frac{1}{2}\}) \cup (X \cap \{\mathbf{x}_{n-1} = 0\}),$$

which is a normal-crossings divisor of X. Denote Q := Cl(S). The semialgebraic sets

$$\begin{split} \mathfrak{T}_1 &:= \mathfrak{Q} \setminus \mathfrak{S} = X \cap \{ \mathtt{x}_{n-2} \leq 0, \mathtt{x}_{n-1} = 0 \} \cap \{ \mathtt{x}_n^2 \leq \frac{1}{4} \}, \\ \mathfrak{T}_2 &:= \mathfrak{S} \setminus \operatorname{Reg}(\mathfrak{S}) = X \cap \{ \mathtt{x}_n^2 = \frac{1}{4} \} \setminus \{ \mathtt{x}_{n-2} \leq 0, \mathtt{x}_{n-1} = 0 \}, \\ \mathfrak{T}_3 &:= \mathfrak{Q} \setminus \operatorname{Reg}(\mathfrak{Q}) = X \cap \{ \mathtt{x}_n^2 = \frac{1}{4} \}, \\ \mathfrak{T}_4 &:= \operatorname{Reg}(\mathfrak{Q}) \setminus \mathfrak{S} = X \cap \{ \mathtt{x}_{n-2} \leq 0, \mathtt{x}_{n-1} = 0 \} \setminus \{ \mathtt{x}_n^2 = \frac{1}{4} \} \neq \emptyset \end{split}$$

are unions of elements of the stratification $\mathfrak{G}(Z)$. In addition, $\operatorname{Reg}(\mathfrak{T}_4)$ is a pure dimensional semialgebraic set of dimension d-1 and

$$\partial \mathfrak{T}_4 = \operatorname{Cl}(\mathfrak{T}_4) \setminus \operatorname{Reg}(\mathfrak{T}_4) = X \cap \{ \mathbf{x}_n^2 = \frac{1}{4} \} \cap \{ \mathbf{x}_{n-2} \le 0, \mathbf{x}_{n-1} = 0 \} \subset \partial \mathfrak{Q}.$$

Thus, we are under the hypothesis of STEP 6 of the Proof of Theorem 1.6. We consider as $N := X \cap \{\mathbf{x}_{n-2} < 0, \mathbf{x}_{n-1} = 0\}$, but we cannot take its Zariski closure $Y := X \cap \{\mathbf{x}_{n-1} = 0\}$, because Y disconnects Reg(S).

5. Folding Nash manifolds

We proved in [FGR, Thm.1.12, Cor.6.5] that a Nash manifold $\Omega \subset \mathbb{R}^n$ with corners is contained as a closed subset in a Nash manifold $M \subset \mathbb{R}^n$ of the same dimension and the Nash closure of its boundary $\partial \Omega$ is a Nash normal-crossings divisor of M. In this section we show that the Nash manifold M can be 'folded' to build the Nash manifold with corners Ω , that is, there exists a surjective Nash map $f: M \to \Omega$ such that its restriction to Ω is a semialgebraic homeomorphism close to the identity, its restriction to $\operatorname{Int}(\Omega)$ is a Nash diffeomorphism close to the identity, and f preserves the natural semialgebraic partition of the boundary $\partial \Omega$ (Theorem 1.7). We use standard tools for smooth manifolds with boundary adapted to the Nash category (equations of the boundary, collars and doubling of smooth manifolds with boundary, etc.). These standard constructions in the Nash setting were mainly developed in [Sh, Ch.VI] (compact case) and in [Fe2, §4] (general case). However, we need to adapt some of the statements and their proofs to obtain constructions compatible with an assigned Nash normal-crossings divisor. We also show that our construction is canonical and does not depend on the order chosen to 'fold M along the facets of Ω' (§5.7). 5.1. Compatible Nash retractions. Let $M \subset \mathbb{R}^n$ be a connected *d*-dimensional Nash manifold and $Y \subset M$ a Nash normal-crossings divisor. Define inductively

$$\begin{split} \operatorname{Sing}_1(Y) &:= \operatorname{Sing}(Y), \\ \operatorname{Sing}_\ell(Y) &:= \operatorname{Sing}_{\ell-1}(\operatorname{Sing}(Y)) \quad \text{for } \ell \geq 2. \end{split}$$

In order to lighten the exposition, we write $\operatorname{Sing}_0(Y) := Y$ and $\operatorname{Sing}_{-1}(Y) := M$. The irreducible components of $\operatorname{Sing}_{\ell}(Y)$ are Nash manifolds for each $\ell \geq 1$ such that $\operatorname{Sing}_{\ell}(Y) \neq \emptyset$. In fact, if $Y_{\ell,1}, \ldots, Y_{\ell,s_{\ell}}$ are the irreducible components of $\operatorname{Sing}_{\ell}(Y)$, then $\operatorname{Sing}_{\ell+1}(Y) = \bigcup_{i \neq j} (Y_{\ell,i} \cap Y_{\ell,j})$. If $\operatorname{Sing}_{\ell}(Y) \neq \emptyset$, then $\dim(\operatorname{Sing}_{\ell}(Y)) = d - \ell - 1$.

For each $\ell \geq 1$ we have $\operatorname{Sing}_{\ell+1}(Y) \subset \operatorname{Sing}_{\ell}(Y)$ and $\operatorname{dim}(\operatorname{Sing}_{\ell+1}(Y)) \leq \operatorname{dim}(\operatorname{Sing}_{\ell}(Y)) - 1$, so there exists an $r \geq 0$ such that $\operatorname{Sing}_k(Y) \neq \emptyset$ for $1 \leq k \leq r$ and $\operatorname{Sing}_k(Y) = \emptyset$ if $k \geq r + 1$.

Definition 5.1. Let Z be an irreducible component of $\text{Sing}_k(Y)$ for some $1 \leq k \leq r$. A Nash retraction $\rho: W \to Z$, where $W \subset M$ is an open semialgebraic neighborhood of Z, is *compatible with* Y if $\rho(Y_i \cap W) = Y_i \cap Z$ for each irreducible component Y_i of Y such that $Y_i \cap Z \neq \emptyset$.

In [FGh1, Prop.4.1] the following result is proved, which is a powerful tool to make constructions compatible with an assigned Nash normal-crossings divisor.

Proposition 5.2 (Compatible Nash retractions, [FGh1, Prop. 4.1]). There exist an open semialgebraic neighborhood $W \subset M$ of Z and a Nash retraction $\rho : W \to Z$ that is compatible with Y. In addition, $\rho(X \cap W) = X \cap Z$ for each irreducible component X of $\operatorname{Sing}_{\ell}(Y)$ such that $X \cap Z \neq \emptyset$ and $\ell \geq 1$.

5.2. Compatible Nash collars for Nash manifolds with boundary. Nash collars for a Nash manifold $H \subset \mathbb{R}^n$ with boundary ∂H have been constructed in [Sh, Lem.VI.1.6] (compact case) and in [Fe2, Lem.4.2](general case). We adapt these constructions to provide Nash collars compatible with a Nash normal-crossings divisor Y that contains ∂H .

Lemma 5.3 (Compatible Nash collars). Let $M \subset \mathbb{R}^n$ be a Nash manifold and $Y \subset M$ a Nash normal-crossings divisor. Let Y_1 be an irreducible component of Y and $\rho : W \to Y_1$ a Nash retraction compatible with Y_1 such that $\rho(X \cap W) = X \cap Y_1$ for each irreducible component X of $\operatorname{Sing}_{\ell}(Y)$ such that $X \cap Y_1 \neq \emptyset$ and $\ell \geq 1$. Let h be a Nash function on W such that $\{h = 0\} = Y_1$ and $d_x h : T_x M \to \mathbb{R}$ is surjective for all $x \in Y_1$. Then there exist an open semialgebraic neighborhood $V \subset W$ of Y_1 and a strictly positive Nash function $\varepsilon : Y_1 \to \mathbb{R}$ such that:

- (i) The Nash map $\varphi := (\rho, \frac{h}{\varepsilon \circ \rho}) : V \to Y_1 \times (-1, 1)$ is a Nash diffeomorphism.
- (ii) $\varphi(Z \cap V) = (Z \cap Y_1) \times (-1, 1)$ for each irreducible component Z of $\operatorname{Sing}_{\ell}(Y)$ such that $Z \notin Y_1$ and $Z \cap Y_1 \neq \emptyset$ and $\ell \geq 0$.

Proof. Define $\phi := (\rho, h) : W \to Y_1 \times \mathbb{R}$. We show first: The derivative $d_x \phi = (d_x \rho, d_x h) : T_x M \to T_x Y_1 \times \mathbb{R}$ is an isomorphism for all $x \in Y_1$. As $\dim(T_x M) = \dim(T_x Y_1 \times \mathbb{R})$, it is enough to prove: $d_x \phi$ is surjective.

As $\phi|_{Y_1} = (\mathrm{id}_{Y_1}, 0)$, we have $d_x \phi|_{T_x Y_1} = (\mathrm{id}_{T_x Y_1}, 0)$, so $T_x Y_1 \times \{0\} \subset \mathrm{im}(d_x \phi)$. In addition $d_x h: T_x M \to \mathbb{R}$ is surjective, so there exists $v \in T_x M$ such that $d_x h(v) = 1$. Thus, $d_x \phi(v) = (d_x \rho(v), 1)$, so $d_x \phi$ is surjective.

Let $W' := \{x \in W : d_x \phi \text{ is an isomorphism}\}$, which is an open semialgebraic neighborhood of Y_1 . Thus, $\phi|_{W'} : W' \to Y_1 \times \mathbb{R}$ is an open map and $\phi(W')$ is an open semialgebraic neighborhood of $Y_1 \times \{0\}$ in $Y_1 \times \mathbb{R}$. As $\phi|_{W'} : W' \to \phi(W')$ is a local homeomorphism and $\phi|_{Y_1} = (\mathrm{id}_{Y_1}, 0)$ is a homeomorphism (onto its image), there exist by [BFR, Lem.9.2] open semialgebraic neighborhoods $W'' \subset W'$ of Y_1 and $U \subset Y_1 \times \mathbb{R}$ of $Y_1 \times \{0\}$ such that $\phi|_{W''} : W'' \to U$ is a semialgebraic homeomorphism.

Consider the strictly positive, continuous semialgebraic map

$$\delta: Y_1 \to (0, +\infty), \ x \mapsto \operatorname{dist}((x, 0), (Y_1 \times \mathbb{R}) \setminus U).$$

By [Sh, Thm.II.4.1] there exists a strictly positive Nash function ε on Y_1 such that $|\varepsilon - \frac{3}{4}\delta| < \frac{1}{4}\delta$, that is, $\frac{1}{2}\delta < \varepsilon < \delta$. Consider the open semialgebraic neighborhood

$$U' := \{(y,t) \in Y_1 \times \mathbb{R} : |t| < \varepsilon(y)\} \subset U$$

of $Y_1 \times \{0\}$ and define $V := (\phi|_{W''})^{-1}(U')$. The restriction $\phi|_V : V \to U'$ is a Nash diffeomorphism. Consequently,

$$\varphi: V \to Y_1 \times (-1, 1), \ x \mapsto \left(\rho(x), \frac{h(x)}{\varepsilon(\rho(x))}\right)$$

is a Nash diffeomorphism, because it is the composition of $\phi|_V$ with the Nash diffeomorphism

$$\psi: U' \to Y_1 \times (-1, 1), \ (y, t) \mapsto \left(y, \frac{t}{\varepsilon(y)}\right).$$

Let $\ell \geq 0$ and Z be an irreducible component of $\operatorname{Sing}_{\ell}(Y)$ such that $Z \not\subset Y_1$ and $Z \cap Y_1 \neq \emptyset$. By Proposition 5.2 $\rho(Z \cap V) = Z \cap Y_1$, so $\varphi(Z \cap V) \subset (Z \cap Y_1) \times (-1, 1)$. Let us prove: $\varphi|_{Z \cap V} : Z \cap V \to (Z \cap Y_1) \times (-1, 1)$ is surjective.

As Y is a Nash normal-crossings divisor, $Z \cap Y_1$ is a Nash manifold, so its connected components C_i are Nash manifolds of dimension e - 1 where $e = \dim(Z)$, because $Z \not\subset Y_1$. Thus, each $C_i \times (-1, 1)$ is a closed connected Nash submanifold of $Y_1 \times (-1, 1)$ of dimension e and, in particular, $C_i \times (-1, 1)$ is an irreducible Nash subset of $Y_1 \times (-1, 1)$ of dimension e.

Let Z_i be a connected component of the Nash manifold $Z \cap V$ such that $C_i \subset Z_i \cap Y_1$. As Z_i is connected, also $\rho(Z_i) \subset Z \cap Y_1$ is connected, so it is contained in one of the connected components of $Z \cap Y_1$. As $C_i \subset Z_i \cap Y_1 \subset \rho(Z_i)$ is a connected component of $Z \cap Y_1$, we deduce $\rho(Z_i) = Z_i \cap Y_1 = C_i$. Then $\varphi(Z_i) \subset C_i \times (-1, 1)$ and $\varphi(Z_i)$ is a Nash subset of $Y_1 \times (-1, 1)$ of dimension e. By the identity principle, we conclude $\varphi(Z_i) = C_i \times (-1, 1)$ because $C_i \times (-1, 1)$ is an irreducible Nash subset of $Y_1 \times (-1, 1)$ of dimension e. Consequently,

$$(Z \cap Y_1) \times (-1, 1) = \bigcup_i (C_i \times (-1, 1)) = \bigcup_i \varphi(Z_i) \subset \varphi(Z \cap V) \subset (Z \cap Y_1) \times (-1, 1),$$

so $\varphi|_{Z\cap V}: Z\cap V \to (Z\cap Y_1) \times (-1,1)$ is surjective, as required.

5.3. Nash equations for the components of the boundary. Let $\Omega \subset \mathbb{R}^n$ be a Nash manifold with corners (Section 2.3). The following result provides suitable Nash equations for the facets of Ω . The proof is strongly inspired by the proof of [Fe2, Lem.4.3], however it needs some modifications and we include it in full detail for the sake of the reader.

Lemma 5.4 (Nash equations for the facets). Let Y_1 be an irreducible component of the Nash normal-crossings divisor Y (which is the Nash closure of $\partial \Omega$ in the Nash envelope M of the Nash manifold with corners Ω). Then, after shrinking M if necessary, there exists a Nash function $h_1: M \to \mathbb{R}$ such that:

- $Y_1 = \{h_1 = 0\}$ and $d_x h_1 : T_x M \to \mathbb{R}$ is surjective for each $x \in Y_1$.
- $H_1 := h_1^{-1}([0, +\infty))$ is a Nash manifold with boundary $\partial H_1 := Y_1$ and interior $\operatorname{Int}(H_1) = h_1^{-1}((0, +\infty))$ that contains \mathfrak{Q} as a closed subset.

Proof. The proof is conducted in several steps:

STEP 1. We construct first an S^2 semialgebraic function h_1^* on M such that $Y_1 \subset \{h_1^* = 0\}$ and $d_x h_1^*(v) > 0$ for each $x \in Y_1 \cap \Omega$ and each non-zero vector $v \in T_x M$ pointing 'inside Ω '.

By Theorem 2.4 there exist finitely many open semialgebraic subsets U_i (for i = 1, ..., r) equipped with Nash diffeomorphisms $u_i := (u_{i1}, ..., u_{id}) : U_i \to \mathbb{R}^d$ such that:

$$\begin{cases} U_i \subset \text{Int}(\mathcal{Q}) \text{ or } U_i \cap \mathcal{Q} = \emptyset, & \text{if } U_i \text{ does not meet } \partial \mathcal{Q}, \\ U_i \cap \mathcal{Q} = \{u_{i1} \ge 0, \dots, u_{ik_i} \ge 0\}, & \text{if } U_i \text{ meets } \partial \mathcal{Q} \text{ (for a suitable } k_i \ge 1). \end{cases}$$

Assume that Y_1 meets U_i exactly for i = 1, ..., s for some $s \leq r$. We reorder the variables u_{ij} in order to guarantee that $Y_1 \cap U_i = \{u_{i1} = 0\}$ and $Q \cap U_i \subset \{u_{i1} \geq 0\}$ for i = 1, ..., s. Let

$$\square$$

 $\{\theta_i\}_{i=1}^{s+1}: M \to [0,1]$ be an \mathcal{S}^2 partition of unity subordinated to the finite covering $\{U_i\}_{i=1}^s \cup U_{s+1} := \{M \setminus Y_1\}$ of M and consider the \mathcal{S}^2 function $h_1^* := \sum_{i=1}^s \theta_i u_{i1}$. It holds $Y_1 \subset \{h_1^* = 0\}$. Fix $x \in Y \cap \Omega$ and let $u \in T$ M be a non-zero vector pointing (inside Ω) that is d $u_i(u) > 0$.

Fix $x \in Y_1 \cap \Omega$ and let $v \in T_x M$ be a non-zero vector pointing 'inside Ω ', that is, $d_x u_{i1}(v) > 0$ if $x \in U_i$. We have $\operatorname{Cl}(\{\theta_i > 0\}) \subset U_i$, $x \notin U_{s+1}$ and $u_{i1}(x) = 0$ if $x \in U_i$, so

$$d_x h_1^* = \sum_{x \in U_i} u_{i1}(x) d_x \theta_i + \sum_{x \in U_i} \theta_i(x) d_x u_{i1} = \sum_{x \in U_i} \theta_i(x) d_x u_{i1} \rightsquigarrow d_x h_1^*(v) = \sum_{x \in U_i} \theta_i(x) d_x u_{i1}(v) > 0$$

because $\sum_{x \in U_i} e_i(x) = 1$, $\theta_i(x) \ge 0$ and $d_i u_{i1}(v) \ge 0$ if $x \in U_i$

because $\sum_{x \in U_i} \theta_i(x) = 1$, $\theta_i(x) \ge 0$ and $d_x u_{i1}(v) > 0$ if $x \in U_i$.

STEP 2. By [BFR, Prop.8.2] there exists a Nash function h'_1 on M close to h^*_1 in the S^2 topology such that $Y_1 \subset \{h'_1 = 0\}$ and $d_x h'_1(v) > 0$ for each $x \in Y_1 \cap Q$ and each non-zero vector $v \in T_x M$ pointing 'inside Q'. We claim: there exists an open semialgebraic neighborhood $W \subset M$ of $Y_1 \cap Q$ such that $\operatorname{Int}(Q) \cap W \subset \{h'_1 > 0\} \cap W$ and $\{h'_1 = 0\} \cap W = Y_1$.

Pick a point $x \in Y_1$ and assume $x \in U_1$. As h'_1 vanishes identically at Y_1 , we may write $h'_1|_{U_1} = u_{11}a_1$, where a_1 is a Nash function on U_1 . Pick $y \in Y_1 \cap U_1 \cap Q$ and observe that $d_yh'_1 = a_1(y)d_yu_{11}$. Let $v \in T_yM$ be a non-zero vector pointing 'inside Q'. As $d_yu_{11}(v) > 0$ and $d_yh'_1(v) > 0$, we deduce $a_1(y) > 0$. Define $W_1 := \{a_1 > 0\} \subset U_1$ and notice that $Y_1 \cap U_1 \cap Q \subset W_1$, $\operatorname{Int}(Q) \cap W_1 \subset \{h'_1 > 0\} \cap W_1$ and $\{h'_1 = 0\} \cap W_1 = Y_1 \cap W_1$. Construct analogously W_2, \ldots, W_s and observe that $W := \bigcup_{i=1}^s W_i$ satisfies the required properties.

Substitute M by $M \setminus (Y_1 \setminus W)$, which is an open semialgebraic subset of M that contains Q as a closed subset. Substitute Y by the Nash closure of ∂Q in the new M and Y_1 by the irreducible component of the new Y that contains the facet $Y_1 \cap Q$.

STEP 3. Next, we construct h_1 . If W = M, it is enough to set $h_1 := h'_1$. Suppose $W \neq M$. Let ε_0 be a (continuous) semialgebraic function whose value is 1 on Y_1 and -1 on $M \setminus W$. Let ε be a Nash approximation of ε_0 such that $|\varepsilon - \varepsilon_0| < \frac{1}{2}$. Then

$$\varepsilon(x) \begin{cases} > \frac{1}{2} & \text{if } x \in Y_1, \\ < -\frac{1}{2} & \text{if } x \in M \setminus W \end{cases}$$

Thus, $\{\varepsilon > 0\} \subset W$ is an open semialgebraic neighborhood of Y_1 in M. By [Sh, Prop.II.5.3] Y_1 is a Nash subset of M. Let f be a Nash equation of Y_1 in M. Substituting f by $\frac{f^2}{\varepsilon^2 + f^2}$, we may assume that f is non-negative and f(x) = 1 if $\varepsilon(x) = 0$. Consider the (continuous) semialgebraic function on M given by

$$\delta(x) := \begin{cases} 1 & \text{if } \varepsilon(x) > 0, \\ \frac{1}{f(x)} & \text{if } \varepsilon(x) \le 0. \end{cases}$$

Let g be a Nash function on M such that $\delta < g$ (see [BCR, Prop.2.6.2], after embedding M in \mathbb{R}^{n+1} as a closed subset). Consider the Nash function $h_1 := h'_1 + f^2 g^2 (h'_1^2 + 1)$ and let us prove that it satisfies the required conditions.

STEP 4. We claim: h_1 is positive on Int(Q).

Let $x \in \text{Int}(\Omega)$. If $h'_1(x) > 0$, then $h_1(x) > 0$. If $h'_1(x) \le 0$, then $x \notin W$ (because $\text{Int}(\Omega) \cap W \subset \{h'_1 > 0\} \cap W$). Thus, $\varepsilon(x) \le 0$ and

$$h_1(x) = h'_1(x) + g^2(x)f^2(x)(h'_1(x) + 1)$$

> $h'_1(x) + \frac{1}{f^2(x)}f^2(x)(h'_1(x) + 1) = h'_1(x) + h'_1(x) + 1 > 0.$

STEP 5. It holds: There exists an open semialgebraic neighborhood $W' \subset W$ of Y_1 such that $\{h_1 = 0\} \cap W' = Y_1$, $Int(\mathfrak{Q}) \cap W' \subset \{h_1 > 0\} \cap W'$ and the differential $d_xh_1 : T_xM \to \mathbb{R}$ is surjective for all $x \in Y_1$.

Recall that $W = \bigcup_{i=1}^{s} W_i$. We have seen in STEP 2 that there exists a Nash function a_1 on U_1 such that $h'_1|_{U_1} = u_{11}a_1$ and $Y_1 \cap U_1 \subset W_1 := \{a_1 > 0\}$. As f vanishes identically at Y_1 , we deduce $f|_{U_1} = u_{11}b_1$ where b_1 is a Nash function on U_1 . Consequently,

$$h_1|_{U_1} = u_{11}a_1 + g^2|_{U_1}u_{11}^2b_1^2(u_{11}^2a_1^2 + 1) = u_{11}(a_1 + g^2|_{U_1}u_{11}b_1^2(u_{11}^2a_1^2 + 1))$$

and $d_x h_1 = a_1(x) d_x u_{11} = d_x h'_1$ for $x \in Y_1 \cap U_1$. Define

$$W_1' := \{a_1 + g^2|_{U_1}u_{11}b_1^2(u_{11}^2a_1^2 + 1) > 0\} \cap W_1,$$

which is an open semialgebraic subset of M. We have $Y_1 \cap U_1 \subset W'_1$ and

 $\operatorname{Int}(\mathfrak{Q}) \cap W_1' \subset \{h_1 > 0\} \cap W_1'.$

We construct analogously W'_2, \ldots, W'_s and observe that the open semialgebraic subset $W' := \bigcup_{i=1}^s W'_i \subset M$ is an open neighborhood of Y_1 that satisfies $\{h_1 = 0\} \cap W' = Y_1$, $\operatorname{Int}(\mathfrak{Q}) \cap W' \subset \{h_1 > 0\} \cap W'$ and $d_x h : T_x M \to \mathbb{R}$ is surjective for all $x \in Y_1$.

Observe that $H_1 := h_1^{-1}([0, +\infty))$ is a Nash manifold with boundary, because it is a semialgebraic set and a smooth manifold with boundary (the latter holds by the implicit function theorem, because 0 is not a critical value of h_1). Consequently, $M' := \{h_1 > 0\} \cup W'$ and $h_1|_{M'}$ satisfy all the required conditions.

Corollary 5.5. Let Y_1, \ldots, Y_ℓ be the irreducible components of Y. After shrinking the manifold M, there exist Nash functions $h_i : M \to \mathbb{R}$ such that $Y_i = \{h_i = 0\}, d_x h_i : T_x M \to \mathbb{R}$ is surjective for each $x \in Y_i$ and $Q = \{h_1 \ge 0, \ldots, h_\ell \ge 0\}$.

The following result is the counterpart of the previous ones.

Lemma 5.6 (Finite intersections of Nash manifolds with boundary). Let $M \subset \mathbb{R}^n$ be a Nash manifold and $Y \subset M$ a Nash normal-crossings divisor. Let Y_1, \ldots, Y_ℓ be the irreducible components of Y and $h_i : M \to \mathbb{R}$ a Nash equation of Y_i such that $d_x h_i : T_x M \to \mathbb{R}$ is surjective for each $x \in Y_i$ and each $i = 1, \ldots, \ell$. Assume that $Q := \{h_1 \ge 0, \ldots, h_\ell \ge 0\}$ is non-empty. Then Q is a Nash manifold with corners and the Nash closure of its boundary ∂Q is Y. In particular, Q is the intersection of the Nash manifolds with boundary $H_i := \{h_i \ge 0\}$ for $i = 1, \ldots, \ell$.

Proof. Observe first that $H_i := \{h_i \ge 0\} \subset M$ is a Nash manifold with boundary, $\partial H_i = Y_i$ for $i = 1, \ldots, \ell$ and $\mathfrak{Q} = \bigcap_{i=1}^{\ell} H_i$. We check next: \mathfrak{Q} is a Nash manifold with corners.

Pick $x \in \mathbb{Q}$. If $h_i(x) > 0$ for $i = 1, \ldots, \ell$, then x belongs to the open semialgebraic subset $\{h_1 > 0, \ldots, h_\ell > 0\}$ of M contained in \mathbb{Q} . Suppose $h_1(x) = 0, \ldots, h_s(x) = 0, h_{s+1}(x) > 0, \ldots, h_\ell(x) > 0$ for some $1 \le s \le \ell$. As Y is a Nash normal-crossings divisor, the linear forms $d_x h_1, \ldots, d_x h_s$ are linearly independent and there exists an open semialgebraic neighborhood $U \subset M$ of x endowed with a Nash diffeomorphism $u := (h_1, \ldots, h_s, u_{s+1}, \ldots, u_d) : U \to \mathbb{R}^d$ such that u(x) = 0 and $Y \cap U = \{h_1 \cdots h_s = 0\} \cap U$. Observe that $\mathbb{Q} \cap U = \{h_1 \ge 0, \ldots, h_s \ge 0\} \cap U$.

The boundary of ${\mathbb Q}$ is

r

$$\partial \mathfrak{Q} = \bigcup_{i=1}^{r} \{ h_1 \ge 0, \dots, h_{i-1} \ge 0, h_i = 0, h_{i+1} \ge 0, \dots, h_{\ell} \ge 0 \}$$

and its Nash closure in M is Y, so Q is a Nash manifold with corners, as required.

5.4. Nash doubles of Nash manifolds with boundary. Doubling a smooth manifold with boundary is a standard tool in differential topology. The Nash construction has been treated in [Sh, Thm.VI.2.1] (compact case) and in [Fe2, 4.B.1, 4.B.2] (general case). We recall next the main statements and refer the reader to [Fe2, 4.B.1, 4.B.2] for the precise details.

Lemma 5.7 (Nash double, [Fe2, 4.B.1]). Let $H \subset \mathbb{R}^n$ be a d-dimensional Nash manifold with boundary ∂H and let $h : M \to \mathbb{R}$ be a Nash equation for ∂H such that $\partial H = \{h = 0\}$, $\operatorname{Int}(H) = \{h > 0\}$ and $d_x h : T_x H \to \mathbb{R}$ is surjective for all $x \in \partial H$. Then, the semialgebraic set $D(H) := \{(x, t) \in H : t \mathbb{R} : t^2 = h(x) = 0\} \subset \mathbb{R}^{n+1}$

$$D(H) := \{(x,t) \in H \times \mathbb{R} : t^2 - h(x) = 0\} \subset \mathbb{R}^n$$

is a d-dimensional Nash manifold that contains $\partial H \times \{0\}$ as the Nash subset $\{t = 0\}$.

Lemma 5.8 ([Fe2, 4.B.2]). The projection $\pi : D(H) \to H$, $(x,t) \mapsto x$ is a surjective Nash map. Fix $\epsilon = \pm$ and denote $H_{\epsilon} := D(H) \cap \{\epsilon t \ge 0\}$. We have the following:




FIGURE 5.1. Nash double of H and the projection π (figure borrowed from [Fe2, Fig.3]).

- (i) The restriction $\pi_{\epsilon} := \pi|_{H_{\epsilon}} : H_{\epsilon} \to H$ is a semialgebraic homeomorphism and the restriction $\pi|_{D(H)\cap\{\epsilon t>0\}} : D(H)\cap\{\epsilon t>0\} \to Int(H)$ is a Nash diffeomorphism.
- (ii) $\pi(x,0) = x$ for all $(x,0) \in \partial H \times \{0\} = D(H) \cap \{t=0\}.$
- (iii) π has local representations $(y_1, \ldots, y_d) \mapsto (y_1^2, y_2, \ldots, y_d)$ at each point of $D(H) \cap \{t = 0\}$.
- (iv) π is open and proper.

Proof. Properties (i), (ii) and (iii) are proved in [Fe2, 4.B.2]. Let us show (iv). We prove first that π is open. Let $A \subset D(H)$ be an open set and let $A_{\epsilon} := A \cap H_{\epsilon}$, which is an open subset of H_{ϵ} . As $\pi_{\epsilon} : H_{\epsilon} \to H$ is a semialgebraic homeomorphism, $\pi(A_{\epsilon})$ is an open subset of H. Thus, $\pi(A) = \pi(A_{+}) \cup \pi(A_{-})$ is an open subset of H, so π is open.

To show that π is proper, pick $K \subset H$ compact and observe that $\pi^{-1}(K) = (\pi_+)^{-1}(K) \cup (\pi_-)^{-1}(K)$. As each $\pi_{\epsilon} : H_{\epsilon} \to H$ is a semialgebraic homeomorphism, $(\pi_{\epsilon})^{-1}(K)$ is compact for $\epsilon = \pm$, so $\pi^{-1}(K)$ is a compact subset of D(H), so π is proper, as required. \Box

5.5. Folding one component of the boundary. Let $M \subset \mathbb{R}^n$ be a *d*-dimensional Nash manifold that contains H as a closed subset and assume that ∂H is a Nash subset of M. Let Y be a Nash normal-crossings divisor of M such that ∂H is a union of irreducible components of Y. Let Y_1, \ldots, Y_r be the irreducible components of Y that meet ∂H but are not contained in ∂H and let Y_{r+1}, \ldots, Y_s be the irreducible components of Y that do not meet ∂H . Let $h: M \to \mathbb{R}$ be a Nash equation of ∂H . Observe that $h_i := h|_{Y_i}$ is a Nash equation of $Y_i \cap \partial H$ such that $Y_i \cap \operatorname{Int}(H) = \{h_i > 0\}$ and $d_x h_i: T_x Y_i \to \mathbb{R}$ is surjective for all $x \in Y_i \cap \partial H$. Thus, $Y_i \cap H$ is a Nash manifold with boundary $Y_i \cap \partial H$ that is contained in Y_i as a closed subset. In addition,

$$D(Y_i \cap H) = \{ (x, t) \in (Y_i \cap H) \times \mathbb{R} : t^2 - h_i(x) = 0 \} = (Y_i \times \mathbb{R}) \cap D(H),$$
(5.1)

is the Nash double of $Y_i \cap H$. We show that there exists a Nash embedding of M into D(H) that maps Y_i into $D(Y_i \cap H) = (Y_i \times \mathbb{R}) \cap D(H)$ for $i = 1, \ldots, r$ and maps Y_j onto $(Y_j \times \mathbb{R}) \cap D(H) \cap \{t \ge 0\}$ for $j = r + 1, \ldots, s$. We begin with the following technical lemma.

Lemma 5.9. Let $k \ge 1$ and $0 < a \le 1$. Consider the C^{2k} semialgebraic function $f_{a,k} := (1 - (t/a)^{2k})^{2k} t + (1 - (1 - (t/a)^{2k})^{2k})\sqrt{t}$. Then:

- $f_{a,k}$ is positive semidefinite and strictly increasing on [0,a].
- The Taylor polynomial of $f_{a,k}$ at t = 0 of degree 2k is t.
- The Taylor polynomials of $f_{a,k}$ and \sqrt{t} of degree 2k-1 coincide at t=a,
- $f_{a,k}(t) \le \sqrt{t}$ on [0,1].

Proof. Write $f := f_{a,k}$ to lighten notations. Using Newton's binomial, we have

$$f = \sum_{\ell=0}^{2k} \binom{2k}{\ell} (-1)^{\ell} \left(\frac{\mathsf{t}}{a}\right)^{2k\ell} \mathsf{t} - \sum_{\ell=1}^{2k} \binom{2k}{\ell} (-1)^{\ell} \left(\frac{\mathsf{t}}{a}\right)^{2k\ell} \sqrt{\mathsf{t}},$$



FIGURE 5.2. Graphs of f_a for $a = \frac{1}{2}$ and k = 2 (left) and k = 6 (right).

so f is a \mathcal{C}^{2k} semialgebraic function and the Taylor polynomial of f at t = 0 of degree 2k is t. In addition,

$$f = (1 - (1 + (t/a) - 1)^{2k})^{2k} t + (1 - (1 - (1 + (t/a) - 1)^{2k})^{2k})\sqrt{t}$$

and using Newton's binomial, we have

$$f = \sqrt{\mathbf{t}} - \left(\frac{\mathbf{t}}{a} - 1\right)^{2k} \left(\sum_{\ell=1}^{2k} \binom{2k}{\ell} \left(\frac{\mathbf{t}}{a} - 1\right)^{\ell-1}\right)^{2k} (\mathbf{t} - \sqrt{\mathbf{t}}),$$

so the Taylor polynomials of f and \sqrt{t} of degree 2k - 1 at t = a coincide.

Define $\sigma := (1 - (t/a)^{2k})^{2k}$ and observe that $\sigma(0) = 1$, $\sigma(a) = 0$ and $\sigma([0, a]) \subset [0, 1]$, so both σ , $1 - \sigma$ are positive semidefinite on [0, a]. Thus, $f = \sigma t + (1 - \sigma)\sqrt{t}$ is positive semidefinite on [0, a]. As $0 < a \le 1$, it holds $\sqrt{t} - t > 0$ on (0, a), so the derivative

$$\begin{split} f' &:= \left(\mathbf{t} - \sqrt{\mathbf{t}} \right) \sigma' + (1 - \sigma) \frac{1}{2\sqrt{\mathbf{t}}} + \sigma \\ &= (\sqrt{\mathbf{t}} - \mathbf{t}) 4k^2 (1 - (\mathbf{t}/a)^{2k})^{2k - 1} (\mathbf{t}/a)^{2k - 1} (1/a) + (1 - \sigma) \frac{1}{2\sqrt{\mathbf{t}}} + \sigma \end{split}$$

is strictly positive on (0, a) and f is strictly increasing on [0, a]. Finally, as $t \leq \sqrt{t}$ on [0, 1], we deduce that $f = \sigma t + (1 - \sigma)\sqrt{t} \le \sqrt{t}$ on [0, 1], as required. \square

We are ready to prove the embedding theorem.

Theorem 5.10 (Nash embedding into the Nash double). After shrinking M if necessary, there exists a Nash embedding $\phi: M \to D(H)$ that maps H onto H_+ and Y_i into $D(Y_i \cap H)$ for $i = 1, \ldots, r$. In addition:

- φ|_{∂H} = id_{∂H} and φ|_H is close to (π|_{H+})⁻¹,
 φ(Y_j) = (Y_j × ℝ) ∩ H₊ for each j = r + 1,...,s,
- There exists a Nash retraction ρ compatible with Y such that $\phi(x)$ is close to $(\rho(x), 0)$ for each $x \in M \setminus H$.
- There exists an open semialgebraic neighborhood $W \subset M$ of ∂H such that $\phi|_{Cl(W) \cup H}$: $Cl(W) \cup H \to D(H)$ is proper.

Proof. The proof is conducted in several steps:

STEP 1. We first construct suitable semialgebraic neighborhoods $U \subset M$ of ∂H and $V \subset D(H)$ of $\partial H \times \{0\}$.

Let $U \subset M$ be an open semialgebraic neighborhood of ∂H equipped with a Nash retraction $\rho: U \to \partial H$ compatible with Y (Proposition 5.2). We may assume that U does not meet the irreducible components of Y that do not meet ∂H . Let $h: M \to \mathbb{R}$ be a Nash equation of ∂H such that $Int(H) = \{h > 0\}$ and $d_x h : T_x H \to \mathbb{R}$ is surjective for all $x \in \partial H$. Define the Nash map $\varphi := (\rho, h) : U \to \partial H \times \mathbb{R}$. By Lemma 5.3 there exists, after shrinking U if necessary, a strictly positive Nash function $\varepsilon : \partial H \to (0, 1)$ such that

$$\varphi(U) = \{(y, s) \in \partial H \times \mathbb{R} : |s| < \varepsilon(y)\},\$$
$$\varphi(U \cap H) = \{(y, s) \in \partial H \times \mathbb{R} : 0 \le s < \varepsilon(y)\},\$$

and $\varphi: U \to \varphi(U)$ is a Nash diffeomorphism such that

$$p(Z \cap U) = \{(y, s) \in (Z \cap \partial H) \times \mathbb{R} : |s| < \varepsilon(y)\}$$

for each irreducible component Z of $\operatorname{Sing}_{\ell}(Y)$ such that $Z \cap \partial H \neq \emptyset$, $Z \not\subset \partial H$ and $\ell \ge 0$. Define $V := \pi^{-1}(U \cap H) = \{(x,t) \in (U \cap H) \times \mathbb{R} : t^2 - h(x) = 0\}.$

STEP 2. Let $V' := \{(y,t) \in \partial H \times \mathbb{R} : |t| < \sqrt{\varepsilon(y)}\}$. We want to prove: The Nash map $\psi : V \to \partial H \times \mathbb{R}, \ (x,t) \mapsto (\rho(x),t)$

is a Nash diffeomorphism onto its image V', such that

 $\psi(Z \cap U \cap H) = \{(y,t) \in (Z \cap \partial H) \times \mathbb{R} : 0 \le t < \sqrt{\varepsilon(y)}\}$

for each irreducible component Z of $\operatorname{Sing}_{\ell}(Y)$ such that $Z \cap \partial H \neq \emptyset$, $Z \not\subset \partial H$ and $\ell \geq 0$.

(1) ψ is injective. If $(x_1, t_1), (x_2, t_2) \in V$ satisfy $\psi(x_1, t_1) = \psi(x_2, t_2)$, then $\rho(x_1) = \rho(x_2)$ and $h(x_1) = t_1^2 = t_2^2 = h(x_2)$. Then we have $\varphi(x_1) = \varphi(x_2)$, so $x_1 = x_2$. Thus, $(x_1, t_1) = (x_2, t_2)$.

(2) $\psi(V) = V'$. Fix a point $(x,t) \in V$. Then, $x = \pi(x,t) \in U$ and $\varphi(x) = (\rho(x), h(x)) \in \varphi(U)$, so $t^2 = h(x) < \varepsilon(\rho(x))$ and $\psi(x,t) \in V'$. Conversely, fix a point $(y,t) \in V'$. As $(y,t^2) \in \varphi(U)$, there exists a point $x \in U$ such that $\varphi(x) = (\rho(x), h(x)) = (y,t^2)$. As $x \in U$ and $h(x) = t^2 \ge 0$, we have $x \in U \cap H$, so $(x,t) \in V$ and $(y,t) = (\rho(x), t) = \psi(x,t) \in \psi(V)$.

(3) The differential $d_z \psi : T_z D(H) \to T_{\rho(z)} H \times \mathbb{R}$ is an isomorphism for each $z \in V$. Write z := (x, t) and notice that

$$T_z D(H) = \{(v, r) \in T_x H \times \mathbb{R} : d_x h(v) - 2tr = 0\}$$

and $d_x\psi(v,r) = (d_x\rho(v),r)$. If $t \neq 0$, then $d_z\psi(v,r) = (d_x\rho(v), \frac{1}{2t}d_xh(v))$. As $d_x\varphi = (d_x\rho, d_xh)$ is an isomorphism, also $d_z\psi$ is an isomorphism. If t = 0, that is, $z = (x,0) \in \partial H \times \mathbb{R}$, then

$$T_z D(H) = \{ (v, r) \in T_x H \times \mathbb{R} : d_x h(v) = 0 \} = T_x \partial H \times \mathbb{R}$$

and $d_z\psi(v,r) = (v,r)$ because $\rho|_{\partial H} = \mathrm{id}_{\partial H}$. Hence, $d_z\psi$ is again an isomorphism.

(4) Let Z be an irreducible component of $\operatorname{Sing}_{\ell}(Y)$ such that $Z \cap \partial H \neq \emptyset$, $Z \not\subset \partial H$ and $\ell \geq 0$. Then

$$\psi(Z \cap U \cap H) = \{(y, t) \in (Z \cap \partial H) \times \mathbb{R} : 0 \le t < \sqrt{\varepsilon(y)}\}.$$

By Proposition 5.2 we have $\rho(Z \cap U) = Z \cap \partial H$. Thus,

$$\psi(Z \cap U \cap H) \subset \{(y,t) \in (Z \cap \partial H) \times \mathbb{R} : 0 \le t < \sqrt{\varepsilon(y)}\}.$$

To prove that the previous inclusion is in fact an equality, it is enough to proceed similarly to the end of the proof of Lemma 5.3.

STEP 3. Let $a : \partial H \to \mathbb{R}$ be a strictly positive Nash function such that $a < \varepsilon$. Recall that $H_{\epsilon} = D(H) \cap \{\epsilon t \ge 0\}$. For $\epsilon = \pm$, define

$$H^{\bullet} := H \setminus \varphi^{-1} \left(\left\{ (y, s) \in \partial H \times \mathbb{R} : |s| < \frac{a(y)}{4} \right\} \right) \supset H \setminus U,$$
$$H^{\bullet}_{\epsilon} := H_{\epsilon} \setminus \psi^{-1} \left(\left\{ (y, s) \in \partial H \times \mathbb{R} : |s| < \frac{\sqrt{a(y)}}{2} \right\} \right) \supset H_{\epsilon} \setminus V.$$

We claim: The restriction $\varpi_{a\epsilon} := \pi|_{H^{\bullet}_{\epsilon}} : H^{\bullet}_{\epsilon} \to H^{\bullet}$, $(x,t) \mapsto x$ is a Nash diffeomorphism, whose inverse map is $\varpi_{a\epsilon}^{-1} : H^{\bullet} \to H^{\bullet}_{\epsilon}$, $x \mapsto (x, \epsilon \sqrt{h(x)})$ for $\epsilon = \pm$.

As $\varpi_{a\epsilon}$ is injective, let us check: *it is surjective*.

Let $x \in H^{\bullet}$. As $x \in \text{Int}(H)$, we have h(x) > 0 and we write $t := \epsilon \sqrt{h(x)}$. It holds that $(x,t) \in D(H)$ and $\pi(x,t) = x$. We want to check that $(x,t) \in H^{\bullet}_{\epsilon}$. If $x \notin U$, then $(x,t) \in H_{\epsilon} \setminus V \subset H^{\bullet}_{\epsilon}$. If $x \in U$, then $\psi(x,t) = (\rho(x),t)$. As $x \in H^{\bullet}$, it holds $\frac{a(\rho(y))}{4} \leq h(x)$, so $\frac{\sqrt{a(\rho(y))}}{2} \leq \sqrt{h(x)} = \epsilon t$. Consequently, $(x,t) \in H^{\bullet}_{\epsilon}$, so $\varpi_{a\epsilon}$ is surjective. In addition, by Lemma 5.8(i) $d_z \varpi_{a\epsilon} = d_z \pi$ is an isomorphism for each $z \in H_{\epsilon}^{\bullet} \subset D(H) \cap \{\epsilon t > 0\}$. Consequently $\varpi_{a\epsilon}$ is a Nash diffeomorphism, as claimed.

STEP 4. We now construct a semialgebraic embedding $\phi_{a,k} : M \to D(H)$ of class \mathcal{C}^{2k-1} , for $k \geq 1$ arbitrarily large. Substitute M by $H \cup U$ and define

$$F_{a,k}: \partial H \times \mathbb{R} \to \partial H \times \mathbb{R}, \ (y,s) \mapsto \begin{cases} (y,s) & \text{if } s < 0, \\ (y,f_{a(y),k}(s)) & \text{if } 0 \le s \le a(y), \\ (y,\sqrt{s}) & \text{if } a(y) < s, \end{cases}$$

where $f_{a(y),k}$ is the C^{2k} semialgebraic function introduced in Lemma 5.9 and we choose k large enough. Denote

$$\begin{split} U' &:= \varphi(U) = \{(y, s) \in \partial H \times \mathbb{R} : \ |s| < \varepsilon(y)\}, \\ V' &= \psi(V) = \{(y, t) \in \partial H \times \mathbb{R} : \ |t| < \sqrt{\varepsilon(y)}\} \text{ (already introduced in STEP 2)} \end{split}$$

The open semialgebraic set $F_{a,k}(U')$ is contained in V', because if $-\varepsilon(y) < -s < 0$, then $-\sqrt{\varepsilon(y)} < -\sqrt{s} < -s < 0$ (recall that $0 < \varepsilon(y) < 1$) and if $0 \le s \le a(y) \le 1$, then $0 \le f_{a(y),k}(s) \le \sqrt{s}$.

The map $F_{a,k}$ is a \mathcal{C}^{2k-1} semialgebraic diffeomorphism, because $f_{a(y),k}: [0, a(y)] \to [0, \sqrt{a(y)}]$ is by Lemma 5.9 a \mathcal{C}^{2k-1} semialgebraic diffeomorphism such that the Taylor polynomial of degree 2k at t = 0 is t and its Taylor polynomial of degree 2k - 1 at t = a(y) coincides with the one of \sqrt{t} .

Define

$$\phi_{a,k}: M \to D(H), \ x \mapsto \begin{cases} \varpi_{a+}^{-1}(x) & \text{if } x \in M \setminus U = H \setminus U \subset H^{\bullet}, \\ (\psi^{-1} \circ F_{a,k} \circ \varphi)(x) & \text{if } x \in U. \end{cases}$$

It holds that $\phi_{a,k}$ is a \mathcal{C}^{2k-1} semialgebraic diffeomorphism onto its image $\phi_{a,k}(M)$, whose \mathcal{C}^{2k-1} semialgebraic inverse is

$$\phi_{a,k}^{-1}:\phi_{a,k}(M)\to M,\ x\mapsto\begin{cases} \varpi_{a+}(y) & \text{if } y\in\phi_{a,k}(M)\setminus V\subset H_+^{\bullet},\\ (\varphi^{-1}\circ F_{a,k}^{-1}\circ\psi)(y) & \text{if } y\in V. \end{cases}$$

In addition, it satisfies:

- $\phi_{a,k}(H) = H_+$ and $\phi_{a,k}|_{\partial H} = \mathrm{id}_{\partial H}$.
- $\phi_{a,k}(Y_j) = (Y_j \times \mathbb{R}) \cap H_+$ for each $j = r+1, \ldots, s$.
- $\phi_{a,k}(M) = \phi_{a,k}(\operatorname{Int}(H)) \cup \phi_{a,k}(U)$ is an open semialgebraic subset of D(H), because $\phi_{a,k}(\operatorname{Int}(H)) = \operatorname{Int}(H_+)$ is an open semialgebraic subset of D(H), $\varphi|_U : U \to U'$, $F_{a,k} : \partial H \times \mathbb{R} \to \partial H \times \mathbb{R}$ and $\psi|_V : V \to V'$ are \mathcal{C}^{2k-1} semialgebraic diffeomorphisms and $F_{a,k}(U') \subset V'$ is an open semialgebraic set.
- $\phi_{a,k}(Y_i) \subset D(Y_i \cap H) = (Y_i \times \mathbb{R}) \cap D(H)$. This inclusion follows from the equality (5.1) and because the retraction ρ is compatible with Y.

It only remains to show that if a is small enough, then $\phi_{a,k}|_H$ is close to $(\pi|_{H_+})^{-1}$ and $\phi_{a,k}$ is close to $(\rho, 0)$ on $M \setminus H$.

Let us check: $\xi(y,s) := \psi \circ (\pi|_{H_+})^{-1} \circ \varphi^{-1}(y,s) = (y,\sqrt{s}).$

Let $x \in U$ be such that $(y,s) = \varphi(x) = (\rho(x), h(x))$, so $(\pi|_{H_+})^{-1}(x) = (x, \sqrt{h(x)})$ and $\psi(x, \sqrt{h(x)}) = (\rho(x), \sqrt{h(x)}) = (y, \sqrt{x})$, so $\xi(y, s) = (y, \sqrt{s})$.

Let $\eta: M \to \mathbb{R}$ be a strictly positive continuous semialgebraic function and let us choose a to guarantee that $\|\phi_{a,k}\|_H - (\pi|_{H_+})^{-1}\| < \eta|_H$. In fact, as ψ, φ are Nash diffeomorphisms, it is enough to check: If $b: \varphi(U) \to \mathbb{R}$ is a strictly positive semialgebraic function, there exists a strictly positive semialgebraic function $a: \partial H \to \mathbb{R}$ such that $a < \varepsilon$ and $\|F_{a,k}\|_{\varphi(U)} - \xi\| < b$.

As $F_{a,k}$ coincides with ξ outside $\{(y,s) \in \partial H \times \mathbb{R} : s \in [0, \varepsilon(y))\}$, we will find $a < \frac{3}{4}\varepsilon < 1$. Let us work on $K := \{(y,s) \in \partial H \times \mathbb{R} : s \in [0, \frac{3}{4}\varepsilon(y)]\}$. We claim: The projection $\pi_1 : K \to \partial H$ is open, proper and surjective. The semialgebraic map π_1 is surjective, because $\partial H \times \{0\} \subset K$. To prove that π_1 is open, let $A \subset K$ be an open set and let $y_0 \in \pi_1(A)$. Let $s_0 \in [0, \frac{3}{4}\varepsilon(y_0)]$ be such that $(y_0, s_0) \in A$. As A is open, there exists $\zeta > 0$ and $B \subset \partial H$ open such that $y_0 \in B$ and $(B \times [s_0 - \zeta, s_0 + \zeta]) \cap K \subset A$. In particular, $(y_0, s) \in A$ and $\pi_1(y_0, s) = y_0$ for each $s \in [\max\{s_0 - \zeta, 0\}, \min\{s_0 + \zeta, \frac{3}{4}\varepsilon(y_0)]$. Let s_1 be the midpoint of the previous interval and observe that $s_1 \in (0, \frac{3}{4}\varepsilon(y_0)) \cap (s_0 - \zeta, s_0 + \zeta)$. Let ζ' be such that $(s_1 - \zeta', s_1 + \zeta') \subset (0, \frac{3}{4}\varepsilon(y_0)) \cap (s_0 - \zeta, s_0 + \zeta)$. As ε is continuous and $s_1 + \zeta' < \frac{3}{4}\varepsilon(y_0)$, we may assume (after shrinking B if necessary) that $\frac{3}{4}\varepsilon(B) \subset [s_1 + \zeta', +\infty)$. Thus, $B \times (s_1 - \zeta', s_1 + \zeta') \subset (B \times [s_0 - \zeta, s_0 + \zeta]) \cap K \subset A$. As $\pi_1(B \times (s_1 - \zeta', s_1 + \zeta')) = B$ is an open neighborhood of y_0 in ∂H , we deduce $\pi_1(A)$ is an open subset of ∂H , so π_1 is an open map.

We show now that π_1 is proper. Let $C \subset \partial H$ be a compact set and let $\lambda := \max\{\frac{3}{4}\varepsilon|_C\}$. Then $\pi_1^{-1}(C) \subset C \times [0, \lambda]$, which is a compact set, so $\pi^{-1}(C)$ is also compact. Thus, π_1 is proper.

As π_1 is open, closed and surjective, by [FG, Const.3.1] the function

$$a: \partial H \to \mathbb{R}, \ y \mapsto \frac{1}{2} \min\left\{b(y,s)^2: \ s \in \left[0, \frac{3}{4}\varepsilon(y)\right]\right\}$$

is strictly positive, continuous and semialgebraic. As the last component of $F_{a,k}$ is strictly increasing on [0, a(y)] and $f_a(a(y)) = \sqrt{a(y)}$, we have

$$\begin{aligned} \|F_{a,k}(y,s) - (y,\sqrt{s})\| &= \begin{cases} 0 < b(y,s) & \text{if } a(y) \le s \le \frac{3}{4}\varepsilon(y), \\ |f_a(s) - \sqrt{s}| = \sqrt{s} - f_a(s) < \sqrt{s} < \sqrt{a(y)} \le b(y,s) & \text{if } 0 \le s < a(y). \end{cases} \end{aligned}$$

Let us check: $\phi_{a,k}$ is close to $(\rho, 0)$ on $M \setminus H$. If we denote $\tau := \psi \circ (\rho, 0) \circ \varphi^{-1}$, it is enough to prove: $||F_{a,k}(y,s) - \tau(y,s)||$ if -a(y) < s < 0.

Observe that $\tau(y, s) = (y, 0)$. Indeed, let $x \in U$ be such that $(y, s) = \varphi(x) = (\rho(x), h(x))$. Thus, $\psi(\rho(x), 0) = (\rho(\rho(x)), 0) = (\rho(x), 0) = (y, 0)$, so $\tau(y, s) = (y, 0)$. In addition, if -a(y) < s < 0, we have (as $a < \frac{3}{4}\varepsilon(y) < 1$)

$$||F_{a,k}(y,s) - \tau(y,s)|| = ||(y,s) - (y,0)|| = -s < a(y) < \sqrt{a(y)} \le b(y,s).$$

STEP 5. Let us see now how to obtain the desired Nash embedding. Recall that Y_1, \ldots, Y_r are the irreducible components of Y that meet ∂H but are not contained in ∂H . Consider the Nash normal-crossings divisor $\overline{Y} := \bigcup_{i=1}^r Y_i \subset M$. Recall that $U \cap \bigcup_{j=r+1}^s Y_j = \emptyset$, so $\phi_{a,k}|_{\bigcup_{j=r+1}^s Y_j} = (\pi|_{H_+})^{-1}|_{\bigcup_{j=r+1}^s Y_j}$. By [BFR, Thm.1.6, Thm 1.7], up to take k big enough, we can approximate the restriction $\phi_{a,k}|_Y : Y \to D(H)$ by a Nash map $\hat{\phi} : Y \to D(H)$ such that $\hat{\phi}|_{\bigcup_{j=r+1}^s Y_j} = (\pi|_{H_+})^{-1}|_{\bigcup_{j=r+1}^s Y_j}, \hat{\phi}|_{\partial H} = (\mathrm{id}_{\partial H}, 0)$ and $\hat{\phi}|_{\overline{Y}} : \overline{Y} \to D(H)$ satisfies $\hat{\phi}(Y_i) \subset D(Y_i \cap H)$ for $i = 1, \ldots, r$. By [BFR, Prop.8.2] we can extend $\hat{\phi}$ to a global Nash map $\phi : M \to D(H)$, that is, up to take k big enough, close to $\phi_{a,k}$ in the \mathcal{C}^1 semialgebraic topology. Thus, by [Sh, Lem.II.1.7] the map ϕ is a Nash embedding. We may assume in addition, $\phi(M) = \phi_{a,k}(M)$, which is an open semialgebraic subset of D(H).

STEP 6. We show: There exists an open semialgebraic neighborhood $W \subset M$ of ∂H such that $\phi|_{\operatorname{Cl}(W)\cup H} : \operatorname{Cl}(W) \cup H \to D(H)$ is proper. Once this is done we conclude that, up to take a smaller semialgebraic function a > 0 if necessary, the Nash embedding $\phi : M \to D(H)$ defined in STEP 5 satisfies all the other required properties.

As $\phi(M)$ is open in D(H), the set $C := D(H) \setminus \phi(M)$ is a closed subset of D(H) that does not intersect ∂H . Let V be an open semialgebraic neighborhood of ∂H in $\phi(M)$ such that $\operatorname{Cl}(V) \cap C = \emptyset$. The set $W := \phi^{-1}(V)$ is an open semialgebraic neighborhood of ∂H in M such that $\pi(\operatorname{Cl}(W)) = \operatorname{Cl}(V)$ because $\phi : M \to \phi(M)$ is a semialgebraic homeomorphism. The restriction $\phi|_{\operatorname{Cl}(W)\cup H} : \operatorname{Cl}(W) \cup H \to D(H)$ satisfies $\phi(\operatorname{Cl}(W) \cup H) = \operatorname{Cl}(V) \cup H_+$ because $\phi(\operatorname{Cl}(W)) = \operatorname{Cl}(V)$ and $\phi(H) = H_+$. As $\operatorname{Cl}(V) \cup H_+$ is closed in D(H), if $K \subset D(H)$ is compact, then the set $K \cap (\operatorname{Cl}(V) \cup H_+)$ is compact. The map $\phi|_{\operatorname{Cl}(W)\cup H} : \operatorname{Cl}(W) \cup H \to \operatorname{Cl}(V) \cup H_+ \subset$ D(H) is a semialgebraic homeomorphism, so $\phi^{-1}(K \cap (\operatorname{Cl}(V) \cup H_+))$ is a compact set. We conclude that the restriction $\phi|_{\operatorname{Cl}(W)\cup H} : \operatorname{Cl}(W) \cup H \to D(H)$ is proper, as required. \Box

As a straightforward consequence of the previous embedding theorem, we have the following:

Corollary 5.11. The composition $f := \pi \circ \phi : M \to H$ is a Nash map with the following properties:

- (i) $f(Y_i) \subset Y_i$ for $i = 1, \ldots, r$ and $f|_{Y_i} = \operatorname{id}_{Y_i}$ for each $i = r + 1, \ldots, s$.
- (ii) $f|_H$ is a semialgebraic homeomorphism close to id_H . Moreover $f|_{Int(H)}$ is a Nash diffeomorphism and $f|_{\partial H} = id_{\partial H}$.
- (iii) $f|_{M\setminus H}: M \setminus H \to H$ is a Nash embedding close to $\rho|_{M\setminus H}$.
- (iv) At each point $x \in \partial H$ the map f has a local presentation of the type

$$(y_1,\ldots,y_d)\mapsto (y_1^2,y_2,\ldots,y_d).$$

(v) There exists an open semialgebraic neighborhood $V \subset M$ of ∂H , such that $f|_{\operatorname{Cl}(V)\cup H}$ is a proper Nash map.

Proof. Statements (i), (ii) and (iii) follow from Theorem 5.10, whereas statement (iv) is a consequence of Lemma 5.8(iii) and the fact that $\phi : M \to \phi(M) \subset D(H)$ is a Nash diffeomorphism onto its image. To prove (v), we use that π is proper (see Lemma 5.8(iv)) and that ϕ is proper on $\operatorname{Cl}(V) \cup H$ where $V \subset M$ is a small semialgebraic neighborhood of ∂H in M.

5.6. Folding all the components of the boundary. We are almost ready to prove Theorem 1.7. Before, we need a technical lemma to confront local models:

Lemma 5.12. Consider the Nash map $\varphi : \mathbb{R}^d :\to \mathbb{R}^d$, $(x_1, \ldots, x_d) \mapsto (x_1^2, x_2, \ldots, x_d)$ and let $\psi_1, \psi_2 : \mathbb{R}^d \to \mathbb{R}^d$ be Nash diffeomorphisms such that $\psi_i(\{\mathbf{x}_j = 0\}) = \{\mathbf{x}_j = 0\}$ for $j = 1, \ldots, s$ and i = 1, 2. Then there exist Nash functions $h_j : \mathbb{R}^d \to \mathbb{R}$ each one strictly positive around $\{\mathbf{x}_j = 0\}$ and a Nash map $g : \mathbb{R}^d \to \mathbb{R}^{d-s}$ such that $g|_{\{\mathbf{x}_1=0,\ldots,\mathbf{x}_s=0\}} : \{\mathbf{x}_1 = 0,\ldots,\mathbf{x}_s = 0\} \to \mathbb{R}^{d-s}$ is a Nash diffeomorphism and $\psi_2 \circ \varphi \circ \psi_1 = ((\mathbf{x}_1h_1)^2, \mathbf{x}_2h_2,\ldots,\mathbf{x}_sh_s, g).$

Proof. As $\psi_i(\{\mathbf{x}_j = 0\}) = \{\mathbf{x}_j = 0\}$ for $j = 1, \dots, s$ and ψ_i is a Nash diffeomorphism, then $\psi_i|_{\{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\}} : \{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\} \rightarrow \{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\}$ is a Nash diffeomorphism. Thus,

$$\psi_2 \circ \varphi \circ \psi_1|_{\{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\}} : \{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\} \to \{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\}$$

is a Nash diffeomorphism. Let $\pi : \mathbb{R}^d \to \mathbb{R}^{d-s}$, $(x_1, \ldots, x_d) \mapsto (x_{s+1}, \ldots, x_d)$ and let $g := \pi \circ \psi_2 \circ \varphi \circ \psi_1$. We have

$$g|_{\{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\}}: \{\mathbf{x}_1=0,\dots,\mathbf{x}_s=0\} \to \mathbb{R}^{d-s}$$

is a Nash diffeomorphism and $\psi_2 \circ \varphi \circ \psi_1 = (g_1, \ldots, g_s, g)$ for some Nash functions $g_i : \mathbb{R}^d \to \mathbb{R}$. As $\psi_i(\{\mathbf{x}_j = 0\}) = \{\mathbf{x}_j = 0\}$, its *j*th component ψ_{ij} is divisible by \mathbf{x}_j and thus, there exists a Nash function $h_{ij} : \mathbb{R}^d \to \mathbb{R}$ that is strictly positive around $\{\mathbf{x}_j = 0\}$ such that $\psi_{ij} = \mathbf{x}_j h_{ij}$ for i = 1, 2 and $j = 1, \ldots, s$. Thus,

$$\begin{split} \psi_{2} \circ \varphi \circ \psi_{1} &= \\ (\psi_{11}^{2}h_{21}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),\psi_{12}h_{22}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),\cdots,\psi_{1s}h_{2s}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),g) \\ &= ((\mathbf{x}_{1}h_{11})^{2}h_{21}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),\mathbf{x}_{2}h_{12}h_{22}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),\ldots \\ &\dots,\mathbf{x}_{s}h_{1s}h_{2s}(\psi_{11}^{2},\psi_{12},\ldots,\psi_{1d}),g). \end{split}$$

It is enough to take $h_1 := h_{11}\sqrt{h_{21}(\psi_{11}^2, \psi_{12}, \dots, \psi_{1d})}$ and $h_j := h_{1j}h_{2j}(\psi_{11}^2, \psi_{12}, \dots, \psi_{1d})$ for $j = 2, \dots, s$.

We next prove Theorem 1.7.

Proof of Theorem 1.7. Let M be a Nash envelope of Q and Y_1, \ldots, Y_ℓ the irreducible components of the Nash closure Y of ∂Q in M (Section 5.3). By Lemma 5.4 there exist, after shrinking Mif necessary, Nash equations $h_i: M \to \mathbb{R}$ of Y_i such that $d_x h_i: T_x M \to \mathbb{R}$ is surjective for each $x \in Y_i$ and $H_i := h_i^{-1}([0, +\infty))$ is a Nash manifold with boundary whose boundary is Y_i for each $i = 1, \ldots, \ell$. By Corollary 5.11 we have (after shrinking M is necessary) for each index i a proper Nash map $f_i: Cl(M) \to H_i$ such that:

- (i) $f_i(Y_j) \subset Y_j$ for $j = 1, \ldots, \ell$ and $f_i|_{Y_i} = \mathrm{id}_{Y_i}$,
- (ii) $f_i|_{H_i} : H_i \to H_i$ is a semialgebraic homeomorphism close to the identity map, whose restriction $f_i|_{\operatorname{Int}(H_i)} : \operatorname{Int}(H_i) \to \operatorname{Int}(H_i)$ is a Nash diffeomorphism and $f_i|_{\partial H_i} = \operatorname{id}_{\partial H_i}$,
- (iii) $f_i|_{M\setminus H_i}: M\setminus H_i \to \operatorname{Int}(H_i)$ is a semialgebraic embedding close to $\rho_i|_{M\setminus H_i}: M\setminus H_i \to \partial H_i$,
- (iv) f_i has local representations $(y_1, \ldots, y_d) \mapsto (y_1^2, y_2, \ldots, y_d)$ at each point $x \in Y_i$.

Let $\{C_k\}_k$ be the family of the connected components of $M \setminus Y$ such that $\operatorname{Cl}(C_k) \cap \mathfrak{Q} = \emptyset$. Then $M \setminus \bigcup_{k=1} \operatorname{Cl}(C_k)$ is an open semialgebraic neighborhood of \mathfrak{Q} in M. Let us substitute M by $M' := M \setminus \bigcup_k \operatorname{Cl}(C_k)$ and Y by $Y \cap M'$. Now, the connected components C of $M \setminus Y$ satisfy $\operatorname{Cl}(C) \cap \mathfrak{Q} \neq \emptyset$, so $\operatorname{Cl}(C) \cap H_i \neq \emptyset$ for each $i = 1, \ldots, \ell$ (because $\mathfrak{Q} = H_1 \cap \cdots \cap H_\ell$ by Lemma 5.6).

Consider the proper Nash map $f := f_{\ell} \circ \cdots \circ f_1 : \operatorname{Cl}(M) \to \operatorname{Cl}(M)$. Let us check that f satisfies the following properties:

- (1) $f(Y_i) \subset Y_i$ for each $i = 1, \ldots, \ell$.
- (2) $f|_{\Omega} : \Omega \to \Omega$ is a semialgebraic homeomorphism close to the identity map, whose restriction $f|_{\mathrm{Int}(\Omega)} : \mathrm{Int}(\Omega) \to \mathrm{Int}(\Omega)$ is a Nash diffeomorphism,
- (3) $f(\operatorname{Cl}(M)) = Q$,
- (4) f has local representations $(y_1, \ldots, y_d) \mapsto (y_1^2, \ldots, y_s^2, y_{s+1}, \ldots, y_d)$ at each point $x \in \partial \Omega$. The integer $s \ge 1$ depends on the point x and corresponds to the number of irreducible components of Y that pass through x.

Property (1) follows straightforwardly from (i).

To prove (2) we show first: $f_i(Q) = Q$ and $f_i(\operatorname{Int}(Q)) = \operatorname{Int}(Q)$ for $i = 1, \ldots, \ell$. Once this is done, as $Q = H_1 \cap \cdots \cap H_\ell$, $f_i|_{H_i} : H_i \to H_i$ is a semialgebraic homomorphism close to the identity map for each i and $f_i|_{\operatorname{Int}(H_i)} : \operatorname{Int}(H_i) \to \operatorname{Int}(H_i)$ is a Nash diffeomorphism, we deduce that $f|_Q : Q \to Q$ is a semialgebraic homeomorphism close to the identity map and its restriction $f|_{\operatorname{Int}(Q)} : \operatorname{Int}(Q) \to \operatorname{Int}(Q)$ is a Nash diffeomorphism.

As f_i is proper and $\Omega = \operatorname{Cl}(\operatorname{Int}(\Omega))$, we have $f_i(\Omega) = f_i(\operatorname{Cl}(\operatorname{Int}(\Omega))) = \operatorname{Cl}(f_i(\operatorname{Int}(\Omega)))$, so it is enough to prove $f_i(\operatorname{Int}(\Omega)) = \operatorname{Int}(\Omega)$.

As the map $f_i|_{\operatorname{Int}(H_i)}$: $\operatorname{Int}(H_i) \to \operatorname{Int}(H_i)$ is a Nash diffeomorphism and $\operatorname{Int}(\mathfrak{Q}) \subset \operatorname{Int}(H_i)$, we have $f_i(\operatorname{Int}(\mathfrak{Q})) \subset \operatorname{Int}(H_i)$. As $f_i(Y_j) \subset Y_j$, we deduce $f_i(Y_j \cap \operatorname{Int}(H_i))$ is a closed Nash submanifold of the Nash manifold $Y_j \cap \operatorname{Int}(H_i)$ and both have the same dimension. Consequently, the image of each connected component of $f_i(Y_j \cap \operatorname{Int}(H_i))$ is a connected component of $Y_j \cap$ $\operatorname{Int}(H_i)$. As f_i is close to the identity map, $f_i(D) = D$ for each connected component of $Y_j \cap \operatorname{Int}(H_i)$. Thus, $f_i(Y_j \cap \operatorname{Int}(H_i)) = Y_j \cap \operatorname{Int}(H_i)$ and $f_i(Y \cap \operatorname{Int}(H_i)) = Y \cap \operatorname{Int}(H_i)$. Observe that $\operatorname{Int}(\mathfrak{Q})$ is a union of connected components of $\operatorname{Int}(H_i) \setminus Y$ because $\mathfrak{Q} \cap Y = \partial \mathfrak{Q}$. As $f_i|_{\operatorname{Int}(H_i)} : \operatorname{Int}(H_i) \to \operatorname{Int}(H_i)$ is a Nash diffeomorphism and $f_i(Y \cap \operatorname{Int}(H_i)) = Y \cap \operatorname{Int}(H_i)$, we deduce that $f_i(\operatorname{Int}(\mathfrak{Q}))$ is also a union of connected components of $\operatorname{Int}(H_i) \setminus Y$. As $f_i|_{\operatorname{Int}(\mathfrak{Q})}$ is close to the identity map, $f_i(C) = C$ for each connected component of $\operatorname{Int}(\mathfrak{Q})$. In particular, $f_i(\operatorname{Int}(\mathfrak{Q})) = \operatorname{Int}(\mathfrak{Q})$, as claimed.

Next, we prove (3). Let us show: $f_i(H_j) \subset H_j$ for each i, j. Once this is proved, we have $f_i(H_j) \subset H_i \cap H_j$ for each i, j (because $f_i(\operatorname{Cl}(M)) \subset H_i$) and:

$$f(\operatorname{Cl}(M)) = (f_{\ell} \circ \cdots \circ f_1)(\operatorname{Cl}(M)) = (f_{\ell} \circ \cdots \circ f_2)(H_1)$$

$$\subset (f_{\ell} \circ \cdots \circ f_3)(H_1 \cap H_2) \subset \cdots \subset H_1 \cap \cdots \cap H_{\ell} = \mathfrak{Q}.$$

As f(Q) = Q, we conclude f(Cl(M)) = Q.

For simplicity we show: $f_2(H_1) \subset H_1$. As f_2 is continuous, it is enough to prove: $f_2(\text{Int}(H_1 \cap H_2)) \subset \text{Int}(H_1 \cap H_2)$ and $f_2(\text{Int}(H_1) \setminus H_2) \subset \text{Int}(H_1)$.

To prove the inclusion $f_2(\text{Int}(H_1 \cap H_2)) \subset \text{Int}(H_1 \cap H_2)$, it is enough to consider the Nash manifold with corners $H_1 \cap H_2$ and to proceed analogously as above with f_i and \mathfrak{Q} .

We show next: $f_2(\text{Int}(H_1) \setminus H_2) \subset \text{Int}(H_1)$. To that end, we prove: $f_2(C) \subset \text{Int}(H_1)$ for each connected component C of $\text{Int}(H_1) \setminus H_2$.

As $f_2|_{M\setminus H_2} : M \setminus H_2 \to \operatorname{Int}(H_2)$ is a Nash embedding and $f_2(Y_1) \subset Y_1$, we deduce that each connected component C of $\operatorname{Int}(H_1) \setminus H_2$ is mapped under f_2 into a connected component of $\operatorname{Int}(H_2) \setminus Y_1$. Let C be a connected component of $\operatorname{Int}(H_1) \setminus H_2 \subset M \setminus (Y_1 \cup Y_2)$. As $M \setminus Y_2 = (M \setminus H_2) \sqcup \operatorname{Int}(H_2)$ and $\operatorname{Int}(H_1) = H_1 \setminus Y_1$, we deduce that $\operatorname{Int}(H_1) \setminus H_2 = (H_1 \setminus \operatorname{Int}(H_2)) \cap (M \setminus (Y_1 \cup Y_2))$ is open and closed in $M \setminus (Y_1 \cup Y_2)$, so C is a connected component of $M \setminus (Y_1 \cup Y_2)$.

CASE 1. $\operatorname{Cl}(C) \cap Y_2 \neq \emptyset$. As C is a connected component of $M \setminus (Y_1 \cup Y_2)$ and $Y_1 \cup Y_2$ is a Nash normal-crossings divisor, there exists a point $x \in \operatorname{Cl}(C) \cap (Y_2 \setminus Y_1)$. Let V be a semialgebraic neighborhood of x in M that does not meet Y_1 . As $x \in \operatorname{Cl}(C)$ and $C \subset \operatorname{Int}(H_1)$, we deduce $V \subset \operatorname{Int}(H_1)$. As $\rho_2(x) = x \in \operatorname{Int}(H_1)$, and $f_2|_{M \setminus H_2} : M \setminus H_2 \to \operatorname{Int}(H_2)$ is close to $\rho_2|_{M \setminus H_2} :$ $M \setminus H_2 \to \partial H_2$, we may assume $f_2(V) \subset \operatorname{Int}(H_1)$. Thus, $f_2(C)$ meets $\operatorname{Int}(H_1)$. As $f_2(C)$ is a connected subset of $\operatorname{Int}(H_2) \setminus Y_1$ and $\operatorname{Int}(H_1) \cap \operatorname{Int}(H_2) = H_1 \cap (\operatorname{Int}(H_2) \setminus Y_1)$ is open and closed in $\operatorname{Int}(H_2) \setminus Y_1$ and $f_2(C)$ meets one of the connected components of $\operatorname{Int}(H_1) \cap \operatorname{Int}(H_2)$, we deduce $f_2(C)$ is contained in one of the connected components of $\operatorname{Int}(H_1) \cap \operatorname{Int}(H_2)$. Consequently, $f_2(C) \subset \operatorname{Int}(H_1) \cap \operatorname{Int}(H_2) \subset \operatorname{Int}(H_1)$.

CASE 2. $\operatorname{Cl}(C) \cap Y_2 = \emptyset$. As $C \subset H_1 \setminus H_2$, we have

$$Cl(C) \cap H_2 = Cl(C) \cap (Y_2 \cup Int(H_2)) = Cl(C) \cap Int(H_2)$$
$$= Cl(C \cap Int(H_2)) \cap Int(H_2) \subset Cl(C \cap H_2) = \emptyset.$$

Observe that C is the closure in $M \setminus (Y_1 \cup Y_2)$ of the union of some connected components of $M \setminus Y$, which is a contradiction because by construction each of the connected components of $M \setminus Y$ meets H_2 .

Thus, $\operatorname{Cl}(C) \cap Y_2 \neq \emptyset$ for each connected component of $\operatorname{Int}(H_1) \setminus H_2$. Consequently, $f_2(C) \subset \operatorname{Int}(H_1)$ for each connected component C of $\operatorname{Int}(H_1) \setminus H_2$, so $f_2(\operatorname{Int}(H_1) \setminus H_2) \subset \operatorname{Int}(H_1)$.

Finally, we prove (4). Pick a point $x \in \partial \Omega$ and assume that x belongs exactly to Y_1, \ldots, Y_s . Recall that the analytic closure of $\partial \Omega_x$ is $Y_{1,x} \cup \cdots \cup Y_{s,x}$ and there exists an open semialgebraic neighborhood $U \subset M$ of x equipped with a Nash diffeomorphism $u : U \to \mathbb{R}^d$ such that u(x) = 0, $u(Y_j \cap U) = \{\mathbf{x}_j = 0\}$ and $u(\Omega \cap U) = \{\mathbf{x}_1 \ge 0, \ldots, \mathbf{x}_s \ge 0\}$.

Observe that $\eta := u \circ f \circ u^{-1} = \eta_{\ell} \circ \cdots \circ \eta_1$ where $\eta_k := u \circ f_k \circ u^{-1} : \mathbb{R}^d \to \mathbb{R}^d$ is either a Nash diffeomorphism or a Nash map that has local representations $(y_1, \ldots, y_d) \mapsto (y_1^2, y_2, \ldots, y_d)$ at each point $x \in Y_k$. In addition, these local representations preserve the local representations of Y_1, \ldots, Y_s , so Y_1, \ldots, Y_s correspond to coordinates' hyperplanes in these (local) coordinates. By Lemma 5.12 we may assume that $\eta = \sigma_s \circ \cdots \circ \sigma_1$ is a composition of Nash maps of the type $\sigma_m : \mathbb{R}^d \to \mathbb{R}^d$,

$$(x_1, \ldots, x_d) \mapsto (x_1 h_{1m}, \ldots, x_{m-1} h_{m-1,m}, x_m^2 h_{mm}^2, x_{m+1} h_{m+1,m}, \ldots, x_s h_{sm}, g_m)$$

where $h_{jm} : \mathbb{R}^d \to \mathbb{R}$ is a Nash function that does not vanish around $\{\mathbf{x}_j = 0\}$ and $g_m : \mathbb{R}^d \to \mathbb{R}^{d-s}$ satisfies $g_m|_{\{\mathbf{x}_1=0,\ldots,\mathbf{x}_s=0\}} : \{\mathbf{x}_1 = 0,\ldots,\mathbf{x}_s = 0\} \to \mathbb{R}^{d-s}$ is a Nash diffeomorphism. Thus, there exist Nash functions $f_j : \mathbb{R}^d \to \mathbb{R}$ that do not vanish around $\{\mathbf{x}_j = 0\}$ and a Nash map $g : \mathbb{R}^d \to \mathbb{R}^{d-s}$ such that $g|_{\{\mathbf{x}_1=0,\ldots,\mathbf{x}_s=0\}} : \{\mathbf{x}_1 = 0,\ldots,\mathbf{x}_s = 0\} \to \mathbb{R}^{d-s}$ is a Nash diffeomorphism satisfying

$$\eta: \mathbb{R}^d \to \mathbb{R}^d, \ (x_1, \dots, x_d) \mapsto (x_1^2 f_1^2, \dots, x_s^2 f_s^2, g).$$

Observe that the map

$$\psi : \mathbb{R}^d \to \mathbb{R}^d, \ (x_1, \dots, x_d) \mapsto (x_1 f_1, \dots, x_s f_s, g)$$

is a Nash diffeomorphism around the origin because the determinant of its Jacobian matrix at the origin

$$J_{\psi}(x) = \begin{bmatrix} f_1(0) & \cdots & 0 & 0 \\ \vdots & \ddots & \vdots & \vdots \\ 0 & \cdots & f_s(0) & 0 \\ \hline * & \cdots & * & J_g(0) \end{bmatrix}$$

is $f_1(0) \cdots f_s(0) J_g(0) \neq 0$. After shrinking the open semialgebraic neighborhood $U \subset M$ of x, there exists a Nash diffeomorphism $\phi : \mathbb{R}^d \to \mathbb{R}^d$ such that $\psi \circ \phi = \mathrm{id}_{\mathbb{R}^d}$. As $\eta = \theta \circ \psi$, where

$$\theta: \mathbb{R}^d \to \mathbb{R}^d, \ (x_1, \dots, x_d) \mapsto (x_1^2, \cdots, x_s^2, x_{s+1}, \dots, x_d),$$

we deduce that $\eta \circ \phi = \theta$, so f has local representation $(y_1, \ldots, y_d) \mapsto (y_1^2, \ldots, y_s^2, y_{s+1}, \ldots, y_d)$ at $x \in \partial \Omega$. Recall that $s \ge 1$ corresponds to the number of irreducible components of Y that passes through x, as required. \Box

5.7. The Nash double of a Nash manifold with corners. Let $\mathcal{Q} \subset \mathbb{R}^n$ be a *d*-dimensional Nash manifold with corners and M a Nash envelope of \mathcal{Q} such that the Nash closure Y of $\partial \mathcal{Q}$ is a Nash normal-crossings divisor of M and $\mathcal{Q} \cap Y = \partial \mathcal{Q}$. Let Y_1, \ldots, Y_ℓ be the irreducible components of Y and $h_i : M \to \mathbb{R}$ Nash equations of Y_i for $i = 1, \ldots, \ell$ such that $d_x h_i : T_x M \to \mathbb{R}$ is surjective for $i = 1, \ldots, \ell$. By Lemma 5.4 the sets $H_i := h_i^{-1}([0, +\infty))$ are Nash manifolds with boundary Y_i that contain \mathcal{Q} as a closed subset. Write (Lemma 5.6) $\mathcal{Q} = H_1 \cap \ldots \cap H_\ell = \{h_1 \geq 0, \ldots, h_\ell \geq 0\} \subset M, t := (t_1, \ldots, t_\ell)$ and define the Nash double of \mathcal{Q} as:

$$D(Q) := \{ (x,t) \in M \times \mathbb{R}^{\ell} : t_1^2 - h_1(x) = 0, \dots, t_{\ell}^2 - h_{\ell}(x) = 0 \}.$$

Observe that $D(\mathfrak{Q})$ only depends on the Nash equations h_1, \ldots, h_ℓ , which are unique up to multiplication by strictly positive Nash functions. Thus, $D(\mathfrak{Q})$ is unique up to Nash diffeomorphism. In addition, the reader can check that $D(\mathfrak{Q})$ is the inverse image of the value 0 under the Nash map $f: M \times \mathbb{R}^{\ell} \to \mathbb{R}^{\ell}, (x, t) \mapsto (t_1^2 - h_1(x), \ldots, t_{\ell}^2 - h_{\ell}(x))$ and that 0 is a regular value for f, so $D(\mathfrak{Q})$ is a *d*-dimensional Nash manifold.

Theorem 5.13 (Nash double of a Nash manifold with corners). The Nash subset $D(\Omega)$ of $M \times \mathbb{R}^{\ell}$ is a d-dimensional Nash manifold that satisfies the following properties:

- (i) The projection $\pi: D(\mathfrak{Q}) \to \mathfrak{Q}, (x,t) \mapsto x$ is proper and surjective.
- (ii) The semialgebraic set Q^ℓ := D(Q) ∩ {t₁ ≥ 0,...,t_ℓ ≥ 0} is a Nash manifold with corners, which is Nash diffeomorphic to Q and the Nash closure of ∂Q^ℓ in D(Q) is the Nash normal-crossings divisor Y^ℓ := U^ℓ_{i=1}{t_i = 0} of D(Q).
- (iii) The restriction $\pi|_{\Omega^{\ell}} : \Omega^{\ell} \to \Omega$ is a semialgebraic homeomorphism.
- (iv) The restriction $\pi|_{\operatorname{Int}(\mathbb{Q}^{\ell})}$: $\operatorname{Int}(\mathbb{Q}^{\ell}) \to \operatorname{Int}(\mathbb{Q})$ is a Nash diffeomorphism.
- (v) There exists an open semialgebraic neighborhood $W \subset D(\Omega)$ of Ω^{ℓ} Nash diffeomorphic to M.

Proof. The proof is conducted in several steps:

STEP 1. Let $(D(H_1), \pi_1)$ be the Nash double of $H_1 := \{h_1 \ge 0\}$. Recall that

 $M^1 := D(H_1) = \{(x,t_1) \in M \times \mathbb{R} : \ t_1^2 - h_1(x) = 0\} \quad \text{and} \quad \pi_1 : D(H_1) \to H_1, \ (x,t_1) \to x.$

The Nash map π_1 is proper by Lemma 5.8(iv). Consider the (d-1)-dimensional Nash submanifolds $Y_1^1 := Y_1 \times \{0\}$ and $Y_i^1 := D(Y_i \cap H_1) = (Y_i \times \mathbb{R}) \cap D(H_1)$ (see (5.1)) of M^1 for $i = 2, \ldots, \ell$. Consider the Nash functions

$$h_i^1: M^1 \to \mathbb{R}, \ (x, t_1) \mapsto h_i^1(x, t_1) := \begin{cases} t_1 & \text{if } i = 1, \\ h_i(x) & \text{if } i = 2, \dots, \ell \end{cases}$$

The tangent space to M^1 at a point $(x, t_1) \in M^1$ is:

$$T_{(x,t_1)}M^1 = \{(v,s_1) \in T_xM \times \mathbb{R} : d_xh_1(v) - 2t_1s_1 = 0\}.$$

We claim: $h_i^1: M^1 \to \mathbb{R}$ is a Nash equation of Y_i^1 in M^1 such that $d_{(x,t_1)}h_i^1: T_{(x,t_1)}M^1 \to \mathbb{R}$ is surjective for each $(x,t_1) \in Y_i^1$ and $i = 1, \ldots, \ell$.

Suppose first i = 1. Then $\{h_1^1 = 0\} = \{h_1(x) = 0, t_1 = 0\} = Y_1 \times \{0\} = Y_1^1$, so h_i^1 is a Nash equation of Y_i^1 in M^1 . In addition, for $(x, 0) \in Y_1^1 = Y_1 \times \{0\}$ we have

$$T_{(x,0)}M^1 = \{(v,s_1) \in T_xM \times \mathbb{R} : d_xh_1(v) = 0\} = T_xY_1 \times \mathbb{R}$$

and $d_{(x,0)}h_1^1: T_xY_1 \times \mathbb{R} \to \mathbb{R}, \ (v,s_1) \mapsto s_1$ is surjective.

Suppose next $i = 2, \ldots, \ell$ and observe that

$$\begin{split} Y_i^1 = \{(x,t_1) \in M \times \mathbb{R} : \ t_1^2 - h_1(x) = 0, h_i(x) = 0\} = \{(x,t_1) \in Y_i \times \mathbb{R} : \ t_1^2 - h_1(x) = 0\}, \\ \text{so } h_i^1 = 0 \text{ is a Nash equation of } Y_i^1 \text{ in } M^1. \text{ Recall that} \end{split}$$

$$T_{(x,t_1)}M^1 = \{(v,s_1) \in T_xM \times \mathbb{R} : d_xh_1(v) - 2t_1s_1 = 0\}$$

and $d_{(x,t_1)}h_i^1: T_{(x,t_1)}M^1 \to \mathbb{R}, \ (v,s_1) \mapsto d_x h_i(v)$. Let us check that it is surjective:

CASE 1: $t_1 \neq 0$. We pick a vector $v \in T_x M$ such that $d_x h_i(v) \neq 0$ and $(v, \frac{d_x h_1(v)}{2t_1}) \in T_{(x,t_1)} M^1$. We have

$$d_{(x,t_1)}h_i^1\left(v, \frac{d_x h_1(v)}{2t_1}\right) = d_x h_i(v) \neq 0.$$

CASE 2: $t_1 = 0$. We have $T_{(x,0)}M^1 = \{(v, s) \in T_xM \times \mathbb{R} : d_xh_1(v) = 0\}$. As $Y_1 \cup Y_i$ is a normal-crossings divisor, d_xh_1 and d_xh_i are linearly independent, so there exists $v \in T_xM$ such that $d_xh_i(v) \neq 0$ and $d_xh_1(v) = 0$. Observe that $(v, 0) \in T_{(x,0)}M^1$ and $d_{(x,t_1)}h_i^1(v, 0) = d_xh_i^1(v) \neq 0$.

Consequently, $d_{(x,t_1)}h_i^1: T_{(x,t_1)}M^1 \to \mathbb{R}$ is surjective (in both cases), as claimed.

By Theorem 5.10 there exist an open semialgebraic neighborhood $N^0 \subset M$ of H_1 and a Nash embedding $\phi_1 : N^0 \to M^1$ such that $\phi_1(H_1) = H_{1+}$ and $W^1 := \phi_1(N^0)$ is an open semialgebraic subset of M^1 that contains the Nash manifold with corners $\phi_1(\Omega)$ as a closed subset. In addition, $\phi_1|_{Y_1} = \operatorname{id}_{Y_1 \times \{0\}}, \phi_1(Y_i) \subset Y_i^1$ for $i = 2, \ldots, \ell$ and $\phi_1|_{H_1} : H_1 \to H_{1+}$ is close to $(\pi_1|_{H_{1+}})^{-1}$. Consider the Nash subset $Y^1 := \bigcup_{i=1}^{\ell} Y_i^1$ of M^1 , whose irreducible components are Nash submanifolds of M^1 . We have $\phi_1(Y) = Y^1 \cap W^1$. In addition, $\pi_{1-}|_{\operatorname{Int}(H_{1-})} : \operatorname{Int}(H_{1-}) \to$ $\operatorname{Int}(H_1)$ is a Nash diffeomorphism and $\pi_{1-}(Y^1 \cap \operatorname{Int}(H_{1-})) = Y \cap \operatorname{Int}(H_1)$. As Y is a Nash normal-crossings divisor of M and $M^1 = D(H_1) = W^1 \cup \operatorname{Int}(H_{1-})$, we conclude that Y^1 is a Nash normal-crossings divisor of M^1 .

Denote $H_1^1 := H_{1+}$ and $H_i^1 := \pi_1^{-1}(H_i) = \{h_i^1 \ge 0\}$ for $i = 2, ..., \ell$. By Lemma 5.6 the intersection

$$Q^{1} := \bigcap_{i=1}^{\ell} H_{i}^{1} = \{(x, t_{1}) \in M \times \mathbb{R} : t_{1}^{2} - h_{1}(x) = 0, t_{1} \ge 0, h_{2}(x) \ge 0, \dots, h_{\ell}(x) \ge 0\} \subset M^{1}$$

is a Nash manifold with corners and the Nash closure in M^1 of ∂Q^1 is Y^1 . Observe that $\pi_1(Q^1) = Q$, the restriction $\pi_1|_{Q^1} : Q^1 \to Q$ is a semialgebraic homeomorphism and the restriction $\pi_1|_{\operatorname{Int}(Q^1)} : \operatorname{Int}(Q^1) \to \operatorname{Int}(Q)$ is a Nash diffeomorphism. We claim: $\phi_1(Q) = Q^1$. As ϕ_1 is a Nash diffeomorphism, it is enough to prove: $\phi_1(H_1 \cap H_i) = H_1^1 \cap H_i^1 = H_{1+} \cap \pi_1^{-1}(H_i)$ for $i = 2, \ldots, \ell$.

As $\phi_1: N^0 \to W^1$ is a Nash diffeomorphism, it is enough to check

$$\phi_1(\operatorname{Int}(H_1) \cap \operatorname{Int}(H_i)) = \operatorname{Int}(H_{1+}) \cap \pi_1^{-1}(\operatorname{Int}(H_i)).$$

Observe that $\operatorname{Int}(H_1) \cap \operatorname{Int}(H_i)$ is a union of connected components of $N^0 \setminus (Y_1 \cup Y_i)$. In addition, $\phi_1(Y_1) = Y_1 \times \{0\}$ and $\phi_1(Y_i) \subset D(Y_i \cap H_1)$, so

$$\phi_1(Y_i \cap \operatorname{Int}(H_1)) \subset D(Y_i \cap H_1) \cap \operatorname{Int}(H_{1+}).$$

As $\phi_1|_{\operatorname{Int}(H_1)}$: $\operatorname{Int}(H_1) \to \operatorname{Int}(H_{1+})$ is a Nash diffeomorphism, $\phi_1(Y_i \cap \operatorname{Int}(H_1))$ is a closed Nash submanifold of the Nash manifold $D(Y_i \cap H_1) \cap \operatorname{Int}(H_{1+})$ of the same dimension. As ϕ_1 is close to $(\pi_1|_{H_1,+})^{-1}$, both Nash manifolds have the same number of connected components and

$$\phi_1(Y_1 \cap \operatorname{Int}(H_1)) = D(Y_i \cap H_1) \cap \operatorname{Int}(H_{1+}).$$

49

Note that $\operatorname{Int}(H_{1+}) \cap \pi_1^{-1}(H_i)$ is a union of connected components of $\operatorname{Int}(H_{1+}) \setminus D(Y_i \cap H_1)$. As ϕ_1 is close to $(\pi_1|_{H_{1+}})^{-1}$, we conclude that the Nash diffeomorphism ϕ_1 provides a bijection between the connected components of $\operatorname{Int}(H_1) \cap \operatorname{Int}(H_i)$ and the connected components of $\operatorname{Int}(H_{1+}) \cap \pi_1^{-1}(H_i)$. Consequently, $\phi_1(\operatorname{Int}(H_1) \cap \operatorname{Int}(H_i)) = \operatorname{Int}(H_{1+}) \cap \pi_1^{-1}(H_i)$, as claimed.

Summarizing, the objects we have constructed in this step are the following:

- The d-dimensional Nash manifold $M^1 := D(H_1)$ and the projection $\pi_1 : D(H_1) \to H_1, (x, t_1) \mapsto x.$
- The Nash manifolds with boundary $H_1^1 := H_{1+}$ and $H_i^1 := \pi_1^{-1}(H_i) = \{h_i^1 \ge 0\}$ for $i = 2, \ldots, \ell$. The boundary of H_1^1 is $Y_1^1 := Y_1 \times \{0\}$ and the boundary of H_i^1 is $Y_i^1 := D(Y_i \cap H_1)$ for $i = 2, \ldots, \ell$.
- The Nash normal-crossings divisor $Y^1 := \bigcup_{i=1}^{\ell} Y_i^1$ of M^1 .
- $\bullet\,$ The Nash equations

$$h_i^1: M^1 \to \mathbb{R}, \ (x, t_1) \mapsto \begin{cases} t_1 & \text{if } i = 1\\ h_i(x) & \text{if } i = 2, \dots, \ell, \end{cases}$$

which satisfy: $d_{(x,t_1)}h_i^1: T_{(x,t_1)}M^1 \to \mathbb{R}$ is surjective for each $(x,t_1) \in Y_i^1$ and $i = 1, \ldots, \ell$.

- The Nash manifold with corners $\Omega^1 := \bigcap_{i=1}^{\ell} H_i^1$, which is mapped by the projection π_1 onto Ω . In addition, the restriction $\pi_1|_{\Omega^1} : \Omega^1 \to \Omega$ is a semialgebraic homeomorphism and the restriction $\pi_1|_{\operatorname{Int}(\Omega^1)} : \operatorname{Int}(\Omega^1) \to \operatorname{Int}(\Omega)$ is a Nash diffeomorphism.
- The open semialgebraic neighborhoods $N^0 \subset M$ of H_1 and $W^1 \subset M^1$ of Q^1 and the Nash diffeomorphism $\phi_1 : N^0 \to W^1$ that maps Q onto Q^1 , Y_1 onto Y_1^1 and Y_i inside Y_i^1 for $i = 2, \ldots, \ell$.

STEP 2. Let $(D(H_2^1), \pi_2)$ be the Nash double of $H_2^1 = \{h_2^1 \ge 0\}$. We have

$$\begin{split} M^2 &:= D(H_2^1) = \{(x,t_1,t_2) \in (M \times \mathbb{R}) \times \mathbb{R} : \ t_1^2 - h_1(x) = 0, t_2^2 - h_2^1(x,t_1) = 0\} = \\ & \{(x,t_1,t_2) \in M \times \mathbb{R}^2 : \ t_1^2 - h_1(x) = 0, t_2^2 - h_2(x) = 0\} \end{split}$$

and $\pi_2 : D(H_2^1) \to H_2^1$, $(x, t_1, t_2) \mapsto (x, t_1)$. The Nash map π_2 is proper by Lemma 5.8(iv). Consider the (d-1)-dimensional Nash submanifolds $Y_1^2 := D(Y_1^1 \cap H_2^1)$, $Y_2^2 = Y_2^1 \times \{0\}$ and $Y_i^2 := D(Y_i^1 \cap H_2^1) = (Y_i^1 \times \mathbb{R}) \cap D(H_2^1)$ (see (5.1)) of M^2 for $i = 3, \ldots, \ell$. Define

$$h_i^2: M^2 \to \mathbb{R}, \ (x, t_1, t_2) \mapsto h_i^2(x, t_1, t_2) = \begin{cases} h_1^1(x, t_1) = t_1 & \text{if } i = 1, \\ t_2 & \text{if } i = 2, \\ h_i^1(x, t_1) = h_i(x) & \text{if } i = 3, \dots, \ell. \end{cases}$$

The tangent space to M^2 at a point $(x, t_1, t_2) \in M^2$ is:

$$T_{(x,t_1,t_2)}M^1 = \{(v,s_1,s_2) \in T_xM \times \mathbb{R}^2 : d_xh_1(v) - 2t_1s_1 = 0, d_xh_2(v) - 2t_2s_2 = 0\}.$$

The reader can check (analogously as we have done in STEP 1): $h_i^2: M^2 \to \mathbb{R}$ is a Nash equation of Y_i^2 in M^2 such that $d_{(x,t_1,t_2)}h_i^2: T_{(x,t_1,t_2)}M^2 \to \mathbb{R}$ is surjective for each $(x,t_1,t_2) \in Y_i^2$ and $i = 1, \ldots, \ell$.

By Theorem 5.10 there exist an open semialgebraic neighborhood $N^1 \subset M^1$ of H_2^1 and a Nash embedding $\phi_2 : N^1 \to M^2$ such that $\phi_2(H_2^1) = H_{2+}^1$ and $W^2 := \phi_2(N^1)$ is an open semialgebraic subset of M^2 that contains the Nash manifold with corners $\phi_2(Q^1)$ as a closed subset. In addition, $\phi_2|_{Y_2^1} = \operatorname{id}_{Y_2^1 \times \{0\}}, \ \phi_2(Y_i^1) \subset Y_i^2$ for $i = 1, 3, \ldots, \ell$ and $\phi_2|_{H_2^1} : H_2^1 \to H_{2+}^1$ is close to $(\pi_2|_{H_{2+}^1})^{-1}$. Consider the Nash subset $Y^2 := \bigcup_{i=1}^{\ell} Y_i^2$ of M^2 , whose irreducible components are Nash submanifolds of M^2 . We have $\phi_2(Y^1 \cap N^1) = Y^2 \cap W^2$. In addition, $\pi_{2-}|_{\operatorname{Int}(H_{2-}^1)} : \operatorname{Int}(H_{2-}^1) \to \operatorname{Int}(H_2^1)$ is a Nash diffeomorphism and $\pi_{2-}(Y^2 \cap \operatorname{Int}(H_{2-}^1)) = Y^1 \cap \operatorname{Int}(H_2^1)$. As Y^1 is a Nash normal-crossings divisor of M^1 and $M^2 = D(H_2^1) = W^2 \cup \operatorname{Int}(H_{2-}^1)$, we conclude that Y^2 is a Nash normal-crossings divisor of M^2 .

Denote $H_2^2 := H_{2+}^1$ and $H_i^2 := \pi_2^{-1}(H_i^1) = \{h_i^2 \ge 0\}$ for $i = 1, 3, ..., \ell$. By Lemma 5.6 the intersection

$$Q^{2} := \bigcap_{i=1}^{\ell} H_{i}^{2} = \{ (x, t_{1}, t_{2}) \in M \times \mathbb{R}^{2} : t_{1}^{2} - h_{1}(x) = 0, t_{2}^{2} - h_{2}(x) = 0, t_{1} \ge 0, t_{2} \ge 0, \\ h_{3}(x) \ge 0, \dots, h_{\ell}(x) \ge 0 \} \subset M^{2}$$

is a Nash manifold with corners and the Nash closure in M^2 of ∂Q^2 is Y^2 . Observe that $\pi_2(Q^2) = Q^1$, the restriction $\pi_2|_{Q^2} : Q^2 \to Q^1$ is a semialgebraic homeomorphism and the restriction $\pi_2|_{\mathrm{Int}(Q^2)} : \mathrm{Int}(Q^2) \to \mathrm{Int}(Q^1)$ is a Nash diffeomorphism. In particular, $(\pi_1 \circ \pi_2)(Q^2) = Q$. The reader can check (analogously as we have done in STEP 1): $\phi_2(Q^1) = Q^2$. Consequently, $(\phi_2 \circ \phi_1)(Q) = Q^2$.

RECURSIVE STEPS. We proceed recursively for $k = 1, \ldots, \ell$ and we construct:

• The *d*-dimensional Nash manifold M^k and the projection

$$\pi_k : M^k \to H_k^{k-1}, \ (x, t_1, \dots, t_k) \mapsto (x, t_1, \dots, t_{k-1}),$$

which is proper by Lemma 5.8(iv).

- The Nash manifolds with boundary H_i^k and the (d-1)-dimensional Nash submanifolds $Y_i^k = \partial H_i^k$ of M^k .
- The Nash normal-crossings divisor $Y^k := \bigcup_{i=1}^{\ell} Y_i^k$ of M^k .
- The Nash equations

$$h_i^k : M^k \to \mathbb{R}, \ (x, t_1, \dots, t_k) \mapsto \begin{cases} t_i & \text{if } i = 1, \dots, k \\ h_i(x) & \text{if } i = k+1, \dots, \ell \end{cases}$$

which satisfy: $d_{(x,t_1,\ldots,t_k)}h_i^k: T_{(x,t_1,\ldots,t_k)}M^k \to \mathbb{R}$ is surjective for each $(x,t_1,\ldots,t_k) \in Y_i^k$ and $i = 1, \ldots, \ell$.

• The Nash manifold with corners contained in ${\cal M}^k$

$$\mathfrak{Q}^k := \bigcap_{i=1}^{\ell} H_i^k = \{ (x, t_1, \dots, t_k) \in M \times \mathbb{R}^k : t_1^2 - h_1(x) = 0, \dots, t_k^2 - h_k(x) = 0, \\
t_1 \ge 0, \dots, t_k \ge 0, h_{k+1}(x) \ge 0, \dots, h_\ell(x) \ge 0 \},$$

which is mapped by the projection π_k onto \mathbb{Q}^{k-1} . Consequently, $(\pi_1 \circ \cdots \circ \pi_k)(\mathbb{Q}^k) = \mathbb{Q}$. In addition, the restriction $\pi_k|_{\mathbb{Q}^k} : \mathbb{Q}^k \to \mathbb{Q}^{k-1}$ is a semialgebraic homeomorphism and the restriction $\pi_k|_{\mathrm{Int}(\mathbb{Q}^k)} : \mathrm{Int}(\mathbb{Q}^k) \to \mathrm{Int}(\mathbb{Q}^{k-1})$ is a Nash diffeomorphism.

• The open semialgebraic neighborhoods $N^{k-1} \subset M^{k-1}$ of H_k^{k-1} and $W^k \subset M^k$ of \mathcal{Q}^k and the Nash diffeomorphism $\phi_k : N^{k-1} \to W^k$ that maps \mathcal{Q}^{k-1} onto \mathcal{Q}^k, Y_k^{k-1} onto Y_k^k and Y_i^{k-1} inside Y_i^k for $i \neq k$. Consequently, $(\phi_k \circ \cdots \circ \phi_1)(\mathcal{Q}) = \mathcal{Q}^k$.

At this point observe that

$$D(Q) = M^{\ell} = \{ (x,t) \in M \times \mathbb{R}^{\ell} : t_1^2 - h_1(x) = 0, \dots, t_{\ell}^2 - h_{\ell}(x) = 0 \}$$

is a d-dimensional Nash manifold and $\mathfrak{Q}^{\ell} = \{t_1 \geq 0, \ldots, t_{\ell}\} \subset D(\mathfrak{Q})$ is a Nash manifold with corners. The Nash closure of the boundary

$$\partial \mathcal{Q}^{\ell} = \bigcup_{i=1}^{\ell} \{ t_1 \ge 0, \cdots, t_{i-1} \ge 0, t_i = 0, t_{i+1} \ge 0, \cdots, t_{\ell} \ge 0 \}$$

is the Nash normal-crossings divisor $Y^{\ell} = \bigcup_{i=1}^{\ell} \{t_i = 0\}$ of $D(\mathcal{Q})$. After shrinking N^0 if necessary, the composition $\phi := \phi_{\ell} \circ \cdots \circ \phi_1 : N^0 \to W := \phi(N^0)$ is a Nash diffeomorphism such that W is an open semialgebraic subset of $D(\mathcal{Q})$ and $\phi(\mathcal{Q}) = \mathcal{Q}^{\ell}$. The composition $\pi := \pi_1 \circ \cdots \circ \pi_{\ell} : D(\mathcal{Q}) \to \mathcal{Q}$ is a proper surjective Nash map such that the restriction $\pi|_{\mathcal{Q}^{\ell}} : \mathcal{Q}^{\ell} \to \mathcal{Q}$ is a semialgebraic homeomorphism and the restriction $\pi|_{\mathrm{Int}(\mathcal{Q}^{\ell})} : \mathrm{Int}(\mathcal{Q}^{\ell}) \to \mathrm{Int}(\mathcal{Q})$ is a Nash diffeomorphism, as required.

6. Applications

In this section we prove several results proposed in the introduction as consequences of Theorems 1.4 and 1.7.

6.1. Weak desingularization of closed semialgebraic sets. We prove next Theorem 1.9.

Proof of Theorem 1.9. By Theorem 1.4 applied to S and [Sh, Rmk.II.1.5] to S and ε we may assume that:

- Q := S is a connected Nash manifolds with corners, which is closed in \mathbb{R}^n .
- The Zariski closure $X \subset \mathbb{R}^n$ of \mathcal{Q} is a *d*-dimensional non-singular real algebraic set.
- The Zariski closure Y of $\partial \Omega$ is a normal-crossings divisor of X.

If S is compact, we may assume in addition by Theorem 1.6 that X is compact and connected.

By [FGR, Thm.1.11, 1.12] there exists an open semialgebraic neighborhood $U \subset X$ of Ω such that:

- The Nash closure Y of ∂Q in M is a Nash normal-crossings divisor of M and $Q \cap Y = \partial Q$.
- For every $x \in \partial \Omega$ the analytic closure of the germ $\partial \Omega_x$ is Y_x .

Choose a strictly positive continuous semialgebraic function $\varepsilon_0 : X \to \mathbb{R}$ and an open semialgebraic neighborhood $U_1 \subset X$ of Ω such that $\operatorname{Cl}(U_1) \subset U$. In case Ω is compact, we may choose ε_0 constant. Define $M := \{x \in U_1 : \operatorname{dist}(x, \Omega) < \varepsilon_0(x)\}$, which is a Nash manifold such that $\Omega \subset M \subset \operatorname{Cl}(M) \subset U$.

By [Sh, Thm.VI.2.1] and its proof there exists a \mathcal{C}^1 semialgebraic function $g: U \to \mathbb{R}$ such that M is diffeomorphic to $g^{-1}((-1, +\infty))$ and $H := g^{-1}([0, +\infty)) \subset M$ is a \mathcal{C}^1 semialgebraic manifold with boundary that contains Ω in its interior. In case Ω is compact, also H is compact. We may assume in addition that the differential $d_xg: T_xM \to \mathbb{R}$ is surjective for each $x \in g^{-1}(0)$ (see Lemma 5.4). As $\operatorname{Cl}(M) \subset U$ is a closed subset of X, there exists a \mathcal{C}^1 semialgebraic extension of $g|_{\operatorname{Cl}(M)}$ to X such that $G^{-1}(0) = g^{-1}(0)$. Let $F: X \to \mathbb{R}$ be a Nash approximation of G in the \mathcal{C}^1 semialgebraic topology (in case X is compact, we may assume F is a polynomial). As G is strictly positive on the closed semialgebraic set Ω , we may assume F is strictly positive on Ω , so $F^{-1}(0) \subset M \setminus \Omega$. In addition, as we consider a \mathcal{C}^1 approximation, we may assume $d_xF: T_xX \to \mathbb{R}$ is surjective for each $x \in F^{-1}(0)$. Thus, $\mathcal{H}'_{\varepsilon_0} := F^{-1}([0, +\infty)) \subset M$ is a Nash manifold with (non-singular) boundary $Z_{\varepsilon_0} := F^{-1}(0)$, which is a Nash submanifold of X of dimension d-1. In case X is compact, also $\mathcal{H}'_{\varepsilon_0}$ is connected, it is contained in one of the connected components $\mathcal{H}_{\varepsilon_0}$ of $\mathcal{H}'_{\varepsilon_0}$. Observe that $\mathcal{H}_{\varepsilon_0}$ is a Nash manifold with boundary that contains Ω in its interior $\operatorname{Int}(\mathcal{H}_{\varepsilon_0})$, which is connected. Thus, $\operatorname{Int}(\mathcal{H}_{\varepsilon_0})$ is a connected component of $X \setminus Z_{\varepsilon_0}$.

In case Ω is non-compact and Z_{ε_0} is not a non-singular algebraic subset of X, we apply [Fe2, Lem.C.1] to assume up to a suitable Nash embedding of X in some affine space \mathbb{R}^p that $X \subset \mathbb{R}^p$ is a non-singular algebraic set and there exists an (algebraic) normal-crossings divisor Y_{ε_0} such that the Nash submanifold Z_{ε_0} is a union of connected components of Y_{ε_0} . By Hironaka's resolution of singularities we may assume that Y_{ε_0} is a non-singular algebraic subset of X.

By Theorem 1.7 there exists (after shrinking M at the beginning if needed) a proper Nash map $h: \operatorname{Cl}(M) \to \mathbb{Q}$ such that $h(M) = h(\mathbb{Q}) = \mathbb{Q}$, the restriction $h|_{\mathcal{H}_{\varepsilon_0} \setminus h^{-1}(\mathfrak{T}_{\varepsilon_0})} : \mathcal{H}_{\varepsilon_0} \setminus h^{-1}(\mathfrak{T}_{\varepsilon_0}) \to \mathbb{Q} \setminus \mathfrak{T}_{\varepsilon_0}$ is a Nash diffeomorphism, where $\mathfrak{T}_{\varepsilon_0} := \{x \in \mathbb{Q} : \operatorname{dist}(x, \partial \mathbb{Q}) < \varepsilon_0\}$. As $\mathbb{Q} \subset \mathcal{H}_{\varepsilon_0} \subset M$, we have $h(\mathcal{H}_{\varepsilon_0}) = \mathbb{Q}$. As $\mathcal{H}_{\varepsilon_0} \subset \operatorname{Cl}(M)$ is closed and $h: \operatorname{Cl}(M) \to \mathbb{Q}$ is proper, $f := h|_{\mathcal{H}_{\varepsilon_0}} : \mathcal{H}_{\varepsilon_0} \to \mathbb{Q}$ is also a proper map, as required.

6.2. An alternative proof to Theorem 1.10. In order to give an alternative proof of Theorem 1.10 based on Theorem 1.9, we need some preliminary results. We start with the following Lemma that extends [Fe2, Lem.2.8] to Nash manifolds with boundary.

Lemma 6.1. Let $H_1 \subset \mathbb{R}^m$ and $H_2 \subset \mathbb{R}^n$ be Nash manifolds with (non-singular) boundary. Let $f : H_1 \to H_2$ be a semialgebraic homeomorphism. Then every continuous semialgebraic map $g : H_1 \to H_2$ close to f in the \mathcal{C}^0 semialgebraic topology, such that $g(\partial H_1) \subset \partial H_2$, is surjective.

Proof. As $f: H_1 \to H_2$ is a semialgebraic homeomorphism, $f(\partial H_1) = \partial H_2$ (use invariance of domain). Consider the Nash doubles $(D(H_i), \pi_i)$ and $H_{i,\epsilon} = D(H_i) \cap \{\epsilon t \ge 0\}$ where $\epsilon = \pm$. Recall that $\pi_{i\epsilon}: H_{i,\epsilon} \to H_i$ is a semialgebraic homeomorphism. As $f(\partial H_1) = \partial H_2$, the map

$$F: D(H_1) \to D(H_2), \ (x,t) \mapsto \begin{cases} (\pi_{2+})^{-1} \circ f \circ \pi_{1+}(x,t) & \text{if } t \ge 0, \\ (\pi_{2-})^{-1} \circ f \circ \pi_{1-}(x,t) & \text{if } t \le 0, \end{cases}$$

is well-defined and semialgebraic. Observe that F is bijective and

$$F^{-1}: D(H_2) \to D(H_1), \ (y,s) \mapsto \begin{cases} (\pi_{1+})^{-1} \circ f^{-1} \circ \pi_{2+}(x,t) & \text{if } t \ge 0, \\ (\pi_{1-})^{-1} \circ f^{-1} \circ \pi_{2-}(x,t) & \text{if } t \le 0. \end{cases}$$

Let us check: F is continuous. Once this is done, the same proof shows that F^{-1} is continuous, so F is a semialgebraic homeomorphism.

As F is continuous on both $H_{1,+}$, $H_{1,-}$ and $\pi_1|_{H_1+\cap H_{1-}} = \pi_1|_{\partial H_1} = \mathrm{id}_{\partial H_1}$, we conclude by the pasting lemma that F is continuous on $D(H_1)$.

Let $g: H_1 \to H_2$ be a continuous semialgebraic map. If $g(\partial H_1) \subset \partial H_2$, the map

$$G: D(H_1) \to D(H_2), \ (x,t) \mapsto (\pi_{2\epsilon})^{-1} \circ g \circ \pi_{1\epsilon}(x,t)$$

is well-defined, continuous and semialgebraic for $\epsilon = \pm$. The proof is analogous to the one for F. Next, by [Fe2, Lem.2.8] there exists a strictly positive continuous semialgebraic function $\varepsilon : D(H_1) \to \mathbb{R}$ such that if $||G - F|| < \varepsilon$, then G is surjective.

For $\epsilon = \pm$, consider the strictly positive continuous semialgebraic functions

$$\varepsilon_{\epsilon} := \varepsilon \circ \pi_{1\epsilon}^{-1} : H_1 \to \mathbb{R}$$

and $\varepsilon^* = \min\{\varepsilon_+, \varepsilon_-\}$. Let $\delta: H_1 \to \mathbb{R}$ be a strictly positive continuous semialgebraic function such that if $f, g: H_1 \to H_2$ are continuous semialgebraic maps such that $||g - f|| < \delta$, then $||(\pi_{2\epsilon})^{-1} \circ f - (\pi_{2\epsilon})^{-1} \circ g|| < \varepsilon^*$ for $\epsilon = \pm$ (see [Sh, Rmk.II.1.5]). Thus, $||(\pi_{2\epsilon})^{-1} \circ f \circ \pi_{1\epsilon} - (\pi_{2\epsilon})^{-1} \circ g \circ \pi_{1\epsilon}|| < \varepsilon^* \circ \pi_{1\epsilon} \le \varepsilon$ for $\epsilon = \pm$. Consequently, $||F - G|| < \varepsilon$, so G is surjective. Taking into account the definition of G, we conclude that g is also surjective, as required. \Box

6.2.1. Preliminary reductions to prove Theorem 1.10. The proof of Theorem 1.10 requires some additional preparation. As the image of a compact semialgebraic set connected by analytic paths under a Nash map is a compact semialgebraic set connected by analytic paths [Fe2, Main Thm.1.4], we are reduced to prove: If $S \subset \mathbb{R}^n$ is a compact semialgebraic set connected by analytic paths of dimension $d \geq 2$, there exists a Nash map $f : \mathbb{R}^d \to \mathbb{R}^n$ such that $f(\overline{\mathbb{B}}_d) = S$.

By Theorem 1.9 we may assume that H := S is a connected compact Nash manifold with smooth boundary. Let M := D(H) be the Nash double of H (which is a connected compact Nash manifold) and consider the proper surjective Nash map $\pi : D(H) \to H$ introduced in Lemma 5.8. Thus, we may assume that M := S is a connected compact Nash manifold. Alternatively, we may use Theorem 1.4 and Theorem 5.13 to obtain a similar reduction.

Consider the simplex $\Delta_d := \{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_d \ge 0, \mathbf{x}_1 + \dots + \mathbf{x}_d \le 1\} \subset \mathbb{R}^d$. In [FU, Cor.2.8] we proved that the closed ball $\overline{\mathcal{B}}_d$ is a polynomial image of the simplicial prism $\Delta_{d-1} \times [0, 1]$. Thus, the proof of Theorem 1.10 is reduced to show the following: If $M \subset \mathbb{R}^n$ is a connected compact Nash manifold of dimension $d \ge 2$, there exists a Nash map $f : \mathbb{R}^d \to \mathbb{R}^n$ such that $f(\sigma \times [a, b]) = M$, where $\sigma \subset \mathbb{R}^{d-1}$ is a (d-1)-dimensional simplex and $[a, b] \subset \mathbb{R}$ a compact interval.

Let $\{U_i\}_{i=1}^r$ be a finite covering of M equipped with Nash diffeomorphisms $u_i: U_i \to \mathbb{R}^d$ such that $M = \bigcup_{i=1}^m u_i^{-1}(\Delta_d)$. Define

$$\Delta'_d := \{\mathbf{x}_1 \ge -2, \dots, \mathbf{x}_d \ge -2, \mathbf{x}_1 + \dots + \mathbf{x}_d \le d+1\}$$

and observe that $\Delta_d \subset \overline{\mathcal{B}}_d \subset \operatorname{Int}(\Delta'_d)$. Let $\psi : \mathbb{R}^d \to \mathbb{R}^d$ be an affine isomorphism such that $\psi(\Delta_d) = \Delta'_d.$

Lemma 6.2. Consider the (d-1)-simplex $\sigma := \{\mathbf{x}_1 \ge 0, \dots, \mathbf{x}_{d-1} \ge 0, \mathbf{x}_1 + \dots + \mathbf{x}_{d-1} \le 1\} \subset \mathbb{C}$ \mathbb{R}^{d-1} . Then the surjective Nash map $f: \sigma \times [0,1] \to \Delta'_d$, $(x,t) \mapsto \psi(((1-t)x,t))$ restricts to a Nash diffeomorphism $f|_{\sigma \times [0,1)} : \sigma \times [0,1) \to \Delta'_d \setminus \{\psi((0,\ldots,0,1))\}.$

Proof. As ψ is an affine isomorphism such that $\psi(\Delta_d) = \Delta'_d$, it is enough to prove that the surjective Nash map $\sigma \times [0,1] \to \Delta_d$, $(x,t) \mapsto ((1-t)x,t)$ restricts to a Nash diffeomorphism

$$\varphi: \sigma \times [0,1) \to \Delta_d \setminus \{(0,\ldots,0,1)\}.$$

This is straightforward, because the Nash map

$$\phi: \Delta_d \setminus \{(0, \dots, 0, 1)\} \to \sigma \times [0, 1), \ (x_1, \dots, x_d) \mapsto \left(\frac{x_1}{1 - x_d}, \dots, \frac{x_{d-1}}{1 - x_d}, x_d\right)$$

we rese of φ .

is the inverse of φ .

We are ready to prove Theorem 1.10.

6.2.2. Proof of Theorem 1.10. We keep the notations quoted above. In view of the contruction made in [CF, §2.3], to obtain a global Nash map $f: \mathbb{R}^d \to \mathbb{R}^n$ such that $f(\sigma \times [0, 2r-1]) = M$, it is enough to prove: The connected compact Nash manifold $M \subset \mathbb{R}^n$ is the image of $\sigma \times [0, 2r-1]$ under a Nash map $f: \sigma \times [0, 2r-1] \to \mathbb{R}^n$.

Let us construct first a surjective continuous semialgebraic function

$$h: \sigma \times [0, 2r - 1] \to M = \bigcup_{i=1}^{r} u_i^{-1}(\Delta_d) = \bigcup_{i=1}^{r} u_i^{-1}(\overline{\mathcal{B}}_d) = \bigcup_{i=1}^{r} u_i^{-1}(\Delta'_d)$$

that is Nash on $\sigma \times (\bigcup_{i=1}^{r} (2(i-1), 2(i-1)+1))$ and satisfies

$$u_i^{-1}(\overline{\mathcal{B}}_d) \subset h(\Delta_d \times (2(i-1), 2(i-1)+1)),$$

for each $i = 1, \ldots, r$.

Let $f: \sigma \times [0,1] \to \Delta'_d$ be the surjective Nash map introduced in Lemma 6.2 and define

$$h_i: \sigma \times [2(i-1), 2(i-1)+1] \to u_i^{-1}(\Delta'_d), \ (x,t) \mapsto u_i^{-1}(f(x,t-2(i-1))),$$

which is a surjective Nash map whose restriction to $\sigma \times \{2(i-1)+1\}$ is constant and its restriction to $\sigma \times [2(i-1), 2(i-1)+1)$ is a Nash diffeomorphism onto its image. Write $\{p_i\} :=$ $h_i(\sigma \times \{2(i-1)+1\})$. Let $\alpha_i : [2(i-1)+1, \frac{1}{2}+2(i-1)+1] \to M$ be a continuous semialgebraic map such that $\alpha_i(2(i-1)+1) = p_i$ and $\alpha_i(\frac{1}{2}+2(i-1)+1) = u_{i+1}^{-1}(0)$. Consider the continuous semialgebraic map

$$g_i: \sigma \times [2(i-1)+1, \frac{1}{2}+2(i-1)+1] \to M, \ (x,t) \mapsto \alpha_i(t).$$

Let b_1, \ldots, b_d be the vertices of Δ'_d different from $\psi(0, \ldots, 0, 1)$. Define $x_d := 1 - x_1 - \cdots - x_{d-1}$ and consider the continuous semialgebraic map

$$g'_{i}: \sigma \times \left[\frac{1}{2} + 2(i-1) + 1, 2(i-1) + 2\right] \to \mathbb{R}^{d},$$
$$(x,t) \mapsto u_{i+1}^{-1} \left(\sum_{i=1}^{d} x_{i} b_{i} 2\left(t - \frac{1}{2} - 2(i-1) - 1\right)\right),$$

which satisfies $g'_i(x, \frac{1}{2} + 2(i-1) + 1) = u_{i+1}^{-1}(0)$ and $g'_i(x, 2(i-1) + 2) = u_{i+1}^{-1}(\sum_{i=1}^d x_i b_i)$.

Define $h: \sigma \times [0, 2r-1] \to M$, as

$$(x,t) \mapsto \begin{cases} h_i(x,t) & \text{if } (x,t) \in \sigma \times [2(i-1), 2(i-1)+1], \\ g_i(x,t) & \text{if } (x,t) \in \sigma \times [2(i-1)+1, \frac{1}{2}+2(i-1)+1], \\ g'_i(x,t) & \text{if } (x,t) \in \sigma \times [\frac{1}{2}+2(i-1)+1, 2(i-1)+2], \\ h_r(x,t) & \text{if } (x,t) \in \sigma \times [2(r-1), 2r-1] \end{cases}$$

for $i = 1, \ldots, r - 1$. Observe that h is a continuous semialgebraic map such that for each $i=1,\ldots,r$:

- the restriction $h|_{\sigma \times (2(i-1),2(i-1)+1)}$ is a Nash diffeomorphism onto its image.
- $h(\sigma \times [2(i-1), 2(i-1)+1]) = u_i^{-1}(\Delta'_d).$ $u_i^{-1}(\overline{\mathcal{B}}_d) \subset u_i^{-1}(\operatorname{Int}(\Delta'_d)) \subset h(\sigma \times (2(i-1), 2(i-1)+1)).$

We conclude that the restriction of h to $\sigma \times (\bigcup_{i=1}^{r} (2(i-1), 2(i-1)+1))$ is Nash and surjective, because $M = \bigcup_{i=1}^{m} u_i^{-1}(\overline{\mathcal{B}}_d)$. In addition, $h_i^{-1}(u_i^{-1}(\overline{\mathcal{B}}_d)) \subset \operatorname{Int}(\sigma) \times (2(i-1), 2(i-1)+1)$, so the sets $h_i^{-1}(u_i^{-1}(\overline{\mathcal{B}}_d))$ (for $i = 1, \ldots, r$) are pairwise disjoint. Let $H : \mathbb{R}^d \to M$ be a continuous semialgebraic extension of h to \mathbb{R}^d . Define $X_1 := \bigcup_{i=1}^r (u_i \circ h_i)^{-1} (\partial \overline{\mathcal{B}}_d)$, which is a Nash subset of $\Omega := \operatorname{Int}(\sigma) \times \bigcup_{i=1}^{r} (2(i-1), 2(i-1)+1)$ and observe that H is Nash on the open semialgebraic set Ω . By [Sh, Thm.II.5.2] there exists a Nash map $F : \mathbb{R}^d \to M$ close to H such that $F|_{(u_i \circ h_i)^{-1}(\overline{\mathcal{B}}_d)} = H|_{(u_i \circ h_i)^{-1}(\partial \overline{\mathcal{B}}_d)}$ for $i = 1, \ldots, r$. As the restriction $H|_{(u_i \circ h_i)^{-1}(\overline{\mathcal{B}}_d)} = h_i|_{h_i^{-1}(u_i^{-1}(\overline{\mathcal{B}}_d))} : h_i^{-1}(u_i^{-1}(\overline{\mathcal{B}}_d)) \to u_i^{-1}(\overline{\mathcal{B}}_d)$ is by Lemma 6.2 a Nash diffeomorphism, the restriction $F|_{(u_i \circ h_i)^{-1}(\overline{\mathcal{B}}_d)}$ is by Lemma 6.1 surjective for $i = 1, \ldots, r$. Thus, $M = \bigcup_{i=1}^{m} u_i^{-1}(\overline{\mathcal{B}}_d) \subset F(\sigma \times [0, 2r-1]) \subset M$, so $F(\sigma \times [0, 2r-1]) = M$, as required.

6.3. Nash structure of orientable compact smooth manifolds. It is well-known that the compact orientable surface of genus q admits a Nash model. We show in the following example how to deduce straightforwardly this fact from Theorem 1.7.

Example 6.3. Let $\mathcal{P}_n \subset \mathbb{R}^2$ be a convex polygon with n edges. As \mathcal{P}_n is convex, there exist polynomials $h_1, \ldots, h_n \in \mathbb{R}[\mathbf{x}, \mathbf{y}]$ of degree 1 such that $\mathcal{P}_n = \{h_1 \ge 0, \ldots, h_n \ge 0\}$. Fix $2 + \frac{1}{2}(1 - 1)$ $(-1)^n \le s \le n$ and let $\mathcal{J} := \{J_k\}_{k=1}^s$ be a partition of the set $\{1, \ldots, n\}$ such that

$$\mathcal{P}_n \cap \{h_i = 0\} \cap \{h_j = 0\} = \emptyset \tag{6.1}$$

for each $i, j \in J_k$ with $i \neq j$. For each $k = 1, \ldots, s$ define the polynomial $h_{J_k} := \prod_{i \in J_k} h_i$. As the partition \mathcal{J} satisfies (6.1), it holds that \mathcal{P}_n is a connected component of the semialgebraic set $\{x \in \mathbb{R}^2 : h_{J_1}(x) \ge 0, \dots, h_{J_s}(x) \ge 0\}$. Thus,

$$D_s(\mathcal{P}_n) := \{ (x,t) \in \mathcal{P}_n \times \mathbb{R}^s : t_1^2 - h_{J_1}(x) = 0, \dots, t_s^2 - h_{J_s}(x) = 0 \} \subset \mathbb{R}^{s+2}$$

is a connected compact Nash surface, which is in addition a connected component of the (maybe singular) real algebraic set

$$X_{s,n} = \{(x,t) \in \mathbb{R}^2 \times \mathbb{R}^s : t_1^2 - h_{J_1}(x) = 0, \dots, t_s^2 - h_{J_s}(x) = 0\} \subset \mathbb{R}^{s+2}.$$

We claim: $D_s(\mathcal{P}_n) \subset \operatorname{Reg}(X_{s,n})$.

Pick $(x,t) \in D_s(\mathcal{P}_n)$ and consider the Jacobian matrix

$$J_{s,n}(x,t) := \begin{pmatrix} 2t_1 & 0 & \cdots & 0 & \frac{\partial h_{J_1}}{\partial \mathbf{x}_1}(x) & \frac{\partial h_{J_1}}{\partial \mathbf{x}_2}(x) \\ 0 & 2t_2 & \cdots & 0 & \frac{\partial h_{J_2}}{\partial \mathbf{x}_1}(x) & \frac{\partial h_{J_2}}{\partial \mathbf{x}_2}(x) \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \cdots & 2t_s & \frac{\partial h_{J_s}}{\partial \mathbf{x}_1}(x) & \frac{\partial h_{J_s}}{\partial \mathbf{x}_2}(x) \end{pmatrix},$$

which has rank $\leq s$. As $(x,t) \in D_s(\mathcal{P}_n)$, then $x \in \mathcal{P}_n$ and there exists at most two indices k, ℓ such that $x \in \{h_{J_k} = 0, h_{J_\ell} = 0\}$. If such is the case, the vectors $\left(\frac{\partial h_{J_k}}{\partial \mathbf{x}_1}(x), \frac{\partial h_{J_k}}{\partial \mathbf{x}_2}(x)\right)$ and $\left(\frac{\partial h_{J_{\ell}}}{\partial \mathbf{x}_1}(x), \frac{\partial h_{J_{\ell}}}{\partial \mathbf{x}_2}(x)\right)$ are linearly independent. If there exists only one index k such that $x \in \{h_{J_k} = 0\}$, the vector $(\frac{\partial h_{J_k}}{\partial x_1}(x), \frac{\partial h_{J_k}}{\partial x_2}(x))$ is non-zero because the partition \mathcal{J} satisfies (6.1). As $t_k^2 - h_{J_k}(x) = 0$ for $k = 1, \ldots, s$, the matrix $J_{s,n}(x,t)$ has rank s, so $(x,t) \in \operatorname{Reg}(X_{s,n})$, as claimed.

As $D_s(\mathcal{P}_n)$ is obtained recursively by doubling orientable Nash manifolds with boundary, $D_s(\mathcal{P}_n)$ is an orientable Nash surface. By the (smooth) classification of surfaces (see for instance [H, Ch.9]) $D_s(\mathcal{P}_n)$ is diffeomorphic to a connected sum of g tori and the genus g completely characterize its diffeomorphism class.

For each $\varepsilon := (\varepsilon_1, \ldots, \varepsilon_s) \in \{-1, 1\}^s$ the set $D_s(\mathcal{P}_n) \cap \{\varepsilon_1 t_1 \ge 0, \ldots, \varepsilon_s t_s \ge 0\}$ is Nash diffeomorphic to \mathcal{P}_n . Thus, $D_s(\mathcal{P}_n)$ is obtained (topologically) by glueing 2^s copies of \mathcal{P}_n , one for each choice of $\varepsilon \in \{-1, 1\}^s$. The polygon \mathcal{P}_n has a (natural) structure of CW complex with n vertices, n edges and 1 face. This CW complex structure induces a CW complex structure on $D_s(\mathcal{P}_n)$ with $2^{s-2}n$ vertices (because each vertex belongs exactly to 4 polygons of the CW complex), $2^{s-1}n$ edges (because each edge belongs exactly to 2 polygons of the CW complex) and 2^s faces. The Euler characteristic of $D_s(\mathcal{P}_n)$ is $2^{s-2}n - 2^{s-1}n + 2^s = 2^{s-2}(4-n)$ and its genus is $g = \frac{2-2^{s-1}(4-n)}{2} = 2^{s-3}(n-4)+1$, that is, $D_s(\mathcal{P}_n)$ is diffeomorphic to the connected sum of $g = 2^{s-3}(n-4) + 1$ tori, whenever the number $2^{s-3}(n-4) + 1$ is a non-negative integer. In particular, the compact orientable surface of genus $g \ge 1$ is diffeomorphic to the Nash surfaces $D_2(\mathcal{P}_{2g+2}) \subset \mathbb{R}^4$ and $D_3(\mathcal{P}_{g+3}) \subset \mathbb{R}^5$. Note that the genus g of the surface $D_s(\mathcal{P}_n)$ depends both on n and s (see Table 1).


	s = 2	s = 3	s = 4	s = 5	s = 6	s = 7
n = 3	—	0	—	_	—	—
n = 4	1	1	1	—	—	—
n = 5	—	2	3	5	—	—
n = 6	2	3	5	9	17	_
n = 7	_	4	7	13	25	49

TABLE 1. The genus of the Nash surface $D_s(\mathcal{P}_n)$ for $n \leq 7$.

By Lemma 2.8 there exist irreducible non-singular real algebraic sets X with at most two connected components that are both Nash diffeomorphic to either $D_2(\mathcal{P}_{2g+2}) \subset \mathbb{R}^4$ or $D_3(\mathcal{P}_{g+3}) \subset \mathbb{R}^5$. Consequently, our procedure provides algebraic sets with at most two connected components (mutually diffeomorphic) such that their connected components are models for the orientable compact smooth surfaces.

6.4. Nash approximation. We make use of Theorem 5.13 to prove Theorem 1.12. Let $\mathcal{Q} \subset \mathbb{R}^n$ be a Nash manifold with corners, let $M \subset \mathbb{R}^n$ be a Nash envelope of \mathcal{Q} and $Y \subset M$ a Nash normal-crossings divisor such that $\mathcal{Q} \cap Y = \partial \mathcal{Q}$. Let $h_1, \ldots, h_\ell \in \mathcal{N}(M)$ be Nash equations for the irreducible components Y_1, \ldots, Y_ℓ of Y, that is, $Y_i := \{h_i = 0\}$ and $d_x h_i : T_x M \to \mathbb{R}$ is surjective for each $x \in Y_i$. We have $\mathcal{Q} = \{h_1 \geq 0, \ldots, h_\ell \geq 0\}$ and consider the Nash double of \mathcal{Q} , that is, the Nash manifold $D(\mathcal{Q}) := \{(x,t) \in M \times \mathbb{R}^\ell : t_1^2 - h_1(x) = 0, \ldots, t_\ell^2 - h_\ell(x) = 0\}$ together with the projection $\pi : D(\mathcal{Q}) \to M, (x, t) \mapsto x$. It holds $\pi(D(\mathcal{Q})) = \mathcal{Q}$ and if we define $\pi_+^{-1} : \mathcal{Q} \to D(\mathcal{Q}), \ x \mapsto (x, \sqrt{h_1(x)}, \ldots, \sqrt{h_\ell(x)})$, then π_+^{-1} is a semialgebraic homeomorphism onto its image $\mathcal{Q}^+ := D(\mathcal{Q}) \cap \{t_1 \geq 0, \ldots, t_\ell \geq 0\}$ such that $\pi_+^{-1} \circ \pi|_{\mathcal{Q}^+} = \mathrm{id}_{\mathcal{Q}^+}$ and $\pi \circ \pi_+^{-1} = \mathrm{id}_{\mathcal{Q}}$.

Proof of Theorem 1.12. By [DK] there exist an open semialgebraic neighborhood $U \subset \mathbb{R}^m$ of S (in which S is closed) and a semialgebraic retraction $\nu : U \to S$. Consider the continuous semialgebraic function $H := \pi_+^{-1} \circ h \circ \nu : U \to D(\Omega)$. Let $\varepsilon_0 : S \to \mathbb{R}$ be a strictly positive continuous semialgebraic function and let $\varepsilon : U \to \mathbb{R}$ be a strictly positive continuous semialgebraic function the following commutative diagram:



As the map $\pi_* : S^0(U, D(\Omega)) \to S^0(U, \Omega), H \mapsto \pi \circ H$ is continuous by [Sh, Rmk.II.1.5], there exists $\delta : U \to \mathbb{R}$ such that if $G \in S^0(U, D(\Omega))$ and $||H - G|| < \delta$, then $||\pi \circ H - \pi \circ G|| < \varepsilon$. Let $G : U \to D(\Omega)$ be a Nash map such that $||H - G|| < \delta$, so $||\pi \circ H - \pi \circ G|| < \varepsilon$. As $H|_{\mathcal{S}} = \pi_+^{-1} \circ h$, we deduce

 $\|h - \pi \circ G|_{\mathfrak{S}}\| = \|\operatorname{id}_{\mathfrak{Q}} \circ h - \pi \circ G|_{\mathfrak{S}}\| = \|\pi \circ \pi_{+}^{-1} \circ h - \pi \circ G|_{\mathfrak{S}}\| = \|\pi \circ H|_{\mathfrak{S}} - \pi \circ G|_{\mathfrak{S}}\| < \varepsilon|_{\mathfrak{S}} = \varepsilon_{0},$ as required.

References

- [A1] S.S. Abhyankar: Resolution of singularities of embedded algebraic surfaces. Pure and Applied Mathematics, 24 Academic Press, New York-London (1966).
- [A2] S.S. Abhyankar: Weighted expansions for canonical desingularization. Lect. Notes Math.910, Berlin Heidelberg New York: Springer (1982).
- [A3] S.S. Abhyankar: Good points of a hypersurface. Adv. in Math. 68 (1988), no. 2, 87–256.
- [AK] S. Akbulut, H. King: A relative Nash theorem. Trans. Amer. Math. Soc. 267 (1981), no. 2, 465–481.
- [BFR] E. Baro, J.F. Fernando, J.M. Ruiz: Approximation on Nash sets with monomial singularities. *Adv. Math.* **262** (2014), 59–114.
- [BM1] E. Bierstone, P.D. Milman: Uniformization of analytic spaces. J. Amer. Math. Soc. 2 (1989), no. 4, 801–836.
- [BM3] E. Bierstone, P.D. Milman: Canonical desingularization in characteristic zero by blow up the maximum strata of a local invariant. *Invent. Math.* **128** (1997), no. 2, 207–302.
- [BM2] E. Bierstone, P.D. Milman: Functoriality in resolution of singularities. *Publ. Res. Inst. Math. Sci.* 44 (2008), no. 2, 609–639.
- [BP] E. Bierstone, A. Parusiński: Global smoothing of a subanalytic set. Duke Math. J. 167 (2018), no.16, 3115–3128.
- [BCR] J. Bochnak, M. Coste, M.-F. Roy: Real algebraic geometry. *Ergeb. Math.* **36**, Springer-Verlag, Berlin (1998).
- [CF] A. Carbone, J.F. Fernando: Surjective Nash maps between semialgebraic sets. Preprint (2023). arXiv:2306.00401
- [C] H. Clemens: Curves in generic hypersurfaces. Ann. Sci. Ecole Norm. Sup. 19 (1986), 629–636.
- [DK] H. Delfs, M. Knebusch: Separation, retractions and homotopy extension in semialgebraic spaces. Pacific J. Math. 114 (1984), no. 1, 47–71.
- [Ef] G.A. Efroymson: The Extension Theorem for Nash Functions. Real Algebraic Geometry and Quadratic Forms (Rennes, 1981). Lecture Notes in Mathematics, vol. 959, Springer, Berlin (1982), 343–357.
- [E] D. Eisenbud: Open Problems in Computational Algebraic Geometry and commutative Algebra, in Computational Algebraic Geometry and Commutative Algebra, Cortona (1991), (ed. D. Eisenbud and L. Robbiano) Cambridge University Press, Cambridge (1993), 49–71.
- [Fe1] J.F. Fernando: On the substitution theorem for rings of semialgebraic functions. J. Inst. Math. Jussieu 14 (2015), no. 4, 857–894.
- [Fe2] J.F. Fernando: On Nash images of Euclidean spaces. Adv. Math. 331 (2018), 627–719.
- [Fe3] J.F. Fernando: On a Nash curve selection lemma through finitely many points. *Preprint RAAG* (2023).
- [FFQU] J.F. Fernando, G. Fichou, R. Quarez, C. Ueno: On regulous and regular images of Euclidean spaces, Q. J. Math. 69 (2018), no. 4, 1327–1351.
- [FG] J.F. Fernando, J.M. Gamboa: On open and closed morphisms between semialgebraic sets. Proc. Amer. Math. Soc. 140 (2012), no. 4, 1207–1219.
- [FGR] J.F. Fernando, J.M. Gamboa, J.M. Ruiz: Finiteness problems on Nash manifolds and Nash sets. J. Eur. Math. Soc. (JEMS) 16 (2014), no. 3, 537–570.
- [FGU] J.F. Fernando, J.M. Gamboa, C. Ueno: Polynomial, regular and Nash images of Euclidean spaces. Ordered algebraic structures and related topics, *Contemp. Math.*, 697, Amer. Math. Soc., Providence, RI (2017), 145–167.
- [FGh1] J.F. Fernando, R. Ghiloni: Differentiable approximation of continuous semialgebraic maps. Selecta Math. (N.S.) 25 (2019), no. 3, Paper No. 46, 30 pp.
- [FGh2] J.F. Fernando, R. Ghiloni: Smooth approximations in PL geometry. Amer. J. Math. 144 (2022), no. 4, 967–1007.
- [FU] J.F. Fernando, C. Ueno: On polynomial images of a closed ball. J. Math. Soc. Japan 75 (2023), no. 2, 679–733.
- $[G] \qquad J.M. \ Gamboa: \ Algebraic images of the real plane. \ Reelle \ algebraische \ Geometrie, \ June, \ 10^{\rm th} 16^{\rm th} \ (1990), \ Oberwolfach.$
- [GS] R. Ghiloni, E. Savi: The topology of real algebraic sets with isolated singularities is determined by the field of rational numbers. *Preprint* (2023). arXiv:2302.04142
- [Ha] J. Harris: Algebraic geometry. A first course. Corrected reprint of the 1992 original. Graduate Texts in Mathematics, 133. Springer-Verlag, New York, (1995).
- [H] M.W. Hirsch: Differential topology. *Graduate Texts in Mathematics*, **33**. Springer–Verlag, New York–Heidelberg–Berlin (1976).
- [Hi1] H. Hironaka: Resolution of singularities of an algebraic variety over a field of characteristic zero. I, II. Ann. of Math. (2) 79 (1964), 109–203; ibid. (2) 79 (1964) 205–326.
- [Hi2] H. Hironaka: Stratification and flatness. Real and complex singularities (*Proc. Ninth Nordic Summer School/NAVF Sympos. Math.*, Oslo, 1976). Sijthoff and Noordhoff, Alphen aan den Rijn (1977), 199–265.
- [HPV] J. Hubbard, P. Papadopol, V. Veselov: A compactification of Hénon mappings in C² as dynamical systems. Acta Math. 184 (2000), no. 2, 203−270.
- [Ko] J. Kollár: Lectures on resolution of singularities. Annals of Mathematics Studies, 166. Princeton University Press, Princeton, NJ, (2007).

- [L] J. Lipman: Introduction to resolution of singularities. Algebraic geometry (Proc. Sympos. Pure Math., 29, Humboldt State Univ., Arcata, Calif., 1974), pp. 187–230. Amer. Math. Soc., Providence, R.I., (1975).
- [M] T. Mostowski: Some properties of the ring of Nash functions. Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 3 (1976), no. 2, 245–266.
- [ORR] E. Outerelo, J. A. Rojo, and J. M. Ruiz, Topología diferencial: un curso de iniciación, Sanz y Torres, S.L., Madrid, (2020).
- [Sh] M. Shiota: Nash manifolds. Lecture Notes in Mathematics, **1269**. Springer-Verlag, Berlin (1987).
- [S] F.P. Di Stefano: Oriented blow-up. Ann. Univ. Ferrara Sez. VII (N.S.) 24 (1979), 25–35.
- [St] J. Stasica: Smooth points of a semialgebraic set. Ann. Pol. Math. 82 (2003), no.2, 149–153.
- [Vi] O. Villamayor: Constructiveness of Hironaka's resolution. Ann. Sci. École Norm. Sup. (4) 22 (1989), no. 1, 1–32.
- [V] C. Voisin: On a conjecture of Clemens on rational curves on hypersurfaces. J. Differential Geom. 44 (1996), 200–214.
- [Wh] H. Whitney: Differentiable manifolds. Ann. of Math. (2) 37 (1936), no. 3, 645–680.
- [Z1] O. Zariski: Algebraic varieties over ground fields of characteristic zero. Amer. J. Math. 62 (1940), 187– 221.
- [Z2] O. Zariski: A simplified proof for the resolution of singularities of an algebraic surface. Ann. of Math.
 (2) 43 (1942), 583–593.
- [Z3] O. Zariski: Reduction of the singularities of algebraic three dimensional varieties. Ann. of Math. (2) 45 (1944), 472–542.

DIPARTIMENTO DI MATEMATICA, VIA SOMMARIVE, 14, UNIVERSITÀ DI TRENTO, 38123 POVO (ITALY)

 $Email \ address: \ \texttt{antonio.carboneQunitn.it}$

DEPARTAMENTO DE ÁLGEBRA, GEOMETRÍA Y TOPOLOGÍA, FACULTAD DE CIENCIAS MATEMÁTICAS, UNIVER-SIDAD COMPLUTENSE DE MADRID, PLAZA DE CIENCIAS 3, 28040 MADRID (SPAIN)

Email address: josefer@mat.ucm.es