On the critical Caffarelli-Kohn-Nirenberg type inequalities involving super-logarithms

Hiroshi Ando, Toshio Horiuchi, Eiichi Nakai

Abstract

We establish the Caffarelli-Kohn-Nirenberg type inequalities involving super-logarithms (infinitely iterated logarithms). As a result the critical Caffarelli-Kohn-Nirenberg type inequalities will be improved, and in certain cases the best constants will be discovered. *

1 Introduction

The main purpose of the present paper is to improve the Caffarelli-Kohn-Nirenberg type inequalities in the critical case studied in [14], which are abbreviated as the critical CKN-type inequalities.

We briefly review the classical CKN-type inequalities. The CKN-type inequalities were introduced by [6] as multiplicative interpolation inequalities, but here we refer to the simple weighted Sobolev inequalities. There is a great deal of research in that case alone. In [14] we also investigated the CKN-type inequalities systematically. Assume that $n \ge 1$, $1 and <math>0 \le 1/p - 1/q \le 1/n$. (1) In the non-critical case that $\gamma \ne 0$, the CKN-type inequalities have the following form:

$$\int_{\mathbf{R}^{n}} |\nabla u(x)|^{p} |x|^{p(1+\gamma)-n} dx \ge S^{p,q,\gamma} \left(\int_{\mathbf{R}^{n}} |u(x)|^{q} |x|^{\gamma q-n} dx \right)^{p/q}, \ u \in C_{c}^{\infty}(\mathbf{R}^{n} \setminus \{0\}). \tag{1.1}$$

Here $S^{p,q;\gamma} = S^{p,q;\gamma}(\mathbf{R}^n)$ is the best constant and given by the following variational problem:

$$S^{p,q;\gamma} = \inf\{E^{p,q;\gamma}[u] : u \in C_{\mathbf{c}}^{\infty}(\mathbf{R}^n \setminus \{0\}) \setminus \{0\}\},\tag{1.2}$$

where

$$E^{p,q;\gamma}[u] = \frac{\int_{\mathbf{R}^n} |\nabla u(x)|^p |x|^{p(1+\gamma)-n} dx}{\left(\int_{\mathbf{R}^n} |u(x)|^q |x|^{\gamma q-n} dx\right)^{p/q}} \quad \text{for } u \in C_c^{\infty}(\mathbf{R}^n \setminus \{0\}) \setminus \{0\}.$$
 (1.3)

We also define the radial best constant $S^{p,q;\gamma}_{\rm rad}=S^{p,q;\gamma}_{\rm rad}({\bf R}^n)$ as follows.

$$S_{\text{rad}}^{p,q;\gamma} = \inf\{E^{p,q;\gamma}[u] : u \in C_{\text{c}}^{\infty}(\mathbf{R}^n \setminus \{0\}) \setminus \{0\}, u \text{ is radial } \}.$$
 (1.4)

(2) In the critical case that $\gamma = 0$, the CKN-type inequalities have the following form: For $\eta > 0$, let B_{η} be the ball $\{x \in \mathbb{R}^n : |x| < \eta\}$.

$$\int_{B_{\eta}} |\nabla u(x)|^{p} |x|^{p-n} dx \ge C^{p,q;R} \left(\int_{B_{\eta}} \frac{|u(x)|^{q}}{|x|^{n} \left(\log(R\eta/|x|) \right)^{1+q/p'}} dx \right)^{p/q}, \quad u \in C_{c}^{\infty}(B_{\eta} \setminus \{0\}), \tag{1.5}$$

Key words and phrases. the critical Caffarelli-Kohn-Nirenberg type inequalities, super-logarithm, infinitely iterated logarithms This research was partially supported by Grant-in-Aid for Scientific Research (No. 20K03670, No. 21K03304).

^{*2010}Mathematics Subject Classification. Primary 35J70; Secondary 35J60.

where p' = p/(p-1), R is a positive number satisfying R > 1 and the best constant $C^{p,q;R} = C^{p,q;R}(B_{\eta})$ is given by the variational problem:

$$C^{p,q;R} = \inf\{F^{p,q;R}[u] : u \in C_c^{\infty}(B_{\eta} \setminus \{0\}) \setminus \{0\}\},$$
(1.6)

where

$$F^{p,q;R}[u] = \frac{\int_{B_{\eta}} |\nabla u(x)|^p |x|^{p-n} dx}{\left(\int_{B_{\eta}} |u(x)|^q |x|^{-n} (\log(R\eta/|x|))^{-1-q/p'} dx\right)^{p/q}} \text{ for } u \in C_c^{\infty}(B_{\eta} \setminus \{0\}) \setminus \{0\}.$$
 (1.7)

We also define the radial best constant

$$C_{\text{rad}}^{p,q;R} = \inf\{F^{p,q;R}[u] : u \in C_{\text{c}}^{\infty}(B_{\eta} \setminus \{0\}) \setminus \{0\}, u \text{ is radial }\}.$$

$$\tag{1.8}$$

Here we remark that the best constants $C^{p,q;R}$ and $C^{p,q;R}_{\mathrm{rad}}$ are independent of $\eta > 0$. To see this it suffices to employ a change of variables given by $x = \eta y$. Roughly speaking if $p \ge n$, the imbedding inequalities (1.5) follow from a generalized rearrangement argument. On the other hand if 1 , then these are established by using the so-called nonlinear potential theory.

Recently M. Sano and F. Takahashi established in [16] the CKN-type inequality with logarithmic weights in two dimensions using the nonlinear potential theory. In [10, 11, 13] we also revisited the CKN inequality and extended them to the case with non-doubling weights. Interestingly, the framework in [11] not only unifies non-critical and critical cases, but also makes it possible to study the critical case in greater depth. For more information on the CKN-type inequalities and relating inequalities see [6, 17, 8, 9]. On the other hand, the super-logarithms have been introduced and developed by [1, 7, 2, 3]. Intuitively, the super-logarithms are defined as concave functions that grow more slowly than all polylogarithms and they are realized as arbitrarily and infinitely iterated logarithm functions. They have been applied to the refinement of the Hardy inequalities by finding infinitely many missing terms in [2, 3] and the analysis of weighted L^p -boundedness of convolution type integral operators in [7].

In this paper we will improve the critical CKN-type inequalities represented by (1.5) by using not only poly-logarithms but also super-logarithms as weights instead of $|x|^{p-n}$ on the left side, and in certain cases the best constants of them will be determined. To do this we use recent results on the CKN-type inequalities from [11] instead of the nonlinear potential theory, and a generalized rearrangement of function from [14].

This paper is organized in the following way: §2 provide the basic and essential knowledge of this paper. In the first §2.1, super-logarithms are reviewed from [1]. Next, §2.2 reviews CKN-type inequalities in the new framework from [11]. Finally, in §2.3 we recall a generalized rearrangement of function from [14]. After these preparations, as preliminary results, §3 gives critical CKN-type inequalities involving poly-logarithms as weights. Then, in §4, we state the critical CKN-type inequalities involving super-logarithms and also give proofs of them. Moreover, §5 gives further results.

2 Preliminaries

2.1 A review on super-logarithms (infinitely iterated logarithms)

We recall poly-logarithm and poly-exponential functions as follows.

$$\begin{cases} \log^{0} r = r, \log^{n} r = \log(\log^{n-1} r), & n \in \mathbb{N}, \\ \exp^{0}(r) = r, \exp^{n}(r) = \exp(\exp^{n-1}(r)), & n \in \mathbb{N}. \end{cases}$$
 (2.1)

First we define two sets of functions, following [1].

Definition 2.1. Let \mathscr{L} be the set of all continuous, increasing and bijective functions f from $(0,\infty)$ to $(-\infty,\infty)$ satisfying

$$\lim_{r\to +0} f(r) = -\infty, \quad f(1) = 0, \quad \lim_{r\to \infty} f(r) = \infty.$$

For example, the logarithmic function $\log r$ is in \mathcal{L} .

Definition 2.2. For a > 1, let \mathscr{F}_a be the set of all continuous, increasing and bijective functions from $[a, \infty)$ to itself.

If $f \in \mathscr{F}_a$, then f(a) = a and $\lim_{u \to \infty} f(u) = \infty$. For a function $f \in \mathscr{F}_a$, let $f^0(u) = u$ and $f^k(u) = f(f^{k-1}(u))$, $k \in \mathbb{N}$. Then f^k is also in \mathscr{F}_a .

We define a function $F \in \mathscr{F}_a$ as

$$F(u) = F_a(u) = a - \log a + \log u \quad (u \ge a).$$
 (2.2)

Then the relation

$$(F^{k}(u))' = \frac{1}{F^{k-1}(u)\cdots F^{1}(u)F^{0}(u)}$$
(2.3)

holds. That is,

$$F^{k}(u) = a + \int_{a}^{u} \frac{dt}{F^{k-1}(t)\cdots F^{1}(t)F^{0}(t)}.$$
 (2.4)

Definition 2.3. For a > 1, let

$$\tilde{F}(u) = \tilde{F}_a(u) = a \prod_{k=0}^{\infty} \frac{F^k(u)}{a} = u \prod_{k=1}^{\infty} \frac{F^k(u)}{a} \quad (u \ge a).$$
 (2.5)

Definition 2.4 (Super-Logarithm). For a > 1, let

$$L(r) = L_a(r) = \int_a^{ar} \frac{1}{\tilde{F}(t)} dt \quad (r \ge 1), \tag{2.6}$$

and let

$$L(r) = -L(1/r) = -\int_{a}^{a/r} \frac{1}{\tilde{F}(t)} dt \quad (0 < r < 1), \tag{2.7}$$

where \tilde{F} is as in (2.5).

Then we have the following.

Theorem 2.1 ([1], Theorem 2.1). *Let* a > 1.

(i) The function \tilde{F} is in \mathscr{F}_a , infinitely differentiable and has the following expression:

$$\tilde{F}(u) = \exp(V(u)), \quad V(u) = \log a + \int_{a}^{u} \left(\sum_{k=0}^{\infty} \frac{1}{\prod_{i=0}^{k} F^{j}(t)}\right) dt.$$
 (2.8)

Further, $\left(\frac{d}{du}\right)^k \frac{\tilde{F}'}{\tilde{F}}$ is bounded for each $k \in \{0\} \cup \mathbf{N}$.

- (ii) The function L is in \mathcal{L} , differentiable on $(0,\infty)$, infinitely differentiable except at 1, and, concave on $[1,\infty)$, Moreover, if $a \geq 2$, then L is concave on $(0,\infty)$.
- (iii) For each $n \in \mathbb{N}$,

$$\lim_{r \to +0} \frac{L(r)}{\log^n(1/r)} = \lim_{r \to \infty} \frac{L(r)}{\log^n r} = 0.$$
 (2.9)

(iv) For
$$r \ge \exp(a)$$
,
$$L(r) \le L(\exp(r)) \le (1+a)L(r).$$

Remark 2.1. We note that the following relations

$$\lim_{r \to \infty} \frac{F^{k}(ar)}{\log^{k} r} = 1, \quad \lim_{r \to \infty} \frac{F^{k}(ar)}{\prod_{j=1}^{\infty} \frac{F^{j}(ar)}{a}} = 0, \quad \lim_{r \to \infty} \frac{a + L(r)}{F^{k}(ar)} = 0, \quad \lim_{r \to \infty} \frac{F^{k}(a + L(r))}{\log^{k}(a + L(r))} = 1$$
 (2.10)

and

$$\begin{cases} L'(r) = \frac{a}{\tilde{F}(ar)} = \frac{1}{r \prod_{j=1}^{\infty} \frac{F^{j}(ar)}{a}} & (r > 1), \\ (\tilde{F}(u))' = \tilde{F}(u) \left(\sum_{k=0}^{\infty} \frac{1}{\prod_{j=0}^{k} F^{j}(u)} \right) & (u > a). \end{cases}$$

$$(2.11)$$

hold for any $k \in \mathbb{N}$ and a > 1.

2.2 A review on the general CKN-type inequalities with non-doubling weights

 $\mathbf{R}_+ = (0, \infty)$. By $C^{0,1}(\mathbf{R}_+)$ we denote the space of all Lipschitz continuous functions on \mathbf{R}_+ . First we define a class of weight functions $W(\mathbf{R}_+)$ which is a slight modification of the one adopted in [11] to suit our purpose.

Definition 2.5. Let us set

$$W(\mathbf{R}_{+}) = \{ w \in C^{0,1}(\mathbf{R}_{+}) : w > 0, \lim_{t \to +0} w(t) = a \text{ for some } a \in [0, \infty] \}.$$
 (2.12)

In the next we define two subclasses of this large class.

Definition 2.6. Let us set

$$P(\mathbf{R}_{+}) = \{ w \in W(\mathbf{R}_{+}) : w^{-1} \notin L^{1}((0, \eta)) \text{ for some } \eta > 0 \}.$$
 (2.13)

$$Q(\mathbf{R}_{+}) = \{ w \in W(\mathbf{R}_{+}) : w^{-1} \in L^{1}((0, \eta)) \text{ for any } \eta > 0 \}.$$
 (2.14)

Example 2.1. $t(\log(1+1/t))^{\alpha} \in P(\mathbf{R}_{+})$ if $\alpha \le 1$ and $t(\log(1+1/t))^{\alpha} \in Q(\mathbf{R}_{+})$ if $\alpha > 1$.

Remark 2.2. From Definitions 2.5 and 2.6 it follows that $W(\mathbf{R}_+) = P(\mathbf{R}_+) \cup Q(\mathbf{R}_+)$ and $P(\mathbf{R}_+) \cap Q(\mathbf{R}_+) = \emptyset$.

We define functions $f_n(t)$ and $G_n(t)$ on $(0, \eta]$ in order to introduce variants of the Hardy potentials.

Definition 2.7. For $\mu > 0$, $\eta > 0$ and $t \in (0, \eta]$ we set the followings:

(i) When $w \in P(\mathbf{R}_+)$,

$$f_{\eta}(t; w, \mu) = \mu + \int_{t}^{\eta} \frac{1}{w(s)} ds.$$
 (2.15)

(ii) When $w \in Q(\mathbf{R}_+)$,

$$f_{\eta}(t;w) = \int_0^t \frac{1}{w(s)} ds.$$
 (2.16)

(iii) For $w \in W(\mathbf{R}_+)$ $G_{\eta}(t; w, \mu) = \mu + \int_{t}^{\eta} \frac{1}{w(s) f_{\eta}(s)} ds. \tag{2.17}$

(iv) $f_{\eta}(t; w, \mu)$ and $f_{\eta}(t; w)$ are abbreviated as $f_{\eta}(t)$. $G_{\eta}(t; w, \mu)$ is abbreviated as $G_{\eta}(t)$.

Now we introduce **the non-degenerate condition (NDC)** as follows. This is an explicit rewrite of the condition which was first introduced in [11] as Definition 3.3:

Definition 2.8 (the non-degenerate condition). Let $\eta > 0$ and $w \in W(\mathbf{R}_+)$. A weight function w is said to satisfy the non-degenerate condition if

$$C_0 := \inf_{0 < t \le \eta} \frac{w(t)}{t} f_{\eta}(t) > 0.$$
 (2.18)

In [11] we have established the *n*-dimensional CKN-type inequalities for p > 1. We cite the following result as the basic knowledge of this paper. Let B_n be the ball $\{x \in \mathbb{R}^n : |x| < \eta\}$.

Theorem 2.2 ([11], Theorem 3.1). Let $n \in \mathbb{N}$, $1 , <math>\mu > 0$, $\eta > 0$ and $0 \le 1/p - 1/q \le 1/n$. Assume that $w \in W(\mathbb{R}_+)$. Moreover assume that if n > 1 and p < q, w satisfies the non-degenerate condition (2.18). Then, we have the followings:

(i) There exists a positive number $C_n = C_n(p,q,\eta,\mu,w)$ such that for any $u \in C_c^{\infty}(B_{\eta} \setminus \{0\})$ we have

$$\int_{B_{\eta}} |\nabla u(x)|^{p} w(|x|)^{p-1} |x|^{1-n} dx \ge C_{n} \left(\int_{B_{\eta}} \frac{|u(x)|^{q} |x|^{1-n} dx}{w(|x|) f_{\eta}(|x|)^{1+q/p'}} \right)^{p/q}$$
(2.19)

where p' = p/(p-1) and $f_{\eta}(t)$ is given by Definition 2.7.

- (ii) If C_n is the best constant, then C_n satisfies the followings:
 - (a) If n = 1, then $C_1 = S^{p,q;1/p'}(\mathbf{R}) = 2^{p/q-1} S^{p,q;1/p'}_{rad}(\mathbf{R})$.
 - (b) If n > 1, then $C_n \ge \min(C_0^p, 1)S^{p,q;1/p'}(\mathbf{R}^n)$ (p < q); $C_n = (1/p')^p$ (p = q).
 - (c) If n > 1, $C_0 \ge 1$ and $1/p' \le \gamma_{p,q}$, then

$$C_n = S^{p,q;1/p'}(\mathbf{R}^n) = S_{\text{rad}}^{p,q;1/p'}(\mathbf{R}^n),$$
 (2.20)

where C_0 is given by (2.18) and $\gamma_{p,q}$ is given by

$$\gamma_{p,q} = \frac{n-1}{1 + q/p'}. (2.21)$$

Here $S^{p,q;1/p'}(\mathbf{R}^n)$ and $S^{p,q;1/p'}_{rad}(\mathbf{R}^n)$ are independent of η . They are called the best constants of the classical CKN-type inequalities and defined in Introduction.

- **Remark 2.3.** (i) Theorem 2.2 was established in [11] as Theorem 3.1 with C^1 -weight functions. It is an easy task to check that it remains valid for weight functions in $W(\mathbf{R}_+)$ ($\subset C^{0,1}(\mathbf{R}_+)$) defined in Definition 2.5.
- (ii) If we consider the inequality (2.19) in the radially symmetric space

$$C_c^{\infty}(B_{\eta} \setminus \{0\})_{\text{rad}} = \{ u \in C_c^{\infty}(B_{\eta} \setminus \{0\}) : u \text{ is radial} \}$$
 (2.22)

in place of $C_c^{\infty}(B_{\eta} \setminus \{0\})$, then the assertion (i) in Theorem 2.2 holds without the non-degenerate condition (2.18). By $C_{n,rad}$ we denote the best constant of the inequality (2.19) in $C_c^{\infty}(B_{\eta} \setminus \{0\})_{rad}$. Then the best constant $C_{n,rad}$ always satisfies $C_{n,rad} = S_{rad}^{p,q;1/p'}(\mathbf{R}^n)$ if $n \ge 2$. In fact, by employing a polar coordinate system, the proof is done in a quite similar way as in [11].

The following lemma gives a sufficient condition for $1/p' \le \gamma_{p,q}$ in Theorem 2.2 (ii) (c).

Lemma 2.3. Assume that n > 1, $0 \le 1/p - 1/q \le 1/n$ and 1 . Then, it holds that

$$1/p' \le \gamma_{p,q}. \tag{2.23}$$

Proof: Let 1/p - 1/q = s $(0 \le s \le 1/n)$. Then q = p/(1 - sp). By (2.21) it suffices to show that

$$1 - 1/p \le \frac{(n-1)(1-sp)}{p(1-s)}.$$

If n > 2, then we have

$$p \le \frac{n-s}{(n-2)s+1} = \frac{1}{n-2} \left(\frac{n+1/(n-2)}{s+1/(n-2)} - 1 \right).$$

The right-hand side takes its minimum (n+1)/2 at s=1/n, hence the assertion follows. If n=2, then the assertion is also clear.

2.3 A generalized rearrangement of functions

We recall a rearrangement of functions with respect to general weight functions instead of Lebesgue measure, which was used in [14] to establish the validities of $S^{p,q;\gamma} = S^{p,q;\gamma}_{rad}$ and $C^{p,q;R} = C^{p,q;R}_{rad}$ under additional conditions.

Definition 2.9. (i) For $g \in L^1_{loc}(\mathbf{R}^n)$ and $g \ge 0$ a.e. on \mathbf{R}^n , let us set for a (Lebesgue) measurable set A

$$\mu_{g}(A) = \int_{A} d\mu_{g} = \int_{A} g(x)dx.$$
 (2.24)

Then μ_g is said to be the measure determined by g.

- (ii) g is said to be admissible, if and only if $g \in L^1_{loc}(\mathbf{R}^n) \cap C(\mathbf{R}^n \setminus \{0\})_{rad}$, $g \ge 0$ on $\mathbf{R}^n \setminus \{0\}$ and g is non-increasing with respect to r = |x|. Here $C(\mathbf{R}^n \setminus \{0\})_{rad} = \{u \in C(\mathbf{R}^n \setminus \{0\}) : u \text{ is radial}\}$.
- (iii) For an admissible g and a Borel set $A \subset \mathbf{R}^n$ satisfying $0 < \mu_1(A) < +\infty$, let us define $r_g[A] > 0$ by $\mu_g(A) = \mu_g(B_{r_g[A]})$. Here the ball $B_{r_g[A]} = \{x \in \mathbf{R}^n : |x| < r_g[A]\}$ is said to be the rearrangement set of A by g.
- (iv) For an admissible g and $u : \mathbf{R}^n \to \mathbf{R}$, we set

$$\mu_g[u](t) = \mu_g(\{|u| > t\}) = \int_{\{|u| > t\}} g(x)dx \qquad \text{for } t \ge 0,$$
(2.25)

$$\mathscr{R}_g[u](x) = \mathscr{R}_g[u](|x|) = \sup\{t \ge 0 : \mu_g[u](t) > \mu_g(B_{|x|})\} \text{ for } x \in \mathbf{R}^n \setminus \{0\}.$$
 (2.26)

Then $\mu_g[u]$ and $\mathcal{R}_g[u]$ are said to be the distribution function of u and the rearrangement function of u with respect to g, respectively.

Direct from this definition we see the next proposition.

Proposition 2.4. Let $1 \le p < \infty$ and assume that g is admissible. Then, for $u : \mathbf{R}^n \to \mathbf{R}$, we have the followings:

- (i) $\mu_g[u](t) = \mu_o[\mathscr{R}_o[u]](t)$ for $t \ge 0$.
- (ii) $\mathscr{R}_{\varrho}[|u|^p](x) = \mathscr{R}_{\varrho}[u](x)^p \text{ for } x \in \mathbf{R}^n \setminus \{0\}.$
- (iii) If u is radially symmetric and non-increasing with respect to r = |x|, then

$$\mathscr{R}_{\varrho}[u](x) = u(x)$$
 for a.e. $x \in \mathbf{R}^n \setminus \{0\}$.

Further we have

Proposition 2.5. Let $1 \le p < \infty$ and assume that g is admissible. Then, for $u, v : \mathbf{R}^n \to \mathbf{R}$, we have the followings:

(i)
$$\int_{\mathbf{R}^n} |u(x)|^p g(x) dx = \int_{\mathbf{R}^n} \mathscr{R}_g[u](x)^p g(x) dx.$$

(ii)
$$\int_{\mathbf{R}^n} |u(x)v(x)|g(x)dx \leq \int_{\mathbf{R}^n} \mathscr{R}_g[u](x) \mathscr{R}_g[v](x)g(x)dx.$$

By $u \in C_{\rm c}^{0,1}({\bf R}^n)$ we denote the space of all Lipschitz continuous functions on ${\bf R}^n$ having compact supports. If $u \in C_{\rm c}^{0,1}({\bf R}^n)$, then u is differentiable for a.e. $x \in {\bf R}^n$ and $|\nabla u| \in L^\infty({\bf R}^n)$. For an admissible g, we see that $\mathscr{R}_g[u]$ for $u \in C_{\rm c}^{0,1}({\bf R}^n)$ is radially symmetric and non-increasing, and hence $\mathscr{R}_g[u]$ is differentiable for a.e. $x \in {\bf R}^n$. Moreover we have the following. For the proof see [14], Proposition 4.5.

Proposition 2.6. Let $1 \le p < \infty$ and assume that g is admissible. Then, for $u \in C_c^{0,1}(\mathbf{R}^n)$ we have

$$\int_{\mathbf{R}^n} |\nabla u(x)|^p \frac{1}{g(x)^{p-1}} dx \ge \int_{\mathbf{R}^n} |\nabla [\mathscr{R}_g[u]](x)|^p \frac{1}{g(x)^{p-1}} dx.$$

From Proposition 2.5 and Proposition 2.6 we have the following:

Proposition 2.7. Let $1 \le p < \infty$ and assume that g is admissible. Moreover assume that $v : \mathbf{R}^n \to \mathbf{R}_+$ and v is radially symmetric and non-increasing. Then, for $u \in C_c^{0,1}(\mathbf{R}^n)$ we have

$$\frac{\displaystyle\int_{\mathbf{R}^n} |\nabla u(x)|^p \frac{1}{g(x)^{p-1}} dx}{\left(\displaystyle\int_{\mathbf{R}^n} |u(x)|^q v(x) g(x) \, dx\right)^{p/q}} \geq \frac{\displaystyle\int_{\mathbf{R}^n} |\nabla [\mathscr{R}_g[u]](x)|^p \frac{1}{g(x)^{p-1}} dx}{\left(\displaystyle\int_{\mathbf{R}^n} \mathscr{R}_g[u](x)^q v(x) g(x) \, dx\right)^{p/q}}.$$

Proof: Since v is non-increasing, we have from Proposition 2.4 (iii) $\mathcal{R}_g[v](x) = v(x)$ for a.e. $x \in \mathbb{R}^n \setminus \{0\}$. Hence, by Proposition 2.5 (ii), we have

$$\int_{\mathbf{R}^n} |u(x)|^q v(x) g(x) dx \le \int_{\mathbf{R}^n} \mathscr{R}_g[|u|^q](x) \mathscr{R}_g[v](x) g(x) dx = \int_{\mathbf{R}^n} \mathscr{R}_g[u](x)^q v(x) g(x) dx.$$

Therefore the assertion holds together with Proposition 2.6.

3 Preliminary results

We begin to improve the critical CKN-type inequality (1.5) by using poly-logarithm functions defined in (2.1). Specifically we adopt the poly-logarithmic weight functions of the form

$$|x|^{p-n} \left(\left(\log^k (R\eta/|x|) \right)^{\alpha} \prod_{j=1}^{k-1} \log^j (R\eta/|x|) \right)^{p-1} \qquad (\alpha \in \mathbf{R}, k \in \mathbf{N})$$
 (3.1)

instead of $|x|^{p-n}$ on the left side of (1.5). Here we adopt the notation $\prod_{j=1}^{0} \log^{j}(R\eta/|x|) = 1$. Then we have the following:

Theorem 3.1. Let $n \in \mathbb{N}$, $1 , <math>\eta > 0$ and $0 \le 1/p - 1/q \le 1/n$. Then, we have the followings:

(i) If $\alpha \neq 1$, $k \in \mathbb{N}$ and $R > \exp^{k-1}(1)$, then there exists a positive number C_n such that we have for any $u \in C_c^{\infty}(B_n \setminus \{0\})$

$$\int_{B_{\eta}} |\nabla u(x)|^{p} |x|^{p-n} \left(\left(\log^{k}(R\eta/|x|) \right)^{\alpha} \prod_{j=1}^{k-1} \log^{j}(R\eta/|x|) \right)^{p-1} dx$$

$$\geq C_{n} |\alpha - 1|^{p-1+p/q} \left(\int_{B_{\eta}} \frac{|u(x)|^{q} dx}{|x|^{n} \left(\log^{k}(R\eta/|x|) \right)^{1+(1-\alpha)q/p'} \prod_{j=1}^{k-1} \log^{j}(R\eta/|x|)} \right)^{p/q} . \tag{3.2}$$

(ii) If $\alpha = 1$, $k \in \mathbb{N} \cup \{0\}$ and $R > \exp^k(1)$, then there exists a positive number C_n such that we have for any $u \in C_c^{\infty}(B_n \setminus \{0\})$

$$\int_{B_{\eta}} |\nabla u(x)|^{p} |x|^{p-n} \left(\prod_{j=1}^{k} \log^{j}(R\eta/|x|) \right)^{p-1} dx$$

$$\geq C_{n} \left(\int_{B_{\eta}} \frac{|u(x)|^{q} dx}{|x|^{n} \left(\log^{k+1}(R\eta/|x|) \right)^{1+q/p'} \prod_{j=1}^{k} \log^{j}(R\eta/|x|)} \right)^{p/q} .$$
(3.3)

- (iii) Assume that C_n is the best constant. Then, C_n satisfies the same assertions in Theorem 2.2 (ii).
- (iv) If 1 < n < p, $\alpha \le 1$ and R is sufficiently large, then $C_n = C_{n,\text{rad}} = S_{\text{rad}}^{p,q;1/p'}(\mathbf{R}^n)$.

Remark 3.1. (i) The inequality (3.3) coincides with (1.5) in the case k = 0.

- (ii) If $\alpha \leq 1$, then the weight functions in the inequalities (3.2) and (3.3) are locally integrable in B_{η} . Hence in this case one can adopt $C_c^{\infty}(B_{\eta})$ as a space of test functions in place of $C_c^{\infty}(B_{\eta} \setminus \{0\})$.
- (iii) If 1 , then it follows from Theorem 3.1 (iii), Lemma 2.3 and Remark 2.3 (ii) that we have (2.20) for a sufficiently large <math>R, that is, $C_n = C_{n,\mathrm{rad}} = S^{p,q;1/p'}(\mathbf{R}^n) = S^{p,q;1/p'}_{\mathrm{rad}}(\mathbf{R}^n)$.

Theorem 3.1 follows from Theorem 2.2 with $w(t) = w_{k,\alpha}(t)$ (= $w_{k,\alpha}(t;\eta)$) below.

Definition 3.1. Let $\alpha \in \mathbb{R}$, $\eta > 0$, $R > \exp^{k-1}(1)$ and $k \in \mathbb{N}$.

(i) We define $w_{k,\alpha}(t;\eta)$ such that $w_{k,\alpha}(t;\eta) \in C^{0,1}(\mathbf{R}_+)$, $w_{k,\alpha}(t;\eta) > 0$ and

$$w_{k,\alpha}(t;\eta) = \begin{cases} t \left(\prod_{j=1}^{k-1} \log^{j}(R\eta/t) \right) \left(\log^{k}(R\eta/t) \right)^{\alpha}, & 0 < t \le \eta, \\ \eta \left(\prod_{j=1}^{k-1} \log^{j} R \right) \left(\log^{k} R \right)^{\alpha}, & t \ge \eta. \end{cases}$$
(3.4)

Here if k = 1 we adopt the notation $\prod_{j=1}^{0} \log^{j}(R\eta/t) = 1$, $\prod_{j=1}^{0} \log^{j} R = 1$.

(ii) $w_{k,\alpha}(t; \eta)$ is abbreviated as $w_{k,\alpha}(t)$.

Then we see that $w_{k,\alpha}(t) \in P(\mathbf{R}_+)$ if $\alpha \leq 1$, and $w_{k,\alpha}(t) \in Q(\mathbf{R}_+)$ if $\alpha > 1$.

For $w(t) = w_{k,\alpha}(t)$ we have the followings:

(i) When $\alpha < 1$, $R > \exp^k(1)$ and $\mu = (\log^k R)^{1-\alpha}/(1-\alpha)$,

$$f_{\eta}(t) = \mu + \int_{t}^{\eta} \frac{1}{w(s)} ds = \frac{1}{1 - \alpha} \left(\log^{k}(R\eta/t) \right)^{1 - \alpha}.$$
 (3.5)

(ii) When $\alpha = 1$, $\mu = \log^{k+1} R$ and $R > \exp^{k+1}(1)$,

$$f_{\eta}(t) = \mu + \int_{t}^{\eta} \frac{1}{w(s)} ds = \log^{k+1}(R\eta/t).$$
 (3.6)

(iii) When $\alpha > 1$ and $R > \exp^k(1)$,

$$f_{\eta}(t) = \int_{0}^{t} \frac{1}{w(s)} ds = \frac{1}{\alpha - 1} \left(\log^{k}(R\eta/t) \right)^{1 - \alpha}.$$
 (3.7)

Then we have

$$\frac{w(t)}{t} f_{\eta}(t) = \begin{cases}
\frac{1}{1 - \alpha} \prod_{j=1}^{k} \log^{j}(R\eta/t), & \alpha < 1, \\
\prod_{j=1}^{k+1} \log^{j}(R\eta/t), & \alpha = 1, \quad 0 < t < \eta, . \\
\frac{1}{\alpha - 1} \prod_{j=1}^{k} \log^{j}(R\eta/t), & \alpha > 1,
\end{cases}$$
(3.8)

Moreover we see that

$$\inf_{0 < t \le \eta} \frac{w(t)}{t} f_{\eta}(t) \ge \begin{cases} \frac{1}{1 - \alpha} \prod_{j=1}^{k} \log^{j} R, & \alpha < 1, \\ \prod_{j=1}^{k+1} \log^{j} R, & \alpha = 1, \\ \frac{1}{\alpha - 1} \prod_{j=1}^{k} \log^{j} R, & \alpha > 1. \end{cases}$$
(3.9)

In particular if $R > \exp^k(1)$, then we have $C_0 := \inf_{0 < t \le \eta} \frac{w(t)}{t} f_{\eta}(t) > 0$ so that $w(t) = w_{k,\alpha}(t)$ satisfies the non-degenerate condition (2.18). Since $C_0 \to \infty$ as $R \to \infty$, $C_0 \ge 1$ holds for a sufficiently large $R > \exp^k(1)$.

Proof of Theorem 3.1: The assertions (i), (ii) and (iii) follow from Theorem 2.2. Hence, it suffices to show the assertion (iv). From Remark 3.1 (ii), we adopt $C_c^{\infty}(B_{\eta})$ as the space of test functions. We use Proposition 2.7 with appropriate g and v. To this end we define g(x) = g(|x|) by

$$g(t)^{1-p} = w_{k,\alpha}(t)^{p-1}t^{1-n} \qquad (t = |x|, t \le \eta),$$
(3.10)

that is

$$g(t) = \frac{t^{(n-p)/(p-1)}}{\left(\prod_{j=1}^{k-1} \log^j(R\eta/t)\right) \left(\log^k(R\eta/t)\right)^{\alpha}} \qquad (1 < n < p, \ \alpha \le 1).$$

We need to check that g(t) is decreasing for a sufficiently large R. By a direct calculation we have

$$g'(t) = \frac{t^{(n-p)/(p-1)-1}}{\left(\prod_{j=1}^{k-1} \log^{j}(R\eta/t)\right) \left(\log^{k}(R\eta/t)\right)^{\alpha}} \left(\frac{n-p}{p-1} + \sum_{j=1}^{k-1} \frac{1}{\prod_{l=1}^{j} \log^{l}(R\eta/t)} + \frac{\alpha}{\prod_{l=1}^{k} \log^{l}(R\eta/t)}\right)$$

$$\leq \frac{t^{(n-p)/(p-1)-1}}{\left(\prod_{j=1}^{k-1} \log^{j}(R\eta/t)\right) \left(\log^{k}(R\eta/t)\right)^{\alpha}} \left(\frac{n-p}{p-1} + \sum_{j=1}^{k-1} \frac{1}{\prod_{l=1}^{j} \log^{l}R} + \frac{\alpha}{\prod_{l=1}^{k} \log^{l}R}\right).$$
(3.11)

Since n < p, we have $g'(t) \le 0$ provided that R is sufficiently large. We define v(x) = v(|x|) for t = |x| by

$$v(t) = \begin{cases} t^{-p'(n-1)} \left(\log^k(R\eta/t) \right)^{-(1-\alpha)(1+q/p')} & (\alpha < 1), \\ t^{-p'(n-1)} \left(\log^{k+1}(R\eta/t) \right)^{-(1+q/p')} & (\alpha = 1). \end{cases}$$
(3.12)

Then we see that

$$v(t)g(t) = \begin{cases} t^{-n} \left(\prod_{j=1}^{k-1} \log^{j}(R\eta/t) \right)^{-1} \left(\log^{k}(R\eta/t) \right)^{-1 - (1-\alpha)q/p'} & (\alpha < 1), \\ t^{-n} \left(\prod_{j=1}^{k} \log^{j}(R\eta/t) \right)^{-1} \left(\log^{k+1}(R\eta/t) \right)^{-1 - q/p'} & (\alpha = 1). \end{cases}$$
(3.13)

In order to apply Proposition 2.7, we need to check if $v'(t) \le 0$. By a direct calculation we have

$$v'(t) = \begin{cases} -p'(n-1)t^{-p'(n-1)-1}(\log^{k}(R\eta/t))^{-(1-\alpha)(1+q/p')-1} \\ \times \left(\log^{k}(R\eta/t) - \frac{(1-\alpha)(1+q/p')}{p'(n-1)} \frac{1}{\prod_{j=1}^{k-1}\log^{j}(R\eta/t)}\right) & (\alpha < 1), \\ -p'(n-1)t^{-p'(n-1)-1}(\log^{k+1}(R\eta/t))^{-1} \\ \times \left(\log^{k+1}(R\eta/t) - \frac{1+q/p'}{p'(n-1)} \frac{1}{\prod_{j=1}^{k}\log^{j}(R\eta/t)}\right) & (\alpha = 1). \end{cases}$$
(3.14)

First we assume that $\alpha \neq 1$. Then we show

$$\log^{k}(R\eta/t) \ge \frac{(1-\alpha)(1+q/p')}{p'(n-1)} \frac{1}{\prod_{j=1}^{k-1} \log^{j}(R\eta/t)} \qquad (0 < t \le \eta).$$

Since $\log^{j}(R\eta/t)$ $(j=1,2,\cdots,k)$ is decreasing, it suffices to check

$$\log^k R \ge \frac{(1-\alpha)(1+q/p')}{p'(n-1)} \frac{1}{\prod_{i=1}^{k-1} \log^j R},$$

and this is valid for a sufficiently large R. Secondly we assume that $\alpha = 1$. Then, in a similar way we see that

$$\log^{k+1} R \ge \frac{1 + q/p'}{p'(n-1)} \frac{1}{\prod_{i=1}^{k} \log^{j} R}$$

holds for a sufficiently large R, hence the assertion follows.

End of the proof: By Proposition 2.7 with (3.10) and (3.12), we can assume that u is radial. Then we see that test function u for Theorem 3.1 can be assumed to be radial. The assertion (iv) therefore comes from Remark 2.3 (ii).

4 Main results

We improve Theorem 3.1 by using the super-logarithm defined in §2.1 as follows. Specifically we adopt the super-logarithmic weight functions of the form

$$|x|^{p-n} \left(\left(\prod_{l=1}^{\infty} \frac{F^l \left(\frac{a\eta}{|x|} \right)}{a} \right) \left(F^k \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{\alpha} \prod_{j=0}^{k-1} F^j \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{p-1}$$
(4.1)

instead of

$$|x|^{p-n} \left(\left(\log^k (R\eta/|x|) \right)^{\alpha} \prod_{j=1}^{k-1} \log^j (R\eta/|x|) \right)^{p-1} \qquad (\alpha \in \mathbf{R}, k \in \mathbf{N})$$

on the left side of (3.2). Then we have the following:

Theorem 4.1. Let $a > 1, k \in \mathbb{N} \cup \{0\}$, $\alpha \in \mathbb{R}$ and $\eta > 0$. Assume that $n \in \mathbb{N}$, $1 and <math>0 \le 1/p - 1/q \le 1/n$. Then, we have the followings:

(i) When $\alpha \neq 1$, there exists a positive number C_n such that we have for any $u \in C_c^{\infty}(B_{\eta} \setminus \{0\})$

$$\int_{B_{\eta}} |\nabla u(x)|^{p} |x|^{p-n} \left(\left(\prod_{l=1}^{\infty} \frac{F^{l} \left(\frac{a\eta}{|x|} \right)}{a} \right) \left(F^{k} \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{\alpha} \prod_{j=0}^{k-1} F^{j} \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{p-1} dx$$

$$\geq C_n |\alpha - 1|^{p-1+p/q} \times$$

$$\left(\int_{B_{\eta}} \frac{|u(x)|^{q} dx}{|x|^{n} \left(\prod_{l=1}^{\infty} \frac{F^{l}\left(\frac{a\eta}{|x|}\right)}{a}\right) \left(F^{k}\left(L\left(\frac{\eta}{|x|}\right) + a\right)\right)^{1 + (1 - \alpha)q/p'} \prod_{j=0}^{k-1} F^{j}\left(L\left(\frac{\eta}{|x|}\right) + a\right)}\right)^{p/q}, \tag{4.2}$$

where if k = 0 we adopt the notation $\prod_{i=0}^{k-1} F^{j}(u) = 1$.

Here $F(u) = F_a(u) = a - \log a + \log u$ $(u \ge a)$, $F^k(u) = F(F^{k-1}(u))$ for $u \ge a > 1$ and L is the super-logarithm defined by Definition 2.4.

(ii) When $\alpha = 1$, there exists a positive number C_n such that we have for any $u \in C_c^{\infty}(B_{\eta} \setminus \{0\})$

$$\int_{B_{\eta}} |\nabla u(x)|^{p} |x|^{p-n} \left(\left(\prod_{l=1}^{\infty} \frac{F^{l} \left(\frac{a\eta}{|x|} \right)}{a} \right) \prod_{j=0}^{k} F^{j} \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{p-1} dx$$

$$\geq C_{n} \left(\int_{B_{\eta}} \frac{|u(x)|^{q} dx}{|x|^{n} \left(\prod_{l=1}^{\infty} \frac{F^{l} \left(\frac{a\eta}{|x|} \right)}{a} \right) \left(F^{k+1} \left(L \left(\frac{\eta}{|x|} \right) + a \right) \right)^{1+q/p'} \prod_{j=0}^{k} F^{j} \left(L \left(\frac{\eta}{|x|} \right) + a \right)} \right)^{p/q} . \tag{4.3}$$

(iii) Assume that C_n is the best constant. Then, C_n satisfies the same assertions in Theorem 2.2 (ii). Further if a > 1 and $a \ge |\alpha - 1|^{1/(k+1)}$, then we have $C_0 \ge 1$.

(iv) If 1 < n < p, $\alpha \le 1$ and a is sufficiently large, then $C_n = C_{n,rad} = S_{rad}^{p,q;1/p'}(\mathbf{R}^n)$.

Remark 4.1. (i) The inequalities (4.2) and (4.3) improve (1.5), (3.2) and (3.3). In particular if k = 0 in (4.3), then the weight in the left-hand side is simply

$$|x|^{p-n}\left(\prod_{l=1}^{\infty}\frac{F^l\left(a\eta/|x|\right)}{a}\right)^{p-1}.$$

Roughly speaking this can be thought of as a formal limit of

$$|x|^{p-n} \left(\prod_{l=1}^k \log^l(R\eta/|x|) \right)^{p-1}$$

as $k \to \infty$.

- (ii) If $\alpha \leq 1$, then the weight functions in the inequalities (4.2) and (4.3) are locally integrable in B_{η} . Hence one can adopt $C_c^{\infty}(B_{\eta})$ as a space of test functions in place of $C_c^{\infty}(B_{\eta} \setminus \{0\})$.
- (iii) If 1 , then it follows from Theorem 4.1 (iii), Lemma 2.3 and Remark 2.3 (ii) that we have (2.20) for <math>a > 1 and $a \ge |\alpha 1|^{1/(k+1)}$, that is, $C_n = C_{n, rad} = S^{p,q;1/p'}(\mathbf{R}^n) = S^{p,q;1/p'}_{rad}(\mathbf{R}^n)$.

Theorem 4.1 will be shown by using Theorem 2.2 with $w(t) = w_{k,\alpha}^{sl}(t)$ $(= w_{k,\alpha}^{sl}(t; \eta))$ below.

Definition 4.1. (i) We define $w_{k,\alpha}^{sl}(t;\eta) \in C^{0,1}(\mathbb{R}_+)$ for $k \in \mathbb{N} \cup \{0\}$ such that

$$w_{k,\alpha}^{sl}(t;\eta) = \begin{cases} t \left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a} \right) \left(\prod_{j=0}^{k-1} F^{j}(L(\eta/t) + a) \right) \left(F^{k}(L(\eta/t) + a) \right)^{\alpha} & (0 < t \le \eta), \\ \eta a^{k+\alpha} & (t \ge \eta). \end{cases}$$

$$(4.4)$$

Here if k = 0 we adopt the notation $\prod_{j=0}^{-1} F^j(L(\eta/t) + a) = 1$.

(ii) $w_{k,\alpha}^{sl}(t; \eta)$ is abbreviated as $w_{k,\alpha}^{sl}(t)$.

Then we have the following:

Lemma 4.2. Let $w(t) = w_{k,\alpha}^{sl}(t) \in W(\mathbf{R}_+)$ be given by (4.4). Then,

(i) The corresponding $f_{\eta}(t)$ $(0 < t < \eta)$ is given by

$$f_{\eta}(t) = \begin{cases} \mu + \int_{t}^{\eta} \frac{1}{w(s)} ds = \frac{1}{1-\alpha} \left(F^{k} (L(\eta/t) + a) \right)^{1-\alpha} & (\alpha < 1, \mu = \frac{1}{1-\alpha} a^{1-\alpha}), \\ \mu + \int_{t}^{\eta} \frac{1}{w(s)} ds = F^{k+1} (L(\eta/t) + a) & (\alpha = 1, \mu = a), \\ \int_{0}^{t} \frac{1}{w(s)} ds = \frac{1}{\alpha - 1} \left(F^{k} (L(\eta/t) + a) \right)^{1-\alpha} & (\alpha > 1). \end{cases}$$
(4.5)

(ii) $\lim_{t\to+0} \frac{w(t)}{t} f_{\eta}(t) = \infty$. In particular if a > 1 and $a \ge |\alpha - 1|^{1/(k+1)}$, then w(t) satisfies the non-degenerate condition (2.18) with $C_0 \ge 1$.

Proof of (i): Let $w(t) = w_{k,\alpha}^{sl}(t)$. First we assume that $\alpha < 1$. Since it holds that by (2.11)

$$\frac{1}{1-\alpha} \frac{d}{dt} \left(F^k(L(t)+a) \right)^{1-\alpha} = \frac{1}{t \left(\prod_{l=1}^{\infty} \frac{F^l(at)}{a} \right) \left(\prod_{j=0}^{k-1} F^j(L(t)+a) \right) \left(F^k(L(t)+a) \right)^{\alpha}}, \tag{4.6}$$

we see that

$$\int_{t}^{\eta} \frac{1}{w(s)} ds = \int_{1}^{\eta/t} \frac{1}{u \left(\prod_{l=1}^{\infty} \frac{F^{l}(au)}{a} \right) \left(\prod_{j=0}^{k-1} F^{j}(L(u)+a) \right) (F^{k}(L(u)+a))^{\alpha}} du \quad (\eta/s = u)$$

$$= \int_{1}^{\eta/t} \frac{1}{1-\alpha} \frac{d}{du} \left(F^{k}(L(u)+a) \right)^{1-\alpha} du$$

$$= \frac{1}{1-\alpha} \left(\left(F^{k}(L(\eta/t)+a) \right)^{1-\alpha} - \left(F^{k}(L(1)+a) \right)^{1-\alpha} \right)$$

$$= \frac{1}{1-\alpha} \left(F^{k}(L(\eta/t)+a) \right)^{1-\alpha} - \mu \quad \left(\mu = \frac{1}{1-\alpha} \left(F^{k}(L(1)+a) \right)^{1-\alpha} = \frac{a^{1-\alpha}}{1-\alpha} \right).$$

Secondly we assume that $\alpha > 1$. By Theorem 2.1 (ii) we have $\lim_{t \to +0} F^k(L(\eta/t) + a) = \infty$. Then in a similar way we have

$$\int_{0}^{t} \frac{1}{w(s)} ds = \frac{1}{\alpha - 1} \left(F^{k} \left(L(\eta/t) + a \right) \right)^{1 - \alpha}. \tag{4.8}$$

Thirdly we assume that $\alpha = 1$. Noting that

$$F^{k+1}\left(L(t)+a\right)=F(F^{k}\left(L(t)+a\right)) \ \ \text{and} \ \ \frac{d}{dt}F^{k+1}\left(L(\eta/t)+a\right)=\frac{1}{t\left(\prod_{l=1}^{\infty}\frac{F^{l}\left(at\right)}{a}\right)\prod_{j=0}^{k}F^{j}\left(L(t)+a\right)},$$

we have

$$\int_{t}^{\eta} \frac{1}{w(s)} ds = F^{k+1} \left(L(\eta/t) + a \right) - \mu \qquad (w(s) = w_{k,1}^{sl}(s), \ \mu = F^{k+1} \left(L(1) + a \right) = a \right). \tag{4.9}$$

Therefore we see that if $\alpha \leq 1$, $w_{k,\alpha}^{sl}(t) \in P(\mathbf{R}_+)$ and if $\alpha > 1$, $w_{k,\alpha}^{sl}(t) \in Q(\mathbf{R}_+)$. By Definition 2.7 we immediately have (4.5) from (4.7), (4.8) and (4.9).

Proof of (ii). Let $w(t) = w_{k,\alpha}^{sl}(t) \in W(\mathbf{R}_+)$. From (4.5) we have

$$\frac{w(t)}{t}f_{\eta}(t) = \begin{cases}
\frac{1}{1-\alpha} \left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a} \right) \prod_{j=0}^{k} F^{j}(L(\eta/t) + a) & (\alpha < 1), \\
\left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a} \right) \left(\prod_{j=0}^{k} F^{j}(L(\eta/t) + a) \right) F^{k+1}(L(\eta/t) + a) & (\alpha = 1), \\
\frac{1}{\alpha - 1} \left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a} \right) \prod_{j=0}^{k} F^{j}(L(\eta/t) + a) & (\alpha > 1).
\end{cases} (4.10)$$

Because $\prod_{l=1}^{\infty} \frac{F^l(a\eta/t)}{a} \ge 1$, $\lim_{t\to +0} F^k(L(\eta/t)+a) = \infty$ and $F^k(L(\eta/t)+a) \ge a$ for $k \in \mathbb{N} \cup \{0\}$ and $0 < t < \eta$, we have $\lim_{t\to +0} \frac{w(t)}{t} f_{\eta}(t) = \infty$ and the estimate

$$\frac{w(t)}{t} f_{\eta}(t) \ge \begin{cases} \frac{a^{k+1}}{|1-\alpha|} & (\alpha \ne 1), \\ a^{k+2} & (\alpha = 1). \end{cases}$$
(4.11)

Therefore, the assertion (ii) is satisfied, if a > 1 and $a \ge |\alpha - 1|^{1/(k+1)}$.

Proof of Theorem 4.1: It suffices to show the assertion (iv). From Remark 4.1 (ii), we adopt $C_c^{\infty}(B_{\eta})$ as the space of test functions. As before we apply Proposition 2.7 with appropriate g and v. Then we see that test function $u \in C_c^{\infty}(B_{\eta})$ for Theorem 4.1 can be assumed to be radial. The assertion (iv) therefore comes from Remark 2.3 (ii).

To this end we define g(x) = g(|x|) by

$$g(t)^{1-p} = w_{k,\alpha}^{sl}(t)^{p-1}t^{1-n} \qquad (t = |x|, t \le \eta).$$
(4.12)

Then

$$g(t) = \frac{t^{(n-p)/(p-1)}}{\left(\prod_{l=1}^{\infty} \frac{F^l(a\eta/t)}{a}\right) \left(\prod_{j=0}^{k-1} F^j(L(\eta/t) + a)\right) \left(F^k(L(\eta/t) + a)\right)^{\alpha}} \qquad (\alpha \le 1).$$

We also define $v_k(x) = v_k(|x|)$ for $k \in \mathbb{N} \cup \{0\}$ and t = |x| by

$$v_k(t) = \begin{cases} t^{-p'(n-1)} \left(F^k \left(L(\eta/t) + a \right) \right)^{-(1-\alpha)(1+q/p')} & (\alpha < 1), \\ t^{-p'(n-1)} \left(F^{k+1} \left(L(\eta/t) + a \right) \right)^{-(1+q/p')} & (\alpha = 1). \end{cases}$$
(4.13)

Then we show the following:

Lemma 4.3. *Let* $k \in \mathbb{N} \cup \{0\}$.

- (i) If a is sufficiently large, then g(t) $(0 < t \le \eta)$ is decreasing.
- (ii) If a is sufficiently large, then $v_k(t)$ $(0 < t \le \eta)$ is decreasing.

Proof of (i): We note the following relations: For any $k \in \mathbb{N}$ and a > 1,

$$\begin{cases} L'(r) = \frac{d}{dr}(a + L(r)) = \frac{a}{\tilde{F}(ar)} = \frac{1}{r \prod_{j=1}^{\infty} \frac{F^{j}(ar)}{a}} & (r > 1), \\ \frac{d}{dr}F^{k}(a + L(r)) = \frac{L'(r)}{k-1} & (r > 1), \\ \prod_{j=0}^{K-1} F^{j}(a + L(r)) & (r > 1), \end{cases}$$

$$\begin{cases} \frac{d}{du}\tilde{F}(u) = \tilde{F}(u) \left(\sum_{k=0}^{\infty} \frac{1}{\prod_{j=0}^{k} F^{j}(u)}\right) & (u > a), \\ \frac{d}{dr} \left(\prod_{j=1}^{\infty} \frac{F^{j}(ar)}{a}\right) = \left(\prod_{j=1}^{\infty} \frac{F^{j}(ar)}{a}\right) \sum_{k=1}^{\infty} \frac{1}{r \prod_{j=1}^{k} F^{j}(ar)} & (r > 1). \end{cases}$$

$$(4.14)$$

By (4.14) we have

$$g'(t) = \frac{g(t)}{t}G(t), \qquad (0 < t < \eta)$$
 (4.15)

where

$$G(t) = \frac{n-p}{p-1} + \sum_{k=1}^{\infty} \frac{1}{\prod_{j=1}^{k} F^{j}(a\eta/t)} + \sum_{j=0}^{k-1} \frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a}\right) \prod_{i=0}^{j} F^{i}(L(\eta/t) + a)} + \alpha \frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a}\right) \prod_{i=0}^{k} F^{i}(L(\eta/t) + a)}.$$

$$(4.16)$$

Then G(t) satisfies

$$G(t) \le \frac{n-p}{p-1} + \sum_{k=1}^{\infty} \frac{1}{a^k} + \sum_{j=0}^{k-1} \frac{1}{a^{j+1}} + |\alpha| \frac{1}{a^{k+1}} < \frac{n-p}{p-1} + \frac{2}{a-1} + |\alpha| \frac{1}{a^{k+1}}.$$

Since p > n, we see that g'(t) < 0 if a is sufficiently large.

Proof of (ii): First we assume that $\alpha = 1$. Then

$$v'_{k}(t) = -p'(n-1)t^{-p'(n-1)-1} \left(F^{k+1} \left(L(\eta/t) + a\right)\right)^{-1+q/p'-1} \times \left(F^{k+1} \left(L(\eta/t) + a\right) - \frac{1+q/p'}{p'(n-1)} \frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^{l} \left(a\eta/t\right)}{a}\right) \prod_{j=0}^{k} F^{j} \left(L(\eta/t) + a\right)}\right).$$
(4.17)

It suffices to show that if a is large, then

$$F^{k+1}(L(\eta/t)+a) \ge \frac{1+q/p'}{p'(n-1)} \frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^l(a\eta/t)}{a}\right) \prod_{j=0}^{k} F^j(L(\eta/t)+a)}.$$
 (4.18)

Since $F^j(L(\eta/t)+a)$ $(j \in \mathbb{N} \cup \{0\})$ and $\prod_{l=1}^{\infty} \frac{F^l(a\eta/t)}{a}$ are decreasing, we have

$$F^{k+1}(L(\eta/t) + a) \ge F^{k+1}(L(1) + a) = a$$

and

$$\frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^{l}(a\eta/t)}{a}\right) \prod_{j=0}^{k} F^{j}(L(\eta/t) + a)} \leq \frac{1}{\prod_{j=0}^{k} F^{j}(L(1) + a)} = \frac{1}{a^{k+1}}.$$

Hence, if

$$a^{k+2} \ge (1 + q/p')/(p'(n-1))$$

is satisfied, then $v_k'(t) < 0$ $(0 < t < \eta)$. Therefore we have the desired result. We proceed to the case that $\alpha \neq 1$. Then we have

$$\begin{split} v_k'(t) &= -p'(n-1)t^{-p'(n-1)-1} \left(F^k \left(L(\eta/t) + a \right) \right)^{-(1-\alpha)(1+q/p')-1} \\ &\times \left(F^k \left(L(\eta/t) + a \right) - \frac{(1-\alpha)(1+q/p')}{p'(n-1)} \frac{1}{\left(\prod_{l=1}^{\infty} \frac{F^l \left(a\eta/t \right)}{a} \right) \prod_{j=0}^{k-1} F^j \left(L(\eta/t) + a \right)} \right), \end{split} \tag{4.19}$$

where we use the notation $\prod_{i=0}^{-1} F^{i}(L(\eta/t) + a) = 1$. In a similar way, if

$$a^{k+1} \ge (1-\alpha)(1+q/p')/(p'(n-1))$$

holds, then we see that $v'_{k}(t) < 0$ (0 < $t < \eta$), and the assertion is proved.

End of the proof of Theorem 4.1 (iv): We have

$$\begin{split} & v_k(t)g(t) = \\ & \left\{ t^{-n} \left(\left(\prod_{l=1}^{\infty} \frac{F^l\left(a\eta/t\right)}{a} \right) \left(\prod_{j=0}^{k-1} F^j\left(L(\eta/t) + a\right) \right) \right)^{-1} \left(F^k\left(L(\eta/t) + a\right) \right)^{-1 - (1-\alpha)q/p'} & (\alpha < 1), \\ & t^{-n} \left(\left(\prod_{l=1}^{\infty} \frac{F^l\left(a\eta/t\right)}{a} \right) \left(\prod_{j=0}^{k} F^j\left(L(\eta/t) + a\right) \right) \right)^{-1} \left(F^{k+1}\left(L(\eta/t) + a\right) \right)^{-(1+q/p')} & (\alpha = 1). \end{split}$$

Then, the best constant C_n is given by

$$C_n = \inf_{u \in C_c^{\infty}(B_{\eta}) \setminus \{0\}} \frac{\int_{B_{\eta}} |\nabla u(x)|^p w_{k,\alpha}^{sl}(|x|)^{p-1} |x|^{1-n} dx}{\left(\int_{B_{\eta}} \frac{|u(x)|^q |x|^{1-n}}{w_{k,\alpha}^{sl}(|x|) f_{\eta}(|x|)^{1+q/p'}} dx\right)^{p/q}} = \inf_{u \in C_c^{\infty}(B_{\eta}) \setminus \{0\}} \frac{\int_{B_{\eta}} |\nabla u(x)|^p \frac{1}{g(x)^{p-1}} dx}{\left(\int_{B_{\eta}} |u(x)|^q v_k(x) g(x) dx\right)^{p/q}}.$$

As functions g(t) and v(t) in Proposition 2.7, we adopt g(t) defined by (4.12) and $v_k(t)$ defined by (4.13) respectively. Noting that $C_c^{\infty}(B_{\eta})$ can be replaced by $C_c^{0,1}(B_{\eta})$ using a density argument, it follows from Proposition 2.7 that u can be assumed radial. Then we can adopt $C_c^{\infty}(B_{\eta})_{\text{rad}}$ as a space of test functions in Theorem 4.1. The assertion (iv) therefore comes from Remark 2.3 (ii).

5 Further results

If p = q holds, then the inequalities (4.2) and (4.3) become the Hardy-type. In this case we can further improve them by adding sharp remainders. To this end we recall and modify the one-dimensional Hardy's inequalities in [10]. The following is an easy corollary to Theorem 3.1 in [10].

Proposition 5.1. Assume that $1 , <math>\eta > 0$, $\mu > 0$ and $w \in W(\mathbf{R}_+)$. Then, there exist a positive number $C = C(w, p, \eta, \mu)$ such that for every $u \in C_c^{\infty}((0, \eta))$, we have

$$\int_0^{\eta} |u'(t)|^p w(t)^{p-1} dt \ge (1/p')^p \int_0^{\eta} \frac{|u(t)|^p dt}{w(t) f_n(t)^p} + C \int_0^{\eta} \frac{|u(t)|^p dt}{w(t) f_n(t)^p G_n(t)^2}, \tag{5.1}$$

where $f_{\eta}(t)$ and $G_{\eta}(t)$ are defined by Definition 2.7 and the coefficient $(1/p')^p$ is sharp.

Then we immediately have the *n*-dimensional version:

Corollary 5.2. Assume that $n \in \mathbb{N}$, $1 , <math>\eta > 0$, $\mu > 0$ and $w \in W(\mathbb{R}_+)$. Then, there exist a positive number $C = C(w, p, \eta, \mu)$ such that for every $u \in C_c^{\infty}(B_{\eta} \setminus \{0\})$, we have

$$\int_{B_{\eta}} |\nabla u(x)|^{p} w(|x|)^{p-1} |x|^{1-n} dx \ge (1/p')^{p} \int_{B_{\eta}} \frac{|u(x)|^{p} |x|^{1-n} dx}{w(|x|) f_{\eta}(|x|)^{p}} + C \int_{B_{\eta}} \frac{|u(x)|^{p} |x|^{1-n} dx}{w(|x|) f_{\eta}(|x|)^{p} G_{\eta}(|x|)^{2}}. \quad (5.2)$$

From Corollary 5.2 with $w(t) = w_{k,1}^{sl}(t)$ we have the following. For the sake of simplicity, the results are limited to case that $\alpha = 1$. By Remark 4.1 (ii) we can adopt $C_c^{\infty}(B_{\eta})$ as the space of test functions. (See also Remark 3.1(ii)).

Proposition 5.3. Let $k \in \mathbb{N} \cup \{0\}$ and $\eta > 0$. Let $w_{k,1}^{sl} \in W(\mathbb{R}_+)$ be given by Definition 4.1. Assume that $n \in \mathbb{N}$, 1 < p and a > 1. Then, there exists a positive number $C = C(k, p, \eta, a)$ such that for any $u \in C_c^{\infty}(B_{\eta})$

$$\int_{B_{\eta}} |\nabla u(x)|^{p} w_{k,1}^{sl}(|x|)^{p-1} |x|^{1-n} dx \ge (1/p')^{p} \int_{B_{\eta}} \frac{|u(x)|^{p} |x|^{1-n} dx}{w_{k,1}^{sl}(|x|) \left(F^{k+1} \left(L(\eta/|x|) + a\right)\right)^{p}} + C \int_{B_{\eta}} \frac{|u(x)|^{p} |x|^{1-n} dx}{w_{k,1}^{sl}(|x|) \left(F^{k+1} \left(L(\eta/|x|) + a\right)\right)^{p} \left(a - \log a + \log \left(F^{k+1} \left(L(\eta/|x|) + a\right)\right)\right)^{2}}.$$
(5.3)

Here, $w_{k,1}^{sl}(t)$ is defined by (4.4) and the coefficient $(1/p')^p$ is sharp.

Proof: By virtue of (2.15) and (2.17) in Definition 2.7, we can calculate $G_{\eta}(t)$ to obtain

$$G_{\eta}(t) = \mu - \log \mu + \log \left(\mu + \int_{t}^{\eta} \frac{1}{w(s)} ds\right).$$
 (5.4)

By (4.9) we have

$$G_{\eta}(t; w_{k,1}^{sl}, \mu) = \mu - \log \mu + \log \left(F^{k+1} \left(L(\eta/t) + a \right) \right), \tag{5.5}$$

where $\mu = F^{k+1}(L(1) + a) = a$. Therefore we have (5.3).

References

- [1] H. Ando, T. Horiuchi and E. Nakai, Construction of slowly increasing functions, Sci. Math. Jpn., Vol.75, No.2 (2012), pp.187–201.
- [2] H. Ando, T. Horiuchi, E. Nakai, Weighted Hardy inequalities with infinitely many sharp missing terms, Math. J. Ibaraki Univ., Vol.46 (2014), pp.9–30.

- [3] H. Ando, T. Horiuchi and E. Nakai, Some properties of slowly increasing functions, Math. J. Ibaraki Univ., Vol.46 (2014), pp.37–49.
- [4] H. Ando, T. Horiuchi, Weighted Hardy's inequalities and the variational problem with compact perturbations, Math. J. Ibaraki Univ., Vol.52 (2020), pp.15-26.
- [5] H. Ando, T. Horiuchi, Generalized weighted Hardy's inequalities with compact perturbations. (preprint)
- [6] L. Caffarelli, R. Kohn, L. Nirenberg, First order interpolation inequalities with weights, Compositio Math., Vol.53 (1984), No. 3, pp.259–275.
- [7] K. Fujiwara, T. Ozawa, Weighted L^p -boundedness of convolution type integral operators associated with bilinear estimates in the Sobolev spaces, J. Math. Soc. Japan, Vol.68, No. 1 (2016) pp.169–191.
- [8] T. Horiuchi, The imbedding theorems for weighted Sobolev spaces, J. Math. Kyoto Univ., Vol.29 (1989), pp. 365–403.
- [9] T. Horiuchi, Best constant in weighted Sobolev inequality with weights being powers of distance from the origin, J. Ineq. and Appl., Vol.1 (1997), pp. 275–292.
- [10] T. Horiuchi, Hardy's inequalities with non-doubling weights and sharp remainders, Sci. Math. Jpn., Vol.85, No.2 (2023), pp.125–147.
- [11] T. Horiuchi, On general Caffarelli-Kohn-Nirenberg type inequalities involving non-doubling weights, arXiv:2210.17173 [math.AP], Sci. Math. Jpn. (in Editions Electronica), e-2022-10, 16 pages.
- [12] T. Horiuchi, Refinement of classical inequalities, Uchida Rokakuho, 2023. (Japanese)
- [13] T. Horiuchi, On general Caffarelli-Kohn-Nirenberg type inequalities involving non-doubling weights (p = 1). (preprint)
- [14] T. Horiuchi, P. Kumlin, On the Caffarelli-Kohn-Nirenberg type inequalities involving Critical and Supercritical Weights, Kyoto J. Math., Vol.52, No.4 (2012), pp. 661–742.
- [15] X. Liu, T. Horiuchi, H. Ando, One dimensional weighted Hardy's inequalities and application. J. Math. Ineq., Vol.14 (2020), No. 4, pp.1203–1222.
- [16] M. Sano, F. Takahashi, On eigenvalue problems involving the critical Hardy potential and Sobolev type inequalities with logarithmic weights in two dimensions. arXiv: 2210. 10282v1 [math.AP] 19 Oct 2022.
- [17] V.G. Maz'ja, Sobolev spaces, Springer, 1985.

Hiroshi Ando, Toshio Horiuchi, Eiichi Nakai Department of Mathematics Faculty of Science Ibaraki University Mito, Ibaraki, 310, Japan

e-mail:

hiroshi.ando.math@vc.ibaraki.ac.jp toshio.horiuchi.math@vc.ibaraki.ac.jp eiichi.nakai.math@vc.ibaraki.ac.jp