# ON INFINITELY DIVISIBLE DISTRIBUTIONS RELATED TO THE RIEMANN HYPOTHESIS

TAKASHI NAKAMURA AND MASATOSHI SUZUKI

Abstract. For the Riemann zeta-function, we introduce a function such that it is a characteristic function of an infinitely divisible distribution on the real line if and only if the Riemann Hypothesis is true.

### 1. INTRODUCTION

The Riemann zeta-function  $\zeta(s)$  is a function of a complex variable  $s = \sigma + it$   $(\sigma, t \in \mathbb{R})$ defined by the Dirichlet series

$$
\zeta(s) := \sum_{n=1}^{\infty} \frac{1}{n^s}
$$

when  $\Re(s) = \sigma > 1$  and extends meromorphically to C. Since  $\zeta(s)$  is holomorphic except for a simple pole at  $s = 1$ , the Riemann xi-function

<span id="page-0-3"></span>
$$
\xi(s) := \frac{1}{2}s(s-1)\pi^{-s/2}\Gamma\left(\frac{s}{2}\right)\zeta(s)
$$
\n(1.1)

isan entire function, where  $\Gamma(s)$  is the Gamma function ([\[9,](#page-4-0) Theorem 2.1]). The Riemann Hypothesis (RH, for short), a famous open problem, claims that all non-real zeros of  $\zeta(s)$  lie on the critical line  $\Re(s) = 1/2$ , which is equivalent to all the zeros of  $\xi(1/2-i\zeta)$ being real. The main result of the present paper states an equivalence condition for the RH in terms of infinitely divisible distributions in probability theory.

A distribution (or probability measure)  $\mu$  on  $\mathbb R$  is infinitely divisible if there exists a distribution  $\mu_n$  on R such that  $\mu = \mu_n * \cdots * \mu_n$  (*n*-fold) for every positive integer n. For any infinitely divisible distribution  $\mu$ , there exists a triplet  $(a, b, \nu)$  consisting of  $a \in \mathbb{R}_{\geq 0}, b \in \mathbb{R}$  and a measure  $\nu$  on  $\mathbb R$  such that the characteristic function

$$
\widehat{\mu}(t) = \int_{-\infty}^{\infty} e^{itx} \mu(dx)
$$

has the Lévy–Khintchine formula

<span id="page-0-0"></span>
$$
\widehat{\mu}(t) = \exp\left[-\frac{1}{2}at^2 + ibt + \int_{-\infty}^{\infty} \left(e^{it\lambda} - 1 - \frac{it\lambda}{1 + \lambda^2}\right)\nu(d\lambda)\right],
$$
\n(1.2)

<span id="page-0-2"></span>
$$
\nu({0}) = 0, \quad \int_{-\infty}^{\infty} \min(1, \lambda^2) \nu(d\lambda) < \infty \tag{1.3}
$$

([\[7,](#page-4-1) Theorem 8.1 and Remark 8.4]). The measure  $\nu$  is referred to as the Lévy measure for  $\mu$ . If the Lévy measure  $\nu$  satisfies  $\int_{|\lambda| \leq 1} |\lambda| \nu(d\lambda) < \infty$ , then [\(1.2\)](#page-0-0) can be rewritten as

<span id="page-0-1"></span>
$$
\widehat{\mu}(t) = \exp\left[-\frac{1}{2}at^2 + ib_0t + \int_{-\infty}^{\infty} (e^{it\lambda} - 1) \nu(d\lambda)\right]
$$
\n(1.4)

 $([7, (8.7)]).$  $([7, (8.7)]).$  $([7, (8.7)]).$ 

Date: Version of June 16, 2023.

<sup>2020</sup> Mathematics Subject Classification. 11M26 60E07 .

Key words and phrases, characteristic function; infinitely divisible distribution; Lévy-Khintchine formula; Riemann zeta-function; Riemann Hypothesis.

The relation between the RH and an infinitely divisible distribution is formulated using the function  $g_{\zeta}(t)$  defined by

<span id="page-1-4"></span>
$$
g_{\zeta}(t) := -4\left(e^{t/2} + e^{-t/2} - 2\right) + \sum_{n \leq e^t} \frac{\Lambda(n)}{\sqrt{n}} (t - \log n)
$$
  

$$
-\frac{t}{2} \left(\psi(1/4) - \log \pi\right) + \frac{1}{4} \left(e^{-t/2} \Phi(e^{-2t}, 2, 1/4) - \Phi(1, 2, 1/4)\right)
$$
\n(1.5)

for non-negative t and  $g_{\zeta}(t) := g_{\zeta}(-t)$  for negative t, where  $\Lambda(n) = \log p$  if n is a power of a prime p and otherwise  $\Lambda(n) = 0$ ,  $\psi(z) = \Gamma'(z)/\Gamma(z)$  is the digamma function, and  $\Phi(z,s,a) = \sum_{n=0}^{\infty} (n+a)^{-s} z^n$  is the Hurwitz–Lerch zeta-function. The function  $g_{\zeta}(t)$ was originally introduced in [\[8,](#page-4-2) (1.1), (1.3), and (1.8)] to study equivalent conditions for the RH in relation to Weil's positivity or Li's criterion, etc.

<span id="page-1-0"></span>Then the main results are stated as follows:

**Theorem 1.1.** Let  $g_{\zeta}(t)$  be the function on the real line defined as above. Then the following two are equivalent:

- (1) the Riemann Hypothesis is true;
- (2)  $\exp(g_{\zeta}(t))$  is a characteristic function of an infinitely divisible distribution on R.

Under the RH, the Lévy measure of the corresponding infinitely divisible distribution is also explicitly described as follows.

<span id="page-1-3"></span>**Theorem 1.2.** Let  $g_{\zeta}(t)$  be as in Theorem [1.1.](#page-1-0) Assume that the RH is true and define

<span id="page-1-1"></span>
$$
\nu_{\zeta}(d\lambda) := \sum_{\gamma} \frac{m_{\gamma}}{\gamma^2} \delta_{-\gamma}(d\lambda),\tag{1.6}
$$

where  $\gamma$  runs through all zeros of  $\xi(1/2 - iz)$  without counting multiplicity,  $m_{\gamma}$  is the multiplicity of  $\gamma$ , and  $\delta_x$  is the delta measure at x. Then the Lévy–Khintchine formula [\(1.4\)](#page-0-1) of  $\exp(g_{\zeta}(t))$  holds for the triplet  $(a, b_0, \nu) = (0, 0, \nu_{\zeta})$ , that is,  $\exp(g_{\zeta}(t))$  is a characteristic function of a compound Poisson distribution.

The above two results are proved in Section [2.](#page-2-0) The core of the proof is the equality

<span id="page-1-2"></span>
$$
g_{\zeta}(t) = \sum_{\gamma} m_{\gamma} \frac{e^{-i\gamma t} - 1}{\gamma^2} \tag{1.7}
$$

proved in [\[8,](#page-4-2) Theorem 1.1 (2)], where the sum and  $m_{\gamma}$  have the same meaning as in  $(1.6)$ . We give a proof of  $(1.7)$  in the appendix (Section [3\)](#page-3-0) for the convenience of readers.

It has long been known that the normalized function  $Z_{\sigma}(t) := \zeta(\sigma + it)/\zeta(\sigma)$  is an infinitely divisible characteristic function for  $\sigma > 1$ . More precisely, the distribution generated by  $Z_{\sigma}(t)$  is compound Poisson on R and

$$
Z_{\sigma}(t) = \exp\left[\int_{-\infty}^{\infty} (e^{it\lambda} - 1) N_{\sigma}(d\lambda)\right], \quad N_{\sigma}(d\lambda) := \sum_{p \in \mathbb{P}} \sum_{r=1}^{\infty} \frac{p^{-r\sigma}}{r} \delta_{-r \log p}(d\lambda),
$$

where $\mathbb P$  denotes the set of all prime numbers ([\[2,](#page-4-3) Example 17.6]). Lin and Hu generalized this result for  $\sum_{n=1}^{\infty} c(n) n^{-s}$ , where  $c(n)$  are completely multiplicative non-negative coefficientsand  $c(n) > 0$  for some  $n \geq 2$  ([\[5,](#page-4-4) Theorem 2]). Other examples of studies dealing with the Riemann zeta-function from a probabilistic point of view, we cite papers reviewed in Biane–Pitman–Yor [\[1\]](#page-4-5), Lagarias–Rains [\[3\]](#page-4-6), and Nakamura [\[6\]](#page-4-7). In particular, Nakamura discovered a direct relation between infinitely divisible distributions and the RH. He proved that the RH is true if and only if  $\Xi_{\sigma}(t) := \xi(\sigma+it)/\xi(\sigma)$  is a characteristic function of a pretended-infinitely divisible distribution on  $\mathbb R$  for every  $1/2 < \sigma < 1$ , where a pretended-infinitely divisible distribution means a distribution satisfies [\(1.2\)](#page-0-0) but does not necessarily assume [\(1.3\)](#page-0-2), and the measure  $\nu$  is a signed measure on R.

The relation between the Riemann zeta-function and probability theory in the above literature is based on the Dirichlet series representation of  $\zeta(s)$  or the integral representation of  $\xi(s)$ . In contrast, the results in the present paper are based on the integral representation of the logarithmic derivative of  $\xi(s)$  in Lemma [2.2](#page-2-1) below. This causes differences from known results.

## 2. Proof of Theorems [1.1](#page-1-0) and [1.2](#page-1-3)

<span id="page-2-2"></span><span id="page-2-0"></span>We start the proof with  $(1.7)$ , and note the following.

**Lemma 2.1.** The right-hand side of  $(1.7)$  converges absolutely and uniformly on any compact subsets of R.

*Proof.* Let K be a compact subset of R. Then  $|e^{-i\gamma t} - 1|$  is uniformly bounded in  $\gamma$  and  $t \in K$  $t \in K$  $t \in K$ , since all zeros of  $\xi(1/2 - iz)$  lie in the horizontal strip  $|\Im(z)| \leq 1/2$  ([\[9,](#page-4-0) §2.12]). Further,  $\sum_{\gamma} m_{\gamma} |\gamma|^{-\alpha}$  converges for any  $\alpha > 1$ , since  $\xi(1/2 - iz)$  is an entire function of orderone  $([9,$  $([9,$  Theorem 2.12]). Hence the sum on the right-hand side of  $(1.7)$  converges absolutely and uniformly for  $t \in K$ .

From the above lemma, the right-hand side of [\(1.7\)](#page-1-2) defines a continuous function on the real line. By the functional equation

<span id="page-2-4"></span>
$$
\xi(s) = \xi(1-s) \tag{2.1}
$$

 $([9, (2.1.13)])$  $([9, (2.1.13)])$  $([9, (2.1.13)])$ , if  $\gamma$  is a zero of  $\xi(1/2-iz)$ , then  $-\gamma$  is also a zero with the same multiplicity. Therefore, the right-hand side of [\(1.7\)](#page-1-2) is an even real-valued function. This is consistent with the definition of  $g_{\zeta}(t)$ .

Proof of  $(1) \Rightarrow (2)$  in Theorem [1.1](#page-1-0) and Theorem [1.2.](#page-1-3) Since (1) is assumed, all zeros  $\gamma$  of  $\xi(1/2 - iz)$  are real. On the other hand,  $\xi(1/2) \neq 0$ , because  $(1 - 2^{1-\sigma})\zeta(\sigma) > 0$  and  $\Gamma(\sigma/2) > 0$  for any positive real numbers  $\sigma$  ([\[9,](#page-4-0) §2.12]). Therefore, [\(1.6\)](#page-1-1) defines a measure on R such that the formula

$$
\exp(g_{\zeta}(t)) = \exp\left[\int_{-\infty}^{\infty} (e^{it\lambda} - 1)\nu_{\zeta}(d\lambda)\right]
$$

holds on  $\mathbb{R}, \nu_{\zeta}(\{0\}) = 0$  and  $\int_{|\lambda| \leq 1} |\lambda| \nu_{\zeta}(d\lambda) < \infty$  by [\(1.7\)](#page-1-2). Moreover,

$$
\int_{-\infty}^{\infty} \min(1, \lambda^2) \, \nu_{\zeta}(d\lambda) < \infty
$$

by the convergence of  $\sum_{\gamma} m_{\gamma} |\gamma|^{-2}$  in the proof of Lemma [2.1.](#page-2-2) Hence, there exists an infinitely divisible distribution  $\mu$  whose characteristic function is given by  $\exp(g_{\zeta}(t))$ with the characteristic triplet  $(0, 0, \nu_{\zeta})$  by [\[7,](#page-4-1) Theorem 8.1 (iii)].

<span id="page-2-1"></span>To prove  $(2) \Rightarrow (1)$ , we need the following integral formula.

Lemma 2.2. The Fourier integral formula

<span id="page-2-5"></span>
$$
\int_0^\infty g_\zeta(t) e^{izt} dt = \frac{1}{z^2} \frac{\xi'}{\xi} \left(\frac{1}{2} - iz\right)
$$
 (2.2)

holds if  $\Im(z) > 1/2$ .

*Proof.* We start with [\(1.7\)](#page-1-2) again. Since  $\xi(s)$  is order one entire function, Hadamard's factorization theorem gives

$$
\xi\left(\frac{1}{2} - iz\right) = e^{a+bz} \prod_{\gamma} \left[ \left(1 - \frac{z}{\gamma}\right) e^{\frac{z}{\gamma}} \right]^{m_{\gamma}}.
$$

Taking the logarithmic derivative of both sides,

<span id="page-2-3"></span>
$$
\frac{\xi'}{\xi}\left(\frac{1}{2} - iz\right) = ib + i\sum_{\gamma} m_{\gamma}\left(\frac{1}{z - \gamma} + \frac{1}{\gamma}\right),\tag{2.3}
$$

where the sum on the right-hand side converges absolutely and uniformly on every compact subset of  $\mathbb C$  outside the zeros  $\gamma$ . Substituting  $z = 0$  in [\(2.3\)](#page-2-3), we have ib =  $(\xi'/\xi)(1/2)$ . On the other hand, taking the logarithmic derivative of  $(2.1)$ ,

<span id="page-3-1"></span>
$$
\frac{\xi'}{\xi}(s) = -\frac{\xi'}{\xi}(1-s).
$$
\n(2.4)

Substituting  $s = 1/2$  in [\(2.4\)](#page-3-1), we get  $(\xi'/\xi)(1/2) = 0$ , thus  $ib = (\xi'/\xi)(1/2) = 0$ . For each term on the right-hand side of [\(2.3\)](#page-2-3), we have

$$
\frac{i}{z^2} \left( \frac{1}{z - \gamma} + \frac{1}{\gamma} \right) = \int_0^\infty \frac{e^{-it\gamma} - 1}{\gamma^2} e^{izt} dt, \quad \Im(z) > \Im(\gamma)
$$

bydirect calculation of the right-hand side, where  $|\Im(\gamma)| \leq 1/2$  ([\[9,](#page-4-0) Theorem 2.12]). Therefore, [\(2.2\)](#page-2-5) is obtained by interchanging summation and integration, which is jus-tified by Lemma [2.1.](#page-2-2)  $\Box$ 

*Proof of (2)*⇒(1) in Theorem [1.1.](#page-1-0) We have  $\exp(g_{\zeta}(t)) = \hat{\mu}(t)$  for some infinitely divisible distribution  $\mu$  on the real line by (2). Then  $\exp(g_{\zeta}(t)) = \hat{\mu}(t) = \hat{\mu}(-t)$ , since  $g_{\zeta}(t)$  is an even function independent of the RH as mentioned before Theorem [1.1.](#page-1-0) The second equality implies  $\widehat{\mu}(t) = \int_{-\infty}^{\infty} \cos(tx) \mu(dx)$ . Therefore,  $g_{\zeta}(t) (= \log \widehat{\mu}(t))$  is real-valued. Moreover,  $g_{\zeta}(t)$  is non-positive on R, since  $|\hat{\mu}(t)| \leq 1$  holds because  $\mu$  is a distribution.

Since  $g_{\zeta}(t)$  is non-positive, the pure imaginary point iy on the horizontal axis  $\Im(z) = y$ of convergence of  $f(z) = \int_0^\infty g_\zeta(t) e^{izt} dt$  is a singularity of  $f(z)$  by [\[10,](#page-4-8) Theorem 5b in Chap. II]. On the other hand,  $(\xi'/\xi)(1/2 - iz)$  has no singularity on the positive imaginary axis  $i\mathbb{R}_{>0}$ , since  $\xi(s)$  is holomorphic and has no zeros on the positive real line  $\mathbb{R}_{>0}$  ([\[9,](#page-4-0) §2.12]). Hence  $f(z)$  converges in the upper half-plane  $\{z \mid \Im(z) > 0\}$  and defines a holomorphic function there. Therefore,  $\xi(1/2 - iz)$  has no zeros in the upper half-plane by [\(2.2\)](#page-2-5), and then it also has no zeros in the lower half-plane  $\{z \mid \Im(z) < 0\}$ <br>by (2.1). Thus all zeros of  $\xi(1/2 - iz)$  are real, it is the RH. by [\(2.1\)](#page-2-4). Thus all zeros of  $\xi(1/2 - iz)$  are real, it is the RH.

# 3. APPENDIX: PROOF OF  $(1.7)$

<span id="page-3-0"></span>We have shown  $(2.2)$  using the right-hand side of  $(1.7)$ . Therefore, it is sufficient to prove that the right-hand side of  $(1.5)$  satisfies  $(2.2)$  by the uniqueness of the Fourier transform. Taking the logarithmic derivative of [\(1.1\)](#page-0-3) and using the series expansion

$$
\frac{\zeta'(s)}{\zeta(s)} = -\sum_{n=2}^{\infty} \frac{\Lambda(n)}{n^s} \quad (\Re(s) > 1)
$$

 $([9, (1.1.8)]),$  $([9, (1.1.8)]),$  $([9, (1.1.8)]),$  we have

$$
\frac{\xi'}{\xi}(s) = \frac{1}{s-1} + \frac{1}{s} - \frac{1}{2}\log \pi + \frac{1}{2}\psi(s/2) - \sum_{n=2}^{\infty} \frac{\Lambda(n)}{n^s}
$$

for  $\Re(s) > 1$ . Writing  $1/2 - iz$  as s, we get

$$
-\int_0^\infty 4(e^{t/2} + e^{-t/2} - 2) e^{izt} dt = \frac{1}{z^2} \left( \frac{1}{s-1} + \frac{1}{s} \right) \text{ if } \Im(z) > 1/2,
$$

$$
\int_0^\infty t e^{izt} dt = -\frac{1}{z^2} \text{ if } \Im(z) > 0, \text{ and}
$$

$$
\int_{\log n}^\infty \frac{(t - \log n)}{\sqrt{n}} e^{izt} dt = -\frac{1}{z^2} n^{-s} \text{ if } \Im(z) > 0
$$

by direct and simple calculation. The third equality leads to

$$
\int_0^\infty \left( \sum_{n \le e^t} \frac{\Lambda(n)}{\sqrt{n}} (t - \log n) \right) e^{izt} dt = \frac{1}{z^2} \left( -\sum_{n=2}^\infty \frac{\Lambda(n)}{n^s} \right) \quad \text{if } \Im(z) > 1/2,
$$

where the condition  $\Im(z) > 1/2$  is due to the convergence of the series on the right-hand side. Hence, the proof is completed if it is proved that

<span id="page-4-11"></span>
$$
\int_0^\infty \frac{1}{4} \left( e^{-t/2} \Phi(e^{-2t}, 2, 1/4) - \Phi(1, 2, 1/4) \right) e^{izt} dt = \frac{1}{2z^2} \left( \psi(s/2) - \psi(1/4) \right) \tag{3.1}
$$

holds for  $\Im(z) > 0$ . To prove this, we recall the series expansion

<span id="page-4-10"></span>
$$
\psi(w) = -\gamma_0 - \sum_{n=0}^{\infty} \left( \frac{1}{w+n} - \frac{1}{n+1} \right),\tag{3.2}
$$

where $\gamma_0$  is the Euler–Mascheroni constant ([\[4,](#page-4-9) (1.3.16)]). Besides, we have

$$
\int_0^\infty \frac{e^{-2(n+\frac{1}{4})t} - 1}{2(n+\frac{1}{4})^2} e^{izt} dt = -\frac{1}{z^2} \left( \frac{1}{s/2 + n} - \frac{1}{n+1/4} \right)
$$

for non-negative integers n and  $\Im(z) > 0$  by direct and simple calculation. Therefore, equality [\(3.2\)](#page-4-10) implies

$$
\int_0^\infty \frac{1}{4} \left( e^{-t/2} \Phi(e^{-2t}, 2, 1/4) - \Phi(1, 2, 1/4) \right) e^{izt} dt
$$
  
=  $\frac{1}{2} \int_0^\infty \left( \sum_{n=0}^\infty \frac{e^{-2(n+\frac{1}{4})t} - 1}{2(n+1/4)^2} \right) e^{izt} dt = -\frac{1}{2z^2} \sum_{n=0}^\infty \left( \frac{1}{s/2 + n} - \frac{1}{n+1/4} \right)$   
=  $-\frac{1}{2z^2} \sum_{n=0}^\infty \left( \frac{1}{s/2 + n} - \frac{1}{n+1} \right) - \frac{1}{2z^2} \sum_{n=0}^\infty \left( \frac{1}{n+1} - \frac{1}{n+1/4} \right)$   
=  $\frac{1}{2z^2} (\psi(s/2) + \gamma_0) - \frac{1}{2z^2} (\psi(1/4) + \gamma_0),$ 

where  $s = 1/2 - iz$ . Hence, we obtain [\(3.1\)](#page-4-11).

Acknowledgments The first and second authors were supported by JSPS KAKENHI Grant Number JP22K03276 and JP17K05163–JP23K03050, respectively. This work was also supported by the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University.

### **REFERENCES**

- <span id="page-4-5"></span>[1] P. Biane, J. Pitman, M. Yor, Probability laws related to the Jacobi theta and Riemann zeta functions, and Brownian excursions, *Bull. Amer. Math. Soc.* (N.S.) 38 (2001), no. 4, 435-465.
- <span id="page-4-3"></span>[2] B. V. Gnedenko, A. N. Kolmogorov, Limit Distributions for Sums of Independent Random Variables (Translated from the Russian by Kai Lai Chung), Addison-Wesley, 1968.
- <span id="page-4-6"></span>[3] J. C. Lagarias, E. Rains, On a two-variable zeta function for number fields, Ann. Inst. Fourier (Grenoble) 53 (2003), no. 1, 1–68.
- <span id="page-4-9"></span>[4] N. N. Lebedev, Special functions and their applications, Revised edition, Dover Publications, Inc., New York, 1972.
- <span id="page-4-7"></span><span id="page-4-4"></span>[5] G. D. Lin, C-Y, Hu, The Riemann zeta distribution, Bernoulli 7 (2001), no. 5, 817–828.
- [6] T. Nakamura, A complete Riemann zeta distribution and the Riemann hypothesis, Bernoulli 21 (2015), no. 1, 604–617.
- <span id="page-4-1"></span>[7] K. Sato, L´evy processes and infinitely divisible distributions, Translated from the 1990 Japanese original, Revised by the author, Cambridge Studies in Advanced Mathematics, 68, Cambridge University Press, Cambridge, 1999.
- <span id="page-4-2"></span>[8] M. Suzuki, Aspects of the screw function corresponding to the Riemann zeta function, to appear in the Journal of the London Mathematical Society.
- <span id="page-4-0"></span>[9] E. C. Titchmarsh, The theory of the Riemann zeta-function, Second edition, Edited and with a preface by D. R. Heath-Brown , The Clarendon Press, Oxford University Press, New York, 1986.
- <span id="page-4-8"></span>[10] V. D. Widder, The Laplace Transform, Princeton Mathematical Series, vol. 6, Princeton University Press, Princeton, N. J., 1941.