RAMSEY NUMBERS OF CONNECTED 4-CLIQUE MATCHING

KRIT KANOPTHAMAKUN AND PANUPONG VICHITKUNAKORN

ABSTRACT. We determine the exact value of the 2-color Ramsey number of a connected 4-clique matching $\mathscr{C}(nK_4)$ which is a set of connected graphs containing n disjoint K_4 . That is, we show that $R_2(\mathscr{C}(nK_4)) = 13n - 3$ for any positive integer $n \geq 3$. The result is an extension of the result by (Roberts, 2017) which gave that result when $n \geq 18$. We also show that the result still holds when n = 2 provided that $R_2(2K_4) \leq 23$.

1. INTRODUCTION AND PRELIMINARIES

The literature on Ramsey-type problems begins in 1930 with Ramsey's original paper [12]. Ramsey's famous theorem makes an analogous assertion regarding the partitioning of r-element subsets of objects into classes. A thorough study of Ramsey's theorem and other partitioning theorems later appears in Graham-Rothschild-Spencer [8].

Definition 1 ([15]). Let $\binom{s}{r}$ denote the set of *r*-element subsets (*r*-sets) of a set S. A set $T \subseteq S$ is homogeneous under a coloring of $\binom{s}{r}$ if all *r*-sets in T receive the same color; it is *i*-homogeneous if that color is *i*.

Let r and p_1, p_2, \ldots, p_k be positive integers. If there is an integer N such that every k-coloring of $\binom{[N]}{r}$ yields an *i*-homogeneous set of size p_i for some *i*, then the smallest integer is the Ramsey number $R(p_1, \ldots, p_k; r)$.

Theorem 2 ([15]). Given a positive integers r and p_1, p_2, \ldots, p_k , there exists an integer N such that every k-coloring of $\binom{[N]}{r}$ yields an i-homogeneous set of size p_i for some i, where $\binom{[N]}{r}$ denote the set of r-subsets of a set [N].

We consider the case r = 2, which is an edge-coloring of graphs. We abbreviate the notation $R(p_1, p_2, \ldots, p_k; r)$ to $R(p_1, p_2, \ldots, p_k)$. Until now, no exact formula is known for general values of p_i and r and few Ramsey numbers have been computed, even for k = 2: R(3,3) = 6, R(3,4) = 9, R(3,5) = 14, R(3,6) = 18, R(3,7) = 23, R(3,8) = 28, R(3,9) = 36, R(4,4) = 18 and R(4,5) = 25. See [11] and references therein.

For k = 3, only two Ramsey numbers are known, which are R(3, 3, 3) = 17 [6] and R(3, 3, 4) = 30 [5]. For k > 3, the exact values of Ramsey numbers are still unknown, see the survey [11] for other bounds on Ramsey numbers R(k, l).

Ramsey's Theorem for r = 2 says that k-edge-coloring of a large enough complete graph G always induces a monochromatic complete subgraph of a given size. If we want a monochromatic copy of non-complete subgraph, it may require a smaller graph G. So we can define the Ramsey number of any graph. In this paper, we follow the notations in [15].

Definition 3. Given simple graphs G_1, G_2, \ldots, G_k , the *(graph) Ramsey number* $R(G_1, \ldots, G_k)$ is the smallest integer n such that every k-coloring of $E(K_n)$ contains a copy of G_i in color i for some i. When $G_i = G$ for all i, we write $R_k(G)$ for $R(G_1, \ldots, G_k)$.

There are a lot of research about Ramsey number for non-complete graphs (see the survey [11]) such as complete bipartite graphs, cycles, paths, stars, etc.

For disconnected graphs, there are only few results. For example, [4] found the exact formula of $R(n_1P_2, n_2P_2, \ldots, n_kP_2)$, and later [9] extended it to a more general case of $R(K_m, n_1P_2, n_2P_2, \ldots, n_kP_2)$. Later, there are more results on general cases of disconnected graphs, such as forest [14] and multiple copies of complete graphs [10, 3, 4].

Other kinds of Ramsey number have been studied as well. For Ramsey numbers $R_2(nK_r)$, the exact formula for any $n \ge 2$ is only known when r is either 2 [4, 9] or 3 [3]. For r = 4, [2] gave the formula when n is large.

Let G be a disconnected graph and $\mathscr{C}(G)$ be the set of all connected graphs containing G. Then $R_k(\mathscr{C}(G))$ is the least integer n such that any k-edge-coloring on K_n contains a monochromatic copy of a graph in $\mathscr{C}(G)$, see [7, 13].

For r = 3, we call a graph in $\mathscr{C}(nK_r)$ a connected triangle matching. $R_2(\mathscr{C}(nK_3)) = 7n-2$ is proved by [7]. For $r \geq 4$, the only result for $R_2(\mathscr{C}(nK_r))$ is the following.

Theorem 4. [13] For $r \ge 4$ and $n \ge R_2(K_r)$ we have

$$R_2(\mathscr{C}(nK_r)) = (r^2 - r + 1)n - r + 1.$$

In this paper, we are only interested in the case when r = 4. Theorem 4 showed that $R_2(\mathscr{C}(nK_4)) = 13n - 3$ when $n \ge R_2(K_4) = 18$. The main goal of this paper is to extend Theorem 4 from $n \ge 18$ to $n \ge 3$.

The paper is structured as follows. In Section 2, we prove Lemma 5, which is the main lemma of the paper. We then construct the proof of the main result, Theorem 7, and divide it into four parts in Sections 3-6. We end the section with another three lemmas which will be used in later sections. In Section 8, we discuss the results and further research.

2. Main results

The following lemma is the main key to proving our main theorem.

Lemma 5. For any positive integers x and y where $y \ge 4$. If a two-edge-coloring of K_{4x+y} with red and blue contains at most x blue copies of K_4 , then there is a

red matching of size r, where

$$r = \begin{cases} y - 3 & \text{for } x \ge y - 4, \\ \lfloor \frac{x + y - 3}{2} \rfloor & \text{for } x \le y - 5. \end{cases}$$

Proof. Let V be the set of vertices of K_{4x+y} . For $x \ge y - 4$, suppose there is no red matching of size y - 3. Let $M = \{u_1v_1, u_2v_2, \ldots, u_mv_m\}$ be a maximum red matching of size $m \le y-4$. Let $P = \{u_1, u_2, \ldots, u_m, v_1, v_2, \ldots, v_m\}$ and $P' = V \setminus P$. We see that all edges between vertices in P' are blue, otherwise, there will be a red matching of size larger than m.

Now, for any $1 \leq i \leq m$, in order to prevent a larger red matching there are no two vertices $w_1, w_2 \in P'$ such that both $u_i w_1$ and $v_i w_2$ are both red. From this observation, there are only two cases:

- (1) If both u_i and v_i are adjacent to some vertices in P' via red edges, then all red edges incident to u_i or v_i must have a common endpoint $w \in P'$. Hence, $u_i w'$ and $v_i w'$ are blue for all $w' \in P' \setminus \{w\}$.
- (2) Either u_i or v_i is adjacent to every vertex in P' via a blue edge. We assume without loss of generality that it is v_i .

We also assume that $\{u_i, v_i\}$ is in the first case when $1 \le i \le k$ and is in the second case when $k + 1 \le i \le m$. Let $W = \{w_1, w_2, \ldots, w_k\}$. We note that w_i 's are not necessarily distinct.

Next we construct a blue copy of mK_4 . For $1 \le i \le k$, a blue copy of K_4 is obtained from v_i , one vertex in W which differs from w_i and two other vertices in $P' \setminus W$. If we cannot choose any vertex from W, we choose any three vertices from W' instead. For $k + 1 \le i \le m$, a blue copy of K_4 is obtained from v_i and three vertices in P'. The construction is feasible, because $|P'| = 4x + y - 2m \ge 4x - y + 8 \ge 3x + 4 > 3x + 1 \ge 3m + 1$.

The number of vertices in P' which are not in those m disjoint blue copies of K_4 is $4x+y-2m-3m = 4x+y-5m \ge 4x+y-5(y-4) \ge 4x-4y+20 = 4(x-y+5)$. These vertices can form other $x - y + 5 \ge x - m + 1$ disjoint blue copies of K_4 . Thus, there are at least x + 1 disjoint blue copies of K_4 , which contradicts to the assumption.

In the case $x \leq y-5$, we follow exactly the same construction. We have $m \leq \lfloor \frac{x+y-3}{2} \rfloor - 1 \leq \frac{x+y-5}{2}$. If x < m, we can construct a blue copy of $(x+1)K_4$ where each K_4 contains one vertex from $\{v_1, v_2, \ldots, v_m\}$ and three vertices from P'. Since $|P'| \geq 3x + 5 > 3(x+1) + 1$, we have a blue copy of $(x+1)K_4$, which is a contradiction. Hence, we may assume $x \geq m$. The construction of the first blue copy of mK_4 is feasible because $|P'| = 4x + y - 2m \geq 4x + y - x - y + 5 = 3x + 5 > 3x + 1 \geq 3m + 1$. The construction of the other x - m + 1 is feasible if $4x+y-2m-3m \geq 4(x-m+1)$, which is equivalent to $y \geq m+4$. From $m \leq \frac{x+y-5}{2}$ and $x \leq y - 5$, we have $y \geq m + 5$. Hence, the construction is feasible.

The lower bound of $R_2(\mathscr{C}(nK_4))$ in Theorem 7 is a consequence of Theorem 1 of [1]. In fact, the result can be directly proved as shown in the following lemma.

Lemma 6. For any integer $n \ge 2$ and $r \ge 3$, $R_2(\mathscr{C}(nK_r)) \ge (r^2 - r + 1)n - r + 1$.

Proof. Let $N = (r^2 - r + 1)n - r$. It can be rewritten as N = (r-1)(rn-1) + (n-1). We can construct a 2-edge-coloring of $G = K_N$ using red and blue without a monochromatic copy of a graph in $\mathscr{C}(nK_r)$ as follows. We form a copy of K_N with a 2-edge-coloring by joining a blue copy of $(r-1)K_{rn-1}$ and a blue copy of K_{n-1} using only red edges. The size of each blue component is at most rn - 1, hence there are at most n - 1 disjoint blue copies of K_r . Thus, this coloring on Gcontains no blue copy of a graph in $\mathscr{C}(nK_r)$.

Any red copy K_r in G must be formed using exactly one vertex from each of the blue components. But, the smallest component, K_{n-1} has only n-1 vertices. Hence, there are at most n-1 disjoint red copies of K_r in G. Thus, there is no red copy of a graph in $\mathscr{C}(nK_r)$.

We prove the upper bound of $R_2(\mathscr{C}(nK_r))$ and obtain the exact Ramsey number in the following theorem.

Theorem 7. For any positive integer $n \ge 3$, $R_2(\mathscr{C}(nK_4)) = 13n - 3$.

Proof. We known that $R_2(\mathscr{C}(nK_4)) \geq 13n - 3$. For the upper bound, we consider a 2-edge-coloring of $G = K_{13n-3}$ with red and blue. We will show that there is a monochromatic subgraph in $\mathscr{C}(nK_4)$, which we will simply call it a *connected* monochromatic copy of nK_4 or n connected disjoint monochromatic copies of K_4 . Since a graph and its complement cannot be both disconnected, at least one color is connected, we suppose that red is connected. If G contains a red copy nK_4 or blue connected copy of nK_4 , then we are done. So, we may assume that there are at most n-1 disjoint red copies of K_4 and no blue connected copy of nK_4 . Since 13n-3 = 18 + 4(3n-5) + n - 1 and R(4,4) = 18, G has at least $3n - 4 + \lfloor \frac{n-1}{4} \rfloor$ disjoint monochromatic copies of K_4 . Since there are at most n-1 disjoint red copies of K_4 , there are at least $3n - 4 + \lfloor \frac{n-1}{4} \rfloor - n + 1 = 2n - 3 + \lfloor \frac{n-1}{4} \rfloor$ disjoint blue copies of K_4 .

Let k be the maximum number of disjoint blue copies of K_4 , and K be the set of such k disjoint blue copies of K_4 . We partition the set of vertices V(G) of G into V_1, V_2, \ldots, V_t and V' in the way that $G[V_i]$ is a blue component of G containing at least one blue copy of K_4 in K and V' is the set of vertices outside $V_1 \cup \cdots \cup V_t$, see Figure 1 for an example. In other words, $V' = V(G) \setminus (V_1 \cup \cdots \cup V_t)$. Let a_i be the number of disjoint blue copies of K_4 from K containing in $G[V_i]$. Since there is no connected disjoint blue copies of K_4 , we have $a_i \leq n-1$. Hence, $t \geq 2$ as $a_1 + \cdots + a_t = k \geq 2n - 3 + \lfloor \frac{n-1}{4} \rfloor > n-1$.

The aim of this proof is to construct a red copy of nK_4 by partitioning G into multiple components where each component consists of one or more blue components $G[V_i]$'s. All edges between any two components will be red. A red



FIGURE 1. An example of a graph G together with V_1 , V_2 and V' where all red edges are omitted.

copy of K_4 will be constructed by choosing a single vertex, a red edge, or a red triangle from different components and combining them into a red copy of K_4 . The key is to show that there are enough vertices in each component to do the construction.

The proof is divided into four cases, Sections 3–6, according to the value of k. In each case, we get a red copy of nK_4 in G, which contradicts to our assumption. Hence, for any 2-edge-coloring of $G = K_{13n-3}$, there exists a monochromatic subgraph in $\mathscr{C}(nK_4)$. Thus, $R_2(\mathscr{C}(nK_4)) \leq 13n - 3$ for $n \geq 3$.

The following three lemmas will be used in Sections 3-6.

Lemma 8. Let $a_1 + \cdots + a_t \ge n - 1 + m$ where $1 \le a_i \le n - 1$ and $m \le \frac{n}{2}$. Then we can partition a_1, \ldots, a_t into two parts where the sum of each part is at least m.

Proof. We can assume without loss of generality that $n - 1 \ge a_1 \ge a_2 \ge \cdots \ge a_t \ge 1$. Since each a_i is at most n - 1, we have $t \ge 2$.

If $a_1 \ge a_2 \ge m$, then $a_1 \ge m$ and $a_2 + \cdots + a_t \ge m$.

If $a_1 \ge m > a_2$, then $a_2 + a_3 + \dots + a_t \ge n - 1 + m - a_1 \ge m$.

If $m > a_1 \ge a_2$, we can partition a_1, \ldots, a_t into two parts whose difference is at most the largest number $a_1 < m$. (We can keep moving the largest number in the larger part to the smaller part until the desired difference is archived.) Let the sum of the smaller part be x. Then the sum of the larger part is at most x+m-1. Since the total sum is at least n-1+m, we have $x + x + m - 1 \ge n - 1 + m$, which simplifies to $x \ge \frac{n}{2} \ge m$.

Hence, we always have two parts whose sum is at least m.

Lemma 9. Let $a_1 + \cdots + a_t \ge 2n - 1 + \lfloor \frac{n}{4} \rfloor$ where $1 \le a_i \le n - 1$. Then we can partition a_1, \ldots, a_t into three parts where the sum of each part is at least $\lceil \frac{n}{4} \rceil$.

Proof. We can assume without loss of generality that $n - 1 \ge a_1 \ge a_2 \ge \cdots \ge a_t \ge 1$. Since each a_i is at most n - 1, we have $t \ge 3$. So, the claim is trivial for

n = 2, 3, 4 because $\left\lceil \frac{n}{4} \right\rceil = 1$. Now, we suppose that $n \ge 5$. If $a_3 \ge \left\lceil \frac{n}{4} \right\rceil$, then we are done.

If $a_2 \ge \left\lceil \frac{n}{4} \right\rceil$ and $a_3 < \left\lceil \frac{n}{4} \right\rceil$, then $a_3 + a_4 + \dots + a_t \ge \left\lfloor \frac{n}{4} \right\rfloor + 1 \ge \left\lceil \frac{n}{4} \right\rceil$. Thus, we can combine a_3, a_4, \dots, a_t into one part, and we are done.

If $a_1 \ge \lfloor \frac{n}{4} \rfloor$ and $a_2 < \lfloor \frac{n}{4} \rfloor$, then $a_2 + a_3 + \cdots + a_t \ge n + \lfloor \frac{n}{4} \rfloor$. From Lemma 8, we can partition a_2, \ldots, a_t into two parts where the sum of each part is at least $\lfloor \frac{n}{4} \rfloor$. Thus, the sum of the smallest part is at least $\lfloor \frac{n}{4} \rfloor$. Therefore, we have three parts whose sum are all at least $\lfloor \frac{n}{4} \rfloor$.

The last case is when $a_1 < \lfloor \frac{n}{4} \rfloor$. Similar to the previous case, we can combine a_1, \ldots, a_t into three parts, where the pairwise difference between the sum of each part is at most the largest number $a_1 \leq \lfloor \frac{n}{4} \rfloor$. Let x be the sum of the smallest part. Then the other parts have their sum at most $x + \lfloor \frac{n}{4} \rfloor - 1$. Then $x + 2(x + \lfloor \frac{n}{4} \rfloor - 1) \geq 2n - 1 + \lfloor \frac{n}{4} \rfloor$, which simplifies to $x \geq \frac{2n - \lfloor n/4 \rfloor}{3} \geq \lfloor \frac{n}{4} \rfloor$. The last inequality comes from the fact that $2n > 4 \lfloor \frac{n}{4} \rfloor$ when $n \geq 5$. Hence, the sum of each part is at least $\lfloor \frac{n}{4} \rfloor$.

Lemma 10. Let $a_1 + \cdots + a_t \ge 3n - 3 + \lfloor \frac{n}{4} \rfloor$ where $1 \le a_i \le n - 1$. Then we can partition a_1, \ldots, a_t into four parts where the sum of each part is at least $\lfloor \frac{n}{4} \rfloor$.

Proof. Let $k = a_1 + a_2 + \cdots + a_t$, where $n - 1 \ge a_1 \ge a_2 \ge \cdots \ge a_t \ge 1$. Since each part is at most n - 1 and k > 3n - 3, we get $t \ge 4$. The claim is obvious when $2 \le n \le 4$.

Let $n \ge 5$. We use the same method as in the proof Lemma 9. Consider the following cases.

- $a_1 \ge a_2 \ge a_3 \ge a_4 \ge \left\lceil \frac{n}{4} \right\rceil$. Then $a_4 + \dots + a_t \ge \left\lceil \frac{n}{4} \right\rceil$.
- $a_1 \ge a_2 \ge a_3 \ge \lceil \frac{n}{4} \rceil > a_4$. Then $a_4 + \dots + a_t \ge 3n 3 + \lceil \frac{n}{4} \rceil (3n 3) \ge \lceil \frac{n}{4} \rceil$.
- $a_1 \ge a_2 \ge \left\lceil \frac{n}{4} \right\rceil > a_3 \ge a_4$. Then $a_3 + \dots + a_t \ge n 1 + \left\lceil \frac{n}{4} \right\rceil$. From Lemma 8, we can combine a_3, \dots, a_t into two parts where each part is at least $\left\lceil \frac{n}{4} \right\rceil$.
- $a_1 \ge \left\lceil \frac{n}{4} \right\rceil > a_2 \ge a_3 \ge a_4$. Then $a_2 + \dots + a_t \ge 2n 2 + \left\lceil \frac{n}{4} \right\rceil$. From Lemma 9, we can combine a_2, \dots, a_t into three parts where each part is at least $\left\lceil \frac{n}{4} \right\rceil$.
- $\left\lceil \frac{n}{4} \right\rceil > a_1 \ge a_2 \ge a_3 \ge a_4$. We can combine a_1, \ldots, a_t into four parts where each part differs from another by at most the size of the largest part $a_1 < \left\lceil \frac{n}{4} \right\rceil$. Let x be the sum of the smallest part. Then the sum of largest part is at most $x + \left\lceil \frac{n}{4} \right\rceil 1$. Consider the sum of the four parts, we have

$$x + 3\left(x + \left\lceil \frac{n}{4} \right\rceil - 1\right) \ge 3n - 3 + \left\lceil \frac{n}{4} \right\rceil.$$

This implies that

$$x \ge \frac{3n - 2\left\lceil \frac{n}{4} \right\rceil}{4} \ge \left\lceil \frac{n}{4} \right\rceil.$$

From all cases, we get four parts where each part is at least $\left|\frac{n}{4}\right|$.

3. Proof of the first case

Let $2n - 3 + \lfloor \frac{n-1}{4} \rfloor \leq k \leq 2n - 2 + \lfloor \frac{n}{4} \rfloor$. We combine V_1, V_2, \ldots, V_t and V' into two disjoint sets, say U_1 and U_2 . So, each U_i is a disjoint union of some sets from V_1, V_2, \ldots, V_t and V'. Let x_1 and x_2 be the number of disjoint blue copies of K_4 containing in $G[U_1]$ and $G[U_2]$, respectively. Since the number of disjoint blue copies of K_4 in $G[V_i]$ is at most n - 1, we can combine these V_1, V_2, \ldots, V_t and V' into U_1 and U_2 so that the difference between x_1 and x_2 is at most n - 1. We note that all edges between U_1 and U_2 are red.

Let Y_1 and Y_2 be the set of vertices in U_1 and U_2 that are not used in any K_4 from K, respectively. We also let $y_1 = |Y_1|$ and $y_2 = |Y_2|$. Since $x_1 + x_2 = k$ and $4x_1 + 4x_2 + y_1 + y_2 = 13n - 3$. Hence,

$$y_1 + y_2 = 13n - 3 - 4k \ge 5n + 5 - 4\left\lfloor\frac{n}{4}\right\rfloor.$$
 (1)

Since the difference between x_1 and x_2 is at most n-1, we get

$$2x_i \ge k - (n-1) \ge 2n - 3 + \left\lfloor \frac{n-1}{4} \right\rfloor - (n-1) = n - 2 + \left\lfloor \frac{n-1}{4} \right\rfloor.$$

We can conclude that

$$x_i \ge \frac{n}{2} - 1 + \frac{1}{2} \left\lfloor \frac{n-1}{4} \right\rfloor.$$
 (2)

This also gives

$$4x_i \ge n. \tag{3}$$

From Eq. (1), y_1 and y_2 cannot be both less than four.

Case 1: Exactly one of y_1 and y_2 is less than four.

We suppose that $y_1 < 4$. Then $y_2 \ge 5n+2-4\lfloor \frac{n}{4} \rfloor \ge 3n+6 = 9+3(n-1)$. Since R(3,4) = 9 and $G[Y_2]$ contains no blue copy of K_4 , there is a red copy of nK_3 in $G[Y_2]$. Since $|U_1| \ge 4x_1 \ge n$ from Eq. (3), we can then construct a red copy of nK_4 using n vertices in U_1 and a red copy of nK_3 in $G[Y_2]$.

Case 2: $y_1, y_2 \ge 4$.

We assume $y_1 \ge y_2$. Eq. (1) gives $y_1 + y_2 \ge 5n + 5 - 4\lfloor \frac{n}{4} \rfloor \ge 4n + 5$. Hence $y_1 \ge 2n+3$. By Eq. (2), we have $x_1 = k - x_2 \le 2n - 2 + \lfloor \frac{n}{4} \rfloor - \frac{n-1}{2} - \frac{1}{2} \lfloor \frac{n-1}{4} \rfloor \le 2n - 2 \le y_1 - 5$. By Lemma 5, $G[U_1]$ contains a red matching of size $r_1 = \lfloor \frac{x_1 + y_1 - 3}{2} \rfloor \ge \lfloor \frac{x_1 + 2n}{2} \rfloor \ge n$. If $G[U_2]$ contains a red matching of size at least n, we can create a red copy of nK_4 using n red edges in $G[U_1]$ and n red edges in $G[U_2]$. So we assume that $G[U_2]$ has no red matching of size n.

Case 2.1: $x_2 \ge y_2 - 4$.

By Lemma 5, $G[U_2]$ contains a red matching of size $y_2 - 3$, which must be less than n. We then have $4 \le y_2 < n + 3$. First, we can construct $y_2 - 3$ disjoint red copies of K_4 from one red matching of size $y_2 - 3$ from $G[U_1]$ and another from $G[U_2]$. Next, we need to construct $n - y_2 + 3$ additional disjoint red copies of K_4 . In Y_1 , apart from vertices in $y_2 - 3$ red copies of K_4 , there are $y_1 - 2(y_2 - 3)$ remaining vertices. From Eq. (1), we have

$$y_1 + y_2 \ge 5n + 5 - 4\left\lfloor \frac{n}{4} \right\rfloor \ge 3n + 9.$$

Hence,

$$y_1 - 2y_2 + 6 \ge 3n - 3y_2 + 15 \ge 9 + 3(n - y_2 + 2).$$

Therefore, there are $n - y_2 + 3$ disjoint red copies of K_3 available in $G[U_1]$. Lastly, we need to show that there are at least $n - y_2 + 3$ remaining vertices in U_2 apart from vertices in $y_2 - 3$ disjoint red copies of K_4 . From Eq. (3), we have

$$4x_2 + y_2 - 2(y_2 - 3) \ge n - y_2 + 3.$$

Hence, there is a red copy of nK_4 in G. Case 2.2: $x_2 \le y_2 - 5$.

Again, by Lemma 5 we get a red matching of size $r_2 = \left\lfloor \frac{x_2 + y_2 - 3}{2} \right\rfloor$, which must be less than n. Again, we can construct a red copy of r_2K_4 by connecting two red matchings of size r_2 from $G[U_1]$ and $G[U_2]$. So, we need additional $n - r_2$ disjoint red copies of K_4 in G. From $r_2 = \lfloor \frac{x_2 + y_2 - 3}{2} \rfloor < n$, we get $x_2 + y_2 \le 2n + 2$. Since $2x_2 \le 2x_2 \le 2n + 2$.

 $x_2 + y_2 - 5 \leq 2n - 3$, we can conclude that $x_2 \leq n - 2$. So,

$$x_1 = k - x_2 \ge 2n - 3 + \left\lfloor \frac{n-1}{4} \right\rfloor - (n-2) \ge n - 1 + \left\lfloor \frac{n-1}{4} \right\rfloor.$$

• $x_1 = n - 1 + \left\lfloor \frac{n-1}{4} \right\rfloor$
Then $x_2 = n - 2$. Hence,

 $n+3 = x_2 + 5 \le y_2 \le 2n+2 - x_2 \le n+4.$

So y_2 can be either n + 3 or n + 4. Since $4x_1 + 4x_2 + y_1 + y_2 =$ 13n-3, we then have

$$y_1 \ge (13n-3) - 4(n-1) - 4\left\lfloor \frac{n-1}{4} \right\rfloor - 4(n-2) - (n+4) \ge 3n+6.$$

Hence,

$$y_1 - 2r_2 \ge 3n + 6 - 3r_2 = 9 + 3(n - r_2 - 1).$$

Thus, there are $n - r_2$ disjoint red copies of K_3 using vertices from U_1 apart from vertices in the r_2 disjoint red copies of K_4 . • $x_1 \ge n + \left\lfloor \frac{n-1}{4} \right\rfloor$.

Recall that we have constructed a red copy of $r_2 K_4$ using $r_2 < n$ red matching in $G[U_1]$. We would like to show further that when $x_1 \ge n + \left| \frac{n-1}{4} \right|$ those r_2 red matching can be constructed without using any vertices in Y_1 . Since each blue component of $G[U_1]$ contains at most n-1 disjoint blue copies of K_4 , we can partition the blue components in $G[U_1]$ into two parts where the smaller part has at least $\lfloor \frac{n-1}{4} \rfloor + 1$ disjoint blue copies of K_4 . We know that edges between these two parts are red and the number of vertices in each part is at least $4(\lfloor \frac{n-1}{4} \rfloor + 1) \ge n$. Hence, we can construct a red matching of size $r_2 < n$ in $G[U_1]$ using no vertices from Y_1 .

We now construct $n-r_2$ disjoint red copies of K_3 in $G[Y_1]$. Since there is no blue copy of K_4 in $G[Y_1]$ and R(3, 4) = 9, we need to show that $y_1 \ge 9 + 3(n - r_2 - 1)$. Consider

$$2y_1 + 3x_2 + 3y_2 - 12 = 3x_2 + 2(y_1 + y_2) + y_2 - 12.$$

From Eq. (1) and $x_2 \leq y_2 - 5$, we have

 $2y_1 + 3x_2 + 3y_2 - 12 \ge 3x_2 + 2(4n+5) + (x_2+5) - 12 = 4x_2 + 8n + 3.$

Since $n \ge 3$ and $x_2 \ge 1$, we then get

$$2y_1 + 3x_2 + 3y_2 - 12 \ge 4x_2 + 8n + 3 \ge 6n + 13 > 6n + 12$$

We can conclude that

$$y_1 + 3r_2 \ge y_1 + 3\left(\frac{x_2 + y_2 - 4}{2}\right) > 3n + 6.$$

Thus, $y_1 > 9 + 3(n - r_2 - 1)$. So there are $n - r_2$ disjoint red copies of K_3 using vertices from U_1 .

From both cases, there are additional $n - r_2$ disjoint red copies of K_3 in $G[U_1]$. The last step is to show that there are also additional $n - r_2$ vertices in U_2 , which is $4x_2 + y_2 - 2r_2 \ge n - r_2$. From $y_2 - 5 \ge x_2$ and Eq. (3), we have

$$4x_2 + y_2 - \frac{x_2 + y_2 - 3}{2} = \frac{7x_2 + y_2 + 3}{2} \ge 4x_2 + 4 \ge n.$$

Since $r_2 = \lfloor \frac{x_2 + y_2 - 3}{2} \rfloor \le \frac{x_2 + y_2 - 3}{2}$, we conclude that
$$4x_2 + y_2 - 2r_2 \ge 4x_2 + y_2 - \frac{x_2 + y_2 - 3}{2} - r_2 \ge n - r_2.$$

This concludes that there is a red copy of nK_4 in G. This finished the proof of the first case.

4. Proof of the second case

Let $2n - 1 + \lfloor \frac{n}{4} \rfloor \leq k \leq 3n - 3$. From Lemma 9, we can combine V_1, V_2, \ldots, V_t and V' into three partitions, say U_1, U_2 and U_3 , where each $G[U_i]$ contains at least $\lfloor \frac{n}{4} \rfloor$ disjoint blue copies of K_4 . In addition, we can partition them so that only one part contains more than n - 1 disjoint blue copies of K_4 . (If the first and the second parts contain more than n-1 disjoint blue copies of K_4 , we can keep repartitioning the second part into two subparts and add the smaller part into the first part until the second part is at most n-1. Note that the larger part is always at least $\frac{n}{2} \ge \left\lceil \frac{n}{4} \right\rceil$.)

For $i \in \{1, 2, 3\}$, we let x_i be the number of disjoint blue copies of K_4 in $G[U_i]$ from K, y_i be the number of vertices in $G[U_i]$ not belong to those disjoint blue copies of K_4 from K. The number of vertices not in any K_4 in K is |V(G-K)| = 13n - 3 - 4k. Since $2n - 1 + \lfloor \frac{n}{4} \rfloor \le k \le 3n - 3$, we have

$$n+9 \le y_1 + y_2 + y_3 \le 5n+10 - 4\left\lfloor \frac{n}{4} \right\rfloor.$$
 (4)

The main goal is to construct a red copy of nK_4 . Since we can divide the graph G into three subgraphs, $G[U_1], G[U_2]$ and $G[U_3]$ where edges between these parts are red, we can construct a red copy of K_4 by choose two vertices whose adjacent by a red edge from one subgraph and choose one vertex each from the other two subgraphs.

Let r_i be the size of a maximum matching in $G[U_i]$. The next two lemmas show that we have a red matching of size $s_1 + s_2 + s_3 = n$ in G, which will be used to construct a red copy of nK_4 . The proofs are postponed to the end of the section.

Lemma 11. $r_1 + r_2 + r_3 \ge n$.

Lemma 12. There exists $s_i \leq r_i$ for $i \in \{1, 2, 3\}$ such that $s_1 + s_2 + s_3 = n$ and $|U_i| \geq s_i + n$ for all i.

Let M be a red matching obtained by choosing s_i edges from a matching of size r_i in $G[U_i]$. We write

$$M = \{u_{11}v_{11}, \dots, u_{1s_1}v_{1s_1}, u_{21}v_{21}, \dots, u_{2s_2}v_{2s_2}, u_{31}v_{31}, \dots, u_{3s_3}v_{3s_3}\}$$

where $u_{ij}, v_{ij} \in U_i$ for any j. From Lemma 12, we can choose distinct vertices w_{ij} in $U_i \setminus \{u_{i1}, v_{i1}, \ldots, u_{is_i}, v_{is_i}\}$ for $1 \leq j \leq n - s_i$. Next, for $1 \leq j \leq s_1$, the four vertices u_{1j}, v_{1j}, w_{2j} and w_{3j} induces s_1 disjoint red copies of K_4 . Similarly, for $1 \leq j \leq s_2$, the four vertices u_{2j}, v_{2j}, w_{1j} and $w_{3(s_1+j)}$ also induces s_2 disjoint red copies of K_4 . Lastly, for $1 \leq j \leq s_3$, the four vertices $u_{3j}, v_{3j}, w_{1(s_2+j)}$ and $w_{2(s_1+j)}$ induces another s_3 disjoint red copies of K_4 . By Lemma 12, the construction is feasible. Since $s_1 + s_2 + s_3 = n$, there is a red copy of nK_4 in G.

Proof of Lemma 11. If there is *i* where $x_i \ge n - 1 + \lfloor \frac{n}{4} \rfloor$, then $x_i > n - 1$. The graph $G[U_i]$ can then be separated into two parts where all edges between the two parts are red and the number of disjoint blue copies of K_4 from K in each part is at least $\lfloor \frac{n}{4} \rfloor$. So the number of vertices in each part is at least $4 \lfloor \frac{n}{4} \rfloor$, hence $r_i \ge 4 \lfloor \frac{n}{4} \rfloor \ge n$. In the rest of the proof, we can then assume that $x_i < n - 1 + \lfloor \frac{n}{4} \rfloor$.

Consider the lower bound of $x_1 + x_2 + x_3 + y_1 + y_2 + y_3$. The lowest possible value occurs when the number of vertices in K is the largest as possible, i.e.,

 $x_1 + x_2 + x_3 = 3n - 3$. So, $y_1 + y_2 + y_3 = n + 9$. We have

$$x_1 + x_2 + x_3 + y_1 + y_2 + y_3 \ge 4n + 6.$$
(5)

Since $x_1 + x_2 + x_3 = k \leq 3n - 3$, all x_1, x_2, x_3 cannot be greater than n - 1. We then separate the proof into 3 cases, according to the number of x_i that is greater than n - 1. We also have $y_1 + y_2 + y_3 = 13n - 3 - 4k \geq n + 9 > 9$. Therefore, y_1, y_2 and y_3 cannot simultaneously be less than four.

Case 1: $x_1, x_2, x_3 \le n - 1$.

Without loss of generality, we assume that $y_3 \leq y_2 \leq y_1$. We consider 3 sub-cases.

Case 1.1: $y_3 \le y_2 < 4 \le y_1$. From Eq. (5), we have

$$x_1 + y_1 \ge 2n + 2.$$

- If x₁ + y₁ ≥ 2n + 3, because x₁ ≤ n − 1, so y₁ ≥ n + 4 ≥ x₁ + 5. By Lemma 5, r₁ ≥ [x₁+y₁-3/2] ≥ [2n/2] = n.
 If x₁ + y₁ = 2n + 2, then this equality happens only when x₁ =
- If $x_1 + y_1 = 2n + 2$, then this equality happens only when $x_1 = n 1$. Thus, $y_1 = n + 3 \le n 1 + 4 = x_1 + 4$. Hence, by Lemma 5, we have $r_1 \ge y_1 3 = n$.

Case 1.2: $y_3 < 4 \le y_2 \le y_1$.

From Eq. (5), we get

$$x_1 + x_2 + y_1 + y_2 \ge (4n+6) - (n-1) - 3 = 3n+4 \tag{6}$$

To apply Lemma 5 to r_1 and r_2 , there are 4 possibilities. In each case, we use Eq. (6) and the inequalities $x_1, x_2 \leq n-1$ when needed.

• Let $x_1 \ge y_1 - 4$ and $x_2 \ge y_2 - 4$. Then

$$r_1 + r_2 \ge y_1 + y_2 - 6 \ge n.$$

• Let $x_1 \le y_1 - 5$ and $x_2 \le y_2 - 5$. Then

$$r_{1} + r_{2} \ge \left\lfloor \frac{x_{1} + y_{1} - 3}{2} \right\rfloor + \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor$$
$$\ge \frac{x_{1} + y_{1} - 4}{2} + \frac{x_{2} + y_{2} - 4}{2}$$
$$\ge \frac{3n - 4}{2}.$$

We know that $\frac{3n-4}{2} \ge n$ when $n \ge 4$. For n = 3, we have $r_1 + r_2 \ge \frac{5}{2}$. So $r_1 + r_2 \ge 3 = n$.

• Let $x_1 \ge y_1 - 4$ and $x_2 \le y_2 - 5$. (The case when $x_1 \le y_1 - 5$ and $x_2 \ge y_2 - 4$ can be treated similarly.) Then

$$r_{1} + r_{2} = y_{1} - 3 + \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor$$

$$\geq y_{1} - 3 + \frac{x_{2} + y_{2} - 4}{2}$$

$$= \frac{x_{1} + y_{1} + x_{2} + y_{2} - (x_{1} - y_{1}) - 10}{2}$$

$$\geq n - \frac{1}{2}.$$

The last inequality follows from $x_1 \leq n-1$ and $y_1 \geq 4$. Since $r_1 + r_2$ is an integer, we have $r_1 + r_2 \geq n$.

Case 1.3: $4 \le y_3 \le y_2 \le y_1$.

We will apply Lemma 5 to r_1, r_2 and r_3 , which have 8 possibilities. They can be separated into 4 groups, by the number of *i* where $x_i \ge y_i - 4$, as follows.

• There is no *i* where $x_i \ge y_i - 4$. Let $x_1 \le y_1 - 5$, $x_2 \le y_2 - 5$ and $x_3 \le y_3 - 5$. Then

$$r_{1} + r_{2} + r_{3} = \left\lfloor \frac{x_{1} + y_{1} - 3}{2} \right\rfloor + \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor + \left\lfloor \frac{x_{3} + y_{3} - 3}{2} \right\rfloor$$
$$\geq \frac{x_{1} + y_{1} - 4}{2} + \frac{x_{2} + y_{2} - 4}{2} + \frac{x_{3} + y_{3} - 4}{2}$$
$$= \frac{x_{1} + y_{1} + x_{2} + y_{2} + x_{3} + y_{3} - 12}{2}$$
$$\geq 2n - 3 \ge n.$$

• There is exactly one *i* where $x_i \ge y_i - 4$. Let $x_1 \le y_1 - 5$ and $x_2 \le y_2 - 5$, but $x_3 \ge y_3 - 4$. Then

$$r_{1} + r_{2} + r_{3} = \left\lfloor \frac{x_{1} + y_{1} - 3}{2} \right\rfloor + \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor + y_{3} - 3$$

$$\geq \frac{x_{1} + y_{1} - 4}{2} + \frac{x_{2} + y_{2} - 4}{2} + y_{3} - 3$$

$$\geq \frac{x_{1} + y_{1} + x_{2} + y_{2} + x_{3} + 2y_{3} - 14}{2}$$

$$\geq \frac{x_{1} + x_{2} + x_{3} + y_{1} + y_{2} + y_{3} - (x_{3} - y_{3}) - 14}{2}$$

$$\geq \frac{4n - 8 - (x_{3} - y_{3})}{2}$$

The last inequality follows from Eq. (5). Since $y_3 \ge 4$ and $x_3 \le n-1$, we have $x_3 - y_3 \le n-5$. Hence, $r_1 + r_2 + r_3 \ge \frac{3n-3}{2} \ge n$.

• There are exactly two *i* where $x_i \ge y_i - 4$. Let $x_1 \le y_1 - 5$, but $x_2 \ge y_2 - 4$ and $x_3 \ge y_3 - 4$. The other two cases can be treated in the same manner. Then

$$r_{1} + r_{2} + r_{3} \ge \left[\frac{x_{1} + y_{1} - 3}{2}\right] + y_{2} - 3 + y_{3} - 3$$

$$\ge \frac{x_{1} + y_{1} - 4}{2} + y_{2} + y_{3} - 6$$

$$\ge \frac{x_{1} + y_{1} + 2y_{2} + 2y_{3} - 16}{2}$$

$$\ge \frac{x_{1} + x_{2} + x_{3} + y_{1} + y_{2} + y_{3} - (x_{2} - y_{2}) - (x_{3} - y_{3}) - 16}{2}$$

$$\ge \frac{4n - 10 - (x_{2} - y_{2}) - (x_{3} - y_{3})}{2}.$$

Since $x_2 - y_2 \le n - 5$ and $x_3 - y_3 \le n - 5$, we get

$$r_1 + r_2 + r_3 \ge \frac{4n - 10 - 2(n - 5)}{2} = n.$$

• Let $x_1 \ge y_1 - 4$, $x_2 \ge y_2 - 4$ and $x_3 \ge y_3 - 4$. From Eq. (5) and $x_1 + x_2 + x_3 \le 3n - 3$, we then get

$$r_1 + r_2 + r_3 \ge y_1 + y_2 + y_3 - 9 \ge n.$$

Case 2: Exactly one of x_1, x_2, x_3 is more than n - 1.

Suppose without loss of generality that $x_1 > n - 1$ and $x_2, x_3 \le n - 1$. Let $x_1 = n - 1 + m$ for some $1 \le m < \lfloor \frac{n}{4} \rfloor$. Thus, we can partition U_1 into two parts where each part has at most n - 1 disjoint blue copies of K_4 and all edges between these two parts are red. Then the smaller part contains at least m disjoint blue copies of K_4 , and so does the larger part. Hence, there is a red matching M of size 4m (between the two sets of m disjoint blue copies of K_4) in $G[U_1]$.

Case 2.1: $y_1 < 4$.

With $x_1 = n - 1 + m$ and $y_1 \leq 3$, Eq. (5) gives

$$x_2 + y_2 + x_3 + y_3 \ge 3n + 4 - m. \tag{7}$$

If either y_2 or y_3 is less than four, then we can suppose that $y_2 < 4$. By Eq. (4), $y_3 \ge n+3$. Since $x_3 \le n-1$, the condition $x_3 \ge y_3 - 4$ occurs only when $x_3 = n-1$ and $y_3 = n+3$. Then by Lemma 5, $r_3 \ge y_3 - 3 = n$.

When $x_3 \leq y_3 - 5$, Lemma 5 gives

$$r_3 \ge \left\lfloor \frac{x_3 + y_3 - 3}{2} \right\rfloor \ge \frac{x_3 + y_3 - 4}{2}.$$

Since $x_2 \leq n-1$ and $y_2 \leq 3$, Eq. (7) gives

$$x_3 + y_3 \ge 2n + 2 - m.$$

Since $r_1 \ge 4m$ with $m \ge 1$, we have

$$r_1 + r_3 \ge 4m + \frac{2n + 2 - m - 4}{2} = \frac{2n - 2 + 7m}{2} > n.$$

Hence, we assume $y_2 \ge 4$ and $y_3 \ge 4$.

To apply Lemma 5 to r_2 and r_3 , we consider 4 possibilities. In each case, we use Eq. (7) and the inequalities $x_2 \leq n-1$ and $x_3 \leq n-1$ when necessary.

• If $x_2 \ge y_2 - 4$ and $x_3 \ge y_3 - 4$, we get $r_2 + r_3 \ge y_2 + y_3 - 6$ $\ge 3n + 4 - m - 2(n - 1) - 6$ = n - m.

So $r_1 + r_2 + r_3 \ge 4m + n - m \ge n$.

• If $x_2 \le y_2 - 5$ and $x_3 \le y_3 - 5$, we get

$$r_{2} + r_{3} \ge \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor + \left\lfloor \frac{x_{3} + y_{3} - 3}{2} \right\rfloor$$
$$\ge \frac{x_{2} + y_{2} - 4}{2} + \frac{x_{3} + y_{3} - 4}{2}$$
$$\ge \frac{x_{2} + y_{2} + x_{3} + y_{3} - 8}{2}$$
$$\ge \frac{3n - 4 - m}{2}.$$

Since $r_1 \ge 4m$ and $m \ge 1$, we have

$$r_1 + r_2 + r_3 \ge 4m + \frac{3n - 4 - m}{2} \ge \frac{3n - 4 + 7m}{2} \ge \frac{3n + 3}{2} \ge n.$$

• If $x_2 \ge y_2 - 4$ and $x_3 \le y_3 - 5$, we get

$$r_{2} + r_{3} \ge y_{2} - 3 + \left\lfloor \frac{x_{3} + y_{3} - 3}{2} \right\rfloor$$
$$\ge y_{2} - 3 + \frac{x_{3} + y_{3} - 4}{2}$$
$$= \frac{x_{2} + y_{2} + x_{3} + y_{3} - (x_{2} - y_{2}) - 10}{2}$$

With $x_2 \leq n-1$, $y_2 \geq 4$ and $m \geq 1$, we have

$$r_{2} + r_{3} \ge \frac{x_{2} + y_{2} + x_{3} + y_{3} - (x_{2} - y_{2}) - 10}{2}$$
$$\ge \frac{3n + 4 - m - (n - 5) - 10}{2}$$
$$\ge n - \frac{m + 1}{2}.$$

Since $r_1 \ge 4m$ and $m \ge 1$, we obtain

$$r_1 + r_2 + r_3 \ge 4m + n - \frac{m+1}{2} \ge n + \frac{7m-1}{4} \ge n.$$

• If $x_2 \leq y_3 - 5$ and $x_3 \geq y_2 - 4$, we can treat this in the same manner as in the previous case.

Case 2.2: $y_1 \ge 4$.

Apart from 2m disjoint blue copies of K_4 which gives a red matching of size 4m in $G[U_1]$, there are $x'_1 = x_1 - 2m = n - m - 1$ disjoint blue copies of K_4 ramained in $G[U_1]$. Let $r'_1 = r_1 - 4m$. We will apply Lemma 5 using x'_1 instead of x_1 , and get a lower bound on r'_1 . Consider the following situations based on the values of y_2 and y_3 .

- (1) Exactly one of y_2 and y_3 is less than 4. We assume without loss
 - of generality that $y_2 \ge 4$ but $y_3 < 4$. Eq. (4) gives

 $y_1 + y_2 \ge n + 6.$

Since $x_3 \leq n-1$, Eq. (5) also gives

 $x_1 + x_2 + y_1 + y_2 \ge 3n + 4.$

- (a) Suppose that $x'_1 \ge y_1 4$. By Lemma 5, $r'_1 \ge y_1 3$. So, $r_1 = r'_1 + 4m \ge y_1 - 3 + 4m$. If $x_2 \ge y_2 - 4$, then $r_2 \ge y_2 - 3$. Hence,
- $r_1 + r_2 \ge y_1 + y_2 + 4m 6 \ge n + 4m \ge n.$

If
$$x_2 \le y_2 - 5$$
, then $r_2 \ge \left\lfloor \frac{x_2 + y_2 - 3}{2} \right\rfloor \ge \frac{x_2 + y_2 - 4}{2}$. Hence,
 $r_1 + r_2 \ge y_1 - 3 + 4m + \frac{x_2 + y_2 - 4}{2}$
 $\ge \frac{x_2 + 2y_1 + y_2 + 8m - 10}{2}$
(8)

From $x_1 + x_2 + y_1 + y_2 \ge 3n + 4$, $x_1 = n - 1 + m$ and $y_1 \ge 4$, we obtain

$$r_1 + r_2 \ge \frac{2n + 7m - 1}{2} > n.$$

(b) Let
$$x'_1 \leq y_1 - 5$$
. By Lemma 5, $r'_1 \geq \left\lfloor \frac{x'_1 + y_1 - 3}{2} \right\rfloor$. So $r_1 = r'_1 + 4m \geq \frac{x'_1 + y_1 - 4}{2} + 4m$.
If $x_2 \geq y_2 - 4$, then $r_2 \geq y_2 - 3$. Hence,
 $r_1 + r_2 \geq \frac{x'_1 + y_1 - 4}{2} + 4m + y_2 - 3$
 $\geq \frac{x'_1 + y_1 + 2y_2 + 8m - 10}{2}$

$$= \frac{n - m - 1 + y_1 + 2y_2 + 8m - 10}{2}$$

$$= \frac{y_1 + 2y_2 + n + 7m - 11}{2}.$$
(9)

Since $y_1 + y_2 \ge n + 6$ and $y_2 \ge 4$, we get $y_1 + 2y_2 \ge n + 10$. Thus,

$$r_1 + r_2 \ge \frac{2n + 7m - 1}{2} \ge n.$$

If
$$x_2 \le y_2 - 5$$
, then $r_2 \ge \lfloor \frac{x_2 + y_2 - 3}{2} \rfloor \ge \frac{x_2 + y_2 - 4}{2}$. Hence,
 $r_1 + r_2 \ge 4m + \frac{x_1' + y_1 - 4}{2} + \frac{x_2 + y_2 - 4}{2}$
 $\ge \frac{8m + x_1' + y_1 + x_2 + y_2 - 8}{2}$
 $\ge \frac{7m + n - 9 + y_1 + x_2 + y_2}{2}$.

Since $x_1 + x_2 + y_1 + y_2 \ge 3n + 4$ and $x_1 = n + m - 1$, we get $y_1 + x_2 + y_2 \ge 2n + 5 - m$. Thus,

$$r_1 + r_2 \ge \frac{3n + 6m - 4}{2} > n.$$

- (2) Both y_2 and y_3 are less than four. From $y_1 + y_2 + y_3 \ge n + 9$, we get $y_1 \ge n + 3$. If $x' \ge x_1 - 4$, then $x' \ge x_1 - 2 \ge n$
 - If $x'_1 \ge y_1 4$, then $r'_1 \ge y_1 3 \ge n$. If $x'_1 \le y_1 - 5$, then $|x'_1 + y_1 - 3| = |2r'_1 + 2|$

$$r'_{1} \ge \left\lfloor \frac{x'_{1} + y_{1} - 3}{2} \right\rfloor \ge \left\lfloor \frac{2x'_{1} + 2}{2} \right\rfloor = x'_{1} + 1 = n - m.$$

Hence,

$$r_1 = 4m + r_1' \ge 3m + n > n.$$

(3) Let both y_2 and y_3 is at least four. ecall that $x'_1 = n - m - 1$ and $r_1 = r'_1 + 4m$. Similar to Eq. (5), we have

$$x_1' + x_2 + x_3 + y_1 + y_2 + y_3 \ge 4n - 2m + 6.$$

From
$$x_1 + x_2 + x_3 \leq 3n - 3$$
, we also get

$$y_1 + y_2 + y_3 \ge n + 9.$$

We apply Lemma 5 to r'_1 , r_2 and r_3 in the same manner as in Case 1.4. Note that the inequality $x'_1 - y_1 \le n - m - 5$ is used when $x'_1 \ge y_1 - 4$, while $x_i - y_i \le n - 5$ is used when $x_i \ge y_i - 4$ when $i \ne 1$.

For simplicity of the following inequality, we let A be the set of all i such that $x_i \ge y_i - 4$ (or $x'_i \ge y_i - 4$ when i = 1). We then have

$$r_{1}' + r_{2} + r_{3} \geq \begin{cases} 2n - m - 3, & A = \emptyset, \\ \frac{3n - m - 3}{2}, & A = \{1\}, \\ \frac{3n - 2m - 3}{2}, & A = \{2\} \text{ or } \{3\}, \\ n - m, & A = \{2, 3\}, \\ n - m, & A = \{2, 3\}, \\ n - \frac{m}{2}, & A = \{1, 2\} \text{ or } \{1, 3\}, \\ n, & A = \{1, 2, 3\}. \end{cases}$$

Hence,

$$r_1 + r_2 + r_3 \ge r'_1 + r_2 + r_3 + 4m \ge n.$$

From the two cases, we can conclude that $r_1 + r_2 + r_3 \ge n$.

Proof of Lemma 12. From the proof of Lemma 11, there are many situations from which the lower bounds on r_i are obtained. In each situation, we choose $s_i = 0$ when the obviouse bound $r_i \ge 0$ is obtained, and choose $s_i \le L_n$ when the lower bound $r_i \ge L_n$ is obtained (either from Lemma 5 or a red matching construction using a blue copy of K_4) in such a way that $s_1 + s_2 + s_3$ holds. We now prove that $|U_i| - s_i \ge n$.

- (1) $x_i \leq n-1$. The lower bound on r_i is obtained from Lemma 5.
 - If $y_i \leq 3$, then $r_i \geq 0$. Then, $|U_i| s_i = 4x_i + y_i$.
 - If $x_i \ge y_i 4$ and $y_i \ge 4$, then $r_i \ge y_i 3$. Then,

$$|U_i| - s_i \ge 4x_i + y_i - (y_i - 3) = 4x_i + 3.$$

• If $x_i \leq y_i - 5$ and $y_i \geq 4$, then $r_i \geq \lfloor \frac{x_i + y_i - 3}{2} \rfloor$. With $y_i \geq x_i + 5$, we get

$$\begin{aligned} |U_i| - s_i &\ge 4x_i + y_i - \left\lfloor \frac{x_i + y_i - 3}{2} \right\rfloor \\ &\ge 4x_i + y_i - \frac{x_i + y_i - 2}{2} \\ &= \frac{7x_i + y_i + 2}{2} \\ &\ge \frac{8x_i + 7}{2}. \end{aligned}$$

From the three cases, we can conclude that $|U_i| - s_i \ge 4x_i \ge n$.

- (2) $n-1 < x_i < n-1 + \lceil \frac{n}{4} \rceil$ and $y_i \ge 4$. We write $x_i = n+m-1$ where $m < \lceil \frac{n}{4} \rceil$. Also, $x'_i = x_i 2m = n-m-1$ and $r_i \ge r'_i + 4m$ where the lower bound on r'_i is obtained from Lemma 5.
 - If $x_i \ge y_i 4$, then

$$|U_i| - s_i \ge 4x_i + y_i - 4m - (y_i - 3) = 4n - 1 \ge n.$$

• If $x_i \leq y_i - 5$, then

$$|U_i| - s_i \ge 4x_i + y_i - 4m - \left\lfloor \frac{x'_i + y_i - 3}{2} \right\rfloor$$

$$\ge 4x_i + y_i - 4m - \frac{x'_i + y_i - 2}{2}$$

$$= \frac{8x_i - x'_i + y_i - 8m + 2}{2}$$

$$\ge \frac{7n + m - 1}{2}$$

$$> n.$$

(3) $n-1 < x_i < n-1 + \lceil \frac{n}{4} \rceil$ and $y_i \leq 3$. Similarly, we write $x_i = n + m - 1$ where $m < \lceil \frac{n}{4} \rceil$. The lower bound $r_i \geq 4m$ is obtained. Then,

$$|U_i| - s_i \ge 4x_i + y_i - 4m \ge 4n + y_i - 4 \ge n.$$

(4)
$$x_i \ge n - 1 + \lceil \frac{n}{4} \rceil$$
. We get $r_i \ge n$. Hence,
 $|U_i| - s_i \ge 4x_i + y_i - n \ge 4n + y_i - 4 \ge n$.

From all four cases, we can conclude that $|U_i| \ge s_i + n$.

5. Proof of the third case

Let $3n - 2 \le k \le 3n - 4 + \lfloor \frac{n}{4} \rfloor$. We have that $n \ge 5$. With the same setup as in Section 4, we let U_1, U_2 and U_3 be the partition of the vertices of G where each $G[U_i]$ contains at least $\lfloor \frac{n}{4} \rfloor$ disjoint blue copies of K_4 from K. In addition,

at most one $G[U_i]$, assumed to be $G[U_1]$, contains more than n-1 disjoint blue copies of K_4 .

Let x_i be the number of disjoint blue copies of K_4 in $G[U_i]$ from K, y_i be the number of vertices in $G[U_i]$ not belong to those disjoint blue copies of K_4 from K. Let r_i be the size of the maximum red matching in $G[U_i]$.

Since $k \ge 3n-2$, at least one of x_1, x_2 and x_3 is larger than n-1. Hence, $x_1 > n-1$ while $x_2 \le n-1$ and $x_3 \le n-1$.

Since $3n - 2 \le k \le 3n - 4 + \left\lceil \frac{n}{4} \right\rceil$, we can write

$$x_1 + x_2 + x_3 = k = 3n + p - 3 \tag{10}$$

for some $1 \leq p < \left\lceil \frac{n}{4} \right\rceil$. Thus,

$$y_1 + y_2 + y_3 = |V(G)| - 4k = 13n - 3 - 4k = n - 4p + 9.$$
(11)

We also have

$$x_1 + x_2 + x_3 + y_1 + y_2 + y_3 = 4n - 3p + 6.$$
(12)

Lastly, Eq. (10) gives $x_1 \ge n + p - 1$. So, we let

$$x_1 = n + m - 1 \tag{13}$$

for some $m \ge p$.

Similar to Section 4, we can show that $r_1 + r_2 + r_3 \ge n$.

Lemma 13. $r_1 + r_2 + r_3 \ge n$.

Using the same method as in Lemma 12, we can show that there is $s_i \leq r_i$ such that $s_1 + s_2 + s_3 = n$ and $|U_i| \geq s_i + n$ for all *i*. In exactly the same way as in Section 4, we can then construct a red copy of nK_4 from a red matching of size *n*.

Proof of Lemma 13. If $x_1 \ge n - 1 + \lceil \frac{n}{4} \rceil$, we can use Lemma 8 to partition the blue connected components in $G[U_1]$ into two parts, where each part has at least $\lceil \frac{n}{4} \rceil$ disjoint blue copies of K_4 . We then construct a red matching of size $4 \lceil \frac{n}{4} \rceil \ge n$ in $G[U_1]$. Thus, $r_1 \ge n$.

We now assume that $n-1 < x_1 < n-1 + \left\lceil \frac{n}{4} \right\rceil$. This implies that $m < \left\lceil \frac{n}{4} \right\rceil$. Hence,

$$1 \le p \le m < \left\lceil \frac{n}{4} \right\rceil. \tag{14}$$

Similarly, we can apply Lemma 8 to Eq. (13) and get a red matching of size 4m in $G[U_1]$. We let

$$x_1' = x_1 - 2m = n - m - 1.$$

Then $G[U_1] - M$ contains a blue copy of x'_1K_4 from K.

In order to apply Lemma 5, we need to consider different cases depending on whether $y_i \ge 4$ and whether $x_i \ge y_i - 4$. We note that for i = 1, we consider $G[U_1] - M$ instead of $G[U_1]$. Hence, we use x'_1 instead of x_1 together with the same y_1 , and let r'_1 be the size of the maximum red matching in $G[U_1] - M$. So

$$r_1 \ge r_1' + 4m.$$

Case 1: Both y_2 and y_3 are less than four. From Eq. (11), we have

$$y_1 \ge n - 4p + 3.$$

This also implies that $y_1 \ge 4$. Case 1.1: $x'_1 \ge y_1 - 4$. We have

$$r_1 \ge r_1' + 4m \ge y_1 - 3 + 4m \ge n.$$

Case 1.2: $x'_1 \le y_1 - 5$. We have

$$r'_{1} \ge \left\lfloor \frac{x'_{1} + y_{1} - 3}{2} \right\rfloor$$
$$\ge \frac{x'_{1} + y_{1} - 4}{2}$$
$$\ge \frac{(n - m - 1) + (n - 4p + 3) - 4}{2}$$
$$= \frac{2n - 4p - m - 2}{2}.$$

Hence,

$$r_1 \ge r'_1 + 4m \ge \frac{2n + 7m - 4p - 2}{2} \ge n.$$

Case 2: Exactly of one y_2 and y_3 is less than four. We may assume that $y_2 \ge 4$ and $y_3 < 4$.

Case 2.1: $y_1 < 4$.

From Eq. (11), we have

$$y_2 \ge n - 4p + 3.$$

Consider the following situations.

• $x_2 \ge y_2 - 4$. We have $r_2 \ge y_2 - 3 \ge n - 4p$. Since $r_1 \ge r'_1 + 4m \ge 4m$, we then get

$$r_1 + r_2 \ge n - 4p + 4m \ge n.$$

• $x_2 \leq y_2 - 5$. We get $r_2 \geq \lfloor \frac{x_2 + y_2 - 3}{2} \rfloor \geq \frac{x_2 + y_2 - 4}{2}$. Since $x_1 = n + m - 1$ and $x_3 \leq n - 1$, Eq. (12) gives

$$x_2 + y_2 = 4n - 3p + 6 - (x_1 + x_3 + y_1 + y_3)$$

$$\ge 2n - 3p - m + 2.$$

Hence,

$$r_1 + r_2 \ge 4m + \frac{2n - 3p - m - 2}{2}$$

 $\ge \frac{2n - 3p + 7m - 2}{2}$
 $> n$

Case 2.2: $y_1 \ge 4$. • $x'_1 \le y_1 - 5$. Then,

$$r'_1 \ge \left\lfloor \frac{x'_1 + y_1 - 3}{2} \right\rfloor \ge \left\lfloor \frac{2x'_1 + 2}{2} \right\rfloor = x'_1 + 1 = n - m.$$

Hence,

 $r_1 \ge r_1' + 4m \ge n.$

• $x'_1 \ge y_1 - 4$ and $x_2 \ge y_2 - 4$. From $y_3 \le 3$, Eq. (11) gives $y_1 + y_2 \ge n - 4p + 6$. From Lemma 5, we can conclude that

$$r_1 + r_2 \ge r'_1 + r_2 + 4m \ge y_1 + y_2 - 6 + 4m \ge n$$

• $x'_1 \ge y_1 - 4$ and $x_2 \le y_2 - 5$. Lemma 5 gives

$$r_{1} + r_{2} \ge y_{1} - 3 + 4m + \left\lfloor \frac{x_{2} + y_{2} - 3}{2} \right\rfloor$$

$$\ge y_{1} - 3 + 4m + \frac{x_{2} + y_{2} - 4}{2}$$

$$= \frac{x_{2} + 2y_{1} + y_{2} + 8m - 10}{2}$$

$$= \frac{x_{1} + x_{2} + y_{1} + y_{2} + (y_{1} - x_{1}) + 8m - 10}{2}.$$
(15)

Since $x_3 \leq n-1$ and $y_3 \leq 3$, Eq. (12) give

 $x_1 + x_2 + y_1 + y_2 \ge 3n - 3p + 4.$

Using $x_1 = n + m - 1$ and $y_1 \ge 4$, we then get $y_1 - x_1 \ge 5 - n - m$. Hence,

$$r_1 + r_2 \ge \frac{(3n - 3p + 4) + (5 - n - m) + 8m - 10}{2}$$
$$= \frac{2n - 3p + 7m - 1}{2}$$
$$\ge n.$$

Case 3: Both y_2 and y_3 are at least four.

Case 3.1: $y_1 \leq 3$. From Eq. (11), we have

$$y_2 + y_3 \ge n - 4p + 6.$$

From Eq. (12), we also have

 $x_2 + x_3 + y_2 + y_3 \ge 4n - 3p + 6 - (n + m - 1) - 3 = 3n - 3p - m + 4.$

We use the two inequalities above together with Lemma 5 in the following situations.

• $x_2 \ge y_2 - 4$ and $x_3 \ge y_3 - 4$. Then

$$r_1 + r_2 + r_3 \ge 4m + y_2 + y_3 - 6 \ge n.$$

•
$$x_2 \le y_2 - 5$$
 and $x_3 \le y_3 - 5$. We have
 $r_1 + r_2 + r_3 \ge 4m + \left\lfloor \frac{x_2 + y_2 - 3}{2} \right\rfloor + \left\lfloor \frac{x_3 + y_3 - 3}{2} \right\rfloor$
 $\ge 4p + \frac{x_2 + y_2 - 4}{2} + \frac{x_3 + y_3 - 4}{2}$
 $\ge \frac{3n - 3p + 7m - 4}{2}$

$$\geq n.$$

• $x_2 \ge y_2 - 4$ and $x_3 \le y_3 - 5$. (Similarly, $x_2 \le y_2 - 5$ and $x_3 \ge y_3 - 4$.) We get

$$r_{1} + r_{2} + r_{3} \ge 4m + y_{2} - 3 + \left\lfloor \frac{x_{3} + y_{3} - 3}{2} \right\rfloor$$
$$\ge 4m + y_{2} - 3 + \frac{x_{3} + y_{3} - 4}{2}$$
$$= \frac{8m + 2y_{2} + x_{3} + y_{3} - 10}{2}$$
$$= \frac{x_{2} + y_{2} + x_{3} + y_{3} + (y_{2} - x_{2}) + 8m - 10}{2}.$$

Since $y_2 \ge 4$ and $x_2 \le n - 1$, we have $y_2 - x_2 \ge 5 - n$. Hence, $r_1 + r_2 + r_3 \ge \frac{4n - 3p + 6 + 5 - n + 8m - 10}{2}$ $= \frac{3n - 3p + 8m + 1}{2}$

$$=\frac{3n-3p+8m+1}{2}$$

$$\geq n.$$

Case 3.2: $y_1 \ge 4$.

If $x'_1 \leq y_1 - 5$, we have

$$r'_1 \ge \left\lfloor \frac{x'_1 + y_1 - 3}{2} \right\rfloor \ge \left\lfloor \frac{2x'_1 + 2}{2} \right\rfloor = x'_1 + 1 = n - m.$$

Hence, $r_1 \ge r'_1 + 4m \ge n$.

So, we will assume that $x'_1 \ge y_1 - 4$. Lemma 5 then gives

 $r_1 \ge r_1' + 4m \ge y_1 + 4m - 3.$

To apply Lemma 5 to r_2 and r_3 , we consider the following situations. • $x_2 \ge y_2 - 4$ and $x_3 \ge y_3 - 4$. We have

$$r_1 + r_2 + r_3 \ge y_1 + y_2 + y_3 - 9 + 4m = n - 4p + 4m \ge n.$$

• $x_2 \le y_2 - 5$ and $x_3 \le y_3 - 5$. Then

$$\begin{aligned} r_1 + r_2 + r_3 &\geq y_1 - 3 + 4m + \left\lfloor \frac{x_2 + y_2 - 3}{2} \right\rfloor + \left\lfloor \frac{x_3 + y_3 - 3}{2} \right\rfloor \\ &\geq y_1 - 3 + 4m + \frac{x_2 + y_2 - 4}{2} + \frac{x_3 + y_3 - 4}{2} \\ &= \frac{2y_1 + x_2 + x_3 + y_2 + y_3 + 8m - 14}{2} \\ &= \frac{x_1 + x_2 + x_3 + y_1 + y_2 + y_3 + (y_1 - x_1) + 8m - 14}{2} \end{aligned}$$

Since $x_1 = n + m - 1$ and $y_1 \ge 4$, Eq. (12) gives

$$r_1 + r_2 + r_3 \ge \frac{3n - 3p + 7m - 3}{2} \ge n.$$

• $x_2 \ge y_2 - 4$ and $x_3 \le y_3 - 5$. Consider

$$r_1 + r_2 + r_3 \ge y_1 - 3 + 4m + y_2 - 3 + \left\lfloor \frac{x_3 + y_3 - 3}{2} \right\rfloor$$
$$\ge y_1 + y_2 + 4m - 6 + \frac{x_3 + y_3 - 4}{2}$$
$$= \frac{x_3 + 2y_1 + 2y_2 + y_3 + 8m - 16}{2}.$$

From Eq. (12), we have

$$r_1 + r_2 + r_3 \ge \frac{2n - 3p + 7m}{2} \ge n.$$

From all three cases, we conclude that $r_1 + r_2 + r_3 \ge n$.

6. Proof of the fourth case

Let $k \geq 3n - 3 + \lfloor \frac{n}{4} \rfloor$. From Lemma 10, we can combine V_1, V_2, \ldots, V_t and V' into four parts, say U_1, U_2, U_3 and U_4 where each $G[U_i]$ contains at least $\lfloor \frac{n}{4} \rfloor$ disjoint blue copies of K_4 from K. Thus, $|U_i| \geq 4 \lfloor \frac{n}{4} \rfloor \geq n$ for all i.

Since all edges between two parts are red, we can construct a red copy of K_4 by choosing one vertex from each part. Each part contains at least n vertices. Therefore, there is a n disjoint red copy nK_4 in G.

7. The case when n = 2

Although most of the techniques in the proof of Theorem 7 can be applied to the case when n = 2, there are still some cases in which such techniques cannot be directly applied. When $n \ge 3$, at the beginning of the proof of Theorem 7, we use R(4,4) = 18 to show that the number of disjoint blue copies of K_4 in K_{23} is at least $2n - 3 + \lfloor \frac{n-1}{4} \rfloor$ disjoint blue copies of K_4 which is larger than n - 1. This implies that there are at least two blue components. When n = 2, there may be some coloring that has only one blue copy of K_4 , hence there is only one blue component. In this case, we will use $R_2(nK_4)$, instead of $R_2(4,4)$. From [2], $R_2(nK_4) = 7n + 4$ when n is sufficiently large. To the authors' knowledge, there is no result on the exact value or the upper bound of $R_2(2K_4)$ for small n. So, if an upper bound of $R_2(2K_4)$ is small enough, we get the exact value of $R_2(\mathscr{C}(2K_4))$ as in the following theorem.

Theorem 14. If $R_2(2K_4) \leq 23$, then $R_2(\mathscr{C}(2K_4)) = 23$.

Proof. From Lemma 6, we have $R_2(\mathscr{C}(2K_4)) \geq 23$. Applying the same setup as in Theorem 7, we consider a 2-edge-coloring of $G = K_{23}$. Since 23 = 18 + 4 + 1, there are at least two monochromatic copies of K_4 in G. Suppose that red is connected and G contains at most one red copy of K_4 . Let k be the number of disjoint blue copies of K_4 in G. Then $1 \leq k \leq 5$. Assume that there is no blue copy of a graph in $\mathscr{C}(2K_4)$. We partition the set of vertices V(G) of G into V_1, V_2, \ldots, V_k and V' in the way that $G[V_i]$ is a blue component of G containing exactly one blue copy of K_4 in K and V' is the set of vertices outside $V_1 \cup \cdots \cup V_k$. Consider the following four cases.

Case 1: k = 1.

Let V'_1 be the set of vertices in V_1 other than those in the blue copy of K_4 . We note that by assumption of V_1 , $G[V'_1]$ contains no blue copy of K_4 .

- $|V_1| = 23$. This means blue is connected. Since $R_2(2K_4) \le 23$, $G[V_1]$ contains a red copy of $2K_4$.
- $|V_1| = 22$. Then $|V'_1| = 18$. Since R(4, 4) = 18, R(3, 4) = 9 but $G[V'_1]$ cannot have another blue copy of K_4 , there must be a red copy of $K_4 + K_3$. Joining the red copy of K_3 with another vertex from V', we obtain a red copy of $2K_4$.

- $16 \leq |V_1| \leq 22$. Since R(3, 4) = 9 and 16 > 12 = 9 + 3, there is a red copy of $2K_3$ in $G[V_1]$. Again, we combine a red copy of K_3 in $G[V_1]$ and one vertex in V', then we get a red copy of $2K_4$.
- $10 \leq |V_1| \leq 15$. Then $|V'| \geq 8$. Since R(2, 4) = 4 and $|V_1|, |V'| > 6$, there are two disjoint red copies of K_2 in both $G[V_1]$ and G[V']. Joining each pair of the copies of K_2 each from $G[V_1]$ and G[V'], we get a red copy of $2K_4$.
- $4 \leq |V_1| \leq 9$. Then $|V'| \geq 14$. Since R(3,4) = 9, G[V'] contains a red copy of $2K_3$. Again, joining each red copy of K_3 in G[V'] with another vertex in V_1 , we get a red copy of $2K_4$.

From all cases, we constructed a red copy of $2K_4$, which is a contradiction.

When $k \ge 2$, let $U_1 = V_1 \cup V'$ and $U_i = V_i$ for all $2 \le i \le k$. Let $y_i = |U_i| - 4$, which is the number of vertices in U_i not in the blue copy of K_4 .

Case 2: k = 2.

There are 8 vertices containing in these two disjoint blue copies of K_4 . Thus, $y_1 + y_2 = 15$ and we may assume that $y_1 \ge y_2$

- $y_1 \ge 12$. Since R(3,4) = 9, there is a red copy of $2K_3$ in $G[U_1]$. Combine this red copy of K_3 in $G[U_1]$ and one vertex in $G[U_2]$, then we obtain a red copy of $2K_4$.
- $10 \leq y_1 \leq 11$. Since R(3,4) = 9, $G[U_1]$ contains a red copy of K_3 . Note that $y_2 = 15 - y_1 \geq 4$. Applying Lemma 5 to $G[U_1] - K_3$ and $G[U_2]$, each of $G[U_1]$ and $G[U_2]$ contains at least one red copy of K_2 . Joining the two disjoint red copies of K_2 , we get a red copy of K_4 . Then joining the red copy of K_3 in $G[U_1]$ with any remaining vertex in U_2 , we get another red copy of K_4 .
- $8 \le y_1 \le 9$. Then $y_2 = 15 y_1 \ge 6$. By Lemma 5, each of $G[U_1]$ and $G[U_2]$ contains two disjoint red copies of K_2 . Joining each pair of red copies of K_2 , we get a red copy of $2K_4$.

Case 3: k = 3.

There are 12 vertices containing in three disjoint blue copies of K_4 . Therefore, $y_1 + y_2 + y_3 = 11$ and we can suppose that $y_1 \ge y_2 \ge y_3$.

- $y_1 \ge 4 > y_2 \ge y_3$. This implies $y_1 \ge 5$. By Lemma 5, $G[U_1]$ contains a red copy of $2K_2$. Construct two disjoint red copies of K_4 where each copy is formed by a red copy of K_2 in $G[U_1]$, a vertex in U_2 and a vertex in U_3 .
- $y_1 \ge y_2 \ge 4 > y_3$. By Lemma 5, each of $G[U_1]$ and $G[U_2]$ contains a red copy of K_2 . The first red copy of K_4 is constructed from one red copy of K_2 in $G[U_1]$, one vertex in U_2 and one vertex in U_3 . The second red copy of K_4 is constructed from one red copy of K_2 in $G[U_2]$, one vertex in U_1 and one vertex in U_3 .

Case 4: $k \ge 4$.

We form two disjoint red copies of K_4 . Each copy is formed by joining one vertex from each $G[U_i]$ for $1 \le i \le 4$.

From four cases, we get two disjoint red copies of K_4 . This contradicts the assumption.

8. CONCLUSION AND DISCUSSION

We have shown that $R_2(\mathscr{C}(nK_4)) = 13n - 3$ for $n \ge 3$ and the formula still holds for n = 2 if $R_2(2K_4) \le 23$.

The exact Ramsey number $R_2(\mathscr{C}(nK_r)) = (r^2 - r + 1)n - r + 1$ when $r \geq 5$ are known when $n \geq R_2(K_r)$ [13]. But, the result may not be true when n is very small. For example, when n = 2 and r = 5, we use the result from [13] and get $R_2(\mathscr{C}(2K_5)) = 21(2) - 4 = 38 < 43 \leq R(5,5)$, which is impossible. It is interesting to know the upper bound of $R_2(\mathscr{C}(nK_r))$, or in particular $R_2(\mathscr{C}(nK_5))$, for small n.

To the authors' knowledge, there is no result on the Ramsey number $R_2(\mathscr{C}(nG))$ for a non-complete graph G. So the results for other basic graphs G such as cycles, paths, trees, or stars will be very interesting.

References

- Stefan A Burr. Ramsey numbers involving graphs with long suspended paths. Journal of the London Mathematical Society, 2(3):405–413, 1981.
- [2] Stefan A Burr. On the ramsey numbers r(G, nH) and r(nG, nH) when n is large. Discrete mathematics, 65(3):215-229, 1987.
- [3] Stefan A Burr, P Erdős, and Joel H Spencer. Ramsey theorems for multiple copies of graphs. Transactions of the American Mathematical Society, 209:87–99, 1975.
- [4] Ernest J Cockayne and Peter J Lorimer. The ramsey number for stripes. Journal of the Australian Mathematical Society, 19(2):252–256, 1975.
- [5] Michael Codish, Michael Frank, Avraham Itzhakov, and Alice Miller. Computing the ramsey number r(4,3,3) using abstraction and symmetry breaking. *Constraints*, 21:375–393, 2016.
- [6] Robert E Greenwood and Andrew Mattei Gleason. Combinatorial relations and chromatic graphs. Canadian Journal of Mathematics, 7:1–7, 1955.
- [7] András Gyárfás and Gábor N Sárközy. Ramsey number of a connected triangle matching. Journal of Graph Theory, 83(2):109–119, 2016.
- [8] András Gyárfás and Gábor Simony. Edge colorings of complete graphs without tricolored triangles. Journal of Graph Theory, 46(3):211–216, 2004.
- [9] Peter Lorimer. The ramsey numbers for stripes and one complete graph. Journal of graph theory, 8(1):177–184, 1984.
- [10] Peter J Lorimer and RJ Segedin. Ramsey numbers for multiple copies of complete graphs. Journal of Graph Theory, 2(1):89–91, 1978.
- [11] Stanislaw Radziszowski. Small ramsey numbers. The electronic journal of combinatorics, 1000:DS1-Aug, 2011.
- [12] Frank P Ramsey. On a problem of formal logic. Classic Papers in Combinatorics, pages 1-24, 1987.
- [13] Barnaby Roberts. Ramsey numbers of connected clique matchings. Electronic Journal of Combinatorics, 24(1), 2017.

- [14] Saul Stahl. On the ramsey number $r(F, K_m)$ where f is a forest. Canadian Journal of Mathematics, 27(3):585–589, 1975.
- [15] Douglas Brent West et al. Introduction to graph theory, volume 2. Prentice hall Upper Saddle River, 2001.

DIVISION OF COMPUTATIONAL SCIENCE, FACULTY OF SCIENCE, PRINCE OF SONGKLA UNIVERSITY, THAILAND

Email address: kritmpsu@hotmail.com

DIVISION OF COMPUTATIONAL SCIENCE, FACULTY OF SCIENCE, PRINCE OF SONGKLA UNIVERSITY, THAILAND

Email address: panupong.v@psu.ac.th