OBSERVABILITY INEQUALITY FROM MEASURABLE SETS AND THE STACKELBERG-NASH GAME PROBLEM FOR DEGENERATE PARABOLIC EQUATIONS

YUANHANG LIU¹, WEIJIA WU^{1,*}, DONGHUI YANG¹, CAN ZHANG² ¹ SCHOOL OF MATHEMATICS AND STATISTICS, CENTRAL SOUTH UNIVERSITY, CHANGSHA 410083, CHINA. ² SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN 430072, CHINA.

ABSTRACT. In this study, we employ the established Carleman estimates and propagation estimates of smallness from measurable sets for real analytic functions, along with the telescoping series method, to establish an observability inequality for the degenerate parabolic equation over measurable subsets in the time-space domain. As a direct application, we formulate a captivating Stackelberg-Nash game problem and provide a proof of the existence of its equilibrium. Additionally, we characterize the set of Stackelberg-Nash equilibria and delve into the analysis of a norm optimal control problem.

1. INTRODUCTION

Observability inequality is a powerful and significant tool for investigating stability and controllability problems in evolution equations. The degenerate parabolic equation, which is a common class of diffusion equations, can describe numerous physical phenomena, such as laminar flow, large ice blocks, solar radiation-climate interactions, and population genetics (for more detailed descriptions, refer to [13]). As a result, the control problems for such equations have gained considerable attention, with studies on controllability and observability problems for some degenerate parabolic equations in [13] and its extensive references. In [11], the authors derived the observability inequality for a kind of degenerate parabolic problems, where the control acts on an open sets, by Carleman estimates based on the choice of suitable weighted functions and Hardy-type inequalities. In [23], the author considered non-smooth general degenerate parabolic equations in non-divergence form with the control acts on an open sets and with degeneracy and singularity occurring in the interior of the spatial domain, in presence of Dirichlet or Neumann boundary conditions. In particular, they obtained the observability inequality for the associated adjoint problem by Carleman estimates. In [8], the authors considered a parabolic problem with degeneracy in the interior of the spatial domain and Neumann boundary conditions with the control acts on an open sets. In particular, they focused on the well-posedness of the problem and on Carleman estimates for the associated adjoint problem. As a consequence, new observability inequalities were established. For other controllability issues related to degenerate parabolic equations, see [10, 14, 17, 22] and references therein.

It is worth noting that the current researches on observability inequalities for degenerate parabolic equations have primarily focused on control exerted on open sets, with little existing results considering control exerted on measurable sets. The reason for these

 $^{^{1} {\}rm Corresponding\ author:\ weijiawu@yeah.net}$

²⁰²⁰ Mathematics Subject Classification. 93B05,93B07.

 $Key\ words\ and\ phrases.$ Observability inequality, measurable sets, Stackelberg-Nash equilibrium, degenerate parabolic equations.

is that in the above works, the main technique used in the arguments, Carleman inequalities, requires to construct suitable Carleman weights: a role for functions which requires smoothness and to have the extreme values in proper regions associated to the control region. The construction of such functions seems to be not possible, when the control region does not an open set. In [29], The authors employed the available Carleman estimates, propagation estimates of smallness from measurable sets for real analytic functions, and the telescoping series method to establish an observability inequality from measurable subsets in the time-space variable for the degenerate parabolic equation with the Grushin operator in certain multidimensional domains. On the contrary, for nondegenerate parabolic equations, there have been extensive findings pertaining to control exerted on measurable sets. The authors in [3, 31] establish the observability inequality of the heat equation for the measurable subsets, and show the null controllability with controls restricted over these sets. In [19], the authors found new quantitative estimates on the space-time analyticity of solutions to linear parabolic evolutions with time-independent analytic coefficients and applied them to obtain observability inequalities for its solutions over measurable sets.

The current paper represents an advancement in the study of observability inequality for measurable subsets within the context of degenerate parabolic equations. However, unlike the approach discussed earlier, which relied on Carleman estimates, this paper builds upon the conventional observability inequality established in [11] and [9] for open subsets. Notably, it is currently not feasible to derive an observability inequality for degenerate parabolic equations from measurable subsets using the Carleman estimates method. Nevertheless, we have made an intriguing observation: When the observation domain is sufficiently distant from the singularity, we can derive an observability inequality with a constant term on the right-hand side, expressed as $e^{\frac{C}{Tk}}$, where C and k are positive parameters, and the solution is real analytic. By employing a telescoping series argument, we are able to obtain the desired result. This novel approach provides fresh insights into the controllability of degenerate parabolic equations, offering alternative perspectives beyond the sole reliance on Carleman estimates.

As an important application, we consider a game problem for degenerate parabolic equations. Differential games were introduced originally by Isaacs (see [26]). Since then, lots of researchers were attracted to establish and improve the related theory. Meanwhile, the theory was applied to a large number of fields. For a comprehensive survey on the differential game theory, we refer to [7, 18, 20, 24] and the references therein. The famous game problem, Stackelberg-Nash game problem, is an interesting question in the framework of mathematical finance. For instance, it is well known that the price of an European call option is governed by a backward PDE. The independent space variable must be interpreted as the stock price and the time variable is in fact the reverse of time. In this regard, it can be interesting to control the solution of the system with the composed action of several agents, each of them corresponding to a different range of values of the space variable. For further information on the modeling and control of phenomena of this kind, see for instance [32, 36]. For more Stackelberg–Nash game problems, we refer the reader to the works in [16, 25]. In this paper, based on the Stackelberg–Nash null controllability, see [4–6], we will focus on the characterizations of the set of Stackelberg–Nash equilibria and a norm optimal control problem. To the best of our knowledge, this paper is the first attempt to obtain the existence of the Stackelberg–Nash equilibria by the Kakutani type fixed point theorems for degenerate parabolic equations. Furthermore, we characterize the sets of Stackelberg–Nash equilibria and consider a norm optimal control problem. After that, we can mention [35], which seems the first time to apply the Kakutani type fixed point theorems to the game problems for heat equations.

The remaining sections of this paper are organized as follows. In Section 2, we present the formulation of the main problems and state the key results: Theorem 2.1, Theorem 2.2, Theorem 2.3, and Theorem 2.7. The proof of Theorem 2.1 is provided in Section 3. In Section 4, we delve into the discussion of a forthcoming Stackelberg-Nash game problem (to be formulated later) and provide the proofs for Theorem 2.2 and Theorem 2.3. These findings are instrumental in the subsequent discussion of a norm optimal control problem (to be formulated later) and the proof of Theorem 2.7 in Section 5.

2. Problems formulation and main results

In this section, we will outline the principal issues investigated and present the pivotal findings of this manuscript. Firstly, let us introduce necessary notations.

Let T > 0 be a fixed positive time constant, and I := (0, 1). Throughout this paper, we denote by $\langle \cdot, \cdot \rangle$ the scalar product in $L^2(I)$ and denote by $\|\cdot\|$ the norm induced by $\langle \cdot, \cdot \rangle$. We denote by $|\cdot|$ the Lebesgue measure on \mathbb{R} .

Let A be an unbounded linear operator on $L^{2}(I)$:

$$\begin{cases} \mathcal{D}(A) := \{ v \in H^{1}_{\alpha}(I) : (x^{\alpha}v_{x})_{x} \in L^{2}(I) \text{ and } BC_{\alpha}(v) = 0 \}, \\ Av := (x^{\alpha}v_{x})_{x}, \ \forall v \in \mathcal{D}(A), \ \alpha \in (0, 2), \end{cases}$$
(2.1)

where

$$H^1_{\alpha}(I) := \left\{ v \in L^2(I) : v \text{ is absolutely continuous in } I, x^{\frac{\alpha}{2}} v_x \in L^2(I) \text{ and } v(1) = 0 \right\},$$

and

$$BC_{\alpha}(v) = \begin{cases} v_{|x=0}, & \alpha \in (0,1), \\ (x^{\alpha}v_x)_{|x=0}, & \alpha \in [1,2), \end{cases}$$

endowed with the norms

,

$$\|v\|_{H^1_{\alpha}(I)}^2 := \|v\|^2 + \|\sqrt{x^{\alpha}}v_x\|^2.$$

The system, that we consider in this paper, is described by the following degenerate equation:

$$\begin{cases} y_t(x,t) - Ay(x,t) = 0, & (x,t) \in I \times (0,T), \\ y(1,t) = BC_{\alpha}(y(\cdot,t)) = 0, & t \in (0,T), \\ y(x,0) = y_0(x), & x \in I, \end{cases}$$
(2.2)

where initial state $y_0 \in L^2(I)$, y is the state variable. By [11], one can check that for all $y_0 \in L^2(I)$, system (2.2) admit a unique solution y in the space $C([0,T];L^2(I)) \cap$ $L^{2}(0,T;H^{1}_{\alpha}(I)).$

The main result of this paper is the following observability inequality for the degenerate parabolic equation (2.2):

Theorem 2.1. Let T > 0, $\alpha \in (0, 2)$. Let $\mu \in (0, 1)$ be defined as

$$\mu = \begin{cases} \frac{3}{4}, & \text{if } \alpha \in (0,2) \setminus \{1\}, \\ \frac{3}{2\gamma} \text{ for any } \gamma \in (0,2), & \text{if } \alpha = 1, \end{cases}$$

$$(2.3)$$

and $D \subset I \times (0,T)$ be a measurable subset with positive measures. Then there exists a constant $C = C(T, I, \alpha, D, \mu) \geq 1$ such that the solution of equation (2.2) satisfies the following observability inequality: for any $y_0 \in L^2(I)$,

$$||y(T)|| \le C \int_{D} |y(x,t)| dx dt.$$
 (2.4)

As an important application of Theorem 2.1, we consider the following degenerate problem with three controls (one leader and two followers), but very similar considerations hold for other systems (degenerated cases or non-degenerated cases) with a higher number of controls.

$$\begin{cases} y_t(x,t) = Ay(x,t) + \chi_{\omega}g + \chi_{\omega_1}u_1 + \chi_{\omega_2}u_2, & (x,t) \in (0,1) \times (0,T), \\ y(1,t) = BC_{\alpha}(y(\cdot,t)) = 0, & t \in (0,T), \\ y(x,0) = y_0(x), & x \in (0,1), \end{cases}$$
(2.5)

where A defined in (2.1), $\omega, \omega_1, \omega_2$ are measurable subsets of (0, 1) with positive measures. ω is the main (leader) control domain and ω_1, ω_2 are the secondary (followers) control domains. The controls $g, u_1, u_2 \in L^{\infty}(0, T; L^2(0, 1))$, where g is the main (leader) control and u_1 and u_2 are the secondary (followers) controls. Initial state is $y_0 \in L^2(0, 1)$. The solution of system (2.5) denote by $y(\cdot; y_0, g, u_1, u_2)$.

We define the following admissible sets of controls:

$$\mathcal{U}_0 := \left\{ g \in L^{\infty}(0,T; L^2(0,1)) : \|g\|_{L^{\infty}(0,T; L^2(0,1))} \le M_0 \right\},\$$

and

$$\mathcal{U}_1 := \{ u_1 \in L^{\infty}(0,T; L^2(0,1)) : \|u_1\|_{L^{\infty}(0,T; L^2(0,1))} \le M_1 \}, \\ \mathcal{U}_2 := \{ u_2 \in L^{\infty}(0,T; L^2(0,1)) : \|u_2\|_{L^{\infty}(0,T; L^2(0,1))} \le M_2 \},$$

where M_0, M_1, M_2 are positive constants. Meanwhile, we introduce the following two functionals: For each i = 1, 2, the functional $J_i : (L^{\infty}(0, T; L^2(0, 1)))^2 \to [0, +\infty)$ is defined by

$$J_i(u_1, u_2) := \|y(T; y_0, g, u_1, u_2) - y_T^i\|_{L^2(G_i)},$$
(2.6)

where $G_i \subset (0, 1)$ (i = 1, 2) be measurable subsets such that $\omega_i \subset G_i$ representing observation domains for the followers and $y_i^T \in L^2(0, 1)$ and

$$y_T^1 \neq y_T^2. \tag{2.7}$$

The mathematical models for the control problems are:

1. The followers u_1 and u_2 assume that the leader g has made a choice and intend to be a Nash equilibrium for the cost functionals $J_i(u_1, u_2), i = 1, 2$. Thus, once the leader g has been fixed:

(P1) Does there exist $(u_1^*, u_2^*) \in \mathcal{U}_1 \times \mathcal{U}_2$ with respect to g so that

$$J_1(u_1^*, u_2^*) \le J_1(u_1, u_2^*)$$
 for all $u_1 \in \mathcal{U}_1$ (2.8)

and

$$J_2(u_1^*, u_2^*) \le J_2(u_1^*, u_2)$$
 for all $u_2 \in \mathcal{U}_2$? (2.9)

We call the problem (P1) as a Stackelberg-Nash game problem. It is a noncooperative game problem of the followers. If the answer to the problem (P1) is yes, we call (u_1^*, u_2^*) a Stackelberg-Nash equilibrium (or an optimal strategy pair, or an optimal control pair) of (P1). We can understand the problem (P1) in the following manner: There are two followers executing their strategies and hoping to achieve their goals y_T^1 and y_T^2 , respectively. If the first follower chooses the strategy u_1^* , then the second follower can execute the strategy u_2^* so that $y(T; y_0, g, u_1^*, u_2^*)$ is closer to y_T^2 ; Conversely, if the second follower chooses the strategy u_2^* , then the first follower can execute the strategy u_1^* so that $y(T; y_0, g, u_1^*, u_2^*)$ is closer to y_T^1 . Roughly speaking, if one follower is deviating from (u_1^*, u_2^*) , then the cost functional of this follower would get larger; and there is no information given if both followers are deviating from the Nash equilibrium (u_1^*, u_2^*) . From the knowledge of game theory, in generally, Nash equilibrium is not unique.

2. Once the Stackelberg-Nash equilibrium (u_1^*, u_2^*) has been identified for each leader g, denoted by $(u_1^*(g), u_2^*(g))$, we consider the following norm optimal control problem:

$$(\mathbf{P2}) \quad N(T, y_0) := \inf_{g \in \mathcal{U}_0} \{ \|g\|_{L^{\infty}(0,T;L^2(0,1))} : y(T; y_0, g, u_1^*(g), u_2^*(g)) = 0 \}.$$
(2.10)

The first main result of the *Stackelberg-Nash game problem* (**P1**) is about the existence of a Stackelberg-Nash equilibrium.

Theorem 2.2. Let $g \in L^{\infty}(0,T; L^2(0,1))$ be given. The problem **(P1)** with respect to the system (2.5) admits a Stackelberg-Nash equilibrium, i.e., there exist at least a pair of $(u_1^*, u_2^*) \in \mathcal{U}_1 \times \mathcal{U}_2$ with respect to g so that (2.8) and (2.9) hold.

The second main result of the *Stackelberg-Nash game problem* (P1) is the characterizations of the set of Stackelberg-Nash equilibria, which concerned with the following bang-bang property of the Stackelberg-Nash equilibria.

Theorem 2.3. Let $g \in L^{\infty}(0,T;L^2(0,1))$ be given and let $(u_1^*,u_2^*) \in U_1 \times U_2$ be a Stackelberg-Nash equilibrium of the problem **(P1)** with respect to g. Then either

$$||u_1^*(t)||_{L^2(G_1)} = M_1$$
, for a.e. $t \in (0,T)$,

or

$$||u_2^*(t)||_{L^2(G_2)} = M_2$$
, for a.e. $t \in (0,T)$.

Remark 2.4. It is noteworthy that in previous studies addressing the Stackelberg-Nash game problem, the scholars have predominantly concentrated on the examination of system controllability, such as exact controllability [4], approximate controllability [16], null controllability [6], and the presence of Stackelberg-Nash equilibrium, while paying no attention to the comprehensive characterization of the set of Stackelberg-Nash equilibrium and their distinctive bang-bang property.

The result of the norm optimal control problem $(\mathbf{P2})$ is concered with the characterization of the leader. To the end, we make some assumptions:

- (A₁). $\omega_i \subset \omega \subset (0, 1)$ for i = 1, 2.
- (A₂). System (2.5) is null controllability, i.e. there exists a control $g \in L^{\infty}(0,T; L^2(0,1))$ such that the corresponding solution y of (2.5) satisfies $y(T; y_0, g, u_1^*(g), u_2^*(g)) = 0$, in (0, 1).

Remark 2.5. The assumption (A_1) is exclusively utilized for the purpose of characterizing the leader g, specifically in Theorem 2.7, which will be introduced later. The assumption (A_2) is well-founded, supported by several references, e.g., [4-6] demonstrating the null controllability of system (2.5). Nevertheless, in the present section, our primary emphasis lies in delineating the properties of the leader g. Therefore, we have omitted the detailed proof of null controllability of system (2.5), directing interested readers to consult the above references for further exploration.

Let us introduce the following adjoint equation corresponding to (2.2):

$$\begin{cases} z_t(x,t) + Az(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z(1,t) = BC_{\alpha}(z(\cdot,t)) = 0, & t \in (0,T), \\ z(x,T) = z_T(x), & x \in (0,1), \end{cases}$$
(2.11)

where $z_T(x) \in L^2(0, 1)$.

Now, for every $T \in \mathbb{R}^+$, let

$$X_T = \{ \chi_{\omega} z(\cdot; z_T) \mid z_T \in L^2(0, 1) \},$$
(2.12)

where $z(\cdot; z_T)$ is the solution of system (2.11) with the terminal value $z_T \in L^2(0, 1)$. It is obviously that X_T is a linear subspace of $L^1(0, T; L^2(0, 1))$. Define

$$Y_T = \overline{X_T}^{\|\cdot\|_{L^1(0,T;L^2(0,1))}}$$
(2.13)

and denote

$$Z_{T} = \left\{ \chi_{\omega} z \in L^{1}(0,T; L^{2}(0,1)) \mid \text{for every } s \in (0,T), \text{ there exists } z_{T,s} \in L^{2}(0,1) \\ \text{such that } z(\cdot) = e^{A(s-\cdot)} z_{T,s} \text{ on } [0,s] \right\}.$$
(2.14)

Lemma 2.6. Suppose (A_2) holds. Let T > 0, $y_0 \in L^2(0,1)$. Let Stackelberg-Nash equilibria $(u_1^*(g), u_2^*(g))$ be given with respect to g. Then there exists at least one $g^* \in L^{\infty}(0,T; L^2(0,1))$ such that $N(T, y_0) = ||g^*||_{L^{\infty}(0,T; L^2(0,1))}$.

The main result of the norm optimal control problem (P2) is stated as follows.

Theorem 2.7. Suppose (A_1) and (A_2) hold. Let $y_0 \in L^2(0,1)$, T > 0. Let $(u_1^*(g^*), u_2^*(g^*))$ be given with respect to g^* in Lemma 2.6 (i.e., g^* is the optimal control for the problem **(P2)**), and

$$J(\chi_{\omega}z) = \frac{1}{2} \left[\int_0^T \|\chi_{\omega}z(t)\| dt \right]^2 + \langle y_0, z(0) \rangle + \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z(t) dx dt$$
$$= \frac{1}{2} \|\chi_{\omega}z\|_{L^1(0,T;L^2(0,1))}^2 + \langle y_0, z(0) \rangle + \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z(t) dx dt,$$

where $\chi_{\omega} z \in Z_T$. Denote a variational problem

$$V(T, y_0) = \inf_{\chi_{\omega} z \in Z_T} J(\chi_{\omega} z).$$

Then there exists a $\chi_{\omega} z_* \in Z_T$ such that

$$V(T, y_0) = J(\chi_\omega z_*).$$

Furthermore, we have the following conclusions:

(I). If
$$\chi_{\omega} z_* \neq 0$$
, then

$$g^*(t) = \frac{\chi_{\omega} z_*(t)}{\|\chi_{\omega} z_*(t)\|} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))}, \text{ for a.e. } t \in (0,T).$$
(II). If $\chi_{\omega} z_* = 0$, then

$$g^*(t) = 0$$
, for a.e. $t \in (0,T)$.

Moreover,

$$V(T, y_0) = -\frac{1}{2}N(T, y_0)^2.$$

3. The proof of Theorem 2.1

The equation (2.2) will be referred to as the controlled system in our discussion, where the control in the form of $\chi_{\omega} u$ is applied on the right-hand side of the equation. Here, $u \in L^2(0,T; L^2(I))$ represents a function, and ω denotes a subset of I.

Let us recall an important result that will be used later, see Section 3.4 in [9].

Lemma 3.1. Let $\alpha \in (0,2)$ and $\omega_0 \subset I$ be an open subset. Let $\mu \in (0,1)$ be defined in (2.3). Then there exists a positive constant $C = C(\alpha, \omega_0, I, \mu)$ such that the solution of (2.2) satisfies the following observability inequality:

$$\int_{I} |y(x,T)|^2 dx \le C e^{CT^{-\frac{\mu}{1-\mu}}} \int_{0}^{T} \int_{\omega_0} |y(x,t)|^2 dx dt, \,\forall y_0 \in L^2(I).$$
(3.1)

Next, we aim to quantify the real analyticity of the solution y of equation (2.2) on the subdomain that is away from the singularity point zero.

Lemma 3.2. Set $\alpha \in (0,2)$. Let ω be a subdomain of I with $0 \notin \bar{\omega}$. Then there are positive constants $C = C(\alpha, I, \omega) \geq 1$ and $\rho = \rho(\alpha, I, \omega), 0 < \rho \leq 1$, such that when $x \in \omega, \forall 0 \leq s < t, a \in \mathbb{N}$ and $\gamma \in \mathbb{N}$, the solution of equation (2.2) satisfies

$$|\partial_x^a \partial_t^\gamma y(x,t)| \le \frac{Ce^{\frac{C}{t-s}} a! \gamma!}{\rho^a ((t-s)/2)^\gamma} \|y(\cdot,s)\|.$$
(3.2)

Proof. We borrow some ideas from [29]. We only need to prove the situation of s = 0. Let $\{e_k\}_{k\geq 1}$ and $\{w_k^2\}_{k\geq 1}$ be respectively the sets of $L^2(I)$ -normalized eigenfunctions and eigenvalues for $-\frac{\partial}{\partial x}\left(x^{\alpha}\frac{\partial}{\partial x}\right)$ with zero lateral Dirichlet boundary conditions; i.e.,

$$\begin{cases} \left(x^{\alpha}(e_k(x))_x\right)_x + w_k^2 e_k(x) = 0, & \text{in } I, \\ e_k = BC_{\alpha}(e_k) = 0, & \text{on } \partial I, \end{cases}$$
(3.3)

here, $0 < w_1 \le w_2 \le \cdots \le w_k \le \cdots$ and $\lim_{k \to +\infty} w_k = +\infty$. Take $y_0 = \sum_{k \ge 1} a_k e_k$ with

$$\sum_{k\geq 1} a_k^2 < +\infty,\tag{3.4}$$

and define

$$y(x, \tau, t) = \sum_{k=1}^{+\infty} a_k e^{-w_k^2 t + w_k \tau} e_k(x)$$
, for $x \in I, t > 0$, and $\tau \in \mathbb{R}$.

Then for all t > 0 and $x \in I$,

$$y(x,0,t) = \sum_{k=1}^{+\infty} a_k e^{-w_k^2 t} e_k(x) = y(x,t)$$

solves equation (2.2) with initial datum y_0 and

$$\partial_t^{\gamma} y(x,\tau,t) = \sum_{k \ge 1} a_k (-w_k^2)^{\gamma} e^{-w_k^2 t + w_k \tau} e_k(x), \text{ for } x \in I, \ \gamma \in \mathbb{N}, \text{ and } \tau \in \mathbb{R}.$$
(3.5)

Moreover, together with (3.3), we have

$$\begin{cases} \partial_{\tau}^{2}(\partial_{t}^{\gamma}y) + \frac{\partial}{\partial x}\left(x^{\alpha}\frac{\partial}{\partial x}(\partial_{t}^{\gamma}y)\right) = 0, & \text{in } I \times \mathbb{R}, \\ \partial_{t}^{\gamma}y = BC_{\alpha}(\partial_{t}^{\gamma}y) = 0, & \text{on } \partial I \times \mathbb{R}. \end{cases}$$
(3.6)

Because $0 \notin \bar{\omega}$, without loss of generality (otherwise use of a finite covering argument), we can assume that there exists $R \leq \frac{1}{8}$, $x_0 \in \omega$ such that

$$\omega \subset D_R(x_0), \ D_{2R}(x_0) \subset I, \ \text{and} \ D_{2R}(x_0) \cap D_R(0) = \emptyset,$$
(3.7)

where

$$D_R(x_0) = \{ x \in \omega : |x - x_0| \le R \}.$$

By the last equality of (3.7), it shows the function x^{α} is real analytic and non-degenerate in $D_{2R}(x_0, 0) \subset I \times \mathbb{R}$. By the real analytic estimates of solutions to linear elliptic equations with real analytic coefficients (cf. for instance, [2], Chapter 3 in [27], and Chapter 5 in [30]), we have that there are constants $C = C(R, \alpha) \geq 1$ and $\rho = \rho(R, \alpha) \in (0, 1)$ such that any solutions to equation (3.6) satisfies,

$$\|\partial_x^a \partial_t^\gamma y(\cdot, \cdot, t)\|_{L^{\infty}(D_R(x_0, 0))} \le \frac{Ca!}{\rho^a} \left(\int_{D_{2R}(x_0, 0)} |\partial_t^\gamma y(x, \tau, t)|^2 dx d\tau \right)^{\frac{1}{2}},$$
(3.8)

where $a \in \mathbb{N}, \gamma \in \mathbb{N}$. Notice that for each t > 0,

$$\int_{D_{2R}(x_0,0)} |\partial_t^{\gamma} y(x,\tau,t)|^2 dx d\tau \leq \int_{-2R}^{2R} \int_{D_{2R}(x_0)} |\partial_t^{\gamma} y(x,\tau,t)|^2 dx d\tau$$

$$\leq \int_{-2R}^{2R} \int_I |\partial_t^{\gamma} y(x,\tau,t)|^2 dx d\tau.$$
(3.9)

By the orthogonality of $\{e_k\}_{k\geq 1}$ in $L^2(I)$, and (3.5), we have

$$\int_{I} |\partial_{t}^{\gamma} y(x,\tau,t)|^{2} dx = \int_{I} \left| \sum_{k \ge 1} a_{k} (-w_{k}^{2})^{\gamma} e^{-w_{k}^{2}t + w_{k}\tau} e_{k}(x) \right|^{2} dx$$

$$= \sum_{k \ge 1} a_{k}^{2} w_{k}^{4\gamma} e^{-2w_{k}^{2}t + 2w_{k}\tau}$$

$$= \sum_{k \ge 1} a_{k}^{2} w_{k}^{4\gamma} e^{-w_{k}^{2}t} e^{-w_{k}^{2}t + 2w_{k}\tau}$$

$$\leq \max_{k \ge 1} \{ w_{k}^{4\gamma} e^{-w_{k}^{2}t} \} \max_{k \ge 1} \{ e^{-w_{k}^{2}t + 2w_{k}\tau} \} \sum_{k > 1} a_{k}^{2}$$

which, along with (3.9), it implies

$$\int_{D_{2R}(x_0,0)} |\partial_t^{\gamma} y(x,\tau,t)|^2 dx d\tau \le \max_{k\ge 1} \{ w_k^{4\gamma} e^{-w_k^2 t} \} \max_{k\ge 1} \{ e^{-w_k^2 t + 4Rw_k} \} \sum_{k\ge 1} a_k^2.$$
(3.10)

On one hand, from Stirling's formula, see page 111 in [28], i.e. for a absolute constant C, $\gamma^{\gamma} \leq Ce^{\gamma}(\gamma)!$, we have

$$\max_{k \ge 1} \{ w_k^{4\gamma} e^{-w_k^2 t} \} \le C \left(\frac{2}{t}\right)^{2\gamma} (\gamma!)^2.$$

On the other hand, we have

$$\max_{k \ge 1} \{ e^{-w_k^2 t + 4Rw_k} \} = e^{\frac{4R^2}{t}}.$$

Therefore, it stands

$$\int_{D_{2R}(x_0,0)} |\partial_t^{\gamma} y(x,\tau,t)|^2 dx d\tau \le C \left(\frac{2}{t}\right)^{2\gamma} (\gamma!)^2 e^{\frac{4R^2}{t}} \sum_{k\ge 1} a_k^2.$$
(3.11)

This together with (3.8), it implies

$$\|\partial_x^a \partial_t^\gamma y(\cdot, t)\|_{L^{\infty}(D_R(x_0, 0))} \leq \frac{Ce^{\frac{2R^2}{t}}a!\gamma!}{\rho^a(t/2)^{\gamma}}\|y_0\|,$$

which, along with the first result of (3.7), we have

$$\|\partial_x^a \partial_t^\gamma y(\cdot, t)\|_{L^{\infty}(\omega)} \le \frac{Ce^{\frac{2R^2}{t}}a!\gamma!}{\rho^a(t/2)^{\gamma}}\|y_0\|$$

Hence, the above inequality implies that (3.2) with s = 0 holds. The proof is completed.

The subsequent two lemmas focus on the propagation of smallness estimates from measurable sets for real analytic functions. This type of observability estimate was initially established in [34] (also refer to [2] and [3] for associated findings). To facilitate subsequent references, one lemma pertains to the one-dimensional scenario, while the other lemma caters to the multidimensional case. For a more streamlined proof, interested readers can refer to Section 3 of [3].

Lemma 3.3. Let $g : [a, a + s] \to \mathbb{R}$, where $a \in \mathbb{R}$ and s > 0 be a real analytic function satisfying

$$\left|\frac{d^k}{dx^k}g(x)\right| \le Mk!(s\rho)^{-k}, \forall x \in [a, a+s], \ \forall k \in \mathbb{N}.$$

where M > 0, $\rho \in (0,1]$. Assume that $F \subset [a, a + s]$ is a measurable subset of positive measure. Then there are $C = C(\rho, |F|/s)$ and $\theta = \theta(\rho, |F|/s)$, $0 < \theta < 1$, such that

$$||g||_{L^{\infty}(a,a+s)} \leq CM^{1-\theta} \left(\frac{1}{|F|} \int_{F} |g(x)| dx\right)^{\theta}.$$

Lemma 3.4. Let ω be a bounded domain in \mathbb{R}^n , $n \ge 1$ and $\bar{\omega} \subset \omega$ be a measurable set of positive measure. Let f be an analytic function in ω satisfying

$$|\partial_x^a f(x)| \le M \rho^{-|a|} |a|!, \ \forall x \in \omega, \ \forall a \in \mathbb{N}^n,$$

where M > 0, $\rho \in (0, 1]$. Then there are $C = C(|\omega|, \rho, |\bar{\omega}|)$ and $\theta = \theta(|\omega|, \rho, |\bar{\omega}|)$, $0 < \theta < 1$, such that

$$||f||_{L^{\infty}(\omega)} \le CM^{1-\theta} \left(\int_{\bar{\omega}} |f(x)| dx \right)^{\theta}.$$

Next, we shall make use of Lemma 3.1 and Lemma 3.2, as well as Lemma 3.3 and Lemma 3.4, to establish an interpolation inequality from measurable sets for any solution y to equation (2.2). For similar results, we refer the reader to [3] or [29].

Theorem 3.5. Let $0 \le t_1 < t_2 < 1$, $\eta \in (0,1)$, $\mu \in (0,1)$ defined in (2.3), $\alpha \in (0,2)$ and $\sigma > 0$. Assume that $E \subset (t_1, t_2)$ is a measurable subset with $|E \cap (t_1, t_2)| \ge \eta(t_2 - t_1)$ such that for all $t \in E$, the measurable subset $D_t \subset \omega$ stands that $|D_t| \ge \sigma$. Then, there are constants $C = C(I, \omega, \alpha, \eta, \sigma, \mu) \ge 1$ and $\theta = \theta(I, \omega, \alpha, \eta, \sigma, \mu) \in (0,1)$ such that the solution to equation (2.2) satisfies

$$\|y(\cdot,t_2)\| \le \left(\int_{t_1}^{t_2} \chi_E \|y(\cdot,t)\|_{L^1(D_t)} dt\right)^{\theta} \left(e^{C(t_2-t_1)^{-\frac{\mu}{1-\mu}}} \|y(\cdot,t_1)\|\right)^{1-\theta}.$$
 (3.12)

Proof. Define

$$\tau = t_1 + \frac{\eta}{3}(t_2 - t_1), \ F = E \cap (\tau, t_2).$$

One can verify that $|F| > \frac{\eta}{2}(t_2 - t_1)$. This shows that $F \subset [\tau, t_2]$ is a measurable subset of positive measure.

From Lemma 3.2, we have that there are constants $C = C(I, \omega, \alpha, \eta, \sigma) \ge 1$ and $\theta = \theta(I, \omega, \alpha, \eta, \sigma) \in (0, 1)$ such that for all $t \in [\tau, t_2]$ and for all $x \in \omega, a \in \mathbb{N}, \gamma \in \mathbb{N}$,

$$|\partial_x^a \partial_t^\gamma y(\cdot, t)| \le \frac{e^{\frac{C}{t_2 - t_1}} a! \gamma!}{\rho^a (\eta(t_2 - t_1)/2)^\gamma} \|y(\cdot, t_1)\|.$$
(3.13)

Now, for simplicity, we write

$$M = e^{\frac{C}{t_2 - t_1}} \|y(\cdot, t_1)\|.$$
(3.14)

Then for all $x \in \omega$ and $\gamma \in \mathbb{N}$, by (3.13) with a = 0, we have

$$\left|\partial_t^{\gamma} y(\cdot, t)\right| \le M\gamma! \left(\frac{\eta(t_2 - t_1)}{2}\right)^{-\gamma} \le M\gamma! \left(\frac{\eta(t_2 - \tau)}{6}\right)^{-\gamma}, \text{ for all } t \in [\tau, t_2].$$
(3.15)

Hence, it follows from Lemma 3.3 that there are constants $C = C(\eta) \ge 1$ and $\theta = \theta(\eta) \in (0, 1)$ such that for all $x \in \omega$,

$$\|y(x,\cdot)\|_{L^{\infty}(\tau,t_{2})} \le CM^{1-\theta} \left(\frac{1}{|F|} \int_{F} |y(x,t)| dt\right)^{\theta}.$$
(3.16)

On the other hand, we obtain from Lemma 3.1 that there exists a constant $C = C(I, \omega, \alpha, \mu)$ such that the following observability inequality holds:

$$\int_{I} |y(x,t_2)|^2 dx \le e^{C(t_2-\tau)^{-\frac{\mu}{1-\mu}}} \int_{\tau}^{t_2} \int_{\omega} |y(x,t)|^2 dx dt.$$
(3.17)

By (3.13) with $a = \gamma = 0$, we have

$$\|y\|_{L^{2}(\tau,t_{2};L^{2}(\omega))} \leq \|y\|_{L^{\infty}(\tau,t_{2};L^{\infty}(\omega))}^{\frac{1}{2}} \|y\|_{L^{1}(\tau,t_{2};L^{1}(\omega))}^{\frac{1}{2}} \leq e^{\frac{C}{t_{2}-t_{1}}} \|y(\cdot,t_{1})\|^{\frac{1}{2}} \|y\|_{L^{1}(\tau,t_{2};L^{1}(\omega))}^{\frac{1}{2}},$$

which along with (3.17), it holds

$$\|y(\cdot,t_{2})\| \leq \|y\|_{L^{1}(\tau,t_{2};L^{1}(\omega))}^{\frac{1}{2}} e^{C(t_{2}-\tau)^{-\frac{\mu}{1-\mu}}} e^{\frac{C}{t_{2}-t_{1}}} \|y(\cdot,t_{1})\|^{\frac{1}{2}} \leq \|y\|_{L^{1}(\tau,t_{2};L^{1}(\omega))}^{\frac{1}{2}} \left(e^{C(t_{2}-t_{1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot,t_{1})\|\right)^{\frac{1}{2}}.$$
(3.18)

It follows by (3.16) that for any $t \in F$,

$$\begin{aligned} \|y\|_{L^{1}(\tau,t_{2};L^{1}(\omega))} &\leq (t_{2}-\tau) \int_{\omega} \|y(x,\cdot)\|_{L^{\infty}(\tau,t_{2})} dx \leq CM^{1-\theta} \int_{\omega} \left(\int_{F} |y(x,t)| dt \right)^{\theta} dx \\ &\leq CM^{1-\theta} \left(\int_{\omega} \int_{F} |y(x,t)| dt dx \right)^{\theta}. \end{aligned}$$

$$(3.19)$$

Finally, by (3.13) with $\gamma = 0$, we infer that for any $t \in F$,

$$|\partial_x^a y(\cdot, t)| \le M a! \rho^{-a}$$
, for all $x \in \omega$.

Since the measurable subset $D_t \subset \omega$ stands that $|D_t| \geq \sigma$, when $t \in F$, we obtain from Lemma 3.4 (with n = 1) that there are constants $C_1 = C_1(\omega, \rho, \sigma) \geq 1$ and $\theta_1 = \theta_1(\omega, \rho, \sigma) \in (0, 1)$ such that

$$\|y(\cdot,t)\|_{L^{\infty}(\omega)} \leq C_1 M^{1-\theta_1} \left(\int_{D_t} |y(x,t)| dx \right)^{\theta_1},$$

which along with (3.19), it implies

$$||y||_{L^{1}(\tau,t_{2};L^{1}(\omega))} \leq CM^{1-\theta} \left[\int_{F} C_{1}M^{1-\theta_{1}} \left(\int_{D_{t}} |y(x,t)| dx \right)^{\theta_{1}} dt \right]^{\theta}$$

$$\leq CC_{1}M^{1-\theta\theta_{1}} \left(\int_{F} \int_{D_{t}} |y(x,t)| dx dt \right)^{\theta\theta_{1}}.$$
(3.20)

Therefore, by (3.18), it implies

$$\|y(\cdot,t_2)\| \le CC_1 M^{1-\theta\theta_1} \left(\int_F \int_{D_t} |y(x,t)| dx dt \right)^{\theta\theta_1} \left(e^{C(t_2-t_1)^{-\frac{\mu}{1-\mu}}} \|y(\cdot,t_1)\| \right)^{\frac{1}{2}}.$$

Set $\theta' = \theta \theta_1 \in (0, 1)$ and by (3.14), we obtain the desired estimate (3.12).

By leveraging Fubini's theorem and the property of Lebesgue density points for measurable subsets, along with the interpolation inequality established in Theorem 3.5, a telescoping series can be constructed with respect to any solution y of (2.2). Subsequently, the desired observability inequality from measurable subsets can be derived directly from this telescoping series (also see [3] and [31]). To begin, we quote the following fact from [31], which pertains to the property of Lebesgue density points and represents an improved version of Lemma 2.1.5 in [21].

Lemma 3.6. Let $E \subset (0,T)$ be a measurable subset of positive measure. Assume that $l \in (0,T)$ be a Lebesgue density point of E. Then, for each $q \in (0,1)$, there exists a sequence $\{\ell_n\}_{n\geq 1} \subset [0,T]$, which monotone decreasing converges to l, such that for any $n \geq 1$,

$$\ell_{n+1} - \ell_{n+2} = q(\ell_n - \ell_{n+1}), \tag{3.21}$$

and

$$|E \cap (\ell_{n+1}, \ell_n)| \ge \frac{\ell_n - \ell_{n+1}}{3}.$$
(3.22)

Now, we are prepared to present the proof of Theorem 2.1 by employing the techniques outlined in Theorem 3.5 and Lemma 3.6, in addition to the telescoping series method.

Proof of Theorem 2.1. Firstly, we can make the assumption, without loss of generality, that $T \in (0, 1]$ and that there exists a subdomain $\omega \subset I$ such that $0 \notin \bar{\omega}$. (Alternatively, we can select a new measurable subset $\tilde{D} \subset (I \setminus \{0\}) \times (T_1, T_2)$ with $0 \leq T_1 < T_2 \leq T$ and $T_2 - T_1 \leq 1$.) Next, for a.e. $t \in (0, T)$, we slice the space

$$D_t = \{ x \in \omega : (x, t) \in D \}.$$
 (3.23)

Define

$$E = \left\{ t \in (0,T) : |D_t| \ge \frac{|D|}{2T} \right\}.$$
 (3.24)

From the Fubini theorem, we have

$$|D| = \int_0^T |D_t| dt = \int_E |D_t| dt + \int_{(0,T)\setminus E} |D_t| dt \le |E| |\omega| + \frac{|D|}{2},$$

which shows $|E| \ge \frac{|D|}{2|\omega|}$, i.e. E is a measurable subset with positive measures. Moreover, we have

$$\chi_E(t)\chi_{D_t}(x) \le \chi_D(x,t), \ a.e.\ (x,t) \in I \times (0,T).$$
(3.25)

Let $l \in (0, T)$ be a Lebesgue density point of E. Then for each $q \in (0, 1)$, which is to be fixed later, by Lemma 3.6, there exists a monotone decreasing sequence $\{\ell_n\}_{n\geq 1}$ such that (3.21) and (3.22) hold. Moreover,

$$\lim_{n \to +\infty} l_n = l. \tag{3.26}$$

We can obtain from Theorem 3.5 that there are positive constants $C = C(I, \omega, \alpha, \eta, \sigma, \mu) \ge 1$ and $\theta = \theta(I, \omega, \alpha, \eta, \sigma, \mu) \in (0, 1)$ such that for all $n \ge 1$,

$$\|y(\cdot, l_n)\| \le \left(\int_{l_{n+1}}^{l_n} \chi_E \|y(\cdot, t)\|_{L^1(D_t)} dt\right)^{\theta} \left(e^{C(l_n - l_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot, l_{n+1})\|\right)^{1-\theta}.$$
 (3.27)

Using the Young inequality with ϵ , i.e., $ab \leq \epsilon a^{(1-\theta)^{-1}} + \epsilon^{-\frac{1-\theta}{\theta}} b^{\theta^{-1}}$, the above inequality leads to

$$\|y(\cdot, l_n)\| \le \epsilon \|y(\cdot, l_{n+1})\| + \epsilon^{-\frac{1-\theta}{\theta}} e^{C(l_n - l_{n+1})^{-\frac{\mu}{1-\mu}}} \int_{l_{n+1}}^{l_n} \chi_E \|y(\cdot, t)\|_{L^1(D_t)} dt.$$
(3.28)

Multiplying the above inequality by $\epsilon^{\frac{1-\theta}{\theta}}e^{-C(\ell_n-\ell_{n+1})^{-\frac{\mu}{1-\mu}}}$, we have

$$\epsilon^{\frac{1-\theta}{\theta}} e^{-C(\ell_n - \ell_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot, l_n)\| - \epsilon^{\frac{1}{\theta}} e^{-C(\ell_n - \ell_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot, l_{n+1})\| \\ \leq \int_{l_{n+1}}^{l_n} \chi_E \|y(\cdot, t)\|_{L^1(D_t)} dt.$$

Choosing $\epsilon = e^{-\theta(\ell_n - \ell_{n+1})^{-\frac{\mu}{1-\mu}}}$ in the above inequality, we get

$$e^{-(C+1-\theta)(\ell_n-\ell_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot,l_n)\| - e^{-(C+1)(\ell_n-\ell_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot,l_{n+1})\| \\ \leq \int_{l_{n+1}}^{l_n} \chi_E \|y(\cdot,t)\|_{L^1(D_t)} dt.$$

Now, fixing $q = \left(\frac{C+1-\theta}{C+1}\right)^{\frac{1-\mu}{\mu}} \in (0,1)$, it follows from (3.21) and the latter inequality that

$$e^{-(C+1-\theta)(\ell_n-\ell_{n+1})^{-\frac{\mu}{1-\mu}}} \|y(\cdot,l_n)\| - e^{-(C+1-\theta)(\ell_{n+1}-\ell_{n+2})^{-\frac{\mu}{1-\mu}}} \|y(\cdot,l_{n+1})\|$$

$$\leq \int_{l_{n+1}}^{l_n} \chi_E \|y(\cdot,t)\|_{L^1(D_t)} dt.$$

To summarize the inequalities earlier from n = 1 to $+\infty$ and noting that (3.26), we conclude that

$$\|y(\cdot, l_1)\| \le e^{(C+1-\theta)(\ell_1-\ell_2)^{-\frac{\mu}{1-\mu}}} \int_l^{l_1} \chi_E(t) \|y(\cdot, t)\|_{L^1(D_t)} dt$$

which along with (3.25) and $||y(\cdot,T)|| \leq C(T)||y(\cdot,l_1)||$, we obtain the desired estimate (2.4). The proof is completed.

4. A Stackelberg-Nash game problem

In this section, we shall discuss the Stackelberg-Nash game problem (P1).

4.1. Existence of Stackelberg-Nash equilibrium. In this subsection, we will prove Theorem 2.2. Its proof needs the Kakutani fixed point theorem quoted from [15].

Lemma 4.1. Let S be a nonempty, compact and convex subset of a locally convex Hausdorff space X. Let $\Phi: S \mapsto 2^S$ (where 2^S denotes the set consisting of all subsets of S) be a set-valued function satisfying:

(i) For each $s \in S$, $\Phi(s)$ is a nonempty and convex subset;

(ii) Graph $\Phi := \{(s, z) : s \in S \text{ and } z \in \Phi(s)\}$ is closed.

Then the set of fixed points of Φ is nonempty and compact, where $s^* \in S$ is called to be a fixed point of Φ if $s^* \in \Phi(s^*)$.

Now, we are in a position to prove Theorem 2.2.

Proof of Theorem 2.2. We first introduce three set-valued functions $\Phi_1 : \mathcal{U}_1 \mapsto 2^{\mathcal{U}_2}, \Phi_2 : \mathcal{U}_2 \mapsto 2^{\mathcal{U}_1} \text{ and } \Phi : \mathcal{U}_1 \times \mathcal{U}_2 \mapsto 2^{\mathcal{U}_1 \times \mathcal{U}_2}$ as follows:

$$\Phi_1 u_1 := \{ u_2 \in \mathcal{U}_2 : J_2(u_1, u_2) \le J_2(u_1, v_2) \text{ for all } v_2 \in \mathcal{U}_2 \}, \ u_1 \in \mathcal{U}_1,$$

$$(4.1)$$

$$\Phi_2 u_2 := \{ u_1 \in \mathcal{U}_1 : J_1(u_1, u_2) \le J_1(v_1, u_2) \text{ for all } v_1 \in \mathcal{U}_1 \}, \ u_2 \in \mathcal{U}_2,$$
(4.2)

and

$$\Phi(u_1, u_2) := \{ (\widetilde{u}_1, \widetilde{u}_2) : \widetilde{u}_1 \in \Phi_2 u_2 \text{ and } \widetilde{u}_2 \in \Phi_1 u_1 \}, \ (u_1, u_2) \in \mathcal{U}_1 \times \mathcal{U}_2.$$
(4.3)

Then we set

 $X := (L^{\infty}(0,T;L^{2}(0,1)))^{2}$ and $S := \mathcal{U}_{1} \times \mathcal{U}_{2}.$

It is clear that X is a locally convex Hausdorff space. The rest of the proof will be carried out by the following four steps.

Step 1. Show that S is a nonempty, compact and convex subset of X.

This fact can be easily checked. We omit the proof here.

Step 2. $\Phi(u_1, u_2)$ is nonempty. We prove that $\Phi(u_1, u_2)$ is nonempty for each $(u_1, u_2) \in S$.

We arbitrarily fix $(u_1, u_2) \in S$. According to (4.1)-(4.3), it suffices to show that $\Phi_1 u_1$ and $\Phi_2 u_2$ are nonempty. For this purpose, we introduce the following auxiliary optimal control problem:

$$(\mathbf{P}_{\mathbf{a}\mathbf{u}}) \qquad \inf_{v_2 \in \mathcal{U}_2} J_2(u_1, v_2).$$

Let

$$d := \inf_{v_2 \in \mathcal{U}_2} J_2(u_1, v_2). \tag{4.4}$$

It is obvious that $d \ge 0$. Let $\{v_{2,n}\}_{n\ge 1} \subseteq \mathcal{U}_2$ be a minimizing sequence so that

$$d = \lim_{n \to \infty} J_2(u_1, v_{2,n}). \tag{4.5}$$

On one hand, since $||v_{2,n}||_{L^{\infty}(0,T;L^2(0,1))} \leq M_2$, there exists a subsequence of $\{v_{2,n}\}_{n\geq 1}$, still denoted by itself, and $v_{2,0} \in \mathcal{U}_2$, so that

$$v_{2,n} \to v_{2,0}$$
 weakly star in $L^{\infty}(0,T;L^2(0,1)).$ (4.6)

We denote by $\bar{y}_n(\cdot) := y(\cdot; y_0, g, u_1, v_{2,n}) - y(\cdot; y_0, g, u_1, v_{2,0})$. According to (2.5), it is clear that

$$\begin{cases} \bar{y}_{n,t}(x,t) = A\bar{y}_n(x,t) + \chi_{\omega_2}(v_{2,n} - v_{2,0}), & (x,t) \in (0,1) \times (0,T), \\ \bar{y}_n(1,t) = BC_\alpha(\bar{y}_n(\cdot,t)) = 0, & t \in (0,T), \\ \bar{y}_n(x,0) = 0, & x \in (0,1). \end{cases}$$
(4.7)

Multiplying the first equation of (4.7) by \bar{y}_n and integrating over (0, 1), one deduces, for all $t \leq T$,

$$\frac{1}{2}\frac{d}{dt}\|\bar{y}_n(t)\|^2 + \|x^{\frac{\alpha}{2}}\bar{y}_{n,x}(t)\|^2 \le \frac{1}{2}\|\bar{y}_n(t)\|^2 + \frac{1}{2}\|(v_{2,n} - v_{2,0})(t)\|^2_{L^2(\omega_2)}.$$
(4.8)

By Gronwall's inequality, we have

$$\sup_{t \in [0,T]} \|\bar{y}_n(t)\|^2 \le e^T \|v_{2,n} - v_{2,0}\|_{L^2(0,T;L^2(\omega_2))}^2.$$
(4.9)

Notice that, from (4.8) and (4.9), it also follows that

$$\|x^{\frac{\alpha}{2}}\bar{y}_{n,x}\|_{L^{2}(0,T;L^{2}(0,1))}^{2} \leq \left[\frac{e^{T}T}{2} + \frac{1}{2}\right] \|v_{2,n} - v_{2,0}\|_{L^{2}(0,T;L^{2}(\omega_{2}))}^{2}.$$
 (4.10)

From these, we obtain

$$\sup_{t \in [0,T]} \left\| \bar{y}_n(t) \right\|^2 + \left\| x^{\frac{\alpha}{2}} \bar{y}_{n,x} \right\|_{L^2(0,T;L^2(0,1))}^2 \le C \left\| v_{2,n} - v_{2,0} \right\|_{L^2(0,T;L^2(\omega_2))}^2, \tag{4.11}$$

where C = C(T) independent of n.

It follows from (4.6) and (4.11) that there exists a subsequence of \bar{y}_n , still denoted by itself, and $\bar{y} \in C([0,T]; L^2(0,1)) \cap L^2(0,T; H^1_{\alpha}(0,1))$ such that

$$\bar{y}_n \to \bar{y}$$
 weakly in $C([0,T]; L^2(0,1)) \cap L^2(0,T; H^1_a(0,1)).$ (4.12)

Since the embedding $H^1_{\alpha}(0,1) \hookrightarrow L^2(0,1)$ is compact, see [33], also see section 6 in [1], it implies that

$$\bar{y}_n \to \bar{y}$$
 strongly in $L^2(0,1)$. (4.13)

Passing to the limit for $n \to \infty$ in (4.7), by (4.6) and (4.12), we have

$$\begin{cases} \bar{y}_t(x,t) = A\bar{y}(x,t), & (x,t) \in (0,1) \times (0,T) \\ \bar{y}(1,t) = BC_\alpha \left(\bar{y}(\cdot,t) \right) = 0, & t \in (0,T), \\ \bar{y}(x,0) = 0, & x \in (0,1), \end{cases}$$

from which we obtain that $\bar{y} = 0$. Hence, on the other hand, we have

$$y(T; y_0, g, u_1, v_{2,n}) \to y(T; y_0, g, u_1, v_{2,0})$$
 strongly in $L^2(0, 1)$. (4.14)

It follows from (4.5), (2.6), (4.14) that

$$d = J_2(u_1, v_{2,0}). (4.15)$$

Noting that $v_{2,0} \in \mathcal{U}_2$, by (4.4), (4.15) and (4.1), we obtain that $v_{2,0} \in \Phi_1 u_1$. This implies that $\Phi_1 u_1 \neq \emptyset$. In the same way, we also have that $\Phi_2 u_2 \neq \emptyset$.

Step 3. Convex subset $\Phi(u_1, u_2)$. We show that $\Phi(u_1, u_2)$ is a convex subset of $\mathcal{U}_1 \times \mathcal{U}_2$ for each $(u_1, u_2) \in \mathcal{U}_1 \times \mathcal{U}_2$.

We arbitrarily fix $(u_1, u_2) \in \mathcal{U}_1 \times \mathcal{U}_2$. According to (4.1)-(4.3), it suffices to prove that $\Phi_1 u_1$ is a convex subset of \mathcal{U}_2 . The convexity of $\Phi_2 u_2$ can be similarly proved. For this purpose, we arbitrarily fix $\tilde{u}_2, \hat{u}_2 \in \Phi_1 u_1$. By (4.1), we get that

$$\widetilde{u}_2, \ \widehat{u}_2 \in \mathcal{U}_2,$$

$$(4.16)$$

and

$$J_2(u_1, \tilde{u}_2) \le J_2(u_1, v_2)$$
 and $J_2(u_1, \tilde{u}_2) \le J_2(u_1, v_2)$ for each $v_2 \in \mathcal{U}_2$. (4.17)

For any $\lambda \in [0, 1]$, by (2.6) and (2.5), we have that

$$\begin{aligned} J_2(u_1,\lambda\widetilde{u}_2+(1-\lambda)\widehat{u}_2) &- [\lambda J_2(u_1,\widetilde{u}_2)+(1-\lambda)J_2(u_1,\widehat{u}_2)] \\ = & \|y(T;y_0,u_1,\lambda\widetilde{u}_2+(1-\lambda)\widehat{u}_2)-y_T^2\|_{L^2(G_2)} - \lambda \|y(T;y_0,u_1,\widetilde{u}_2)-y_T^2\|_{L^2(G_2)} \\ &- (1-\lambda)\|y(T;y_0,u_1,\widehat{u}_2)-y_T^2\|_{L^2(G_2)} \\ = & \|\lambda[y(T;y_0,u_1,\widetilde{u}_2)-y_T^2]+(1-\lambda)[y(T;y_0,u_1,\widehat{u}_2)-y_T^2]\|_{L^2(G_2)} \\ &- \lambda \|y(T;y_0,u_1,\widetilde{u}_2)-y_T^2\|_{L^2(G_2)} - (1-\lambda)\|y(T;y_0,u_1,\widehat{u}_2)-y_T^2\|_{L^2(G_2)} \\ \leq & 0 \end{aligned}$$

This, along with (4.16) and (4.17), yields that

$$\lambda \widetilde{u}_2 + (1-\lambda)\widehat{u}_2 \in \mathcal{U}_2$$

and

$$J_2(u_1, \lambda \widetilde{u}_2 + (1-\lambda)\widehat{u}_2) \leq J_2(u_1, v_2)$$
 for each $v_2 \in \mathcal{U}_2$,

which indicate that $\lambda \tilde{u}_2 + (1 - \lambda) \hat{u}_2 \in \Phi_1 u_1$ (see (4.1)). Hence, $\Phi_1 u_1$ is a convex subset of \mathcal{U}_2 .

Step 4. Prove that Graph Φ is closed.

It suffices to show that if $(u_{n,1}, u_{n,2}) \in \mathcal{U}_1 \times \mathcal{U}_2, \widetilde{u}_{n,1} \in \Phi_2 u_{n,2}, \ \widetilde{u}_{n,2} \in \Phi_1 u_{n,1}, (u_{n,1}, u_{n,2}) \to (u_1, u_2)$ in X and $(\widetilde{u}_{n,1}, \widetilde{u}_{n,2}) \to (\widetilde{u}_1, \widetilde{u}_2)$ in X, then

$$(u_1, u_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \ \widetilde{u}_1 \in \Phi_2 u_2 \text{ and } \widetilde{u}_2 \in \Phi_1 u_1.$$
 (4.18)

Indeed, on one hand, by (4.1) and (4.2), we can easily check that

 $(u_1, u_2) \in \mathcal{U}_1 \times \mathcal{U}_2, \ \widetilde{u}_1 \in \mathcal{U}_1 \text{ and } \widetilde{u}_2 \in \mathcal{U}_2.$ (4.19)

On the other hand, according to $\tilde{u}_{n,1} \in \Phi_2 u_{n,2}$, (4.2) and (2.6), it is obvious that for each $v_1 \in \mathcal{U}_1$,

$$\|y(T; y_0, g, \widetilde{u}_{n,1}, u_{n,2}) - y_T^1\|_{L^2(G_1)} \le \|y(T; y_0, g, v_1, u_{n,2}) - y_T^1\|_{L^2(G_1)}.$$
(4.20)

Since $(\tilde{u}_{n,1}, u_{n,2}) \to (\tilde{u}_1, u_2)$ weakly star in $(L^{\infty}(0, T; L^2(0, 1)))^2$, by similar arguments as those to get (4.14), there exists a subsequence of $\{n\}_{n>1}$, still denoted by itself, so that

$$(y(T; y_0, g, \widetilde{u}_{n,1}, u_{n,2}), y(T; y_0, g, v_1, u_{n,2}))$$

 $\rightarrow (y(T; y_0, g, \widetilde{u}_1, u_2), y(T; y_0, g, v_1, u_2))$ strongly in $(L^2(0, 1))^2$,

which, implies that

 $(y(T; y_0, g, \tilde{u}_{n,1}, u_{n,2}), y(T; y_0, g, v_1, u_{n,2}))$ (4.21)

$$\rightarrow (y(T; y_0, g, \widetilde{u}_1, u_2), y(T; y_0, g, v_1, u_2))$$
 strongly in $(L^2(G_1))^2$.

Passing to the limit for $n \to \infty$ in (4.20), by (4.21), we get that for each $v_1 \in \mathcal{U}_1$,

$$||y(T; y_0, g, \widetilde{u}_1, u_2) - y_T^1||_{L^2(G_1)} \le ||y(T; y_0, g, v_1, u_2) - y_T^1||_{L^2(G_1)}.$$

This, together with (2.6), (4.2) and the second conclusion in (4.19), implies that $\tilde{u}_1 \in \Phi_2 u_2$. Similarly, $\tilde{u}_2 \in \Phi_1 u_1$. Hence, (4.18) follows.

Step 5. Finish the proof.

According to Steps 1-4 and Lemma 4.1, there exists a pair of $(u_1^*, u_2^*) \in \mathcal{U}_1 \times \mathcal{U}_2$ so that $(u_1^*, u_2^*) \in \Phi(u_1^*, u_2^*)$, which, combined with (4.1)-(4.3), indicates that (u_1^*, u_2^*) is a Stackelberg-Nash equilibrium of the problem **(P1)**.

In summary, we end the proof of Theorem 2.2.

4.2. Decomposition and Characterization of Stackelberg-Nash Equilibria. In this subsection, we first give some conclusions that will be used later. By Theorem 2.1, we have the observability inequality for adjoint equation (2.11).

Corollary 4.2. Let $\alpha \in (0, 2)$, $\mu \in (0, 1)$ and $\omega \times E \subset I \times (0, T)$ be measurable subsets with positive measures. Then there exists a constant $C = C(T, I, \alpha, \omega, E, \mu) \ge 1$ such that the solution of (2.11) satisfies the following observability inequality: for any $z_T \in L^2(0, 1)$,

$$||z(0)|| \le C \int_E ||\chi_\omega z(t)|| dt.$$
 (4.22)

Lemma 4.3. Let $g \in L^{\infty}(0,T; L^2(0,1))$ be given. Assume ω, ω_0 are measurable subsets of (0,1) with positive measures. Let $\omega_0 \subset G$, where G is a measurable subset of (0,1) and $\zeta(\cdot; \zeta_0, g, u)$ be the solution to the following system:

$$\begin{cases} \zeta_t(x,t) = A\zeta(x,t) + \chi_{\omega}g + \chi_{\omega_0}u, \quad (x,t) \in (0,1) \times (0,T), \\ \zeta(1,t) = BC_{\alpha}(\zeta(\cdot,t)) = 0, \qquad t \in (0,T), \\ \zeta(x,0) = \zeta_0(x), \qquad x \in (0,1), \end{cases}$$
(4.23)

where $u \in \mathcal{U}$ with

$$\mathcal{U} := \left\{ u \in L^{\infty}(0,T; L^{2}(0,1)) : \|u\|_{L^{\infty}(0,T; L^{2}(0,1))} \le M \right\},\$$

with M > 0. Consider the following problem:

$$\inf_{u \in \mathcal{U}} \|\zeta(T;\zeta_0, g, u) - \zeta_T\|_{L^2(G)},\tag{4.24}$$

where $\zeta_T \in L^2(0,1)$ is a given function. Then there exists $u^* \in \mathcal{U}$ such that

$$\|\zeta(T;\zeta_0,g,u^*) - \zeta_T\|_{L^2(G)} = \inf_{u \in \mathcal{U}} \|\zeta(T;\zeta_0,g,u) - \zeta_T\|_{L^2(G)}.$$

Moreover, if $\zeta(T;\zeta_0,g,u^*) \neq \zeta_T$, then u^* enjoys the following form

$$u^{*}(t) = M \frac{\chi_{\omega_{0}} z(t)}{\|\chi_{\omega_{0}} z(t)\|_{L^{2}(G)}}, \text{ for a.e. } t \in (0,T).$$

$$(4.25)$$

where z is a solution solving the following equation:

$$\begin{cases} z_t(x,t) + Az(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z(1,t) = BC_\alpha(z(\cdot,t)) = 0, & t \in (0,T), \\ z(x,T) = \zeta_T - \zeta(T;\zeta_0, g, u^*), & x \in (0,1). \end{cases}$$
(4.26)

Proof. We carry out this proof step by step.

Step 1. We show the existence of the problem (4.24).

Note that $\zeta(\cdot; \zeta_0, g, u) = \zeta(\cdot; \zeta_0, g, 0) + \zeta(\cdot; 0, 0, u)$, then the problem (4.24) is equivalent to the following problem:

$$\inf_{u \in \mathcal{U}} \|\zeta(T; 0, 0, u) - (\zeta_T - \zeta(T; \zeta_0, g, 0))\|_{L^2(G)}.$$

Let $\{u_n\}_{n\geq 1} \subseteq \mathcal{U}$ be the minimal sequence of problem (4.24), i.e.,

$$\|\zeta(T;0,0,u_n) - (\zeta_T - \zeta(T;\zeta_0,g,0))\|_{L^2(G)} \to \inf_{u \in \mathcal{U}} \|\zeta(T;0,0,u) - (\zeta_T - \zeta(T;\zeta_0,g,0))\|_{L^2(G)}.$$
(4.27)

Since $\{u_n\}_{n\geq 1} \subseteq \mathcal{U}$, there exists a subsequence of $\{u_n\}_{n\geq 1}$, still denoted by itself, and $u^* \in L^{\infty}(0,T;L^2(0,1))$ such that

$$u_n \to u^*$$
 weakly star in $L^{\infty}(0,T;L^2(0,1))$.

Note that $\zeta(\cdot; 0, 0, u_n)$ is the solution of the following system

$$\begin{cases} \zeta(t;0,0,u_n)_t(x) = A\zeta(t;0,0,u_n)(x) + \chi_{\omega_0}u_n, \quad (x,t) \in (0,1) \times (0,T) \\ \zeta(t;0,0,u_n)(1) = BC_\alpha(\zeta(t;0,0,u_n)(\cdot)) = 0, \qquad t \in (0,T), \\ \zeta(0;0,0,u_n)(x) = 0, \qquad x \in (0,1). \end{cases}$$

By similar arguments as those to get (4.14), there exists a subsequence of $\{n\}_{n\geq 1}$, still denoted by itself and $\zeta^* \in C([0,T]; L^2(0,1)) \cap L^2(0,T; H^1_a(0,1))$ such that

$$\begin{aligned} \zeta(\cdot; 0, 0, u_n) \to \zeta^* & \text{weakly in } C([0, T]; L^2(0, 1)) \cap L^2(0, T; H^1_a(0, 1)) \\ & \text{and strongly in } L^2(0, 1). \end{aligned}$$
(4.28)

Passing to the limit for $n \to \infty$ in (4.28), we obtain that ζ^* is the solution of the following system

$$\begin{cases} \zeta_t^*(x,t) = A\zeta^*(x,t) + \chi_{\omega_0} u^*, & (x,t) \in (0,1) \times (0,T), \\ \zeta^*(1,t) = BC_\alpha \left(\zeta^*(\cdot,t)\right) = 0, & t \in (0,T), \\ \zeta^*(x,0) = 0, & x \in (0,1). \end{cases}$$

This means that $\zeta^* = \zeta(\cdot; 0, 0, u^*)$. Now, letting $n \to \infty$ in (4.27) we obtain that

$$\|\zeta(T;0,0,u^*) - (\zeta_T - \zeta(T;\zeta_0,g,0))\|_{L^2(G)} = \inf_{u \in \mathcal{U}} \|\zeta(T;0,0,u) - (\zeta_T - \zeta(T;\zeta_0,g,0))\|_{L^2(G)}.$$

This proves the existence of problem (4.24). In other words, u^* is the optimal control for the problem (4.24).

Step 2. We characterize the optimal control u^* .

Denote $\hat{\zeta}_T = \zeta_T - \zeta(T; \zeta_0, g, 0)$. Since u^* is an optimal control, i.e., for each $u \in \mathcal{U}$ we have

$$\|\zeta(T;0,0,u^*) - \hat{\zeta}_T\|_{L^2(G)} \le \|\zeta(T;0,0,u) - \hat{\zeta}_T\|_{L^2(G)}.$$
(4.29)

Set

$$u_{\lambda} := u^* + \lambda(u - u^*), \quad \lambda \in [0, 1].$$

It is obvious that $u_{\lambda} \in \mathcal{U}$. Then, by (4.29), we see

$$\begin{aligned} &\|\zeta(T;0,0,u^*)\|_{L^2(G)}^2 - 2\langle\zeta(T;0,0,u^*),\hat{\zeta}_T\rangle_{L^2(G)} \\ \leq &\|\zeta(T;0,0,u_\lambda)\|_{L^2(G)}^2 - 2\langle\zeta(T;0,0,u_\lambda),\hat{\zeta}_T\rangle_{L^2(G)}. \end{aligned}$$

After some calculations, the above inequality holds

$$\|\zeta(T;0,0,u-u^*)\|_{L^2(G)}\lambda^2 + 2\lambda\langle\zeta(T;0,0,u-u^*),\zeta(T;0,0,u^*) - \hat{\zeta}_T\rangle_{L^2(G)} \ge 0,$$

which, implies that for all $u \in \mathcal{U}$

$$\langle \zeta(T; 0, 0, u - u^*), \zeta(T; 0, 0, u^*) - \hat{\zeta}_T \rangle_{L^2(G)} \ge 0.$$
 (4.30)

Noting that $\hat{\zeta}_T = \zeta_T - \zeta(T; \zeta_0, g, 0)$, and plugging it into (4.26), then multiplying the first equation of (4.26) by $\zeta(\cdot; 0, 0, u - u^*)$ and integrating over $G \times (0, T)$, one deduces,

$$-\langle \zeta(T;0,0,u-u^*), \zeta(T;0,0,u^*) - \hat{\zeta}_T \rangle_{L^2(G)} = \int_0^T \langle u(t) - u^*(t), \chi_{\omega_0} z(t) \rangle_{L^2(G)} dt.$$

This, along with (4.30) stands that

$$\int_0^T \langle u(t) - u^*(t), \chi_{\omega_0} z(t) \rangle_{L^2(G)} dt \le 0, \text{ for each } u \in \mathcal{U},$$

which, implies

$$\int_0^T \langle \chi_{\omega_0} z(t), u^*(t) \rangle_{L^2(G)} dt = \max_{u(\cdot) \in \mathcal{U}} \int_0^T \langle \chi_{\omega_0} z(t), u(t) \rangle_{L^2(G)} dt.$$

This is equivalent to the following condition:

$$\langle \chi_{\omega_0} z(t), u^*(t) \rangle_{L^2(G)} = \max_{u(\cdot) \in \mathcal{U}} \langle \chi_{\omega_0} z(t), u(t) \rangle_{L^2(G)}, \text{ for } a.e. \ t \in (0, T).$$
 (4.31)

By the condition $\zeta(T;\zeta_0,g,u^*) \neq \zeta_T$ and noting that $\hat{\zeta}_T - \zeta(T;0,0,u^*) = \zeta_T - \zeta(T;\zeta_0,g,0) - \zeta(T;0,0,u^*) = \zeta_T - \zeta(T;\zeta_0,g,u^*)$, it holds $\hat{\zeta}_T - \zeta(T;0,0,u^*) \neq 0$. This, together with (4.22) in Corollary 4.2, implies

$$\|\chi_{\omega_0} z(t)\| \neq 0$$
, for *a.e.* $t \in (0, T)$.

By (4.31), we see

$$\langle \chi_{\omega_0} z(t), u^*(t) \rangle_{L^2(G)} \le M \| \chi_{\omega_0} z(t) \|.$$

Denote

$$u_0(t) = M \frac{\chi_{\omega_0} z(t)}{\|\chi_{\omega_0} z(t)\|} \text{ for a.e. } t \in (0,T).$$

It implies $u_0 \in \mathcal{U}$. Also, we have

$$\langle \chi_{\omega_0} z(t), u_0(t) \rangle_{L^2(G)} = \left\langle \chi_{\omega_0} z(t), M \frac{\chi_{\omega_0} z(t)}{\|\chi_{\omega_0} z(t)\|} \right\rangle_{L^2(G)} = M \|\chi_{\omega_0} z(t)\|_{L^2(G)}, \text{ for a.e. } t \in (0,T)$$

which, along with (4.31) again, it holds

$$\langle \chi_{\omega_0} z(t), u^*(t) \rangle_{L^2(G)} = M \| \chi_{\omega_0} z(t) \|_{L^2(G)}$$
 for a.e. $t \in (0, T)$.

Therefore, we have (4.25). This completes the proof.

Next, we shall obtain that the set of Stackelberg-Nash equilibria breaks down into three disjoint parts.

Let N be the set of Stackelberg-Nash equilibria of problem (P1) with respect to the system (2.5), and let

$$N_{0} = \left\{ (u_{1}^{*}, u_{2}^{*}) \in N : y(T, y_{0}, g, u_{1}^{*}, u_{2}^{*}) \neq y_{T}^{1}, y(T, y_{0}, g, u_{1}^{*}, u_{2}^{*}) \neq y_{T}^{2} \right\},\$$

$$N_{1} = \left\{ (u_{1}^{*}, u_{2}^{*}) \in N : y(T, y_{0}, g, u_{1}^{*}, u_{2}^{*}) = y_{T}^{1} \right\},\$$

$$N_{2} = \left\{ (u_{1}^{*}, u_{2}^{*}) \in N : y(T, y_{0}, g, u_{1}^{*}, u_{2}^{*}) = y_{T}^{2} \right\}.$$

$$(4.32)$$

Clearly, N_0, N_1, N_2 are three disjoint subsets of N and $N = N_0 \cup N_1 \cup N_2$.

Proposition 4.4. Let $g \in L^{\infty}(0,T; L^{2}(0,1))$ be given and $(u_{1}^{*}, u_{2}^{*}) \in N$. Then either $\|u_{1}^{*}\|_{L^{2}(G_{1})} = M_{1}$, for a.e. $t \in (0,T)$,

or

$$|u_2^*||_{L^2(G_2)} = M_2$$
, for a.e. $t \in (0,T)$.

Proof. Let $(u_1^*, u_2^*) \in N$. Then we have

$$J_1(u_1^*, u_2^*) \le J_1(u_1, u_2^*), \ \forall u_1 \in \mathcal{U}_1,$$

$$J_2(u_1^*, u_2^*) \le J_1(u_1^*, u_2), \ \forall u_2 \in \mathcal{U}_2.$$

Similar to the proof of Lemma 4.3, we obtain:

Case 1: if $y(T, y_0, g, u_1^*, u_2^*) \neq y_T^1$, then

$$u_1^*(t) = M_1 \frac{\chi_{\omega_1} z(t)}{\|\chi_{\omega_1} z(t)\|_{L^2(G_1)}}, \text{ for a.e. } t \in (0,T),$$

where z is a solution solving the following equation:

$$\begin{cases} z_t(x,t) + Az(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z(1,t) = BC_{\alpha}(z(\cdot,t)) = 0, & t \in (0,T), \\ z(x,T) = y_T^1 - y(T, y_0, g, u_1^*, u_2^*), & x \in (0,1). \end{cases}$$

One can easily check that

$$||u_1^*||_{L^2(G_1)} = M_1$$
, for a.e. $t \in (0,T)$.

Case 2: if $y(T, y_0, g, u_1^*, u_2^*) \neq y_T^2$, then

$$u_2^*(t) = M_2 \frac{\chi_{\omega_2} z(t)}{\|\chi_{\omega_2} z(t)\|_{L^2(G_2)}}, \text{ for a.e. } t \in (0,T),$$

where z is a solution solving the following equation:

$$\begin{cases} z_t(x,t) + Az(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z(1,t) = BC_{\alpha}(z(\cdot,t)) = 0, & t \in (0,T), \\ z(x,T) = y_T^2 - y(T,y_0,g,u_1^*,u_2^*), & x \in (0,1). \end{cases}$$

One can easily check that

$$||u_2^*||_{L^2(G_2)} = M_2$$
, for a.e. $t \in (0,T)$.

The proof is completed.

Remark 4.5. If Stackelberg-Nash equilibria $(u_1^*, u_2^*) \in N_0$, we have

$$u_1^*(t) = M_1 \frac{\chi_{\omega_1} z(t)}{\|\chi_{\omega_1} z(t)\|_{L^2(G_1)}}, \text{ for a.e. } t \in (0,T),$$
$$u_2^*(t) = M_2 \frac{\chi_{\omega_2} z(t)}{\|\chi_{\omega_2} z(t)\|_{L^2(G_2)}}, \text{ for a.e. } t \in (0,T).$$

Proposition 4.6. Let $(u_1^*, u_2^*) \in N_1$, *i.e.*,

$$u_2^*(t) = M_2 \frac{\chi_{\omega_2} z(t)}{\|\chi_{\omega_2} z(t)\|_{L^2(G_2)}}, \text{ for a.e. } t \in (0,T),$$
(4.33)

where z is a solution for adjoint equation (4.26) with the terminal condition z(T) = $y_T^2 - y(T; y_0, g, u_1, u_2^*), \ u_1 \in \mathcal{U}_1.$ Denote

$$A = \left\{ u_1 \in \mathcal{U}_1 \mid \begin{array}{c} y(T, y_0, g, u_1, u_2^*) = y_T^1, \\ \|y(T; y_0, g, u_1, u_2^*) - y_T^2\|_{L^2(G_2)} \le \|y(T; y_0, g, u_1, u_2) - y_T^2\|_{L^2(G_2)}, \\ \forall u_2 \in \mathcal{U}_2. \end{array} \right\},$$

and

$$B = \{ u_1 \in \mathcal{U}_1 : y(T, y_0, g, u_1, u_2^*) = y_T^1 \}.$$

Then, A = B.

Proof. It is clear that $A \subseteq B$. We only need to show $B \subseteq A$.

Let $u_1 \in B$. Then there exists an element $u_2 \in \mathcal{U}_2$ such that the following problem holds:

$$\begin{cases} y(T, y_0, g, u_1, u_2) = y_T^1, \\ \|y(T; y_0, g, u_1, u_2) - y_T^2\|_{L^2(G_2)} \le \|y(T; y_0, g, u_1, w_2) - y_T^2\|_{L^2(G_2)}, \ \forall w_2 \in \mathcal{U}_2. \end{cases}$$

$$(4.34)$$

For every $w_2 \in \mathcal{U}_2$ and $\lambda \in (0, 1)$, set

$$w_{\lambda} = u_2 + \lambda(w_2 - u_2)$$

then $w_{\lambda} \in \mathcal{U}_2$. Then for all $\lambda \in (0, 1)$, we have

$$||y(T; y_0, g, u_1, u_2) - y_T^2||_{L^2(G_2)} \le ||y(T; y_0, u_1, w_\lambda) - y_T^2||_{L^2(G_2)}$$

which shows that for all $\lambda \in (0, 1)$,

$$\|y(T; y_0, g, u_1, u_2) - y_T^2\|_{L^2(G_2)}^2 \le \|y(T; y_0, g, u_1, w_\lambda) - y_T^2\|_{L^2(G_2)}^2.$$

This implies that for all $\lambda \in (0, 1)$,

$$2\langle y(T; y_0, g, u_1, w_{\lambda}) - y(T; y_0, g, u_1, u_2), y_T^2 \rangle_{L^2(G_2)} \\ \leq \langle y(T; y_0, g, u_1, w_{\lambda}) - y(T; y_0, g, u_1, u_2), y(T; y_0, u_1, w_{\lambda}) + y(T; y_0, g, u_1, u_2) \rangle_{L^2(G_2)}$$

In other words, for all $\lambda \in (0, 1)$, we have

$$2\lambda \langle y(T;0,0,0,w_2-u_2), y_T^2 \rangle_{L^2(G_2)} \leq \langle \lambda y(T;0,0,0,w_2-u_2), 2y(T;y_0,g,u_1,u_2) + \lambda y(T;0,0,0,w_2-u_2) \rangle_{L^2(G_2)}.$$

Letting $\lambda \to 0^+$, it holds

$$\langle y(T;0,0,0,w_2-u_2), y_T^2 \rangle_{L^2(G_2)} \le \langle y(T;0,0,0,w_2-u_2), y(T;y_0,g,u_1,u_2) \rangle_{L^2(G_2)}.$$

This shows for all $w_2 \in \mathcal{U}_2$, it stands

$$\langle y(T;0,0,0,w_2-u_2), y_T^2 - y(T;y_0,g,u_1,u_2) \rangle_{L^2(G_2)} \leq 0.$$

In other words, for all $w_2 \in \mathcal{U}_2$, we have

$$\langle y(T; 0, 0, 0, w_2 - u_2), y_T^2 - y_T^1 \rangle_{L^2(G_2)} \leq 0.$$

From these, we obtain

$$\left\langle \int_{0}^{T} e^{A(T-t)} \chi_{\omega_{2}}(w_{2}-u_{2}) dt, y_{T}^{2}-y_{T}^{1} \right\rangle_{L^{2}(G_{2})} \leq 0.$$

This implies that for all $w_2 \in \mathcal{U}_2$,

$$\int_{0}^{T} \left\langle \chi_{\omega_{2}}(w_{2}-u_{2}), e^{A(T-t)}(y_{T}^{2}-y_{T}^{1}) \right\rangle_{L^{2}(G_{2})} dt \leq 0.$$

Therefore, we obtain for all $w_2 \in \mathcal{U}_2$,

$$\left\langle (w_2 - u_2), \chi_{\omega_2} e^{A(T-t)} (y_T^2 - y_T^1) \right\rangle_{L^2(0,T; L^2(G_2))} \le 0.$$

This is equivalent to the following condition: for *a.e.* $t \in (0,T)$

$$\langle \chi_{\omega_2} e^{A(T-t)} (y_T^2 - y_T^1), u_2(t) \rangle_{L^2(G_2)} = \max_{w_2(\cdot) \in \mathcal{U}_2} \langle \chi_{\omega_2} e^{A(T-t)} (y_T^2 - y_T^1), w_2(t) \rangle_{L^2(G_2)}.$$

By the same line as the proof of Lemma 4.3, it holds

$$u_2^*(t) = M_2 \frac{\chi_{\omega_2} z(t)}{\|\chi_{\omega_2} z(t)\|_{L^2(G_2)}}, \text{ for a.e. } t \in (0,T),$$

which, together with (4.33), shows

 $y(T, y_0, g, u_1, u_2^*) = y_T^1,$

$$\|y(T; y_0, g, u_1, u_2^*) - y_T^2\|_{L^2(G_2)} \le \|y(T; y_0, g, u_1, w_2) - y_T^2\|_{L^2(G_2)}, \ \forall w_2 \in \mathcal{U}_2.$$

This implies $u_1 \in A$. Hence, we have $B \subseteq A$. The proof is completed.

Proposition 4.7. Let $(u_1^*, u_2^*) \in N_2$, *i.e.*,

$$u_1^*(t) = M_1 \frac{\chi_{\omega_1} z(t)}{\|\chi_{\omega_1} z(t)\|_{L^2(G_1)}}, \text{ for a.e. } t \in (0,T),$$
(4.35)

where z is a solution for adjoint equation (4.26) with the terminal condition $z(T) = y_T^1 - y(T; y_0, g, u_1^*, u_2), u_2 \in \mathcal{U}_2.$

Denote

$$\bar{A} = \left\{ u_2 \in \mathcal{U}_2 \mid \begin{array}{c} y(T, y_0, g, u_1^*, u_2) = y_T^2, \\ \|y(T; y_0, g, u_1^*, u_2) - y_T^1\|_{L^2(G_1)} \leq \|y(T; y_0, g, u_1, u_2) - y_T^1\|_{L^2(G_1)}, \\ \forall u_1 \in \mathcal{U}_1. \end{array} \right\},$$

and

$$\bar{B} = \left\{ u_1 \in \mathcal{U}_1 : y(T, y_0, g, u_1^*, u_2) = y_T^2 \right\}.$$

Then, $\bar{A} = \bar{B}$.

Proof. Since the proof is similar to that of Proposition 4.6, we here omit the detail. \Box

Remark 4.8. Under the assumption of Proposition 4.6, if $u_1 \in A$, then $u_1 \in U_1$, and $y(T, y_0, g, u_1, u_2^*) = y_T^1$,

$$\|y(T;y_0,g,u_1,u_2^*) - y_T^2\|_{L^2(G_2)} \le \|y(T;y_0,g,u_1,u_2) - y_T^2\|_{L^2(G_2)}, \ \forall u_2 \in \mathcal{U}_2.$$

Thus,

$$\|y(T; y_0, g, u_1, u_2^*) - y_T^1\|_{L^2(G_1)} = 0 \le \|y(T; y_0, g, w_1, u_2^*) - y_T^1\|_{L^2(G_1)}, \ \forall w_1 \in \mathcal{U}_1,$$

 $\|y(T;y_0,g,u_1,u_2^*) - y_T^2\|_{L^2(G_2)} \le \|y(T;y_0,g,u_1,u_2) - y_T^2\|_{L^2(G_2)}, \ \forall u_2 \in \mathcal{U}_2.$

These imply $(u_1, u_2^*) \in N$. i.e.,

$$A = \{u_1 \in \mathcal{U}_1 : (u_1, u_2^*) \in N, y(T, y_0, g, u_1, u_2^*) = y_T^1\}.$$

Therefore, Proposition 4.6 tells us: if

$$u_{2}^{*} = M_{2} \frac{\chi_{\omega_{2}} e^{A(T-\cdot)} (y_{T}^{2} - y_{T}^{1})}{\|\chi_{\omega_{2}} e^{A(T-\cdot)} (y_{T}^{2} - y_{T}^{1})\|_{L^{2}(G_{2})}},$$

then

 $A = \{u_1 \in \mathcal{U}_1 : (u_1, u_2^*) \in N, y(T, y_0, g, u_1, u_2^*) = y_T^1\} = \{u_1 \in \mathcal{U}_1 : y(T, y_0, g, u_1, u_2^*) = y_T^1\}.$ Similarly, Proposition 4.7 tells us: if

$$u_1^* = M_1 \frac{\chi_{\omega_1} e^{A(T-\cdot)} (y_T^1 - y_T^2)}{\|\chi_{\omega_1} e^{A(T-\cdot)} (y_T^1 - y_T^2)\|_{L^2(G_1)}},$$

then

$$\bar{A} = \{u_2 \in \mathcal{U}_2 : (u_1^*, u_2) \in N, y(T, y_0, g, u_1^*, u_2) = y_T^2\} = \{u_2 \in \mathcal{U}_2 : y(T, y_0, g, u_1^*, u_2) = y_T^2\}.$$

From Remark 4.8, we obtain the following further characterizations of N_1 and N_2 .

Proposition 4.9. Let $g \in L^{\infty}(0,T;L^2(0,1))$ be given. Then we have the following characterizations:

$$N_{1} = \begin{cases} (u_{1}^{*}, u_{2}^{*}) \in (\mathcal{U}_{1} \times \mathcal{U}_{2}) \\ u_{2}^{*} = M_{2} \frac{\chi_{\omega_{2}} e^{A(T-\cdot)} (y_{T}^{2} - y_{T}^{1})}{\|\chi_{\omega_{2}} e^{A(T-\cdot)} (y_{T}^{2} - y_{T}^{1})\|_{L^{2}(G_{2})}}. \end{cases}$$

and

$$N_{2} = \left\{ (u_{1}^{*}, u_{2}^{*}) \in (\mathcal{U}_{1} \times \mathcal{U}_{2}) \middle| \begin{array}{c} y(T, y_{0}, g, u_{1}^{*}, u_{2}^{*}) = y_{T}^{2}, \\ u_{1}^{*} = M_{1} \frac{\chi_{\omega_{1}} e^{A(T-\cdot)}(y_{1}^{1} - y_{T}^{2})}{\|\chi_{\omega_{1}} e^{A(T-\cdot)}(y_{1}^{1} - y_{T}^{2})\|_{L^{2}(G_{1})}}. \end{array} \right\}.$$

In conclusion, Theorem 2.3 is a direct consequence of (4.32), Proposition 4.4 and Proposition 4.9.

5. The NORM OPTIMAL CONTROL PROBLEM

In this section, we discuss the norm optimal control problem (P2). At first, we study the existence, i.e., Lemma 2.6. Then the leader control g is characterized in terms of the solution of a variational problem, i.e., Theorem 2.7.

Proof of Lemma 2.6. Let $\{g_n\}_{n\geq 1} \subset \mathcal{U}_0$ be a minimizing sequence of problem (**P2**) corresponding a Stackelberg-Nash equilibrium sequence $(u_{1,n}^*(g_n), u_{2,n}^*(g_n)) \in N = N_0 \cup N_1 \cup N_2, N \subset \mathcal{U}_1 \times \mathcal{U}_2$ such that

$$||g_n||_{L^{\infty}(0,T;L^2(0,1))} \to \inf_{g \in \mathcal{U}_0} ||g||_{L^{\infty}(0,T;L^2(0,1))}.$$

Since $\{g_n\}_{n=1}^{\infty} \subset L^{\infty}(0,T; L^2(0,1))$ is bounded, then there exists a subsequence, still denoted by itself, and $g^* \in L^{\infty}(0,T; L^2(0,1))$ such that

$$g_n \to g^*$$
 weakly star in $L^{\infty}(0,T;L^2(\Omega))$ (5.1)

with

$$\|g^*\|_{L^{\infty}(0,T;L^2(\Omega))} \le \inf_{g \in \mathcal{U}_0} \|g\|_{L^{\infty}(0,T;L^2(0,1))}.$$
(5.2)

Note that $\|u_1^*(g_n)\|_{L^{\infty}(0,T;L^2(0,1))} \leq M_1, \|u_2^*(g_n)\|_{L^{\infty}(0,T;L^2(0,1))} \leq M_2$, then there exists a subsequence of $\{(u_1^*(g_n), u_2^*(g_n))\}_{n=1}^{\infty} \subset (L^{\infty}(0,T;L^2(0,1)))^2$, still denoted by itself, and $(u_1^*, u_2^*) \in (L^{\infty}(0,T;L^2(0,1)))^2$, such that

$$u_1^*(g_n) \to u_1^*, \text{ and } u_2^*(g_n) \to u_2^* \text{ weakly star in } L^{\infty}(0, T; L^2(0, 1)).$$
 (5.3)

According to $y(T; y_0, g_n, u_1^*(g_n), u_2^*(g_n)) = 0$, we get $y(T; y_0, g^*, u_1^*, u_2^*) = 0$ by (5.1) and (5.3). By (5.2), we only need to show $u_1^* = u_1^*(g^*), u_2^* = u_2^*(g^*)$.

It is obviously that there exists a subsequence of $\{(u_{1,n}^*(g_n), u_{2,n}^*(g_n))\}_{n=1}^{\infty}$, by extracting a subsequence (the subsequence is also a minimizing sequence of Problem **(P2)**), still denoted by itself, such that the sequence $\{(u_{1,n}^*(g_n), u_{2,n}^*(g_n))\}_{n=1}^{\infty}$ is contained in N_0, N_1 or N_2 . This implies the following three cases:

Case 1. If $(u_{1,n}^*(g_n), u_{2,n}^*(g_n)) \in N_0$, i.e., $y_T^1 \neq 0 \neq y_T^2$, by Remark 4.5 and (A_2) , we have

$$u_{1,n}^*(g_n) = M_1 \frac{\chi_{\omega_1} z_{1,n}}{\|\chi_{\omega_1} z_{1,n}\|_{L^2(G_1)}}, \quad u_{2,n}^*(g_n) = M_2 \frac{\chi_{\omega_2} z_{2,n}}{\|\chi_{\omega_2} z_{2,n}\|_{L^2(G_2)}},$$

where $z_{i,n}$ are the solutions of following systems

$$\begin{cases} z_{i,n,t}(,t) + Az_{i,n}(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z_{i,n}(1,t) = BC_{\alpha}(z_{i,n}(\cdot,t)) = 0, & t \in (0,T), \\ z_{i,n}(T) = y_{T}^{i}, & x \in (0,1), \end{cases}$$
(5.4)

for i = 1, 2, respectively.

Note that $z_{i,n}$ are independent on n, then $u_{1,n}^*(g_n) = u_1^*, u_{2,n}^*(g_n) = u_2^*$, i.e.,

$$u_{1,n}^*(g_n) = M_1 \frac{\chi_{\omega_1} z_1}{\|\chi_{\omega_1} z_1\|_{L^2(G_1)}}, \quad u_{2,n}^*(g_n) = M_2 \frac{\chi_{\omega_2} z_2}{\|\chi_{\omega_2} z_2\|_{L^2(G_2)}},$$

where (i = 1, 2)

$$\begin{aligned} & (x,t) + Az_i(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ & z_i(1,t) = BC_\alpha (z_i(\cdot,t)) = 0, & t \in (0,T), \\ & z_i(T) = y_T^i, & x \in (0,1). \end{aligned}$$

Hence,

$$u_1^*(g^*) = M_1 \frac{\chi_{\omega_1} z_1}{\|\chi_{\omega_1} z_1\|_{L^2(G_1)}}, \quad u_2^*(g^*) = M_2 \frac{\chi_{\omega_2} z_2}{\|\chi_{\omega_2} z_2\|_{L^2(G_2)}}$$

by (5.3) and Remark 4.5.

Case 2. If $(u_{1,n}^*(g_n), u_{2,n}^*(g_n)) \in N_1$, i.e., $y_T^1 = 0$, by Proposition 4.6 and assumption (A_2) , we have

$$u_{2,n}^*(g_n) = M_2 \frac{\chi_{\omega_2} z_{2,n}}{\|\chi_{\omega_2} z_{2,n}\|_{L^2(G_2)}},$$

where $z_{2,n}$ is the solution of following system

$$\begin{cases} z_{2,n,t}(x,t) + Az_{2,n}(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z_{2,n}(1,t) = BC_{\alpha} (z_{2,n}(\cdot,t)) = 0, & t \in (0,T), \\ z_{2,n}(x,T) = y_T^2, & x \in (0,1). \end{cases}$$

Similar to the proof of Case 1, we have

$$u_{2,n}^*(g_n) = M_2 \frac{\chi_{\omega_2} z_2}{\|\chi_{\omega_2} z_2\|_{L^2(G_2)}} = u_2^*(g^*),$$

where z_2 is the solution of following system

$$\begin{cases} z_{2,t}(x,t) + Az_2(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z_2(1,t) = BC_{\alpha}(z_2(\cdot,t)) = 0, & t \in (0,T), \\ z_2(x,T) = y_T^2, & x \in (0,1). \end{cases}$$

By Remark 4.8 and $y(T; y_0, g^*, u_1^*, u_2^*(g^*)) = 0 = y_T^1$, we have $(u_1^*, u_2^*(g^*)) \in N_1$, i.e., $u_1^* = u_1^*(g^*).$

Case 3. If $(u_{1,n}^*(g_n), u_{2,n}^*(g_n)) \in N_2$, i.e., $y_T^2 = 0$, the rest proof is similar to Case 2, then we also have $u_1^* = u_1^*(g^*), u_2^* = u_2^*(g^*)$. \square

This completes the proof.

Remark 5.1. From the proof of Lemma 2.6, we see that under the assumption (A_2) , if there exists a Stackelberg-Nash equilibrium $(u_1^*(g), u_2^*(g)) \in N$, then $(u_1^*, u_2^*) \in N_0$ or $(u_1^*(g), u_2^*) \in N_1 \text{ or } (u_1^*, u_2^*(g)) \in N_2.$

Remark 5.2. Unfortunately, due to the influences of the functions $u_1^*(g^*)$ and $u_2^*(g^*)$ with respect to g^* , where g^* is an optimal control of problem (P2), the confirmation of the bang-bang property of the optimal control g^* remains elusive, thereby preventing the attainment of the uniqueness of the norm optimal control of problem (P2).

Lemma 5.3. For any $T \in \mathbb{R}^+$, let Y_T and Z_T be defined as (2.13) and (2.14) respectively. Then

$$Y_T = Z_T. (5.5)$$

Proof. The proof will be accomplished by following two steps.

Step 1: Show that $Y_T \subset Z_T$.

Indeed, for any $\psi \in Y_T$ and noting the embedding $H^1_{\alpha}(0,1) \hookrightarrow L^2(0,1)$ is compact and similar to the proof of (4.12) and (4.13), there exists $\{z_{T,n}\}_{n\geq 1} \subset L^2(0,1)$ such that

$$\chi_{\omega} z(\cdot; z_{T,n}) \to \psi \text{ strongly in } L^1(0,T;L^2(0,1)), \tag{5.6}$$

where $z(\cdot; z_{T,n})$ is the solution of system (2.11) with the terminal data $z_{T,n} \in L^2(0,1)$. Hence

$$\|\chi_{\omega} z(\cdot; z_{T,n})\|_{L^1(0,T;L^2(0,1))} \le C.$$
(5.7)

Choose $\{T_n\}_{n\geq 1}$ with $T_n \uparrow T$ and $T_n \geq \frac{T}{2}$ for all $n \in \mathbb{N}$. We consider the following three cases.

Case a). For T_1 , there exists a constant $C_1 = C(C_0, T_2) > 0$ such that

 $\|z(T_2; z_{T,n})\| \le C_1 \|\chi_{\omega} z(\cdot; z_{T,n})\|_{L^1(T_2,T;L^2(0,1))} \le C_1 C,$

where we used the observability inequality (4.22). Hence there exists a subsequence $\{z_{T,1n}\}_{n\geq 1}$ of $\{z_{T,n}\}_{n\geq 1}$ and $\hat{z}_{T,1} \in L^2(0,1)$ such that

$$z(T_2; z_{T,1n}) \to \hat{z}_{T,1}$$
 weakly in $L^2(0, 1)$. (5.8)

Denote $z_{1n} = z(\cdot; z_{T,1n})$. Then z_{1n} is the solution to the following system

$$\begin{cases} z_{1n,t}(x,t) + Az_{1n}(x,t) = 0, & (x,t) \in (0,1) \times (0,T_2) \\ z_{1n}(1,t) = BC_{\alpha} (z_{1n}(\cdot,t)) = 0, & t \in (0,T_2), \\ z_{1n} (x,T) = z_{T,1n}, & x \in (0,1). \end{cases}$$

Let $\hat{z}_1 \in L^1(0, T_2; L^2(\Omega)) \cap C([0, T]; L^2(\Omega))$ be the solution to the following system

$$\begin{cases} \partial_t \hat{z}_1(x,t) + A \hat{z}_1(x,t) = 0, & (x,t) \in (0,1) \times (0,T_2) \\ \hat{z}_1(1,t) = B C_\alpha (\hat{z}_1(\cdot,t)) = 0, & t \in (0,T_2), \\ \hat{z}_1(T_2) = \hat{z}_{T,1}, & x \in (0,1). \end{cases}$$

Similar to the proof of (5.6), by (5.8), for every $\delta \in (\frac{T}{2}, T)$, we have

$$\chi_{\omega} z_{1n} \to \chi_{\omega} \hat{z}_1$$
 strongly in $L^1(0, T_2; L^2(0, 1)) \cap C([0, T_2 - \delta); L^2(0, 1)),$

and

$$\chi_{\omega} z_{1n} \to \chi_{\omega} \hat{z}_1 \text{ strongly in } L^1(0, T_1; L^2(0, 1)) \cap C([0, T_1]; L^2(0, 1)).$$
 (5.9)

This, together with (5.6), implies that

$$\psi = \chi_{\omega} \hat{z}_1 \text{ in } L^1(0, T_1; L^2(0, 1)),$$
(5.10)

where $\hat{z}_1(\cdot) = e^{A(T_1 - \cdot)} z_{T,1}$ and $z_{T,1} = e^{A(T_2 - T_1)} \hat{z}_{T,1}$.

Case b). Along the same way as case a), there exist a subsequence $\{z_{T,2n}\}$ of $\{z_{T,1n}\}$ and $\hat{z}_2 \in L^1(0,T_3;L^2(0,1)) \cap C([0,T_3];L^2(0,1))$ such that

$$\begin{cases} \partial_t \hat{z}_2(x,t) + A \hat{z}_2(x,t) = 0, & (x,t) \in (0,1) \times (0,T_3) \\ \hat{z}_2(1,t) = B C_\alpha (\hat{z}_2(\cdot,t)) = 0, & t \in (0,T_3), \\ \hat{z}_2(T_3) = \hat{z}_{T,2}, & x \in (0,1), \end{cases}$$

and

 $\chi_{\omega} z_{2n} = \chi_{\omega} z(\cdot; z_{T,2n}) \to \chi_{\omega} \hat{z}_2$ strongly in $L^1(0, T_2; L^2(0, 1)) \cap C([0, T_2]; L^2(0, 1)).$

These combine with (5.6), (5.9) and (5.10) imply that

$$\chi_{\omega}\hat{z}_2|_{[0,T_1]} = \chi_{\omega}\hat{z}_1,$$

and

$$\psi = \chi_{\omega} \hat{z}_2$$
 in $L^1(0, T_2; L^2(0, 1))$

where $\hat{z}_2(\cdot) = e^{A(T_2 - \cdot)} z_{T,2}$ and $z_{T,2} = e^{A(T_3 - T_2)} \hat{z}_{T,2}$.

Case c). Similarly to a) and b), for every $k \in \mathbb{N}$, we can find a subsequence $\{z_{T,kn}\}_{n\geq 1}$ of $\{z_{T,k-1,n}\}_{n\geq 1}$ such that

- $\hat{z}_k \in L^1(0, T_{k+1}; L^2(0, 1)) \cap C([0, T_{k+1}]; L^2(0, 1)),$
- $\hat{z}_{k+1}|_{[0,T_k]} = \hat{z}_k,$
 - \hat{z}_k satisfies

$$\begin{cases} \partial_t \hat{z}_k(x,t) + A\hat{z}_k(x,t) = 0, & (x,t) \in (0,1) \times (0,T_{k+1}), \\ \hat{z}_k(1,t) = BC_\alpha (\hat{z}_k(\cdot,t)) = 0, & t \in (0,T_{k+1}), \\ \hat{z}_k(T_{k+1}) = \hat{z}_{T,k}, & x \in (0,1). \end{cases}$$

• $\psi = \chi_{\omega} \hat{z}_k$ in $L^1(0, T_k; L^2(0, 1))$, where $\hat{z}_k(\cdot) = e^{A(T_k - \cdot)} z_{T,k}$ and $z_{T,k} = e^{A(T_{k+1} - T_k)} \hat{z}_{T,k}$.

Finally, define

$$\hat{z}(t) = \hat{z}_k(t)$$
 for $t \in [0, T_k]$ and $k \in \mathbb{N}$.

Such \hat{z} is well defined over [0, T) since $\hat{z}_{k+1}|_{[0, T_k]} = \hat{z}_k$, which satisfies $\hat{z} \in C([0, T); L^2(\Omega)) \cap L^1(0, T; L^2(\Omega))$, and

$$\begin{cases} \partial_t \hat{z}(x,t) + A\hat{z}(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ \hat{z}(1,t) = BC_{\alpha} \left(\hat{z}(\cdot,t) \right) = 0, & t \in (0,T) \,. \end{cases}$$

Furthermore,

$$\psi = \chi_{\omega} \hat{z}$$
 in $L^1(0,T;L^2(0,1)),$

where $\hat{z}(\cdot) = e^{A(s-\cdot)} z_{T,s}$ with $z_{T,s} \in L^2(0,1)$ and $s \in (0,T)$. From this we obtain $Y_T \subset Z_T$. Step 2: Show that $Z_T \subset Y_T$.

In fact, by the definition of Z_T , for any $\xi \in Z_T$, there exists z such that

$$\xi = \chi_{\omega} z \in L^1(0,T;L^2(0,1))$$

and for every $s \in (0, T)$ there exists $z_{T,s} \in L^2(0, 1)$ such that

$$z(\cdot) = e^{A(s-\cdot)} z_{T,s}$$
 on $[0,s]$.

Let $\{T_n\}_{n\geq 1} \subset (\frac{T}{2}, T)$ with $T_n \uparrow T$. Then, for every $n \in \mathbb{N}$, there exists $z_{T,n} \in L^2(0,1)$ such that

$$z(\cdot) = e^{A(T_n - \cdot)} z_{T,n}$$
 on $[0, T_n]$

This implies that for all $k \ge n$,

$$z_{T,n} = z(T_n) = e^{A(T_k - T_n)} z_{T,k}.$$
(5.11)

For any $l \in \mathbb{N}$, assume that $z_l(\cdot) = z(\cdot; z_{T,l})$, i.e., z_l is the solution to the following system

$$\begin{cases} z_{l,t}(x,t) + Az_{l}(x,t) = 0, & (x,t) \in (0,1) \times (0,T), \\ z_{l}(1,t) = BC_{\alpha}(z_{l}(\cdot,t)) = 0, & t \in (0,T), \\ z_{l}(x,T) = z_{T,l}, & x \in (0,1). \end{cases}$$

Then for every fixed $n \in \mathbb{N}$ and all $l, k \ge n$ with $k \ge l$, we have

$$\begin{aligned} \|z_{k}(T_{n}) - z_{l}(T_{n})\| \\ &= \left\| e^{A(T-T_{n})} z_{T,k} - e^{A(T-T_{n})} z_{T,l} \right\| = \left\| e^{A(T-T_{k})} e^{A(T_{k}-T_{n})} z_{T,k} - e^{A(T-T_{l})} e^{A(T_{l}-T_{n})} z_{T,l} \right\| \\ &= \left\| e^{A(T-T_{k})} z_{T,n} - e^{A(T-T_{l})} z_{T,n} \right\| = \left\| e^{A(T-T_{k})} \left(z_{T,n} - e^{A(T_{k}-T_{l})} z_{T,n} \right) \right\| \\ &\leq \left\| z_{T,n} - e^{A(T_{k}-T_{l})} z_{T,n} \right\|, \end{aligned}$$

where we used (5.11) in the third equality above and the operator A generates a C_0 semigroup of linear contractions in the last inequality above, see [12]. This implies that $\{z_k(T_n)\}_{k\geq n}$ is a Cauchy sequence in $L^2(0, 1)$. Furthermore,

$$||z_k(T_n) - z_{T,n}|| = \left\| e^{A(T - T_n)} z_{T,k} - z_{T,n} \right\| = \left\| e^{A(T - T_k)} z_{T,n} - z_{T,n} \right\| \to 0 \text{ as } k \to \infty,$$

from which and $z_{T,n} = z(T_n)$ and similar to the proof of (5.6) we obtain that

 $\chi_{\omega} z_k \to \chi_{\omega} z$ strongly in $L^1(0, T_n; L^2(0, 1))$ as $k \to \infty$ for every $n \in \mathbb{N}$. (5.12)

Now, it remains to show that

$$\chi_{\omega} z_k \to \chi_{\omega} z$$
 strongly in $L^1(0,T;L^2(0,1)).$ (5.13)

Let \tilde{z} satisfy

$$\begin{cases} \partial_t \tilde{z}(x,t) + A\tilde{z}(x,t) = 0, & (x,t) \in (0,1) \times (-T,T), \\ \tilde{z}(1,t) = BC_\alpha(\tilde{z}(\cdot,t)) = 0, & t \in (-T,T), \end{cases}$$

 24

and

$$\tilde{z} = z \text{ in } (0,1) \times (0,T).$$

This is a backward equation which must admit solution $\tilde{z},$ and

$$\tilde{z} \in C([-T,T); L^2(0,1)) \cap L^1(-T,T; L^2(0,1)).$$

By the definitions of \tilde{z} and z, we have

$$z_k(t) = \tilde{z}(t - (T - T_k))$$
 on $t \in (0, T)$.

By the absolutely continuity of $\chi_{\omega}\tilde{z}$, (i.e., for any $\varepsilon > 0$, there exists $\delta(\varepsilon) > 0$ such that for all $a, b \in [-T, T]$ with a < b and $|a - b| \le \delta(\varepsilon)$) we have

$$\|\chi_{\omega}\tilde{z}\|_{L^1(a,b;L^2(0,1))} < \varepsilon, \tag{5.14}$$

and by the continuity of \tilde{z} , for $\varepsilon > 0$ as above, there exists $\eta(\varepsilon) > 0$ such that for all $a, b \in [-T, T - \frac{\delta(\varepsilon)}{2}]$ with a < b and $|a - b| < \eta(\varepsilon)$,

$$\|\tilde{z}(a) - \tilde{z}(b)\| < \varepsilon. \tag{5.15}$$

Let $k_0 = k_0(\varepsilon)$ be such that for all $k \ge k_0$,

$$0 < T - T_k \leq \eta(\varepsilon)$$
 for all $k \geq k_0$.

This, together with (5.12), (5.14) and (5.15), implies that

$$\begin{split} &\|\chi_{\omega}z - \chi_{\omega}z_{k}\|_{L^{1}(0,T;L^{2}(0,1))} \\ &= \|\chi_{\omega}\tilde{z}(\cdot) - \chi_{\omega}\tilde{z}(\cdot - (T - T_{k}))\|_{L^{1}(0,T;L^{2}(0,1))} \\ &\leq \|\chi_{\omega}\tilde{z}(\cdot) - \chi_{\omega}\tilde{z}(\cdot - (T - T_{k}))\|_{L^{1}(0,T - \frac{\delta(\varepsilon)}{2};L^{2}(0,1))} + \|\chi_{\omega}\tilde{z}(\cdot - (T - T_{k}))\|_{L^{1}(T - \frac{\delta(\varepsilon)}{2},T;L^{2}(0,1))} \\ &+ \|\chi_{\omega}\tilde{z}\|_{L^{1}(T - \frac{\delta(\varepsilon)}{2},T;L^{2}(0,1))} \\ &\leq \|\tilde{z}(\cdot) - \tilde{z}(\cdot - (T - T_{k}))\|_{L^{1}(0,T - \frac{\delta(\varepsilon)}{2};L^{2}(0,1))} + \|\chi_{\omega}\tilde{z}(\cdot - (T - T_{k}))\|_{L^{1}(T - \frac{\delta(\varepsilon)}{2},T;L^{2}(0,1))} \\ &+ \|\chi_{\omega}\tilde{z}\|_{L^{1}(T - \frac{\delta(\varepsilon)}{2},T;L^{2}(0,1))} \\ &\leq \left(T - \frac{\delta(\varepsilon)}{2}\right)\varepsilon + 2\varepsilon. \end{split}$$

This gives (5.13). We have thus proved $Z_T \subset Y_T$. The Lemma 5.3 is then proved by combination of Steps 1 and 2.

We are now ready to prove Theorem 2.7.

Proof of Theorem 2.7. We carry out this proof by four steps. Step 1. The solvability of $V(T, y_0)$.

Let z with $\chi_{\omega} z \in Z_T$. From Lemma 5.3, we have

$$\begin{split} J(\chi_{\omega}z) \\ =& \frac{1}{2} \left\| \chi_{\omega}z \right\|_{L^{1}(0,T;L^{2}(0,1))}^{2} + \langle y_{0}, z(0) \rangle + \int_{0}^{T} \int_{0}^{1} \sum_{i=1}^{2} \chi_{\omega_{i}} u_{i}^{*}(g^{*}) z dx dt \\ \geq & \frac{1}{2} \left\| \chi_{\omega}z \right\|_{L^{1}(0,T;L^{2}(0,1))}^{2} - \left\| y_{0} \right\| \| z(0) \| - \max\{M_{1},M_{2}\} \| \chi_{\omega}z \|_{L^{1}(0,T;L^{2}(0,1))} \\ \geq & \frac{1}{2} \left\| \chi_{\omega}z \right\|_{L^{1}(0,T;L^{2}(0,1))}^{2} - C \| y_{0} \| \| \chi_{\omega}z \|_{L^{1}(0,T;L^{2}(0,1))} - \max\{M_{1},M_{2}\} \| \chi_{\omega}z \|_{L^{1}(0,T;L^{2}(0,1))}, \end{split}$$
where we used observability inequality (4.22) and (A_{1}), which implies that

$$\lim_{\|\chi_{\omega} z\|_{L^{1}(0,T;L^{2}(0,1))} \to \infty} J(\chi_{\omega} z) = \infty$$

Since $J(\chi_{\omega} z)$ is convex and continuous, therefore there exists z_* with $\chi_{\omega} z_* \in Z_T$ such that

$$V(T, y_0) = \frac{1}{2} \left\| \chi_{\omega} z_* \right\|_{L^1(0,T; L^2(0,1))}^2 + \langle y_0, z_*(0) \rangle + \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z_* dx dt.$$
(5.16)

Step 2. Construct a control \bar{g} .

Case (a). If $z_{*,T} \in L^2(0,1) \setminus \{0\}$, noting that $z_*(t) = e^{A(T-t)} z_{*,T}$ for all $t \in [0,T]$, from observability inequality (4.22), we obtain

$$z_*(t) \neq 0$$
 for a.e. $t \in (0, T)$. (5.17)

Hence, for all $h \in \mathbb{R}$, we have

$$\begin{split} 0 &\leq \frac{1}{2} \left\| \chi_{\omega}(z_{*} + hz) \right\|_{L^{1}(0,T;L^{2}(0,1))}^{2} + \langle y_{0}, (z_{*} + hz)(0) \rangle + \int_{0}^{T} \int_{0}^{1} \sum_{i=1}^{2} \chi_{\omega_{i}} u_{i}^{*}(g^{*})(z_{*} + hz) dx dt \\ &- \frac{1}{2} \left\| \chi_{\omega} z_{*} \right\|_{L^{1}(0,T;L^{2}(0,1))}^{2} - \langle y_{0}, z_{*}(0) \rangle - \int_{0}^{T} \int_{0}^{1} \sum_{i=1}^{2} \chi_{\omega_{i}} u_{i}^{*}(g^{*}) z_{*} dx dt \\ &= \frac{1}{2} \int_{0}^{T} \frac{2h \int_{0}^{1} \chi_{\omega} z_{*}(t) \chi_{\omega} z(t) dx + h^{2} \int_{0}^{1} \chi_{\omega} z(t)^{2} dx}{\| \chi_{\omega} z_{*}(t) + h \chi_{\omega} z(t) \| + \| \chi_{\omega} z_{*}(t) \|} dt \\ &+ h \langle y_{0}, z(0) \rangle + h \int_{0}^{T} \int_{0}^{1} \sum_{i=1}^{2} \chi_{\omega_{i}} u_{i}^{*}(g^{*}) z dx dt. \end{split}$$

Passing to the limit as $h \to 0$, we arrive at the Euler-Lagrange equation: for all z with $\chi_{\omega} z \in Z_T$,

$$\int_{0}^{T} \frac{\int_{0}^{1} \chi_{\omega} z_{*}(t) z(t) dx}{\|\chi_{\omega} z_{*}(t)\|} dt \cdot \int_{0}^{T} \|\chi_{\omega} z_{*}(t)\| dt + \langle y_{0}, z(0) \rangle + \int_{0}^{T} \int_{0}^{1} \sum_{i=1}^{2} \chi_{\omega_{i}} u_{i}^{*}(g^{*}) z dx dt = 0.$$
(5.18)

In particular,

$$\|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))}^2 + \langle y_0, z_*(0) \rangle + \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z_* dx dt = 0.$$
(5.19)

By (5.17), let

$$\bar{g}(t) = \frac{\chi_{\omega} z_*(t)}{\|\chi_{\omega} z_*(t)\|} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))} \text{ for a.e. } t \in (0,T),$$
(5.20)

then we have

$$\|\bar{g}\|_{L^{\infty}(0,T;L^{2}(0,1))} = \|\chi_{\omega} z_{*}\|_{L^{1}(0,T;L^{2}(0,1))}.$$
(5.21)

Case (b). If $z_{*,T} = 0$, similar to Case (a), we arrive at the Euler-Lagrange equation: for all z with $\chi_{\omega} z \in Z_T$,

$$\langle y_0, z(0) \rangle + \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z dx dt = 0.$$
 (5.22)

Let

$$\bar{g}(t) = 0$$
 for a.e. $t \in (0, T)$. (5.23)

Step 3. Show that
$$g^*(t) = \frac{\chi_{\omega} z_*(t)}{\|\chi_{\omega} z_*\|} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))}$$
 or $g^*(t) = 0$.

Case (i). If (5.20) holds, then multiplying z with $\chi_{\omega} z \in X_T \subset Z_T$ on both sides of the first equation of the following system

$$\begin{cases} y_t(x,t) = Ay(x,t) + \chi_{\omega}\bar{g} + \chi_{\omega_1}u_1^*(g^*) + \chi_{\omega_2}u_2^*(g^*), & (x,t) \in (0,1) \times (0,T), \\ y(1,t) = BC_{\alpha}(y(\cdot,t)) = 0, & t \in (0,T), \\ y(x,0) = y_0(x), & x \in (0,1), \end{cases}$$

and integrating with respect to $x \in (0, 1)$ and $t \in (0, T)$, it produces

$$\langle y(T; y_0, \bar{g}, u_1^*(g^*), u_2^*(g^*)), z_T \rangle - \langle y(0), z(0) \rangle - \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z dx dt$$

$$= \int_0^T \frac{\int_\omega z_*(t) z(t) dx}{\|\chi_\omega z_*(t)\|} dt \cdot \|\chi_\omega z_*\|_{L^1(0,T;L^2(0,1))},$$

which, along with (5.18), it implies that

$$y(T; y_0, \bar{g}, u_1^*(g^*), u_2^*(g^*)) = 0.$$

From the proof of Lemma 2.6, we know that $u_1^*(\bar{g}) = u_1^*(g^*)$ or $u_2^*(\bar{g}) = u_2^*(g^*)$. Case 1. If $u_1^*(\bar{g}) = u_1^*(g^*)$, which, along with Remark 4.8 and $y(T; y_0, \bar{g}, u_1^*(\bar{g}_1), u_2^*(g^*)) = 0$, implies that $u_2^*(\bar{g}) = u_2^*(g^*)$.

Case 2. If $u_2^*(\bar{g}) = u_2^*(g^*)$, similar to Case 1, we also have $u_1^*(\bar{g}) = u_1^*(g^*)$. From these discussions, we obtain $u_1^*(\bar{g}) = u_1^*(g^*)$ and $u_2^*(\bar{g}) = u_2^*(g^*)$.

Next, since g^* is the optimal control of the problem (P2), then we have

$$y(T; y_0, g^*, u_1^*(g^*), u_2^*(g^*)) = 0$$

Since $Z_T = Y_T$, similar to the proof of (5.6), assume $\{z_{T,n}\}_{n\geq 1} \subset L^2(0,1)$ be such that

$$\chi_{\omega} z(\cdot; z_{T,n}) \to \chi_{\omega} z_* \text{ strongly in } L^1(0,T; L^2(0,1)),$$
(5.24)

and

$$\chi_{\omega_i} z(\cdot; z_{T,n}) \to \chi_{\omega_i} z_* \text{ strongly in } L^1(0,T; L^2(0,1)), \text{ for } i = 1, 2.$$
Multiplying $z(\cdot; z_{T,n})$ on both sides of the first equation of the following system
$$(5.25)$$

$$\begin{cases} y_t(x,t) = Ay(x,t) + \chi_{\omega}g^* + \chi_{\omega_1}u_1^*(g^*) + \chi_{\omega_2}u_2^*(g^*), & (x,t) \in (0,1) \times (0,T), \\ y(1,t) = BC_{\alpha}(y(\cdot,t)) = 0, & t \in (0,T), \\ y(x,0) = y_0(x), & x \in (0,1), \end{cases}$$

and integrating with respect to $x \in (0, 1)$ and $t \in (0, T)$, it produces

$$-\langle y(0), z(0; z_{T,n}) \rangle - \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z(t; z_{T,n}) dx dt = \int_0^T \langle g^*(t), \chi_\omega z(t; z_{T,n}) \rangle dt.$$
(5.26)

By the same arguments as the proof of Lemma 5.3, there exists a subsequence of $\{z_{T,n}\}_{n\geq 1}$, still denoted by itself, such that

$$z(0; z_{T,n}) \to z_*(0)$$
 weakly in $L^2(0, 1)$. (5.27)

By (5.24)-(5.27), letting $n \to \infty$, we obtain that

$$-\langle y(0), z_*(0) \rangle - \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z_*(t) dx dt = \int_0^T \langle g^*(t), \chi_{\omega} z_*(t) \rangle dt,$$

which, along with (5.19) and (5.21), we have

$$\|\bar{g}\|_{L^{\infty}(0,T;L^{2}(0,1))} = \|\chi_{\omega}z_{*}\|_{L^{1}(0,T;L^{2}(0,1))} \le \|g^{*}\|_{L^{\infty}(0,T;L^{2}(0,1))}.$$
(5.28)

This implies that

$$\|\bar{g}\|_{L^{\infty}(0,T;L^{2}(0,1))} = \|g^{*}\|_{L^{\infty}(0,T;L^{2}(0,1))} = N(T,y_{0}).$$

Therefore, \bar{g} is also an optimal control of problem (P2). Replacing g^* by \bar{g} and together with (5.20), we have

$$g^*(t) = \frac{\chi_{\omega} z_*(t)}{\|\chi_{\omega} z_*(t)\|} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))} \text{ for a.e. } t \in (0,T).$$

Case (ii). If (5.23) holds, then multiplying z with $\chi_{\omega} z \in X_T \subset Z_T$ on both sides of the first equation of the following system

$$\begin{cases} y_t(x,t) = Ay(x,t) + \chi_{\omega_1} u_1^*(g^*) + \chi_{\omega_2} u_2^*(g^*), & (x,t) \in (0,1) \times (0,T), \\ y(1,t) = BC_\alpha(y(\cdot,t)) = 0, & t \in (0,T), \\ y(x,0) = y_0(x), & x \in (0,1), \end{cases}$$

and integrating with respect to $x \in (0, 1)$ and $t \in (0, T)$, it produces

$$\langle y(T; y_0, 0, u_1^*(g^*), u_2^*(g^*)), z_T \rangle - \langle y(0), z(0) \rangle - \int_0^T \int_0^1 \sum_{i=1}^2 \chi_{\omega_i} u_i^*(g^*) z dx dt = 0$$

which, along with (5.22), it implies that

$$y(T; y_0, 0, u_1^*(g^*), u_2^*(g^*)) = 0.$$

Similar to the proof of Case (i), we obtain $u_1^*(0) = u_1^*(g^*)$ and $u_2^*(0) = u_2^*(g^*)$. This shows that

$$y(T; y_0, 0, u_1^*(0), u_2^*(0)) = 0$$

Therefore, by the definition of $N(T, y_0)$, which defined in (2.10), $\bar{g} = 0$ is also an optimal control of problem (**P2**). Replacing g^* by \bar{g} and together with (5.23), we have

$$g^*(t) = 0$$
 for a.e. $t \in (0, T)$.

Step 4. Conclude that $V(T, y_0) = -\frac{1}{2}N(T, y_0)^2$. By Step 3, it shows that

Case (I). If

$$g^*(t) = \frac{\chi_{\omega} z_*(t)}{\|\chi_{\omega} z_*(t)\|} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))} \text{ for a.e. } t \in (0,T),$$

then

$$N(T, y_0) = \|\chi_{\omega} z_*\|_{L^1(0,T; L^2(0,1))},$$

which, together with (5.19) and (5.16), deduces that

$$V(T, y_0) = -\frac{1}{2} \|\chi_{\omega} z_*\|_{L^1(0,T;L^2(0,1))}^2 = -\frac{1}{2} N(T, y_0)^2.$$

Case (II). If

 $g^*(t) = 0$, for a.e. $t \in (0,T)$,

then

$$V(T, y_0) = N(T, y_0)$$

From these, we have

$$V(T, y_0) = -\frac{1}{2}N(T, y_0)^2.$$

The proof is completed.

Acknowledgement

The first three authors are supported by the National Natural Science Foundation of China under grant 11871478, the Science Technology Foundation of Hunan Province.

The last author is supported by the National Natural Science Foundation of China under grant 11971363, and by the Fundamental Research Funds for the Central Universities under grant 2042023kf0193.

References

- F. Alabau-Boussouira, P. Cannarsa, and G. Fragnelli. Carleman estimates for degenerate parabolic operators with applications to null controllability. *Journal of Evolution Equations*, 6:161–204, 2006.
- [2] J. Apraiz and L. Escauriaza. Null-control and measurable sets. ESAIM: Control, Optimisation and calculus of variations, 19(1):239–254, 2013.
- [3] J. Apraiz, L. Escauriaza, G. Wang, and C. Zhang. Observability inequalities and measurable sets. *Journal of the European Mathematical Society*, 16(11):2433–2475, 2014.
- [4] F. Araruna, E. Fernández-Cara, and M. Santos. Stackelberg-nash exact controllability for linear and semilinear parabolic equations. ESAIM: Control, Optimisation and Calculus of Variations, 21(3):835–856, 2015.
- [5] F. D. Araruna, E. Fernández-Cara, S. Guerrero, and M. C. Santos. New results on the stackelberg–nash exact control of linear parabolic equations. *Systems & Control Letters*, 104:78–85, 2017.
- [6] F. D. Araruna, B. Araújo, and E. Fernández-Cara. Stackelberg-nash null controllability for some linear and semilinear degenerate parabolic equations. *Mathematics of Control, Signals, and Systems*, 30:1–31, 2018.
- [7] T. Başar and G. J. Olsder. Dynamic Noncooperative Game Theory. SIAM, 1998.
- [8] I. Boutaayamou, G. Fragnelli, and L. Maniar. Carleman estimates for parabolic equations with interior degeneracy and neumann boundary conditions. *Journal d'Analyse Mathématique*, 135:1–35, 2018.
- [9] R. Buffe and K. D. Phung. A spectral inequality for degenerate operators and applications. *Comptes Rendus Mathematique*, 356(11-12):1131–1155, 2018.
- [10] P. Cannarsa, P. Martinez, and J. Vancostenoble. Null controllability of degenerate heat equations. Advance Differential Equations, 10:153–190, 2005.
- [11] P. Cannarsa, P. Martinez, and J. Vancostenoble. Carleman estimates for a class of degenerate parabolic operators. SIAM Journal on Control and Optimization, 47(1): 1–19, 2008.
- [12] P. Cannarsa, J. Tort, and M. Yamamoto. Unique continuation and approximate controllability for a degenerate parabolic equation. *Applicable analysis*, 91(8):1409– 1425, 2012.
- [13] P. Cannarsa, P. Martinez, and J. Vancostenoble. Global Carleman Estimates For Degenerate Parabolic Operators With Applications, volume 239. American Mathematical Society, 2016.
- [14] P. Cannarsa, R. Ferretti, and P. Martinez. Null controllability for parabolic operators with interior degeneracy and one-sided control. SIAM Journal on Control and Optimization, 57(2):900–924, 2019.
- [15] D. Charalambos and B. Aliprantis. Infinite Dimensional Analysis: A Hitchhiker's Guide. Springer-Verlag Berlin and Heidelberg GmbH & Company KG, 2013.
- [16] J. Díaz and J. Lions. On the approximate controllability of stackelberg-nash strategies. In Ocean Circulation and Pollution Control—A Mathematical and Numerical Investigation: A Diderot Mathematical Forum, pages 17–27. Springer, 2004.
- [17] R. Du. Approximate controllability of a class of semilinear degenerate systems with boundary control. *Journal of Differential Equations*, 256(9):3141–3165, 2014.

- [18] R. J. Elliott and N. J. Kalton. The Existence Of Value In Differential Games, volume 126. American Mathematical Soc., 1972.
- [19] L. Escauriaza, S. Montaner, and C. Zhang. Observation from measurable sets for parabolic analytic evolutions and applications. *Journal de Mathématiques Pures et Appliquées*, 104(5):837–867, 2015.
- [20] L. C. Evans and P. E. Souganidis. Differential games and representation formulas for solutions of hamilton-jacobi-isaacs equations. *Indiana University mathematics journal*, 33(5):773–797, 1984.
- [21] H. O. Fattorini. Infinite Dimensional Linear Control Systems: The Time Optimal And Norm Optimal Problems. Elsevier, 2005.
- [22] G. Floridia. Approximate controllability for nonlinear degenerate parabolic problems with bilinear control. *Journal of Differential Equations*, 257(9):3382–3422, 2014.
- [23] G. Fragnelli. Interior degenerate/singular parabolic equations in nondivergence form: well-posedness and carleman estimates. *Journal of Differential Equations*, 260(2): 1314–1371, 2016.
- [24] A. Friedman. Dfferential Games. Wiley, New York, 1971.
- [25] F. Guillén-González, F. Marques-Lopes, and M. Rojas-Medar. On the approximate controllability of stackelberg-nash strategies for stokes equations. *Proceedings of the American Mathematical Society*, 141(5):1759–1773, 2013.
- [26] R. P. Isaacs. Dfferential Games. Wiley, New York, 1965.
- [27] F. John. Plane Waves And Spherical Means Applied To Partial Differential Equations. Courier Corporation, 2004.
- [28] S. G. Krantz and H. R. Parks. A Primer Of Real Analytic Functions. Springer Science & Business Media, 2002.
- [29] H. Liu and C. Zhang. Observability from measurable sets for a parabolic equation involving the grushin operator and applications. *Mathematical Methods in the Applied Sciences*, 40(10):3821–3832, 2017.
- [30] C. B. Morrey Jr. Multiple Integrals In The Calculus Of Variations. Springer Science & Business Media, 2009.
- [31] K. D. Phung and G. Wang. An observability estimate for parabolic equations from a measurable set in time and its applications. *Journal of the European Mathematical Society*, 15(2):681–703, 2013.
- [32] S. M. Ross. An Elementary Introduction To Mathematical Finance. Cambridge University Press, 2011.
- [33] H. Sun and D. Yang. The fundamental gap of a kind of sub-elliptic operator. Proceedings of the Royal Society of Edinburgh Section A: Mathematics, pages 1–32, 2022.
- [34] S. Vessella. A continuous dependence result in the analytic continuation problem. 1999.
- [35] L. Wang, D. Yang, and Z. Yu. A game problem for heat equation. arXiv preprint arXiv:1802.08412, 2018.
- [36] P. Wilmott, S. Howson, S. Howison, J. Dewynne, et al. The Mathematics Of Financial Derivatives: A Student Introduction. Cambridge university press, 1995. Email address: liuyuanhang97@163.com

Email address: weijiawu@yeah.net

Email address: donghyang@outlook.com

Email address: zhangcansx@163.com