# ON TORSION-FREENESS OF KÄHLER DIFFERENTIAL SHEAVES

NILKANTHA DAS AND SUMIT ROY

ABSTRACT. Let X be a normal algebraic variety over an algebraically closed field k of characteristic zero. We prove that the Kähler differential sheaf of X is torsion-free if and only if any regular section of the ideal sheaf of the first order deformation of X inside  $X \times_k X$ , defined outside the singular locus of  $X \times_k X$ , extends regularly to the singular locus.

### 1. INTRODUCTION

Let k be an algebraically closed field of characteristic zero, and X be an algebraic variety over k. Among other few objects that come naturally with the structure of X, the sheaf of Kähler differentials  $\Omega_X := \Omega_{X/k}$  and its higher exterior powers are the central ones. For example, the algebraic de Rahm cohomology theory can be defined using the complex of regular differential forms (cf. [9]). There are several open problems that predict the structure of a variety in terms of the sheaf of Kähler differentials. For example, the Berger conjecture: a curve is non-singular if and only if its Kähler differential sheaf is torsion-free ([1, 11]), and the Zariski-Lipman conjecture: a normal variety X is non-singular if and only if the dual of  $\Omega_X$  is locally free. The Zariski-Lipaman conjecture has been studied extensively by several mathematicians over the last few decades (cf. [2, 3, 4, 5, 12, 15, 16]). The study of the various properties of  $\Omega_X$  is an active area of research till date.

It is well known that an algebraic variety X is non-singular if and only if  $\Omega_X$  is locally free. It is also well known that the torsion-free sheaves form a larger class than the class of locally free sheaves, and the reflexive sheaves form a larger class than the torsion-free ones. Therefore for singular varieties, it is quite natural to ask for conditions on X for which  $\Omega_X$  is reflexive or at least torsion-free, or if there is an equivalent criterion for  $\Omega_X$  to be torsion-free or reflexive. As mentioned in [17], these kind of questions are quite interesting. Reflexivity of differential forms has some interesting geometrical consequences (for more details, see [6, 13]). Lipman in [15, Proposition 8.1] proved that if X is a normal and local complete intersection, then  $\Omega_X$  is torsion-free, and it is reflexive if and only if X is regular in codimension 2. An alternative proof of this result can be found in [14, Corolloary 9.8]. In general,  $\Omega_X$  is not torsion-free, even if we consider X to be normal. For example, the Kähler differential sheaf of the affine cone over the twisted cubic has torsion elements (cf. [7]). In [7], Greb and Rollenske developed an equivalent criterion for the differential sheaf of the affine cone over a smooth projective variety to be torsion-free. They proved that the differential sheaf of the affine cone over a smooth projectively normal variety is torsion-free if and only if the first infinitesimal neighbourhood of the variety is projectively normal. Lipman in [15, Lemma, p. 897] also gave an equivalent cohomological criterion for a coherent sheaf to be torsion-free and reflexive, respectively. It is worthwhile to note that the above criterion works for any coherent sheaf. Nothing special happens if we restrict our attention to  $\Omega_X$ . It is tempting to

<sup>2020</sup> Mathematics Subject Classification. 14F10, 13N05, 14B05.

Key words and phrases. Kähler differential, torsion-free, normal variety.

### N. DAS AND S. ROY

ask for special equivalent criterion for  $\Omega_X$  to be torsion-free, or reflexive.

The goal of the present note is to give an equivalent criterion for  $\Omega_X$  to be torsion-free when X is a normal variety. We have proved the following

**Theorem 1.1.** Let X be a normal algebraic variety over an algebraically closed field k of characteristic zero. Then the Kähler differential sheaf  $\Omega_{X/k}$  is torsion-free if and only if the natural morphism

$$\mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I}^2$$

is surjective, where  $\mathcal{I}$  is the ideal sheaf given by the diagonal embedding  $\Delta : X \longrightarrow X \times_k X$ and j is the inclusion map of the non-singular locus of  $X \times_k X$  inside  $X \times_k X$ .

It is worthwhile to note that  $\mathcal{I}^2$  is the ideal sheaf of the first order deformation of the image of X inside  $X \times_k X$  under the diagonal morphism  $\Delta$ . Theorem 1.1 can be rephrased as follows: the Kähler differential sheaf  $\Omega_{X/k}$  is torsion-free if and only if for an open subset U of  $X \times_k X$ , a section of  $\mathcal{I}^2$  defined outside the singular locus of U, can be extended to a section of it over U. Loosely speaking, torsion-freeness of  $\Omega_X$  is equivalent to an extension problem of certain sections defined outside the singular locus of  $X \times_k X$ .

Let us phrase the problem slightly differently. The torsion submodule of  $\Omega_X$  is precisely the kernel of the natural map

$$\Omega_X \longrightarrow \Omega_X^{\vee \vee},$$

where  $\Omega_X^{\vee\vee}$  is the double dual of  $\Omega_X$ . Thus  $\Omega_X$  is torsion-free if and only if the above morphism is injective. Therefore Theorem 1.1 is now converted to prove the injectivity of the above morphism. This observation will be used in the proof of Theorem 1.1.

**Notation:** Throughout this note we assume k is an algebraically closed field of characteristic zero. An algebraic variety over k stands for an integral separated scheme of finite type over k. Given an algebraic variety X/k, the Kähler differential sheaf  $\Omega_{X/k}$  is denoted by  $\Omega_X$  for the notational convenience. Given an algebraic variety X, the diagonal map  $X \longrightarrow X \times_k X$  is denoted by  $\Delta_X$  (or simply by  $\Delta$  if there is no cause of confusion).

# 2. Preliminaries

In this section, we recall some basic facts that we need while proving Theorem 1.1. Although the following result is well-known, we will sketch a proof for the convenience of the reader.

**Lemma 2.1.** Let k be an algebraically closed field, and X, Y be two algebraic varieties over k. If both X and Y are normal, so is their product  $X \times_k Y$ .

*Proof.* Recall that a morphism of schemes is said to be normal if it is flat and for every fiber of the morphism is geometrically normal. Since k is algebraically closed, X is a normal variety if and only if X is geometrically normal over k. Also  $X \to \operatorname{spec} k$  is always flat. Therefore, X is a normal variety if and only if  $X \to \operatorname{spec} k$  is normal. By the normality theorem (cf. [8], [19, Lemma 33.10.5]), we conclude that  $X \times_k Y$  is a normal scheme over k. We are done.  $\Box$ 

As a consequence of the above result, we get if X is a normal variety over k, then so is  $X \times_k X$ .

Next, we move towards an extension problem of sections of certain kind of coherent ideal sheaves of a normal scheme. It is well-known that a reflexive sheaf  $\mathcal{F}$  on a normal scheme X is normal. That is, given an open subset V of X and an open subset  $U \subseteq V$  that contain all the codimension 1 points of V, any section of  $\mathcal{F}$  defined over U can be extended to a section


over V. Though the ideal sheaves are not reflexive in general, they enjoy the same extension property for some special kind of open subsets U.

**Proposition 2.2.** Let  $\mathcal{I}$  be an ideal sheaf of a normal integral locally Noetherian scheme X over k such that it defines an integrally closed subscheme Y of X. Assume U is an open subset of X containing all the codimension 1 points of X and  $U \cap Y \neq \emptyset$ . If the inclusion of U inside X is denoted by j, the the natural morphism

$$\mathcal{I} \longrightarrow j_* j^* \mathcal{I} \tag{2.1}$$

of sheaves on X is an isomorphism.

*Proof.* If X is an algebraic curve, the result is straightforward. Form now onwards, we assume dim  $X \ge 2$ . First, we will show that the map in eq. (2.1) is injective. Consider the following commutative diagram



Note that X is assumed to be normal and codimension of  $X \setminus U$  is  $\geq 2$ , Hartog's phenomenon immediately yields that the natural map

$$\mathcal{O}_X \longrightarrow j_*\mathcal{O}_U$$

is an isomorphism. That is, the lower horizontal morphism in the diagram is an isomorphism. Since the vertical arrows are injective, it follows that the morphism in eq. (2.1) is injective. Surjectivity of eq. (2.1) is quite subtle. Observe that  $\mathcal{I}_U$  (equivalently,  $j^*\mathcal{I}$ ) defines a closed subscheme of U whose support is  $V := U \cap Y$ . Next we will show that  $j_*j^*\mathcal{I}$  defines a closed subscheme of X whose support is Y, and subsequently, we show that  $\mathcal{I} = j_*j^*\mathcal{I}$ . Observe that  $(V, \mathcal{O}_U/\mathcal{I}_U)$  is a locally closed subscheme of X. Now applying [18, Proposition 2.3.11], we conclude that the ideal sheaf  $\overline{\mathcal{I}}$  of the scheme theoretic closure of V is given by

$$\overline{\mathcal{I}}(W) = \ker \left( \mathcal{O}_X(W) \longrightarrow \mathcal{O}_V(V \cap W) \right).$$

Now Hartog's phenomenon immediately yields that  $\mathcal{O}_X(W) = \mathcal{O}_X(W \cap U)$ . On the other hand,  $\mathcal{O}_V(V \cap W)$  is the same as  $\frac{\mathcal{O}_X}{I}\Big|_U (Y \cap W \cap U)$  which is again the same as  $\mathcal{O}_Y(W \cap U)$ .

Thus the above kernel is the same as  $(j_*j^*\mathcal{I})(W)$ . Therefore, we conclude that  $\overline{\mathcal{I}} = j_*j^*\mathcal{I}$ . On the other hand, observe that the underlying topological space of the scheme theoretic closure of V is the closure of V in X. Since Y is irreducible, V is a non-empty open subset of Y, and hence, dense in Y. It follows that the closure of V in X is Y itself. Thus we conclude that  $j_*j^*\mathcal{I}$  defines a closed subscheme of X whose support is Y. Now observe that Y is assumed to be a reduced closed subscheme. By the uniqueness of reduced closed subscheme structure with a given support (cf. [18, Proposition 2.3.11]), it follows that  $j_*j^*\mathcal{I} \subseteq \mathcal{I}$ . Since the morphism in eq. (2.1) is injective, it follows that  $\mathcal{I} = j_*j^*\mathcal{I}$ . This completes the proof.  $\Box$ 

**Remark 1.** In light of criterion (iii) of [10, Proposition 1.6], one might think that  $\mathcal{I}$  is a reflexive  $\mathcal{O}_X$ -module. But this is not the case. In fact, criterion (iii) of [10, Proposition 1.6] is demanding something more than that of Proposition 2.2. Indeed, if we consider Y is of codimension  $\geq 2$ , we can't apply Proposition 2.2 for  $U = X \setminus Y$  as  $U \cap Y = \emptyset$ . This is a place where criterion (iii) of [10, Proposition 1.6] may fails.

**Remark 2.** It is worthwhile to note that the hypothesis  $U \cap Y \neq \emptyset$  is important. If we remove this hypothesis, Proposition 2.2 is not true anymore. For example, consider Y is of codimension  $\geq 2$ , and  $U = X \setminus Y$ . Then  $j_*j^*\mathcal{I}$  is the same as  $j_*\mathcal{O}_U$ . The latter is the same as  $\mathcal{O}_X$ , thanks to the Hartog's phenomenon. Therefore, the morphism in eq. (2.1) becomes the natural injection of the ideal sheaf  $\mathcal{I}$  inside  $\mathcal{O}_X$ , but not an isomorphism. In fact, if we drop this hypothesis from Proposition 2.2, it is immediate to conclude that  $\mathcal{I}$  is reflexive. Now applying [10, Corollary 1.5] over the short exact sequence

$$0 \longrightarrow \mathcal{I} \longrightarrow \mathcal{O}_X \longrightarrow \mathcal{O}_X / \mathcal{I} \longrightarrow 0,$$

we conclude that Ass Y consists points of codimension 0 and 1 in X. Thus we get a restriction on Y.

## 3. Proof of the theorem

We have already noticed that given an algebraic variety X,  $\Omega_X$  is torsion-free if and only if the natural induced map

$$\Omega_X \longrightarrow \Omega_X^{\vee \vee} \tag{3.1}$$

of sheaves is injective. We will first find an alternative criterion of injectivity of the map in eq. (3.1). The following result will be useful in our purpose.

**Proposition 3.1.** Let U be the set of all non-singular points of a normal algebraic variety X over k. Let us denote the natural inclusion of U in X by i. Then the morphism in eq. (3.1) is injective if and only if the natural morphism of sheaves of  $\mathcal{O}_X$ -modules

$$\Omega_X \longrightarrow i_* i^* \Omega_X \tag{3.2}$$

is injective.

*Proof.* If dim X = 1, the result is immediate. Thus we assume dim  $X \ge 2$ . Let us denote S to be the space of singular points of X. Then codimension of S is at least 2 in X. Consider the following commutative diagram:

First note that  $i^*\Omega_X = \Omega_U$ . Now observe that  $i^*\Omega_X^{\vee\vee}$  is the same as  $\Omega_U^{\vee\vee}$ . This follows from the general fact that given two sheaves  $\mathcal{F}$  and  $\mathcal{G}$  of  $\mathcal{O}_X$ -modules, the sheaf  $i^* \mathscr{H}_{om\mathcal{O}_X}(\mathcal{F},\mathcal{G})$ is the same as  $\mathscr{H}_{om\mathcal{O}_U}(i^*\mathcal{F},i^*\mathcal{G})$ . Now U being the non-singular locus of X, the sheaf  $\Omega_U$  is locally free, and hence the natural map

$$\Omega_U \longrightarrow \Omega_U^{\vee \vee}$$

is an isomorphism. The functionality of  $i_*$  immediately yields that the lower horizontal arrow of the diagram (3.3) is an isomorphism.

Next we will show that the right vertical arrow of the diagram is an isomorphism. Note that  $\Omega_X^{\vee}$  is coherent, and thus  $\Omega_X^{\vee\vee}$  is reflexive by [10, Corollary 1.2]. Now apply [10, Proposition 1.6] to conclude the isomorphism.

We get both the lower horizontal arrow and the right vertical arrow are isomorphisms. The commutativity of the diagram (3.3) gives us that injectivity of (3.1) and (3.2) are equivalent.

Now we are in a position to prove our main result. For the convenience of the reader we repeat the full statement.

**Theorem 3.2.** Let X be a normal algebraic variety over k, and Y be the singular locus of X. If i denotes the inclusion map of  $U := X \setminus Y$  in X, then the natural morphism  $\Omega_X \longrightarrow i_* i^* \Omega_X$ is injective if and only if the natural map

$$\mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I}^2$$

is surjective, where j is the inclusion of  $U \times_k U$  in  $X \times_k X$ .

*Proof.* If X is a normal algebraic curve, it is always non-singular. Thus U = X, and both i, j are the identity maps. The theorem follows immediately.

From now onward we will assume dim  $X \ge 2$ . Let

$$\Delta_X: X \longrightarrow X \times_k X$$

be the diagonal morphism. Then we get the following commutative diagram

$$U \xrightarrow{\Delta_U} U \times_k U$$

$$i \bigvee_{i} \xrightarrow{\Delta_X} X \times_k X$$

$$(3.4)$$

Let us denote the map  $i \times_k i$  by j for the sake of notational simplicity. As X is separated,  $\Delta_X(X)$  (respectively  $\Delta_U(U)$ ) is closed in  $X \times_k X$  ( $U \times_k U$ , respectively). Note that  $X \times_k X$ is also a normal variety by Lemma 2.1. Let  $\mathcal{I}$  be the ideal sheaf of  $\Delta_X(X)$  in  $X \times_k X$ . Then  $j^*\mathcal{I}$  will be the ideal sheaf of  $\Delta_U(U)$  in  $U \times_k U$ .

Note that  $(X \times_k X)$  is a normal variety, and a straightforward dimension calculation shows that the codimension

$$\operatorname{codim}_{X \times_k X} ((X \times_k X) \setminus (U \times_k U)) \ge 2.$$

Thus the natural morphism

$$\mathcal{O}_{X \times_k X} \longrightarrow j_* j^* \mathcal{O}_{X \times_k X}$$

is an isomorphism, thanks to the Hartog's phenomenon. On the other hand,  $\Delta_X(X)$ , being isomorphic to X, is a reduced and irreducible closed subscheme of  $X \times_k X$ . Also  $\Delta_X(X) \cap$  $(U \times_k U)$  is the same as  $\Delta_U(U)$ , which is non-empty. It follows from Proposition 2.2 that the natural morphism of ideals

$$\mathcal{I} \longrightarrow j_* j^* \mathcal{I}$$

is an isomorphism.

Now consider the following commutative diagram:

Due to Hartog's phenomenon, the lower horizontal arrow is an isomorphism. Being ideal of the respective sheaves, both the vertical arrows are injective as well. It follows that the map

$$\mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I}^2$$

is always injective.

Note that as  $\mathcal{O}_{X \times_k X}$ -module the sheaf  $\Omega_X$  is the same as  $\mathcal{I}/\mathcal{I}^2$ . Similarly, as  $\mathcal{O}_{U \times_k U}$ module the sheaf  $\Omega_U$  is the same as  $j^* \mathcal{I}/j^* \mathcal{I}^2$ . Now let us analyse the sheaf  $i_* i^* \Omega_X$ , i.e.,  $i_* \Omega_U$ . The pushforward sheaf  $i_* \Omega_{U/k}$  is isomorphic to  $j_* (j^* \mathcal{I}/j^* \mathcal{I}^2)$  as  $\mathcal{O}_{X \times_k X}$ -module. Consider the following short exact sequence of  $\mathcal{O}_{U \times_k U}$ -modules:

$$0 \longrightarrow j^* \mathcal{I}^2 \longrightarrow j^* \mathcal{I} \longrightarrow j^* \mathcal{I} / j^* \mathcal{I}^2 \longrightarrow 0.$$

Applying  $j_*$  we get the following exact sequence of  $\mathcal{O}_{X \times_k X}$ -modules:

$$0 \longrightarrow j_* j^* \mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I} \longrightarrow j_* \left( j^* \mathcal{I} / j^* \mathcal{I}^2 \right).$$

Now consider the following diagram of exact sequences of  $\mathcal{O}_{X \times_k X}$ -modules:

Note that the middle vertical arrow is an isomorphism and the left vertical one is injective. Applying the snake lemma, we conclude that the morphism

$$\mathcal{I}/\mathcal{I}^2 \longrightarrow j_* \left( j^* \mathcal{I}/j^* \mathcal{I}^2 \right)$$

is injective if and only if

$$\mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I}^2$$

is an isomorphism. On the other hand,

$$\Omega_X \longrightarrow i_* i^* \Omega_X$$

is injective if and only if

$$(\Delta_X)_* \Omega_X \longrightarrow (\Delta_X)_* i_* \Omega_U$$

is injective as well. And the latter morphism is nothing but the rightmost vertical arrow in eq. (3.5) as follows from the commutativity of the diagram in eq. (3.4). This completes the proof.

**Remark 3.** In the above proof, we note that the morphism

$$\mathcal{I}^2 \longrightarrow j_* j^* \mathcal{I}^2$$

between the ideal sheaves is always injective and we also note that this morphism is not always an isomorphism. Therefore the Proposition 2.2 can't be applicable. In fact, the ideal sheaf  $\mathcal{I}^2$  gives us a closed subscheme of  $X \times_k X$  whose support is  $\Delta_X(X)$  which is irreducible, but the quotient sheaf  $\mathcal{O}_{X \times_k X}/\mathcal{I}^2$  is not reduced. This violates the hypothesis of Proposition 2.2.

## Acknowledgment

The authors would like to thank Sönke Rollenske for carefully going through an earlier draft and pointing out subtle issues. Both the authors are supported by the INSPIRE faculty fellowship (Ref No.: IFA21-MA 161 and IFA22-MA 186) funded by the DST, Govt. of India.

#### References

- [1] R. Berger. Differentialmoduln eindimensionaler lokaler Ringe. Math. Z., 81:326–354, 1963.
- [2] I. Biswas, R. V. Gurjar, and S. U. Kolte. On the Zariski-Lipman conjecture for normal algebraic surfaces. J. Lond. Math. Soc. (2), 90(1):270–286, 2014.
- [3] H. Flenner. Extendability of differential forms on nonisolated singularities. Invent. Math., 94(2):317–326, 1988.
- [4] P. Graf. The generalized Lipman-Zariski problem. Math. Ann., 362(1-2):241-264, 2015.
- [5] P. Graf and S. J. Kovács. An optimal extension theorem for 1-forms and the Lipman-Zariski conjecture. Doc. Math., 19:815–830, 2014.
- [6] D. Greb, S. Kebekus, and T. Peternell. Reflexive differential forms on singular spaces. Geometry and cohomology. J. Reine Angew. Math., 697:57–89, 2014.
- [7] D. Greb and S. Rollenske. Torsion and cotorsion in the sheaf of Kähler differentials on some mild singularities. *Math. Res. Lett.*, 18(6):1259–1269, 2011.
- [8] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV. Inst. Hautes Études Sci. Publ. Math., (32):361, 1967.
- [9] R. Hartshorne. Algebraic de Rham cohomology. Manuscripta Math., 7:125–140, 1972.
- [10] R. Hartshorne. Stable reflexive sheaves. Math. Ann., 254(2):121–176, 1980.
- [11] C. Huneke, S. Maitra, and V. Mukundan. Torsion in differentials and Berger's conjecture. Res. Math. Sci., 8(4):Paper No. 60, 15, 2021.
- [12] C. Jörder. A weak version of the Lipman-Zariski conjecture. Math. Z., 278(3-4):893–899, 2014.
- [13] S. Kebekus. Pull-back morphisms for reflexive differential forms. Adv. Math., 245:78–112, 2013.
- [14] E. Kunz. Kähler differentials. Advanced Lectures in Mathematics. Friedr. Vieweg & Sohn, Braunschweig, 1986.
- [15] J. Lipman. Free derivation modules on algebraic varieties. American Journal of Mathematics, 87(4):874– 898, 1965.
- [16] C. Miller and S. Vassiliadou. (Co)torsion of exterior powers of differentials over complete intersections. J. Singul., 19:131–162, 2019.
- [17] D. Mumford. The red book of varieties and schemes, volume 1358 of Lecture Notes in Mathematics. Springer-Verlag, Berlin, expanded edition, 1999. Includes the Michigan lectures (1974) on curves and their Jacobians, With contributions by Enrico Arbarello.
- [18] D. Mumford and T. Oda. Algebraic geometry. II, volume 73 of Texts and Readings in Mathematics. Hindustan Book Agency, New Delhi, 2015.
- [19] T. Stacks Project Authors. Stacks Project. https://stacks.math.columbia.edu, 2018.

STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE, 203 B.T. ROAD, KOLKATA 700 108, INDIA. *Email address:* dasnilkantha17@gmail.com

STAT-MATH UNIT, INDIAN STATISTICAL INSTITUTE, 203 B.T. ROAD, KOLKATA 700 108, INDIA. *Email address:* sumit.roy061@gmail.com