Connectivity of graphs that do not have the edge-Erdős-Pósa property

Henning Bruhn and Raphael Steck

July 24, 2023

Abstract

We show that we can assume graphs that do not have the edge-Erdős-Pósa property to be connected. Then we strengthen this result to 2connectivity under the additional assumptions of a minor-closed property and a generic counterexample.

A class \mathcal{F} has the *edge-Erdős-Pósa property* if there exists a function $f : \mathbb{N} \to \mathbb{R}$ such that for every graph G and every integer k, there are k edgedisjoint graphs in G each isomorphic to some graph in \mathcal{F} or there is an edge set $X \subseteq E(G)$ of size at most f(k) meeting all subgraphs in G isomorphic to some graph in \mathcal{F} . The edge set X is called the *hitting set*. If we replace vertices with edges in the above definition, that is, we look for a vertex hitting set or vertexdisjoint graphs, then we obtain the *vertex-Erdős-Pósa property*. The class \mathcal{F} that is studied in this paper arises from taking minors: For a fixed graph H, we define the set

 $\mathcal{F}_H = \{ G \,|\, H \text{ is a minor of } G \}.$

In other words, \mathcal{F}_H is the set of *H*-expansions. The vertex-Erdős-Pósa property for \mathcal{F}_H is well understood: Robertson and Seymour [1] proved that the class \mathcal{F}_H has the vertex-Erdős-Pósa property if and only if *H* is planar. This implies that the vertex-Erdős-Pósa property is closed under taking minors, which in turn implies that

If for a graph H the class \mathcal{F}_H has the vertex-Erdős-Pósa property, then so does the class \mathcal{F}_C for every component C of H. (1)

For the edge-Erdős-Pósa property, it is not known whether it is minor-closed or not, and it is not at all clear whether it should be. Thus, we tackle (1) for the edge-Erdős-Pósa property. We show that

Theorem 1. If for a graph H the class \mathcal{F}_H has the edge-Erdős-Pósa property, then so does \mathcal{F}_C for every component C of H.

Interestingly, Robertson and Seymour proved their result about the vertex-Erdős-Pósa property of planar graphs in two steps: First, they proved it for every connected planar graph. In a second step, they lifted the connectivity requirement. Thus Theorem 1 might also provide some help in verifying for which graphs H the class \mathcal{F}_H has the edge-Erdős-Pósa property. For example, it might help to prove that \mathcal{F}_H does not have the edge-Erdős-Pósa property if H has large treewidth (for larger or arbitrary maximum degree of H).

To this end, we also attempt a strengthening of Theorem 1 which allows us to not only focus on connected, but 2-connected graphs H. This, however, is not achieved in full generality, and we prove it only by imposing some additional assumptions (see Theorem 3).

For a graph H, a connected graph $G \subseteq H$ with at least one vertex $v \in V(G)$ with $d_H(v) \geq 3$ and an integer $r \in \mathbb{N}$, we define G^{\times} to be the following graph: Starting with the empty graph, we add a copy of every vertex $v \in V(G)$ with $d_H(v) \geq 3$ to G^{\times} . For every non-trivial path P of length l in G between two vertices such vertices, we add r internally disjoint paths of length $\max\{l, 2\}$ between their corresponding copies in G^{\times} . Finally, for every path P of length l in G between a vertex u with $d_G(u) = 1$, $d_H(u) \leq 2$ and its closest vertex $v \in V(G)$ with $d_H(v) \geq 3$, we add r paths of length l which are disjoint except for the copy of v. See Figure 1 for an example.




Figure 1: Construction of G^{\times} , with $u \in V(H) \setminus V(G)$.

Let us check that for every edge set X of size at most r-1, $G^{\times} - X$ contains a G-expansion. Every $v \in V(G)$ with $d_H(v) \geq 3$ can be mapped to its copy $v' \in V(G^{\times})$. Every u-v path between two such vertices can be mapped to one of its copies in G^{\times} that is disjoint from X. For every vertex $u \in V(G)$ with $d_G(u) = 1$ and $d_H(u) \leq 2$, there is vertex $v \in V(G)$ with $d_H(v) \geq 3$ that is closest to u. Among all copies of the u-v path, we pick a copy P' that is disjoint from X and map P to P' such that v is mapped to v'. The prerequisite of G being connected and containing a vertex v with $d_H(v) \geq 3$ implies that every $v \in V(G)$ with $d_G(v) = d_H(v) = 2$ lies on a path in G between vertices of degree other than 2 in H, with at least one endvertex of the path having degree at least 3 in H. We conclude that the above mapping yields a G-expansion in $G^{\times} - X$.

For $r \geq 3$, the number of vertices of degree at least 3 in G^{\times} is the same as the number of vertices v with $d_H(v) \geq 3$ in G.

Remark 2. Let H be a connected graph for which \mathcal{F}_H does not have the edge-Erdős-Pósa property. Then H contains vertices of degree at least 3.

Proof. Suppose H would only contain vertices of degree 2 or less. Since H is connected, H must be a cycle, a path or an isolated vertex. However, for all of those graphs, \mathcal{F}_H is already known to have the edge-Erdős-Pósa property.

For two graphs A and B, we define \leq by

$$A \preceq B \Leftrightarrow A$$
 is a minor of B .

Similarly, we define $\not\preceq$ by

$$A \not\preceq B \Leftrightarrow A$$
 is not a minor of B .

1 1-Connectivity

Proof of Theorem 1. Let A be some component of H such that \mathcal{F}_A does not have the edge-Erdős-Pósa property. Thus, there exists an integer $k \in \mathbb{N}$ such that for every $r \in \mathbb{N}$, there exists a graph A_r^* such that A_r^* neither contains k edge-disjoint expansions of A nor an edge set X of size at most r-1 such that $A_r^* - X$ contains no expansion of A. We separate the other components of H into two disjoint sets \mathcal{B} and \mathcal{C} , which we define by

 $\mathcal{B} = \{B \text{ component of } H \mid A \not\preceq B\} \text{ and}$ $\mathcal{C} = \{C \text{ component of } H \mid A \preceq C\} \setminus \{A\}.$

To prove that \mathcal{F}_H does not have the edge-Erdős-Pósa property, let r be given. We prove that there exists a graph H^* that contains neither k edge-disjoint H-expansions nor an edge set X of size at most r-1 such that $H^* - X$ contains no H-expansion. We define H^* to be the disjoint union of

- $A^* = A_r^*$,
- for every $B \in \mathcal{B}$: r distinct copies of B and
- for every $C \in \mathcal{C}$: one C^{\times} .

First we show that H^* does not contain an edge-hitting set meeting all A-expansions. Let $X \subseteq E(H^*)$ be an edge set of size at most r-1. We claim that $H^* - X$ still contains an H-expansion: Indeed, there is an A-expansion in $A^* - X$ by choice of A^* . For every $B \in \mathcal{B}$, at least one of the r copies of B in H is disjoint from X. Thus, there is a B-expansion in that copy. Finally, for every $C \in \mathcal{C}$, there is a C-expansion in $C^{\times} - X$ by construction of C^{\times} . Together, this yields an H-expansion in $H^* - X$.

We claim

Every H-expansion in
$$H^*$$
 contains an A-expansion in A^* . (2)

Note that (2) finishes the proof of the theorem: Indeed, by choice of A^* , there can be no k edge-disjoint A-expansions in A^* . To prove the claim, consider an expansion H' in H^* , and suppose that (2) is false for H'.

Since every component of H' is connected, it must be contained in a single component of H^* . Further note that every expansion of a $C \in \mathcal{C}$ in some component of H^* contains an expansion of A by definition of \mathcal{C} . Thus, by definition of \mathcal{B} , no A-expansion (and thus no C-expansion for any $C \in \mathcal{C}$) can be contained in some copy of some $B \in \mathcal{B}$. On top of that, if an A-expansion (or a C-expansion for any $C \in \mathcal{C}$) is embedded in A^* , this proves the above claim. Thus, suppose all A-expansion in H' (and thus all C-expansions for every $C \in \mathcal{C}$) are contained in $\bigcup_{\substack{C \in \mathcal{C} \\ C \in \mathcal{C}}} C^{\times}$. Every H-expansion in H^* contains at least one vertex of degree ≥ 3 in H^* for every vertex of degree ≥ 3 in H. However, by construction of C^{\times} , $\bigcup_{C \in \mathcal{C}} C^{\times}$ contains no more vertices of degree ≥ 3 than $\bigcup_{C \in \mathcal{C}} C$.

Since A contains vertices of degree ≥ 3 by Remark 2 and the branch sets of all vertices in $A \cup \left(\bigcup_{C \in \mathcal{C}} C\right)$ must be contained in $\bigcup_{C \in \mathcal{C}} C^{\times}$, this is a contradiction. Thus (2) holds, proving the theorem.

2 2-Connectivity

If we want to prove that for some graph H, \mathcal{F}_H does not have the edge-Erdős-Pósa property, the above theorem implies that it suffices to check the components of H individually. Thus, without loss of generality, we can assume that His 1-connected. To improve this to 2-connectivity of H, we need two additional assumptions.

First, we define a graph property \mathcal{P} to be *hereditary* if $\overline{\mathcal{P}}$ is closed under taking minors, that is, for every graph H without property \mathcal{P} , no minor of Hhas property \mathcal{P} . An example for a hereditary property would be large treewidth: If a graph H does not have treewidth at least t, then no minor of H has treewidth at least t.

Second, for a block A of a graph H, let S be the set of cutvertices of H that lie in A. We say that there is a generic counterexample for A if there exists some $k \in \mathbb{N}$ such that for every $r \in \mathbb{N}$, there is a graph A^* with the following properties: There are no k edge-disjoint A-expansions in A^* . Furthermore, for every $s \in S$, there is a $s' \in V(A^*)$ such that for every edge set X of size at most r - 1, there is an embedding of A in A^* such that for every vertex $s \in S$, the branch set B_s contains s'. (Together, this implies that \mathcal{F}_A does not have the edge-Erdős-Pósa property.) Every known construction that shows that some class \mathcal{F}_A does not have the edge-Erdős-Pósa property is a generic counterexample.

Theorem 3. Let \mathcal{P} be a hereditary graph property and let H be a graph that contains a block with property \mathcal{P} . Furthermore, for every block A of H with property \mathcal{P} , let there be a generic counterexample.

Then \mathcal{F}_H does not have the edge-Erdős-Pósa property.

An example for an application of Theorem*3 could be the following. Suppose we want to show that:

For every graph H of treewidth at least 10^{100} , \mathcal{F}_H does not have the edge-Erdős-Pósa property. (3)

Having large treewidth is a hereditary graph property. Assume we were able to show that for every 2-connected graph H of treewidth at least 10^{100} , \mathcal{F}_H does not have the edge-Erdős-Pósa property. Then we will most likely do that by giving a generic counterexample. Thus we can apply Theorem 3 to drop the connectivity requirement and we obtain (3). Now let us prove Theorem 3.

Proof. We can assume H to be 1-connected: Indeed, if H contains a block A with property \mathcal{P} , then there is a component Q of H that contains A. Furthermore, if all block of H with property \mathcal{P} allow for a generic counterexample, then this includes the blocks of Q. If we are able to prove that \mathcal{F}_Q does not have the edge-Erdős-Pósa property, then using Theorem 1, we conclude that \mathcal{F}_H does not have the edge-Erdős-Pósa property.

We consider the block tree T of H. Let $T_{\mathcal{P}}$ be the minimal subtree of T that contains all blocks with property \mathcal{P} . We pick A to be a leaf of $T_{\mathcal{P}}$. Note that A is a block of H that has the property \mathcal{P} . Observe that A is not a trivial block, i.e. A is not a K_2 due to Remark 2.

We define

$$\mathcal{C} = \{ C \text{ block of } H \mid A \preceq C \} \setminus \{A\}$$

Since \mathcal{P} is hereditary, it holds for every $C \in \mathcal{C}$ that C has the property \mathcal{P} , too. Let $T_{\mathcal{C}}$ be the minimal subtree of T that contains all blocks of $\mathcal{C} \cup \{A\}$. We observe that $T_{\mathcal{C}}$ is a subgraph of $T_{\mathcal{P}}$. Thus,

A is a leaf of
$$T_{\mathcal{C}}$$
. (4)

We define

$$\mathcal{B} = \{B \text{ block of } H \mid A \not\preceq B\} \cap V(T_{\mathcal{C}}) \text{ and}$$
$$\mathcal{D} = \{D \text{ component of } \bigcup_{\substack{B \text{ block of } H \\ B \notin V(T_{\mathcal{C}})}} B\}.$$

Note that since T is a tree, $T - T_C$ is a forest. For each component T' in $T - T_C$, V(T') are the blocks and cutvertices of one element of \mathcal{D} .

To show that H does not have the edge-Erdős-Pósa property, let $r \geq 3$ be some integer. We define our counterexample graph H^* to be the union of

- A^* ,
- for every $C \in \mathcal{C}$: one C^{\times} ,
- for every $D \in \mathcal{D}$: r distinct copies of D,
- for every non-trivial $B \in \mathcal{B}$: one B^{\times} and
- for every component P of $\bigcup_{\substack{B \in \mathcal{B} \\ B=K_2}} B$: one P^{\times} .

We pick the above graphs to be disjoint except for those vertices which are copies of the same vertex $v \in H$, which we identify with each other in H^* , too. In A^* , we pick the vertex s' for every $s \in S$ and identify it with all copies of s. Note that for all blocks $B \in \mathcal{B}$, there is a path P in H whose endvertices are in a block in $\{A\} \cup \mathcal{C}$ and P contains an edge of B. With A and C being non-trivial blocks for all $C \in \mathcal{C}$, their cutvertices have degree at least 3 in H. Thus the union of all trivial blocks in B is a collection of paths P whose endvertices are in non-trivial blocks and have degree at least 3 in H. We denote the union of all P^{\times} and all B^{\times} for every non-trivial $B \in \mathcal{B}$ by \mathcal{B}^{\times} . Note that every vertex v' in $\mathcal{B}^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}$ with $d_{H^*}(v') \geq 3$ is the copy of some vertex v in $\bigcup_{B \in \mathcal{B}} B \cup \bigcup_{C \in \mathcal{C}} C$ with $d_H(v) \geq 3$.

Now we show that H^* does not contain an edge-hitting set meeting all A-expansions. Let $X \subseteq E(H^*)$ be an edge set of size at most r-1. We claim that $H^* - X$ still contains an embedding of H: Indeed, we can embed A in $A^* - X$ such that for every $s \in S$, its branch set B_s contains s' by choice of A^* . For every $D \in \mathcal{D}$, at least one of the r copies of D is disjoint from X. Thus, we

can embed D in that copy. For every non-trivial $B \in \mathcal{B} \cup \mathcal{C}$, we can embed B in $B^{\times} - X$ by construction of B^{\times} . We observed above that trivial blocks of B are contained in some paths between vertices of degree at least 3, which can be embedded in their copy in $H^* - X$. Together, this yields an embedding of H in $H^* - X$ by construction of H^* .

It remains to show that there are no k edge-disjoint embeddings of H in H^* . For this, we claim:

Every H-expansion in H^* contains an A-expansion in A^* . (5)

Note that (5) proves the theorem: Indeed, by choice of A^* , there can be no k edge-disjoint embeddings of A in A^* . Let us prove (5). Way say that a block B of H is embedded in a block B^* of H^* , when for every $v \in V(B)$ with $d_H(v) \geq 3$, the branch set B_v contains a vertex $v^* \in V(B^*)$ with $d_{B^*} \geq 3$. In this sense, every block B of H is embedded in a single block B^* of H^* . Note that every embedding of a $C \in C$ in some block of H^* contains an embedding of A by definition of C. Thus neither A nor any $C \in C$ can be embedded in any copy of some $D \in \mathcal{D}$. Additionally, if some $C \in C$ is embedded in A^* , this includes an embedding of A in A^* , which was what we wanted. Thus, we may assume that neither A nor any $C \in C$ is embedded in A^* .

We conclude that $A \cup \bigcup_{C \in \mathcal{C}} C$ is embedded in $\mathcal{B}^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}$. Let $B \in \mathcal{B}$. By definition of \mathcal{B} , B is on the unique path in T that connects two blocks $C_1, C_2 \in \mathcal{C} \cup \{A\}$. Since A is a leaf of $T_{\mathcal{C}}$ by (4) and we assumed that neither C_1 nor C_2 are embedded in A^* , B cannot be embedded in A^* . For every $D \in \mathcal{D}$, there is a cutvertex separating D from all $C \in \{A\} \cup \mathcal{C}$. Therefore, B cannot be embedded in $\mathcal{B}^{\times} \cup \bigcup C^{\times}$.

be embedded in D. We conclude that B is embedded in $B^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}$. To sum up, $A \cup \bigcup_{B \in \mathcal{B}} B \cup \bigcup_{C \in \mathcal{C}} C$ is embedded in $\mathcal{B}^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}$. However, the number of vertices v in $\bigcup_{B \in \mathcal{B}} B \cup \bigcup_{C \in \mathcal{C}} C$ with $d_H(v) \ge 3$ is the same as the number of vertices v' in $\mathcal{B}^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}$ with $d_{H^*}(v') \ge 3$. By by Remark 2, A contains at least one vertex v with $d_A(v) \ge 3$. Since

By by Remark 2, A contains at least one vertex v with $d_A(v) \ge 3$. Since A is 2-connected, we conclude that it must contain at least two vertices v with $d_A(v) \ge 3$. Since A is a leaf of T_C , it shares exactly one vertex with $\bigcup_{B \in \mathcal{B}} B \cup B \cup B \in \mathcal{B}$

 $\bigcup_{C \in \mathcal{C}} C. \text{ Thus, } A \setminus \left(\bigcup_{B \in \mathcal{B}} B \cup \bigcup_{C \in \mathcal{C}} C \right) \text{ contains at least one vertex } v \text{ with } d_H(v) \geq d_A(v) \geq 3. \text{ But then it is impossible to embed } A \cup \bigcup_{B \in \mathcal{B}} B \cup \bigcup_{C \in \mathcal{C}} C \text{ in } \mathcal{B}^{\times} \cup \bigcup_{C \in \mathcal{C}} C^{\times}.$ Thus, Claim (2) holds, proving the theorem.

References

- Neil Robertson and P.D Seymour, Graph minors. V. Excluding a planar graph, Journal of Combinatorial Theory, Series B 41 (1986), no. 1, 92 – 114.
- [2] Raphael Steck and Arthur Ulmer, On the edge-Erdős-Pósa property of Ladders, 2020.