Subcubic graphs of large treewidth do not have the edge-Erdős-Pósa property

Henning Bruhn and Raphael Steck

June 16, 2023

Abstract

We show that subcubic graphs of treewidth at least 2500 do not have the edge-Erdős-Pósa property.

1 Introduction

Menger's theorem provides a strong duality between packing and covering for paths: In every graph G, there are either k disjoint paths between predefined sets $A, B \subseteq V(G)$, or there is a set $X \subseteq V(G)$ of size at most k such that G - Xcontains no A-B path. Relaxed versions of this result exist for many sets of graphs, and we call this duality the *Erdős-Pósa property*. In this article, we focus on the edge variant: A class \mathcal{F} has the *edge-Erdős-Pósa property* if there exists a function $f : \mathbb{Z}_+ \to \mathbb{R}$ such that for every graph G and every integer k, there are k edge-disjoint subgraphs of G each isomorphic to some graph in \mathcal{F} or there is an edge set $X \subseteq E(G)$ of size at most f(k) meeting all subgraphs of G isomorphic to some graph in \mathcal{F} . The edge set X is called the *hitting set*. If we replace vertices with edges in the above definition, that is, if we look for a vertex hitting set or vertex-disjoint graphs, then we obtain the *vertex-Erdős-Pósa property*. The class \mathcal{F} that is studied in this article arises from taking minors: For a fixed graph H, we define the set $\mathcal{F}_H = \{G : H \text{ is a minor of } G\}$. Any graph $G \in \mathcal{F}_H$ is called an H-expansion.

The vertex-Erdős-Pósa property for \mathcal{F}_H is well understood: Robertson and Seymour [7] proved that the class \mathcal{F}_H has the vertex-Erdős-Pósa property if and only if H is planar. While both the vertex- and the edge-Erdős-Pósa property are false for all non-planar graphs H (see for example [6]), the situation is much more mysterious for planar graphs. For some simple planar graphs H such as long cycles [2] or K_4 [1], \mathcal{F}_H still has the edge-Erdős-Pósa property, while for some others, for example subcubic trees of large pathwidth [3], it does not. For most planar graphs, it is unknown whether the edge-Erdős-Pósa property holds or not. For an overview of results on the Erdős-Pósa-property, we recommend the website of Jean-Florent Raymond [5].

We partially fill this gap by proving that for every subcubic graph of large treewidth H, \mathcal{F}_H does not have the edge-Erdős-Pósa property. Note that while it was known that large walls do not have the edge-Erdős-Pósa property (claimed without proof in [3]), this does not imply our main result as, unlike the vertex-Erdős-Pósa property, is not known whether the edge variant is closed under taking minors.

Theorem 1. For subcubic graphs H of treewidth at least 2500, \mathcal{F}_H does not have the edge-Erdős-Pósa property.

To prove Theorem 1, we only use treewidth to deduce that H contains a large wall, for which we use the linear bound provided by Grigoriev [4]. So in fact, we show the following theorem:

Theorem 2. For subcubic graphs H that contain a wall of size 250×250 , \mathcal{F}_H does not have the edge-Erdős-Pósa property.

There is room for improvement in the theorem. Requiring the graph H to be subcubic simplifies the argument considerably, but we suspect it is not necessary. Moreover, we believe that with a more careful but somewhat tedious analysis the wall size could be dropped to about 30×30 . Still, this seems unlikely to be close to be best possible. Indeed, walls of size 6×4 do not have the edge-Erdős-Pósa property [8]. (Whether graphs containing 6×4 -walls have the property is not known.)

2 Construction

There is only one known tool to prove that a set \mathcal{F}_H of *H*-expansions that satisfies the vertex-Erdős-Pósa property does not have the edge-Erdős-Pósa property: The *Heinlein Wall*, after [3], shown at size 5 in Figure 1.



Figure 1: A Heinlein Wall of size 5.

For any integer $n \in \mathbb{Z}_+$, we define $[n] = \{1, \ldots, n\}$. A Heinlein Wall W of size $r \in \mathbb{Z}_+$ is the graph consisting of the following:

- For every $j \in [r]$, let $P^j = u_1^j \dots u_{2r}^j$ be a path of length 2r 1 and for $j \in \{0\} \cup [r]$, let z_j be a vertex. Moreover, let a^* , b^* be two further vertices.
- For every $i, j \in [r]$, add the edges $z_{j-1}u_{2i-1}^j, z_ju_{2i}^j, z_{i-1}z_i, a^*u_1^j$ and $b^*u_{2r}^j$.

We define $c^* = z_0$ and $d^* = z_r$. We call the vertices a^*, b^*, c^* and d^* terminals of W, while the vertices $z_j, j \in \{0\} \cup [r]$ are called *bottleneck vertices*. Additionally, we define $W^0 = W - \{a^*, b^*, c^*, d^*\}$.

An (a^*-b^*, c^*-d^*) linkage is the vertex-disjoint union of an a^*-b^* path with a c^*-d^* path. We need an easy observation:

Lemma 3 (Bruhn et al [3]). There are no two edge-disjoint (a^*-b^*, c^*-d^*) linkages in a Heinlein Wall.

For $m, n \in \mathbb{Z}_+$, an elementary grid of size $m \times n$ is a graph with vertices $v_{i,j}$ for all $i \in [m], j \in [n]$ and edges $v_{i,j}v_{i+1,j} \forall i \in [m-1], j \in [n]$ as well as $v_{i,j}v_{i,j+1} \forall i \in [m], j \in [n-1]$. A grid is a subdivision of an elementary grid.

A wall is the subcubic variant of a grid. We define an elementary wall as an elementary grid with every second vertical edge removed. That is, an elementary wall of size $m \times n$ is an elementary grid of size $(m + 1) \times (2n + 2)$ with every edge $v_{i,2j}v_{i+1,2j}$, $i \in [m], i$ is odd, $j \in [n+1]$ and every edge $v_{i,2j-1}v_{i+1,2j-1}$, $i \in [m], i$ is even, $j \in [n + 1]$ being removed. Additionally, we remove all vertices of degree 1 and their incident edges. The i^{th} row of an elementary wall is the induced subgraph on $v_{i,1}, \ldots, v_{i,2n+2}$ for $i \in [m + 1]$ (ignore the vertices that have been removed); this is a path. There is a set of exactly n + 1 disjoint paths between the first row and the $(m + 1)^{th}$ row. These paths are the *columns* of an elementary wall. The *bricks* of an elementary wall are its 6-cycles. (See Figure 2)



Figure 2: An elementary wall of size 8×8 .

A wall is defined as the subdivision of an elementary wall. However, elementary walls have some vertices of degree 2 on the outer face of the wall. As we never want to distinguish between graphs that only differ by subdivision of edges, we avoid some annoying technicalities by slightly modifying the above definition. We define a wall' of size $m \times n$ as the subdivision of an elementary wall of size $m \times n$ with all degree 2 vertices being contracted. (See Figure 3)



Figure 3: An elementary wall of size 4×3 and a wall' of the same size.

Throughout, we will use this slightly modified definition of a wall. The key properties of a wall, such as large treewidth and planarity, carry over to a wall'. The definition of rows, columns and bricks in an elementary wall carries over to a wall' in a natural way (with some truncation of the first and last row and column). For brevity of notation, we define an *n*-wall' as a wall' of size $n \times n$.

The outercycle of a wall' W is the cycle C contained in W that contains the first and last row and first and last column. Two vertices u, v of W are *d*-apart in W if every u-v path in W, every u-C path and every v-C path in W intersects at least d+1 rows or at least d+1 columns of W. We extend the definition to bricks by saying that two bricks B_1, B_2 of W are *d*-apart in W if every pair of one vertex from B_1 and one vertex from B_2 is *d*-apart in W. Note that if v_1, v_2 are *d*-apart and if v_1 lies in the brick B_1 , and v_2 in the brick B_2 then B_1, B_2 are (d-2)-apart.

Note, furthermore, that if W is part of a planar graph G then there are no shortcuts in G. That is, if u, v are d-apart in W then there is also no u-v path in G that meets fewer than d + 1 rows and columns of W, and the same holds true for paths from u or v to the outercycle.

To apply Menger's theorem, for $n \in \mathbb{Z}_+$ and vertex sets A and B in a graph G, we define an *n*-separator as a vertex set $X \subseteq V(G)$ of size $|X| \leq n$ such that there is no A-B path in G-X. We will usually apply this for one side being a single vertex, that is $A = \{a\}$, in which case we additionally require that $a \notin X$.

3 Large treewidth results

How do we prove our main result? Let H be a planar subcubic graph of treewidth ≥ 2500 . Given a size r of a hypothetical hitting set, we show that there is a graph Z that neither contains two edge-disjoint subdivisions of H, nor admits an edge set U of size $|U| \leq r$ such that Z - U is devoid of subdivisions of H. That then proves that \mathcal{F}_H does not have the edge-Erdős-Pósa property.

Since H has treewidth ≥ 2500 , it contains a grid-minor of size at least 501×501 [4] and thus a wall' M of size at least 250×250 . We pick two edges e_1 and e_2 of M such that both of them are incident with a branch vertices of degree 3 of M and such that

every pair of one endvertex from e_1 and one endvertex from e_2 (1) is 70-apart in M.

As H is planar and M large enough it is possible to find such edges e_1, e_2 . We denote the endvertex of e_1 that is also a branch vertices of degree 3 of M by a, and the other endvertex by b (which may, or not, be a branch vertex, too). For e_2 , we call its endvertices c and d, where c is chosen to be a branch vertex of degree 3 of M.

Given a positive integer r, we define Z as follows:

- start with a copy of $H \{e_1, e_2\}$, where we denote the copy of a vertex h of H by h^* ;
- replace every edge g^*h^* in the copy of $H \{e_1, e_2\}$ by 2r internally disjoint g^*-h^* paths of length 2; and
- add a Heinlein wall W of size 2r, where the terminals a^*, b^* of W are identified with the endvertices of e_1 , and where the terminals c^*, d^* are identified with the endvertices of e_2 .



Figure 4: Construction of the counterexample graph Z

A depiction of Z can be seen in Figure 4.

We extend the mapping $V(H) \to V(Z)$ defined by $h \mapsto h^*$ to sets of vertices in $H - \{e_1, e_2\}$: for a vertex set $J \subseteq V(H)$, we set $J^* = \{h^* : h \in J\}$.

To better to distinguish between H and Z, we use the first half of the alphabet (a-m) for vertices, vertex sets and graphs that are part of H, while the second half of the alphabet (o-z) is reserved for objects belonging to Z. Starred letters of the first half (a^*-m^*) are used for vertices and objects in Z that have counterparts in H. We define M^* to be an arbitrary subdivision of $M - \{e_1, e_2\}$ in Z such that the set of its branch vertices is precisely $(V(M))^*$ and such that each subdivided edge of M^* consists of one of the 2r paths originating from multiplying the corresponding edge of $M - \{e_1, e_2\}$. Note that M^* is a wall' except for e_1, e_2 , and note that M^* is disjoint from W^0 .

Let us first prove the first half of Theorem 1: there is no small edge hitting set in Z.

Lemma 4. For every edge set U in Z of size $|U| \le r$, the graph Z - U contains a subdivision of H.

Proof. As for every edge $gh \in E(H) \setminus \{e_1, e_2\}$, the vertices g^* and h^* are linked by 2r internally disjoint paths, we may easily find a subdivision of $H - \{e_1, e_2\}$ in Z - U. Moreover, U is too small to meet all (a^*-b^*, c^*-d^*) linkages in the Heinlein wall W. Thus, the subdivision of $H - \{e_1, e_2\}$ can be extended to one of H in Z - U.

The harder part of Theorem 1 is to prove that there can be no two edgedisjoint subdivisions of H in Z. We will prove:

Lemma 5. Every subdivision of H in Z contains an (a^*-b^*, c^*-d^*) linkage in W.

Recall that, by Lemma 3, any two such linkages share an edge. Thus, once we have shown the above lemma, we then have finished the proof of the theorem.

When we talk about a subdivision of H in Z, we implicitly assume that an *embedding* of H into Z is fixed: a function Φ that maps every vertex of Hto the corresponding branch vertex in Z, and that maps every edge of H to the corresponding subdivided edge in Z. We will extend such an embedding Φ to subgraphs of H in the obvious way. In particular, $\Phi(H)$ then denotes the subdivision of H in Z. For the remainder of this article, we assume Φ to be a fixed embedding of H in Z. We will prove Lemma 5 for this fixed embedding of H. The main difficulty is that we do not know how H embeds in Z. In order to get some control on what is mapped where by Φ , we concentrate on a set of vertices that are well connected to large walls'. We will later see that only a small number of them can be mapped into W. We define a 3-fan from a vertex v to a set S as the union of three non-trivial paths from v to S that are disjoint except for their first vertex v. Set

 $B = \{h \in V(H) : \text{ there is 10-wall' } M' \text{ and} \\ a \text{ 3-fan from } h \text{ to the branch vertices of degree 3 of } M' \}.$

Note that M is also a 10-wall'.

Lemma 6. Not all branch vertices of any 10-wall' can be contained in W.

Proof. It is easy to check that a Heinlein Wall has pathwidth at most 5, and thus also treewidth at most 5. Therefore, it cannot contain a 10-wall' since the latter has treewidth at least 10. \Box

As we are only ever interested in branch vertices of degree 3, we will call those proper branch vertices. Moreover, a proper branch vertex of M^* is the image under the *-map of a proper branch vertex of M. Note that every proper branch vertex of every 10-wall' M' is in B: Indeed, every proper branch vertex in M' is connected to its three adjacent proper branch vertices of M', and those paths form the desired 3-fan. In particular, this implies that every proper branch vertex of M is in B. Recall that by choice of e_1 and e_2 , this includes aand c. For b and d, we do not know, but the following lemma helps to deal with them.

Lemma 7. Let $h^* \in V(Z - W)$ and let $T \subseteq Z$ be a 3-fan from h^* to the union of the proper branch vertices of M^* with $\{b^*, d^*\}$. Then there is also a 3-fan from h to proper branch vertices of M in H.

Proof. To prove the lemma, we need to show two things: First, we need to find a 3-fan that is disjoint from W^0 so we can pull it back to H. Second, we need to get rid of b and d and find a 3-fan that connects h with proper branch vertices of M only.

Since the terminals a^* and c^* of W are proper branch vertices of M^* and since $h^* \notin V(W)$, we can shorten the 3-fan T to obtain a 3-fan that is disjoint from W^0 but still connects h^* with proper branch vertices of M^* or b^* or d^* if necessary. Since this 3-fan is disjoint from W^0 , we can find a corresponding 3-fan F in H that connects h with proper branch vertices of M or b or d.

By Menger's theorem, we may assume that h can be separated in H from the proper branch vertices of M by a set $K \subseteq V(H) \setminus \{h\}$ of at most two vertices; otherwise we are done. In particular, the 3-fan F has to contain at least one of b and d; let us say it contains b. Moreover, the h-b path L_b in F cannot meet K as K already has to meet the two other paths in the 3-fan F. We are done if b is a proper branch vertex itself. Thus we may assume that there is a unique subdivided edge E of M that contains b in its interior. One endvertex of E is a. The set K also has to separate b from the endvertices of E (as we can reach b from h via L_b without meeting K), which implies $K \subseteq V(E)$, and $a \in K$ as b is

a neighbour of a. This implies $a \in V(F)$. Now consider the h-a path L_a in F, and observe that L_a is internally disjoint from K as $a \in K$. Furthermore, since $b \notin V(L_a)$, the penultimate vertex of L_a is a neighbour $g \neq b$ of a. Then, as H is subcubic, g lies on a subdivided edge E' of M that is not E. By extending hL_ag along E' to the endvertex of E' that is not a, we obtain a path from h to a proper branch vertex of M that avoids E. Since $K \subseteq V(E)$, that path also avoids K, a contradiction.

The next lemma gives us control over Φ , at least for the set B.

Lemma 8. $\Phi(B) \subseteq B^* \cup V(W)$.

Proof. Consider a vertex $z \in \Phi(B) \setminus V(W)$. First observe that, by definition of B, every vertex in $\Phi(B)$ has degree at least 3 in Z. Thus, for z there is a vertex h of H with $z = h^*$. We will show that $h \in B$, which then implies $z = h^* \in B^*$.

As $h^* \in \Phi(B)$, there is a vertex $g \in B$ with $h^* = \Phi(g)$. Since $g \in B$, there is a 3-fan in $\Phi(H)$ connecting h^* to the set of proper branch vertices of a 10-wall' $R \subseteq \Phi(H)$. We define O to be the union of this fan and R. If O is disjoint from the proper branch vertices of M^* and also disjoint from b^* and d^* , then it is also disjoint from W^0 and we can find a corresponding wall' and fan in H, implying that $h \in B$. (When pulling back from Z to H, paths between proper branch vertices of R can become shorter, so that the resulting graph in H may be missing some of the required degree 2 vertices to be considered a wall; this is precisely the reason why we make do with walls'.) Therefore, we conclude that O contains some proper branch vertex of M^* (and thus potentially also a part of W^0) or that O contains b^* or d^* .

Next, suppose that there is no 2-separator that separates h^* from all proper branch vertices of M^* and from b^* and d^* in Z. By Menger's theorem, there is thus a 3-fan from h^* to the proper branch vertices of M^* or b^* or d^* . We apply Lemma 7 to obtain a 3-fan in H from h to proper branch vertices of M only, which proves $h \in B$.

We conclude that there is a 2-separator $\{x, y\} \subseteq V(Z-h^*)$ that separates h^* from all proper branch vertices of M^* and all terminals of W. As every vertex of degree 3 in O is connected via three internally disjoint paths to h^* , we deduce that there is an x-y path P in O that contains all vertices that are separated by $\{x, y\}$ from h^* in O and such that all interior vertices of P have degree 2 in O. As O contains a proper branch vertex of M^* or a terminal, the path xPymust contain a vertex from $(V(M))^*$. Pick p, q to be the first respectively the last vertex of $(V(M))^*$ on P, and choose a p-q path Q in M^* . Note that Q is disjoint from O - pPq since $O \cap M^* \subseteq pPq$. Moreover, note that Q is disjoint from W^0 as M^* is disjoint from W^0 . Replacing pPq by Q, we obtain a new graph O' that is the union of a 10-wall' R' with a 3-fan from h^* to the branch vertices of R' that is disjoint from W^0 . We then also find in H a 3-fan from hto the branch vertices of a 10-wall', which again leads to $h \in B$.

With the next two lemmas we show that, with only a few exceptions, a vertex in B is mapped to a vertex in B^* under Φ .

Lemma 9. $|B^* \setminus \Phi(B)| = |\Phi(B) \cap (V(W) \setminus B^*)|$

Proof. By Lemma 8, we have

$$\begin{aligned} |\Phi(B) \cap B^*| + |\Phi(B) \cap (V(W) \setminus B^*)| &= |\Phi(B)| = |B| = |B^*| \\ &= |B^* \cap \Phi(B)| + |B^* \setminus \Phi(B)|. \end{aligned}$$

Lemma 10. $|B^* \setminus \Phi(B)| \le 52$.

Proof. By Lemma 9, it suffices to show that $|\Phi(B) \cap (V(W) \setminus B^*)| \leq 52$. We show that $|\Phi(B) \cap V(W^0)| \leq 48$, which proves the above claim since $V(W) \setminus B^*$ may differ from $V(W^0)$ only in the 4 terminals of W.

Let $z \in \Phi(B) \cap V(W^0)$, and let h be such that $z = \Phi(h)$. By definition of B, h has a 3-fan to proper branch vertices of a 10-wall' M' in H. By Lemma 6, some proper branch vertex of M' needs to be mapped outside W under Φ . Then, however, there is a 3-fan $T_z \subseteq \Phi(H)$ from z to a set of vertices in Z - W. This 3-fan must contain at least three terminals of W, and thus at least one of a^* and b^* .

Since $z \in V(W^0)$, it lies in one or possibly two blocks of $W - \{a^*, b^*\}$. We say that a block O of $W - \{a^*, b^*\}$ owns a vertex $z \in \Phi(B) \cap V(W^0)$ if z is incident in $\Phi(H)$ with at least two edges of O. As each $z \in \Phi(B) \cap V(W^0)$ has degree 3, every vertex in $\Phi(B) \cap V(W^0)$ is owned by exactly one block of $W - \{a^*, b^*\}$.

Now, assume that the block O owns z. If z is not a bottleneck vertex, then the three paths in T_z cannot all leave O through its two bottleneck vertices: one such path traverses an edge between O and a^* or b^* . The same happens if z is a bottleneck vertex: then the two paths in T_z with an edge in O cannot both leave O through the remaining bottleneck vertex. Therefore, whenever a block O owns a vertex in $\Phi(B)$, there must be an edge between O and $\{a^*, b^*\}$ in $\Phi(H)$. As a^*, b^* both have degree at most 3 in $\Phi(H)$, at most six blocks may own vertices in $\Phi(B)$.

How many vertices in $\Phi(B)$ may be owned by a block O of $W - \{a^*, b^*\}$? Every $z \in \Phi(B) \cap V(W^0)$ that is not a bottleneck vertex must have a bottleneck vertex as its neighbour in $\Phi(H)$ since z has degree 3, see Figure 1. As each bottleneck vertex has degree at most 3 in $\Phi(H)$, we conclude that each block contains at most six non-bottleneck vertices of $\Phi(B)$. Together with the two bottleneck vertices, we obtain ≤ 8 vertices of $\Phi(B)$ per block. As at most six blocks may own vertices in $\Phi(B)$, we obtain at most 48 vertices in blocks of $W - \{a^*, b^*\}$. Together with the terminals, this yields $|\Phi(B) \cap (V(W) \setminus B^*)| \leq$ 52.

Define B_M to be the set of all vertices in H that send a 3-fan to proper branch vertices of M. We note that B_M contains all proper branch vertices of M, and $B_M \subseteq B$.

Lemma 11. Let h^* be a vertex in Z - W with a 3-fan $T \subseteq Z$ to vertices in B^*_M . Then $h \in B_M$.

Proof. Suppose there is a set X of at most two vertices that separates h^* from all proper branch vertices of M^* in Z. Because X cannot separate h^* from all three endvertices of T, there exists a path P in Z - X between h^* and some vertex $g^* \in B_M^*$. As there is, by definition, a 3-fan from g to proper branch

vertices of M in H, there is also 3-fan from g^* to proper branch vertices of M^* in Z, and then, as $|X| \leq 2$, also a path Q from g^* to a proper branch vertex of M^* in Z - X. However, $P \cup Q$ is disjoint from X but contains a path from h^* to a proper branch vertex of M^* , which is impossible.

Therefore, by Menger's theorem, there is a 3-fan T' from h^* to proper branch vertices of M^* . By Lemma 7, we obtain $h \in B_M$.

In conjunction with Lemma 10, the next lemma will be used to repair M, that is to prove that $\Phi(H)$ contains most proper branch vertices of M^* and sufficient subdivided edges in between them.

Lemma 12. Let $g, h \in B_M$, and let L be a g-h path in $H - e_1 - e_2$. Let P be a g^* -h^{*} path in Z such that $V(P) \cap (V(H))^* = (V(L))^*$ and such that P is disjoint from $B^* \setminus \Phi(B)$. For every vertex $i^* \in V(P)$ that is a terminal, we furthermore require that i^* has degree 2 in $\Phi(H) - W^0$. Let S be the set of all B_M^* -paths in Z that are disjoint from the interior of P and that have at most one endvertex with P in common. Then there is a g^* -h^{*} path Q in $\Phi(H) - W^0$ that is internally disjoint from every path in S.

Proof. We do induction on the number n of internal vertices of P that lie in B_M^* . Because it is shorter, we start with the induction step. Thus, assume that n > 0, ie that P contains an internal vertex $k^* \in B_M^*$. We split the path P into $P_1 = g^* Pk^*$ and $P_2 = k^* Ph^*$, and observe that both paths have fewer than n internal vertices in B_M^* . As subpaths of P, the paths P_1 and P_2 still satisfy the conditions of the lemma. Now induction yields a g^*-k^* path $Q_1 \subseteq \Phi(H) - W^0$ and a k^*-h^* path $Q_2 \subseteq \Phi(H) - W^0$. Let Q be a g^*-h^* path contained in $Q_1 \cup Q_2 \subseteq \Phi(H) - W^0$.

Consider a path $S \in S$, and suppose that S meets Q in an internal vertex of Q. We first note that S cannot contain k^* as any path in S is disjoint from the interior of P. Thus, S meets an internal vertex of Q_1 or of Q_2 , say of Q_1 . This, however, is impossible as S is disjoint from the interior of P_1 , and may have at most one endvertex with P_1 in common. Therefore, Q is as desired, and we have proved the induction step.

It remains to establish the induction start. Then, n = 0, which implies that:

No internal vertex of P lies in
$$B_M^*$$
. (2)

As P is disjoint from $B^* \setminus \Phi(B)$, we get $g^* \in \Phi(B)$. Thus, there is a 10-wall' R and a 3-fan from g^* to proper branch vertices of R in $\Phi(H)$. We denote by O the union of R and this 3-fan. Note that O is a subgraph of $\Phi(H)$.

Let us prove that:

For any neighbour g_0 of g in $H - e_1 - e_2$, the vertex g_0^* lies in O. (3)

Indeed, since H is subcubic and since g^* has degree 3 in O it follows that for every neighbour g_0 of g in H, we have $g_0^* \in V(O)$ — unless g^* is a terminal. Then, since $g \in B_M$, g has degree 2 in $H - e_1 - e_2$, and by assumption, g^* has degree 2 in $\Phi(H) - W^0$: again, for every neighbour g_0 of g in $H - e_1 - e_2$, the vertex g_0^* lies in O.

Let g_1 be the neighbour of g in $H - e_1 - e_2$ that lies in L, the g-h path in H. It now follows from (3) that:

For the neighbour g_1 of g in L it holds that $g_1^* \in V(O \cap P)$. (4)

Among all vertices in $O \cap P$, pick k^* to be closest to h^* on P. Note that since $g_1^* \in V(P)$, it is a candidate for k^* . Thus we immediately have $k^* \neq g^*$.

Next, we claim:

$$k^* \in B_M^* \tag{5}$$

Suppose not. In particular, $k^* \neq h^*$ as $h^* \in B_M^*$. By Lemma 11, there are two vertices $x_1, x_2 \neq k^*$ that separate k^* from B_M^* . As k^*Ph^* is a $k^*-B_M^*$ path, one of x_1, x_2 lies in k^*Ph^* , say x_1 . By choice of k^* , the subpath k^*Ph^* meets O only in k^* , which implies that $x_1 \notin V(O)$. In O there are two internally disjoint k^*-g^* paths. Since $x_1 \notin V(O)$, one of the two internally disjoint k^*-g^* paths in O is disjoint from x_1, x_2 unless $x_2 = g^*$. We thus conclude $x_2 = g^*$.

Next, as $g^* \in B_M^*$ by assumption, it follows that there exists a 3-fan T from g^* to B_M^* in Z. Since H is subcubic, for two neighbours g_1, g_2 of g in $H - e_1 - e_2$, g_1^* and g_2^* lie on different paths of the 3-fan T from g^* to B_M^* in Z. That is, there are disjoint paths P_1, P_2 , where P_1 is a $g_1^* - B_M^*$ path and P_2 is a $g_2^* - B_M^*$ path, both disjoint from $x_2 = g^*$. As O is 2-connected, there are paths in O from k^* to g_1^* and g_2^* that avoid $\{x_1, x_2\}$ (recall that $x_1 \notin V(O)$). Since P_1 and P_2 are disjoint, at least one of them is disjoint from x_1 . Thus there is a $k^* - B_M^*$ path in $Z - \{x_1, x_2\}$, a contradiction. This proves (5).

With (2) we get that $k^* = h^*$, which implies $h^* \in V(O)$. We claim that:

There is a g^*-h^* path Q in O whose second vertex in $(V(H))^*$ is g_1^* . (6)

Since O is 2-connected and $g_1^* \in V(O)$ by (4), there is a $g_1^*-h^*$ path Q' in O that is disjoint from g^* . Since g_1 is a neighbour of g in $H - e_1 - e_2$, there is a $g^*-g_1^*$ path Q'' of length 2 in O, which by construction of Z is internally disjoint from Q'. Combining those to $Q = Q'' \cup Q'$ thus yields the desired g^*-h^* path Q in O. This proves (6).

Note that $Q \subseteq \Phi(H)$. Thus, to finish the proof we need to show that Q is disjoint from W^0 ; and that Q is internally disjoint from every $S \in \mathcal{S}$.

Suppose that the interior of Q meets either W^0 or some path in S, and let q be the first vertex in the interior of Q where that happens. Next, among all vertices in $g^*Qq \cap P$, pick ℓ^* to be the one closest to h^* on P.



Figure 5: Situation in Lemma 12. Vertices in B_M^* in black.

We observe that ℓ^* must be an internal vertex of P. Indeed, $\ell^* \neq h^*$ as q is an internal vertex of Q, and $\ell^* \neq g^*$ by (6). From (2) it follows that $\ell^* \notin B_M^*$, and from Lemma 11 it follows that there is a set $Y = \{y_1, y_2\}$ of at most two vertices that separates ℓ^* from B_M^* in Z.

As the paths $g^*Q\ell^*$ and ℓ^*Ph^* meet only in ℓ^* by choice of ℓ^* , it follows that one vertex in Y, y_1 say, lies in ℓ^*Ph^* and the other, y_2 , in $g^*Q\ell^*$. Now, the path ℓ^*Qq meets $g^*Q\ell^*$ and ℓ^*Ph^* also only in ℓ^* and thus is disjoint from Y. As a consequence, q cannot lie in W^0 as every vertex in W^0 sends a 3-fan to B^*_M . Therefore, q lies on a path $S \in \mathcal{S}$.

Note that as $\ell^* Qq$ is disjoint from Y and as the endvertices of S lie in B_M^* , it follows that both vertices in Y must lie on S. If y_1 lies in S then, as $y_1 \in V(P)$ and as P is internally disjoint from S, the vertex y_1 must be an endvertex of P, ie, $y_1 = h^*$. As S is a B_M^* -path, it follows that y_1 is an endvertex of S. That $y_2 \in V(g^*Q\ell^*)$ lies in S implies, too, that y_2 must be an endvertex of S: Indeed, q was the first internal vertex on Q to lie in S, and thus $y_2 = g^*$, which lies in B_M^* . But now, S has both endvertices with P in common, which is not allowed for a path in S. We have obtained the final contradiction that proves the lemma.

We are done if we find an (a^*-b^*, c^*-d^*) linkage in $\Phi(H) \cap W$. The next lemma tells us that if there is no such linkage then we obtain two different paths between the terminals, one inside the Heinlein wall and one outside.

Lemma 13. Either there is an (a^*-b^*, c^*-d^*) linkage in $\Phi(H) \cap W$, or there is an $\{a^*, b^*\}-\{c^*, d^*\}$ path in $\Phi(H) \cap W$ whose endvertices are in the same component of $\Phi(H) - W^0$.

Proof. We proceed by case distinction. First, consider the case that there is a $v \in \Phi(B)$ such that v lies in W^0 or such that v is a terminal with degree at least 2 in $\Phi(H) \cap W$.

As $v \in \Phi(B)$, there is a 3-fan T in $\Phi(H)$ from v to the proper branch vertices of some 10-wall' $R \subseteq \Phi(H)$. Moreover, as R is too large to fit into W by Lemma 6, there must be some proper branch vertex w of R outside W. Thus, $T \cup R$ contains three internally disjoint v-w paths $P_1, P_2, P_3 \subseteq \Phi(H)$.

By definition of v, there are three terminals that are incident with an edge in $(P_1 \cup P_2 \cup P_3) - W^0$. Therefore, $P_1 \cup P_2 \cup P_3$ contains an $\{a^*, b^*\} - \{c^*, d^*\}$ path that lies in $\Phi(H) \cap W$. Moreover, the endvertices of that path are connected in $\Phi(H) - W^0$ via $P_1 \cup P_2 \cup P_3 - W^0$.

Second, we consider the case when $\Phi(B) \cap V(W^0) = \emptyset$ and when every terminal in $\Phi(B)$ has degree at most 1 in $\Phi(H) \cap W$. We claim that

$$\Phi(B) \cap (V(W) \setminus B^*) = \emptyset \tag{7}$$

Since $\Phi(B) \cap V(W^0) = \emptyset$ and since $\{a^*, c^*\} \subseteq B^*$, the claim (7) can only be violated if $b^* \in \Phi(B) \setminus B^*$ or if $d^* \in \Phi(B) \setminus B^*$. While b^* and d^* are not exchangeable, they are largely symmetric for the purpose of the proof of (7). Therefore, we only concentrate on b^* and consider the case that $b^* \in \Phi(B)$ and then show that this implies $b^* \in B^*$. The proof for d^* is similar.

From $b^* \in \Phi(B)$ it follows that there are three paths P_1, P_2, P_3 in $\Phi(H)$ from b^* to proper branch vertices of some 10-wall' $R \subseteq \Phi(H)$ such that P_1, P_2, P_3 are disjoint except for b^* . Note that all proper branch vertices of R lie in $Z - W^0$ as $\Phi(B)$ is disjoint from W^0 . Therefore, R may only intersect W^0 in at most two paths. (Here, we also use that every terminal in $\Phi(B)$ has degree at most 1 in $\Phi(H) \cap W$.) Let Q_1, Q_2 be the paths in R between proper branch vertices of R that are incident with W^0 (if they exist at all).

Let $P \in \{P_1, P_2, P_3, Q_1, Q_2\}$, and observe that $P \subseteq \Phi(H)$. As the endvertices of P are either proper branch vertices of R or b^* , it follows that they lie in

 $V(H)^*$. We denote them by g^* and h^* . Moreover, as we assume $b^* \in \Phi(B)$ it follows that $\Phi(H)$ contains three internally disjoint disjoint g^*-h^* paths. Only two of these may intersect W^0 . As a consequence, the endvertices g^* and h^* are contained in the same component of $\Phi(H) - W^0$. Therefore, if $P \cap W$ contains exactly one non-trivial $\{a^*, b^*\} - \{c^*, d^*\}$ path Q, then, with the help of $P - W^0$, we see that the endvertices of Q are in the same component of $\Phi(H) - W^0$. As, moreover, $P \cap W$ contains a path between the endvertices of Q, we have found a path as in the statement of the lemma and are done.

If, on the other hand, $P \cap W$ contains two non-trivial $\{a^*, b^*\} - \{c^*, d^*\}$ paths, we can use e_1 and e_2 to find a g-h path I_P in H with $V(I_P)^* \subseteq V(P)$. If $P \cap W$ contains an a^*-b^* path or a c^*-d^* path (or both), we can again use e_1 or e_2 to find a g-h path I_P in H with $V(I_P)^* \subseteq V(P)$. (If $P \cap W$ contains both an a^*-b^* path and a c^*-d^* path, we can actually stop as then we have the desired linkage.) Finally, if $P \cap W$ contains no non-trivial path then, too, we easily find a g-h path I_P in H with $V(I_P)^* = V(P) \cap V(H)^*$.

Since R intersects W^0 only in Q_1 and Q_2 (if these exist at all), using I_{Q_1} and I_{Q_2} , we find a 10-wall' M' in H such that $V(M')^* \subseteq V(R)$. In the same way, we note that the b-M' paths $I_{P_1}, I_{P_2}, I_{P_3}$ in H satisfy $V(I_{P_j})^* \subseteq V(P_j)$ for j = 1, 2, 3. In particular, $I_{P_1}, I_{P_2}, I_{P_3}$ are pairwise disjoint except for b. In total, we have found a 3-fan from b to a 10-wall', which implies that $b \in B$ and thus $b^* \in B^*$. This proves (7).

By Lemma 9, it follows from (7) and $|B| = |\Phi(B)|$ that

$$B^* = \Phi(B). \tag{8}$$

In particular, the terminals a^* and c^* lie in $\Phi(B)$, which implies that there is, for every terminal $v \in \{a^*, c^*\}$, a 3-fan $T \subseteq \Phi(H)$ from v to proper branch vertices of some 10-wall' $R \subseteq \Phi(H)$. Note that all proper branch vertices of Rlie in $\Phi(B)$ and thus outside W^0 . Therefore, there is for every $v \in \{a^*, c^*\}$ a path Q_v that starts in v, that ends in another terminal and that is completely contained in W. Moreover, via the 3-fan T, there is a path between the endvertices in $\Phi(H) - W^0$. (Observe that the paths Q_{a^*}, Q_{c^*} do not have to be disjoint, nor distinct.)

If Q_{a^*} ends in $\{c^*, d^*\}$ or if Q_{c^*} ends in $\{a^*, b^*\}$, we observe that Q_{a^*} or Q_{c^*} is an $\{a^*, b^*\} - \{c^*, d^*\}$ path in $\Phi(H) \cap W$ whose endvertices are in the same component of $\Phi(H) - W^0$ and we are done. Thus we may assume that Q_{a^*} is an a^*-b^* path and Q_{c^*} is a c^*-d^* path. If Q_{a^*} is disjoint from Q_{c^*} , they form an (a^*-b^*, c^*-d^*) linkage in $\Phi(H) \cap W$, which was what we wanted. Thus, we may assume that Q_{a^*} intersects Q_{c^*} , which implies that $Q_{a^*} \cup Q_{c^*}$ contains an a^*-c^* path P. We then apply Lemma 12 to a^*, c^* in the role of g^*, h^* , and to some a-c path L in $M - e_1 - e_2$. Note that $(V(L))^*$ is automatically disjoint from $B^* \setminus \Phi(B)$, as the latter set is empty, by (8). The path we obtain from the lemma then shows that the endvertices of P lie in the same component of $\Phi(H) - W^0$, and we are done.

In the next lemma we will use planarity arguments. To this end, if G is a planar graph that is drawn in the plane, ie, if $G \subseteq \mathbb{R}^2$, then we define the *interior* int(G) as the set $\mathbb{R}^2 \setminus F$, where F is the outer face (the unbounded face) of G.

We have reached the final lemma that concludes the proof of Theorem 2.

Lemma 5. $\Phi(H)$ contains an (a^*-b^*, c^*-d^*) linkage in W.

Proof. Suppose that $\Phi(H) \cap W$ does not contain any (a^*-b^*, c^*-d^*) linkage. Then Lemma 13 yields $v_1 \in \{a^*, b^*\}$, $v_2 \in \{c^*, d^*\}$ and v_1-v_2 paths P and Q such that $P \subseteq \Phi(H) \cap W$ and $Q \subseteq \Phi(H) - W^0$.

Set $D = \{h \in B : h^* \notin \Phi(B)\}$ and observe that Lemma 10 implies that $|D| \leq 52$. Every vertex is incident with at most one row and one column of the wall' M. Thus, there is a wall' $M' \subseteq M - D - \{a, b, c, d\}$ that contains all but at most 56 rows and columns of M, and that is disjoint from D and from the terminals a, b, c, d. We write $M'^* \subseteq Z$ for the subwall' of M^* that contains all images of the branch vertices of M' under *.

As all proper branch vertices of M' are in B_M and as M'^* is disjoint from $B^* \setminus \Phi(B)$ we can apply Lemma 12 to every (subdivided) edge gh of M' to see that there is a g^*-h^* path in $\Phi(H)-W^0$. Moreover, as such a path is a B^*_M -path (and thus in S with respect to the lemma), the obtained paths are all internally disjoint. Replacing the subdivided edges of M'^* one by one in this way, we obtain a wall' R in $\Phi(H) - W^0$ whose proper branch vertices are identical with those from M'^* . In particular, for every row (resp. for every column) of M'^* there is a row (resp. a column) of R with the same proper branch vertices.

We note for later that

$$R \subseteq \Phi(H) - W \tag{9}$$

We make a second observation. The graph $Z - W^0$ is planar as H is planar, and, in what follows, we consider a fixed drawing of $Z - W^0$. Then, the interior int(S) of any brick S of M^* is well-defined. We may assume that $Z - W^0$ is drawn in such a way that no brick interior contains the outercycle of M^* . We use this to observe that if S' is a brick of M'^* and if S is the corresponding brick of R with the same proper branch vertices then any vertex in int(S') lies in the interior int(S) or in the interior of a brick of R that is adjacent to S, ie, that shares a subdivided edge with S.

Recall the v_1-v_2 path Q contained in $\Phi(H) - W^0$. We claim:

Q meets R, and if q_1 is its first and q_2 its last vertex in R then q_1, q_2 are 8-apart in R. (10)

As each pair of one vertex from $\{a, b\}$ and one of $\{c, d\}$ is 70-apart in M, it follows that v_1, v_2 are 70-apart in M^* . (Recall that M^* is a subdivision of $M - e_1 - e_2$ in $Z - W^0$.) As every path in M^* from v_1 or from v_2 to the outercycle of M^* meets at least 70 rows or columns, and as M'^* contains all but 56 rows and all but 56 columns of M^* it follows that there are bricks S'_1, S'_2 of M'^* such that $v_i \in int(S'_i)$ for i = 1, 2.

Consider a path $Q' \subseteq M'^*$ from a vertex of S'_1 to a vertex of S'_2 and suppose that Q' meets M'^* fewer than 10 times. Then follow Q, which is a path in $Z - W^0$, from v_1 to the first vertex in S'_1 , then along S'_1 to the first vertex of Q', then along Q' to S'_2 , from there to the last vertex of Q in S'_2 and along Qto v_2 . The resulting v_1-v_2 path $Q'' \subseteq Z - W^0$ meets fewer than 14 rows and columns of M'^* (each of the bricks S'_1 and S'_2 may contribute at most two more rows and columns). As M'^* contains all but 56 rows and columns of M^* we see that Q'' meets fewer than 70 rows and columns of M^* , which is impossible as v_1, v_2 are 70-apart in M^* . In a similar way, we see that each path from v_1 or from v_2 to the outercycle of M'^* meets 10 rows or columns of M'^* . Therefore, S'_1, S'_2 are 10-apart in M'^* . As we had observed that the interior of each brick of R is contained in the interior of the corresponding brick in M'^* together with the interiors of adjacent bricks, it follows that there are bricks S_1 and S_2 of R such that $v_i \in int(S_i)$ for i = 1, 2 and such that S_1, S_2 are 8-apart in R.

As a consequence, the path Q, which is entirely contained in the plane graph $Z - W^0$, meets R (in at least eight vertices). Denote by q_1 the first vertex of Q in R, and let q_2 be the last vertex of Q in R. Then q_1 lies in the brick S_1 , and q_2 lies in S_2 . Therefore, q_1, q_2 are 8-apart in R. This proves (10).

Recall the v_1-v_2 path P contained in $\Phi(H) \cap W$. As H is planar, and as as $Q \cup P \cup R \subseteq \Phi(H)$, it follows that $Q \cup P \cup R$ is planar, too. Consider $q_1Qv_1 \cup P \cup v_2Qq_2$: this is a q_1-q_2 path that meets the wall' R only in its endvertices since $P \subseteq W$, while R is disjoint from W, by (9). However, q_1, q_2 are 8-apart, by (10). Clearly, this is impossible in a planar graph. The final contradiction proves the lemma.

References

- Henning Bruhn and Matthias Heinlein, K₄-subdivisions have the edge-Erdős-Pósa property, SIAM Journal on Discrete Mathematics 35 (2021), no. 1, 392–430.
- [2] Henning Bruhn, Matthias Heinlein, and Felix Joos, Long cycles have the edge-Erdős-Pósa property, Combinatorica 39 (2019), 1–36.
- [3] Henning Bruhn, Matthias Heinlein, and Felix Joos, The edge-Erdős-Pósa property, Combinatorica 41 (2020), 147 – 173.
- [4] Alexander Grigoriev, Tree-width and large grid minors in planar graphs, Discrete Mathematics & Theoretical Computer Science 13 (2011), 13–20.
- [5] Jean-Florent Raymond, Dynamic Erdős-Pósa listing, https://perso.limos.fr/~jfraymon/Erd%c5%91s-P%c3%b3sa/, Accessed: 2023-05-04.
- [6] Jean-Florent Raymond and Dimitrios M. Thilikos, Recent techniques and results on the Erdős–Pósa property, Discrete Applied Mathematics 231 (2017), 25 – 43, Algorithmic Graph Theory on the Adriatic Coast.
- [7] Neil Robertson and P.D Seymour, Graph minors. V. Excluding a planar graph, Journal of Combinatorial Theory, Series B 41 (1986), no. 1, 92 – 114.
- [8] Raphael Steck, On the edge-erdős-pósa property of walls, 2023.