Decomposition of an Integrally Convex Set into a Minkowski Sum of Bounded and Conic Integrally Convex Sets

Kazuo Murota* and Akihisa Tamura†

June 2023 / October 2023

Abstract

Every polyhedron can be decomposed into a Minkowski sum (or vector sum) of a bounded polyhedron and a polyhedral cone. This paper establishes similar statements for some classes of discrete sets in discrete convex analysis, such as integrally convex sets, L^{\natural} -convex sets, and M^{\natural} -convex sets.

Keywords: Discrete convex analysis, Integrally convex set, L^{\natural} -convex set, M^{\natural} -convex set, Minkowski sum, Characteristic cone.

^{*}The Institute of Statistical Mathematics, Tokyo 190-8562, Japan; and Faculty of Economics and Business Administration, Tokyo Metropolitan University, Tokyo 192-0397, Japan, murota@tmu.ac.jp

[†]Department of Mathematics, Keio University, Yokohama 223-8522, Japan, aki-tamura@math.keio.ac.jp

1 Introduction

As is well known, every polyhedron can be decomposed into a Minkowski sum (or vector sum) of a bounded polyhedron and a polyhedral cone. The objective of this paper is to establish similar decomposition theorems with additional features of integrality and discrete convexity using concepts from discrete convex analysis [\[3,](#page-20-0) [8,](#page-20-1) [9,](#page-20-2) [10,](#page-20-3) [11\]](#page-21-0). Emphasis is laid on integrally convex sets. This notion in discrete convex analysis is equivalent, via convex hull, to that of box-integer polyhedra in the theory of polyhedra [\[19,](#page-21-1) [20\]](#page-21-2) (see Proposition [2.2](#page-3-0) for the precise statement).

Integral convexity is a fundamental concept introduced by Favati–Tardella [\[2\]](#page-20-4) for functions on the integer lattice \mathbb{Z}^n , and integrally convex sets are defined in [\[9,](#page-20-2) Section 3.4] as the set version of integral convexity; see Section [2.2](#page-3-1) for the precise definition. Integral convexity encompasses almost all kinds of discrete convexity proposed so far, such as L^{\natural} convexity, M^{\natural} -convexity, M^{\natural} -convexity, and multimodularity [\[9\]](#page-20-2). A discrete fixed point theorem was formulated by Iimura–Murota–Tamura [\[4\]](#page-20-5) in terms of integrally convex sets (see also [\[10,](#page-20-3) Section 11.9], [\[11,](#page-21-0) Section 13.1] for expositions). Mathematical properties of integrally convex sets and functions have been clarified in recent studies (Moriguchi–Murota [\[5\]](#page-20-6), Moriguchi–Murota–Tamura–Tardella [\[7\]](#page-20-7), Murota–Tamura [\[15,](#page-21-3) [16\]](#page-21-4)). The reader is referred to Murota–Tamura [\[17\]](#page-21-5) for a recent comprehensive survey on integral convexity.

For any sets $S_1, S_2 \subseteq \mathbb{R}^n$, we denote their *Minkowski sum* (or *vector sum*) by $S_1 + S_2$, that is,

$$
S_1 + S_2 = \{x + y \mid x \in S_1, y \in S_2\}.
$$

Let $P \subseteq \mathbb{R}^n$ be a polyhedron. A fundamental fact in the theory of polyhedra says that it can be represented as $P = Q + C$ with a bounded polyhedron Q and a polyhedral cone C (see Section [2.1](#page-2-0) for details). In this decomposition, the cone *C* is uniquely determined from *P*, coinciding with the characteristic (or recession) cone of *P*, whereas there is some degree of freedom in the choice of *Q*. We are interested in integrality and discrete convexity in this decomposition, and our contribution consists of two phases.

In the first phase we consider a box-integer polyhedron *P* and impose an additional condition that *Q* and *C* be box-integer polyhedra. Our first main result, Theorem [3.3,](#page-5-0) states that this is indeed possible. Furthermore, it is shown in Theorem [3.4](#page-6-0) that if P is an L^{\natural} -convex (resp., M^{\natural} -convex) polyhedron, then we can impose that *Q* and *C* be L^{\natural} -convex (resp., M^{\natural} convex). A technical challenge in establishing Theorem [3.3](#page-5-0) for box-integer polyhedra stems from the lack of 'outer description' of box-integer polyhedra in terms of inequality systems. In contrast, inequality systems are available for L^{\natural} -convex and M^{\natural} -convex polyhedra, which makes the proof of Theorem [3.4](#page-6-0) shorter and more transparent.

In the second phase we are concerned with discrete sets $S \subseteq \mathbb{Z}^n$. Our second main result, Theorem [3.6,](#page-8-0) states that an integrally convex set *S* can be represented as $S = T + G$ with a bounded integrally convex set *T* and a 'conic' integrally convex set *G*. Furthermore, it is shown in Theorem [3.7](#page-8-1) that if S is an L^{\nmid} -convex (resp., M^{\nmid} -convex) set, then we can impose that *T* and *G* be L^{\sharp} -convex (resp., M^{\sharp} -convex). A technical challenge in the second phase is to overcome the well-known difficulty of discreteness in the Minkowski summation. Namely, for discrete sets $S_1, S_2 \subseteq \mathbb{Z}^n$, the Minkowski sum $S_1 + S_2$ may possibly be different from $(\overline{S_1} + \overline{S_2}) \cap \mathbb{Z}^n$ (see Figure [1](#page-4-0) of Example [2.1](#page-4-1) for a concrete example). The possibility of $(\overline{S_1} + \overline{S_2}) \cap \mathbb{Z}^n \neq S_1 + S_2$ prevents us to derive the decomposition theorem for integrally convex sets as a corollary of Theorem [3.3](#page-5-0) for box-integer polyhedra.

This paper is organized as follows. Section [2](#page-2-1) is devoted to preliminaries on polyhedra and

integrally convex sets. The main results are described in Section [3.](#page-4-2) Section [3.1](#page-4-3) deals with subsets of \mathbb{R}^n such as box-integer polyhedra, L^{\natural}-convex polyhedra, and M^{\natural}-convex polyhedra, while Section [3.2](#page-7-0) treats subsets of \mathbb{Z}^n such as integrally convex sets, L^{\ti}-convex sets, and M^{\ti}convex sets. The proofs are given in Section [4,](#page-8-2) and Section [5](#page-20-8) concludes the paper.

2 Preliminaries

2.1 Polyhedra

A subset P of \mathbb{R}^n is called a *polyhedron* if it is described by a finite number of linear inequalities, that is, $P = \{x \mid Ax \leq b\}$ for some matrix *A* and a vector *b*. In this paper we always assume that a polyhedron is nonempty. A subset Q of \mathbb{R}^n is called a *polytope* if it is the convex hull of a finite number of points, that is, $Q = \overline{S}$ for a finite subset *S* of \mathbb{R}^n , where \overline{S} denotes the convex hull of *S* . It is known that a polytope is nothing but a bounded polyhedron. A subset *C* of \mathbb{R}^n is called a *cone* if $d \in C$ implies $\lambda d \in C$ for all $\lambda \ge 0$. We follow [\[19,](#page-21-1) [20\]](#page-21-2) for terminology about polyhedra.

Let *P* be a polyhedron. The *characteristic cone* of *P*, denoted by char.cone *P*, is the polyhedral cone given by

$$
char.cone P = \{d \mid x + d \in P \text{ for all } x \text{ in } P\}. \tag{2.1}
$$

The characteristic cone is also called the *recession cone*. The following are basic facts about the characteristic cone:

$$
d \in \text{char.cone } P
$$

\n
$$
\iff \text{there is an } x \text{ in } P \text{ such that } x + \lambda d \in P \text{ for all } \lambda \ge 0,
$$
 (2.2)

$$
d \in \text{char.cone } P
$$

$$
\iff \text{for all } x \text{ in } P \text{, it holds that } x + \lambda d \in P \text{ for all } \lambda \ge 0,
$$
 (2.3)

$$
P + \text{char.cone } P = P,\tag{2.4}
$$

If
$$
P = \{x \mid Ax \le b\}
$$
, then char.cone $P = \{d \mid Ad \le 0\}$. (2.5)

The following is a fundamental theorem, stating that a polyhedron can be decomposed into a Minkowski sum of a polytope and a cone.

Proposition 2.1 (Decomposition theorem for polyhedra).

(1) *Every polyhedron P can be represented as P* = *Q*+*C with some polytope Q and polyhedral cone C.*

(2) *If P* = *Q* + *C, with Q a polytope and C a polyhedral cone, then P is a polyhedron and* $C = \text{char.cone } P$.

It is emphasized that the choice of the polytope Q in $P = Q + C$ is not unique, while C is uniquely determined by *P* as stated in (2).

A polyhedron is said to be *rational* if it is described by a finite number of linear inequalities with rational coefficients. A polyhedron *P* is an *integer polyhedron* if $P = P \cap \mathbb{Z}^n$, i.e., if it coincides with the convex hull of the integer points contained in it, or equivalently, if *P* is rational and each face of *P* contains an integer vector. A polyhedron *P* is called *boxinteger* if $P \cap \{x \in \mathbb{R}^n \mid l \le x \le u\}$ is an integer polyhedron for each choice of integer vectors $l, u \in \mathbb{Z}^n$ with $l \leq u$ ([\[20,](#page-21-2) Section 5.15]). We call a subset *B* of \mathbb{R}^n an *integral box* if $B = \{x \in \mathbb{R}^n \mid l \le x \le u\}$ for some integer vectors $l, u \in \mathbb{Z}^n$ with $l \le u$.

2.2 Integrally convex sets

In this section we introduce the concept of integrally convex sets, as defined in [\[9,](#page-20-2) Section 3.4], and discuss subtleties related to the Minkowski sum of integrally convex sets. The reader is referred to Murota–Tamura [\[17\]](#page-21-5) for technical details of integral convexity including the most recent results.

For $x \in \mathbb{R}^n$ the *integral neighborhood* of *x* is defined by

$$
N(x) = \{ z \in \mathbb{Z}^n \mid |x_i - z_i| < 1 \ (i = 1, 2, \dots, n) \}. \tag{2.6}
$$

It is noted that strict inequality " \lt " is used in this definition and $N(x)$ admits an alternative expression

$$
N(x) = \{ z \in \mathbb{Z}^n \mid \lfloor x_i \rfloor \le z_i \le \lceil x_i \rceil \ (i = 1, 2, \dots, n) \},\tag{2.7}
$$

where, for $t \in \mathbb{R}$ in general, $\lfloor t \rfloor$ denotes the largest integer not larger than t (rounding-down to the nearest integer) and $\lceil t \rceil$ is the smallest integer not smaller than t (rounding-up to the nearest integer). That is, $N(x)$ consists of all integer vectors *z* between $|x| = (|x_1|, |x_2|, \ldots, |x_n|)$ and $[x] = ([x_1], [x_2], \ldots, [x_n])$.

Let *S* be a subset of \mathbb{Z}^n and recall that \overline{S} denotes the convex hull of *S*. As is well known, *S* coincides with the set of all convex combinations of (finitely many) elements of *S* . For any real vector $x \in \mathbb{R}^n$, we call the convex hull of $S \cap N(x)$ the *local convex hull* of *S* around *x*. A nonempty set $S \subseteq \mathbb{Z}^n$ is said to be *integrally convex* if the union of the local convex hulls $S \cap N(x)$ over $x \in \mathbb{R}^n$ is convex. In other words, a set $S \subseteq \mathbb{Z}^n$ is called integrally convex if

$$
\overline{S} = \bigcup_{x \in \mathbb{R}^n} \overline{S \cap N(x)}.
$$
\n(2.8)

This condition is equivalent to saying that every point *x* in the convex hull of *S* is contained in the convex hull of $S \cap N(x)$, i.e.,

$$
x \in \overline{S} \implies x \in \overline{S \cap N(x)}.\tag{2.9}
$$

Obviously, every subset of $\{0, 1\}^n$ is integrally convex.

We say that a set $S \subseteq \mathbb{Z}^n$ is *hole-free* if

$$
S = \overline{S} \cap \mathbb{Z}^n. \tag{2.10}
$$

It is known that an integrally convex set is hole-free; see [\[17,](#page-21-5) Proposition 2.2] for a formal proof. It is also known that the convex hull of an integrally convex set is a polyhedron (Murota–Tamura [\[15,](#page-21-3) Section 4.1]). However, no characterization is known about the inequality systems to describe integrally convex sets.

The concept of integrally convex sets is closely related (or essentially equivalent) to that of box-integer polyhedra as follows.

Proposition 2.2 ([\[13,](#page-21-6) Section 2.2]). *If a set* $S \subseteq \mathbb{Z}^n$ *is integrally convex, then its convex hull* \overline{S} is a box-integer polyhedron and $S = \overline{S} \cap \mathbb{Z}^n$. Conversely, if P is a box-integer polyhedron, *then* $P \cap \mathbb{Z}^n$ *is an integrally convex set and* $P = \overline{P \cap \mathbb{Z}^n}$ *.*

Minkowski summation is an intriguing operation in discrete setting. For two (discrete) sets $S_1, S_2 \subseteq \mathbb{Z}^n$, in general, we have

$$
\overline{S_1 + S_2} = \overline{S_1} + \overline{S_2}
$$
 (2.11)

1: Minkowski sum of discrete sets Figure 1: Minkowski sum of discrete sets

(see, e.g., $[9,$ Proposition 3.17(4)]). In contrast, the naive looking relation

$$
S_1 + S_2 = (\overline{S_1 + S_2}) \cap \mathbb{Z}^n \tag{2.12}
$$

is not always true, as Example [2.1](#page-4-1) below shows.

Example 2.1 ([\[9,](#page-20-2) Example 3.15]). The Minkowski sum of $S_1 = \{(0,0), (1,1)\}$ and $S_2 =$ $(S_1 + S_2)$. That is, the Minkowski sum $S_1 + S_2$ has a 'hole' at (1, 1). See Figure [1.](#page-4-0) of M vex sets is M vex ([11, Section 4.6], [11, Theorem 6.15], [13, Theorem 3.13]). $\{(1, 0), (0, 1)\}\$ is equal to $S_1 + S_2 = \{(1, 0), (0, 1), (2, 1), (1, 2)\}\$, for which $(1, 1) \in (\overline{S_1 + S_2}) \setminus \overline{S_2}$

It may be said that if (2.12) is true for some class of discrete convex sets, this equality for two M^{\natural} -convex sets, since the Minkowski sum of two M^{\natural} -convex sets remains to be M^{\natural} -convex ([\[9,](#page-20-2) Section 4.6], [\[12,](#page-21-7) Section 3.5]). The identity (2.12) also holds for two L^{\natural} -convex sets, since the Minkowski sum of two L^{\natural}-convex sets is integrally convex [\[9,](#page-20-2) Theorem 8.42], although it is not necessarily L^{\t*}-convex. captures a certain essence of the discrete convexity in question. For example, [\(2.12\)](#page-4-4) is true

For the Minkowski sum of integrally convex sets S_1 and S_2 , we observe the following.

- $S_1 + S_2$ may have a 'hole', that is, [\(2.12\)](#page-4-4) may fail (see Example [2.1\)](#page-4-1).
- ω wyski (see Exemple 2.1) • $S_1 + S_2$ may not be integrally convex (see Example [2.1\)](#page-4-1).
- $\overline{S_{\alpha}}$ \cap \mathbb{Z}^n may not be integrally convex (see Example 2.2 below) • $\overline{(S_1 + S_2)} \cap \mathbb{Z}^n$ may not be integrally convex (see Example [2.2](#page-4-5) below).

Example 2.2. Consider $S_1 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ and $S_2 = \{(0, 0, 0), (1, 1, 1)\}$. Their Let $S = (S_1 + S_2) \cap \mathbb{Z}^2$ and consider $x = [(1, 0, 0) + (1, 1, 2)]/2 = (1, 1/2, 1)$ belonging to \overline{S} . We have $N(x) = {(1, 0, 1), (1, 1, 1)}, N(x) ∩ S = {(1, 1, 1)},$ and $x \notin N(x) ∩ S$. Thus the condition [\(2.9\)](#page-3-2) for integral convexity of *S* is violated. This example also shows that the Minkowski sum of box-integer polyhedra is not necessarily box-integer; see also Remark [3.1.](#page-5-1) Minkowski sum is given by $S_1 + S_2 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (2, 1, 1), (1, 2, 1), (1, 1, 2)\}.$ Let $S = (S_1 + S_2) \cap \mathbb{Z}^3$ and consider $x = [(1, 0, 0) + (1, 1, 2)]/2 = (1, 1/2, 1)$ belonging to

Discrepancy between $S_1 + S_2$ and $\overline{S_1 + S_2}$ has attracted considerable attention in (ordinary) convex analysis, leading to the Shapley–Folkman theorem, which has applications in economics, optimization, etc. A recent paper [\[18\]](#page-21-8) of the present authors shows a Shapley– Folkman-type theorem for integrally convex sets.

, in particular, then ≤ ⌊ 3 Results

3.1 Decomposition of box-integer polyhedra

In this section we describe our first main result (Theorem [3.3\)](#page-5-0), a decomposition theorem for box-integer polyhedra. The proof of this theorem relies on the following technical results (in their equivalent reformulations in Propositions [4.1](#page-10-0) and [4.4;](#page-14-0) see Figure [3](#page-10-1) in Section [4\)](#page-8-2).

Proposition 3.1. *The characteristic cone of a box-integer polyhedron is generated by* {−1, 0, +1}*-vectors.*

Proposition 3.2. *The characteristic cone of a box-integer polyhedron is box-integer.*

The proofs of these propositions are quite long and involved, probably because no characterization is known about inequality systems to describe box-integer polyhedra. The proofs of Propositions [3.1](#page-5-2) and [3.2](#page-5-3) are given in Sections [4.1](#page-10-2) and [4.2,](#page-14-1) respectively. Our decomposition theorem for box-integer polyhedra is as follows.

Theorem 3.3. *Every box-integer polyhedron P can be represented as*

$$
P = Q + C \tag{3.1}
$$

п

with a bounded box-integer polyhedron Q and a box-integer polyhedral cone C.

Proof. By Proposition [2.1,](#page-2-2) we can decompose *P* as $P = \hat{Q} + C$, where \hat{Q} is a polytope and *C* is the characteristic cone of *P*. The cone *C* is box-integer by Proposition [3.2.](#page-5-3) Take a bounded integral box *B* containing \hat{Q} and define $Q := P \cap B$, which is a bounded box-integer polyhedron. Since

$$
Q = P \cap B = (\hat{Q} + C) \cap B \supseteq \hat{Q} \cap B = \hat{Q},
$$

we obtain

$$
Q + C \supseteq \hat{Q} + C = P.
$$

The reverse inclusion $Q + C \subseteq P$ follows from $Q \subseteq P$ and $P + C = P$ in [\(2.4\)](#page-2-3) as $Q + C \subseteq$ $P + C = P$.

Remark 3.1. In view of Proposition [2.1\(](#page-2-2)2) we may be tempted to imagine that if *Q* is a bounded box-integer polyhedron and *C* is a box-integer polyhedral cone, then $Q + C$ is a box-integer polyhedron. But this is not the case. A counterexample can be constructed from Example [2.2.](#page-4-5) Let *Q* be the convex hull of $S_1 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}\$ and *C* be the polyhedral cone generated by $S'_2 = \{(1, 1, 1)\}\$, that is, $C = \{\lambda(1, 1, 1) \mid \lambda \ge 0\}$. Both *Q* and *C* are box-integer, but $Q + C$ is not. Indeed, $T = (Q + C) \cap \mathbb{Z}^3$ is not integrally convex, because $x = [(1, 0, 0) + (0, 0, 1)]/2 + (1, 1, 1)/2 = (1, 1/2, 1) \in \overline{T}$, $N(x) = {(1, 0, 1), (1, 1, 1)}$, *N*(*x*) ∩ *T* = {(1, 1, 1)}, and *x* ∉ *N*(*x*) ∩ *T*.

Remark 3.2. By Proposition [3.1,](#page-5-2) a box-integer cone is generated by $\{-1, 0, +1\}$ -vectors, but the latter property does not characterize a box-integer cone. Consider the cone *C* generated by $(1, 1, 0, 1), (0, 1, 1, 1), (1, 0, 1, 1),$ that is,

$$
C = \{x \mid x = \alpha_1(1, 1, 0, 1) + \alpha_2(0, 1, 1, 1) + \alpha_3(1, 0, 1, 1), \alpha_1, \alpha_2, \alpha_3 \ge 0\}
$$

=
$$
\{x \mid x = (\alpha_1 + \alpha_3, \alpha_1 + \alpha_2, \alpha_2 + \alpha_3, \alpha_1 + \alpha_2 + \alpha_3) : \alpha_1, \alpha_2, \alpha_3 \ge 0\}.
$$

For $\alpha_1 = \alpha_2 = \alpha_3 = 1/2$, we have $x = (1, 1, 1, 3/2)$ and $N(x) = \{(1, 1, 1, 2), (1, 1, 1, 1)\}\$. But $(1, 1, 1, 2) \notin C$ and $(1, 1, 1, 1) \notin C$, and hence $N(x) \cap C = \emptyset$. This shows that $C \cap \mathbb{Z}^4$ is not integrally convex, and hence *C* is not box-integer.

Theorem [3.3](#page-5-0) can be adapted to some classes of integer polyhedra treated in discrete convex analysis, such as L^{\natural}-convex and M^{\natural}-convex polyhedra. An L^{\natural}-convex polyhedron is, by definition, an integer polyhedron obtained as the convex hull of an L^{\sharp}-convex set. It is known that an L^{\natural} -convex polyhedron *P* can be described as

$$
P = \{x \in \mathbb{R}^n \mid l_i \le x_i \ (i \in I), x_j \le u_j \ (j \in J), x_j - x_i \le d_{ij} \ ((i, j) \in E)\}
$$
(3.2)

for some *I*, *J* ⊆ {1, 2, . . . , *n*}, *E* ⊆ {1, 2, . . . , *n*} × {1, 2, . . . , *n*}, l_i ∈ ℤ (i ∈ *I*), u_j ∈ ℤ (j ∈ *J*), and $d_{ij} \in \mathbb{Z}$ ((*i*, *j*) \in *E*), and the converse is also true. An *L*^{\uparrow}-*convex cone* means an L^{\uparrow}-convex polyhedron that is a cone. An *M*♮ *-convex polyhedron* is a synonym of an integral generalized polymatroid, and hence an M♮ -convex polyhedron *P* is described as

$$
P = \{x \in \mathbb{R}^n \mid \mu(X) \le x(X) \le \rho(X) \, (\forall X \subseteq \{1, 2, \dots, n\})\},\tag{3.3}
$$

where $x(X) = \sum_{i \in X} x_i$, for a (strong or paramodular) pair of an integer-valued supermodular function μ and an integer-valued submodular function ρ (cf., [\[3,](#page-20-0) Section 3.5(a)], [\[9,](#page-20-2) Section 4.7]); μ and ρ are allowed to take $-\infty$ and $+\infty$, respectively. An M^{\sharp} -*convex cone* is defined in an obvious manner. Other kinds of polyhedra (such as L_2^{\natural} -convex polyhedron, M_2^{\natural} -convex polyhedron, and multimodular polyhedron) are defined similarly from the corresponding notions for sets of integer vectors. More precisely, an L_2^{\natural} -convex set is defined as the Minkowski sum of two L^{\natural} -convex sets and an L_2^{\natural} -convex polyhedron is the convex hull of an L_2^{\natural} -convex set, implying that an L_2^{\natural} -convex polyhedron can also be defined as the Minkowski sum of two L^{\natural} -convex polyhedra. Similarly, an M_2^{\natural} -convex set is defined as the intersection of two M^{\natural} -convex sets and an M_2^{\natural} -convex polyhedron is the convex hull of an M_2^{\natural} -convex set; then it is known (cf., e.g., [\[9,](#page-20-2) Theorem 4.22]) that an M_2^{\dagger} -convex polyhedron can also be defined as the intersection of two M^{\†}-convex polyhedra.

The adaptation of Theorem [3.3](#page-5-0) to specific classes is given in Theorem [3.4](#page-6-0) below. It should be clear that, although L^{\natural} -convex polyhedra, etc., constitute subclasses of box-integer polyhedra, Theorem [3.3](#page-5-0) does not imply the corresponding statements for these subclasses. It is worth noting that the proofs for these special cases do not rely on Theorem [3.3](#page-5-0) and that they are shorter and simpler because of the inequality descriptions known for these special cases (see Murota [\[9\]](#page-20-2), Moriguchi–Murota [\[6,](#page-20-9) Table 1], Murota–Tamura [\[17,](#page-21-5) Table 1]).

Theorem 3.4.

(1) Every L^{\natural} -convex polyhedron P can be represented as $P = Q + C$ with a bounded L^{\natural} -convex *polyhedron Q and an L*♮ *-convex cone C.*

(2) *Every* L_2^{\natural} -convex polyhedron P can be represented as $P = Q + C$ with a bounded L_2^{\natural} -convex *polyhedron Q and an L*♮ 2 *-convex cone C.*

(3) *Every* M^{\natural} -convex polyhedron P can be represented as $P = Q + C$ with a bounded M^{\natural} *convex polyhedron Q and an M*♮ *-convex cone C. Similarly for an M-convex polyhedron P, with Q and C being M-convex.*

(4) *Every* M_2^{\natural} -convex polyhedron P can be represented as $P = Q + C$ with a bounded M_2^{\natural} *convex polyhedron Q and an* M_2^{\natural} -convex cone C. Similarly for an M₂-convex polyhedron P, *with Q and C being M₂-convex.*

(5) *Every multimodular polyhedron P can be represented as* $P = Q + C$ *with a bounded multimodular polyhedron Q and a multimodular cone C.*

Proof. (1) The proof of Theorem [3.3](#page-5-0) can be adapted to an $L^{\frac{1}{2}}$ -convex polyhedron on the basis of the following properties of an L^{\sharp} -convex polyhedron.

- 1. The characteristic cone of an L^{\natural} -convex polyhedron is L^{\natural} -convex.
- 2. The intersection of an L^{\natural} -convex polyhedron with an integral box is L^{\natural} -convex.

We can prove the first statement by making use of the fact that an L^{\sharp} -convex polyhedron *P* ⊆ \mathbb{R}^n is described as [\(3.2\)](#page-5-4). It follows from (3.2) and [\(2.5\)](#page-2-4) that the characteristic cone of

P is given by $C = \{x \mid 0 \le x_i \ (i \in I), x_i \le 0 \ (j \in J), x_i - x_i \le 0 \ ((i, j) \in E) \}$, which is also an L^{\natural} -convex polyhedron. The second statement also follows from [\(3.2\)](#page-5-4). We consider the decomposition $P = \hat{Q} + C$ in Proposition [2.1,](#page-2-2) take a bounded integral box *B* containing \hat{Q} , and define $Q := P \cap B$, for which we can show $P = Q + C$ as in the proof of Theorem [3.3.](#page-5-0)

(2)–(5) These cases are proved in Section [4.4](#page-18-0) by using a unified proof scheme consistent with the case of (1). \Box

Remark 3.3. Theorem [3.4\(](#page-6-0)1) gives a decomposition of an L^{\nexists} -convex polyhedron. However, we cannot obtain a similar statement for an L-convex polyhedron, simply because there is no bounded L-convex polyhedron. Note that an L-convex polyhedron *P* has the invariance in the direction of $1 = (1, 1, \ldots, 1)$ in the sense that $x \in P$ implies $x + \lambda 1 \in P$ for all $\lambda \in \mathbb{R}$. Similarly, there is no bounded L_2 -convex polyhedron.

Remark 3.4. In each case of Theorem [3.4,](#page-6-0) the polyhedron *P* is necessarily an integer polyhedron. Recall that we have defined P to be an L^{\natural} -convex polyhedron if it is the convex hull of an L^{\natural} -convex set $S \subseteq \mathbb{Z}^n$). In the literature of discrete convex analysis, the notion of L^{\natural}-convexity is generalized to non-integer polyhedra (Murota–Shioura [\[14\]](#page-21-9)). An L^{\natural}-convex polyhedron (not necessarily integral) is described by [\(3.2\)](#page-5-4) with $l_i \in \mathbb{R}$ ($i \in I$), $u_j \in \mathbb{R}$ ($j \in J$), and $d_{ij} \in \mathbb{R}$ ((*i*, *j*) $\in E$). For an L^{\sharp}-convex polyhedron *P* in this generalized sense, we also obtain the decomposition $P = Q + C$. Similar generalizations are possible for M^{\ti}-convex polyhedra, etc., in (2)–(5) of Theorem [3.4.](#page-6-0)

3.2 Decomposition of integrally convex sets

Theorem [3.3](#page-5-0) for box-integer polyhedra can be rephrased for integrally convex sets as follows.

Corollary 3.5. The convex hull \overline{S} of an integrally convex set $S \subseteq \mathbb{Z}^n$) can be represented as

$$
\overline{S} = Q + C \tag{3.4}
$$

 $$

Proof. Since *S* is integrally convex, \overline{S} is a box-integer polyhedron by Proposition [2.2.](#page-3-0) By Theorem [3.3](#page-5-0) applied to \overline{S} we obtain the decomposition [\(3.4\)](#page-7-1), where Q is a bounded boxinteger polyhedron and *C* is a box-integer cone. Then $Q \cap \mathbb{Z}^n$ and $C \cap \mathbb{Z}^n$ are integrally convex by Proposition [2.2.](#page-3-0)

While the decomposition $\overline{S} = Q + C$ in [\(3.4\)](#page-7-1) is defined via embedding of *S* into \mathbb{R}^n , our second main result (Theorem [3.6](#page-8-0) below) establishes a decomposition of an integrally convex set *S* directly within \mathbb{Z}^n . We emphasize the difference between $\overline{S} = Q + C$ and

$$
S = (Q \cap \mathbb{Z}^n) + (C \cap \mathbb{Z}^n). \tag{3.5}
$$

We can show " $(3.5) \Rightarrow (3.4)$ $(3.5) \Rightarrow (3.4)$ $(3.5) \Rightarrow (3.4)$ " as

$$
\overline{S} = \overline{(Q \cap \mathbb{Z}^n) + (C \cap \mathbb{Z}^n)} = \overline{Q \cap \mathbb{Z}^n} + \overline{C \cap \mathbb{Z}^n} = Q + C,
$$

where $\overline{S_1 + S_2} = \overline{S_1} + \overline{S_2}$ in [\(2.11\)](#page-3-3) is used. However, the converse "[\(3.5\)](#page-7-2) \Leftarrow [\(3.4\)](#page-7-1)" is not always true (see Example [3.1](#page-8-3) below). Thus, [\(3.5\)](#page-7-2) is (strictly) stronger than [\(3.4\)](#page-7-1).

To state the theorem we need to introduce a terminology. We call a set $G \subseteq \mathbb{Z}^n$ a *conic set* if its convex hull \overline{G} is a cone. An integrally convex set *G* is conic if and only if $G = C \cap \mathbb{Z}^n$ for some box-integer cone *C*.

Theorem 3.6. *Every integrally convex set* $S \subseteq \mathbb{Z}^n$ *) can be represented as*

$$
S = T + G \tag{3.6}
$$

with a bounded integrally convex set T and a conic integrally convex set G.

Proof. The proof, to be given in Section [4.3,](#page-16-0) is based on propositions equivalent to Proposi-tions [3.1](#page-5-2) and [3.2.](#page-5-3) \Box

Example 3.1. We compare the decompositions in Corollary [3.5](#page-7-3) and Theorem [3.6](#page-8-0) for a simple two-dimensional example. Let *S* be an infinite subset of \mathbb{Z}^2 depicted at the top left of Figure [2,](#page-9-0) which can be described, e.g., as $S = \{x \in \mathbb{Z}^2 \mid x_1 + x_2 \ge 1, |x_1 - x_2| \le 1\}$. This set *S* is integrally convex, and the convex hull \overline{S} is a box-integer polyhedron described as $\overline{S} = \{x \in S\}$ $\mathbb{R}^2 | x_1 + x_2 \ge 1$, $|x_1 - x_2| \le 1$. Let *Q* be the line segment connecting (1,0) and (0, 1) and *C* be the semi-infinite line starting at $(0, 0)$ and emanating in the direction of $(1, 1)$. Both *Q* and *C* are box-integer, and we obtain the decomposition $\overline{S} = Q + C$ in Corollary [3.5.](#page-7-3) The semi-infinite line *C* is, in fact, the characteristic cone of \overline{S} . Both $Q \cap \mathbb{Z}^2$ and $C \cap \mathbb{Z}^2$ are integrally convex, but the identity $S = (Q \cap \mathbb{Z}^2) + (C \cap \mathbb{Z}^2)$ in [\(3.5\)](#page-7-2) fails, because of the 'holes' in $(Q \cap \mathbb{Z}^2) + (C \cap \mathbb{Z}^2)$ at $x = (t, t)$ for integers $t \geq 1$. With the choice of *T* = $(Q ∩ \mathbb{Z}^2) ∪ {(1, 1)} = {(1, 0), (0, 1), (1, 1)}$ and $G = C ∩ \mathbb{Z}^2 = {(t, t) | t ≥ 0, t ∈ \mathbb{Z}}$, we obtain the decomposition $S = T + G$ in Theorem [3.6.](#page-8-0) Here both *T* and *G* are integrally convex.

Theorem [3.6](#page-8-0) can be adapted to some classes of discrete convex sets in discrete convex analysis, such as L^{\natural} -convex and M^{\natural} -convex sets (see Murota [\[9\]](#page-20-2) for definitions of these concepts). The corresponding statements for these subclasses are given in Theorem [3.7](#page-8-1) below. It is emphasized that Theorem [3.7](#page-8-1) does not follow from Theorem [3.6](#page-8-0) (for general inte-grally convex sets) nor from Theorem [3.4](#page-6-0) (for L^{\natural}-convex polyhedra, etc.). Note that we have $S, T, G \subseteq \mathbb{Z}^n$ in Theorem [3.7,](#page-8-1) whereas $P, Q, C \subseteq \mathbb{R}^n$ in Theorem [3.4.](#page-6-0)

Theorem 3.7.

(1) *Every* L^{\natural} -convex set *S* can be represented as $S = T + G$ with a bounded L^{\natural} -convex set *T and a conic L*♮ *-convex set G.*

(2) *Every* L_2^{\natural} -convex set *S* can be represented as $S = T + G$ with a bounded L_2^{\natural} -convex set *T and a conic L*♮ 2 *-convex set G.*

(3) *Every* M^{\natural} -convex set S can be represented as $S = T + G$ with a bounded M^{\natural} -convex set T *and a conic M*♮ *-convex set G. Similarly for an M-convex set S , with T and G being M-convex.* (4) *Every* M_2^{\natural} -convex set *S* can be represented as $S = T + G$ with a bounded M_2^{\natural} -convex *set T and a conic M*♮ 2 *-convex set G. Similarly for an M*2*-convex set S , with T and G being M*2*-convex.*

(5) *Every multimodular set S can be represented as S* = *T* +*G with a bounded multimodular set T and a conic multimodular set G.*

Proof. The proof is given in Section [4.5.](#page-19-0) □

4 Proofs

The structure of the proofs (dependence among propositions and theorems) is shown in the diagram in Figure [3.](#page-10-1)

Figure 2: $Q + C = \overline{S}$, $(Q \cap \mathbb{Z}^2) + (C \cap \mathbb{Z}^2) \neq S$, and $T + G = S$

Figure 3: Dependence among propositions and theorems

S is integrally convex

S is not integrally convex

Figure 4: Necessity of integral convexity in Proposition [4.2](#page-10-3)

4.1 Proof of Proposition [3.1](#page-5-2)

In this section we prove Proposition [3.1,](#page-5-2) stating that the characteristic cone of a box-integer polyhedron is generated by $\{-1, 0, +1\}$ -vectors. By Proposition [2.2,](#page-3-0) this statement can be rephrased (equivalently) in terms of integral convexity as follows.

Proposition 4.1. Let $S \subseteq \mathbb{Z}^n$ be an integrally convex set. The characteristic cone C of its *convex hull* \overline{S} is generated by vectors in $\{-1,0,+1\}^n$. In particular, C is an integer polyhedron.

Proof. Take any $d \in C$ with $||d||_{\infty} = 1$. Proposition [4.2](#page-10-3) below shows that there exist $d^1, d^2, \ldots, d^h \in N(d)$ such that $d \in \{d^1, d^2, \ldots, d^h\}$, where $N(d)$ denotes the integral neighborhood of *d* defined in [\(2.6\)](#page-3-4). We have $N(d) \subseteq \{-1, 0, +1\}^n$ since $||d||_{\infty} = 1$. □

 $\frac{1}{2}$ **Proposition 4.2.** *Let* $S \subseteq \mathbb{Z}^n$ *be an integrally convex set,* $x^0 \in S$ *, and* $d \in \mathbb{R}^n$ *with* $||d||_{\infty} = 1$ *. If*

$$
x^0 + \lambda d \in \overline{S} \quad \text{for all } \lambda \ge 0,
$$
\n
$$
(4.1)
$$

there exist $d^1, d^2, \ldots, d^h \in N(d)$ *such that* $d \in \{d^1, d^2, \ldots, d^h\}$ *and*

$$
x^{0} + kd^{j} \in S \qquad (j = 1, 2, ..., h; k = 1, 2, ...). \tag{4.2}
$$

The condition [\(4.1\)](#page-10-4) is equivalent to saying that *d* belongs to the characteristic cone of converse is not true because [\(4.2\)](#page-10-5) imposes an additional requirement of integrality. The role *S*. The condition [\(4.2\)](#page-10-5) implies that each d^j belongs to the characteristic cone of \overline{S} , but the

Figure 5: Notations in Lemma [4.3](#page-11-0)

of integral convexity of *S* is illustrated in Figure [4.](#page-10-6) In the left panel, the set *S* is integrally convex, while S is not integrally convex in the right, where d^1 does not meet the condition in [\(4.2\)](#page-10-5).

To prove Proposition [4.2,](#page-10-3) we need the following general lemma concerning a set of $\{0, 1\}$ -vectors. Figure [5](#page-11-1) illustrates this lemma when $X = \{0, 1\}^2$, where we think of *R* and *B* as sets of 'red' and 'black' points, respectively, which are disjoint by [\(4.3\)](#page-11-2).

Lemma 4.3. Let $X = \{0, 1\}^m$. For any $R \subseteq X$ and $d \in \overline{X} \setminus \overline{R}$, there exists some $B \subseteq X$ that *satisfies the following conditions:*

$$
R \subseteq X \setminus B \qquad (i.e., \ R \cap B = \emptyset), \tag{4.3}
$$

$$
d \notin \overline{X \setminus B},\tag{4.4}
$$

$$
\overline{B} \cap (\overline{X \setminus B}) = \emptyset. \tag{4.5}
$$

Moreover, the elements of B can be ordered as $B = \{d^1, d^2, \ldots, d^l\}$ *(where* $l = |B|$ *) so as to satisfy*

$$
\overline{\{d^1, d^2, \dots, d^i\}} \cap \overline{\{d^i, d^{i+1}, \dots, d^l\} \cup (X \setminus B)} = \{d^i\} \quad \text{for } i = 1, 2, \dots, l. \tag{4.6}
$$

Proof. We first point out that [\(4.6\)](#page-11-3) is a refinement of [\(4.5\)](#page-11-4). Indeed, (4.6) for $i = l$ reads $\overline{B} \cap \{d^l\} \cup (X \setminus B) = \{d^l\}.$ Since $d^l \notin \overline{X \setminus B}$, this implies $\overline{B} \cap (\overline{X \setminus B}) = \emptyset$ in [\(4.5\)](#page-11-4).

In the (special) case where the given vector *d* belongs to *X*, *d* is an extreme point of \overline{X} and hence we can take $B = \{d\}$ to meet the requirements [\(4.3\)](#page-11-2), [\(4.4\)](#page-11-5), and [\(4.6\)](#page-11-3). In the following we assume $d \notin X$.

The given subset *R* may be empty or nonempty. Suppose first that $R \neq \emptyset$. Since $d \notin \overline{R}$, the point *d* can be separated from \overline{R} by a hyperplane. More precisely, there exists a hyperplane $H = \{x \mid a^\top x = \delta\}$, where $\delta \in \mathbb{R}$, $a \in \mathbb{R}^n$, and $a^\top x = \sum_{i=1}^n a_i x_i$, such that the (open) half spaces $H^+ := \{x \mid a^\top x > \delta\}$ and $H^- := \{x \mid a^\top x < \delta\}$ contain *d* and \overline{R} , respectively. It follows from *d* ∈ *H*⁺ and \overline{R} ⊆ *H*[−] that *B* := *H*⁺ ∩ *X* meets the requirements of [\(4.3\)](#page-11-2) and [\(4.4\)](#page-11-5). Indeed, *B* ⊆ *X* \ *R* in [\(4.3\)](#page-11-2) follows from *B* ⊆ *H*⁺ and *R* ⊆ *H*⁻, and *d* ∉ $\overline{X \setminus B}$ in [\(4.4\)](#page-11-5) follows from *d* ∈ *H*⁺ and *X* \ *B* ⊆ *H*[−] ∪ *H*. To meet [\(4.6\)](#page-11-3), we perturb the vector *a* so that $a^{\top}x$ are distinct for $x \in X \cup \{d\}$, and number the elements of $B = \{d^1, d^2, \dots, d^l\}$ so that $a^{\top}d^1 > a^{\top}d^2 > \dots > a^{\top}d^l$. In the remaining (rather exceptional) case where $R = \emptyset$, we choose a vector *a* for which $a^{\dagger} x$ are distinct for $x \in X \cup \{d\}$, and define $\delta := a^{\top}d - \varepsilon$ with a sufficiently small positive ε . Using such (a, δ) we define *H*, *H*⁺, *H*⁻, and *B* = *H*⁺ \cap *X*. The rest of the argument is the same as in the case of $R \neq \emptyset$.

We are ready to begin the proof of Proposition [4.2.](#page-10-3) Without loss of generality, we may assume *d* ≥ 0, since integral convexity of *S* is preserved under coordinate inversions x_i → − x_i for *i* in an arbitrary subset of $\{1, 2, \ldots, n\}$.

Let *X* := *N*(*d*). Then $X \subseteq \{0, 1\}^n$ and $d \in \overline{X}$. Up to a permutation of coordinates, *X* is equal to a set of the form $\{1\}^p \times \{0\}^q \times \{0, 1\}^m$ $(p + q + m = n; p, q, m \ge 0)$, so that we may identify *X* with $\{0, 1\}^m$. Define

$$
R := \{ d' \in X \mid x^0 + kd' \in S \ (k = 1, 2, \ldots) \},\tag{4.7}
$$

or equivalently, $R := X \cap \text{char.cone } \overline{S}$. Then we have $d \in \overline{R}$ if and only if there exist $d^1, d^2, \ldots, d^h \in N(d)$ satisfying $d \in \{d^1, d^2, \ldots, d^h\}$ and [\(4.2\)](#page-10-5). That is, our goal is to show $d \in \overline{R}$. To prove this by contradiction, we assume $d \notin \overline{R}$.

We have $R \subseteq X$ and $d \in \overline{X} \setminus \overline{R}$, where *X* can be identified with $\{0, 1\}^m$. This allows us to use Lemma [4.3](#page-11-0) to obtain $B = \{d^1, d^2, ..., d^l\}$ ($\subseteq X$) satisfying [\(4.3\)](#page-11-2)–[\(4.6\)](#page-11-3). Let $j^* \in \{0, 1, ..., l\}$ be the (uniquely determined) number such that

$$
x^{0} + d^{i} \notin S \quad (i = 1, 2, ..., j^{*}), \qquad x^{0} + d^{j^{*}+1} \in S,
$$
\n(4.8)

where $j^* = 0$ if $x^0 + d^1 \in S$, and $j^* = l$ if $x^0 + d^i \notin S$ for all $i = 1, 2, ..., l$. Using this index j^* we define $B^* = \{d^1, d^2, \dots, d^{j^*}\}\$. Note that $B^* = \emptyset$ if $j^* = 0$, and $B^* = B$ if $j^* = l$.

Let $y := x^0 + d$. By the assumption [\(4.1\)](#page-10-4), namely, $d \in \text{char.cone } \overline{S}$, we have $y \in \overline{S}$, which, in turn, implies $y \in N(y) \cap S$ by integral convexity of *S*. It follows from $N(d) = X$ and the definition of B^* that^{[1](#page-12-0)}

$$
N(y) \cap S = N(x^0 + d) \cap S = x^0 + \{d' \in X \mid x^0 + d' \in S\} \subseteq x^0 + (X \setminus B^*).
$$

Hence $y \in \overline{N(y) \cap S} \subseteq x^0 + \overline{X \setminus B^*}$, that is, $d \in \overline{X \setminus B^*}$. On the other hand, $d \notin \overline{X \setminus B}$ as shown in [\(4.4\)](#page-11-5). Thus we obtain

$$
d \in \overline{X \setminus B^*}, \qquad d \notin \overline{X \setminus B}.\tag{4.9}
$$

If $B^* = B$, these two assertion contradict each other, and we are done. If B^* is a proper subset of *B*, we cannot derive a contradiction from [\(4.9\)](#page-12-1).

We overcome this difficulty as follows. Although the definition of *R* in [\(4.7\)](#page-12-2) refers to x^0 , it is, in fact, independent of the initial point x^0 , as seen from the alternative expression $R = X \cap \text{char.cone } \overline{S}$. The set *B* is also independent of x^0 , whereas B^* , defined via [\(4.8\)](#page-12-3), varies with x^0 , that is, $B^* = B^*(x^0)$. Our strategy is to show that, if $B^*(x^0) \neq B$, we can choose another initial point x^1 satisfying $B^*(x^0) \subsetneq B^*(x^1)$. By repeating this process, we can increase B^* until $B^* = B$. Then we obtain a contradiction from [\(4.9\)](#page-12-1), to complete the proof of Proposition [4.2.](#page-10-3)

Since d^{j^*+1} ∈ *B* and $R \cap B = \emptyset$ (cf. [\(4.3\)](#page-11-2)), we have $d^{j^*+1} \notin R$, while $x^0 + d^{j^*+1} \in S$ by [\(4.8\)](#page-12-3). Therefore, there exists a positive integer $k^* \geq 1$ such that

$$
x^{0} + kd^{j^{*}+1} \in S \quad (k = 1, 2, ..., k^{*}), \qquad x^{0} + (k^{*} + 1)d^{j^{*}+1} \notin S. \tag{4.10}
$$

This integer k^* represents the maximum number of steps from x^0 toward d^{j^*+1} to stay in *S*. We define $x^1 := x^0 + k^* d^{j^*+1}$, which is a point in *S*. We shall show $B^*(x^0) \subsetneq B^*(x^1)$ by proving

$$
x^{1} + d^{i} \notin S \quad (i = 1, 2, ..., j^{*}), \tag{4.11}
$$

$$
x^1 + d^{j^*+1} \notin S. \tag{4.12}
$$

The second property [\(4.12\)](#page-12-4) is easy to prove. Namely,

$$
x^{1} + d^{j^{*}+1} = (x^{0} + k^{*}d^{j^{*}+1}) + d^{j^{*}+1} = x^{0} + (k^{*} + 1)d^{j^{*}+1} \notin S
$$

using the definition of k^* in [\(4.10\)](#page-12-5). To prove [\(4.11\)](#page-12-6), we consider a sequence of intermediate points, say, *x'*, *x''*,... between *x*⁰ and *x*¹, where *x'* := *x*⁰ + *d*^{*j**+1}, *x''* := *x*⁰ + 2*d*^{*j**+1}, etc.

¹For any vector *x* and set *Y*, we use abbreviation $x + Y$ for $\{x\} + Y$.

Claim 4.1. *For* $x' = x^0 + d^{j^*+1}$ *we have* $x' \in S$ *and*

$$
x' + d^i \notin S \quad (i = 1, 2, \dots, j^*). \tag{4.13}
$$

Proof. First, we see $x' \in S$ from [\(4.10\)](#page-12-5). To prove [\(4.13\)](#page-13-0), fix *i* ($1 \le i \le j^*$) and define \hat{d} := $(d^{j^*+1} + d^i)/2$. We have

$$
\hat{d} = (d^{j^*+1} + d^i)/2 \in \overline{\{d^{j^*+1}, d^i\}} \subseteq \overline{B^* \cup \{d^{j^*+1}\}}.
$$

Since $\hat{d} \neq d^{j^*+1}$ (which is equivalent to $d^i \neq d^{j^*+1}$) and

$$
\overline{B^* \cup \{d^{j^*+1}\} \cap X \setminus B^*}
$$
\n
$$
= \{d^1, d^2, \dots, d^{j^*+1}\} \cap \{d^{j^*+1}, \dots, d^l\} \cup (X \setminus B)
$$
\n
$$
= \{d^{j^*+1}\}\
$$

by (4.6) , we have

$$
\hat{d} \notin \overline{X \setminus B^*}.\tag{4.14}
$$

On the other hand, it follows from the definition of B^* that

$$
x^{0} + \overline{X \setminus B^{*}} \supseteq \overline{N(x^{0} + \hat{d}) \cap S}.
$$
 (4.15)

Combining (4.14) and (4.15) we obtain

$$
x^0 + \hat{d} \notin \overline{N(x^0 + \hat{d}) \cap S}.
$$
\n(4.16)

If $x' + d^i \in S$ were true, we would obtain

$$
x^{0} + \hat{d} = x^{0} + \frac{1}{2}(d^{j^{*}+1} + d^{i}) = \frac{1}{2}x^{0} + \frac{1}{2}(x' + d^{i}) \in \overline{S},
$$
\n(4.17)

which is a contradiction to [\(4.16\)](#page-13-3), since *S* is integrally convex. Therefore, we must have $x' + d^i \notin S$, proving [\(4.13\)](#page-13-0).

For the second intermediate point $x'' = x^0 + 2d^{j^*+1} = x' + d^{j^*+1}$, we can prove

 $x'' \in S$, $x'' + d^i \notin S$ $(i = 1, 2, ..., j^*)$

in a similar manner, by replacing (x^0, x') in the proof of Claim [4.1](#page-13-4) by (x', x'') . Continuing in this way, we can show the statement [\(4.11\)](#page-12-6) at the new initial point x^1 where $B^*(x^1)$ is strictly larger than $B^*(x^0)$.

If $B^*(x^1) = B$, we are done, with a contradiction from [\(4.9\)](#page-12-1). Otherwise, we repeat the same procedure to obtain a (finite) sequence x^0, x^1, \ldots, x^s of initial points such that the associated *B*^{*} increases to *B*, i.e., $B^*(x^0) \subsetneq B^*(x^1) \cdots \subsetneq B^*(x^s) = B$. This completes the proof of Proposition [4.2.](#page-10-3)

4.2 Proof of Proposition [3.2](#page-5-3)

In this section we prove Proposition [3.2,](#page-5-3) stating that the characteristic cone of a box-integer polyhedron is box-integer. By Proposition [2.2,](#page-3-0) this statement can be rephrased (equivalently) in terms of integral convexity as follows.

Proposition 4.4. Let $S \subseteq \mathbb{Z}^n$ be an integrally convex set. The characteristic cone C of its *convex hull* \overline{S} *has the property that* $C \cap \mathbb{Z}^n$ *is integrally convex.*

We begin the proof of Proposition [4.4](#page-14-0) by observing that the convex hull \overline{S} can be represented as $\overline{S} = Q + C$ with a bounded box-integer polyhedron O and a polyhedral cone C. Indeed, by Proposition [2.1,](#page-2-2) we can decompose \overline{S} as $\overline{S} = \hat{Q} + C$, where \hat{Q} is a polytope and *C* is the characteristic cone of \overline{S} . Take a bounded integral box *B* containing \hat{Q} and define *Q* := \overline{S} ∩ *B*, which is a bounded box-integer polyhedron. Since $Q = \overline{S}$ ∩ *B* = (\hat{Q} + *C*) ∩ *B* ⊇ $Q \cap B = Q$ ⁿ, we obtain $Q + C \supseteq Q + C = \overline{S}$. The reverse inclusion $Q + C \subseteq \overline{S}$ follows from $\overline{Q} \subseteq \overline{S}$ and \overline{S} + $C = \overline{S}$ (cf. [\(2.4\)](#page-2-3)) as $Q + C \subseteq \overline{S} + C = \overline{S}$.

We prove Proposition [4.4](#page-14-0) by contradiction. Namely, we assume that $C \cap \mathbb{Z}^n$ is not integrally convex and derive a contradiction to the integral convexity of *S* . We shall construct a point $y^* \in \overline{S}$ with the property $y^* \notin \overline{N(y^*) \cap S}$. We start with an arbitrary $x^0 \in Q \cap \mathbb{Z}^n$ and find a point $y^0 \in x^0 + C$ with some properties (Claim [4.2](#page-14-2) below). We consider a system of inequalities describing $x^0 + C$. With reference to the inequalities tight at y^0 , we find a vertex x^* of *Q*. Then the point *y*^{*} is constructed as $y^* = y^0 + (x^* - x^0)$ in [\(4.29\)](#page-15-0) below.

Recalling that *Q* is a nonempty integer polyhedron, take any $x^0 \in Q \cap \mathbb{Z}^n$ and define

$$
D := x^0 + C, \qquad D_I := D \cap \mathbb{Z}^n.
$$

By Proposition [4.1,](#page-10-0) *C* is an integer polyhedron, which implies that *D* is an integer polyhedron and $D = D_I$. The set D_I is not integrally convex as a consequence of the assumption that $C \cap \mathbb{Z}^n$ is not integrally convex.

Claim 4.2. *There exists* $y^0 \in D$ *that satisfies the following conditions:*

$$
y^0 \notin \overline{N(y^0) \cap D_I},\tag{4.18}
$$

$$
y^0 \text{ is a vertex of } \overline{N(y^0)} \cap D, \tag{4.19}
$$

y 0 is a relative interior point of *N*(*y* 0 (4.20)

Proof. Since D_I is not integrally convex, there exists $z \in \overline{D_I}$ such that $z \notin \overline{N(z) \cap D_I}$. Take such *z* with the smallest dimension of $\overline{N(z)}$. Note that $\overline{N(z)}$ is an integral box of the form $\{x \in \mathbb{R}^n \mid l \leq x \leq u\}$ for some $l, u \in \mathbb{Z}^n$ with $\|u - l\|_{\infty} \leq 1$ and the dimension of $\overline{N(z)}$ is equal to the number of indices *i* satisfying $u_i - l_i = 1$.

The set $\overline{N(z)} \cap D$ is a bounded polyhedron, and $(\overline{N(z)} \cap D) \setminus \overline{N(z) \cap D_1} \neq \emptyset$ since $z \in$ $(N(z) \cap D) \setminus N(z) \cap D_I$. Hence there is a vertex *v* of $N(z) \cap D$ not contained in $N(z) \cap D_I$ (see Figure [6\)](#page-15-1). The vertex *v* is a relative interior point of $\overline{N(z)}$, because, otherwise, we would have dim $\overline{N(v)} < \dim \overline{N(z)}$ while $v \notin \overline{N(v) \cap D_i}$ from $v \notin \overline{N(z) \cap D_i} \supseteq \overline{N(v) \cap D_i}$, a contradiction to our choice of *z*. Since *v* is a relative interior point of $\overline{N(z)}$, we have $N(v) = N(z)$. Let $y^0 := v$, which satisfies the three conditions [\(4.18\)](#page-14-3)–[\(4.20\)](#page-14-4).

Consider a (non-redundant) system of inequalities describing D . Since y^0 is a vertex of *N*(y ⁰) ∩ *D* lying in the relative interior of *N*(y ⁰), at least one inequality is tight (i.e., satisfied in equality). Enumerate all such inequalities as

$$
a_1^\top x \le \beta_1, \quad a_2^\top x \le \beta_2, \quad \dots, \quad a_k^\top x \le \beta_k,\tag{4.21}
$$

 $\frac{1}{2}$ (is the triangular region) Figure 6: Notations in the proof of Claim [4.2](#page-14-2) (The square represents $\overline{N(z)}$ and the triangle is $\overline{N(z)}$ ∩ *D*)

cone, all the inequalities in [\(4.21\)](#page-14-5) are also tight at x^0 , that is, $a_i^T x^0 = \beta_i$ for $i = 1, 2, ..., k$. \mathcal{L} where $k \geq 1$. By definition we have a_i^{\dagger} $i_j^T y^0 = \beta_i$ for $i = 1, 2, ..., k$. Since $D = x^0 + C$ and *C* is a $i^{\top} x^0 = \beta_i$ for $i = 1, 2, ..., k$.

 $and \beta = \sum_{i=1}^{k} \mu_i \beta_i$ satisfy **Claim 4.3.** *There exist some positive coefficients* $\mu_1, \mu_2, \ldots, \mu_k > 0$ *such that* $a = \sum_{i=1}^k \mu_i a_i$

$$
a^{\top}x^0 = \beta, \quad a^{\top}y^0 = \beta,\tag{4.22}
$$

$$
a^{\top} x \le \beta \qquad (\forall x \in D), \tag{4.23}
$$

$$
a^{\top} x \le 0 \qquad (\forall x \in C), \tag{4.24}
$$

$$
a^{\top}x \neq \beta \qquad (\forall x \in N(y^0) \setminus D_I). \tag{4.25}
$$

Proof. [\(4.22\)](#page-15-2)–[\(4.24\)](#page-15-3) hold for any $\mu_1, \mu_2, \ldots, \mu_k > 0$. (4.22) is immediate from the tightness $a_i^{\top} x^0 = a_i^{\top} y^0 = \beta_i$ for $i = 1, 2, ...$ shown as follows. Since *y*⁰ is a vertex, the intersection of $\overline{N(y^0)}$ and the hyperplanes $a_i^{\top} x = \beta_i$ $(i = 1, 2, \ldots, k)$ consists of a single vector y^0 , that is, for $x \in \overline{N(y^0)}$, we have $a_i^{\top} x = \beta_i$ for all $a_i^T x \neq \beta_i$. It then follows that (4.25) holds for randomly channels $a_i^{\top} x^0 = a_i^{\top}$ $i_j^{\text{T}} y^0 = \beta_i$ for $i = 1, 2, ..., k$. [\(4.23\)](#page-15-4) holds since the inequalities in [\(4.21\)](#page-14-5) are valid for *D*. [\(4.24\)](#page-15-3) follows from [\(4.22\)](#page-15-2) and [\(4.23\)](#page-15-4) because $D = x^0 + C$ and *C* is a cone. [\(4.25\)](#page-15-5) can be i^{\top} *x* = β_i i^{\top} *x* = β_i for all $i = 1, 2, \ldots, k$ if and only if $x = y^0$. Therefore, for each $x \in N(y^0) \setminus D_I$, there is some *i* with $\hat{u}_i^{\top} x \neq \beta_i$. It then follows that [\(4.25\)](#page-15-5) holds for randomly chosen $\mu_1, \mu_2, \dots, \mu_k > 0$.

Claim 4.4. Claim 4.4.

$$
y^0 \notin \overline{N(y^0) \cap \{x \in \mathbb{Z}^n \mid a^\top x \le \beta\}}.\tag{4.26}
$$

Proof. Recall from [\(4.23\)](#page-15-4) that $\{x \in \mathbb{Z}^n \mid a^\top x \leq \beta\} \supseteq D_I$. Using notation $E := \{x \in \mathbb{Z}^n \mid a^\top x \leq \beta\}$ β \setminus *D_I*, we can rewrite [\(4.26\)](#page-15-6) as

$$
y^{0} \notin \overline{N(y^{0}) \cap (E \cup D_{I})} = \overline{(N(y^{0}) \cap D_{I}) \cup (N(y^{0}) \cap E)}.
$$
 (4.27)

 $N(y^0) \cap E$ by [\(4.25\)](#page-15-5). Then [\(4.27\)](#page-15-7) follows. We have $y^0 \notin \overline{N(y^0) \cap D_1}$ in [\(4.18\)](#page-14-3) and $a^{\dagger}y^0 = \beta$ in [\(4.22\)](#page-15-2), whereas $a^{\dagger}x < \beta$ for all $x \in$

Let β^* denote the maximum value of $a^{\top}x$ over *Q*, that is,

$$
\beta^* := \max\{a^{\top} x \mid x \in \mathcal{Q}\}.
$$
 (4.28)

Since Q is a bounded integer polyhedron, we may assume that this maximum is attained by an integer vector $x^* \in Q \cap \mathbb{Z}^n$. Define $y^* \in \mathbb{R}^n$ by

$$
y^* := y^0 + (x^* - x^0). \tag{4.29}
$$

 $\frac{1}{\sqrt{2}}$ We have *y*^{*} ∈ \overline{S} , since *y*^{*} = *x*^{*} + (*y*⁰ − *x*⁰) ∈ *Q* + (*D* − *x*⁰) = *Q* + *C* = \overline{S} . Claim 4.5.

$$
y^* \notin \overline{N(y^*) \cap \{x \in \mathbb{Z}^n \mid a^\top x \le \beta^*\}}.\tag{4.30}
$$

Proof. Recall from [\(4.26\)](#page-15-6) that

 $y^0 \notin \overline{N(y^0) \cap \{x \in \mathbb{Z}^n \mid a^{\top}x \leq \beta\}}.$

By adding $x^* - x^0$ to the left-hand side, we obtain $y^* = y^0 + (x^* - x^0)$. On the right-hand side, we have $N(y^0) + (x^* - x^0) = N(y^0 + x^* - x^0) = N(y^*)$, where the first equality is true by $x^* - x^0 \in \mathbb{Z}^n$. Since $a^{\top} x^* = \beta^*$ by the definition of x^* and $a^{\top} x^0 = \beta$ by [\(4.22\)](#page-15-2), we also have

$$
\{x \mid a^{\top}x \le \beta\} + (x^* - x^0) = \{x + (x^* - x^0) \mid a^{\top}x \le \beta\}
$$

=
$$
\{z \mid a^{\top}(z - x^* + x^0) \le \beta\} = \{z \mid a^{\top}z \le \beta^*\}.
$$

Thus we obtain (4.30) .

Claim 4.6.

$$
S \subseteq \{x \in \mathbb{Z}^n \mid a^\top x \le \beta^*\}.
$$
\n
$$
(4.31)
$$

Proof. We have $Q \subseteq \{x \mid a^\top x \leq \beta^*\}$ by the definition [\(4.28\)](#page-15-8) of β^* , whereas $C \subseteq \{x \mid a^\top x \leq 0\}$ by [\(4.24\)](#page-15-3). Therefore, every $x \in \overline{S} = Q + C$ satisfies $a^{\top} x \leq \beta^*$. В последните последните под на применения и последните и последните под на применения и последните и последн
В последните последните последните под на применения и последните последните последните последните последните

It follows from [\(4.30\)](#page-16-1) and [\(4.31\)](#page-16-2) that $y^* \notin \overline{N(y^*) \cap S}$, whereas $y^* \in \overline{S}$. This is a contradiction to the integral convexity of *S* , completing the proof of Proposition [4.4.](#page-14-0)

4.3 Proof of Theorem [3.6](#page-8-0)

In this section we prove Theorem [3.6,](#page-8-0) stating that every integrally convex set $S \subseteq \mathbb{Z}^n$ can be represented as $S = T + G$ with a bounded integrally convex set T and a conic integrally convex set *G*.

By Proposition [2.1,](#page-2-2) the convex hull \overline{S} of *S* can be represented as

$$
\overline{S} = \hat{Q} + C \tag{4.32}
$$

with a polytope \hat{Q} and the characteristic cone *C* of \overline{S} . By Proposition [4.4,](#page-14-0) $C \cap \mathbb{Z}^n$ is integrally convex. With reference to the polytope \hat{Q} , define

$$
l_i := \lfloor \min\{x_i \mid x \in \hat{Q}\} \rfloor, \qquad u_i := \lceil \max\{x_i \mid x \in \hat{Q}\} \rceil
$$

for $i = 1, 2, \ldots, n$. The numbers l_i, u_i are (finite) integers with $l_i \leq u_i$, since \hat{Q} is a nonempty and bounded polyhedron.

Let $\{d^1, d^2, \ldots, d^L\}$ be a generating set of cone *C*, where we may assume $d^j \in \{-1, 0, +1\}^n$ by Proposition [4.1.](#page-10-0) With reference to the number *L* of the generators of *C*, define a bounded integral box *B* by

$$
B := \{x \in \mathbb{R}^n \mid l_i - L \le x_i \le u_i + L \ (i = 1, 2, \dots, n)\}
$$

and put $Q := \overline{S} \cap B$, which is a bounded box-integer polyhedron containing \hat{Q} . We have $\overline{S} = Q + C$, since $\overline{S} = \hat{Q} + C \subseteq Q + C \subseteq \overline{S} + C = \overline{S}$.

Define

$$
T := Q \cap \mathbb{Z}^n = \overline{S} \cap B \cap \mathbb{Z}^n = S \cap B, \qquad G := C \cap \mathbb{Z}^n, \tag{4.33}
$$

which are, respectively, a bounded integrally convex set and a conic integrally convex set. In the following we show $S = T + G$ by a sequence of claims.

Claim 4.7. $S \supseteq T + G$.

Proof. For two (discrete) sets $S_1, S_2 \subseteq \mathbb{Z}^n$, in general, we have

$$
(\overline{S_1} + \overline{S_2}) \cap \mathbb{Z}^n \supseteq S_1 + S_2.
$$

Using this for $(S_1, S_2) = (T, G)$ as well as $\overline{S} = O + C = \overline{T} + \overline{G}$, we obtain

$$
S = \overline{S} \cap \mathbb{Z}^n = (\overline{T} + \overline{G}) \cap \mathbb{Z}^n \supseteq T + G.
$$

To show the reverse inclusion $S \subseteq T + G$, take any $z \in S$. By $\overline{S} = \hat{Q} + C$ in [\(4.32\)](#page-16-3), there exist real vectors $\hat{x} \in \hat{O}$ and $\hat{d} \in C$ satisfying

$$
z = \hat{x} + \hat{d}.
$$

The vector \hat{d} can be represented as a nonnegative combination of the generators $\{d^1, d^2, \ldots, d^L\}$ of *C* as

$$
\hat{d} = \sum_{j=1}^{L} \lambda_j d^j, \qquad \lambda_j \ge 0 \ \ (j = 1, 2, ..., L).
$$

With reference to this expression, define vectors d^* and x^* by

$$
d^* := \sum_{j=1}^L \lfloor \lambda_j \rfloor \, d^j,\tag{4.34}
$$

$$
x^* := \hat{x} + \sum_{j=1}^{L} (\lambda_j - \lfloor \lambda_j \rfloor) d^j,
$$
 (4.35)

for which we have

$$
x^* + d^* = \hat{x} + \hat{d} = z.
$$
 (4.36)

Claim 4.8. d^* ∈ *G*.

Proof. [\(4.34\)](#page-17-0) shows $d^* \in C$. We also have $d^* \in \mathbb{Z}^n$, since $\lfloor \lambda_j \rfloor \in \mathbb{Z}$ and $d^j \in \{-1, 0, +1\}^n$ for $j = 1, 2, \ldots, L$ by Proposition [4.1.](#page-10-0) Therefore, $d^* \in C \cap \mathbb{Z}^n = G$.

Claim 4.9. x^* ∈ *T*.

Proof. Since $T = \overline{S} \cap B \cap \mathbb{Z}^n$ (see [\(4.33\)](#page-16-4)), it suffices to show (i) $x^* \in \mathbb{Z}^n$, (ii) $x^* \in \overline{S}$, and (iii) $x^* \in B$. We have $x^* \in \mathbb{Z}^n$, since $z \in \mathbb{Z}^n$, $d^* \in \mathbb{Z}^n$, and $x^* = z - d^*$ by [\(4.36\)](#page-17-1). We have $x^* \in \overline{S}$, since $x^* \in \hat{Q} + C$ by [\(4.35\)](#page-17-2) and $\hat{Q} + C = \overline{S}$ by [\(4.32\)](#page-16-3). Finally, we show $x^* \in B$. For the first term \hat{x} on the right-hand side of [\(4.35\)](#page-17-2), we have $l \leq \hat{x} \leq u$ since $\hat{x} \in \hat{Q}$. Each component of the second term $\sum_{j=1}^{L} (\lambda_j - \lambda_j) d^j$ lies between $-L$ and $+L$, since $0 \leq \lambda_j - \lambda_j$ < 1 and $d^{j} \in \{-1, 0, +1\}$ ⁿ for $j = 1, 2, ..., L$ by Proposition [4.1.](#page-10-0) Therefore, $x^* \in B$. □

The inclusion $S \subseteq T + G$ follows from Claims [4.8](#page-17-3) and [4.9,](#page-17-4) while $S \supseteq T + G$ is already shown in Claim [4.7.](#page-17-5) This completes the proof of Theorem [3.6.](#page-8-0)

4.4 Proof of Theorem [3.4](#page-6-0)

In this section we prove Theorem [3.4](#page-6-0) for polyhedra P ($\subseteq \mathbb{R}^n$) with particular discrete convexities such as L^{\natural} -convexity, M^{\natural} -convexity, etc. The proof for the case (1) of L^{\natural} -convex polyhedra has already been given in Section [3.1,](#page-4-3) right after Theorem [3.4.](#page-6-0) Here we present a unified proof scheme for all cases including $L^{\frac{1}{2}}$ -convex polyhedra. We use a generic name "A-convex" to mean any of L^{\natural} -convex, L_2^{\natural} -convex, M^{${\natural}$}-convex, M-convex, M₂⁻convex, M₂convex, and multimodular.

The unified proof scheme is as follows. Let *P* be an A-convex polyhedron. By Proposi-tion [2.1,](#page-2-2) we can decompose *P* as $P = \hat{Q} + C$, where \hat{Q} is a polytope and *C* is the characteristic cone of *P*. We assume that

The characteristic cone of an A-convex polyhedron is A-convex. (4.37)

There exists a bounded A-convex polyhedron *Q* satisfying $\hat{Q} \subseteq Q \subseteq P$. (4.38)

By \hat{Q} ⊆ Q ⊆ P in [\(4.38\)](#page-18-1), we have $P = \hat{Q} + C$ ⊆ $Q + C$ ⊆ $P + C = P$. This shows $P = Q + C$, where *O* is a bounded A-convex polyhedron by (4.38) and *C* is an A-convex cone by (4.37) .

The first assumption (4.37) is met by each discrete convexity in (1) – (5) . Indeed, a polyhedron *P* with such discrete convexity can be described as $P = \{x \mid Ax \leq b\}$, where a necessary and sufficient condition on (*A*, *b*) for that discrete convexity of *P* is known. For example, an L^{\natural} -convex polyhedron is described by [\(3.2\)](#page-5-4) and an M^{\natural} -convex polyhedron by [\(3.3\)](#page-6-1); see Murota [\[9\]](#page-20-2), Moriguchi–Murota [\[6,](#page-20-9) Table 1], and Murota–Tamura [\[17,](#page-21-5) Table 1] for other cases. This enables us to prove that the characteristic cone $C = \{d \mid Ad \leq 0\}$ is also endowed with the same kind of discrete convexity.

For the second assumption [\(4.38\)](#page-18-1), we consider $Q := P \cap B$ for a bounded integral box *B* containing \hat{Q} , expecting that Q is endowed with A-convexity as a consequence of the assumed A-convexity of *P*. This construction is indeed valid for all discrete convexities in question, with the exception of L_2^{\natural} -convexity in (2) (see Remark [4.1](#page-18-3) below).

In Case (2) of an L_2^{\natural} -convex polyhedron *P*, we construct an L_2^{\natural} -convex (integer) polyhedron *Q* as follows. Let $P = P_1 + P_2$ with two L^{\sharp}-convex polyhedra P_1 and P_2 . Enumerate all vertices of the polytope \hat{Q} as $\{z^1, z^2, \ldots, z^m\}$, where $z^j \in \mathbb{R}^n$ for $j = 1, 2, \ldots, m$. By $z^j \in \hat{Q} \subseteq P_1 + P_2$, each z^j can be expressed as $z^j = x^j + y^j$ with $x^j \in P_1$ and $y^j \in P_2$. Take integral boxes B_1 and B_2 satisfying $\{x^1, x^2, \ldots, x^m\} \subseteq B_1$ and $\{y^1, y^2, \ldots, y^m\} \subseteq B_2$, respectively, and define $Q_1 := P_1 \cap B_1$, $Q_2 := P_2 \cap B_2$, and $Q := Q_1 + Q_2$. Then Q_1 and Q_2 are L^{\natural} -convex (integer) polyhedra, and hence Q is an L_2^{\natural} -convex (integer) polyhedron. Then we have

$$
\hat{Q} = \overline{\{z^1, z^2, \dots, z^m\}} = \overline{\{x^1 + y^1, x^2 + y^2, \dots, x^m + y^m\}}
$$
\n
$$
\subseteq \overline{\{x^1, x^2, \dots, x^m\}} + \overline{\{y^1, y^2, \dots, y^m\}}
$$
\n
$$
\subseteq (P_1 \cap B_1) + (P_2 \cap B_2) = Q_1 + Q_2 = Q
$$

and $Q = Q_1 + Q_2 \subseteq P_1 + P_2 = P$. Thus we obtain $\hat{Q} \subseteq Q \subseteq P$.

This completes the proof of Theorem [3.4.](#page-6-0)

Remark 4.1. The intersection of an L_2^{\dagger} -convex polyhedron with an integral box is not necessarily L_2^{\natural} -convex. For example, let P_1 be the line segment connecting (0, 0, 0) and (1, 1, 0) and P_2 be the one connecting (0, 0, 0) and (0, 1, 1). Then $P = P_1 + P_2 \subseteq \mathbb{R}^3$ is an L_2^{\natural} -convex polyhedron, which is a parallelogram lying on the plane $x_1 - x_2 + x_3 = 0$ in \mathbb{R}^3 . For the unit box $B = \{x \mid 0 \le x_i \le 1 \ (i = 1, 2, 3)\}\$, the intersection $P \cap B$ is a triangle with vertices at $(0, 0, 0), (0, 1, 1),$ and $(1, 1, 0)$. This triangle is not L_2^{\dagger} -convex.

Remark 4.2. Here is an alternative proof of Theorem [3.4\(](#page-6-0)2) that relies on (1) for an L^{\sharp} convex polyhedron. Let $P = P_1 + P_2$ with L^{\natural} -convex polyhedra P_1 and P_2 . By (1) we have $P_i = Q_i + C_i$ with a bounded L^{\dagger} -convex polyhedron Q_i and an L^{\dagger} -convex cone C_i , where *i* = 1, 2. Then $P = (Q_1 + Q_2) + (C_1 + C_2)$, where $Q_1 + Q_2$ is a bounded L_2^{\dagger} -convex polyhedron and $C_1 + C_2$ is an L_2^{\natural} -convex cone.

4.5 Proof of Theorem [3.7](#page-8-1)

In this section we prove Theorem [3.7](#page-8-1) for discrete sets $S \subseteq \mathbb{Z}^n$ with particular discrete convexities such as L^{\natural} -convexity, M^{\natural} -convexity, etc. The proof relies on Theorem [3.6](#page-8-0) for integrally convex sets. Just as in Section [4.4,](#page-18-0) we present a unified proof scheme by using a generic name "A-convex" to mean any of L^{\natural} -convex, L_2^{\natural} -convex, M^{\natural} -convex, M-convex, $\textmd{M}_{2}^{\natural}$ $_{2}^{\pi}$ -convex, M₂-convex, and multimodular.

The unified proof scheme is as follows. Let *S* be an A-convex set. This implies that *S* is an integrally convex set. By Theorem [3.6](#page-8-0) we can decompose *S* as $S = \hat{T} + G$, where \hat{T} is a bounded integrally convex set and *G* is a conic integrally convex set. We have $G = C \cap \mathbb{Z}^n$ for the characteristic cone *C* of the convex hull \overline{S} of *S*, where \overline{S} is an A-convex polyhedron. We assume that

The characteristic cone of an A-convex polyhedron is A-convex. (4.39)

There exists a bounded A-convex set *T* satisfying $\hat{T} \subseteq T \subseteq S$. (4.40)

By $\hat{T} \subseteq T \subseteq S$ in [\(4.40\)](#page-19-1), we have $S = \hat{T} + G \subseteq T + G \subseteq S + G \subseteq S$, where the last inclusion follows from $\overline{S + G} = \overline{S} + \overline{G} = \overline{S} + C = \overline{S}$, $S + G \subseteq \overline{S + G} \cap \mathbb{Z}^n$, and $\overline{S} \cap \mathbb{Z}^n = S$. Therefore, $S = T + G$, where *T* is a bounded A-convex set by [\(4.40\)](#page-19-1) and *G* is a conic A-convex set by [\(4.39\)](#page-19-2).

The first assumption [\(4.39\)](#page-19-2), which is the same as [\(4.37\)](#page-18-2), is met by each discrete convexity in (1)–(5), as explained in the proof of Theorem [3.4](#page-6-0) in Section [4.4.](#page-18-0) Recall that the inequality representations are used here.

For the second assumption [\(4.40\)](#page-19-1), we consider $T := S \cap B$ for a bounded integral box *B* containing \hat{T} , expecting that T is endowed with A-convexity as a consequence of the assumed A-convexity of *S* . This construction is indeed valid for all discrete convexities in question (see [\[9,](#page-20-2) [12\]](#page-21-7)), with the exception of L_2^{\dagger} -convexity in (2) (see Remark [4.3](#page-19-3) below).

In Case (2) of an L_2^{\natural} -convex set *S*, we construct *T* as follows. Represent *S* as $S =$ $S_1 + S_2$ with two L^{\natural}-convex sets S_1 and S_2 . Enumerate all members of the finite set \hat{T} as $\hat{T} = \{z^1, z^2, \dots, z^m\}$. Each $z^j \in \hat{T} \subseteq S = S_1 + S_2$ can be expressed as $z^j = x^j + y^j$ with $x^j \in S_1$ and $y^j \in S_2$. Take integral boxes B_1 and B_2 satisfying $\{x^1, x^2, \ldots, x^m\} \subseteq B_1$ and $\{y^1, y^2, \ldots, y^m\} \subseteq B_2$, respectively, and define $T_1 := S_1 \cap B_1$, $T_2 := S_2 \cap B_2$, and $T := T_1 + T_2$. Then T_1 and T_2 are L^{\natural} -convex, and hence T is L_2^{\natural} -convex. We have $\hat{T} \subseteq T$, since $x^j \in S_1 \cap B_1$ and *y*^{*j*} ∈ *S*₂ ∩ *B*₂ imply that $z^j = x^j + y^j$ ∈ (*S*₁ ∩ *B*₁) + (*S*₂ ∩ *B*₂) = *T*₁ + *T*₂ = *T*. Finally we note $T = T_1 + T_2 \subseteq S_1 + S_2 = S$, to obtain $\hat{T} \subseteq T \subseteq S$.

This completes the proof of Theorem [3.7.](#page-8-1)

Remark 4.3. The intersection of an L_2^{\natural} -convex set with an integral box is not necessarily L_2^{\natural} -convex. For example, consider an L_2^{\natural} -convex set $S = S_1 + S_2$ given by two L^{\natural} convex sets $S_1 = \{(0, 0, 0), (1, 1, 0)\}\$ and $S_2 = \{(0, 0, 0), (0, 1, 1)\}\$. That is, $S = S_1 + S_2 =$ { $(0, 0, 0)$, $(0, 1, 1)$, $(1, 1, 0)$, $(1, 2, 1)$ }. For $B = \{x \mid 0 \le x_i \le 1 \ (i = 1, 2, 3)\}$ we have $S \cap B = \{(0, 0, 0), (0, 1, 1), (1, 1, 0)\}\text{, which is not } L_2^{\natural}\text{-convex.}$

Remark 4.4. Here is an alternative proof of Theorem [3.7\(](#page-8-1)2) that relies on (1) for an L^{\sharp} convex set. Let $S = S_1 + S_2$ with L^{\natural} -convex sets S_1 and S_2 . By (1) we have $S_i = T_i + G_i$ with a bounded L^{\natural} -convex set T_i and a conic L^{\natural} -convex set G_i , where $i = 1, 2$. Then $S =$ $(T_1 + T_2) + (G_1 + G_2)$, where $T_1 + T_2$ is a bounded L_2^{\dagger} -convex set and $G_1 + G_2$ is a conic L_2^\natural ^q-convex set. Note that $G_1 + G_2 = G_1 + G_2$ is a cone.

5 Conclusion

Our proofs given in Sections [4.1](#page-10-2)[–4.3](#page-16-0) are long and primitive based on the very definition of integral convexity. On the other hand, it is known (Chervet–Grappe–Robert [\[1\]](#page-20-10)) that a polyhedral cone is box-integer if and only if it is box-TDI. It is left for future investigation to find shorter or more transparent proofs, possibly making use of this equivalence.

Acknowledgement. This work was supported by JSPS/MEXT KAKENHI JP23K11001 and JP21H04979.

References

- [1] Chervet P., Grappe, R., Robert, L.-H.: Box-total dual integrality, box-integrality, and equimodular matrices. Mathematical Programming, Ser. A 188, 319–349 (2021)
- [2] Favati, P., Tardella, F.: Convexity in nonlinear integer programming. Ricerca Operativa 53, 3–44 (1990)
- [3] Fujishige, S.: Submodular Functions and Optimization, 2nd edn. Annals of Discrete Mathematics 58, Elsevier, Amsterdam (2005)
- [4] Iimura, T., Murota, K., Tamura, A.: Discrete fixed point theorem reconsidered. Journal of Mathematical Economics 41, 1030–1036 (2005)
- [5] Moriguchi, S., Murota, K.: Projection and convolution operations for integrally convex functions. Discrete Applied Mathematics 255, 283–298 (2019)
- [6] Moriguchi, S., Murota, K.: Note on the polyhedral description of the Minkowski sum of two L-convex sets. Japan Journal of Industrial and Applied Mathematics 40, 223–263 (2023)
- [7] Moriguchi, S., Murota, K., Tamura, A., Tardella, F.: Scaling, proximity, and optimization of integrally convex functions. Mathematical Programming 175, 119–154 (2019)
- [8] Murota, K.: Discrete convex analysis. Mathematical Programming 83, 313–371 (1998)
- [9] Murota, K.: Discrete Convex Analysis. Society for Industrial and Applied Mathematics, Philadelphia (2003)
- [10] Murota, K.: Recent developments in discrete convex analysis. In: Cook, W., Lovász, L., Vygen, J. (eds.) Research Trends in Combinatorial Optimization, Chapter 11, pp. 219– 260. Springer, Berlin (2009)
- [11] Murota, K.: Discrete convex analysis: A tool for economics and game theory. Journal of Mechanism and Institution Design 1, 151–273 (2016)
- [12] Murota, K.: A survey of fundamental operations on discrete convex functions of various kinds. Optimization Methods and Software 36, 472–518 (2021)
- [13] Murota, K.: On basic operations related to network induction of discrete convex functions. Optimization Methods and Software 36, 519–559 (2021)
- [14] Murota, K., Shioura, A.: Extension of M-convexity and L-convexity to polyhedral convex functions. Advances in Applied Mathematics 25, 352–427 (2000)
- [15] Murota, K., Tamura, A.: Integrality of subgradients and biconjugates of integrally convex functions. Optimization Letters 14, 195–208 (2020)
- [16] Murota, K., Tamura, A.: Discrete Fenchel duality for a pair of integrally convex and separable convex functions. Japan Journal of Industrial and Applied Mathematics 39, 599–630 (2022)
- [17] Murota, K., Tamura, A.: Recent progress on integrally convex functions. Japan Journal of Industrial and Applied Mathematics 40, 1445–1499 (2023)
- [18] Murota, K., Tamura, A.: Shapley–Folkman-type theorem for integrally convex sets. arXiv: http://arxiv.org/abs/[2305.15125](http://arxiv.org/abs/2305.15125) (2023)
- [19] Schrijver, A.: Theory of Linear and Integer Programming. Wiley, New York (1986)
- [20] Schrijver, A.: Combinatorial Optimization—Polyhedra and Efficiency. Springer, Heidelberg (2003)

Contents

