

A note on secant defective varieties and Clifford modules

Oliver Nash
Imperial College London

June 2023

Abstract

We generalise a construction of Landsberg, which associates certain Clifford algebra representations to Severi varieties. We thus obtain a new proof of Russo's Divisibility Property for LQEL varieties.

1 Introduction

The geometry of secant-defective varieties is surprisingly rich. In the early 20th Century, the subject captured the attention of several members of the Italian School of Algebraic Geometry and important results appear in numerous beautiful old papers, such as those of Scorza [15], Severi [16], and Terracini [17].

In the 1980s the subject enjoyed a renaissance, largely due to a series of breakthroughs made by Zak [18]. Zak's solution of Hartshorne's linear normality conjecture [3] led to his classification of maximally-secant-deficient varieties, which he named Severi varieties. He showed that there are exactly four Severi varieties, that they correspond to normed division algebras, and that their dimensions are 2, 4, 8, 16. The hardest part of the classification was establishing the dimension restriction.

In 1996, an intriguing paper of Landsberg [8] appeared in which he showed that the extrinsic geometry of a Severi variety induces certain Clifford algebra representations. Using the classification of Clifford modules, it is then trivial to see that the dimensions of the Severi varieties must take the values already established by Zak.

The main purpose of this note is to show that Landsberg's Clifford modules may be generalised. Severi varieties are examples of a class of varieties

introduced by Russo [13] in 2009, called LQEL varieties and we show that Landsberg’s construction works in this more general setting.

Actually the generalisation is only an extremely mild extension of Landsberg’s results. However we believe it is worth highlighting because, together with the classification of Clifford modules, it provides a new proof of Russo’s Divisibility Property for LQEL varieties (corollary 2.8).

2 Secant defective varieties and Clifford modules

We shall follow Landsberg [6, 8] closely and so recall his constructions and notation¹.

Our primary object of concern is a subvariety of projective space. We write:

$$X \subseteq \mathbb{P}^{n+a},$$

to indicate that the variety X is n -dimensional and that the embedding has codimension a . We work over \mathbb{C} throughout, and assume that X is smooth, irreducible, and non-degenerate², with secant deficiency $\delta \geq 1$.

2.1 Second fundamental form

We recall [2] that the second fundamental form of an embedding $X \subseteq \mathbb{P}^{n+a}$ is a section of $\text{Hom}(S^2TX, N)$, where S^2TX is the symmetric square of the tangent bundle and N is the normal bundle. Thus for $x \in X$, we have a symmetric bilinear map:

$$II^x : S^2T_xX \rightarrow N_x.$$

When we have chosen a point $x \in X$ and there is no possibility of confusion, we will write T for T_xX , N for N_x and II for II^x . Taking the transpose, we also regard the second fundamental form as a linear system of quadrics:

$$II^* : N^* \rightarrow S^2T^*.$$

A key observation is that global properties of X are visible infinitesimally via the second fundamental form, and exceptional global properties tend to produce linear systems of quadrics with exceptional properties.

¹Note that [6] (which we follow) adopts slightly different conventions than [8]. For example $\text{Ann}(v)$ in [8] corresponds to $\mathbb{P}\text{Ann}(v)$ in [6].

²Not contained in a hyperplane.

For example, if X has a smooth dual variety, then at a general point II^* is a linear system of quadrics of *constant rank*, and from this follows Landman's parity theorem for dual-deficient varieties (see [6, Theorem 12.4.8 and Corollary 12.4.10]).

We shall show that if X has the exceptional property that the Gauss map of a general tangential projection has zero-dimensional fibres, then its second fundamental form can be used to construct certain Clifford modules, and from this follows Russo's Divisibility Property for secant-deficient varieties (see [13, Theorem 2.8] or [14, Theorem 4.2.8]).

2.2 Tangential projections

We now assume X is non-linear³ make two definitions to fix notation:

Definition 2.1. *Let $x \in X$ and let $\mathbb{T}_x X \subseteq \mathbb{P}^{n+a}$ be the embedded tangent space at x . Away from $\mathbb{T}_x X$ we define a rational map:*

$$\begin{aligned} \pi_x : X &\rightarrow \mathbb{P}N, \\ y &\mapsto [(y, \mathbb{T}_x X)], \end{aligned}$$

where N is the fibre of normal bundle of X at x . The map π_x is known as the tangential projection map at x .

We recommend [14, §2.3.2, §3.3] for a valuable discussion of tangential projections.

Definition 2.2. *Let $x \in X$ and II be the second fundamental form at x . Away from $\text{Baseloc } II^*$ we define the rational map:*

$$\begin{aligned} ii : \mathbb{P}T &\rightarrow \mathbb{P}N, \\ [v] &\mapsto [II(v, v)]. \end{aligned}$$

We recall that the closures of the images of π_x and ii coincide and have dimension $n - \delta$ (see e.g., [14, Proposition 2.3.5] and its proof). Let this $(n - \delta)$ -dimensional irreducible subvariety be:

$$Z \subseteq \mathbb{P}N,$$

and let:

$$\text{Sec}(X) \subseteq \mathbb{P}^{n+a},$$

be the $(2n + 1 - \delta)$ -dimensional secant variety of X , then we note for future reference that:

$$\text{codim } Z = \text{codim } \text{Sec}(X). \tag{1}$$

³Note that this is automatic if $\text{Sec}(X) \neq \mathbb{P}^{n+a}$ since X is non-degenerate.

2.3 Second fundamental form of a tangential projection

The following is essentially a restatement of [8, Lemma 6.6].

Given a general point $x \in X$ and a general⁴ vector $v \in T$, let $z = ii([v])$. It follows from the definition of ii that there is a natural commutative diagram:

$$\begin{array}{ccc} T & \xrightarrow{II_v} & N \\ \downarrow & & \downarrow \\ T_{[v]}\mathbb{P}T & \xrightarrow{ii_*[v]} & T_z\mathbb{P}N \end{array}$$

where $ii_*[v]$ is the derivative of ii at $[v]$ and II_v is the map:

$$II_v : w \mapsto II(v, w).$$

We thus have natural exact sequences:

$$0 \rightarrow \langle v, \ker II_v \rangle \rightarrow T \xrightarrow{\rho_T} T_z Z \rightarrow 0, \quad (2)$$

and:

$$0 \rightarrow II_v(T) \rightarrow N \xrightarrow{\rho_N} N_z Z \rightarrow 0,$$

where $N_z Z$ is the normal bundle of $Z \subseteq \mathbb{P}N$ at z . The maps ρ_T and ρ_N fit into the following commutative diagram:

$$\begin{array}{ccc} S^2 T & \xrightarrow{II} & N \\ \rho_T \otimes \rho_T \downarrow & & \downarrow \rho_N \\ S^2 T_z Z & \xrightarrow{\tilde{II}} & N_z Z \end{array} \quad (3)$$

where \tilde{II} is the second fundamental form of Z at z .

2.4 Singular locus of the second fundamental form

Griffiths and Harris noticed that the quadrics of the second fundamental form are all singular along the fibres of the Gauss map. In fact they proved [2, (2.6)]:

$$\text{Singloc}(N^*) = T_x F,$$

⁴Thus v is non-zero and $[v] \notin \text{Baseloc } II^*$.

where F is the fibre of the Gauss map through x and for any $A \subseteq N^*$, $\text{Singloc}(A)$ is the intersection of all the singular loci:

$$\text{Singloc}(A) = \bigcap_{f \in A} \{v \in T \mid v \lrcorner II^*(f) = 0\}. \quad (4)$$

A key insight of Landsberg [8] was that there are natural subsystems $A \subseteq N^*$ for which $\text{Singloc}(A)$ captures more delicate geometric features of X . Indeed it follows from (3) that there is a natural exact sequence:

$$0 \rightarrow \ker \rho_T \rightarrow \text{Singloc}(\rho_N^*(N_z^*Z)) \rightarrow \text{Singloc}(N_z^*Z) \rightarrow 0, \quad (5)$$

and so for $v \in T$, Landsberg defined:

$$\begin{aligned} \text{Ann}(v) &= \rho_N^*(N_z^*Z) \\ &= II_v(T)^\perp \\ &= \{f \in N^* \mid v \lrcorner II^*(f) = 0\}, \end{aligned}$$

and studied the middle term $\text{Singloc}(\text{Ann}(v))$ appearing in (5).

We note for future reference that:

$$\begin{aligned} \dim \text{Ann}(v) &= \text{codim } Z \\ &= \text{codim Sec}(X) \quad \text{by (1)}. \end{aligned} \quad (6)$$

2.5 Clifford modules

The simplest class of secant-deficient varieties are those for which the Gauss map of a general tangential projection has zero-dimensional fibres. For emphasis we state a key consequence of this property:

Lemma 2.3. *Let $X \subseteq \mathbb{P}^{n+a}$ be a smooth, irreducible, non-degenerate variety of secant deficiency $\delta \geq 1$. Let $x \in X$, $v \in T_x X$ be general and let $Z \subseteq \mathbb{P}N$ be the closure of the tangential projection at x . Then the Gauss map of Z has zero-dimensional fibres if and only if:*

$$\langle v, \ker II_v \rangle = \text{Singloc}(\text{Ann}(v)). \quad (7)$$

Proof. Applying Griffiths and Harris's result [2, (2.6)] to Z , we know that the Gauss map of Z has zero-dimensional fibres if and only if the third term in (5) vanishes. Bearing in mind (2), the conclusion is clear. \square

Remark 2.4. *The Scorza Lemma [14, Theorem 3.3.3] tells us that the varieties satisfying the conditions of lemma 2.3 are LQEL varieties. Moreover we have examples:*

- the quadratic Veronese embeddings $\nu_2(\mathbb{P}^n) \subseteq \mathbb{P}^{n(n+3)/2}$ for $n \geq 2$ ($\delta = 1$),
- the binary Segre embeddings $\mathbb{P}^n \times \mathbb{P}^m \subseteq \mathbb{P}^{mn+m+n}$ for $m+n \geq 3$ ($\delta = 2$),
- the rank-2 Plücker embeddings $G(2, n) \subseteq \mathbb{P}^{(n-2)(n+1)/2}$ for $n \geq 5$ ($\delta = 4$),
- the 16-dimensional Severi variety in \mathbb{P}^{26} ($\delta = 8$),

as well as their linear projections. We recommend [14] for further discussion.

Remark 2.5. In [14, Definition 3.3.1], given general points $x, y \in X$ and general $p \in \text{Sec}(X)$ on the line \overline{xy} , Russo defines the contact locus $\Gamma_p \subseteq X$ as:

$$\Gamma_p = \overline{\{x \in X \mid T_x X \subseteq T_p \text{Sec}(X)\}},$$

and notes that by Terracini's lemma:

$$\Sigma_p \subseteq \Gamma_p,$$

where Σ_p is the entry locus of X with respect to $p \in \text{Sec}(X)$.

Let $Z = \overline{\pi_x(X)}$ be the closure of the tangential projection at x and $F \subseteq Z$ be the fibre of the Gauss map of Z through $z = \pi_x(y)$. As argued by Russo in the proof of [14, Lemma 3.3.2] the irreducible components of $\pi_x^{-1}(F)$ and Γ_p through y coincide. We should thus have a natural exact sequence of tangent spaces:

$$0 \rightarrow T_y \Sigma_p \rightarrow T_y \Gamma_p \rightarrow T_z F \rightarrow 0. \quad (8)$$

The line \overline{yp} naturally determines a vector $v \in T_y X$, and we expect:

$$\text{Singloc}(\text{Ann}(v)) = T_y \Gamma_p,$$

so that (8) can be interpreted as (5). Given this, [14, Lemma 3.3.2 (2)] would provide an alternate proof of lemma 2.3 above.

We come at last to our main point:

Theorem 2.6. Let $X \subseteq \mathbb{P}^{n+a}$ be an smooth, irreducible, non-degenerate variety of secant deficiency $\delta \geq 1$ such that $\text{Sec}(X) \neq \mathbb{P}^{n+a}$. Suppose that the Gauss map of the tangential projection at a general point x has zero-dimensional fibres, and let $v \in T_x X$ be general. Then $T/\text{Singloc}(\text{Ann}(v))$ carries a natural Clifford module structure over $\ker II_v$.

Proof. Let $Z = \overline{\pi_x(X)} \subseteq \mathbb{P}N$ be the closure of the image of the tangential projection at x .

The result we need is exactly [8, Lemma 6.26] except that we have made no assumption about Z being a cone (instead assuming that its Gauss map has zero-dimensional fibres) and we do not assume that Z is a hypersurface. In view of (1), Z is a hypersurface if and only if $\text{Sec}(X)$ is. However since the linear projection from a linear subspace which does not meet $\text{Sec}(X)$ is an isomorphism, we may select a maximal such subspace and project down to the case that $\text{Sec}(X)$ is a hypersurface; the lemma then applies, and our proof is complete.

For the benefit of readers who wish to compare with [6], we provide a reference for the argument as presented there. The key equation is [6, (12.22) page 374]:

$$q_{\epsilon\kappa}^{n+j} q_{\delta i}^{n+k} + q_{\delta k}^{n+j} q_{\epsilon i}^{n+k} = -2q_{\epsilon\delta}^{n+1} \delta_j^i \quad \forall \epsilon, \delta, j, k, i.$$

The key assumption required for the derivation is $S = 0$ where:

$$S = \dim \text{Singloc Ann}(v) - \dim \langle v, \ker II_v \rangle,$$

which follows from lemma 2.3. □

Remark 2.7. *We can restate theorem 2.6 without referring to the second fundamental form as follows.*

Let X be as in theorem 2.6 and let $p \in \text{Sec}(X)$ and $x \in X$ be general points. Let $Q \subseteq X$ be the irreducible component of the p -entry locus through x and let $Q' \subseteq Q$ be the corresponding⁵ irreducible component of the tangent locus through x . Then $T_x Q'$ carries a non-degenerate quadratic form and the fibre $N_{Q|X}^x$ of the normal bundle of Q in X is a Clifford module for the Clifford algebra $Cl(T_x Q')$.

We emphasise the following corollary:

Corollary 2.8. *Let X be as in theorem 2.6 then:*

$$2^{\lfloor \frac{\delta-1}{2} \rfloor} \mid n - \delta.$$

Proof. The result follows immediately from theorem 2.6 together with the classification of Clifford modules. Indeed if there exists a k -dimensional Clifford module of a non-degenerate l -dimensional complex quadratic form, then:

$$p \mid k,$$

⁵See [11, Lemma 2.4].

where $p = 2^{\lfloor \frac{l}{2} \rfloor}$. The reason is that the Clifford algebra of the quadratic form is the matrix algebra $\mathbb{C}^{p \times p}$ if l is even or $\mathbb{C}^{p \times p} \oplus \mathbb{C}^{p \times p}$ if l is odd (see e.g., [1, Table 1]) and the natural action of $\mathbb{C}^{p \times p}$ on \mathbb{C}^p is its unique irreducible representation. \square

Remark 2.9. *The divisibility condition established in corollary 2.8 was first proved by Russo and appeared in [13, Theorem 2.8] (see also [14, Theorem 4.2.8]). The proof involved an inductive study of the Hilbert scheme of lines through a general point of an LQEL variety.*

A second proof (due to the author) appeared as [11, Corollary 2.6]. This proof was topological and the key was a calculation in topological K -theory.

We now have a third proof (albeit with slightly different assumptions) and this time it is Clifford module theory that is the key.

It would be interesting to explore the relationship between the second and third proofs given the deep connections between K -theory and Clifford modules identified by Atiyah, Bott, and Shapiro in [1]. The first step should be to argue that Landsberg's construction actually provides bundles of representations of Clifford algebras, as x varies over a general tangent locus.

Remark 2.10. *Note that the proof of corollary 2.8 shows that the 2 which appears in the expression $(\delta - 1)/2$ corresponds to the mod-2 periodicity of Morita equivalence classes of complex Clifford algebras. Thus it is the same 2 which appears in complex Bott periodicity.*

Remark 2.11. *A quite different connection between Clifford modules and Severi varieties arises in the context of 'Clifford structures', introduced by Moroianu and Semmelmann in [10]. The Severi varieties appear in the classification of parallel non-flat even Clifford structures in [10] (see also [12]). It might be interesting to explore whether these Clifford structures have any relationship to Landsberg's Clifford modules.*

3 A remark about the $\delta \leq 8$ problem

Let $X \subseteq \mathbb{P}^{n+a}$ be a smooth, irreducible, non-degenerate, subvariety with $\text{Sec}(X) \neq \mathbb{P}^{n+a}$. It follows from Zak's proof of Hartshorne's linear normality conjecture that the secant deficiency satisfies:

$$\delta \leq \left\lfloor \frac{n}{2} \right\rfloor. \quad (9)$$

In the course of their exposition [9] of Zak's work, Lazarsfeld and Van de Ven highlighted that all known examples of X as above satisfy $\delta \leq 8$. They thus

posed the problem to investigate whether δ could be arbitrarily large (see [9, §1f, page 19]). In view of (9), any X with $\delta > 8$ must have dimension $n \geq 18$.

Very little progress has been made on this problem in the 40 years since it was first posed. Kaji [7] has shown that any variety with $\delta > 8$ must be non-homogeneous but otherwise the problem remains completely open: all known examples still satisfy $\delta \leq 8$ and the 16-dimensional Severi variety remains the only variety known to achieve $\delta = 8$. The problem remains open even for the very special class of LQEL varieties (see [14, chapter 4.4, page 113] as well as [4, Remark 3.8] and [5, Conjecture 4.5]).

We mention this problem here, to highlight that by combining known results, one may sharpen (9) slightly as follows:

Lemma 3.1. *Let $X \subseteq \mathbb{P}^{n+a}$ be a smooth, irreducible, non-degenerate subvariety with $\text{Sec}(X) \neq \mathbb{P}^{n+a}$. Suppose $n \geq 17$, then:*

$$\delta \leq \left\lfloor \frac{n-1}{2} \right\rfloor.$$

Proof. For a general tangential projection of X , let $\tilde{\gamma}$ be the dimension of the fibre of its Gauss map and $\tilde{\xi}$ its dual deficiency.

Suppose first that $\tilde{\gamma} = 0$. We may assume $\delta \geq 1$ (otherwise there is nothing to prove) so by Scorza's Lemma [14, Theorem 3.3.3] X is an LQEL variety. By [14, Corollary 4.4.11]:

$$\delta \leq \left\lfloor \frac{n+8}{3} \right\rfloor \leq \left\lfloor \frac{n-1}{2} \right\rfloor$$

since $n \geq 17$.

It remains to consider the case $\tilde{\gamma} \geq 1$. By [14, Theorem 5.4.1, Lemma 3.3.2]:

$$\delta \leq \frac{n-\tilde{\xi}}{2} \leq \frac{n-\tilde{\gamma}}{2} \leq \frac{n-1}{2},$$

as required⁶. □

Note that lemma 3.1 increases the dimension restriction on a variety with $\delta > 8$ slightly to $n \geq 19$. One might hope to make further progress by studying varieties for which $\tilde{\gamma} = 1$ and then arguing as in lemma 3.1 but with three cases corresponding to whether $\tilde{\gamma}$ is 0, 1, or at least 2.

⁶We note in passing that one could instead deal with the case $\tilde{\gamma} \geq 1$ using results of Landsberg. Indeed after projecting to ensure $\text{Sec}(X)$ is a hypersurface, one could apply [8, Corollary 7.3] (equivalently, the inequality at the bottom of page 373 of [6]).

References

- [1] M. F. Atiyah, R. Bott, and A. Shapiro. Clifford modules. *Topology*, 3(suppl. 1):3–38, 1964.
- [2] Phillip Griffiths and Joseph Harris. Algebraic geometry and local differential geometry. *Ann. Sci. École Norm. Sup. (4)*, 12(3):355–452, 1979.
- [3] Robin Hartshorne. Varieties of small codimension in projective space. *Bull. Amer. Math. Soc.*, 80:1017–1032, 1974.
- [4] Paltin Ionescu and Francesco Russo. Varieties with quadratic entry locus. II. *Compos. Math.*, 144(4):949–962, 2008.
- [5] Paltin Ionescu and Francesco Russo. On dual defective manifolds. *Math. Res. Lett.*, 21(5):1137–1154, 2014.
- [6] Thomas A. Ivey and Joseph M. Landsberg. *Cartan for beginners*, volume 175 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2016. Differential geometry via moving frames and exterior differential systems, Second edition [of MR2003610].
- [7] Hajime Kaji. Homogeneous projective varieties with degenerate secants. *Trans. Amer. Math. Soc.*, 351(2):533–545, 1999.
- [8] J. M. Landsberg. On degenerate secant and tangential varieties and local differential geometry. *Duke Math. J.*, 85(3):605–634, 1996.
- [9] R. Lazarsfeld and A. Van de Ven. *Topics in the geometry of projective space*, volume 4 of *DMV Seminar*. Birkhäuser Verlag, Basel, 1984. Recent work of F. L. Zak, With an addendum by Zak.
- [10] Andrei Moroianu and Uwe Semmelmann. Clifford structure on Riemannian manifolds. *Adv. Math.*, 228(2):940–967, 2011.
- [11] Oliver Nash. K -theory, LQEL manifolds and Severi varieties. *Geom. Topol.*, 18(3):1245–1260, 2014.
- [12] Maurizio Parton and Paolo Piccinni. The even Clifford structure of the fourth Severi variety. *Complex Manifolds*, 2(1):89–104, 2015.
- [13] Francesco Russo. Varieties with quadratic entry locus. I. *Math. Ann.*, 344(3):597–617, 2009.

- [14] Francesco Russo. *On the geometry of some special projective varieties*, volume 18 of *Lecture Notes of the Unione Matematica Italiana*. Springer, Cham; Unione Matematica Italiana, Bologna, 2016.
- [15] Gaetano Scorza. Sulle varietà a quattro dimensioni di sr ($r \geq 9$) i cui s_4 tangenti si tagliano a due a due. *Rend. Circ. Mat. Palermo*, 27:148–178, 1909.
- [16] Francesco Severi. Intorno ai punti doppi impropri di una superficie generale dello spazio a quattro dimensioni, e a' suoi punti tripli apparenti. *Rendiconti del Circolo Matematico di Palermo*, 15(1):33–51, 1901.
- [17] Alessandro Terracini. Sulle v_k per cui la varietà degli $s_h(h+1)$ -seganti ha dimensione minore dell'ordinario. *Rend. Circ. Mat. Palermo*, 31:392–396, 1911.
- [18] F. L. Zak. *Tangents and secants of algebraic varieties*, volume 127 of *Translations of Mathematical Monographs*. American Mathematical Society, Providence, RI, 1993. Translated from the Russian manuscript by the author.