THE MINIMUM VALUE OF THE FIRST DYNAMICAL DEGREES OF AUTOMORPHISMS OF COMPLEX SIMPLE ABELIAN VARIETIES WITH FIXING DIMENSIONS

YUTARO SUGIMOTO

ABSTRACT. We consider the minimum value of the first dynamical degrees, which are larger than 1, of automorphisms for prime dimensional complex simple abelian varieties. Also, we calculate the minimum value of the first dynamical degrees, which are larger than 1, of automorphisms of complex simple abelian varieties with fixing the dimensions from 2 to 10.

1. INTRODUCTION

Let X be a compact Kähler manifold and $f: X \dashrightarrow X$ be its dominant meromorphic map. The first dynamical degree of f is defined by

$$\lambda_1(f) = \lim_{n \to +\infty} ||(f^n)^* : \mathrm{H}^{1,1}(X) \to \mathrm{H}^{1,1}(X)||^{\frac{1}{n}}$$

where $|| \cdot ||$ is a norm and $H^{1,1}(X)$ is the (1,1)-Dolbeault cohomology. Especially if X is a complex simple abelian variety and f is an automorphism,

$$\lambda_1(f) = \lim_{n \to +\infty} ||(f^n)^* : \mathrm{H}^{1,1}(X) \to \mathrm{H}^{1,1}(X)||^{\frac{1}{n}}$$
$$= \lim_{n \to +\infty} ||(f^*)^n : \mathrm{H}^{1,1}(X) \to \mathrm{H}^{1,1}(X)||^{\frac{1}{n}}$$
$$= \rho(f^* : \mathrm{H}^{1,1}(X) \to \mathrm{H}^{1,1}(X))$$

and we only consider this case in this paper. It is known that $\lambda_1(f) \ge 1$ (cf. [DS05] or [DS17]) and if g = 1, then $\lambda_1(f) = 1$ (cf. [DS17, Definition 4.1]).

In our past study, we prove the next theorem.

Theorem 1.1 (cf. [Sug23, Theorem 7.2]). For an integer $g \ge 2$, the set

$$S_g := \left\{ \begin{array}{c} \text{the first dynamical degrees of automorphisms} \\ \text{of simple abelian varieties over } \mathbb{C} \text{ whose dimension is } g \end{array} \right\} \setminus \{1\}$$

has the minimum value.

Denote the minimum value of S_g by m(g) and in this paper, we study m(g) for some g in detail.

In Section 5, we consider the minimum value for prime dimensional simple abelian varieties.

Theorem A. For a prime number p, we get the next result about m(p).

$$\begin{cases} m(p) = 4\cos^2\left(\frac{\pi}{2p+1}\right) & (if \ 2p+1 \ is \ also \ a \ prime) \\ 4 + 4^{-2p-3} < m(p) < 4.3264 = (2.08)^2 & (otherwise) \end{cases}$$

Remark 1.2. p is called a Sophie Germain prime if p and 2p + 1 are both prime numbers, and it is conjectured that there are infinitely many Sophie Germain primes (cf. [Sho09, Chapter 5.5.5]). Conversely, since there are infinitely many prime numbers p such that $p \equiv 1 \pmod{3}$, there are infinitely many prime numbers p which are not Sophie Germain primes.

In Section 6, we consider the minimum value for lower dimensions.

Theorem B. For g = 2 to 10, the value m(g) is the following.

dimension g	minimum value $m(g)$
1	$(all\ 1)$
2	$4\cos^2(\pi/5) = 2.6180\cdots$
3	$4\cos^2(\pi/7) = 3.2469\cdots$
4	$2\cos(\pi/5) = 1.6180\cdots$
5	$4\cos^2(\pi/11) = 3.6825\cdots$
6	$2\cos(\pi/7) = 1.8019\cdots$
7	$4.0333\cdots$
8	$2\cos(\pi/5) = 1.6180\cdots$
9	$1.1509 \cdots^2 = 1.3247 \cdots$
10	$1.1762 \cdots^2 = 1.3836 \cdots$

Acknowledgements. The author thanks Professor Keiji Oguiso for suggesting the direction of this paper.

2. Algebraic preliminaries

This section is devoted to the algebraic preparations for the following sections. Some of the results are also used in our past study (cf. [Sug23]).

Definition 2.1 (cf. [Lan83, Chapter 1.2]). A CM-field is a number field K which satisfies the following equivalent conditions.

- (1) K is a totally complex field, and is a quadratic extension of some totally real number field.
- (2) K is closed under the complex conjugation and all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$ commute with the complex conjugation, and K is not real.

For a number field K, denote its ring of integer by \mathcal{O}_K and define

$$U_K := \{ x \in \mathcal{O}_K \setminus \{0\} \mid x^{-1} \in \mathcal{O}_K \}$$

as the group of algebraic units.

The next lemma follows from the definition of CM-fields.

Lemma 2.2 (cf. [Lan83, Chapter 1.2]). Let L be a totally real number field.

- (1) If L' is a totally complex quadratic extension of L, then for $x \in U_{L'}$, $|x|^2$ is an element of U_L .
- (2) Fix $a \in U_L \setminus \{0\}$. Then the next two conditions are equivalent.
 - (i) There exist a totally complex quadratic extension L'/L and $x \in U_{L'}$ which satisfies $|x|^2 = a$.

(ii) The conjugate elements of $a \in L$ are all positive (i.e., a is totally positive). Moreover, these conditions are equivalent to the totally realness of $L(\sqrt{a})$.

Proof. We start with the proof of (1). By the assumption, $x, \frac{1}{x} \in \mathcal{O}_{L'}$. Since [L' : L] = 2 and L is totally real, $x\overline{x}, \frac{1}{x\overline{x}} \in \mathcal{O}_L$. Thus, $|x|^2 = x\overline{x} \in U_L$.

Next, consider the proof of (2). If all conjugate elements of $a \in L$ are positive, then by defining $L' := L(\sqrt{-a})$, L' is a totally complex number field. This implies (ii) \Rightarrow (i).

Assume the condition (i). Fix a \mathbb{Q} -embedding $\sigma' : L \to \mathbb{C}$ and take a \mathbb{Q} -embedding $\sigma : L' \to \mathbb{C}$ with $\sigma' = \sigma \circ i$ where $i : L \to L'$ is the natural inclusion. By Definition 2.1, the complex conjugation commutes with σ' and so

$$\sigma'(a) = \sigma'(|x|^2) = \sigma'(x\overline{x}) = \sigma'(x)\sigma'(\overline{x}) = \sigma'(x)\overline{\sigma'(x)} = |\sigma'(x)|^2.$$

Thus, all the conjugates of a is positive and this implies (i) \Rightarrow (ii) and the proof is concluded.

In this paper, we denote $\zeta_n = \cos\left(\frac{2\pi}{n}\right) + i \cdot \sin\left(\frac{2\pi}{n}\right)$ for a positive integer n. Denote the minimal polynomial of ζ_n over \mathbb{Q} by $\Phi_n(x) \in \mathbb{Z}[x]$ and called a cyclotomic polynomial. Now the degree of $\Phi_n(x)$ is Euler's totient function $\varphi(n)$. Let $\Psi_n(x)$ be the minimal polynomial of $2\cos\left(\frac{2\pi}{n}\right)$ over \mathbb{Q} (define $\Psi_4(x) = x$ for n = 4). The constant term of $\Psi_n(x)$ is calculated in [ACR16] as below.

Proposition 2.3 (cf. [ACR16]). The absolute value of the constant term of $\Psi_n(x)$ is equal to 1 except the following cases.

- (i) $|\Psi_n(0)| = 0$ for n = 4
- (ii) $|\Psi_n(0)| = 2$ for $n = 2^m$, with $m \in \mathbb{Z}_{>0} \setminus \{2\}$
- (iii) $|\Psi_n(0)| = p$ for $n = 4p^k$, with $k \in \mathbb{Z}_{>0}$, p is an odd prime number

Also for $n \geq 3$, the polynomial $\Psi_n(x)$ satisfies

$$\Phi_n(x) = x^{\frac{\varphi(n)}{2}} \Psi_n\left(x + \frac{1}{x}\right)$$

and $\Psi_n(x)$ is of degree $\frac{\varphi(n)}{2}$.

The next lemma, which follows from Kronecker's theorem, is useful to determine the minimum value of the first dynamical degrees.

Lemma 2.4. Let $P(x) = x^n + a_1 x^{n-1} + \cdots + a_n \in \mathbb{Z}[x]$ be an irreducible monic polynomial whose roots are all real. If the maximal absolute value of the roots of P(x) is less than 2, then P(x) is a polynomial which satisfies $x^n P\left(x + \frac{1}{x}\right) = \Phi_N(x)$ for some $N \in \mathbb{Z}_{\geq 3}$. This implies $P(x) = \Psi_N(x)$. Also, the roots of P(x) can be written as $\zeta_N^m + \frac{1}{\zeta_N^m} = 2\cos(\frac{2m\pi}{N})$ for some $m \in \mathbb{Z}_{>0}$ with m < N, gcd(N, m) = 1. Moreover, the maximal absolute value of the roots of P(x) can be written as

$$\begin{array}{c} 2\cos\left(\frac{2\pi}{N}\right) & (if \ N \ is \ even) \\ 2\cos\left(\frac{\pi}{N}\right) & (if \ N \ is \ odd) \end{array}$$

Let L/K be a field extension of number fields with [L : K] = n and \mathcal{O}_L and \mathcal{O}_K be their ring of integers. Then \mathcal{O}_L and \mathcal{O}_K are Dedekind domains. For a non-zero prime ideal $\mathfrak{p} \subset \mathcal{O}_K$, there is the prime ideal factorization $\mathfrak{p}\mathcal{O}_L = \mathfrak{P}_1^{e_1} \cdots \mathfrak{P}_g^{e_g}$ since \mathcal{O}_L is a Dedekind domain. We say that \mathfrak{P}_i is over \mathfrak{p} . Now L/K is a separable extension of degree n and so

$$n = \sum_{i=1}^{g} e_i f_i.$$

Moreover, if L/K is a Galois extension, by Hilbert's ramification theory (cf. [Neu99, Chapter I, §9]),

$$e_1 = \dots = e_g = e, f_1 = \dots = f_g = f$$

hold and this implies n = efg.

Remark 2.5 (cf. [Neu99, Chapter I, Proposition 8.3]). Let L/K be a field extension of number fields and let $\theta \in \mathcal{O}_L$ be a primitive element for L/K (i.e., $L = K(\theta)$).

Denote the conductor of $\mathcal{O}_K[\theta]$ by $\mathfrak{J} \subset \mathcal{O}_L$, in other words,

$$\mathfrak{J} = \{ \alpha \in \mathcal{O}_L \mid \alpha \mathcal{O}_L \subset \mathcal{O}_K[\theta] \}.$$

Under this condition, let $\mathfrak{p} \subset \mathcal{O}_K$ be a prime ideal which is relatively prime to \mathfrak{J} (i.e., $\mathfrak{p}\mathcal{O}_L + \mathfrak{J} = \mathcal{O}_L$). Denote the minimal polynomial of θ over \mathcal{O}_K by $p(X) \in \mathcal{O}_K[X]$ and denote the image of p(X) via the map $\mathcal{O}_K[X] \to (\mathcal{O}_K/\mathfrak{p})[X]$ by $\overline{p}(X)$. $\overline{p}(X)$ is decomposed as

$$\overline{p}(X) = \overline{p_1}(X)^{e_1} \cdots \overline{p_g}(X)^{e_g}$$

in $(\mathcal{O}_K/\mathfrak{p})[X]$ for some $p_i(X) \in \mathcal{O}_K[X]$. Then, by defining

$$\mathfrak{P}_i := \mathfrak{p}\mathcal{O}_L + p_i(\theta)\mathcal{O}_L,$$

 $\mathfrak{P}_i \subset \mathcal{O}_L$ are all different prime ideals over \mathfrak{p} and

$$\mathfrak{p}\mathcal{O}_L=\mathfrak{P}_1^{e_1}\cdots\mathfrak{P}_g^{e_g}$$

holds and this is the prime ideal factorization of \mathfrak{p} in L. Also, $f_i = [\mathcal{O}_L/\mathfrak{P}_i : \mathcal{O}_K/\mathfrak{p}] = \deg(p_i(X))$ holds.

Let L/K be an extension of number fields with [L:K] = n again. Let $\{\sigma_i\}_{1 \le i \le n}$ be the set of all K-embeddings $L \hookrightarrow \mathbb{C}$.

For $a_1, \ldots, a_n \in L$, the discriminant over L/K is defined as

$$\Delta_{L/K}(a_1,\ldots,a_n) := \det \begin{pmatrix} \sigma_1(a_1) & \sigma_1(a_2) & \ldots & \sigma_1(a_n) \\ \sigma_2(a_1) & \sigma_2(a_2) & \ldots & \sigma_2(a_n) \\ \vdots & \vdots & \ddots & \vdots \\ \sigma_n(a_1) & \sigma_n(a_2) & \ldots & \sigma_n(a_n) \end{pmatrix}^2$$

Then, $\Delta_{L/K}(a_1, \ldots, a_n) \in K$ and $\Delta_{L/K}(a_1, \ldots, a_n) \neq 0$ if and only if $a_1, \ldots, a_n \in L$ is linearly independent over K. Now, the next theorem holds.

Lemma 2.6 (cf. [Neu99, Chapter I, Lemma 2.9]). Let L/K be an extension of number fields with [L:K] = n and $a_1, \ldots, a_n \in \mathcal{O}_L$ be a basis of L/K. Define $\delta := \Delta_{L/K}(a_1, \ldots, a_n)$ and then

$$\mathcal{O}_L \subset \frac{a_1}{\delta} \mathcal{O}_K \oplus \cdots \oplus \frac{a_n}{\delta} \mathcal{O}_K$$

Remark 2.7. The conductor defined in Remark 2.5 contains $\delta = \Delta_{L/K}(1, \theta, \dots, \theta^{n-1})$, where n = [L : K], by Lemma 2.6.

Definition 2.8 (cf. [Voi21, Definition 3.2.1, Definition 8.4.1]). Let *B* be a finite-dimensional algebra over \mathbb{Q} . An anti-involution is a \mathbb{Q} -linear map $\phi : B \to B$ such that

(1) $\phi(1) = 1$

(2)
$$\phi(\phi(x)) = x \quad (x \in B)$$

(3) $\phi(xy) = \phi(y)\phi(x) \quad (x, y \in B)$

An anti-involution $\phi : B \to B$ is called positive anti-involution over \mathbb{Q} if and only if $\operatorname{Tr}_{B/\mathbb{Q}}(\phi(x)x) > 0$ for all $x \in B$.

Also, in this paper, we adopt the next definition of Salem numbers.

Definition 2.9 ([BDGPS92, Chapter 5.2]). A Salem number is a real algebraic integer λ greater than 1 whose other conjugates have modulus at most equal to 1, at least one having a modulus equal to 1.

This implies that all Salem numbers have degrees at least 4.

3. Endomorphism algebras of simple abelian varieties

Let X be a g-dimensional simple abelian variety and then its endomorphism algebra $B := \operatorname{End}_{\mathbb{Q}}(X) = \operatorname{End}(X) \otimes_{\mathbb{Z}} \mathbb{Q}$ is a division ring. B is a finitely generated algebra over \mathbb{Q} with a center K, which is a field. Also, B admits the Rosati involution and it is a positive anti-involution $\phi : B \to B$ over \mathbb{Q} . Now ϕ can be restricted to K and define $K_0 := \{x \in K \mid \phi(x) = x\}$. Then K_0 is a totally real number field and $[K : K_0]$ is 1 or 2. This deduction gives the classification of simple abelian varieties by their endomorphism algebras as below (cf. [BL04, Chapter 5.5]). Denote $[B : K] = d^2$, $[K : \mathbb{Q}] = e$ and $[K_0 : \mathbb{Q}] = e_0$.

TABLE 1. Classification of X by its endomorphism algebra

X	$B = \operatorname{End}_{\mathbb{Q}}(X)$	K	d	e_0	restriction
Type 1	K	totally real	1	e	$e \mid g$
Type 2	totally indefinite quaternion algebra over K	totally real	2	e	$2e \mid g$
Type 3	totally definite quaternion algebra over K	totally real	2	e	$2e \mid g$
Type 4	division ring with center K	CM-field	d	$\frac{e}{2}$	$\frac{d^2e}{2} \mid g$

An endomorphism $f: X \to X$ can be identified as an element $\alpha \in B$ with the minimal polynomial $p(x) \in \mathcal{O}_K[x]$ over K and the minimal polynomial $P(x) \in \mathbb{Z}[x]$ over \mathbb{Q} . If f is an automorphism, the constant term of p(x) is in U_K and the constant term of P(x) is in $\{\pm 1\}$. The degree of p(x) is at most d and the degree of P(x) is at most $de \leq 2g$. Also, the first dynamical degree of the endomorphism f is the square of the maximal absolute value of the roots of $P(x) \in \mathbb{Z}[x]$ (cf. [Sug23, Section 3]).

Conversely, for a division algebra in Table 1 with some property, we have a simple abelian variety whose endomorphism algebra is the same with the division algebra by the following propositions (cf. [BL04, Chapter 9]). The same methods are also used in [Sug23].

Proposition 3.1 (Type 1). Let K be a totally real number field with $[K : \mathbb{Q}] = e$ and fix a \mathbb{Z} -order \mathcal{O} . Then, for any ineger $m \in \mathbb{Z}_{>0}$, there exists an em-dimensional simple abelian variety X with an isomorphism $K \xrightarrow{\simeq} \operatorname{End}_{\mathbb{Q}}(X)$ which induces an injective ring homomorphism $\mathcal{O} \hookrightarrow \operatorname{End}(X)$. **Proposition 3.2** (Type 2). Let B be a totally indefinite quaternion algebra over a totally real number field K with $[K : \mathbb{Q}] = e$ and fix a \mathbb{Z} -order \mathcal{O} . Assume there exists a positive anti-involution $\phi : B \to B$ over \mathbb{Q} and fix a positive integer $m \in \mathbb{Z}_{>0}$. Then there exists a 2em-dimensional simple abelian variety X with an isomorphism $B \xrightarrow{\simeq} \operatorname{End}_{\mathbb{Q}}(X)$ which induces an injective ring homomorphism $\mathcal{O} \hookrightarrow \operatorname{End}(X)$.

Proposition 3.3 (Type 4). Let B be a central simple division algebra over a CM-field K with $[B:K] = d^2$, $[K:\mathbb{Q}] = e = 2e_0$ and fix a \mathbb{Z} -order \mathcal{O} . Assume there exists a positive anti-involution $\phi: B \to B$ over \mathbb{Q} of the second kind (i.e., $\phi|_K \neq \mathrm{id}_K$) and fix a positive integer $m \in \mathbb{Z}_{>0}$. Then there exists a d^2e_0m -dimensional abelian variety X with an injective ring homomorphism $B \hookrightarrow \mathrm{End}_{\mathbb{Q}}(X)$ which induces an injective ring homomorphism $\mathcal{O} \hookrightarrow$ $\mathrm{End}(X)$.

In addition, assume one of the next conditions.

- (1) $dm \geq 3$
- (2) dm = 2 and $e_0 \ge 2$

Then the above X can be taken as a simple abelian variety and $B \simeq \operatorname{End}_{\mathbb{Q}}(X)$.

Remark 3.4. The construction of simple abelian varieties of Type 3 is also written in [BL04, Chapter 9], but it is not used in this paper, so we omit here.

The construction of a division ring is due to the next theorems.

Theorem 3.5 ([DH22, Theorem 2.5]). Let F be a totally real number field and $K = F(\sqrt{a})$ a quadratic extension for $a \in \mathcal{O}_F$. Then there exists a prime number p such that the quaternion algebra $B = \left(\frac{a,p}{F}\right)$ is divisional.

Theorem 3.6 (cf. [Pie82, Chapter 15.1, Corollary d]). Let E/F be a cyclic extension of fields with [E:F] = n. Define $G = \operatorname{Gal}(E/F)$ and denote its generator by σ . Fix $u \in F^{\times}$ and take a symbol v as $v^n = u$. Define $B = \bigoplus_{i=0}^{n-1} v^i E$ with the multiplication on B as $e \cdot v = v \cdot \sigma(e)$ for $e \in E$. Assume that the order of u in $F^{\times}/\operatorname{N}_{E/F}(E^{\times})$ is exactly n.

Then B is a division ring and also a central simple algebra over F with $[B:F] = n^2$.

The next lemma is used for searching $u \in F^{\times}$ with $u \notin N_{E/F}(E^{\times})$.

Lemma 3.7 ([Sug23, Lemma 8.9]). Let E/F be a Galois extension of number fields with [E:F] = n and let $\mathfrak{q} \subset \mathcal{O}_E$ be a prime ideal and take $\alpha \in E^{\times} \setminus U_E$.

Assume $\sigma(\mathfrak{q}) = \mathfrak{q}$ for all $\sigma \in \operatorname{Gal}(E/F)$. Then, the multiplicity of \mathfrak{q} of the prime ideal factorization of the ideal generated by $N_{E/F}(\alpha)$ in \mathcal{O}_E is a multiple of n.

Also, the construction of a positive anti-involution is reduced to the following theorems in this paper.

Theorem 3.8 (cf. [BL04, Theorem 5.5.3]). Let B be a totally indefinite quaternion algebra of finite dimension over \mathbb{Q} with center a totally real number field K. Assume that B is divisional. Then, \mathbb{Q} -linear map $\phi : B \to B$ is a positive anti-involution over \mathbb{Q} if and only if it can be written as

$$\phi(x) = c^{-1}\overline{x}c$$

where $c \in B \setminus K$ with $c^2 \in K$ totally negative and $x \mapsto \overline{x}$ is the quaternion conjugation.

Theorem 3.9 (cf. [Mum70, Chapter 21], [BL04, Theorem 5.5.6]). Let B be a division algebra of finite dimension over \mathbb{Q} with center a CM-field K. Assume that there exists an anti-involution $\phi: B \to B$ of the second kind. Then there exists a positive anti-involution $\phi': B \to B$ over \mathbb{Q} of the second kind.

4. The minimum value of the first dynamical degrees of automorphisms

First we consider the general results for considering the minimum value of the first dynamical degrees of automorphisms. The flow is repeatedly used in the following sections. Let X be a g-dimensional simple abelian variety and consider the automorphisms and its first dynamical degrees for each type in Table 1. Define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Since B = K is a totally real number field, so an automorphism $f : X \to X$ can be identified as an element $\alpha \in U_K$ and its conjugates are all real. The degree of α divides e and by the restricton in Table 1, it also divides $g = \dim(X)$. Assuming that the first dynamical degree is less than 4, then the conjugates of α are all in the interval (-2, 2). Thus, by Lemma 2.4, it is reduced to the deduction of cyclotomic polynomials.

Conversely, let K be a totally real number field with $[K : \mathbb{Q}] = e$ and take $\alpha \in U_K$ and $m \in \mathbb{Z}_{>0}$. Then by Proposition 3.1, there exists an *em*-dimensional simple abelian variety X with the endomorphism algebra $K = \operatorname{End}_{\mathbb{Q}}(X)$ with $\mathcal{O}_K \subset \operatorname{End}(X)$. Thus for the endomorphism $f : X \to X$ which corresponds to α , the condition $\alpha \in U_K$ implies $\frac{1}{\alpha} \in \mathcal{O}_K$ and so $f : X \to X$ is an automorphism.

Therefore, the minimum value of the first dynamical degrees for this type is caluculated by

 $\min \left\{ \begin{array}{l} \text{the square of the maximal absolute value of the conjugates} \\ \text{of an algebraic unit, whose conjugates are all real, of degree divides } g \end{array} \right\} \setminus \{1\}$

Type 2, Type 3

An automorphism f corresponds to an element $\alpha \in B$ and since B is a quaternion algebra over a totally real number field K, α has the minimal polynomial $p(x) \in \mathcal{O}_K[x]$ over K of degree 1 or 2.

If the degree is 1, then $\alpha \in U_K$ and so the deduction about the minimum value is reduced to the case of Type 1.

If the degree is 2, then it can be written as $p(x) = x^2 + ax + b \in \mathcal{O}_K[x]$ with $b \in U_K$. Now the minimal poynomial of α over \mathbb{Q} satisfies

$$P(x)^m = \prod_{i=1}^e (x^2 + \sigma_i(a)x + \sigma_i(b)) \in \mathbb{Z}[x]$$

for some $m \in \mathbb{Z}_{>0}$, where $\{\sigma_i\}$ is the set of all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$.

The first dynamical degree of f is the square of the maximal absolute value of the roots of P(x) and we consider the following three cases.

• there exists $1 \leq i \leq e$ such that $|\sigma_i(b)| > 1$

If $|\sigma_i(b)| > 1$, one of the roots of $x^2 + \sigma_i(a)x + \sigma_i(b)$ has absolute value not less than $\sqrt{|\sigma_i(b)|}$ and so the first dynamical degree is not less than $|\sigma_i(b)|$. $\sigma_i(b)$ are all real and so by assuming that the first dynamical degree is less than 2, $\sigma_i(b)$ are all in the interval

(-2, 2) and the remaining is same as the case of Type 1.

• $\sigma_i(b) = 1$ for all $1 \le i \le e$

The roots of $x^2 + \sigma_i(a)x + 1$ have modulus 1 if and only if $-2 \leq \sigma_i(a) \leq 2$. Thus, for small first dynamical degrees except 1, $\max\{|\sigma_i(a)|\}$ must be larger than 2 and close to 2. • $\sigma_i(b) = -1$ for all $1 \le i \le e$

The roots of $x^2 + \sigma_i(a)x - 1$ have modulus 1 if and only if $\sigma_i(a) = 0$. Thus, for the small first dynamical degrees except 1, max{ $|\sigma_i(a)|$ } must be larger than 0 and close to 0. Since $\prod_{i=1}^{e} \sigma_i(a) \in \mathbb{Z}$, max{ $|\sigma_i(a)|$ } ≥ 1 for $a \neq 0$ and the equality is always achieved by a = 1. For this case, the minimum value of the maximal absolute value of the conjugates is $\frac{1+\sqrt{5}}{2}$. Thus, $\frac{3+\sqrt{5}}{2}$ is a lower bound of the first dynamical degrees for this case.

The realizability would be considered independently in the following sections.

Type 4

For the case d = 1, B = K is a CM-field and an automorphism of X corresponds to $\alpha \in U_K$.

If $\alpha \in K_0$, the deduction is reduced to the case of Type 1.

If $\alpha \in K \setminus K_0$, by Lemma 2.2, for denoting $a = |\alpha|^2 \in U_{K_0}$, $K_0(\sqrt{a})$ is a totally real number field. The first dynamical degree of α is the square of the maximal absolute value of the conjugate elements of α , and so this is equal to the maximal absolute value of the conjugates of $a \in K_0$. In order to search the small first dynamical degree, assume the maximal absolute value of the conjugate elements of \sqrt{a} is less than 2. If a = 1, then the first dynamical degree would be 1 and so we assume $a \neq 1$. Then, as in Lemma 2.4, it can be written as $\sqrt{a} = \zeta_N^m + \frac{1}{\zeta_N^m} (m, N \in \mathbb{Z}_{>0})$ with m < N, gcd(N, m) = 1 and we can assume $N \neq 1, 2, 3, 4, 6$. Thus, $a = \left(\zeta_N^m + \frac{1}{\zeta_N^m}\right)^2 = \zeta_N^{2m} + \frac{1}{\zeta_N^{2m}} + 2$ and so

$$[\mathbb{Q}(a):\mathbb{Q}] = \begin{cases} \frac{1}{2}\varphi(\frac{N}{2}) & (N:\text{even}) \\ \frac{1}{2}\varphi(N) & (N:\text{odd}) \end{cases}$$

Now $K_0 \supset \mathbb{Q}(a) \supset \mathbb{Q}$ and so we get the candidates of N from this condition.

Conversely, let K be a CM-field with $[K:\mathbb{Q}] = e = 2e_0$ and take $\alpha \in U_K$ and fix $m \in \mathbb{Z}_{>0}$. Now K admits the complex conjugate as a positive anti-involution of the second kind. If $m \in \mathbb{Z}_{\geq 3}$ (or $e \geq 4$ and $m \in \mathbb{Z}_{\geq 2}$) holds, then by Proposition 3.3, there exist an e_0m -dimensional simple abelian variety X and an automorphism $f: X \to X$ which corresponds to α .

The case $d \ge 2$ would be considered independently in the following sections.

5. The minimum value of the first dynamical degrees for simple abelian VARIETIES WITH PRIME DIMENSIONS

Define

$$m(g) := \min \left\{ \begin{array}{c} \text{the first dynamical degrees of automorphisms} \\ \text{of simple abelian varieties over } \mathbb{C} \text{ whose dimension is } g \end{array} \right\} \setminus \{1\}.$$

We prove the next result in this section.

Theorem 5.1. For a prime number p, we get the next result about m(p).

$$\begin{cases} m(p) = 4\cos^2\left(\frac{\pi}{2p+1}\right) & (2p+1 \text{ is also a prime}) \\ 4 + 4^{-2p-3} < m(p) < (2.08)^2 & (otherwise) \end{cases}$$

Remark 5.2. Especially, $(2.08)^2$ is an upper bound of m(p) for any prime number p.

First, we consider the case p = 2. In other words, we prove the next theorem.

Proposition 5.3.

$$m(2) = 4\cos^2\left(\frac{\pi}{5}\right)$$

Proof. We consider by dividing into the type of simple abelian varieties in Table 1.

Define B, K, K_0, d, e and e_0 as in Section 3 for each type of X.

Type 1

Let X be a 2-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 2 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, the conjugates of α are all in the interval (-2, 2). If the degree of α is 1, then $\alpha = \pm 1$ and so the first dynamical degree is 1 and so we assume the degree of α is 2. Denote the minimal polynomial of α over \mathbb{Q} by P(x), whose degree is 2. By Lemma 2.4, $x^2P(x+\frac{1}{x})$ is an irreducible cyclotomic polynomial and its degree is 4. Cyclotomic polynomials $\Phi_N(x)$ have degree 4 only for N = 5, 8, 10, 12. By comparing with Proposition 2.3, since P(x) has constant term ± 1 , N = 5, 10 are only possible. Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1 for this type.

Type 2, Type 3

Let X be a 2-dimensional simple abelian variety of Type 2 or 3 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$. Now $[K : \mathbb{Q}] = 1$ by the restriction in Table 1 and so $K = \mathbb{Q}$.

Denote the minimal polynomial of α over \mathbb{Q} by P(x). If the degree of P(x) is 1, then $\alpha \in U_{\mathbb{Q}} = \{\pm 1\}$ and so the first dynamical degree is 1.

If the degree of P(x) is 2, then it can be written as $P(x) = x^2 + ax + b \in \mathbb{Z}[x] (b \in \{\pm 1\})$ and consider the following 3 cases as in Section 4. Let σ_1 be the Q-embedding $\mathbb{Q} \hookrightarrow \mathbb{C}$.

• there exists $1 \le i \le e$ such that $|\sigma_i(b)| > 1$

This case would not be occured for this condition.

• $\sigma_i(b) = 1$ for all $1 \le i \le e$

Since $\max\{|\sigma_i(a)|\}$ must be larger than 2 and close to 2, a = 3 provides the minimum value of the maximal absolute value of the conjugates of α . Thus, $\left(\frac{3+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

• $\sigma_i(b) = -1$ for all $1 \le i \le e$

a = 1 provides the minimum value of the first dynamical degrees as in Section 4. Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound of the first dynamical degrees except 1 for these types.

Type 4

Let X be a 2-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$. Now d = 1 and $[K : \mathbb{Q}] = 2$ or 4 by the restriction in Table 1 and then $[K_0 : \mathbb{Q}] = 1$ or 2, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $\left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. If a = 1, then the first dynamical degree would be 1 and so we assume $a \neq 1$. Also, by assuming that the maximal absolute value of the conjugates of \sqrt{a} is less than 2, it can be written as $\sqrt{a} = \zeta_N^m + \frac{1}{\zeta_N^m} (m, N \in \mathbb{Z}_{>0})$ with m < N, $\gcd(N, m) = 1$ and we can assume $N \neq 1, 2, 3, 4, 6$. Thus, $a = \left(\zeta_N^m + \frac{1}{\zeta_N^m}\right)^2 = \zeta_N^{2m} + \frac{1}{\zeta_N^{2m}} + 2$ and so

ame
$$N \neq 1, 2, 3, 4, 6$$
. Thus, $a = \left(\zeta_N^m + \frac{1}{\zeta_N^m}\right)^m = \zeta_N^m + \frac{1}{\zeta_N^2} + 2$ and so
$$\left[\mathbb{Q}(a) : \mathbb{Q}\right] = \begin{cases} \frac{1}{2}\varphi(\frac{N}{2}) & (N : \text{even})\\ \frac{1}{2}\varphi(N) & (N : \text{odd}) \end{cases}.$$

Now $K_0 \supset \mathbb{Q}(a) \supset \mathbb{Q}$ and so $[\mathbb{Q}(a) : \mathbb{Q}] = 1$ or 2, and this implies N = 5, 8, 10, 12, 16, 20, 24. By comparing with Proposition 2.3, since P(x) has constant term ± 1 , N = 5, 10, 24 are only possible. Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1. Thus, $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound of the first dynamical degrees except 1 for this type.

Therefore, the minimum value of the first dynamical degrees is $\frac{3+\sqrt{5}}{2} = 4\cos^2(\frac{\pi}{5})$ and realized on Type 1.

Next, we consider the case that p is a Sophie Germain prime except for p = 2.

Proposition 5.4. For a prime number $p \neq 2$ such that 2p + 1 is also a prime,

$$m(p) = 4\cos^2\left(\frac{\pi}{2p+1}\right)$$

Proof. By the restriction in Table 1, it suffices to consider only for Type 1 and 4. Define B, K, K_0, d, e and e_0 as in Section 3 for each type of X.

Type 1

Let X be a p-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or p by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, the conjugates of α are all in the interval (-2, 2). If the degree of α is 1, then $\alpha = \pm 1$ and so the first dynamical degree is 1 and so we assume the degree of α is 1. Denote the minimal polynomial of α over \mathbb{Q} by P(x), whose degree is p. By Lemma 2.4, $x^p P(x + \frac{1}{x})$ is an irreducible cyclotomic polynomial and its degree is 2p. Cyclotomic polynomials $\Phi_N(x)$ have degree 2p only for N = 2p + 1, 4p + 2 (but for p = 3, N = 7, 9, 14, 18). By comparing with Proposition 2.3, since P(x) has constant term $\pm 1, N = 2p + 1, 4p + 2$ are possible (but for p = 3, N = 7, 9, 14, 18 are possible). Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{2p+1}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{2p+1}\right)$ except 1 for this type.

Type 4

Let X be a p-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$. Now d = 1 and $[K : \mathbb{Q}] = 2$ or 2p by the restriction in Table 1 and then $[K_0 : \mathbb{Q}] = 1$ or p, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{2p+1}\right)$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. If a = 1, then the first dynamical degree would be 1 and so we assume $a \neq 1$. Also, by assuming that the maximal absolute value of the conjugates of \sqrt{a} is less than 2, it can be written as $\sqrt{a} = \zeta_N^m + \frac{1}{\zeta_N^m} (m, N \in \mathbb{Z}_{>0})$ with m < N, gcd(N, m) = 1 and we can assume $N \neq 1, 2, 3, 4, 6$. Thus, $a = \left(\zeta_N^m + \frac{1}{\zeta_N^m}\right)^2 = \zeta_N^{2m} + \frac{1}{\zeta_N^m} + 2$ and so

$$[\mathbb{Q}(a):\mathbb{Q}] = \begin{cases} \frac{1}{2}\varphi(\frac{N}{2}) & (N:\text{even})\\ \frac{1}{2}\varphi(N) & (N:\text{odd}) \end{cases}$$

Now $K_0 \supset \mathbb{Q}(a) \supset \mathbb{Q}$ and so $[\mathbb{Q}(a) : \mathbb{Q}] = 1$ or p, and this implies N = 8, 12, 2p + 1, 4p + 2, 8p + 4 (but for p = 3, N = 7, 8, 9, 12, 14, 18, 28, 36). By comparing with Proposition 2.3, since P(x) has constant term $\pm 1, N = 2p + 1, 4p + 2$ are only possible (but for p = 3, N = 7, 9, 14, 18 are possible). Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{2p+1}\right)$ except 1. Thus, $4\cos^2\left(\frac{\pi}{2p+1}\right)$ is a lower bound of the first dynamical degrees except 1 for this type.

Therefore, the minimum value of the first dynamical degrees is $4\cos^2(\frac{\pi}{2p+1})$ and realized on Type 1.

Next, we consider the case that p is not a Sophie Germain prime. First, we prove the next lemma.

Lemma 5.5. The first dynamical degree of an automorphism of a simple abelian variety is not equal to n^2 for an integer $n \in \mathbb{Z}_{\geq 2}$. Especially, The first dynamical degree of an automorphism is not equal to 4.

Proof. Suppose that there is an automorphism $f: X \to X$ of a simple abelian variety X which has the first dynamical degree n^2 . Let B be the endomorphism algebra of X. Let $\alpha \in B$ be the element corresponding to $f: X \to X$ and denote the minimal polynomial of α over \mathbb{Q} by $P(x) \in \mathbb{Z}[x]$. Since f is an automorphism, the constant term of P(x) is ± 1 .

By the assumption that the first dynamical degree of f is n^2 , the maximal absolute value of the roots of P(x) is n, and denote this root by z. Now |z| = n, $|\frac{1}{z}| = \frac{1}{n}$ and $\frac{1}{z}$ is an algebraic integer, and so $\frac{1}{z} \cdot \frac{1}{\overline{z}} = \frac{1}{n^2}$ is also an algebraic integer, a contradiction.

Proposition 5.6. For a prime number p such that 2p + 1 is not a prime,

 $m(p) = \min \left\{ \begin{array}{l} \text{the maximal absolute value of the conjugates of an} \\ \text{algebraic unit, whose conjugates are all real and positive, of degree } p \end{array} \right\}$ and also m(p) is larger than $4 + 4^{-2p-3}$.

Remark 5.7. The minimum value of the right side exists since the coefficients of the minimal polynomial is restricted by the maximal absolute value of its roots by using the triangle inequality. As a result, this equation holds for all prime numbers.

Proof of Proposition 5.6. By the restriction in Table 1, it suffices to consider only for Type 1 and 4. Define B, K, K_0, d, e and e_0 as in Section 3 for each type of X.

Type 1

Let X be a p-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or p by the restriction in Table 1.

If α has degree 1, then $\alpha = \pm 1$ and the first dynamical degree is 1 and so we assume α has degree p. Thus, by the deduction in Section 4, the set of the first dynamical degrees except 1 for this type is

 $\mathcal{A} := \left\{ \begin{array}{l} \text{the square of the maximal absolute value of the conjugates} \\ \text{of an algebraic unit, whose conjugates are all real, of degree } p \end{array} \right\}.$

Type 4

Let X be a p-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$. Now d = 1 and $[K : \mathbb{Q}] = 2$ or 2p by the restriction in Table 1 and then $[K_0 : \mathbb{Q}] = 1$ or p, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the set of the first dynamical degrees except 1 is in

 $\mathcal{A} := \left\{ \begin{array}{l} \text{the square of the maximal absolute value of the conjugates} \\ \text{of an algebraic unit, whose conjugates are all real, of degree } p \end{array} \right\},$

by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. If a has degree 1, then a = 1 and so the first dynamical degree would be 1 and so we assume a has degree p. The first dynamical degree of f is the square of the maximal absolute value of the conjugates of α and it is equal to the maximal absolute value of the conjugates of a. Now a is a totally positive algebraic unit of degree p, the first dynamical degrees except 1 are contained in

 $\mathcal{A}' := \left\{ \begin{array}{l} \text{the maximal absolute value of the conjugates of an} \\ \text{algebraic unit, whose conjugates are all real and positive, of degree } p \end{array} \right\}.$

Let *a* be an element of \mathcal{A}' . If *a* is an element of \mathcal{A} , *a* is realizable as the first dynamical degree of an automorphism of a simple abelian variety of Type 1. If *a* is not an element of \mathcal{A} , then \sqrt{a} is of degree 2p. *a* is a totally positive algebraic unit of degree *p* and let $K = \mathbb{Q}(\sqrt{-a}), K_0 = \mathbb{Q}(a), d = 1, e = 2p, e_0 = p$ and m = 1. Now *K* is a CM-field by Lemma 2.2. *K* admits the complex conjugate as a positive anti-involution of the second kind, so by Proposition 3.3, there exist a *p*-dimensional abelian variety *X* and an

Let r be an algebraic unit, whose conjugates are all real, of degree p. We can assume that the maximal absolute value of the conjugates of r is |r|. Then, r^2 is an algebraic unit, whose conjugates are all real and positive, and the maximal absolute value of the conjugate of r^2 is $r^2 = |r|^2$. Since $\mathbb{Q}(r) \supset \mathbb{Q}(r^2) \supset \mathbb{Q}$ and $[\mathbb{Q}(r) : \mathbb{Q}(r^2)] = 1$ or 2, the degree of r^2 is p. Thus, $\mathcal{A} \subset \mathcal{A}'$ and so

$$m(p) = \min(\mathcal{A}').$$

It remains to show $m(p) > 4 + 4^{-2p-3}$. If m(p) < 4, then there exists a totally real algebraic unit r of degree p, whose conjugates are all inside the interval (0, 4). Thus, r-2 is also a totally real algebraic integer and so by Lemma 2.4, the minimal polynomial of r-2 is written as $\Psi_N(x)$ for some $N \ge 3$. The degree of this polynomial is $p = \frac{1}{2}\varphi(N)$, but this would not be occured by the assumption that 2p + 1 is not a prime. Therefore, by composing with Lemma 5.5, m(p) > 4. Let r' be an algebraic unit of degree p whose conjugates are all real and positive, and at most one conjugate of r' is larger than 4. Then r'-2 is a totally real algebraic integer and its maximal absolute value of conjugates is larger than 2. Thus, by [SZ65, Theorem 2], the maximal absolute value of the conjugates of r' - 2 is larger than $2 + 4^{-2p-3}$, and it concludes that at least one conjugate of r' is larger than $4 + 4^{-2p-3}$.

Lemma 5.8. Let p be an odd prime number, a be a totally positive algebraic unit of degree p and X be a p-dimensional abelian variety. Assume \sqrt{a} is of degree 2p and denote $\alpha = \sqrt{-a}$. If $\alpha \in \operatorname{End}_{\mathbb{Q}}(X)$, then X is a simple abelian variety.

Proof. Let $f(x) \in \mathbb{Z}[x]$ be the minimal polynomial of a and so its constant term is ± 1 . Then the minimal polynomial of α over \mathbb{Q} is $g(x) = -f(-x^2) \in \mathbb{Z}[x]$ of degree 2p.

Assume X is not a simple abelian variety. By Poincare's complete reducibility theorem ([BL04, Theorem 5.3.7]), there is an isogeny

$$X \to X_1^{n_1} \times \dots \times X_r^{n_r},$$

where X_i are simple abelian varieties and not isogenious each other. Denote dim $(X_1) = g_1$ and then $p = \sum_{i=1}^r g_i n_i$. Moreover, by its corollary (cf. [BL04, Corollary 5.3.8]),

$$\operatorname{End}_{\mathbb{Q}}(X) \simeq \operatorname{M}_{n_1}(\operatorname{End}_{\mathbb{Q}}(X_1)) \oplus \cdots \oplus \operatorname{M}_{n_r}(\operatorname{End}_{\mathbb{Q}}(X_r)),$$

where $\operatorname{End}_{\mathbb{Q}}(X_i)$ are division rings.

 $\alpha \in \operatorname{End}_{\mathbb{Q}}(X)$ is decomposed as $\alpha = \alpha_1 + \cdots + \alpha_r$ with $\alpha_i \in \operatorname{M}_{n_i}(\operatorname{End}_{\mathbb{Q}}(X_i))$. Each α_i has the minimal polynomial $p_i(x)$ over \mathbb{Q} and its degree is at most $2g_in_i$. The minimal polynomial $g(x) \in \mathbb{Z}[x]$ of α is a factor of $p_1(x) \cdots p_r(x)$ and by the inequality

$$2p = \deg(g(x)) \le \sum_{i=1}^{r} \deg(p_i(x)) = \sum_{i=1}^{r} 2g_i n_i = 2p,$$

this implies r = 1. Since X is not simple, the case $g_1 = 1$, $n_1 = p$ is only possible. For this case, the minimal polynomial of α_1 is g(x).

Now

$$\operatorname{End}_{\mathbb{Q}}(X) = \operatorname{M}_p(\operatorname{End}_{\mathbb{Q}}(X_1))$$

and since $\alpha_1 \in M_p(\operatorname{End}_{\mathbb{Q}}(X_1))$ has the minimal polynomial of degree 2p over \mathbb{Q} , $\operatorname{End}_{\mathbb{Q}}(X_1) \neq \mathbb{Q}$. By Table 1, X_1 is of Type 4 and $\operatorname{End}_{\mathbb{Q}}(X_1)$ is a totally complex quadratic algebra over \mathbb{Q} . Denote $\operatorname{End}_{\mathbb{Q}}(X) = \mathbb{Q}(\sqrt{-D})$ for some $D \in \mathbb{Z}_{>0}$ a square-free integer. The polynomial g(x) is written as

$$g(x) = (x^{p} + (a_{1} + b_{1}\sqrt{-D})x^{p-1} + \dots + (a_{p} + b_{p}\sqrt{-D}))(x^{p} + (a_{1} - b_{1}\sqrt{-D})x^{p-1} + \dots + (a_{p} - b_{p}\sqrt{-D}))$$
(*)

 $=g_1(x)g_2(x)$

with $a_i + b_i \sqrt{-D} \in \mathcal{O}_{\mathbb{Q}(\sqrt{-D})}$. If $D \equiv 1, 2 \pmod{4}$, then $a_i, b_i \in \mathbb{Z}$ and if $D \equiv 3 \pmod{4}$, then $a_i, b_i \in \frac{1}{2}\mathbb{Z}$. Now the roots of g(x) are in $\mathbb{R}i$ and this implies $g_1(x) = (-1)^p g_2(-x)$. Thus, $a_p + b_p \sqrt{-D} = -a_p + b_p \sqrt{-D}$ and so $a_p = 0$. By considering the constant term of the equation (*),

$$\pm 1 = a_p^2 + b_p^2 D = b_p^2 D$$

and so D = 1.

Thus, it can be said that $\mathbb{Q}(\alpha) \supset \mathbb{Q}(i)$ and therefore $\mathbb{Q}(\alpha) \supset \mathbb{Q}(\sqrt{a}) \supset \mathbb{Q}(a)$. By the assumption, $[\mathbb{Q}(\sqrt{a}) : \mathbb{Q}(a)] = 2$ and this implies $\mathbb{Q}(\alpha) = \mathbb{Q}(\sqrt{a})$. This contradicts to $\mathbb{Q}(\alpha) \not\subset \mathbb{R}$ and $\mathbb{Q}(\sqrt{a}) \subset \mathbb{R}$, and so X is a simple abelian variety. \square

Next, for proving Theorem 5.1, we would like to prove $m(p) < (2.08)^2$ for a prime number p which is not a Sophie Germain prime. This result follows from the next proposition.

Proposition 5.9. For any prime number p > 3, there exists a monic irreducible polynomial $T(x) \in \mathbb{Z}[x]$ of degree p, whose constant term is ± 1 and whose roots are all real and inside the interval (-2.08, 2.08).

Remark 5.10. The proposition holds also for p = 2, 3, but for simplifying the proof, we assume p > 3.

For proving this proposition, we use the next lemmas.

Lemma 5.11. Define the polynomial

$$P_n(x) = \frac{x^{n-2}(x^3 - x - 1) + (x^3 + x^2 - 1)}{x - 1}, \ Q_m(x) = x^{m-3}(x^3 - x - 1) - (x^3 + x^2 - 1)$$

for $n \geq 3$ and $m \geq 4$, The next properties hold.

- (1) For any $n \ge 3$, there is at most one root of mudulus larger than 1, counted with multiplicity, among the roots of $P_n(x)$, and it is a real root.
- (2) For any $m \ge 4$, there is at most one root of mudulus larger than 1, counted with multiplicity, among the roots of $Q_m(x)$, and it is a real root.

Proof. $x^3 - x - 1$ is the minimal polynomial of (the smallest) Pisot number and by the proof of [BDGPS92, Theorem 6.4.1], the lemma holds.

Lemma 5.12. The cyclotomic polynomials which divide $Q_m(x) \in \mathbb{Z}[x]$ for some $m \ge 4$ are only $\Phi_2(x), \Phi_8(x), \Phi_{12}(x), \Phi_{18}(x), \Phi_{30}(x)$.

Proof. For $N \ge 1$, if $\Phi_N(x)$ divides $Q_m(x)$ for some $m \ge 4$, then $Q_m(\zeta_N) = 0$ for some $m \ge 4$. Thus,

$$\zeta_N^{3-m} = \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = -\zeta_N^{-3} \frac{1 + \zeta_N - \zeta_N^3}{1 + \zeta_N^{-1} - \zeta_N^{-3}} \tag{**}$$

and for N = 1 to 8, left side calculated as

$$N = 1: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = -1, \qquad N = 2: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = 1, N = 3: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = -\zeta_3^2, \qquad N = 4: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = \frac{4 + 3i}{5}, N = 5: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = -\zeta_5^3, \qquad N = 6: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = \frac{11 + 5\sqrt{3}i}{14}, N = 7: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = \frac{3 + \sqrt{7}i}{4}, \qquad N = 8: \frac{\zeta_N^3 - \zeta_N - 1}{\zeta_N^3 + \zeta_N^2 - 1} = \zeta_8.$$

Thus, N = 2, 8 are only possible. For N > 8, the equation (**) implies

$$\arg(1+\zeta_N-\zeta_N^3)=-\frac{\pi}{2N}, -\frac{\pi}{N}, -\frac{3\pi}{2N}, \dots$$

and this is equivalent to

$$\angle PAQ = \frac{\pi}{2N}, \frac{\pi}{N}, \frac{3\pi}{2N}, \dots$$

in Figure 1, Figure 2 or Figure 3. By some coordinate calculation, it is also equivalent to

$$\frac{2\sin\left(\frac{2\pi}{N}\right)\cos\left(\frac{4\pi}{N}\right)}{1+2\sin\left(\frac{2\pi}{N}\right)\sin\left(\frac{4\pi}{N}\right)} = \tan\left(\frac{\pi}{2N}\right), \tan\left(\frac{\pi}{N}\right), \tan\left(\frac{3\pi}{2N}\right), \dots$$

(1) $9 \le N < 18$

See Figure 1. For this case, point A is outside of the unit circle, so by the inscribed angle theorem, $\angle PAQ < \frac{2\pi}{N}$ and so $\angle PAQ = \frac{\pi}{2N}, \frac{\pi}{N}, \frac{3\pi}{2N}$. By observing the equations

$$\frac{2\sin(\theta)\cos(2\theta)}{1+2\sin(\theta)\sin(2\theta)} = \tan\left(\frac{\theta}{4}\right), \tan\left(\frac{\theta}{2}\right), \tan\left(\frac{3\theta}{4}\right)$$

for $\frac{\pi}{9} < \theta \le \frac{2\pi}{9}$, then $\theta = \frac{\pi}{6}$ and this corresponds to the case N = 12. (2) N = 18


See Figure 2. For this case, point A is on the unit circle, so by the inscribed angle theorem, $\angle PAQ = \frac{2\pi}{N}$, and this is the case.

(3) 18 < N

See Figure 3. For this case, point A is inside of the unit circle, so by the inscribed angle theorem, $\angle PAQ > \frac{2\pi}{N}$. Also, by $\angle PBA = \frac{2\pi}{N}$ and AB < 1 < AP, $\angle PAQ = \angle PBA + \angle APB < 2\angle PBA = \frac{4\pi}{N}$ and so $\angle PAQ = \frac{5\pi}{2N}, \frac{3\pi}{N}, \frac{7\pi}{2N}$. By observing the equations

$$\frac{2\sin(\theta)\cos(2\theta)}{1+2\sin(\theta)\sin(2\theta)} = \tan\left(\frac{5\theta}{4}\right), \tan\left(\frac{3\theta}{2}\right), \tan\left(\frac{7\theta}{4}\right)$$

for $0 < \theta < \frac{\pi}{9}$, then $\theta = \frac{\pi}{15}$ and this corresponds to the case N = 30.






Figure 1. 9 < N < 18

FIGURE 2. N = 18




Figure 3. 18 < N

Remark 5.13. For 0 < t < 2, the equation

$$\frac{2\sin(\theta)\cos(2\theta)}{1+2\sin(\theta)\sin(2\theta)} = \tan(t\theta)$$

has at most one root in the interval $(0, \frac{\pi}{4})$ (see Appendix B) and so the way for searching the suitable $N \in \mathbb{Z}_{>0}$ is returned to substituting integers into the equations in the proof.

Remark 5.14. By using the equation (**) in the proof of the lemma, it can be said that

 $Q_m(x) \text{ is divided by } \Phi_2(x) = x + 1 \iff m \equiv 1 \pmod{2}$ $Q_m(x) \text{ is divided by } \Phi_8(x) = x^4 + 1 \iff m \equiv 2 \pmod{8}$ $Q_m(x) \text{ is divided by } \Phi_{12}(x) = x^4 - x^2 + 1 \iff m \equiv 1 \pmod{12}$ $Q_m(x) \text{ is divided by } \Phi_{18}(x) = x^6 - x^3 + 1 \iff m \equiv 17 \pmod{18}$ $Q_m(x) \text{ is divided by } \Phi_{30}(x) = x^8 + x^7 - x^5 - x^4 - x^3 + x + 1 \iff m \equiv 24 \pmod{30}$

Lemma 5.15. The cyclotomic polynomials which divide $P_n(x) \in \mathbb{Z}[x]$ for some $n \ge 10$ are only $\Phi_2(x), \Phi_3(x), \Phi_5(x), \Phi_8(x), \Phi_{12}(x), \Phi_{18}(x), \Phi_{30}(x)$.

Proof. The proof is proceeding as the proof of Lemma 5.12. For $1 \le N \le 8$, N = 2, 3, 5, 8 are only possible. Also for N > 8, the only different point is the candidates for $\angle PAQ$ are changed to $\angle PAQ = \frac{\pi}{N}, \frac{2\pi}{N}, \frac{3\pi}{N}$ and N = 12, 18, 30 are possible.

Remark 5.16. As in Remark 5.14, it can be said that

 $P_n(x) \text{ is divided by } \Phi_2(x) = x + 1 \quad \Longleftrightarrow \quad n \equiv 1 \pmod{2}$ $P_n(x) \text{ is divided by } \Phi_3(x) = x^2 + x + 1 \quad \Longleftrightarrow \quad n \equiv 0 \pmod{3}$ $P_n(x) \text{ is divided by } \Phi_5(x) = x^4 + x^3 + x^2 + x + 1 \quad \Longleftrightarrow \quad n \equiv 4 \pmod{5}$ $P_n(x) \text{ is divided by } \Phi_8(x) = x^4 + 1 \quad \Longleftrightarrow \quad n \equiv 5 \pmod{8}$ $P_n(x) \text{ is divided by } \Phi_{12}(x) = x^4 - x^2 + 1 \quad \Longleftrightarrow \quad n \equiv 6 \pmod{12}$ $P_n(x) \text{ is divided by } \Phi_{18}(x) = x^6 - x^3 + 1 \quad \Longleftrightarrow \quad n \equiv 7 \pmod{18}$ $P_n(x) \text{ is divided by } \Phi_{30}(x) = x^8 + x^7 - x^5 - x^4 - x^3 + x + 1 \quad \Longleftrightarrow \quad n \equiv 8 \pmod{30}$

Remark 5.17. Similar results for Lemma 5.15 and Remark 5.16 are stated in [GHM09, Theorem 1.1].

Lemma 5.18. For each $n \ge 10$, the roots of $P_n(x)$ are all simple. Also, for each $m \ge 4$, the roots of $Q_m(x)$ are all simple.

Proof. Every $P_n(x)$ and $Q_m(x)$ has at most one root of modulus larger than 1, so this root and its conjugates are all simple. The simplicity of the other roots can be checked by using Remark 5.16 and Remark 5.14 for each cyclotomic polynomial.

Remark 5.19. Assume $n \ge 10$ and $m \ge 4$. Dividing $P_n(x)$ (resp. $Q_m(x)$) by its cyclotomic factor as many as possible, the remaining polynomial is constant or the polynomial which has only one root of modulus larger than 1 by Lemma 5.11.

By Lemma 5.18, the dividing operation is ended at most 7 times (resp. 5 times) and the degree decreases at most 29 (resp. 23). Thus, by Table A.1 and Table A.2 in Appendix A, the remaining polynomials have degree at least 4 and so these are Salem polynomials. Write $P_n(x) = C_n(x)S_n(x)$ (resp. $Q_m(x) = C'_m(x)S'_m(x)$) where $C_n(x)$ (resp. $C'_m(x)$) is a product of cyclotomic polynomials and $S_n(x)$ (resp. $S'_m(x)$) is a Salem polynomial.

Lemma 5.20. Assume $n \ge 10$ and $m \ge 4$. Let s_n be a Salem number whose minimal polynomial is $S_n(x)$ and let s'_m be a Salem number whose minimal polynomial is $S'_m(x)$. Then, the sequence $\{s_n\}_{n\ge 10}$ strictly increasing and the sequence $\{s'_m\}_{m\ge 4}$ strictly decreasing. Also, the limit of these sequence is the same and it is $1.3247\cdots$, the smallest Pisot number.

Proof. $1.3247\cdots$ is the real root of $x^3 - x - 1 \in \mathbb{Z}[x]$ and this result is proved as in the proof of [BDGPS92, Theorem 6.4.1].

For proceeding the proof, we need the next notation.

Definition 5.21. For a Salem polynomial S(x) of degree 2d, the trace polynomial of S(x) is a polynomial T(x) of degree d such that $S(x) = x^d T(x + \frac{1}{x})$.

Remark 5.22. By the definition of Salem numbers, the roots of T(x) are all real and it has only one root outside of the interval (-2, 2). Thus, for proving Proposition 5.9, we need a Salem polynomial S(x) of degree 2p and the associated trace polynomial with constant term ± 1 .

Lemma 5.23. For a Salem polynomial S(x) and its associated trace polynomial T(x), the constant term of T(x) is ± 1 if and only if |S(i)| = 1.

Proof. By the relation $S(x) = x^d T(x + \frac{1}{x})$, it is immediate that

the constant term of T(x) is $\pm 1 \iff T(0) = \pm 1 \iff |S(i)| = 1$

For $n \ge 10$ and $m \ge 4$, let $T_n(x)$ be the trace polynomial of $S_n(x)$ and let $T'_m(x)$ be the trace polynomial of $S'_m(x)$. By using Remark 5.16 and Remark 5.14, we get the condition for that when the constant term of $T_n(x)$ or $T'_m(x)$ is ± 1 .

Lemma 5.24. Assume $n \ge 10$ and $m \ge 4$. Then,

the constant term of $T_n(x)$ is $\pm 1 \iff n \equiv 0, 3, 4, 5, 6, 7, 8, 11, 12, 13, 15,$

 $16, 18, 19, 20, 21, 23 \pmod{24}$

the constant term of $T'_m(x)$ is $\pm 1 \iff m \equiv 1, 2, 3, 7, 10, 11, 13, 15, 18, 19, 23 \pmod{24}$ Proof. By calculating,

$$|\Phi_{2}(i)| = \sqrt{2}, |\Phi_{3}(i)| = 1, |\Phi_{5}(i)| = 1, |\Phi_{8}(i)| = 2, |\Phi_{12}(i)| = 3, |\Phi_{18}(i)| = 1, |\Phi_{30}(i)| = 1, |\Phi_{30}(i)| = 1, |\Phi_{12}(i)| = 1, |\Phi_{1$$

$$|P_n(i)| = \begin{cases} 2\sqrt{2} & (n \equiv 1 \pmod{4}) \\ 3 & (n \equiv 2 \pmod{4}) \\ \sqrt{2} & (n \equiv 3 \pmod{4}) \end{cases}, \ |Q_m(i)| = \begin{cases} 3\sqrt{2} & (n \equiv 1 \pmod{4}) \\ 2 & (n \equiv 2 \pmod{4}) \\ \sqrt{2} & (n \equiv 3 \pmod{4}) \end{cases}$$

By Lemma 5.23, it is enough to calculate |S(i)| and so by using Remark 5.16 and Remark 5.14, cyclicity of order 24 can be found and the proof is done.

In Appendix A, Table A.3 is enumerating the degree of $S_n(x)$ for n, which satisfies the condition in Lemma 5.24. Table A.4 is enumerating the degree of $S'_m(x)$ for m, which satisfies the condition in Lemma 5.24. Both have cyclicity of order 360 and so only $10 \le n \le 373$ and $7 \le m \le 378$ are filled in.

Proof of Proposition 5.9. Now all the double of an odd integer, larger than 3, would be appeared as the degree of $S_n(x)$ in Table A.3 or $S'_m(x)$ in Table A.4 by the cyclicity. Thus, for a prime number p > 3, 2p is appeared as the degree of $S_n(x)$ or $S'_m(x)$ with the conditions in Lemma 5.24.

If 2p is appeared as the degree of $S_n(x)$ for some n, then the associated trace polynomial $T_n(x)$ has degree p and its roots are all totally real algebraic intregers.

By Lemma 5.20, the maximal absolute value of the roots of $S_n(x)$ is less than $1.3247\cdots$ and so the maximal absolute value of the roots of $T_n(x)$ is less than $1.3247\cdots + \frac{1}{1.3247\cdots} = 2.0795\cdots < 2.08$.

The first 2p, which is not appeared as the degree of $S_n(x)$ is 26 and it is appeared as the degree of $S'_{27}(x)$ and the associated trace polynomial $T'_{27}(x)$ has degree 13. By Lemma 5.20 again, the maximal absolute value of $S'_m(x)$ is strictly decreasing, and so it suffices to show that the maximal absolute value of $T'_{27}(x)$ is less than 2.08. The Salem number for $S'_{27}(x)$ is $1.3255\cdots$ and so the maximal absolute value of $T'_{27}(x)$ is $1.3255\cdots + \frac{1}{1.3255\cdots} =$ $2.0799\cdots < 2.08$. Thus, the proof is concluded.

Proof of Theorem 5.1. Theorem 5.1 is concluded by composing Proposition 5.3, Proposition 5.4, Proposition 5.6 and Proposition 5.9. \Box

6. The minimum value of the first dynamical degrees for simple abelian varieties with lower dimensions

In this section, we calculate the minimum value of the first dynamical degrees except 1 with fixing the dimension of simple abelian varieties g = 2 to 10. The result is the following.

dimension g	minimum value
1	(all 1)
2	$4\cos^2(\pi/5) = 2.6180\cdots$
3	$4\cos^2(\pi/7) = 3.2469\cdots$
4	$2\cos(\pi/5) = 1.6180\cdots$
5	$4\cos^2(\pi/11) = 3.6825\cdots$
6	$2\cos(\pi/7) = 1.8019\cdots$
7	$4.0333 \cdots^{*1}$
8	$2\cos(\pi/5) = 1.6180\cdots$
9	$1.1509\cdots^2 = 1.3247\cdots^{*2}$
10	$1.1762 \cdots^2 = 1.3836 \cdots^{*3}$

Remark 6.1. If g = 1, then the first dynamical degree of an automorphism is the last dynamical degree and it is always 1.

The cases g = 2, 3, 5 are already done and the result is compatible with Theorem 5.1.

We use a computer algebra for g = 6 to 10. The calculation would be proceeding as dividing into the type of simple abelian varieties.

6.1. dimension g = 4

Let X be a 4-dimensional simple abelian variety and define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Let X be a 4-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 2 or 4 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, the conjugates of α are all in the interval (-2, 2). If the degree of α is 1, then $\alpha = \pm 1$ and so the first dynamical degree is 1 and so we assume the degree of α is 2 or 4. Denote the minimal polynomial of α over \mathbb{Q} by P(x), whose degree is r = 2 or 4. By Lemma 2.4, $x^r P(x + \frac{1}{x})$ is an irreducible cyclotomic polynomial and its degree is 2r = 4 or 8. Cyclotomic polynomials $\Phi_N(x)$ have degree 4 or 8 only for N = 5, 8, 10, 12, 15, 16, 20, 24, 30. By comparing with Proposition 2.3, since P(x) has constant term ± 1 , N = 5, 10, 15, 24, 30 are only possible. Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right) \exp(11)$. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2 \exp 1$ for this type.

 $^{^{*1}4.0333\}cdots$ is the maximal absolute value of the roots of $x^7 - 14x^6 + 77x^5 - 211x^4 + 301x^3 - 210x^2 + 56x - 10x^2 + 10x^$

 $^{^{*2}1.1509\}cdots$ is the absolute value of a complex root of $x^3 - x^2 + 1$

 $^{^{*3}1.1762\}cdots$ is the Lehmer number

Type 2, Type 3

Let X be a 4-dimensional simple abelian variety of Type 2 or 3 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$. Now $[K : \mathbb{Q}] = 1$ or 2 by the restriction in Table 1.

Denote the minimal polynomial of α over K by p(x) and the minimal polynomial of α over \mathbb{Q} by P(x). If the degree of p(x) is 1, then $\alpha \in U_K$ and so it is reduced to Type 1 of 2-dimensional case. Thus, $\frac{3+\sqrt{5}}{2}$ is a lower bound except 1 for this case.

If the degree of p(x) is 2, then it is written as $p(x) = x^2 + ax + b \in \mathcal{O}_K[x](b \in U_K)$ and consider the following 3 cases as in Section 4. Let $\{\sigma_i\}$ be the set of all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$.

• there exists $1 \leq i \leq e$ such that $|\sigma_i(b)| > 1$

The maximal absolute value of all conjugates of $b \in U_K$ is at least $\frac{1+\sqrt{5}}{2}$ and so the maximal absolute value of the roots of P(x) is at least $\sqrt{\frac{1+\sqrt{5}}{2}}$. This is achived by $p(x) = x^2 + \frac{1+\sqrt{5}}{2}$ and $P(x) = x^4 + x^2 - 1$. Thus, $\frac{1+\sqrt{5}}{2}$ is a lower bound except 1 for this case. • $\sigma_i(b) = 1$ for all $1 \le i \le e$

Since $\max\{|\sigma_i(a)|\}$ must be larger than 2 and close to 2, $a = \sqrt{5}$ provides the minimum value of the maximal absolute value by some calculation. For this case, the maximal absolute value of the roots of $P(x) = x^4 - 3x^2 + 1$ is $\frac{1+\sqrt{5}}{2}$ and $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

• $\sigma_i(b) = -1$ for all $1 \le i \le e$

a = 1 provides the minimum value of the first dynamical degrees as in Section 4. Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

Thus, $\frac{1+\sqrt{5}}{2}$ is the minimum value of the first dynamical degrees except 1 for these types by the next theorem.

Theorem 6.2. Define $\alpha = \sqrt{-\frac{1+\sqrt{5}}{2}}$. There exists a prime number p where $B := \left(\frac{\alpha^2, p}{\mathbb{Q}(\sqrt{5})}\right)$ is a divisional totally indefinite quaternion algebra by Theorem 3.5. On this condition, there is a positive anti-involotion on B over \mathbb{Q} . Moreover, on the case Type 2 (d = 2, e = 2, m = 1), there is an automorphism of a 4-dimensional simple abelian vaeriety which corresponds to $\alpha = \sqrt{-\frac{1+\sqrt{5}}{2}}$.

Proof. By Theorem 3.8, it is enough to find $c \in B \setminus \mathbb{Q}(\sqrt{5})$ such that $c^2 \in \mathbb{Q}(\sqrt{5})$ is totally negative.

Denote c = xi + yij $(x, y \in \mathbb{Q}(\sqrt{5}))$ where 1, i, j, ij is an $\mathbb{Q}(\sqrt{5})$ -basis of B with $i^2 = \alpha^2$ and $j^2 = p$. Then $c^2 = x^2 \cdot \left(-\frac{1+\sqrt{5}}{2}\right) - y^2 \cdot \left(-\frac{1+\sqrt{5}}{2}\right) \cdot p = \left(\frac{1+\sqrt{5}}{2}\right) (y^2p - x^2)$ by calculation. Take a pair (u, w) of integers which are larger than p, which satisfies a Pell's equation $u^2 - 5w^2 = 1$. By substituting $x = u + w\sqrt{5}, y = 1$, then,

$$c^{2} = \left(\frac{1+\sqrt{5}}{2}\right)(p-(u+w\sqrt{5})^{2}) < 0.$$

Now $1 = u^2 - 5w^2 = (u + w\sqrt{5})(u - w\sqrt{5})$ and so $0 < (u - w\sqrt{5}) < 1$.

Thus, the another conjugate of $c^2 \in \mathbb{Q}(\sqrt{5})$ is

$$\left(\frac{1-\sqrt{5}}{2}\right)(p-(u-w\sqrt{5})^2) < 0.$$

Therefore, $c^2 \in \mathbb{Q}(\sqrt{5})$ is totally negative and so the division ring B has a positive antiinvolution over \mathbb{Q} .

Now there is the order $\mathcal{O} = \mathcal{O}_{\mathbb{Q}(\sqrt{5})} \oplus \mathcal{O}_{\mathbb{Q}(\sqrt{5})} i \oplus \mathcal{O}_{\mathbb{Q}(\sqrt{5})} j \oplus \mathcal{O}_{\mathbb{Q}(\sqrt{5})} i j$ in B, so by Proposition 3.2, there exists a 4-dimensional simple abelian variety whose endomorphism ring contains \mathcal{O} . Thus, the automorphism corresponding to $i \in \mathcal{O}$ corresponds to $\alpha = \sqrt{-\frac{1+\sqrt{5}}{2}}$ and so the proof is concluded. \Box

Type 4

Let X be a 4-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$.

• If d = 1, $[K : \mathbb{Q}] = 2$ or 4 or 8 by the restriction in Table 1 and $\alpha \in U_K$.

Now $[K_0 : \mathbb{Q}] = 1$ or 2 or 4, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $\left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1, by the deduction in Type 1. If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. If a = 1, then the first dynamical degree would be 1 and so we assume $a \neq 1$. Also, by assuming that the maximal absolute value of the conjugates of \sqrt{a} is less than 2, it can be written as $\sqrt{a} = \zeta_N^m + \frac{1}{\zeta_N^m} (m, N \in \mathbb{Z}_{>0})$ with m < N, gcd(N, m) = 1 and we can assume $N \ge 3$. Thus, $a = \left(\zeta_N^m + \frac{1}{\zeta_N^m}\right)^2 = \zeta_N^{2m} + \frac{1}{\zeta_N^{2m}} + 2$ and so

$$[\mathbb{Q}(a):\mathbb{Q}] = \begin{cases} \frac{1}{2}\varphi(\frac{N}{2}) & (N:\text{even}) \\ \frac{1}{2}\varphi(N) & (N:\text{odd}) \end{cases}$$

Now $K_0 \supset \mathbb{Q}(a) \supset \mathbb{Q}$ and so $[\mathbb{Q}(a) : \mathbb{Q}] = 1$ or 2 or 4, and this implies N = 5, 8, 10, 12, 15, 16, 20, 24, 30, 32, 40, 48, 60. By comparing with Proposition 2.3, since P(x) has constant term ± 1 , N = 5, 10, 15, 24, 30, 40, 48, 60 are only possible. Thus, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos(\frac{\pi}{5})$ except 1.

Thus, $4\cos^2\left(\frac{\pi}{5}\right)$ is the minimum value of the first dynamical degrees except 1 for this case.

• If d = 2, $[K : \mathbb{Q}] = 2$ and denote $K = \mathbb{Q}(\sqrt{-D})$ for some square-free integer D > 0.

Denote the minimal polynomial of α over K by $p(x) \in \mathcal{O}_K[x]$ and the minimal polynomial of α over \mathbb{Q} by P(x).

If the degree of p(x) is 1, then $\alpha \in U_K$ and so the maximal absolute value of the conjugates is always 1.

For the case that p(x) is 2, it can be written as $p(x) = x^2 + sx + t \in \mathcal{O}_K[x]$ with $t \in U_K$. By Dirichlet's unit theorem, U_K is generated by some cyclotomic element ζ which has the smallest positive angle and denote $t = \zeta^n$.

If the degree of P(x) is 2, then it can be written as $x^2 + sx \pm 1 \in \mathbb{Z}[x]$ and the minimum value of the maximal absolte value of the roots of P(x) is $\frac{1+\sqrt{5}}{2}$ except 1.

Otherwise, P(x) has degree 4 of the form $(x^2 + sx + \zeta^n)(x^2 + \overline{s}x + \zeta^{-n}) \in \mathbb{Z}[x]$. The roots of this polynomial are $\alpha, \frac{\zeta^n}{\alpha}, \overline{\alpha}, \frac{\zeta^{-n}}{\overline{\alpha}}$ and so the maximal absolute value of these roots is $\max\{|\alpha|, \frac{1}{|\alpha|}\}$. Assuming that $\max\{|\alpha|, \frac{1}{|\alpha|}\} < 1.5$, then $s \in \mathcal{O}_K$ satisfies $|s| < 1.5 + \frac{1}{1.5} = \frac{13}{6}$. For the case $D \neq 1, 3, U_K = \{\pm 1\}$ and so t is ± 1 and this implies $s \notin \mathbb{Z}$.

• If $-D \equiv 1 \pmod{4}$, then $\mathcal{O}_{\mathbb{Q}(\sqrt{-D})} = \mathbb{Z}\left[\frac{1+\sqrt{-D}}{2}\right]$. By denoting $s = p + q\left(\frac{1+\sqrt{-D}}{2}\right) \in \mathcal{O}_{\mathbb{Q}(\sqrt{-D})} \ (p,q \in \mathbb{Z}),$ $|s|^2 = \left(p + \frac{q}{2}\right)^2 + \frac{q^2D}{4} = \frac{(2p+q)^2 + q^2D}{4} < \frac{169}{36}.$

Thus, if $D \neq 3$, then $q \neq 0$ and so D = 7, 11, 15.

• If $-D \equiv 2, 3 \pmod{4}$, then $\mathcal{O}_{\mathbb{Q}(\sqrt{-D})} = \mathbb{Z}\left[\sqrt{-D}\right]$. By denoting $s = p + q\sqrt{-D} \in \mathcal{O}_{\mathbb{Q}(\sqrt{-D})} \ (p, q \in \mathbb{Z})$,

$$|s|^2 = p^2 + q^2 D < \frac{169}{36}.$$

Thus, if $D \neq 1$, then $q \neq 0$ and so D = 2.

For any integer m, the absolute values of the roots of $(x^2 + sx + t)(x^2 + \overline{s}x + \overline{t})$ and $(x^2 + s\zeta^m x + t\zeta^{2m})(x^2 + \overline{s}\zeta^{-m}x + \overline{t}\zeta^{-2m})$ are the same and so the maximal absolute value is the same too. Also, $s \neq 0$ by assuming that the maximal absolute value of the roots is larger than 1.

Thus, only the cases $0 \le \arg(s) < \arg(\zeta)$ are worth to consider and they are enumerated in Table A.5 and the minimum of the square of the maximal absolute value of the roots of P(x) is $1.3122\cdots^2 = 1.7220\cdots$ except 1.

Therefore, the minimum value of the first dynamical degrees of automorphisms of 4dimensional simple abelian varieties is $\frac{1+\sqrt{5}}{2} = 1.6180\cdots$ except 1, and realized on a simple abelian variety of Type 2.

6.2. dimension g = 6

Let X be a 6-dimensional simple abelian variety and define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Let X be a 6-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 2 or 3 or 6 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, by using similar deduction as in the 4-dimensional case, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1 for this type.

Type 2, Type 3

Let X be a 6-dimensional simple abelian variety of Type 2 or 3 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$. Now $[K : \mathbb{Q}] = 1$ or 3 by the restriction in Table 1.

Denote the minimal polynomial of α over K by p(x) and the minimal polynomial of α over \mathbb{Q} by P(x). If the degree of p(x) is 1, then $\alpha \in U_K$ and so it is reduced to Type 1 of 3-dimensional case. Thus, $4\cos^2\left(\frac{\pi}{7}\right)$ is a lower bound except 1 for this case.

If the degree of p(x) is 2, it can be written as $p(x) = x^2 + ax + b \in \mathcal{O}_K[x](b \in U_K)$ and consider the following 3 cases as in Section 4. Let $\{\sigma_i\}$ be the set of all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$.

• there exists $1 \le i \le e$ such that $|\sigma_i(b)| > 1$

The maximal absolute value of all conjugates of $b \in U_K$ is at least $2\cos\left(\frac{\pi}{7}\right)$ and so the maximal absolute value of the roots of P(x) is at least $\sqrt{2\cos\left(\frac{\pi}{7}\right)}$. This is achived by $p(x) = x^2 - 2\cos\left(\frac{2\pi}{7}\right)$ and $P(x) = (x^2 - 2\cos\left(\frac{2\pi}{7}\right))(x^2 - 2\cos\left(\frac{4\pi}{7}\right))(x^2 - 2\cos\left(\frac{6\pi}{7}\right))$. Thus, $2\cos\left(\frac{\pi}{7}\right) = 1.8019\cdots$ is a lower bound except 1 for this case. • $\sigma_i(b) = 1$ for all $1 \le i \le e$

 $\max\{|\sigma_i(a)|\}\$ must be larger than 2 and close to 2. By using a computer algebra, totally real algebraic integers of degree 3 whose conjugates have modulus less than 2.25 are enumerated in Table A.6. $a = 2.1149\cdots$ provides the minimum of the maximal absolute value. For this case, the maximal absolute value of the roots of P(x) is $1.4012\cdots$ and so $1.4012\cdots^2 = 1.9635\cdots$ is a lower bound except 1 for this case.

•
$$\sigma_i(b) = -1$$
 for all $1 \le i \le e$

a = 1 provides the minimum value of the first dynamical degrees as in Section 4. Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

Thus, $2\cos\left(\frac{\pi}{7}\right) = 1.8019\cdots$ is the minimum value of the first dynamical degrees except 1 for these types by the next theorem.

Theorem 6.3. Define $\alpha := \sqrt{\zeta_7 + \frac{1}{\zeta_7}}$. Theorem 3.5 implies that there exists a prime number p where $B := \begin{pmatrix} \frac{\zeta_7 + \frac{1}{\zeta_7}, p}{\mathbb{Q}(\zeta_7 + \frac{1}{\zeta_7})} \end{pmatrix}$ is a divisional totally indefinite quaternion algebra. On this condition, there is a positive anti-involotion on B over \mathbb{Q} . Moreover, there is an automorphism of a 6-dimensional simple abelian variety of Type 2 (d = 2, e = 3, m = 1) which corresponds to $\alpha = \sqrt{\zeta_7 + \frac{1}{\zeta_7}}$.

Proof. By Theorem 3.8, it is enough to find $c \in B \setminus \mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)$ such that $c^2 \in \mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)$ is totally negative.

Denote $c = xi + yij \in B$ where 1, i, j, ij is an $\mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)$ -basis of B with $i^2 = \zeta_7 + \frac{1}{\zeta_7}$ and $j^2 = p$. Then $c^2 = x^2 \cdot \left(\zeta_7 + \frac{1}{\zeta_7}\right) - y^2 \cdot \left(\zeta_7 + \frac{1}{\zeta_7}\right) \cdot p = \left(\zeta_7 + \frac{1}{\zeta_7}\right) (x^2 - y^2 p)$ by calculation. By substituting $x = a + b\left(\zeta_7 + \frac{1}{\zeta_7}\right), y = 1$ with $a, b \in \mathbb{Z}$, then,

$$c^{2} = \left(\zeta_{7} + \frac{1}{\zeta_{7}}\right) \left(\left(a + b\left(\zeta_{7} + \frac{1}{\zeta_{7}}\right)\right)^{2} - p \right).$$

The other conjugates of $c^2 \in \mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)$ are

$$\left(\zeta_7^2 + \frac{1}{\zeta_7^2}\right) \left(\left(a + b\left(\zeta_7^2 + \frac{1}{\zeta_7^2}\right)\right)^2 - p \right)$$

and

$$\left(\zeta_7^3 + \frac{1}{\zeta_7^3}\right) \left(\left(a + b\left(\zeta_7^3 + \frac{1}{\zeta_7^3}\right)\right)^2 - p \right)$$

Take (a, b) as a > 0, b < 0, $0 < a + b\left(\zeta_7 + \frac{1}{\zeta_7}\right) < 1$ and then by $a \to \infty$, $b \to -\infty$, it holds that $c^2 < 0$, $\left|a + b\left(\zeta_7^2 + \frac{1}{\zeta_7^2}\right)\right| \to \infty$ and $\left|a + b\left(\zeta_7^3 + \frac{1}{\zeta_7^3}\right)\right| \to \infty$. Thus, all the conjugates of c^2 can be negative and so it defines a positive anti-involution on B over \mathbb{Q} ,

Now there is the order $\mathcal{O} = \mathcal{O}_{\mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)} \oplus \mathcal{O}_{\mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)} i \oplus \mathcal{O}_{\mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)} j \oplus \mathcal{O}_{\mathbb{Q}\left(\zeta_7 + \frac{1}{\zeta_7}\right)} i j$ in B, so by Proposition 3.2, there exists a 6-dimensional simple abelian variety whose endomorphism ring contains \mathcal{O} . Thus, the automorphism corresponding to $i \in \mathcal{O}$ corresponds to $\alpha = \sqrt{\zeta_7 + \frac{1}{\zeta_7}}$ and so the proof is concluded. \Box

Type 4

Let X be a 6-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$.

Now d = 1 and $[K : \mathbb{Q}] = 2$ or 4 or 6 or 12 by the restriction in Table 1 and $\alpha \in U_K$.

Now $[K_0 : \mathbb{Q}] = 1$ or 2 or 3 or 6, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{7}\right)$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. By using similar deduction as in the 4-dimensional case, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{5}\right)$ except 1.

Therefore, the minimum value of the first dynamical degrees of automorphisms of 6dimensional simple abelian varieties is $2\cos\left(\frac{\pi}{7}\right)$ except 1, and realized on a simple abelian variety of Type 2.

6.3. dimension g = 7

By using a computer algebra,

 $\min \left\{ \begin{array}{l} \text{the maximal absolute value of the conjugates of an} \\ \text{algebraic unit, whose conjugates are all real and positive, of degree } p \end{array} \right\}$

is $4.0333\cdots$, the maximum root of the polynomial $x^7 - 14x^6 + 77x^5 - 211x^4 + 301x^3 - 210x^2 + 56x - 1$ by Table A.7. By Proposition 5.6, $m(p) = 4.0333\cdots$ and it is realized on a simple abelian variety of Type 4.

6.4. dimension g = 8

Let X be an 8-dimensional simple abelian variety and define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Let X be an 8-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 2 or 4 or 8 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, by using similar deduction as in the 4-dimensional case, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1 for this type.

Type 2, Type 3

Let X be an 8-dimensional simple abelian variety of Type 2 or 3 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$. Now $[K : \mathbb{Q}] = 1$ or 2 or 4 by the restriction in Table 1.

Denote the minimal polynomial of α over K by p(x) and the minimal polynomial of α over \mathbb{Q} by P(x). If the degree of p(x) is 1, then $\alpha \in U_K$ and so it is reduced to Type 1 of 4-dimensional case. Thus, the minimum value is $\frac{3+\sqrt{5}}{2}$ for this case.

If the degree of p(x) is 2, it can be written as $p(x) = x^2 + ax + b \in \mathcal{O}_K[x](b \in U_K)$ and consider the following 3 cases as in Section 4. Let $\{\sigma_i\}$ be the set of all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$.

• there exists $1 \le i \le e$ such that $|\sigma_i(b)| > 1$

The maximal absolute value of all conjugates of $b \in U_K$ is at least $2\cos\left(\frac{\pi}{5}\right)$ and so the maximal absolute value of the roots of P(x) is at least $\sqrt{2\cos\left(\frac{\pi}{5}\right)}$. This is achived as in 4-dimensional case. Thus, $2\cos\left(\frac{\pi}{5}\right) = 1.6180\cdots$ is a lower bound except 1 for this case.

This is realizable for the case of Type 2 (d = 2, e = 2, m = 2) as in Theorem 6.2.

• $\sigma_i(b) = 1$ for all $1 \le i \le e$

 $\max\{|\sigma_i(a)|\}\$ must be larger than 2 and close to 2. By using a computer algebra, totally real algebraic integers of degree 4 whose conjugates have modulus less than 2.1 are enumerated in Table A.8. $a = 2.0614\cdots$ provides the minimum of the maximal absolute value. For this case, the maximal absolute value of the roots of P(x) is $1.2806\cdots$ and so $1.2806\cdots^2 = 1.6400\cdots$ is a lower bound except 1 for this case.

• $\sigma_i(b) = -1$ for all $1 \le i \le e$

a = 1 provides the minimum value of the first dynamical degrees as in Section 4. Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

Therefore, $2\cos\left(\frac{\pi}{5}\right)$ is the minimum value of the first dynamical degrees for these types. Type 4

Let X be an 8-dimensional simple abelian variety of Type 4 and $f \in End(X)$ be an automorphism of X which corresponds to $\alpha \in B$.

• If d = 1, $[K : \mathbb{Q}] = 2$ or 4 or 8 or 16 by the restriction in Table 1 and $\alpha \in U_K$.

Now $[K_0 : \mathbb{Q}] = 1$ or 2 or 4 or 8, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $\left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. By using similar deduction as in the 4-dimensional case, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1.

Thus, $4\cos^2\left(\frac{\pi}{5}\right)$ is the minimum value of the first dynamical degrees except 1 for this case.

• If d = 2, $[K : \mathbb{Q}] = 2$ or 4 is only possible by the restriction in Table 1.

If $[K : \mathbb{Q}] = 2$, then it is reduced to the case of 4-dimensional, and $1.3122\cdots^2 = 1.7220\cdots$ is a lower bound of the first dynamical degrees except 1.

For the case $[K : \mathbb{Q}] = 4$, let p(x) be the minimal polynomial of α over K and let P(x) be the minimal polynomial of α over \mathbb{Q} . If the degree of p(x) is 1, then $\alpha \in U_K$ and so $\frac{3+\sqrt{5}}{2}$ is a lower bound except 1 as the case of 2-dimensional.

For the case that the degree of p(x) is 2, it can be written as $p(x) = x^2 + sx + t \in \mathcal{O}_K[x]$ with $t \in U_K$. If max{modulus of conjugates of t} > 1, then the maximal absolute value of the roots of P(x) is at least $\sqrt{\max\{\text{modulus of conjugates of }t\}}$ and so $\sqrt{\frac{1+\sqrt{5}}{2}}$ is a lower bound. Thus, $\frac{1+\sqrt{5}}{2}$ is a lower bound of the first dynamical degrees except 1 for this case. For the case max{modulus of conjugates of t} = 1, by $[K : \mathbb{Q}] = 4$,

$$t \in \{1, -1, \zeta_3, \zeta_3^2, i, -i, \zeta_5, \zeta_5^2, \zeta_5^3, \zeta_5^4, \zeta_6, \zeta_6^5, \zeta_8, \zeta_8^3, \zeta_8^5, \zeta_8^7, \zeta_{10}, \zeta_{10}^3, \zeta_{10}^7, \zeta_{10}^9, \zeta_{12}, \zeta_{12}^5, \zeta_{12}^7, \zeta_{12}^{11}\}$$

For considering the absolute value of the roots of P(x), by the correspondence

$$\begin{split} (x^2 + sx + \zeta_3) \leftrightarrow (x^2 + s\zeta_3x + 1) \leftrightarrow (x^2 + s\zeta_3^2x + \zeta_3^2), \\ (x^2 + sx + \zeta_5) \leftrightarrow (x^2 + s\zeta_5x + \zeta_5^3) \leftrightarrow (x^2 + s\zeta_5^2x + 1) \\ & \leftrightarrow (x^2 + s\zeta_5^3x + \zeta_5^2) \leftrightarrow (x^2 + s\zeta_5^4x + \zeta_5^4), \\ (x^2 + sx + \zeta_6) \leftrightarrow (x^2 + s\zeta_6x - 1) \leftrightarrow (x^2 + s\zeta_6^2x + \zeta_6^5), \\ (x^2 + sx + \zeta_{10}) \leftrightarrow (x^2 + s\zeta_{10}x + \zeta_{10}^3) \leftrightarrow (x^2 + s\zeta_{10}^2x - 1) \\ & \leftrightarrow (x^2 + s\zeta_{10}^3x + \zeta_{10}^7) \leftrightarrow (x^2 + s\zeta_{10}^4x + \zeta_{10}^9), \\ (x^2 + sx + i) \leftrightarrow (x^2 + six - i), \\ (x^2 + sx + \zeta_8) \leftrightarrow (x^2 + s\zeta_8x + \zeta_8^3) \leftrightarrow (x^2 + s\zeta_8^2x + \zeta_8^5) \leftrightarrow (x^2 + s\zeta_8^3x + \zeta_8^7), \\ (x^2 + sx + \zeta_{12}) \leftrightarrow (x^2 + s\zeta_{12}^2x + \zeta_{12}^5) \leftrightarrow (x^2 + s\zeta_{12}^3x + \zeta_{12}^7) \leftrightarrow (x^2 + s\zeta_{12}^5x + \zeta_{11}^{11}), \end{split}$$

it suffices to consider the cases $t = 1, -1, i, \zeta_8, \zeta_{12}$. Denote $K_0 = \mathbb{Q}(\sqrt{D})$ where $D \in \mathbb{Z}_{>0}$ is a square-free integer. $s \in \mathcal{O}_K$ is of degree at most 2 over K_0 .

(1) Case t = 1

It can be written as $P(x) = (x^2 + sx + 1)(x^2 + s_1x + 1)(x^2 + s_2x + 1)(x^2 + s_3x + 1)$ where $s = s_0, s_1, s_2, s_3$ are conjugates over \mathbb{Q} . Let

$$R(X) = (X^{2} + (a + b\sqrt{D})X + a' + b'\sqrt{D})(X^{2} + (a - b\sqrt{D})X + a' - b'\sqrt{D})$$

= X⁴ + 2aX³ + (a² - b²D + 2a')X² + (2aa' - 2bb'D)X + a'² - b'²D (6.1)

be a polynomial, with $a + b\sqrt{D}$, $a' + b'\sqrt{D} \in \mathcal{O}_{\mathbb{Q}(\sqrt{D})}$, which has $s = s_0, s_1, s_2, s_3$ as the roots. For N = 1.28, all roots of P(x) have modulus less than N if and only if $s = s_0, s_1, s_2, s_3$ are all inside the domain

$$E := \left\{ z \in \mathbb{C} \mid \operatorname{Re}(z)^2 \left(N - \frac{1}{N} \right)^2 + \operatorname{Im}(z)^2 \left(N + \frac{1}{N} \right)^2 < \left(N + \frac{1}{N} \right)^2 \left(N - \frac{1}{N} \right)^2 \right\}$$

The quartic polynomials, whose roots are all in E, are showed in Table A.9 and considering the polynomial is of the form (6.1), or not. Thus, $1.2720 \cdots = \sqrt{\frac{1+\sqrt{5}}{2}}$ is the minimum value and so $\frac{1+\sqrt{5}}{2}$ is a lower bound except 1 for this case.

(2) Case t = -1

It can be written as $P(x) = (x^2 + sx - 1)(x^2 + s_1x - 1)(x^2 + s_2x - 1)(x^2 + s_3x - 1)$ where $s = s_0, s_1, s_2, s_3$ are conjugates over \mathbb{Q} .

For N = 1.28, all roots of P(x) have modulus less than N if and only if $s = s_0, s_1, s_2, s_3$ are all inside the domain

$$E_1 := \left\{ z \in \mathbb{C} \mid \operatorname{Re}(z)^2 \left(N + \frac{1}{N} \right)^2 + \operatorname{Im}(z)^2 \left(N - \frac{1}{N} \right)^2 < \left(N + \frac{1}{N} \right)^2 \left(N - \frac{1}{N} \right)^2 \right\}$$

The quartic polynomials, whose roots are all in E_1 , are showed in Table A.10 and considering the polynomial is of the form (6.1), or not. Thus, $1.2720 \cdots = \sqrt{\frac{1+\sqrt{5}}{2}}$ is the minimum value and so $\frac{1+\sqrt{5}}{2}$ is a lower bound except 1 for this case.

(3) Case t = i

It can be written as $P(x) = (x^2 + sx + i)(x^2 + tx + i)(x^2 + \overline{s}x - i)(x^2 + \overline{t}x - i)$ where $s, t, \overline{s}, \overline{t}$ are conjugates over \mathbb{Q} . Especially, s, t are conjugates over $\mathbb{Q}(i)$.

z = s, t is the roots of

$$R_1(z) = z^2 + (a+bi)z + (c+di) \in \mathbb{Z}[i][z].$$
(6.2)

For N = 1.28, all roots of P(x) have modulus less than N if and only if s, t are inside the domain

$$E_2 := \{ z \in \mathbb{C} \mid z \cdot \zeta_8 \in E_1 \}$$

The quadratic polynomials of the form (6.2), whose roots are all in E_2 , are showed in Table A.11. Thus, $1.2720 \cdots = \sqrt{\frac{1+\sqrt{5}}{2}}$ is the minimum value except 1 and so $\frac{1+\sqrt{5}}{2}$ is a lower bound except 1 for this case.

(4) Case $t = \zeta_8$

 $K = \mathbb{Q}(\zeta_8)$ and $K_0 = \mathbb{Q}\left(\zeta_8 + \frac{1}{\zeta_8}\right) = \mathbb{Q}(\sqrt{2})$. It can be written as $P(x) = (x^2 + sx + \zeta_8)(x^2 + tx + \zeta_8^3)(x^2 + \bar{t}x + \zeta_8^5)(x^2 + \bar{s}x + \zeta_8^7)$ where s, t, \bar{t}, \bar{s} are conjugates over \mathbb{Q} . Especially, s, t are conjugates over $\mathbb{Q}(\sqrt{2}i)$.

z = s, t is the roots of

$$R_2(z) = z^2 + (a + b\sqrt{2}i)z + (c + d\sqrt{2}i) \in \mathbb{Z}[\sqrt{2}i][z].$$
(6.3)

For N = 1.28, all roots of P(x) have modulus less than N if and only if s is inside the domain

$$E_3 := \left\{ z \in \mathbb{C} \mid z \cdot \zeta_{16}^3 \in E_1 \right\}$$

and t is inside the domain

$$E_4 := \{ z \in \mathbb{C} \mid z \cdot \zeta_{16} \in E_1 \}.$$

The quadratic polynomials of the form (6.3), with the roots $s, t \in \mathbb{Q}(\zeta_8)$ such that $s \in E_3, t \in E_4$ and $(x^2 + sx + \zeta_8)(x^2 + tx + \zeta_8^3) \in \mathbb{Z}[\sqrt{2}i][z]$, are showed in Table A.12. Thus, 1.28² is a lower bound except 1 for this case.

(5) Case $t = \zeta_{12}$

 $K = \mathbb{Q}(\zeta_{12}) \text{ and } K_0 = \mathbb{Q}\left(\zeta_{12} + \frac{1}{\zeta_{12}}\right) = \mathbb{Q}(\sqrt{3}). \text{ It can be written as } P(x) = (x^2 + sx + \zeta_{12})(x^2 + tx + \zeta_{12}^5)(x^2 + tx + \zeta_{12}^7)(x^2 + sx + \zeta_{12}^{11}) \text{ where } s, t, \overline{t}, \overline{s} \text{ are conjugates over } \mathbb{Q}.$ Especially, s, t are conjugates over $\mathbb{Q}(i)$.

z = s, t is the roots of

$$R_3(z) = z^2 + (a+bi)z + (c+di) \in \mathbb{Z}[i][z].$$
(6.4)

For N = 1.28, all roots of P(x) have modulus less than N if and only if s is inside the domain

$$E_5 := \left\{ z \in \mathbb{C} \mid z \cdot \zeta_{24}^5 \in E_1 \right\}$$

and t is inside the domain

$$E_6 := \{ z \in \mathbb{C} \mid z \cdot \zeta_{24} \in E_1 \} \,.$$

The quadratic polynomials of the form (6.4), with the roots $s, t \in \mathbb{Q}(\zeta_{12})$ such that $s \in E_5, t \in E_6$ and $(x^2 + sx + \zeta_{12})(x^2 + tx + \zeta_{12}^5) \in \mathbb{Z}[i][z]$, are showed in Table A.13. Thus, 1.28² is a lower bound except 1 for this case.

Therefore, the minimum value of the first dynamical degrees of automorphisms of 8dimensional simple abelian varieties is $\frac{1+\sqrt{5}}{2}$ except 1, and realized on a simple abelian variety of Type 2.

6.5. dimension g = 9

Let X be a 9-dimensional simple abelian variety and define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Let X be a 9-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 3 or 9 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, by using similar deduction as in the 4-dimensional case, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{7}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{7}\right)$ except 1 for this type.

Type 4

Let X be a 9-dimensional simple abelian variety of Type 4 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$.

• If d = 1, $[K : \mathbb{Q}] = 2$ or 6 or 18 by the restriction in Table 1 and $\alpha \in U_K$.

Now $[K_0 : \mathbb{Q}] = 1$ or 3 or 9, respectively. Denote the minimal polynomial of α over \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{7}\right)$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. By using similar deduction as in the 4-dimensional case, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{7}\right)$ except 1. • If d = 3, $[K : \mathbb{Q}] = 2$ is only possible by the restriction in Table 1.

Denote the minimal polynomial of α over K by $p(x) \in \mathcal{O}_K[x]$ and the minimal polynomial of α over \mathbb{Q} by P(x).

If the degree of p(x) is 1, then $\alpha \in U_K$ and so the maximal absolute value of the conjugates is always 1.

If the degree of p(x) is 3, then it can be written as $p(x) = x^3 + sx^2 + tx + u \in \mathcal{O}_K[x]$ with $u \in U_K$. By Dirichlet's unit theorem, U_K contains only cyclotomic elements and so by $[K : \mathbb{Q}] = 2$,

$$u \in \{1, -1, i, -i, \zeta_6, \zeta_6^2, \zeta_6^4, \zeta_6^5\}$$

For considering the absolute value of the roots of P(x), by the correspondence

it suffices to consider the cases $u = 1, \zeta_6$. Denote $K = \mathbb{Q}(\sqrt{-D})$ where $D \in \mathbb{Z}_{>0}$ is a squarefree integer. Write $p(x) = x^3 + (a + b\sqrt{-D})x^2 + (a' + b'\sqrt{-D})x + u \in \mathcal{O}_K[x]$ and then $P(x) = (x^3 + (a + b\sqrt{-D})x^2 + (a' + b'\sqrt{-D})x + u)(x^3 + (a - b\sqrt{-D})x^2 + (a' - b'\sqrt{-D})x + \overline{u}).$ For N = 1.3, all roots of P(x) have modulus less than N if and only if all roots of p(x)have modulus less than N.

- (1) Case u = 1, b = 0, b' = 0 $p(x) = x^3 + ax^2 + a'x + 1 \in \mathbb{Z}[x]$ and by Table A.14, the minimum value of the maximal absolute value of the conjugates is $1.1509\cdots$. Thus, $1.1509\cdots^2 = 1.3247\cdots$ is a lower bound of the first dynamical degrees except 1 for this case.
- (2) Case u = 1, (b, b') ≠ (0, 0)
 Assuming that all roots of p(x) have modulus less than N = 1.3, the minimum value of the maximal absolute value of the conjugates is 1.1509 ··· by Table A.15. Thus, 1.1509 ···² = 1.3247 ··· is a lower bound of the first dynamical degrees except 1 for this case.
- (3) Case $u = \zeta_6$

Assuming that all roots of p(x) have modulus less than N = 1.3, the minimum value of the maximal absolute value of the conjugates is $1.2167\cdots$ by Table A.16. Thus, $1.2167\cdots^2 = 1.4803\cdots$ is a lower bound of the first dynamical degrees except 1 for this case.

Therefore, $1.3247 \cdots$ is a lower bound of the first dynamical degrees except 1.

It remains to show that this is the minimum value.

Denote the real root of $x^3 - x^2 + 1 \in \mathbb{Z}[x]$ by λ and the other complex roots by z, \overline{z} . Define $E := \mathbb{Q}(\lambda, z), F := \mathbb{Q}(\sqrt{-23})$, and $K := \mathbb{Q}(\lambda)$.

Lemma 6.4. E contains F as a subfield. Moreover, E/F is a Galois extension of degree 3 and it is cyclic.

Proof. $E \supset K \supset \mathbb{Q}$ is a sequence of field extensions and it is clear that E/\mathbb{Q} is a Galois extension, $[K : \mathbb{Q}] = 3$ and $[E : K] \leq 2$. $E \not\subset \mathbb{R}$ and $K \subset \mathbb{R}$ implies $E \neq K$ and so [E : K] = 2 and $[E : \mathbb{Q}] = 6$.

The equation

$$x^{3} - x^{2} + 1 = (x - \lambda)(x^{2} + (\lambda - 1)x + \lambda^{2} - \lambda)$$

implies

$$z, \overline{z} = \frac{1 - \lambda \pm \sqrt{-3\lambda^2 + 2\lambda + 1}}{2}$$

and by

$$(3\lambda^2 - 2\lambda)\sqrt{-3\lambda^2 + 2\lambda + 1} = \sqrt{(3\lambda^2 - 2\lambda)^2(-3\lambda^2 + 2\lambda + 1)}$$

= $\sqrt{-23}$,

 $\mathbb{Q}(\lambda, \sqrt{-23}) = \mathbb{Q}(\lambda, z) = E$ and so $E \supset F$.

Now $[F : \mathbb{Q}] = 2$ and so E/F is a Galois extension of degree 3. The Galois group $\operatorname{Gal}(E/F)$ is a group of order 3 and so it is cyclic.

Define

over \mathbb{Q} and then σ is a generator of $\operatorname{Gal}(E/F)$.

Lemma 6.5. There exists $u \in F^{\times}$ such that the order of u in $F^{\times}/N_{E/F}(E^{\times})$ is 3.

Proof. First consider the prime factorization of the prime ideal $3\mathbb{Z}$ in E and F.

 σ

For the field extension F/\mathbb{Q} and the primitive element $\theta = \sqrt{-23}$, the conductor \mathfrak{J} of $\mathbb{Z}[\sqrt{-23}]$ in $\mathcal{O}_F = \mathbb{Z}\left[\frac{1+\sqrt{-23}}{2}\right]$ contains 2 and so $3\mathbb{Z}$ and \mathfrak{J} are relatively prime. By using Remark 2.5, there is a prime ideal factorization

$$\mathcal{BO}_F = \mathfrak{p}_1\mathfrak{p}_2 \quad (\mathfrak{p}_1, \ \mathfrak{p}_2 \subset \mathcal{O}_F \text{ are prime ideals}),$$

which corresponds to

$$X^{2} + 23 = (X+1)(X+2) \in (\mathbb{Z}/3\mathbb{Z})[X].$$

Now F/\mathbb{Q} is a Galois extension and so $[\mathcal{O}_F/\mathfrak{p}_1 : \mathbb{Z}/3\mathbb{Z}] = 1$. For the field extension E/F and the primitive element $\theta' = \lambda$, by using Remerk 2.7, the conductor \mathfrak{J}' of $\mathcal{O}_F[\lambda]$ in \mathcal{O}_E contains

$$\delta = \Delta_{E/F}(1,\lambda,\lambda^2) = \det \begin{pmatrix} 1 & \lambda & \lambda^2 \\ 1 & z & z^2 \\ 1 & \overline{z} & \overline{z}^2 \end{pmatrix}^2$$
$$= (\lambda - z)^2 (\lambda - \overline{z})^2 (z - \overline{z})^2 = -23$$

by using the equation of Vandermonde's determinant and the discriminant of a polynomial for $x^3 - x^2 + 1 \in \mathbb{Z}[x]$. This implies \mathfrak{p}_1 is relatively prime to \mathfrak{J}' and so by using Remark 2.5,

 $\mathfrak{p}_1\mathcal{O}_E = \mathfrak{q} \quad (\mathfrak{q} \subset \mathcal{O}_F \text{ is a prime ideal}),$

which corresponds to

$$X^3 - X^2 + 1$$
 is irreducible in $(\mathcal{O}_F/\mathfrak{p}_1)[X] = (\mathbb{Z}/3\mathbb{Z})[X].$

The pair $(p = 3, \mathfrak{p} = \mathfrak{p}_1, \mathfrak{q})$ satisfies the following conditions.

• \mathfrak{p} is over $p\mathbb{Z}$ and \mathfrak{q} is over \mathfrak{p} .

- $\sigma(\mathbf{q}) = \mathbf{q}$ for all $\sigma \in \operatorname{Gal}(E/F)$.
- the multiplicity of \mathfrak{q} of the prime ideal factorization of the ideal $p\mathcal{O}_E$ is not a multiple of 3.

Thus, by Lemma 3.7, there is no $\alpha \in E^{\times} \setminus U_E$ such that $N_{E/F}(\alpha) = p$. Moreover, since p is not an unit of \mathbb{Z} , there is no $\alpha \in U_E$ such that $N_{E/F}(\alpha) = p$. Thus, $p \notin N_{E/F}(E^{\times})$ and the order of p in $F^{\times}/N_{E/F}(E^{\times})$ is 3.

Remark 6.6. Take a symbol v with $v^3 = u$ and define $B = \bigoplus_{i=0}^2 v^i E$ as in Theorem 3.6, and then B is a division ring and also a central simple algebra over F with [B:F] = 9.

Also, the order $\mathcal{O} := \bigoplus_{i=0}^2 v^i \mathcal{O}_E \subset B$ contains both λ and $\frac{1}{\lambda}$.

Theorem 6.7. For the division algebra B in Remark 6.6, there is a positive anti-involution of the second kind on B over \mathbb{Q} .

Proof. Define

$$\phi: \begin{array}{ccc} B & \to & B \\ v^i \cdot a & \mapsto & v^i \cdot (\sigma^i \circ \tau)(a) & (a \in E) \end{array}$$

over \mathbb{Q} , where τ is the complex conjugation map.

By using the equation $\sigma \circ \tau \circ \sigma = \tau$,

$$\begin{split} \phi(1) &= 1, \\ \phi(\phi(v^i \cdot a)) = \phi(v^i \cdot \sigma^i \circ \tau(a)) = v^i \cdot (\sigma^i \circ \tau \circ \sigma^i \circ \tau)(a) = v^i \cdot a, \\ \phi(v^i \cdot a \cdot v^{i'} \cdot a') &= \phi(v^i v^{i'} \cdot \sigma^{i'}(a)a') = \phi(v^{i+i'} \cdot \sigma^{i'}(a)a') \\ &= v^{i+i'} \cdot (\sigma^{i+i'} \circ \tau)(\sigma^{i'}(a)a') \\ &= v^{i+i'} \cdot (\sigma^i \circ \tau)(a) \cdot (\sigma^{i+i'} \circ \tau)(a') \end{split}$$

and

$$\phi(v^{i'} \cdot a')\phi(v^i \cdot a) = v^{i'} \cdot (\sigma^{i'} \circ \tau)(a') \cdot v^i \cdot (\sigma^i \circ \tau)(a)$$
$$= v^{i'}v^i \cdot (\sigma^{i+i'} \circ \tau)(a') \cdot (\sigma^i \circ \tau)(a)$$
$$= u^{i+i'} \cdot (\sigma^i \circ \tau)(a) \cdot (\sigma^{i+i'} \circ \tau)(a')$$

hold and so ϕ is the anti-involution map over \mathbb{Q} . Also, the restriction map $\phi|_F$ is the complex conjugation map and so ϕ is of the second kind. Thus, by composing with Theorem 3.9, there exists a positive anti-involution map over \mathbb{Q} of the second kind. \Box

By Proposition 3.3, there is a 9-dimensional simple abelian variety with an automorphism corresponding to λ for the case d = 3, e = 2, $e_0 = 1$ and m = 1.

The first dynamical degree of the automorphism is the square of the maximal absolute value of the conjugates of λ and it is $1.1509 \cdots^2 = 1.3247 \cdots$.

Therefore, the minimum value of the first dynamical degrees of automorphisms of 9dimensional simple abelian varieties is $1.3247\cdots$ except 1, and realized on a simple abelian variety of Type 4.

Remark 6.8. By some deduction, we can check that $1.3247 \cdots$ is the real root of $x^3 - x - 1 \in \mathbb{Z}[x]$ and it is the smallest Pisot number (cf. [BDGPS92, Theorem 7.2.1]).

6.6. dimension g = 10

Let X be a 10-dimensional simple abelian variety and define B, K, K_0, d, e and e_0 as in Section 3.

Type 1

Let X be a 10-dimensional simple abelian variety of Type 1 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in U_K$ as in Section 4. Now $[K : \mathbb{Q}] = 1$ or 2 or 5 or 10 by the restriction in Table 1.

Assuming that the first dynamical degree is less than 4, by using similar deduction as in the 4-dimensional case, the minimum value of the maximal absolute value of the roots of P(x) is $2\cos\left(\frac{\pi}{5}\right)$ except 1. The realizability is concerned in Section 4. Thus, the minimum value of the first dynamical degrees is $4\cos^2\left(\frac{\pi}{5}\right) = \left(\frac{1+\sqrt{5}}{2}\right)^2$ except 1 for this type.

Type 2, Type 3

Let X be a 10-dimensional simple abelian variety of Type 2 or 3 and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$. Now $[K : \mathbb{Q}] = 1$ or 5 by the restriction in Table 1.

Denote the minimal polynomial of α over K by p(x) and the minimal polynomial of α over \mathbb{Q} by P(x). If the degree of p(x) is 1, then $\alpha \in U_K$ and so it is reduced to Type 1 of 5-dimensional case. Thus, $4\cos^2\left(\frac{\pi}{11}\right)$ is a lower bound except 1 for this case.

If the degree of p(x) is 2, it can be written as $p(x) = x^2 + ax + b \in \mathcal{O}_K[x](b \in U_K)$ and consider the following 3 cases as in Section 4. Let $\{\sigma_i\}$ be the set of all \mathbb{Q} -embeddings $K \hookrightarrow \mathbb{C}$.

• there exists $1 \le i \le e$ such that $|\sigma_i(b)| > 1$

The maximal absolute value of all conjugates of $b \in U_K$ is at least $2\cos\left(\frac{\pi}{11}\right)$ and so the maximal absolute value of the roots of P(x) is at least $\sqrt{2\cos\left(\frac{\pi}{11}\right)}$. Thus, $2\cos\left(\frac{\pi}{11}\right) = 1.9189\cdots$ is a lower bound except 1 for this case.

• $\sigma_i(b) = 1$ for all $1 \le i \le e$

 $\max\{|\sigma_i(a)|\}\$ must be larger than 2 and close to 2. By using a computer algebra, totally real algebraic integers of degree 5 whose conjugates have modulus less than 2.1 are enumerated in Table A.17. $a = 2.0264\cdots$ provides the minimum of the maximal absolute value. For this case, the maximal absolute value of the roots of P(x) is the Lehmer number $1.1762\cdots$ and so $1.1762\cdots^2 = 1.3836\cdots$ is a lower bound except 1 for this case.

• $\sigma_i(b) = -1$ for all $1 \le i \le e$

a = 1 provides the minimum value of the first dynamical degrees as in Section 4. Thus, $\left(\frac{1+\sqrt{5}}{2}\right)^2$ is a lower bound except 1 for this case.

Thus, $1.1762 \cdots^2 = 1.3836 \cdots$ is a lower bound of the first dynamical degrees except 1 for these types. The realizability of $1.1762 \cdots^2 = 1.3836 \cdots$ as the first dynamical degree is due to [DH22, Theorem 2.1 (i)].

Type 4

Let X be a 10-dimensional simple abelian variety and $f \in \text{End}(X)$ be an automorphism of X which corresponds to $\alpha \in B$.

Now d = 1 and $[K : \mathbb{Q}] = 2$ or 4 or 10 or 20 by the restriction in Table 1 and $\alpha \in U_K$. Now $[K_0 : \mathbb{Q}] = 1$ or 2 or 5 or 10, respectively. Denote the minimal polynomial of α over

 \mathbb{Q} by P(x).

If $\alpha \in K_0$, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{5}\right)$ except 1, by the deduction in Type 1.

If $\alpha \in K \setminus K_0$, by denoting $a = |\alpha|^2 \in U_{K_0}$, as in Section 4, $\mathbb{Q}(\sqrt{a})$ is a totally real number field. By using similar deduction as in the 4-dimensional case, the minimum value of the square of the maximal absolute value of the roots of P(x) is $4\cos^2\left(\frac{\pi}{5}\right)$ except 1.

Therefore, the minimum value of the first dynamical degrees of automorphisms of 10dimensional simple abelian varieties is $1.3836\cdots$ except 1, and realized on a simple abelian variety of Type 2.

APPENDIX A. TABLES

The tables we use in this paper are lined up in this section. On creating from Table A.6 to Table A.17, we use Python as a programming language. The program is available at https://github.com/sugi0000/maximal-absolute-value.

degree of $P_n(x)$	cyclotomic factor of $P_n(x)$	degree of $S_n(x)$
10		10
11	$\Phi_2(x)$	10
12	$\Phi_3(x)$	10
13	$\Phi_2(x), \Phi_8(x)$	8
14	$\Phi_5(x)$	10
15	$\Phi_2(x), \Phi_3(x)$	12
16		16
17	$\Phi_2(x)$	16
18	$\Phi_3(x), \Phi_{12}(x)$	12
19	$\Phi_2(x), \Phi_5(x)$	14
20		20
21	$\Phi_2(x), \Phi_3(x), \Phi_8(x)$	14
22		22
23	$\Phi_2(x)$	22
24	$\Phi_3(x), \Phi_5(x)$	18
25	$\Phi_2(x), \Phi_{18}(x)$	18
26		26
27	$\Phi_2(x), \Phi_3(x)$	24
28		28
29	$\Phi_2(x), \Phi_5(x), \Phi_8(x)$	20
30	$\Phi_3(x), \Phi_{12}(x)$	24
31	$\Phi_2(x)$	30
32		32
33	$\Phi_2(x), \Phi_3(x)$	30

TABLE A.1. The degree of $S_n(x)$ for small n

degree of $Q_m(x)$	cyclotomic factor of $Q_m(x)$	degree of $S'_m(x)$
4		4
5	$\Phi_2(x)$	4
6		6
7	$\Phi_2(x)$	6
8		8
9	$\Phi_2(x)$	8
10	$\Phi_8(x)$	6
11	$\Phi_2(x)$	10
12		12
13	$\Phi_2(x), \Phi_{12}(x)$	8
14		14
15	$\Phi_2(x)$	14
16		16
17	$\Phi_2(x), \Phi_{18}(x)$	10
18	$\Phi_8(x)$	14
19	$\Phi_2(x)$	18
20		20
21	$\Phi_2(x)$	20
22		22
23	$\Phi_2(x)$	22
24	$\Phi_{30}(x)$	16
25	$\Phi_2(x), \Phi_{12}(x)$	20
26	$\Phi_8(x)$	22
27	$\Phi_2(x)$	26
28		28
29	$\Phi_2(x)$	28
30		30
31	$\Phi_2(x)$	30

TABLE A.2. The degree of $S'_m(x)$ for small m

TABLE A.3. The degree of $S_n(x)$

degree of	degree of	degree of	degree of	degree of	degree of	degree of	degree of	degree of	degree of
$P_n(x)$	$S_n(x)$	$P_n(x)$	$S_n(x)$	$P_n(x)$	$S_n(x)$	$P_n(x)$	$S_n(x)$	$P_n(x)$	$S_n(x)$
$\frac{1}{n(x)}$ 11	$\frac{D_n(x)}{10}$	$\frac{1}{n}(x)$ 90	$\frac{D_n(x)}{84}$	168	166	$\frac{1}{247}$	$\frac{D_n(x)}{246}$	$\frac{1}{327}$	$\frac{D_n(x)}{324}$
11	10	91	90	100	168	248	240	328	324
13	8	92	92	171	172	240	240	330	324
15	12	93	86	172	168	251	250	331	324
16	12	95	94	173	164	252	230	332	332
10	10	95	94 94	174	174	255	248	333	326
18	12	90	94 92	175	174 176	255	252	335	334
20	20	100	92 100	170	170	258	250	336	334
20	14	100	96	179	174	259	232	339	332
21 23	22	101	90	180	176	259	248	340	340
23	18	102	102	181	170	260	200	340	336
24 27	24	103	102	183	180	263	262	341	336
27	24 28	104	100	184	180	263	262	343	342
28	28	107	106	180	180	264	258	343	342
30	24	109	100	188	180	268	268	347	346
31	30	111	108	189	178	269	260	348	346
32	32	112	112	191	190	270	264	349	334
35	34	114	104	192	190	271	270	351	348
36	34	115	108	195	192	272	272	352	352
37	32	116	116	196	196	275	274	354	344
39	32	117	110	197	192	276	274	355	354
40	40	119	114	198	192	277	266	356	356
42	36	120	118	199	194	279	272	357	350
43	36	123	120	200	200	280	280	359	354
44	40	124	120	203	202	282	276	360	358
45	38	125	120	204	198	283	282	363	360
47	46	126	120	205	194	284	280	364	360
48	46	127	126	207	204	285	278	365	360
51	48	128	120	208	208	287	286	366	360
52	52	131	130	210	204	288	286	367	360
53	48	132	130	211	210	291	288	368	360
54	44	133	122	212	212	292	292	371	370
55	54	135	132	213	206	293	288	372	370
56	56	136	136	215	214	294	284	373	368
59	54	138	132	216	214	295	288		
60	58	139	134	219	212	296	296		
61	50	140	140	220	220	299	294		
63	60	141	134	221	216	300	298		
64	60	143	142	222	216	301	296		
66	60	144	138	223	216	303	300		
67	66	147	144	224	220	304	300		
68	60	148	148	227	226	306	300		
69	58	149	140	228	226	307	306		
71	70	150	144	229	220	308	300		
72	70	151	144	231	228	309	298		
75	72	152	152	232	232	311	310		
76	76	155	154	234	224	312	310		
77	72	156	154	235	234	315	312		
78	72	157	152	236	236	316	316		
79	68	159	152	237	230	317	312		
80	80	160	160	239	234	318	312		
83	82	162	156	240	238	319	314		
84	78	163	162	243	240	320	320		
85	80	164	160	244	240	323	322		
87	84	165	158	245	240	324	318		
88	88	167	166	246	240	325	320		

TABLE A.4. The degree of $S'_m(x)$

| degree of |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
| $Q_m(x)$ | $S'_m(x)$ |
| 7 | 6 | 95 | 94 | 181 | 176 | 267 | 266 | 355 | 354 |
| 10 | 6 | 97 | 92 | 183 | 182 | 271 | 270 | 359 | 352 |
| 11 | 10 | 98 | 94 | 186 | 182 | 274 | 270 | 361 | 356 |
| 13 | 8 | 99 | 98 | 187 | 186 | 275 | 274 | 362 | 358 |
| 15 | 14 | 103 | 102 | 191 | 190 | 277 | 272 | 363 | 362 |
| 18 | 14 | 106 | 102 | 193 | 188 | 279 | 278 | 367 | 366 |
| 19 | 18 | 107 | 100 | 194 | 190 | 282 | 278 | 370 | 366 |
| 23 | 22 | 109 | 104 | 195 | 194 | 283 | 282 | 371 | 370 |
| 25 | 20 | 111 | 110 | 199 | 198 | 287 | 280 | 373 | 368 |
| 26 | 22 | 114 | 102 | 202 | 198 | 289 | 284 | 375 | 374 |
| 27 | 26 | 115 | 114 | 203 | 202 | 290 | 286 | 378 | 374 |
| 31 | 30 | 119 | 118 | 205 | 200 | 291 | 290 | | |
| 34 | 30 | 121 | 116 | 207 | 206 | 295 | 294 | | |
| 35 | 28 | 122 | 118 | 210 | 206 | 298 | 294 | | |
| 37 | 32 | 123 | 122 | 211 | 210 | 299 | 298 | | |
| 39 | 38 | 127 | 126 | 215 | 208 | 301 | 296 | | |
| 42 | 38 | 130 | 126 | 217 | 212 | 303 | 302 | | |
| 43 | 42 | 131 | 130 | 218 | 214 | 306 | 302 | | |
| 47 | 46 | 133 | 128 | 219 | 218 | 307 | 306 | | |
| 49 | 44 | 135 | 134 | 223 | 222 | 311 | 310 | | |
| 50 | 46 | 138 | 134 | 226 | 222 | 313 | 308 | | |
| 51 | 50 | 139 | 138 | 227 | 226 | 314 | 310 | | |
| 55 | 54 | 143 | 136 | 229 | 224 | 315 | 314 | | |
| 58 | 54 | 145 | 140 | 231 | 230 | 319 | 318 | | |
| 59 | 58 | 146 | 142 | 234 | 222 | 322 | 318 | | |
| 61 | 56 | 147 | 146 | 235 | 234 | 323 | 316 | | |
| 63 | 62 | 151 | 150 | 239 | 238 | 325 | 320 | | |
| 66 | 62 | 154 | 150 | 241 | 236 | 327 | 326 | | |
| 67 | 66 | 155 | 154 | 242 | 238 | 330 | 326 | | |
| 71 | 64 | 157 | 152 | 243 | 242 | 331 | 330 | | |
| 73 | 68 | 159 | 158 | 247 | 246 | 335 | 334 | | |
| 74 | 70 | 162 | 158 | 250 | 246 | 337 | 332 | | |
| 75 | 74 | 163 | 162 | 251 | 244 | 338 | 334 | | |
| 79 | 78 | 167 | 166 | 253 | 248 | 339 | 338 | | |
| 82 | 78 | 169 | 164 | 255 | 254 | 343 | 342 | | |
| 83 | 82 | 170 | 166 | 258 | 254 | 346 | 342 | | |
| 85 | 80 | 171 | 170 | 259 | 258 | 347 | 346 | | |
| 87 | 86 | 175 | 174 | 263 | 262 | 349 | 344 | | |
| 90 | 86 | 178 | 174 | 265 | 260 | 351 | 350 | | |
| 91 | 90 | 179 | 172 | 266 | 262 | 354 | 342 | | |

D	(p,q)	t	$(x^2 + sx + t)(x^2 + \overline{s}x + \overline{t})$	maximal absolute value
7	(-2,1)	1	$x^4 - 3x^3 + 6x^2 - 3x + 1$	$2.1022\cdots$
7	(-2,1)	-1	$x^4 - 3x^3 + 2x^2 + 3x + 1$	$2.1889\cdots$
7	(-1,1)	1	$x^4 - x^3 + 4x^2 - x + 1$	$1.8832\cdots$
7	(-1,1)	-1	$x^4 - x^3 + x + 1$	$1.3722\cdots$
7	(0,1)	1	$x^4 + x^3 + 4x^2 + x + 1$	$1.8832\cdots$
7	(0, 1)	-1	$x^4 + x^3 - x + 1$	$1.3722\cdots$
7	(1, 1)	1	$x^4 + 3x^3 + 6x^2 + 3x + 1$	$2.1022\cdots$
7	(1, 1)	-1	$x^4 + 3x^3 + 2x^2 - 3x + 1$	$2.1889\cdots$
11	(-1,1)	1	$x^4 - x^3 + 5x^2 - x + 1$	$2.1537\cdots$
11	(-1,1)	-1	$x^4 - x^3 + x^2 + x + 1$	$1.4675\cdots$
11	(0, 1)	1	$x^4 + x^3 + 5x^2 + x + 1$	$2.1537\cdots$
11	(0, 1)	-1	$x^4 + x^3 + x^2 - x + 1$	$1.4675\cdots$
15	(-1,1)	1	$x^4 - x^3 + 6x^2 - x + 1$	$2.3869\cdots$
15	(-1,1)	-1	$x^4 - x^3 + 2x^2 + x + 1$	$1.6180\cdots$
15	(0, 1)	1	$x^4 + x^3 + 6x^2 + x + 1$	$2.3869\cdots$
15	(0, 1)	-1	$x^4 + x^3 + 2x^2 - x + 1$	$1.6180\cdots$
2	(-1,1)	1	$x^4 - 2x^3 + 5x^2 - 2x + 1$	$2.0322\cdots$
2	(-1,1)	-1	$x^4 - 2x^3 + x^2 + 2x + 1$	$1.8039\cdots$
2	(0, 1)	1	$x^4 + 4x^2 + 1$	$1.9318\cdots$
2	(0, 1)	-1	$x^4 + 1$	1
2	(1, 1)	1	$x^4 + 2x^3 + 5x^2 + 2x + 1$	$2.0322\cdots$
2	(-1,1)	-1	$x^4 + 2x^3 + x^2 - 2x + 1$	$1.8039\cdots$
3	(1, 1)	1	$x^4 + 3x^3 + 5x^2 + 3x + 1$	$1.7220\cdots$
3	(1, 1)	ω^{*1}	$x^4 + 3x^3 + 4x^2 + 3x + 1$	1
3	(1, 1)	ω^2	$x^4 + 3x^3 + 2x^2 + 1$	$1.7220\cdots$
3	(1, 1)	$\omega^3 = -1$	$x^4 + 3x^3 + x^2 - 3x + 1$	$2.0758\cdots$
3	(1, 1)	ω^4	$x^4 + 3x^3 + 2x^2 - 3x + 1$	$2.1889\cdots$
3	(1, 1)	ω^5	$x^4 + 3x^3 + 4x^2 + 1$	$2.0758\cdots$
3	(1, 0)	ω	$x^4 + 2x^3 + 2x^2 + x + 1$	$1.3122\cdots$
3	(1, 0)	ω^2	$x^4 + 2x^3 - x + 1$	$1.5392\cdots$
3	(1, 0)	ω^4	$x^4 + 2x^3 - x + 1$	$1.5392\cdots$
3	(1, 0)	ω^5	$x^4 + 2x^3 + 2x^2 + x + 1$	$1.3122\cdots$
3	(2, 0)	ω	$x^4 + 4x^3 + 5x^2 + 2x + 1$	$1.9318\cdots$
3	(2, 0)	ω^2	$x^4 + 4x^3 + 3x^2 - 2x + 1$	$2.2966\cdots$
3	(2, 0)	ω^4	$x^4 + 4x^3 + 3x^2 - 2x + 1$	$2.2966\cdots$
3	(2,0)	ω^5	$x^4 + 4x^3 + 5x^2 + 2x + 1$	$1.9318\cdots$
1	(1, 1)	1	$x^4 + 2x^3 + 4x^2 + 2x + 1$	$1.7000\cdots$
1	(1, 1)	i	$x^4 + 2x^3 + 2x^2 + 2x + 1$	1
1	(1, 1)	-1	$x^4 + 2x^3 - 2x + 1$	$1.7000\cdots$
1	(1, 1)	-i	$x^4 + 2x^3 + 2x^2 - 2x + 1$	$1.9318\cdots$
1	(1, 0)	i	$x^4 + 2x^3 + x^2 + 1$	$1.4425\cdots$
1	(1, 0)	-i	$x^4 + 2x^3 + x^2 + 1$	$1.4425\cdots$
1	(2,0)	i	$x^4 + 4x^3 + 4x^2 + 1$	$2.1474\cdots$
1	(2,0)	-i	$x^4 + 4x^3 + 4x^2 + 1$	$2.1474\cdots$

TABLE A.5. Calculation of the maximal absolute value of the roots for the case of Type 4 of 4-dimensional

*1 Denote $\omega = \frac{1+\sqrt{3}i}{2}$

whose conjugates have modulus < 2.25 algebraic integer polynomial

TABLE A.6. totally real, cubic algebraic integer, which is larger than 2,

algebraic integer	polynomial
$\boxed{2.1149\cdots}$	$x^3 - 4x - 1$
$2.1700\cdots$	$x^3 - x^2 - 3x + 1$
$2.1986\cdots$	$x^3 - x^2 - 4x + 3$
$2.2143\cdots$	$x^3 - 4x - 2$
$2.2469\cdots$	$x^3 - 2x^2 - x + 1$

TABLE A.7. totally positive, septic algebraic unit, which is larger than 4, whose conjugates are in (0, 4.04)

algebraic unit	polynomial
$4.0333\cdots$	$x^{7} - 14x^{6} + 77x^{5} - 211x^{4} + 301x^{3} - 210x^{2} + 56x - 1$
$4.0341\cdots$	$x^7 - 14x^6 + 76x^5 - 200x^4 + 259x^3 - 146x^2 + 24x - 1$

TABLE A.8. totally real, quartic algebraic integer, which is larger than 2, whose conjugates have modulus < 2.1

algebraic integer	minimal polynomial
$2.0614\cdots$	$x^4 - 4x^2 - x + 1$
$2.0743\cdots$	$x^4 - 5x^2 + 3$
$2.0952\cdots$	$x^4 - x^3 - 3x^2 + x + 1$

TABLE A.9. irreducible quartic polynomial, all roots are in E, one root is not inside [-2, 2], identifying R(X) with R(-X)

condition (6.1)	polynomial $R(X)$	polynomial $x^4 R(x + \frac{1}{x})$	maximal absolute value
N	$X^4 - 2X^3 - 2X^2 + 5X - 1$	$ x^8 - 2x^7 + 2x^6 - x^5 + x^4 -x^3 + 2x^2 - 2x + 1 $	$1.2150\cdots$
N	$X^4 - X^3 - 4X^2 + 2X + 5$	$x^8 - x^7 - x^5 + 3x^4 - x^3 - x + 1$	$1.1837\cdots$
N	$X^4 - X^3 - 4X^2 + 4X + 2$	$x^8 - x^7 + x^5 + x^3 - x + 1$	$1.2744\cdots$
N	$X^4 - X^3 - 3X^2 + X + 3$	$ \begin{array}{c} x^8 - x^7 + x^6 - 2x^5 + 3x^4 \\ -2x^3 + x^2 - x + 1 \end{array} $	$1.2522\cdots$
N	$X^4 - X^3 - 3X^2 + 3X - 1$	$x^8 - x^7 + x^6 - x^4 + x^2 - x + 1$	$1.1705\cdots$
N	$X^4 - X^3 - 3X^2 + 3X + 1$	$x^8 - x^7 + x^6 + x^4 + x^2 - x + 1$	$1.2196\cdots$
N	$X^4 - X^3 - 2X^2 + 1$	$ \begin{array}{c} x^8 - x^7 + 2x^6 - 3x^5 + 3x^4 \\ -3x^3 + 2x^2 - x + 1 \end{array} $	$1.2408\cdots$
N	$X^4 - X^3 - 2X^2 + X + 2$	$\begin{array}{c} x^8 - x^7 + 2x^6 - 2x^5 + 4x^4 \\ -2x^3 + 2x^2 - x + 1 \end{array}$	$1.2474\cdots$
N	$X^4 - X^3 - 2X^2 + 2X - 1$	$\begin{array}{c} x^8 - x^7 + 2x^6 - x^5 + x^4 \\ - x^3 + 2x^2 - x + 1 \end{array}$	$1.2722\cdots$
N	$X^4 - 5X^2 + 7$	$x^8 - x^6 + 3x^4 - x^2 + 1$	$1.2406\cdots$
Ν	$X^4 - 4X^2 - X + 3$	$x^8 - x^5 + x^4 - x^3 + 1$	$1.1692\cdots$
Y	$X^4 - 4X^2 - 1$	$x^8 - 3x^4 + 1$	$1.2720\cdots$
N	$X^4 - 4X^2 + 5$	$x^8 + 3x^4 + 1$	$1.2720\cdots$
N	$X^4 - 3X^2 + 3$	$x^8 + x^6 + 3x^4 + x^2 + 1$	$1.2406\cdots$

condition (6.1)	polynomial $R(X)$	polynomial $x^4 R(x - \frac{1}{x})$	maximal absolute value
N	$X^4 - X^3 + 3X^2 - 2X + 1$	$\begin{array}{c} x^8 - x^7 - x^6 + x^5 + x^4 \\ -x^3 - x^2 + x + 1 \end{array}$	$1.2245\cdots$
N	$X^4 - X^3 + 4X^2 - 3X + 1$	$x^8 - x^7 - x^4 + x + 1$	$1.2306\cdots$
N	$X^4 - X^3 + 4X^2 - 3X + 2$	$x^8 - x^7 + x + 1$	$1.2612\cdots$
N	$X^4 - X^3 + 4X^2 - 2X + 2$	$x^8 - x^7 + x^5 - x^3 + x + 1$	$1.2397\cdots$
N	$X^4 - X^3 + 4X^2 - 2X + 3$	$x^8 - x^7 + x^5 + x^4 - x^3 + x + 1$	$1.1837\cdots$
Ν	$X^4 - X^3 + 5X^2 - 4X + 3$	$\begin{array}{c} x^8 - x^7 + x^6 - x^5 + x^4 \\ + x^3 + x^2 + x + 1 \end{array}$	$1.2788\cdots$
N	$X^4 - X^3 + 5X^2 - 3X + 4$	$x^8 - x^7 + x^6 + x^2 + x + 1$	$1.2272\cdots$
N	$X^4 - X^3 + 5X^2 - 3X + 5$	$x^8 - x^7 + x^6 + x^4 + x^2 + x + 1$	$1.2734\cdots$
N	$X^4 - X^3 + 6X^2 - 3X + 8$	$x^8 - x^7 + 2x^6 + 2x^4 + 2x^2 + x + 1$	$1.2474\cdots$
N	$X^4 + 2X^2 - X + 1$	$x^8 - 2x^6 - x^5 + 3x^4 + x^3 - 2x^2 + 1$	$1.2553\cdots$
N	$X^4 + 3X^2 - X + 1$	$x^8 - x^6 - x^5 + x^4 + x^3 - x^2 + 1$	$1.1932\cdots$
N	$X^4 + 3X^2 - X + 2$	$x^8 - x^6 - x^5 + 2x^4 + x^3 - x^2 + 1$	$1.2461\cdots$
N	$X^4 + 3X^2 + 3$	$x^8 - x^6 + 3x^4 - x^2 + 1$	$1.2406\cdots$
N	$X^4 + 4X^2 - X + 1$	$x^8 - x^5 - x^4 + x^3 + 1$	$1.2512\cdots$
N	$X^4 + 4X^2 - X + 2$	$x^8 - x^5 + x^3 + 1$	$1.2331\cdots$
N	$X^4 + 4X^2 - X + 3$	$x^8 - x^5 + x^4 + x^3 + 1$	$1.2450\cdots$
Y	$X^4 + 4X^2 - 1$	$x^8 - 3x^4 + 1$	$1.2720\cdots$
N	$X^4 + 4X^2 + 5$	$x^8 + 3x^4 + 1$	$1.2720\cdots$
Ν	$X^4 + 5X^2 + 7$	$x^8 + x^6 + 3x^4 + x^2 + 1$	$1.2406\cdots$

TABLE A.10. irreducible quartic polynomial, all roots are in E_1 , one root is not written as si with $s \in [-2, 2]$, identifying R(X) with R(-X)

TABLE A.11. quadratic polynomial of the form (6.2), all roots are in E_2 , identifying $R_1(z)$ with $R_1(-z)$

polynomial $R_1(z)$	polynomial $x^2 R_1(x + \frac{i}{x})$	maximal
polynomial $n_1(z)$	polynomial $x R_1(x + \frac{1}{x})$	absolute value
$z^2 - (2+2i)z + 2i$	$x^4 - (2+2i)x^3 + 4ix^2 + (2-2i)x - 1$	1
$z^2 - (1+i)z - i$	$x^4 - (1+i)x^3 + ix^2 + (1-i)x - 1$	1
$z^2 + (-1 - i)z$	$x^4 - (1+i)x^3 + 2ix^2 + (1-i)x - 1$	1
$z^2 - 1 - 2i$	$x^4 - x^2 - 1$	$1.2720\cdots$
$z^2 - 4i$	$x^4 - 2ix^2 - 1$	1
$z^2 - 3i$	$x^4 - ix^2 - 1$	1
$z^2 - 2i$	$x^4 - 1$	1
z^2-i	$x^4 + ix^2 - 1$	1
z^2	$x^4 + 2ix^2 - 1$	1
$z^2 + 1 - 2i$	$x^4 + x^2 - 1$	$1.2720\cdots$

TABLE A.12. quadratic polynomial of the form (6.3), with the roots $s, t \in \mathbb{Q}(\zeta_8)$ satisfy $s \in E_3$, $t \in E_4$ and $(x^2 + sx + \zeta_8)(x^2 + tx + \zeta_8^3) \in \mathbb{Z}[\sqrt{2}i][x]$, identifying $R_2(z)$ with $R_2(-z)$

polynomial $R_2(z)$	polynomial $(x^2 + sx + \zeta_8)(x^2 + tx + \zeta_8^3)$	maximal absolute value
$z^2 - (2 + \sqrt{2}i)z + \sqrt{2}i$	$x^4 + (2 + \sqrt{2}i)x^3 + 2\sqrt{2}ix^2 + (-2 + \sqrt{2}i)x - 1$	1
$z^2 - \sqrt{2}iz - \sqrt{2}i$	$x^4 + \sqrt{2}ix^3 + \sqrt{2}ix - 1$	1
z^2	$x^4 + \sqrt{2}ix^2 - 1$	1

TABLE A.13. quadratic polynomial of the form (6.4), with the roots $s, t \in \mathbb{Q}(\zeta_{12})$ satisfy $s \in E_5$ and $t \in E_6$ and $(x^2 + sx + \zeta_{12})(x^2 + tx + \zeta_{12}^5) \in \mathbb{Z}[i][x]$, identifying $R_3(z)$ with $R_3(-z)$

polynomial $R_3(z)$	polynomial $(x^2 + sx + \zeta_{12})(x^2 + tx + \zeta_{12}^5)$	maximal absolute value
$\boxed{z^2 - (2+i)z + i}$	$x^4 + (2+i)x^3 + 2ix^2 - (2-i)x - 1$	1
$z^2 - (1+2i)z + i$	$x^4 + (1+2i)x^3 + 2ix^2 - (1-2i)x - 1$	1
$z^2 - (1-i)z - 2i$	$x^4 + (1-i)x^3 - ix^2 - (1+i)x - 1$	1
z^2	$x^4 + ix^2 - 1$	1

TABLE A.14. cubic polynomial with constant term 1, with coefficients in \mathbb{Z} , whose roots have modulus less than 1.3, at least one root has modulus > 1

polynomial $p(x)$	maximal absolute value
$x^3 - x^2 + 1$	$1.1509\cdots$
$x^3 + x + 1$	$1.2106\cdots$

TABLE A.15. cubic polynomial with constant term 1, with coefficients in $\mathcal{O}_{\mathbb{Q}(\sqrt{-D})}$, whose roots have modulus less than 1.3, at least one root has modulus> 1, identifying p(x) with $\overline{p(x)}$

polynomial $p(x)$	maximal absolute value
$x^{3} + \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)x^{2} + 1$	$1.1509\cdots$
$x^3 - \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)x + 1$	$1.2106\cdots$
$x^3 + x^2 + (1 - i)x + 1$	$1.2328\cdots$
$x^3 - ix^2 - x + 1$	$1.2878\cdots$
$x^3 - ix + 1$	$1.2966\cdots$
$x^3 - (1+i)x^2 + ix + 1$	$1.2969\cdots$

TABLE A.16. cubic polynomial with constant term $\frac{1}{2} + \frac{\sqrt{3}}{2}i$, with coefficients in $\mathcal{O}_{\mathbb{Q}(\sqrt{-3})}$, whose roots have modulus less than 1.3, at least one root has modulus> 1

polynomial $p(x)$	maximal absolute value
$x^{3} - \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)x^{2} + \left(-\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2167\cdots$
$x^{3} - \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)x^{2} + \left(-\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2167\cdots$
$x^{3} + x^{2} + x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2167\cdots$
$x^3 - x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2746\cdots$
$x^{3} + \left(\frac{1}{2} - \frac{\sqrt{3}}{2}i\right)x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2746\cdots$
$x^{3} + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)x + \left(\frac{1}{2} + \frac{\sqrt{3}}{2}i\right)$	$1.2746\cdots$

TABLE A.17. totally real, quintic algebraic integer, which is larger than 2, whose conjugates have modulus < 2.1

algebraic integer	minimal polynomial
$2.0264\cdots$	$x^5 + x^4 - 5x^3 - 5x^2 + 4x + 3$
$2.0384\cdots$	$x^5 - 5x^3 + 4x - 1$
$2.0431\cdots$	$x^5 - 5x^3 - x^2 + 5x + 1$
$2.0541\cdots$	$x^5 - 6x^3 + 8x - 1$
$2.0665\cdots$	$x^5 - 6x^3 - x^2 + 8x + 3$
$2.0850\cdots$	$x^{5} - x^{4} - 5x^{3} + 4x^{2} + 5x - 3$
$2.0911\cdots$	$x^5 - x^4 - 5x^3 + 4x^2 + 4x - 1$

Appendix B. The proof of the fact in Remark 5.13

In Remark 5.13, we presented that for 0 < t < 2, the equation

$$\frac{2\sin(x)\cos(2x)}{1+2\sin(x)\sin(2x)} = \tan(tx)$$

has at most one root in the interval $(0, \frac{\pi}{4})$. We prove this fact in this section. Denote

$$f(x) = 1 + 2\sin(x)\sin(2x), \ g(x) = 2\sin(x)\cos(2x).$$

Now

$$\frac{g(x)}{f(x)} > 0$$
 and $\tan(tx) > 0$

for $0 < x < \frac{\pi}{4}$. In the *xy* coordinate plane, the curve $C : y = \frac{g(x)}{f(x)}$ and $D_t : y = \tan(tx)$ both through (x, y) = (0, 0). Assume *C* and D_t intersect at least one point for $0 < x < \frac{\pi}{4}$. Let $(x, y) = (x_1, y_1)$ be an intersection point on the very right of *C* and D_t with $0 < x < \frac{\pi}{4}$. Denote the line passing through (0, 0) and (x_1, y_1) by *L*. First,

$$\left(\frac{d}{dx}\right)^2 (\tan(tx)) = \frac{2t^2 \sin(tx) \cos(tx)}{\cos^4(tx)} > 0$$

for $0 < x < \frac{\pi}{4}$ and so $\tan(tx)$ is a convex downward function on the interval $(0, \frac{\pi}{4})$. Thus, L is over D_t for $0 < x < x_1$ and L is under D_t for $x_1 < x < \frac{\pi}{4}$. Next we show that Lintersects with C at only (x_1, y_1) transversally. By some calculation,

$$\begin{pmatrix} \frac{d}{dx} \end{pmatrix} (g'(x)f(x) - f'(x)g(x)) = g''(x)f(x) - f''(x)g(x)$$

= -10sin(x)cos(2x) - 8cos(x)sin(2x)
- 16sin(x)cos(x)sin^2(2x) - 16sin(x)cos(x)cos^2(2x)
< 0

for $0 < x < \frac{\pi}{4}$ and so

g'(x)f(x) - f'(x)g(x)

is strictly decreasing on the interval $(0, \frac{\pi}{4})$. Now by

$$f'(x) = 2\cos(x)\sin(2x) + 4\sin(x)\cos(2x) > 0, \ f(x) > 0$$

for $0 < x < \frac{\pi}{4}$, f(x) is strictly increasing on the interval $(0, \frac{\pi}{4})$ and so

$$\left(\frac{d}{dx}\right)\left(\frac{g(x)}{f(x)}\right) = \frac{g'(x)f(x) - f'(x)g(x)}{f(x)^2}$$

is strictly decreasing on $\left\{x \in \left(0, \frac{\pi}{4}\right) \mid g'(x)f(x) - f'(x)g(x) > 0\right\}$. Thus, by $\frac{y_1}{x_1} > 0$, there exists at most one root of the equation

$$\left(\frac{d}{dx}\right)\left(\frac{g(x)}{f(x)}\right) = \frac{y_1}{x_1}$$

for $0 < x < \frac{\pi}{4}$. By the mean value theorem, L and C intersects at only $(x, y) = (x_1, y_1)$ for $0 < x < \frac{\pi}{4}$ and it is transversally. Also by g'(0)f(0) - f'(0)g(0) > 0,

$$\left(\frac{d^2}{dx^2}\right)\left(\frac{g(x)}{f(x)}\right) < 0,$$

on $(0, \epsilon)$ for some $\epsilon > 0$ and so $C : y = \frac{g(x)}{f(x)}$ is a convex upward function on $(0, \epsilon)$. Thus, C is over L for $0 < x < x_1$ and C is under L for $x_1 < x < \frac{\pi}{4}$.

Therefore, C and D_t intersects at only (x_1, y_1) and so the proof is concluded.

References

- [ACR16] Adigaa, C., Cangul, I.N., Ramaswamy, H.N.: On the constant term of the minimal polynomial of $\cos \frac{2\pi}{n}$ over \mathbb{Q} , Filomat, **30**(2016), No. 4, 1097-1102.
- [BDGPS92] Bertin, M.J., Decomps-Guilloux, A., Grandet-Hugot, M., Pathiaux-Delefosse, M., Schreiber, J.P.: Pisot and Salem numbers, Birkhäuser Verlag, Basel, 1992.
- [BL04] Birkenhake, C., Lange, H.: Complex abelian varieties, second ed., Grundlehren der mathematischen Wissenschaften, **302**, Springer-Verlag, Berlin, 2004.
- [DH22] Dang, N.-B., Herrig, T.: Dynamical degrees of automorphisms on abelian varieties, Advances in Mathematics, **395**(2022), 108082.
- [DS05] Dinh, T.-C., Sibony, N.: Une borne supérieure pour l'entropie topologique d'une application rationnelle, Ann. of Math., 161(2005), 1637–1644.
- [DS17] Dinh, T.-C., Sibony, N. : Equidistribution problems in complex dynamics of higher dimension, International Journal of Mathematics, 28(2017), No. 07, 1750057.
- [GHM09] Gross, B.H., Hironaka, E., McMullen, C.T.: Cyclotomic factors of Coxeter polynomials, Journal of Number Theory, **129**(2009), 1034-1043.

- [Lan83] Lang, S. : Complex Multiplication, Grundlehren der mathematischen Wissenschaften, 255, Springer-Verlag, New York, 1983.
- [Mum70] Mumford, D. : *Abelian Varieties*, Tata institute of Fundamental Research studies in Mathematics, **5**, Oxford University Press, 1970.
- [Neu99] Neukirch, J. : *Algebraic number theory*, Grundlehren der mathematischen Wissenschaften, **322**, Springer-Verlag, Berlin, 1999.
- [Pie82] Pierce, R.S.: Associative Algebras, Graduate Texts in Mathematics, 88, Springer, New York, 1982.
- [Sho09] Shoup, V. : A Computational Introduction to Number Theory and Algebra, Second edition, Cambridge University Press, 2009.
- [Sug23] Sugimoto, Y.: Dynamical degrees of automorphisms of complex simple abelian varieties and Salem numbers, 2023., arXiv:2302.02271.
- [SZ65] Schinzel, A., Zassenhaus, H. : A refinement of two theorems of Kronecker, Michigan Math. J., 12(1965), 81-85.
- [Voi21] Voight, J.: Quaternion Algebras, Graduate Texts in Mathematics, 288, Springer, 2021.