### ON THE HILBERT DEPTH OF MONOMIAL IDEALS

SILVIU BĂLĂNESCU<sup>1</sup>, MIRCEA CIMPOEAȘ<sup>2</sup> AND CHRISTIAN KRATTENTHALER<sup>3</sup>

ABSTRACT. Let  $S = K[x_1, \ldots, x_n]$  be the ring of polynomials over a field K. Given two monomial ideals  $0 \subset I \subsetneq J \subset S$ , we present a new method to compute the Hilbert depth of J/I. As an application, we show that if  $u \in S$  is a monomial regular of S/I, then hdepth $(S/I) \ge$  hdepth $(S/(I, u)) \ge$  hdepth(S/I) - 1.

Also, we reprove the formula of the Hilbert depth of a squarefree Veronese ideal.

#### 1. INTRODUCTION

Let K be a field and  $S = K[x_1, \ldots, x_n]$  the polynomial ring over K. Let M be a  $\mathbb{Z}^n$ -graded S-module. A Stanley decomposition of M is a direct sum

$$\mathcal{D}: M = \bigoplus_{i=1}^r m_i K[Z_i],$$

as a  $\mathbb{Z}^n$ -graded K-vector space, where  $m_i \in M$  is homogeneous with respect to  $\mathbb{Z}^n$ -grading,  $Z_i \subset \{x_1, \ldots, x_n\}$  such that  $m_i K[Z_i] = \{um_i : u \in K[Z_i]\} \subset M$  is a free  $K[Z_i]$ -submodule of M. We define sdepth( $\mathcal{D}$ ) = min\_{i=1,...,r} |Z\_i| and

 $sdepth(M) = \max\{sdepth(\mathcal{D}) : \mathcal{D} \text{ is a Stanley decomposition of } M\}.$ 

The number sdepth(M) is called the *Stanley depth* of M.

Herzog, Vlădoiu and Zheng showed in [11] that sdepth(M) can be computed in a finite number of steps if M = I/J, where  $J \subset I \subset S$  are monomial ideals. In [20], Rinaldo gave a computer implementation of this algorithm, in the computer algebra system CoCoA (cf. [7]).

In [1], Apel restated a conjecture firstly given by Stanley in [24], namely that

 $sdepth(M) \ge depth(M),$ 

for any  $\mathbb{Z}^n$ -graded S-module M.

<sup>&</sup>lt;sup>1</sup>Silviu Bălănescu, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, E-mail: silviu.balanescu@stud.fsa.upb.ro

<sup>&</sup>lt;sup>2</sup>Mircea Cimpoeaş, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, Romania and Simion Stoilow Institute of Mathematics, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania, E-mail: mircea.cimpoeas@upb.ro, mircea.cimpoeas@imar.ro

<sup>&</sup>lt;sup>3</sup>Christian Krattenthaler, Fakultät für Mathematik, Universität Wien Oskar-Morgenstern-Platz 1 A-1090 Vienna, Austria, E-mail: Christian.Krattenthaler@univie.ac.at

<sup>2020</sup> Mathematics Subject Classification. 05A18, 06A07, 13C15, 13P10, 13F20.

Key words and phrases. Stanley depth, Hilbert depth, depth, monomial ideal, squarefree Veronese.

## SILVIU BĂLĂNESCU<sup>1</sup>, MIRCEA CIMPOEAȘ<sup>2</sup> AND CHRISTIAN KRATTENTHALER<sup>3</sup>

This conjecture proves to be false, in general, for M = S/I and M = J/I, where  $0 \neq I \subsetneq J \subset S$  are monomial ideals; see [8], but remains open for M = I.

Now, we consider the ring of polynomial S, with the standard grading. Let M be a finitely generated graded S-module. In [25], Uliczka introduced a new invariant associated to M, called Hilbert depth. More precisely, the Hilbert depth of M, denoted by hdepth(M), is the maximal depth of a finitely generated graded S-module N with the same Hilbert series as M. He also proved that

hdepth(M) = max{
$$r : (1-t)^r H_M(t)$$
 is non-negative }.

A subsequent number of papers studied several properties of this invariant. In [3] it was noticed that

$$sdepth(M) \leq hdepth(M),$$

for any multigraded S-module M. For a friendly introduction into the thematic of Stanley depth and Hilbert depth, we refer the reader to [12].

Given two squarefree monomial ideals  $0 \subset I \subsetneq J \subset S$ , for all  $0 \leq j \leq n$ , we let  $\alpha_j(J/I) :=$  the number of squarefree monomials  $u \in S$  such that  $u \in J \setminus I$  and  $\deg(u) = j$ . Also, let

$$\beta_k^q(J/I) = \sum_{j=0}^k (-1)^{k-j} \binom{q-j}{k-j} \alpha_j(J/I) \text{ for all } 0 \le k \le q \le n.$$

In Teorem 2.4 we prove that

 $\mathbf{2}$ 

$$hdepth(J/I) = \max\{q : \beta_k^q(J/I) \ge 0 \text{ for all } 0 \le k \le q\}.$$

In particular, in Proposition 2.8, we reprove the fact that  $hdepth(J/I) \ge sdepth(J/I)$  for any monomial ideals  $0 \subset I \subsetneq J \subset S$ , not necessarily squarefree.

In Theorem 2.22 we show that, if I is a monomial ideal and  $u \in S$  is a monomial regular of S/I, then

$$hdepth(S/I) \ge hdepth(S/(I, u)) \ge hdepth(S/I) - 1,$$

and, moreover, these inequalities are sharp. However, if I is a monomial complete intersection ideal, minimally generated by m monomials, then hdepth(S/I) = sdepth(S/I) = depth(S/I) = n - m. In Remark 2.29 we note that, in this case, we may have hdepth $(I) > \text{sdepth}(I) = n - \left\lfloor \frac{m}{2} \right\rfloor$ .

In Section 3, as an application of our new method for computing the Hilbert depth, we tackle the case of  $J_{n,m}$ , the squarefree Veronese ideal of degree m, i.e., the monomial ideal generated by all the squarefree monomials of degree m in S. In Proposition 3.4 we note that hdepth $(S/J_{n,m}) = m - 1$ . Ge et al. proved in [10] that

hdepth
$$(J_{n,m}) = m + \left\lfloor \frac{n-m}{m+1} \right\rfloor$$
.

We give an alternative proof of this result, which makes use of a transformation formula for hypergeometric series; see Theorem 3.10.

#### 2. Main results

First, we fix some notations,

We denote  $[n] := \{1, 2, ..., n\}$ . For a subset  $C \subset [n]$ , we denote  $x_C := \prod_{j \in C} x_j$ . For two subsets  $C \subset D \subset [n]$ , we denote  $[C, D] := \{A \subset [n] : C \subset A \subset D\}$ , and we call it the *interval* bounded by C and D. Let  $I \subset J \subset S$  be two square-free monomial ideals. We let

$$\mathbf{P}_{J/I} := \{ C \subset [n] : x_C \in J \setminus I \} \subset 2^{[n]}.$$

A partition of  $P_{J/I}$  is a decomposition

$$\mathcal{P}: \mathbf{P}_{J/I} = \bigcup_{i=1}^{r} [C_i, D_i],$$

into disjoint intervals. If  $\mathcal{P}$  is a partition of  $P_{J/I}$ , we let  $\operatorname{sdepth}(\mathcal{P}) := \min_{i=1,\ldots,r} |D_i|$ . The Stanley depth of  $P_{J/I}$  is

 $sdepth(P_{J/I}) := \max\{sdepth(\mathcal{P}) : \mathcal{P} \text{ is a partition of } P_{J/I}\}.$ 

Herzog, Vlădoiu and Zheng proved in [11] that

$$\operatorname{sdepth}(J/I) = \operatorname{sdepth}(\mathcal{P}_{J/I}).$$

Let  $\mathbf{P} := \mathbf{P}_{J/I}$ , where  $I \subset J \subset S$  are square-free monomial ideals. For any k with  $0 \le k \le n$ , we denote

$$P_k := \{A \in P : |A| = k\} \text{ and } \alpha_k(J/I) = \alpha_k(P) = |P_k|.$$

For any q and k with  $0 \le k \le q \le n$ , we consider:

$$\beta_k^q(J/I) = \sum_{j=0}^k (-1)^{k-j} \binom{q-j}{k-j} \alpha_j(J/I).$$
(2.1)

Using the inverse relation [21, Equation (7) on p. 50 with q = 0], from (2.1) we get

$$\alpha_k(J/I) = \sum_{j=0}^k \binom{q-j}{k-j} \beta_j^q(J/I), \text{ for } 0 \le k \le q.$$

$$(2.2)$$

Let  $\mathcal{P}$ :  $P_{J/I} = \bigcup_{i=1}^{r} [C_i, D_i]$  be a Stanley decomposition of  $P_{J/I}$  with  $q := \text{sdepth}(\mathcal{P}) = \text{sdepth}(J/I)$ . By refining  $\mathcal{P}$ , we can assume that if  $|C_i| < d$  then  $|D_i| = q$ ; see [20, Lemma 3.4]. With this assumption it is easy to see that

$$\beta_k^q(J/I) = |\{i : |C_i| = k\}| \ge 0 \text{ for all } 0 \le k \le q.$$
(2.3)

**Lemma 2.1.** With the above notations, for every  $0 \le r \le n$ , we have that:

$$(1-t)^{r}H_{J/I}(t) = \beta_{0}^{r}(J/I) + \beta_{1}^{r}(J/I)t + \dots + \beta_{r}^{r}(J/I)t^{r} + \alpha_{r+1}(J/I)\frac{t^{r+1}}{1-t} + \dots + \alpha_{n}(J/I)\frac{t^{n-r}}{(1-t)^{n-r}}$$

*Proof.* First, let us consider the case J = S. Let  $\Delta$  be the Stanley Reisner simplicial complex associated to I and assume that  $\dim(\Delta) = d-1$ . We denote  $f = (f_{-1}, f_0, \ldots, f_{d-1})$  the f-vector of  $\Delta$ . It is easy to see that

$$\alpha_j(S/I) = \begin{cases} f_{j-1}(\Delta), & 0 \le j \le d\\ 0, & d+1 \le j \le n \end{cases}$$

It follows that

$$H_{S/I}(t) = \sum_{j=0}^{d} f_{j-1}(\Delta)t^{j}(1-t)^{-j} = \sum_{j=0}^{n} \alpha_{j}(S/I)t^{j}(1-t)^{-j}.$$

Multiplying the above identity with  $(1-t)^r$  and using (2.1) we get the required conclusion.

The general case follows from the case J = S and the obvious facts:

$$H_{J/I}(t) = H_{S/I}(t) - H_{S/J}(t), \ \alpha_j(J/I) = \alpha_j(S/I) - \alpha_j(S/J), \text{ for all } 0 \le j \le n,$$

and, therefore,  $\beta_k^r(J/I) = \beta_k^r(S/I) - \beta_k^r(S/J)$  for all  $0 \le k \le r \le n$ .

We recall the definition of the Hilbert depth of a module.

**Definition 2.2.** Let M be a finitely generated graded S-module. The *Hilbert depth* of M is the number

$$hdepth(M) = \max \left\{ r : \begin{array}{ll} \text{There exists a f.g. graded } S\text{-module } N \\ \text{with } H_M(t) = H_N(t) \text{ and } depth(N) = r \end{array} \right\}.$$

Also, we recall the following result:

**Theorem 2.3.** ([25, Theorem 3.2]) Let M be a finitely generated graded S-module. Then hdepth $(M) = \max\{r : (1-t)^q H_M(t) \text{ is non-negative }\}.$ 

Now, we can prove our first main result:

**Theorem 2.4.** Let  $I \subset J \subset S$  be two square-free monomial ideals. Then:

$$hdepth(J/I) := \max\{q : \beta_k^q(J/I) \ge 0 \text{ for } 0 \le k \le q\}.$$

*Proof.* It follows from Lemma 2.1 and Theorem 2.3.

A simple, but very useful lemma is the following:

**Lemma 2.5.** Let  $0 \subset I \subsetneq J \subset S$  be two square-free monomial ideals. Then:

- (1) hdepth $(J/I) \le \max\{k : \alpha_k(J/I) > 0\}.$
- (2) hdepth $(J/I) \ge \min\{k : \alpha_k(J/I) > 0\}.$

*Proof.* (1) Let  $m := \max\{k : \alpha_k(J/I) > 0\}$ . If m = n then there is nothing to prove. Suppose m < n. From (2.2), we have

$$0 = \alpha_{m+1}(J/I) = \sum_{j=0}^{m+1} \beta_j^{m+1}(J/I).$$
(2.4)

If hdepth $(J/I) \ge m + 1$ , from (2.4) it follows that  $\beta_j^{m+1}(J/I) = 0$  for all j with  $0 \le j \le m + 1$ . Therefore I = J, a contradiction.

(2) Here, the proof is similar.

In order to extend the method of computing Hilbert depth given in Theorem 2.4 to quotients of arbitrary monomial ideals, we can use the well-known procedure of polarization; see for instance [13, Section 1.5].

Let  $I \subset J \subset S$  be two monomial ideals. Let  $x^g := \operatorname{lcm}(u : u \in G(I) \cup G(J))$ , where  $g = (g_1, \ldots, g_n) \in \mathbb{N}^n$  and  $x^g = x_1^{g_1} \cdots x_n^{g_n}$ . We consider the polynomial ring

$$R := S[x_{ij} : 1 \le i \le n, 2 \le j \le g_i]$$

Note that we added  $N = \sum_{i=1}^{n} \max\{0, g_i - 1\}$  new variables, i.e.,  $\dim(R) = \dim(S) + N$ .

If  $u = x_1^{a_1} \cdots x_n^{a_n}$  is a monomial such that  $u \mid x^g$ , that is,  $a_i \leq g_i$  for  $1 \leq i \leq n$ , we define the square-free monomial

$$u^{p} = x_{1}^{\min\{a_{1},1\}} x_{12} \cdots x_{1,a_{1}} x_{2}^{\min\{a_{2},1\}} x_{22} \cdots x_{2,a_{2}} \cdots x_{n}^{\min\{a_{n},1\}} x_{n2} \cdots x_{n,a_{n}}.$$

The polarizations of I and J are the square-free monomial ideals

 $I^p = (u^p : u \in G(I)) \subset R \text{ and } J^p = (u^p : u \in G(J)) \subset R.$ 

**Proposition 2.6.** With the above notations:

$$\operatorname{hdepth}(J/I) := \operatorname{hdepth}(J^p/I^p) - N.$$

*Proof.* Since J/I is obtained from  $J^p/I^p$  by factorization with a regular sequence consisting of N linear forms, we have that

$$H_{J/I}(t) = (1-t)^N H_{J^p/I^p}(t).$$

Now, the conclusion follows from Theorem 2.3.

As a direct consequence of [14, Theorem 4.3], we have the following result:

**Proposition 2.7.** With the above notations:

$$\operatorname{sdepth}(J/I) = \operatorname{sdepth}(J^p/I^p) - N.$$

Now, we can reprove the following well known fact:

**Proposition 2.8.** Let  $I \subset J \subset S$  be two monomial ideals. Then

$$\operatorname{sdepth}(J/I) \leq \operatorname{hdepth}(J/I).$$

*Proof.* From Proposition 2.6 and Proposition 2.7 we can reduce to the squarefree case. The conclusion follows from (2.3) and Theorem 2.4.

*Example* 2.9. Let  $I = (x_1^2, x_1 x_2^2) \subset S = K[x_1, x_2]$ . Then  $I^p = (x_1 x_{12}, x_1 x_2 x_{22}) \subset R = S[x_{12}, x_{22}]$ . For simplicity, we denote  $x_3 := x_{12}, x_4 := x_{22}$ , and thus  $I^p = (x_1 x_3, x_1 x_2 x_4) \subset S[x_1 x_2 x_2 x_3 x_3 x_3 x_3 x_3 x_4]$ 

 $R = K[x_1, x_2, x_3, x_4]$ . We consider  $P = P_{R/I^p}$ . We denote  $\alpha_j := \alpha_j(R/I^p)$  and  $\beta_k^q := \beta_k^q(R/I^p)$  for all j, k and q. It is easy to see that

$$\alpha_0 = 1, \ \alpha_1 = 4, \ \alpha_2 = 5, \ \text{and} \ \alpha_3 = 1.$$

For q = 2, we have

$$\beta_0^2 = \alpha_0 = 1, \ \beta_1^2 = \alpha_1 - \binom{2}{1}\beta_0^2 = 2, \ \beta_2^2 = \alpha_2 - \binom{2}{2}\beta_0^2 - \binom{1}{1}\beta_1^2 = 5 - 2 - 1 = 2.$$

For q = 3, we have

$$\beta_0^3 = \alpha_0 = 1, \ \beta_1^3 = \alpha_1 - \binom{3}{1}\beta_0^2 = 1, \ \beta_2^3 = \alpha_2 - \binom{3}{2}\beta_0^3 - \binom{2}{1}\beta_1^3 = 5 - 3 - 2 = 0.$$

Moreover, we have

$$\beta_3^3 = \alpha_3 - \binom{3}{3}\beta_0^3 - \binom{2}{2}\beta_1^3 = 1 - 1 - 1 = -1 < 0.$$

It follows that  $hdepth(R/I^p) = 2$  and thus  $hdepth(S/I) = hdepth(R/I^p) - 2 = 0$ .

The following result can be seen as the counterpart of [11, Lemma 3.6] in the framework of Hilbert depth:

**Lemma 2.10.** Let  $I \subsetneq J \subset S$  be two monomial ideals. Let  $\overline{I} = I\overline{S}$  and  $\overline{J} = J\overline{S}$  be the extensions of I and J in the ring  $\overline{S} := S[x_{n+1}] = K[x_1, \ldots, x_{n+1}]$ . Then

$$hdepth(J/I) = hdepth(J/I) + 1$$

*Proof.* Since  $x_{n+1}$  is regular on  $(\overline{J}/\overline{I})$  and  $(\overline{J}/\overline{I})/x_{n+1}(\overline{J}/\overline{I}) \cong J/I$ , we have that

$$H_{J/I}(t) = (1-t)H_{\overline{J}/\overline{I}}(t).$$

The conclusion follows from Theorem 2.3.

We also recall the following result:

**Proposition 2.11.** Let  $0 \to U \to M \to N \to 0$  be a short exact sequence of finitely generated graded S-modules. Then:

 $hdepth(M \oplus N) \ge \min\{hdepth(M), hdepth(N)\}.$ 

*Proof.* Since  $H_M(t) = H_U(t) + H_N(t)$ , the conclusion follows from Theorem 2.3.

As a particular case, we get:

**Corollary 2.12.** Let  $I \subsetneq J \subset S$  be two monomial ideals. Then:

- (1) hdepth $(S/I) \ge \min\{\text{hdepth}(S/J), \text{hdepth}(J/I)\}.$
- (2)  $\operatorname{hdepth}(J) \ge \min\{\operatorname{hdepth}(I), \operatorname{hdepth}(J/I)\}.$

**Lemma 2.13.** Let  $I \subset J \subset S' := K[x_1, \ldots, x_m]$  be two monomial ideals and  $u \in S[x_{m+1}, \ldots, x_n]$  a monomial. Then

hdepth(JS/IS) = hdepth(u(JS/IS)).

*Proof.* Since  $H_{u(JS/IS)}(t) = t^{\deg(u)} H_{JS/IS}(t)$ , the conclusion follows from Theorem 2.3.

As a direct consequence of Lemma 2.13 we get:

**Corollary 2.14.** Let  $I \subset S$  be a monomial ideal and  $u \in S$  a monomial such that  $u \notin I$  and I = u(I : u). Then hdepth(I : u) = hdepth(I).

**Lemma 2.15.** Let  $I \subset S$  be a monomial ideal and  $u \in S$  a monomial with  $u \notin I$ . Then hdepth $(S/I) \geq \min\{\text{hdepth}(S/(I:u)), \text{hdepth}(S/(I,u))\}.$ 

*Proof.* Using the short exact sequence

$$0 \to S/(I:u) \xrightarrow{\cdot u} S/I \to S/(I,u) \to 0,$$

we deduce that

$$H_{S/I}(t) = t^{\deg u} \cdot H_{S/(I:u)}(t) + H_{S/(I,u)}(t)$$

The conclusion follows from Theorem 2.3.

We recall the following well-known results.

**Proposition 2.16.** Given a monomial ideal  $I \subset S$  and  $u \in S$  a monomial which is not contained in I, we have the following:

- (1)  $\operatorname{sdepth}(S/(I:u)) = \operatorname{sdepth}(S/I)$  if I = u(I:u); see [4, Theorem 1.1(1)];
- (2)  $\operatorname{sdepth}(S/(I:u)) \ge \operatorname{sdepth}(S/I)$ ; see [17, Proposition 1.3] (arXiv version);
- (3)  $\operatorname{sdepth}(I:u) \ge \operatorname{sdepth}(I); see [5, \operatorname{Proposition} 2.7(2)];$
- (4)  $\operatorname{depth}(S/(I:u)) \ge \operatorname{depth}(S/I)$ ; see [19, Corollary 1.3].

It is natural to ask if similar formulae hold for Hilbert depth. This is not the case, as the following example shows.

Example 2.17. (1) Let  $I = (x_1x_2, x_2x_3, x_3x_4, x_4x_5) \subset S = K[x_1, \ldots, x_6]$ . By straightforward computations, using Theorem 2.4, we get

hdepth(S/I) = 3 and hdepth $(S/x_6I) = 4$ .

This shows that results similar to (1), (2) and (4) of Proposition 2.16 do not hold in the framework of Hilbert depth.

Let  $I' = I \cap S'$ , where  $S' = K[x_1, \ldots, x_5]$ . From Lemma 2.10, Corollary 2.14 and the straightforward computation of hdepth(I'), we have

 $hdepth(I) = hdepth(x_6I) = hdepth(I') + 1 = 5.$ 

(2) Let  $I = (x_1x_2, x_2x_3, x_3x_4, x_4x_5, x_5x_1)$  and  $J = (x_1x_2, x_2x_3, x_3x_4, x_4x_5, x_5x_1x_6)$  be ideals in  $S = K[x_1, \ldots, x_6]$ . It is clear that  $(J : x_6) = I$ . By straightforward computations we get

hdepth(I) = 5 > hdepth(J) = 4,

which shows that the results (3) and (4) of Proposition 2.16 do not hold in the framework of Hilbert depth.

We recall the following result.

**Theorem 2.18.** Let  $I \subset S$  be a monomial ideal minimally generated by m monomials. Then:

- (1) sdepth $(S/I) \ge n m$ ; see [4, Proposition 1.2];
- (2) sdepth(I)  $\geq \max\{1, n \lfloor \frac{m}{2} \rfloor\}$ ; see [16, Theorem 2.3].

**Corollary 2.19.** Let  $I \subset S$  be a monomial ideal minimally generated by m monomials. Then:

- (1)  $\operatorname{hdepth}(S/I) \ge n m$ .
- (2) hdepth(I)  $\geq \max\{1, n \lfloor \frac{m}{2} \rfloor\}.$

*Proof.* This follows from Theorem 2.18 and Proposition 2.8.

**Lemma 2.20.** Let  $1 \le m < n$ , and let  $I' \subset S' := K[x_1, \ldots, x_m]$  be a square-free monomial ideal. Let  $u = x_{m+1} \cdots x_n$  and  $s = \deg(u) = m - n$ . Let  $q \ge 0$  be an integer. Then

$$\beta_k^{q+s}(S/(I,u)) = \begin{cases} \beta_k^q(S'/I') - \beta_{k-s}^q(S'/I'), & \text{if } 0 \le k \le q, \\ \sum_{\ell=0}^{k-q-1} \binom{k-q-1}{\ell} \alpha_{q+1+\ell}(S'/I') - \beta_{k-s}^d(S'/I'), & \text{if } q+1 \le k \le q+s, \end{cases}$$

where  $\beta_j^q(S'/I') = 0$  for j < 0.

*Proof.* Let  $k \leq s$ . Since  $S/(I', u) \cong S/I'S \oplus (I, u)/I$  it follows that

$$\alpha_k(S/(I', u)) = \alpha_k(S/I'S) - \alpha_k((I', u)/I'S).$$
(2.5)

Let  $v \in S$  be a square-free monomial. Then  $v \in (I', u) \setminus I'$  if and only if  $v \mid u$  and u = vwwith  $w \in S' \setminus I'$ . It follows that  $\alpha_k((I', u)/I'S) = \alpha_{k-s}(S'/I')$ , and therefore from (2.5) we get

$$\alpha_k(S/(I',u)) = \alpha_k(S/I'S) - \alpha_{k-s}(S'/I') \text{ for } 0 \le k \le d,$$

$$(2.6)$$

where  $\alpha_j(S'/I') = 0$  for j < 0. If k < s then  $\alpha_k(S/(I', u)) = \alpha_k(S/I'S)$  and thus  $\beta_k^q(S/(I', u)) = \beta_k^q(S/I'S)$ . Now, assume  $k \ge s$ . From (2.1) it follows that

$$\beta_k^{q+s}(S/(I',u)) = \beta_q^{d+s}(S/I'S) - \sum_{j=s}^k (-1)^{k-j} \binom{q+s-j}{k-j} \alpha_{j-s}(S'/I') \text{ for } 0 \le k \le q.$$
(2.7)

Using induction on  $s \ge 1$ , we can easily get

$$\beta_k^{q+s}(S/I'S) = \begin{cases} \beta_k^q(S'/I'), & 0 \le k \le q, \\ \sum_{\ell=0}^{k-q-1} \binom{k-q-1}{\ell} \alpha_{q+1+\ell}(S'/I'), & q+1 \le k \le q+s. \end{cases}$$
(2.8)

On the other hand, using the substitution m := j - s, we get

$$\sum_{j=s}^{k} (-1)^{k-j} {q+s-j \choose k-j} \alpha_{j-s} (S'/I') = \sum_{m=0}^{k-s} (-1)^{(k-s)-m} {q-m \choose (k-s)-m} \alpha_m (S'/I') = \beta_{k-s}^q (S'/I'), \quad (2.9)$$

for  $s \le k \le q$ . From (2.7), (2.8) and (2.9) we get the required conclusion.

We recall the following result.

**Theorem 2.21** ([18, Theorem 1.1]). Let  $I \subset S$  be a monomial ideal and  $u \in S$  a monomial, regular of S/I. Then:

$$\operatorname{sdepth}(S/(I, u)) = \operatorname{sdepth}(S/I) - 1.$$

It is natural to ask if a similar result holds in the framework of Hilbert depth. The following result is the best we can expect.

**Theorem 2.22.** Let  $I \subset S$  be a monomial ideal and  $u \in S$  a monomial, regular of S/I. Then:

- (1)  $\operatorname{hdepth}(S/I) \ge \operatorname{hdepth}(S/(I, u)) \ge \operatorname{hdepth}(S/I) 1$ . In particular, if u is a variable then  $\operatorname{hdepth}(S/(I, u)) = \operatorname{hdepth}(S/I) 1$ .
- (2)  $\operatorname{hdepth}((I, u)) \ge \min\{\operatorname{hdepth}(I), \operatorname{hdepth}(S/I)\}.$

*Proof.* (1) Without loss of generality, we may assume that I = I'S, where  $I' \subset S' = K[x_1, \ldots, x_m]$  is a square-free monomial ideal, and  $u = x_{m+1} \cdots x_n$ . Let j := n - m. We use induction on  $j \ge 1$ . If j = 1 then m = n - 1,  $u = x_n$  and

$$S/(I, u) \cong S/(I', x_n) \cong S'/I'.$$

Thus, the result follows from Lemma 2.10.

Now, assume  $j \ge 2$ . For  $0 \le j \le n - m$  we let  $I_j := I + (x_{m+j+1} \cdots x_n) \subset S$ . We have the inclusions

$$I = I_0 \subset I_1 \subset \cdots \subset I_{n-m-1} \subset I_{n-m} = S.$$

Therefore, we have

$$S/I \cong (I_1/I_0) \oplus (I_2/I_1) \oplus \dots \oplus (I_{n-m}/I_{n-m-1}).$$
 (2.10)

Since  $I_{j+1}/I_j \cong S/(I, x_{m+j+1})$  for all j with  $0 \le j \le n - m - 1$ , from Theorem 2.10 it follows that

$$hdepth(I_{j+1}/I_j) = hdepth(S/I) - 1 \text{ for } 0 \le j \le n - m - 1.$$
 (2.11)

From (2.10), (2.11) and Proposition 2.11 it follows that

$$hdepth(S/(I, u)) \ge hdepth(S/I) - 1.$$

Thus, in order to complete the proof, we have to show the other inequality.

Let q := hdepth(S'/I'). From Lemma 2.20, we have

$$\beta_k^{q+1+s}(S/(I,u)) = \beta_k^{q+1}(S'/I') - \beta_{k-s}^{q+1}(S'/I') \text{ for } 0 \le k \le q+1.$$
(2.12)

Let  $j_0 := \min\{\beta_j^{q+1}(S'/I') < 0\}$ . From (2.12) it follows that

$$\beta_{j_0}^{q+1+s}(S/(I,u)) < 0,$$

and therefore  $hdepth(S/(I, u)) \le q + s = hdepth(S/I)$ , as required.

(2) We write  $(I, u) = I \oplus (I, u)/I$ . Since (I, u)/I = u(S/I), the result follows from Propositions 2.11 and Lemma 2.13.

Remark 2.23. Let  $I' \subset S' = K[x_1, \ldots, x_m]$  be a square-free monomial ideal and  $u = x_{m+1} \cdots x_n$ , as in the proof of Theorem 2.22. Let I = I'S. Let q := hdepth(S'/I') and assume that

$$\alpha_{q+1}(S'/I') < \beta_{q+1-\deg(u)}^q(S'/I').$$

From Lemma 2.20 it follows that

$$\beta_{q+1}^{q+s}(S/(I,u)) = \alpha_{q+1}(S'/I') - \beta_{q+1-\deg(u)}^q(S'/I') < 0,$$

hence  $hdepth(S/(I, u)) \leq hdepth(S/I) - 1$ . Therefore, from Theorem 2.22, it follows that hdepth(S/(I, u)) = hdepth(S/I) - 1. However, in general this equality does not hold, as the following example shows.

*Example 2.24.* Let  $I' = (x_1, x_2) \cap (x_3, x_4) \cap (x_5, x_6, x_7) \subset S' = K[x_1, x_2, \dots, x_7]$ . We also consider the ideal I := I'S and the monomial  $u = x_8x_9 \in S$ , where  $S = K[x_1, x_2, \dots, x_9]$ .

Using CoCoA [7], we computed hdepth(S'/I') = 3 and hdepth(S/(I, u)) = 5. Therefore, we have hdepth(S/I) = hdepth(S/(I, u)).

Let  $I \subset S$  be a square-free monomial ideal with  $G(I) = \{u_1, u_2, \ldots, u_m\}$ . For any nonempty subset  $J \subset [m] = \{1, 2, \ldots, m\}$ , we let

 $u_J := \operatorname{lcm}(u_j : j \in J) \text{ and } d_J := \operatorname{deg}(u_J).$ 

Furthermore, we denote  $u_{\emptyset} := 1$  and  $d_{\emptyset} := 0$ . As usual, if j < 0 and  $m \ge 0$  we define  $\binom{m}{i} := 0$ .

,

Theorem 2.25. With the above notations, we have

(1) 
$$\alpha_k(I) = \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|-1} \binom{n-d_J}{k-d_J}$$
  
(2)  $\alpha_k(S/I) = \sum_{J \subset [m]} (-1)^{|J|} \binom{n-d_J}{k-d_J}.$ 

*Proof.* (1) We use the inclusion-exclusion principle. Let  $P = P_I \subset 2^{[n]}$ . We consider the subsets

$$P_j = \{ A \in P : u_j : x_A = \prod_{i \in A} x_i \} \text{ for } 1 \le j \le m.$$
(2.13)

Moreover, if  $J \subset [m]$  is nonempty, we denote  $P_J = \bigcap_{j \in J} P_j$ . In particular,  $P_{\{j\}} = P_j$  for  $1 \leq j \leq m$ . From (2.13) it follows that

$$P_J = \left\{ A \in P : u_J : x_A = \prod_{i \in A} x_i \right\} \text{ for } \emptyset \neq J \subset [m].$$

$$(2.14)$$

Since  $P = P_1 \cup P_2 \cup \cdots \cup P_m$ , it follows that

$$|P| = \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|-1} |P_J|.$$
(2.15)

On the other hand, since  $\deg(u_J) = d_J$ , the number of square-free monomials of degree k in  $P_J$  is  $\binom{n-d_J}{k-d_J}$ , that is,  $\alpha_k(P_J) = \binom{n-d_J}{k-d_J}$ . Therefore, the required formula follows from (2.14) and (2.15).

(2) This follows from (1) and the fact that  $\alpha_k(S/I) = \binom{n}{k} - \alpha_k(I) = \binom{n-d_{\emptyset}}{k-d_{\emptyset}} - \alpha_k(I)$ .  $\Box$ 

Remark 2.26. Let  $0 \subset I \subset J \subset S$  be two square-free monomial ideals. From the decomposition  $S/I = S/J \oplus J/I$ , it follows that

$$\alpha_k(J/I) = \alpha_k(S/I) - \alpha_k(S/J) \text{ for } 0 \le k \le n.$$

Therefore, applying Theorem 2.25, we can write  $\alpha_k(J/I)$  in combinatorial terms of the degrees of the least common multiple of the minimal monomial generators of I and J, respectively.

**Corollary 2.27.** Let  $I \subset S$  be a square-free monomial complete intersection with  $G(I) = \{u_1, u_2, \ldots, u_m\}$ . Let  $d_j = \deg(u_j)$  for all j with  $1 \leq j \leq n$ . For a nonempty subset  $J \subset [m]$  we let  $d_J := \sum_{j \in J} d_j$ . Then we have

(1) 
$$\alpha_k(I) = \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|-1} \binom{n-d_J}{k-d_J},$$
  
(2)  $\alpha_k(S/I) = \sum_{J \subset [m]} (-1)^{|J|} \binom{n-d_J}{k-d_J}.$ 

*Proof.* It is enough to notice that  $u_J = \text{lcm}(u_j : j \in J) = \prod_{j \in J} u_j$  and thus  $d_J = \sum_{j \in J} d_j$ , as required. Subsequently we apply Theorem 2.25.

We recall the following result.

**Theorem 2.28.** Let  $I \subset S$  be a monomial complete intersection, minimally generated by *m* monomials. Then:

- (1)  $\operatorname{sdepth}(S/I) = n m$ ; see Theorem 2.21;
- (2) sdepth(I) =  $n \lfloor \frac{m}{2} \rfloor$ ; see [23, Theorem 2.4].

*Remark* 2.29. If  $\mathbf{m} = (x_1, \ldots, x_n)$  is the maximal graded ideal of *S*, then the following identities are well known:

$$hdepth(\mathbf{m}) = sdepth(\mathbf{m}) = \left\lceil \frac{n}{2} \right\rceil.$$

It is natural to ask if this result can be generalized to the case of monomial complete intersections, that is, if hdepth(I) = sdepth(I) for any square-free monomial complete intersection  $I \subset S$  minimally generated by m monomials? The answer is no.

Let  $I = (x_1, x_2, x_3, x_4, x_5x_6, x_7x_8) \subset S = K[x_1, \dots, x_8]$ . We have

hdepth
$$(I) = 6 > 8 - \left\lfloor \frac{6}{2} \right\rfloor = 5 = \text{sdepth}(I).$$

Hence, a result similar to Theorem 2.28(2) does not hold in the framework of Hilbert depth.

Note that, according to Theorems 2.18 and 2.28, the case of monomial complete intersections  $I \subset S$  gives minimal values for sdepth(S/I) and sdepth(I) in terms of the number of minimal monomial generators of I. In the following proposition, we note that a similar result holds for hdepth(S/I).

**Proposition 2.30.** Let  $I \subset S$  be a monomial complete intersection, minimally generated by m monomials. Then hdepth(S/I) = n - m.

*Proof.* This result can be easily deduced from the special form of the Hilbert series of S/I. Here we present an alternative proof: Without any loss of generality, we may assume that I is squarefree and, moreover, that I has the minimal system of generators  $G(I) = \{u_1, \ldots, u_m\}$  with  $d_j = \deg(u_j)$  for  $1 \le j \le m$ . Furthermore, since I is minimally generated by the monomials  $u_1, \ldots, u_m$  with disjoint supports, it is easy to see that the maximal degree of a squarefree monomial u which is not contained in I is n - m. Therefore, we have

$$\alpha_k(S/I) = 0 \text{ and } \alpha_k(I) = \binom{n}{k} \text{ for } n - m + 1 \le k \le n.$$
 (2.16)

In particular, from Lemma 2.5, it follows that

hdepth $(S/I) \le \max\{k : \alpha_k(S/I) > 0\} = n - m.$ 

On the other hand,  $hdepth(S/I) \ge sdepth(S/I) = n - m$ . Hence hdepth(S/I) = n - m.  $\Box$ 

Our computer experiments lead us to conjecture the following striking identity.

**Conjecture 2.31.** Let  $I \subset S$  be a monomial complete intersection, minimally generated by m monomials. Then

$$\beta_k^{n-m+1}(S/I) + \beta_{n-m+1-k}^{n-m+1}(S/I) = 0 \text{ for } 0 \le k \le n-m+1.$$

**Proposition 2.32.** Let  $I \subset S$  be a monomial complete intersection, minimally generated by m monomials. Then

$$\beta_{n-m+1}^{n-m+1}(S/I) = -1.$$

In particular, Conjecture 2.31 holds for k = 0.

*Proof.* From (2.1) we have

$$\beta_{n-m+1}^{n-m+1}(S/I) = \sum_{j=0}^{n-m+1} (-1)^{n-m+1-j} \alpha_j(S/I).$$

Therefore, using (2.16) and changing j into k, we obtain

$$\beta_{n-m+1}^{n-m+1}(S/I) = (-1)^{m-1} \sum_{k=0}^{n} (-1)^{n-k} \alpha_k(S/I).$$
(2.17)

From (2.17) and Corollary 2.27 it follows that

$$\beta_{n-m+1}^{n-m+1}(S/I) = (-1)^{m-1} \sum_{k=0}^{n} (-1)^{n-k} \left( \binom{n}{k} + \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|} \binom{n-d_J}{k-d_J} \right), \qquad (2.18)$$

where 
$$d_J = \sum_{j \in J} d_j$$
. Since  $\sum_{k=0}^n (-1)^{n-k} {n \choose k} = (1-1)^n = 0$ , from (2.18) we get  

$$\beta_{n-m+1}^{n-m+1}(S/I) = (-1)^{m-1} \sum_{k=0}^n (-1)^{n-k} \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|} {n-d_J \choose k-d_J}$$

$$= (-1)^{m-1} \sum_{\emptyset \neq J \subset [m]} (-1)^{|J|} \sum_{k=0}^n (-1)^{n-k} {n-d_J \choose k-d_J}.$$
(2.19)

If  $J \neq [m]$ , then  $d_J < n$  and therefore

$$\sum_{k=0}^{n} (-1)^{n-k} \binom{n-d_J}{k-d_J} = \sum_{k=d_J}^{n} (-1)^{n-k} \binom{n-d_J}{k-d_J} = \sum_{\ell=0}^{n-d_J} (-1)^{n-d_J-\ell} \binom{n-d_J}{\ell} = 0. \quad (2.20)$$

On the other hand, if J = [m], then  $d_J = n$  and therefore

$$\sum_{k=0}^{n} (-1)^{n-k} \binom{n-d_J}{k-d_J} = \binom{0}{0} = 1.$$
 (2.21)

From (2.19), (2.20) and (2.21) it follows that

$$\beta_{n-m+1}^{n-m+1}(S/I) = (-1)^{m-1} \cdot (-1)^m = (-1)^{2m-1} = -1 = -\beta_0^{n-m+1}(S/I).$$

Hence, the proof is complete.

# 3. Square-Free Veronese ideals

**Definition 3.1.** Let  $n \ge m \ge 1$  be two integers. Let  $J_{n,m}$  be the ideal in  $S := K[x_1, \ldots, x_n]$  generated by all square-free monomials of degree m.  $J_{n,m}$  is called the square-free Veronese ideal of degree m.

*Example* 3.2. If n = 4 and m = 2 then there are exactly  $\binom{4}{2} = 6$  square-free monomials of degree 2 in  $S := K[x_1, x_2, x_3, x_4]$ . Moreover, we have

$$J_{4,2} = (x_1x_2, x_1x_3, x_1x_4, x_2x_3, x_2x_4, x_3x_4) \subset S = K[x_1, x_2, x_3, x_4].$$

Lemma 3.3. With the above notations, we have the following:

(1) 
$$\alpha_k(S/J_{n,m}) = \begin{cases} \binom{n}{k}, & \text{if } k < m, \\ 0, & \text{if } m \le k \le n \end{cases}$$
  
(2)  $\alpha_k(J_{n,m}) = \begin{cases} 0, & \text{if } k < m, \\ \binom{n}{k}, & \text{if } m \le k \le n. \end{cases}$ 

*Proof.* (1) We have  $P_{S/J_{n,m}} = \{A \subset [n] : x_A = \prod_{j=1}^n x_j \notin J_{n,m}\}$ . Since  $J_{n,m}$  is generated by all square-free monomials of degree m, the required conclusion follows.

(2) This follows from the fact that  $P_{J_{n,m}} = 2^{[n]} \setminus P_{S/J_{n,m}}$  and (1).

Proposition 3.4. With the above notations, we have

$$hdepth(S/J_{n,m}) = sdepth(S/J_{n,m}) = m - 1.$$

*Proof.* From [6, Theorem 1.1(2)] it follows that  $\operatorname{sdepth}(S/J_{n,m}) = m - 1$ . On the other hand, from Lemma 2.5 and Lemma 3.3 it follows that  $\operatorname{hdepth}(S/J_{n,m}) \leq m - 1$ . The conclusion follows from Proposition 2.6.

**Proposition 3.5.** We have hdepth $(J_{n,m}) \leq m + \lfloor \frac{n-m}{m+1} \rfloor$ . In particular, if  $n \leq 2m$  then hdepth $(J_{n,m}) = m$ .

*Proof.* If n = m then there is nothing to prove, so we may assume n > m.

Since  $\alpha_1(J_{n,m}) = \cdots = \alpha_{m-1}(J_{n,m}) = 0$ , from Lemma 2.5 we get hdepth $(J_{n,m}) \ge m$ . Let d be an integer with  $m < d \le n$ . From Lemma 3.3 it follows that

$$\beta_m^d(J_{n,m}) = \binom{n}{m} \text{ and } \beta_{m+1}^d(J_{n,m}) = \binom{n}{m+1} - (d-m)\binom{n}{m}.$$
(3.1)

If  $\binom{n}{m+1} < \binom{n}{m}$ , i.e.,  $m \ge \lfloor \frac{n+1}{2} \rfloor$ , then, according to (3.1), it follows that  $\beta_{m+1}^d(J_{n,m}) < 0$  and thus hdepth $(J_{n,m}) = m$ .

Now, assume this is not the case. From (3.1) it follows that

$$\beta_{m+1}^d(J_{n,m}) < 0$$
 if and only if  $d > m + \frac{n-m}{m+1}$ 

Therefore, we get the required formula.

**Proposition 3.6.** Let  $m \ge 1$  and  $n \ge 2m + 1$  be two integers and let  $q := \lfloor \frac{n-m}{m+1} \rfloor$ . For  $1 \le t \le q$  we have

$$\beta_{m+t}^{m+q}(J_{n,m}) = \sum_{j=0}^{t} (-1)^{t-j} \binom{q-j}{t-j} \binom{n}{m+j}.$$

*Proof.* Since  $n \ge 2m+1$  it follows that  $q \ge 1$ . Let  $d := m + \lfloor \frac{n-m}{m+1} \rfloor = m+q$ . From Lemma 3.3(2) and (2.1) we have  $\beta_m^d(J_{n,m}) = \binom{n}{m}$  and

$$\beta_k^{m+q}(J_{n,m}) = \sum_{\ell=m}^k (-1)^{k-\ell} \binom{m+q-\ell}{k-\ell} \binom{n}{\ell} \text{ for } m+1 \le k \le m+q.$$
(3.2)

By letting t = k - m and  $j = \ell - m$ , the conclusion follows from (3.2).

We want to prove that

$$b(n,m,q,t) := \beta_{m+t}^{m+q}(J_{n,m}) = \sum_{j=0}^{t} (-1)^{t-j} \binom{q-j}{t-j} \binom{n}{m+j}$$
(3.3)

is non-negative for  $m, q \ge 1, 1 \le t \le q$ , and  $n \ge mq + m + q$ . We shall show something stronger, see Theorems 3.7 and 3.9 below.

**Theorem 3.7.** Let t be even. Then  $b(n, m, q, t) \ge 0$  for all non-negative integers n, m, q, t. Proof. We do a simultaneous induction on n and m.

For the start of the induction, we need to verify the claim for m = 0 and for n = 0. Indeed, by the Chu–Vandermonde summation formula, we have

$$b(n, 0, q, t) = \sum_{j=0}^{t} (-1)^{t-j} {q-j \choose t-j} {n \choose j}$$
$$= \sum_{j=0}^{t} {-q+t-1 \choose t-j} {n \choose j}$$
$$= {-q+t-1+n \choose t}.$$

For  $-q + t - 1 + n \ge 0$  this last binomial coefficient is non-negative, regardless whether t is even or odd. Moreover, it is also non-negative for -q + t - 1 + n < 0 if t is even.

On the other hand, if t is even, then we have  $b(0, m, q, t) = \delta_{m,0} {q \choose t}$ , where  $\delta_{m,0}$  is 1 if m = 0 and 0 otherwise. Clearly, this shows that b(0, m, q, t) is non-negative.

For the induction step, we observe that, by the standard three-term recurrence for binomial coefficients, we have

$$b(n, m, q, t) = b(n - 1, m, q, t) + b(n - 1, m - 1, q, t).$$

This relation completes the induction.

Lemma 3.8. We have

$$b(n,m,q,t) = \sum_{j=0}^{t} (-1)^{t-j} \frac{(m+t-j-1)! \, n! \, (n-m-q+j-1)! \, (q-j)!}{(m-1)! \, (n-m)! \, (n-m-q-1)! \, (t-j)! \, (m+t)! \, (q-t)!}.$$

*Proof.* We write the binomial sum b(n, m, q, t) in standard hypergeometric notation (cf. [22])

$${}_{r}F_{s}\begin{bmatrix}a_{1},\ldots,a_{r}\\b_{1},\ldots,b_{s};z\end{bmatrix} = \sum_{k=0}^{\infty}\frac{(a_{1})_{k}\cdots(a_{r})_{k}}{k!\,(b_{1})_{k}\cdots(b_{s})_{k}}z^{k}$$

where  $(\alpha)_k := \alpha(\alpha + 1) \cdots (\alpha + k - 1), k \ge 1$ , and  $(\alpha)_0 := 1$ . We obtain

$$b(n,m,q,t) = (-1)^t \binom{q}{t} \binom{n}{m} {}_3F_2 \begin{bmatrix} m-n,1,-t\\-q,m+1 \end{bmatrix}.$$

We now apply the transformation formula (see [9, (3.1.1)])

$$_{3}F_{2}\begin{bmatrix}a,b,-m\\d,e\end{bmatrix} = \frac{(e-b)_{m}}{(e)_{m}} _{3}F_{2}\begin{bmatrix}-m,b,d-a\\d,1+b-e-m\end{bmatrix},$$

where m is a non-negative integer. After some manipulation, one obtains the claimed expression.

**Theorem 3.9.** Let t be odd. Then  $b(n, m, q, t) \ge 0$  for all non-negative integers n, m, q, t with

$$n \ge \max\{mq + m + q - (t-1)(m+1), mq + m + q - \frac{q(m-1)(t-1)}{t}\}.$$
(3.4)

## SILVIU BĂLĂNESCU<sup>1</sup>, MIRCEA CIMPOEAȘ<sup>2</sup> AND CHRISTIAN KRATTENTHALER<sup>3</sup>

*Proof.* Clearly,  $b(n, m, q, 0) = \binom{n}{m} \ge 0$ . Furthermore, for q < t, we have

$$b(n,m,q,t) = \sum_{j=q+1}^{t} (-1)^{t-j} \binom{q-j}{t-j} \binom{n}{m+j} = \sum_{j=q+1}^{t} \binom{t-q-1}{t-j} \binom{n}{m+j} > 0.$$

Hence, we may assume  $t \ge 1$  and  $q \ge t$  from now on.

We use the expression from the lemma. We investigate the growth properties of the summand (without sign)

$$f(n,m,q,t,j) = \frac{(m+t-j-1)!\,n!\,(n-m-q+j-1)!\,(q-j)!}{(m-1)!\,(n-m)!\,(n-m-q-1)!\,(t-j)!\,(m+t)!\,(q-t)!}.$$

We claim that, under the assumption (3.4), f(n, m, q, t, j) is (weakly) monotone increasing in j. Indeed, we have

$$\frac{f(n,m,q,t,j)}{f(n,m,q,t,j-1)} = \frac{(n-m-q+j-1)(t-j+1)}{(q-j+1)(m+t-j)}$$

This will be at least 1 if

$$n \ge 1 - 2j + 2m + 2q + \frac{(q-t)(m-1)}{t-j+1}.$$
(3.5)

We consider the right-hand side as a function in j. Differentiation of the right-hand side yields

$$-2 + \frac{(q-t)(m-1)}{(t-j+1)^2}$$

Equating this to zero, we arrive at a quadratic equation in j with two real solutions, one less than t+1, one greater than t+1 (here we use that  $t \ge 1$  and  $q \ge t$ ). Since the function (in j) on the right-hand side of (3.5) tends to  $+\infty$  as  $j \to -\infty$  and as  $j \to (t+1)^-$ , it is convex for  $j \in [1, t]$ . The maximum on [1, t] is therefore found as the greater of the values of the right-hand side of (3.5) at the boundary points j = 1 and j = t. Indeed, these two values are the ones of which the maximum is taken on the right-hand side of (3.4).

Now we pair the summands in the sum that defines b(n, m, q, t),

$$b(n, m, q, t) = (f(n, m, q, t, t) - f(n, m, q, t, t - 1)) + (f(n, m, q, t, t - 2) - f(n, m, q, t, t - 3)) + \cdots$$

Given the just proved monotonicity of the summand, we see that each pair produces a non-negative value, proving non-negativity of b(n, m, q, t) under the assumption (3.4).  $\Box$ 

Now, we can reprove the main result of [10].

**Theorem 3.10.** ([10, Theorem 1.2]) Let  $n \ge m \ge 1$ . We have that:

hdepth
$$(J_{n,m}) = m + \left\lfloor \frac{n-m}{m+1} \right\rfloor.$$

*Proof.* Let  $q = \lfloor \frac{n-m}{m+1} \rfloor$ . From Proposition 3.5 we have hdepth $(J_{n,m}) \leq m+q$ . The other inequality follows from Theorem 3.7, Theorem 3.9, (3.3) and Proposition 3.6.

Remark 3.11. In [6] we proposed the conjecture that  $\operatorname{sdepth}(J_{n,m}) = m + \lfloor \frac{n-m}{m+1} \rfloor$ ; see [6, Conjecture 1.6], and we proved that it holds for  $n \leq 3m$ ; see [6, Theorem 1.1] and [6, Corollary 1.5]. Keller et al. improved this result to  $n \leq 5m+3$ ; see [15, Theorem 1.1]. Note that  $n \leq 5m+3$  means  $q \leq 3$ . In Theorem 3.10 we reproved that  $\operatorname{hdepth}(J_{n,m}) = m + \lfloor \frac{n-m}{m+1} \rfloor$  for any  $n \geq m \geq 1$ . However, this does not imply that a similar result holds for sdepth, but it shows that such a result is credible.

Acknowledgements. We gratefully acknowledge the use of the computer algebra system Cocoa (cf. [7]) for our experiments.

Mircea Cimpoeaş was supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, project number PN-III-P1-1.1-TE-2021-1633, within PNCDI III.

**Data availability.** Data sharing not applicable to this article as no data sets were generated or analyzed during the current study.

**Conflict of interest.** The authors have no relevant financial or non-financial interests to disclose.

#### References

- J. Apel, On a conjecture of R. P. Stanley; Part II quotients modulo monomial ideals, J. Alg. Comb. 17 (2003), 57–74.
- [2] C. Biro, D. M. Howard, M. T. Keller, W. T. Trotter, S. J. Young, Interval partitions and Stanley depth, J. Combin. Theory, Ser. A 117(4) (2010), 475–482.
- [3] W. Bruns, C. Krattenthaler, J. Uliczka, Stanley decompositions and Hilbert depth in the Koszul complex, Journal of Commutative Algebra vol.2, no.3 (2010), 327–357.
- [4] M. Cimpoeaş, Stanley depth of monomial ideals with small number of generators, Central European J. Math. 7(4) (2009), 629–634.
- [5] M. Cimpoeaş, Several inequalities regarding Stanley depth, Romanian J. Math. Computer Sci. 2(1) (2012), 28–40.
- [6] M. Cimpoeaş, Stanley depth of squarefree Veronese ideals, An. Şt. Univ. Ovidius Constanţa 21(3) (2013), 67–71.
- [7] CoCoATeam, CoCoA: a system for doing Computations in Commutative Algebra, Available at http://cocoa.dima.unige.it.
- [8] A. M. Duval, B. Goeckneker, C. J. Klivans, J. L. Martine, A non-partitionable Cohen-Macaulay simplicial complex, Adv. in Math. 299 (2016), 381–395.
- [9] G. Gasper and M. Rahman, *Basic hypergeometric series*, Encyclopedia of Mathematics And Its Applications 35, Cambridge University Press, Cambridge, 1990.
- [10] M. Ge, J. Lin, Y. Wang, Hilbert series and Hilbert depth of squarefree Veronese ideals, Journal of Algebra 344 (2011), 260–267.
- [11] J. Herzog, M. Vlădoiu, X. Zheng, How to compute the Stanley depth of a monomial ideal, J. Algebra 322(9) (2009), 3151–3169.
- [12] J. Herzog, A survey on Stanley depth, In Monomial Ideals, Computations and Applications, Springer, 2013, pp. 3–45.
- [13] J. Herzog, J. Hibi, Monomial ideals, Graduate Texts in Maths, Springer, London, 2010.
- B. Ichim, L. Katthän, and J. J. Moyano-Fernandez, The behavior of Stanley depth under polarization, J. Combinatorial Theory, Ser. A, 135 (2015), 332–347.

- [15] M. T. Keller, W. T. Trotter, Y.-H. Shen, S. J. Young, On the Stanley depth of squarefree Veronese ideals, J. Algebr. Comb. 33 (2011), 313–324.
- [16] R. Okazaki, A lower bound of Stanley depth of monomial ideals, J. Commut. Algebra 3(1), (2011), 83–88.
- [17] D. Popescu, An inequality between depth and Stanley depth, Bull. Math. Soc. Sci. Math. Roumanie 52(100) (2009), 377–382, arXiv:0905.4597v2.
- [18] A. Rauf, Stanley Decompositions, Pretty Clean Filtrations and Reductions Modulo Regular Elements, Bull. Math. Soc. Sc. Math. Roumanie, 50(98) (2007), 347–354.
- [19] A. Rauf, Depth and sdepth of multigraded module, Commun. Algebra, 38(2) (2010), 773–784.
- [20] G. Rinaldo, An algorithm to compute the Stanley depth of monomial ideals, Matematiche 63(2) (2008), 243–256.
- [21] J. Riordan, Combinatorial identities, J. Wiley, New York, 1968.
- [22] L. J. Slater, Generalized hypergeometric functions, Cambridge University Press, Cambridge, 1966.
- [23] Y. Shen, Stanley depth of complete intersection monomial ideals and upper-discrete partitions, J. Algebra 321 (2009), 1285–1292.
- [24] R. P. Stanley, Linear Diophantine equations and local cohomology, Invent. Math. 68 (1982), 175–193.
- [25] J. Uliczka, Remarks on Hilbert series of graded modules over polynomial rings, Manuscripta math. 132 (2010), 159–168.