Second order quantitative bounds for unadjusted generalized Hamiltonian Monte Carlo

Evan Camrud, Alain Durmus, Pierre Monmarché, Gabriel Stoltz

May 14, 2024

Abstract

This paper provides a convergence analysis for generalized Hamiltonian Monte Carlo samplers, a family of Markov Chain Monte Carlo methods based on leapfrog integration of Hamiltonian dynamics and kinetic Langevin diffusion, that encompasses the unadjusted Hamiltonian Monte Carlo method. Assuming that the target distribution π satisfies a log-Sobolev inequality and mild conditions on the corresponding potential function, we establish quantitative bounds on the relative entropy of the iterates defined by the algorithm, with respect to π . Our approach is based on a perturbative and discrete version of the modified entropy method developed to establish hypocoercivity for the continuous-time kinetic Langevin process. As a corollary of our main result, we are able to derive complexity bounds for the class of algorithms at hand. In particular, we show that the total number of iterations to achieve a target accuracy $\varepsilon > 0$ is of order $d/\varepsilon^{1/4}$, where d is the dimension of the problem. This result can be further improved in the case of weakly interacting mean field potentials, for which we find a total number of iterations of order $(d/\varepsilon)^{1/4}$.

1 Introduction

We consider in this paper the problem of sampling from a target distribution π . This problem is ubiquitous in various fields such as statistical physics [36], statistics [24], and machine learning [2]. However, in most applications, the distribution π has a density with respect to a dominating measure known up to an intractable multiplicative constant. Markov chain Monte Carlo methods are now a family of popular algorithms for solving this problem. They consist in designing a Markov chain associated with a Markov kernel for which π is an invariant distribution. One of the best known MCMC instances is the family of Metropolis–Hastings algorithms [40, 31]. In the case where the target distribution has a smooth and positive density with respect to the Lebesgue measure on \mathbb{R}^d , denoted by π still, another class of MCMC algorithms is based on discretizations of continuous-time stochastic dynamics [27, 26, 45]. Famous examples of such MCMC methods are the Unadjusted Langevin Algorithm (ULA) [44] and Stochastic Gradient Langevin Dynamics [51], which are based on the overdamped Langevin diffusion.

Here we consider numerical schemes based on Hamiltonian-type dynamics, *i.e.*, ideal Hamiltonian Monte Carlo and kinetic (or underdamped) Langevin diffusion. Although these two dynamics are different, they have many features in common. For example, both define an extended process on \mathbb{R}^{2d} that has the product of π and the standard Gaussian distribution as invariant measure, denoted hereafter by μ . Also, the infinitesimal generators of these extended processes differ only in their symmetric parts, while their antisymmetric parts are the same and correspond to the Hamiltonian dynamics associated with the potential U associated with π , *i.e.*, $\pi \propto \exp(-U)$. From this observation it follows that common discretization strategies have been employed for both dynamics. Among these methods, those based on splitting techniques [9, 34] are particularly attractive since they come with valuable properties and important convergence guarantees.

In this paper, we are particularly interested in the family of splitting methods known as generalized Hamiltonian Monte Carlo (gHMC). This method has been shown to yield weak second order errors. More precisely, denoting by P_{ω} the Markov kernel associated with gHMC, where ω are the hyperparameters of the algorithm including the step size $\delta > 0$, the refreshment rate η and the integration time T > 0, the following convergence bound holds under suitable conditions (see e.g. [1] for the Langevin case $T = \delta$): for any sufficiently regular function $f : \mathbb{R}^{2d} \to \mathbb{R}$ and initial distribution ν_0 , there exist $C_f \ge 0$ and $c_1 > 0$ depending on the parameters ω only through the friction $\gamma = (1 - \eta)/T$ (c_1 being independent of f, ν_0) such that for any number of iterations $k \in \mathbb{N}$, $|\nu_0 P_{\omega}^k f - \pi(f)| \le C_f (e^{-c_1 Tk} + \delta^2)$. This result illustrates the advantages of using well-chosen splitting strategies compared to traditional Euler schemes, for which similar conclusions can be drawn but with a larger second term on the right hand side of the previous inequality, typically of order δ instead of δ^2 . While weak error bounds already provide significant convergence guarantees, another line of research is concerned with establishing quantitative bounds for MCMC algorithms, paying particular attention to the dimension dependence [15, 20].

Regarding the kinetic Langevin algorithm, existing works [37, 13, 52, 16] analyze for most of them a modification of the Euler scheme. For this particular algorithm, [37] shows that, when π satisfies a log-Sobolev inequality and under additional conditions on U, and denoting by ν_k the distribution of the k-th iterate of the algorithm starting from ν_0 with step size $\delta > 0$, there are constants $C_{\nu_0} \ge 0$ and $c_1 > 0$, independent of δ , such that for any $k \in \mathbb{N}$, $\mathcal{H}(\nu_k | \pi) \le C_{\nu_0}(e^{-c_1\delta k} + \delta^2)$. where \mathcal{H} denotes the relative entropy or Kullback– Leibler divergence. Note that the Pinsker inequality then implies the same type of bounds for the total variation distance, but with a second term of order δ .

Quantitative bounds for splitting schemes of the kinetic Langevin diffusion are scarcer up to our knowledge. Higher order quantitative bounds are established in [42] in this context for the case when the potential U is convex. On the other hand, regarding unadjusted HMC, combining results from [6] and [18] implies that, for a fixed integration time T > 0, there exists $c_1 > 0$ (depending on the parameters of the dynamics only through T) such that, for any initial distribution ν_0 , $k \in \mathbb{N}$, $\mathbf{W}_1(\tilde{\nu}_k, \pi) \leq C_{\nu_0}(e^{-c_1k} + \delta^2)$, where $C_{\nu_0} \geq 0$ is a constant, $\tilde{\nu}_k$ is the distribution of the k-th iterates of unadjusted HMC starting from ν_0 , and \mathbf{W}_1 denotes the Wasserstein distance of order 1. This result once again highlights the improved accuracy of the leapfrog integrator. Finally, the analysis of gHMC in the Wasserstein distance has been conducted in [25], but only for the strongly convex scenario.

The main contribution of this paper is to extend and generalize to the non-convex scenario the results we just mentioned and to analyze gHMC. In particular, we show that gHMC achieves a higher order accuracy than traditional Euler schemes in relative entropy, under the condition that π satisfies a log-Sobolev inequality, and additional relatively mild assumptions on the potential energy function U. Roughly speaking, we establish a bound of the form $\mathcal{H}(\nu_0 \mathbf{P}^k | \mu) \leq C_{\nu_0} e^{-c_1 k} + C_2 \delta^4$ for some explicit constants $C_{\nu_0}, C_2 \geq 0$ and $c_1 > 0$. Our approach is based on a perturbative argument of the modified entropy approach initiated in [49], and more precisely to its recent discrete-time variation in [41] for idealized gHMC. From our main result, we derive bounds on the total number of iterations k and a step size δ to achieve $\mathcal{H}(\nu_0 \mathbf{P}^k | \mu) \leq \varepsilon$ for some $\varepsilon > 0$, where two cases can be distinguished: in the general case, we obtain a bound on the number of gradient computations of order $\mathcal{O}(d/\varepsilon^{1/4})$; while in a weakly interacting mean-field regime, an improved bound of order $\mathcal{O}((d/\varepsilon)^{1/4})$ can be achieved.

The paper is organized as follows. The family of (unadjusted) generalized Hamiltonian Monte Carlo samplers is introduced in Section 2.1. Our main assumptions and results are stated in Section 2.2. They are discussed and compared to previous works in Section 2.3. Section 3 is devoted to the proof of Theorem 1. Other more technical proofs are postponed to Appendices.

Notation. We denote by |x| the Euclidean norm of $x \in \mathbb{R}^d$. We denote by $C^k(\mathbb{R}^d)$ the set of functions from \mathbb{R}^d to \mathbb{R} with continuous derivatives up to order k. For $f \in C^k(\mathbb{R}^d)$, ∇f stands for the gradient of f and $D^k f$ for the k-th derivative of f. In addition, $|D^k f(x)|$ is the multilinear operator norm of $D^k f(x)$ with respect to the Euclidean norm on \mathbb{R}^d and $||D^k f||_{\infty} = \sup_{x \in \mathbb{R}^d} |D^k f(x)|$. For a $d \times d$ matrix \mathbf{A} , $|\mathbf{A}|$ stands for the operator norm on \mathbb{R}^d and $||D^k f||_{\infty} = \sup_{x \in \mathbb{R}^d} |D^k f(x)|$. For a $d \times d$ matrix \mathbf{A} , $|\mathbf{A}|$ stands for the operator norm on \mathbb{R}^d we with respect to the Euclidean norm. For a differentiable map $\Phi : \mathbb{R}^d \to \mathbb{R}^p$ and $x \in \mathbb{R}^d$ we write $\nabla \Phi(x) = (\partial_{x_i} \Phi_j(x))_{i \in [\![1,d]\!], j \in [\![1,p]\!]}$ (where i stands for the line and j for the column). This notation is such that $\nabla (\Phi \circ \Psi) = \nabla \Psi \nabla \Phi \circ \Psi$ and is consistent in the case p = 1 (since $\nabla \Phi$ is in that case the gradient of Φ). Notice that $\nabla \Phi$ is the *transpose* of what is most commonly named the Jacobian matrix of Φ .

Leb stands for the Lebesgue measure, $N(0, I_d)$ for the *d*-dimensional standard Gaussian distribution and $\mathcal{P}(\mathbb{R}^d)$ for the set of distribution of \mathbb{R}^d endowed with its Borel σ -field denoted by $\mathcal{B}(\mathbb{R}^d)$. We write $\nu \ll \mu$ when ν is absolutely continuous with respect to μ . We define the relative entropy (or Kullback-Leibler [KL-] divergence) and the Fisher information of $\nu_1 \in \mathcal{P}(\mathbb{R}^d)$ with respect to $\nu_2 \in \mathcal{P}(\mathbb{R}^d)$ respectively as

$$\mathcal{H}(\nu_1|\nu_2) = \begin{cases} \int \ln\left(\frac{\mathrm{d}\nu_1}{\mathrm{d}\nu_2}\right) \mathrm{d}\nu_2 & \text{if } \nu_1 \ll \nu_2 \\ +\infty & \text{otherwise} \end{cases}, \quad \mathcal{I}(\nu_1|\nu_2) = \begin{cases} \int \left|\nabla \ln\left(\frac{\mathrm{d}\nu_1}{\mathrm{d}\nu_2}\right)\right|^2 \mathrm{d}\nu_1 & \text{if } \nu_1 \ll \nu_2 \\ +\infty & \text{otherwise} \end{cases}$$

where for a measurable function $f : \mathbb{R}^d \to \mathbb{R}, |\nabla f|$ is defined for any $z \in \mathbb{R}^d$ as

$$|\nabla f|(z) = \lim_{r \downarrow 0} \sup \left\{ \frac{|f(z) - f(y)|}{|z - y|} : y \in \mathbb{R}^d, \ 0 < |y - z| \le r \right\}.$$

If f is continuously differentiable, note that $|\nabla f|(z)$ is simply the norm of $\nabla f(z)$ and therefore our notation is consistent.

We write $[\![i, j]\!]$ the integer interval $\{i, \ldots, j\}$ for $i, j \in \mathbb{Z}$.

2 Non-asymptotic bounds for splitting schemes for Hamiltonian type dynamics

2.1 Splitting schemes for Hamiltonian type dynamics – gHMC

Recall that we assume that the target distribution π admits a positive density with respect to the Lebesgue measure $\pi \propto \exp(-U)$. In addition, we suppose the following condition on U.

H1. The potential $U \in C^4(\mathbb{R}^d)$ is such that $e^{-U} \in L^1(\mathbb{R}^d)$, and there exists L > 0 such that $|\nabla^2 U(x)| \leq L$ for any $x \in \mathbb{R}^d$.

We now introduce more formally the two continuous dynamics that are considered in this work: Hamiltonian dynamics and kinetic (also referred to as underdamped) Langevin dynamics. As previously mentioned, these two dynamics leave the extended target distribution $\mu = \pi \otimes N(0, I_d)$ invariant.

Ideal Hamiltonian Monte Carlo. The Hamiltonian dynamics associated with the potential U defines the differential flow $(\psi_t)_{t\geq 0}$ for any $t \in \mathbb{R}_+$ and $x, v \in \mathbb{R}^d$ as $\psi_t(x, v) = (x_t, v_t)$, where $(x_t, v_t)_{t\geq 0}$ is the solution of the Hamiltonian differential equation:

$$\partial_t x_t = v_t$$
, $\partial_t v_t = -\nabla U(x_t)$, with $(x_0, v_0) = (x, v)$. (1)

As discussed in [43] for instance, the flow $(\psi_t)_{t\geq 0}$ preserves μ , *i.e.*, if (X_0, V_0) has distribution μ then so does $\psi_t(X_0, V_0)$ for all $t \geq 0$. However, the trajectory $(\psi_t(X_0, V_0))_{t\geq 0}$ is of course not ergodic for μ as, starting from any fixed initial conditions $x_0, v_0 \in \mathbb{R}^d$, it remains in the corresponding level set $\{(x, v) \in \mathbb{R}^{2d} : H(x, v) = H(x_0, v_0)\}$ of the Hamiltonian function H defined by

$$H(x,v) = U(x) + \frac{1}{2} |v|^2$$
.

To address this issue, one can add a velocity randomization (or refreshment) at deterministic times multiple of T > 0. To formalize this procedure, consider the Markov operator D_{η} given, for $\eta \in [0, 1)$ and any measurable and bounded function $f : \mathbb{R}^{2d} \to \mathbb{R}$, by

$$D_{\eta}f(x,v) = \int_{\mathbb{R}^d} f\left(x, \eta v + \sqrt{1-\eta^2}g\right) \varphi(g) dg ,$$

where φ stands for the density with respect to Leb of N(0, I_d). Then, under mild assumptions, the resulting kernel

$$\mathbf{K}_{\eta,\mathbf{T}}f(x,v) = \mathbf{D}_{\eta}[f \circ \psi_{\mathbf{T}}](x,v) , \qquad (2)$$

is ergodic with respect to μ . Recently, [41] used hypocoercivity techniques from [49] to show exponential convergence of $K_{\eta,T}$ in a modified entropy with respect to μ .

Underdamped Langevin dynamics. The second process which shares some important features with Hamiltonian dynamics is the underdamped Langevin diffusion:

$$d\mathbf{X}_s = \mathbf{V}_s \, \mathrm{d}s \,, \qquad \mathrm{d}\mathbf{V}_s = -\nabla U(\mathbf{X}_s) \, \mathrm{d}s - \gamma \mathbf{V}_s \, \mathrm{d}s + \sqrt{2\gamma} \, \mathrm{d}B_s \,, \tag{3}$$

where $(B_s)_{s\geq 0}$ is a *d*-dimensional Brownian motion and $\gamma > 0$ is a damping parameter. It can easily be shown that under mild assumptions on U, the Hamiltonian dynamics with a suitably scaled refreshment weakly converges to (3): for any $s \geq 0$, $x, v \in \mathbb{R}^d$ and smooth and bounded function $f : \mathbb{R}^{2d} \to \mathbb{R}$, it holds $\lim_{\delta \to 0} K_{\eta_{\delta},\delta}^{[s/\delta]} f(x,v) = \mathbf{P}_s f(x,v)$ where

$$\eta_{\delta} = \mathrm{e}^{-\delta\gamma} \,, \tag{4}$$

and $(\mathbf{P}_s)_{s\geq 0}$ is the Markov semigroup associated with (3).

Discretizations of underdamped Langevin and ideal HMC. We now present a family of splitting schemes which encompasses discretizations for both Hamiltonian and underdamped Langevin dynamics. This family of algorithms will be referred to as generalized Hamiltonian Monte Carlo (gHMC). It is based on the Verlet discretization of (1) defined at time $k\delta$, for $k \in \mathbb{N}$ and a given stepsize $\delta > 0$, by $(x_k, v_k) = \Phi_{\delta}^k(x_0, v_0)$ where $\Phi_{\delta}^k = \Phi_{\delta}^{k-1} \circ \Phi_{\delta}$, with Φ_{δ}^0 is the identity function and

$$\Phi_{\delta} = \Phi_{\delta/2}^{(v)} \circ \Phi_{\delta}^{(x)} \circ \Phi_{\delta/2}^{(v)} , \quad \Phi_{\delta/2}^{(v)}(x,v) = (x, v - (\delta/2)\nabla U(x)) , \quad \Phi_{\delta}^{(x)}(x,v) = (x + \delta v, v) .$$

It is well-known that the integrator Φ_{δ} is symplectic and reversible (see [30]). Its inverse reads

$$\Phi_{\delta}^{-1} = \mathbf{R} \circ \Phi_{\delta} \circ \mathbf{R} , \qquad \mathbf{R}(x, v) = (x, -v) .$$
(5)

The gHMC algorithm then consists in the composition of a (possibly partial) velocity refreshment with a K-step Verlet scheme. Setting $\boldsymbol{\omega} = (K, \delta, \eta)$, this corresponds to the inexact Markov chain Monte Carlo (MCMC) method with Markov kernel specified by

$$\mathbf{P}_{\boldsymbol{\omega}} = \mathbf{D}_{\eta} \mathbf{V}_{\delta}^{K} ,$$

where V_{δ} corresponds to the deterministic kernel

$$V_{\delta}((x,v),\cdot) = \delta_{\Phi_{\delta}(x,v)}(\cdot)$$
, for $x, v \in \mathbb{R}^d$.

For $\eta = 0$ and $K\delta = T$ for an integration time T > 0, the kernel P_{ω} corresponds to the usual unadjusted Hamiltonian Monte Carlo algorithm, which is a discretized version of the ideal Hamiltonian dynamics $K_{T,\eta}$ defined by (2). On the other hand, with K = 1 and the damping parameter η_{δ} defined in (4), the kernel P_{ω} corresponds to a splitting scheme whose invariant probability measure is correct at second order in δ (this corresponds in fact to the so-called GLA scheme [7]; see also [34, 36] for other splitting schemes). Alternatively, taking $\eta = 0$ and K = 1 leads to the Euler–Maruyama scheme with step size $\delta^2/2$ for the overdamped Langevin process.

Normalization and rescaling. When H1 is satisfied, we introduce

$$\mathbf{T} = \delta K \sqrt{L} , \qquad \gamma = (1 - \eta) / \mathbf{T} , \qquad (6)$$

which are respectively, up to a suitable rescaling, the physical time of integration of the Hamiltonian dynamics by the Verlet integrator, and the strength of the damping. As discussed in [25], upon rescaling the process so that L = 1 (see the discussion in Section 2.2.1), the time needed for the process to forget its initial velocity is of order $1/\gamma$, so that this quantity is also the typical distance covered by one of the coordinates of the process in a single ballistic run in flat parts of the space.

2.2 Main result

We present in this section our main result, under additional assumptions regarding π and U (see Section 2.2.1). We next provide in Section 2.2.2 sufficient conditions under which these additional assumptions are satisfied.

2.2.1 Uniform convergence in time

Additional assumptions. Our first assumption is that π satisfies a log-Sobolev inequality with constant $C_{\text{LS}} \ge 0$, *i.e.*

$$\forall \nu \in \mathcal{P}(\mathbb{R}^d), \qquad \mathcal{H}(\nu|\pi) \leqslant C_{\mathrm{LS}} \mathcal{I}(\nu|\pi) . \tag{7}$$

Various sufficient conditions for this inequality to hold have been derived, see [4]. It holds for instance when U is uniformly convex outside some compact set. Various estimates are available for C_{LS} , e.g. in mean-field, convex or low-temperature cases, which can capture more specific information on the target π than uniform bounds on the curvature $\langle x - y, \nabla U(x) - \nabla U(y) \rangle / |x - y|^2$ that are used in direct coupling methods. Notably, in non-convex cases, for a fixed dimension d, up to polynomial terms in the other parameters, the log-Sobolev constant is of order e^{c_*} where c_* is the so-called critical height of U (see [41, Section 5] and references within).

Part of our analysis consists in controlling some numerical errors of the leapfrog integrator, *i.e.*, for example differences of the Hamiltonian function evaluated at the dynamics at times 0 and s > 0 starting from the k-th iterates of the gHMC chain. To this end, we need additional regularity conditions on U. More specifically, we require uniform bounds on the third and fourth derivatives of U, as made precise in the following condition.

H2. There exist two norms N_3 and N_4 on \mathbb{R}^d such that, for all $x, y, z \in \mathbb{R}^d$,

$$\left(\nabla^2 U(x+y) - \nabla^2 U(x)\right) z \bigg| \leqslant \mathcal{N}_3(y) \mathcal{N}_3(z) , \qquad (8)$$

$$\left|\nabla U(x+y) - \nabla U(x) - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x+y)\right) y\right| \leq \mathcal{N}_4^3(y) \,. \tag{9}$$

Using a third order Taylor expansion of ∇U , it can be shown that (9) follows from a uniform bound on the third and fourth derivatives of U (see Proposition 2 below). H2 may be relaxed, i.e, if one assumes only bounded third derivatives for U, our main result still holds but with worse dependencies with respect to the step size, see Remark 15 below.

Bounding the left-hand sides in (8) and (9) with respect to the Euclidean distance on \mathbb{R}^d may lead to sub-optimal dependencies with respect to the dimension d in Theorem 1 below (for instance when U is separable, see Theorem 3 below), which is why we allow in H2 some flexibility in the choice of the norms. A specific choice of norms is given in Proposition 2, and the relevance of this choice for the dimensionality dependence is discussed in Remark 4.

As made precise in Section 3.4, the numerical errors that need to be controlled at a point $(x, v) \in \mathbb{R}^{2d}$ can be bounded by the function

$$\mathbf{M}(x,v) = L^2 |v|^2 + L^2 |\nabla U(x)|^2 + \mathbf{N}_3^4(v) + \mathbf{N}_3^4(\nabla U(x)) + \mathbf{N}_4^6(v) + \mathbf{N}_4^6(\nabla U(x)).$$
(10)

In order to control the expectation of these errors, one needs uniform moment bounds for the Markov kernel P_{ω} . These estimates typically follow from a Lyapunov condition, but for clarity we postpone this analysis to Theorem 3 below and, for the time being, state it as the next assumption.

H 3. There exist $\rho, C_1, C_2 > 0$ and $\mathfrak{W} : \mathbb{R}^{2d} \to \mathbb{R}_+$ such that, for all $(x, v) \in \mathbb{R}^{2d}$, $k \in \{0, \ldots, K\}$ and $n \in \mathbb{N}$,

$$\mathbf{P}^{n}_{\boldsymbol{\omega}} \mathbf{D}_{\eta} \mathbf{V}^{k}_{\delta} \mathbf{M}(x, v) \leqslant C_{1} \mathbf{e}^{-\rho n \mathsf{T}} \mathfrak{W}(x, v) + C_{2},$$

where T is defined in (6).

In practice, this can be established by designing a suitable Lyapunov function \mathfrak{W} such that $\mathbf{M} \leq C_1 \mathfrak{W}$ and

$$\mathbf{P}_{\boldsymbol{\omega}} \mathfrak{W} \leqslant \mathbf{e}^{-\rho \mathsf{T}} \mathfrak{W} + C_2' \mathsf{T} , \qquad (11)$$

for some constants $C_1, C'_2, \rho > 0$. As we will see in Theorem 3, the constants C_1, C_2, ρ typically only depend on the parameters $\boldsymbol{\omega}$ through γ defined in (6). In particular, they are uniform over all sufficiently small step-sizes δ . For the continuous-time kinetic Langevin diffusion and its discretizations, such Lyapunov functions have been designed in [39, 47, 19] under coercitivity conditions on U. We show in Appendix B that appropriate modifications of these Lyapunov functions can be used to establish a drift condition (11) for $P_{\boldsymbol{\omega}}$.

Main convergence result. We are now in position to state our main result. For the sake of clarity, we only consider the case where L = 1 in H1, the general case being obtained by rescaling, as discussed at the end of this section.

Theorem 1. Assume that H1-H2 hold with L = 1, and consider $\boldsymbol{\omega} = (K, \delta, \eta)$ with $K \in \mathbb{N}^*$, $\delta > 0$ and $\eta \in [0, 1)$ such that H3 holds and $T = K\delta \leq 1/10$. Furthermore, assume that π satisfies the log-Sobolev inequality (7). Consider an initial condition ν_0 such that $\mathcal{I}(\nu_0|\mu) < +\infty$ and $\int_{\mathbb{R}^{2d}} |z|^p d\nu_0(z) < +\infty$ for any p > 0. Then, for any $n \in \mathbb{N}$,

$$\mathcal{H}\left(\nu_{0}\mathcal{P}_{\boldsymbol{\omega}}^{n}|\boldsymbol{\mu}\right) \leqslant (1+\kappa \mathsf{T})^{-n} \left[\mathcal{H}(\nu_{0}|\boldsymbol{\mu})+2a\,\mathcal{I}(\nu_{0}|\boldsymbol{\mu})\right] \\ +\delta^{4}M\left[n\mathsf{T}\theta^{n\mathsf{T}}C_{1}\nu_{0}(\mathfrak{W})+C_{2}(\kappa^{-1}+\mathsf{T})\right],\tag{12}$$

where $\theta = \max(e^{-\rho}, (1 + \kappa T)^{-1/T})$ and

$$a = \frac{\gamma}{7 + 3(\gamma + 3)^2}, \qquad \kappa = \frac{a}{3\max(C_{\rm LS}, 1) + 6a}, \qquad M = 9 + \frac{1}{a}.$$
 (13)

The first term on the right hand side of (12) encodes the long-time convergence of the idealized HMC chain [41] (and goes to 0 as $n \to \infty$). The Fisher term $\mathcal{I}(\nu_0|\mu)$ appearing there is due to the fact Theorem 1 is proven using a modified entropy (defined in (21)) instead of \mathcal{H} . The second line of (12), of order δ^4 , corresponds to the numerical error due to the Verlet integration, which is decomposed into two terms. The first one comes from the fact that the initial condition is not invariant under P_{ω} and highlights the exponential forgetting of this initial condition since $\theta < 1$. The second term involving C_2 accounts for the numerical error at stationarity, *i.e.* the bias between the invariant probability measure of the discrete process and the reference probability measure μ .

Since, in the case L = 1, the physical integration time of the Hamiltonian dynamics is T, the total physical time of the simulation after *n* iterations is *n*T. The constant κ is therefore the convergence rate per unit physical time. In view of (13), κ depends on the parameters ω only through the friction γ , with a dependency of order min (γ, γ^{-1}) similarly to what is obtained for the continuous-time Langevin diffusion (see for instance [17, 28, 33]). From this observation, it follows that the convergence per gradient computation is $\kappa T/K = \kappa \delta$. At the expense of more involved expressions, we provide an improved version of Theorem 1 in Theorem 5 below. In particular, we provide a slightly sharper value of κ , which is optimal in the overdamped case since, in the continuous time limit $\delta \to 0$, one obtains (see Remark 6)

$$\mathcal{H}(\hat{\nu}_0 \mathbf{Q}_s | \pi) \leqslant e^{-s/C_{\rm LS}} \mathcal{H}(\hat{\nu}_0 | \pi) \,, \tag{14}$$

where $(Q_t)_{t\geq 0}$ is the semi-group associated to the overdamped Langevin diffusion. This shows that our result is sharp since it *implies* that π satisfies a log-Sobolev inequality with constant $C_{\rm LS}$ (see e.g., [4, Theorem 5.2.1]).

Rescaling. As mentioned above, following [41, Section 1.3], it is not restrictive to assume that L = 1 in **H1**. Indeed, if $Z_n = (X_n, V_n)$ is an unadjusted gHMC chain with potential energy function U and parameters K, δ, η , then $(\sqrt{L}X_n, V_n)$ is an unadjusted gHMC chain associated with the potential energy function $\tilde{U}(x) = U(x/\sqrt{L})$ (so that $\nabla \tilde{U}$ is 1-Lipschitz under **H1**) and parameters $K, \delta\sqrt{L}, \eta$. Moreover, if π satisfies a log-Sobolev inequality with constant C_{LS} , then the change of variable $x \mapsto x/\sqrt{L}$ implies that the rescaled target measure $\tilde{\pi} \propto e^{-\tilde{U}}$ satisfies a log-Sobolev inequality with constant LC_{LS} . Similarly, using the same change of variable, the relative entropy is invariant by scaling, namely, if (X, V)has distribution $\nu \in \mathcal{P}(\mathbb{R}^{2d})$ and $\tilde{\nu}$ stands for the law of $(\sqrt{L}X, V)$, then $\mathcal{H}(\nu|\mu) = \mathcal{H}(\tilde{\nu}|\tilde{\mu})$, where $\tilde{\mu} = \tilde{\pi} \otimes N(0, I_d)$; while the Fisher information satisfies

$$\mathcal{I}\left(\tilde{\nu}|\tilde{\mu}\right) = \int_{\mathbb{R}^{2d}} \frac{L^{-1} |\nabla_x h|^2 + |\nabla_v h|^2}{h} \mathrm{d}\mu \,, \text{ where } h = \mathrm{d}\nu/\mathrm{d}\mu \,.$$

2.2.2 Sufficient conditions for H₂ and H₃

We now provide practical conditions on U to establish **H2** and **H3**. We assume that $d = d_0 q$ for some $d_0, q \ge 1$ and, decomposing $x = (x_1, \ldots, x_{d_0}) \in \mathbb{R}^d = (\mathbb{R}^q)^{d_0}$, that U is of the following form:

$$U(x) = \sum_{i=1}^{d_0} U^{(q)}(x_i) + \frac{\epsilon}{d_0} \sum_{i,j=1}^{d_0} W^{(q)}(x_i - x_j) , \qquad (15)$$

where $U^{(q)}$ and $W^{(q)}$ are potentials defined on \mathbb{R}^q , and $\epsilon > 0$. Such mean field potentials are commonly encountered in various applications (see e.g., [10]). Furthermore, in line with the approach taken in [8], we consider this formulation as a representative example to demonstrate the dimension dependence in weakly correlated cases. Let us however emphasize that (15) is not a restrictive setting since it always holds with the choice $d_0 = 1$, $U^{(q)} = U$, and $W^{(q)} = 0$. However, as discussed below, this mean-field formulation allows us to precisely determine the dependence of C_1 and C_2 in **H** 3 on the dimension d, in particular by distinguishing the roles of d_0 and q. For conditions under which π satisfies a log-Sobolev inequality with a constant C_{LS} independent of d_0 for U given by (15) (provided ϵ is sufficiently small), see [29, Theorem 8]. First, as proved in Appendix A, we can readily obtain the following.

Proposition 2. Assume that $d = d_0 q$ for some $d_0, q \ge 1$ and that U is of the form (15) where $U^{(q)}, W^{(q)} \in C^4(\mathbb{R}^q)$ have uniformly bounded third and fourth derivatives. Then H^2 holds with

$$N_3(x) = L_3 \left(\sum_{i=1}^{d_0} |x_i|^4 \right)^{1/4}, \qquad N_4(x) = L_4 \left(\sum_{i=1}^{d_0} |x_i|^6 \right)^{1/6},$$

where

$$\begin{split} L_3^4 &= (1+\epsilon) \left\| \mathbf{D}^3 U^{(q)} \right\|_{\infty}^2 + 16\epsilon (1+\epsilon) \left\| \mathbf{D}^3 W^{(q)} \right\|_{\infty}^2 \,, \\ L_4^6 &= \left(\frac{1+\epsilon}{144} \left\| \mathbf{D}^4 U^{(q)} \right\|_{\infty}^2 + \frac{4\epsilon (1+\epsilon)}{9} \left\| \mathbf{D}^4 W^{(q)} \right\|_{\infty}^2 \right) q \end{split}$$

As a consequence, assuming furthermore that $\nabla U^{(q)}(0) = 0 = \nabla W^{(q)}(0)$, the function $\mathbf{M}(x, v)$ defined in (10) is bounded as

$$\mathbf{M}(x,v) \leqslant \sum_{i=1}^{d_0} \sum_{\ell \in \{2,4,6\}} L_{\ell}^{\ell}(r_{\ell}|x_i|^{\ell} + |v_i|^{\ell}),$$

with $r_2 = 1$, $r_\ell = 2^{\ell-1} \|\nabla^2 U^{(q)}\|_{\infty}^{\ell} + 2^{3\ell} \epsilon^{\ell} \|\nabla^2 W^{(q)}\|_{\infty}^{\ell}$ for $\ell \in \{4, 6\}$ and $L_2 = L$.

To establish moment bounds, we consider the following assumption.

H 4. The dimension is $d = d_0 q$ for some $d_0, q \ge 1$ and U is of the form (15) where $U^{(q)}, W^{(q)} \in C^2(\mathbb{R}^q)$ have uniformly bounded second derivatives and are such that $\nabla U^{(q)}(0) = \nabla W^{(q)}(0) = 0$. In addition, there exist m > 0 and $M \ge 0$ such that for any $x_1 \in \mathbb{R}^q$,

$$\left\langle x_1, \nabla U^{(q)}(x_1) \right\rangle \ge \mathbf{m} \left| x_1 \right|^2 - \mathbf{M} .$$
 (16)

Note that this assumption is more general than assuming that $U^{(q)}$ is strongly convex outside a compact set. For instance, a potential such as $|x_1|^2 + C \cos |x_1|$ satisfies (16) for any $C \in \mathbb{R}$, but is not strongly convex when $|C| \ge 2$.

Theorem 3. Suppose that H_4 is satisfied, and define

$$\mathfrak{W}(x,v) = \sum_{\ell \in \{2,4,6\}} \sum_{i=1}^{d_0} \left(|x_i|^\ell + |v_i|^\ell \right) \,. \tag{17}$$

For any $\gamma_0 > 0$, there exist $\overline{T}, \overline{\delta}, \overline{C}_1, \overline{C}_2, \overline{\epsilon}, \rho > 0$, which depend on $\|\nabla^2 U^{(q)}\|_{\infty}$, $\|\nabla^2 W^{(q)}\|_{\infty}$, m, M, γ_0 (but not on q, d_0), such that the following holds: for any $\epsilon \in (0, \overline{\epsilon}]$ and $\boldsymbol{\omega} = (K, \delta, \eta)$ with $\delta \in (0, \overline{\delta}]$, $\eta \in [0, 1)$ such that $(1 - \eta)/(K\delta) = \gamma_0$ and $\delta K \leq \overline{T}$, it holds, for any $(x, v) \in \mathbb{R}^{2d}$, $k \in \{0, \ldots, K\}$ and $n \in \mathbb{N}$,

$$\mathbf{P}^{n}_{\omega} \mathbf{D}_{\eta} \mathbf{V}^{k}_{\delta} \mathfrak{W}(x, v) \leqslant \overline{C}_{1} \mathrm{e}^{-\rho n K \delta} \mathfrak{W}(x, v) + \overline{C}_{2} d_{0} q^{3}.$$

Explicit expressions for $\overline{C}_1, \overline{C}_2, \overline{\epsilon}, \rho$ are given in Appendix B, where a more detailed result, Theorem 18, is stated and proven. As explained in Appendix B, Theorem 3 is a straightforward corollary of Theorem 18.

Remark 4. In order to determine more precisely the dimension dependence of C_1, C_2 in **H**³, obtained for \mathfrak{W} given by (17) by combining Proposition 2 with Theorem 3, we treat $\|D^k U^{(q)}\|_{\infty}$ and $\|D^k W^{(q)}\|_{\infty}$ for $k \in \{2,3,4\}$ as constants independent from q. This motivates defining $R = q^{-1} \max\{L_{\ell}^{\ell} \max(r_{\ell}, 1) : \ell \in \{2,4,6\}\}$, which can be considered to be uniform in d_0 and q in this context, so that the expressions $C_1 = R\overline{C}_1q$, $C_2 = R\overline{C}_2d_0q^4$ capture the dimension dependence of C_1, C_2 in terms of q, d_0 .

2.3 Discussion and related works

Complexity bounds. Regarding the dependence in the dimension of the constants appearing in H3, we distinguish two cases in view of Remark 4. In the first one, which we call the weakly interacting case, we consider that C_2 and $\nu_0(\mathfrak{W})$ in (12) are $\mathcal{O}(d)$. This corresponds in Remark 4 to the case where q is fixed, so that d is proportional to d_0 . In the second case, which we call the general case, we consider that C_2 and $\nu_0(\mathfrak{W})$ are $\mathcal{O}(d^4)$. This corresponds in Remark 4 to the case $d_0 = 1$ and d = q. In both settings, we consider that L, C_1 and γ are constants independent of the dimension, and that $\mathcal{H}(\nu_0|\mu)$ and $\mathcal{I}(\nu_0|\mu)$ are $\mathcal{O}(d)$ (see e.g. [48, Lemma 1]). Moreover, the convergence rate κ is of order $1/C_{\text{LS}}$, with C_{LS} large, so that $(1 + \kappa T)^{-n}$ is of order $e^{-\alpha n K \delta/C_{\text{LS}}}$ for some constant $\alpha > 0$. In this context, Theorem 1 gives the following complexity bounds for a given error tolerance $\varepsilon > 0$ (in relative entropy):

- in the general case, upon choosing $\delta = \mathcal{O}(\varepsilon^{1/4}d^{-1})$ (so that the second term on the right hand side of (12) is of order ε), we get that $\mathcal{H}(\nu_0 \mathbf{P}^n_{\omega}|\mu) \leq \varepsilon$ after a number $nK = \mathcal{O}(C_{\mathrm{LS}}d\varepsilon^{-1/4}\ln(d\varepsilon^{-1}))$ of computations of the gradient ∇U .
- in the weakly interacting case, a similar reasoning suggests choosing $\delta = \mathcal{O}(\varepsilon^{1/4} d^{-1/4})$, in which cas $\mathcal{H}(\nu_0 \mathbf{P}^n_{\boldsymbol{\omega}} | \mu) \leq \varepsilon$ after a number $nK = \mathcal{O}(C_{\mathrm{LS}}(d/\varepsilon)^{1/4} \ln(d\varepsilon^{-1}))$ of computations of the gradient ∇U .

Note that, in both cases, the Pinsker and Talagrand inequalities respectively imply that $\|\mu - \nu_0 \mathbf{P}^n_{\omega}\|_{\text{TV}}$ and $\mathbf{W}_2(\mu, \nu_0 \mathbf{P}^n_{\omega})$ are of order $\mathcal{O}(\varepsilon^{1/2})$, where $\|\cdot\|_{\text{TV}}$ and \mathbf{W}_2 respectively denote the total variation distance and Wasserstein distance of order 2. In view of this, the dependence of the number of gradient evaluation on $\varepsilon^{1/2}$, namely $nK = \mathcal{O}(\varepsilon^{-1/4})$ (omitting the logarithmic term), highlights that gMHC is a second-order splitting scheme. This is consistent with existing weak error estimates; see [1] and references therein.

Error estimates on the invariant probability measure. For a fixed set of parameters $\boldsymbol{\omega}$, if $\nu_0 P^n_{\boldsymbol{\omega}}$ converges weakly to a stationary distribution $\mu_{\boldsymbol{\omega}}$, Theorem 1 implies that $\|\boldsymbol{\mu} - \boldsymbol{\mu}_{\boldsymbol{\omega}}\|_{\text{TV}}$ and $\mathbf{W}_2(\boldsymbol{\mu}, \boldsymbol{\mu}_{\boldsymbol{\omega}})$ are both of order $\mathcal{O}(\delta^2)$ (using the lower semicontinuity of the entropy function, see [50, Theorem 29.1]). This is indeed the expected scaling of the bias on the invariant probability measure of the Markov chain, since the latter one is a composition of two evolution operators, namely D_{η} which exactly preserves μ , and V_{δ} which is a second order approximation of the Hamiltonian evolution operator exactly preserving μ .

Comparison with previous works. There is an abundant literature on the convergence of Markov chain Monte Carlo algorithms based on discretization of continuous-time dynamics, see for instance [39, 47, 16, 12, 42, 3, 38]. Making a complete survey and a detailed description of these works is beyond the scope of the paper. Here, we focus on works establishing quantitative bounds for gHMC or some of its particular instances, and convergence in the entropic sense for discretizations of the kinetic Langevin diffusion.

• Entropy methods for discretization of the kinetic Langevin diffusion. An adaptation of the modified entropy strategy from [49] for continuous-time processes has been conducted by [37] for a slight modification of the Euler–Maruyama scheme for the kinetic Langevin process. More precisely, the scheme considered in [37] (introduced in [13] in the machine learning community, and referred to as the Stochastic exponential Euler scheme in [19], where earlier apparitions in the physics literature are mentioned [11, 23, 46]) is given by the solution of

$$\mathrm{d}\mathbf{X}_{t}^{\delta} = \mathbf{V}_{t}^{\delta}\mathrm{d}t , \qquad \mathrm{d}\mathbf{V}_{t}^{\delta} = -\nabla U(\mathbf{X}_{\delta\left[t/\delta\right]}^{\delta})\mathrm{d}t - \gamma \mathbf{V}_{t}^{\delta}\mathrm{d}t + \sqrt{2\gamma}\mathrm{d}B_{t} , \qquad (18)$$

which is similar to (3), except that the force $-\nabla U$ is constant within time intervals $[k\delta, (k+1)\delta)$ for $k \in \mathbb{N}$. Hence, in [37], the approximate entropy dissipation is established through a continuous-time derivation. In contrast, we follow the discrete time computations of [41], which focuses on idealized gHMC, where the exact Hamiltonian dynamics is followed instead of the Verlet integration. In particular, we are not considering in this work the underdamped Langevin diffusion process as a continuous-time reference, since we also cover unadjusted HMC. Moreover, in terms of dependence with respect to the stepsize δ , the schemes of [37] are only of first order (see the discussion in [37, Section 3.3], in particular Equation (14)), and we obtain an improvement from $\sqrt{\varepsilon}$ to $\varepsilon^{1/4}$ in the complexity. Concerning the dimension d, the stepsize in [37] scales as $d^{-1/2}L_H^{-1}$, where L_H is the Lipschitz constant of $\nabla^2 U$ in the Frobenius norm, *i.e.*, U is supposed to satisfy $\|\nabla^2 U(x) - \nabla^2 U(y)\|_F \leq L_H |x-y|$ for all $x, y \in \mathbb{R}^d$. Assuming that L_H is independent of d thus corresponds to the weakly interacting case, while the assumption that the derivatives of U are bounded uniformy in d (in terms of the Euclidean norm) leads to L_H of order \sqrt{d} . We get an improvement from $d^{-1/2}$ to $d^{-1/4}$ in the first case, and the same dependence d^{-1} in the second. Notice that, in order to take advantage of an higher order numerical scheme, we have to assume that U has bounded first four derivatives, in contrast to [37] which only assumes that U has bounded first three derivatives. As discussed in Remark 15, if we consider this second condition on U, a result similar to [37] and with the same dependence on ε and d can be derived. The main takeaway of this observation is that gHMC can achieve better accuracy when U is sufficiently smooth, and does not yield a worst complexity than the discretization (18) considered in [37] under the same conditions. Finally, up to our knowledge, our result is the first one based on entropy methods for unadjusted HMC, and for splitting discretization schemes of the underdamped Langevin diffusion.

- **HMC.** Analyses of HMC for strongly convex potentials U have been conducted in [38, 6]. For non-convex potentials, the works [6, 18, 5, 8] have established several non-asymptotic convergence bounds for position HMC (*i.e.*, $\eta = 0$), both in Wasserstein 1 distance and total variation norm. In the mean-field case in particular, [8, Theorem 10] shows that, after $\mathcal{O}(\sqrt{d_0/\varepsilon'})$ gradient evaluations, the law of the chain is at distance at most ε' to π in terms of the \mathbf{W}_{ℓ_1} Wasserstein distance associated to the ℓ_1 norm $||x - y||_{\ell_1} = \sum_{i=1}^{d_0} |x_i - y_i|$. To compare this result with ours, we can bound this distance by $\sqrt{d_0}$ times the standard \mathbf{W}_2 Wasserstein distance (associated with the Euclidean distance), whose square is bounded by the relative entropy times the log-Sobolev constant of π (which, from the results of [29], is independent of d_0 in the weakly interacting case where ϵ is small enough, which is the regime where this complexity is obtained in [8]). In other words, we should take $\varepsilon' = \sqrt{d_0 \varepsilon}$ in the results of [8] for a fair comparison. In that case, the number of gradient computations required in [8, Theorem 10] is $\mathcal{O}((d_0/\varepsilon)^{1/4})$ in the weakly interacting case, which is the same as our results. The approach in [5, 6, 8] is based on reflection couplings and concave modifications of the distance (building upon the method developed for continuous-time diffusion processes, in particular [21] for the overdamped Langevin process and [22] for the underdamped one). In particular, it does not provide a result for the relative entropy or the \mathbf{W}_2 Wasserstein distance, and the bound on the convergence rate is expressed in terms of global bounds on $\langle x - y, \nabla U(x) - \nabla U(y) \rangle / |x - y|^2$.
- Splitting schemes for kinetic Langevin diffusion for strongly convex potential U. Splitting schemes for Langevin diffusion with strongly convex potential energy functions have been investigated in [42, 35]. Their conclusions on the complexity for the corresponding gHMC method are similar to ours. Finally, the analysis of the general gHMC methodology have been conducted in [25] under the same strong convexity condition. Once again, the same conclusions can be drawn.

As a conclusion, up to our knowledge, none of the existing works cover our results, which give simple second order explicit estimates in relative entropy uniformly over a class of unadjusted gHMC samplers and for non-convex potential energy functions U.

3 Proof of Theorem 1

The proof of Theorem 1 is an easy consequence of a more detailed result given in Section 3.1, stated for a modified entropy function mixing a relative entropy and some Fisher information similar to the one considered for the time continuous Langevin dynamics. Instead of time derivatives, we have to estimate the discrete-time evolution of the two parts of the modified entropy along the two steps V_{δ} and D_{η} of the chain. The technical details of this analysis are postponed to Section 3.2, where this evolution is shown to involve, on the one hand, the Jacobian matrix of the Verlet map and, on the other hand, some numerical error terms (due to the fact μ is not invariant by V_{δ}). The study of these terms is performed respectively in Sections 3.3 and 3.4.

3.1 A more detailed approximate decay result for a modified entropy

Relative density and modified entropy. In the remainder of the present Section 3, for an initial distribution $\nu_0 \in \mathcal{P}(\mathbb{R}^{2d})$ which satisfies the conditions in Theorem 1, parameters $\boldsymbol{\omega} = (K, \delta, \eta)$ and $n \in \mathbb{N}$, we denote by

$$h_n = \frac{\mathrm{d}\left(\nu_0 \mathrm{P}^n_{\omega}\right)}{\mathrm{d}\mu} , \qquad \mathcal{H}(h_n) = \mathcal{H}(\nu_0 \mathrm{P}^n_{\omega}|\mu) , \qquad \mathcal{I}(h_n) = \mathcal{I}(\nu_0 \mathrm{P}^n_{\omega}|\mu) ,$$

where \mathcal{H} and \mathcal{I} denote (with some abuse of notation) the relative entropy and Fisher information with respect to μ respectively. More generally, we denote by $\mathcal{H}(h) = \mathcal{H}(\nu|\mu)$ and $\mathcal{I}(h) = \mathcal{I}(\nu|\mu)$ for any probability density $h : \mathbb{R}^{2d} \to \mathbb{R}$ with respect to μ , so that $\nu = h\mu$ is a probability measure. In addition, for $\tilde{\mathbf{A}} \in \mathbb{R}^{2d \times 2d}$, or possibly a matrix field $z \mapsto \tilde{\mathbf{A}}(z)$ on \mathbb{R}^{2d} , define

$$\mathcal{I}_{\tilde{\mathbf{A}}}(\nu|\mu) = \begin{cases} \int_{\mathbb{R}^{2d}} \left| \tilde{\mathbf{A}} \nabla \ln \left(\frac{\mathrm{d}\nu}{\mathrm{d}\mu} \right) \right|^2 \mathrm{d}\nu & \text{if } \nu \ll \mu, \\ +\infty & \text{otherwise} \end{cases}.$$

Note that if $\nu \ll \mu$ and $h = d\nu/d\mu$,

$$\mathcal{I}_{\tilde{\mathbf{A}}}(\nu|\mu) = \int_{\mathbb{R}^{2d}} \frac{|\tilde{\mathbf{A}}\nabla h|^2}{h} \,\mathrm{d}\mu \;. \tag{19}$$

Following the convention above, we write $\mathcal{I}_{\tilde{\mathbf{A}}}(h) = \mathcal{I}_{\tilde{\mathbf{A}}}(\nu|\mu)$, for any probability density $h : \mathbb{R}^{2d} \to \mathbb{R}$ with respect to μ with $\nu = h\mu$.

The main result of this section, from which we deduce Theorem 1, establishes the dissipation in time of a suitable modified entropy up to some numerical error. More precisely, we consider, for $\nu \in \mathcal{P}(\mathbb{R}^{2d})$ and $\tilde{a} > 0$,

$$\mathcal{L}_{\tilde{a}}(\nu|\mu) = \mathcal{H}(h) + \tilde{a}\mathcal{I}_{\mathbf{A}}(h) , \qquad \mathbf{A} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1\\ 1 & 1 \end{pmatrix} .$$
(20)

When $\nu \ll \mu$ with $h = d\nu/d\mu$,

$$\mathcal{L}_{\tilde{a}}(\nu|\mu) = \int_{\mathbb{R}^{2d}} h \ln(h) \,\mathrm{d}\mu + \tilde{a} \int_{\mathbb{R}^{2d}} \frac{|\nabla_x h + \nabla_v h|^2}{h} \,\mathrm{d}\mu \,, \tag{21}$$

and, following the same convention as previously, we write $\mathcal{L}_{\tilde{a}}(h) = \mathcal{L}_{\tilde{a}}(\nu|\mu)$, for any probability density $h : \mathbb{R}^{2d} \to \mathbb{R}$ with respect to μ with $\nu = h\mu$. With this choice, $\mathcal{L}_{\tilde{a}}$ involves a mixed derivative term $\langle \nabla_x h, \nabla_v h \rangle$ in the Fisher part, which has been crucially used in hypocoercive studies of the continuous-time kinetic Langevin process [47, 32, 49]. More specifically, (21) is exactly the modified entropy introduced in continuous-time settings in [49, Section 6].

Precise statement of the convergence result. We are now ready to state the main result of this section.

Theorem 5. Assume H1-H2 with L = 1, and consider $\boldsymbol{\omega} = (K, \delta, \eta)$ with $K \in \mathbb{N}^*$, $\delta > 0$ and $\eta \in (0, 1)$ such that H3 holds and $\mathbb{T} = K\delta \leq 1/10$. Furthermore, assume that π satisfies the log-Sobolev inequality (7). Consider an initial condition ν_0 such that $\mathcal{I}(\nu_0|\mu) < +\infty$ and for all p > 0, $\int_{\mathbb{R}^{2d}} |z|^p d\nu_0(z) < +\infty$. Then, for any $n \in \mathbb{N}$,

$$(1 + \kappa \mathsf{T}) \mathcal{L}_a(h_{n+1}) \leq \mathcal{L}_a(h_n) + \mathsf{T}\delta^4 M \left(C_1 \mathrm{e}^{-\rho n t} \nu_0 \left(\mathfrak{W} \right) + C_2 \right) , \qquad (22)$$

where a, κ, M are given in (13).

Moreover, a refined estimate can be obtained as follows: define

$$m_1 = 2 - 3T$$
, $m_2 = \frac{\gamma}{\eta} + 2 + 4T$, $m_3 = \frac{\gamma(1+\eta)}{\eta^2} \left(\frac{1}{2\tilde{a}} + 1\right) - 2 - 4T$,

and

$$\lambda = \frac{m_1 + m_3}{2} - \sqrt{\left(\frac{m_1 + m_3}{2}\right)^2 - m_1 m_3 + m_2^2}$$
(23)

Then, for any $\tilde{a} > 0$ such that $\lambda > 0$, for any $\varepsilon \in (0,1)$ and $n \in \mathbb{N}$,

$$(1 + \tilde{\kappa} \mathsf{T}) \,\mathcal{L}_{\tilde{a}}(h_{n+1}) \leqslant \mathcal{L}_{\tilde{a}}(h_n) + \mathsf{T}\delta^4 \tilde{M} \left(C_1 \mathrm{e}^{-\rho n t} \nu_0 \left(\mathfrak{W} \right) + C_2 \right) \,, \tag{24}$$

where

$$\tilde{\kappa} = \frac{\tilde{a}\lambda(1-\varepsilon)(1-3\mathrm{T})}{\max(LC_{\mathrm{LS}},1)+2\tilde{a}}, \qquad \tilde{M} = \frac{1}{\varepsilon\lambda}\left(\frac{4}{15\tilde{a}}+24\tilde{a}\right) + \tilde{a}\left((33\lambda+4)\mathrm{T} + \frac{\lambda\varepsilon}{240}\right).$$

Note that this result is only stated for $\eta > 0$. As explained in the proof of Theorem 1, the case $\eta = 0$ is obtained by a limiting procedure. The expression for λ in (23) admits a

limit when $\eta \to 0$; more precisely, this limit is $2 - 3T - 2\tilde{a}/[T(1+2\tilde{a})]$ which can be obtained from

$$\lambda = \frac{m_1 m_3 - m_2^2}{\frac{m_1 + m_3}{2} + \sqrt{\left(\frac{m_1 + m_3}{2}\right)^2 - m_1 m_3 + m_2^2}}$$

the numerator and denominator being respectively $\eta^{-2}\gamma(2-3T)(1+1/(2\tilde{a}))$ and $\eta^{-2}\gamma[(1+1/(2\tilde{a}))-\gamma]$ at dominant order in η , with $\gamma = T^{-1}$ when $\eta = 0$ in view of (6).

Remark 6. We discuss in this remark how to obtain (14) from the refined convergence result (24). First, since L is any upper bound of $\|\nabla^2 U\|_{\infty}$, we can assume without loss of generality that $LC_{LS} \ge 1$. Upon rescaling, it can therefore be assumed that L = 1 and $C_{LS} \ge$ 1 (see the discussion at the end of Section 2.2.1). We consider an initial condition of the form $\nu_0 = \hat{\nu}_0 \otimes N(0, I_d)$. Having in mind that the Euler-Maruyama discretization of overdamped Langevin dynamics corresponds to one-step HMC with full resampling of the momenta at each step with effective step-size $\delta^2/2$, we set K = 1 and $\eta = 0$, and therefore consider $n = 2s/\delta^2$ for some time s > 0. It follows from (6) that $T = \delta$. Choosing for instance $\varepsilon = \delta$ (or any other choice ensuring that ε vanishes as $\delta \to 0$) and $\tilde{a} = \alpha \delta$, it is then easily seen that λ is at dominant order $2(1 - \alpha)$, so that the convergence rate $\tilde{\kappa}$ is of order $2\delta\alpha(1 - \alpha)/C_{LS}$. This suggests to choose $\alpha = 1/2$ to maximize the convergence rate. Note that the factor $(1 + \tilde{\kappa}T)^{-n}$ then converges to s/C_{LS} as $\delta \to 0$ with these choices, which leads to (14) as claimed.

Theorem 5 is proved below. First, let us deduce Theorem 1 from Theorem 5:

Proof of Theorem 1. In the case $\eta > 0$, an easy induction based on (22) in Theorem 5 implies that, for any $n \ge 1$,

$$\mathcal{L}_{a}(h_{n}) \leq (1+\kappa \mathsf{T})^{-n} \mathcal{L}_{a}(h_{0}) + \mathsf{T}\delta^{4}M \sum_{k=1}^{n} (1+\kappa \mathsf{T})^{-k} \left[C_{1} \mathrm{e}^{-\rho(n-k)\mathsf{T}} \nu_{0}\left(\mathfrak{W}\right) + C_{2} \right]$$

The proof then follows from the bounds

$$\sum_{k=1}^{n} (1+\kappa T)^{-k} \leqslant \frac{1+\kappa T}{\kappa T}, \qquad \sum_{k=1}^{n} (1+\kappa T)^{-k} e^{-\rho(n-k)T} \leqslant n \left[\max\left((1+\kappa T)^{-1/T}, e^{-\rho} \right) \right]^{nT},$$

as well as $\mathcal{H}(h_n) \leq \mathcal{L}_a(h_n)$ and $\mathcal{L}_a(h_0) \leq \mathcal{H}(h_0) + 2a\mathcal{I}(h_0)$.

The case $\eta = 0$ is obtained by letting η go to 0 in the previous case.¹ Indeed, it is clear that the right hand side of (12) converges. For the left hand side, we use the lower semi-continuity of the relative entropy (see [50, Theorem 29.1]) with respect to the weak convergence of probability measures. The convergence of $\nu_0 P^n_{\omega}$ in the Wasserstein sense

¹We mention that the argument in the case $\eta = 0$ in [41] is not correct, an inequality is used in the wrong sense.

(hence in the weak sense) as $\eta \to 0$ follows from the moment bounds of **H** ³ together with a synchronous coupling argument, i.e. using the same Gaussian variables in the randomization steps of two chains, one with $\eta = 0$ and one with $\eta' > 0$ (so that the expected distance between the chains goes to zero as η' goes to zero).

Proof of Theorem 5. Before providing the proof of Theorem 5, let us recall the key steps of the proof of the similar result established in [49] for the continuous-time kinetic Langevin process, as the structure of the proof is similar. Denoting by $(\hat{h}_t)_{t\geq 0}$ the analogue of $(h_n)_{n\in\mathbb{N}}$ in this continuous-time context, the first observation is that

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{H}(\hat{h}_t) = -\int_{\mathbb{R}^{2d}} \frac{|\nabla_v \hat{h}_t|^2}{\hat{h}_t} \,\mathrm{d}\mu\,.$$
(25)

This entropy dissipation is not sufficient to conclude to an exponential decay, as it can vanish for non-constant densities \hat{h}_t (and thus in particular it cannot be upper bounded by $-\kappa \mathcal{H}(\hat{h}_t)$ for some $\kappa > 0$). A key observation is then that, when $\nabla^2 U$ is bounded,

$$\frac{\mathrm{d}}{\mathrm{d}t} \left(\int_{\mathbb{R}^{2d}} \frac{|\nabla_x \hat{h}_t + \nabla_v \hat{h}_t|^2}{\hat{h}_t} \,\mathrm{d}\mu \right) \leqslant -c \int_{\mathbb{R}^{2d}} \frac{|\nabla_x \hat{h}_t|^2}{\hat{h}_t} \,\mathrm{d}\mu + C \int_{\mathbb{R}^{2d}} \frac{|\nabla_v \hat{h}_t|^2}{\hat{h}_t} \,\mathrm{d}\mu \tag{26}$$

for some constants c, C > 0. The last term can be controlled thanks to the entropy dissipation (25). As a consequence, for a small enough, considering \mathcal{L}_a given by (21),

$$\frac{\mathrm{d}}{\mathrm{d}t}\mathcal{L}_a(\hat{h}_t) \leqslant -ac\mathcal{I}(\hat{h}_t)\,,\tag{27}$$

and the conclusion follows from

$$\mathcal{L}_a(\tilde{h}_t) \leq (\max(C_{\mathrm{LS}}, 1) + 2a)\mathcal{I}(\tilde{h}_t),$$

where we used that μ satisfies a log-Sobolev inequality with constant max(C_{LS} , 1) as the tensor product of π and N(0, I_d).

The proof of Theorem 5 is fundamentally based on the same ingredients, but with discrete-time evolutions along the two steps V_{δ} and D_{η} of the chain, instead of the time derivatives of (25) and (26). This requires various estimates given in Section 3.2.

Proof of Theorem 5. We start by general estimates, and then specify at the very end particular choices leading respectively to (22) and (24). Fix $\tilde{a} > 0$ and $n \in \mathbb{N}$. In this proof, we consider the matrix **A** given in (20). We write the proof for $\eta > 0$, the case $\eta = 0$ being obtained by passing to the limit $\eta \to 0$ in the final expressions, as discussed after Theorem 5.

The estimates on the evolution of h_n are based on a dual formulation where we consider the evolution of an arbitrary bounded measurable function g:

$$\int_{\mathbb{R}^{2d}} h_{n+1} g \,\mathrm{d}\mu = \int_{\mathbb{R}^{2d}} h_n \mathcal{P}_{\omega} g \,\mathrm{d}\mu \,,$$

in other words $h_{n+1} = P^*_{\omega}h_n$ where we denote by Q^* the adjoint of a bounded linear operator Q on $L^2(\mu)$. The first task is to identify the action of the adjoints of the operators constituting P_{ω} . A simple calculation shows that the damping step is reversible in the probabilistic sense, namely $D^*_{\eta} = D_{\eta}$. Since the Verlet map Φ^K_{δ} satisfies $|\det \nabla \Phi^K_{\delta}| \equiv 1$ (as a consequence of symplecticity [30]),

$$\begin{split} \int_{\mathbb{R}^{2d}} h\left(\mathbf{V}_{\delta}^{K}g\right) \mathrm{d}\mu &= \int_{\mathbb{R}^{2d}} h(z)g\left(\Phi_{\delta}^{K}(z)\right)\mu(z) \,\mathrm{d}z = \int_{\mathbb{R}^{2d}} h\left(\Phi_{\delta}^{-K}(z)\right)g(z)\mu\left(\Phi_{\delta}^{-K}(z)\right) \,\mathrm{d}z \\ &= \int_{\mathbb{R}^{2d}} g\left[\left(\mathbf{V}_{\delta}^{K}\right)^{*}h\right] \mathrm{d}\mu\,, \end{split}$$

which allows to identify the action of the adjoint operator as

$$\left(\mathbf{V}_{\delta}^{K}\right)^{*}h = \left(h \circ \Phi_{\delta}^{-K}\right)\mathbf{e}^{\Box H}, \qquad \Box H = H - H \circ \Phi_{\delta}^{-K}.$$
(28)

In addition, using the reversibility property (5), we obtain that $\Phi_{\delta}^{-K} = (\Phi_{\delta}^{-1})^{\circ K}$ with

$$\Phi_{\delta}^{-1}(x,v) = \left(x - \delta v - \frac{\delta^2}{2}\nabla U(x) , v + \frac{\delta}{2}\left[\nabla U(x) + \nabla U\left(x - \delta v - \frac{\delta^2}{2}\nabla U(x)\right)\right]\right).$$
(29)

From these observations and since $\mathbf{P}^*_{\omega} = (\mathbf{V}^K_{\delta})^* \mathbf{D}^*_{\eta}$, we consider the decomposition

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) - \mathcal{L}_{\tilde{a}}(h_n) = \mathcal{L}_{\tilde{a}}(\mathbf{P}^*_{\boldsymbol{\omega}}h_n) - \mathcal{L}_{\tilde{a}}(h_n) = \mathbf{\Delta}_1 + \mathbf{\Delta}_2 + \tilde{a}\mathbf{\Delta}_3 + \tilde{a}\mathbf{\Delta}_4$$

where

$$\begin{aligned} \boldsymbol{\Delta}_1 &= \mathcal{H}\left((\mathbf{V}_{\delta}^K)^* \mathbf{D}_{\eta} h_n\right) - \mathcal{H}(\mathbf{D}_{\eta} h_n) , \qquad \boldsymbol{\Delta}_2 &= \mathcal{H}(\mathbf{D}_{\eta} h_n) - \mathcal{H}(h_n) \\ \boldsymbol{\Delta}_3 &= \mathcal{I}_{\mathbf{A}}\left((\mathbf{V}_{\delta}^K)^* \mathbf{D}_{\eta} h_n\right) , \qquad \boldsymbol{\Delta}_4 &= -\mathcal{I}_{\mathbf{A}}(h_n) . \end{aligned}$$

We bound these four terms as follows. By Corollary 12, for any $\varepsilon_1 > 0$,

$$\mathbf{\Delta}_{1} \leqslant 4\varepsilon_{1} K \delta \mathcal{I} \left(\mathbf{D}_{\eta} h_{n} \right) + E_{\varepsilon_{1}} W_{n}$$

where for conciseness we write $W_n = C_1 e^{-\rho n T} \nu_0(\mathfrak{W}) + C_2$ (recall the notation from **H3**), and, by Corollaries 14 and 17,

$$E_{\varepsilon_1} = \sum_{j=0}^{K-1} \left[6\delta^7 j^2 \varepsilon_1 + \varepsilon_1 \int_0^\delta 2s^6 ds + \frac{1}{30\varepsilon_1} \delta^5 \right]$$

$$\leqslant 2\varepsilon_1 \delta^7 K^3 + \frac{2}{7} \varepsilon_1 K \delta^7 + \frac{1}{30\varepsilon_1} K \delta^5 \leqslant \frac{1}{30} \left(\varepsilon_1 + \frac{1}{\varepsilon_1} \right) K \delta^5 ,$$

where we used that $K\delta \leq 1/10$.

By Lemma 7,

$$\boldsymbol{\Delta}_2 \leqslant -\frac{1}{2}(\eta^{-2} - 1)\mathcal{I}_{\mathbf{B}_v}(\mathbf{D}_{\eta}h_n) , \qquad \boldsymbol{\Delta}_4 \leqslant -\mathcal{I}_{\mathbf{A}\mathbf{B}_{\eta}^{-1}}(\mathbf{D}_{\eta}h_n) ,$$

where

$$\mathbf{B}_{v} = \begin{pmatrix} 0 & 0\\ 0 & 1 \end{pmatrix}, \qquad \mathbf{B}_{\eta}^{-1} = \begin{pmatrix} 1 & 0\\ 0 & 1/\eta \end{pmatrix}.$$
(30)

Finally, using Lemma 8 and Corollary 14 and setting

$$\Psi = (\nabla \Phi_{\delta}^{-K}) \circ \Phi_{\delta}^{K} , \qquad (31)$$

it holds, for any $\varepsilon_2 > 0$,

$$\begin{split} \mathbf{\Delta}_{3} &\leqslant (1+\varepsilon_{2})\mathcal{I}_{\mathbf{A}\Psi}(\mathbf{D}_{\eta}h_{n}) + 2(1+\varepsilon_{2}^{-1})|\mathbf{A}|^{2}K^{2}\delta^{6}W_{n} \\ &\leqslant \mathcal{I}_{\mathbf{A}\Psi}(\mathbf{D}_{\eta}h_{n}) + \varepsilon_{2}|\mathbf{A}|^{2}\|\Psi\|_{\infty}^{2}\mathcal{I}(\mathbf{D}_{\eta}h_{n}) + 2(1+\varepsilon_{2}^{-1})|\mathbf{A}|^{2}K^{2}\delta^{6}W_{n} \\ &\leqslant \mathcal{I}_{\mathbf{A}\Psi}(\mathbf{D}_{\eta}h_{n}) + 3\varepsilon_{2}\mathcal{I}(\mathbf{D}_{\eta}h_{n}) + 4(1+\varepsilon_{2}^{-1})K^{2}\delta^{6}W_{n} \,, \end{split}$$

where we used $|\mathbf{A}| = \sqrt{2}$ and $\|\Psi\|_{\infty}\|^2 \leq 3/2$, the latter bound following from Equation (40) in Lemma 11 and $\mathbf{T} = K\delta \leq 1/10$.

Therefore, by combining the inequalities above, we get

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) - \mathcal{L}_{\tilde{a}}(h_n) \leqslant -\frac{\eta^{-2} - 1}{2} \mathcal{I}_{\mathbf{B}_v}(\mathbf{D}_{\eta}h_n) + \tilde{a}\mathcal{I}_{\mathbf{A}\Psi}(\mathbf{D}_{\eta}h_n) - \tilde{a}\mathcal{I}_{\mathbf{A}\mathbf{B}_{\eta}^{-1}}(\mathbf{D}_{\eta}h_n) + (4\varepsilon_1 K\delta + 3\tilde{a}\varepsilon_2) \mathcal{I}(\mathbf{D}_{\eta}h_n) + \left[\frac{\varepsilon_1 + \varepsilon_1^{-1}}{30} K\delta^5 + 4\tilde{a}(1 + \varepsilon_2^{-1})K^2\delta^6\right] W_n.$$
(32)

The sum of the terms in the first line of the right hand side is equal to

$$-\tilde{a}\mathcal{I}_{\mathbf{S}^{1/2}}(\mathbf{D}_{\eta}h_{n}) = -\tilde{a}\int_{\mathbb{R}^{2d}} \frac{\langle \nabla\left(\mathbf{D}_{\eta}h_{n}\right), \mathbf{S}\nabla\left(\mathbf{D}_{\eta}h_{n}\right)\rangle}{\mathbf{D}_{\eta}h_{n}} \,\mathrm{d}\mu \,, \tag{33}$$

with, for $z \in \mathbb{R}^{2d}$,

$$\mathbf{S}(z) = (2\tilde{a})^{-1}(\eta^{-2} - 1)\mathbf{B}_v^{\top}\mathbf{B}_v - (\mathbf{A}\Psi(z))^{\top}\mathbf{A}\Psi(z) + (\mathbf{A}\mathbf{B}_{\eta}^{-1})^{\top}\mathbf{A}\mathbf{B}_{\eta}^{-1},$$

where Ψ is defined in (31).

A key step of the proof is now to establish a positive lower bound on **S** uniformly in z, for \tilde{a} small enough, which is the analogue in our context of (27). We introduce the $d \times d$ block decompositions

$$\Psi(z) = \begin{pmatrix} \psi_{11}(z) & \psi_{12}(z) \\ \psi_{21}(z) & \psi_{22}(z) \end{pmatrix}, \qquad \mathbf{S}(z) = \begin{pmatrix} s_{11}(z) & s_{12}(z) \\ s_{21}(z) & s_{22}(z) \end{pmatrix},$$

so that

$$s_{11} = \mathbf{I}_d - (\psi_{11} + \psi_{21})^\top (\psi_{11} + \psi_{21}) , \quad s_{12} = s_{21} = \eta^{-1} \mathbf{I}_d - (\psi_{11} + \psi_{21})^\top (\psi_{12} + \psi_{22}) ,$$

$$s_{22} = \left(\frac{\eta^{-2} - 1}{2\tilde{a}} + \eta^{-2}\right) \mathbf{I}_d - (\psi_{22} + \psi_{12})^\top (\psi_{22} + \psi_{12}) .$$

Using Lemma 11 and $T \leq 1/10$, it holds, for any $x \in \mathbb{R}^d$,

$$\begin{aligned} |(\psi_{11} + \psi_{21})x| &\leq \left| \left((1 - \mathsf{T}) \, \mathsf{I}_d - \frac{\mathsf{T}^2}{2} \alpha \right) x \right| + \frac{2\mathsf{T}^3}{3} |x| &\leq \left(1 - \mathsf{T} + \mathsf{T}^2 \right) |x| \\ |(\psi_{22} + \psi_{12})x| &\leq \left| \left(\mathsf{I}_d + \mathsf{T}\beta - \frac{\mathsf{T}^2}{2} \zeta \right) x \right| + \frac{2\mathsf{T}^3}{3} |x| &\leq \left(1 + \mathsf{T} + \mathsf{T}^2 \right) |x| , \end{aligned}$$

and similarly

$$|(\psi_{11} + \psi_{21})x - x| \leq (\mathsf{T} + \mathsf{T}^2) |x|, \qquad |(\psi_{22} + \psi_{12})x - x| \leq (\mathsf{T} + \mathsf{T}^2) |x|,$$

so that, for any $x, y \in \mathbb{R}^d$,

$$\begin{split} \langle x, s_{11}x \rangle &\geq \left[1 - \left(1 - \mathsf{T} + \mathsf{T}^2 \right)^2 \right] |x|^2 \geqslant \left(2\mathsf{T} - 3\mathsf{T}^2 \right) |x|^2 \;, \\ \langle y, s_{22}y \rangle &\geq \left[\frac{\eta^{-2} - 1}{2\tilde{a}} + \eta^{-2} - \left(1 + \mathsf{T} + \mathsf{T}^2 \right)^2 \right] |y|^2 \geqslant \left(\frac{\eta^{-2} - 1}{2\tilde{a}} + \eta^{-2} - 1 - 2\mathsf{T} - 4\mathsf{T}^2 \right) |y|^2 \;, \\ \langle x, s_{12}y \rangle &= \left(\eta^{-1} - 1 \right) x \cdot y + x \left(\mathsf{I}_d - (\psi_{11} + \psi_{21})^\top (\psi_{12} + \psi_{22}) \right) y \\ &= \left(\eta^{-1} - 1 \right) x \cdot y + \langle (\mathsf{I}_d - (\psi_{11} + \psi_{21})) x, (\psi_{12} + \psi_{22}) y \rangle + \langle x, (\mathsf{I}_d - (\psi_{12} + \psi_{22})) y \rangle \\ &\geq - \left(\eta^{-1} - 1 + \left(2 + \mathsf{T} + \mathsf{T}^2 \right) \left(\mathsf{T} + \mathsf{T}^2 \right) \right) |x||y| \\ &\geqslant - \left(\eta^{-1} - 1 + 2\mathsf{T} + 4\mathsf{T}^2 \right) |x||y| \,. \end{split}$$

Note that the last right hand sides in the previous three inequalities are of order T at dominant order in T (using (6) to write $\eta = 1 - \gamma T$). This motivates factoring out T, and writing, for any $x, y \in \mathbb{R}^d$ and $z \in \mathbb{R}^{2d}$,

$$\begin{pmatrix} x \\ y \end{pmatrix}^{\top} \mathbf{S}(z) \begin{pmatrix} x \\ y \end{pmatrix} \ge \mathsf{T}\lambda \left(|x|^2 + |y|^2 \right) , \qquad (34)$$

where λ is the smallest eigenvalue of the symmetric 2 \times 2 matrix

$$\begin{pmatrix} m_1 & m_2 \\ m_2 & m_3 \end{pmatrix} := \frac{1}{\mathsf{T}} \begin{pmatrix} 2\mathsf{T} - 3\mathsf{T}^2 & -(\eta^{-1} - 1 + 2\mathsf{T} + 4\mathsf{T}^2) \\ -(\eta^{-1} - 1 + 2\mathsf{T} + 4\mathsf{T}^2) & \frac{\eta^{-2} - 1}{2\tilde{a}} + \eta^{-2} - 1 - 2\mathsf{T} - 4\mathsf{T}^2 \end{pmatrix}$$

$$= \mathbf{B}_{\eta}^{-1} \begin{pmatrix} 2 - 3\mathsf{T} & -(\gamma + \eta(2 + 4\mathsf{T})) \\ -(\gamma + \eta(2 + 4\mathsf{T})) & \gamma(1 + \eta) \left(\frac{1}{2\tilde{a}} + 1\right) - \eta^{2}(2 + 4\mathsf{T}) \end{pmatrix} \mathbf{B}_{\eta}^{-1},$$

recalling the notation $\gamma = (1 - \eta)/T$. This provides the definitions given in (35). The expression (23) for λ then follows from

$$\lambda = \frac{m_1 + m_3}{2} - \sqrt{\left(\frac{m_1 + m_3}{2}\right)^2 - m_1 m_3 + m_2^2},$$

which is positive provided \tilde{a} is small enough. Alternatively, to get the simpler expressions (13), we proceed as follows: using that $\eta \leq 1$ and $T \leq 1/10$, we bound

$$m_1 \ge \frac{17}{10}, \qquad |m_2| \le \frac{1}{\eta} \left(\gamma + \frac{12}{5}\right), \qquad m_3 \ge \frac{1}{\eta^2} \left(\frac{\gamma}{2\tilde{a}} - \frac{12}{5}\right),$$
(35)

to get that

$$\begin{pmatrix} x \\ y \end{pmatrix}^{\top} \begin{pmatrix} m_1 & m_2 \\ m_2 & m_3 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = m_1 |x|^2 + 2m_2 |x| |y| + m_3 |y|^2 \ge \frac{1}{2} m_1 |x|^2 + \left(m_3 - \frac{2m_2^2}{m_1} \right) |y|^2 \ge \frac{17}{20} \left(|x|^2 + |y|^2 \right) ,$$

i.e., $\lambda \ge 17/20$ if we choose \tilde{a} such that $m_3 \ge m_1/2 + 2m_2^2/m_1$. This condition is satisfied for any $\eta \in (0, 1]$ when it is satisfied for $\eta = 1$, in which case it reads

$$\frac{\gamma}{2\tilde{a}} \ge \frac{17}{20} + \frac{12}{5} + \frac{20}{17} \left(\gamma + \frac{12}{5}\right)^2 \,.$$

This holds in particular with the choice of $\tilde{a} = a$ in (13).

In both cases, at this point, plugging the lower bound (34) in (33), we have determined $\lambda > 0$ such that

$$\tilde{a}\mathcal{I}_{\mathbf{S}^{1/2}}(\mathbf{D}_{\eta}h_n) \leqslant -\tilde{a}\lambda \mathtt{T}\mathcal{I}(\mathbf{D}_{\eta}h_n)$$
.

Combining this result with (32) yields

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) - \mathcal{L}_{\tilde{a}}(h_n) \leqslant -\rho' \mathcal{I}\left(\mathcal{D}_{\eta}h_n\right) + \left[\frac{\varepsilon_1 + \varepsilon_1^{-1}}{30}K\delta^5 + 4\tilde{a}(1 + \varepsilon_2^{-1})K^2\delta^6\right] W_n, \quad (36)$$

for

$$\rho' = \tilde{a}\lambda \mathbf{T} - 4\varepsilon_1 \mathbf{T} - 3\tilde{a}\varepsilon_2 . \tag{37}$$

Now, applying Lemma 8 with $\tilde{\mathbf{A}} = \Psi^{-1} \circ \Phi_{\delta}^{-K}$ and *h* replaced by $D_{\eta}h_n$ (see Remark 9), we obtain that, for any $\varepsilon_3 > 0$

$$(1+\varepsilon_3)\mathcal{I}(\mathbf{D}_{\eta}h_n) \geqslant \mathcal{I}_{\Psi^{-1}\circ\Phi_{\delta}^{-K}}(\mathbf{P}_{\omega}^*h_n) - (1+\varepsilon_3^{-1}) \left\|\Psi^{-1}\circ\Phi_{\delta}^{-K}\right\|_{\infty}^2 \operatorname{Er}_1^{\delta,K}(\mathbf{D}_{\eta}h_n).$$

Thanks to Lemma 11, and more specifically to (40), for any $z, u \in \mathbb{R}^{2d}$,

$$\begin{split} |\Psi^{-1}(z)u|^2 &\geqslant \left(1 + \mathrm{T} + \mathrm{T}^2/2 + \mathrm{T}^3/3\right)^{-2} |u|^2 \\ &\geqslant (1 + 5\mathrm{T}/2)^{-1} |u|^2 \\ &\geqslant (1 - 5\mathrm{T}/2) |u|^2 \,, \end{split}$$

so that $\mathcal{I}_{\Psi^{-1} \circ \Phi_{\delta}^{-K}}(\mathbb{P}_{\omega}^*h_n) \ge (1 - 5T/2) \mathcal{I}(\mathbb{P}_{\omega}^*h_n)$. Using Lemma 11 again, we have, for any $z, u \in \mathbb{R}^{2d}$,

$$|u| \leq |\Psi(z)u| + |\Psi(z)u - u| \leq |\Psi(z)u| + (T + T^2/2 + T^3/3) |u|,$$

and thus

$$\left|\Psi^{-1}(z)u\right| \le |u| + \left(\mathsf{T} + \mathsf{T}^2/2 + \mathsf{T}^3/3\right) \left|\Psi^{-1}(z)u\right| \le |u| + \frac{1}{2} \left|\Psi^{-1}(z)u\right|,$$

so that $\|\Psi^{-1} \circ \Phi_{\delta}^{-K}\|_{\infty} \leq 2$. Plugging these bounds in (36) yields

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) - \mathcal{L}_{\tilde{a}}(h_n) \leqslant -\frac{\rho'}{1+\varepsilon_3} \left[\left(1 - \frac{5\mathrm{T}}{2} \right) \mathcal{I}(h_{n+1}) - 4(1+\varepsilon_3^{-1}) \mathrm{Er}_1^{\delta,K}(\mathrm{D}_{\eta}h_n) \right] \\ + \left[\frac{\varepsilon_1 + \varepsilon_1^{-1}}{30} K \delta^5 + 4\tilde{a}(1+\varepsilon_2^{-1}) K^2 \delta^6 \right] W_n \,.$$

We next use that μ satisfies a log-Sobolev inequality with constant max $(1, C_{\text{LS}})$ (by tensorization from the log-Sobolev inequalities satisfied by π and N $(0, I_d)$), and the equality $|\mathbf{A}|^2 = 2$, to write

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) \leqslant \left(\max(1, C_{\mathrm{LS}}) + 2\tilde{a} \right) \mathcal{I}(h_{n+1}) \,.$$

Choosing $\varepsilon_3 = T/4$, using Corollary 14 to bound the remaining numerical error term, setting

$$M = 8\rho'\left(1 + \frac{4}{\mathrm{T}}\right)\mathrm{T} + \frac{\varepsilon_1 + \varepsilon_1^{-1}}{30} + 4\tilde{a}(1 + \varepsilon_2^{-1})\mathrm{T},$$

and noting that $(1 - 5T/2)/(1 + T/4) \ge 1 - 3T$ and $\rho'(1 - 3T) > 0$, we can conclude that

$$\mathcal{L}_{\tilde{a}}(h_{n+1}) - \mathcal{L}_{\tilde{a}}(h_n) \leq -\rho'(1-3\mathsf{T})\mathcal{I}(h_{n+1}) + \mathsf{T}\delta^4 M W_n$$
$$\leq -\frac{\rho'(1-3\mathsf{T})}{\max(1,C_{\mathrm{LS}}) + 2\tilde{a}}\mathcal{L}_{\tilde{a}}(h_{n+1}) + \mathsf{T}\delta^4 M W_n.$$

Let us finally discuss how to obtain (22) and (24):

• to obtain (22) with the simpler expressions of κ and M in (13), we first note that a in (13) is smaller than 1/10, as can be seen for instance by distinguishing the case $\gamma \leq 3$, for which $a \leq \gamma/34$, and the case $\gamma \geq 3$, for which $a \leq (\gamma + 3)/[3(\gamma + 3)^2])$. We next rely on the expression (37) for ρ' , with the choices $\varepsilon_1 = a/20$ and $\varepsilon_2 = T/20$, so that, replacing λ by 17/20,

$$\rho' = a\mathsf{T}\left(\lambda - \frac{1}{5} - \frac{3}{20}\right) = \frac{a\mathsf{T}}{2}.$$

The value of κ is then obtained as

$$\kappa = \frac{\rho'(1-3\mathsf{T})}{\max(1,C_{\rm LS})+2a} = \frac{a(1-3\mathsf{T})}{2\left[\max(1,C_{\rm LS})+2a\right]} \ge \frac{a}{3\left[\max(1,C_{\rm LS})+2a\right]}$$

In the expression of M, we simply bound $\rho' \leq 1/200$, $a \leq 1/10$, $T \leq 1/10$.

• To get the sharper inequality (24), we keep a free parameter $\varepsilon \in (0, 1)$ and choose $\varepsilon_1 = \tilde{a}\varepsilon\lambda/8$, $\varepsilon_2 = T\varepsilon\lambda/6$ which implies that $\rho' = \tilde{a}\lambda T(1-\varepsilon)$ (which immediately leads to the claimed value for κ) and

$$M = 8\tilde{a}\lambda T \left(T + 4\right) + \frac{\tilde{a}\lambda\varepsilon}{240} + \frac{4}{15\tilde{a}\lambda\varepsilon} + 4\tilde{a}\left(T + \frac{6}{\varepsilon\lambda}\right),$$

and using that $T \leq 1/10$ gives the claimed result.

This allows to conclude the proof of Theorem 5.

In the next sections, we derive the technical results that we used in the proof of Theorem 5. In Section 3.2, we bound $\mathcal{L}_{\tilde{a}}(h_{n+1})$ in terms of h_n , the Jacobian matrix of the (reverse) Verlet map Φ_{δ}^{-K} and some numerical error terms. The latters are studied respectively in Sections 3.3 and 3.4.

3.2 Evolution of entropies under elementary steps

In this section, we compute the evolution of the relative entropy and Fisher-like terms (19) along the damping and Verlet steps. The velocity randomization step is the same as in [41, Section 4.1.1], where the following is established.

Lemma 7 (Dissipation from velocity randomization). Recalling from (30) the notation

$$\mathbf{B}_{v} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \qquad \mathbf{B}_{\eta} = \begin{pmatrix} 1 & 0 \\ 0 & \eta \end{pmatrix},$$

then, for all $\eta \in (0,1)$ and all measurable h with $\mathcal{H}(h) < +\infty$,

$$\mathcal{H}(\mathbf{D}_{\eta}h) - \mathcal{H}(h) \leqslant -\left(\frac{\eta^{-2}-1}{2}\right)\mathcal{I}_{\mathbf{B}_{v}}(\mathbf{D}_{\eta}h).$$

and, for any matrix $\tilde{\mathbf{A}}$,

$$\mathcal{I}_{\tilde{\mathbf{A}}\mathbf{B}_{\eta}^{-1}}(\mathbf{D}_{\eta}h) \leqslant \mathcal{I}_{\tilde{\mathbf{A}}}(h).$$

Proof. The first inequality is equivalent to

$$0 \leq \mathcal{H}(\mathbf{D}_{\eta}h) + \left(\frac{\eta^{-2} - 1}{2}\right) \mathcal{I}_{\mathbf{B}_{v}}(\mathbf{D}_{\eta}h) \leq \mathcal{H}(h) < +\infty,$$

which is established in [41, Equation (45)] (it is sufficient to prove the bound for a smooth positive h with compact support and pass to the limit).

The second point follows from Jensen's inequality and the convexity of $(t, x) \mapsto |x|^2 / t$ on $\mathbb{R}^*_+ \times \mathbb{R}^d$ (using [14, Proposition 2.3 (ii)]) which imply

$$\int_{\mathbb{R}^{2d}} \frac{|\tilde{\mathbf{A}}\nabla h|^2}{h} \mathrm{d}\mu = \int_{\mathbb{R}^{2d}} \mathcal{D}_\eta \left(\frac{|\tilde{\mathbf{A}}\nabla h|^2}{h}\right) \mathrm{d}\mu \geqslant \int_{\mathbb{R}^{2d}} \frac{|\mathcal{D}_\eta \tilde{\mathbf{A}}\nabla h|^2}{\mathcal{D}_\eta h} \mathrm{d}\mu = \int_{\mathbb{R}^{2d}} \frac{|\tilde{\mathbf{A}}\mathbf{B}_\eta^{-1}\nabla \mathcal{D}_\eta h|^2}{\mathcal{D}_\eta h} \mathrm{d}\mu,$$

where the last equality comes from $\nabla_v(\mathbf{D}_\eta h) = \eta \mathbf{D}_\eta(\nabla_v h)$ and $\mathbf{D}_\eta(\nabla_x h) = \nabla_x(\mathbf{D}_\eta h)$. \Box

Lemma 8 (Fisher inequality on Verlet step). For all matrices $\tilde{\mathbf{A}}$, all smooth h with $\mathcal{I}(h) < +\infty$, and all $\varepsilon > 0$,

$$\mathcal{I}_{\tilde{\mathbf{A}}}\left((\mathbf{V}_{\delta}^{K})^{*}h\right) \leqslant (1+\varepsilon)\mathcal{I}_{\tilde{\mathbf{A}}\Psi}(h) + (1+\varepsilon^{-1})\left|\tilde{\mathbf{A}}\right|^{2}\operatorname{Er}_{1}^{\delta,K}(h),$$

with $\Psi = (\nabla \Phi_{\delta}^{-K}) \circ \Phi_{\delta}^{K}$ and the error term

$$\operatorname{Er}_{1}^{\delta,K}(h) = \int_{\mathbb{R}^{2d}} \left| (\nabla \Box H) \circ \Phi_{\delta}^{K} \right|^{2} h \, \mathrm{d}\mu \, .$$

Proof. By an approximation argument, it can be assumed that h is bounded below by a positive constant and globally Lipschitz. Recalling the action of $(V_{\delta}^{K})^{*}$ from (28),

$$\begin{split} \int_{\mathbb{R}^{2d}} \frac{\left|\tilde{\mathbf{A}}\nabla(\mathbf{V}_{\delta}^{K})^{*}h\right|^{2}}{(\mathbf{V}_{\delta}^{K})^{*}h} \,\mathrm{d}\mu &= \int_{\mathbb{R}^{2d}} \frac{\left|\tilde{\mathbf{A}}\nabla\left((h\circ\Phi_{\delta}^{-K})\mathrm{e}^{\Box H}\right)\right|^{2}}{(h\circ\Phi_{\delta}^{-K})\mathrm{e}^{\Box H}} \,\mathrm{d}\mu \\ &= \int_{\mathbb{R}^{2d}} \frac{\left|\tilde{\mathbf{A}}\left[\nabla(\Phi_{\delta}^{-K})(\nabla h)\circ\Phi_{\delta}^{-K}-(h\circ\Phi_{\delta}^{-K})\nabla\Box H\right]\mathrm{e}^{\Box H}\right]^{2}}{(h\circ\Phi_{\delta}^{-K})\mathrm{e}^{\Box H}} \,\mathrm{d}\mu \\ &= \int_{\mathbb{R}^{2d}} \frac{\left|\tilde{\mathbf{A}}\left[\nabla(\Phi_{\delta}^{-K})(\nabla h)\circ\Phi_{\delta}^{-K}-(h\circ\Phi_{\delta}^{-K})\nabla\Box H\right]\right|^{2}}{h\circ\Phi_{\delta}^{-K}} \mathrm{e}^{\Box H} \,\mathrm{d}\mu \\ &= \int_{\mathbb{R}^{2d}} \frac{\left|\tilde{\mathbf{A}}\left[\left(\nabla(\Phi_{\delta}^{-K})\circ\Phi_{\delta}^{K}\right)\nabla h-h(\nabla\Box H)\circ\Phi_{\delta}^{K}\right]\right|^{2}}{h} \,\mathrm{d}\mu \,, \end{split}$$

where we used the change of variable Φ_{δ}^{-K} in the last line. Introducing $\Psi = \nabla(\Phi_{\delta}^{-K}) \circ \Phi_{\delta}^{K}$, we therefore obtain, with a discrete Cauchy–Schwarz inequality: for any $\varepsilon > 0$,

$$\int_{\mathbb{R}^{2d}} \frac{|\tilde{\mathbf{A}}\nabla(\mathbf{V}_{\delta}^{K})^{*}h|^{2}}{(\mathbf{V}_{\delta}^{K})^{*}h} \,\mathrm{d}\mu \leqslant (1+\varepsilon) \int_{\mathbb{R}^{2d}} \frac{|\tilde{\mathbf{A}}\Psi\nabla h|^{2}}{h} \,\mathrm{d}\mu + \left(1+\frac{1}{\varepsilon}\right) \left|\tilde{\mathbf{A}}\right|^{2} \int_{\mathbb{R}^{2d}} \left|\nabla\Box H \circ \Phi_{\delta}^{K}\right|^{2} h \,\mathrm{d}\mu \,,$$

which is indeed the claimed estimate.

Remark 9. If the constant matrix $\tilde{\mathbf{A}}$ is replaced by a matrix field $z \mapsto \tilde{\mathbf{A}}(z)$, a straightforward adapation of the proof yields

$$\mathcal{I}_{\tilde{\mathbf{A}}}\left((\mathbf{V}_{\delta}^{K})^{*}h\right) \leqslant (1+\varepsilon)\mathcal{I}_{(\tilde{\mathbf{A}}\circ\Phi_{\delta}^{K})\Psi}(h) + (1+\varepsilon^{-1})\left\|\tilde{\mathbf{A}}\right\|_{\infty}^{2}\operatorname{Er}_{1}^{\delta,K}(h)$$

with $\left\|\tilde{\mathbf{A}}\right\|_{\infty} = \sup_{z \in \mathbb{R}^{2d}} |\tilde{\mathbf{A}}(z)|.$

Lemma 10 (Entropy inequality on Verlet step). For all smooth h with $\mathcal{I}(h) < +\infty$ and all $\varepsilon > 0$,

$$\mathcal{H}\left((\mathbf{V}_{\delta}^{K})^{*}h\right) - \mathcal{H}(h) \leqslant \sum_{j=0}^{K-1} \left[\frac{\varepsilon}{2} \int_{0}^{\delta} \mathcal{I}\left((\mathbf{V}_{\delta}^{j}\mathbf{V}_{s}^{1})^{*}h\right) ds + \frac{1}{2\varepsilon} \mathrm{Er}_{2}\left((\mathbf{V}_{\delta}^{j})^{*}h\right)\right],$$

with the error term

$$\operatorname{Er}_{2}(h) = \int_{0}^{\delta} \left(\int_{\mathbb{R}^{2d}} |\partial_{s} \Phi_{s} - F_{H} \circ \Phi_{s}|^{2} h \, d\mu \right) \mathrm{d}s \,,$$

where

$$F_H(x,v) = \begin{pmatrix} v \\ -\nabla U(x) \end{pmatrix} .$$
(38)

Proof. In view of the decomposition

$$\mathcal{H}\left((\mathbf{V}_{\delta}^{K})^{*}h\right) - \mathcal{H}(h) = \sum_{j=0}^{K-1} \left[\mathcal{H}\left((\mathbf{V}_{\delta}^{j+1})^{*}h\right) - \mathcal{H}\left((\mathbf{V}_{\delta}^{j})^{*}h\right)\right],$$

we simply have to establish the result for K = 1.

For $z \in \mathbb{R}^{2d}$ and $t \in [0, \delta]$, let $z_t = \Phi_t(z)$. For f_0 an initial (smooth positive) distribution, denote by f_t the law of z_t when $z \sim f_0$ and $h_t = f_t/\mu$. By a change a variable we see that, for any smooth function g on \mathbb{R}^{2d} ,

$$\int_{\mathbb{R}^{2d}} g(z) f_t(z) \, \mathrm{d}z = \int_{\mathbb{R}^{2d}} g(z_t) \, f_0(z) \, \mathrm{d}z = \int_{\mathbb{R}^{2d}} g(z) f_0\left(\Phi_t^{-1}(z)\right) \, \mathrm{d}z \,,$$

i.e.
$$f_t(z) = f_0\left(\Phi_t^{-1}(z)\right)$$
, and

$$\int_{\mathbb{R}^{2d}} g(z)\partial_t f_t(z) \,\mathrm{d}z = \frac{\mathrm{d}}{\mathrm{d}t} \left(\int_{\mathbb{R}^{2d}} g\left(z_t\right) f_0(z) \,\mathrm{d}z\right) = \int_{\mathbb{R}^{2d}} \partial_t \Phi_t(z) \cdot \nabla g(\Phi_t(z)) f_0(z) \,\mathrm{d}z$$

$$= \int_{\mathbb{R}^{2d}} j_t(z) \cdot \nabla g(z) f_t(z) \,\mathrm{d}z, \qquad (39)$$

with $j_t = \partial_t \Phi_t \circ \Phi_t^{-1}$. Then,

$$\int_{\mathbb{R}^{2d}} f_{\delta} \ln h_{\delta} - \int_{\mathbb{R}^{2d}} f_0 \ln h_0 = \int_0^{\delta} \frac{\mathrm{d}}{\mathrm{d}s} \left(\int_{\mathbb{R}^{2d}} f_s \ln h_s \, \mathrm{d}s \right),$$

and

$$\frac{\mathrm{d}}{\mathrm{d}s} \left(\int_{\mathbb{R}^{2d}} f_s \ln h_s \right) = \int_{\mathbb{R}^{2d}} \partial_s h_s \,\mathrm{d}\mu + \int_{\mathbb{R}^{2d}} \left(\ln h_s \right) \partial_s f_s \,.$$

The first term of the right hand side vanishes as it is the time derivative of the integral of h_s , which is 1 for any $s \ge 0$. Combining the latter equality with (39) then gives

$$\frac{\mathrm{d}}{\mathrm{d}s} \left(\int_{\mathbb{R}^{2d}} f_s \ln h_s \right) = \int_{\mathbb{R}^{2d}} j_s \cdot \nabla(\ln h_s) f_s \,.$$

Since $F_H \cdot \nabla \mu = 0$ and F_H is divergence free,

$$\int_{\mathbb{R}^{2d}} F_H \cdot \nabla(\ln h_s) f_s = \int_{\mathbb{R}^{2d}} F_H \cdot \nabla h_s \, \mathrm{d}\mu = -\int_{\mathbb{R}^{2d}} h_s \left(F_H \cdot \nabla \mu + \mu \operatorname{div} F_H \right) = 0.$$

Thus, we can add this term to the previous equality to get, for any $\varepsilon > 0$,

$$\begin{aligned} \frac{\mathrm{d}}{\mathrm{d}s} \left(\int_{\mathbb{R}^{2d}} f_s \ln h_s \right) &= \int_{\mathbb{R}^{2d}} \left(j_s(z) - F_H(z) \right) \cdot \nabla \ln h_s(z) f_s(z) \, \mathrm{d}z \\ &\leqslant \frac{1}{2\varepsilon} \int_{\mathbb{R}^{2d}} \left| j_s(z) - F_H(z) \right|^2 f_s(z) \, \mathrm{d}z + \frac{\varepsilon}{2} \int_{\mathbb{R}^{2d}} \left| \nabla \ln h_s(z) \right|^2 f_s(z) \, \mathrm{d}z \\ &= \frac{1}{2\varepsilon} \int_{\mathbb{R}^{2d}} \left| \partial_s \Phi_s - F_H \circ \Phi_s \right|^2 h_s \, \mathrm{d}\mu + \frac{\varepsilon}{2} \int_{\mathbb{R}^{2d}} \left| \nabla \ln h_s(z) \right|^2 f_s(z) \, \mathrm{d}z \,, \end{aligned}$$

which gives the announced result once integrated over $s \in [0, \delta]$.

At this stage, gathering Lemmas 7, 8 and 10 yields a bound on $\mathcal{L}_a(h_{n+1})$ involving h_n , the matrix field Ψ of Lemma 8 and the numerical error terms of Lemmas 8 and 10. It remains to understand these parts. We give some estimates on Ψ in Section 3.3 and analyse the numerical errors in Section 3.4.

3.3 Jacobian matrix of the Verlet integrator

As mentioned above, this section focuses on the matrix field $\Psi = (\nabla \Phi_{\delta}^{-K}) \circ \Phi_{\delta}^{K}$ appearing in Lemma 8. The following result is the analogue for the Verlet scheme of the first part of [41, Lemma 1] for the Hamiltonian dynamics.

Lemma 11. Under H1 with L = 1, for all $z \in \mathbb{R}^{2d}$, there exist $d \times d$ matrices α, β, ζ with norms less than 1 such that, recalling the notation $T = K\delta$,

$$\left| \Psi(z) - \begin{pmatrix} \mathbf{I}_d - \frac{\mathbf{T}^2}{2} \alpha & \mathbf{T}\beta \\ -\mathbf{T} & \mathbf{I}_d - \frac{\mathbf{T}^2}{2} \zeta \end{pmatrix} \right| \leqslant \frac{1}{3} \mathbf{T}^3 \, .$$

In particular,

$$|\Psi(z)| \leq 1 + \mathsf{T} + \frac{1}{2}\mathsf{T}^2 + \frac{1}{3}\mathsf{T}^3$$
 (40)

Proof. Denote by \mathcal{M}_1 the set of $d \times d$ matrices with operator norm bounded by 1. Note that the product of two elements in \mathcal{M}_1 is still in \mathcal{M}_1 . Fix $z \in \mathbb{R}^{2d}$. For $k \ge 1$, set $z_k = \Phi_{\delta}^{-1}(z_{k-1})$, with $z_0 = \Phi_{\delta}^K(z)$. For $k \ge 0$, set $Q_k = \nabla^2 U(x_k)$, where $z_k = (x_k, v_k)$. By assumption, $Q_k \in \mathcal{M}_1$ for all $k \ge 0$.

Recall that, for $F : \mathbb{R}^d \to \mathbb{R}^d$, we use the convention $\nabla F = (\partial_i F_j)_{1 \leq i,j \leq d}$ where *i* stands for the row and *j* for the column. Hence, from (29), for all $k \geq 0$,

$$\nabla(\Phi_{\delta}^{-1})(z_k) = \begin{pmatrix} \mathrm{I}_d - \frac{\delta^2}{2}Q_k & \frac{\delta}{2}\left[Q_k + \left(\mathrm{I}_d - \frac{\delta^2}{2}Q_k\right)\right)Q_{k+1}\right] \\ -\delta & \mathrm{I}_d - \frac{\delta^2}{2}Q_{k+1} \end{pmatrix}.$$

In particular,

$$\left| \nabla(\Phi_{\delta}^{-1})(z_k) - \begin{pmatrix} \mathrm{I}_d - \frac{\delta^2}{2} Q_k & \frac{\delta}{2} \left[Q_k + Q_{k+1} \right] \\ -\delta & \mathrm{I}_d - \frac{\delta^2}{2} Q_{k+1} \end{pmatrix} \right| \leqslant \frac{\delta^3}{4} \,. \tag{41}$$

For all $k \in [[1, K]]$, let us determine by induction a constant $C_k \ge 0$ and matrices $\alpha_k, \beta_k, \zeta_k \in \mathcal{M}_1$ such that

$$\left| \nabla(\Phi_{\delta}^{-k})(z_0) - \begin{pmatrix} \mathbf{I}_d - \frac{(\delta k)^2}{2} \alpha_k & \delta k \beta_k \\ -\delta k & \mathbf{I}_d - \frac{(\delta k)^2}{2} \zeta_k \end{pmatrix} \right| \leqslant C_k \,. \tag{42}$$

For k = 1, this is given by (41) (applied at k = 0) with $C_1 = \delta^3/4$. Suppose that the result is true for some $k \ge 1$. In particular, using that $\delta k \le 1/10$,

$$|\nabla(\Phi_{\delta}^{-k})(z_0)| \leq \left| \begin{pmatrix} \mathbf{I}_d - \frac{(\delta k)^2}{2} \alpha_k & 0\\ 0 & \mathbf{I}_d - \frac{(\delta k)^2}{2} \zeta_k \end{pmatrix} \right| + \left| \begin{pmatrix} 0 & \delta k \beta_k \\ -\delta k & 0 \end{pmatrix} \right| + C_k$$

$$\leqslant 1 + \frac{21}{20}\delta k + C_k \,. \tag{43}$$

We next use $\nabla \Phi_{\delta}^{-k-1}(z_0) = \nabla \Phi_{\delta}^{-1}(z_k) \nabla \Phi_{\delta}^{-k}(z_0)$. At dominant order, the product of the terms on the right hand side of the previous equality is

$$\begin{pmatrix} \mathbf{I}_d - \frac{\delta^2}{2} Q_k & \frac{\delta}{2} [Q_k + Q_{k+1}] \\ -\delta & \mathbf{I}_d - \frac{\delta^2}{2} Q_{k+1} \end{pmatrix} \begin{pmatrix} \mathbf{I}_d - \frac{(\delta k)^2}{2} \alpha_k & \delta k \beta_k \\ -\delta k & \mathbf{I}_d - \frac{(\delta k)^2}{2} \zeta_k \end{pmatrix}$$
$$= \begin{pmatrix} \mathbf{I}_d - \frac{\delta^2 (k+1)^2}{2} \alpha_{k+1} & \delta (k+1) \beta_{k+1} \\ -\delta (k+1) & \mathbf{I}_d - \frac{\delta^2 (k+1)^2}{2} \zeta_{k+1} \end{pmatrix} + \Re,$$

with

$$\begin{aligned} \alpha_{k+1} &= \frac{1}{(k+1)^2} \left(Q_k + k^2 \alpha_k + k(Q_k + Q_{k+1}) \right) \,, \qquad \beta_{k+1} = \frac{1}{2(k+1)} \left(Q_k + Q_{k+1} + 2k\beta_k \right) \,, \\ \zeta_{k+1} &= \frac{1}{(k+1)^2} \left(2k\beta_k + Q_{k+1} + k^2\zeta_k \right) \,, \end{aligned}$$

which are all three in \mathcal{M}_1 , and a remainder term

$$\mathfrak{R} = \begin{pmatrix} \frac{\delta^4 k^2}{4} Q_k \alpha_k & -\frac{\delta^3 k}{4} \left(2Q_k \beta_k + k(Q_k + Q_{k+1})\zeta_k \right) \\ \frac{\delta^3 k}{2} \left(Q_{k+1} + k\alpha_k \right) & \frac{\delta^4 k^2}{4} Q_{k+1}\zeta_k \end{pmatrix}$$

.

Using that $\delta \leq 1/10$,

$$\begin{aligned} |\Re| &= \frac{\delta^4 k^2}{4} \left| \begin{pmatrix} Q_k \alpha_k & 0\\ 0 & Q_{k+1} \zeta_k \end{pmatrix} \right| + \frac{\delta^3 k}{2} \left| \begin{pmatrix} 0 & -Q_k \beta_k - k(Q_k + Q_{k+1})/2\zeta_k \\ Q_{k+1} + k\alpha_k & 0 \end{pmatrix} \right| \\ &\leqslant \frac{\delta^4 k^2}{4} + \frac{\delta^3 k}{2} (k+1) \leqslant \frac{21}{40} \delta^3 k (k+1). \end{aligned}$$
(44)

Then,

$$\begin{aligned} (\star) &:= \left| \nabla \Phi_{\delta}^{-k-1}(z_{0}) - \begin{pmatrix} \mathrm{I}_{d} - \delta^{2}(k+1)^{2}\alpha_{k+1}/2 & \delta(k+1)\beta_{k+1} \\ -\delta(k+1) & \mathrm{I}_{d} - \delta^{2}(k+1)^{2}\zeta_{k+1}/2 \end{pmatrix} \right| \\ &\leqslant |\Re| + \left| \begin{bmatrix} \nabla(\Phi_{\delta}^{-1})(z_{k}) - \begin{pmatrix} \mathrm{I}_{d} - \delta^{2}Q_{k}/2 & \delta\left[Q_{k} + Q_{k+1}\right]/2 \\ -\delta & \mathrm{I}_{d} - \delta^{2}Q_{k+1}/2 \end{pmatrix} \right] \nabla \Phi_{\delta}^{-k}(z_{0}) \right| \\ &+ \left| \begin{pmatrix} \mathrm{I}_{d} - \delta^{2}Q_{k}/2 & \delta\left[Q_{k} + Q_{k+1}\right]/2 \\ -\delta & \mathrm{I}_{d} - \delta^{2}Q_{k+1}/2 \end{pmatrix} \left[\nabla \Phi_{\delta}^{-k}(z_{0}) - \begin{pmatrix} \mathrm{I}_{d} - (\delta k)^{2}\alpha_{k}/2 & \delta k\beta_{k} \\ -\delta & \mathrm{I}_{d} - (\delta k)^{2}\zeta_{k}/2 \end{pmatrix} \right] \right| \end{aligned}$$

$$\leq \frac{21}{40}\delta^{3}k(k+1) + \frac{\delta^{3}}{4}|\nabla\Phi_{\delta}^{-k}(z)| + \left(1 + \frac{21}{20}\delta\right)C_{k},$$

where we used (44) to bound $|\Re|$, (41) for the second term and the induction hypothesis (42) for the third one (with the same computation as in (43) for the factor $1 + \frac{21}{20}\delta$). In view of (43), we finally obtain

$$(\star) \leq C_{k+1} := \frac{21}{40} \delta^3 k(k+1) + \frac{\delta^3}{4} \left(1 + \frac{21}{20} \delta k + C_k \right) + \left(1 + \frac{21}{20} \delta \right) C_k$$

$$\leq \frac{21}{40} \delta^3 (k+1)^2 + \left(1 + \frac{11}{10} \delta \right) C_k .$$

which allows to obtain the induction hypothesis (42) at the next step. From this, since $C_1 \leq 21\delta^3/40$,

$$C_K \leqslant \frac{21}{40} \delta^3 \sum_{k=0}^{K-1} (k+1)^2 \left(1 + \frac{11}{10} \delta\right)^{K-1-k} \leqslant \frac{21}{40 \times 3} (\delta K)^3 e^{11\delta K/10} \leqslant \frac{1}{3} (\delta K)^3.$$

which gives the desired estimate.

In particular, thanks to the estimate on $|\Psi|$ of Lemma 11 and to Lemma 8 we can bound the Fisher term appearing in Lemma 10 to get the following somewhat more explicit estimate.

Corollary 12 (Improved entropy inequality on Verlet step). For any $\varepsilon > 0$,

$$\begin{split} \mathcal{H}\left((\mathbf{V}_{\delta}^{K})^{*}h\right) - \mathcal{H}(h) &\leqslant 4\varepsilon K \delta \mathcal{I}\left(h\right) \\ &+ \sum_{j=0}^{K-1} \left[3\delta\varepsilon \mathrm{Er}_{1}^{\delta,j}\left(h\right) + \varepsilon \int_{0}^{\delta} \mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \mathrm{d}s + \frac{1}{2\varepsilon} \mathrm{Er}_{2}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \right] \,. \end{split}$$

Proof. For $j \in [0, K-1]$ and $s \in [0, \delta]$, writing $\Psi_s = \nabla \Phi_s^{-1} \circ \Phi_s$ and $\Psi_j = \nabla \Phi_{\delta}^{-j} \circ \Phi_{\delta}^{j}$, applying twice Lemma 8 (with $\tilde{\mathbf{A}} = I_{2d}$ and $\varepsilon = 1$) and using the bound (40) (with $s \leq K\delta \leq 1/10$) on $|\Psi_s|$ and $|\Psi_j|$, which implies that these quantities are smaller than 3/2 in sup norm,

$$\begin{split} \mathcal{I}\left((\mathbf{V}_{\delta}^{j}\mathbf{V}_{s}^{1})^{*}h\right) &\leqslant 2\mathcal{I}_{\Psi_{s}}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) + 2\mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \\ &\leqslant 2\|\Psi_{s}\|_{\infty}^{2}\mathcal{I}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) + 2\mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \\ &\leqslant 3\mathcal{I}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) + 2\mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \\ &\leqslant 6\mathcal{I}_{\Psi_{j}}(h) + 6\mathrm{Er}_{1}^{\delta,j}(h) + 2\mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) \\ &\leqslant 8\mathcal{I}(h) + 6\mathrm{Er}_{1}^{\delta,j}(h) + 2\mathrm{Er}_{1}^{s,1}\left((\mathbf{V}_{\delta}^{j})^{*}h\right) . \end{split}$$

Plugging this bound in the result of Lemma 10 gives the claimed estimate.

3.4 Numerical error

The goal of this section is to bound the numerical errors appearing in Lemmas 8 and 10. At various places, we use the following inequality (obtained by convexity): for $\alpha \ge 1$ and nonnegative sequences $(a_i)_{1 \le i \le I}$ and $(w_i)_{1 \le i \le I}$, it holds

$$\left(\sum_{i=1}^{I} w_i a_i\right)^{\alpha} \leqslant \left(\sum_{i=1}^{I} w_i\right)^{\alpha-1} \sum_{i=1}^{I} w_i a_i^{\alpha}.$$
(45)

3.4.1 Error in the Fisher term

Lemma 13. Under H1 with L = 1 and H2, it holds, for any $z \in \mathbb{R}^d$,

$$\begin{split} |\nabla \Box H(z)|^2 &\leqslant 2\delta^6 K \sum_{k=0}^{K-1} \left[N_4^6(v_k) + N_4^6(\nabla U(x_k)) + N_3^4(v_k) + N_3^4(\nabla U(x_k)) + |v_k|^2 + |\nabla U(x_k)|^2 \right], \end{split}$$

where $(x_k, v_k) = \Phi_{\delta}^{-k}(z)$ for $k \in [[0, K]]$.

Proof. In view of the decomposition

$$\Box H = H - H \circ \Phi_{\delta}^{-K} = \sum_{k=1}^{K} \left(H \circ \Phi_{\delta}^{-k+1} - H \circ \Phi_{\delta}^{-k} \right) \,,$$

which leads to the upper bound

$$|\nabla \Box H|^2 \leqslant K \sum_{k=1}^{K} \left| \nabla \left[H \circ \Phi_{\delta}^{-k+1} - H \circ \Phi_{\delta}^{-k} \right] \right|^2.$$

it suffices to establish the estimate for K = 1. Recall $H(x, v) = U(x) + |v|^2/2$ and recall from (29) that

$$(x_1, v_1) = \Phi_{\delta}^{-1}(x, v)$$

= $\left(x - \delta v - \frac{\delta^2}{2} \nabla U(x), v + \frac{\delta}{2} \left[\nabla U(x) + \nabla U \left(x - \delta v - \frac{\delta^2}{2} \nabla U(x)\right) \right] \right).$

Hence, starting with the gradient with respect to the velocities,

$$\nabla_{v} \left(H - H \circ \Phi_{\delta}^{-1} \right) (x, v)$$
$$= v - \left[-\delta \nabla U(x_{1}) + \left(I - \frac{\delta^{2}}{2} \nabla^{2} U(x_{1}) \right) v_{1} \right]$$

$$= \frac{\delta}{2} \left(\nabla U(x_1) - \nabla U(x) \right) + \frac{\delta^2}{2} \nabla^2 U(x_1) v + \frac{\delta^3}{4} \nabla^2 U(x_1) \left(\nabla U(x) + \nabla U(x_1) \right) \\ = \frac{\delta}{2} \left(\nabla U(x_1) - \nabla U(x) + \nabla^2 U(x_1) (x - x_1) \right) + \frac{\delta^3}{4} \nabla^2 U(x_1) \nabla U(x_1) \,.$$

Notice that (8) implies that

$$\left|\nabla U(x+y) - \nabla U(x) - \nabla^2 U(x)y\right| = \left|\int_0^1 \left(\nabla^2 U(x+sy) - \nabla^2 U(x)\right)y \,\mathrm{d}s\right| \le \frac{1}{2} N_3^2(y) \,. \tag{46}$$

Using this bound and the assumption $\|\nabla^2 U\|_{\infty} \leq L = 1$, we have thus obtained that

$$\begin{aligned} \left| \nabla_{v} \left(H - H \circ \Phi_{\delta}^{-1} \right) (x, v) \right| &\leq \frac{\delta}{4} \mathrm{N}_{3}^{2} (x - x_{1}) + \frac{\delta^{3}}{4} |\nabla U(x_{1})| \\ &\leq \frac{\delta^{3}}{2} \mathrm{N}_{3}^{2} (v) + \frac{\delta^{5}}{8} \mathrm{N}_{3}^{2} (\nabla U(x)) + \frac{\delta^{3}}{4} \left(|\nabla U(x)| + |x_{1} - x| \right) \\ &\leq \frac{\delta^{3}}{2} \mathrm{N}_{3}^{2} (v) + \frac{\delta^{5}}{8} \mathrm{N}_{3}^{2} (\nabla U(x)) + \frac{\delta^{4}}{4} |v| + \left(\frac{\delta^{3}}{4} + \frac{\delta^{5}}{8} \right) |\nabla U(x)| \,. \end{aligned}$$

With $\delta \leqslant 1/10$ and a Cauchy–Schwarz inequality,

$$|\nabla_{v} \left(H - H \circ \Phi_{\delta}^{-1} \right) (x, v)|^{2} \leqslant \delta^{6} \left(\frac{1}{2} N_{3}^{2}(v) + \frac{1}{800} N_{3}^{2} (\nabla U(x)) + \frac{1}{40} |v| + \frac{201}{800} |\nabla U(x)| \right)^{2} \\ \leqslant \frac{\delta^{6}}{3} \left(N_{3}^{4}(v) + N_{3}^{4} (\nabla U(x)) + |v|^{2} + |\nabla U(x)|^{2} \right).$$
(47)

Turning to the gradient with respect to the positions,

$$\begin{aligned} \nabla_x \left(H - H \circ \Phi_{\delta}^{-1} \right) (x, v) \\ &= \nabla U(x) - \left[\left(I - \frac{\delta^2}{2} \nabla^2 U(x) \right) \nabla U(x_1) + \frac{\delta}{2} \left(\nabla^2 U(x) + \left(I - \frac{\delta^2}{2} \nabla^2 U(x) \right) \nabla^2 U(x_1) \right) v_1 \right] \\ &= (*) + (**) + (***), \end{aligned}$$

with, organizing terms depending on their orders in δ , and using $\delta v_1 = x - x_1 + \delta \nabla U(x_1)/2$,

$$(*) = \nabla U(x) - \nabla U(x_1) - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x_1) \right) (x - x_1),$$

$$(**) = \frac{\delta^2}{4} \left(\nabla^2 U(x) - \nabla^2 U(x_1) \right) \nabla U(x_1),$$

$$(***) = \frac{\delta^3}{4} \nabla^2 U(x) \nabla^2 U(x_1) v_1.$$

Using the bound (9) on the fourth derivative of U and (45),

$$|(*)| \leq N_4^3(x - x_1) \leq \delta^3 \left(N_4(v) + \delta N_4(\nabla U(x)) \right)^3$$

$$\leqslant \delta^3 \left(1+\delta\right)^2 \left(N_4^3(v)+\delta N_4^3(\nabla U(x))\right) \,.$$

From the bound (8) on the third derivative of U and that $\|\nabla^2 U\|_{\infty} \leq L = 1$

$$\begin{aligned} |(**)| &\leq \frac{\delta^2}{4} |(\nabla^2 U(x) - \nabla^2 U(x_1)) \nabla U(x)| + \frac{\delta^2}{2} |\nabla U(x_1) - \nabla U(x)| \\ &\leq \frac{\delta^2}{4} N_3(x - x_1) N_3(\nabla U(x)) + \frac{\delta^2}{2} |x_1 - x| \\ &\leq \frac{\delta^3}{8} N_3^2(v) + \frac{\delta^3 + \delta^4}{8} N_3^2(\nabla U(x)) + \frac{\delta^3}{2} |v| + \frac{\delta^4}{4} |\nabla U(x)|. \end{aligned}$$

Finally,

$$|(***)| \leq \frac{\delta^{3}}{4} |v_{1}| \leq \frac{\delta^{3}}{4} |v| + \frac{\delta^{4}}{4} |\nabla U(x)| + \frac{\delta^{4}}{8} |x_{1} - x|$$
$$\leq \frac{2\delta^{3} + \delta^{5}}{8} |v| + \frac{4\delta^{4} + \delta^{6}}{16} |\nabla U(x)|.$$

As a conclusion, using that $\delta \leq 1/10$ and Cauchy–Schwarz inequality,

$$\begin{split} |\nabla_x \left(H - H \circ \Phi_{\delta}^{-1} \right) (x, v)|^2 \\ &\leqslant \delta^6 \Big[(1 + \delta)^2 \left(N_4^3(v) + \delta N_4^3(\nabla U(x)) \right) + \frac{1}{8} N_3^2(v) \\ &\quad + \frac{1 + \delta}{8} N_3^2(\nabla U(x)) + \left(\frac{3}{4} + \frac{\delta^2}{8} \right) |v| + \delta \left(\frac{1}{2} + \frac{\delta^2}{16} \right) |\nabla U(x)| \Big]^2 \\ &\leqslant \frac{5}{3} \delta^6 \Big[N_4^6(v) + N_4^6(\nabla U(x)) + N_3^4(v) + N_3^4(\nabla U(x)) + |v|^2 + |\nabla U(x)|^2 \Big] \,, \end{split}$$

which, combined with (47), gives the desired conclusion.

Integrating the bound of Lemma 13 with respect to $h\mu$ immediately yields the following (recall the definitions of the error in Lemma 8 and of the function **M** in (10)).

Corollary 14. Under H_1 with L = 1 and H_2 , for any smooth positive relative density h,

$$\operatorname{Er}_{1}^{\delta,K}(h) \leqslant 2\delta^{6}K \sum_{k=0}^{K-1} \int_{\mathbb{R}^{2d}} \left(\mathbf{M} \circ \Phi_{\delta}^{k} \right) h \mathrm{d}\mu$$

In particular, assuming moreover H^3 , and writing $h_n = (\mathbb{P}^n_{\omega})^* h_0$ for $n \in \mathbb{N}$,

$$\operatorname{Er}_{1}^{\delta,K}(\mathrm{D}_{\eta}h_{n}) \leq 2\delta^{6}K^{2}\left(C_{1}\mathrm{e}^{-\rho n\mathsf{T}}\nu_{0}(\mathfrak{W})+C_{2}\right) ,$$

$$\operatorname{Er}_{1}^{\delta,1}\left((\mathrm{V}_{\delta}^{j})^{*}\mathrm{D}_{\eta}h_{n}\right) \leq 2\delta^{6}\left(C_{1}\mathrm{e}^{-\rho n\mathsf{T}}\nu_{0}(\mathfrak{W})+C_{2}\right) ,$$

for all $j \in [0, K]$.

Remark 15. If in H_2 we only assume (8) but not (9), from (8), using that $\nabla U(x+y) - \nabla U(x) = \int_0^1 \nabla^2 U(x+sy) y \, ds$ (and distinguishing the integration on [0, 1/2] and on [1/2, 1]), we deduce that for all $x, y \in \mathbb{R}^d$,

$$\left|\nabla U(x+y) - \nabla U(x) - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x+y)\right) y\right| \leqslant \frac{1}{4} \mathcal{N}_3^2(y) \,.$$

We can thus use this inequality instead of (9) in Lemma 13 (which is the only place where (9) intervenes) and replace N_4^6 by N_3^4 in the definition (10) of **M**. Following then the rest of the proof, we see that Theorem 1 still holds in this case, but with δ^4 replaced by δ^2 as the error terms in Corollary 14 are of order δ^4 instead of δ^6 .

3.4.2 Error on the entropy term

It suffices to consider the case K = 1, as the general case is deduced from this one by summing up estimates. Recall that F_H is defined in (38).

Lemma 16. Under H1 with L = 1 and H2, for all $z \in \mathbb{R}^d$ and $s \in [0, \delta]$,

$$|\partial_s \Phi_s(z) - F_H(\Phi_s(z))|^2 \leq s^4 \left(\frac{1}{3}|v|^2 + \frac{1}{7}|\nabla U(x)|^2 + \frac{1}{7}N_3^4(v) + \frac{1}{140}N_3^4(\nabla U(x))\right)$$

Proof. Decomposing $z_s = \Phi_s(x, v)$ as

$$z_s = \begin{pmatrix} x_s \\ v_s \end{pmatrix} = \begin{pmatrix} x + sv - \frac{s^2}{2} \nabla U(x) \\ v - \frac{s}{2} \left[\nabla U(x) + \nabla U \left(x + sv - \frac{s^2}{2} \nabla U(x) \right) \right] \end{pmatrix},$$

we compute

$$\begin{aligned} (\partial_s \Phi_s)(z) - F_H(z_s) &= \begin{pmatrix} v - s \nabla U(x) \\ -\frac{1}{2} \left(\nabla U(x) + \nabla U(x_s) \right) - \frac{s}{2} \nabla^2 U(x_s) \frac{dx_s}{ds} \end{pmatrix} - \begin{pmatrix} v_s \\ -\nabla U(x_s) \end{pmatrix} \\ &= \frac{1}{2} \begin{pmatrix} s \left(\nabla U(x_s) - \nabla U(x) \right) \\ \nabla U(x_s) - \nabla U(x) - s \nabla^2 U(x_s) \left(v - s \nabla U(x) \right) \end{pmatrix}. \end{aligned}$$

As a consequence, using (46) and $\|\nabla^2 U\|_{\infty} \leq 1$,

$$\begin{aligned} |(\partial_s \Phi_s)(z) - F_H(z_s)|^2 &\leqslant \frac{s^2}{4} |x_s - x|^2 + \frac{1}{2} |\nabla U(x_s) - \nabla U(x) - \nabla^2 U(x_s)(x_s - x)|^2 \\ &+ \frac{1}{2} |x_s - x - s(v - s \nabla U(x))|^2 \\ &\leqslant \frac{s^2}{4} |x_s - x|^2 + \frac{1}{8} N_3^4(x_s - x) + \frac{s^4}{8} |\nabla U(x)|^2 \,. \end{aligned}$$

Using that $s \leq \delta \leq 1/10$, we bound

$$N_3^4(x_s - x) \leqslant \left(sN_3(v) + \frac{s}{20}N_3(\nabla U(x))\right)^4 \leqslant \frac{21^3}{20^3}s^4\left(N_3^4(v) + \frac{1}{20}N_3^4(\nabla U(x))\right)$$

and similarly $|x_s - x|^2 \leq \frac{21}{20s^2} (|v|^2 + |\nabla U(x)|^2/20)$, which allows to conclude.

Again, integrating the previous bound as

$$\operatorname{Er}_{2}(\delta,h) = \int_{0}^{\delta} \left(\int_{\mathbb{R}^{2d}} |\partial_{s}\Phi_{s} - F_{H} \circ \Phi_{s}|^{2} h \, d\mu \right) \, \mathrm{d}s \leqslant \frac{1}{3} \int_{0}^{\delta} \left(\int_{\mathbb{R}^{2d}} s^{4} \mathbf{M} \, h \, d\mu \right) \, \mathrm{d}s \, \mathrm{d}s$$

immediately yields the following result (recall the definitions of the Error 2 in Lemma 10 and of the function \mathbf{M} in (10)).

Corollary 17. Under H_1 with L = 1 and H_2 , for any smooth positive relative density h,

$$\operatorname{Er}_2(h) \leqslant \frac{\delta^5}{15} \int_{\mathbb{R}^{2d}} \mathbf{M} \, h \, d\mu$$

In particular, assuming moreover $H_{\mathfrak{s}}^{\mathfrak{s}}$, and writing $h_n = (\mathbb{P}^n_{\omega})^*$ for $n \in \mathbb{N}$ then, for any $j \in [\![0, K-1]\!]$,

$$\operatorname{Er}_{2}((\operatorname{V}_{\delta}^{j})^{*}\operatorname{D}_{\eta}h_{n}) \leqslant \frac{\delta^{5}}{15} \left(C_{1}(1-\rho K\delta)^{n}\nu_{0}(\mathfrak{W})+C_{2}\right) \,.$$

Acknowledgments

The works of P.M. and G.S. are supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (project EMC2, grant agreement No 810367); and by the Agence Nationale de la Recherche, under grant SWIDIMS (ANR-20-CE40-0022) for P.M., and ANR-19-CE40-0010-01 (QuAMProcs) and ANR-21-CE40-0006 (SINEQ) for G.S. A.O.D. would like to thank the Isaac Newton Institute for Mathematical Sciences for support and hospitality during the programme "The mathematical and statistical foundation of future data-driven engineering" where work on this paper was undertaken. This work was supported by EPSRC grant number EP/R014604/1.

References

- Assyr Abdulle, Gilles Vilmart, and Konstantinos C Zygalakis. Long time accuracy of Lie–Trotter splitting methods for Langevin dynamics. SIAM Journal on Numerical Analysis, 53(1):1–16, 2015.
- [2] Christophe Andrieu, Nando De Freitas, Arnaud Doucet, and Michael I Jordan. An introduction to MCMC for machine learning. *Machine learning*, 50:5–43, 2003.

- [3] Christophe Andrieu, Alain Durmus, Nikolas Nüsken, and Julien Roussel. Hypocoercivity of piecewise deterministic Markov process-Monte Carlo. The Annals of Applied Probability, 31(5):2478 – 2517, 2021.
- [4] Dominique Bakry, Ivan Gentil, and Michel Ledoux. Analysis and geometry of Markov diffusion operators, volume 348 of Grundlehren der Mathematischen Wissenschaften /Fundamental Principles of Mathematical Sciences]. Springer, Cham, 2014.
- [5] Nawaf Bou-Rabee and Andreas Eberle. Mixing time guarantees for unadjusted Hamiltonian Monte Carlo. *Bernoulli*, 29(1):75 104, 2023.
- [6] Nawaf Bou-Rabee, Andreas Eberle, and Raphael Zimmer. Coupling and convergence for Hamiltonian Monte Carlo. The Annals of Applied Probability, 30(3):1209 – 1250, 2020.
- [7] Nawaf Bou-Rabee and Houman Owhadi. Long-run accuracy of variational integrators in the stochastic context. *SIAM J. Numer. Anal.*, 48:278–297, 2010.
- [8] Nawaf Bou-Rabee and Katharina Schuh. Convergence of unadjusted Hamiltonian Monte Carlo for mean-field models. *Electronic Journal of Probability*, 28(none):1 – 40, 2023.
- [9] Giovanni Bussi and Michele Parrinello. Accurate sampling using Langevin dynamics. *Physical Review E*, 75(5):056707, 2007.
- [10] Louis-Pierre Chaintron and Antoine Diez. Propagation of chaos: A review of models, methods and applications. II. Applications. *Kinetic and Related Models*, 15(6):1017– 1173, 2022.
- [11] Subrahmanyan Chandrasekhar. Stochastic problems in physics and astronomy. Reviews of modern physics, 15(1):1, 1943.
- [12] Yuansi Chen, Raaz Dwivedi, Martin J. Wainwright, and Bin Yu. Fast mixing of Metropolized Hamiltonian Monte Carlo: Benefits of multi-step gradients. *Journal of Machine Learning Research*, 21(92):1–72, 2020.
- [13] Xiang Cheng, Niladri S Chatterji, Peter L Bartlett, and Michael I Jordan. Underdamped Langevin MCMC: A non-asymptotic analysis. In *Conference on learning* theory, pages 300–323. PMLR, 2018.
- [14] Patrick L Combettes. Perspective functions: Properties, constructions, and examples. Set-Valued and Variational Analysis, 26:247–264, 2018.
- [15] Arnak S. Dalalyan. Theoretical guarantees for approximate sampling from smooth and log-concave densities. *Journal of the Royal Statistical Society. Series B (Statistical Methodology)*, pages 651–676, 2017.

- [16] Arnak S. Dalalyan and Lionel Riou-Durand. On sampling from a log-concave density using kinetic langevin diffusions. *Bernoulli*, 26(3):1956–1988, 2020.
- [17] Jean Dolbeault, Axel Klar, Clément Mouhot, and Christian Schmeiser. Exponential rate of convergence to equilibrium for a model describing fiber lay-down processes. *Appl. Math. Res. eXpress*, 2013(2):165–175, 2013.
- [18] Alain Durmus and Andreas Eberle. Asymptotic bias of inexact Markov chain Monte Carlo methods in high dimension. to appear in Annals of Applied Probability, 2024.
- [19] Alain Durmus, Aurélien Enfroy, Éric Moulines, and Gabriel Stoltz. Uniform minorization condition and convergence bounds for discretizations of kinetic Langevin dynamics. arXiv e-prints, page arXiv:2107.14542, July 2021.
- [20] Alain Durmus and Éric Moulines. Nonasymptotic convergence analysis for the unadjusted Langevin algorithm. Annals of Applied Probability, 27(3):1551–1587, 2017.
- [21] Andreas Eberle. Reflection coupling and Wasserstein contractivity without convexity. Comptes Rendus Mathematique, 349(19):1101–1104, 2011.
- [22] Andreas Eberle, Arnaud Guillin, and Raphael Zimmer. Couplings and quantitative contraction rates for Langevin dynamics. *The Annals of Probability*, 47(4):1982 – 2010, 2019.
- [23] Donald L. Ermak and Helen Buckholz. Numerical integration of the Langevin equation: Monte Carlo simulation. Journal of Computational Physics, 35(2):169–182, 1980.
- [24] Andrew Gelman, John B Carlin, Hal S Stern, David B Dunson, Aki Vehtari, and Donald B Rubin. *Bayesian data analysis.* CRC press, 2013.
- [25] Nicolaï Gouraud, Pierre Le Bris, Adrien Majka, and Pierre Monmarché. HMC and underdamped Langevin united in the unadjusted convex smooth case. arXiv preprint arXiv:2202.00977, 2022.
- [26] Ulf Grenander. Tutorial in pattern theory. Division of Applied Mathematics, Brown University, Providence, 1983.
- [27] Ulf Grenander and Michael I. Miller. Representations of knowledge in complex systems. J. Roy. Statist. Soc. Ser. B, 56(4):549–603, 1994. With discussion and a reply by the authors.
- [28] Martin Grothaus and Patrik Stilgenbauer. Hilbert space hypocoercivity for the Langevin dynamics revisited. *Methods Funct. Anal. Topology*, 22(2):152–168, 2016.

- [29] Arnaud Guillin, Wei Liu, Liming Wu, and Chaoen Zhang. Uniform Poincaré and logarithmic Sobolev inequalities for mean field particle systems. *The Annals of Applied Probability*, 32(3):1590 – 1614, 2022.
- [30] Ernst Hairer, Gerhard Wanner, and Christian Lubich. Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations, volume 31 of Springer Series in Computational Mathematics. Springer-Verlag, 2006.
- [31] Wilfred Keith Hastings. Monte Carlo sampling methods using Markov chains and their applications. *Biometrika*, 57(1):97–109, 04 1970.
- [32] Frédéric Hérau. Short and long time behavior of the Fokker-Planck equation in a confining potential and applications. J. Funct. Anal., 244(1):95–118, 2007.
- [33] Alessandra Iacobucci, Stefano Olla, and Gabriel Stoltz. Convergence rates for nonequilibrium Langevin dynamics. Ann. Math. Québec, 43(1):73–98, 2019.
- [34] Benedict Leimkuhler and Charles Matthews. Rational Construction of Stochastic Numerical Methods for Molecular Sampling. Applied Mathematics Research eXpress, 2013(1):34–56, 06 2012.
- [35] Benedict Leimkuhler, Daniel Paulin, and Peter A Whalley. Contraction and convergence rates for discretized kinetic Langevin dynamics. *arXiv preprint arXiv:2302.10684*, 2023.
- [36] Tony Lelievre and Gabriel Stoltz. Partial differential equations and stochastic methods in molecular dynamics. *Acta Numerica*, 25:681–880, 2016.
- [37] Yi-An Ma, Niladri S. Chatterji, Xiang Cheng, Nicolas Flammarion, Peter L. Bartlett, and Michael I. Jordan. Is there an analog of Nesterov acceleration for gradient-based MCMC? *Bernoulli*, 27(3):1942 – 1992, 2021.
- [38] Oren Mangoubi and Aaron Smith. Mixing of Hamiltonian Monte Carlo on strongly log-concave distributions: Continuous dynamics. The Annals of Applied Probability, 31(5):2019 – 2045, 2021.
- [39] Jonathan C. Mattingly, Andrew M. Stuart, and Desmond J. Higham. Ergodicity for sdes and approximations: locally Lipschitz vector fields and degenerate noise. *Stochastic Processes and their Applications*, 101(2):185–232, 2002.
- [40] Nicholas Metropolis, Arianna W Rosenbluth, Marshall N Rosenbluth, Augusta H Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6):1087–1092, 1953.
- [41] Pierre Monmarché. An entropic approach for Hamiltonian Monte Carlo: the idealized case. to appear in Annals of Applied Probability, 2024.

- [42] Pierre Monmarché. High-dimensional MCMC with a standard splitting scheme for the underdamped Langevin diffusion. *Electronic Journal of Statistics*, 15(2):4117 – 4166, 2021.
- [43] Radford M. Neal. MCMC using Hamiltonian dynamics. Handbook of Markov Chain Monte Carlo, 54:113–162, 2010.
- [44] Gareth O. Roberts and Richard L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341–363, 1996.
- [45] Peter J. Rossky, Jimmie D. Doll, and Harold L. Friedman. Brownian dynamics as smart monte carlo simulation. *The Journal of Chemical Physics*, 69(10):4628–4633, 1978.
- [46] Robert D. Skeel and Jesüs A. Izaguirre. An impulse integrator for Langevin dynamics. Molecular Physics, 100(24):3885–3891, 2002.
- [47] Denis Talay. Stochastic Hamiltonian systems: exponential convergence to the invariant measure, and discretization by the implicit Euler scheme. Markov Process. Related Fields, 8(2):163–198, 2002.
- [48] Santosh Vempala and Andre Wibisono. Rapid convergence of the unadjusted Langevin algorithm: Isoperimetry suffices. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, Advances in Neural Information Processing Systems, volume 32. Curran Associates, Inc., 2019.
- [49] Cédric Villani. Hypocoercivity. Mem. Amer. Math. Soc., 202(950):iv+141, 2009.
- [50] Cédric Villani. Optimal transport. Old and new, volume 338 of Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]. Springer-Verlag, Berlin, 2009.
- [51] Max Welling and Yee Whye Teh. Bayesian learning via stochastic gradient langevin dynamics. In Proceedings of the 28th International Conference on Machine Learning (ICML-11), pages 681–688, 2011.
- [52] Shunshi Zhang, Sinho Chewi, Mufan Li, Krishna Balasubramanian, and Murat A Erdogdu. Improved discretization analysis for underdamped langevin monte carlo. In *The Thirty Sixth Annual Conference on Learning Theory*, pages 36–71. PMLR, 2023.

A Proof of Proposition 2

Note first that, for $j \neq i$ and defining $w(x_i, x_j) = W^{(q)}(x_i - x_j)$, it holds

$$\nabla_{x_i, x_j}^2 w(x_i, x_j) = -\frac{1}{d_0} \nabla^2 W^{(q)}(x_i - x_j) \,,$$

while

$$\nabla_{x_i}^2 w(x_i, x_j) = \frac{1}{d_0} \sum_{j=1}^{d_0} \nabla^2 W^{(q)}(x_i - x_j) \,.$$

Therefore,

$$\begin{split} \sum_{i=1}^{d_0} \left| \sum_{j=1}^{d_0} \left(\nabla_{x_i, x_j}^2 w(x_i + y_i, x_j + y_j) - \nabla_{x_i, x_j}^2 w(x_i, x_j) \right) z_j \right|^2 \\ &= \frac{1}{d_0^2} \sum_{i=1}^{d_0} \left| \sum_{j \neq i} \left(\nabla^2 W^{(q)}(x_i + y_i - x_j - y_j) - \nabla^2 W^{(q)}(x_i - x_j) \right) (z_i - z_j) \right|^2 \\ &= \frac{1}{d_0^2} \sum_{i=1}^{d_0} \left| \sum_{j=1}^{d_0} \left(\nabla^2 W^{(q)}(x_i + y_i - x_j - y_j) - \nabla^2 W^{(q)}(x_i - x_j) \right) (z_i - z_j) \right|^2. \end{split}$$

We can now compute

$$\begin{split} \left| \left(\nabla^2 U(x+y) - \nabla^2 U(x) \right) z \right|^2 \\ &= \sum_{i=1}^{d_0} \left| \sum_{j=1}^{d_0} \left(\nabla_{x_i, x_j}^2 U(x+y) - \nabla_{x_i, x_j}^2 U(x) \right) z_j \right|^2 \\ &\leq (1+\epsilon) \sum_{i=1}^{d_0} \left| \left(\nabla^2 U^{(q)}(x_i+y_i) - \nabla^2 U^{(q)}(x_i) \right) z_i \right|^2 \\ &+ \frac{\epsilon^2 (1+\epsilon^{-1})}{d_0^2} \sum_{i=1}^{d_0} \left| \sum_{j=1}^{d_0} \left(\nabla^2 W^{(q)}(x_i+y_i-x_j-y_j) - \nabla^2 W^{(q)}(x_i-x_j) \right) (z_i-z_j) \right|^2 \\ &\leq (1+\epsilon) \left\| D^3 U^{(q)} \right\|_{\infty}^2 \sum_{i=1}^{d_0} |y_i|^2 |z_i|^2 + \frac{\epsilon(1+\epsilon)}{d_0} \left\| D^3 W^{(q)} \right\|_{\infty}^2 \sum_{i=1}^{d_0} \sum_{j=1}^{d_0} |y_i-y_j|^2 |z_i-z_j|^2 \\ &\leq (1+\epsilon) \left(\left\| D^3 U^{(q)} \right\|_{\infty}^2 + 16\epsilon \left\| D^3 W^{(q)} \right\|_{\infty}^2 \right) \sqrt{\sum_{i=1}^{d_0} |y_i|^4 \sum_{j=1}^{d_0} |z_j|^4} \,, \end{split}$$

where we used

$$\frac{1}{d_0} \sum_{i=1}^{d_0} \sum_{j=1}^{d_0} |y_i - y_j|^2 |z_i - z_j|^2 \leq \frac{4}{d_0} \sum_{i=1}^{d_0} \sum_{j=1}^{d_0} \left(|y_i|^2 + |y_j|^2 \right) \left(|z_i|^2 + |z_j|^2 \right)$$
$$\leq 8 \sum_{i=1}^{d_0} |y_i|^2 |z_i|^2 + \frac{8}{d_0} \sum_{i=1}^{d_0} |y_i|^2 \sum_{j=1}^{d_0} |z_j|^2$$

$$\leq 16 \sqrt{\sum_{i=1}^{d_0} |y_i|^4 \sum_{j=1}^{d_0} |z_j|^4},$$

to conclude the proof of (8). Following similar computations,

$$\begin{split} \left| \nabla U(x+y) - \nabla U(x) - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x+y) \right) y \right|^2 \\ &= \left| \left(\int_0^1 \nabla^2 U(x+sy) \, \mathrm{d}s - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x+y) \right) \right) y \right|^2 \\ &= \sum_{i=1}^{d_0} \left| \sum_{j=1}^{d_0} \left(\int_0^1 \nabla^2_{x_i,x_j} U(x+sy) \, \mathrm{d}s - \frac{1}{2} \left(\nabla^2_{x_i,x_j} U(x) + \nabla^2_{x_i,x_j} U(x+y) \right) \right) y_j \right|^2 \\ &\leqslant (1+\epsilon) \sum_{i=1}^{d_0} \left| \left(\int_0^1 \nabla^2 U^{(q)}(x_i+sy_i) \, \mathrm{d}s - \frac{1}{2} \left(\nabla^2 U^{(q)}(x_i) + \nabla^2 U^{(q)}(x_i+y_i) \right) \right) y_i \right|^2 \\ &+ \frac{\epsilon + \epsilon^2}{d_0} \sum_{i,j=1}^{d_0} \left| \left(\int_0^1 \nabla^2 W^{(q)}(x_i-x_j+s(y_i-y_j)) \, \mathrm{d}s \right. \\ &\left. - \frac{1}{2} \left(\nabla^2 W^{(q)}(x_i-x_j) + \nabla^2 W^{(q)}(x_i+y_i-x_j-y_j) \right) \right) (y_i-y_j) \right|^2 \, . \end{split}$$

Using that, for all $f \in C^2([0,1], \mathbb{R})$, integrating twice by parts,

$$\int_0^1 f(s) ds - \frac{f(0) + f(1)}{2} = \frac{1}{2} \int_0^1 s(1-s) f''(s) ds,$$

fixing $i \in [\![1, d_0]\!]$, $x_i, y_i \in \mathbb{R}^q$ and denoting by \mathbf{e}_{ℓ} the ℓ -th canonical vector of \mathbb{R}^q ,

$$\begin{aligned} \left| \mathbf{e}_{\ell} \cdot \left(\int_{0}^{1} \nabla^{2} U^{(q)}(x_{i} + sy_{i}) \, \mathrm{d}s - \frac{1}{2} \left(\nabla^{2} U^{(q)}(x_{i}) + \nabla^{2} U^{(q)}(x_{i} + y_{i}) \right) \right) y_{i} \right| \\ &= \frac{1}{2} \left| \int_{0}^{1} s(1 - s) \mathrm{D}^{4} U^{(q)}(x_{i} + sy_{i}) \{ \mathbf{e}_{\ell}, y_{i}, y_{i}, y_{i} \} \, \mathrm{d}s \right| \\ &\leq \frac{1}{12} \| \mathrm{D}^{4} U^{(q)} \|_{\infty} |y_{i}|^{3} \, . \end{aligned}$$

The terms involving $W^{\left(q\right)}$ are treated in a similar manner and we get

$$\begin{aligned} \left| \nabla U(x+y) - \nabla U(x) - \frac{1}{2} \left(\nabla^2 U(x) + \nabla^2 U(x+y) \right) y \right|^2 \\ \leqslant (1+\epsilon) \sum_{i=1}^{d_0} \frac{q}{144} \left\| \mathbf{D}^4 U^{(q)} \right\|_{\infty}^2 |y_i|^6 + \frac{\epsilon+\epsilon^2}{d_0} \sum_{i,j=1}^{d_0} \frac{q}{144} \left\| \mathbf{D}^4 W^{(q)} \right\|_{\infty}^2 |y_i - y_j|^6 \end{aligned}$$

$$\leq \left(\frac{1+\epsilon}{144} \left\| \mathbf{D}^{4} U^{(q)} \right\|_{\infty}^{2} + \frac{4(\epsilon+\epsilon^{2})}{9} \left\| \mathbf{D}^{4} W^{(q)} \right\|_{\infty}^{2} \right) q \sum_{i=1}^{d_{0}} |y_{i}|^{6} \,,$$

which concludes the proof of (9).

Finally, to bound **M**, for $\ell \in \{2, 4, 6\}$, using that $\nabla U^{(q)}(0) = 0 = \nabla W^{(q)}(0)$, we write

$$\begin{split} |\nabla_{x_i} U(x)|^{\ell} &= \left| \nabla U^{(q)}(x_i) + \frac{2\epsilon}{d_0} \sum_{j=1}^{d_0} \nabla W^{(q)}(x_i - x_j) \right|^{\ell} \\ &\leq 2^{\ell-1} |\nabla U^{(q)}(x_i)|^{\ell} + \frac{2^{2\ell-1}\epsilon^{\ell}}{d_0} \sum_{j=1}^{d_0} \left| \nabla W^{(q)}(x_i - x_j) \right|^{\ell} \\ &\leq 2^{\ell-1} \left\| \nabla^2 U^{(q)} \right\|_{\infty}^{\ell} |x_i|^{\ell} + \frac{2^{2\ell-1}\epsilon^{\ell}}{d_0} \left\| \nabla^2 W^{(q)} \right\|_{\infty}^{\ell} \sum_{j=1}^{d_0} |x_i - x_j|^{\ell} . \end{split}$$

Using $|x_i - x_j|^{\ell} \leq 2^{\ell-1}(|x_i|^{\ell} + |x_j|^{\ell})$ in the last term, we end up with

$$\sum_{i=1}^{d_0} |\nabla_{x_i} U(x)|^{\ell} \leq \left(2^{\ell-1} \left\| \nabla^2 U^{(q)} \right\|_{\infty}^{\ell} + 2^{3\ell} \epsilon^{\ell} \left\| \nabla^2 W^{(q)} \right\|_{\infty}^{\ell} \right) \sum_{i=1}^{d_0} |x_i|^{\ell} \,,$$

which leads to the claimed result.

B Uniform-in-time moment bounds

In all this section, which is devoted to the proof of Theorem 3, assumption H 4 is enforced. We denote by $L_1 = \max(\|\nabla^2 U^{(q)}\|_{\infty}, \|\nabla^2 W^{(q)}\|_{\infty})$. In fact, using a rescaling as in Section 2.2.1, we assume without loss of generality that $L_1 \leq 1$. The structure of the proof is the following. After providing some useful preliminary estimates, we describe the class of Lyapunov functions we consider, state the main result in Theorem 18, and discuss that it implies Theorem 3. The remainder of the section is then devoted to the proof of Theorem 18. We start by making precise in Lemma 19 how the Lyapunov function evolves over one Verlet step; and then over several steps in Lemma 21, relying on Lemma 20 which quantifies how much prefactors change when performing several steps. We can then consider the addition of the fluctuation-dissipation step, by first incorporating the momentum update in Lemma 22, then averaging over the momentum updates in Lemma 25, relying on some moment bounds provided by Lemma 24. The proof of Theorem 18 can then be concluded.

Let us emphasize that our proofs are written both for kinetic Langevin dynamics, for which $\eta > 0$ (more precisely $1 - \eta \ge 0$ is of order δ), and for HMC, for which $\eta = 0$. This second case requires more precise estimates.

Preliminary estimates. We state a few consequences of H4 which will be useful in the remainder of this section. We additionally assume without loss of generality that $U^{(q)}(0) = 0$. First, (16) and the inequality

$$\left\langle x_1, \nabla U^{(q)}(x_1) \right\rangle \leqslant \frac{1}{2\mathfrak{m}} \left| \nabla U^{(q)}(x_1) \right|^2 + \frac{\mathfrak{m}}{2} |x_1|^2,$$

imply that

$$\forall x_1 \in \mathbb{R}^q, \qquad \left|\nabla U^{(q)}(x_1)\right|^2 \ge 2\mathfrak{m}\left(\frac{\mathfrak{m}}{2}|x_1|^2 - \mathfrak{M}\right). \tag{48}$$

In addition, since $\nabla U^{(q)}$ is L_1 -Lipschitz and $\nabla U^{(q)}(0) = 0$, the inequality $\langle x_1, \nabla U^{(q)}(x_1) \rangle \leq L_1 |x_1|^2$, and the drift condition (16) together imply, by taking the limit $|x_1| \to +\infty$, that

$$\mathbf{m} \leqslant L_1 \leqslant 1 \ . \tag{49}$$

Finally, since $\nabla U^{(q)}(0) = \nabla W^{(q)}(0) = 0$, we obtain that, for any $x_1 \in \mathbb{R}^q$

$$\left|\nabla U^{(q)}(x_1)\right| \le |x_1| , \qquad \left|\nabla W^{(q)}(x_1)\right| \le |x_1| , \qquad U^{(q)}(x_1) \le \frac{|x_1|^2}{2} .$$
 (50)

Construction of Lyapunov functions. Our analysis is based on the following Lyapunov functions. Under **H4**, for $\boldsymbol{\omega} = (K, \delta, \eta) \in \mathbb{N}^* \times \mathbb{R}^*_+ \times [0, 1)$ and $\ell \in \mathbb{N}^*$, we define $\mathfrak{M}^{(\ell)}_{\boldsymbol{\omega}} : \mathbb{R}^{2d} \to \mathbb{R}$ as: for any $x = (x_1, \ldots, x_{d_0})$ and $v = (v_1, \ldots, v_{d_0}) \in \mathbb{R}^d$,

$$\mathfrak{W}_{\boldsymbol{\omega}}^{(\ell)}(x,v) = \sum_{i=1}^{d_0} \left(\mathfrak{W}_{\boldsymbol{\omega},i}(x,v)\right)^{\ell} , \quad \mathfrak{W}_{\boldsymbol{\omega},i}(x,v) = \frac{\gamma_0^2}{2} |x_i|^2 + |v_i|^2 + \eta \gamma_0 \langle x_i, v_i \rangle + U^{(q)}(x_i) ,$$

where (recall (6)) $\gamma_0 = \frac{1-\eta}{\delta K}$. Note that since $\eta \in [0,1)$, using Young's inequality $\langle \mathbf{a}, \mathbf{b} \rangle \leq (\varepsilon/2) |\mathbf{a}|^2 + |\mathbf{b}|^2 / (2\varepsilon)$ with $\varepsilon = 2/3$ for the left inequality and $\varepsilon = 1$ for the right one, we get

$$\frac{1}{6} \left(\gamma_0^2 |x|^2 + |v|^2 \right) + \sum_{i=1}^{d_0} U^{(q)}(x_i) \leqslant \mathfrak{W}^{(1)}_{\omega}(x,v) \leqslant \frac{3}{2} \left(\gamma_0^2 |x|^2 + |v|^2 \right) + \sum_{i=1}^{d_0} U^{(q)}(x_i) .$$
(51)

By (51) and the fact $0 \leq U^{(q)}(x_1) \leq |x_1|^2/2$, the function \mathfrak{W} defined in (17) is equivalent to $\mathfrak{W}^{(3)}_{\boldsymbol{\omega}}$, more precisely there exists A > 0 which depends only on γ_0 such that, for all $(x, v) \in \mathbb{R}^{2d}$,

$$A^{-1}\left(d_0 + \mathfrak{W}^{(3)}_{\boldsymbol{\omega}}(x,v)\right) \leqslant d_0 + \mathfrak{W}(x,v) \leqslant A\left(d_0 + \mathfrak{W}^{(3)}_{\boldsymbol{\omega}}(x,v)\right).$$
(52)

Statement of the main result. The main result in this section is the following.

Theorem 18. Assume that H_4^4 holds with $L_1 = 1$ and let $\delta, T, \bar{\eta} > 0$. Then, for any $\ell \in \mathbb{N}$, $\epsilon > 0$, $\omega = (K, \delta, \eta)$, $\delta > 0$, $K \in \mathbb{N}^*$, $\eta \in [0, \bar{\eta})$ satisfying

$$\epsilon \leqslant \epsilon_1 \wedge \epsilon_2 , \qquad \delta \leqslant \delta_1 \wedge \delta_2 \wedge 1/[7 \times 8] \wedge \delta , \qquad K\delta \leqslant T_1 \wedge T_2 \wedge T_3 \wedge \overline{T} ,$$

for any $x, v \in \mathbb{R}^d$ and $k \in \{0, \ldots, K\}$,

$$P^{n}_{\omega} D_{\eta} V^{k}_{\delta} \mathfrak{W}^{(\ell)}_{\omega}(x,v) \leqslant C^{\ell}_{1,\ell} (1 - \rho_{\omega} K \delta)^{\ell n} \mathfrak{W}^{(\ell)}_{\omega}(x,v) + d_{0} (C_{2,\ell} q^{\ell} + C_{3,\ell}) \sup_{s \in [0,\bar{T}]} \frac{s}{1 - (1 - \rho_{\omega} s/8)^{\ell}}, \quad (53)$$

where $\epsilon_1, \epsilon_2, T_1, T_2, T_3, \delta_1, \delta_2$ are defined in (83)-(84)-(85)-(87)-(88),

$$\rho_{\omega} = 2^{5} [\gamma_{0} \wedge \{\eta \gamma_{0} + \gamma_{0}^{2} K \delta / 2\} \mathfrak{m}(2^{6} 3)^{-1}], \qquad (54)$$

and $C_{1,\ell}, C_{2,\ell}, C_{3,\ell}$ are given in (89). These constants do not depend on d_0, q and only depend on δ, K, η through γ_0 and ρ_{ω} .

The remainder of Section B is devoted to the proof of this result. Before proceeding with this proof, let us notice that Theorem 18 implies Theorem 3 in view of (52) and, as can be checked on their expressions, the constants $\epsilon_1, \epsilon_2, T_1, T_2, T_3, \delta_1, \delta_2, C_{1,\ell}, C_{2,\ell}, C_{3,\ell}$ appearing in Theorem 18 depend only on $\mathfrak{m}, \mathfrak{M}, \overline{T}, \overline{\delta}, \gamma_0$ and not on q, d_0, ϵ .

Evolution over one Verlet step. In order to prove Theorem 18, we will work with a Lyapunov function slightly more general than $\mathfrak{W}_{\omega,i}$. Consider, for any $\mathbf{o} = (a, b, c, b_0, e, f) \in \mathbb{R}^4$, $i \in \{1, \ldots, d_0\}$, and $x = (x_1, \ldots, x_q), v = (v_1, \ldots, v_q) \in \mathbb{R}^d$, the function

$$\mathfrak{V}_{\mathbf{o},i}(x,v) = a |x_i|^2 + b |v_i|^2 + c \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i) + e \frac{\epsilon}{d_0} \sum_{j=1}^{d_0} \left[|x_i - x_j|^2 + |v_i - v_j|^2 \right] + f$$

Notice that the structure of $\mathfrak{V}_{\mathbf{o},i}$ is similar to the one of $\mathfrak{W}_{\omega,i}$ except that we have added the parameters \mathbf{o} to be more flexible. That way, we will bound the value of $\mathfrak{V}_{\mathbf{o},i}$ after some Verlet steps in terms of $\mathfrak{V}_{\mathbf{o},i}$ with different parameters $\mathbf{\tilde{o}}$, so that the general forms of the function is preserved and we only have to keep track of the evolution of the coefficients \mathbf{o} .

Specifically, the following result makes precise how the Lyapunov function evolves over one Verlet step, by bounding it with a similar Lyapunov function with modified coefficients. Note that the coefficient b_0 in front of $U^{(q)}$ in the expression of $\mathfrak{V}_{\mathbf{o},i}$ is fixed, so that this parameter of the Lyapunov function does not change over the iterations. **Lemma 19.** Assume that H_4 holds and that $\epsilon \leq m/4$. Then, for any $x, v \in \mathbb{R}^{2d}$, $\delta \in (0, m/40]$, $i \in \{1, \ldots, d_0\}$ and $\mathbf{o} = (a, b, c, b_0, e, f) \in \mathbb{R}^6_+$ with $b_0 \leq b$, it holds

$$V_{\delta}\mathfrak{V}_{\mathbf{o},i}(x,v) \leqslant \mathfrak{V}_{\tilde{\mathbf{o}},i}(x,v)$$

where $\tilde{\mathbf{o}} = (\tilde{a}, \tilde{b}, \tilde{c}, b_0, \tilde{e}, \tilde{f})$ with

$$\begin{split} \tilde{a} &= a \left(1 - \frac{\mathbf{m}\delta^2}{2} \right) + 8b\delta^2 - b_0 \frac{\delta^2 \mathbf{m}^2}{2} - \frac{\delta \mathbf{m}c}{2} + \delta(b - b_0) \;, \\ \tilde{b} &= 2\delta^2 a + (1 + \epsilon\delta + 31\delta^2)b + 2b_0\delta^2 + \delta(1 + 10\delta)c + \delta(b - b_0) \;, \\ \tilde{c} &= 2\delta a + c \;, \\ \tilde{e} &= e + \delta(2a\delta + 2b + 2\delta b_0 + 2c + 7e) \;, \\ \tilde{f} &= f + a\delta^2 \mathbf{M} + 2b_0\delta^2 \mathbf{m} \mathbf{M} + \delta \mathbf{M}c \;. \end{split}$$

Proof. Note first that the assumption $\epsilon \leq m/4$ and the bound $m \leq 1$ from (49) imply that $\epsilon \leq 1/4$. Let $x, v \in \mathbb{R}^d$, $i \in \{1, \ldots, d_0\}$, $\delta \leq m/40$ and $\mathbf{o} = (a, b, c, b_0, e, f)$. To alleviate the notation, we introduce, for $j \in \{1, \ldots, d_0\}$, the partial derivative of U with respect to the j-th variable, namely the function $\psi_j : \mathbb{R}^d \to \mathbb{R}^d$ defined for any $\tilde{x} = (\tilde{x}_1, \ldots, \tilde{x}_{d_0}) \in \mathbb{R}^d$ as

$$\psi_j(\tilde{x}) = \nabla U^{(q)}(\tilde{x}_j) + \frac{\epsilon}{d_0} \sum_{j'=1}^{d_0} \nabla W^{(q)}(\tilde{x}_j - \tilde{x}_{j'}) .$$

With this notation, it holds by definition of V_{δ} that

$$V_{\delta}\mathfrak{V}_{\mathbf{o},i}(x,v) = \mathfrak{V}_{\mathbf{o},i}(y,p) , \qquad (55)$$

where $y = (y_1, \ldots, y_{d_0})$ and $p = (p_1, \ldots, p_{d_0})$ with, for $j \in \{1, \ldots, d_0\}$,

$$y_j = x_j + \delta v_j - \frac{\delta^2}{2} \psi_j(x) , \qquad p_j = v_j - \frac{\delta}{2} \{ \psi_j(x) + \psi_j(y) \}$$

We start by giving some useful preliminary estimates. First, by a discrete Cauchy–Schwarz inequality and (50),

$$|\psi_{i}(x) - \psi_{j}(x)|^{2} \leq 2 \left| \nabla U^{(q)}(x_{i}) - \nabla U^{(q)}(x_{j}) \right|^{2} + \frac{2\epsilon^{2}}{d_{0}} \sum_{k=1}^{d_{0}} \left| \nabla W^{(q)}(x_{k} - x_{i}) - \nabla W^{(q)}(x_{k} - x_{j}) \right|^{2} \leq 2 \left(1 + \epsilon^{2} \right) |x_{i} - x_{j}|^{2}.$$
(56)

Next, by two successive discrete Cauchy–Schwarz inequalities and (50), we have, for any $j \in \{1, \ldots, d_0\}$,

$$|y_j - x_j|^2 \leq \frac{\delta^4}{2} |\psi_j(x)|^2 + 2\delta^2 |v_j|^2$$

$$\leq 2 \left(\delta^{4} \left| \nabla U^{(q)}(x_{j}) \right|^{2} + \frac{\delta^{4} \epsilon^{2}}{d_{0}} \sum_{j'=1}^{d_{0}} \left| \nabla W^{(q)}(x_{j} - x_{j'}) \right|^{2} + \delta^{2} |v_{j}|^{2} \right)$$

$$\leq 2 \left(\delta^{4} |x_{j}|^{2} + \frac{\delta^{4} \epsilon^{2}}{d_{0}} \sum_{j'=1}^{d_{0}} |x_{j} - x_{j'}|^{2} + \delta^{2} |v_{j}|^{2} \right) .$$
(57)

In view of (50) and discrete Cauchy–Schwarz inequalities, it holds, for any $\tilde{x} = (\tilde{x}_1, \ldots, \tilde{x}_{d_0}), \tilde{y} = (y_1, \ldots, y_{d_0}) \in \mathbb{R}^{d_0}$ and $i \in \{1, \ldots, d_0\}$,

$$\begin{aligned} |\psi_{i}(\tilde{x}) - \psi_{i}(\tilde{y})|^{2} &\leq 2 |\tilde{x}_{i} - \tilde{y}_{i}|^{2} + \frac{2\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |\tilde{x}_{i} - \tilde{x}_{j} - (\tilde{y}_{i} - \tilde{y}_{j})|^{2} \\ &\leq 2 \left(1 + 2\epsilon^{2} \right) |\tilde{x}_{i} - \tilde{y}_{i}|^{2} + \frac{4\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |\tilde{x}_{j} - \tilde{y}_{j}|^{2} , \end{aligned}$$
(58)

as well as

$$|\psi_i(\tilde{x})|^2 \leq 2 |\tilde{x}_i|^2 + \frac{2\epsilon^2}{d_0} \sum_{j=1}^{d_0} |\tilde{x}_i - \tilde{x}_j|^2 .$$
(59)

As a result, using (57) to bound all the terms on right hand side of (58), we obtain

$$\begin{aligned} |\psi_{i}(x) - \psi_{i}(y)|^{2} &\leq 4 \left(1 + 2\epsilon^{2} \right) \left(\delta^{4} |x_{i}|^{2} + \delta^{4} \frac{\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \delta^{2} |v_{i}|^{2} \right) \\ &+ \frac{8\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} \left(\delta^{4} |x_{j}|^{2} + \frac{\delta^{4}\epsilon^{2}}{d_{0}} \sum_{j'=1}^{d_{0}} |x_{j} - x_{j'}|^{2} + \delta^{2} |v_{j}|^{2} \right) \\ &\leq 4 \left(1 + 2\epsilon^{2} \right) \left(\delta^{4} |x_{i}|^{2} + \delta^{4} \frac{\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \delta^{2} |v_{i}|^{2} \right) \\ &+ \frac{16\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} \left(\delta^{4} |x_{i}|^{2} + (1 + 2\epsilon^{2})\delta^{4} |x_{i} - x_{j}|^{2} + \delta^{2} |v_{i}|^{2} + \delta^{2} |v_{j} - v_{i}|^{2} \right) \\ &\leq 30 \left(\delta^{4} |x_{i}|^{2} + \frac{\delta^{4}\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \delta^{2} |v_{i}|^{2} + \frac{\epsilon^{2}\delta^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |v_{i} - v_{j}|^{2} \right) , \tag{60}$$

where we used $20 + 40\epsilon^2 \leq 30$ in the last inequality, and the second inequality has been obtained by using

$$\frac{1}{d_0} \sum_{j=1}^{d_0} |x_j|^2 \leq 2 |x_i|^2 + \frac{2}{d_0} \sum_{j=1}^{d_0} |x_j - x_i|^2,$$

and a similar inequality for v, as well as the inequality $|x_j - x_i + x_i - x_{j'}|^2 \leq 2|x_j - x_i|^2 + 2|x_i - x_{j'}|^2$ to write

$$\frac{1}{d_0^2} \sum_{j=1}^{d_0} \sum_{j'=1}^{d_0} |x_j - x_{j'}|^2 = \frac{1}{d_0^2} \sum_{j=1}^{d_0} \sum_{j'=1}^{d_0} |x_j - x_i + x_i - x_{j'}|^2 \le \frac{4}{d_0} \sum_{j=1}^{d_0} |x_j - x_i|^2 .$$

We can now turn to estimating the various terms in $\mathfrak{V}_{\tilde{\mathbf{0}},i}(x,v)$. In view of (57),

$$\begin{aligned} |y_i|^2 &= |x_i|^2 + 2\left\langle x_i, \delta v_i - \frac{\delta^2}{2}\psi_i(x) \right\rangle + |y_i - x_i|^2 \\ &\leq |x_i|^2 - \delta^2\left\langle x_i, \nabla U^{(q)}(x_i) \right\rangle + \delta^2\epsilon |x_i|^2 + \frac{\delta^2\epsilon}{d_0} \sum_{j=1}^{d_0} |x_i - x_j|^2 \\ &+ 2\delta\left\langle x_i, v_i \right\rangle + 2\left(\delta^4 |x_i|^2 + \frac{\delta^4\epsilon^2}{d_0} \sum_{j=1}^{d_0} |x_i - x_j|^2 + \delta^2 |v_i|^2\right) \,, \end{aligned}$$

where $\langle x_i, \nabla W^{(q)}(x_i - x_j) \rangle$ is bounded with a discrete Cauchy–Schwarz inequality and (50). Therefore, using first the drift condition (16) and the inequality $2\delta^2 \epsilon \leq 1$, and then $\max(\delta, \epsilon) \leq m/4 \leq 1/4$,

$$\begin{aligned} |y_{i}|^{2} &\leq \left(1 + \delta^{2} \epsilon + 2\delta^{4}\right) |x_{i}|^{2} + 2\delta \langle x_{i}, v_{i} \rangle + 2\delta^{2} |v_{i}|^{2} \\ &- \delta^{2} \left\langle x_{i}, \nabla U^{(q)}(x_{i}) \right\rangle + \frac{\delta^{2} \epsilon (1 + 2\delta^{2} \epsilon)}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \\ &\leq \left(1 - \delta^{2} \mathbf{m} + \delta^{2} \epsilon + 2\delta^{4}\right) |x_{i}|^{2} + 2\delta \langle x_{i}, v_{i} \rangle + 2\delta^{2} |v_{i}|^{2} + \delta^{2} \mathbf{M} + \frac{2\delta^{2} \epsilon}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \\ &\leq \left(1 - \frac{\delta^{2} \mathbf{m}}{2}\right) |x_{i}|^{2} + 2\delta \langle x_{i}, v_{i} \rangle + 2\delta^{2} |v_{i}|^{2} + \delta^{2} \mathbf{M} + \frac{2\delta^{2} \epsilon}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} . \end{aligned}$$

$$\tag{61}$$

Similarly, using a discrete Cauchy–Schwarz inequality to bound $\delta \langle v_i, \psi_i(x) - \psi_i(y) \rangle$,

$$\begin{aligned} |p_i|^2 &= |v_i|^2 - \delta \langle v_i, \psi_i(x) + \psi_i(y) \rangle + \frac{\delta^2}{4} |\psi_i(x) + \psi_i(y)|^2 \\ &\leqslant |v_i|^2 - 2\delta \langle v_i, \psi_i(x) \rangle + \frac{\delta^2}{2} |v_i|^2 + \frac{1}{2} |\psi_i(x) - \psi_i(y)|^2 \\ &+ \frac{\delta^2}{2} \left[4 |\psi_i(x)|^2 + |\psi_i(x) - \psi_i(y)|^2 \right]. \end{aligned}$$

Using

$$2\left\langle v_i, \frac{1}{d_0} \sum_{j=1}^{d_0} \nabla W^{(q)}(x_i - x_j) \right\rangle \leq |v_i|^2 + \frac{1}{d_0} \sum_{j=1}^{d_0} |x_i - x_j|^2,$$

as well as (59), it follows that

$$\begin{aligned} |p_{i}|^{2} &\leqslant |v_{i}|^{2} - 2\delta \left\langle v_{i}, \nabla U^{(q)}(x_{i}) \right\rangle + \left(\delta\epsilon + \frac{\delta^{2}}{2}\right) |v_{i}|^{2} + \frac{1+\delta^{2}}{2} |\psi_{i}(x) - \psi_{i}(y)|^{2} \\ &+ 4\delta^{2} \left[|x_{i}|^{2} + \frac{\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \right] + \frac{\delta\epsilon}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \\ &\leqslant \left(1 + \delta\epsilon + \frac{\delta^{2}}{2}\right) |v_{i}|^{2} + 4\delta^{2} |x_{i}|^{2} - 2\delta \left\langle v_{i}, \nabla U^{(q)}(x_{i}) \right\rangle + \frac{\delta\epsilon(1 + 4\delta\epsilon)}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \\ &+ |\psi_{i}(x) - \psi_{i}(y)|^{2} \\ &\leqslant \left[1 + \delta\epsilon + \frac{\delta^{2}}{2} + 30\delta^{2}\right] |v_{i}|^{2} - 2\delta \left\langle v_{i}, \nabla U^{(q)}(x_{i}) \right\rangle + 4\delta^{2} \left[1 + 8\delta^{2}\right] |x_{i}|^{2} \\ &+ \frac{\delta\epsilon(1 + 4\delta\epsilon + 30\delta^{3}\epsilon)}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \frac{30\epsilon^{2}\delta^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |v_{i} - v_{j}|^{2} \end{aligned} \tag{62}$$

$$&\leqslant \left(1 + \delta\epsilon + 31\delta^{2}\right) |v_{i}|^{2} - 2\delta \left\langle v_{i}, \nabla U^{(q)}(x_{i}) \right\rangle + 8\delta^{2} |x_{i}|^{2} \\ &+ \frac{2\delta\epsilon}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \frac{30\epsilon^{2}\delta^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |v_{i} - v_{j}|^{2} \end{cases} \tag{63}$$

where we used (60) to obtain the third inequality and that $\delta \leq 1/40$ and $\epsilon \leq 1/4$ to simplify the bounds in the last inequality. Next,

$$\begin{split} \langle y_{i}, p_{i} \rangle &= \left\langle x_{i} + \delta v_{i} - \frac{\delta^{2}}{2} \psi_{i}(x), v_{i} - \frac{\delta}{2} \left(\psi_{i}(x) + \psi_{i}(y) \right) \right\rangle \\ &= \left\langle x_{i}, v_{i} \right\rangle - \frac{\delta}{2} \left\langle x_{i}, \psi_{i}(x) + \psi_{i}(y) \right\rangle + \delta \left| v_{i} \right|^{2} - \frac{\delta^{2}}{2} \left\langle v_{i}, \psi_{i}(x) \right\rangle \\ &- \frac{\delta^{2}}{2} \left\langle v_{i}, \psi_{i}(x) + \psi_{i}(y) \right\rangle + \frac{\delta^{3}}{4} \left\langle \psi_{i}(x), \psi_{i}(x) + \psi_{i}(y) \right\rangle \\ &\leqslant \left\langle x_{i}, v_{i} \right\rangle - \delta \left\langle x_{i}, \psi_{i}(x) \right\rangle + \frac{\delta}{2} \left| x_{i} \right| \left| \psi_{i}(x) - \psi_{i}(y) \right| + \delta \left| v_{i} \right|^{2} - \frac{3\delta^{2}}{2} \left\langle v_{i}, \psi_{i}(x) \right\rangle \\ &+ \frac{\delta^{2}}{2} \left| v_{i} \right| \left| \psi_{i}(x) - \psi_{i}(y) \right| + \frac{\delta^{3}}{2} \left| \psi_{i}(x) \right|^{2} + \frac{\delta^{3}}{4} \left| \psi_{i}(x) \right| \left| \psi_{i}(x) - \psi_{i}(y) \right| \\ &\leqslant \left\langle x_{i}, v_{i} \right\rangle - \delta \left\langle x_{i}, \psi_{i}(x) \right\rangle + \frac{1}{4} \left(\delta^{2} \left| x_{i} \right|^{2} + \left| \psi_{i}(x) - \psi_{i}(y) \right|^{2} \right) - \frac{3\delta^{2}}{2} \left\langle v_{i}, \psi_{i}(x) \right\rangle \\ &+ \delta \left(1 + \frac{\delta}{4} \right) \left| v_{i} \right|^{2} + \frac{5\delta^{3}}{8} \left| \psi_{i}(x) \right|^{2} + \frac{\delta^{2}(2 + \delta)}{8} \left| \psi_{i}(x) - \psi_{i}(y) \right|^{2}. \end{split}$$

Let us emphasize that the term $-\delta \langle x_i, \psi_i(x) \rangle$ is crucial to obtained a dissipation of order δ in the position variables. This motivates the inclusion of a cross term $\langle x_i, v_i \rangle$ in the Lyapunov function. We next use (50), (59) and (60) and the drift condition (16) to write

$$\begin{split} \langle y_{i}, p_{i} \rangle &\leqslant \langle x_{i}, v_{i} \rangle - \delta \left\langle x_{i}, \nabla U^{(q)}(x_{i}) \right\rangle - \frac{3\delta^{2}}{2} \left\langle v_{i}, \nabla U^{(q)}(x_{i}) \right\rangle \\ &+ \delta \left(|x_{i}| + \frac{3\delta}{2} |v_{i}| \right) \frac{\epsilon}{d_{0}} \sum_{j=1}^{d_{0}} \left| \nabla W^{(q)}(x_{i} - x_{j}) \right| + \delta \left(1 + \frac{\delta}{4} \right) |v_{i}|^{2} \\ &+ \frac{\delta^{2}}{4} |x_{i}|^{2} + \frac{7\delta^{3}}{8} |\psi_{i}(x)|^{2} + \frac{2 + \delta^{2}(2 + \delta)}{8} |\psi_{i}(x) - \psi_{i}(y)|^{2} \\ &\leqslant \langle x_{i}, v_{i} \rangle - \delta \mathbf{m} |x_{i}|^{2} + \delta \mathbf{M} + \frac{3\delta^{2}}{4} \left(|v_{i}|^{2} + \left| \nabla U^{(q)}(x_{i}) \right|^{2} \right) + \delta \left(1 + \frac{\delta}{4} \right) |v_{i}|^{2} \\ &+ \delta \epsilon \left(\frac{1}{2} |x_{i}|^{2} + \frac{9\delta^{2}}{8} |v_{i}|^{2} + \frac{1}{d_{0}} \sum_{j=1}^{d_{0}} \left| \nabla W^{(q)}(x_{i} - x_{j}) \right|^{2} \right) + \frac{\delta^{2}}{4} |x_{i}|^{2} \\ &+ \frac{7\delta^{3}}{4} \left(|x_{i}|^{2} + \frac{\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} \right) \\ &+ \frac{30\delta^{2}[2 + \delta^{2}(2 + \delta)]}{8} \left(\delta^{2} |x_{i}|^{2} + \frac{\delta^{2}\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + |v_{i}|^{2} + \frac{\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |v_{i} - v_{j}|^{2} \right) \\ &\leqslant \langle x_{i}, v_{i} \rangle - \frac{\delta \mathbf{m}}{2} |x_{i}|^{2} + \delta \mathbf{M} + \delta(1 + 10\delta) |v_{i}|^{2} \\ &+ \frac{2\delta\epsilon}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} + \frac{8\delta^{2}\epsilon^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |v_{i} - v_{j}|^{2} , \end{split}$$

$$\tag{64}$$

where we used in the last inequality that $\epsilon \leq m/4$ and $\delta \leq m/40 \leq 1/40$.

Using first a Taylor expansion and (recalling (50) and (59))

$$\left\langle \nabla U^{(q)}(x_i), \psi_i(x) \right\rangle \leqslant \frac{1}{2} \left(\left| \nabla U^{(q)}(x_i) \right|^2 + |\psi_i(x)| \right) \leqslant \frac{1}{2} \left(3 |x_i|^2 + \frac{2\epsilon^2}{d_0} \sum_{j=1}^{d_0} |x_i - x_j| \right),$$

then (57) and (48), it follows

$$U^{(q)}(y_i) \leq U^{(q)}(x_i) + \left\langle \nabla U^{(q)}(x_i), y_i - x_i \right\rangle + \frac{1}{2} |y_i - x_i|^2$$

$$\leq U^{(q)}(x_i) + \left\langle \nabla U^{(q)}(x_i), \delta v_i - \frac{\delta^2}{2} \psi_i(x) \right\rangle + \frac{1}{2} |y_i - x_i|^2$$

$$\leq U^{(q)}(x_i) - \frac{\delta^2}{2} \left| \nabla U^{(q)}(x_i) \right|^2 + \delta \left\langle \nabla U^{(q)}(x_i), v_i \right\rangle$$

$$+ \frac{\delta^2 \epsilon}{2} \left(2 |x_i|^2 + \frac{2}{d_0} \sum_{j=1}^{d_0} |x_i - x_j|^2 \right) + \frac{1}{2} |y_i - x_i|^2$$

$$\leq U^{(q)}(x_{i}) - \frac{\delta^{2} \mathbf{m}^{2}}{2} |x_{i}|^{2} + \delta^{2} \mathbf{m} \mathbf{M} + \delta \left\langle \nabla U^{(q)}(x_{i}), v_{i} \right\rangle$$

$$+ \delta^{2} \left(\epsilon + \delta^{2}\right) |x_{i}|^{2} + \delta^{2} |v_{i}|^{2} + \frac{\epsilon \delta^{2} (1 + \delta^{2} \epsilon)}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2}$$

$$\leq U^{(q)}(x_{i}) - \frac{\delta^{2} \mathbf{m}^{2}}{4} |x_{i}|^{2} + \delta^{2} \mathbf{m} \mathbf{M} + \delta \left\langle \nabla U^{(q)}(x_{i}), v_{i} \right\rangle + \delta^{2} |v_{i}|^{2}$$

$$+ \frac{2\epsilon \delta^{2}}{d_{0}} \sum_{j=1}^{d_{0}} |x_{i} - x_{j}|^{2} .$$

$$(65)$$

, Using first $|\mathbf{a} + \mathbf{b}|^2 \leq (1 + \delta) |\mathbf{a}|^2 + (1 + \delta^{-1}) |\mathbf{b}|^2$, then (56) as well as $\delta \leq m/40 \leq 1/40$ and $\epsilon \leq m/4 \leq 1/4$, we obtain

$$|y_{i} - y_{j}|^{2} = \left| x_{i} - x_{j} - \frac{\delta^{2}}{2} \left(\psi_{i}(x) - \psi_{j}(x) \right) + \delta(v_{i} - v_{j}) \right|^{2}$$

$$\leq (1 + \delta) \left| x_{j} - x_{j} \right|^{2} + \delta^{2} \left(1 + \delta^{-1} \right) \left| v_{i} - v_{j} - \frac{\delta}{2} \left(\psi_{i}(x) - \psi_{j}(x) \right) \right|^{2}$$

$$\leq (1 + \delta) \left| x_{i} - x_{j} \right|^{2} + 2\delta(1 + \delta) \left| v_{i} - v_{j} \right|^{2} + \frac{\delta^{3}(1 + \delta)}{2} \left| \psi_{i}(x) - \psi_{j}(x) \right|^{2}$$

$$\leq \left[1 + \delta + \delta^{3}(1 + \delta) \left(1 + \epsilon^{2} \right) \right] \left| x_{i} - x_{j} \right|^{2} + 2\delta(1 + \delta) \left| v_{i} - v_{j} \right|^{2}$$

$$\leq (1 + 2\delta) \left| x_{i} - x_{j} \right|^{2} + 4\delta \left| v_{i} - v_{j} \right|^{2} . \tag{66}$$

Finally, using twice (56),

$$\begin{split} |p_{i} - p_{j}|^{2} &\leq (1+\delta) |v_{i} - v_{j}|^{2} + \frac{\delta^{2} (1+\delta^{-1})}{4} |\psi_{i}(y) + \psi_{i}(x) - \psi_{j}(y) - \psi_{j}(x)|^{2} \\ &\leq (1+\delta) |v_{i} - v_{j}|^{2} + \frac{\delta(1+\delta)}{2} \left(|\psi_{i}(x) - \psi_{j}(x)|^{2} + |\psi_{i}(y) - \psi_{j}(y)|^{2} \right) \\ &\leq (1+\delta) |v_{i} - v_{j}|^{2} + \delta(1+\delta) \left(1+\epsilon^{2} \right) \left(|x_{i} - x_{j}|^{2} + |y_{i} - y_{j}|^{2} \right) \\ &\leq \delta(1+\delta) \left(1+\epsilon^{2} \right) \left(4 |x_{i} - x_{j}|^{2} + 3\delta^{2} |v_{i} - v_{j}|^{2} + \frac{3\delta^{4}}{4} |\psi_{i}(x) - \psi_{j}(x)|^{2} \right) \\ &+ (1+\delta) |v_{i} - v_{j}|^{2} \\ &\leq \delta(1+\delta) \left(1+\epsilon^{2} \right) \left(4 + \frac{3\delta^{2}}{2} \left(1+\epsilon^{2} \right) \right) |x_{i} - x_{j}|^{2} + (1+2\delta) |v_{i} - v_{j}|^{2} \\ &\leq 5\delta |x_{i} - x_{j}|^{2} + (1+2\delta) |v_{i} - v_{j}|^{2} \ . \end{split}$$

Combining this last inequality with (61)-(63)-(64)-(65)-(66) in (55), and (recalling $b \ge b_0$) $-2\delta(b-b_0)\langle v_i, \nabla U^{(q)}(x_i)\rangle \leqslant \delta(b-b_0)\left(|v_i|^2+|x_i|^2\right),$

completes the proof.

Quantifying how prefactors change over several steps. In view of Lemma 19, controlling the evolution of the Lyapunov function along several Verlet steps amounts to a discrete Grönwall-type result for the sequence of parameters defined recursively by the relation $\mathbf{o} \mapsto \tilde{\mathbf{o}}$ of Lemma 19. Notice that, among these parameters, b_0 is fixed and f does not influence the other parameters, so that we focus first on the relations intertwining the parameters a, b, c, e. We state an abstract result, independent from the specific constants appearing in Lemma 19, except that it is crucial to keep track of the order of the various parts in terms of the step size δ . More specifically, Lemma 19 reads $\tilde{\mathbf{o}} = (I + \delta A + \delta^2 B)\mathbf{o}$ for some matrices A, B and thus, iterating this k times, as long as $k\delta$ is small enough, it is sufficient to keep the lowest order terms in $k\delta$ and δ in $(I + \delta A + \delta^2 B)^k$. This is essentially the content of the next lemma.

Lemma 20. Consider the four real sequences $(a_k)_{k\in\mathbb{N}}, (b_k)_{k\in\mathbb{N}}, (c_k)_{k\in\mathbb{N}}, (e_k)_{k\in\mathbb{N}}$ such as $a_0, b_0, c_0, e_0 > 0$ and defined by

$$\begin{aligned} a_{k+1} &= (1 - \delta^2 C_{a,1})a_k - \delta c_k C_{a,2} + \delta^2 b_k C_{a,3} + \delta e_k C_{a,4} + C_{a,5} \delta(b_k - b_0) , \\ b_{k+1} &= (1 + C_{b,1} \delta + C_{b,2} \delta^2)b_k + \delta^2 C_{b,3} a_k + \delta c_k C_{b,4} + \delta e_k C_{b,5} + \delta(b_k - b_0) , \\ c_{k+1} &= c_k + C_{c,1} \delta a_k , \quad e_{k+1} &= e_k + C_{e,1} \delta(a_k \delta + b_k + c_k + e_k) , \end{aligned}$$

for some non-negative real constants $\{C_{\Box, \bigtriangleup} : \Box \in \{a, b, c, e\}, \bigtriangleup \in \{1, 2, 3, 4, 5\}\}$ and $\delta > 0$. Then, for any $\delta > 0$ and $k \in \mathbb{N}$ satisfying

$$\delta \leqslant \tilde{\delta}_1 \wedge \tilde{\delta} , \quad k\delta \leqslant \tilde{T}_1 \wedge \tilde{T}_2 \wedge \tilde{T} , \qquad (67)$$

for $\tilde{T}, \tilde{\delta} > 0$, with $\tilde{\delta}_1 = [C_{b,2}]^{-1} \wedge [a_0 C_{a,2} C_{c,1} / \{8(\tilde{C}_{e,1} C_{a,4} + \tilde{C}_{b,3}(\tilde{\delta} C_{a,3} + C_{a,5}))\}] \wedge [4C_{a,1}]^{-1} \wedge [4C_{c,1} C_{a,2} \tilde{C}_{a,3}]^{-1}$ $\tilde{T}_1 = 4^{-1} \{\tilde{C}_{b,1}^{-1} \wedge \tilde{C}_{b,2}^{-1/2} \wedge C_{e,1}^{-1} \wedge 1\},$ $\tilde{T}_2 = 4^{-1} \{[C_{a,2} / (4C_{b,4} C_{a,5})] \wedge [a_0 C_{a,2} C_{c,1} (2C_{a,5} \tilde{C}_{b,2} b_0)^{-1}] \wedge [a_0 (2C_{a,2} c_0)^{-1}] \wedge [2C_{c,1} C_{a,2}]^{-1/2}\},$ assuming moreover that

$$2e_0C_{a,4} \leqslant c_0C_{a,2}/4 , \qquad C_{a,5}(C_{b,1}b_0 + C_{b,5}e_0) + \tilde{C}_{e,1}C_{a,4} \leqslant C_{a,2}C_{c,1}a_0/8 , \qquad (68)$$

it holds

$$\begin{aligned} a_0/2 &\leqslant a_k \leqslant a_0 + \tilde{C}_{a,3}k\delta^2 , \quad b_0 \leqslant b_k \leqslant 2b_0 + \tilde{C}_{b,3}k\delta^2 , \quad e_k \leqslant 2e_0 + \tilde{C}_{e,1}k\delta + \tilde{C}_{e,2}k\delta^2 , \\ c_0 + k\delta a_0 C_{c,1}/2 &\leqslant c_0 + k\delta a_0 C_{c,1} + (\delta k)^2 \delta C_{c,1} \tilde{C}_{a,3} \\ b_k &\leqslant b_0 (1 + k\delta \tilde{C}_{b,1} + (k\delta)^2 \tilde{C}_{b,2}) + \delta^2 k \tilde{C}_{b,3} \\ e_k &\leqslant e_0 (1 + k\delta C_{e,1}) + \tilde{C}_{e,1}k\delta + \tilde{C}_{e,2}k\delta^2 \\ a_k &\leqslant a_0 (1 - k\delta \tilde{C}_{a,1} - (k\delta)^2 \tilde{C}_{a,2}) + \delta^2 k \tilde{C}_{a,3} \\ a_k &\geqslant a_0 (1 - k\delta (C_{a,2}c_0/a_0 + \delta C_{a,1}) - (k\delta)^2 C_{a,2} C_{c,1} (1 + \delta \tilde{T} \tilde{C}_{a,3})) \\ c_k &\geqslant c_0 + C_{c,1}k\delta a_0 - (k\delta)^2 C_{c,1} (C_{a,2}c_0 + \delta C_{a,1}a_0) - (k\delta)^3 C_{c,1}^2 C_{a,2}a_0 (1 + \delta \tilde{T} \tilde{C}_{a,3}) , \end{aligned}$$
(69)

where

$$\begin{split} \tilde{C}_{a,1} &= c_0 C_{a,2} / (2a_0) \;, \quad \tilde{C}_{a,2} = a_0 [C_{a,1} \wedge C_{a,2} C_{c,1} / 16] \;, \\ \tilde{C}_{a,3} &= 2b_0 C_{a,3} + a_0 C_{a,1} \tilde{T} (\tilde{C}_{a,1} + \tilde{T} \tilde{C}_{a,2}) \;, \\ \tilde{C}_{b,1} &= C_{b,1} + c_0 C_{b,4} / b_0 + e_0 C_{b,5} / b_0 \;, \\ \tilde{C}_{b,2} &= 4 [C_{b,1} \tilde{C}_{b,1} + \tilde{C}_{b,1} + C_{b,4} (C_{c,1} a_0 + \tilde{C}_{e,1} C_{b,5})) / b_0] \;, \\ \tilde{C}_{b,3} &= 3 \{ b_0 C_{b,2} + a_0 C_{b,3} + \tilde{T} (C_{b,3} \tilde{C}_{a,3} + b_0 C_{b,2} \tilde{C}_{b,1} + C_{b,5} \tilde{C}_{e,2} + \tilde{T} \tilde{C}_{a,3} C_{c,1} C_{b,4}) \} \\ \tilde{C}_{e,1} &= 2 C_{e,1} (2b_0 + c_0 + \tilde{T} (C_{e,1} e_0 + C_{c,1} a_0 + \delta \tilde{C}_{b,3})) \\ \tilde{C}_{e,2} &= 2 C_{e,1} (a_0 + \tilde{T} \delta \tilde{C}_{a,3} + \tilde{T}^2 C_{c,1} \tilde{C}_{a,3})) \;. \end{split}$$

Proof. The proof is by induction. Equation (69) is trivially true for k = 0. Suppose that it is true for $k \in \mathbb{N}$ and suppose $\delta, k+1$ satisfy (67). Then by definition, the induction hypothesis implies that $c_0 + (k+1)\delta a_0 C_{c,1}/2 \leq c_{k+1} \leq c_0 + (k+1)\delta a_0 C_{c,1} + (k+1)^2 \delta^3 C_{c,1} \tilde{C}_{a,3}$.

Similarly, it follows that

$$\begin{split} b_{k+1} &\leqslant [(1+k\delta\tilde{C}_{b,1}+(k\delta)^2\tilde{C}_{b,2})b_0 + \tilde{C}_{b,3}\delta^2k](1+C_{b,1}\delta + C_{b,2}\delta^2) \\ &+ \delta^2 C_{b,3}(a_0 + \tilde{C}_{a,3}\delta^2k) + \delta C_{b,4}(c_0 + \delta k C_{c,1}a_0 + (\delta k)^2\delta C_{c,1}\tilde{C}_{a,3}) \\ &+ \delta C_{b,5}(2e_0 + \delta k\tilde{C}_{e,1} + \delta^2k\tilde{C}_{e,2}) \\ &\leqslant b_0 + \delta b_0(k\tilde{C}_{b,1} + C_{b,1} + c_0C_{b,4}/b_0 + 2C_{b,5}e_0/b_0) \\ &+ \delta^2kb_0\{\tilde{C}_{b,2}k + k\delta\tilde{C}_{b,2}C_{b,1} + C_{b,1}\tilde{C}_{b,1} + \tilde{C}_{b,1} \\ &+ (C_{b,4}/b_0)(C_{c,1}a_0 + \tilde{C}_{e,1}C_{b,5}) + k\delta(\delta\tilde{C}_{b,2}C_{b,2} + \tilde{C}_{b,2})\} \\ &+ \delta^2(k\tilde{C}_{b,3} + \tilde{C}_{b,3}(k\delta C_{b,1} + k\delta^2C_{b,2}) + b_0C_{b,2} + a_0C_{b,3} \\ &+ \tilde{T}^2\tilde{C}_{a,3}C_{c,1}C_{b,4} + \tilde{T}C_{b,3}\tilde{C}_{a,3} + \tilde{T}b_0C_{b,2}\tilde{C}_{b,1} + \tilde{T}C_{b,5}\tilde{C}_{e,2}) \\ &\leqslant b_0(1 + (k+1)\delta\tilde{C}_{b,1} + k(k+2)\delta^2\tilde{C}_{b,2}) + \tilde{C}_{b,3}(k+1)\delta^2 \;, \end{split}$$

where we have used that $k\delta \leq 4^{-1}[C_{b,1}^{-1} \wedge 1], \ \delta \leq C_{b,2}^{-1}, \ \tilde{C}_{b,1} \geq C_{b,1} \ \text{and} \ \tilde{C}_{b,2} \geq C_{b,2}$ by definition. In addition, $b_0 \leq b_{k+1} \leq 2b_0 + \delta^2 k \tilde{C}_{b,3}$ since $\delta, k+1$ satisfy (67). Similarly, using that $k\delta \leq (4C_{e,1})^{-1}$, we have

$$\begin{split} e_{k+1} &\leqslant \left[(1+k\delta C_{e,1})e_0 + \delta k\tilde{C}_{e,1} + \delta^2 k\tilde{C}_{e,2} \right] (1+C_{e,1}\delta) \\ &+ C_{e,1}\delta^2(a_0 + \tilde{C}_{a,3}k\delta^2) + C_{e,1}\delta(2b_0 + \delta^2 k\tilde{C}_{b,3}) + C_{e,1}\delta(c_0 + k\delta C_{c,1}a_0 + (k\delta)^2 \delta C_{c,1}\tilde{C}_{a,3}) \\ &\leqslant (1+(k+1)\delta C_{e,1})e_0 + \delta(k\tilde{C}_{e,1} + C_{e,1}(2b_0 + c_0 + \tilde{T}C_{e,1}e_0 + \tilde{T}\tilde{C}_{e,1} + \tilde{T}C_{c,1}a_0 + \delta\tilde{T}\tilde{C}_{b,3})) \\ &+ \delta^2(k\tilde{C}_{e,2} + C_{e,1}(k\delta\tilde{C}_{e,2} + a_0 + k\delta^2\tilde{C}_{a,3} + (k\delta)^2 C_{c,1}\tilde{C}_{a,3})) \;, \end{split}$$

which concludes the proof of the upper bound for e_{k+1} using $k\delta \leq 1/(4C_{e,1})$.

Using the induction hypothesis, in particular the lower bounds on c_k and the definition of $\tilde{C}_{b,1}$, we get

$$a_{k+1} \leq a_0 - \delta(a_0 k C_{a,1} + C_{a,2} c_0 - 2C_{a,4} e_0) - \delta^2 a_0 C_{a,1}$$

$$\begin{split} &+ \delta^2 (k\tilde{C}_{a,3} + 2b_0C_{a,3} + a_0C_{a,1}\tilde{T}(\tilde{C}_{a,1} + \tilde{T}\tilde{C}_{a,2})) \\ &- \delta^2 k [k\tilde{C}_{a,2}a_0 + a_0C_{c,1}C_{a,2}/2 - C_{a,5}\tilde{C}_{b,1}b_0 - \tilde{C}_{e,1}C_{a,4} \\ &- \delta kC_{a,5}\tilde{C}_{b,2}b_0 - \delta^2 C_{a,3}\tilde{C}_{b,3} - \delta C_{a,4}\tilde{C}_{e,1} - \delta C_{a,5}\tilde{C}_{b,3}] \\ \leqslant a_0 - \delta (a_0k\tilde{C}_{a,1} + C_{a,2}c_0 - 2C_{a,4}e_0 - c_0C_{b,4}C_{a,5}\delta k) - \delta^2 a_0C_{a,1} \\ &+ \delta^2 (k\tilde{C}_{a,3} + 2b_0C_{a,3} + a_0C_{a,1}\tilde{T}(\tilde{C}_{a,1} + \tilde{T}\tilde{C}_{a,2})) \\ &- \delta^2 k [k\tilde{C}_{a,2}a_0 + a_0C_{c,1}C_{a,2}/2 - C_{a,5}(C_{b,1}b_0 + C_{b,5}e_0) - \tilde{C}_{e,1}C_{a,4} \\ &- \delta kC_{a,5}\tilde{C}_{b,2}b_0 - \delta^2 C_{a,3}\tilde{C}_{b,3} - \delta C_{a,4}\tilde{C}_{e,1} - \delta C_{a,5}\tilde{C}_{b,3}] \,. \end{split}$$

Using the condition (68), $\delta, k+1$ satisfy (67), we get that $a_{k+1} \leq a_0 - a_0(k+1)\delta \tilde{C}_{a,1} - a_0(k+2)k\delta^2 \tilde{C}_{a,2} - a_0 C_{a,1}\delta^2 + \delta^2 k \tilde{C}_{a,3} \leq a_0 - a_0(k+1)\delta \tilde{C}_{a,1} - a_0(k+1)^2 \delta^2 \tilde{C}_{a,2} + \tilde{C}_{a,3}\delta^2(k+1)$. The lower bound for a_{k+1} proceeds similarly using the upper bound on c_k given by the induction hypothesis. Finally, the conditions $(k+1)\delta \leq 8^{-1}\{[a_0(C_{a,2}c_0)^{-1}] \wedge [C_{c,1}C_{a,2}]^{-1/2}\}, \delta \leq [4C_{a,1}]^{-1} \wedge [4C_{c,1}C_{a,2}\tilde{C}_{a,3}]^{-1}$ and $(k+1)\delta \leq 1/2$ imply that $a_0/2 \leq a_{k+1} \leq a_0 + \delta^2 k \tilde{C}_{a,3}$. The last inequality for c_{k+1} in (69) uses the induction hypothesis $a_k \geq a_0(1-k\delta(C_{a,2}c_0/a_0+\delta C_{a,1}) - (k\delta)^2 C_{a,2}C_{c,1}(1+\delta \tilde{T}\tilde{C}_{a,3}))$ and $c_{k+1} = c_k + C_{c,1}\delta a_k$.

Evolution over several Verlet steps. Combining Lemmas 19 and 20 leads to the following, which gives a control of the expected value of the Lyapunov function after a full trajectory of Verlet integrator.

Lemma 21. Assume that H_4 holds. Let $\mathbf{o} = (a_0, b_0, c_0, 2b_0, e_0, f_0) \in \mathbb{R}^6_+, \ \overline{T}, \overline{\delta} > 0$ and denote

$$\begin{split} \bar{C}_{a,1} &= \mathsf{m}/(3\times 4) \;, \quad \bar{C}_{a,2} = \mathsf{m}/(3\times 2^3) \;, \\ \bar{C}_{a,3} &= 16b_0 + a_0\bar{T}(\mathsf{m}/2)^2(1+\bar{T}) \;, \\ \bar{C}_{b,1} &= \epsilon + (c_0/b_0)(1+10\bar{\delta}) \;, \quad \bar{C}_{b,2,0} = 4[(1+\epsilon)\bar{C}_{b,1} + (1+10\bar{\delta})(2a_0/b_0)] \;, \\ \bar{C}_{b,3,0} &= 3\{31b_0 + 2a_0 + \bar{T}(2\bar{C}_{a,3} + 31b_0\bar{C}_{b,1} + 2\bar{C}_{a,3}\bar{T}(1+10\bar{\delta}))\} \;, \\ \bar{C}_{b,2} &= \bar{C}_{b,2,0} + c_0\mathsf{m} + a_0\mathsf{m}\bar{T} \;, \quad \bar{C}_{b,3} = \bar{C}_{b,3,0} + \bar{C}_{a,3} \;, \\ \bar{C}_{e,1} &= 14[2b_0 + c_0 + \bar{T}(14e_0 + 2a_0 + \bar{\delta}\bar{C}_{b,3})] \;, \quad \bar{C}_{e,2} = 14(a_0 + \bar{T}\bar{C}_{a,3}(\bar{\delta} + 2\bar{T})) \;, \\ \bar{C}_{f,1} &= a_0\mathsf{M}\bar{\delta} + \bar{C}_{a,3}\mathsf{M}\bar{\delta}\bar{T} + 4b_0\mathsf{m}\mathsf{M}\bar{\delta} + \mathsf{M}(c_0 + \bar{C}_{b,3,0}\bar{T}^2\bar{\delta} + 2\bar{T}a_0 + 2\bar{C}_{a,3}\bar{T}^2\bar{\delta}) \;. \end{split}$$

Let $\epsilon \leq m/4$ and suppose that

$$\epsilon b_0 \leqslant a_0 \mathbf{m}/2^3 . \tag{70}$$

Then, for any $\delta > 0$, $\eta \in [0,1)$, and $K \in \mathbb{N}$ satisfying $\delta \leq \overline{\delta}_1 \wedge \overline{\delta}$ and $K\delta \leq \overline{T}_1 \wedge \overline{T}_2 \wedge \overline{T}$, and

$$\delta\eta\bar{C}_{a,3} \leqslant \frac{c_0\mathfrak{m}}{3\times 2^2} , \quad 2\delta\frac{(1-\eta)}{\delta K}\bar{C}_{a,3} \leqslant \frac{a_0\mathfrak{m}}{3\times 2^3} , \tag{71}$$

where

$$\bar{\delta}_{1} = 31^{-1} \wedge [\mathfrak{m}/40] \wedge [a_{0}2^{-3}/(\bar{C}_{b,3}(1+8\bar{\delta}))] \wedge [1/(2\mathfrak{m})] \wedge [1/(8\mathfrak{m}\bar{C}_{a,3})],$$

$$\bar{T}_{1} = 4^{-1} \left[31^{-1} \wedge \bar{C}_{b,1}^{-1} \wedge \bar{C}_{b,2}^{-1/2} \right],$$

$$\bar{T}_{2} = 4^{-1} \{ [2^{-3}\mathfrak{m}/(1+10\bar{\delta})] \wedge [8^{-1}a_{0}/(\bar{C}_{b,2}b_{0})] \wedge a_{0}\mathfrak{m}(2c_{0})^{-1} \wedge 1/(7\mathfrak{m})^{1/2} \}.$$
(72)

 $it\ holds$

(a) For any $i \in \{1, \ldots, d_0\}$, $x, v \in \mathbb{R}^d$, $\mathcal{V}_{\delta}^K \mathfrak{V}_{\mathbf{o},i}(x, v) \leq \mathfrak{V}_{\mathbf{o}_K,i}(x, v)$, where $\mathbf{o}_K = (a_K, b_K, c_K, 2b_0, e_K, f_K)$

$$\begin{split} a_{K} &= a_{0} - K\delta\bar{C}_{a,1}c_{0} - (K\delta)^{2}\bar{C}_{a,2}a_{0} , \qquad b_{K} = b_{0}(1 + K\delta\bar{C}_{b,1} + (K\delta)^{2}\bar{C}_{b,2}) + \delta^{2}K\bar{C}_{b,3} , \\ c_{K} &= c_{0} + 2K\delta a_{0} , \quad e_{K} = e_{0}(1 + 17K\delta) + \bar{C}_{e,1}K\delta + \bar{C}_{e,2}K\delta^{2} , \\ f_{K} &= f_{0} + K\delta\bar{C}_{f,1} . \end{split}$$

(b) In addition for any $k \in \{0, \ldots, K\}$, $i \in \{1, \ldots, d_0\}$, $x, v \in \mathbb{R}^d$, $V^k_{\delta} \mathfrak{V}_{\mathbf{o},i}(x, v) \leq \mathfrak{V}_{\mathbf{o}_k,i}(x, v)$, where $\mathbf{o}_k = (a_k, b_k, c_k, 2b_0, e_k, f_k)$

$$\begin{aligned} a_k &= a_0 + 2\bar{C}_{a,3}\delta^2 k , \quad b_k &= 2b_0 + \delta^2 k\bar{C}_{b,3} , \\ c_k &= c_0 + 2k\delta a_0 , \quad e_k &= 2e_0 + \bar{C}_{e,1}k\delta + \bar{C}_{e,2}k\delta^2 , \quad f_k &= f_0 + k\delta\bar{C}_{f,1} \end{aligned}$$

Proof. (a) Let $i \in \{1, \ldots, d_0\}$, $\delta \leq \overline{\delta}_1$, $K \in \mathbb{N}$, $K\delta \leq \overline{T} \wedge \overline{T}_1 \wedge \overline{T}_2$ and $x, v \in \mathbb{R}^d$. Let $(a_0, b_0, c_0, e_0, f_0) \in \mathbb{R}^6_+$. By Lemma 19, we have starting from $\tilde{\mathbf{o}}_0 = \mathbf{o}_0 = (a_0, b_0, c_0, 2b_0, e_0, f_0)$, for any $k \in \mathbb{N}$,

$$\mathbf{V}_{\delta}^{k+1}\mathfrak{V}_{\tilde{\mathbf{o}}_{0},i}(x,v) \leqslant \mathbf{V}_{\delta}\mathfrak{V}_{\tilde{\mathbf{o}}_{k},i}(x,v) \leqslant \mathfrak{V}_{\tilde{\mathbf{o}}_{k+1},i}(x,v)$$

where $\tilde{\mathbf{o}}_{k+1} = (\tilde{a}_{k+1}, \tilde{b}_{k+1}, \tilde{c}_{k+1}, b_0, \tilde{e}_{k+1}, \tilde{f}_{k+1})$ are defined by the recursion

$$\begin{split} \tilde{a}_{k+1} &= \tilde{a}_k (1 - \mathbf{m} \delta^2 / 2) + 8 \tilde{b}_k \delta^2 - \delta \mathbf{m} \tilde{c}_k / 2 + \delta |\tilde{b}_k - b_0| \\ \tilde{b}_{k+1} &= 2 \delta^2 \tilde{a}_k + (1 + \epsilon \delta + 31 \delta^2) \tilde{b}_k + \delta (1 + 10 \delta) \tilde{c}_k + \delta |\tilde{b}_k - b_0| \\ \tilde{c}_{k+1} &= 2 \delta \tilde{a}_k + \tilde{c}_k , \quad \tilde{e}_{k+1} &= \tilde{e}_k + 7 \delta (\tilde{a}_k \delta + \delta b_0 + \tilde{b}_k + \tilde{c}_k + \tilde{e}_k) \\ \tilde{f}_{k+1} &= \tilde{f}_k + \tilde{a}_k \delta^2 \mathbf{M} + 2 \tilde{b}_k \mathbf{m} \mathbf{M} \delta^2 + \delta \mathbf{M} \tilde{c}_k . \end{split}$$

Applying Lemma 20, we get that for any $k \in \{0, \ldots, K\}$,

$$\begin{split} \tilde{a}_k &\leqslant a_0 (1 - \mathfrak{m} k \delta c_0 / (4a_0) - \mathfrak{m} (k\delta)^2 / 2^3) + (16b_0 + a_0 (\mathfrak{m} / 2)^2 \bar{T} (1 + \bar{T})) \delta^2 k \\ \tilde{b}_k &\leqslant b_0 (1 + k \delta \bar{C}_{b,1} + (k\delta)^2 \bar{C}_{b,2,0}) + \bar{C}_{b,3,0} k \delta^2 \\ |\tilde{c}_k - c_0 - 2k\delta a_0| &\leqslant (k\delta)^2 c_0 \mathfrak{m} + a_0 \mathfrak{m} (k\delta)^3 + 2\bar{C}_{a,3} \delta^2 k \\ \tilde{e}_k &\leqslant e_0 (1 + 7k\delta) + k \delta \bar{C}_{e,1} + k \delta^2 \bar{C}_{e,2} , \quad \tilde{f}_k \leqslant f_0 + \delta k \bar{C}_{f,1} . \end{split}$$

Using that $|[c_k - c_0 - 2\delta ka_0] \langle v, x \rangle| \leq 2^{-1} |c_k - c_0 - 2\delta ka_0| [|x|^2 + |v|^2]$, (71) and $K\delta \leq 3 \times 2^{-3}$ completes the proof.

(b) The proof of the second statement follows the same lines and is omitted.

Introducing momentum updates. In the next lemma, we state pointwise estimates that control the effect on the Lyapunov function of a momentum update for a given realization g of a Gaussian variable. After raising these estimates to some powers, the expectation with respect to the Gaussian distribution will be taken afterwards (Lemma 25 below).

Lemma 22. Assume that H_4 holds and that $\epsilon \leq m/4$ and consider the notations introduced in Lemma 21. Then, for any $i \in \{1, \ldots, d_0\}$, $\boldsymbol{\omega} = (\delta, K, \eta)$, $\delta > 0$, $K \in \mathbb{N}$, $\eta \in (0, 1)$, $\mathbf{o} = (a_0, b_0, c_0, 2b_0, e_0, f_0) \in \mathbb{R}^6_+$, satisfying $\delta \leq \overline{\delta}_1$, $K\delta \leq \overline{T} \wedge \overline{T}_1 \wedge \overline{T}_2$ with $\overline{T} > 0$, (70)-(71),

$$c_{0}(\eta - 1) + 2\eta a_{0}\delta K = 0 , \quad c_{0} \leqslant \sqrt{2a_{0}b_{0}} , \quad c_{0}/b_{0} \leqslant \eta(1 - \eta)/(\delta K) , \bar{C}_{b,1} - c_{0}/b_{0} \leqslant (1 - \eta)/[4\delta K] , \quad \delta K\bar{C}_{b,2} + \delta\bar{C}_{b,3}/b_{0} \leqslant (1 - \eta)/[4\delta K] ,$$

$$(73)$$

 $it\ holds$

(a) For any
$$x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}), g = (g_1, \ldots, g_{d_0}) \in \mathbb{R}^d$$
,

$$\begin{aligned} \mathbf{V}_{\delta}^{K} \mathfrak{V}_{\mathbf{o},i}(x,\eta v + \tilde{\eta}g) &\leq (1 - \bar{\rho}_{\omega} K \delta) [a_{0} |x_{i}|^{2} + b_{0} |v_{i}|^{2} + c_{0} \langle x_{i}, v_{i} \rangle + 2b_{0} U^{(q)}(x_{i})] + f_{K} \\ &+ 2\eta \tilde{\eta} b_{K} \langle v_{i}, g_{i} \rangle + c_{K} \tilde{\eta} \langle x, g \rangle + \tilde{\eta}^{2} b_{K} |g_{i}|^{2} + e_{K} \frac{\epsilon}{d_{0}} \sum_{j=1}^{d_{0}} [|x_{i} - x_{j}|^{2} + |v_{i} - v_{j} + g_{i} - g_{j}|^{2}] \end{aligned}$$

where $b_K \leq 2b_0 + \delta^2 K \bar{C}_{b,3}$, $e_K \leq 2e_0 + (K\delta) \bar{C}_{e,1} + K \delta^2 \bar{C}_{e,2}$,

$$\tilde{\eta}^2 = 1 - \eta^2$$

$$\bar{\rho}_{\omega} = \left[(c_0 + K\delta a_0) \bar{C}_{a,2} / (4a_0) \right] \wedge \left[(1 - \eta) / (4K\delta) \right] \wedge \left[(c_0 + K\delta a_0) \bar{C}_{a,2} / (4b_0 a_0) \right].$$

(b) In addition for any $k \in \{1, \ldots, K\}$, $x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}), g = (g_1, \ldots, g_{d_0}) \in \mathbb{R}^d$,

$$\begin{aligned} \mathbf{V}_{\delta}^{k}\mathfrak{V}_{\mathbf{o},i}(x,\eta v+\tilde{\eta}g) &\leqslant \bar{C}_{1}[a_{0}|x_{i}|^{2}+b_{0}|v_{i}|^{2}+c_{0}\langle x_{i},v_{i}\rangle+2b_{0}U^{(q)}(x_{i})]+f_{K} \\ &+2\eta\tilde{\eta}b_{K}\langle v_{i},g_{i}\rangle+c_{K}\tilde{\eta}\langle x,g\rangle+\tilde{\eta}^{2}b_{K}|g_{i}|^{2}+e_{K}\frac{\epsilon}{d_{0}}\sum_{j=1}^{d_{0}}[|x_{i}-x_{j}|^{2}+|v_{i}-v_{j}+g_{i}-g_{j}|^{2}],\end{aligned}$$

where

$$\bar{C}_{1} = [\{a_{0}+2c_{0}+\bar{T}a_{0}+\bar{T}^{2}c_{0}m/8+a_{0}m\bar{T}^{3}/4+2\bar{C}_{a,3}\bar{\delta}\bar{T}\}/a_{0}] \vee [\{2b_{0}+2c_{0}+\bar{T}a_{0}+\bar{\delta}\bar{T}\bar{C}_{b,3}\}/b_{0}] .$$
Remark 23. Note that the first line in (73) is in particular satisfied choosing $b_{0} = 1$, $a_{0} = (1-\eta)^{2}/[2(\delta K)^{2}]$ and $c_{0} = \eta(1-\eta)/(\delta K)$.

Proof. (a) Let $\delta > 0, K \in \mathbb{N}, \eta \in (0,1), \mathbf{o} = (a_0, b_0, c_0, 2b_0, e_0, f_0) \in \mathbb{R}^6_+$, satisfying $\delta \leq \bar{\delta}_1 \wedge \bar{\delta}, K\delta \leq \bar{T} \wedge \bar{T}_1 \wedge \bar{T}_2, (70)$ -(71)-(73). For any $x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}), g = (g_1, \ldots, g_{d_0}) \in \mathbb{R}^d$, using Lemma 21-(a), we get

$$\begin{aligned} \mathbf{V}_{\delta}^{K} \mathfrak{V}_{\mathbf{o},i}(x,\eta v + \tilde{\eta}g) &\leq a_{K} |x_{i}|^{2} + b_{K} \eta^{2} |v_{i}|^{2} + c_{K} \eta \langle x_{i}, v_{i} \rangle + 2b_{0} U^{(q)}(x_{i}) + f_{K} \\ &+ 2\eta \tilde{\eta} b_{K} \langle v_{i}, g_{i} \rangle + \tilde{\eta}^{2} b_{K} |g_{i}|^{2} + c_{K} \tilde{\eta} \langle x_{i}, g_{i} \rangle + \frac{\epsilon e_{K}}{d_{0}} \sum_{j=1}^{d_{0}} [|x_{i} - x_{j}|^{2} + |v_{i} - v_{j} + g_{i} - g_{j}|^{2}] . \end{aligned}$$

Note that by the first line in (73) and using $c_K = c_0 + 2K\delta a_0$,

$$\eta c_K \langle x, v \rangle = c_0 \langle x, v \rangle . \tag{74}$$

Second using the second line in (73), we have

$$b_{K}\eta^{2} = (1 + \bar{C}_{b,1}K\delta + \bar{C}_{b,2}(K\delta)^{2} + \bar{C}_{b,3}K\delta^{2}/b_{0})b_{0}(1 - (1 + \eta)(1 - \eta))$$

$$\leq b_{0}[1 - (1 - \eta) - \eta(1 - \eta) + C_{1}\delta K + \delta K(\delta K\bar{C}_{2,b} + \delta\bar{C}_{3,b}/b_{0})]$$

$$\leq b_{0}[1 - (1 - \eta) - \eta(1 - \eta) + c_{0}\delta K/b_{0} + (1 - \eta)/4 + \delta K(\bar{C}_{2,b} + \delta\bar{C}_{3,b}/b_{0})]$$

$$\leq b_{0}(1 - \delta K(1 - \eta)/[2\delta K]).$$
(75)

Then using (74)-(75), $U(x) \leq |x|^2$ under **H** 4 and since U(0) = 0, we get setting $\bar{C}_{a,4} = K\delta\bar{C}_{a,2}(c_0 + \delta K a_0)/a_0$,

$$\begin{split} \mathbf{V}_{\delta}^{K} \mathfrak{V}_{\mathbf{o},i}(x,\eta v+\tilde{\eta}g) &\leqslant (1-\bar{C}_{a,4}/2)a_{0} \left|x_{i}\right|^{2} + \delta K(1-\eta) \left|v_{i}\right|^{2}/[2K\delta] + c_{0} \left\langle x_{i},v_{i}\right\rangle \\ &+ (1-\bar{C}_{a,4}/(4b_{0}))2b_{0}U^{(q)}(x_{i}) + f_{K} + 2\eta\tilde{\eta}b_{K} \left\langle v_{i},g_{i}\right\rangle + \tilde{\eta}^{2}b_{K} \left|g_{i}\right|^{2} + c_{K}\tilde{\eta} \left\langle x_{i},g_{i}\right\rangle \\ &+ \frac{\epsilon e_{K}}{d_{0}} \sum_{j=1}^{d_{0}} \left[\left|x_{i}-x_{j}\right|^{2} + \left|v_{i}-v_{j}+g_{i}-g_{j}\right|^{2}\right] \\ &\leqslant (1-\bar{C}_{a,4}/2)a_{0} \left|x_{i}\right|^{2} + \delta K(1-\eta) \left|v_{i}\right|^{2}/[2K\delta] + c_{0}(1-\bar{\rho}_{\omega}K\delta) \left\langle x_{i},v_{i}\right\rangle + \bar{\rho}_{\omega}K\delta c_{0} \left\langle x,v\right\rangle \\ &+ (1-\bar{C}_{a,4}/(4b_{0}))2b_{0}U^{(q)}(x_{i}) + f_{K} + 2\eta\tilde{\eta}b_{K} \left\langle v_{i},g_{i}\right\rangle + \tilde{\eta}^{2}b_{K} \left|g_{i}\right|^{2} + c_{K}\tilde{\eta} \left\langle x_{i},g_{i}\right\rangle \\ &+ \frac{\epsilon e_{K}}{d_{0}} \sum_{j=1}^{d_{0}} \left[\left|x_{i}-x_{j}\right|^{2} + \left|v_{i}-v_{j}+g_{i}-g_{j}\right|^{2}\right], \end{split}$$

which completes the proof using the Cauchy-Schwarz inequality and the condition $c_0 \leq 2\sqrt{a_0b_0}$.

(b) The proof of the second statement follows the same lines using Lemma 21-(b) instead of Lemma 21-(a).

Averaging over momentum updates. We first give a technical result on certain Gaussian moments.

Lemma 24. Let $a \in \mathbb{R}_+$, $\ell \in \mathbb{N}^*$ and G be a d-dimensional zero-mean Gaussian random variable with covariance matrix identity. Then, for any $x \in \mathbb{R}^d$,

$$\mathbb{E}[(a + \langle G, x \rangle)^{\ell}] \leqslant a^{\ell - 2\lfloor \ell/2 \rfloor} \left(a^2 + \tilde{\mathbf{m}}_{\ell} |x|^2\right)^{\lfloor \ell/2 \rfloor}, \text{ where } \tilde{\mathbf{m}}_{\ell} = e^2(\ell/2)^2 \lfloor \ell/2 \rfloor^{-1} \mathbf{m}_{2\lfloor \ell/2 \rfloor}^{1/\lfloor \ell/2 \rfloor},$$
(76)

and $\mathbf{m}_{\ell'}$ is the ℓ' -th moment of the zero-mean one-dimensional Gaussian distribution with variance 1.

Proof. Expanding $(a + \langle G, x \rangle)^{\ell}$ and using that $\langle G, x \rangle$ is a zero-mean Gaussian random variable with variance $|x|^2$, we get

$$\mathbb{E}[(a+\langle G,x\rangle)^{\ell}] = \sum_{k=0}^{\ell} {\ell \choose k} a^{\ell-k} \mathbb{E}[\langle G,x\rangle^{k}] = \sum_{k=0}^{\lfloor \ell/2 \rfloor} {\ell \choose 2k} |x|^{2k} a^{\ell-2k} \mathbf{m}_{2k}^{2k/2k}$$
$$\leq a^{\ell-2\lfloor \ell/2 \rfloor} \sum_{k=0}^{\lfloor \ell/2 \rfloor} {\ell \choose 2k} |x|^{2k} a^{2(\lfloor \ell/2 \rfloor - k)} \mathbf{m}_{2\lfloor \ell/2 \rfloor}^{2k/2\lfloor \ell/2 \rfloor},$$

where we apply Jensen inequality for the last step. Using that $\binom{\ell}{2k} \leq (e\ell/(2k))^{2k} \leq (e^2\ell/2)^k (\ell/(2k))^k \leq (e^2(\ell/2)^2 \lfloor \ell/2 \rfloor^{-1})^k \binom{\lfloor \ell/2 \rfloor}{k}$, we get

$$\mathbb{E}[(a + \langle G, x \rangle)^{\ell}] \leq a^{\ell - 2\lfloor \ell/2 \rfloor} \sum_{k=0}^{\lfloor \ell/2 \rfloor} (e^{2}(\ell/2)^{2} \lfloor \ell/2 \rfloor^{-1})^{k} {\binom{\lfloor \ell/2 \rfloor}{k}} |x|^{2k} a^{2(\lfloor \ell/2 \rfloor - k)} \mathbf{m}_{2\lfloor \ell/2 \rfloor}^{2k/2\lfloor \ell/2 \rfloor}$$
$$\leq a^{\ell - 2\lfloor \ell/2 \rfloor} \left(a^{2} + e^{2}(\ell/2)^{2} \lfloor \ell/2 \rfloor^{-1} |x|^{2} \mathbf{m}_{2\lfloor \ell/2 \rfloor}^{1/\lfloor \ell/2 \rfloor} \right)^{\lfloor \ell/2 \rfloor},$$

which completes the proof.

Define for any $x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}) \in \mathbb{R}^d$ and $\ell \in \mathbb{N}^*$,

$$\mathfrak{V}_{\mathbf{o}}^{(\ell)}(x,v) = \sum_{i=1}^{d_0} \mathfrak{V}_{\mathbf{o},i}^{\ell}(x,v) \; .$$

We are now ready to control the evolution of the Lyapunov function $\mathfrak{V}_{\mathbf{0},i}^{\ell}$ for any $\ell \in \mathbb{N}^*$ over one full transition of the gHMC chain (notice that we are essentially interested in $\ell = 3$, since this is the dominant order in (52)).

Lemma 25. Assume that \mathbf{H}_{4} holds and that $\epsilon \leq \mathfrak{m}/4$. Then, for any $\boldsymbol{\omega} = (K, \delta, \eta), K \in \mathbb{N}$, $\delta > 0, \eta \in (0, 1), \mathbf{o} = (a_{0}, b_{0}, c_{0}, 2b_{0}, e_{0}, f_{0}) \in \mathbb{R}^{6}_{+}$, satisfying $\delta \leq \overline{\delta}_{1} \wedge \overline{\delta}, K\delta \leq \overline{T} \wedge \overline{T}_{1} \wedge \overline{T}_{2}$ with $\overline{T}, \overline{\delta} > 0, (70)$ -(71)-(73), it holds:

(a) For any $x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}) \in \mathbb{R}^d$ and $\ell \in \mathbb{N}^*$,

$$P_{\omega}\mathfrak{V}^{\ell}_{\mathbf{o},i}(x,v) \leq (1 - \bar{\rho}_{\omega}K\delta/4)^{\ell} [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^{\ell}$$

$$+ 2^{\ell} \lfloor \ell/2 \rfloor \bar{E}_{\ell} [(a_0 \bar{D}_2^x + b_0 (\bar{D}_2^v + 2\bar{D}_2^x) + c_0 (\bar{D}_2^x)^{1/2} (\bar{D}_2^v)^{1/2})^2 + \bar{E}_{\ell}]^{\lfloor \ell/2 \rfloor - 1} + 12^{\ell-1} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} [\tilde{\eta}^{2\ell} b_K^{\ell} q^{\ell} \mathbf{m}_{2\ell} + 8^{\ell-1} \epsilon^{\ell} e_K^{\ell} q^{\ell} \mathbf{m}_{2\ell} + f_K^{\ell}] + 12^{\ell-1} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} \epsilon^{\ell} e_K^{\ell} \left[d_0^{-1} \sum_{j=1}^{d_0} |x_i - x_j|^2 + |v_i - v_j|^2 \right]^{\ell} ,$$

where

$$\bar{D}_{1}^{x} = \frac{2^{3/2} (c_{K} \tilde{\eta})^{2} \tilde{\mathbf{m}}_{\ell}}{a_{0}^{2} \bar{\rho}_{\omega} K \delta(\sqrt{2} - 1)} , \quad \bar{D}_{1}^{v} = \frac{8\sqrt{2} (\tilde{\eta} \eta b_{K})^{2} \tilde{\mathbf{m}}_{\ell}}{\bar{\rho}_{\omega} K \delta(\sqrt{2} - 1)} , \quad (77)$$

$$\bar{D}_{2}^{x} = [2\bar{D}_{1}^{x}] \lor 4 \sqrt{\frac{\bar{D}_{1}^{v} \tilde{\mathbf{m}}_{\ell} (\eta \tilde{\eta} b_{K})^{2} \sqrt{2}}{\bar{\rho}_{\omega} K \delta a_{0}^{2} (\sqrt{2} - 1)}} , \quad \bar{D}_{2}^{v} = [2\bar{D}_{1}^{v}] \lor 2 \sqrt{\frac{\bar{D}_{1}^{x} \tilde{\mathbf{m}}_{\ell} (\tilde{\eta} c_{K})^{2} \sqrt{2}}{\bar{\rho}_{\omega} K \delta b_{0}^{2} (\sqrt{2} - 1)}} , \quad \bar{E}_{\ell} = [4(\tilde{\eta} \eta b_{K})^{2} \bar{D}_{2}^{v} + 2(c_{K} \tilde{\eta})^{2} \bar{D}_{2}^{x}] \tilde{\mathbf{m}}_{\ell} ,$$

where $\tilde{\mathbf{m}}_{\ell}$ is defined in (76) and $\tilde{\eta}^2 = 1 - \eta^2$. (b) In addition, if for $\ell \in \mathbb{N}^*$

$$\epsilon \leqslant \ell^{1/(\ell-1)} (1 - \bar{\rho}_{\omega} K \delta/4) , \quad 24e_K \epsilon (1 + 4/(\bar{\rho}_{\omega} K \delta))^{(\ell-1)/\ell} \leqslant \bar{\rho}_{\omega} K \delta/8 , \tag{78}$$

for any $x = (x_1, \ldots, x_{d_0}), v = (v_1, \ldots, v_{d_0}) \in \mathbb{R}^d$,

$$\begin{split} \mathbf{P}_{\boldsymbol{\omega}} \mathfrak{V}_{\mathbf{o}}^{(\ell)}(x,v) &\leqslant (1-\bar{\rho}_{\boldsymbol{\omega}}K\delta/8)^{\ell} \sum_{i=1}^{d_0} |a_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^{\ell} \\ &+ 4^{\ell} \lfloor \ell/2 \rfloor \, \bar{E}_{\ell} [(a_0 \bar{D}_1^x + b_0 (\bar{D}_1^v + 2\bar{D}_1^x) + c_0 (\bar{D}_1^x)^{1/2} (\bar{D}_1^v)^{1/2})^2 + \bar{E}_{\ell}]^{\lfloor \ell/2 \rfloor - 1} \\ &+ 12^{\ell-1} d_0 (1 + 4/(\bar{\rho}_{\boldsymbol{\omega}}K\delta))^{\ell-1} [\tilde{\eta}^{2\ell} b_K^{\ell} q^{\ell} \mathbf{m}_{2\ell} + 8^{\ell-1} \epsilon^{\ell} e_K^{\ell} q^{\ell} \mathbf{m}_{2\ell} + f_K^{\ell}] \,. \end{split}$$

(c) Finally, for any $k \in \{0, ..., K\}$, $x = (x_1, ..., x_{d_0}), v = (v_1, ..., v_{d_0}) \in \mathbb{R}^d$ and $\ell \in \mathbb{N}^*,$

$$\begin{aligned} \mathbf{D}_{\eta} \mathbf{V}_{\delta}^{k} \mathfrak{V}_{\mathbf{o}}^{(\ell)}(x,v) &\leqslant \bar{C}_{2}^{\ell} \sum_{i=1}^{d_{0}} |a_{i}|^{2} + b_{0} |v_{i}|^{2} + c_{0} \langle x_{i}, v_{i} \rangle + 2b_{0} U^{(q)}(x_{i})]^{\ell} \\ &+ 12^{\ell-1} d_{0} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} [\tilde{\eta}^{2\ell} b_{K}^{\ell} q^{\ell} \mathbf{m}_{2\ell} + 8^{\ell-1} \epsilon^{\ell} e_{K}^{\ell} q^{\ell} \mathbf{m}_{2\ell} + f_{K}^{\ell}] , \end{aligned}$$

where

$$\bar{C}_2 = 8(1 + \bar{\rho}_{\omega}K\delta/4)[\bar{C}_1 + ([4(\eta\tilde{\eta}b_K)^2b_0^{-1}] \vee [(c_K\tilde{\eta})^2a_0^{-1}])] + 24(1 + 4/(\bar{\rho}_{\omega}K\delta))^{(\ell-1)/\ell}\epsilon e_K .$$
(79)

Proof. Let $\delta \leq \bar{\delta}_1 \wedge \bar{\delta}$, $K\delta \leq \bar{T} \wedge \bar{T}_1 \wedge \bar{T}_2$ with $\bar{T}, \bar{\delta} > 0$, satisfying (70)-(71)-(73). (a) Consider $x, v \in \mathbb{R}^d$ and $\ell \in \mathbb{N}^*$. Using Lemma 22-(a) and Jensen inequality, we have, for any $g = (g_1, \ldots, g_{d_0}) \in \mathbb{R}^d$,

$$\begin{aligned} \mathbf{V}_{\delta}^{K} \mathfrak{V}_{\mathbf{o},i}^{\ell}(x,\eta v+\tilde{\eta}g) \\ &\leqslant (1+\bar{\rho}_{\omega}K\delta/4)^{\ell-1} [A+2\eta\tilde{\eta}b_{K}\left\langle v_{i},g_{i}\right\rangle +c_{K}\tilde{\eta}\left\langle x_{i},g_{i}\right\rangle]^{\ell} \end{aligned}$$

$$+ 12^{\ell-1} (1 + 4/\bar{\rho}_{\omega}K\delta)^{\ell-1} [\tilde{\eta}^{2\ell}b_{K}^{\ell} |g|^{2\ell} + \epsilon^{\ell}e_{K}^{\ell}d_{0}^{-1}\sum_{j=1}^{d_{0}} |g_{i} - g_{j}|^{2\ell} + f_{K}^{\ell}] + 12^{\ell-1} (1 + 4/\bar{\rho}_{\omega}K\delta)^{\ell-1}\epsilon^{\ell}e_{K}^{\ell} \left[d_{0}^{-1}\sum_{j=1}^{d_{0}} [|x_{i} - x_{j}|^{2} + |v_{i} - v_{j}|^{2} \right]^{\ell} ,$$

where we have set $A = (1 - \bar{\rho}_{\omega} K \delta) [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]$. Then, taking expectation and Lemma 24 imply that

$$P_{\omega}\mathfrak{V}^{\ell}_{\mathbf{o},i}(x,v) \leq (1+\bar{\rho}_{\omega}K\delta/4)^{\ell-1}A^{\ell-2\lfloor\ell/2\rfloor}B^{\lfloor\ell/2\rfloor}$$

$$+12^{\ell-1}(1+4/\bar{\rho}_{\omega}K\delta)^{\ell-1}[\tilde{\eta}^{2\ell}b^{\ell}_{K}q^{\ell}\mathbf{m}_{2\ell}+8^{\ell-1}\epsilon^{\ell}e^{\ell}_{K}q^{\ell}\mathbf{m}_{2\ell}+f^{\ell}_{K}]$$

$$+12^{\ell-1}(1+4/\bar{\rho}_{\omega}K\delta)^{\ell-1}\epsilon^{\ell}e^{\ell}_{K}d^{-1}\sum_{j=1}^{d_{0}}[|x_{i}-x_{j}|^{2}+|v_{i}-v_{j}|^{2}]^{\ell}.$$

$$(80)$$

with $B = A^2 + [4(\eta \tilde{\eta} b_K)^2 |v_i|^2 + (c_K \tilde{\eta})^2 |x_i|^2] \tilde{\mathbf{m}}_{\ell}$. For $|x_i|^2 \ge \bar{D}_1^x$ and $|v_i|^2 \ge \bar{D}_1^v$, we get using $c_0 \langle x_i, v_i \rangle \le (a_0 |x_i|^2 + b_0 |v_i|^2) / \sqrt{2}$ since $c_0 \le \sqrt{2a_0b_0}$ by the first line in (73),

$$\begin{split} B &\leqslant (1 - \bar{\rho}_{\omega} K \delta/2) [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^2 \\ &- \bar{\rho}_{\omega} K \delta(\sqrt{2} - 1) 2^{-1/2} [a_0 |x_i|^2 + b_0 |v_i|^2]^2 + [4(\eta \tilde{\eta} b_K)^2 |v_i|^2 + (c_K \tilde{\eta})^2 |x_i|^2] \tilde{\mathbf{m}}_{\ell} \\ &\leqslant (1 - \bar{\rho}_{\omega} K \delta/2) [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^2 . \end{split}$$

Similarly, for $|x_i|^2 \geqslant \bar{D}_2^x$ and $|v_i|^2 \leqslant \bar{D}_1^v$ or $|x_i|^2 \leqslant \bar{D}_1^x$ and $|v_i|^2 \geqslant \bar{D}_2^v$, we get
 $B \leqslant (1 - \bar{\rho}_{\omega} K \delta/2) [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^2 . \end{split}$

It yields for $|x_1|^2 \ge \bar{D}_2^x$ or $|v_i|^2 \ge \bar{D}_2^v$, $(1+\bar{\rho}_{\omega}K\delta/4)^{\ell-1}A^{\ell-2\lfloor\ell/2\rfloor}B^{\lfloor\ell/2\rfloor}$ $\leq (1 - \bar{\rho}_{\omega} K \delta/4)^{\ell} [a_0 |x_i|^2 + b_0 |v_i|^2 + c_0 \langle x_i, v_i \rangle + 2b_0 U^{(q)}(x_i)]^{\ell}.$

Therefore, the proof is completed for $|x_1|^2 \ge \bar{D}_2^x$ or $|v_i|^2 \ge \bar{D}_2^v$. It remains to consider the case where $|x_1|^2 \le \bar{D}_2^x$ and $|v_i|^2 \le \bar{D}_2^v$. Using that $(t+s)^{\ell'} - t^{\ell'} \le \ell' s(t+s)^{\ell'-1}$, for $s,t \ge 0$ and $\ell' \ge 1$, we get for $|x_1|^2 \le \bar{D}_2^x$.

and $|v_i|^2 \leqslant \bar{D}_2^v$,

$$B^{\lfloor \ell/2 \rfloor} - A^{2 \lfloor \ell/2 \rfloor} \leqslant (A^2 + \bar{E}_\ell)^{\lfloor \ell/2 \rfloor} - A^{2 \lfloor \ell/2 \rfloor} \leqslant \lfloor \ell/2 \rfloor \bar{E}_\ell B^{\lfloor \ell/2 \rfloor - 1}$$

Combining this inequality and (81), we obtain

$$P_{\omega}\mathfrak{V}_{\mathbf{o},i}^{\ell}(x,v) \leq (1+\bar{\rho}_{\omega}K\delta/4)^{\ell-1}A^{\ell} + (1+\bar{\rho}_{\omega}K\delta/4)^{\ell-1}\lfloor\ell/2\rfloor\bar{E}_{\ell}B^{\lfloor\ell/2\rfloor}$$

$$+ 12^{\ell-1}(1+4/\bar{\rho}_{\omega}K\delta)^{\ell-1}[\tilde{\eta}^{2\ell}b_{K}^{\ell}q^{\ell}\mathbf{m}_{2\ell} + 8^{\ell-1}\epsilon^{\ell}e_{K}^{\ell}q^{\ell}\mathbf{m}_{2\ell} + f_{K}^{\ell}]$$

$$+ 12^{\ell-1}(1+4/\bar{\rho}_{\omega}K\delta)^{\ell-1}\epsilon^{\ell}e_{K}^{\ell}d_{0}^{-1}\sum_{j=1}^{d_{0}}[|x_{i}-x_{j}|^{2} + |v_{i}-v_{j}|^{2}]^{\ell},$$

$$(81)$$

and the proof is completed using the definition of A, $(1 + \bar{\rho}_{\omega}K\delta/4)^{\ell-1}(1 - \bar{\rho}_{\omega}K\delta)^{\ell} \leq (1 - \bar{\rho}_{\omega}K\delta/2)^{\ell}, B \leq (a_0\bar{D}_2^x + b_0(\bar{D}_2^v + 2\bar{D}_2^x) + c_0(\bar{D}_2^x)^{1/2}(\bar{D}_2^v)^{1/2})^2 + \bar{E}_{\ell}$ since $U(x) \leq |x|^2$ under **H4** and $(1 + \bar{\rho}_{\omega}K\delta/4)^{\ell-1} \leq 2^{\ell}$.

(b) By Young's inequality, we have

$$\begin{split} \mathbf{P}_{\boldsymbol{\omega}}\mathfrak{V}_{\mathbf{o}}^{(\ell)}(x,v) &\leqslant (1-\bar{\rho}_{\boldsymbol{\omega}}K\delta/4)^{\ell}\sum_{i=1}^{d_{0}}|a_{i}|^{2}+b_{0}|v_{i}|^{2}+c_{0}\langle x_{i},v_{i}\rangle+2b_{0}U^{(q)}(x_{i})]^{\ell} \\ &+2^{\ell}\left\lfloor \ell/2\right\rfloor d_{0}\bar{E}_{\ell}[(a_{0}\bar{D}_{2}^{x}+b_{0}(\bar{D}_{2}^{v}+2\bar{D}_{2}^{x})+c_{0}(\bar{D}_{2}^{x})^{1/2}(\bar{D}_{2}^{v})^{1/2})^{\ell}+\bar{E}_{\ell}^{\ell}] \\ &+12^{\ell-1}(1+4/(\bar{\rho}_{\boldsymbol{\omega}}K\delta))^{\ell-1}d_{0}[\tilde{\eta}^{2\ell}b_{K}^{\ell}q^{\ell}\mathbf{m}_{2\ell}+8^{\ell-1}\epsilon^{\ell}e_{K}^{\ell}q^{\ell}\mathbf{m}_{2\ell}+f_{K}^{\ell}] \\ &+2\times24^{\ell-1}(1+4/(\bar{\rho}_{\boldsymbol{\omega}}K\delta))^{\ell-1}\epsilon^{\ell}e_{K}^{\ell}\sum_{i=1}^{d_{0}}[|x_{i}|^{2}+|v_{i}|^{2}]^{\ell} \,\,. \end{split}$$

Then the proof is completed using

 $a_{0} |x_{i}|^{2} + b_{0} |v_{i}|^{2} \leq \sqrt{2}/(\sqrt{2}-1) \{a_{0} |x_{i}|^{2} + b_{0} |v_{i}|^{2} + c_{0} \langle x_{i}, v_{i} \rangle \} \leq 4 \{a_{0} |x_{i}|^{2} + b_{0} |v_{i}|^{2} + c_{0} \langle x_{i}, v_{i} \rangle \},$ (82) the fact that by (78)

$$\begin{aligned} (1 - \bar{\rho}_{\omega} K \delta/4)^{\ell} &+ 2 \times 4 \times 24^{\ell-1} [a_0^{-1} \vee b_0^{-1}]^{\ell} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} \epsilon^{\ell} e_K^{\ell} \\ &\leqslant (1 - \bar{\rho}_{\omega} K \delta/4)^{\ell} + \ell (1 - \bar{\rho}_{\omega} K \delta/4)^{\ell-1} [24 e_K \epsilon [a_0^{-1} \vee b_0^{-1}] (1 + 4/(\bar{\rho}_{\omega} K \delta))^{(\ell-1)/\ell}] \\ &\leqslant (1 - \bar{\rho}_{\omega} K \delta/4 + 24 e_K \epsilon [a_0^{-1} \vee b_0^{-1}] (1 + 4/(\bar{\rho}_{\omega} K \delta))^{(\ell-1)/\ell})^{\ell} , \end{aligned}$$

and (78) again.

(c) Following the same lines as in the proof (a) using Lemma 22-(b) instead of Lemma 22(a) we have using (82)

$$\begin{split} \mathcal{D}_{\eta} \mathcal{V}_{\delta}^{k} \mathfrak{V}_{\mathbf{o},i}^{\ell}(x,v) &\leqslant \bar{C}_{2,0}^{\ell} [a_{0} \, |x_{i}|^{2} + b_{0} \, |v_{i}|^{2} + c_{0} \, \langle x_{i}, v_{i} \rangle + 2b_{0} U^{(q)}(x_{i})]^{\ell} \\ &\quad + 2^{\ell} \, \lfloor \ell/2 \rfloor \, \bar{E}_{\ell} [(a_{0} \bar{D}_{2}^{x} + b_{0} (\bar{D}_{2}^{v} + 2\bar{D}_{2}^{x}) + c_{0} (\bar{D}_{2}^{x})^{1/2} (\bar{D}_{1}^{v})^{1/2})^{\ell} + \bar{E}_{\ell}^{\ell}] \\ &\quad + 12^{\ell-1} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} [\tilde{\eta}^{2\ell} b_{K}^{\ell} q^{\ell} \mathbf{m}_{2\ell} + 8^{\ell-1} \epsilon^{\ell} e_{K}^{\ell} q^{\ell} \mathbf{m}_{2\ell} + f_{K}^{\ell}] \\ &\quad + 12^{\ell-1} (1 + 4/(\bar{\rho}_{\omega} K \delta))^{\ell-1} \epsilon^{\ell} e_{K}^{\ell} d_{0}^{-1} \sum_{j=1}^{d_{0}} \left[|x_{i} - x_{j}|^{2} + |v_{i} - v_{j}|^{2} \right]^{\ell} \,, \end{split}$$

with

$$\bar{C}_{2,0} = 8(1 + \bar{\rho}_{\omega}K\delta/4)[\bar{C}_1 + ([4(\eta\tilde{\eta}b_K)^2\tilde{\mathbf{m}}_{\ell}b_0^{-1}] \vee [(c_K\tilde{\eta})^2\tilde{\mathbf{m}}_{\ell}a_0^{-1}])].$$

The proof is then easily completed.

Conclusion. We can finally conclude this section by providing the proof of Theorem 18. Indeed, it only remains to chose carefully the initial parameters **o** so that Lemma 25 reads $P_{\omega}\mathfrak{V}_{\mathbf{o}}^{(\ell)} \leq (1 - \rho_{\omega}K\delta)^{\ell}\mathfrak{V}_{\mathbf{o}}^{(\ell)} + C$ for some constant C (with ρ_{ω} given by (54)), and then (53) will follow by induction on n and the equivalence between $\mathfrak{W}_{\omega,i}$ and $\mathfrak{V}_{\mathbf{o}}^{(\ell)}$.

Proof of Theorem 18. Taking

$$a_0 = \gamma_0^2/2$$
, $b_0 = 1$, $c_0 = \eta \gamma_0$, $\gamma_0 = (1 - \eta)/(\delta K)$, $f_0 = 0$, $e_0 = 0$.

As mentioned in Remark 23, the first condition in (73) is satisfied for this case. In addition, for this choice of parameter, the constants appearing in Lemma 21 become:

$$\begin{split} C_{a,1} &= \mathsf{m}/(3\times 2^5) \;, \quad C_{a,2} = \mathsf{m}/(3\times 2^6) \;, \\ C_{a,3} &= 20 + \gamma_0^2 \bar{T}(\mathsf{m}/(3\times 2^5) + \bar{T}\mathsf{m}/(3\times 2^6))/2 \;, \\ C_{b,1} &= \epsilon + \eta \gamma_0 (1+10\bar{\delta}) \;, \quad C_{b,2,0} = 4[(1+\epsilon)C_{b,1} + (1+10\bar{\delta})(\gamma_0^2 + C_{e,1}\epsilon)] \;, \\ C_{b,3,0} &= 17 + \gamma_0^2 + \bar{T}(2C_{a,3} + 17C_{b,1} + \epsilon C_{e,2} + 20\bar{T}(1+10\bar{\delta})) \;, \\ C_{b,2} &= C_{b,2,0} + \eta \gamma_0 \mathsf{m}/8 + a_0 \mathsf{m}\bar{T}/4 \;, \quad C_{b,3} = C_{b,3,0} + 20 + 2C_{a,3} \;, \\ C_{e,1} &= 37[2 + \eta \gamma_0 + \bar{T}(17\eta \gamma_0 + \gamma_0^2 + \bar{\delta}C_{b,3})] \;, \quad C_{e,2} = 37(\gamma_0^2 + \bar{T}C_{a,3}(\bar{\delta}C_{a,3} + 2\bar{T})) \;, \\ C_{f,1} &= \gamma_0^2 \mathsf{M}\bar{\delta}/2 + \bar{C}_{a,3}\mathsf{M}\bar{\delta}\bar{T} + \mathsf{m}\mathsf{M}\bar{\delta}/2 + \mathsf{M}(\eta \gamma_0 + \bar{T}\gamma_0^2/2) \;. \end{split}$$

Then, with these notations, $\delta_1, \bar{T}_1, \bar{T}_2$ in (72) can be written as

$$\delta_{1} = 17^{-1} \wedge [\mathfrak{m}/40] \wedge [\gamma_{0}^{2}2^{-6}/(48\epsilon C_{e,1} + C_{b,3}(1+10\bar{\delta}))] ,$$

$$T_{1} = 4^{-1} \left[17^{-1} \wedge C_{b,1}^{-1} \wedge C_{b,2,0}^{-1/2} \right] ,$$

$$T_{2} = 4^{-1} \{ [2^{-5}/(1+10\bar{\delta})] \wedge [2^{4}\gamma_{0}^{2}/C_{b,2}] \wedge \gamma_{0}/2^{4} \wedge 2 \} .$$
(83)

and the condition (70)-(71) are equivalent to

$$\epsilon \leq \epsilon_1 , \qquad \epsilon_1 = \gamma_0^2 m 2^{-6} / [1 + 48C_{e,1}] , \qquad (84)$$

and

$$\delta \leqslant \delta_2 , \qquad \delta_2 = \gamma_0 \mathfrak{m}(3 \times 2^8)^{-1} / [20 + \gamma_0^2 \bar{T}(C_{a,1} + \bar{T}C_{a,2})/2] . \tag{85}$$

In addition,

$$b_K \leqslant \overline{b}$$
, $\overline{b} = 2 + \overline{T}\overline{\delta}C_{b,3}$, $e_K \leqslant K\delta\overline{e}$, $\overline{e} = (C_{e,1} + \overline{\delta}C_{e,2})$. (86)

Furthermore, the second line of (73) is equivalent to $\delta \leqslant 1/(7\times 8)$ and

$$K\delta \leqslant T_3$$
, $T_3 = 4^{-1}\gamma_0 / [(1-\eta^2)^{1/2}C_{b,1} + C_{b,2}]$. (87)

Then, the proof is an easy consequence of Lemma 25 with ρ_{ω} given by (54). Condition (78) is equivalent using (86) to

$$\epsilon \leqslant \epsilon_2 , \quad \epsilon_2 = \left[\ell^{1/(\ell-1)} (1 - \bar{\rho}_{\omega} K \delta/4) \right] \wedge \left[(8 \times 24)^{-1} \bar{\rho}_{\omega} / \{ \bar{e} (1 + 4/(\rho_{\omega} \bar{T}))^{(\ell-1)/\ell} \} \right] . \tag{88}$$

and (77)-(79) can be written using (86) and $1 + \eta \leq 2$, as

$$D_{1} = 16\gamma_{0}\rho_{\omega}^{-1}\{\bar{\eta}^{2} + \bar{T}^{2}\gamma_{0}^{2}\}, \quad D_{2} = 32 \times 2\bar{\eta}^{2}\gamma_{0}/\rho_{\omega},$$

$$E_{\ell} = \delta K F_{\ell}, \quad F_{\ell} = 8\gamma_{0}[\bar{\eta}^{2}(2 + C_{b,3}\bar{\delta}\bar{T})D_{2} + (\gamma_{0}^{2}\bar{\eta}^{2} + \gamma_{0}^{4}\bar{T}^{2})D_{1}]\tilde{\mathbf{m}}_{\ell},$$

$$C_{2} = (1 + \rho_{\omega}\bar{T}/4)[C_{1} + ([8\gamma_{0}\bar{T}\bar{\eta}^{2}(2 + C_{b,3}\bar{\delta}\bar{T})] \vee [16\gamma_{0}\bar{T}(\bar{\eta}^{2} + \gamma_{0}^{2}\bar{T}^{2})])]$$

$$+ 24\bar{T}(C_{e,1} + \bar{\delta}C_{e,2})(4 + \rho_{\omega})^{(\ell-1)/\ell} ,$$

$$C_{1} = \left[2\{(1 + \bar{T} + \bar{T}^{3}m/4)\gamma_{0}^{2}/2 + 2\bar{\eta}(1 + \bar{T}^{2}m/8)\gamma_{0} + 2C_{a,3}\bar{\delta}\bar{T}\}/\gamma_{0}^{2}\right]$$

$$\vee \left[2 + 2\gamma_{0}\bar{\eta} + \bar{T}\gamma_{0}^{2}/2 + \bar{\delta}\bar{T}C_{b,3}\right] ,$$

where $\tilde{\mathbf{m}}_{\ell}$ is defined in (76). It completes the proof of (53) with

$$C_{1,\ell} = C_2 ,$$

$$C_{2,\ell} = 2^{\ell} \lfloor \ell/2 \rfloor F_{\ell} [(a_0 D_1 + b_0 (D_2 + 2D_1) + c_0 D_1^{1/2} D_2^{1/2})^{\ell} + \bar{T}^{\ell} F_{\ell}^{\ell}] ,$$

$$C_{3,\ell} = 2 \times 12^{\ell-1} \rho_{\omega}^{1-\ell} (4 + \rho_{\omega} \bar{T})^{\ell-1} [4^{\ell} \gamma_0^{\ell} (2 + \bar{\delta} \bar{T} C_{b,3}) \mathbf{m}_{2\ell} + 8^{\ell-1} \epsilon^{\ell} (C_{e,1} + \bar{\delta} C_{e,2})^{\ell} \mathbf{m}_{2\ell} + C_{f,1}] ,$$
(89)

(89) where $\mathbf{m}_{\ell'}$ is the ℓ' -th moment of the zero-mean one-dimensional Gaussian distribution with variance 1.