Minimum ℓ -degree thresholds for rainbow perfect matching in k-uniform hypergraphs

Jie You

Center for Applied Mathematics Tianjin University Tianjin 300072, P. R. China

E-mail: <u>yj_math@tju.edu.cn</u>

Abstract

Given $n \in k\mathbb{N}$ elements set V and k-uniform hypergraphs $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ on V. A rainbow perfect matching is a collection of pairwise disjoint edges $E_1 \in \mathcal{H}_1, \ldots, E_{n/k} \in \mathcal{H}_{n/k}$ such that $E_1 \cup \cdots \cup E_{n/k} = V$. In this paper, we determine the minimum ℓ -degree condition that guarantees the existence of a rainbow perfect matching for sufficiently large n and $\ell \geq k/2$.

1 Introduction

1.1 Background

Finding a spanning subgraph in a given graph or hypergraph is a fundamental problem in graph theory. In particular, it is desirable to fully characterize all graphs or hypergraphs that contain a spanning copy of a specific subgraph. For instance, Tutte's theorem [35] provides a characterization of all graphs that contain a perfect matching. However, for many hypergraphs, such a characterization is unlikely to exist due to the NP-completeness of the decision problem of whether a hypergraph \mathcal{H} contains a given subgraph \mathcal{F} . In fact, Garey and Johnson [10] showed that the decision problem of whether a k-uniform hypergraph contains a perfect matching is NP-complete for $k \geq 3$.

Given a set V of size n and an integer $k \geq 2$, we use $\binom{V}{k}$ to denote the family of all k-element subsets (k-subsets, for short) of V. A subfamily $\mathcal{H} \subseteq \binom{V}{k}$ is called a k-uniform hypergraph (or k-graph in short). For $\mathcal{H} \subseteq \binom{V}{k}$, we often use $V(\mathcal{H})$ to denote its vertex set V and use \mathcal{H} to denote its edge set. Define the complement of \mathcal{H} as $\overline{\mathcal{H}} := \binom{V}{k} \setminus \mathcal{H}$. Given $A \subseteq V$, let $\mathcal{H}[A]$ denote the sub k-graph of \mathcal{H} induced by A, namely, $\mathcal{H}[A] := \mathcal{H} \cap \binom{A}{k}$. Define $\mathcal{H} - A := \mathcal{H}[V(\mathcal{H}) \setminus A]$.

For $S \in \binom{V}{\ell}$ with $0 \leq \ell \leq k-1$, define the link graph of $S \mathcal{N}_{\mathcal{H}}(S) := \{T: S \cup T \in \mathcal{H}\}$, and let $\deg_{\mathcal{H}}(S)$ be the cardinality of $\mathcal{N}_{\mathcal{H}}(S)$. The minimum ℓ -degree $\delta_{\ell}(\mathcal{H})$ of \mathcal{H} is the minimum of $\deg_{\mathcal{H}}(S)$ over all ℓ -element subsets S of $V(\mathcal{H})$. Clearly, $\delta_0(\mathcal{H})$ is the number of edges in \mathcal{H} . We refer to $\delta_1(\mathcal{H})$ as the minimum vertex degree of \mathcal{H} and $\delta_{k-1}(\mathcal{H})$ as the minimum codegree of \mathcal{H} . We often omit the subscript \mathcal{H} when the context is clear.

Given a partition $V = A \cup B$, let $\mathcal{H}^0(A, B)$ (or $\mathcal{H}^1(A, B)$) denote the k-graph on V with edge set consists of all the edges intersect A in an even (or odd) number of vertices. Clearly, $\mathcal{H}^i(A, B)$ contains perfect matching if $n \in k\mathbb{N}$ and in/k has some parity with |A|.

Let $n \in k\mathbb{N}$. Define

 $ext(n,k) := \left\{ \mathcal{H}^i(A,B) \colon A \cup B = V, in/k \text{ has different parity with } |A| \right\}.$

Let

$$\delta(n,k,\ell) = \max_{\mathcal{H} \in ext(n,k)} \delta_{\ell}(\mathcal{H}).$$

Treglown and Zhao determine the minimum ℓ -degree condition that guarantees the existence of a perfect matching in k-graph.

Theorem 1.1 ([33, 34]). Given integers k, ℓ such that $k \geq 3$ and $k/2 \leq \ell \leq k-1$, there exists $n_0 \in \mathbb{N}$ such that the following holds. Suppose that \mathcal{H} is a k-graph on n vertices with $n \in k\mathbb{N}$ and $n \geq n_0$ satisfying $\delta_{\ell}(\mathcal{H}) > \delta(n, k, \ell)$, then \mathcal{H} contains a perfect matching.

Given k-uniform hypergraphs $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ on V, a rainbow perfect matching is a collection of pairwise disjoint edges $E_1 \in \mathcal{H}_1, \ldots, E_{n/k} \in \mathcal{H}_{n/k}$ such that $E_1 \cup \cdots \cup E_{n/k} = V$. Lu, Wang and Yu give the co-degree threshold for rainbow perfect matchings in k-graphs.

Theorem 1.2 ([23]). Given integers k, ℓ such that $k \geq 3$ and $k/2 \leq \ell \leq k-1$ and $n \in k\mathbb{N}$, there exists $n_0 \in \mathbb{N}$ such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on n vertices with $n \geq n_0$ and $n \in k\mathbb{N}$ satisfying $\delta_{k-1}(\mathcal{H}_i) > \delta(n, k, k-1)$ for $i = 1, \ldots, n/k$. Then $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ admit a rainbow matching.

In this paper, we determine the minimum ℓ -degree condition that guarantees the existence of a rainbow perfect matching in $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ for sufficiently large $n \in k\mathbb{N}$ and $\ell \geq k/2$, which is a generalisation of Theorem 1.1 and 1.2.

Theorem 1.3 (Main Result). Given integers k, ℓ such that $k \geq 3$ and $k/2 \leq \ell \leq k-1$ and $n \in k\mathbb{N}$, there exists $n_0 \in \mathbb{N}$ such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on n vertices with $n \geq n_0$ and $n \in k\mathbb{N}$ satisfying $\delta_{\ell}(\mathcal{H}_i) > \delta(n, k, \ell)$ for $i = 1, \ldots, n/k$. Then $\{\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}\}$ admits a rainbow matching.

It is shown in [13] that $\delta(n, k, \ell)$ takes its maximum value at $\mathcal{H}^i(A, B)$ for some A, B, i satisfy $-1 \leq |A| - |B| \leq 1$ and $2 \nmid in/k + |A|$, and denote one of the extremal graph as \mathcal{H}_{ext} . By letting $\mathcal{H}_1 = \ldots = \mathcal{H}_{n/k} = \mathcal{H}_{ext}$. We infer that there is no rainbow perfect matching in $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$. It implies that the minimum ℓ -degree condition in Theorem 1.3 is best possible.

It seems hard to compute the precise values of $\delta(n, k, \ell)$ for $\ell \leq k - 2$. In [33], it is showed that

$$\delta(n,k,k-1) = \begin{cases} n/2 - k + 2 & \text{if } k/2 \text{ is even and } n/k \text{ is odd} \\ n/2 - k + 3/2 & \text{if } k \text{ is odd and } (n-1)/2 \text{ is odd} \\ n/2 - k + 1/2 & \text{if } k \text{ is odd and } (n-1)/2 \text{ is even} \\ n/2 - k + 1 & \text{otherwise.} \end{cases}$$
(1.1)

The following proposition allows us to infer ℓ' minimum degree from ℓ minimum degree for $\ell' \leq \ell$. The proof is straightforward and omitted here. This result has also used in [33,34].

Proposition 1.4. Let $0 \leq \ell' \leq \ell < k$ and let \mathcal{H} be a k-graph. If $\delta_{\ell}(\mathcal{H}) \geq x \binom{n-\ell}{k-\ell}$ for some $0 \leq x \leq 1$, then $\delta_{\ell'}(\mathcal{H}) \geq x \binom{n-\ell'}{k-\ell'}$.

By Proposition 1.4, $\delta(n, k, \ell)$ are only known to be $(1/2 - o(1))\binom{n-\ell}{k-\ell}$.

We often write $0 < a_1 \ll a_2$ to mean that there are increasing functions f such that $a_1 \leq f(a_2)$. Throughout the paper, we omit floors and ceilings whenever this does not affect the argument.

1.2 **Proof overview**

We say a hypergraph is a (1, r)-graph if its vertex set can be partitioned into two parts $X \cup V$, such that r|X| = |V|, and every edge intersects X in exactly one vertex and intersects V in r vertices. Moreover, we say a subset $U \subseteq X \cup V$ is balanced if $r|U \cap X| = |U \cap V|$.

In [24], Lu, Yu, and Yuan introduced a (1, k)-graphs and reduced finding rainbow matchings in $\mathcal{H}_1, ..., \mathcal{H}_{n/k}$ to finding matchings in this (1, k)-graph. Define $\mathcal{T}(\mathcal{H}_1, ..., \mathcal{H}_{n/k})$ be the hypergraph with vertex set $X \cup V$ where $X = \{x_1, ..., x_{n/k}\}$ and edges set $\bigcup_{i=1}^t \{\{x_i\} \cup E : E \in \mathcal{H}_i\}$. It is clear that $\mathcal{H}_1, ..., \mathcal{H}_{n/k}$ contain a rainbow matching if and only if $\mathcal{T}(\mathcal{H}_1, ..., \mathcal{H}_{n/k})$ contains a perfect matching.

Let $\epsilon > 0$ and suppose that \mathcal{H} and \mathcal{H}' are r-graphs on $V(\mathcal{H})$. We say that \mathcal{H} is ϵ -close to \mathcal{H}' if \mathcal{H} can be made a copy of \mathcal{H}' by adding and deleting at most $\epsilon |V(\mathcal{H})|^r$ edges.

Denote $\mathcal{T}(\mathcal{H}_{ext}, \ldots, \mathcal{H}_{ext})$ by \mathcal{T}_{ext} , where there are $n/k \; H_{ext}$'s. As a common approach to obtain exact results, Theorem 1.3 is proven by distinguishing an *extremal* case from a *non-extremal* case and solve them separately.

Theorem 1.5 (Non-extremal Case). Given k, ℓ such that $k \geq 3$ and $k/2 \leq \ell \leq k-1$ and $\epsilon > 0$, there exists n_0 such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on vertex set V with $|V| = n \geq n_0$ and $n \in k\mathbb{N}$. Let $\mathcal{T} = \mathcal{T}(\mathcal{H}_1, \ldots, \mathcal{H}_{n/k})$. If \mathcal{T} is not ϵ -close to \mathcal{T}_{ext} and $\deg_{\ell}(\mathcal{H}_i) > \delta(n, k, \ell)$ for every $1 \leq i \leq n/k$, then \mathcal{T} contains a perfect matching.

Theorem 1.6 (Extremal Case). Given k, ℓ such that $k \ge 3$ and $k/2 \le \ell \le k-1$. There exists $\epsilon > 0$ and n_0 such that for $n \ge n_0$ the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on vertex set V with $|V| = n \ge n_0$ and $n \in k\mathbb{N}$. Let $\mathcal{T} = \mathcal{T}(\mathcal{H}_1, \ldots, \mathcal{H}_{n/k})$. If \mathcal{T} is ϵ -close to \mathcal{T}_{ext} and $\deg_{\ell}(\mathcal{H}_i) > \delta(n, k, \ell)$ for every $1 \le i \le n/k$, then \mathcal{T} contains a perfect matching.

In the packing problem, the absorb method is a commonly used technique, which was first introduced by Rödl, Ruciński and Szemerédi [30], and subsequently employed by many researchers (see [13, 21, 23, 25, 28, 33, 34, 37, etc]). To establish this method, we need to prove two lemmas: the Absorbing Lemma and the Almost Cover Lemma. The Absorbing Lemma involves finding a matching \mathcal{A} in \mathcal{T} of an appropriate size, such that for any sufficiently small subset $U \subset X \cup V$, where U is a balanced set, there exists a matching \mathcal{Q} with $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$. The matching \mathcal{A} is referred to as an absorb set. The Almost Cover Lemma involves finding a matching in \mathcal{T} that covers almost all the vertices. The proof of the Almost Cover Lemma follows the process outlined in [25]. By excluding an absorb set in advance, applying the Almost Cover Lemma to the remaining graph, and then absorbing the uncovered vertices into the absorb set, we can prove Theorem 1.3.

In order to establish the Absorbing Lemma, we require a counting lemma. Given a balanced (k + 1)-set, this lemma counts the number of size-2 matchings that can absorb this set. By applying the Frankl-Kupavskii concentration inequality, we can then prove the Absorbing Lemma. It should be noted that the counting lemma valid under the extra condition that \mathcal{T} is not close to the extremal hypergraph \mathcal{T}_{ext} . This is why we separate the proof into extremal and non-extremal cases.

For the extremal case, we mainly use a result of the author joint with Wang in [37]. This result states that if every vertex in an r-partite r-graph is "good", then a perfect matching exists. Here, the term "good" refers to a binary relation between graphs that is similar to, but stronger than, the concept of "close" as previously defined. The precise definition of "good" will be provided later, following the earlier explanation of "close". Therefore, we have two remaining tasks: The first one is to remove the vertices that are not "good" using a matching. Then we get a hypergraph with all the vertices are "good". The second one is to divide the remaining hypergraph into two (k + 1)-partite (k + 1)-graphs. By showing that vertices in these two (k + 1)-partite (k + 1)-graphs inherit the "good" property, we can apply the result in [37] to complete the proof for the extremal case.

The proofs of Theorems 1.5 and 1.6 are given in Sections 2 and 3, respectively.

2 Non-extremal Case

In this section, we deal with the non-extremal case by following the absorbing method initiated by Rödl, Ruciński and Szemerédi [30].

These following two lemmas will be used in the Non-extremal Case. The proofs of these two lemmas will be respectively deferred to Sections 2.1 and 2.2.

Lemma 2.1 (Almost cover). Given $\xi > 0$ and k, ℓ such that $k \ge 3$ and $k/2 \le \ell \le k-1$, there exists n_0 such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on vertex set V with $|V| = n \ge n_0$ and $n \in k\mathbb{N}$. Let $\mathcal{T} = \mathcal{T}(\mathcal{H}_1, \ldots, \mathcal{H}_{n/k})$. If for every $1 \le i \le n/k$,

$$\delta_{\ell}(\mathcal{H}_i) \ge \left(\frac{k-\ell}{k} - \frac{1}{k^{k-\ell}} + \xi\right) \binom{n-\ell}{k-\ell},$$

then \mathcal{T} contains a matching cover all but at most \sqrt{n} vertices.

Lemma 2.2 (Absorbing). Given $\epsilon > 0$ and k, ℓ such that $k \ge 3$ and $k/2 \le \ell \le k-1$, there exists γ and n_0 such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on vertex set V with $|V| = n \ge n_0$ and $n \in k\mathbb{N}$. Let $\mathcal{T} = \mathcal{T}(\mathcal{H}_1, \ldots, \mathcal{H}_{n/k})$. If \mathcal{T} is not ϵ -close to \mathcal{T}_{ext} , and $\delta_\ell(\mathcal{H}_i) > \delta(n, k, \ell)$ for every $1 \le i \le n/k$. Then there exists a matching \mathcal{A} in \mathcal{T} with $|V(\mathcal{A})| \le \gamma n$ such that for every balanced set $U \subset V(\mathcal{T}) \setminus V(\mathcal{A})$ with $|U| \le \gamma^8 n$, there exists a matching \mathcal{Q} in \mathcal{T} such that $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$.

We are now ready to prove Theorem 1.5.

Proof of Theorem 1.5. Using Lemma 2.2, we can find a matching \mathcal{A} in \mathcal{T} with $|V(\mathcal{A})| \leq \gamma n$. Removing this matching from \mathcal{T} , we obtain a (1, k)-graph \mathcal{T}' on the vertex set $X' \cup V'$, where $\mathcal{T}' = \mathcal{T} - V(\mathcal{A}), V' = V \setminus V(\mathcal{A})$, and $X' = X \setminus V(\mathcal{A})$. For $i \in X'$, let $\mathcal{H}'_i = \mathcal{H}_i - V(\mathcal{A})$. Let $n' = |V'| \geq n - \gamma n > n/2$. Recall that $\delta(n, k, \ell) = (1/2 - o(1)) \binom{n-\ell}{k-\ell}$, we have

$$\delta_{\ell}(\mathcal{H}'_{i}) \geq \delta(n,k,\ell) - |V(\mathcal{A})| \binom{n'-\ell-1}{k-\ell-1}$$
$$> \left(\frac{1}{2} - o(1) - \gamma k\right) \binom{n'-\ell}{k-\ell}$$
$$> \left(\frac{k-\ell}{k} - \frac{1}{k^{k-\ell}} + \xi\right) \binom{n'-\ell}{k-\ell},$$

when n is sufficiently large and $\gamma k < \xi = \frac{1}{2k^{k-\ell}}$.

Then by Lemma 2.1, there exists a matching \mathcal{M} in \mathcal{T}' cover all but at most \sqrt{n} vertices. Let $U = V(\mathcal{T}') \setminus V(\mathcal{M})$. Since $\sqrt{n} \ll \gamma^8 n$, by Lemma 2.2, there exists a matching \mathcal{Q} such that $V(\mathcal{Q}) = V(\mathcal{A}) \cup U$. Thus $\mathcal{M} \cup \mathcal{Q}$ forms a perfect matching of \mathcal{T} .

2.1 Proof of Almost Cover Lemma

The Almost Cover Lemma is a rainbow version of Lemma 2 in [25], and the proof is very similar. For the sake of completeness, we include the full proof as well.

For two hypergraphs \mathcal{H} and \mathcal{H}' , let $N(\mathcal{H}, \mathcal{H}')$ be the number of copies of \mathcal{H}' in \mathcal{H} . Same as in [25], the following results are needed in the proof.

Lemma 2.3 ([7]). For every integer $r \ge 2$, every d > 0, and every r-partite r-graph \mathcal{H}' , there exist c > 0 and n_0 such that for every r-graph \mathcal{H} on $n \ge n_0$ vertices and $|\mathcal{H}| \ge dn^r$, we have $N(\mathcal{H}, \mathcal{H}') \ge cn^{|V(\mathcal{H}')|}$.

Fact 2.4 ([25]). For all integer $k \ge 1$, $n \ge 2$, and $1 \le t \le n-1$, the maximum number of edges in a k-partite k-graph with n vertices in each class and no matching of t + 1 is tn^{k-1} .

Lemma 2.5 ([27]). Let M_1, \ldots, M_N be matchings each of size t in a r-partite r-uniform hypergraph. If $N > (t-1)t^r$, then there exists i_1, \ldots, i_t and pairwise disjoint $e_{i_1} \in M_{i_1}, \ldots, e_{i_t} \in M_{i_t}$.

Proof of Lemma 2.1. Let \mathcal{M} be a matching in \mathcal{T} that maximizes the size $|\mathcal{M}|$. Assume to the contrary that $n - |V(\mathcal{M})| \ge \sqrt{n}$, and let $X_1 = X \setminus V(\mathcal{M})$ and $V_1 = V \setminus V(\mathcal{M})$.

The proof strategy involves finding a matching Q in T such that Q intersects at most |Q| - 1 elements in \mathcal{M} . By replacing these elements with Q, we increase the size of \mathcal{M} . It is precisely due to the use of this strategy that we can safely assume $n - |V(\mathcal{M})| = \sqrt{n}$; otherwise, we can augment \mathcal{M} by adding any (k + 1)-balanced sets from $X_1 \times V_1$ until $n - |V(\mathcal{M})| = \sqrt{n}$ is achieved.

For each $S \in X_1 \times \binom{V_1}{\ell}$, define

$$L_{S}(\mathcal{M}) := \left\{ T \in \binom{V(\mathcal{M})}{k-\ell} : S \cup T \in \mathcal{T}, \forall e \in \mathcal{M}, |T \cap e| \le 1 \right\}.$$

Note that at most $o(n^{k-\ell})$ edges in \mathcal{T} that contain S and intersect any edge in \mathcal{M} with more than one vertex. Moreover, at most $\sqrt{n}\binom{n-\ell-1}{k-\ell-1}$ edges intersecting at least one vertex outside $V(\mathcal{M})$. Due to the minimum ℓ -degree condition, we have a lower bound

$$|L_S(\mathcal{M})| > \left(\frac{k-\ell}{k} - \frac{1}{k^{k-\ell}} + \xi - o(1)\right) \binom{n-\ell}{k-\ell}.$$
(2.1)

Clearly, each element in $L_S(\mathcal{M})$ intersect exactly $k - \ell$ edges in \mathcal{M} . Then

$$L_{S}(\mathcal{M}) = \bigcup_{\mathcal{E} \in \binom{\mathcal{M}}{k-\ell}} L_{S}(\mathcal{E}).$$

$$(2.2)$$

We break the family $\binom{\mathcal{M}}{k-\ell}$ into two parts $\binom{\mathcal{M}}{k-\ell} = A(S) \cup B(S)$, where $A(S) = \{\mathcal{E} \in \binom{\mathcal{M}}{k-\ell} : |L_S(\mathcal{E})| \le (k-\ell)k^{k-\ell-1}-1\}$ and $B(S) = \binom{\mathcal{M}}{k-\ell} \setminus A(S)$.

The equation (2.2) and the trivial bounds $|L_S(\mathcal{E})| \leq k^{k-\ell}, |A(S)| \leq {|\mathcal{M}| \choose k-\ell}, |\mathcal{M}| \leq n/k$ imply that

$$|L_S(\mathcal{M})| \le k^{k-\ell} |B(S)| + ((k-\ell)k^{k-\ell-1} - 1) \binom{|\mathcal{M}|}{k-\ell}$$
$$\le \left(\frac{|B(S)|}{\binom{|\mathcal{M}|}{k-\ell}} + \frac{k-\ell}{k} - \frac{1}{k^{k-\ell}}\right) \binom{n-\ell}{k-\ell},$$

Together with the lower bound (2.1), then

$$|B(S)| > (1 - o(1)) \binom{|\mathcal{M}|}{k - \ell}.$$
 (2.3)

According to Fact 2.4, for any $\mathcal{E} \in B(S)$, the maximum matching size of $L_S(\mathcal{E})$ is at least $k - \ell$, and $k - \ell$ achieved only when $L_S(\mathcal{E})$ is isomorphic to a $(k - \ell)$ -partite $(k - \ell)$ -graph on these $k(k - \ell)$ vertices that consists of ℓ isolated vertices belonging to the same vertex part, along with all the remaining possible edges. Let us denote the set of \mathcal{E} such that maximum matching size of $L_S(\mathcal{E})$ is $k - \ell$ as $B_1(S)$, and $B_2(S) = B(S) \setminus B_1(S)$.

Starting from the minimum degree condition, we derived (2.3). Now, we will establish the following two claims based on the maximality of $|\mathcal{M}|$.

Claim 2.6. For at most $\xi|X_1|\binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$, we have $|B_2(S)| \ge \frac{\xi}{3} \binom{|\mathcal{M}|}{k-\ell}$.

Proof of Claim. Let us assume, for the sake of contradiction, that there are at least $\xi|X_1|\binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$ such that $|B_2(S)| \geq \frac{\xi}{3} \binom{|\mathcal{M}|}{k-\ell}$. Then, by averaging, there exists $\mathcal{E}_0 \in \binom{\mathcal{M}}{k-\ell}$ such that $\mathcal{E}_0 \in B_2(S)$ for at least $\frac{\xi^2}{3}|X_1|\binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$. From these sets, recall that $|V_1| = \sqrt{n}$ and $|X_1| = \sqrt{n}/k$ we can choose a collection of pairwise disjoint sets S_1, S_2, \ldots, S_N where $N > (k - \ell + 1)^{r+1}$. According to Lemma 2.5, by rearranging the indices there exist pairwise disjoint elements $T_1 \in L_{S_1}(\mathcal{E}_0), T_2 \in L_{S_2}(\mathcal{E}_0), \ldots, T_{k-\ell+1} \in L_{S_{k-\ell+1}}(\mathcal{E}_0)$.

Thus $\mathcal{M} \setminus \mathcal{E}_0 \cup \{S_1 \cup T_1, \dots, S_{k-\ell+1} \cup T_{k-\ell+1}\}$ forms a matching of \mathcal{T} of size larger than \mathcal{M} . This is a contradiction with the maximality of $|\mathcal{M}|$.

Claim 2.7. For at most $\xi |X_1| \binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$, we have $|B_1(S)| \geq \frac{\xi}{3} \binom{|\mathcal{M}|}{k-\ell}$.

Proof of Claim. Assume that there are at least $\xi |X_1| \binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$ such that $|B_1(S)| \geq \frac{\xi}{3} \binom{|\mathcal{M}|}{k-\ell}$. Then, there are at least $\frac{\xi^2}{6} \binom{|\mathcal{M}|}{k-\ell}$ elements $\mathcal{E} \in \binom{\mathcal{M}}{k-\ell}$ such that each $\mathcal{E} \in B_1(S)$ for at least $\frac{\xi^2}{6} |X_1| \binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$. From these $\frac{\xi^2}{6} \binom{|\mathcal{M}|}{k-\ell}$ sets $\mathcal{E} \in \binom{\mathcal{M}}{k-\ell}$, it is possible to choose three sets, $\mathcal{E}_1, \mathcal{E}_2$, and \mathcal{E}_3 , that each pair of sets has precisely one common element, and the intersection between any two sets is distinct. Indeed, by Lemma 2.3 such configuration exist.

Without loss of generality, assume that $\mathcal{E}_1 = \{f_1, \ldots, f_{k-\ell-1}, e_1\}$, $\mathcal{E}_2 = \{e_1, \ldots, e_{k-\ell-1}, g_1\}$ and $\mathcal{E}_3 = \{g_1, \ldots, g_{k-\ell-1}, f_1\}$. It follows from the definition that, for each i = 1, 2, 3, there exist $\frac{\xi^2}{6}|X_1|\binom{|V_1|}{\ell}$ sets $S \in X_1 \times \binom{V_1}{\ell}$ such that $\mathcal{E}_i \in B_1(S)$. Among these S, let us arbitrarily choose S^1, S^2, S^3 satisfying $\mathcal{E}_i \in B_1(S^i)$. As mentioned previously, it is known that $L_{S^i}(\mathcal{E}_i)$ is isomorphic to a $(k - \ell)$ -partite $(k - \ell)$ -graph with ℓ isolated vertices. Importantly, these isolated vertices are all contained within the same partite set. Therefore, it is guaranteed that there exist two of $L_{S^i}(\mathcal{E}_i)$ where the isolated vertices are not in the same element of \mathcal{M} . Without loss of generality, let us assume that the isolated vertices of $L_{S^1}(\mathcal{E}_1)$ and $L_{S^2}(\mathcal{E}_2)$ are not in the same element of \mathcal{M} . Since $L_{S^1}(\mathcal{E}_1)$ and $L_{S^2}(\mathcal{E}_2)$ contain all the remaining possible edges, it follows that $L_{S^1}(\mathcal{E}_1) \cup L_{S^2}(\mathcal{E}_2)$ contains a matching $T_1^1, \ldots, T_{k-\ell}^1, T_1^2, \ldots, T_{k-\ell}^2$ such that $T_i^1 \in L_{S^1}(\mathcal{E}_1)$ and $T_i^2 \in L_{S^2}(\mathcal{E}_2)$ for $i = 1, 2 \ldots, k - \ell$.

Recall that $\mathcal{E}_i \in B_1(S)$ for at least $\frac{\xi^2}{3} |X_1| {|V_1| \choose \ell}$ sets $S \in X_1 \times {|V_1| \choose \ell}$. Moreover $|V_1| = \sqrt{n}$ and $|X_1| = \sqrt{n/k}$, from these sets we can choose a collection of pairwise disjoint sets $S_1^1, S_2^1, \ldots, S_N^1, S_1^2, S_2^2, \ldots, S_N^2$ where $\mathcal{E}_i \in B_1(S_j^i)$ for i = 1, 2 and $j = 1, \ldots, N$, $N > (k - \ell + 1)^{r+1}$. According to Lemma 2.5, by rearranging the indices there exist pairwise disjoint elements $T_j^i \in L_{S_s^i}(\mathcal{E}_i)$ for $j = 1, \ldots, k - \ell$ and i = 1, 2.

Note that $|\mathcal{E}_1 \cup \mathcal{E}_2| = 2(k-\ell) - 1$, thus $(\mathcal{M} \setminus (\mathcal{E}_1 \cup \mathcal{E}_2)) \cup \{S_j^i \cup T_j^i\}_{1 \le j \le k-\ell, i=1,2}$ forms a matching in \mathcal{T} of size $|\mathcal{M}| - |\mathcal{E}_1 \cup \mathcal{E}_2| + 2(k-\ell) = |\mathcal{M}| + 1$. This is a contradiction with the maximality of $|\mathcal{M}|$.

The (2.3) implies that for each $S \in X_1 \times {\binom{V_1}{\ell}}$, we have $|B_1(S)| + |B_2(S)| > (1 - o(1)) {\binom{|\mathcal{M}|}{k-\ell}}$. Therefore either half of $S \in X_1 \times {\binom{V_1}{\ell}}$ satisfy $B_1(S) > \frac{1 - o(1)}{2} {\binom{|\mathcal{M}|}{k-\ell}}$ or half of $S \in X_1 \times {\binom{V_1}{\ell}}$ satisfy $B_2(S) > \frac{1 - o(1)}{2} {\binom{|\mathcal{M}|}{k-\ell}}$, which contradict with at least one of Claim 2.6 and Claim 2.7. This concludes the proof of Lemma 2.1.

2.2 Proofs of the Absorbing Lemma.

Let A be the vertex set of 1 (or 2) edges in \mathcal{T} . For a balanced (k+1)-set E, we say A is an 1- (or 2-)absorber for E if there exists a matching \mathcal{Q} in \mathcal{T} such that $V(\mathcal{Q}) = V(E) \cup V(A)$.

We prove Absorbing Lemma via the following counting lemma, which itself is proved in Section 2.3.

Lemma 2.8 (Counting). Given $\epsilon > 0$ and k, ℓ such that $k \ge 3$ and $k/2 \le \ell \le k - 1$, there exists γ and n_0 such that the following holds. Suppose that $\mathcal{H}_1, \ldots, \mathcal{H}_{n/k}$ are k-graphs on vertex set V with $|V| = n \ge n_0$ and

 $n \in k\mathbb{N}$. Let $\mathcal{T} = \mathcal{T}(\mathcal{H}_1, \ldots, \mathcal{H}_{n/k})$. If $\delta_{\ell}(\mathcal{H}_i) > \delta(n, k, \ell)$ for every $1 \leq i \leq n/k$ and \mathcal{T} is not ϵ -close to \mathcal{T}_{ext} . Then for each balanced (k+1)-set E, there exists at least $\gamma^5 n^{2(k+1)}$ 2-absorbers.

We also need a concentration inequality due to Frankl and Kupavskii.

Lemma 2.9 (Frankl-Kupavskii Concentration Inequality, [8]). Suppose that m, k, t are integers and $m \geq tk$. Let $\mathcal{G} \subset \binom{[m]}{k}$ be a family, and $\theta = |\mathcal{G}| / \binom{[m]}{k}$. Let η be the random variable equal to the size of the intersection of \mathcal{G} with a t-matching \mathcal{B} of k-sets, chosen uniformly at random. Then $\mathbb{E}[\eta] = \theta t$ and, for any positive γ , we have

$$\Pr[|\eta - \theta t| \ge 2\gamma \sqrt{t}] \le 2e^{-\gamma^2/2}.$$
(2.4)

Proof of Lemma 2.2. By Lemma 2.8, for each balanced (k + 1)-set E, there are at least $\gamma^5 n^{2(k+1)}$ 2-absorbers of E. We denote the family of absorbers by $\mathcal{A}(E)$.

Let $m = \gamma n/2(k+1)$. Let $\mathcal{M}' \subseteq {V(\mathcal{T}) \choose 2(k+1)}$ be a matching of size m chosen uniformly at random. By Theorem 2.9, we have

$$\Pr\left(\left|\left|\mathcal{M}'\cap\mathcal{A}(E)\right| - \frac{\gamma^5 n^{2(k+1)}}{\binom{n+n/k}{2(k+1)}}m\right| > 2\gamma^6 m\right) < 2e^{-\frac{\gamma^{12}}{2}m}, \quad \text{for all } E.$$

Since $2e^{-\frac{\gamma^{12}}{2}m} < \frac{1}{n^{2(k+1)}}$ for sufficiently large *n*, by the union bound, with probability more than 0, we can choose \mathcal{M}' such that for all E,

$$|\mathcal{M}' \cap \mathcal{A}(E)| > \frac{\gamma^5 n^{2(k+1)}}{\binom{n+n/k}{2(k+1)}} m - 2\gamma^6 m > \gamma^7 n.$$
(2.5)

Removing all non-absorbing 2(k+1)-sets in \mathcal{M}' , we get a matching \mathcal{M} .

In the following, for any balanced set U of size at most $(k+1)\gamma^8 n$, we are tring to absorb U by \mathcal{M} . First, part U into balanced (k+1)-sets $E_1, E_2, \ldots, E_j, 1 \leq j \leq \gamma^8 n$. By (2.5), for each $i \in [j]$ there are at least $\gamma^7 n$ 2-absorbers for E_i in \mathcal{M} . Let us absorb E_i by element in \mathcal{M} step by step, in the *i*-th step, there are at least

$$\gamma^7 n - |U \cap A_1 \cup \ldots \cup A_{i-1}| \binom{V(\mathcal{H})}{2k+1} > \gamma^7 n - 3(k+1)\gamma^8 n \binom{n+n/k}{2k+1} > 0$$

2-absorbers in \mathcal{M} disjoint to $U \cup A_1 \cup \ldots \cup A_{i-1}$, choose one of it and denote by A_i . Thus, we obtain $\{A_1, A_2, \ldots, A_j\} \subset \mathcal{M} - U$ and A_i absorbs E_i . Note that \mathcal{M} is obtained by removing all non-absorbing 2(k+1)-sets in \mathcal{M}' , thus \mathcal{M} is consisted by 2-absorbers. Since each absorber is a matching in \mathcal{T} of size 2, thus \mathcal{M} is a matching in \mathcal{T} with $|V(\mathcal{M})| \leq \gamma n$ and \mathcal{M} absorb any balanced set U with $|U| \leq (k+1)\gamma^8 n$.

2.3 Proof of Counting Lemma

We follow the similar approach as [34].

For any given balanced (k + 1)-set E, and a partition $E = \{x\} \cup L \cup R$ such that $x \in X$, $|L| = \lceil k/2 \rceil$, and $|R| = \lfloor k/2 \rfloor$. Let $\ell = \binom{n}{\lceil k/2 \rceil}$ and $r = \binom{n}{\lfloor k/2 \rfloor}$. Denote $\mathcal{F} = \mathcal{N}_{\mathcal{T}}(x)$. By Proposition 1.4, the degree assumption $\delta_{\ell}(\mathcal{F}) > \delta(n, k, \ell)$ implies that

$$\deg_{\mathcal{F}}(L) > (\frac{1}{2} - \frac{1}{2}\gamma)r, \quad \deg_{\mathcal{F}}(R) > (\frac{1}{2} - \frac{1}{2}\gamma)\ell, \tag{2.6}$$

for any $\gamma > 0$ and sufficiently large n.

Claim 2.10 ([34]). Given $\epsilon > 0$, there exists $\gamma > 0$ and n_0 such that the following holds. Let \mathcal{H} be a k-graph on V, $|V| = n > n_0$. If \mathcal{H} is not ϵ -close to \mathcal{H}_{ext} , then one of the following holds.

(a) For any $L \in \binom{V}{\lceil k/2 \rceil}$, there are at least $(\frac{1}{2} + \gamma)\ell L' \in \binom{V}{\lceil k/2 \rceil}$ such that $|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| \ge \gamma r$. (b) $|\{R' \in \binom{V}{\lfloor k/2 \rceil} : |\mathcal{N}_{\mathcal{H}}(R')| \ge (\frac{1}{2} + \gamma)\ell\}| \ge 2\gamma r$.

Claim 2.11. For any $x' \in X \setminus \{x\}$. Let $\mathcal{H} = \mathcal{N}_{\mathcal{T}}(x')$. If there are at least $\gamma \ell/2$ $L' \in \mathcal{N}_{\mathcal{F}}(R)$ such that $|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| \geq \gamma r$, then there are at least $\frac{\gamma^3}{2}n^k$ 1-absorbers A for the set E with $|A \cap X| = \{x'\}$.

Proof. We first choose a $L' \in \mathcal{N}_{\mathcal{F}}(R)$ disjoint to $L \cup R$ such that $|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| \ge \gamma r$, the number of choices is at least

$$\gamma \ell/2 - |L \cup R| \binom{n}{\lceil k/2 \rceil - 1} > \frac{\gamma}{3} \ell$$

Then we choose a $R' \in \mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')$ disjoint to L, R, L', the number of choices is at least

$$|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| - |L \cup L' \cup R| \binom{n}{\lfloor k/2 \rfloor - 1} > \frac{\gamma}{2}r$$

Therefore $\{x'\} \cup L' \cup R'$ forms a 1-absorbers of E, since $\{x'\} \cup L' \cup R'$, $\{x\} \cup L' \cup R$, and $\{x'\} \cup L \cup R'$ are edges of \mathcal{T} . The number of such absorbers is at least $\frac{\gamma}{3}\ell\frac{\gamma}{2}r > \gamma^3 n^k$.

Claim 2.12. Given $x_1, x_2 \in X \setminus \{x\}$, there are at least $\gamma^3 n^k$ 1-absorbers or at least $\gamma^4 n^{2k}$ 2-absorbers A satisfy $|A \cap X| \subseteq \{x_1, x_2\}$ for E.

Proof. If case (a) in Claim 2.10 holds for $\mathcal{H} = \mathcal{N}_{\mathcal{T}}(x_1)$ or $\mathcal{N}_{\mathcal{T}}(x_1)$. Without lose of generality, assume holds for $\mathcal{H} = \mathcal{N}_{\mathcal{T}}(x_1)$. Then there are at least $(\frac{1}{2} + \gamma)\ell L' \in \binom{V}{\lfloor k/2 \rfloor}$ such that $|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| \geq \gamma r$. Therefore, the number of $L' \in \mathcal{N}_{\mathcal{F}}(R)$ such that $|\mathcal{N}_{\mathcal{H}}(L) \cap \mathcal{N}_{\mathcal{H}}(L')| \geq \gamma r$ is at least

$$|\mathcal{N}_{\mathcal{F}}(R)| + (\frac{1}{2} + \gamma)\ell - \ell > \frac{\gamma\ell}{2}.$$

By Claim 2.11, there are at least $\gamma^3 n^k$ 1-absorbers A for E such that $|A \cap X| = \{x_1\}$. We are done.

Now we assume that case (b) in Claim 2.10 holds for $\mathcal{H} = \mathcal{N}_{\mathcal{T}}(x_1)$. Let

$$\mathcal{R} := \left\{ R' \in \binom{V}{\lfloor k/2 \rfloor} : |\mathcal{N}_{\mathcal{H}}(R')| \ge (\frac{1}{2} + \gamma)\ell \right\}.$$

Furthermore, we can assume that

$$|\mathcal{R} \cap \mathcal{N}_{\mathcal{H}}(L)| < \frac{\gamma r}{2} \tag{2.7}$$

and

$$|\{L' \in \mathcal{N}_{\mathcal{F}}(R) \colon |\mathcal{N}_{\mathcal{H}}(L') \cap \mathcal{N}_{\mathcal{H}}(L)| \ge \gamma r\}| < \frac{\gamma \ell}{2}$$
(2.8)

If (2.8) does not hold, then according to Claim 2.11, there exist at least $\frac{\gamma^3}{2}n^k$ 1-absorbers A for E such that $|A \cap E| = \{x_1\}$. On the other hand, if (2.7) does not hold, then we have $|\{R' \in \mathcal{N}_{\mathcal{F}}(L) : |\mathcal{N}_{\mathcal{H}}(R') \cap \mathcal{N}_{\mathcal{H}}(R)| \ge \gamma \ell\}| > \gamma r/2$, which can be viewed as a mirror case of (2.8).

First, choose $L' \in \mathcal{N}_{\mathcal{F}}(R)$ disjoint to $L \cup R$ such that $|\mathcal{N}_{\mathcal{H}}(L') \cap \mathcal{N}_{\mathcal{H}}(L)| < \gamma r$. By (2.6) and (2.8) the number of choices is at least

$$(\frac{1}{2} - \frac{\gamma}{2})\ell - \frac{\gamma\ell}{2} > \frac{\gamma\ell}{3}.$$

By (2.6),(2.7), and the fact that $|\mathcal{N}_{\mathcal{H}}(L') \cap \mathcal{N}_{\mathcal{H}}(L)| < \gamma r$, we can conclude that due to the definition $|\mathcal{R}| > 2\gamma r$, the following holds:

$$|\mathcal{N}_{\mathcal{H}}(L') \cap \mathcal{R}| \ge \frac{\gamma r}{2}.$$
(2.9)

Second, choose $R' \in \mathcal{N}_{\mathcal{H}}(L') \cap \mathcal{R}$ disjoint to $L \cup R \cup L'$. By (2.9) the number of choices is at least $\frac{\gamma r}{3}$. Denote $\mathcal{G} = \mathcal{N}_{\mathcal{T}}(x_2)$. Third, choose $R'' \in \mathcal{N}_{\mathcal{G}}(L)$ disjoint to $L \cup R \cup L' \cup R'$, the number of choices is at least $\frac{1}{3}r$. By (2.6), for each $R'' \in \mathcal{N}_{\mathcal{G}}(L)$,

$$|\mathcal{N}_{\mathcal{G}}(R'') \cap \mathcal{N}_{\mathcal{H}}(R')| \ge (\frac{1}{2} - \frac{1}{2}\gamma)\ell + (\frac{1}{2} + \gamma)\ell - \ell = \frac{\gamma}{2}\ell.$$

Last, choose $L'' \in \mathcal{N}_{\mathcal{G}}(R'') \cap \mathcal{N}_{\mathcal{H}}(R')$ disjoint to $L \cup R \cup L' \cup R' \cup R''$, the number of choices is at least $\frac{\gamma}{3}\ell$. Thus, $\{x_1\} \cup L' \cup R', \{x_2\} \cup L'' \cup R'' \in \mathcal{T}$, and $\{x\} \cup L' \cup R, \{x_1\} \cup L'' \cup R', \{x_2\} \cup L \cup R'' \in \mathcal{T}$. Therefore, $\{\{x_1\} \cup L' \cup R', \{x_2\} \cup L'' \cup R''\}$ is a 2-absorber of E. The choice number is at least

$$\frac{\gamma}{3}\ell\frac{\gamma}{3}r\frac{1}{3}r\frac{\gamma}{3}\ell > \gamma^4 n^{2k}$$

There are $\binom{n/k-1}{2}$ choices for $\{x_1, x_2\}$, by Claim 2.12 either for half of choices $\{x_1, x_2\} \in \binom{X \setminus \{x\}}{2}$ there are at least $\gamma^4 n^k$ 2-absorbers such that $|A \cap X| = \{x_1, x_2\}$, or half of choices of $\{x_1, x_2\}$ there are at least $\gamma^3 n^k$ 1-absorbers such that $|A \cap X| \in \{x_1, x_2\}$. The former implies that there are at least

$$\binom{n/k-1}{2}\gamma^4n^{2k}>\gamma^5n^{2(k+1)}$$

2-absorbers for E in \mathcal{T} . The later implies that there are at least

$$\binom{n/k-1}{2}\gamma^3 n^k/n > \gamma^4 n^{k+1}$$

1-absorbers for E in \mathcal{T} ,

Claim 2.13. If there are at least m 1-absorbers for E in \mathcal{T} , then there are at least $m\gamma n^{k+1}$ 2-absorbers for E in \mathcal{T} .

Proof. By the degree assumption, for each $x \in X$, the size of $\mathcal{N}_{\mathcal{T}}(x)$ is at least

$$\deg_{\mathcal{T}}(x) \ge \binom{n}{\ell} \left(\frac{1}{2} - \gamma\right) \binom{n-\ell}{k-\ell} / \binom{k}{\ell} > \frac{1}{3} \binom{n}{k}$$

Let \mathcal{A} be the family consists of all 1-absorbers for E. For any $A \in \mathcal{A}$, the number of edges in \mathcal{T} that disjoint $A \cup E$ is at least

$$(|X|-2)\left(\deg_{\mathcal{T}}(x) - |A \cup E|\binom{n-1}{k-1}\right) > \frac{1}{4}\binom{n+1}{k+1}.$$

Each such edge together with A forms a 2-absorber of E. Therefore, there are at least $m_{\frac{1}{4}}\binom{n+1}{k+1} > m\gamma n^{k+1}$ 2-absorbers for E.

By Claim 2.13, in either case there are at least $\gamma^5 n^{2(k+1)}$ 2-absorbers for E in \mathcal{T} .

3 Extremal Case - Proof of Theorem 1.6

Let $\mathcal{Q}, \mathcal{Q}'$ be two *r*-graph on $V(\mathcal{Q})$. We say $x \in V(\mathcal{Q})$ is α -good in \mathcal{Q} with respect to \mathcal{Q}' if $\deg_{\mathcal{Q}'\setminus\mathcal{Q}}(x) \leq \alpha\binom{|V(\mathcal{Q})|-1}{r-1}$. Moreover, we say \mathcal{Q} is α -good with respect to \mathcal{Q}' if every vertex is α -good.

The following proposition shows that subgraphs inherit the property "good".

Proposition 3.1 ([33]). Given reals $0 < \alpha' < 1$ and $0 \le c < 1$. Let $\alpha := \alpha'/c^{r-|S|}$. Suppose that S is α' -good in \mathcal{Q} with respect to \mathcal{Q}' . Let \mathcal{Q}'' be a subgraph of \mathcal{Q}' on $U \subset V(\mathcal{Q})$ such that $S \subset U$ and $|U| \ge cn$. Then S is α -good in $\mathcal{Q}[U]$ with respect to \mathcal{Q}'' .

Proposition 3.2. Given real $0 < \epsilon \leq 1$ and integer $1 \leq j \leq r-1$. Let $\epsilon' := \sqrt{r^r \epsilon}$. Suppose that Q is ϵ -close to Q'. Then the number of not ϵ' -good vertex is at most $\epsilon'|V(Q)|$.

Proof. Let *m* be the number of vertices that are not ϵ' -good. Since \mathcal{Q} is ϵ -close to \mathcal{Q}' , there are at most $\epsilon |V(\mathcal{Q})|^r$ edges in $\mathcal{Q}' \setminus \mathcal{Q}$. It follows that

$$m\epsilon' \binom{|V(\mathcal{Q})|-1}{r-1} \leq r|\mathcal{Q}' \setminus \mathcal{Q}| \leq r\epsilon|V(\mathcal{Q})|^r.$$

Then $m \leq r^r \epsilon |V(\mathcal{Q})|/\epsilon'$. By setting $\epsilon' = \sqrt{r^r \epsilon}$, we conclude that $m \leq \epsilon' |V(\mathcal{Q})|$.

By letting $\epsilon' = \sqrt{(k+1)^{k+1}\epsilon}$ and using Proposition 3.2, there are at most $\epsilon'(n+k/n)$ vertices in $X \cup V$ are not ϵ' -good with respect to \mathcal{T}_{ext} . Denote U by the family consists of all vertices not ϵ' -good. The following lemma allow us to find a matching \mathcal{M} such that covering all the vertices not ϵ' -good. Moreover, we will show later that $\mathcal{T} - V(\mathcal{M})$ contains a perfect matching.

Lemma 3.3. Given $k \ge 2$ and $\epsilon > 0$. There exists n_0 such that the following holds. Suppose that \mathcal{T} is a (1, k)-graph on $X \cup V$, and $|V| = k|X| = n \ge n_0$. If \mathcal{T} is ϵ -close to \mathcal{T}_{ext} , and $\delta_\ell(\mathcal{N}_\mathcal{T}(x)) > \delta(n, k, \ell)$ for each $x \in X$. Then there exists a matching \mathcal{M} in \mathcal{T} of size at most |U|+1 such that the following hold. (i) Let $\mathcal{T}' = \mathcal{T} - V(\mathcal{M})$, $\mathcal{T}'_{ext} = \mathcal{T} - V(\mathcal{M})$, then \mathcal{T}' is $\epsilon' 2^k$ -good with respect to \mathcal{T}'_{ext} , (ii) Recall that $\mathcal{T}_{ext} = \mathcal{T}(\mathcal{H}^i(A, B), \ldots, \mathcal{H}^i(A, B))$ for some $i \in \{0, 1\}$ and A, B. Let $A' = A \setminus V(\mathcal{M})$, $X' = X \setminus V(\mathcal{M})$, then

$$i|X'| \equiv |A'| \pmod{2}.\tag{3.1}$$

Proof. We claim that there exists $E \in \mathcal{T} \setminus \mathcal{T}_{ext}$ such that

$$|E \cap A| \equiv i + 1 + |E \cap U| \pmod{2}. \tag{3.2}$$

Move the vertices in $U \cap A$ and $U \cap B$ to the other part, we get a new partition $V = A_1 \cup B_1$ where $A_1 = (A \setminus U) \cup (B \cap U)$ and $B_1 = (B \setminus U) \cup (A \cap U)$. Fix any $x \in X \setminus U$, by the degree assumption $\delta_{\ell}(\mathcal{N}_{\mathcal{T}}(x)) > \delta(n, k, \ell) \geq \delta_{\ell}(\mathcal{H}^i(A_1, B_1))$, there exists $E_1 \in \binom{V}{k}$ such that $E_1 \in \mathcal{N}_{\mathcal{T}}(x) \setminus \mathcal{H}^i(A_1, B_1)$. Thus

$$|E_1 \cap A| - |E_1 \cap A \cap U| + |E_1 \cap B \cap U| = |E_1 \cap A_1| \equiv i + 1 \pmod{2}.$$

Therefore, $\{x\} \cup E_1 = E$ satisfies (3.2).

For $v_i \in U$, we greedily to find $E_i \in \mathcal{T}$ such that $E_i \cap U = \{v_i\}$ and $E \setminus U, E_1, \ldots, E_i$ are pairwise disjoint. This is possible since in the *i*-th step there are at most $(k+1)|U| \leq \epsilon'(k+1)n$ vertices used in $(E \setminus U) \cup (U \setminus \{v_i\}) \cup E_1 \cup \ldots \cup E_{i-1}$. Then for *n* sufficient large at least

$$\epsilon'(n+n/k)^k - \epsilon'(k+1)n\binom{|X \cup V| - 2}{k-1} \ge \frac{1}{2}\epsilon' n^k \ge 1$$

elements in $\mathcal{N}_{\mathcal{T}}(v_i) \setminus \mathcal{T}_{ext}$ do not intersect $(E \setminus U) \cup (U \setminus \{v_i\}) \cup E_1 \cup \ldots \cup E_{i-1}$. Let $\mathcal{M}' = \{E_1, \ldots, E_{|U|}\}$. Since $E_i \in \mathcal{N}_{\mathcal{T}}(v_i) \setminus \mathcal{T}_{ext}$ for $i \in |U|$, then

$$|E_i \cap A| \equiv i+1 \pmod{2}. \tag{3.3}$$

If

$$(n/k - |U|)i \equiv |A \setminus V(\mathcal{M}')| \pmod{2},\tag{3.4}$$

then by letting $\mathcal{M} = \mathcal{M}'$, the (ii) follows.

Now assume (3.4) does not hold. Let $\mathcal{M} = \mathcal{M}'' \cup \{E\}$ where $\mathcal{M}'' = \mathcal{M}' \setminus \{E_i : v_i \in E \cap U\}$. Therefore, by (3.2), we have

$$|A \setminus V(\mathcal{M})| \equiv |A \setminus V(\mathcal{M}'')| - (i+1+|E \cap U|) \pmod{2}.$$

Then by (3.3), it follows that

$$|A \setminus V(\mathcal{M}'')| \equiv |A \setminus V(\mathcal{M}')| + |E \cap U|(i+1) \pmod{2}.$$

Since (3.4) not holds, we obtain that

$$|A \setminus V(\mathcal{M})| \equiv |A \setminus V(\mathcal{M}')| + |E \cap U|(i+1) - (i+1+|E \cap U|) \pmod{2}$$

$$\equiv (n/k - |U|)i + 1 + (i+1+|E \cap U|) + |E \cap U|(i+1) \pmod{2}$$

$$\equiv (n/k - |U| + |E \cap U| - 1)i \pmod{2}.$$

Then, the (ii) follows.

Moreover, by Proposition 3.1 each v is $\epsilon' 2^k$ -good with respect to \mathcal{T}'_{ext} , the (i) follows.

By using Lemma 3.3, we are left with a "good" sub-hypergraph which contains no "bad" vertices.

In the following, we are going to divide \mathcal{T}' into two (k+1)-partite (k+1)-graph with large minimum vertex degree. Recall that $\mathcal{T}_{ext} = \mathcal{T}(\mathcal{H}^i(A, B), \ldots, \mathcal{H}^i(A, B))$ for some A, B, i when $||A| - |B|| \leq 1$ and $2 \nmid in/k + |A|$. Let \mathcal{M} be a matching in \mathcal{T} described in 3.3. Let $X' = X \setminus V(\mathcal{M}), A' = A \setminus V(\mathcal{M}), B' = B \setminus V(\mathcal{M})$, and $V' = A' \cup B' = V \setminus V(\mathcal{M}), n' = |V'|$. Note that $V(\mathcal{M}) \leq (|U| + 1)(k + 1) \leq 2\epsilon' kn$, then

$$n/2 - 2\epsilon' kn \le |A'|, |B'| \le n/2 + 2\epsilon' kn.$$

$$(3.5)$$

We are looking for a partition of $V' = (S_1 \cup \cdots \cup S_k) \cup (T_1 \cup \cdots \cup T_k) \cup E$ and a partition of $X' = Y_1 \cup Y_2 \cup Y_3$ satisfy some conditions. According to the parity of *i* and *k*, we distinguish four cases.

(1) If i = 0 and k is even.

Let |A'| = rk + s for some $0 \le s \le k - 1$. By Lemma 3.3,

$$|A'| = rk + s \equiv in_1/k = 0 \pmod{2}.$$
(3.6)

Since k is even, we have 2 | s. Let $Y_3 \cup E$ be an edge in \mathcal{T}' intersect A' exactly s elements. Such an edge exists since \mathcal{T}' is $\epsilon' 2^k$ -good with respect to \mathcal{T}'_{ext} .

Partition $X' = Y_1 \cup Y_2 \cup Y_3$, $A' \setminus E = T_1 \cup \dots \cup T_k$ and $B' \setminus E = S_1 \cup \dots \cup S_k$ such that $|Y_1| = |T_1| = \dots = |T_k| = r$, and $|Y_2| = |S_1| = \dots = |S_k| = \frac{n_1}{k} - r - 1$.

(2) If i = 0 and k is odd.

Let |A'| = r(k-1) + s for some $0 \le s \le k-2$. By Lemma 3.3,

$$|A'| = r(k-1) + s \equiv in_1/k = 0 \pmod{2}$$

Since k is odd, we have 2 | s. Let $Y_3 \cup E$ be an edge in \mathcal{T}' intersect A' exactly s elements. Such an edge exists since \mathcal{T}' is $\epsilon' 2^k$ -good with respect to \mathcal{T}'_{ext} .

Partition $X' = Y_1 \cup Y_2 \cup Y_3$, $A' \setminus E = T_1 \cup \cdots \cup T_{k-1}$ and $B' \setminus E = S_1 \cup \cdots \cup S_k \cup T_k$ such that $|Y_1| = |T_1| = \ldots = |T_k| = r$, and $|Y_2| = |S_1| = \ldots = |S_k| = \frac{n_1}{k} - r - 1$.

(3) If i = 1 and k is even.

Let $|A'| = n_1/k + r(k-2) + s$ for some $0 \le s \le k-3$. By Lemma 3.3,

$$|A'| = \frac{n_1}{k} + r(k-2) + s \equiv \frac{in_1}{k} = \frac{n_1}{k} \pmod{2}.$$

Since k is even, we have s is even.

Let $Y_3 \cup E$ be union of two edges in \mathcal{T}' one intersect A' exactly 1 element and the other intersect A' exactly s - 1 elements. Such edges exists since \mathcal{T}' is $\epsilon' 2^k$ -good with respect to \mathcal{T}'_{ext} .

Partition $X' = Y_1 \cup Y_2 \cup Y_3$, $A' \setminus E = T_1 \cup \dots \cup T_{k-1} \cup S_k$ and $B' \setminus E = S_1 \cup \dots \cup S_{k-1} \cup T_k$ such that $|Y_1| = |T_1| = \dots = |T_k| = r$, and $|Y_2| = |S_1| = \dots = |S_k| = \frac{n_1}{k} - r - 2$.

(4) If i = 1 and k is odd.

We can seen A' as B' and B' as A'. This case is same as case (2).

In both four cases, the complete (k + 1)-partite (k + 1)-graph on $(Y_1, T_1, T_2, \ldots, T_k)$ and $(Y_2, S_1, S_2, \ldots, S_k)$ are subgraphs of \mathcal{T}'_{ext} . Since (3.5) and $k \geq 3$, those vertex sets of size at least $n_1/3$. Then by Proposition 3.1, the induced subgraph of \mathcal{T}' on $(Y_1, T_1, T_2, \ldots, T_k)$ and $(Y_2, S_1, S_2, \ldots, S_k)$ are $\epsilon' 3^k$ -good with respect to the complete (k + 1)-partite (k + 1)-graph on the same vertex set.

In fact, for a single vertex the property of being "good" is equivalent to having a large degree. The next lemma due to author joint with Wang allow us to find a perfect matching in r-part r-graph with large degree.

Lemma 3.4 ([37]). For every integer $r \ge 2$, there exists $\alpha > 0$ and n_0 such that the following holds. Suppose that \mathcal{F} is a r-partite r-graph with each part of size $n \ge n_0$ vertices, and $\delta(\mathcal{F}) > (1-\alpha)n^{r-1}$. Then \mathcal{F} contains a perfect matching.

By using Lemma 3.4, we get a perfect matching in the two (k + 1)-partite (k + 1)-graphs, which together with $Y_3 \cup E$ forms a perfect matching in \mathcal{T}' , further imply a perfect matching in \mathcal{T} . The proof of Theorem 1.6 is completed.

References

- J. d. O. Bastos, G. O. Mota, M. Schacht, J. Schnitzer, and F. Schulenburg. Loose Hamiltonian cycles forced by large (k - 2)-degree—approximate version. SIAM Journal on Discrete Mathematics, 31(4):2328-2347, 2017.
- [2] J. d. O. Bastos, G. O. Mota, M. Schacht, J. Schnitzer, and F. Schulenburg. Loose Hamiltonian cycles forced by large (k-2)-degree-sharp version. Contributions to Discrete Mathematics, 61:101-106, 2017.
- [3] E. Buss, H. Hàn, and M. Schacht. Minimum vertex degree conditions for loose Hamilton cycles in 3-uniform hypergraphs. Journal of Combinatorial Theory, Series B, 103(6):658–678, 2013.
- [4] A. Czygrinow and T. Molla. Tight codegree condition for the existence of loose Hamilton cycles in 3-graphs. SIAM Journal on Discrete Mathematics, 28(1):67–76, 2014.
- [5] G. A. Dirac. Some theorems on abstract graphs. Proceedings of the London Mathematical Society, 3(1):69– 81, 1952.

- [6] P. Erdős, C. Ko, and R. Rado. Intersection theorems for systems of finite sets. The Quarterly Journal of Mathematics, 12(1):313–320, 1961.
- [7] P. Erdős. On extremal problems of graphs and generalized graphs. Israel Journal of Mathematics, 2(3):183– 190, 1964.
- [8] P. Frankl and A. Kupavskii. The Erdős Matching Conjecture and concentration inequalities. Journal of Combinatorial Theory, Series B, 157:366–400, 2022.
- [9] F. Garbe and R. Mycroft. Hamilton cycles in hypergraphs below the Dirac threshold. *Journal of Combi*natorial Theory, Series B, 133:153–210, 2018.
- [10] M. R. Garey and D. S. Johnson. Computers and intractability, volume 174. freeman San Francisco, 1979.
- [11] R. Glebov, Y. Person, and W. Weps. On extremal hypergraphs for Hamiltonian cycles. *European Journal of Combinatorics*, 33(4):544–555, 2012.
- [12] P. Hall. On Representatives of Subsets. Journal of the London Mathematical Society, 1(1):26-30, 1935.
- [13] H. Hàn, J. Han, and Y. Zhao. Minimum degree thresholds for Hamilton (k/2)-cycles in k-uniform hypergraphs. Journal of Combinatorial Theory, Series B, 153:105–148, 2022.
- [14] H. Hàn and M. Schacht. Dirac-type results for loose Hamilton cycles in uniform hypergraphs. Journal of Combinatorial Theory, Series B, 100(3):332–346, 2010.
- [15] J. Han and Y. Zhao. Minimum codegree threshold for Hamilton l-cycles in k-uniform hypergraphs. Journal of Combinatorial Theory, Series A, 132:194–223, 2015.
- [16] J. Han and Y. Zhao. Minimum vertex degree threshold for loose Hamilton cycles in 3-uniform hypergraphs. Journal of Combinatorial Theory, Series B, 114:70–96, 2015.
- [17] J. Han and Y. Zhao. Forbidding Hamilton cycles in uniform hypergraphs. Journal of Combinatorial Theory, Series A, 143:107–115, 2016.
- [18] S. Janson, T. Luczak, and A. Rucinski. Random Graphs. John Wiley & Sons, Inc., 2000.
- [19] G. Y. Katona and H. A. Kierstead. Hamiltonian chains in hypergraphs. Journal of Graph Theory, 30(3):205– 212, 1999.
- [20] P. Keevash, D. Kühn, R. Mycroft, and D. Osthus. Loose Hamilton cycles in hypergraphs. Discrete Mathematics, 311(7):544–559, 2011.
- [21] D. Kühn, R. Mycroft, and D. Osthus. Hamilton ℓ-cycles in uniform hypergraphs. Journal of Combinatorial Theory, Series A, 117(7):910–927, 2010.
- [22] D. Kühn and D. Osthus. Hamilton cycles in graphs and hypergraphs: an extremal perspective. 2014.
- [23] H. Lu, Y. Wang and X. Xing. Co-degree threshold for rainbow perfect matchings in uniform hypergraphs. arXiv:2111.00372
- [24] H. Lu, X. Yu and X. Yuan. Rainbow matchings for 3-uniform hypergraphs. Journal of Combinatorial Theory, Series A, 183: 105489, 2021.
- [25] K. Markström and A. Ruciński. Perfect matchings (and Hamilton cycles) in hypergraphs with large degrees. European Journal of Combinatorics, 32(5):677–687, 2011.
- [26] J. Moon and L. Moser. On Hamiltonian bipartite graphs. Israel Journal of Mathematics, 1(3):163–165, 1963.
- [27] C. Pohoata, L. Sauermann, and D. Zakharov. Sharp bounds for rainbow matchings in hypergraphs[J]. arXiv preprint arXiv:2212.07580, 2022.

- [28] C. Reiher, V. Rödl, A. Ruciński, M. Schacht, and E. Szemerédi. Minimum vertex degree condition for tight Hamiltonian cycles in 3-uniform hypergraphs. *Proceedings of the London Mathematical Society*, 119(2):409– 439, 2019.
- [29] V. Rödl and A. Ruciński. Dirac-type questions for hypergraphs a survey (or more problems for endre to solve). In *Bolyai Society Mathematical Studies*, pages 561–590. Springer Berlin Heidelberg, 2010.
- [30] V. Rödl, A. Ruciński, and E. Szemerédi. A Dirac-type theorem for 3-uniform hypergraphs. Combinatorics, Probability and Computing, 15(1-2):229, 2006.
- [31] V. Rödl, A. Ruciński, and E. Szemerédi. Dirac-type conditions for Hamiltonian paths and cycles in 3uniform hypergraphs. Advances in Mathematics, 227(3):1225–1299, 2011.
- [32] V. Rödl, E. Szemerédi, and A. Ruciński. An approximate Dirac-type theorem for k-uniform hypergraphs. Combinatorica, 28(2):229–260, 2008.
- [33] A. Treglown and Y. Zhao. Exact minimum degree thresholds for perfect matchings in uniform hypergraphs. Journal of Combinatorial Theory, Series A, 119(7):1500–1522, 2012.
- [34] A. Treglown and Y. Zhao. Exact minimum degree thresholds for perfect matchings in uniform hypergraphs II. Journal of Combinatorial Theory, Series A, 120(7):1463–1482, 2013.
- [35] W. T. Tutte. The factorization of linear graphs. Journal of the London Mathematical Society, 1(2):107–111, 1947.
- [36] J. Wang. A note on minimum degree condition for Hamilton (a, b)-cycles in hypergraphs. Discrete Mathematics, 346(1):113120, 2023.
- [37] J. Wang and J. You. Minimum degree thresholds for Hamilton $(\ell, k \ell)$ -cycles in k-uniform hypergraphs. arXiv:2302.04845
- [38] Y. Zhao. Recent advances on Dirac-type problems for hypergraphs. Recent trends in combinatorics, pages 145–165, 2016.