# WILF INEQUALITY IS PRESERVED UNDER GLUING OF SEMIGROUPS

SRISHTI SINGH AND HEMA SRINIVASAN

ABSTRACT. Wilf Conjecture on numerical semigroups is an inequality connecting the Frobenius number, embedding dimension and the genus of the semigroup. The conjecture is still open in general. We prove that the Wilf inequality is preserved under gluing of numerical semigroups. If the numerical semigroups minimally generated by  $A = \{a_1, \ldots, a_p\}$  and  $B = \{b_1, \ldots, b_q\}$  satisfy the Wilf inequality, then so does their gluing which is minimally generated by  $C = k_1 A \sqcup k_2 B$ . We discuss the extended Wilf's Conjecture in higher dimensions and prove an analogous result.

## 1. INTRODUCTION

A numerical semigroup  $\langle A \rangle$  is a submonoid of N minimally generated by  $A = (a_1, \ldots a_p)$  where  $gcd(a_1, \ldots, a_p) = 1$ . Such a semigroup will contain all but finitely many positive integers. The largest positive integer not in the numerical semigroup  $\langle A \rangle$  is called the Frobenius number  $F_A$ . We denote by  $[n_A]$  the set of  $n_A$  elements in  $\langle A \rangle$  that are less than  $F_A$ . Thus, the number of gaps is  $F_A - n_A$ . Moreover,  $n_A \geq 1$ as  $0 \in \langle A \rangle$  for every numerical semigroup A.

Wilf Conjecture, named after a question posed by H. Wilf in 1978, is an inequality connecting the three invariants of a numerical semigroup, the minimal number of generators or the embedding dimension, the Frobenius number, and the number of gaps. To be precise, the Wilf inequality for a numerical semigroup  $A$ , minimally generated by  $p$  elements, is  $pn_A \ge F_A + 1$ . When  $\langle A \rangle$  is symmetric,  $F_A + 1 = 2n_A$ , hence, Wilf Conjecture is true for all symmetric semigroups, i.e. for all Gorenstein semigroups or semigroup rings of type 1. Further, it is known from **[Fr]**, that for any numerical semigroup of type t,  $(t + 1)n_A \geq F_A + 1$ . Thus, Wilf Conjecture is true whenever the type is less than the embedding dimension. In particular, all numerical semigroups generated by arithmetic sequence must satisfy Wilf inequality as their type is less than the embedding dimension. However, it is possible for the type of a numerical semigroup to be any positive number when the embedding dimension is four or more. For an example, see  $\mathbf{Fr}$ . Although it has attracted much attention, Wilf Conjecture is still unsolved. A survey of known results are in [De]. The numerical semigroups which satisfy Wilf inequality are called Wilf Semigroups. In this note, we will prove that Wilf inequailty is preserved under gluing, an important construction for generating more numerical semigroups. That is, if  $\langle A \rangle$  and  $\langle B \rangle$  are two Wilf numerical semigroups, then so is any  $\langle C \rangle$  which is a gluing of  $\langle A \rangle$  and  $\langle B \rangle$ . We will prove this in the next section. Further, we will also discuss the extension to higher dimensions in section 3.

### 2. Gluing preserves Wilf

**Definition 2.1.** Let  $A = \{a_1, \ldots, a_p\}$  and  $B = \{b_1, \ldots, b_q\}$  minimally generate two numerical semigroups  $\langle A \rangle$  and  $\langle B \rangle$ . A numerical semigroup  $\langle C \rangle$  is said to be the gluing of  $\langle A \rangle$  and  $\langle B \rangle$  if the set of minimal generators of C is a disjoint union  $k_1(a_1, \ldots, a_p) \sqcup k_2(b_1, \ldots, b_q)$  where  $k_1 \in \langle B \rangle$  but not in B, and  $k_2 \in \langle A \rangle$  but not in A.

Since  $\langle C \rangle$  is a numerical semigroup,  $k_1$  and  $k_2$  must be relatively prime. In the context of numerical semigroups, this type of decomposition was first introduced by Delorme  $(|\mathbf{De}|)$  to characterize complete intersections. He proved that gluing of any two complete intersections results in a complete intersection and more impressively, all complete intersections of embedding dimension  $\geq$  3 must be the gluing of two other complete intersections. It was further developed by Rosales as a gluing of numerical semigroups and extensively studied in [ros], [GS1], and [GS2]. Many invariants of  $\langle C \rangle$  can be computed from the invariants of  $\langle A \rangle$  and  $\langle B \rangle$ . These formulae for the invariants can be derived from the minimal graded resolution of the semigroup ring  $k[C]$  ([GS1]). We write  $\langle C \rangle = \langle A \rangle \Join \langle B \rangle$  to denote that C is a gluing of A and B. The following formula for  $F_C$  can be derived from the regularity formula of the graded Betti numbers of  $k[A]$  and  $k[B]$  as in [GS1] but here we give a direct and simple direct proof, for the sake of completeness.

<span id="page-1-0"></span>**Theorem 2.2.** Let  $A = \{a_1, \ldots, a_p\}$  and  $B = \{b_1, \ldots, b_q\}$  minimally generate two numerical semigroups  $\langle A \rangle$  and  $\langle B \rangle$  and let  $C = k_1A \sqcup k_2B$ so that  $\langle C \rangle = \langle A \rangle \Join \langle B \rangle$ . Then

$$
F_C = k_1 F_A + k_2 F_B + k_1 k_2
$$

*Proof.* First of all, since  $k_1, k_2$  are relatively prime, for any positive integer  $t \geq 1$ , we can find positive integers m and n, with  $0 < n < k_1$ such that  $mk_1 - nk_2 = t$ . Then  $k_1F_A + k_2F_B + k_1k_2 + t = k_1(F_A + m) +$  $k_2(F_B - n+k_1) \in \langle C \rangle$ . We will now show that  $k_1F_A + k_2F_B + k_1k_2 \notin \langle C \rangle$ .

Suppose  $k_1 F_A + k_2 F_B + k_1 k_2 = k_1 \sum_{i=1}^p r_i a_i + k_2 \sum_{j=1}^q s_j b_j$  for some  $r_i, s_j \in \mathbb{N}, i \in \{1, ..., p\}, j \in \{1, ..., q\}.$  Then we have  $k_1(F_A - \sum_{i=1}^p r_i a_i) =$  $k_2(\sum_{j=1}^q s_j b_j - k_1 - F_B).$ 

Since  $k_1$  and  $k_2$  are relatively prime and  $F_A \notin \langle A \rangle$ , we must have  $F_A - \sum_{i=1}^{p} r_i a_i = k_2 t$  for some non zero integer t. But then  $F_A =$  $\sum_i^p$  $\sum_{i=1}^{i=1}$  $r_{i=1}^{p} r_i a_i + t k_2 \in \langle A \rangle$  if  $t > 0$ . So,  $t < 0$ . Hence,  $F_B = -t k_1 - k_1 +$ <br> $r_{j=1}^{q} s_j b_j = \sum_{j=1}^{q} s_j b_j + (-1-t) k_1 \in \langle B \rangle$ , which is not possible either. So,  $k_1F_A + k_2F_B + k_1k_2 \notin \langle C \rangle$ . Thus,  $F_C = k_1F_A + k_2F_B + k_1k_2$ .

The next theorem states that if two numerical semigroups A and B satisfy Wilf Conjecture, so does every gluing of A and B.

<span id="page-2-0"></span>**Theorem 2.3.** Suppose  $\langle A \rangle$  and  $\langle B \rangle$  satisfy Wilf inequality. Then so does  $\langle C \rangle = \langle A \rangle \Join \langle B \rangle$ 

*Proof.* Suppose  $A = \{a_1, \ldots, a_p\}$  and  $B = \{b_1, \ldots, b_q\}$  minimally generate two numerical semigroups  $\langle A \rangle$  and  $\langle B \rangle$ , and let  $C = k_1 A \sqcup k_2 B$ so that  $\langle C \rangle = \langle A \rangle \bowtie \langle B \rangle$ .

Thus,  $k_2 \in \langle A \rangle$ , not in A, and  $k_1 \in \langle B \rangle$  and not in B, and  $(k_1, k_2) =$ 1. By [\(2.2\)](#page-1-0),  $F_C = k_1 F_A + k_2 F_B + k_1 k_2$ . Without loss of generality, we may take  $p \leq q$ .

To begin with, we have by a theorem of Froberg et al  $(\mathbf{Fr})$  which states  $(t + 1)n<sub>C</sub> \ge F<sub>C</sub> + 1$  for all numerical semigroups of type t. By [GS1], the type of  $\langle C \rangle$  is the product of types of  $\langle A \rangle$  and  $\langle B \rangle$ . When the embedding dimension is 1 or 2, the type is 1 as it is a complete intersection and when the embedding dimension is 2, the type is either 1 or 2 ([He]). Thus, if  $p \leq 2$  and  $q \leq 2$ , then C is also a complete intersection and  $F_C < 2n_C \le (p+q)n_C$  as required.

If  $p \leq q = 3$ , then again since  $t \leq 2$ , we get that

$$
F_C < (t+1)n_C \le 3n_C \le (p+q)n_C.
$$

Thus, we just need to consider the case when  $q \geq 4$ .

Now, for all  $x \in [nA], 1 \le i \le k_1, k_1x + k_2(F_B + i) \in [n_C]$ . Further,

$$
k_1x + k_2(F_B + i) = k_1x' + k_2(F_B + j) \implies k_1(x - x') = k_2(j - i).
$$

But  $k_1, k_2$  are relatively prime, and  $j - i \leq k_1$ . So  $j = i, x = x'$ . Hence, all these aforementioned elements are distinct in  $[n_C]$ .

Similarly, for  $1 \leq i \leq |k_2/2|, 1 \leq j \leq |k_1/2|$ , the elements  $k_1(F_A +$  $i)+k_2(F_B + j) \in [n_C]$ , and all these  $\lfloor k_2/2 \rfloor \lfloor k_1/2 \rfloor$  elements are distinct. So,

$$
n_C \ge k_1 n_A + \lfloor k_2/2 \rfloor \lfloor k_1/2 \rfloor \ge k_1 n_A + (k_1 - 1)(k_2 - 1)/4
$$

Similarly,

$$
n_C \ge k_2 n_B + \lfloor k_2/2 \rfloor \lfloor k_1/2 \rfloor \ge k_2 n_B + (k_1 - 1)(k_2 - 1)/4
$$

Now suppose  $q \ge p \ge 2$ , so that  $F_A$  and  $F_B$  are both positive integers. Hence,

$$
F_C = k_1 F_A + k_2 (F_B + k_1) = k_1 (F_A + 1) + k_2 (F_B + 1) + k_1 k_2 - k_1 - k_2
$$
  
=  $k_1 (F_A + 1) + k_2 (F_B + 1) + (k_1 - 1)(k_2 - 1) - 1$ 

Since A and B satisfy the Wilf inequality,  $F_A + 1 \leq pn_A$  and  $F_B + 1 \leq$  $qn_B$ . It now follows that

$$
F_C = k_1(F_A + 1) + k_2(F_B + 1) + (k_1 - 1)(k_2 - 1) - 1
$$
  
\$\leq k\_1 p n\_A + k\_2 q n\_B + (k\_1 - 1)(k\_2 - 1) - 1\$

Next,

$$
\frac{F_C}{n_C} = \frac{k_1(F_A + 1)}{n_C} + \frac{k_2(F_B + 1) + (k_1 - 1)(k_2 - 1) - 1}{n_C}
$$

Recall that  $n_C \ge k_1 n_A$ , and  $n_C \ge k_2 n_B + (k_1 - 1)(k_2 - 1)/4$ . Hence,

$$
\frac{F_C}{n_C} \le \frac{k_1 p n_A}{k_1 n_A} + \frac{k_2 q n_B + (k_1 - 1)(k_2 - 1) - 1}{k_2 n_B + (k_1 - 1)(k_2 - 1)/4}
$$

But

$$
\frac{k_2qn_B + (k_1 - 1)(k_2 - 1) - 1}{k_2n_B + \frac{(k_1 - 1)(k_2 - 1)}{4}} \le q \frac{k_2n_B + \frac{(k_1 - 1)(k_2 - 1) - 1}{q}}{k_2n_B + \frac{(k_1 - 1)(k_2 - 1)}{4}} < q
$$

since  $q \geq 4$ .

Thus,  $F_C < (p+q)n_C$ , as long as  $p \geq 2$ . It remains to show the inequality if  $p = 1$  and  $q \ge 4$ . In that case,

$$
(q+1)n_C - F_C \ge (q+1)(k_2n_B + (k_1 - 1)(k_2 - 1)/4) - k_2F_B - k_1(k_2 - 1)
$$
  
=  $k_2(qn_B - F_B - 1) + (k_1k_2 - k_2 - k_1 + 1)\left(\frac{q+1}{4} - 1\right) + k_2n_B - 1$   
> 0,

since  $qn_B \geq F_B + 1$ ,  $\frac{q+1}{4} \geq 1$  and  $k_2 n_B - 1 \geq 2 - 1 > 0$ . So, we have that Wilf Conjecture holds for  $\langle C \rangle$  as well.

# 3. Gluing of Affine Semigroups

We now want to explore the same question in higher dimension, namely for subsemigroups of  $\mathbb{N}^n$  as opposed to in  $\mathbb{N}$ . A finitely generated submonoid of  $\mathbb{N}^n$  is also called an *affine semigroup*. Thus, if A is an  $n \times p$  matrix in N with columns  $a_i, 1 \leq i \leq n$ , representing elements of  $\mathbb{N}^n$ , the monoid generated by the columns of A in  $\mathbb{N}^n$  is an affine semigroup  $\langle A \rangle$ . As before without loss of generaliity, we may assume that the entries in A are relatively prime as  $\langle A \rangle$  and  $\langle dA \rangle$  are

isomorphic as monoids. The question then becomes, given two numerical semigroups  $\langle A \rangle$  and  $\langle B \rangle$  in  $\mathbb{N}^n$ ,  $n \geq 2$ , is there a generalized notion of this notable conjecture by Wilf, and, if so, does a gluing of  $\langle A \rangle$  and  $\langle B \rangle$  satisfy this notion if  $\langle A \rangle$  and  $\langle B \rangle$  do?

The concept of gluing of numerical semigroups in higher dimensions has been elucidated in [GS2] and [GS22]. Consider finite subsets  $A =$  ${a_1, ..., a_p}, B = {b_1, ..., b_q}$  and  $C = k_1 A \sqcup k_2 B$  in  $\mathbb{N}^n$  where  $n \geq 2$ , and  $k_1$  and  $k_2$  are positive integers. Let  $\langle A \rangle$ ,  $\langle B \rangle$ ,  $\langle C \rangle$  be the semigroups generated by the aforementioned sets, and  $k[A] \cong k[\mathbf{x}_1, ..., \mathbf{x}_p]/I_A, k[B] \cong$  $k[\mathbf{y}_1, ..., \mathbf{y}_q]/I_B$ , and  $k[C] \cong k[\mathbf{x}_1, ..., \mathbf{x}_p, \mathbf{y}_1, ..., \mathbf{y}_q]/I_C$  be their respective semigroup rings. Set  $R := k[\mathbf{x}_1, ..., \mathbf{x}_p, \mathbf{y}_1, ..., \mathbf{y}_q].$ 

**Definition 3.1.**  $\langle C \rangle$  is said to be a gluing of  $\langle A \rangle$  and  $\langle B \rangle$ , written  $\langle C \rangle = \langle A \rangle \otimes \langle B \rangle$ , if  $I_C = I_A R + I_B R + \langle \rho \rangle$  with  $\rho = \mathbf{x}^{\mathbf{c}} - \mathbf{y}^{\mathbf{d}}$  for some  $\mathbf{c} \in \mathbb{N}^{\mathbf{p}}$  and  $\mathbf{d} \in \mathbb{N}^{\mathbf{q}}$ .

Although any two numerical semigroups can be glued, it is not true in hgher dimensions. An affine semigroup  $\langle A \rangle \in \mathbb{N}^n$  is said to be nondegenerate if they A spans  $\mathbb{Q}^n$ . Two non-degenerate semigroups in dimension  $\geq 2$  cannot be glued. Precise conditions when such gluings exist can be found in [GS22].

To extend the Wilf inequality to higher dimensions, it is necessary to have some finiteness to the smeigroup, so as to have an analogue of Frobenius number and the number of gaps. There is a notion of Generalised Numerical Semigroup, namely, if  $H(S) := \mathbb{N}^n \backslash S$  is finite, we call S a Generalised Numerical Semigroup (GNS). Further, if  $\mathcal{H}(S) := (\text{cone}(C) \backslash C) \cap \mathbb{N}^n$  is finite, where  $\text{cone}(C)$  is the cone of S in  $\mathbb{Q}_{\geq 0}^n$ , then we call S a C semigroup. Thus, A is a C semigroup if and only if there are only finitely many elements in  $\mathbb{N}^n$  such that no integral multiple of them are in  $\langle A \rangle$ .

Suppose S is a C semigroup (or GNS). Let  $\prec$  be a monomial ordering satisfying the following condition:

<span id="page-4-0"></span>(1) 
$$
\mathbf{a} \in \mathbb{N}^n \implies |\{\mathbf{b} \in \mathbb{N}^n \mid \mathbf{b} \prec \mathbf{a}\}| < \infty
$$

Let  $F(S)$ , called the Frobenius Element of S, denote the maximal element of the set  $\mathcal{H}(S)$  (respectively,  $H(S)$ ) with respect to  $\prec$ . Set  $\mathcal{N}(S) = |\mathcal{H}(S)| + |\{\mathbf{x} \in S \mid \mathbf{x} \prec F(S)\}|$  (respectively,  $\mathcal{N}(S) =$  $|H(S)| + |\{x \in S \mid x \prec F(S)\}|$ . This exists due to the restriction applied on ≺. The Extended Wilf Conjecture can be formulated in [wilfext] as follows

$$
|\{\mathbf{x} \in S \mid \mathbf{x} \prec F(S)\}| \cdot e(S) \ge \mathcal{N}(S) + 1
$$

First note that if  $\langle A \rangle$  and  $\langle B \rangle$  are generalized numerical semigroups, and  $\langle C \rangle = k_1 \langle A \rangle \otimes k_2 B$ , then  $\langle C \rangle$  cannot be a GNS. To see this, we will first show that if  $S = {\mathbf{s}_1, ..., \mathbf{s}_t}$  is a GNS in  $\mathbb{N}^n$ , then rank  $S = n$ , where S is the  $n \times t$  matrix that is, a GNS must be non-degenerate.  $[\mathbf{s}_1 \cdots \mathbf{s}_t]$ . For any  $\mathbf{x} \in \mathbb{N}^n$ , if the rank of S is strictly less than *n*, then one of the rows  $R_i, 1 \leq i \leq n$  of the matrix S, say the *nth*, can be written as  $\sum_{i=1}^{n-1} r_i R_i$  for some  $r_i \in \mathbb{Q}$ . Then any element in  $(t_1, \ldots, t_n)^T \in \mathbb{N}^n$  must satisfy  $t_n = \sum_{i=1}^{n-1} r_i t_i$ . Thus, there are infinitely many elements in  $\mathbb{N}^n$  that will be missing from §. This implies that  $\mathbb{N}^n\setminus\langle A\rangle$  is not finite, contradicting the fact that  $\langle A\rangle$  is a GNS.

It now follows from [gimenez2022gluing] Corollary 1.6, which states that if  $\langle A \rangle$  and  $\langle B \rangle$  can be glued, then rank $A$  + rank $B = n + 1$ , where A is the  $n \times p$  matrix  $[\mathbf{a}_1 \cdots \mathbf{a}_p]$ , and B is the  $n \times q$  matrix  $[\mathbf{b}_1 \cdots \mathbf{b}_q]$ , that two generalized numerical semigroups  $\langle A \rangle$  and  $\langle B \rangle$  cannot be glued in  $\mathbb{N}^n$  to obtain another GNS if  $n \geq 2$ .

Thus to generalize the theorem [2.3](#page-2-0) we will consider gluings of  $\mathcal C$ semigroups. First, let us glue a non degenerate  $\mathcal C$  semigroup  $A$  to a completely degenerate C semigroup B as explained in  $[GS2]$ . We will describe a gluing of a semigroup  $\langle A \rangle$  and  $\langle \mathbf{b} \rangle$  for some  $\mathbf{b} \in \mathbb{N}^n$ , and show that if  $\langle A \rangle$  is a C semigroup that satisfies the Extended Wilf Conjecture then so is the gluing. In fact, we can generalize this to gluing a non degenerate  $\alpha$ , (thus  $\langle A \rangle$  may even be GNS) with a very degenerate  $\langle B \rangle$ .

<span id="page-5-0"></span>**Theorem 3.2.** ([GS2] Proposition 4.1) Let  $A = {\bf{a_1}, ..., a_p} \subseteq \mathbb{N}^n$  be a non-degenerate affine semigroup. Suppose that  $\mathbf{b} = (b_1, ..., b_n) \in \langle A \rangle$ and  $B = \mathbf{b} \begin{bmatrix} u_1 & \cdots & u_q \end{bmatrix}$  for some  $u_1, ..., u_q \in \mathbb{N}$ . Set  $C := k_1 A \cup k_2 B$ , for positive integers  $k_1, k_2$  such that  $k_1 \in \langle u_1, ..., u_q \rangle$ . Then C is a gluing of  $\langle A \rangle$  and  $\langle B \rangle$ . Without loss of generality, we may assume that  $\gcd(b_1, ..., b_n) = 1$ ,  $\gcd(u_1, ..., u_q) = 1$ , and  $\gcd(a_{ij}, 1 \leq i \leq n, 1 \leq j \leq q)$  $p$ ) = 1. If this does not occur, we can factor out the gcd in A or B and simplify to get another semigroup with the same semigroup ring.

**Theorem 3.3.** If  $\langle A \rangle$  is a C semigroup satisfying the Extended Wilf Conjecture then so is the gluing  $\langle C \rangle$  as obtained above [3.2](#page-5-0)

*Proof.* Suppose  $\langle A \rangle$  is a C semigroup so that  $\mathcal{H}(A)$  is finite. Let  $\mathbf{x} \in$  $\mathcal{H}(C)$ . So  $\mathbf{x} \in \mathbb{N}^n$ , and  $\mathbf{x} \in \text{cone}(C) \backslash C$ . Write  $\mathbf{x} = y_1 k_1 \mathbf{a_1} + ...$  $y_p k_1 a_p + z_1 k_2 u_1 b + \ldots + z_q k_2 u_q b$ , for some  $y_i, z_j \in \mathbb{Q}_{\geq 0}, 1 \leq i \leq p, 1 \leq j$  $j \leq q$ . Since  $\mathbf{x} \notin \langle C \rangle$ , at least one of the  $y_i$  or  $z_j$  belongs to Q\N. But  $b \in \langle A \rangle$ , so  $\mathbf{x} \in \mathcal{H}(A)$ . Thus,  $\mathcal{H}(C) \subseteq \mathcal{H}(A)$  is finite. So  $\langle C \rangle$  is a C semigroup.

We can now conclude that  $F(A)$  and  $F(C)$  exist. Moreover, the sets  $\{x \in \langle A \rangle \mid x \prec F(A)\},\$  and  $\{y \in \langle C \rangle \mid y \prec F(C)\}\$ , are both finite due to the condition [\(1\)](#page-4-0) on  $\prec$ .

Observe that since  $\langle A \rangle$  satisfies the Extended Wilf Conjecture, it satisfies the equivalent inequality

$$
\frac{|\mathcal{H}(A)|}{|\{\mathbf{x} \in \langle A \rangle \mid \mathbf{x} \prec F(A)\}|} < e(A) - 1
$$

First note that since  $\langle C \rangle$  is a gluing of  $\langle A \rangle$  and  $\langle B \rangle$ , we have that  $e(C) = e(A) + e(B) = p + q$ . We will show that  $\langle C \rangle$  satisfies

$$
\frac{|\mathcal{H}(C)|}{|\{\mathbf{x} \in \langle C \rangle \mid \mathbf{x} \prec F(C)\}|} < e(C) - 1
$$

We have,  $\mathcal{H}(C) \subset \mathcal{H}(A)$  and hence  $|\mathcal{H}(C)| \leq |\mathcal{H}(A)|$ . Next, we claim that  $k_1 F(A) \notin \langle C \rangle$ . Suppose otherwise. So

$$
k_1F(A) = k_1\mathbf{x} + k_2(z_1u_1 + \dots + z_qu_q)\mathbf{b}
$$

for some  $z_1, ..., z_q \in \mathbb{N}, \mathbf{x} \in \langle A \rangle$ . This implies that  $k_1$  must divide  $k_2(z_1u_1+\ldots+z_qu_q)b_i$  for all  $i\in[n]$ . Let  $gcd(k_1,b_i)=d_i$  for  $i\in[n]$ . Then  $k_1 = k_i d_i$  for some  $k_i \in \mathbb{N}$ , and  $b_i = b'_i d_i$  for some  $b'_i \in \mathbb{N}$ ,  $1 \le i \le n$ . Next, since  $gcd(k_i, b_i) = 1$ ,  $k_i$  must divide  $k_2(z_1u_1 + ... + z_qu_q)$ . So  $k_i l_i = k_2(z_1 u_1 + \dots + z_q u_q)$  for some  $l_i \in \mathbb{N}, 1 \le i \le n$ . On multiplying both sides of the equation by  $d_i$ , we obtain

<span id="page-6-0"></span>(2) 
$$
k_1 l_i = k_2 (z_1 u_1 + \dots + z_q u_q) d_i
$$

But  $d_i | b_i$  for all  $i \in [n]$  and  $gcd(b_1, ..., b_n) = 1$ , so  $gcd(d_1, ..., d_n) = 1$ . There exist  $m_1, ..., m_n \in \mathbb{Z}$  such that  $m_1d_1 + ... + m_nd_n = 1$ . For each  $i \in [n]$ , multiply [\(2\)](#page-6-0) by  $m_i$  and add all n equations to get

$$
k_1 \sum_{i=1}^n m_i l_i = k_2 (z_1 u_1 + \dots + z_q u_q) \sum_{i=1}^n m_i d_i = k_2 (z_1 u_1 + \dots + z_q u_q) > 0
$$

so that  $k_1 l = k_2(z_1 u_1 + ... + z_q u_q) \in \mathbb{N}$  for  $l = \sum_{i=1}^n m_i l_i \in \mathbb{N}$ . But this implies that  $F(A) = \mathbf{x} + l\mathbf{b} \in \langle A \rangle$ , a contradiction. So  $k_1F(A) \notin$  $\langle C \rangle$ . Moreover,  $F(A) \in \mathcal{H}(A)$ , so  $F(A) \in \text{cone}(A)$  which implies  $k_1F(A) \in \text{cone}(k_1A) \subseteq \text{cone}(C)$ . Thus,  $k_1F(A) \in \mathcal{H}(C)$ . It follows that  $F(C) \geq k_1 F(A)$ . Next, observe that for all  $\mathbf{x} \in \langle A \rangle$  and  $\mathbf{x} \prec F(A)$ , we have  $k_1\mathbf{x} \in \langle C \rangle$ , and  $k_1\mathbf{x} \prec k_1F(A) \preccurlyeq F(C)$ . Note also that each of these elements is distinct in  $\langle C \rangle$  since  $k_1\mathbf{x} = k_1\mathbf{y}$  for some  $\mathbf{x}, \mathbf{y} \in \langle A \rangle$ implies  $\mathbf{x} = \mathbf{y}$ . Thus, for every element  $\mathbf{x} \in \langle A \rangle$  such that  $\mathbf{x} \prec F(A)$ , we obtain an element  $\mathbf{z} \in \langle C \rangle$  such that  $\mathbf{z} \prec F(C)$ . It follows that

$$
|\{\mathbf{x} \in \langle C \rangle \mid \mathbf{x} \prec F(C)\}| \ge |\{\mathbf{x} \in \langle A \rangle \mid \mathbf{x} \prec F(A)\}|
$$

Thus,

$$
\frac{|\mathcal{H}(C)|}{|\{\mathbf{x} \in \langle C \rangle \mid \mathbf{x} \prec F(C)\}|} \le \frac{|\mathcal{H}(A)|}{|\{\mathbf{x} \in \langle A \rangle \mid \mathbf{x} \prec F(A)\}|} < e(A) - 1 < p + q - 1
$$
\nWe have shown that  $\langle C \rangle$  satisfies the Extended Wilf Conjecture.

An interesting observation is that although Wilf inequality in N is seemingly sharp, it is not the case for the extended inequality in  $\mathbb{N}^n$ as seen in the proof above. The inequality is in fact easier to obtain in this special higher dimensional gluing. A stronger inequality may work in higher dimensions. It is also possible to have a gluing of  $\langle A \rangle$ and  $\langle B \rangle$  when both are necessarily degenerate under some conditions. This leads us to conjecture that if  $C = A \Join B$ , with  $\langle A \rangle$ ,  $\langle B \rangle$  and  $\langle C \rangle$ all C semigroups, then if  $\langle A \rangle$  and  $\langle B \rangle$  are Wilf, then so is  $\langle C \rangle$ .

## Declarations

- *Ethical Approval:* Not applicable.
- Competing interests: There are no competing interests, of either financial or personal nature.
- Availability of data and materials: Data sharing not applicable to this article as no data-sets were generated or analyzed during the current study.

### **REFERENCES**

- [1] P. Gimenez and H. Srinivasan, The structure of the minimal free resolution of semigroup rings obtained by gluing, J. Pure Appl. Alg.  $223$  (2019), 1411–1426.
- [2] P. Gimenez and H. Srinivasan, Gluing semigroups: when and how. Semigroup Forum 101 (2020). No.3, 603-618.
- [3] P. Gimenez and H. Srinivasan, On gluing semigroups in  $N^n$  and the consequences, Res. in Math. Sci. 9 (2022). No.2 1-14.
- [4] C. Delorme, Sous-monoïdes d'intersection complète de N, Ann. Sci. École Norm. Sup. (4) 9 (1976), 145–154.
- [5] M. Delgado, Conjecture of Wilf A survey, arXiv:1902.03461 (2019)
- [6] R. Fröberg, C. Gottlieb and R. Häggkvist On numerical semigroup, Semigroup Forum
- [7] J. Herzog, Generators and relations of abelian semigroups and semigroup rings, Manuscr. Math. 3 (1970), 175–193.
- [8] J. C. Rosales, On presentations of subsemigroups of  $\mathbb{N}^n$ , Semigroup Forum 55 (1997), 152–159.

University of Missouri, Columbia, MO 65211, USA