On Permutation Trinomials of the type $X^{q^2-q+1} + AX^{q^2} + BX$ over \mathbb{F}_{q^3}

Daniele Bartoli^{*}, and Francesco Ghiandoni[†]

June 19, 2023

Abstract

Necessary and sufficient conditions on $A, B \in \mathbb{F}_{q^3}^*$ for $f(X) = X^{q^2-q+1} + AX^{q^2} + BX$ being a permutation polynomial of \mathbb{F}_{q^3} are investigated via a connection with algebraic varieties over finite fields.

MSC: 05A05 11T06 11T55

Keywords: Permutation polynomials, algebraic varieties, finite fields, permutation trinomials

1 Introduction

Let q be a prime power and denote by \mathbb{F}_q the finite field with q elements. A polynomial $f(x) \in \mathbb{F}_q[x]$ is a *permutation polynomial* (PP) of \mathbb{F}_q if it is a bijection of \mathbb{F}_q into itself. Permutation polynomials were first studied by Hermite and Dickson; see [13,20].

In general, simple structures or additional extraordinary properties are required by applications of PPs in other areas of mathematics and engineering, such as cryptography, coding theory, or combinatorial designs. In this case, permutation polynomials meeting these criteria are usually difficult to find. For a deeper introduction to the connections of PPs with other fields of mathematics we refer to [9,24,39] and the references therein.

Permutation polynomials of monomial or binomial type have been widely investigated in the last decades. Although much less is known for polynomials having more than 2 terms, specific families of trinomials and quadrinomials were considered; see for instance [1-8, 12, 14-16, 18-38, 40-58], where in most of the cases q is a square.

When investigating the permutational properties of a polynomial, a fruitful connection with algebraic curves is provided by the following observation (we refer to [17] for a comprehensive introduction on algebraic curves). Given a polynomial $f(x) \in \mathbb{F}_q[x]$, let us consider the curve \mathcal{C}_f with affine equation

$$C_f : \frac{f(X) - f(Y)}{X - Y} = 0.$$
(1)

^{*}Dipartimento di Matematica e Informatica, Università degli Studi di Perugia, Perugia, Italy. daniele.bartoli@unipg.it

[†]Dipartimento di Matematica e Informatica "Ulisse Dini", Università degli studi di Firenze, Firenze, Italy, francesco.ghiandoni@unifi.it

A standard approach to the problem of deciding whether f(x) is a PP is the investigation of the set of \mathbb{F}_q -rational points of \mathcal{C}_f . In fact, if $a \neq b$ are two distinct elements of \mathbb{F}_q such that f(a) = f(b), then the \mathbb{F}_q -rational point (a, b) belongs to \mathcal{C}_f and does not lie on the line X - Y = 0. On the other hand, if (a, b) is an \mathbb{F}_q -rational point of \mathcal{C}_f not lying on X - Y = 0, then f(a) = f(b) and so f(x) is not a PP.

In general to decide whether an algebraic curve C_f defined over \mathbb{F}_q possesses suitable \mathbb{F}_q rational points is a hard task. A standard way to bypass such a problem is to prove the existence of absolutely irreducible components defined over \mathbb{F}_q of C and then apply to such a component the celebrated Hasse-Weil Theorem (or refinements of it); see [28].

Unfortunately, this machinery fails whenever the degree of C is too large with respect to q, namely larger than roughly $\sqrt[4]{q}$.

To overcome such a problem, for specific families of polynomials of large degree, one can investigate higher dimensional varieties attached to the starting polynomial f.

In this paper we consider polynomials of the form

$$f_{A,B}(X) = X^{q^2 - q + 1} + AX^{q^2} + BX \in \mathbb{F}_{q^3}[X],$$
(2)

where A, B are nonzero elements of \mathbb{F}_{q^3} and q is even. Such a family has been partially investigated in [40], where the authors proved the following result.

Theorem 1.1. If $A^{q^2+q+1} = B^{1+q+q^2}$ and $A^q B \in \mathbb{F}_q \setminus \{0,1\}$, then $f_{A,B}(X) = X^{q^2-q+1} + AX^{q^2} + BX$ is a *PP* over \mathbb{F}_{q^3} .

Exploiting a connection with algebraic surfaces (see Section 3) and generalizations of Hasse-Weil Theorem to higher dimensions, we are able to prove that, for q large enough and apart from a few exceptions, the conditions on parameters A, B expressed in Theorem 1.1 are also sufficient.

Main Theorem. Let $q = 2^m$, $m \ge 19$. Then $f_{A,B} = X^{q^2-q+1} + AX^{q^2} + BX \in \mathbb{F}_{q^3}[X]$ is a PP over $\mathbb{F}_{q^3}[X]$ if and only if one of the following conditions holds:

- $A^{1+q+q^2} = B^{1+q+q^2}$ and $A^q B \in \mathbb{F}_q \setminus \{0,1\};$
- $A^q B = 1$ and $B^{1+q+q^2} \neq 1$.

2 Preliminaries

Let $q = 2^m$, $m \in \mathbb{N}$, and denote by \mathbb{F}_q the finite field with q elements. As a notation, $\mathbb{P}^r(\mathbb{F}_q)$ and $\mathbb{A}^r(\mathbb{F}_q)$ (or \mathbb{F}_q^r) denote the projective and the affine space of dimension $r \in \mathbb{N}$ over the finite field \mathbb{F}_q respectively. A variety, and more specifically a curve or a surface (i.e. a variety of dimension 1 or 2 respectively), is described by a certain set of equations with coefficients in \mathbb{F}_q . We say that a variety \mathcal{V} is *absolutely irreducible* if there are no varieties \mathcal{V}' and \mathcal{V}'' defined over the algebraic closure of \mathbb{F}_q (denoted by $\overline{\mathbb{F}_q}$) and different from \mathcal{V} such that $\mathcal{V} = \mathcal{V}' \cup \mathcal{V}''$. If a variety $\mathcal{V} \subseteq \mathbb{P}^r(\mathbb{F}_q)$ is defined by $F_i(X_0, \ldots, X_r) = 0$, for $i = 1, \ldots, s$, an \mathbb{F}_q -rational point of \mathcal{V} is a point $(x_0 : \cdots : x_r) \in \mathbb{P}^r(\mathbb{F}_q)$ such that $F_i(x_0, \ldots, x_r) = 0$, for $i = 1, \ldots, s$. The set of the \mathbb{F}_q -rational points of \mathcal{V} is usually denoted by $\mathcal{V}(\mathbb{F}_q)$. If $s = 1, \mathcal{V}$ is called a hypersurface and it is absolutely irreducible if the corresponding polynomial $F(X_1, \ldots, X_r)$ is absolutely irreducible, i.e. it possesses no non-trivial factors over $\overline{\mathbb{F}_q}$. Moreover, we say that \mathcal{V} is a variety of degree d(and write $\deg(\mathcal{V}) = d$) if $d = \#(\mathcal{V} \cap H)$, where $H \subseteq \mathbb{P}^r(\overline{\mathbb{F}_q})$ is a general projective subspace of dimension r - s. To determine the degree of a variety is generally not straighforward; however an upper bound to $\deg(\mathcal{V})$ is given by $\prod_{i=1}^s \deg(F_i)$. We also recall that the Frobenius map $\Phi_q : x \mapsto x^q$ is an automorphism of \mathbb{F}_{q^3} and generates the group $Gal(\mathbb{F}_{q^3}/\mathbb{F}_q)$ of automorphisms of \mathbb{F}_{q^3} that fixes \mathbb{F}_q pointwise. The Frobenius automorphism induces also a collineation of $\mathbb{P}^r(\mathbb{F}_q^3)$ and an automorphism of $\mathbb{F}_{q^3}[X_0, \ldots, X_r]$.

A crucial point in our investigation of permutation trinomials over \mathbb{F}_{q^3} is to prove the existence of suitable \mathbb{F}_q -rational points in algebraic surfaces \mathcal{V} attached to each permutation trinomial. This is reached by proving the existence of absolutely irreducible \mathbb{F}_q -rational components in \mathcal{V} and lower bounding the number of their \mathbb{F}_q -rational points. To this end, generalizations of Lang-Weil type bounds for algebraic varieties are needed. To ensure the existence of a suitable \mathbb{F}_q -rational point of \mathcal{V} , we report the following result.

Theorem 2.1. [11, Theorem 7.1] Let $V \subseteq \mathbb{A}^n(\mathbb{F}_q)$ be an absolutely irreducible variety defined over \mathbb{F}_q of dimension r > 0 and degree δ . If $q > 2(r+1)\delta^2$, then the following estimate holds:

$$\left| \#(\mathcal{V}(\mathbb{A}^{n}(\mathbb{F}_{q}))) - q^{r} \right| \leq (\delta - 1)(\delta - 2)q^{r-1/2} + 5\delta^{13/3}q^{r-1}.$$
(3)

3 Connection with algebraic surfaces

The following proposition gives the explicit link between permutation polynomial and algebraic curves.

Proposition 3.1. If $f_{A,B}(X) = X^{q^2-q+1} + AX^{q^2} + BX \in \mathbb{F}_{q^3}[X]$ is a PP on \mathbb{F}_{q^3} then the curve

$$\mathcal{C}_{f_{A,B}}:\varphi_{f_{A,B}}:=\frac{Y^q(X^{q^2+1}+AX^{q^2+q}+BX^{q+1})+X^q(Y^{q^2+1}+AY^{q^2+q}+BY^{q+1})}{X^qY^q(X+Y)}=0$$

has no affine \mathbb{F}_{q^3} -rational points off XY(X+Y) = 0. Moreover, if f(X) has no trivial roots in \mathbb{F}_{q^3} , then the converse is also true.

Proof. If $f_{A,B}$ is a PP on \mathbb{F}_{q^3} , then for all $x, y \in \mathbb{F}_{q^3}$, $x \neq 0 \neq y$ and $x \neq y$,

$$0 \neq x^{q}y^{q}(f_{A,B}(x) + f_{A,B}(y)) = y^{q}(x^{q^{2}+1} + Ax^{q^{2}+q} + Bx^{q+1}) + x^{q}(y^{q^{2}+1} + Ay^{q^{2}+q} + By^{q+1})$$

and so $C_{f_{A,B}}$ has no affine \mathbb{F}_{q^3} -rational points off XY(X+Y) = 0. The converse also holds.

Unfortunately, $\deg(C_{f_{A,B}}) = q^2 - 1 > \sqrt[4]{q^3}$ so Hasse-Weil type theorems do not ensure the existence of \mathbb{F}_{q^3} -rational points for the curve $C_{f_{A,B}}$.

We overcome this problem by considering a link between $C_{f_{A,B}}$ and a suitable surface in $\mathbb{P}^5(\mathbb{F}_{q^3})$ of small degree, in order to apply Theorem 2.1.

Define $\mathcal{C}: X^q Y^q (X+Y) \varphi_{f_{A,B}}(X,Y) = 0$. Let $\left\{\xi, \xi^q, \xi^{q^2}\right\}$ be a normal basis of \mathbb{F}_{q^3} over \mathbb{F}_q . Write $X = x_0 \xi + x_1 \xi^q + x_2 \xi^{q^2}$ and

$$Y = y_0 \xi + y_1 \xi^q + y_2 \xi^{q^2}$$
, where $x_i, y_i \in \mathbb{F}_q$, $i = 0, 1, 2$.

Clearly, the invertible map

$$\Lambda: \mathbb{F}_{q^3}^2 \to \mathbb{F}_q^6, \qquad (X, Y) \mapsto (x_0, x_1, x_2, y_0, y_1, y_2) \tag{4}$$

induces a surjective map between the set of affine \mathbb{F}_{q^3} -rational points of \mathcal{C} and the set of projective \mathbb{F}_{q^-} rational points of a suitable surface $\mathcal{V} \subseteq \mathbb{P}^5(\mathbb{F}_q)$ of equation

$$f_1(x_0, x_1, x_2, y_0, y_1, y_2)\xi + f_2(x_0, x_1, x_2, y_0, y_1, y_2)\xi^q + f_3(x_0, x_1, x_2, y_0, y_1, y_2)\xi^{q^2} = 0$$

i.e.

$$\mathcal{V}: \left\{ \begin{array}{l} f_1(x_0, x_1, x_2, y_0, y_1, y_2) = 0\\ f_2(x_0, x_1, x_2, y_0, y_1, y_2) = 0\\ f_3(x_0, x_1, x_2, y_0, y_1, y_2) = 0, \end{array} \right.$$

where $f_i \in \mathbb{F}_q[x_0, x_1, x_2, y_0, y_1, y_2]$ and $\deg(f_i) = 3$, for i = 1, 2, 3. Setting

$$X = x_0\xi + x_1\xi^q + x_2\xi^{q^2} =: X_0 Y = y_0\xi + y_1\xi^q + y_2\xi^{q^2} =: Y_0$$

$$X^q = x_2\xi + x_0\xi^q + x_1\xi^{q^2} =: X_1 Y^q = y_2\xi + y_0\xi^q + y_1\xi^{q^2} =: Y_1$$

$$X^{q^2} = x_1\xi + x_2\xi^q + x_0\xi^{q^2} =: X_2 Y^{q^2} = y_1\xi + y_2\xi^q + y_0\xi^{q^2} =: Y_2,$$

it follows that the map $\Theta: (x_0: x_1: x_2: y_0: y_1: y_2) \mapsto (X_0: X_1: X_2: Y_0: Y_1: Y_2)$ is a projectivity of $\mathbb{P}^5(\mathbb{F}_{q^3})$. Moreover the Frobenius automorphism Φ_q induces a collineation of $\mathbb{P}^5(\mathbb{F}_{q^3})$ that fixes \mathcal{V} , so the surface $\Theta^{-1}(\mathcal{V})$ is fixed by the collineation

$$\Psi_q: \mathbb{P}^5(\mathbb{F}_{q^3}) \to \mathbb{P}^5(\mathbb{F}_{q^3}), \quad (u_0: u_1: u_2: v_0: v_1: v_2) \mapsto (u_2^{q^2}: u_0^{q^2}: u_1^{q^2}: v_2^{q^2}: v_0^{q^2}: v_1^{q^2}).$$

We will denote by Ψ_q also the automorphism of $\mathbb{F}_{q^3}[X_0, X_1, X_2, Y_0, Y_1, Y_2]$ given by

$$H(X_0, X_1, X_2, Y_0, Y_1, Y_2) \mapsto H^q(X_1, X_2, X_0, Y_1, Y_2, Y_0),$$

where H^q denotes the polynomial obtained raising the coefficients of H to the power q. Observing that

$$X^{q}Y^{q}(X+Y)\varphi_{f_{A,B}} = Y^{q}(X^{q^{2}}X+AX^{q^{2}}X^{q}+BX^{q}X) + X^{q}(Y^{q^{2}}Y+AY^{q^{2}}Y^{q}+BY^{q}Y)$$

=: $\overline{F}(X, X^{q}, X^{q^{2}}, Y, Y^{q}, Y^{q^{2}}),$

with $\overline{F} \in \mathbb{F}_{q^3}[X_0, X_1, X_2, Y_0, Y_1, Y_2]$, we get that the surface $\Theta^{-1}(\mathcal{V})$ is defined by

$$\left\{ \begin{array}{l} \overline{F}=0\\ \Psi_q(\overline{F})=0\\ \Psi_q^2(\overline{F})=0, \end{array} \right.$$

i.e.

$$\Theta^{-1}(\mathcal{V}): \begin{cases} Y_1(X_0X_2 + AX_1X_2 + BX_0X_1) = X_1(Y_0Y_2 + AY_1Y_2 + BY_0Y_1) \\ Y_2(X_0X_1 + A^qX_0X_2 + B^qX_1X_2) = X_2(Y_0Y_1 + A^qY_0Y_2 + B^qY_1Y_2) \\ Y_0(X_1X_2 + A^{q^2}X_0X_1 + B^{q^2}X_0X_2) = X_0(Y_1Y_2 + A^{q^2}Y_0Y_1 + B^{q^2}Y_0Y_2). \end{cases}$$
(5)

Again the composition

$$\Theta \circ \Lambda : \mathbb{F}_{q^3}^2 \to \mathbb{F}_q^6, \quad (X, Y) \mapsto (X_0, X_1, X_2, Y_0, Y_1, Y_2)$$

induces a surjective map between the set of affine \mathbb{F}_{q^3} -rational points of \mathcal{C} and the set of points of $\Theta^{-1}(\mathcal{V})$ in $\mathbb{P}^{5}(\mathbb{F}_{q^{3}})$ which are fixed by Ψ_{q} . Furthermore, one can observe that every component of $\Theta^{-1}(\mathcal{V})$ which is fixed by Ψ_q corresponds to an \mathbb{F}_q -rational component of \mathcal{V} and viceversa.

We are now interested in studying the existence of absolutely irreducible components of $\Theta^{-1}(\mathcal{V})$ which are fixed by Ψ_q .

Clearly, from $\varphi_{f_{A,B}}(X,0) = \varphi_{f_{A,B}}(0,Y) = \varphi_{f_{A,B}}(X,X) = 0$ one deduces that the planes

$$\mathcal{U}_{1}: \begin{cases} X_{2} = 0 \\ X_{1} = 0 \\ X_{0} = 0 \end{cases} \qquad \mathcal{U}_{2}: \begin{cases} Y_{2} = 0 \\ Y_{1} = 0 \\ Y_{0} = 0 \end{cases} \qquad \mathcal{U}_{3}: \begin{cases} X_{2} = Y_{2} \\ X_{1} = Y_{1} \\ X_{0} = Y_{0} \end{cases} \tag{6}$$

are absolutely irreducible components of $\Theta^{-1}(\mathcal{V})$. It may be noticed that points on XY(X+Y) = 0correspond via the map $\Theta \circ \Lambda$ to points on $\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3$ and vice versa. It is not hard to see that also the surface

$$\mathcal{U}: \begin{cases} X_2 = 0\\ X_0 = 0\\ Y_0 Y_2 + A Y_1 Y_2 + B Y_0 Y_1 = 0 \end{cases}$$

is an absolutely irreducible component of $\Theta^{-1}(\mathcal{V})$. Moreover, since $\varphi_{f_{A,B}}(X,Y) = \varphi_{f_{A,B}}(Y,X), \Theta^{-1}(\mathcal{V})$ is fixed by the projectivity

$$\Sigma: (X_0, X_1, X_2, Y_0, Y_1, Y_2) \mapsto (Y_0, Y_1, Y_2, X_0, X_1, X_2)$$

over $\mathbb{P}^5(\mathbb{F}_{q^3})$, as well as by Ψ_q . Therefore $\mathcal{U}, \Sigma(\mathcal{U}), \Psi_q(\mathcal{U}), \Sigma(\Psi_q(\mathcal{U})), \Psi_q^2(\mathcal{U}), \Sigma(\Psi_q^2(\mathcal{U}))$ are six distinct components of $\Theta^{-1}(\mathcal{V})$ of degree 2, which are not fixed by Ψ_q . Thus $\Theta^{-1}(\mathcal{V})$ splits in at least ten components

$$\Theta^{-1}(\mathcal{V}) \supseteq \left(\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3 \cup \mathcal{U} \cup \Sigma(\mathcal{U}) \cup \Psi_q(\mathcal{U}) \cup \Sigma(\Psi_q(\mathcal{U})) \cup \Psi_q^2(\mathcal{U}) \cup \Sigma(\Psi_q^2(\mathcal{U}))\right) \cup \mathcal{W},\tag{7}$$

where \mathcal{W} is a surface fixed by Ψ_q and of degree at most 27 - 3 - 12 = 12.

From now on we will investigate the absolutely irreducibility of \mathcal{W} . From the first equation of System (5) we obtain $X_2 = X_1 \cdot \frac{Y_2(Y_0 + AY_1) + BY_1(X_0 + Y_0)}{Y_1(X_0 + AX_1)}$. By replacing X_2 in the second equation of (5), we get $X_1 = 0$ (from which we obtain $\mathcal{U}_1 \cup \Psi_{q^2}(\mathcal{U})$) or

$$X_{1} = \frac{[Y_{2}(Y_{0} + AY_{1}) + BY_{1}(X_{0} + Y_{0})][A^{q}Y_{2}(X_{0} + Y_{0}) + Y_{1}(Y_{0} + B^{q}Y_{2})] + X_{0}^{2}Y_{1}Y_{2}}{Y_{2}[AX_{0}Y_{1} + B^{q}(Y_{2}(Y_{0} + AY_{1}) + BY_{1}(X_{0} + Y_{0}))]}$$

Replacing X_1 and X_2 as rational functions in X_0, Y_0, Y_1, Y_2 in the last equation of the system, we obtain $X_0 = Y_0$ (i.e. the component \mathcal{U}_3) or

$$\mathcal{W}: \left\{ \begin{array}{l} X_2 = \frac{[Y_2(Y_0 + AY_1) + BY_1(X_0 + Y_0)][A^qY_2(X_0 + Y_0) + Y_1(Y_0 + B^qY_2)] + X_0^2Y_1Y_2}{Y_1[(A^{q+1} + B^q)X_0Y_2 + A^{q+1}Y_0Y_2 + AB^qY_1Y_2 + AY_0Y_1]} \\ X_1 = \frac{[Y_2(Y_0 + AY_1) + BY_1(X_0 + Y_0)][A^qY_2(X_0 + Y_0) + Y_1(Y_0 + B^qY_2)] + X_0^2Y_1Y_2}{Y_2[AX_0Y_1 + B^q(Y_2(Y_0 + AY_1) + BY_1(X_0 + Y_0))]} \\ G(X_0, Y_0, Y_1, Y_2) = 0, \end{array} \right.$$

with $G \in \mathbb{F}_{q^3}[X_0, Y_0, Y_1, Y_2]$ homogeneous. By MAGMA computations [10], we get $\deg(G(X_0, Y_0, Y_1, Y_2)) = 8$ and

$$G_*(X_0, Y_0, Y_1) = G(X_0, Y_0, Y_1, 1) = \alpha(Y_0, Y_1)X_0^3 + \beta(Y_0, Y_1)X_0^2 + \gamma(Y_0, Y_1)X_0 + \delta(Y_0, Y_1),$$
(8)

where $\alpha, \beta, \gamma, \delta \in \mathbb{F}_{q^3}[Y_0, Y_1]$.

If α is not zero, then $\deg_{X_0}(G_*) = 3$ and G_* is not absolutely irreducible if and only if there exists a factor $\epsilon(Y_0, Y_1)X_0 + \sigma(Y_0, Y_1)$ of G_* , where $\epsilon(Y_0, Y_1) \mid \alpha(Y_0, Y_1)$ and $\sigma(Y_0, Y_1) \mid \delta(Y_0, Y_1)$. By MAGMA computations, it results that

$$\begin{aligned} \alpha(Y_0,Y_1) &= (A^q B + 1)(A^{1+q+q^2} + AB^{q^2} + A^q B + A^{q^2}B^q + B^{1+q+q^2} + 1)Y_0Y_1^2, \\ \delta(Y_0,Y_1) &= (Y_0Y_1 + B^q Y_1Y_2 + A^q Y_0Y_2)_*^2(AY_1Y_2 + BY_0Y_1 + Y_0Y_2)_*^2 =: M_*^2N_*^2. \end{aligned}$$

Thus $\epsilon(Y_0, Y_1) = Y_0^i Y_1^j$ and $\sigma(Y_0, Y_1) = \lambda M_*^\ell N_*^k$, with $0 \le i \le 1, 0 \le j, \ell, k \le 2$, for some $\lambda \in \mathbb{F}_{q^3} \setminus \{0\}$. Also $\epsilon(Y_0, Y_1) X_0 + \sigma(Y_0, Y_1) \mid G_*$ if and only if $G_*(\frac{\lambda M_*^\ell N_*^k}{Y_0^i Y_1^j}, Y_0, Y_1) = 0$ in $\mathbb{F}_{q^3}(Y_0, Y_1)$, that is (clearing the denominators)

$$\alpha M_*^{3\ell} N_*^{3k} + \beta M_*^{2\ell} N_*^{2k} Y_0^i Y_1^j + \gamma M_*^\ell N_*^k Y_0^{2i} Y_1^{2j} + \delta Y_0^{3i} Y_1^{3j} = 0$$
(9)

in $\mathbb{F}_{q^3}[Y_0, Y_1]$.

We first investigate the case $\alpha = 0$.

Proposition 3.2. Let $A, B \in \mathbb{F}_{q^3}$. Then $f_{A,B}$ is not a permutation polynomial over \mathbb{F}_{q^3} in any of the following cases.

(i) $A^qB + 1 = 0$ and $A^{1+q+q^2} = 1;$

(ii) $A^qB + 1 \neq 0$ and $A^{1+q+q^2} + AB^{q^2} + A^qB + A^{q^2}B^q + B^{1+q+q^2} + 1 = 0.$

Proof. We want to find a non-trivial root of $f_{A,B}$ in \mathbb{F}_{q^3} ; the statement will follow.

(i) Clearly, $f_{A,B}$ has a non-trivial root in \mathbb{F}_{q^3} if and only if there exists $u \in \mathbb{F}_{q^3}, u^{1+q+q^2} = 1$, such that

$$u^q + Au^{q+1} + A^{-q} = 0$$

(see [40, Theorem 1]), i.e. such that

$$(Au)^{q} + (Au)^{q+1} + 1 = 0. (10)$$

It is well-known that all the q + 1 solutions of the equation $y^{q+1} + y^q + 1 = 0$ are elements of \mathbb{F}_{q^3} and satisfy $y^{1+q+q^2} = 1$ (see for istance [39]). Thus consider $\overline{u} = A^{-1}\overline{y}$, where $\overline{y}^{q+1} + \overline{y}^q + 1 = 0$; such \overline{u} satisfies $\overline{u}^{1+q+q^2} = 1$ if and only if $A^{1+q+q^2} = 1$. From the surjectivity of the map $\gamma : \mathbb{F}_{q^3} \to \{x \in \mathbb{F}_{q^3} : x^{1+q+q^2} = 1\}, x \mapsto x^{q-1}$, we get the statement.

(ii) Again, f has a non-trivial root in \mathbb{F}_{q^3} if and only if there exists $u \in \mathbb{F}_{q^3}, u^{1+q+q^2} = 1$, such that

$$u^q + Au^{q+1} + B = 0; (11)$$

(see [40]).

In [40, Theorem 1], the authors proved that if u is a solution of (11) with $u^{1+q+q^2} = 1$, then $(A + B^{1+q})u = A^q B + 1$. Moreover, from

$$\left\{ \begin{array}{l} A^{1+q+q^2} + AB^{q^2} + A^qB + A^{q^2}B^q + B^{1+q+q^2} + 1 = 0 \\ A + B^{1+q} = 0 \end{array} \right.$$

it follows that $A^{1+q+q^2} = B^{1+q+q^2} = 1$, a contradiction to $A^q B + 1 \neq 0$. So $u = \frac{A^q B + 1}{A + B^{1+q}}$ is an element of $\mathbb{F}_{q^3}^*$. We now prove that such u has the required properties. We have that $u^{1+q+q^2} = 1$ if and only if $(A^q B + 1)^{1+q+q^2} = (A + B^{1+q})^{1+q+q^2}$ that is $B^{1+q+q^2}(A^{1+q+q^2} + AB^{q^2} + A^q B + A^{q^2} B^q + B^{1+q+q^2} + 1) = 0$. Finally, by replacing $u = \frac{A^q B + 1}{A + B^{1+q}}$ in Equation (11) we obtain

$$\begin{pmatrix} \frac{A^{q}B+1}{A+B^{1+q}} \end{pmatrix}^{q} + A \cdot \left(\frac{A^{q}B+1}{A+B^{1+q}} \right)^{q+1} + B$$

$$= \frac{A^{q^{2}}B^{q}+1}{A^{q}+B^{q+q^{2}}} \left[1 + \frac{A(A^{q}B+1)}{A+B^{1+q}} \right] + B$$

$$= \frac{(A^{q^{2}}B^{q}+1)(B^{1+q}+A^{1+q}B)}{(A+B^{1+q})(A^{q}+B^{q+q^{2}})} + B$$

$$= \frac{B^{1+q}(A^{1+q+q^{2}}+AB^{q^{2}}+A^{q}B+A^{q^{2}}B^{q}+B^{1+q+q^{2}}+1)}{(A+B^{1+q})(A^{q}+B^{q+q^{2}})}$$

$$= 0.$$

The case $A^q B = 1$ and $A^{1+q+q^2} \neq 1$ is investigated in Proposition 4.1. In the rest of this section we assume $\alpha \neq 0$.

Proposition 3.3. Let $\alpha \neq 0$. Then $f_{i,j,\ell,k} := \lambda Y_0^i Y_1^j X_0 + M_*^\ell N_*^k \nmid G_*$ in any of the following cases:

- (a) $\ell + k \ge 3;$
- (b) $\ell + k = 2$ and i + j < 2;
- (c) $\ell + k = 0;$
- (d) $\ell + k = 1$ and $i + j \ge 2$.

Proof. By MAGMA computations, we get $\deg(\beta) = 5$, $\deg(\gamma) = 7$ (if β, γ are not zero) and $\deg(\delta) = 8$. By a direct check the leading term of the polynomial in Equation (9) appears in $\alpha M_*^{3\ell} N_*^{3k}$ in cases (a) and (b), and in $\delta Y_0^{3i} Y_1^{3j}$ in cases (c) and (d). The claim follows.

We consider now all the pending cases,

• $\ell + k = 2$ and i + j = 2, 3;

• $\ell + k = 1$ and i + j = 0, 1.

Proposition 3.4. Let $\alpha \neq 0$. If G_* is reducible, then a factor $\epsilon X_0 + \sigma$ of G_* equals one among

- $Y_0 Y_1 X_0 + \lambda M_* N_*,$
- $X_0 + \lambda N_*$,
- $Y_1X_0 + \lambda M_*$.

Proof. It is readily seen that a polynomial $\epsilon(Y_0, Y_1)X_0 + \sigma(Y_0, Y_1)$ divides G_* if and only if

$$\widetilde{G}(Y_0, Y_1) := \epsilon(Y_0, Y_1)^3 G_* \left(\frac{\sigma(Y_0, Y_1)}{\epsilon(Y_0, Y_1)}, Y_0, Y_1 \right)$$

vanishes. We distinguish the following cases:

- $Y_1^2 X_0 + \lambda N_* (Y_0, Y_1)^2$. MAGMA computations show that the coefficient of $Y_0^5 Y_1^7$ in $\widetilde{G}(Y_0, Y_1)$ is $\lambda A^{1+q^2} (A^q B + 1) \neq 0$, so $Y_1^2 X_0 + \lambda N_* \nmid G_*$.
- $Y_0 Y_1 X_0 + \lambda N_*^2$.

The coefficient of $Y_0^6 Y_1^5$ in $\widetilde{G}(Y_0, Y_1)$ is $\lambda AB^q (A^q B + 1)(A + B^{q+1})^{q^2}$, that is zero if and only if $A + B^{1+q} = 0$. By replacing $A = B^{q+1}$ in $\widetilde{G}(Y_0, Y_1)$ we get that the coefficient of $Y_0 Y_1^7$ in $\widetilde{G}(Y_0, Y_1)$ is $\lambda^3 B^{6(1+q)} (B^{1+q+q^2} + 1)^3$. By replacing $B^{1+q+q^2} = 1$ in $\widetilde{G}(Y_0, Y_1)$, MAGMA computations show that the coefficient of $Y_0^7 Y_1^7$ in $\widetilde{G}(Y_0, Y_1)$ is $B^2 \neq 0$. Thus $Y_0 Y_1 X_0 + \lambda N_*^2 \nmid G_*$.

- $Y_0 Y_1^2 X_0 + \lambda N_*^2$. Analogous to the case $\epsilon = Y_0 Y_1$ and $\sigma = \lambda N_*^2$.
- $Y_1^2 X_0 + \lambda M_* N_*$. The coefficients of $Y_0^7 Y_1^8$ in $\widetilde{G}(Y_0, Y_1)$ reads $\lambda^3 B^3 (A^q B + 1) (A^{1+q+q^2} + A B^{q^2} + A^q B + A^{q^2} B^q + B^{1+q+q^2} + 1) \neq 0$. So $Y_1^2 X_0 + \lambda M_* N_* \nmid G_*$.
- $Y_0Y_1^2X_0 + \lambda M_*N_*$. Analogous to the case $\epsilon = Y_0Y_1$ and $\sigma = \lambda N_*^2$.
- $Y_1^2 X_0 + \lambda M_*^2$. The coefficient of $Y_0^5 Y_1^7$ in $\widetilde{G}(Y_0, Y_1)$ is $\lambda B^{q+q^2}(A^q B + 1) \neq 0$. Thus $Y_1^2 X_0 + \lambda M_*^2 \nmid G_*$.
- $Y_0Y_1X_0 + \lambda M_*^2$. Analogous to the case $\epsilon = Y_0Y_1$ and $\sigma = \lambda N_*^2$.
- $Y_0Y_1^2X_0 + \lambda M_*^2$. Analogous to the case $\epsilon = Y_0Y_1$ and $\sigma = \lambda N_*^2$.
- $Y_1X_0 + \lambda N_*$. The coefficient of $Y_0Y_1^7$ in $\widetilde{G}(Y_0, Y_1)$ reads $\lambda A^3(A^qB + 1)^q \neq 0$, so $Y_1X_0 + \lambda N_* \nmid G_*$.

- $Y_0X_0 + \lambda N_*$. The coefficient of Y_0^7 in $\widetilde{G}(Y_0, Y_1)$ is $A^{2q} \neq 0$. Thus $Y_0X_0 + \lambda N_* \nmid G_*$.
- $X_0 + \lambda M_*$. The coefficient of $Y_0 Y_1^3$ in $\widetilde{G}(Y_0, Y_1)$ reads $\lambda B^{3q} (A^q B + 1)^{q^2} \neq 0$, so $X_0 + \lambda M_* \nmid G_*$.
- $Y_0X_0 + \lambda M_*$ The coefficient of Y_0^7 in $\widetilde{G}(Y_0, Y_1)$ is $A^{2q} \neq 0$. Thus $Y_0X_0 + \lambda M_* \nmid G_*$.

The claim now follows from Proposition 3.3.

Remark 3.5. For $A^q B \in \mathbb{F}_q^* \setminus \{1\}$ and $A^{1+q+q^2} = B^{1+q+q^2}$, it is already known that $f(X) = X^{q^2-q+1} + AX^{q^2} + BX$ is a PP over \mathbb{F}_{q^3} for each q (see [40, Theorem 1]). In these cases \mathcal{W} decomposes in three components and $G_* = (A^q B + 1)^2 (Y_0 Y_1 X_0 + \lambda_1 M_* N_*) (X_0 + \lambda_2 N_*) (Y_1 X_0 + \lambda_3 M_*)$, for suitable $\lambda_1, \lambda_2, \lambda_3 \in \mathbb{F}_{q^3}$. In what follows we will prove that for q large enough, also the converse is true.

Proposition 3.6. Let $\alpha \neq 0$. If $X_0 + \lambda N_* | G_*$ or $Y_1 X_0 + \lambda M_* | G_*$, then $A^q B \in \mathbb{F}_q^* \setminus \{1\}$ and $A^{1+q+q^2} = B^{1+q+q^2}$.

Proof. MAGMA computations show that the coefficient of Y_1^4 in $\widetilde{G}(Y_0, Y_1) = G_*(\lambda N_*, Y_0, Y_1)$ is $A^2B^q((A^{q+1} + B^q)\lambda + B^q)$, so $\widetilde{G}(Y_0, Y_1) = 0$ yields $(A^{q+1} + B^q)\lambda = B^q$, and in particular $A^{q+1} + B^q \neq 0$. By replacing $\lambda = \frac{B^q}{A^{q+1} + B^q}$ in \widetilde{G} , the coefficients of $Y_0^2Y_1^2$ and $Y_0^2Y_1^3$ in \widetilde{G} are $A^{1+q}(A^{q+1} + B^q)(A^{1+2q} + B^{2q+q^2})$ and $A^{1+q}B^q(A^{q+1} + B^q)(A^{1+q+q^2} + A^qB + A^{q^2}B^q + B^{1+q+q^2})$, respectively. Therefore, $\widetilde{G} = 0$ implies $A^{1+2q} + B^{2q+q^2} = 0 = A^{1+q+q^2} + A^qB + A^{q^2}B^q + B^{1+q+q^2}$, from which it follows that $B^q(A^{q+1} + B^q)(A^qB + A^{q^2}B^q) = 0$, so $A^qB \in \mathbb{F}_q^* \setminus \{1\}$ and $A^{1+q+q^2} = B^{1+q+q^2}$. The case $Y_1X_0 + \lambda N_*, X_0 \mid G_*$ is analogous. □

Proposition 3.7. Suppose that \mathcal{W} does not contain an absolutely irreducible component fixed by Ψ_q . If $X_0Y_0Y_1 + \lambda M_*N_* \mid G_*$, then there is another factor $\epsilon X_0 + \sigma$ that divides G_* . In particular $A^q B \in \mathbb{F}_q^* \setminus \{1\}$ and $A^{1+q+q^2} = B^{1+q+q^2}$.

Proof. First, we observe that $Y_0^2 \nmid \alpha(Y_0, Y_1)$, so $X_0Y_0Y_1 + \lambda MN$ cannot be a repeated factor of G_* In particular

$$\mathcal{W}_{1}: \begin{cases} X_{2} = \frac{[BX_{0}Y_{1} + N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2} + M(Y_{0},Y_{1},Y_{2})] + X_{0}^{2}Y_{1}Y_{2}}{Y_{1}[(A^{q+1} + B^{q})X_{0}Y_{2} + A \cdot M(Y_{0},Y_{1},Y_{2})]} \\ X_{1} = \frac{[BX_{0}Y_{1} + N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2} + M(Y_{0},Y_{1},Y_{2})] + X_{0}^{2}Y_{1}Y_{2}}{Y_{2}[(A + B^{q+1})X_{0}Y_{1} + B^{q} \cdot N(Y_{0},Y_{1},Y_{2})]} \\ X_{0} = \lambda \cdot \frac{M(Y_{0},Y_{1},Y_{2})N(Y_{0},Y_{1},Y_{2})}{Y_{0}Y_{1}Y_{2}} \end{cases}$$

is a non-repeated absolutely irreducible component of \mathcal{W} over \mathbb{F}_{q^3} . Thus there are two possible factorization of G_* :

(i) $G_* = (X_0 Y_0 Y_1 + \lambda M_* N_*) p(X_0, Y_0, Y_1)$ where $p = \frac{\alpha(Y_0, Y_1)}{Y_0 Y_1^2} X_0^2 Y_1 + \dots + \frac{1}{\lambda} M_*(Y_0, Y_1) N_*(Y_0, Y_1)$ is absolutely irreducible of degree two in X_0 .

(ii) $G_* = (X_0Y_0Y_1 + \lambda M_*N_*)(\epsilon_1X_0 + \sigma_1)(\epsilon_2X_0 + \sigma_2)$ with $\epsilon_1 \neq Y_0Y_1 \neq \epsilon_2$.

We want to prove that a factorization of type (i) cannot exist, namely that $p(Y_0, Y_1)$ can't be absolutely irreducible; from this the claim will follow.

Suppose by way of contradiction that p is absolutely irreducible; then

$$\mathcal{W}_{2}: \begin{cases} X_{2} = \frac{[BX_{0}Y_{1} + N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2} + M(Y_{0},Y_{1},Y_{2})] + X_{0}^{2}Y_{1}Y_{2}}{Y_{1}[(A^{q+1} + B^{q})X_{0}Y_{2} + A \cdot M(Y_{0},Y_{1},Y_{2})]} \\ X_{1} = \frac{[BX_{0}Y_{1} + N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2} + M(Y_{0},Y_{1},Y_{2})] + X_{0}^{2}Y_{1}Y_{2}}{Y_{2}[(A + B^{q+1})X_{0}Y_{1} + B^{q} \cdot N(Y_{0},Y_{1},Y_{2})]} \\ \frac{\alpha(Y_{0},Y_{1})}{Y_{0}Y_{1}^{2}}X_{0}^{2}Y_{1}Y_{2} + \dots + \frac{1}{\lambda}M(Y_{0},Y_{1},Y_{2})N(Y_{0},Y_{1},Y_{2}) = 0 \end{cases}$$

is an absolutely irreducible component of \mathcal{W} defined over \mathbb{F}_{q^3} and $\mathcal{W} = \mathcal{W}_1 \cup \mathcal{W}_2$. Consider the collineation Ψ_q of $\mathbb{P}^5(\mathbb{F}_{q^3})$ introduced before. Since \mathcal{V}_1 is not fixed by Ψ_q so $\Psi_q(\mathcal{V}_1) = \mathcal{V}_2$. As Ψ_q is a collineation of $\mathbb{P}^5(\mathbb{F}_{q^3})$, \mathcal{V}_1 and \mathcal{V}_2 have the same degree. Moreover, if p is absolutely irreducible, then it is not a square as a polynomial, so there exists a subspace $\pi : \begin{cases} Y_0 = hY_2 \\ Y_1 = kY_2 \end{cases}$ of $\mathbb{P}^5(\mathbb{F}_{q^3})$ of dimension 3 not contained in the hyperplanes $Y_0 = 0$ nor $Y_1 = 0$ (i.e. $h \neq 0 \neq k$) such that $p(X_0, h, k)$ has 2 distinct roots, i.e. $|\mathcal{V}_2 \cap \pi| = 2$, while $|\mathcal{V}_1 \cap \tilde{\pi}| = 1$, where $\Psi_q(\tilde{\pi}) = \pi$, $\tilde{\pi} : \begin{cases} Y_2 = h^{q^2}Y_1 \\ Y_0 = k^{q^2}Y_1 \end{cases}$, a contradiction. So there exists a factor $\epsilon X_0 + \sigma$ of G_* different from $X_0Y_0Y_1 + \lambda M_*N_*$. By Proposition 3.4 we have that one between $X_0 + \lambda N_*$ and $Y_1X_0 + \lambda M_*$ divides G_* ; the claim now follows from Proposition 3.6.

We can now prove our main result.

Theorem 3.8. Let $q = 2^m$, with $m \ge 19$. Assume that $f(X) = X^{q^2-q+1} + AX^{q^2} + BX \in \mathbb{F}_{q^3}[X]$ is a PP over \mathbb{F}_{q^3} and let $A^qB + 1 \ne 0 \ne A^{1+q+q^2} + AB^{q^2} + A^qB + A^{q^2}B^q + B^{1+q+q^2} + 1$. Then $A^{q^2+q+1} = B^{1+q+q^2}$ and $A^qB \in \mathbb{F}_q \setminus \{0, 1\}$.

Proof. Assume by way of contradiction that there exists an absolutely irreducible components component \mathcal{W}_1 of \mathcal{W} fixed by Ψ_q .

Then \mathcal{W}_1 corresponds via the projectivity Θ to an absolutely irreducible \mathbb{F}_q -rational component \mathcal{V}_1 of \mathcal{V} , with $\dim(\mathcal{V}_1) = \dim(\mathcal{W}_1) \leq \dim(\mathcal{W}) \leq 12$ (see Equation (7)). So, by Theorem 2.1 it follows that

$$\#(\mathcal{V}_1(\mathbb{F}_q^5)) \ge q^2 - 110q^{3/2} - 5 \cdot 12^{13/3}q.$$
(12)

Moreover, \mathcal{V}_1 intersects each of the planes

$$\Theta(\mathcal{U}_1): \begin{cases} x_2 = 0 \\ x_1 = 0 \\ x_0 = 0 \end{cases} \qquad \Theta(\mathcal{U}_2): \begin{cases} y_2 = 0 \\ y_1 = 0 \\ y_0 = 0 \end{cases} \qquad \Theta(\mathcal{U}_3): \begin{cases} x_2 = y_2 \\ x_1 = y_1 \\ x_0 = y_0 \end{cases}$$
(13)

in at most 36 points. Since

$$q^2 - 110q^{3/2} - 5 \cdot 12^{13/3}q > 36$$

is satisfied for $q \geq 2^{19}$, we deduce the existence of \mathbb{F}_q -rational points on \mathcal{V} off $\Theta(\mathcal{U}_1 \cup \mathcal{U}_2 \cup \mathcal{U}_3)$, which corresponds to points on $\mathcal{C}_{f_{A,B}}$ off $X^q Y^q (X + Y) = 0$, a contradiction to our hypothesis via Proposition 3.1.

Thus, no absolutely irreducible components component of \mathcal{W} is fixed by Ψ_q .

In particular, \mathcal{W} is reducible and so is G_* . By Propositions 3.4, 3.6 and 3.7, $A^{q^2+q+1} = B^{1+q+q^2}$ and $A^q B \in \mathbb{F}_q \setminus \{0,1\}.$

4 Case $A^q B = 1$

Set without restriction $B^{1+q+q^2} \neq 1$. Let us consider again the surface

$$\mathcal{W}: \begin{cases} X_{2} = \frac{[BX_{0}Y_{1}+N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2}+M(Y_{0},Y_{1},Y_{2})]+X_{0}^{2}Y_{1}Y_{2}}{Y_{1}[(A^{q+1}+B^{q})X_{0}Y_{2}+A\cdot M(Y_{0},Y_{1},Y_{2})]}\\ X_{1} = \frac{[BX_{0}Y_{1}+N(Y_{0},Y_{1},Y_{2})][A^{q}X_{0}Y_{2}+M(Y_{0},Y_{1},Y_{2})]+X_{0}^{2}Y_{1}Y_{2}}{Y_{2}[(A+B^{q+1})X_{0}Y_{1}+B^{q}\cdot N(Y_{0},Y_{1},Y_{2})]} \\ G(X_{0},Y_{0},Y_{1},Y_{2}) = 0. \end{cases}$$
(14)

Replacing $A = \frac{1}{Bq^2}$, $A^q = \frac{1}{B}$, and $A^{q^2} = \frac{1}{B^q}$ in G we get

$$G = (B^{2+q+2q^2})^{-1} \cdot \tilde{M} \cdot \tilde{N} \cdot [(B^{1+q+q^2}+1)Y_2X_0 + \tilde{M}][(B^{1+q+q^2}+1)Y_1X_0 + B^q\tilde{N}],$$
(15)

where $\tilde{N} := B^{1+q^2}Y_0Y_1 + B^{q^2}Y_0Y_2 + Y_1Y_2$ and $\tilde{M} := \Psi_q(\tilde{N}) = BY_0Y_1 + Y_0Y_2 + B^{1+q}Y_1Y_2$.

Proposition 4.1. If $A^q B = 1$ and $A^{1+q+q^2} \neq 1$, then $f_{A,B}$ is a PP over \mathbb{F}_{q^3} .

Proof. Recall that the map

$$\Theta \circ \Lambda : (X,Y) \mapsto (X,X^q,X^{q^2},Y,Y^q,Y^{q^2}) = (X_0,X_1,X_2,Y_0,Y_1,Y_2)$$

induces a surjection between the set of affine \mathbb{F}_{q^3} -rational points of \mathcal{C} and the set of points of $\Theta^{-1}(\mathcal{V})$ in $\mathbb{P}^5(\mathbb{F}_{q^3})$ which are fixed by Ψ_q . Consider a point $P = (\overline{X_0}, \overline{X_1}, \overline{X_2}, \overline{Y_0}, \overline{Y_1}, \overline{Y_2}) \in \Theta^{-1}(\mathcal{V})$ such that $\Psi_q(P) = P$, with $\overline{X_i Y_j} \neq 0$ and $\overline{X_i} \neq \overline{Y_i}$, for i, j = 0, 1, 2.

Now, $P \in \mathcal{W}$ and G(P) = 0 yields one of the following.

- 1. $\tilde{N}(P) = 0$, which is equivalent to $\tilde{M}(P) = 0$. In this case, $(A^{q+1} + B^q)\overline{X_0X_2Y_1Y_2} = 0$ or $(A + B^{q+1})\overline{X_0X_1Y_1Y_2} = 0$, a contradiction to $\overline{X_iY_j} \neq 0$ for i, j = 0, 1, 2.
- 2. $(B^{1+q+q^2}+1)\overline{Y_2X_0} = \tilde{M}$. From the first equation in (14) we get

$$[B\overline{X_0Y_1} + \tilde{N}(\overline{Y_0}, \overline{Y_1}, \overline{Y_2})][A^q\overline{X_0Y_2} + \tilde{M}(\overline{Y_0}, \overline{Y_1}, \overline{Y_2})] + \overline{X_0}^2\overline{Y_1Y_2} = 0,$$

which yields, by the second equation in (14), $\overline{X_1} = 0$, a contradiction to $\overline{X_i Y_j} \neq 0$ for i, j = 0, 1, 2.

3. $(B^{1+q+q^2}+1)\overline{Y_1X_0} = B^q \tilde{N}$. As in the previous case, a contradiction arises.

Finally, from Equation (10) (Proposition 3.2(i)), it follows that $f_{A,B}$ has no trivial root in \mathbb{F}_{q^3} . The claim follows from Proposition 3.1.

Acknowledgments

This research was supported by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

References

- AKBAL, Y. L. R. M., TEMÜR, B. G., AND ONGAN, P. N. A short note on permutation trinomials of prescribed type. *Comm. Algebra* 48, 4 (2020), 1608–1612.
- [2] BAI, T., AND XIA, Y. A new class of permutation trinomials constructed from Niho exponents. Cryptogr. Commun. 10, 6 (2018), 1023–1036.
- [3] BARTOLI, D. On a conjecture about a class of permutation trinomials. *Finite Fields Appl. 52* (2018), 30–50.
- [4] BARTOLI, D. Permutation trinomials over \mathbb{F}_{q^3} . Finite Fields Appl. 61 (2020), 101597, 19.
- [5] BARTOLI, D., AND TIMPANELLA, M. A family of permutation trinomials over \mathbb{F}_{q^2} . Finite Fields Appl. 70 (2021), Paper No. 101781, 16.
- [6] BARTOLI, D., AND TIMPANELLA, M. On trinomials of type $X^{n+m}(1 + AX^{m(q-1)} + BX^{n(q-1)})$, n, m odd, over \mathbb{F}_{q^2} , $q = 2^{2s+1}$. Finite Fields Appl. 72 (2021), Paper No. 101816, 31.
- [7] BARTOLI, D., AND ZINI, G. On permutation trinomials of type $x^{2p^s+r} + x^{p^s+r} + \lambda x^r$. Finite Fields Appl. 49 (2018), 126–131.
- [8] BHATTACHARYA, S., AND SARKAR, S. On some permutation binomials and trinomials over \mathbb{F}_{2^n} . Des. Codes Cryptogr. 82, 1-2 (2017), 149–160.
- [9] BLACKBURN, S. R., ETZION, T., AND PATERSON, K. G. Permutation polynomials, de Bruijn sequences, and linear complexity. J. Combin. Theory Ser. A 76, 1 (1996), 55–82.
- [10] BOSMA, W., CANNON, J., AND PLAYOUST, C. The Magma algebra system. I. The user language. J. Symbolic Comput. 24, 3-4 (1997), 235–265. Computational algebra and number theory (London, 1993).
- [11] CAFURE, A., AND MATERA, G. Improved explicit estimates on the number of solutions of equations over a finite field. *Finite Fields Appl.* 12, 2 (2006), 155–185.
- [12] DENG, H., AND ZHENG, D. More classes of permutation trinomials with Niho exponents. Cryptogr. Commun. 11, 2 (2019), 227–236.
- [13] DICKSON, L. E. The analytic representation of substitutions on a power of a prime number of letters with a discussion of the linear group. Ann. of Math. 11, 1-6 (1896/97), 65–120.

- [14] DING, C., QU, L., WANG, Q., YUAN, J., AND YUAN, P. Permutation trinomials over finite fields with even characteristic. SIAM J. Discrete Math. 29, 1 (2015), 79–92.
- [15] DING, Z., AND ZIEVE, M. E. Determination of a class of permutation quadrinomials, 2022.
- [16] FERNANDO, N. A note on permutation binomials and trinomials over finite fields. New Zealand J. Math. 48 (2018), 25–29.
- [17] FULTON, W. Algebraic curves. Advanced Book Classics. Addison-Wesley Publishing Company, Advanced Book Program, Redwood City, CA, 1989. An introduction to algebraic geometry, Notes written with the collaboration of Richard Weiss, Reprint of 1969 original.
- [18] GUPTA, R., GAHLYAN, P., AND SHARMA, R. K. New classes of permutation trinomials over \mathbb{F}_{q^3} . Finite Fields Appl. 84 (2022), Paper No. 102110, 21.
- [19] GUPTA, R., AND SHARMA, R. K. Some new classes of permutation trinomials over finite fields with even characteristic. *Finite Fields Appl.* 41 (2016), 89–96.
- [20] HERMITE, C. Sur les fonctions de sept lettres. C. R. Acad. Sci. Paris 57 (1863).
- [21] HOU, X.-D. A class of permutation trinomials over finite fields. Acta Arith. 162, 1 (2014), 51–64.
- [22] HOU, X.-D. Determination of a type of permutation trinomials over finite fields. Acta Arith. 166, 3 (2014), 253–278.
- [23] HOU, X.-D. Determination of a type of permutation trinomials over finite fields, II. Finite Fields Appl. 35 (2015), 16–35.
- [24] HOU, X.-D. Permutation polynomials over finite fields—a survey of recent advances. *Finite Fields Appl. 32* (2015), 82–119.
- [25] HOU, X.-D. A survey of permutation binomials and trinomials over finite fields. In *Topics in finite fields*, vol. 632 of *Contemp. Math. Amer. Math. Soc.*, Providence, RI, 2015, pp. 177–191.
- [26] HOU, X.-D. On a class of permutation trinomials in characteristic 2. Cryptogr. Commun. 11, 6 (2019), 1199–1210.
- [27] HOU, X.-D. On the Tu-Zeng permutation trinomial of type (1/4, 3/4). Discrete Math. 344, 3 (2021), Paper No. 112241, 14.
- [28] HOU, X.-D., AND IEZZI, A. An application of the Hasse-Weil bound to rational functions over finite fields. Acta Arith. 195, 2 (2020), 207–216.
- [29] HOU, X.-D., TU, Z., AND ZENG, X. Determination of a class of permutation trinomials in characteristic three. *Finite Fields Appl.* 61 (2020), 101596, 27.
- [30] LI, K., QU, L., AND CHEN, X. New classes of permutation binomials and permutation trinomials over finite fields. *Finite Fields Appl.* 43 (2017), 69–85.

- [31] LI, K., QU, L., CHEN, X., AND LI, C. Permutation polynomials of the form $cx + \text{Tr}_{q^l/q}(x^a)$ and permutation trinomials over finite fields with even characteristic. *Cryptogr. Commun.* 10, 3 (2018), 531–554.
- [32] LI, K., QU, L., LI, C., AND CHEN, H. On a conjecture about a class of permutation quadrinomials. *Finite Fields Appl. 66* (2020), 101690, 20.
- [33] LI, K., QU, L., LI, C., AND FU, S. New permutation trinomials constructed from fractional polynomials. Acta Arith. 183, 2 (2018), 101–116.
- [34] LI, N. On two conjectures about permutation trinomials over $\mathbb{F}_{3^{2k}}$. Finite Fields Appl. 47 (2017), 1–10.
- [35] LI, N., AND HELLESETH, T. Several classes of permutation trinomials from Niho exponents. Cryptogr. Commun. 9, 6 (2017), 693–705.
- [36] LI, N., AND HELLESETH, T. New permutation trinomials from Niho exponents over finite fields with even characteristic. *Cryptogr. Commun.* 11, 1 (2019), 129–136.
- [37] LI, N., AND HU, Q. A conjecture on permutation trinomials over finite fields of characteristic two. Adv. Math. Commun. 13, 3 (2019), 505–512.
- [38] LIU, Q., AND SUN, Y. Several classes of permutation trinomials from Niho exponents over finite fields of characteristic 3. J. Algebra Appl. 18, 4 (2019), 1950069, 12.
- [39] MULLEN, G. L., Ed. *Handbook of finite fields*. Discrete Mathematics and its Applications (Boca Raton). CRC Press, Boca Raton, FL, 2013.
- [40] PANG, T., XU, Y., LI, N., AND ZENG, X. Permutation polynomials of the form $x^d + L(x^s)$ over \mathbb{F}_{q^3} . Finite Fields Appl. 76 (2021), Paper No. 101906, 22.
- [41] PENG, J., ZHENG, L., WU, C., AND KAN, H. Permutation polynomials $x^{2^{k+1}+3} + ax^{2^k+2} + bx$ over $F_{2^{2k}}$ and their differential uniformity. *Sci. China Inf. Sci.* 63, 10 (2020), Paper No. 209101, 3.
- [42] QIN, X., AND YAN, L. Constructing permutation trinomials via monomials on the subsets of μ_{q+1} . Appl. Algebra Engrg. Comm. Comput. 34, 2 (2023), 321–334.
- [43] SHARMA, R. K., AND GUPTA, R. Determination of a type of permutation binomials and trinomials. Appl. Algebra Engrg. Comm. Comput. 31, 1 (2020), 65–86.
- [44] TU, Z., AND ZENG, X. Two classes of permutation trinomials with Niho exponents. *Finite Fields Appl. 53* (2018), 99–112.
- [45] TU, Z., AND ZENG, X. A class of permutation trinomials over finite fields of odd characteristic. Cryptogr. Commun. 11, 4 (2019), 563–583.
- [46] TU, Z., ZENG, X., AND HELLESETH, T. New permutation quadrinomials over $\mathbb{F}_{2^{2m}}$. Finite Fields Appl. 50 (2018), 304–318.

- [47] TU, Z., ZENG, X., LI, C., AND HELLESETH, T. A class of new permutation trinomials. *Finite Fields Appl. 50* (2018), 178–195.
- [48] WANG, L., WU, B., YUE, X., AND ZHENG, Y. Further results on permutation trinomials with Niho exponents. *Cryptogr. Commun.* 11, 5 (2019), 1057–1068.
- [49] WANG, Y., ZHA, Z., AND ZHANG, W. Six new classes of permutation trinomials over F_{33k}. Appl. Algebra Engrg. Comm. Comput. 29, 6 (2018), 479–499.
- [50] WANG, Y., ZHANG, W., AND ZHA, Z. Six new classes of permutation trinomials over \mathbb{F}_{2^n} . SIAM J. Discrete Math. 32, 3 (2018), 1946–1961.
- [51] WU, D., YUAN, P., DING, C., AND MA, Y. Permutation trinomials over \mathbb{F}_{2^m} . Finite Fields Appl. 46 (2017), 38–56.
- [52] WU, G., AND LI, N. Several classes of permutation trinomials over \mathbb{F}_{5^n} from Niho exponents. Cryptogr. Commun. 11, 2 (2019), 313–324.
- [53] ZHA, Z., HU, L., AND FAN, S. Further results on permutation trinomials over finite fields with even characteristic. *Finite Fields Appl.* 45 (2017), 43–52.
- [54] ZHENG, L., KAN, H., AND PENG, J. Two classes of permutation trinomials with Niho exponents over finite fields with even characteristic. *Finite Fields Appl.* 68 (2020), 101754, 14.
- [55] ZHENG, L., KAN, H., PENG, J., AND TANG, D. Two classes of permutation trinomials with Niho exponents. *Finite Fields Appl.* 70 (2021), Paper No. 101790, 15.
- [56] ZHU, X., ZENG, X., AND CHEN, Y. Some binomial and trinomial differentially 4-uniform permutation polynomials. Internat. J. Found. Comput. Sci. 26, 4 (2015), 487–497.
- [57] ZIEVE, M. E. On some permutation polynomials over \mathbb{F}_q of the form $x^r h(x^{(q-1)/d})$. Proc. Amer. Math. Soc. 137, 7 (2009), 2209–2216.
- [58] ZIEVE, M. E. A note on the paper arxiv:2207.13335, 2022.