## A Numerically Robust and Stable Time-Space Pseudospectral Approach for Generalized Burgers-Fisher Equation

Harvindra Singh<sup>1</sup>, L.K. Balyan<sup>1\*</sup>, A.K. Mittal<sup>2</sup>, Parul Saini<sup>1</sup>

<sup>1\*</sup>Discipline of Mathematics, PDPM IIITDM Jabalpur, Jabalpur, 482005, Madhya Pradesh, India.

<sup>2</sup>Department of Mathematics, VIT Chennai, Chennai, 600127, Tamil Nadu, India.

\*Corresponding author(s). E-mail(s): lokendra.balyan@gmail.com; Contributing authors: singh.harvindrahs@gmail.com; avinash.mittal10@gmail.com; iniaslurap1212@gmail.com;

#### Abstract

In this article, we present the time-space Chebyshev pseudospectral method (TS-CPsM) to approximate a solution to the generalised Burgers-Fisher (gBF) equation. The Chebyshev-Gauss-Lobatto (CGL) points serve as the foundation for the recommended method, which makes use of collocations in both the time and space directions. Further, using a mapping, the non-homogeneous initial-boundary value problem is transformed into a homogeneous problem and a system of algebraic equations is obtained. The numerical approach known as Newton-Raphson is implemented in order to get the desired results for the system. The proposed method's stability analysis has been performed. Different researchers' considerations on test problems have been explored to illustrate the robustness and practicality of the approach presented. The approximate solutions we found using the proposed method are highly accurate and are significantly better than the existing results.

**Keywords:** Pseudospectral method, Chebyshev-Gauss-Lobbato(CGL) points, Burgers equation, Fisher equation, Burgers-Fisher equation, Error analysis, 3D-plots.

### 1 Introduction

The nonlinearity of equations has been a puzzling problem for researchers for decades. Numerous studies have been conducted and are being conducted to address this concern and help science move forward. Some of the most popular non-linear equations are Burgers and Fishers equations which are of great importance due to their wide range of applications in mathematical physics. In 1915, Harry Bateman initially proposed the Burgers equation

$$U_{\tilde{t}} + UU_{\tilde{\eta}} - \mu U_{\tilde{\eta}\tilde{\eta}} = 0 \tag{1}$$

which was further investigated in 1948 by Johannes Martinus Burgers [1], [2]. The applications of Burgers equation can be seen in the context of traffic flow [3], fluid mechanics [4], gas dynamics [5], and quantum field [6]. Fishers equation

$$U_{\tilde{t}} - \mu U_{\tilde{\eta}\tilde{\eta}} = \sigma_2 U(1 - U) \tag{2}$$

referred to in certain circles as KPP(Kolmogorov-Petrovsky-Piskunov) equation or Fisher-KPP equation is a kind of equation that takes into account both reactions and diffusions which was introduced by R. A. Fisher in 1937 [7]. The Fishers equation (reaction-diffusion equation) appears in the modelling of a wide range of scientific and physical phenomena, such as Brownian motion [8], ecology [9], combustion [10], nuclear reactor theory [11], population dynamics [7], physiology, etc. A nonlinear equation is obtained by combining these two equations 1 and 2 called Burgers-Fisher equation [12] and is given by

$$U_{\tilde{t}} + \sigma_1 U U_{\tilde{\eta}} - \mu U_{\tilde{\eta}\tilde{\eta}} = \sigma_2 U (1 - U). \tag{3}$$

This equation consists of the effects of non-linear advection, linear diffusion, and nonlinear logistic reactions. This equation is a preliminary model that may be used to describe the interaction between the reaction mechanism, the diffusive transport, and the convective effect [13]. The Burgers-Fisher equation is useful for solving a wide variety of issues involving applied mathematics and physics. Its aplications include gas dynamics, traffic flow, number theory, elasticity, heat conduction, and financial mathematics [14–16]. The generalized Burgers-Fisher equation, abbreviated as the gBF equation, is the most comprehensive representation of the Burgers-Fisher equation and is given by

$$U_{\tilde{t}} + \sigma_1 U^{\delta} U_{\tilde{\eta}} - \mu U_{\tilde{\eta}\tilde{\eta}} = \sigma_2 U(1 - U^{\delta}), \quad 0 \le \tilde{\eta} \le 1, \quad \tilde{t} \ge 0, \tag{4}$$

where  $\sigma_1 \geq 0$ ,  $\mu \geq 0$ , and  $\sigma_2 \geq 0$  represent the advection coefficient of the non-linear advection term, the diffusion coefficient of linear diffusion term, and the reaction coefficient of non-linear reaction term, respectively. Literature shows a very high range of applications of Eq. 4 in the field of physics, including chemical physics, plasma physics, and fluid physics. Applications are not limited and are reported in various other fields too, such as nonlinear optics, capillary gravity waves, etc.

Due to its extreme non-linearity, the generalized Burgers-Fisher equation has served as a model for numerous academics to try out new numerical and analytical techniques. In 2004, an explicit solution was given by Kaya and El-Sayed via a numerical simulation for the gBF equation [17]. Further, a b-spline quasi-interpolation method was proposed by Zhu and Kang [18] in 2010 to approximate the gBF equation. Cubic b-spline collocation and fourth-order b -spline collocation method [19], [20], homotopy perturbation method(HPM) [21], Adomian decomposition method(ADM) [22], collocation-based radial basis functions method (CBRBF) [23], Exp-function method [24], Lattice Boltzmann model [25], etc. have been used by several researchers to find the numerical as well as the analytical solutions of the gBF equation.

Further, a compact finite difference method and a polynomial-based quadrature method were proposed by Sari *et al.* in order to approximate the gBF equation [26], [27]. A local discontinuous Galerkin (LDG) methods and optimal homotopy asymptotic method (OHAM) have been employed by Zhang *et al.* [28] and Nawaz *et al.* [29], respectively. Galerkin finite element method was employed by Yadav and Jiwari [30] to approximate and analyze the Burgers-Fisher equation. Malik *et al.* [31] numerically simulate the gBF equation using Exp-function method.

The most recent work on the gBF equation is referred to [32-39]. The research articles [40-44] and the references therein provide further information on the pseudospectral approach for both smooth and non-smooth functions.

Within the past few years, the spectral method has gained popularity as a reliable scientific computing tool for numerically solving linear and strongly non-linear partial differential equations [45–48]. The method's high-order accuracy has led to its widespread use. Interpolation and non-interpolation are the broad categories into which the spectral method has been classified. The pseudospectral method, also known as the spectral collocation method, depends on the interpolation methodology. The pseudospectral method forms a class of techniques that are efficient and highly accurate and having the potential for rapidly convergent approximations and low error estimation. This article will examine the gBF equation. In 2006, an approximation of the gBF equation using a modified version of the pseudospectral approach was given by Javidi [49] with tremendous accuracy in which the pseudospectral method was applied in the space direction only. For the numerical approximation of the gBF equation, we offer a spectrally accurate enhanced version of the Chebyshev pseudospectral approach that takes place in time as well as in space, and the results obtained using the suggested approach have proved that time is just as significant as it utilizes in space. To the best of the author's knowledge, this is the first work to employ the pseudospectral technique for the approximation of the gBF equation in both time and space.

The rest of the article is arranged as follows: We provide an in-depth explanation of the suggested approach as well as a clear image of how the discretization process works in Sect. 2. Analysis of the method's stability for the gBF equation is conducted in Sect. 3. The accuracy and efficiency of the suggested technique are shown in Sect. 4. Lastly, the observations and the concluding remarks are given in Sect. 5.

# 2 Chebyshev pseudospectral discretization of the gBF equation in both time and space

In this section, we discuss the Chebyshev pseudospectral approach to approximate the gBF equation in both time and space. We look for a pseudospectral approximation  $U(\eta, t)$  in the form of a finite linear combination of a product of Chebyshev polynomials with spectral coefficients such that

$$I_N U = \sum_{\alpha=0}^N \sum_{\beta=0}^N \phi_\alpha(\eta) \phi_\beta(t) \Theta_{\alpha\beta},$$
(5)

where, in space and time directions, Chebyshev polynomials are  $\phi_{\alpha}(\eta)$  and  $\phi_{\beta}(t)$  respectively and spectral coefficients  $\Theta_{\alpha\beta}$  are given by

$$\Theta_{\alpha\beta} = \frac{1}{\hat{h}_{\alpha}\hat{h}_{\beta}} \sum_{l=0}^{N} \sum_{m=0}^{N} \phi_{\alpha}(\eta_{l})\phi_{\beta}(t_{m})w_{l}w_{m}U(\eta_{\alpha}, t_{\beta}), \tag{6}$$

where  $\hat{h}_p = \frac{\epsilon_p}{2}\pi$  is a discrete normalizing factor,  $\epsilon_0 = 2$  and  $\epsilon_p = 1$  for  $p \ge 1$ . Where  $p = \{\alpha, \beta\}$  and w denotes the weight function. Also, we can write

$$I_{N}U = \sum_{\alpha=0}^{N} \sum_{\beta=0}^{N} \phi_{\alpha}(\eta)\phi_{\beta}(t)\Theta_{\alpha\beta},$$
  

$$= \sum_{\alpha=0}^{N} \sum_{\beta=0}^{N} \phi_{\alpha}(\eta)\phi_{\beta}(t)\frac{1}{\hat{h}_{\alpha}\hat{h}_{\beta}}\sum_{l=0}^{N} \sum_{m=0}^{N} \phi_{\alpha}(\eta_{l})\phi_{\beta}(t_{m})w_{l}w_{m}U(\eta_{\alpha}, t_{\beta}),$$
  

$$= \sum_{\alpha=0}^{N} \sum_{\beta=0}^{N} \left(\frac{1}{\hat{h}_{\alpha}\hat{h}_{\beta}}\sum_{l=0}^{N} \sum_{m=0}^{N} w_{l}\phi_{\alpha}(\eta_{l})\phi_{\alpha}(\eta)w_{m}\phi_{\beta}(t_{m})\phi_{\beta}(t)\right)U(\eta_{\alpha}, t_{\beta}),$$
  

$$= \sum_{\alpha=0}^{N} \sum_{\beta=0}^{N} \mathcal{L}_{\alpha}(\eta)\mathcal{L}_{\beta}(t)U(\eta_{\alpha}, t_{\beta}),$$
(7)

where  $\mathcal{L}$  denotes the Lagrange polynomials which can be written as

$$\mathcal{L}_p(z) = \frac{1}{\hat{h}_p} \sum_{r=0}^N w_r \phi_p(z_r) \phi_p(z) = \frac{(-1)^{p+1}(1-z^2)}{c_p N^2(z-z_p)} \frac{\partial}{\partial z} \phi_N(z),$$

where,  $p = \{\alpha, \beta\}, r = \{l, m\}, z = \{\eta, t\}$ , and

$$c_p = \begin{cases} 2, & p = 0 \text{ and } N, \\ 1, & 0$$

Λ	
4	

Here, the Lagrange polynomial satisfies the delta function i.e.  $\mathcal{L}_p(z_\beta) = \delta_{p\beta}$ . The differential matrix  $\mathcal{D}$  of first order is given by,

$$\mathcal{D}_{\alpha\beta} = \begin{cases} -\frac{2N^2+1}{6}, & \alpha = \beta = 0, \\ \frac{c_{\alpha}}{c_{\beta}} \frac{(-1)^{\alpha+\beta}}{z_{\alpha}-z_{\beta}}, & \alpha \neq \beta, \\ -\frac{z_{\alpha}}{2(1-z_{\alpha}^2)}, & \alpha = \beta \in [1,...,N-1], \\ \frac{2N^2+1}{6}, & \alpha = \beta = N \end{cases}$$
(8)

It is possible to write the spectral approximation given in Eq. 7 as a direct product

$$I_N U = [\mathcal{L}_0(\eta) \mathcal{L}_0(t), \dots \mathcal{L}_0(\eta) \mathcal{L}_N(t), \dots, \mathcal{L}_N(\eta) \mathcal{L}_0(t), \dots, \mathcal{L}_N(\eta) \mathcal{L}_N(t)] U,$$
  

$$= [\{\mathcal{L}_0(\eta), \dots, \mathcal{L}_N(\eta)\} \otimes \{\mathcal{L}_0(t), \dots, \mathcal{L}_N(t)\}] U,$$
  

$$= (\psi_{[0:N]}(\eta) \otimes \psi_{[0:N]}(t))^T U,$$
  

$$= (K_{[0:L]}(\eta, t))^T U.$$
(9)

where

$$\boldsymbol{U} = [U_{00}, ..., U_{0N} | U_{10}, ..., U_{1N} | ... | U_{N0}, ..., U_{NN} ]^T,$$
  
$$\psi_{[0:N]}(z) = [\mathcal{L}_0(z), ..., \mathcal{L}_N(z)]^T, \quad \text{and} \quad \boldsymbol{L} = (N+1) \times (N+1) - 1.$$

The first and second-order derivatives of U with respect to  $`\eta`$  are now determined using Eq. 9

$$\frac{\partial U(\eta, t)}{\partial \eta} = (K_{[0:L]}(\eta, t))^T (\mathcal{D}'_{[0:N,0:N]} \otimes I_{N+1}) \boldsymbol{U},$$
(10)

$$\frac{\partial^2 U(\eta, t)}{\partial \eta^2} = (K_{[0:L]}(\eta, t))^T (\mathcal{D}''_{[0:N,0:N]} \otimes I_{N+1}) U,$$
(11)

where  $\mathcal{D}'_{[0:N,0:N]}$  and  $\mathcal{D}''_{[0:N,0:N]}$  denotes the first and the second order differential matrices. Moreover, the first derivative with respect to 't' is given by

$$\frac{\partial U(\eta, t)}{\partial t} = (K_{[0:L]}(\eta, t))^T (I_{N+1} \otimes \mathcal{D}^t_{[0:N,0:N]}) \boldsymbol{U}.$$
(12)

Now, we will discretize the gBF equation based on discussed pseudospectral method and in order to apply the method the idea is to convert the space and the time intervals into [-1, 1] and therefore to serve the purpose we use the following transformation for any  $\tilde{v} \in [I_L, I_R]$ 

$$\tilde{v} \rightarrow \frac{I_R - I_L}{2}v + \frac{I_R + I_L}{2}.$$

By using the transformations in both space and time, we obtain the following generalized Burgers-Fisher equation in new time and space intervals

$$U_t + A(U^{\delta+1})_{\eta} - BU_{\eta\eta} - C(U - U^{\delta+1}) = 0,$$
(13)

with the initial condition

$$U(\eta, -1) = h(\eta), \qquad \eta \in [-1, 1],$$
 (14)

and the associated boundary conditions

$$U(-1,t) = g_1(t), \quad U(1,t) = g_2(t), \quad t \in [-1,1],$$
(15)

where

$$A = \frac{\sigma_1 T}{(\delta + 1)(\eta_R - \eta_L)}, \ B = \frac{2T}{(\eta_R - \eta_L)^2}, \ \text{and} \ C = \frac{\sigma_2 T}{2}.$$
 (16)

Further, to make it easier to solve, a mapping is presented for changing a system's initial and boundary values from non-homogeneous to homogeneous,

$$\Omega(\eta, t) = \frac{(1-t)}{2}h(\eta) + \frac{(1-\eta)}{2}g_1(t) + \frac{(1+\eta)}{2}g_2(t) - \frac{(1-t)(1-\eta)}{4}g_1(-1) - \frac{(1-t)(1+\eta)}{4}g_2(-1),$$
(17)

where at the corners of the domain the initial-boundary conditions are supposed to satisfy  $h(-1) = g_1(-1)$  and  $h(1) = g_2(-1)$ . Now, we have transformed our non-homogeneous initial-boundary value problem into homogeneous one. Let us define a variable V, such that

$$U(\eta, t) = V(\eta, t) + \Omega(\eta, t).$$
(18)

Using Eq. 18, Eqs. 13-16 can be written as

$$(V(\eta, t) + \Omega(\eta, t))_t + A[(V(\eta, t) + \Omega(\eta, t))^{\delta+1}]_\eta - B(V(\eta, t) + \Omega(\eta, t))_{\eta\eta} - C[(V(\eta, t) + \Omega(\eta, t)) - (V(\eta, t) + \Omega(\eta, t))^{\delta+1}] = 0,$$
(19)

with the following homogeneous initial and boundary conditions

$$V(\eta, -1) = 0, \quad \eta \in [-1, 1],$$
(20)

$$V(-1,t) = 0, V(1,t) = 0, t \in [-1,1],$$
 (21)

Using Binomial expansion in Eq. 19, we get

$$(V(\eta,t) + \Omega(\eta,t))_{t} + A[^{\delta+1}c_{0}V^{\delta+1}(\eta,t)\Omega^{0}(\eta,t) + {}^{\delta+1}c_{1}V^{\delta}(\eta,t)\Omega^{1}(\eta,t) + \dots + {}^{\delta+1}c_{\delta}V^{1}(\eta,t)\Omega^{\delta}(\eta,t) + {}^{\delta+1}c_{\delta+1}V^{0}(\eta,t)\Omega^{\delta+1}(\eta,t)]_{\eta} - B(V(\eta,t) + \Omega(\eta,t))_{\eta\eta} - C(V(\eta,t) + \Omega(\eta,t)) + C[{}^{\delta+1}c_{0}V^{\delta+1}(\eta,t)\Omega^{0}(\eta,t) + {}^{\delta+1}c_{1}V^{\delta}(\eta,t)\Omega^{1}(\eta,t) + \dots + {}^{\delta+1}c_{\delta}V^{1}(\eta,t)\Omega^{\delta}(\eta,t) + {}^{\delta+1}c_{\delta+1}V^{0}(\eta,t)\Omega^{\delta+1}(\eta,t)] = 0.$$

$$(22)$$

Now, as described in detail at the beginning of this section, we look for an approximation that can be defined in vector-vector product form, therefore, the spectral

approximation  $V(\eta, t)$  is defined as

$$\mathbf{V}(\eta, t) = (\psi_{[1:N-1]}(\eta) \otimes \psi_{[1:N]}(t))^T V, 
\mathbf{V}(\eta, t) = (K_{[1:(N-1)N]}(\eta, t))^T V,$$
(23)

where V is a  $(N-1) \times N$  vector, given by

$$V = [V_{11}, ..., V_{1N} | V_{21}, ..., V_{2N} | ... | V_{(N-1)1}, ..., V_{(N-1)N}]^T.$$

Additionally, we can define the following derivatives of  $V(\eta, t)$ 

$$\boldsymbol{V}_t(\eta, t) = (K_{[1:(N-1)N]}(\eta, t))^T (I_N \otimes \mathcal{D}^t) V,$$
(24)

$$\boldsymbol{V}_{\eta}(\eta, t) = (K_{[1:(N-1)N]}(\eta, t))^{T} (\mathcal{D}^{\eta'} \otimes I_{N}) V, \qquad (25)$$

$$\boldsymbol{V}_{\eta\eta}(\eta,t) = (K_{[1:(N-1)N]}(\eta,t))^T (\mathcal{D}^{\eta^{\prime\prime}} \otimes I_N) V, \qquad (26)$$

where  $\mathcal{D}^t = \mathcal{D}^t_{[1:N,1:N]}$  is the first order differential matrix w.r.t. 't', and  $\mathcal{D}^{\eta'} = \mathcal{D}^{\eta'}_{[1:N-1,1:N-1]}$  and  $\mathcal{D}^{\eta''} = \mathcal{D}^{\eta''}_{[1:N-1,1:N-1]}$  are the first and the second order differential matrices with respect to '\eta', respectively. By using Eqs. 23-26 in Eq. 22, we obtain

$$(K_{[1:(N-1)N]}(\eta,t))^{T} \Big\{ (I_{N-1} \otimes \mathcal{D}^{t})V + A(\mathcal{D}^{\eta'} \otimes I_{N})V^{\delta+1} + A^{\delta+1}c_{1} \left[ (\operatorname{diag}(\Omega_{\eta}(\eta,t))) + \operatorname{diag}(\Omega(\eta,t))(\mathcal{D}^{\eta} \otimes I_{N}) \right]V^{\delta} + \ldots + A^{\delta+1}c_{\delta} \left[ (\operatorname{diag}((\Omega^{\delta}(\eta,t))_{\eta})) + \operatorname{diag}(\Omega^{\delta}(\eta,t))(\mathcal{D}^{\eta} \otimes I_{N}) \right]V - B(\mathcal{D}^{\eta''} \otimes I_{N})V + C[-(I_{N-1} \otimes I_{N})V + (I_{N-1} \otimes I_{N})V^{\delta+1} + {}^{\delta+1}c_{1} \left(\operatorname{diag}(\Omega(\eta,t)))V^{\delta} + \ldots + {}^{\delta+1}c_{\delta} \left(\operatorname{diag}(\Omega^{\delta}(\eta,t)))V \right] \Big\} + \Omega_{t}(\eta,t) + A(\Omega^{\delta+1}(\eta,t))_{\eta} - B\Omega_{\eta\eta}(\eta,t) - C\Omega(\eta,t) + C\Omega^{\delta+1}(\eta,t) = 0.$$
(27)

Further, in Eq. 27, using Chebyshev Gauss Lobbatto points in  $\eta$  and t directions such that  $(\eta, t) = (\eta_i, t_j)$ , i = 1, ..., N - 1, and j = 1, ..., N, and using the fact that the Lagrange polynomial satisfies the properties of a delta function and hence  $(K_{[1:(N-1)N]}(\eta_i, t_j))^T = [\psi_{[1:N-1]}(\eta_i) \otimes \psi_{[1:N]}(t_j)]^T$  gives a vector  $V_r^T$  of order  $(N - 1) \times N$  in which except the  $r^{th}$  element rest are zero. Therefore, we obtain

$$\begin{aligned} (I_{N-1} \otimes \mathcal{D}^{t})V + A(\mathcal{D}^{\eta'} \otimes I_{N})V^{\delta+1} + A^{\delta+1}c_{1} \left[ (\operatorname{diag}(\Omega_{\eta}(\eta_{i}, t_{j}))) \right. \\ &+ \operatorname{diag}(\Omega(\eta_{i}, t_{j}))(\mathcal{D}^{\eta} \otimes I_{N}) V^{\delta} + \ldots + A^{\delta+1}c_{\delta} \left[ (\operatorname{diag}(\delta\Omega^{\delta-1}(\eta_{i}, t_{j})\Omega_{\eta}(\eta_{i}, t_{j}))) \right. \\ &+ \operatorname{diag}(\Omega^{\delta}(\eta_{i}, t_{j}))(\mathcal{D}^{\eta} \otimes I_{N}) V - B(\mathcal{D}^{\eta''} \otimes I_{N})V \\ &+ C[-(I_{N-1} \otimes I_{N})V + (I_{N-1} \otimes I_{N})V^{\delta+1} + {}^{\delta+1}c_{1} \left(\operatorname{diag}(\Omega(\eta_{i}, t_{j})))V^{\delta} \right] \end{aligned}$$

$$+ \dots + {}^{\delta+1}c_{\delta} \left(\operatorname{diag}(\Omega^{\delta}(\eta_i, t_j)))V\right] + \Omega_t(\eta_i, t_j) + (\delta + 1)A(\Omega^{\delta}(\eta_i, t_j))(\Omega_\eta(\eta_i, t_j)) - B\Omega_{\eta\eta}(\eta_i, t_j) - C\Omega(\eta_i, t_j) + C\Omega^{\delta+1}(\eta_i, t_j) = 0.$$

$$(28)$$

In conclusion, the equation system 28 can be converted into the matrix form as follows

$$(I_{N-1} \otimes \mathcal{D}^{t})V + A(\mathcal{D}^{\eta'} \otimes I_{N})V^{\delta+1} + A^{\delta+1}c_{1} \ [(\operatorname{diag}(\mathbf{Q})) + \operatorname{diag}(\mathbf{P})(\mathcal{D}^{\eta} \otimes I_{N})]V^{\delta} + \dots + A^{\delta+1}c_{\delta} \ [(\operatorname{diag}(\delta\mathbf{P}^{\delta-1}\mathbf{Q}) + \operatorname{diag}(\mathbf{P}^{\delta})(\mathcal{D}^{\eta} \otimes I_{N})]V - B(\mathcal{D}^{\eta''} \otimes I_{N})V + C[-(I_{N-1} \otimes I_{N})V + (I_{N-1} \otimes I_{N})V^{\delta+1} + {}^{\delta+1}c_{1} \ (\operatorname{diag}(\mathbf{P}))V^{\delta} + \dots + {}^{\delta+1}c_{\delta} \ (\operatorname{diag}(\mathbf{P}^{\delta}))V] + \mathbf{S} + (\delta+1)A\mathbf{P}^{\delta}\mathbf{Q} - B\mathbf{R} - C\mathbf{P} + C\mathbf{P}^{\delta+1} = 0,$$
(29)

where

$$V = [V_{11}, ..., V_{1N} | V_{21}, ..., V_{2N} | ... | V_{(N-1)1}, ..., V_{(N-1)N}]^{T},$$
  

$$\mathbf{P} = [\Omega(\eta_{1}, t_{1}), ..., \Omega(\eta_{1}, t_{N}) | \Omega(\eta_{2}, t_{1}), ..., \Omega(\eta_{2}, t_{N}) | ... | \Omega(\eta_{N-1}, t_{1}), ..., \Omega(\eta_{N-1}, t_{N})]^{T},$$
  

$$\mathbf{Q} = [\Omega_{\eta}(\eta_{1}, t_{1}), ..., \Omega_{\eta}(\eta_{1}, t_{N}) | \Omega_{\eta}(\eta_{2}, t_{1}), ..., \Omega_{\eta}(\eta_{2}, t_{N}) | ... | \Omega_{\eta}(\eta_{N-1}, t_{1}), ..., \Omega_{\eta}(\eta_{N-1}, t_{N})]^{T},$$
  

$$\mathbf{R} = [\Omega_{\eta\eta}(\eta_{1}, t_{1}), ..., \Omega_{\eta\eta}(\eta_{1}, t_{N}) | \Omega_{\eta\eta}(\eta_{2}, t_{1}), ..., \Omega_{\eta\eta}(\eta_{2}, t_{N}) | ... | \Omega_{\eta\eta}(\eta_{N-1}, t_{1}), ..., \Omega_{\eta\eta}(\eta_{N-1}, t_{N})]^{T},$$
  

$$\mathbf{S} = [\Omega_{t}(\eta_{1}, t_{1}), ..., \Omega_{t}(\eta_{1}, t_{N}) | \Omega_{t}(\eta_{2}, t_{1}), ..., \Omega_{t}(\eta_{2}, t_{N}) | ... | \Omega_{t}(\eta_{N-1}, t_{1}), ..., \Omega_{t}(\eta_{N-1}, t_{N})]^{T}$$
  
(30)

Eq. 29 is the discretized form of Eqs. 19-21, using the Chebyshev pseudospectral method.

Finally, we have

$$G(V) = 0. \tag{31}$$

The solution to the nonlinear equation system  $31~{\rm may}$  be obtained by employing the Newton-Raphson technique

$$V_{n+1} = V_n - [J_n]^{-1} G(V_n), (32)$$

with the following Jacobian matrix

$$J = \begin{bmatrix} \left| \frac{\partial G_{11}}{\partial V_{11}} \cdots \frac{\partial G_{11}}{\partial V_{1N}} & \frac{\partial G_{11}}{\partial V_{21}} \cdots \frac{\partial G_{11}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{11}}{\partial V_{(N-1)1}} \cdots \frac{\partial G_{11}}{\partial V_{(N-1)N}} \right| \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & |\cdots| & \vdots & \ddots & \ddots \\ \frac{\partial G_{1N}}{\partial V_{11}} & \cdots & \frac{\partial G_{1N}}{\partial V_{1N}} & \frac{\partial G_{1N}}{\partial V_{21}} \cdots & \frac{\partial G_{1N}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{1N}}{\partial V_{(N-1)1}} & \cdots & \frac{\partial G_{1N}}{\partial V_{(N-1)N}} \\ \frac{\partial G_{21}}{\partial V_{11}} & \cdots & \frac{\partial G_{21}}{\partial V_{1N}} & \frac{\partial G_{21}}{\partial V_{21}} \cdots & \frac{\partial G_{21}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{21}}{\partial V_{(N-1)1}} & \cdots & \frac{\partial G_{21}}{\partial V_{(N-1)N}} \\ \vdots & \vdots & \ddots & \ddots & \ddots & \ddots & |\cdots| & \ddots & \ddots \\ \frac{\partial G_{2N}}{\partial V_{11}} & \cdots & \frac{\partial G_{2N}}{\partial V_{2N}} & \frac{\partial G_{2N}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{2N}}{\partial V_{(N-1)1}} & \cdots & \frac{\partial G_{2N}}{\partial V_{(N-1)N}} \\ \vdots & \vdots \\ \frac{\partial G_{(N-1)1}}{\partial V_{11}} & \cdots & \frac{\partial G_{(N-1)1}}{\partial V_{1N}} & \frac{\partial G_{(N-1)1}}{\partial V_{21}} & \cdots & \frac{\partial G_{(N-1)1}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{(N-1)1}}{\partial V_{(N-1)1}} & \cdots & \frac{\partial G_{(N-1)N}}{\partial V_{(N-1)N}} \\ \vdots & \ddots & \ddots & \ddots & \ddots & \ddots & |\cdots| & \frac{\partial G_{(N-1)N}}{\partial V_{(N-1)N}} & \frac{\partial G_{(N-1)N}}{\partial V_{21}} & \cdots & \frac{\partial G_{(N-1)N}}{\partial V_{2N}} & |\cdots| & \frac{\partial G_{(N-1)N}}{\partial V_{(N-1)1}} & \cdots & \frac{\partial G_{(N-1)N}}{\partial V_{(N-1)N}} \\ \end{bmatrix} \right]$$

where the order of the Jacobian matrix is  $(N-1)N \times (N-1)N$ .

## 3 Stability analysis of the gBF equation

In order to conduct a study of the method's stability for the gBF equation, we make use of the relation between the collocation points and the Gauss-type integration formulas. These quadrature formulae have the characteristics that let us make the transition from summing to integrating. The gBF equation is given by

$$U_t(\eta, t) + A(U^{\delta+1}(\eta, t))_\eta - BU_{\eta\eta}(\eta, t) - CF(U(\eta, t)) = 0.$$
 (33)

Multiplying Eq. 33 by  $U(\eta_{\alpha}, t)$  with the weight function  $w(\eta_{\alpha}) = \frac{1}{\sqrt{(1-\eta_{\alpha}^2)}}$  and summing, we have

$$\sum_{\alpha=0}^{N} U(\eta_{\alpha}, t) w(\eta_{\alpha}) (U_t(\eta_{\alpha}, t) + A(U^{\delta+1}(\eta_{\alpha}, t))_{\eta} - BU_{\eta\eta}(\eta_{\alpha}, t) - CF(U(\eta_{\alpha}, t))) = 0,$$
(34)

where A, B, and C are constants. Eq. 34 can be written as

$$\frac{1}{2} \frac{d}{dt} \sum_{\alpha=0}^{N} U^{2}(\eta_{\alpha}, t) w(\eta_{\alpha}) = -A \sum_{\alpha=0}^{N} U(\eta_{\alpha}, t) (U^{\delta+1}(\eta_{\alpha}, t))_{\eta} w(\eta_{\alpha}) 
+ B \sum_{\alpha=0}^{N} U(\eta_{\alpha}, t) U_{\eta\eta}(\eta_{\alpha}, t) w(\eta_{\alpha}) + C \sum_{\alpha=0}^{N} U(\eta_{\alpha}, t) F(U(\eta_{\alpha}, t)) w(\eta_{\alpha}).$$
(35)

The first two terms of the right hand side in Eq. 35 can be shown easily non-positive

using integration by parts as follows: Let  $wU = z \Rightarrow U_{\eta} = \frac{(wU)_{\eta}}{w} - \frac{w_{\eta}U}{w} = \frac{z_{\eta}}{w} + z(\frac{1}{w})_{\eta}$ , which leads to the following result

$$\sum_{\alpha=0}^{N} U(\eta_{\alpha}, t) U_{\eta\eta}(\eta_{\alpha}, t) w(\eta_{\alpha}) = \int_{-1}^{1} w(\eta) U(\eta, t) U_{\eta\eta}(\eta, t) d\eta$$
$$= -\int_{-1}^{1} z_{\eta} (\frac{z_{\eta}}{w} + z(\frac{1}{w})_{\eta}) d\eta$$
$$= -\int_{-1}^{1} \frac{z_{\eta}^{2}}{w} d\eta + \frac{1}{2} \int_{-1}^{1} z^{2} (\frac{1}{w})_{\eta\eta} d\eta \leq 0.$$
(36)

Hence the Eq. 35 becomes

$$\frac{1}{2}\frac{d}{dt}\sum_{\alpha=0}^{N}U^{2}(\eta_{\alpha},t)w(\eta_{\alpha}) \leq C\sum_{\alpha=0}^{N}U(\eta_{\alpha},t)F(U(\eta_{\alpha},t))w(\eta_{\alpha}).$$
(37)

Since, we have  $F(U(\eta,t))=U(\eta,t)(1-U^{\delta}(\eta,t))$  and hence the above Eq. 37 transformed into the following equation

$$\begin{split} \frac{1}{2} \frac{d}{dt} \sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha}) &\leq C \sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha}) - C \sum_{\alpha=0}^{N} U^{\delta+2}(\eta_{\alpha}, t) w(\eta_{\alpha}), \\ \Rightarrow \frac{d}{dt} \sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha})}{\sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha})} &\leq 2C - 2C \frac{\sum_{\alpha=0}^{N} U^{\delta+2}(\eta_{\alpha}, t) w(\eta_{\alpha})}{\sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha})} \leq 2C, \\ \Rightarrow \log(\sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha})) \leq 2Ct + \log(k). \end{split}$$

Further simplifying, we get

$$\sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha}) \le k \exp(2Ct).$$

Finally using the initial condition, we obtain

$$\sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t) w(\eta_{\alpha}) \leq \exp(2Ct) \sum_{\alpha=0}^{N} U^2(\eta_{\alpha}, t_0) w(\eta_{\alpha}),$$

hence conclude the result.

## 4 Numerical findings and discussion

In order to illustrate the performance, we consider Eq. 4 with the finite spatial and time intervals with the following initial condition

$$U(\tilde{\eta}, 0) = \left(\frac{1}{2} + \frac{1}{2} \tanh(U_1 \tilde{\eta})\right)^{\frac{1}{\delta}}, \quad 0 \le \tilde{\eta} \le 1,$$
(38)

and boundary conditions

$$U(0,\tilde{t}) = \left(\frac{1}{2} + \frac{1}{2}\tanh(U_1 U_2 \tilde{t})\right)^{\frac{1}{\delta}}, \quad \tilde{t} \ge 0,$$
(39)

$$U(1,\tilde{t}) = \left(\frac{1}{2} + \frac{1}{2}\tanh(U_1(1 - U_2\tilde{t}))\right)^{\frac{1}{\delta}}, \quad \tilde{t} \ge 0.$$
(40)

The exact solution [13] of Eq. 4 is

$$U(\tilde{\eta}, \tilde{t}) = \left(\frac{1}{2} + \frac{1}{2} \tanh\left(U_1(\tilde{\eta} - U_2\tilde{t})\right)\right)^{\frac{1}{\delta}}, \quad 0 \le \tilde{\eta} \le 1, \quad \tilde{t} \ge 0,$$
(41)

where  $U_1$  and  $U_2$  are given by

$$U_1 = \frac{-\sigma_1\delta}{2(\delta+1)}, \quad U_2 = \left(\frac{\sigma_1}{\delta+1} + \sigma_2\frac{\delta+1}{\sigma_1}\right).$$

The accuracy and efficiency of the technique have been verified for the distinct values of constants  $\sigma_1$  and  $\sigma_2$  with different sets of time and various values of  $\delta$ , all of which yielded excellent and encouraging numerical results. To express the error,  $L_{\infty}$ -norm has been calculated at different grid points which can be defined as

$$L_{\infty} = \max\{|U_{i,j}^{exa} - U_{i,j}^{num}| : i = 1, 2, ..., N - 1, \ j = 1, 2, ..., N\}$$

where  $U^{exa}$  and  $U^{num}$  denotes the exact and approximated solutions respectively.

#### 4.1 Numerical examples for the gBF equation

#### Example 1.

In this example, gBF equation 4 is approximated for particular small and negative values  $\sigma_1 = 0.1$ , and  $\sigma_2 = -0.0025$  using TS-CPsM. Increasing values of  $\delta$  make the equation highly non-linear, which is taken care of by the proposed method with excellent accuracy. At a very small number of grid points, we have obtained highly accurate results at different times in terms of the error infinity norm, which are efficient compared to those existing in the literature [18]. The calculated results have been verified for different sets of values of  $\delta$  and t, and the same is reported in Table 1. Results show that for a very small number of grid points N = 6 in both space and

time directions, we achieved very good accuracy. Figure 1 shows a comparison of the numerical and exact solutions for  $\delta = 1, 2, 4$ , and 8 at various time values. A solid line indicates an exact solution, whereas a block indicates a numerical result at a certain time step. Further, 3D graphs for exact vs approximate solution at time T = 0.3 for  $\delta = 1, 2, 4$ , and 8, respectively, are given in Figure 2.




Fig. 1 Numerical vs Exact solution of  $U(\eta, t)$  for Example 1 when different time T = 0.1, 0.2, 0.3, 0.4 and 0.5 for  $\delta = 1, 2, 4$ , and 8, respectively.

#### Example 2.

In the second example of the queue the considered Eq. 4 is approximated using TS-CPsM for a new set of positive and equal values  $\sigma_1 = 1$ , and  $\sigma_2 = 1$ . As may be seen from Table 2, with an increase in grid points, the accuracy is getting higher and higher. Here, we found highly accurate results for 16 grid points in both space and




Fig. 2 3D graphs for exact (left) vs approximate (right) solution for Example 1 at time T=0.3 for  $\delta=1,\,2,\,4,$  and 8, respectively.

**Table 1** Numerical solution proposed forExample 1 with different time t for various values of<br/> $\delta$  and grid points N.

	NT 4	NL C	
t	N=4	N=6	DCOL [10]
	TS-CPsM	TS-CPsM	BSQI [18]
0.1			
$\delta = 1$	9.5696e-13	5.5511e-17	1.3240e-11
$\delta = 2$	1.4242e-12	4.5519e-15	2.8470e-10
$\delta = 4$	9.9620e-13	1.1102e-16	3.9917e-10
$\delta = 8$	4.7429e-13	1.1102e-16	-
0.2			
$\delta = 1$	1.0758e-12	5.5511e-17	1.7803e-11
$\delta = 2$	1.6033e-12	2.2204e-16	3.8795e-10
$\delta = 4$	1.1265e-12	1.1102e-16	5.4380e-10
$\delta = 8$	5.4279e-13	1.1102e-16	-
0.3			
$\delta = 1$	1.1146e-12	5.5511e-17	1.9426e-11
$\delta = 2$	1.6627 e-12	1.1102e-16	4.2465e-10
$\delta = 4$	1.1720e-12	1.1102e-16	5.9517e-10
$\delta = 8$	5.6999e-13	1.1102e-16	-
0.4			
$\delta = 1$	1.1343e-12	5.5511e-17	2.0008e-11
$\delta = 2$	1.6921e-12	1.1102e-16	4.3759e-10
$\delta = 4$	1.1960e-12	1.1102e-16	6.1323e-10
$\delta = 8$	5.8642e-13	1.1102e-16	-
0.5			
$\delta = 1$	1.1478e-12	5.5511e-17	2.0216e-11
$\delta = 2$	1.7099e-12	1.1102e-16	4.0205e-10
$\delta = 4$	1.2114e-12	1.1102e-16	6.1941e-10
$\delta = 8$	5.9841e-13	1.1102e-16	-

time; however, the literature says that a large number of grid points in the temporal direction is required to get good results. A further observation is that there is a slight decay in accuracy as the delta values are increased for a large time, though the results are still comparably better and quite acceptable. Graphical outcomes of approximated solutions with the exact solutions for different sets of values of time with varying  $\delta$  are shown in Figure 3. A solution that is exact is shown by a solid line, whereas numerical results at discrete times are shown as blocks. Further, 3D graphs for exact vs approximate solution at time T = 0.6 for  $\delta = 1, 2, 4$ , and 8, respectively, are given in Figure 4.

#### Example 3.

In this demonstration, a different combination of the variables  $\sigma_1 = 1$  and  $\sigma_2 = 0$ is taken into account, which results in the transformation of the gBF equation into the more straightforward Burgers equation. The importance and use of the equation have already been discussed in the introduction part. The error infinity norm for increasing number of grid points is shown in Table 3 for a different set of values of  $\delta$  and time t. Figure 5 shows a comparison of the numerical and exact solutions for  $\delta = 1$ , 2, and 4. A solid line indicates an exact solution, whereas a block indicates a numerical result at a certain time step. Further, 3D graphs for exact vs approximate solution at time T = 1.5 for  $\delta = 1$ , 2, and 4, respectively, is given in Figure 6.

	N=4	N=6	N=8	N=10	N=12	N=14	N=16		
	TS-CP <sub>s</sub> M	$TS-CP_{SM}$	$TS-CP_{SM}$	$TS-CP_{SM}$	$TS-CP_{SM}$	$TS-CP_{SM}$	TS-CPsM	BSQI [18]	SCM [49]
t=0.2									
$\delta = 1$	1.094e-07	1.363e-10	1.665e-13	2.220e-16	1.110e-16	1.110e-16	1.110e-16	5.558e-7	8.749e-12
$\delta = 2$	1.430e-07	2.267e-10	3.968e-13	8.882e-16	1.110e-16	1.110e-16	2.220e-16	2.561e-6	5.812e-11
$\delta = 4$	1.722e-07	1.145e-09	6.861e-12	3.775e-14	2.220e-16	1.110e-16	1.110e-16	1.762e-6	1.397e-10
$\delta = 8$	2.574e-06	8.626e-08	2.139e-09	4.481e-11	8.373e-13	1.421e-14	2.220e-16	I	2.972e-10
t=0.4									
$\delta = 1$	1.008e-07	2.416e-10	6.424e-13	1.499e-15	1.110e-16	1.110e-16	1.110e-16	9.055e-7	1.667e-1
$\delta = 2$	4.296e-07	3.601e-09	2.289e-11	1.376e-13	6.661e-16	2.220e-16	2.220e-16	4.243e-6	6.388e-11
$\delta = 4$	3.209e-06	1.508e-07	4.579e-09	1.098e-10	2.264e-12	4.152e-14	6.661 e - 16	4.174e-7	1.365e-10
$\delta = 8$	5.131e-05	3.319e-06	7.566e-08	1.537e-08	1.609e-09	9.949e-11	4.360e-12	ı	2.169e-10
t=0.6									
$\delta = 1$	3.683e-07	2.995e-09	$1.816e{-}11$	9.320e-14	4.441e-16	1.110e-16	1.110e-16	2.188e-6	2.274e-11
$\delta = 2$	5.567e-07	1.280e-08	4.287e-10	1.083e-11	2.0140e-13	3.220e-15	2.220e-16	3.569e-6	6.190e-1
$\delta = 4$	2.722e-05	2.085e-06	9.336e-08	2.570e-09	$3.230e{-}11$	$3.884e{-}12$	2.531e-13	2.424e-6	9.537e-11
$\delta = 8$	5.763e-05	1.164e-05	3.690e-06	3.129e-07	1.654e-08	5.834e-09	6.217e-10		2.722e-11
t=0.8									
$\delta = 1$	9.945e-07	1.384e-08	1.356e-10	1.047e-12	6.939e-15	1.110e-16	1.110e-16	2.933e-6	2.655e-11
$\delta = 2$	3.882e-06	2.548e-07	1.141e-08	3.577e-10	8.942e-12	1.873e-13	3.331e-15	1.465e-6	6.153e-11
$\delta = 4$	6.466e-05	5.138e-06	1.557e-07	3.309e-08	3.627e-09	2.205e-10	7.847e-12	2.358e-6	5.230e-11
$\delta = 8$	8.497e-05	5.466e-05	7.401e-06	1.286e-06	4.016e-07	1.778e-08	8.669e-09		5.573e-12
t=1.0									
$\delta = 1$	1.811e-06	3.257e-08	3.246e-10	2.000e-12	6.017e-14	1.499e-15	1.110e-16	3.015e-6	2.813e-11
$\delta = 2$	1.296e-05	1.245e-06	6.765e-08	2.454e-09	6.029e-11	7.717e-13	3.853e-14	5.542e-6	5.320e-11
$\delta = 4$	8.701e-05	5.130e-06	1.653e-06	2.581e-07	1.563e-08	6.608e-10	1.886e-10	1.444e-6	2.533e-1
$\delta = 8$	1.431e-04	8 879e-05	0 084-06	6 545-06	1.0350-07	9 768a-07	1 00.9 - 08		1 111a-19

**Table 2** Numerical solution proposed for Example 2 with different time t for various values of  $\delta$  and grid points N.







Fig. 3 Numerical vs Exact solution of  $U(\eta, t)$  for Example 2 when different time T = 0.2, 0.4, 0.6, 0.8 and 1 for  $\delta = 1, 2, 4$ , and 8, respectively.

#### Example 4.

Consideration of a different set of time values has been found in the literature [18], [50] for the parameters  $\sigma_1 = 1$  and  $\sigma_2 = 1$  and the approximation for the same set is made and compared in this example. Table 4 displays the error infinity norms for varying  $\delta$  and t. As can be seen from this table, with an increase in the grid points N, the error is decreasing and the results are getting more accurate. A comparison with the existing results shows the efficiency and robustness of the proposed technique. The comparison between the exact and approximated results is depicted in Figure 7 for varying  $\delta$  at various time values t.










Fig. 4 3D graphs for exact (left) vs approximate (right) solution for Example 2 at time T=0.6 for  $\delta=1,\,2,\,4,$  and 8, respectively.




Fig. 5 Numerical vs Exact solution for Example 3 when different time T = 0.001, 0.5, 1, 1.5, 2 and 2.5 for  $\delta = 1, 2, \text{ and } 4$ , respectively.




Fig. 6 3D graphs for exact(left) vs approximate(right) solution for Example 3 at time T = 1.5 for  $\delta = 1, 2, 4$ , and 8, respectively. 19

	N=4	N=6	N=8	N=10	
	TS-CPsM	TS-CPsM	TS-CPsM	TS-CPsM	SCM [49]
t=0.001					
$\delta = 1$	2.7715e-09	1.1814e-11	3.5527 e- 14	1.1102e-16	6.5918e-13
$\delta = 2$	5.8538e-09	4.3071e-11	1.9740e-13	7.7716e-16	4.0404e-13
$\delta = 4$	5.8482e-09	6.1648e-11	4.0001e-13	1.7764e-15	1.8770e-13
t = 0.5					
$\delta = 1$	1.1039e-07	1.3468e-10	1.5776e-13	2.2204e-16	1.7366e-13
$\delta = 2$	1.9814e-07	3.6723e-10	7.2031e-13	1.5543e-15	4.5848e-13
$\delta = 4$	1.9663e-07	4.7425e-10	1.3300e-12	3.7748e-15	1.9439e-13
t=1					
$\delta = 1$	1.1644e-07	1.4060e-10	1.6886e-13	2.2204e-16	3.5012e-13
$\delta = 2$	1.8895e-07	3.3341e-10	6.5703e-13	1.4433e-15	5.0652e-13
$\delta = 4$	1.8534e-07	4.4473e-10	1.2240e-12	3.6637e-15	1.9994e-13
t = 1.5					
$\delta = 1$	1.1146e-07	1.3011e-10	1.5110e-13	2.2204e-16	5.1602e-13
$\delta = 2$	1.7174e-07	2.9088e-10	5.3968e-13	1.1102e-15	5.4759e-13
$\delta = 4$	1.6998e-07	4.0300e-10	1.0719e-12	3.1086e-15	2.0447e-13
t=2					
$\delta = 1$	1.0571e-07	1.1338e-10	2.2149e-13	6.1062e-16	7.9885e-13
$\delta = 2$	1.4803e-07	2.2619e-10	3.9735e-13	7.7716e-15	5.8118e-13
$\delta = 4$	1.5113e-07	3.4518e-10	8.9762e-13	2.5535e-15	2.0795e-13
t = 2.5					
$\delta = 1$	9.9698e-08	2.3955e-10	1.2607e-12	5.5511e-15	6.6683e-13
$\delta = 2$	1.1989e-07	1.7146e-10	3.2796e-13	6.6613e-16	6.0701e-13
$\delta = 4$	1.2930e-07	2.7381e-10	6.8279e-13	1.9984e-15	2.1038e-13

**Table 3** Numerical solution proposed for Example 3 with different time t for various values of  $\delta$  and grid points N.

## 5 Conclusion

Using the time-space Chebyshev pseudospectral method (TS-CPsM), numerical solutions for the gBF equation with various values of advection and reaction coefficients are reported with tremendous accuracy in this study. It has been observed that for a very small number of grid points N and for large values of non-linear factor  $\delta$ , the approximated results for the considered equation using the suggested technique have been proven to be highly competent with exact solutions and comparably much better with those existing in the literature, including [17], [18], [22], [49], [51], and others referenced. The proposed method's stability analysis has been performed. Four different cases for the considered problem have been discussed in order to demonstrate the effectiveness and reliability of the method, and all of them give excellent and encouraging numerical results.

**Acknowledgments.** The Ministry of Human Resource Development of India provided financial assistance for this study, which the first author gratefully acknowledges.

Conflict of interest. The authors declare no conflict of interests.

	N=4			N=8			N=16		
	$TS-CP_{SM}$	BSQI [18]	SDD [50]	$TS-CP_{SM}$	BSQI [18]	SDD [50]	TS-CPsM	BSQI [18]	SDD [50]
t=0.3									
$\delta = 1$	1.096e-07	1.350e-6	1.476e-5	1.503e-13	5.773e-6	4.165e-11	1.110e-16	2.218e-6	1.184e-10
$\delta = 2$	1.665e-07		ı	3.332e-12	ı	·	2.220e-16	ı	·
$\delta = 4$	4.247e-07	1.230e-5	9.159e-6	1.765e-10	5.750e-5	1.285e-10	1.110e-16	3.274e-5	2.119e-10
$\delta = 8$	2.133e-05	1.638e-6	3.956e-6	4.988e-08	3.919e-6	8.851e-12	7.383e-14	4.151e-7	2.119e-10
t=0.9									
$\delta = 1$	1.400e-06	9.770e-6	4.110e-5	2.494e-10	4.384e-5	9.668e-11	1.110e-16	2.480e-5	5.109e-10
$\delta = 2$	7.276e-06		ı	3.124e-08	ı	ı	8.549e-15	ı	,
$\delta = 4$	7.917e-05	3.017e-6	1.189e-5	5.254e-07	1.342e-5	3.891e-11	2.268e-11	7.292e-6	7.681e-10
$\delta = 8$	1.058e-04	1.116e-7	4.334e-6	4.914e-06	4.923e-7	6.158e-12	2.730e-08	2.843e-7	7.016e-10

**Table 4** Numerical solution proposed for Example 4 with different time t for various values of  $\delta$  and grid points N.




Fig. 7 Numerical vs Exact solution for Example 4 when time T = 0.3 and 0.9 for  $\delta = 1, 2, 4$ , and 8, respectively.

## References

- Bateman, H.: Some recent researches on the motion of fluids. Monthly Weather Review 43(4), 163–170 (1915)
- [2] Burgers, J.M.: A mathematical model illustrating the theory of turbulence. Advances in Applied Mechanics, vol. 1, pp. 171–199. Elsevier (1948)
- [3] Greenshields, B.D., Bibbins, J., Channing, W., Miller, H.: A study of traffic capacity. In: Highway Research Board Proceedings, vol. 14, pp. 448–477 (1935). Washington, DC
- [4] Murray, J.: On burgers' model equations for turbulence. Journal of Fluid

Mechanics **59**(2), 263–279 (1973)

- [5] Hirsh, R.S.: Higher order accurate difference solutions of fluid mechanics problems by a compact differencing technique. Journal of computational physics 19(1), 90–109 (1975)
- Yepez, J.: An efficient quantum algorithm for the one-dimensional burgers equation. arXiv preprint quant-ph/0210092 (2002)
- [7] Fisher, R.A.: The wave of advance of advantageous genes. Annals of eugenics 7(4), 355–369 (1937)
- [8] Bramson, M.D.: Maximal displacement of branching brownian motion. Communications on Pure and Applied Mathematics 31(5), 531–581 (1978)
- [9] Britton, N.F., et al.: Reaction-diffusion Equations and Their Applications to Biology. Academic Press, ??? (1986)
- [10] Frank-Kamenetskii, D.A.: Diffusion and Heat Exchange in Chemical Kinetics vol. 2171. Princeton University Press, ??? (2015)
- [11] Canosa, J.: Diffusion in nonlinear multiplicative media. Journal of Mathematical Physics 10(10), 1862–1868 (1969)
- [12] Ablowitz, M.J., Fuchssteiner, B., Kruskal, M.: Topics in Soliton Theory and Exactly Solvable Nonlinear Equations: Proceedings of the Conference on Nonlinear Evolution Equations, Solitons and the Inverse Scattering Transform. World Scientific, ??? (1987)
- [13] Wang, X., Lu, Y.: Exact solutions of the extended burgers-fisher equation. Chinese Physics Letters 7(4), 145–147 (1990)
- [14] Chandrasekaran, P., Ramasami, E.: Painleve analysis of a class of nonlinear diffusion equations. International Journal of Stochastic Analysis 9(1), 77–86 (1996)
- [15] Chen, H., Zhang, H.: New multiple soliton solutions to the general burgers-fisher equation and the kuramoto-sivashinsky equation. Chaos, Solitons & Fractals 19(1), 71–76 (2004)
- [16] Lu, J., Yu-Cui, G., Shu-Jiang, X.: Some new exact solutions to the burgers–fisher equation and generalized burgers–fisher equation. Chinese Physics 16(9), 2514 (2007)
- [17] Kaya, D., El-Sayed, S.M.: A numerical simulation and explicit solutions of the generalized burgers–fisher equation. Applied Mathematics and computation 152(2), 403–413 (2004)

- [18] Zhu, C.-G., Kang, W.-S.: Numerical solution of burgers-fisher equation by cubic b-spline quasi-interpolation. Applied Mathematics and Computation 216(9), 2679–2686 (2010)
- [19] Mittal, R., Tripathi, A.: Numerical solutions of generalized burgers-fisher and generalized burgers-huxley equations using collocation of cubic b-splines. International Journal of Computer Mathematics 92(5), 1053–1077 (2015)
- [20] Singh, A., Dahiya, S., Singh, S.: A fourth-order b-spline collocation method for nonlinear burgers–fisher equation. Mathematical Sciences 14, 75–85 (2020)
- [21] Rashidi, M., Ganji, D., Dinarvand, S.: Explicit analytical solutions of the generalized burger and burger-fisher equations by homotopy perturbation method. Numerical Methods for Partial Differential Equations: An International Journal 25(2), 409–417 (2009)
- [22] Ismail, H.N., Raslan, K., Abd Rabboh, A.A.: Adomian decomposition method for burger's-huxley and burger's-fisher equations. Applied mathematics and computation 159(1), 291–301 (2004)
- [23] Khattak, A.J.: A computational meshless method for the generalized burger'shuxley equation. Applied Mathematical Modelling 33(9), 3718–3729 (2009)
- [24] Xu, Z.-h., Xian, D.-q.: Application of exp-function method to generalized burgersfisher equation. Acta Mathematicae Applicatae Sinica, English Series 26(4), 669– 676 (2010)
- [25] Zhang, J., Yan, G.: A lattice boltzmann model for the burgers-fisher equation. Chaos: An Interdisciplinary Journal of Nonlinear Science 20(2), 023129 (2010)
- [26] Sari, M., Gürarslan, G., Dağ, İ.: A compact finite difference method for the solution of the generalized burgers-fisher equation. Numerical Methods for Partial Differential Equations: An International Journal 26(1), 125–134 (2010)
- [27] Sari, M.: Differential quadrature solutions of the generalized burgers-fisher equation with a strong stability preserving high-order time integration. Mathematical and Computational Applications 16(2), 477–486 (2011)
- [28] Zhang, R., Yu, X., Zhao, G.: The local discontinuous galerkin method for burger's-huxley and burger's-fisher equations. Applied Mathematics and Computation 218(17), 8773–8778 (2012)
- [29] Nawaz, R., Ullah, H., Islam, S., Idrees, M.: Application of optimal homotopy asymptotic method to burger equations. Journal of Applied Mathematics 2013 (2013)

- [30] Yadav, O.P., Jiwari, R.: Finite element analysis and approximation of burgers'fisher equation. Numerical Methods for Partial Differential Equations 33(5), 1652–1677 (2017)
- [31] Malik, S.A., Qureshi, I.M., Amir, M., Malik, A.N., Haq, I.: Numerical solution to generalized burgers'-fisher equation using exp-function method hybridized with heuristic computation. PLoS One 10(3), 0121728 (2015)
- [32] Nadeem, M., Li, F., Ahmad, H.: Modified laplace variational iteration method for solving fourth-order parabolic partial differential equation with variable coefficients. Computers & Mathematics with Applications 78(6), 2052–2062 (2019)
- [33] Loyinmi, A.C., Akinfe, T.K.: Exact solutions to the family of fisher's reactiondiffusion equation using elzaki homotopy transformation perturbation method. Engineering Reports 2(2), 12084 (2020)
- [34] Akinfe, T.K., Loyinmi, A.C.: A solitary wave solution to the generalized burgersfisher's equation using an improved differential transform method: A hybrid scheme approach. Heliyon 7(5), 07001 (2021)
- [35] Mohanty, R.K., Sharma, S.: A high-resolution method based on off-step nonpolynomial spline approximations for the solution of burgers-fisher and coupled nonlinear burgers' equations. Engineering Computations 37(8), 2785–2818 (2020)
- [36] Kumar Verma, A., Kayenat, S.: On the stability of micken's type nsfd schemes for generalized burgers fisher equation. Journal of Difference Equations and Applications 25(12), 1706–1737 (2019)
- [37] Li, J.: Geometric properties and exact travelling wave solutions for the generalized burger-fisher equation and the sharma-tasso-olver equation. Journal of Nonlinear Modeling and analysis 1(1), 1–10 (2019)
- [38] Onyejekwe, O.O., Minale, B., Habtamu, F., Amha, T., Tamiru, G., Mengistu, B., Demiss, Y., Alemseged, N., *et al.*: Numerical discrete-domain integral formulations for generalized burger-fisher equation. Applied Mathematics **11**(03), 137 (2020)
- [39] Zhang, H., Xia, Y., N'gbo, P.-R.: Global existence and uniqueness of a periodic wave solution of the generalized burgers–fisher equation. Applied Mathematics Letters 121, 107353 (2021)
- [40] Mittal, A.K., Balyan, L.K.: Numerical solutions of time and space fractional coupled burgers equations using time-space chebyshev pseudospectral method. Mathematical Methods in the Applied Sciences 44(4), 3127-3137 (2021)
- [41] Mittal, A.: Spectrally accurate approximate solutions and convergence analysis

of fractional burgers' equation. Arabian Journal of Mathematics 9(3), 633–644 (2020)

- [42] Balyan, L., Mittal, A., Kumar, M., Choube, M.: Stability analysis and highly accurate numerical approximation of fisher's equations using pseudospectral method. Mathematics and Computers in Simulation 177, 86–104 (2020)
- [43] Saini, P., Balyan, L., Kumar, A., Singh, G.: Comparative analysis of postprocessing on spectral collocation methods for non-smooth functions. Signal, Image and Video Processing, 1–9 (2022)
- [44] Saini, P., Balyan, L., Kumar, A., Singh, G.: Modification of chebyshev pseudospectral method to minimize the gibbs oscillatory behaviour in resynthesizing process. Circuits, Systems, and Signal Processing 41(11), 6238–6265 (2022)
- [45] Hesthaven, J.S., Gottlieb, S., Gottlieb, D.: Spectral Methods for Time-dependent Problems vol. 21. Cambridge University Press, ??? (2007)
- [46] Boyd, J.P.: Chebyshev and Fourier Spectral Methods. Courier Corporation, ??? (2001)
- [47] Trefethen, L.N.: Spectral methods in matlab, volume 10 of software, environments, and tools. Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA 24, 57 (2000)
- [48] Canuto, C., Hussaini, M., Quarteroni, A., Thomas Jr, A.: others. Spectral methods in fluid dynamics. Springer (2012)
- [49] Javidi, M.: Modified pseudospectral method for generalized burger's-fisher equation. In: International Mathematical Forum, vol. 1, pp. 1555–1564 (2006). Citeseer
- [50] Golbabai, A., Javidi, M.: A spectral domain decomposition approach for the generalized burger's-fisher equation. Chaos, Solitons & Fractals 39(1), 385–392 (2009)
- [51] Ismail, H.N., Abd Rabboh, A.A.: A restrictive padé approximation for the solution of the generalized fisher and burger–fisher equations. Applied Mathematics and Computation 154(1), 203–210 (2004)