COMPLETIONS OF IMPLICATIVE ASSEMBLIES

ALEXANDRE MIQUEL AND KRZYSZTOF WORYTKIEWICZ

ABSTRACT. We continue our work on implicative assemblies by investigating under which circumstances a subset $M \subseteq \mathscr{S}$ gives rise to a lex full subcategory \mathbf{Asm}_M of the quasitopos \mathbf{Asm}_A of all assemblies such that $(\mathbf{Asm}_M)_{\text{reg}/\text{lex}} \simeq \mathbf{Asm}_A$. We establish a characterisation. Furthermore, this latter is relevant to the study of \mathbf{Asm}_M 's ex/lex-completion.

Contents

1. Introduction	2
2. Regular Completions	2
2.1. Memjog on Regular Categories	2
2.2. The Universal Property of a Regular Completion	3
3. Some Implicative Remarks	5
3.1. Assemblies as Indexed Families.	5
3.2. Stability under finite Limits	6
3.3. Implicative Existence	6
3.4. Valuations	6
4. Regular Completions of Implicative Assemblies	8
4.1. Density	8
4.2. Compactness	9
4.3. Generators	13
5. Exact Completions	13
6. Exact Completions of Implicative Assemblies	15
6.1. The Exact Completion of $\mathbf{Asm}_{\mathcal{A}}$	15
6.2. Tripos-to-Topos Construction	15
6.3. The Exact Completion of \mathbf{Asm}_M in Context	16
References	24

$\mathbf{2}$

A. MIQUEL AND K. WORYTKIEWICZ

1. INTRODUCTION

In this note we investigate under which circumstances a subset $M \subseteq \mathscr{S}$ gives rise to a lex full subcategory Asm_M of the quasitopos Asm_A of all assemblies such that $(\operatorname{Asm}_M)_{\operatorname{reg}/\operatorname{lex}} \simeq \operatorname{Asm}_A$. It is well-known that we have such a situation in the effective topos $\mathcal{E}ff$ [6, 5] where there is the subcategory $\operatorname{PAss} \subseteq \omega \operatorname{Set} \subseteq \mathcal{E}ff$ whose objects are known as *partitioned assemblies* (their existence predicates are singletons [12, 13]). We establish a characterisation for the implicative case and the case of partitioned assemblies turns out to be quite easy going. We next observe that $(\operatorname{Asm}_A)_{\operatorname{ex}/\operatorname{reg}}$ is always a topos, which is a rather immediate consequence of the material in [9]. This observation implies that $(\operatorname{Asm}_M)_{\operatorname{ex}/\operatorname{lex}}$ is a topos as well. We then construct a functor

$\mathcal{K}: (\mathbf{Asm}_M)_{\mathrm{ex}/\mathrm{lex}} \to \mathbf{Set}[\mathcal{A}]$

the latter being the topos constructed from an implicative tripos [10, 4] by means of the tripos-to-topos construction. It turns out that \mathcal{K} may not always be an equivalence, although it is always the case when \mathcal{A} is compatible with joins. There are undoubtly connections to the material in [8], we only (very) recently became aware of the latter work. Then again our focus is quite different as we specifically investigate categorical aspects of implicative algebras. The reader is referred to [10, 4] for basic material about implicative algebras and implicative assemblies.

2. Regular Completions

2.1. Memjog on Regular Categories.

The notion of regular category axiomatises image factorisations, as a matter of fact an abelian category is an additive regular category. It is thus not really surprising that regular categories have been studied for some time [1].

Definition 2.1.

- 1. A regular epi is an epi which is a coequaliser (in any category).
- 2. A *lex category* is a category with finite limits.

Remark 2.1. In a lex category, a regular epi is a coequaliser if its kernel pair.

Definition 2.2. A lex category \mathbb{D} is regular if

- i. it has coequalisers of kernel pairs;
- ii. regular epis are stable under pullback.

Despite notorious non-examples like **Top** (always troubles with this one..) or **Cat**, regular categories abound in nature: **Set**, **Grp** (all groups), **Ab**, any LCCC with coequalisers (thus any quasitopos), any topos etc etc. As mentioned, any abelian category is regular as well. This is the reason why in the context of regular categories we tend to call *exact sequence* a coequaliser diagram of a kernel pair. For, the datum

$$X' \xrightarrow{e_0} X \xrightarrow{e} X''$$

(with (e_0, e_1) the kernel pair of e) in an abelian category yields the short exact sequence

$$0 \longrightarrow X' \xrightarrow{\langle e_0, e_1 \rangle} X \oplus X \xrightarrow{[e, -e]} X'' \longrightarrow 0$$

and vice versa. An equivalent definition of a regular category is then as above but with condition *ii*. replaced by requiring that exact sequences are stable under pullback.

Given any morphism $f: X \to Y$ in a regular category we can construct the diagram


$$\operatorname{Ker}(f) \xrightarrow{f_0} X \xrightarrow{\overline{f}} \stackrel{\operatorname{Im}(f)}{\xrightarrow{\iota_f}} Y$$

with $\text{Im}(f) \cong X/\text{Ker}(f)$, \bar{f} the coequaliser of f's kernel pair (f_0, f_1) and ι_f given by universal property (it is an exercise in diagram chasing to prove that ι_f is mono). In an abelian category it is just the first isomorphism theorem.

Notation. Given a morphism f in a regular category, we shall write $f = \iota_f \circ \overline{f}$ for its factorisation with ι_f mono and \overline{f} regular epi.

Regular epis and monos form an orthogonal (thus functorial) factorisation system ($\mathcal{E}_{reg}, \mathcal{M}$). Another equivalent definition of a regular category is as a lex category with pullback-stable image factorisations.

Example 2.1. $\operatorname{Asm}_{\mathcal{A}}$ is a quasitopos, hence regular. It is nonetheless handy to know how to calculate image factorisations. Let $f : X \to Y$ be a morphism in $\operatorname{Asm}_{\mathcal{A}}$. Its kernel pair is given by the pullback square




Recall that a pullback square in $\operatorname{Asm}_{\mathcal{A}}$ is given on carriers by the forgetful functor $\Gamma : \operatorname{Asm}_{\mathcal{A}} \to \operatorname{Set}$, with existence and projections inherited from the product in $\operatorname{Asm}_{\mathcal{A}}$, so $\mathsf{E}_{\operatorname{Ker}(f)}(x, x') = x \sqcap x'$ while p_1 and p_2 are tracked by $(\lambda z.z \ (\lambda xy.x))^{\mathcal{A}}$ and $(\lambda z.z \ (\lambda xy.y))^{\mathcal{A}}$, respectively. Let $[x] := f^{-1}(f(x))$ be the inverse image of f(x). The coequaliser diagram

$$\operatorname{Ker}(f) \xrightarrow{p_1} X \xrightarrow{\overline{f}} \operatorname{Im}(f)$$

is given by

- i. $\underline{\mathrm{Im}(f)} = X/\sim$ where \sim is the equivalence relation generated by $f(x) \sim f(x')$, so $\underline{\mathrm{Im}(f)}$ is the set $\overline{\{[x]|x \in X\}}$. The existence predicate is $\mathsf{E}_{\mathrm{Im}(f)}([x]) = \underset{x' \in [x]}{\exists} \mathsf{E}_X(x')$;
- ii. $\overline{f}: x \mapsto [x]$ is the canonical projection, tracked by $(\lambda xz.zx)^{\mathcal{A}}$.

The image factorisation



is in turn given by universal property of Im(f). Specifically, we have $\iota_f([x]_f) = f(x)$ independently of the choice of representant x, tracked by $(\lambda z. z\tau)^{\mathcal{A}}$ with τ some tracker of f. It can be seen without invoking category theory that ι_f is injective as a map, thus a mono in $\text{Asm}_{\mathcal{A}}$.

2.2. The Universal Property of a Regular Completion.

Definition 2.3.

- 1. A *lex functor* is a functor commuting with finite limits.
- 2. Let \mathbb{D} and \mathbb{D}' be regular categories. An *exact functor* $R : \mathbb{D} \to \mathbb{D}'$ is a lex functor which preserves regular epis.


Equivalently, a lex functor preserving exact sequences is exact (whence the name). An exact functor obviously preserves image factorisations.

Notation. We shall write

- Lex for the 2-category of lex categories, lex funtors and their natural transformations;

- Reg for the 2-category of regular categories, exact funtors and their natural transformations;
- $|-|: \mathbf{Reg} \to \mathbf{Lex}$ for the forgetful 2-functor.

It is part of the lore that |-| has a left biadjoint $(-)_{\operatorname{reg}/\operatorname{lex}}$. Given a lex category \mathbb{C} we thus have a lex functor $\boldsymbol{y}: \mathbb{C} \to \mathbb{C}_{\operatorname{reg}/\operatorname{lex}}$ such that for any lex functor $L: \mathbb{C} \to \mathbb{D}$ with \mathbb{D} a regular category there is a unique exact functor $\mathcal{U}: \mathbb{C}_{\operatorname{reg}/\operatorname{lex}} \to \mathbb{D}$ such that the triangle



commutes (up-to equivalence). The objects of $\mathbb{C}_{reg/lex}$ are morphisms of \mathbb{C} . A morphism

$$\begin{bmatrix} l \end{bmatrix} : \begin{bmatrix} X \\ \downarrow f \\ Y \end{bmatrix} \to \begin{bmatrix} X' \\ \downarrow f' \\ Y' \end{bmatrix}$$

in \mathbb{C}_{reg} is an equivalence class [l] of morphisms $l: X \to X'$ such that $f' \circ l$ coequalises f's kernel pair, with $[l_1] = [l_2]$ if $f' \circ l_1 = f' \circ l_2$.

The insertion functor $\boldsymbol{y}: \mathbb{C} \to \mathbb{C}_{\mathrm{reg}}$ lex is given by $X \mapsto \begin{bmatrix} X \\ \downarrow \mathrm{id} \\ X \end{bmatrix}$

Given a morphism


$$[f]: \begin{bmatrix} X\\ \downarrow \mathrm{id}\\ X \end{bmatrix} \to \begin{bmatrix} Y\\ \downarrow \mathrm{id}\\ Y \end{bmatrix}$$

in the image of y, it is obvious that f is the only member of [f] and easy to see that

$$\operatorname{Im}([f]) = \begin{bmatrix} X \\ \downarrow f \\ Y \end{bmatrix}$$

(where f is seen as an object of $\mathbb{C}_{\text{reg}/\text{lex}}$). It would be fair to say that this is the idea behind the construction as the latter freely adds images.

A frequent use-case is when $\mathbb{D} \in \mathbf{Reg}$ and $\mathbb{C} \subset \mathbb{D}$ is a lex subcategory. The canonical functor \mathcal{U} making the diagram




commute is particularly easy to describe in this situation:

$$\begin{aligned} \mathcal{U} : \mathbb{C}_{\operatorname{reg}/\operatorname{lex}} & \longrightarrow & \mathbb{D} \\ \begin{bmatrix} X \\ \downarrow f \\ Y \end{bmatrix} & \mapsto & \operatorname{Im}(f) \end{aligned}$$

with action on morphisms given by universal property as follows. Assume a morphism

$$\begin{bmatrix} l \end{bmatrix} : \begin{bmatrix} X \\ \downarrow f \\ Y \end{bmatrix} \to \begin{bmatrix} X' \\ \downarrow f' \\ Y' \end{bmatrix}$$

We then have the diagram



in \mathbb{D} . As $f' \circ l$ coequalises the kernel pair (f_0, f_1) of f and $\iota_{f'}$ is mono, $\overline{f'}$ coequalises (f_0, f_1) as well so \tilde{l} is given by universal property of \bar{f} . It is easy to see that the assignment $[l] \mapsto \tilde{l}$ is well defined and functorial. Moreover, \mathcal{U} is always faithful in this situation. Finally, we have $\mathbb{C}_{\text{reg}/\text{lex}} \simeq \mathbb{D}$ if and only if \mathcal{U} is (part of) an equivalence of categories.

Definition 2.4. \mathbb{D} is a *regular completion* of \mathbb{C} if \mathcal{U} is an equivalence of categories.

In practice we thus need to check if \mathcal{U} is essentially surjective and full. It is well-known that if \mathbb{D} is a *regular* completion of \mathbb{C} , closing \mathbb{C} under isomorphisms yields \mathbb{D} 's subcategory of projective objects.

3. Some Implicative Remarks

3.1. Assemblies as Indexed Families.

Notation. Let \mathcal{A} be an implicative algebra.

- 1. $\mathbb{P}[\mathcal{A}]$ stands for the implicative tripos over \mathcal{A} and $\mathbb{P}[\mathcal{A}]_X$ for its fibre over X.
- 2. Let $Q \subseteq \mathscr{S}$ be a subset of \mathscr{S} . We shall write $\mathbb{P}[Q]$ for the full subcategory of $\mathbb{P}[\mathcal{A}]$ of families of elements of Q.
- 3. $\mathbb{P}[Q]_X := \mathbb{P}[\mathcal{A}]_X \cap \mathbb{P}[Q].$

Definition 3.1. We call $\mathbb{P}[Q]_X$ the *Q*-fibre over *X*.

Proposition 3.1. The assignment

$$\begin{array}{rccc} \Xi: \mathbf{Asm}_{\mathcal{A}} & \longrightarrow & \mathbb{P}[\mathscr{S}] \\ & X & \mapsto & \mathsf{E}_{X} \\ & f \downarrow & & \downarrow (f, \mathsf{E}_{X} \sqsubseteq_{X} f^{*} \mathsf{E}_{Y}) \\ & Y & \mapsto & \mathsf{E}_{Y} \end{array}$$

is an isomorphism of categories with inverse

$$\begin{aligned} \Xi^{-1} : \mathbb{P}[\mathscr{S}] &\longrightarrow \mathbf{Asm}_{\mathcal{A}} \\ (u_x)_{x \in X} &\mapsto (X, u) \\ (f, u \sqsubseteq_X f^* v) \downarrow & \downarrow f \\ (v_y)_{y \in Y} &\mapsto (Y, v) \end{aligned}$$

Proof. Ξ_0 and Ξ_0^{-1} are obviously inverse bijections (on classes). Furthermore, the maps

$$\begin{aligned} \Xi_{X,Y} : \mathbf{Asm}_{\mathcal{A}}(X,Y) &\longrightarrow & \mathbb{P}[\mathscr{S}] \left(\mathsf{E}_X,\mathsf{E}_Y\right) \\ f &\mapsto & (f,\mathsf{E}_X \sqsubseteq_X f^*\mathsf{E}_Y) \end{aligned}$$

and

$$\begin{aligned} \Xi_{u,v}^{-1} : \mathbb{P}[\mathscr{S}]((u_x)_{x \in X}, (v_y)_{y \in Y}) &\longrightarrow & \mathbf{Asm}_{\mathcal{A}}(X, Y) \\ (f, u \sqsubseteq_X f^* v) &\mapsto & f \end{aligned}$$

are inverse bijections since

$$f \text{ is tracked} \iff \bigwedge_{x \in X} (\mathsf{E}_X(x) \to \mathsf{E}_Y(f(x))) \in \mathscr{S}$$
$$\iff \bigwedge_X (\mathsf{E}_X \to_X f^*\mathsf{E}_Y) \in \mathscr{S}$$
$$\iff \mathsf{E}_X \sqsubseteq_X f^*\mathsf{E}_Y$$

The ismorphism above is obviously rather shallow. Still, it is quite practical to be able to take either point of view according to a given situation.

3.2. Stability under finite Limits.

Definition 3.2. Let $M \subseteq \mathscr{S}$. The category Asm_M of *M*-assemblies is the full subcategory of Asm_A with objects assemblies valued in M.

Remark 3.1. $\operatorname{Asm}_M \cong \mathbb{P}[M]$

Recall that in an implicative algebra, the product \sqcap (with respect to the entailment preorder) is given by the second-order encoding of pairs.

Definition 3.3. A subset $M \subseteq \mathscr{S}$ is *algebraic* if if it is closed under \sqcap .

Remark 3.2. \mathbf{Asm}_M is closed under finite limits if M is algebraic.

Example 3.1. $\mathcal{A} = (\mathfrak{P}(P), \subseteq, \rightarrow, \mathscr{S} = \mathfrak{P}_{\bullet}(P))$ with P a combinatory algebra. The subset

$$M := \{\{p\} | p \in P\}$$

of the separator is manifestly algebraic as in this case we have $\{p_1\} \sqcap \{p_2\} = \{\langle p_1, p_2 \rangle\}.$

3.3. **Implicative Existence.** Before we proceed further we need to compile some further useful facts about the operation of existence in implicative algebras.

Remark 3.3. Recall that the implicative existence is given by second-order encoding as

$$\exists U = \bigwedge_{c \in \mathcal{A}} \bigwedge_{u \in U} ((u \to c) \to c)$$

- 1. We have $\vdash \lambda z.zu : \forall c : \mathcal{A}.\forall u : U.(u \to c) \to c$, hence $(\lambda z.zu)^{\mathcal{A}} \preccurlyeq \exists U$ for all $u \in U$, so in particular $\exists U \in \mathscr{S}$ if $U \cap \mathscr{S} \neq \emptyset$.
- 2. A consequence of the above item is that if we have $\chi := a \to \exists U$ then for any $a' \preccurlyeq a$ there is an $u \in U$ such that $\chi a' \preccurlyeq (\lambda z. zu)^A$. We systematically exploit this fact, in particular when building trackers.

3.4. Valuations.

Definition 3.4. Let X be a set and $\emptyset \neq M \subseteq \mathscr{S}$. An *M*-valuation of X (or just valuation if M and X are understood) is map $\nu : X \to \mathfrak{P}_{\bullet}(M)$.

Remark 3.4. Let X be a set and $M \subseteq \mathscr{S}$. We have the predicate

$$\begin{array}{cccc} \mathbf{x}: \mathfrak{P}_{\bullet}\left(M\right) & \longrightarrow & \mathscr{S} \\ U & \mapsto & \exists U \end{array}$$

in $\mathbb{P}[\mathscr{S}]_{\mathfrak{P}_{\bullet}(M)}$. Given a valuation $\nu: X \to \mathfrak{P}_{\bullet}(M)$, we have the predicate $\nu^{*}(\mathrm{ex}) \in \mathbb{P}[\mathscr{S}]_{X}$ where

 \mathbf{SO}

$$\nu^*(\mathbf{ex}) = \left(\exists (\nu(x))\right)_{x \in X}$$

Notation. Let $\nu: X \to \mathfrak{P}_{\bullet}(M)$ be a valuation. We shall use the notation $\exists \nu := \nu^*(ex)$.

ez

Lemma 3.1. Any morphism $f: X \to Y$ in Asm_M gives rise to the valuation

 ν

$$f: \underline{\mathrm{Im}}(f) \longrightarrow \mathfrak{P}_{\bullet}(M)$$
$$[x] \mapsto \{\mathsf{E}_X(x') | x' \in [x]\}$$

and we have $\mathsf{E}_{\mathrm{Im}(f)} = \exists \nu_f$.

Notation. $[x] := f^{-1}(f(x)).$

Proof. We have $\underline{\mathrm{Im}(f)} = \{[x] | x \in X\}$ where $\mathsf{E}_{\mathrm{Im}(f)}([x]) = \underset{x' \in [x]}{\exists} \mathsf{E}_X(x')$, hence $(\exists \nu_f)_{[x]} = (\nu_f^* \operatorname{ex})_{[x]}$ $= \exists_{x' \in [x]} \mathsf{E}_X(x')$

Proposition 3.2. Let X be a set and $\nu : X \to \mathfrak{P}_{\bullet}(M)$ be a valuation. Let $\hat{X} \in \mathbf{Asm}_M$ be the assembly given by

- $\underline{\hat{X}} := \sum_{x \in X} \nu(x);$ $- \mathsf{E}_{\hat{X}}(x,m) := m \text{ for all } x \in X \text{ and } m \in \nu(x).$

and $\check{X} \in \mathbf{Asm}_M$ be the assembly given by

 $- \underline{\check{X}} := X;$


$$- \mathsf{E}_{\check{X}}(x) := m_0 \text{ for all } x \in X \text{ and some } m_0 \in M.$$

The obvious surjective map $g_{\nu}: \underline{\hat{X}} \twoheadrightarrow \underline{\check{X}}$ is tracked by $(\lambda x.m_0)^{\mathcal{A}}$ and we have the isomorphism

$$\begin{aligned} [-]_{\nu} : (X, \exists \nu) & \xrightarrow{\cong} & (\mathrm{Im}(g_{\nu}), \exists g_{\nu}) \\ x & \mapsto & g_{\nu}^{-1}(x) \end{aligned}$$

in Asm_A .

Proof. Consider the image factorisation



We have $Im(g_{\nu}) = \{g_{\nu}^{-1}(x) | x \in X\}$ as

$$[x_m] = g_{\nu}^{-1}(g_{\nu}(x_m)) = g_{\nu}^{-1}(x)$$

for all $m \in \nu(x)$, hence the bijection

$$\begin{array}{cccc} [-]_{\nu} : X & \stackrel{\cong}{\longrightarrow} & \operatorname{Im}(g_{\nu}) \\ & x & \mapsto & g_{\nu}^{-1}(x) \end{array}$$

We have


$$\nu_{g_{\nu}}(g_{\nu}^{-1}(x)) = \{\mathsf{E}_{\hat{X}}(x_m) | x_m \in g_{\nu}^{-1}(x)\} \\
= \{m \in M | x_m \in g_{\nu}^{-1}(x)\} \\
= \nu(x)$$

hence

$$\mathsf{E}_{\mathrm{Im}(g_{\nu})}([x]_{\nu}) = (\exists \nu_{g_{\nu}})_{g_{\nu}^{-1}(x)}$$
 Lemma 3.1
$$= \exists (\nu_{g_{\nu}}(g_{\nu}^{-1}(x)))$$
$$= \exists \nu(x)$$

and therefore

$$\begin{split} & \bigwedge_{x \in X} \left(\left(\exists \nu \right)_x \to \mathsf{E}_{\mathrm{Im}(g_{\nu})}([x]_{\nu}) \right) &= & \bigwedge_{x \in X} \left(\left(\exists \nu \right)_x \to \left(\exists \nu \right)_x \right) \\ &= & \bigwedge_{x \in X} \left(\mathsf{E}_{\mathrm{Im}(g_{\nu})}([x]_{\nu}) \to \left(\exists \nu \right)_x \right) \end{split}$$



But $\bigwedge_{x \in X} ((\exists \nu)_x \to (\exists \nu)_x) \in \mathscr{S}$, hence $[-]_{\nu}$ and $[-]_{\nu}^{-1}$ are tracked.

4. Regular Completions of Implicative Assemblies

4.1. Density.

Definition 4.1. A non-empty subset $M \subseteq \mathscr{S}$ of the separator is *dense* in \mathscr{S} (or just *dense* if \mathscr{S} is understood) if there is a valuation $\nu : \mathscr{S} \to \mathfrak{P}_{\bullet}(M)$ such that $\mathrm{id}_{\mathscr{S}} \cong_{\mathscr{S}} \exists \nu$.

Theorem 4.1. Let $M \subseteq \mathscr{S}$ be an algebraic subset. The following are equivalent

- i. M is dense;
- ii. for any $X \in \mathbf{Set}$ and any $u \in \mathbb{P}[\mathscr{S}]_X$ there is a valuation $\nu : X \to \mathfrak{P}_{\bullet}(M)$ such that $u \cong_X \exists \nu$;
- iii. the functor $U: (\mathbf{Asm}_M)_{\mathrm{reg}/\mathrm{lex}} \to \mathbf{Asm}_{\mathcal{A}}$ is essentially surjective.

Proof. We need to insist that M is algebraic only in order to be able to talk about $(\mathbf{Asm}_M)_{\text{reg/lex}}$, otherwise it is *deus ex machina* behind the stage.

$$(i) \Longrightarrow (ii)$$
 There is a valuation $\nu : \mathscr{S} \to \mathfrak{P}_{\bullet}(M)$ of \mathscr{S} such that $\mathrm{id}_{\mathscr{S}} \cong_{\mathscr{S}} \exists \nu$. Let $u \in \mathbb{P}[\mathscr{S}]_X$. We have
 $\mathrm{id}_{\mathscr{S}} = i_s^*(\mathrm{id}_{\mathcal{A}})$

with $i_{\mathscr{S}}: \mathscr{S} \to \mathcal{A}$ the insertion map, hence

$$u = u^{*}(\mathrm{id}_{\mathcal{A}}) \qquad \mathrm{id}_{\mathcal{A}} \in \mathbb{P}\left[\mathcal{A}\right]_{\mathcal{A}} \text{ is a split generic object} \\ = (i_{\mathscr{S}} \circ u)^{*}(\mathrm{id}_{\mathcal{A}}) \\ = u^{*}(i_{s}^{*}(\mathrm{id}_{\mathcal{A}})) \qquad \mathbb{P}[\mathcal{A}] \to \mathbf{Set} \text{ is a split fibration} \\ = u^{*}(\mathrm{id}_{\mathscr{S}}) \\ \cong_{X} u^{*}(\exists \nu) \qquad \text{functoriality} \\ = \exists (\nu \circ u)$$

so the valuation $\nu \circ u : X \to \mathfrak{P}_{\bullet}(M)$ witnesses the fact that M is dense.

 $(ii) \Longrightarrow (iii)$ Let $X \in \mathbf{Asm}_M$. Assume a valuation $\nu : X \to \mathfrak{P}_{\bullet}(M)$ such that $\mathsf{E}_X \cong_X \exists \nu$. There is a

morphism g_{ν} such that $\Xi(\operatorname{Im}(g_{\nu})) = \exists \nu$ (c.f. Lemma 3.1). We furthermore have $\Xi(X) = \mathsf{E}_X$, hence the isomorphism

$$\Xi^{-1}(\mathsf{E}_X \cong_X \exists \nu) : X \xrightarrow{\cong} \operatorname{Im}(g_{\nu})$$

 $(iii) \Longrightarrow (i)$ We have $\Xi^{-1}(\mathrm{id}_{\mathscr{S}}) = (\mathscr{S}, \mathrm{id}_{\mathscr{S}})$. There is by hypothesis a morphism $f: X \to Y$ in Asm_M such that $\mathscr{S} \cong \mathrm{Im}(f)$. Let $k: \mathscr{S} \xrightarrow{\cong} \mathrm{Im}(f)$ be a witnessing iso. We have

$$\Xi(k) = (k, \mathrm{id} \sqsubseteq_{\mathscr{S}} k^* \exists \nu_f)$$

= $(k, \mathrm{id} \sqsubseteq_{\mathscr{S}} \exists (\nu_f \circ k))$

and

$$\Xi(k^{-1}) = (k^{-1}, \exists \nu_f \sqsubseteq_{\operatorname{Im}(f)} (k^{-1})^* \operatorname{id}_{\mathscr{S}})$$
$$= (k^{-1}, \exists \nu_f \sqsubseteq_{\operatorname{Im}(f)} k^{-1})$$

Notice that k^{-1} : Im $(f) \to \mathscr{S}$ has the "right type" here in the sense that $k^{-1} \in \mathbb{P}[\mathscr{S}]_{\mathrm{Im}(f)}$. Reindexing the vertical morphism $\exists \nu_f \sqsubseteq_{\mathrm{Im}(f)} k^{-1}$ by k yields

$$k^* \left(\exists \nu_f \sqsubseteq_{\mathrm{Im}(f)} k^{-1} \right) = k^* \left(\exists \nu_f \right) \sqsubseteq_{\mathscr{S}} k^*(k^{-1})$$

= $\exists (\nu_f \circ k) \sqsubseteq_{\mathscr{S}} \mathrm{id}_{\mathscr{S}}$

We thus have $\operatorname{id}_{\mathscr{S}} \cong_{\mathscr{S}} \exists (\nu_f \circ k)$ by virtue of the valuation $\nu_f \circ k$.

8

Example 4.1. In example 3.1 we considered an implicative algebra $\mathcal{A} = (\mathfrak{P}(P), \subseteq, \rightarrow, \mathscr{S})$ constructed from an applicative structure P and showed that the set $M := \{\{p\} | p \in P\}$ of all singletons is algebraic. Now $\mathscr{S} = \mathfrak{P}_{\bullet}(P)$ here so we have the *trivial valuation*

$$\begin{array}{rcl} \nu : \mathscr{S} & \longrightarrow & \mathfrak{P}_{\bullet}\left(M\right) \\ & U & \mapsto & \{\{m\} | m \in U\} \end{array}$$

which yields $\exists \nu_U = U$, hence

$$\begin{split} & \bigwedge_{U \in \mathscr{S}} \left(U \to \left(\exists \, \nu \right)_U \right) &= \bigcap_{U \in \mathscr{S}} \left(U \to U \right) \\ &= \bigwedge_{U \in \mathscr{S}} \left(\left(\exists \, \nu \right)_U \to U \right) \end{split}$$

But $\bigcap_{U \in \mathscr{S}} (U \to U) \neq \emptyset$ as it contains id, hence id $\cong_{\mathscr{S}} \exists \nu$.

Definition 4.2. An implicative structure \mathcal{A} is *compatible with joins* provided

$$\bigwedge_{a \in A} (a \to b) = \left(\bigvee_{a \in A} a \right) \to b$$

for all $A \subseteq A$ and $b \in A$. An implicative algebra is compatible with joins if it is the case for the underlying implicative structure.

Remark 4.1. In an implicative algebra \mathcal{A} compatible with joins the operations \exists and γ are the same.

Proposition 4.1. *et* \mathcal{A} *be an implicative algebra compatible with joins. An algebraic subset* $M \subseteq \mathscr{S}$ *is dense provided any element of* \mathscr{S} *is a join of elements of* M.

Proof. For each $a \in \mathcal{A}$ we can choose a subset $M_a \subseteq M$ such that $a = \Upsilon M_a$ by virtue of (i), hence there is the trivial valuation

$$\nu: \mathcal{A} \longrightarrow \mathfrak{P}_{\bullet}(M)$$
$$a \mapsto M_a$$

We have

$$(\exists \nu)_a = \exists M_a = \bigvee M_a \quad \text{compatible with joins} = a$$

hence


$$\begin{split} & \bigwedge_{a \in A} \left(a \to \left(\exists \, \nu \right)_a \right) &= & \bigwedge_{a \in A} \left(a \to a \right) \\ &= & \bigwedge_{a \in A} \left(\left(\exists \, \nu \right)_a \to a \right) \end{split}$$

But id $\preccurlyeq \bigwedge_{a \in \mathcal{A}} (a \to a)$ hence $\bigwedge_{a \in \mathcal{A}} (a \to a) \in \mathscr{S}$, which implies that M is dense.

4.2. Compactness.

Lemma 4.1. Let $M \subseteq \mathscr{S}$ and let $g: A \to B$ be a morphism in Asm_M . The following are equivalent


i. any morphism $k: X \to \text{Im}(g)$ in $\text{Asm}_{\mathcal{A}}$ with $X \in \text{Asm}_M$ admits a lift




ii. any morphism $l \circ \overline{f} : X \to \operatorname{Im}(g)$ in $\operatorname{Asm}_{\mathcal{A}}$ where

 $-f: X \to Y$ is a morphism in Asm_M ;

 $-l: \operatorname{Im}(f) \to \operatorname{Im}(g)$ is a morphism in $\operatorname{Asm}_{\mathcal{A}}$ admits a lift



Proof. The implication $(i) \Rightarrow (ii)$ is trivial. Assume (ii) and let $k : X \to \text{Im}(g)$ be a morphism in $\text{Asm}_{\mathcal{A}}$. We have the diagram



in $\mathbf{Asm}_{\mathcal{A}}$ with

- $-X, B \in \mathbf{Asm}_M;$
- $\exists!$ the unique lift of the lower square (given that regular epis and monos form an orthogonal factorisation system)

hence $k = (\exists !) \circ \overline{\iota_q \circ k}$ lifts.

Definition 4.3. $M \subseteq \mathscr{S}$ is *compact* if for any $u \in \mathbb{P}[M]_X$ and any valuation $\nu : X \to \mathfrak{P}_{\bullet}(M)$ such that $u \sqsubseteq_X \exists \nu$


there is $b \in \mathbb{P}[M]_X M$ with $b_x \in \nu(x)$ for all $x \in X$ such that

$$u \sqsubseteq_X b \sqsubseteq_X \exists \nu$$

Lemma 4.2. Let $M \subseteq \mathscr{S}$. Assuming choice the following are equivalent

i. M is compact;

ii. for any morphisms $g: A \to B$ in Asm_M and $k: X \to \operatorname{Im}(g)$ in Asm_A with $X \in \operatorname{Asm}_M$ the diagram



admits a lift.

Proof. $(i) \Rightarrow (ii)$ Let $g : A \to B$ be a morphism in Asm_M and $k : X \to \operatorname{Im}(g)$ be a morphism in Asm_A with $X \in \operatorname{Asm}_M$. The induced valuation $\nu_g : \operatorname{Im}(g) \to \mathfrak{P}_{\bullet}(M)$ verifies

$$\exists \nu_g = \mathsf{E}_{\mathrm{Im}(g)} \quad (\star)$$

(c.f. Lemma 3.1). The map $\nu := \nu_{\bar{g}} \circ k$ is a valuation as well. Furthermore

$$\begin{split} & \bigwedge_{x \in X} \left(\mathsf{E}_X(x) \to \exists \, \nu(x) \right) &= & \bigwedge_{x \in X} \left(\mathsf{E}_X(x) \to \exists \, \nu_g(k(x)) \right) \\ &= & \bigwedge_{x \in X} \left(\mathsf{E}_X(x) \to \mathsf{E}_{\mathrm{Im}(g)}(k(x)) \right) \\ &\in & \mathscr{S} \end{split}$$

since k is tracked, hence $\mathsf{E}_X \sqsubseteq_X \exists \nu$. M is discrete by hypothesis, hence there is $b \in \mathbb{P}[M]_X M$ with $b_x \in \nu(x)$ for all $x \in X$ such that

$$\mathsf{E}_X \sqsubseteq_X b \sqsubseteq_X \exists \nu$$

We have

$$\Xi (\mathsf{E}_X \sqsubseteq_X b) = (X, \mathsf{E}_X) \xrightarrow{\mathrm{id}} (X, b)$$

. .

On the other hand

$$(\exists \nu) (x) = \exists (\nu_{\bar{g}} \circ k)(x) = \exists \nu_{\bar{g}}(k(x))$$

for all $x \in X$, hence $\exists \nu = k^* \exists \nu_{\bar{g}}$ which entails

(

$$\begin{split} \Xi \left(b \ \sqsubseteq_X \ \exists \nu \right) &= \ \Xi \left(b \ \sqsubseteq_X \ k^* \ \exists \nu_{\bar{g}} \right) \\ &= \ (X, b) \xrightarrow{k} \left(\operatorname{Im}(f), \exists \nu_g \right) \end{split}$$

This in turn entails that k factors as

$$(X, \mathsf{E}_X) \xrightarrow{\mathrm{id}} (X, b) \xrightarrow{k} (\mathrm{Im}(f), \exists \nu_g)$$


in $\operatorname{Asm}_{\mathcal{A}}$. Now $b_x \in \nu(x) = \{\mathsf{E}_A(a) | a \in k(x)\}$ for each $x \in X$, hence there is $\tilde{a} \in k(x)$ such that $b_x = \mathsf{E}_A(\tilde{a})$. Notice that while the value $\mathsf{E}_A(\tilde{a})$ of b_x is determined, \tilde{a} itself does not need to be unique. Let

$$\hat{A}_x := \{\tilde{a} \in k(x) | b_x = \mathsf{E}_A(\tilde{a})\}$$


be the set of those \tilde{a} . Assuming choice there is a map $l: X \to A$ such that $l(x) \in \tilde{A}_x$ for all $x \in X$. Furthermore

$$\begin{split} & \bigwedge_{x \in X} \left(b_x \to \mathsf{E}_A(l(x)) \right) &= & \bigwedge_{x \in X} \left(\mathsf{E}_A(\tilde{a}) \to \mathsf{E}_A(\tilde{a}) \right) \\ & \in & \mathscr{S} \end{split}$$

so l is tracked. Finally we have $\bar{g} \circ l = k$ as maps since $k(x) = \bar{g}(a)$ for any $a \in k(x)$ hence the lift



 $(ii) \Rightarrow (i)$ Let $u \in \mathbb{P}[M]_X$ and $\nu : X \to \mathfrak{P}_{\bullet}(M)$ be a valuation such that $\bigwedge_{x \in X} (u_x \to (\exists \nu)_x) \in \mathscr{S}$. We thus have $u \sqsubseteq_X \exists \nu$ and further $\Xi^{-1} (u \sqsubseteq_X \exists \nu) = \mathrm{id} : (X, u) \to (X, \exists \nu)$. Composing the latter with $[-]_{\nu}$ (c.f Proposition 3.2) yields the morphism $[-]_{\nu} : (X, u) \to (\mathrm{Im}(g_{\nu}), \mathsf{E}_{\mathrm{Im}(g_{\nu})})$ in $\mathbf{Asm}_{\mathcal{A}}$. The latter has a lift



Let $b \in \mathbb{P}[M]_X$ be given by $b_x := \mathsf{E}_{\hat{X}}(l(x))$. We have by construction $l(x) = x_m$ for some $m \in \nu(x)$, hence

$$\mathsf{E}_{\hat{X}}(l(x)) = \mathsf{E}_{\hat{X}}(x_m)$$
$$= m$$

so $b_x \in \nu(x)$ for all $x \in X$. Now l is tracked so

$$\begin{split} & \bigwedge_{x \in X} (u_x \to b_x) &= & \bigwedge_{x \in X} \left(\mathsf{E}_X \to \mathsf{E}_{\hat{X}}(l(x)) \right) \\ & \in & \mathscr{S} \end{split}$$


Theorem 4.2. Let $M \subseteq \mathscr{S}$ be an algebraic subset. Assuming choice the following are equivalent

- *i.* M is compact;
- ii. $U : \operatorname{Asm}_{M_{\operatorname{reg}}} \to \operatorname{Asm}_{\mathcal{A}}$ is full.

Proof. M being compact is equivalent to the assertion that any morphism $l \circ \overline{f} : X \to \text{Im}(g)$ where

- i. $f: X \to Y$ and $g: A \to B$ are morphisms in \mathbf{Asm}_M ;
- ii. $l: \operatorname{Im}(f) \to \operatorname{Im}(g)$ is a morphism in $\operatorname{Asm}_{\mathcal{A}}$

admits a lift



(c.f. Lemma 4.1 and 4.2), which is obviously equivalent to the assertion that $U : \operatorname{Asm}_{M_{\operatorname{reg}}} \to \operatorname{Asm}_{\mathcal{A}}$ is full (c.f. Section 2).

Example 4.2. We observed that the set of all singletons $M := \{\{p\} | p \in P\}$ is algebraic (c.f. Example 3.1) and dense (c.f. Example 4.1). Taking it from there we now show that M is compact as well. Let $u \in \mathbb{P}[M]_X$ be a family of singletons $u_x = \{\check{u}_x\}$ and $\nu : X \to \mathfrak{P}_{\bullet}(M)$ be a valuation. Suppose $\bigwedge_{x \in X} (u_x \to (\exists \nu)_x) \in \mathscr{S}$. Unravelling yields

$$\begin{split} & \bigwedge_{x \in X} (u_x \to (\exists \, \nu)_x) &= \bigcap_{x \in X} \left\{ \{\check{u}_x\} \to \bigcup_{s \in \nu(x)} s \right\} \\ &= \bigcap_{x \in X} \left\{ c \in P | c \cdot \check{u}_x \in \bigcup \nu(x) \right\} \\ &= \bigcap_{x \in X} \left\{ c \in P | c \cdot \check{u}_x \in \nu(x) \right\} \quad \text{ elements of } \nu(x) \text{ are singletons} \\ &\neq \varnothing \end{split}$$

Let $c_0 \in \bigcap_{x \in X} \{c \in P | c \cdot \check{u}_x \in \nu(x)\}$ and further $b \in \mathbb{P}[M]_X$ given by $b_x := \{c \cdot \check{u}_x\}$. We have $b_x \in \nu(x)$ for all $x \in X$ and further

$$\begin{split} & \bigwedge_{x \in X} (u_x \to b_x) &= \bigcap_{x \in X} (u_x \to b_x) \\ &= \bigcap_{x \in X} (\{\check{u}_x\} \to \{c_0 \cdot \check{u}_x\}) \\ &\neq \varnothing \end{split}$$

4.3. Generators.

Definition 4.4. An algebraic subset $M \subseteq \mathcal{S}$ is a generator of \mathcal{S} if it is dense and compact.

Theorem 4.3. Let $M \subseteq \mathscr{S}$ be an algebraic subset. The following are equivalent

i. M is a generator of \mathscr{S} ;

ii. $\operatorname{Asm}_{\mathcal{A}}$ is the regular completion of Asm_{M} .

Proof. Theorem 4.1 and 4.2.

Corollary 4.1. If one and thus both of the equivalent assertions of Theorem 4.3 hold, the closure of Asm_M under isomorphisms is Asm_A 's subcategory of projective objects.

Corollary 4.2. If M and M' are generators then $\operatorname{Asm}_M \simeq \operatorname{Asm}_{M'}$.

5. EXACT COMPLETIONS

Definition 5.1. Let \mathbb{D} be a regular category

1. An equivalence relation in \mathbb{D} is *effective* if it is a kernel pair.

2. \mathbb{D} is *exact* if every equivalence relation is effective.

Example 5.1. Any topos among others.

An exact category in the above sense is sometimes called *Barr-exact* as there is quite a different notion of exactness due to Quillen. We won't need the distinction here as the only exact categories in sight are those in the sense of Barr.

Notation. We shall write

- Ex for the 2-category of exact categories, exact funtors and their natural transformations;
- $|-|: \mathbf{Ex} \to \mathbf{Reg}$ for the forgetful 2-functor.

It is part of the lore that the forgetful 2-functor |-| has a left biadjoint $(-)_{ex/reg}$: $\mathbf{Reg} \to \mathbf{Ex}$. This entails that the forgetful 2-functor $|-|: \mathbf{Ex} \to \mathbf{Lex}$ has a left biadjoint $(-)_{ex/lex}: \mathbf{Lex} \to \mathbf{Ex}$ as well since biadjoints compose.

Let \mathbb{C} be a lex category.

Definition 5.2. We call $\mathbb{C}_{e_{x/lex}} \mathbb{C}$'s *exact completion*.

There is an embedding $\boldsymbol{y} : \mathbb{C} \to \mathbb{C}_{\text{ex/lex}}$ enjoying a universal property structurally identical to the one of regular completions (c.f. Section 2). There is a well-known direct construction of $\mathbb{C}_{\text{ex/lex}}$ [3] which we now recall.

Let \mathbb{C} be a category.

Definition 5.3. Let \mathcal{Q} be a category $s, t : Q_1 \rightrightarrows Q_0$ with two objects and two parallel morphisms. A quiver in \mathbb{C} is a functor $\mathcal{Q} \rightarrow \mathbb{C}$.

As the only quivers of interest here are those internal to a (typically lex) category \mathbb{C} , we shall dispense with the qualifier "in \mathbb{C} ".


Notation. Let X be a quiver.

1. $X(x, x') := s_X, t_X^{-1}(x, x').$

2. We shall write $e: x \rightsquigarrow x'$ for an edge $e \in X(x, x')$.


Definition 5.4. Let X, Y be quivers.

1. A quiver morphism $f: X \to Y$ is given by morphisms $f_0: X_0 \to Y_0$ and $f_1: X_1 \to Y_1$ such that



commutes.

2. Let $f, g: X \to Y$ be quiver morphisms. A homotopy from f to g is a morphism $h: X_0 \to Y_1$ such that the morphism $\langle f_0, g_0 \rangle: X_0 \to Y_0 \times Y_0$ has a lift



with respect to s_Y, t_Y .

Notation. We shall write $h: f \Rightarrow g$ for a homotopy h.

Remark 5.1. Let $f, g: X \to Y$ be quiver morphisms. A homotopy $h: f \Rightarrow g$ is just a family

$$(h(x): f_0(x) \rightsquigarrow g_0(x))_{x \in X_0}$$

of edges of Y_1 . Notice that for any such edge there is an edge $\sigma_Y(h(x)) : g_0(x) \rightsquigarrow f_0(x)$ in the opposite direction.

Suppose now \mathbb{C} is lex.

Definition 5.5. A pseudo-groupoid X in \mathbb{C} is a quiver $s_X, t_X : X_1 \rightrightarrows X_0$ equipped with structural morphisms

- $\begin{array}{l} \ \rho_X : X_0 \to X_1 \text{ such that} \\ \text{i. } s_X \circ \rho_X = \text{id}; \\ \text{ii. } t_X \circ \rho_X = \text{id}; \\ \ \sigma_X : X_1 \to X_1 \text{ such that} \\ \text{i. } s_X \circ \sigma_X = t_X; \\ \text{ii. } t_X \circ \sigma_X = s_X; \\ \ \tau_X : X_1 \circledast X_1 \to X_1 \text{ with } X_1 \circledast X_1 \text{ given by pullback of } s_X \text{ along } t_X, \text{ such that} \end{array}$
 - i. $s_X \circ \tau_X = s_X \circ p_0;$
 - A A A 10)
 - ii. $t_X \circ \tau_X = t_X \circ p_1$.

Entities we call pseudo-groupoids are also (and perhaps better) known as *pseudo-equivalence relations* [2, 3]. As the only pseudo-groupoids of interest here are those internal to a lex category \mathbb{C} , we shall once again dispense with the qualifier "in \mathbb{C} ".

Notice that ρ_X is a local section of $s_X, t_X : X_1 \to X_0 \times X_0$ with respect to the diagonal. The homotopy relation is stable by composition while composition of quiver morphisms among pseudo-groupoids is associative up-to homotopy, so pseudo-groupoids organise themselves in the category $\mathbf{PGrpd}(\mathbb{C})$ along with homotopy classes of quiver morphisms among them. We have

$$\mathbb{C}_{\text{ex/lex}} \simeq \mathbf{PGrpd}(\mathbb{C})$$

[2, 3]. Quite unsurprisingly, the functor $\boldsymbol{y} : \mathbb{C} \to \mathbb{C}_{ex/lex}$ turns then out to be the one sending an object $X \in \mathbb{C}$ on the *discrete* pseudo-groupoid id, id : $X \rightrightarrows X$. As pointed out in [14], pseudo-groupoids give rise to "groupoidal" bicategories internal to \mathbb{C} , with homotopies as 2-cells. So if we restrain from quotienting by the

homotopy relation, we end up with a tricategory where these data organise themselves (taking into account coherence conditions). The homotopy 1-category of this tricategory is then $\mathbf{PGrpd}(\mathbb{C})$ (see also [7]).


6. EXACT COMPLETIONS OF IMPLICATIVE ASSEMBLIES

6.1. The Exact Completion of $Asm_{\mathcal{A}}$. In this subsection we address the ex/reg completion of $Asm_{\mathcal{A}}$ as well as the ex/lex completion of $Asm_{\mathcal{M}}$, exploiting concepts and results found in [9].

Definition 6.1.

1. A chaotic situation is given by two lex categories **S** and \mathbb{C} connected by an adjunction $|-| \dashv \Delta : \mathbf{S} \to \mathbb{C}$.

2. A morphism $f: X \to Y$ in \mathbb{C} is a *pre-embedding* if the square



is a pullback.

3. An object $\Upsilon \in \mathbb{C}$ is *generic* if for every $X \in \mathbb{C}$ there is a pre-embedding $X \to \Upsilon$.

(as a word of caution, we use the symbol Δ for ∇ in op.cit.). A generic object in the above sense is not the same as (although related to) the homonymous fibred artefact. The notion of chaotic situation is an abstract version of what happens when we have a forgetful functor from a concrete category into **Set**.

Proposition 6.1 (Menni). Let \mathbb{C} have a chaotic situation. The following are equivalent

- *i.* \mathbb{C} has a generic object;
- ii. \mathbb{C} has a generic mono.

This is Prop 8.1.8 in op.cit.

Theorem 6.1 (Menni). Let \mathbb{D} be a regular LCCC with generic mono. Then $\mathbb{D}_{ex/reg}$ is a topos.

This is Theorem 11.3.3 in op.cit.

Remark 6.1. (S, id_S) is a generic object in the sense of Definition 6.1.

Theorem 6.2. $(Asm_{\mathcal{A}})_{ex/reg}$ is a topos.

Proof. Proposition 6.1, Theorem 6.1 and Remark 6.1.

Corollary 6.1. Let M be a generator. Then

$$(\mathbf{Asm}_M)_{\mathrm{ex}/\mathrm{lex}} \simeq ((\mathbf{Asm}_M)_{\mathrm{reg}/\mathrm{lex}})_{\mathrm{ex}/\mathrm{reg}} \simeq (\mathbf{Asm}_\mathcal{A})_{\mathrm{ex}/\mathrm{reg}}$$

is a topos.

6.2. Tripos-to-Topos Construction.

Definition 6.2. An *implicative set* (X, \approx) is a set equipped with a *non-standard equality*, that is a predicate $|-\approx -|: X \times X \to \mathcal{A}$ which is

- i. symmetric: $\Vdash \bigwedge_{x,x' \in X} (|x \approx x'| \rightarrow |x' \approx x|);$
- $\text{ii. } transitive: \Vdash \bigwedge_{x,x',x'' \in X} (|x \approx x'| \sqcap |x' \approx x''| \rightarrow |x \approx x''|).$

Implicative sets are the objects of $\mathbf{Set}[\mathcal{A}]$. Notice that we do not necessarily have *reflexivity*, that is $|x \approx x| \in S$ for all $x \in X$. This is in a sense the main point of the construction, as $|x \approx x|$ is interpreted as an *existence* predicate. Accordingly, we call ghosts elements $x \in X$ such that $|x \approx x| \notin S$.

Notation.

$$\begin{split} &- \mathsf{S}_{\mathsf{X}} := \bigwedge_{x,x' \in X} \left(| \, x \approx x' \, | \rightarrow | \, x' \approx x \, | \right); \\ &- \mathsf{T}_{\mathsf{X}} := \bigwedge_{x,x',x'' \in X} \left(| \, x \approx x' \, | \, \sqcap \ | \, x' \approx x'' \, | \rightarrow | \, x \approx x'' \, | \right); \\ &- \mathsf{E}_{X}^{\approx} := | \, x \approx x \, |. \end{split}$$

Definition 6.3. Let (X, \approx) and (Y, \approx) be implicative sets. A functional relation

$$F: (X, \approx) \to (Y, \approx)$$

is a predicate $F: X \times Y \to \mathcal{A}$ which is

i. extensional: $\Vdash \bigwedge_{x,x' \in X} \bigwedge_{y,y' \in Y} (F(x,y) \sqcap |x \approx x'| \sqcap |y \approx y'| \rightarrow F(x',y'));$

ii. strict: $\Vdash \bigwedge_{x \in X} \bigwedge_{y \in Y} (F(x, y) \to \mathsf{E}_X^{\approx}(x) \sqcap \mathsf{E}_Y^{\approx}(y));$

- iii. single-valued: $\Vdash \bigwedge_{x \in X} \bigwedge_{y,y' \in Y} (F(x,y) \sqcap F(x,y') \rightarrow | y \approx y' |);$
- $\text{iv. } total: \Vdash {\textstyle \bigwedge}_{x \in X} \bigg({\sf E}_X^\approx(x) \to \underset{y \in Y_0}{\exists} \left({\sf E}_Y^\approx(y) \sqcap F(x,y) \right) \bigg).$

Two functional relations $F, G: (X, \approx) \to (Y, \approx)$ are *equivalent* if they are isomorphic in $\mathbb{P}[\mathcal{A}]_{X \times Y}$.

Notation.

$$\begin{split} &- \operatorname{Ext}_{\mathcal{F}} := \bigwedge_{x,x' \in X} \bigwedge_{y,y' \in Y} \left(F(x,y) \sqcap |x \approx x'| \sqcap |y \approx y'| \to F(x',y') \right); \\ &- \operatorname{Str}_{\mathcal{F}} := \bigwedge_{x \in X} \bigwedge_{y \in Y} \left(F(x,y) \to \mathsf{E}_X^{\approx}(x) \sqcap \mathsf{E}_Y^{\approx}(y) \right); \\ &- \operatorname{Sv}_{\mathcal{F}} := \bigwedge_{x \in X} \bigwedge_{y,y' \in Y} \left(F(x,y) \sqcap F(x,y') \to |y \approx y'| \right); \\ &- \operatorname{Tot}_{\mathcal{F}} := \bigwedge_{x \in X} \left(\mathsf{E}_X^{\approx}(x) \to \underset{y \in Y_0}{\exists} \left(\mathsf{E}_Y^{\approx}(y) \sqcap F(x,y) \right) \right). \end{split}$$

Proposition 6.2. Two functional relations $f, g: (X, \approx) \to (Y, \approx)$ are equivalent if and only if

$$\Vdash \bigwedge_{x \in X, y \in Y} (F(x, y) \to F(x, y))$$

6.3. The Exact Completion of Asm_M in Context.

Let $M \subseteq \mathscr{S}$ be a generator. We already observed that $(\mathbf{Asm}_M)_{e_X/le_X}$ is a topos (c.f. Corollary 6.1), so the question is now if it is the same topos as $\mathbf{Set}[\mathcal{A}]$.

Notation. In what follows we make systematic use λ -terms, some of them with relatively long arguments lists containing dummy arguments. We shall use the symbol \diamond for dummies in order to improve readability.

6.3.1. A fully faithful functor \mathcal{K} .

Remark 6.2. A pseudo-groupoid X induces a valuation

$$\nu_X : X_0 \times X_0 \longrightarrow \mathfrak{P}_{\bullet}(M)$$
$$(x, x') \mapsto \{\mathsf{E}_{X_1}(e) | e \in X(x, x')\}$$

Lemma 6.1. Let X be a pseudo-groupoid. X_0 equipped with the relation

$$|x \approx_X x'| := \mathsf{E}_{X_0}(x) \sqcap \mathsf{E}_{X_0}(x') \sqcap \exists \nu_X(x, x')$$

is an implicative set.

Proof. We have

$$\mathsf{S}_{\mathsf{X}} \quad = \quad \bigwedge_{x,x' \in X_0} \left(\mathsf{E}_{X_0}(x) \sqcap \mathsf{E}_{X_0}(x') \sqcap \exists \, \nu_X(x,x') \to \mathsf{E}_{X_0}(x') \sqcap \mathsf{E}_{X_0}(x) \sqcap \exists \, \nu_X(x',x) \right)$$

Let $e: x \rightsquigarrow x' \in X_1$ be an edge. We then have the following subquiver of X



with the "opposite" edge $\sigma_X(e): x' \rightsquigarrow x$. Let ξ be a tracker of σ_X . We then have

$$\lambda uvw.u \sqcap v \sqcap (\lambda z.z (\xi (wI)))^{\mathcal{A}} \preccurlyeq \mathsf{S}_{\mathsf{X}}$$

and therefore $\Vdash S_{\mathrm{X}}.$ Next we have

$$\begin{aligned} \mathsf{T}_{\mathbf{X}} &= \bigwedge_{x,x',x'' \in X_{0}} (\\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{X_{0}}(x') \sqcap \exists \nu_{X}(x,x') \sqcap \\ & \mathsf{E}_{X_{0}}(x') \sqcap \mathsf{E}_{X_{0}}(x'') \sqcap \exists \nu_{X}(x',x'') \\ & \to \\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{X_{0}}(x'') \sqcap \exists \nu_{X}(x,x'') \end{aligned}$$

Let $e_1: x \rightsquigarrow x'$ and $e_2: x' \rightsquigarrow x''$ be edges. We then have the following subquiver of X

with the "composed" edge $\tau_X(e_1, e_2) : x \rightsquigarrow x''$. Let χ be a tracker of τ_X . We then have

$$\lambda u \diamond w \diamond v w'. u \sqcap v \sqcap (\lambda z. \chi ((w\mathbf{I}) \sqcap (w'\mathbf{I})))^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{T}_{\mathbf{X}}$$

and therefore $\Vdash \mathsf{T}_{\mathsf{X}}$.

Let $f: X \to Y$ be a morphism of pseudo-groupoids. The relation \mathfrak{R}_f from (X, \approx_X) to (Y, \approx_Y) is given by $\mathfrak{R}_f(x, y) := \mathsf{E}_{X_0}(x) \sqcap \mathsf{E}_{Y_0}(y) \sqcap \exists \nu_Y(y, f_0(x))$

Remark 6.3. Let $(x, y) \in X \times Y$. We have $\Vdash \mathfrak{R}_f(x, y)$ if and only if there is an edge $e: y' \rightsquigarrow f_0(x)$ in Y and $\mathfrak{R}_f(x, y) = \bot$ otherwise.

Remark 6.4. Let $\mathsf{Rf} := \bigwedge_{x \in X_0, y \in Y_0} (\mathfrak{R}_f(x, y) \to |y| \approx_Y |f_0(x)|)$. We have

$$\begin{array}{rcl} \mathsf{Rf} & = & \displaystyle{\bigwedge_{x \in X_0, y \in Y_0}}(\\ & & \mathsf{E}_{X_0}(x) \sqcap \mathsf{E}_{Y_0}(y) \sqcap \exists \, \nu_Y(y, f_0(x)) \\ & \rightarrow \\ & & \mathsf{E}_{Y_0}(y) \sqcap \mathsf{E}_{Y_0}(f_0(x)) \sqcap \exists \, \nu_Y(y, f_0(x)) \\ & & \end{pmatrix} \\ \end{array}$$

Suppose ξ tracks f_0 . Then

 $(\lambda uvw.v \sqcap (\xi u) \sqcap w)^{\mathcal{A}} \preccurlyeq \mathsf{Rf}$


and therefore $\Vdash Rf$.

Lemma 6.2. \mathfrak{R}_f is extensional.

Proof. We have

$$\begin{split} \mathsf{Ext}_{\mathfrak{R}_{f}} &= \bigwedge_{x,x' \in X_{0}} \bigwedge_{y,y' \in Y_{0}} (\\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{Y_{0}}(y) \sqcap \exists \nu_{Y}(y,f_{0}(x)) \sqcap \\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{X_{0}}(x') \sqcap \exists \nu_{X}(x,x') \sqcap \\ & \mathsf{E}_{Y_{0}}(y) \sqcap \mathsf{E}_{Y_{0}}(y') \sqcap \exists \nu_{Y}(y,y') \\ & \rightarrow \\ & \mathsf{E}_{X_{0}}(x') \sqcap \mathsf{E}_{Y_{0}}(y') \sqcap \exists \nu_{Y}(y',f_{0}(x')) \end{split}$$

Let $x, x'' \in X_0$ be vertices of X and $y, y' \in Y_0$ be vertices of Y. Let $e : x \rightsquigarrow x'$ be an edge of X. Let $e_1 : y \rightsquigarrow f_0(x)$ and $e_2 : y \rightsquigarrow y'$ be edges of Y (notice that besides their "types" we did not make any hypotheses on the vertices and edges in question). We then have the following subquiver of Y



Suppose χ tracks f_1 , κ tracks σ_Y and ϖ tracks τ_Y . Then

$$(\lambda \diamond \diamond w \diamond vw' \diamond v'w''.v \sqcap v' \sqcap \lambda z.z \, (\varpi \left((\varpi \left(\kappa \left(w''\mathbf{I} \right) \right) \sqcap \left(w\mathbf{I} \right) \sqcap \left(\chi \left(w'\mathbf{I} \right) \right) \right))))^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Ext}_{\mathfrak{R}_{f}}$$

and therefore $\Vdash \mathsf{Ext}_{\mathfrak{R}_f}$.

Lemma 6.3. \Re_f is strict.

Proof. We have

$$\begin{array}{lll} \mathsf{Str}_{\mathfrak{R}_{f}} & = & \displaystyle \bigwedge_{x \in X_{0}} \bigwedge_{y \in Y_{0}} (\\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{Y_{0}}(y) \sqcap \exists \, \nu_{Y}(y, f_{0}(x)) \\ & \rightarrow \\ & \mathsf{E}_{X_{0}}(x) \sqcap \exists \, \nu_{X}(x, x) \sqcap \\ & \mathsf{E}_{Y_{0}}(y) \sqcap \exists \, \nu_{Y}(y, y) \end{array}$$

Suppose κ tracks ρ_X while κ' tracks ρ_Y . Then

$$(\lambda uv \diamond . u \sqcap \lambda z. z(\kappa u) \sqcap v \sqcap \lambda z. z(\kappa' v))^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Str}_{\mathfrak{R}_f}$$


and therefore $\Vdash \mathsf{Str}_{\mathfrak{R}_f}$.

Lemma 6.4. \mathfrak{R}_f is single valued.

Proof. We have

$$\begin{split} \mathsf{Sv}_{\mathfrak{R}_{f}} &= \bigwedge_{x \in X_{0}} \bigwedge_{y, y' \in Y_{0}} (\\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{Y_{0}}(y) \sqcap \exists \, \nu_{Y}(y, f_{0}(x)) \sqcap \\ & \mathsf{E}_{X_{0}}(x) \sqcap \mathsf{E}_{Y_{0}}(y') \sqcap \exists \, \nu_{Y}(y', f_{0}(x)) \\ & \rightarrow \\ & \mathsf{E}_{Y_{0}}(y) \sqcap \mathsf{E}_{Y_{0}}(y') \sqcap \exists \, \nu_{Y}(y, y') \\ &) \end{split}$$

Let $e: y \rightsquigarrow f_0(x)$ and $e': y' \rightsquigarrow f_0(x)$ be edges of Y. We then have the subquiver of Y



Suppose ξ tracks σ_Y and ϖ tracks τ_Y . Then

$$(\lambda \diamond vw \diamond v'w'.v \sqcap v' \sqcap \lambda z.z \left(\varpi \left((w\mathbf{I}) \sqcap \left(\xi \left(w'\mathbf{I} \right) \right) \right) \right)^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Sv}_{\mathfrak{R}_f}$$

and therefore $\Vdash \mathsf{Sv}_{\mathfrak{R}_f}$.

Lemma 6.5. \Re_f is total.

Proof. We have

$$\begin{aligned} \mathsf{Tot}_{\mathfrak{R}_{f}} &= \bigwedge_{x \in X_{0}} (\\ & \mathsf{E}_{X_{0}}(x) \sqcap \exists \nu_{X}(x, x) \\ & \to \\ & \exists \\ y \in Y_{0}} \left(\mathsf{E}_{Y_{0}}(y) \sqcap \exists \nu_{Y}(y, y) \sqcap \mathsf{E}_{X_{0}}(x) \sqcap \exists \nu_{Y}(y, f_{0}(x)) \right) \\ &) \end{aligned}$$

Suppose χ tracks f_0 while κ tracks ρ_Y . We know $y \in Y_0$ that fits, namely $f_0(x)$. Then

 $(\lambda uv.(\chi u) \sqcap \lambda z.\kappa(\chi u) \sqcap u \sqcap \lambda z.\kappa(\chi u))^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Tot}_{\mathfrak{R}_f}$

and therefore $\Vdash \mathsf{Tot}_{\mathfrak{R}_f}$.

Proposition 6.3. The assignment

$$\begin{array}{rcl} \mathcal{K} : \left(\mathbf{Asm}_{M}\right)_{\mathrm{ex}\,/\,\mathrm{lex}} & \longrightarrow & \mathbf{Set}[\mathcal{A}] \\ & X & \mapsto & \left(X_{0}, \approx_{X}\right) \end{array}$$

extends to a faithful functor with action on morphisms $f: X \to Y$ given by

$$\mathcal{K}(f)(x,y) := \mathfrak{R}_f(x,y)$$

Proof. (X_0, \approx_X) is an implicative set by Lemma 6.1 while \mathfrak{R}_f is a functional relation by Lemmas 6.2, 6.3, 6.4 and 6.5. Let $g: X \to Y$ be a further morphism of pseudo-groupoids and $h: f \Rightarrow g$ be a homotopy. There is thus an edge $h(x): f_0(x) \rightsquigarrow g_0(x)$ for every vertex $x \in X_0$ (c.f. Remark 5.1), which entails $\Vdash \exists \nu_Y(f_0(x), g_0(x))$ and therefore

$$\Vdash |f_0(x) \approx_Y g_0(x)|$$

We further have $\Vdash \bigvee_{x \in X_0} \bigvee_{y \in Y_0} (\mathfrak{R}_f(x, y) \to |y \approx_Y f_0(x)|)$ (c.f. Remark 6.4). Now $\mathfrak{R}_f(x, y)$ is either \bot (so it implies anything) or $\Vdash \mathfrak{R}_f(x, y)$ (c.f. Remark 6.3). In the latter case we have $\Vdash |y \approx_Y f_0(x)|$ by (the implicative) modus ponens and further $\Vdash |y \approx_Y g_0(x)|$ by symmetry and transitivity, hence

$$\Vdash \mathfrak{R}_f \to \mathfrak{R}_c$$

which entails that \mathcal{K} is well-defined on morphisms (c.f. Proposition 6.2). It is easy to see that \mathcal{K} preserves identities and composition.

Let $u, v: X \to Y$ be morphisms of pseudo-groupoids. Suppose $\mathcal{K}(u) = \mathcal{K}(v)$, that is

$$\Vdash \mathfrak{R}_u \leftrightarrow \mathfrak{R}_v \quad (\star)$$

Given $x \in X_0$ (*) implies $\Vdash |u_0(x)| \approx Y_0 |v_0(x)|$, so there is an edge $e_x : u_0(x) \rightsquigarrow v_0(x)$. All these edges are indexed by X and organise themselves in a homotopy $h : u \Rightarrow v$, hence \mathcal{K} is faithful. \Box

Notation. Still seeking the greater good by making things human-readable, we shall also add the let x = e in macro to our λ -calculus.

Proposition 6.4. The functor $\mathcal{K} : (\mathbf{Asm}_M)_{\text{ex / lex}} \to \mathbf{Set}[\mathcal{A}]$ is full.

Proof. Let $X, Y \in (\mathbf{Asm}_M)_{\text{ex/lex}}$ and $F : \mathcal{K}(X) \to \mathcal{K}(Y)$ be a functional relation. Unravelling the definition yields in this case

$$\mathbb{H}\mathsf{Sv}_{\mathsf{F}} = \bigwedge_{x \in X_0} \bigwedge_{y, y' \in Y_0} \left(F(x, y) \sqcap F(x, y') \to \mathsf{E}_{Y_0}(y) \sqcap \mathsf{E}_{Y_0}(y') \sqcap \exists \nu_Y(y, y') \right)$$

$$\mathbb{H}\mathsf{Tot}_{\mathsf{F}} = \bigwedge_{x \in X_0} \left(\left(\mathsf{E}_{X_0}(x) \sqcap \exists \nu_X(x, x) \right) \to \underset{y \in Y_0}{\exists} \left(\mathsf{E}_{Y_0}(y) \sqcap \exists \nu_Y(y, y) \sqcap F(x, y) \right) \right)$$

As implicative sets in the image of \mathcal{K} do not have ghosts, F can be presented as a map

$$\begin{array}{rcl} \tilde{F}: X_0 & \longrightarrow & \mathfrak{P}(Y_0) \\ & x & \mapsto & \{y \in Y_0 | F(x,y) \in \mathscr{S}\} \end{array}$$

Assuming choice we can extract a map $f_0: X_0 \to Y_0$ from \tilde{F} . Suppose κ tracks ρ_X . For any $x \in X_0$ there is y_x (in general not unique) such that

$$\begin{split} \mathbf{let} \ t &= \lambda p.\mathsf{Tot}_{\mathsf{F}}(p \sqcap (\kappa p)) \ \mathbf{in} \\ & (\lambda q. \langle \boldsymbol{\pi}_0, \boldsymbol{\pi}_2 \rangle \left((tq) \mathsf{I} \right))^{\mathcal{A}} \quad \preccurlyeq \quad \bigwedge_{x \in X_0} \left(\mathsf{E}_{X_0}(x) \to \left(\mathsf{E}_{Y_0}(y_x) \sqcap F(x, y_x) \right) \right) \end{split}$$

thus

$$\Vdash \bigwedge_{x \in X_0} (\mathsf{E}_{X_0}(x) \to (\mathsf{E}_{Y_0}(y_x) \sqcap F(x, y_x)))$$

But $\Vdash F(x, f_0(x))$ for all $x \in X$ by definition of f_0 , hence

let
$$t' = \lambda r. Sv_F ((\pi_1 r) \sqcap F(x, f_0(x)))$$
 in

$$(\lambda s. \boldsymbol{\pi}_1(t's))^{\mathcal{A}} \quad \preccurlyeq \quad \bigwedge_{x \in X_0} \left(\mathsf{E}_{Y_0}(y_x) \sqcap F(x, y_x) \right) \to \mathsf{E}_{Y_0}(f_0(x))$$

thus

$$\Vdash \bigwedge_{x \in X_0} \left(\mathsf{E}_{Y_0}(y_x) \sqcap F(x, y_x) \right) \to \mathsf{E}_{Y_0}(f_0(x))$$

But then

$$\Vdash \bigwedge_{x \in X_0} \left(\mathsf{E}_{X_0}(x) \to \mathsf{E}_{Y_0}(f_0(x)) \right)$$

so f_0 is tracked.

Let $e: x \rightsquigarrow x'$ be an edge of X, so we have $\Vdash |x \approx_X x'|$. We have $\Vdash F(x, f_0(x))$ and $\Vdash F(x', f_0(x'))$ by definition of f_0 , hence $\Vdash F(x, f_0(x'))$ by extensionality and therefore $\Vdash |f_0(x) \approx_Y f_0(x')|$ by single-valuedness. But then $Y(f_0(x), f_0(x')) \neq \emptyset$, so given $e: x \rightsquigarrow x'$ we have

$$((f_0 \times f_0) \circ s_X, t_X)(e) = (s_Y, t_Y \circ f_1)(e)$$

for any map such that

$$f_1(e) \in Y(f_0(x), f_0(x'))$$

Assuming choice such a map always exists, what remains to be shown is that it is tracked. Consider the map

$$\begin{aligned} \phi : \mathcal{A} & \longrightarrow & \mathcal{A} \\ a & \mapsto & \begin{cases} \mathsf{E}_{Y_1}(f_1(e)) & \text{ if there are } x, x' \in X_0 \text{ and } e \in X(x, x') \text{ such that } a = \mathsf{E}_{X_1}(e) \\ \top & \text{ otherwise} \end{cases}$$

We have

$$\lambda \phi = \bigwedge_{a \in \mathcal{A}} (a \to \phi(a))$$

Notice that $\lambda \phi \in \mathscr{S}$ as either $a \in \mathscr{S}$ and $\phi(a) \in \mathscr{S}$ or $\phi(a) = \top$. Furthermore, we have

$$(\lambda p. (\boldsymbol{\lambda}\phi) p)^{\mathcal{A}} \quad \preccurlyeq \quad \bigwedge_{e \in X_1} (\mathsf{E}_{X_1}(e) \to \mathsf{E}_{Y_1}(f_1(e)))$$

hence

$$\Vdash \bigwedge_{e \in X_1} \left(\mathsf{E}_{X_1}(e) \to \mathsf{E}_{Y_1}(f_1(e)) \right)$$

6.3.2. Is K always essentially surjective?

Let (X, \approx) be an implicative set and $\nu : X \times X \to \mathfrak{P}_{\bullet}(M)$ be a valuation such that $\approx^+ \cong_{X \times X} \exists \nu$ (such a valuation always exists since M is dense). Assuming choice there is a map $c : X^+ \to M$ such that $c(x) \in \nu(x, x)$ for all $x \in X^+$.

Notation.

- 1. Equ_X ⁺ := { $(x, x') \in X \times X$ | $|x \approx x'| \in \mathscr{S}$ } (the set of valid equalitites on X);
- 2. $X^+ := \{x \in X | \mathsf{E}^{\approx}_X(x) \in \mathscr{S}\}$ (the set of X's non-ghosts);
- $3. \approx^+ := \approx_{|_{\mathrm{Equ}_X} +};$

4. $\mathsf{E}_{X}^{+} := (\mathsf{E}_{X}^{\approx})_{|_{X}^{+}}$.

Lemma 6.6. The M-assemblies

$$\begin{array}{l} - \ \hat{X}_1 \ given \ by \ \underline{\hat{X}_1} := \sum_{(x,x') \in \operatorname{Equ}_X^+} \nu(x,x') \ and \ \mathsf{E}_{\hat{X}_1}((x,x'),m) := c(x) \sqcap m \sqcap c(x'); \\ - \ \hat{X}_0 \ given \ by \ \underline{\hat{X}_0} := X^+ \ and \ \mathsf{E}_{\widehat{X}_0}(x) = c(x) \end{array}$$

along with the maps $s := \pi_0 \circ \pi_0$ and $t := \pi_1 \circ \pi_0$ form a pseudo-groupoid \hat{X} .

Proof. \widehat{X}_1 is an *M*-assembly since *M* is algebraic. The source and target maps *s* and *t* are tracked by π_0 and π_2 , respectively. We have the local section of *s*, *t*

$$\begin{array}{ccc} \rho: & \underline{\hat{X}_0} \longrightarrow & \underline{\hat{X}_1} \\ x & \mapsto & ((x,x), c(x)) \end{array}$$

tracked by id. Next we have the iso

$$\approx^+ \cong_{X \times X} \exists \nu \quad (\star)$$

by density, which entails

$$\Vdash \mathsf{S}'_{\mathsf{X}} := \bigwedge_{x,x' \in X^+} \left(\exists \, \nu(x,x') \to \exists \, \nu(x',x) \right)$$

Let $x, x' \in X^+$ and $m \in \nu(x, x')$. We then have $(\mathsf{S}'_X(\lambda z. zm)^{\mathcal{A}}) \mathbf{I} \in \nu(x', x)$, hence the "twist" map

$$\sigma: \quad \underline{\widehat{X_1}} \longrightarrow \quad \underline{\widehat{X_1}} \\ ((x, x'), m) \quad \mapsto \quad \left((x', x), \left(\mathsf{S}'_{\mathsf{X}}(\lambda z. zm)^{\mathcal{A}} \right) \mathsf{I} \right)$$

tracked by $(\lambda umv.v \sqcap ((\mathsf{S}'_X \lambda z.zm) \mathbf{I}) \sqcap u)^{\mathcal{A}}$. But (\star) also entails

$$\Vdash \mathsf{T}'_{\mathsf{X}} := \bigwedge_{x,x',x''} \left(\exists \, \nu(x,x') \sqcap \exists \, \nu(x',x'') \to \exists \, \nu(x,x'') \right)$$

Let $x, x', x'' \in X^+$, $m \in \nu(x, x')$ and $n \in \nu(x', x'')$. We then have

$$\left(\mathsf{T}'_{\mathsf{X}}(\lambda z.zm)^{\mathcal{A}}(\lambda z.zn)^{\mathcal{A}}\right)\mathsf{I} \in \nu(x,x'')$$

hence the "connecting" map

$$\tau: \widehat{\underline{X}_1} \circledast \widehat{\underline{X}_1} \longrightarrow \widehat{\underline{X}_1}$$

$$(((x, x'), m), ((x', x''), n)) \mapsto ((x, x''), (\mathsf{T}'_X(\lambda z. zm)^{\mathcal{A}}(\lambda z. zn)^{\mathcal{A}}) \mathsf{I})$$

tracked by $(\lambda u \diamond \diamond mnv.u \sqcap (\mathsf{T}'_{\mathsf{X}}(\lambda z.zm)^{\mathcal{A}}(\lambda z.zn)^{\mathcal{A}}) \amalg \sqcap v)^{\mathcal{A}}.$

We thus have $\mathcal{K}(\hat{X}) = (X^+, \asymp)$ where

$$\mid x \asymp x' \mid = \begin{cases} c(x) \sqcap c(x') \sqcap \exists \nu(x, x') & (x, x') \in \mathrm{Equ}_{\mathrm{X}} + \\ \bot & \text{otherwise} \end{cases}$$

as $\nu = \nu_{B_X}$ by construction (c.f. Lemma 6.6). There are no ghosts here.

Lemma 6.7. The relation $K: (X^+, \asymp) \to (X, \approx)$ given by

$$\begin{array}{rccc} K: X^+ \times X & \longrightarrow & \mathcal{A} \\ (x, x') & \mapsto & x \approx x' \end{array}$$

is functional.

Proof. We have

$$\mathsf{Ext}_{\mathsf{K}} \quad = \quad \bigwedge_{x_1, x_2 \in X^+} \bigwedge_{x_1', x_2' \in X} \left(| \, x_1 \approx x_1' \, | \, \sqcap \mid x_1 \asymp x_2 \mid \sqcap | \, x_1' \approx x_2' \mid \rightarrow \mid x_2 \approx x_2' \mid \right)$$

Suppose $(x_1, x_2) \notin \operatorname{Equ}_X^+$. We then have $|x_1 \asymp x_2| = \bot$ so $|x_1 \approx x_1'| \sqcap |x_1 \asymp x_2| \sqcap |x_1' \approx x_2'| \rightarrow |x_2|$

$$x_1 \approx x_1' \mid \Box \mid x_1 \asymp x_2 \mid \Box \mid x_1' \approx x_2' \mid \rightarrow \mid x_2 \approx x_2' \mid = \top$$

in those cases, hence

$$\mathsf{Ext}_{\mathsf{K}} \quad = \quad \bigwedge_{(x_1, x_2) \in \mathrm{Equ}_X^+} \bigwedge_{x_1', x_2' \in X} \left(| \, x_1 \approx x_1' \, | \, \sqcap \mid x_1 \asymp x_2 \mid \sqcap | \, x_1' \approx x_2' \, | \, \rightarrow \mid x_2 \approx x_2' \, | \right)$$

But $\mathsf{Ext}_{\mathrm{K}}$ is equivalent to

$$\mathsf{Ext}'_{\mathsf{K}} := \bigwedge_{(x_1, x_2) \in \mathrm{Equ}_X^+} \bigwedge_{x_1', x_2' \in X} (|x_1 \approx x_1'| \, \sqcap c(x_1) \sqcap c(x_2) \sqcap |x_1 \approx x_2| \sqcap |x_1' \approx x_2'| \to |x_2 \approx x_2'|)$$

by density and furthermore

$$(\lambda u \diamond \diamond v w. \mathsf{T}_{\mathsf{X}} (\mathsf{S}_{\mathsf{X}} v) (\mathsf{T}_{\mathsf{X}} u v))^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Ext}'_{\mathsf{K}}$$

Next we have

$$\mathsf{Str}_{\mathcal{K}} \hspace{.1 in} = \hspace{.1 in} \bigwedge_{x \in \mathrm{Equ}_{\mathcal{X}}} + \underset{x' \in X}{\bigwedge} \left(| \hspace{.06cm} x \approx x' \hspace{.05cm} | \hspace{.06cm} \to | \hspace{.06cm} x \asymp x| \sqcap | \hspace{.06cm} x' \approx x' \hspace{.05cm} | \right)$$

But $\mathsf{Str}_{\mathrm{K}}$ is equivalent to

$$\mathsf{Str}'_{\mathcal{K}} \quad = \quad \bigwedge_{x \in \mathrm{Equ}_{\mathcal{K}}} \, \bigwedge_{+ \, x' \in X} \left(| \, x \approx x' \, | \to c(x) \sqcap \mathsf{E}^+_X(x) \sqcap \mathsf{E}^\approx_X(x') \right)$$

by density. Consider the map

$$\begin{array}{cccc} d: \mathcal{A} & \longrightarrow & \mathcal{A} \\ a & \mapsto & \left\{ \begin{array}{ccc} c(x) & & \text{there is } x \in X^+ \text{such that } a = \mathsf{E}_X^+(x) \\ \top & & \text{otherwise} \end{array} \right. \end{array}$$

We have $\boldsymbol{\lambda} d \in \mathscr{S}$ and furthermore

$$\begin{split} \mathbf{let} \ \varepsilon &= \lambda p. \mathsf{T}_{\mathsf{X}} p \left(\mathsf{S}_{\mathsf{X}} p \right) \ \mathbf{in} \\ \mathbf{let} \ \varepsilon' &= \lambda q. \left(\boldsymbol{\lambda} d \right) \left(\varepsilon q \right) \mathbf{in} \\ & \left(\lambda r. (\varepsilon' r) \sqcap \left(\varepsilon r \right) \sqcap \left(\mathsf{T}_{\mathsf{X}} \left(\mathsf{S}_{\mathsf{X}} p \right) p \right) \right)^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Str}_{\mathsf{K}}' \end{split}$$

Next we obviously have

$$\Vdash \mathsf{Sv}_{\mathsf{K}} \quad = \quad \bigwedge_{x_1 \in X^+} \bigwedge_{x_1', x_2' \in X} \left(| \, x \approx x_1' \, | \, \sqcap \, | \, x \approx x_2' \, | \, \rightarrow \, | \, x_1' \approx x_2' \, | \right)$$

Finally we have

$$\begin{aligned} \mathsf{Tot}_{\mathcal{K}} &= \bigwedge_{x \in X^+} \left(|x \asymp x| \to \underset{x' \in X}{\exists} (\mathsf{E}_X^{\approx}(x') \sqcap K(x, x')) \right) \\ &= \bigwedge_{x \in X^+} \left(\left(c(x) \sqcap \exists \nu(x, x) \right) \to \underset{x' \in X}{\exists} (\mathsf{E}_X^{\approx}(x') \sqcap |x \approx x'|) \right) \end{aligned}$$

and furthermore $\Vdash \xi := \int_{x \in X} \left(\exists \nu(x, x) \to \mathsf{E}_X^+(x) \right)$ by density, hence $(\lambda \diamond p.\xi p)^{\mathcal{A}} \preccurlyeq \mathsf{Tot}_{\mathsf{K}}$

Proposition 6.5. The morphism $[K] : (X^+, \asymp) \to (X, \approx)$ in **Set**[\mathcal{A}] is is internally injective.

Proof. Let

$$\begin{split} & \mathsf{Inj} \quad := \quad \bigwedge_{x_1, x_2 \in X^+} \bigwedge_{x \in X} \left(K(x_1, x) \sqcap K(x_2, x) \to \mid x_1 \asymp x_2 \mid \right) \\ & = \quad \bigwedge_{x_1, x_2 \in X^+} \bigwedge_{x \in X} \left(\mid x_1 \approx x \mid \sqcap \mid x_2 \approx x \mid \to c(x_1) \sqcap c(x_2) \sqcap \exists \, \nu(x_1, x_2) \right) \end{split}$$

Consider again the map

$$\begin{array}{rccc} d: \mathcal{A} & \longrightarrow & \mathcal{A} \\ a & \mapsto & \left\{ \begin{array}{cc} c(x) & & \text{there is } x \in X^+ \text{such that } a = \mathsf{E}_X^+(x) \\ \top & & \text{otherwise} \end{array} \right. \end{array}$$

used in the proof of Lemma 6.7. We have in particular $\lambda d \in \mathscr{S}$. Furthermore

$$\Vdash \chi := \bigwedge_{x_1, x_2 \in X^+} \left(|x_1 \approx x_2| \to \exists \nu(x_1, x_2) \right)$$

by density, hence

$$\begin{split} \mathbf{let} \ & j = \lambda p q. \mathsf{T}_{\mathsf{X}} p \left(\mathsf{S}_{\mathsf{X}} q \right) \ \mathbf{in} \\ \mathbf{let} \ & e_1 = \lambda p. \mathsf{T}_{\mathsf{X}} \left(\mathsf{S}_{\mathsf{X}} p \right) p \ \mathbf{in} \\ \mathbf{let} \ & e_2 = \lambda p. \mathsf{T}_{\mathsf{X}} p \left(\mathsf{S}_{\mathsf{X}} p \right) \ \mathbf{in} \\ \mathbf{let} \ & \delta = \lambda p. \left(\boldsymbol{\lambda} d \right) p \ \mathbf{in} \\ & \left(\lambda p q. \mathbf{let} \ \varepsilon = j p q \ \mathbf{in} \left(\left(\delta(e_1 \varepsilon) \right) \sqcap \left(\delta(e_2 \varepsilon) \right) \sqcap \left(\chi \varepsilon \right) \right) \right)^{\mathcal{A}} \quad \preccurlyeq \quad \mathsf{Inj} \end{split}$$

Alas, [K] may fail to be internally surjective. For internal surjectivity means in our case that

$$\bigwedge_{x' \in X} \left(\mathsf{E}_X^{\approx}(x') \to \underset{x \in X^+}{\exists} K(x, x') \right) \qquad (\star)$$

is in \mathscr{S} , yet in general we don't have control over what happens on ghosts. Notice that if \mathcal{A} is compatible with joins, x' being a ghost means $\mathsf{E}_X^{\approx}(x') = \bot$, hence factors indexed by ghosts evaluate to \top so we can "track" (\star). [K] is then internally surjective as well, an exemple for this phenomenon being the well-known equivalence **PAss** $\simeq \mathcal{E}ff$ [12, 9]. On the other hand if we co-restrict \mathcal{K} to "ghostless" implicative sets, we get an equivalence of categories.

Theorem 6.3. Let $\mathbf{Set}[\mathcal{A}]^+$ be the full subcategory of $\mathbf{Set}[\mathcal{A}]$ with objects implicative sets (X, \approx) such that $\mathsf{E}_X^{\approx}(x) \in \mathscr{S}$ for all $x \in X$. We have

$$(\mathbf{Asm}_M)_{\mathrm{ex}\,/\,\mathrm{lex}} \ \simeq \ \mathbf{Set}[\mathcal{A}]^+$$

Proof. The image of the functor $\mathcal{K} : (\mathbf{Asm}_M)_{\mathrm{ex/lex}} \to \mathbf{Set}[\mathcal{A}]$ is contained in $\mathbf{Set}[\mathcal{A}]^+$. Its co-restriction $\mathcal{K}^+ : (\mathbf{Asm}_M)_{\mathrm{ex/lex}} \to \mathbf{Set}[\mathcal{A}]^+$ is thus full (c.f. Proposition 6.4), faithful (c.f. Proposition 6.3 and internally injective (c.f. Proposition 6.5). Let

$$\mathsf{Surj} \quad := \quad \bigwedge_{x' \in X} \left(\mathsf{E}^\approx_X(x') \to \underset{x \in X^+}{\exists} K(x,x') \right)$$

Suppose $\mathsf{E}_X^{\approx}(x')$. We then have $x' \in X^+$ so in particular $K(x', x') = \mathsf{E}_X^{\approx}(x')$, thus id $\preccurlyeq \mathsf{Surj}$. Hence [K] is iso, thus \mathcal{K}^+ is an equivalence of categories.

Wrapping it up, the following picture begins to emerge



All the categories involved but Asm_M and Asm_A are topoi. In particular, we see that Asm_A is a subcategory of $\operatorname{Set}[\mathcal{A}]$ yet the embedding is not the trivial one. As a matter of fact, assemblies embed as kernel pairs of their projective covers. A characterisation of those implicative algebras for which we have $(\operatorname{Asm}_M)_{ex/lex} \simeq \operatorname{Set}[\mathcal{A}]$ is an open question. We may have more to say about this circle of ideas in subsequent elaborations.

References

- [1] Michael Barr, Pierre Grillet, Donovan Van Osdol, and Michael Barr. Exact categories. Springer, 1971.
- [2] Aurelio Carboni. Some free constructions in realizability and proof theory. Journal of pure and applied algebra, 103(2):117– 148, 1995.
- Magno. Celia The exact [3] Aurelio Carboni and free category on \mathbf{a} left exact one. Journal of the Australian Mathematical Society, 33(3):295–301, 1982.
- [4] Félix Castro, Alexandre Miquel, and Krzysztof Worytkiewicz. Implicative assemblies. Submitted, 2023.
- [5] Martin Hyland. The effective topos. In Studies in Logic and the Foundations of Mathematics, volume 110, pages 165–216. Elsevier, 1982.
- [6] Martin Hyland, Peter Johnstone, and Andrew Pitts. Tripos theory. In Mathematical Proceedings of the Cambridge philosophical society, volume 88, pages 205–232. Cambridge University Press, 1980.
- [7] Yoshiki Kinoshita and John Power. Category theoretic structure of setoids. *Theoretical Computer Science*, 546:145–163, 2014.
- [8] Maria Emilia Maietti and Giuseppe Rosolini. Unifying exact completions. Applied Categorical Structures, 23:43–52, 2015.
- [9] Matías Menni. Exact completions and toposes. PhD thesis, 2000.
- [10] Alexandre Miquel. Implicative algebras: a new foundation for realizability and forcing. Mathematical Structures in Computer Science, 30(5):458–510, 2020.
- [12] Edmund Robinson and Giuseppe Rosolini. Colimit completions and the effective topos. The Journal of Symbolic Logic, 55(2):678–699, 1990.
- [13] Giuseppe Rosolini, Fabio Pasquali, and Maria Emilia Maietti. Elementary quotient completions, church's thesis, and particular assemblies. Logical Methods in Computer Science, 15, 2019.
- [14] Michael Shulman. The derivator of setoids. Cahiers de topologie et géométrie différentielle catégoriques, 2023.

Instituto de Matemática y Estadística Rafael Laguardia (IMERL), Facultad de Ingeniería, Universidad de la República, Julio Herrera y Reissig 565, C.P. 11300 Montevideo, Uruguay

LABORATOIRE DE MATHÉMATIQUES (LAMA), UNIVERSITÉ SAVOIE MONT BLANC, BÂTIMENT LE CHABLAIS, CAMPUS SCIEN-TIFIQUE, 73376 LE BOURGET DU LAC, FRANCE