The uniform supertrees with the extremal spectral radius^{*}

Guanglong Yu^a Lin Sun^{$a\dagger$} Hailiang Zhang^b Gang Li^c

^aDepartment of Mathematics, Lingnan Normal University, Zhanjiang, Guangdong, 524048, P.R.China

^bDepartment of Mathematics, Taizhou University, Linhai, Zhejiang, 317000, P.R.China

^cCollege of Mathematics and Systems Science, Shandong University of Science and Technology,

Qingdao, Shandong, 266590, P.R.China

Abstract

For a hypergraph $\mathcal{G} = (V, E)$ consisting of a nonempty vertex set $V = V(\mathcal{G})$ and an edge set $E = E(\mathcal{G})$, its adjacency matrix $\mathcal{A}_{\mathcal{G}} = [(\mathcal{A}_{\mathcal{G}})_{ij}]$ is defined as $(\mathcal{A}_{\mathcal{G}})_{ij} = \sum_{e \in E_{ij}} \frac{1}{|e|-1}$, where $E_{ij} = \{e \in E : i, j \in e\}$. The spectral radius of a hypergraph \mathcal{G} , denoted by $\rho(\mathcal{G})$, is the maximum modulus among all eigenvalues of $\mathcal{A}_{\mathcal{G}}$. In this paper, among all k-uniform $(k \geq 3)$ supertrees with fixed number of vertices, the supertrees with the maximum, the second maximum and the minimum spectral radius are completely determined, respectively.

AMS Classification: 05C50

Keywords: Spectral radius; supertree; hypergraph

1 Introduction

In the past twenty years, some different connectivity hypermatrices (or tensors) had been defined and been developed to explore spectral hypergraph theory [1]-[5], [9], [12]-[20], [24], [27]-[32], [35, 36]. Using different hypermatrices for general hypergraphs, many interesting spectral properties have been studied that many properties of spectral graph theory have been extended to spectral hypergraph theory. A lot of interesting results have emerged and the spectra of hypergraphs have been further studied [6, 7, 8, 10, 21, 22, 25, 34], [39]-[42]. In [2], A. Banerjee introduced an adjacency matrix and use its spectrum so that some spectral and structural properties of hypergraphs are revealed. In this paper, we go on studying the spectra of hypergraphs according to the adjacency matrix introduced in [2].

Now we recall some notations and definitions related to hypergraphs. For a set S, we denote by |S| its cardinality. A hypergraph $\mathcal{G} = (V, E)$ consists of a nonempty vertex set $V = V(\mathcal{G})$ and an edge set $E = E(\mathcal{G})$, where each edge $e \in E(\mathcal{G})$ is a subset of $V(\mathcal{G})$ containing at least two vertices. The cardinality $n = |V(\mathcal{G})|$ is called the order; $m = |E(\mathcal{G})|$ is called the edge number of hypergraph \mathcal{G} . Denote by t-set a set with size (cardinality) t. We say that a hypergraph \mathcal{G} is uniform if its every edge has the same size, and call it k-uniform if its every edge has size k (i.e. every edge is a k-subset). It is known that a 2-uniform graph is always called a ordinary graph or graph for short.

For a hypergraph \mathcal{G} , we define $\mathcal{G} - e \ (\mathcal{G} + e)$ to be the hypergraph obtained from \mathcal{G} by deleting the edge $e \in E(\mathcal{G})$ (by adding a new edge e if $e \notin E(\mathcal{G})$); for an edge subset $B \subseteq E(\mathcal{G})$, we define $\mathcal{G} - B$ to be the hypergraph obtained from \mathcal{G} by deleting each edge $e \in B$; for a vertex subset $S \subseteq V(\mathcal{G})$, we define $\mathcal{G} - S$ to be the hypergraph obtained from \mathcal{G} by deleting all the vertices in S and deleting the edges incident with any vertex in S. For two k-uniform hypergraphs $\mathcal{G}_1 = (V_1, E_1)$ and $\mathcal{G}_2 = (V_2, E_2)$, we say the two graphs are *isomorphic* if there is a bijection f from V_1 to V_2 , and there is a bijection g from E_1 to E_2 that maps each edge $\{v_1, v_2, \ldots, v_k\}$ to $\{f(v_1), f(v_2), \ldots, f(v_k)\}$.

^{*}Supported by National natural science foundation of China (NSFC) (Nos. Nos. 12171222, 12101285, 12071411), Natural science foundation of Guangdong province (No. 2021A1515010254), Foundation of Lingnan Normal University(ZL1923).

[†]Corresponding authors, E-mail addresses: yglong01@163.com (G. Yu), sunlin@lingnan.edu.cn (L. Sun).

In a hypergraph, two vertices are said to be *adjacent* if both of them are contained in an edge. Two edges are said to be *adjacent* if their intersection is not empty. An edge e is said to be *incident* with a vertex v if $v \in e$. The *neighbor set* of vertex v in hypergraph \mathcal{G} , denoted by $N_{\mathcal{G}}(v)$, is the set of vertices adjacent to v in \mathcal{G} . The *degree* of a vertex v in \mathcal{G} , denoted by $deg_{\mathcal{G}}(v)$ (or deg(v) for short), is the number of the edges incident with v. For a hypergraph \mathcal{G} , among all of its vertices, we denote by $\Delta(\mathcal{G})$ (or Δ for short) the *maximal degree*, and denote by $\delta(\mathcal{G})$ (or δ for short) the *minimal degree* respectively. A vertex of degree 1 is called a *pendant vertex*. A *pendant edge* is an edge with at most one vertex of degree more than one and other vertices in this edge being all pendant vertices.

In a hypergraph, a hyperpath of length q (q-hyperpath) is defined to be an alternating sequence of vertices and edges $v_1e_1v_2e_2\cdots v_qe_qv_{q+1}$ such that (1) $v_1, v_2, \ldots, v_{q+1}$ are all distinct vertices; (2) e_1, e_2, \ldots, e_q are all distinct edges; (3) $v_i, v_{i+1} \in e_i$ for $i = 1, 2, \ldots, q$; (4) $e_i \cap e_{i+1} = v_{i+1}$ for $i = 1, 2, \ldots, q - 1$; (5) $e_i \cap e_j = \emptyset$ if $|i - j| \ge 2$. If there is no discrimination, a hyperpath is sometimes written as $e_1e_2\cdots e_{q-1}e_q, e_1v_2e_2\cdots v_qe_q$ or $v_1e_1v_2e_2\cdots v_qe_q$. A hypercycle of length q (q-hypercycle) $v_1e_1v_2e_2\cdots v_{q-1}e_{q-1}v_qe_qv_1$ is obtained from a hyperpath $v_1e_1v_2e_2\cdots v_{q-1}e_{q-1}v_q$ by adding a new edge e_q between v_1 and v_q where $e_q \cap e_1 = \{v_1\}, e_q \cap e_{q-1} =$ $\{v_q\}, e_q \cap e_j = \emptyset$ if $j \ne 1, q - 1$ and $|q - j| \ge 2$. The length of a hyperpath P (or a hypercycle C), denoted by L(P) (or L(C)), is the number of the edges in P (or C). A hypergraph \mathcal{G} is connected if there exists a hyperpath from v to u for all $v, u \in V$, and \mathcal{G} is called acyclic if it contains no hypercycle.

Recall that a tree is an ordinary graph which is 2-uniform, connected and acyclic. A supertree is similarly defined to be a hypergraph which is both connected and acyclic. Clearly, in a supertree, its each pair of the edges have at most one common vertex. Therefore, the edge number of a k-uniform supertree of order n is $m = \frac{n-1}{k-1}.$

Let G = (V, E) be an ordinary graph (2-uniform). For every $k \ge 3$, the kth power of G, denoted by $G^k = (V^k, E^k)$, is defined as the k-uniform hypergraph with the edge set $E^k = \{e \cup \{v_{e_1}, v_{e_2}, \ldots, v_{e_{k-2}}\} : e \in E\}$ and the vertex set $V^k = V \cup (\cup_{e \in E} \{v_{e_1}, v_{e_2}, \ldots, v_{e_{k-2}}\})$, where $V \cap (\cup_{e \in E} \{v_{e_1}, v_{e_2}, \ldots, v_{e_{k-2}}\}) = \emptyset$, $\{v_{e_1}, v_{e_2}, \ldots, v_{e_{k-2}}\} \cap \{v_{f_1}, v_{f_2}, \ldots, v_{f_{k-2}}\} = \emptyset$ for $e \ne f$, $e, f \in E$. The kth power of an ordinary tree is called a hypertree. Obviously, a hypertree is a supertree.

Denote by $\mathcal{P}(n,k)$ the k-uniform hyperpath of order n. A k-uniform superstar of order n, denoted by $\mathcal{S}^*(n,k)$ (see Fig. 1.1), is a supertree in which all edges intersect at just one common vertex. A k-uniform double hyperstar of order n, denote by $\mathcal{S}(n,k;l_1,l_2)$ where $l_1, l_2 \geq 1$ (see Fig. 1.1), is a supertree obtained by attaching l_1 pendant edges at vertex u_1 of an edge e, and attaching l_2 pendant edges at the other vertex u_2 of edge e, where $u_1 \neq u_2$.



Fig. 1.1. $\mathcal{S}^*(n,k)$ and $\mathcal{S}(n;2,2)$

Let $E_{ij} = \{e \in E : i, j \in e\}$. The adjacency matrix $\mathcal{A}_{\mathcal{G}} = [(\mathcal{A}_{\mathcal{G}})_{ij}]$ of a hypergraph \mathcal{G} is defined as

$$(\mathcal{A}_{\mathcal{G}})_{ij} = \sum_{e \in E_{ij}} \frac{1}{|e| - 1}.$$

It is easy to find that $\mathcal{A}_{\mathcal{G}}$ is symmetric if there is no requirement for direction on hypergraph \mathcal{G} , and find that

 $\mathcal{A}_{\mathcal{G}}$ is very convenient to be used to investigate the spectrum of a hypergraph even without the requirement for edge uniformity. The *spectral radius* $\rho(\mathcal{G})$ of a hypergraph \mathcal{G} is defined to be the spectral radius $\rho(\mathcal{A}_{\mathcal{G}})$, which is the maximum modulus among all eigenvalues of $\mathcal{A}_{\mathcal{G}}$. In spectral theory of hypergraphs, the spectral radius is an index that attracts much attention due to its fine properties [4, 7, 8, 17, 20, 22, 25, 33, 35, 37, 41].

We assume that the hypergraphs throughout this paper are simple, i.e. $e_i \neq e_j$ if $i \neq j$, and assume the hypergraphs throughout this paper are undirected. In this paper, among all k-uniform $(k \ge 3)$ supertrees with fixed number of vertices, the supertrees with the maximum, the second mainum and the minimum spectral radius are completely determined respectively, getting the following result:

Theorem 1.1 Let \mathcal{G} be a k-uniform $(k \ge 3)$ supertree of order n. Then $\rho(\mathcal{G}) \le \rho(\mathcal{S}^*(n,k))$ with equality if and only if $\mathcal{G} \cong \mathcal{S}^*(n,k)$.

Theorem 1.2 Let \mathcal{G} be a k-uniform $(k \geq 3)$ supertree of order n and with $m(\mathcal{G}) \geq 3$ satisfying that $\mathcal{G} \ncong \mathcal{S}^*(n,k)$. Then $\rho(\mathcal{G}) \leq \rho(\mathcal{S}(n,k;\frac{n-1}{k-1}-2,1))$ with equality if and only if $\mathcal{G} \cong \mathcal{S}(n,k;\frac{n-1}{k-1}-2,1)$.

Theorem 1.3 Let \mathcal{G} be a k-uniform $(k \ge 3)$ supertree of order n. Then $\rho(\mathcal{P}(n,k)) \le \rho(\mathcal{G})$ with equality if and only if $\mathcal{G} \cong \mathcal{P}(n,k)$.

Corollary 1.4 Suppose T^k $(k \ge 3)$ of order n is the kth power of ordinary tree T. Then (1) $\rho(T^k) \le S^*(n,k)$ with equality if and only if $T^k \cong S^*(n,k)$. (2) $\rho(\mathcal{P}(n,k)) \le \rho(T^k)$ with equality if and only if $T^k \cong \mathcal{P}(n,k)$.

The layout of this paper is as follows: section 2 introduces some basic knowledge and working lemmas; section 3 represents our results.

2 Preliminary

For the requirements in the narrations afterward, we need some prepares. For a hypergraph \mathcal{G} with vertex set $\{v_1, v_2, \ldots, v_n\}$, a vector on \mathcal{G} is a vector $X = (x_{v_1}, x_{v_2}, \ldots, x_{v_n})^T \in \mathbb{R}^n$ that entry x_{v_i} is mapped to vertex v_i for $i \leq i \leq n$.

From [26], by the famous Perron-Frobenius theorem, for $\mathcal{A}_{\mathcal{G}}$ of a connected uniform hypergraph \mathcal{G} of order n, we know that there is unique one positive eigenvector $X = (x_{v_1}, x_{v_2}, \ldots, x_{v_n})^T \in \mathbb{R}^n_{++}$ (\mathbb{R}^n_{++} means the set of positive real vectors of dimension n) corresponding to $\rho(\mathcal{G})$, where $\sum_{i=1}^n x_{v_i}^2 = 1$ and each entry x_{v_i} is mapped to each vertex v_i for $i \leq i \leq n$. We call such an eigenvector X the principal eigenvector of \mathcal{G} .

Let A be an irreducible nonnegative $n \times n$ real matrix (with every entry being real number) with spectral radius $\rho(A)$. The following extremal representation (Rayleigh quotient) will be useful:

$$\rho(A) = \max_{X \in R^n, X \neq 0} \frac{X^T A X}{X^T X},$$

and if a vector X satisfies that $\frac{X^TAX}{X^TX} = \rho(A)$, then $AX = \rho(A)X$.

Lemma 2.1 Let A be an irreducible nonnegative square real matrix with order n and spectral radius ρ , $Y \in (R^n_+ \setminus \{0\}^n)$ be a nonnegative vector $(R^n_+$ means the set of nonnegative real vectors of dimension n, $\{0\}^n = \{(0, 0, \dots, 0)^T\}$). If $AY \ge \rho Y$, then $AY = \rho Y$.

Proof. Note the relation between the spectral radius and Rayleigh quotient for an irreducible nonnegative square real matrix. It follows that $\frac{Y^T A Y}{Y^T Y} = \rho$, and $AY = \rho Y$. Thus the result follows. This completes the proof. \Box

Lemma 2.2 [38] Let A be an irreducible nonnegative square symmetric real matrix with order n and spectral radius ρ , $Y \in (R_+^n \setminus \{0\}^n)$ be a nonnegative vector. If there exists $r \in R_+$ such that $AY \leq rY$, then $\rho \leq r$. Similarly, if there exists $r \in R_+$ such that $AY \geq rY$, then $\rho \geq r$.

3 Main results

Let X be an eigenvector of a connected k-uniform hypergraph G. For the simplicity, we let $x_e = \sum_{i=1}^{n} x_{v_i} x_{v_j}$ for an edge $e = \{v_1, v_2, \dots, v_k\}$.

Lemma 3.1 Let $e_1 = \{v_{1,1}, v_{1,2}, ..., v_{1,k}\}, e_2 = \{v_{2,1}, v_{2,2}, ..., v_{2,k}\}, ..., e_j = \{v_{j,1}, v_{j,2}, ..., v_{j,k}\}$ be some edges in a connected k-uniform hypergraph $\mathcal{G}; v_{u,1}, v_{u,2}, ..., v_{u,t}$ be vertices in \mathcal{G} that t < k. For $1 \le i \le j$, $\{v_{u,1}, v_{u,2}, ..., v_{u,t}\} \notin e_i, e'_i = (e_i \setminus \{v_{i,1}, v_{i,2}, ..., v_{i,t}\}) \cup \{v_{u,1}, v_{u,2}, ..., v_{u,t}\}$ satisfying that $e'_i \notin E(\mathcal{G})$. Let $\mathcal{G}' = \mathcal{G} - \sum e_i + \sum e'_i$. If in the principal eigenvector X of \mathcal{G} , for $1 \le i \le j$, $x_{v_{i,1}} \le x_{v_{u,1}}, x_{v_{i,2}} \le x_{v_{u,2}}, ..., and$ $x_{v_{i,t}} \le x_{v_{u,t}}$, then $\rho(\mathcal{G}') > \rho(\mathcal{G})$.

Proof. Note that $X^T(\mathcal{A}_{\mathcal{G}'} - \mathcal{A}_{\mathcal{G}})X = \frac{2}{k-1}\sum_{i=1}^{\infty} (x_{e'_i} - x_{e_i}) \geq 0$. It follows that $\rho(\mathcal{G}') \geq \rho(\mathcal{G})$. If $\rho(\mathcal{G}') = \rho(\mathcal{G})$, then $\rho(\mathcal{G}') = X^T \mathcal{A}_{\mathcal{G}'} X = X^T \mathcal{A}_{\mathcal{G}} X = \rho(\mathcal{G})$. It follows that $\mathcal{A}_{\mathcal{G}'} X = \rho(\mathcal{G}) X = \rho(\mathcal{G}) X = \mathcal{A}_{\mathcal{G}} X$. Without loss of generality, suppose $v_{u,1} \notin e_1$. Then $(\mathcal{A}_{\mathcal{G}'} X)_{v_{u,1}} - (\mathcal{A}_{\mathcal{G}} X)_{v_{u,1}} \geq \frac{x_{v_{1,k}}}{k-1} > 0$, which contradicts $\mathcal{A}_{\mathcal{G}'} X = \mathcal{A}_{\mathcal{G}} X$. Consequently, it follows that $\rho(\mathcal{G}') > \rho(\mathcal{G})$. This completes the proof. \Box

Proof of Theorem 1.1. Suppose \mathcal{T} is a k-uniform supertree of order n satisfying that $\rho(\mathcal{T}) = \max\{\rho(\mathcal{G}) : \mathcal{G}\}$ is a k-uniform supertree of order $n\}$. Let X be the principal eigenvector of \mathcal{T} and $x_u = \max\{x_v : v \in V(\mathcal{T})\}$. Suppose that $\mathcal{T} \ncong \mathcal{S}^*(n, k)$. Then in \mathcal{T} , there exist edges not incident with vertex u. Suppose e' is not incident with vertex u. Note that \mathcal{T} is connected. Then there is a hyperpath $P = ue_1v_1e_2v_2\cdots v_te'$ from u to e'. Let $e'_2 = (e_2 \setminus \{v_1\}) \cup \{u\}$, and $\mathcal{T}' = \mathcal{T} - e_2 + e'_2$. Then by Lemma 3.1, it follows that $\rho(\mathcal{T}') > \rho(\mathcal{T})$, which contradicts the maximality of $\rho(\mathcal{T})$. Hence, it follows that $\mathcal{T} \cong \mathcal{S}^*(n, k)$. This completes the proof. \Box **Proof of Theorem 1.2.** Let $\Lambda = \{\mathcal{G} : \mathcal{G}$ be a k-uniform supertree of order n and $\mathcal{G} \ncong \mathcal{S}^*(n, k)\}$. Suppose $\mathcal{T} \in \Lambda$ satisfies that $\rho(\mathcal{T}) = \max\{\rho(\mathcal{G}) : \mathcal{G} \in \Lambda\}$. Let X be the principal eigenvector of \mathcal{T} and $x_u = \max\{x_v : v \in V(\mathcal{T})\}$. Note that $\mathcal{T} \ncong \mathcal{S}^*(n, k)$, and \mathcal{T} is connected. Then in \mathcal{T} , there exist a hyperpath $P = ue_1v_1e_2$.

Assertion Except edges e_1 , e_2 , any one of other edges is incident with vertex u. Otherwise, suppose one edge e_t is not incident with vertex u. Note that \mathcal{T} is connected. Then there is a hyperpath $\mathcal{P} = ue_{a_1}v_{a_1}e_{a_2}v_{a_2}\cdots v_{a_t}e_t$ from u to e_t .

Claim 1 If $e_1 \in E(\mathcal{P})$, then $e_{a_1} = e_1$. Otherwise, suppose $e_{a_i} = e_1$ where $1 < i \le t$. Then $ue_{a_1}v_{a_1}e_{a_2}v_{a_2}\cdots v_{a_i}e_{a_i}u$ contains cycle, which contradicts that \mathcal{T} is a supertree.

In the same way, we get the following Claim 2.

Claim 2 If $e_2 \in E(\mathcal{P})$, then $e_{a_1} = e_1, e_{a_2} = e_2$.

Therefore, there 3 cases to consider, which are: (1) $e_{a_1} = e_1$, $e_2 \notin E(\mathcal{P})$; (2) $e_{a_1} = e_1$, $e_{a_2} = e_2$; (3) $e_1 \notin E(\mathcal{P}), e_2 \notin E(\mathcal{P})$. For the case that $e_{a_1} = e_1, e_2 \notin E(\mathcal{P})$, let $e'_{a_2} = (e_{a_2} \setminus \{v_{a_1}\}) \cup \{u\}$, and $\mathcal{T}' = \mathcal{T} - e_{a_2} + e'_{a_2}$, where $\mathcal{T}' \in \Lambda$. Then by Lemma 3.1, it follows that $\rho(\mathcal{T}') > \rho(\mathcal{T})$, which contradicts the maximality of $\rho(\mathcal{T})$. In the same way, for the cases (2) and (3), we can get a supertree \mathcal{T}' where $\mathcal{T}' \in \Lambda$, such that $\rho(\mathcal{T}') > \rho(\mathcal{T})$ which contradicts the maximality of $\rho(\mathcal{T})$. Thus, our assertion holds.

From the above assertion, it follows that $\mathcal{T} \cong \mathcal{S}(n,k;\frac{n-1}{k-1}-2,1)$. This completes the proof. \Box

Lemma 3.2 [23] Let A be an irreducible nonnegative square matrix with order n and spectral radius ρ . Let s_i^A be the ith row sum, $s_A = \min\{s_i^A : 1 \le i \le n\}$, and $S_A = \max\{s_i^A : 1 \le i \le n\}$. Then $s_A \le \rho \le S_A$ with either one equality if and only if A is regular (all of the row sums of A are equal).

From Lemma 3.2, combining with hypergraph, we can get the following corollary naturally.

Corollary 3.3 For a connected hypergraph \mathcal{G} , we have $\delta \leq \rho(\mathcal{G}) \leq \Delta$ with either one equality if and only if \mathcal{G} is regular, where δ is the minimum degree, Δ is the maximum degree.

Using Lemma 3.2, we can get an improvement for Lemma 2.2.

Lemma 3.4 Let A be an irreducible nonnegative square symmetric real matrix with order n and spectral radius ρ , $y \in R_{++}^n$ be a positive vector. If there exists $r \in R_+$ such that $Ay \leq ry$, then $\rho \leq r$ with equality if and only if Ay = ry. Similarly, if there exists $r \in R_+$ such that $Ay \geq ry$, then $\rho \geq r$ with equality if and only if Ay = ry.

Proof. Using Lemma 2.2 gets that $\rho \leq r$ if $Ay \leq ry$; $\rho \geq r$ if $Ay \geq ry$. Next we prove the conclusion for $\rho = r$. We first prove the conclusion that $\rho = r$ if and only if Ay = ry under the condition $Ay \leq ry$. Suppose

$$y = (y_1, y_2, \dots, y_n)^T.$$
 Let $B = \begin{pmatrix} \frac{y_1}{y_1} & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{y_2} & 0 & \vdots & 0 \\ \vdots & & & & \\ 0 & 0 & \vdots & 0 & \frac{1}{y_2} \end{pmatrix} A \begin{pmatrix} y_1 & 0 & 0 & \dots & 0 \\ 0 & y_2 & 0 & \vdots & 0 \\ \vdots & & & & \\ 0 & 0 & \vdots & 0 & y_n \end{pmatrix}.$ Denote by $\rho(B)$ the

 $\begin{pmatrix} 0 & 0 & : & 0 & \frac{1}{y_n} \end{pmatrix}$ $\begin{pmatrix} 0 & 0 & : & 0 & y_n \end{pmatrix}$ spectral radius of *B*. Note that the eigenvalues of *B* are the same to the eigenvalues of *A*; $\rho = r$ means $\rho(B) = r$. Note that $Ay \leq ry$ means $S_B \leq r$; $\rho(B) = r$ means all of the row sums of *B* equals *r* by Lemma 3.2, which implies that Ay = ry. As a result, it follows that under the condition that $Ay \leq ry$, if $\rho = r$, then Ay = ry. Conversely, if Ay = ry, then all of the row sums of *B* equals *r*, and then $\rho(B) = r = \rho$.

In the same way, we get that $\rho = r$ if and only if Ay = ry under the condition that $Ay \ge ry$. This completes the proof. \Box

Lemma 3.5 (1) Suppose c > 0, d > 0, a - c > 0, b - d > 0. If $\frac{a}{b} \ge \frac{c}{d}$, then $\frac{a-c}{b-d} \ge \frac{a}{b}$ with equality if and only if $\frac{a}{b} = \frac{c}{d}$. Moreover, if $\frac{a}{b} > \frac{c}{d}$, then $\frac{a-c}{b-d} > \frac{a}{b}$.

(2) Suppose c > 0, d > 0, a - c > 0, b - d > 0. If $\frac{a}{b} \ge \frac{c}{d}$, then $\frac{a+c}{b+d} \le \frac{a}{b}$ with equality if and only if $\frac{a}{b} = \frac{c}{d}$. Moreover, if $\frac{a}{b} > \frac{c}{d}$, then $\frac{a+c}{b+d} < \frac{a}{b}$.

(3) Suppose $\frac{a}{b} \ge 1$, b > c > 0. Then $\frac{a-c}{b-c} \ge \frac{a}{b}$.

Proof. (1) From $\frac{a}{b} \geq \frac{c}{d}$, it follows that $ab - bc \geq ab - ad$, which induces $\frac{a-c}{b-d} \geq \frac{a}{b}$. In the same way, we get that $\frac{a-c}{b-d} > \frac{a}{b}$ if $\frac{a}{b} > \frac{c}{d}$. Then (1) follows.

(2) is proved as (1). (3) is a corollary following from (1). This completes the proof. \Box



Fig. 3.1. e_0 , e_1 , e_2 in \mathcal{G}

Lemma 3.6 Let \mathcal{G} be a hypergraph with spectral radius ρ , e_0 , e_1 , e_2 be three edges in \mathcal{G} with $e_0 = \{v_1, v_2, \ldots, v_{k-1}, v_k\}$, satisfying that $\deg_{\mathcal{G}}(v_2) = \deg_{\mathcal{G}}(v_3) = \cdots \deg_{\mathcal{G}}(v_{k-1}) = 1$ $(k \ge 3)$, $e_1 \cap e_0 = \{v_1\}$, $e_2 \cap e_0 = \{v_k\}$ (see Fig. 3.1.). Let X be the principal eigenvector of hypergraph \mathcal{G} . Then $x_{v_2} = x_{v_3} = \cdots = x_{v_{k-1}} = \frac{x_{v_1} + x_{v_k}}{(k-1)\rho - (k-3)} < \min\{x_{v_1}, x_{v_k}\}$.

Proof. For $2 \le i \le k-1$, we prove $x_{v_i} < \min\{x_{v_1}, x_{v_k}\}$ by contradiction. Suppose that $\min\{x_{v_1}, x_{v_k}\} = x_{v_1}$, and $x_{v_z} \ge \min\{x_{v_1}, x_{v_k}\}$ for some $2 \le z \le k-1$. Let $e'_1 = (e_1 \setminus \{v_1\}) \cup \{v_z\}$ and $\mathcal{G}_1 = \mathcal{G} - e_1 + e'_1$. Using Lemma 3.1 gets $\rho(\mathcal{G}_1) > \rho(\mathcal{G})$. But it contradicts $\rho(\mathcal{G}_1) = \rho(\mathcal{G})$ because $\mathcal{G}_1 \cong \mathcal{G}$. As a result, for $2 \le i \le k-1$, it follows that $x_{v_i} < \min\{x_{v_1}, x_{v_k}\}$.

Note that $\rho x_{v_2} = \frac{1}{k-1}(x_{v_1} + x_{v_3} + \sum_{i=4}^k x_{v_i}), \ \rho x_{v_3} = \frac{1}{k-1}(x_{v_1} + x_{v_2} + \sum_{i=4}^k x_{v_i}).$ It follows that $(\rho + \frac{1}{k-1})(x_{v_2} - x_{v_3}) = 0$. Note that $\rho > 1$ by Corollary 3.3. Then we get $x_{v_2} = x_{v_3}$. Proceeding like this, we get that $x_{v_2} = x_{v_3} = \cdots = x_{v_{k-1}}$. Thus from $\rho x_{v_2} = \frac{1}{k-1}((k-3)x_{v_3} + x_{v_1} + x_{v_k})$, it follows that $x_{v_2} = \frac{x_{v_1} + x_{v_k}}{(k-1)\rho - (k-3)}$. Thus the result follows. This completes the proof. \Box

Similar to Lemma 3.6, we get the following Lemma 3.7.

Lemma 3.7 Let \mathcal{G} be a hypergraph with spectral radius ρ , $e = \{u, v_1, v_2, \dots, v_{k-1}\}$ be a pendant edge in \mathcal{G} ($k \ge 2$), where $deg_{\mathcal{G}}(u) \ge 2$. Then in the principal eigenvector X of \mathcal{G} , $x_{v_1} = x_{v_2} = \cdots = x_{v_{k-1}} = \frac{x_u}{(k-1)\rho-(k-2)} < x_u$.



Fig. 3.2. $\mathcal{G}_0, \mathcal{G}_1, \mathcal{G}_2$

Lemma 3.8 Suppose \mathcal{G} is a connected hypergraph with spectral radius ρ and principal eigenvector X. e_1 , e_2 , e_3 , e_4 are edges in \mathcal{G} , where $|e_1|, |e_2|, |e_3|, |e_4| \ge 3$, $e_1 \cap e_2 = \{v_1\}$, $e_1 \cap e_4 = \{v_2\}$, $e_2 \cap e_3 = \{v_3\}$, $deg_{\mathcal{G}}(v_1) = deg_{\mathcal{G}}(v_2) = deg_{\mathcal{G}}(v_3) = 2$, $deg_{\mathcal{G}}(v) = 1$ for $v \in (e_1 \cup e_2) \setminus \{v_1, v_2, v_3\}$.

(1) Let $e' \subset (e_1 \cup e_2)$ satisfy that $\{v_2, v_3\} \subseteq e'$, $e' \notin E(\mathcal{G})$. Let $\mathcal{G}_0 = \mathcal{G} - e_1 - e_2 + e'$ and t = |e'| (see Fig. 3.2).

 $(1.1) If t \ge \max\{|e_1|, |e_2|\}, x_{v_1} \ge x_{v_2}, x_{v_1} \ge x_{v_3}, then \ \rho(\mathcal{G}_0) \le \rho(\mathcal{G}) \text{ with equality if and only if } |e'| = |e_1| = |e_2| and x_{v_1} = x_{v_2} = x_{v_3}. Moreover, if t > \max\{|e_1|, |e_2|\}, x_{v_1} \ge x_{v_2}, x_{v_1} \ge x_{v_3}, then \ \rho(\mathcal{G}_0) < \rho(\mathcal{G}).$

(1.2) If $t \leq \max\{|e_1|, |e_2|\}$, $x_{v_1} \leq x_{v_2}$, $x_{v_1} \leq x_{v_3}$, then $\rho(\mathcal{G}_0) \geq \rho(\mathcal{G})$ with equality if and only if $|e'| = |e_1| = |e_2|$ and $x_{v_1} = x_{v_2} = x_{v_3}$. Moreover, if $t < \max\{|e_1|, |e_2|\}$, $x_{v_1} \leq x_{v_2}$, $x_{v_1} \leq x_{v_3}$, then $\rho(\mathcal{G}_0) > \rho(\mathcal{G})$.

(2) Let $e_1^{'} = (e_1 \setminus \{v_1\}) \cup \{u\}, \ e_2^{'} = (e_2 \setminus \{v_1\}) \cup \{u\}, \ e^{'} = \{v_1, \ u_1, \ u_2, \ \dots, \ u_{t-2}, \ u\}$ where $u \notin V(\mathcal{G}), u_i \notin V(\mathcal{G})$ for $1 \le i \le t-2, \ \mathcal{G}_1 = \mathcal{G} - e_1 + e_1^{'} + e^{'}, \ \mathcal{G}_2 = \mathcal{G} - e_2 + e_2^{'} + e^{'}$ (see Fig. 3.2).

(2.1) If $t \leq \min\{|e_1|, |e_2|\}$, $x_{v_1} \geq x_{v_2}$, $x_{v_1} \geq x_{v_3}$, then $\rho(\mathcal{G}_1) \geq \rho(\mathcal{G})$, $\rho(\mathcal{G}_2) \geq \rho(\mathcal{G})$ with either equality holding if and only if $|e^{'}| = |e_1| = |e_2|$, $x_{v_1} = x_{v_2} = x_{v_3} = x_u$ and $x_z = x_\omega$ for $z, \omega \in (e_1 \cup e_2 \cup e^{'}) \setminus \{v_1, v_2, u\}$. Moreover, if $t < \min\{|e_1|, |e_2|\}$, $x_{v_1} \geq x_{v_2}$, $x_{v_1} \geq x_{v_3}$, then $\rho(\mathcal{G}_1) > \rho(\mathcal{G})$, $\rho(\mathcal{G}_2) > \rho(\mathcal{G})$.

 $(2.2) If t \ge \max\{e_1, e_2\}, x_{v_1} \le x_{v_2}, x_{v_1} \le x_{v_3}, then \ \rho(\mathcal{G}_1) \le \rho(\mathcal{G}), \ \rho(\mathcal{G}_2) \le \rho(\mathcal{G}) \text{ with either equality holding if and only if } |e^{'}| = |e_1| = |e_2|, x_{v_1} = x_{v_2} = x_{v_3} = x_u \text{ and } x_z = x_\omega \text{ for } z, \omega \in (e_1 \cup e_2 \cup e^{'}) \setminus \{v_1, v_2, u\}. Moreover, if t > \max\{e_1, e_2\}, x_{v_1} \le x_{v_2}, x_{v_1} \le x_{v_3}, then \ \rho(\mathcal{G}_1) < \rho(\mathcal{G}), \ \rho(\mathcal{G}_2) < \rho(\mathcal{G}).$

Proof. (1.1) Suppose $e_1 = \{v_1, v_{\alpha(1,1)}, v_{\alpha(1,2)}, \dots, v_{\alpha(1,j_1-2)}, v_2\}, e_2 = \{v_1, v_{\alpha(2,1)}, v_{\alpha(2,2)}, \dots, v_{\alpha(2,j_2-2)}, \dots, v_$

 v_3 }. By Lemma 3.6, we have $x_{v_{\alpha(1,w)}} = x_{v_{\alpha(1,z)}} < \min\{x_{v_1}, x_{v_2}\}$ for $1 \le w < z \le j_1 - 2$, $x_{v_{\alpha(2,w)}} = x_{v_{\alpha(2,z)}} < \min\{x_{v_1}, x_{v_3}\}$ for $1 \le w < z \le j_2 - 2$. Let Y be a vector on \mathcal{G}_0 satisfying that

$$\begin{cases} y_v = \min\{x_z : z \in (e_1 \cup e_2) \setminus \{v_1, v_2, v_3\}\}, & v \in e^{'} \setminus \{v_2, v_3\}\\ y_v = x_v, & others. \end{cases}$$

Note that $|e'| \ge \max\{|e_1|, |e_2|\}, x(v_1) \ge x(v_2), x(v_1) \ge x(v_3)$. Without loss of generality, suppose $\min\{x_v : v \in (e_1 \cup e_2)\} = x_{v_{\alpha(2,1)}}$. For $v \in (e' \setminus \{v_2, v_3\})$, noting that $deg_{\mathcal{G}_0}(v) = 1$ and $x_{v_{\alpha(2,1)}} < \min\{x_{v_1}, x_{v_3}\}$, we have

$$(\mathcal{A}_{\mathcal{G}_{0}}Y)_{v} = \frac{(t-3)y_{v} + y_{v_{2}} + y_{v_{3}}}{t-1} = \frac{(t-3)x_{v_{\alpha(2,1)}} + x_{v_{2}} + x_{v_{3}}}{t-1}$$
$$= \frac{(j_{2}-3)x_{v_{\alpha(2,1)}} + x_{v_{2}} + x_{v_{3}} + (t-j_{2})x_{v_{\alpha(2,1)}}}{j_{2}-1+t-j_{2}}$$
$$\leq \frac{(j_{2}-3)x_{v_{\alpha(2,1)}} + x_{v_{1}} + x_{v_{3}} + (t-j_{2})x_{v_{\alpha(2,1)}}}{j_{2}-1+t-j_{2}}$$
$$\leq \frac{(j_{2}-3)x_{v_{\alpha(2,1)}} + x_{v_{1}} + x_{v_{3}}}{j_{2}-1} \qquad (by \ Lemma \ 3.5)$$
$$= \rho x_{v_{\alpha(2,1)}} = \rho y_{v}.$$

In the same way, we get

$$(\mathcal{A}_{\mathcal{G}_{0}}Y)_{v_{2}} = \frac{(t-2)x_{v_{\alpha(2,1)}} + y_{v_{3}}}{t-1} = \frac{(t-2)x_{v_{\alpha(2,1)}} + x_{v_{3}}}{t-1} \le \rho x_{v_{2}} = \rho y_{v_{2}};$$
$$(\mathcal{A}_{\mathcal{G}_{0}}Y)_{v_{3}} = \frac{(t-2)x_{v_{\alpha(2,1)}} + y_{v_{2}}}{t-1} = \frac{(t-2)x_{v_{\alpha(2,1)}} + x_{v_{2}}}{t-1} \le \rho x_{v_{3}} = \rho y_{v_{3}};$$

for $v \in (V(\mathcal{G}_0) \setminus e')$, $(\mathcal{A}_{\mathcal{G}_0}Y)_v = (\mathcal{A}_{\mathcal{G}}X)_v = \rho y_v$. By lemma 2.2, it follows that $\rho(\mathcal{G}_0) \leq \rho(\mathcal{G})$. Note that Y is positive. Combining Lemma 3.4, we find that if $\rho(\mathcal{G}_0) = \rho(\mathcal{G})$, then $\mathcal{A}_{\mathcal{G}_0}Y = \rho Y$. Thus it follows that $|e'| = |e_1| = |e_2|$ and $x_{v_1} = x_{v_2} = x_{v_3}$. Conversely, if $|e'| = |e_1| = |e_2|$ and $x_{v_1} = x_{v_2} = x_{v_3}$, then it can be checked as above that for $v \in (e' \setminus \{v_2, v_3\})$, $(\mathcal{A}_{\mathcal{G}_0}Y)_v = \rho y_v$; $(\mathcal{A}_{\mathcal{G}_0}Y)_{v_2} = \rho y_{v_2}$; $(\mathcal{A}_{\mathcal{G}_0}Y)_{v_3} = \rho y_{v_3}$; for $v \in (V(\mathcal{G}_0) \setminus e')$, $(\mathcal{A}_{\mathcal{G}_0}Y)_v = (\mathcal{A}_{\mathcal{G}}X)_v = \rho y_v$. Thus it follows that $\rho(\mathcal{G}_0) = \rho(\mathcal{G})$.

If $t > \max\{|e_1|, |e_2|\}, x_{v_1} \ge x_{v_2}, x_{v_1} \ge x_{v_3}$, combining Lemma 3.5, as the above proof, we get $\rho(\mathcal{G}_0) < \rho(\mathcal{G})$. From the above narrations, then (1.1) follows as desired.

(1.2) Let Y be a vector on \mathcal{G}_0 satisfying that

$$\begin{cases} y(v) = \max\{x_z : z \in (e_1 \cup e_2) \setminus \{v_1, v_2, v_3\}\}, & v \in e' \setminus \{v_2, v_3\}\\ y(v) = x_v, & others. \end{cases}$$

Then (1.2) is proved as (1.1).

(2.1) For both \mathcal{G}_1 and \mathcal{G}_2 , let Y be a vector satisfying that

$$\begin{cases} y(u) = x_{v_1} \\ y(v) = \max\{x_z : z \in (e_1 \cup e_2) \setminus \{v_1, v_2, v_3\}\}, & v \in e^{'} \setminus \{v_1, u\} \\ y(v) = x_v, & others. \end{cases}$$

Then (2.1) is proved as (1.1).

(2.2) For both \mathcal{G}_1 and \mathcal{G}_2 , let Y be a vector satisfying that

$$\begin{cases} y(u) = x_{v_1} \\ y(v) = \min\{x_z : z \in (e_1 \cup e_2) \setminus \{v_1, v_2, v_3\}\}, & v \in e^{'} \setminus \{v_1, u\} \\ y(v) = x_v, & others. \end{cases}$$

Then (2.2) is proved as (1.1).

Thus the result follows. This completes the proof. \Box

Lemma 3.9 Let e be a new edge not containing in connected hypergraph \mathcal{G} . Let $\mathcal{G}' = \mathcal{G} + e$. If \mathcal{G}' is also connected, then $\rho(\mathcal{G}') > \rho(\mathcal{G})$.

Proof. Let X be the *principal eigenvector* of \mathcal{G} , Y be a vector on \mathcal{G}' satisfying that

$$\begin{cases} y_v = x_v, & v \in V(\mathcal{G}) \\ y_v = 0, & others. \end{cases}$$

Then $Y^T \mathcal{A}_{\mathcal{G}'} Y - X^T \mathcal{A}_{\mathcal{G}} X \ge 0$, $Y^T Y = X^T X$. It follows that $\rho(\mathcal{G}') \ge \rho(\mathcal{G})$. Suppose that $\rho(\mathcal{G}') = \rho(\mathcal{G})$. Then $\rho(\mathcal{G}') = Y^T \mathcal{A}_{\mathcal{G}'} Y = X^T \mathcal{A}_{\mathcal{G}} X = \rho(\mathcal{G})$, and then Y is a principal eigenvector of \mathcal{G}' . If there exists $y_v = 0$, then we get a contradiction because the principal eigenvector of \mathcal{G}' is positive by Perron-Frobenius theorem.

Suppose Y is positive next. Note that Y = X if Y is positive. It follows that $V(\mathcal{G}') = V(\mathcal{G})$ now. Denote by $e = \{v_1, v_2, \ldots, v_k\}$. Then $e \subseteq V(\mathcal{G})$, and

$$\rho(\mathcal{G}')y_{v_1} = (\mathcal{A}_{\mathcal{G}'}Y)_{v_1} = (\mathcal{A}_{\mathcal{G}'}X)_{v_1} + \frac{1}{k-1}\sum_{i=2}^k x_{v_i} = \rho(\mathcal{G})x_{v_1} + \frac{1}{k-1}\sum_{i=2}^k x_{v_i} > \rho(\mathcal{G})x_{v_1},$$

which contradicts $\rho(\mathcal{G}') = \rho(\mathcal{G})$. As a result, we get that $\rho(\mathcal{G}') > \rho(\mathcal{G})$. This completes the proof. \Box

Denote by $\mathcal{G}(\mathcal{D}v; p, q; v_{p+q}\mathcal{H})$ the k-uniform connected hypergraph obtained from k-uniform hypergraph \mathcal{D} and k-uniform hypergraph \mathcal{H} by adding a pendant path P_1 with length p at vertex v of \mathcal{D} , and adding a path P_2 with length q between vertex v and vertex v_{p+q} of \mathcal{H} , where \mathcal{D} and \mathcal{H} are two disjoint, $V(P_1) \cap V(\mathcal{D}) = \{v\}$, $V(P_2) \cap V(\mathcal{D}) = \{v\}, V(P_2) \cap V(\mathcal{H}) = \{v_{p+q}\}$ (see two examples in Fig. 3.3). In particular, if $H = v_{p+q}$, we denote by $\mathcal{G}(\mathcal{D}v; p, q; v_{p+q})$ for $\mathcal{G}(\mathcal{D}v; p, q; v_{p+q}\mathcal{H})$ for short.



Lemma 3.10 If p, q > 0, then $\rho(\mathcal{G}(\mathcal{D}v; p, q; v_{p+q}\mathcal{H})) > \rho(\mathcal{G}(\mathcal{D}v; 0, p+q; v_{p+q}\mathcal{H}))$.

Proof. Let X be the principal eigenvector of the uniform hypergraph $\mathcal{G}((\mathcal{D}v; 0, p + q; v_{p+q}\mathcal{H}))$. Assume that in \mathcal{D} , the edges incident with v are $\varepsilon_1, \varepsilon_2, \ldots, \varepsilon_\eta$. Let $e'_{p+1} = (e_{p+1} \setminus \{v_p\}) \cup \{v\}, \mathcal{G}_1 = \mathcal{G}((\mathcal{D}v; 0, p + q; v_{p+q}\mathcal{H})) - e_{p+1} + e'_{p+1}$. If $x_v \ge x_{v_p}$, then $\rho(\mathcal{G}_1) > \rho(\mathcal{G}(\mathcal{D}v; 0, p + q; v_{p+q}\mathcal{H}))$ by Lemma 3.1. Let $\varepsilon'_i = (\varepsilon_i \setminus \{v\}) \cup \{v_p\}$ for $1 \le i \le \eta, \mathcal{G}_2 = \mathcal{G}((\mathcal{D}v; 0, p + q; v_{p+q}\mathcal{H})) - \sum_{i=1}^{\eta} \varepsilon_i + \sum_{i=1}^{\eta} \varepsilon'_i$. If $x_v \le x_{v_p}$, then $\rho(\mathcal{G}_2) > \rho(\mathcal{G}(\mathcal{D}v; 0, p+q; v_{p+q}\mathcal{H}))$ by Lemma 3.1. Note that both G_1 and G_2 are isomorphic to $\mathcal{G}(\mathcal{D}v; p, q; v_{p+q}\mathcal{H})$. Thus we get that $\rho(\mathcal{G}(\mathcal{D}v; p, q; v_{p+q}\mathcal{H})) > \rho(\mathcal{G}(\mathcal{D}v; 0, p+q; v_{p+q}\mathcal{H}))$. This completes the proof. \Box

Fig. 3.4. $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$

Lemma 3.11 Let X be the principal eigenvector of the uniform hypergraph $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$. Denote by $\mathcal{P} = v_0 e_1 v_1 e_2 v_2 \cdots e_t v_t$ the pendant path from vertex v_0 to vertex v_t in $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$ and $e_i = \{v_{i-1}, v_{a(i,1)}, v_{a(i,2)}, \dots, v_{a(i,k-2)}, v_i\}$ for $1 \le i \le t$ (see Fig. 3.4). Suppose $x_{v_p} = \max\{x_{v_i} : 0 \le i \le t\}$. Then

(1) $p \le t - 1$.

Let t = p + q. Moreover, we have

(2) if p > 0, then $x_{v_i} \le x_{v_{i+1}}$ for $0 \le i \le p-1$, $x_{v_i} \ge x_{v_{i+1}}$ for $p \le i \le t-1$; if p = 0, then $x_{v_i} \ge x_{v_{i+1}}$ for $0 \le i \le t-1$.

(3) if p > 0 and there exists $\omega \leq p$ and $\eta \leq q$ such that $x_{v_{n-\omega}} \geq x_{v_{n+n}}$, then

 $(3.1) if \omega \leq \eta, then x_{v_{p-\omega+i}} \geq x_{v_{p+\eta-i}} \text{ for } 0 \leq i \leq \omega, x_{v_{a(p-\omega+i,1)}} \geq x_{v_{a(p+\eta-i+1,1)}} \text{ for } 1 \leq i \leq \omega.$

(3.2) if
$$\omega \ge \eta$$
, then $x_{v_{p-\omega+i}} \ge x_{v_{p+\eta-i}}$ for $0 \le i \le \eta - 1$, $x_{v_j} = x_{v_p}$ for $p - \omega + \eta \le j \le p$, and

 $x_{v_{a(p-\omega+i,1)}} \ge x_{v_{a(p+\eta-i+1,1)}}$ for $1 \le i \le \eta$.

(4) if p > 0 and there exists $\omega \leq p$ and $\eta \leq q$ such that $x_{v_{p-\omega}} \leq x_{v_{p+\eta}}$, then

(4.1) if $\omega \leq \eta$, then $x_{v_{p-\omega+i}} \leq x_{v_{p+\eta-i}}$ for $0 \leq i \leq \omega - 1$, $x_{v_j} = x_{v_p}$ for $p+1 \leq j \leq p+\eta-\omega$, and $x_{v_{a(p-\omega+i,1)}} \leq x_{v_{a(p+\eta-i+1,1)}}$ for $1 \leq i \leq \omega$.

(4.2) if $\omega \ge \eta$, then $x_{v_{p-\omega+i}} \le x_{v_{p+\eta-i}}$ for $0 \le i \le \eta$, $x_{v_{a(p-\omega+i,1)}} \le x_{v_{a(p+\eta-i+1,1)}}$ for $1 \le i \le \eta$.

(5) if p > 0 and there exists $\omega \leq p, \eta \leq q$ such that $x_{v_{p-\omega}} = x_{v_{p+\eta}}$, then

(5.1) if $\omega \leq \eta$, then $x_{v_{p-\omega+i}} = x_{v_{p+\eta-i}}$ for $0 \leq i \leq \omega - 1$, $x_{v_j} = x_{v_p}$ for $p+1 \leq j \leq p+\eta-\omega$, and $x_{v_{a(p-\omega+i,1)}} = x_{v_{a(p+\eta-i+1,1)}}$ for $1 \leq i \leq \omega$.

(5.2) if $\omega \ge \eta$, then $x_{v_{p-\omega+i}} = x_{v_{p+\eta-i}}$ for $0 \le i \le \eta - 1$, $x_{v_j} = x_{v_p}$ for $p - \omega + \eta \le j \le p$, $x_{v_{a(p-\omega+i,1)}} = x_{v_{a(p+\eta-i+1,1)}}$ for $1 \le i \le \eta$.

Proof. (1) By Lemma 3.7, it follows that $x_{v_t} < x_{v_{t-1}}$. Thus $p \le t - 1$.

(2) Using Lemma 3.6 gets that for $1 \le i \le t - 1$,

$$x_{v_{a(i,1)}} = x_{v_{a(i,2)}} = \dots = x_{v_{a(i,k-2)}} < \min\{x_{v_{i-1}}, x_{v_i}\}.$$

Note that $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$ is uniform and $p \leq t - 1$.

Case 1 p > 0.

Claim $x_{v_0} \leq x_{v_1} \leq x_{v_2} \leq \cdots \leq x_{v_p}$. If p = 1, this claim hold naturally.

For $p \geq 2$, we prove this claim by contradiction. Suppose that x_{v_z} $(0 \leq z \leq p-1)$ is the first vertex from 0 to p-1 such that $x_{v_z} > x_{v_{z+1}}$. Then there exists $z+1 \leq \zeta \leq p-1$ such that $x_{v_z} > x_{v_{z+1}} \geq \cdots \geq x_{v_\zeta} \leq x_{v_{\zeta+1}}$. Let $e' = \{v_{\zeta}, u_1, u_2, \ldots, u_{k-2}, u\}$, $e'_{\zeta} = (e_{\zeta} \setminus \{v_{\zeta}\}) \cup \{u\}$, where $u \notin V(\mathcal{G})$, $u_i \notin V(\mathcal{G})$ for $1 \leq i \leq t-2$, $\mathcal{G}_1 = \mathcal{G} - e_{\zeta} + e'_{\zeta} + e'$, $\mathcal{G}_2 = \mathcal{G}_1 - \{v_{a(t,1)}, v_{a(t,2)}, \ldots, v_{a(t,k-2)}, v_t\}$. By Lemma 3.8 and Lemma 3.9, we get that $\rho(\mathcal{G}_2) < \rho(\mathcal{G}_1) \leq \rho(\mathcal{G})$, which contradicts $\rho(\mathcal{G}) = \rho(\mathcal{G}_2)$ because $\mathcal{G}_2 \cong \mathcal{G}$. Thus our claim holds.

In the same way, it is proved that $x_{v_p} \ge x_{v_{p+1}} \ge x_{v_{p+2}} \ge \cdots \ge x_{v_{t-1}}$. Combining Lemma 3.7, we get that $x_{v_p} \ge x_{v_{p+1}} \ge x_{v_{p+2}} \ge \cdots \ge x_{v_{t-1}} > x_{v_t}$.

Case 2 p = 0. As Case 1, it is proved that $x_{v_i} \ge x_{v_{i+1}}$ for $0 \le i \le t - 1$. Thus (2) follows. (3) If $\omega \le \eta$, we let Y be a vector on $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$ satisfying that

$$\begin{cases} y_{v_{p-\omega+i}} = \max\{x_{v_{p-\omega+i}}, x_{v_{p+\eta-i}}\} & 0 \le i \le \eta; \\ y_{v_{a(p-\omega+i,j)}} = \max\{x_{v_{a(p-\omega+i,1)}}, x_{v_{a(p+\eta-i+1,1)}}\} & 1 \le i \le \omega, 1 \le j \le k-2; \\ y_{v} = x_{v} & others. \end{cases}$$

As the proof of Lemma 3.8, it is proved that $\mathcal{A}(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))Y \ge \rho(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))Y$. Using Lemma 2.1 gets that $\mathcal{A}(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))Y = \rho(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))Y$. Note that $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$ is connected. Then $\mathcal{A}(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))$ is irreducible. Consequently, it follows that the dimension of the eigenspace of the eigenvalue $\rho(\mathcal{G}(\mathcal{D}v_0; 0, t; v_t))$ is one. Then Y = lX for some l > 0. Then (3.1) follows.

(3.2), (4.1) and (4.2) are proved in the same way. (5) follows from (3) or (4).

This completes the proof. \Box

Lemma 3.12 $L_0, L_1, L_2, \ldots, L_f$ $(f \ge 1)$ are positive integers satisfying $L_1 \le L_2 \le \cdots \le L_f \le L_0 - 1$ and $\sum_{i=0}^{f} L_i = t$. For an integer $\mu > 0$, if $t - \mu \ge L_0$, then there exists some $1 \le j \le f$ such that $t - \mu > L_1 + \cdots + L_j$, but $L_0 + L_1 + \cdots + L_j > t - \mu$.

Proof. If $\sum_{i=1}^{f} L_i < t - \mu$, then $t = \sum_{i=0}^{f} L_i > t - \mu$ follows naturally. Thus the result holds.

Suppose $\sum_{i=1}^{f} L_i \ge t - \mu$. Note that $t - \mu \ge L_0$ and $L_1 \le L_2 \le \cdots \le L_f \le L_0 - 1$. Then $f \ge 2$ now, and there exists some $1 \le g \le f - 1$ such that $L_1 + \cdots + L_g + L_{g+1} \ge t - \mu$, but $L_1 + \cdots + L_g < t - \mu$. Note that $L_{g+1} \le L_0 - 1$. Then it follows that $L_0 + L_1 + \cdots + L_g > t - \mu$, but $t - \mu > L_1 + \cdots + L_g$. This completes the proof. \Box



Fig. 3.5. $\mathcal{G}_1, \mathcal{G}_2$

Lemma 3.13 \mathcal{D} is a k-uniform hypergraph where $v_0 \in V(\mathcal{D})$. Both $\mathbf{P} = v_0 e_1 v_1 e_2 v_2 \cdots e_t v_t$ and $\mathbf{P}_0 = v_0 \tilde{e}_1 u_1 \tilde{e}_2 u_2 \cdots \tilde{e}_s u_s$ are k-uniform hyperpaths where $\tilde{e}_1 = \{v_0, v_{\varphi(1,1)}, v_{\varphi(1,2)}, \dots, v_{\varphi(1,k-2)}, u_1\}, V(\mathcal{D}) \cap V(\mathbf{P}) = \{v_0\}, V(\mathcal{D}) \cap V(\mathbf{P}_0) = \{v_0\}.$ $\mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_f$ $(1 \leq f \leq k-2)$ are k-uniform hyperpaths attached respectively at vertices $v_{\varphi(1,1)}, v_{\varphi(1,2)}, \dots, v_{\varphi(1,f)}$ in \tilde{e}_1 satisfying $1 \leq L(\mathbf{P}_1) \leq L(\mathbf{P}_2) \leq \cdots \leq L(\mathbf{P}_f) \leq L(\mathbf{P}_0) - 1$, $\sum_{i=0}^{f} L(\mathbf{P}_i) = t, V(\mathcal{D}) \cap V(\mathbf{P}_i) = \emptyset$ for $1 \leq i \leq f$. K-uniform hypergraph \mathcal{G}_1 is a $\mathcal{G}(\mathcal{D}v_0; 0, t; v_t)$ consisting of \mathcal{D} and \mathbf{P}_i ; k-uniform hypergraph \mathcal{G}_2 consists of \mathcal{D} and $\mathbf{P}_0, \mathbf{P}_1, \mathbf{P}_2, \dots, \mathbf{P}_f$ (see Fig. 3.5). Then $\rho(\mathcal{G}_1) < \rho(\mathcal{G}_2)$.

Proof. For brevity, we denote by $L_i = L(\mathbf{P}_i)$ for $0 \le i \le f$. Without loss of generality, we suppose f = 3 next.

In \mathcal{G}_1 , denote by $e_i = \{v_{i-1}, v_{a(i,1)}, v_{a(i,2)}, \dots, v_{a(i,k-2)}, v_i\}$ for $1 \le i \le t-1$, $e_t = \{v_{t-1}, v_{a(t,1)}, v_{a(t,2)}, \dots, v_{a(t,k-2)}, v_{a(t,k-1)}\}$ where $v_{a(t,k-1)} = v_t$. Assume that in \mathcal{D} , the edges incident with v_0 are $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_\eta$. Let X be the principal eigenvector of \mathcal{G}_1 , and $x_{v_p} = \max\{v_i : 0 \le i \le t\}$. By Lemma 3.11, we know that $p \le t-1$. By Lemma 3.6 and Lemma 3.7, we know that $x_{v_{a(i,j)}} = x_{v_{a(i,z)}} < \min\{x_{v_{i-1}}, x_{v_i}\}$ for $2 \le j, z \le k-2$ where $1 \le i \le t-1$, $x_{v_{a(t,j)}} = x_{v_{a(t,z)}}$ for $2 \le j, z \le k-2$ where $1 \le i \le t-1$, $x_{v_{a(t,j)}} = x_{v_{a(t,z)}}$ for $2 \le j, z \le k-2$ where $1 \le i \le t-1$, $x_{v_{a(t,j)}} = x_{v_{a(t,z)}}$ for $2 \le j, z \le k-1$. By Lemma 3.11, we know that if p > 0, then $x_{v_i} \le x_{v_{i+1}}$ for $0 \le i \le p-1$; if $p \ge 0$, then $x_{v_i} \ge x_{v_{i+1}}$ for $p \le i \le t-1$.

Case 1 p > 0.

Subcase 1.1 $t-p \ge L_0$ (see Fig. 3.6). By Lemma 3.12, there exists $1 \le j \le 3$ such that $t-p > L_1 + \cdots + L_j$, but $L_0 + L_1 + \cdots + L_j > t - p$. Without loss of generality, we suppose j = 2. Now $t - L_1 - L_2 > p$, $t - L_0 - L_1 - L_2 < p$. Note that $L_3 = t - L_0 - L_1 - L_2$. For brevity and convenience, we let $\xi = t - L_1$, $\eta = t - L_1 - L_2$.

$$\underbrace{\mathcal{D} \quad v_0}_{e_1} \underbrace{v_1}_{e_{L_3}} \underbrace{v_{L_3}}_{e_{L_3}} \underbrace{v_{\eta-1}}_{e_{\eta}} \underbrace{v_{\eta}}_{e_{\eta}} \underbrace{v_{\xi-1}}_{e_{\eta}} \underbrace{v_{\xi}}_{e_{\xi+1}} \underbrace{v_{t-1}}_{e_{\xi+1}} \underbrace{v_{t-1}}_{e_{t}} \underbrace{v_{t}}_{e_{t}} \underbrace{v_{t-1}}_{e_{t}} \underbrace{v_{t}}_{e_{t}} \underbrace{v_{t-1}}_{e_{t}} \underbrace{v_{t}}_{e_{t}} \underbrace{v_{t}} \underbrace{v_{t}}_{e_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace{v_{t}} \underbrace$$

Fig. 3.6.
$$G_1 (t - p \ge L_0)$$

Subcase 1.1.1 $x_{v_{\eta}} \ge x_{v_{L_3}}$. By Lemma 3.11, we know that $x_{v_{\eta-1}} \ge x_{v_{L_3+1}}$.

Subcase 1.1.1.1 $x_{v_0} \leq x_{v_{a(\eta,1)}}, x_{v_{L_3}} \leq x_{v_{a(\eta,2)}}, x_{v_{\xi}} \leq x_{v_{a(\eta,3)}}$. Let $\varepsilon_i' = (\varepsilon_i \setminus \{v_0\}) \cup \{v_{a(\eta,1)}\}$ for $1 \leq i \leq \eta$, $e_{L_3}' = (e_{L_3} \setminus \{v_{L_3}\}) \cup \{v_{a(\eta,2)}\}, e_{\xi+1}' = (e_{\xi+1} \setminus \{v_{\xi}\}) \cup \{v_{a(\eta,3)}\}; \mathcal{S}_i = \sum_{v \in (\varepsilon_i \setminus \{v_0\})} x_v$ for $i = 1, 2, ..., \eta$, $\mathbb{S}_{L_3} = \sum_{v \in e_{L_3} \setminus \{v_{L_3}\}} x_v, \mathbb{S}_{\xi+1} = \sum_{v \in (e_{\xi+1} \setminus \{v_{\xi}\})} x_v.$

Let

$$\mathcal{G}_{1}^{'} = \mathcal{G}_{1} - \sum_{i=1}^{\eta} \varepsilon_{i} + \sum_{i=1}^{\eta} \varepsilon_{i}^{'} - e_{L_{3}} + e_{L_{3}}^{'} - e_{\xi+1} + e_{\xi+1}^{'}.$$

Then

$$X^{T}\mathcal{A}(\mathcal{G}_{1}')X - X^{T}\mathcal{A}(\mathcal{G}_{1})X = \frac{2}{k-1}\{(x_{v_{a(\eta,1)}} - x_{v_{0}})\sum_{i=1}^{\eta}\mathcal{S}_{i} + (x_{v_{a(\eta,2)}} - x_{v_{L_{3}}})\mathbb{S}_{L_{3}} + (x_{v_{a(\eta,3)}} - x_{v_{\xi}})\mathbb{S}_{\xi+1}\} \ge 0.$$

It follows that $\rho(\mathcal{G}_1') \geq \rho(\mathcal{G}_1)$. Suppose $\rho(\mathcal{G}_1') = \rho(\mathcal{G}_1)$. Then $\rho(\mathcal{G}_1') = X^T \mathcal{A}(\mathcal{G}_1') X = X^T \mathcal{A}(\mathcal{G}_1) X = \rho(\mathcal{G}_1)$. Hence X is also the principal eigenvector of \mathcal{G}_1' and $\mathcal{A}(\mathcal{G}_1') X = \mathcal{A}(\mathcal{G}_1) X$. But a contradiction comes immediately because $(\mathcal{A}(\mathcal{G}_1')X)_{v_{a(\eta,1)}} > (\mathcal{A}(\mathcal{G}_1)X)_{v_{a(\eta,1)}}$. Thus it follows that $\rho(\mathcal{G}_1') > \rho(\mathcal{G}_1)$. Note that $\mathcal{G}_1' \cong \mathcal{G}_2$. Then $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

 $\begin{aligned} \mathbf{Subcase \ 1.1.1.2} \ x_{v_0} > x_{v_{a(\eta,1)}}, \ x_{v_{L_3}} &\leq x_{v_{a(\eta,2)}}, \ x_{v_{t-L_1}} \leq x_{v_{a(\eta,3)}}. \ \text{Let} \ e_1' = (e_1 \setminus \{v_0\}) \cup \{v_{a(\eta,1)}\}, \ e_{\eta}' = (e_{\eta} \setminus \{v_{a(\eta,1)}\}) \cup \{v_0\}, \ e_{L_3}' = (e_{L_3} \setminus \{v_{L_3}\}) \cup \{v_{a(\eta,2)}\}, \ e_{\xi+1}' = (e_{\xi+1} \setminus \{v_{\xi}\}) \cup \{v_{a(\eta,3)}\}; \ \mathbb{S}_1 = \sum_{v \in (e_1 \setminus \{v_0\})} x_v, \\ \mathbb{S}_\eta = \sum_{v \in (e_\eta \setminus \{v_{a(\eta,1)}\})} x_v, \ \mathbb{S}_{L_3} = \sum_{v \in e_{L_3} \setminus \{v_{L_3}\}} x_v, \ \mathbb{S}_{\xi+1} = \sum_{v \in (e_{\xi+1} \setminus \{v_{\xi}\})} x_v. \ \text{Note} \ x_{v_\eta} \geq x_{v_{t-L_0-L_1-L_2}} \geq x_{v_0}, \\ x_{\eta-1} \geq x_{t-L_0-L_1-L_2+1} = x_{L_3+1}. \ \text{Using Lemma 3.6, we get} \ x_{v_{a(1,j)}} < x_{a(L_3+1,j)} < x_{v_{a(\eta,j)}} < \min\{x_{v_{\eta-1}}, x_{v_\eta}\} \\ \text{for} \ 1 \leq j \leq k-2. \ \text{As a result, it follows that} \ \mathbb{S}_1 \leq \mathbb{S}_{\eta}. \end{aligned}$

Let

$$\mathcal{G}_{1}^{'} = \mathcal{G}_{1} - e_{1} + e_{1}^{'} - e_{\eta} + e_{\eta}^{'} - e_{L_{3}} + e_{L_{3}}^{'} - e_{\xi+1} + e_{\xi+1}^{'}.$$

Then

$$X^{T}\mathcal{A}(\mathcal{G}_{1}^{'})X - X^{T}\mathcal{A}(G_{1})X = \frac{2}{k-1} \{ (x_{v_{0}} - x_{v_{a(\eta,1)}}) \mathbb{S}_{\eta} - (x_{v_{0}} - x_{v_{a(\eta,1)}}) \mathbb{S}_{1} + (x_{v_{a(\eta,2)}} - x_{v_{L_{3}}}) \mathbb{S}_{L_{3}} + (x_{v_{a(\eta,3)}} - x_{v_{\xi}}) \mathbb{S}_{\xi+1} \} \ge 0.$$

Thus it follows that $\rho(\mathcal{G}_1) \geq \rho(\mathcal{G}_1)$. Note that $(\mathcal{A}(\mathcal{G}_1)X)_{v_\eta} > (\mathcal{A}(\mathcal{G}_1)X)_{v_\eta}$ and $\mathcal{G}_1' \cong \mathcal{G}_2$. As Subcase 1.1.1.1, we get that $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

Subcase 1.1.1.3 $x_{v_0} \leq x_{v_{a(\eta,1)}}, x_{v_{L_3}} > x_{v_{a(\eta,2)}}, x_{v_{\xi}} \leq x_{v_{a(\eta,3)}}$. Let $\varepsilon'_i = (\varepsilon_i \setminus \{v_0\}) \cup \{v_{a(\eta,1)}\}$ for $1 \leq i \leq \eta$, $e'_{\eta} = (e_{\eta} \setminus \{v_{a(\eta,2)}\}) \cup \{v_{L_3}\}, e'_{L_3+1} = (e_{L_3+1} \setminus \{v_{L_3}\}) \cup \{v_{a(\eta,2)}\}, e'_{\xi+1} = (e_{\xi+1} \setminus \{v_{\xi}\}) \cup \{v_{a(\eta,3)}\}.$ Let

$$G_{1}^{'} = G_{1} - \sum_{i=1}^{\eta} \varepsilon_{i} + \sum_{i=1}^{\eta} \varepsilon_{i}^{'} - e_{L_{3}+1} + e_{L_{3}+1}^{'} - e_{\eta} + e_{\eta}^{'} - e_{\xi+1} + e_{\xi+1}^{'}.$$

As Subcase 1.1.1.1 and Subcase 1.1.1.2, we get that $\rho(\mathcal{G}_1) > \rho(\mathcal{G}_1)$, and $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

In the same way, for the subcases: (i) $x_{v_0} \leq x_{v_{a(\eta,1)}}, x_{v_{L_3}} \leq x_{v_{a(\eta,2)}}, x_{v_{\xi}} > x_{v_{a(\eta,3)}}$; (ii) $x_{v_0} > x_{v_{a(\eta,1)}}, x_{v_{L_3}} > x_{v_{a(\eta,2)}}, x_{v_{\xi}} \leq x_{v_{a(\eta,3)}}$; (iii) $x_{v_0} > x_{v_{a(\eta,1)}}, x_{v_{L_3}} > x_{v_{a(\eta,2)}}, x_{v_{\xi}} > x_{v_{a(\eta,3)}}$; (iv) $x_{v_0} \leq x_{v_{a(\eta,1)}}, x_{v_{L_3}} > x_{v_{a(\eta,2)}}, x_{v_{\xi}} > x_{v_{a(\eta,3)}}$; (iv) $x_{v_0} \leq x_{v_{a(\eta,1)}}, x_{v_{L_3}} > x_{v_{a(\eta,2)}}, x_{v_{\xi}} > x_{v_{a(\eta,3)}}$; (v) $x_{v_0} > x_{v_{a(\eta,1)}}, x_{v_{L_3}} \leq x_{v_{a(\eta,2)}}, x_{v_{\xi}} > x_{v_{a(\eta,3)}}$, we get that $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

As a result, from the above narrations for Subcase 1.1.1 that $x_{v_{\eta}} \ge x_{v_{L_3}}$, we get $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

Subcase 1.1.2 $x_{v_{\eta}} < x_{v_{L_3}}$. By Lemma 3.11, we know that $x_{v_{\eta-1}} \leq x_{v_{L_3+1}}$. By considering the comparisons between x_{v_0} and $x_{v_{a(L_3+1,1)}}$, between $x_{v_{\xi}}$ and $x_{v_{a(L_3+1,2)}}$, between $x_{v_{\eta}}$ and $x_{v_{a(L_3+1,3)}}$, as Subcase 1.1.1, we get that $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.

Subcase 1.2 $t - p < L_0$ (see Fig. 3.7). Let $\omega = L_1 + L_2$, $\varphi = L_1 + L_2 + L_3 + 1$. By considering the comparisons between x_{v_0} and $x_{v_{a(\varphi,1)}}$, between $x_{v_{L_1}}$ and $x_{v_{a(\varphi,2)}}$, between $x_{v_{\omega}}$ and $x_{v_{a(\varphi,3)}}$, as Subcase 1.1, we get that $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$.



Case 2 p = 0 (see Fig. 3.8). Let $\kappa = L_0 + L_1$, $\varsigma = L_0 + L_1 + L_2$. By considering the comparisons between $x_{v_{a(1,1)}}$ and $x_{v_{L_0}}$, between $x_{v_{a(1,2)}}$ and $x_{v_{\kappa}}$, between $x_{v_{a(1,3)}}$ and $x_{v_{\varsigma}}$, as Case 1, we get that $\rho(\mathcal{G}_2) > \rho(\mathcal{G}_1)$. This completes the proof. \Box

Proof of Theorem 1.3. For k-uniform supertrees of order n, using Lemma 3.10 and Lemma 3.13 repeatedly gets the result. This completes the proof. \Box

Proof of Corollary 1.4. Note that $S^*(n,k)$ is the *k*th power of the ordinary star $S^*(\frac{n-1}{k-1}+1,2)$, $\mathcal{P}(n,k)$ is the *k*th power of the ordinary path $P(\frac{n-1}{k-1}+1,2)$. Then the result follows from Theorem 1.1 and Theorem 1.3. This completes the proof. \Box

References

- [1] A. Banerjee, A. Char, B. Mondal, Spectra of general hypergraphs, Linear Algebra Appl. 518 (2017) 14-30.
- [2] A. Banerjee, On the spectrum of hypergraphs, Linear Algebra Appl. 614 (2021) 82-110.
- [3] A. Banerjee, A. Sarkar, On the spectral radius of some linear hypergraphs, https://doi.org/10.48550/arXiv.2303.14545.
- [4] K. Chang, K. Pearson, T. Zhang, Perron-Frobenius Theorem for nonnegative tensors, Commun. Math. Sci. 6 (2008) 507-520.
- [5] K. Chang, K. Pearson, T. Zhang, On eigenvalue problems of real symmetric tensors, J. Math. Anal. Appl. 350 (2009) 416-422.
- [6] J. Cooper, A. Dutle, Spectra of uniform hypergraphs. Linear Algebra Appl. 2012; 436: 3268-3292.
- [7] Y. Fan, H. Yang, J. Zheng, High-ordered spectral characterization of unicyclic graphs, Discuss. Math. Graph Theory, in press, https://doi.org/10.7151/dmgt.2489.
- [8] S. Friedland, S. Gaubert, L. Han, Perron-Frobenius theorem for nonnegative multilinear forms and extensions, Linear Algebra Appl. 438 (2013), 738-749.
- [9] D. Gao, Duality Principles in Nonconvex Systems, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2000.
- [10] H. Guo, B. Zhou, On the spectral radius of uniform hypertrees, Linear Algebra Appl. 558 (2018) 236-249.
- Y. Hou, A. Chang, J. Cooper, Spectral Extremal Results for Hypergraphs, Electronic Journal of Combinatorics 28 (2021) DOI: https://doi.org/10.37236/9018.
- [12] K. Huang, M. Xue, M. Lu, Tensor Analysis, second ed., Tsinghua University Publisher, Beijing, 2003 (in Chinese).
- [13] S. Hu, L. Qi, J. Shao, Cored hypergraphs, power hypergraphs and their Laplacian eigenvalues, Linear Algebra Appl. 439 (2013) 2980-2998.
- [14] M. Khan, Y. Fan, On the spectral radius of a class of non-odd-bipartite even uniform hypergraphs. Linear Algebra Appl. 480 (2015) 93-106.
- [15] K. Li, A. Huang, Tensor Analysis and Their Applications, Scientific Publisher, Beijing, 2004 (in Chinese).
- [16] G. Li, L. Qi, G. Yu, The Z-eigenvalues of a symmetric tensor and its application to spectral hypergraph theory, Numer. Linear Algebra Appl. 20 (2013) 1001-1029.

- [17] H. Li, J. Shao, L. Qi, The extremal spectral radii of k-uniform supertrees, J. Comb. Optim. 32 (2016), 741-764.
- [18] L. Lim, Singular values and eigenvalues of tensors: a variational approach, in: Proceedings of the IEEE International Workshopon Computational Advances in Multi-Sensor Adaptive Processing, CAMSAP 05, 1 (2005) 129-132.
- [19] L. Lim, Eigenvalues of tensors and some very basic spectral hypergraph theory, matrix computations and scientific computing seminar, April 16, 2008. http://www.stat.uchicago.edu/lekheng/work/mcsc2.
- [20] H. Lin, B. Zhou, B. Mo, Upper bounds for H and Z-spectral radii of uniform hypergraphs, Linear Algebra Appl. 510 (2016) 205-221.
- [21] L. Liu, L. Kang, S. Bai, Bounds on the spectral radius of uniform hypergraphs, Discrete Appl. Math. 259 (2019) 160-169.
- [22] L. Lu, S. Man, Connected hypergraphs with small spectral radius, Linear Algebra Appl. 509 (2016) 206-227.
- [23] H. Minc, Nonnegative Matrices, John Wiley & Sons Inc., New York, 1988.
- [24] M. Ng, L. Qi, G. Zhou, Finding the largest eigenvalue of a nonnegative tensor, SIAM J. Matrix Anal. Appl. 31 (3) (2009) 1090-1099.
- [25] C. Ouyang, L. Qi, X. Yuan, The first few unicyclic and bicyclic hypergraphs with largest spectral radii, Linear Algebra Appl. 527 (2017) 141-162.
- [26] O. Perron, Zur theorie dor matrizan, Math Ann., 64 (1907), 248-263.
- [27] L. Qi, Eigenvalues of a real supersymmetric tensor, J. Symbolic Comput. 40 (2005) 1302-1324.
- [28] L. Qi, Symmetric nonnegative tensors and copositive tensors, Linear Algebra Appl. 439 (2013) 228-38.
- [29] L. Qi, Z. Luo, Tensor Analysis: Spectral Theory and Special Tensors, SIAM, 2017.
- [30] J. Shao, A general product of tensors with applications. Linear Algebra Appl. 439 (2013) 2350-2366.
- [31] Y. Talpaert, Tensor Analysis and Continuum Mechanics, Kluwer Academic Publishers, Dordrecht, The Netherlands, 2002.
- [32] J. Shao, H. Shan, L. Zhang, On some properties of the determinants of tensors, Linear Algebra Appl. 439 (2013) 3057-3069.
- [33] L. Su, L. Kang, The matching polynomials and spectral radii of uniform supertrees, Electron. J. Comb. 25(4) DOI: 10.37236/7839
- [34] P. Xiao, L. Wang, Y. Lu, The maximum spectral radii of uniform supertrees with given degree sequences, Linear Algebra Appl. 523 (2017) 33-45.
- [35] Y. Yang, Q. Yang, Further results for Perron-Frobenius theorem for nonnegative tensors, SIAM J. Matrix Anal. Appl. 31 (5) (2010) 2517-2530.
- [36] Y. Yang, Q. Yang, Further results for perron-frobenious theorem for nonnegative tensors II, SIAM J. Matrix Anal. Appl. 32 (4) (2011) 1236-1250.
- [37] C. Lv, L. You, X. Zhang, A sharp upper bound on the spectral radius of a nonnegative k-uniform tensor and its applications to (directed) hypergraphs, J. Inequal. Appl. 1 (2020) DOI: 10.1186/s13660-020-2305-2.
- [38] G. Yu, J. Shu, Yuan Hong, Bounds of the spectral radii of $K_{2,3}$ -minor free graphs, Electron. J. Linear Algebra 23 (2012), 171-179.
- [39] G. Yu, C. Yan, L. Sun, Y. Wu, H. Zhang, Spectral radius and matching number of the unicyclic hypergraph, Linear Algebra Appl. 610 (2021) 571-590.
- [40] X. Yuan, J. Shao, H. Shan, Ordering of some uniform supertrees with larger spectral radii, Linear Algebra Appl. 495 (2016) 206-222.
- [41] J. Zhang, J. Li, H. Guo, Uniform hypergraphs with the first two smallest spectral radii, Linear Algebra Appl. 594 (2020) 71-80.
- [42] J. Zhou, L. Sun, W. Wang, C. Bu, Some spectral properties of uniform hypergraphs, Electron. J. Comb. 21 (2014) P4.24.