Exponential Tail Bounds on Queues: A Confluence of Non-Asymptotic Heavy Traffic and Large Deviations

Prakirt Raj Jhunjhunwala^a, Daniela Hurtado-Lange^b, Siva Theja Maguluri^c

^a Georgia Institute of Technology, North Avenue, Atlanta, 30332 GA, USA
 ^b William and Mary, Williamsburg, VA 23185, USA
 ^c Georgia Institute of Technology, North Avenue, 30332 GA, Atlanta, USA

Abstract

In general, obtaining the exact steady-state distribution of queue lengths is not feasible. Therefore, our focus is on establish bounds for the tail probabilities of queue lengths. Specifically, we examine queueing systems under Heavy-Traffic (HT) conditions and provide exponentially decaying bounds for the probability $\mathbb{P}(\epsilon q > x)$, where ϵ is the HT parameter denoting how far the load is from the maximum allowed load. Our bounds are not limited to asymptotic cases and are applicable even for finite values of ϵ , and they get sharper as $\epsilon \to 0$. Consequently, we derive non-asymptotic convergence rates for the tail probabilities. Unlike other approaches such as moment bounds based on drift arguments and bounds on Wasserstein distance using Stein's method, our method yields sharper tail bounds. Furthermore, our results offer bounds on the exponential rate of decay of the tail, given by $-\frac{1}{x} \log \mathbb{P}(\epsilon q > x)$ for any finite value of x. These can be interpreted as non-asymptotic versions of Large Deviation (LD) results. To obtain our results, we use an exponential Lyapunov function to bound the moment generating function of queue lengths and apply Markov's inequality.

We demonstrate our approach by presenting tail bounds for: (i) a continuous time Join-the-shortest queue (JSQ) load balancing system, (ii) a discrete time single-server queue and (iii) an M/M/n queue. We not only bridge the gap between classical-HT and LD regimes but also explore the large system HT regimes for JSQ and M/M/n systems. In these regimes, both the system size and the system load increase simultaneously. Our results also close a gap in the existing literature on the limiting distribution of JSQ in the super-NDS (a.k.a. super slowdown) regime. This contribution is of an independent interest. Here, a key ingredient is a more refined characterization of state space collapse for JSQ system, achieved by using an exponential Lyapunov function designed to approximate the ℓ_{∞} norm.

Keywords: Classical heavy traffic, Large deviations, Tail probabilities, Join-the-shortest queue, Transform Method, exponential Lyapunov function

1. Introduction

Queueing models are used to study performance of many systems such as cloud computing, data centers, ride hailing, call centers etc. In general, obtaining the complete distribution of queue lengths in these systems is intractable. Therefore, a common approach is to study asymptotic regimes. There are several popular regimes such as Heavy Traffic (HT), large scale regime, or Large Deviations (LD). In the HT regime, the system is loaded close to its maximum capacity while keeping the number of servers fixed. In the large-systems regime, the system's load is fixed, but the number of servers is increased to infinity. And in the LD limit one studies the probability of rare events, that is, the tail probability for large thresholds. Recently, the Many-Server Heavy-Traffic (Many-Server-HT) regime has gained more popularity, where the system is

Email addresses: prakirt@gatech.edu (Prakirt Raj Jhunjhunwala), dahurtadolange@wm.edu (Daniela Hurtado-Lange), siva.theja@gatech.edu (Siva Theja Maguluri)

loaded to maximum capacity while simultaneously increasing the number of servers. The system's behavior varies greatly depending on how quickly the load increases relative to the number of servers. As such one employs very different analysis techniques to study queueing systems in different regimes.

In the study of HT asymptotics, one typically scales the queue lengths using a parameter that represents the system's load. By denoting the load as $1 - \epsilon$, the HT limit is achieved when ϵ approaches zero. Most of the literature focuses on systems that satisfy the so-called Complete Resource Pooling (CRP) condition and behave like a single-server queue in the limit. For such systems, it is well-known that the scaled queue length follows an exponential distribution in the HT limit, which gives the tail probabilities of the limiting system. However, the rate of convergence of the tail probabilities (of the pre-limit system) to the corresponding HT value remains unknown.

Most real world systems involve Service Level Agreements (SLA), where customers are promised a specific level of service, including the maximum delay they can expect. Motivated by this, in this paper, we focus on establishing sharp bounds on the tail probabilities of scaled queue length of the pre-limit system, i.e., for $\epsilon > 0$. In particular, we get non-asymptotic bounds of the form

$$\mathbb{P}(\epsilon q > x) \le \kappa(\epsilon, x) e^{-\theta(\epsilon)x},$$

where q represents the total queue length in steady state. Here, $\theta(\epsilon)$ gives the decay rate of the tail probability of the pre-limit system, and $\theta(\epsilon)$ converges to the correct HT value as $\epsilon \to 0$. Recent results show the rate of convergence to HT in terms of the mean, moments, or Wasserstein's distance (for references on each of these, see Section 1.3). These methods focus on the entire distribution of the queue lengths and drown the tail. For example, consider the second moment, and suppose ϵq converges in distribution to the random variable Υ . Then, from existing results, one obtains that $|\mathbb{E}[\epsilon^2 q^2] - \mathbb{E}[\Upsilon^2]|$ is $O(\epsilon)$, which gives a valid bound. From these results, one can obtain bounds in terms of tail probability of the form $|\mathbb{P}(\epsilon q > x) - \mathbb{P}(\Upsilon > x)| \leq O(\epsilon)$. However, these are not very informative as the tail probability itself can be much smaller than $O(\epsilon)$. Therefore, the rate of convergence of tail probabilities cannot be obtained using the existing methodologies. In this work, we correctly characterize $\theta(\epsilon)$ to obtain the rate of convergence of tail probability to the corresponding HT value. Our results are non-asymptotic in the sense that they are valid whenever ϵ is small, and not just when $\epsilon \to 0$. Also, our results are precise when ϵ gets closer to 0, recovering the HT results.

Our work bridges the gap between the LD, HT, and Many-Server-HT regimes. When one studies the LD regime, the goal is to find the exponential rate at which the tail probability decays, which is precisely given by $\theta(\epsilon)$. As such, our tail bounds can be used to recover the non-asymptotic LD results. Thus, our tail bounds are at a confluence of non-asymptotic HT and non-asymptotic LD. This extends the understanding of tail behavior beyond the classical HT regime. To the best of our knowledge, such comprehensive LD results have not been previously reported in the existing literature.

1.1. Main contribution

We illustrate our methodology by providing results for three well-studied systems, viz, a load-balancing system under Join the Shortest Queue (hereafter referred to as the JSQ system), a discrete-time Single-Server Queue (SSQ), and a multi-server system with a single queue (M/M/n queue). Our contributions for each of these systems are mentioned below.

1.1.1. JSQ system

We consider a continuous-time system with n servers, each of them with its own queue. Jobs arrive according to a Poisson process with rate λ_n , and are routed to the server with the shortest queue (breaking ties at random). Further, the service times are exponentially distributed with rate μ . In the context of the JSQ system, we consider the Many-Server-HT regime, where the system size n grows to infinity while the HT parameter ϵ_n approaches zero. Specifically, we consider $\epsilon_n = n^{-\alpha}$ with $\alpha > 1$ constant, and take the limit as $n \to \infty$. For the JSQ system, we have the following two contributions.



Figure 1: Visual representation of the condition $n\epsilon_n \log \frac{1}{\epsilon_n} < \delta$, where δ is a small constant.

• Tail probability: In Theorem 2, we show that the tail probability of the steady-state scaled total queue length satisfies

$$\frac{1}{1-\epsilon_n}e^{-\theta_n x} \le \mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) \le 2ex\Big(1+\kappa_2 n\epsilon_n \log \frac{1}{\epsilon_n}\Big)e^{-\theta_n x},$$

where $\theta_n := \frac{1}{\epsilon_n} \log \frac{1}{1-\epsilon_n}$. The lower bound is valid for all values of n and ϵ_n , while the upper bound holds when the term $n\epsilon_n \log \frac{1}{\epsilon_n}$ is sufficiently small. The upper bound leverages the State Space Collapse (SSC) property of the JSQ system, where the *n*-dimensional state vector collapses to a onedimensional subspace, so the JSQ system behaves like an SSQ. The SSC property holds only when the load is sufficiently large, and so we need $n\epsilon_n \log \frac{1}{\epsilon_n}$ to be sufficiently small. This is satisfied for sufficiently large n, when $\epsilon_n = n^{-\alpha}$ with $\alpha > 1$.

Figure 1 visually illustrates the condition discussed above. Region 1 represents values of n and α where $n\epsilon_n \log \frac{1}{\epsilon_n}$ is small, and our tail bounds for the JSQ system hold within this region. As α approaches 1, satisfying the condition $n\epsilon_n \log \frac{1}{\epsilon_n} < \delta$ becomes more challenging, leading to a regime change consistent with existing literature [1]. In Region 2, we believe that the HT approximation fails to accurately describe the system dynamics. Finally, Region 3 corresponds to the range $\alpha \in [0, 1]$, where the results and analysis techniques significantly differ from those presented in this paper.

We also obtain an LD result for the JSQ system, i.e., we have

$$\lim_{x \to \infty} \frac{1}{x} \log \mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) = -\theta_n.$$

Our LD result concerning the scaled total queue length is asymptotically precise as ϵ_n approaches zero in Many-Server-HT settings (provided that $\alpha > 1$).

Importantly, our work provides an exact characterization of the decay rate of the tail probabilities for the pre-limit system. This is possible because we establish both upper and lower bounds on the tail probability. As a result, our research represents a significant advancement compared to existing literature. Thus, for JSQ, our result not only bridges the gap between HT and LD, but also connects these to (some) many server regimes.

• Limiting distribution: In Theorem 1, we show that as $n \to \infty$, all the (scaled) queue lengths become identical and are exponentially distributed with mean 1, i.e., in the Many-Server-HT (i.e., $\epsilon_n = n^{-\alpha}$ with $\alpha > 1$), we prove that

$$n\epsilon_n \mathbf{q} \stackrel{d}{\to} \Upsilon \mathbf{1},$$

where **q** is the queue length vector in steady state, Υ is an exponentially distributed random variable, and **1** is a vector of all ones. Prior work [2] establishes such a result only for $\alpha > 2$ and leaves the regime $1 < \alpha \leq 2$ open. This result thus fills the gap in the literature and in conjunction with all the prior work [2], and completes our understanding of JSQ in all of the Many-Server-HT regimes. A key step to this end is in establishing SSC for the JSQ system especially in the regimes $\alpha \in (1, 2]$. Our approach in establishing SSC result for the JSQ system uses a novel Lyapunov function, and differs from the approach based on the drift arguments of [3] as used in prior work, [4, 5, 6]. Intuitively, instead of working with the ℓ_2 distance of the queue length vector from the one dimensional subspace, our key idea is that one should work with the ℓ_{∞} distance which gives sharper bounds. However, since the ℓ_{∞} distance is not easily amenable to drift arguments, we work with a Lyapunov function that can be interpreted as a smooth approximation of the ℓ_{∞} distance.

1.1.2. Discrete time Single-Server Queue

We consider a discrete-time SSQ with general arrival and service distributions, characterized by mean arrival rate λ_{ϵ} and mean service rate μ_{ϵ} , respectively. We introduce the parameter ϵ to represent the HT condition, where $\lambda_{\epsilon} = \mu_{\epsilon}(1-\epsilon)$. Additionally, we define σ_{ϵ}^2 as the sum of the variances of the arrival and service processes.

In Theorem 5, we obtain the following tail bound and LD result for the pre-limit system:

$$\mathbb{P}(\epsilon q > x) \le e\theta_{\epsilon} x e^{-\theta_{\epsilon}(1-\kappa_{1}\epsilon)x}, \qquad \qquad \lim_{x \to \infty} \frac{1}{x} \log \mathbb{P}(\epsilon q > x) \le -\theta_{\epsilon}(1-\kappa_{1}\epsilon)$$

Here, $\theta_{\epsilon} = \frac{2\mu_{\epsilon}}{\sigma_{\epsilon}^2}$ and κ_1 are defined in terms of the system parameters and moments of the arrival and service processes. Also, the constant κ_1 depend on the thrid moments of arrival and service. This result exhibit the influence of a general arrival and service distribution on the decay rate of the tail probability under non-asymptotic HT conditions and also provide a non-asymptotic LD result for the steady-state queue length.

1.1.3. M/M/n system

An M/M/n queue is a multi-server system with a single queue, where the inter-arrival and service times (of each server) are exponentially distributed, with rates λ_n and μ , respectively. To parameterize the Many-Server-HT regime, we use $\lambda_n = n\mu(1-\epsilon_n)$. In this case, we consider two separate quantities: (i) the number of waiting customers, denoted by w_n , and (ii) the number of idle servers, denoted by r_n ; and provide tail probabilities for both quantities under proper scaling.

Our approach provides results in three different Many-Server-HT regimes, viz. Sub-Halfin-Whitt (Sub-HW) regime when $\alpha \in (0, \frac{1}{2})$, Halfin-Whitt (HW) regime when $\alpha = \frac{1}{2}$, and Super-Halfin-Whitt (Super-HW) regime when $\alpha \in (\frac{1}{2}, \infty)$. A brief overview of our results for the M/M/n system is presented in Table 1 below. More details are provided in Section 4.

The limiting distribution of the scaled number of waiting customers and the scaled number of idle servers were established in [7] so we skip mentioning that here. However, one can easily verify that our results recover the limiting distribution as $n \to \infty$. Note that the probability of there being any idle server, i.e., $\mathbb{P}(r_n > 0)$ goes to zero in the Super-HW regime. As such, the tail bound on r_n is useful only in Sub-HW and HW regime. Similarly, the tail bound on w_n is valid only in HW and Super-HW regimes.

	Sub-HW	Halfin-Whitt	Super-HW
$\epsilon_n = n^{-\alpha}$ with	$\alpha \in \left(0, \frac{1}{2}\right)$	$\alpha = \frac{1}{2}$	$\alpha > \frac{1}{2}$
$\mathbb{P}(\epsilon_n w_n > x w_n > 0)$	$O(e^{- heta_n x})$		
$\mathbb{P}(\eta_n \tilde{r}_n > x r_n > 0)$	$O(e^{-\frac{1}{2}x^2})$		
$\mathbb{P}(w_n > 0)$	$O(n^{\alpha - \frac{1}{2}} e^{-n\epsilon_n})$	$\rightarrow p < 1$	$\rightarrow 1$
$\mathbb{P}(r_n > 0)$	$\rightarrow 1$	$\rightarrow 1 - p < 1$	$O(n^{\frac{1}{2}-\alpha})$

Table 1: A brief overview of our results for the M/M/n queue. Here, $\tilde{r}_n := r_n - n\epsilon_n$. The scaling parameter for w_n is the HT parameter ϵ_n , while the scaling parameter for \tilde{r}_n is $\eta_n := \frac{1}{\sqrt{n(1-\epsilon_n)}}$. Also, p is a constant given in Theorem 8.b.

1.2. Key aspects of our approach

The key idea of this paper is to leverage the existence of the Moment Generating Function (MGF) of the scaled queue length process in the vicinity of zero to obtain the decay rate of the tail probability. Let's consider a non-negative random variable X and a constant a > 0 such that $\mathbb{E}[e^{aX}] \leq \kappa$. By applying Markov's inequality, we can establish that $\mathbb{P}(X > x) \leq \kappa e^{-ax}$. This implies that the tail of X decays exponentially with rate a. Thus, determining the appropriate decay rate of the tail probability reduces to finding the largest a such that $\mathbb{E}[e^{aX}]$ is bounded by a constant.

We adopt this approach to derive tail bounds for the scaled steady-state queue-length process. We establish an upper bound on the MGF of the scaled queue lengths, denoted as $\mathbb{E}[e^{\theta \epsilon q}]$, for a large range of θ . To determine the correct range of θ , we approximate the behavior of the actual system by considering its HT counterpart. By doing so, we also obtain the rate of convergence of the tail probabilities to their corresponding values in the HT regime.

The use of exponential Lyapunov functions to study queue-length behavior was introduced as a transform method to obtain HT results in [6]. To get tight pre-limit results, we use the same exponential test function, but we bound the error terms in a more refined manner. As such, our technique is divided into three steps: (i) derive the MGF (or an upper bound) of the scaled queue length, $\mathbb{E}[e^{\theta \epsilon q}]$, for a large range of values of θ , (ii) approximate the MGF with its HT counterpart, and (iii) use Markov's inequality and optimize over the values of θ . In the process, we encounter three multiplicative terms that are required to characterize the tail probability.

The first term is called the SSC violation. As mentioned before, JSQ system satisfies SSC in HT. However, in non-asymptotic HT conditions, SSC is not fully satisfied. As such, the term SSC violation accounts for the level to which SSC is violated in a pre-limit system. The second one is the pre-limit tail. When the HT parameter ϵ is greater than zero, the decay rate of the tail probability deviates slightly from its HT value, and we refers to this correct decay rate as the pre-limit tail. The third term is referred to as the pre-exponent error. Our approach uses Markov's inequality to compute bounds tail probabilities from the MGF. However, Markov's inequality incurs a cost while obtaining the tail bound, which is captured by the pre-exponent error.

As a consequence, we easily obtain a LD result after characterizing a bound on the tail probability. Mathematically, as $\mathbb{E}[e^{aX}] \leq \kappa$ implies $\mathbb{P}(X > x) \leq \kappa e^{-ax}$, we also get $\lim_{x\to\infty} \frac{1}{x} \log \mathbb{P}(X > x) \leq -a$. While characterizing the LD result, the pre-limit error term plays a key role. The other two error terms, namely the pre-exponent error and the SSC error, vanish as $x \to \infty$. Therefore, the pre-limit error term is essential for accurately characterizing the LD behavior of the system.

1.3. Related work

Most of the literature studying the HT behavior of various queues uses a methodology frequently called 'diffusion limits.' A non-exhaustive sample of articles using this method is [8, 9, 10, 11, 12]. Under this approach, the scaled system is shown to converge to a Reflected Brownian Motion (RBM) in HT, and the

α	Regime	Reference
$\alpha \downarrow 0$	Mean Field	[22, 23, 24, 25, 15]
$\alpha \in \left(0, \frac{1}{2}\right)$	Sub-Halfin-Whitt	[26, 27, 28]
$\alpha = \frac{1}{2}$	Halfin-Whitt	[29, 30, 31, 32, 33]
$\alpha \in \left(\frac{1}{2}, 1\right)$	Super-Halfin-Whitt	[34, 35]
$\alpha = 1$	Non-Degenerate Slowdown	[1]
$\alpha \in (1,2]$	Super Slowdown	This Work
$\alpha > 2$		[2]
$\alpha = \infty$	Classic-HT	[4, 36, 37, 10, 6, 38, 9]

Table 2: Literature review of asymptotic regimes for Load balancing system.

steady-state behavior of this RBM is studied. The final step is the so-called interchange-of-limit proof which is usually remarkably challenging.

More recently, there has been an increased number of papers using alternative methods that do not require the interchange-of-limits step. These are Stein's method [13, 14, 15, 16], the BAR approach [17, 18, 19], and the drift method [4, 6, 20]. All these methods have something in common with the work we present in this paper and, at the same time, are substantially different.

In the first approach, i.e., the Stein's method, one derives bounds for the Wasserstein distance [21] between the pre-limit system and the limiting distribution, enabling one to obtain the convergence rates. Similar to this, we also establish bounds that depict the similarity between the pre-limit system and the limiting distribution. However, our focus lies in directly characterizing the tail probability. In the second method, i.e., the BAR approach, the goal is to establish a certain equation, called Basic Adjoint Relationship (BAR), in terms of the MGF of the steady state distribution of the limiting distribution. In contrast, we direct work with the MGF of the pre-limit system.

In the third method, i.e., the drift method, the main idea is to use steady-state conditions and carefully chosen test functions to compute bounds that are tight in heavy traffic. Within this method, the use of exponential test functions contributed to the development of the transform method [6, 20]. Our work in this paper is inspired by the transform method in the sense that using exponential test functions and careful manipulation of the queue-length dynamics are essential to obtain the results. In contrast to the transform method, which only focuses on the limiting distribution, we carefully compute the error terms also to obtain the rate of convergence of the tail probabilities. To the best of our knowledge, the rate of convergence to heavy traffic in terms of tail probabilities has been not been known before.

The literature on Many-Server-HT asymptotics is sub-divided in multiple categories depending on how fast the load increases with respect to the number of servers. Using the parameterization $\epsilon_n = n^{-\alpha}$ with $\alpha > 0$ introduced above, different regimes are obtained depending on the value of α . The literature on Many-Server-HT with $\alpha \in (0, 1]$ is vast and uses different analysis techniques. We direct the readers to [35] and references therein for more details on Many-Server-HT with $\alpha \in (0, 1]$. Closest to our work, in [2], it is proved that for $\alpha > 2$ the scaled total queue length is exponentially distribution in limit as $n \to \infty$. In our work, we close the gap between the result in [2] and the Non-Degenerate Slowdown (NDS) [39] regime ($\alpha = 1$), and we obtain the HT behavior for all $\alpha > 1$.

Large Deviations (LD) is a popular regime when one needs to study the performance of a control policy for routing or scheduling jobs. A comprehensive report on the application of large deviation theory to queueing problems can be found in [40]. In [41], it was proved that the if the sequence of arrivals and services follow a certain Large Deviation Principle (LDP), then, one can obtain the decay rate of the tail probability of the associated queue length. This argument was used in [42, 43] to prove a LDP for the queue length process of a single server process, with further generalization in [44]. Recently, there has been

Exponential Tail Bounds on Queues

quite some work on establishing a LDP for the JSQ system, see [45, 46, 47] and the references therein. In contrast to the existing literature, our work provides a LD result as a simple closed form expression in a non-asymptotic LD regime. Notably, we also provide an upper bound on the pre-exponent term, which is typically absent in previous works. Moreover, our work considers the JSQ system in a Many-Server regime, and incorporates the crucial phenomenon of State Space Collapse. Such a phenomenon was considered in [46], but only for a JSQ system of size 2.

In addition to proving a LDP, an intriguing research area involves the use of Lyapunov functions to develop policies that minimize the probability of queue overflow [48, 49, 50]. The focus in [48, 49] is on minimizing the decay rate of tail probability, and non-asymptotic large deviation results are not provided. Another related research field examines queues with Gaussian input, which are particularly challenging to analyze in most scenarios [51, 52]. These are related to ours in the sense that all compute tight bounds that characterize the probability of rare events. However, the methodologies are different, and they focus on the LD regime only. In this paper, we obtain LD results that are closely connected to the tail probabilities of the queue lengths in HT and Many-Server-HT.

2. Join-the-shortest queue system

2.1. Model

We consider a continuous-time queueing system consisting of n SSQs in parallel, each serving jobs according to first-come-first-serve. At any time t, let $\mathbf{q}(t)$ denote the queue length vector, where $q_i(t)$ is the queue length of i^{th} queue. For the ease of notation, we use $\overline{q}(t)$ to denote the total queue length at time t, i.e., $\overline{q}(t) = \sum_{i=1}^{n} q_i(t)$. Jobs arrive to the system according to a Poisson process with rate λ_n , and service times are exponentially distributed with rate μ .

When a job arrives, it is dispatched according to JSQ, that is, the job is sent to the queue with index

$$i^*(t) \in \arg\min q_i(t),$$

breaking ties uniformly at random. Under the JSQ policy, the queue-length process $\{\mathbf{q}(t)\}_{t\geq 0}$ is a Continuous-Time Markov Chain (CTMC). Further, it is well-known that the queue-length process is stable (positive recurrent) if the arrival rate is strictly smaller than the total service rate $(\lambda_n < n\mu)$. In this work, we assume that the system satisfies the stability condition. Under this assumption, the steady-state distribution of the queue-length process $\{\mathbf{q}(t)\}_{t=0}^{\infty}$ exists, and we denote by π_n . We use \mathbf{q} to denote the steady-state queue length vector, that is, \mathbf{q} follows the distribution π_n , and $\overline{q} := \sum_{i=1}^n q_i$. The system load is $\rho_n := \frac{\lambda_n}{n\mu}$, and we define $\epsilon_n = 1 - \rho_n$. Then, the system approaches HT as $\epsilon_n \to 0$.

The system load is $\rho_n := \frac{\lambda_n}{n\mu}$, and we define $\epsilon_n = 1 - \rho_n$. Then, the system approaches HT as $\epsilon_n \to 0$. We consider the JSQ system in HT and Many-Server-HT. In (classical) HT, the system size n is a constant, and the HT parameter ϵ_n does not depend on n. Then, we drop the subscript n and take the limit $\epsilon \to 0$. In the Many-Server-HT, the load and the number of servers increase together. Then, we consider $\epsilon_n = n^{-\alpha}$ and take the limit as $n \to \infty$. In this work, in case of Many-Server-HT we only consider the case when $\alpha > 1$, which is also known as super-slowdown regime. In both regimes, we aim to provide a tail bound on the total queue length, i.e., a bound on $\mathbb{P}(\epsilon_n \overline{q} > x)$.

2.2. Results for JSQ system

It is well known that the HT distribution of the scaled steady-state total queue length $\epsilon \bar{q}$ converges to an exponential random variable as $\epsilon \to 0$ [6]. Further, as shown in [2], the result extends to Many-Server-HT, where $\epsilon_n \bar{q}$ converges to an exponential random variable in distribution if $\alpha > 2$.

In Theorem 1, we complete the result by demonstrating that $\epsilon_n \overline{q}$ converges in distribution to an exponential random variable $\alpha > 1$. Our result enhances the understanding of the behavior of the scaled queue length in Many-Server-HT under a broader range of traffic conditions, encompassing values of $\alpha \in (1, 2]$.

Theorem 1. Consider the JSQ system as presented in Section 2.1. Suppose the system satisfies the condition $\lambda_n = n\mu(1 - n^{-\alpha})$, i.e., $\epsilon_n = n^{-\alpha}$, where $\alpha > 1$. Then, for any $\theta < 1$, we have

$$\lim_{n \to \infty} \mathbb{E}[e^{\theta n^{-\alpha} \sum_{i=1}^{n} q_i}] = \frac{1}{1 - \theta}, \qquad n^{1 - \alpha} \mathbf{q} \stackrel{d}{\to} \Upsilon \mathbf{1}, \quad as \quad n \to \infty,$$

where Υ is an exponential random variable with mean 1.

Theorem 1 provides the limiting distribution of the steady-state scaled queue length vector as $n \to \infty$ for all $\alpha > 1$. The proof of Theorem 1 relies on the fact that the JSQ system satisfies SSC, where the *n*dimensional state vector of the system collapses to a one-dimensional subspace. More precisely, as $n \to \infty$, we have that $n^{1-\alpha}q_i \approx n^{-\alpha} \sum_{j=1}^n q_j$ for all $i \in \{1, 2, \ldots, n\}$. The correct characterization of SSC in the Many-Server-HT regime is crucial in proving Theorem 1 and in completing the result of [2] for all $\alpha > 1$. More details on SSC for the JSQ system in Many-Server-HT are provided in Proposition 3 in Section 2.3. A proof sketch for Theorem 1 is provided in Section 2.4 and the details are provided in Appendix B.

Next, we provide the tail bound of the scaled steady-state total queue length for the JSQ system. In Theorem 2, we provide a bound on the tail probability $\mathbb{P}(\epsilon_n \overline{q} > x)$ when the system size is finite.

Theorem 2. Consider the JSQ system as presented in Section 2.1. Suppose the system satisfies the condition $\lambda_n = n\mu(1 - \epsilon_n)$, where ϵ_n is small enough such that $n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \frac{\theta_{\perp}}{4}$, and let $\kappa_2 := \frac{4\epsilon\kappa_{\perp}^2}{\theta_{\perp}}$, where κ_{\perp} and θ_{\perp} are constants given in Proposition 3. Suppose $\theta_n := \frac{1}{\epsilon_n} \log \frac{1}{1-\epsilon_n}$ Then, for all $x > 1 - \epsilon_n$ we have

$$\mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) \le \Big[2ex\big(1 - \kappa_2 n\epsilon_n \log \epsilon_n\big)\Big]e^{-\theta_n x}.$$
(1)

Further, for any $n \ge 1$ and $\epsilon_n \in (0, 1)$, we have the lower bound

$$\mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) \ge \frac{1}{1-\epsilon_n} e^{-\theta_n x}.$$
(2)

As a consequence, we have the following large deviation result.

$$\lim_{x \to \infty} -\frac{1}{x} \log \mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) = \theta_n = \frac{1}{\epsilon_n} \log \frac{1}{1 - \epsilon_n} \in \Big(1, \frac{1}{1 - \epsilon_n}\Big).$$
(3)

Theorem 2 establishes the exponential decay of the tail of the total queue length for a JSQ system. The bounds presented in Eq. (1) and (2) are valid for both the HT and Many-Server-HT regimes. Further, the result in Theorem 2 is consistent with the fact that the distribution of the scaled steady-state total queue length, i.e., $\epsilon_n \bar{q}$, converges to an exponential random variable in distribution, as n grows to ∞ .

In Theorem 2, we are able to characterize the exact tail decay rate of the continuous time JSQ system. Our result implies that, in Many-Server-HT with $\alpha > 1$ and when the term $n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right)$ is small enough, the decay rate of the JSQ system exactly matches the tail decay rate of an SSQ. This is a significant advancement compared to existing literature. Previous work primarily focused on comparing the behavior of the JSQ system with an SSQ in the limiting traffic condition, specifically as $\epsilon_n \to 0$. In contrast, our work examines the behavior of a pre-limit JSQ system and directly compares it to the corresponding SSQ.

Remark 1. Tail probabilities are often better characterized by multiplicative rather than additive errors. For instance, a bound of the form $\left|\mathbb{P}(\epsilon q > x) - \exp\left(-\frac{2\mu x}{\sigma^2}\right)\right| \leq O(\epsilon)$ can become very loose for large values of x. An interesting example is presented in [53, Section 2.1], where the Central Limit Theorem provides an approximation of N i.i.d. random variables by a normal distribution, but the error in the approximation can be as large as $O(1/\sqrt{N})$, which is much larger than the tail itself. Thus, even if the tail probability converges to an exponentially decaying tail in the limit, the error in the approximation may still decay slowly. Therefore, concentration-inequality-type bounds (as in Theorem 5) better characterize the tail probability.

Additionally, Eq. (3) represents a significant result in the form of an LD principle. As $\epsilon_n \to 0$ in HT or Many-Server-HT, we have $\theta_n \to 1$, and so, our LD result for $\epsilon_n \overline{q}$ is asymptotically precise in both, the HT and Many-Server-HT settings (provided that $\alpha > 1$).

2.2.1. Discussion on terms in Theorem 2:

Our bounds on tail probability on JSQ system, presented in Theorem 2, can be decomposed into terms as discussed below.

• SSC violation: For the JSQ system, the SSC violation term is given by $(1 - \kappa_2 n \epsilon_n \log \epsilon_n)$. In nonasymptotic HT conditions (i.e., when $\epsilon > 0$ in HT, or $n < \infty$ in Many-Server-HT), the SSC property is not fully satisfied. This introduces an additional multiplicative term in the tail probability bound, which is captured by $1 - \kappa_2 n \epsilon_n \log \epsilon_n$, and reflects the extent to which SSC is violated.

To get the SSC violation term below a certain threshold δ in HT, we need $\epsilon \log\left(\frac{1}{\epsilon}\right)$ to be on the order of $O\left(\frac{\delta}{n}\right)$. Similarly, in the Many-Server-HT scenario, it is required that $n^{1-\alpha} \log n$ is on the order of $O(\delta)$, or alternatively, n must be at least $\Omega\left(\exp\left(\frac{1}{\alpha-1}\log\frac{1}{\delta}\right)\right)$, or $n \sim \Omega\left(\delta^{-\frac{1}{\alpha-1}}\right)$ in magnitude. Satisfying such conditions becomes increasingly challenging as α approaches 1. This is also shown in Fig. 1 in Section 1.1. Furthermore, it is important to acknowledge that such conditions cannot be met for $\alpha \leq 1$. These observations provide an intuitive argument for the failure of the SSC property (as presented in this paper) when $\alpha \leq 1$. Essentially, for $\alpha \leq 1$, the notion of SSC deviates significantly from the one considered in this paper.

It is worth noting that although a large system size, represented by n, is required for the SSC error to be small, it remains bounded. Specifically, in the case of Many-Server-HT, it can be shown that the term $n\epsilon_n \log \frac{1}{\epsilon_n}$ is bounded by $\frac{\alpha}{e(\alpha-1)}$. Consequently, the SSC error is of order at most $O\left(\frac{1}{\alpha-1}\right)$. Therefore, it is reasonable to expect that even for moderate system sizes, the tail probability, $\mathbb{P}(\epsilon \overline{q} > x)$, exhibits exponential decay.

• **Pre-limit tail:** The pre-limit tail denotes the actual decay rate of the tail probability of $\epsilon_n \overline{q}$ under nonasymptotic HT condition, i.e., $n < \infty$. For the continuous-time JSQ system, we exactly characterize the pre-limit tail, which is given by $\theta_n = \frac{1}{\epsilon_n} \log \frac{1}{1-\epsilon_n}$. In super-slowdown regime, as ϵ_n goes to zero, the tail of $\epsilon_n \overline{q}$ matches that of an exponential distribution with mean 1, as $\lim_{n\to\infty} \theta_n = 1$. Further, note that, the deviation of the pre-limit tail from the corresponding HT value is given by $|\theta_n - 1|$, which is of order $O(\epsilon_n)$.

In the case of continuous-time JSQ, the arrivals and service times follow an exponential distribution, which is characterized by a single parameter, i.e., the arrival or service rate. The specific properties of the exponential distribution allow for a more accurate analysis of the tail probability. Intuitively, this distinction is the reason why we are able to precisely characterize the pre-limit tail for the JSQ system. In case of general arrival and service distributions, as presented in the case of the discrete-time SSQ, we do a second-order approximation to approximate the tail probability of the scaled queue length with its HT counterpart. Consequently, we obtain an upper bound on the pre-limit tail. More arguments are provided in Section 3.

• **Pre-exponent error:** In the context of the JSQ system, the pre-exponent error is represented by the expression 2ex. This error term arises from using Markov's Inequality to obtain tail-probability bounds using MGF. To clarify this error term, consider a random variable X that follows an exponential distribution with rate λ . In this case, the MGF of X is given by $\mathbb{E}[\exp(\theta X)] = \frac{1}{1-\theta/\lambda}$ for all $\theta < \lambda$. As shown in Lemma 7 of Appendix A, by Applying Markov's Inequality to the MGF and optimizing over the value of θ , we obtain

$$\mathbb{P}(X > x) \le e\lambda x e^{-\lambda x}$$

The upper bound differs from the actual tail of X by a multiplicative factor of $e\lambda x$, which arises from using Markov's Inequality. We acknowledge that it may be possible to eliminate the Markov-Inequality error by employing more complex techniques. However, we have chosen to rely solely on Markov's Inequality for our analysis to maintain simplicity.

2.3. State Space Collapse for JSQ system

Next, we mathematically specify SSC for the JSQ system with $\lambda_n = n\mu(1 - n^{-\alpha})$ and $\alpha > 1$.

Proposition 3. Consider the JSQ system as presented in Section 2.1. Suppose the system satisfies the condition $\lambda_n = n\mu(1 - \epsilon_n)$, and let $\kappa_{\perp} = 128$ and $\theta_{\perp} = 1/96$. Then, for $\epsilon_n \leq 1/2$, and $\theta \in (0, \theta_{\perp})$, we have that for all $i \in \{1, 2, ..., n\}$,

$$\mathbb{E}_{\pi_n} \Big[\sum_{i=1}^n e^{-\theta q_{\perp i}} + \sum_{i=1}^n e^{\theta q_{\perp i}} \Big] \le \kappa_{\perp} n, \qquad \qquad \mathbb{E}_{\pi_n} \Big[e^{\theta |q_{\perp i}|} \Big] \le \kappa_{\perp}, \qquad (4)$$

where $q_{\perp i} = q_i - \frac{1}{n} \sum_{j=1}^n q_j$.

As mentioned before, an essential component in establishing the limiting distribution (Theorem 1) and the tail probability (Theorem 2) is to characterize SSC accurately. For the JSQ system, when SSC is achieved, the queue-length vector \mathbf{q} collapses to a subspace where all its coordinates become identical, implying that the asymptotic behavior of the JSQ system closely resembles that of an SSQ.

In Proposition 3, we establish a specific notion of SSC by demonstrating that the MGF of the deviations of the individual queue lengths and their average, referred to as the "perpendicular component" and denoted by $\mathbf{q}_{\perp} = \mathbf{q} - \frac{1}{n} \bar{q} \mathbf{1}_n$, remains uniformly bounded within an interval around zero for all values of $\epsilon_n \leq 1/2$. In particular, this result implies that all the moments of the perpendicular component remain uniformly bounded, even in the limit as the system approaches HT conditions ($\epsilon_n \rightarrow 0$). In comparison, the elements of the queue length vector \mathbf{q} are on the order of $\frac{1}{\epsilon_n}$. Consequently, as ϵ_n becomes small, we observe that $\epsilon_n \mathbf{q} \approx \frac{\epsilon_n}{n} \bar{q} \mathbf{1}_n$ (or equivalently, $\epsilon_n \mathbf{q}_{\perp} \approx \mathbf{0}_n$).

Intuition behind the choice of the Lyapunov function. In order to establish Proposition 3, we employ the Lyapunov function $\sum_{i=1}^{n} e^{-\theta q_{\perp i}} + \sum_{i=1}^{n} e^{\theta q_{\perp i}}$, demonstrating its negative drift. It is worth noting that this Lyapunov function differs from the one considered in [2], namely $\exp(\theta \|\mathbf{q}_{\perp}\|_2)$, which yields the SSC result in the Many-Server-HT regime for $\alpha > 2$. However, $\exp(\theta \|\mathbf{q}_{\perp}\|_2)$ proves inadequate for establishing SSC in the case of $\alpha \in (1, 2]$. This limitation arises because, to show SSC in Many-Server-HT for $\alpha \in (1, 2]$, it is necessary to obtain a bound for the MGF of $\|\cdot\|_{\infty}$ -norm of the perpendicular component, denoted as $\|\mathbf{q}_{\perp}\|_{\infty}$. Intuitively, for an *n*-dim vector \mathbf{x} , if $\|\mathbf{x}\|_{\infty}$ is O(1), then it implies that $\|\mathbf{x}\|_2$ is $O(\sqrt{n})$. However, vice-versa need not hold. As such, a bound on ℓ_{∞} norm is sharper than a bound on ℓ_2 norm, and it plays a significant role while proving SSC for $\alpha \in (1, 2]$. Furthermore, using $\exp(\theta \|\mathbf{q}_{\perp}\|_{\infty})$ as the Lyapunov function proves challenging due to the non-smooth nature of the $\|\cdot\|_{\infty}$ -norm. Consequently, we use the Lyapunov function $\sum_{i=1}^{n} e^{-\theta q_{\perp i}} + \sum_{i=1}^{n} e^{\theta q_{\perp i}}$ as a suitable alternative. This choice represents a smooth approximation of $\exp(\theta \|\mathbf{q}_{\perp}\|_{\infty})$. Similar exponential function was also used in the context of graphical allocation of balls in bins in [54, 55].

2.4. Proof of Theorem 1 and Theorem 2

In this section, provide the proof sketches for Theorem 1 and Theorem 2. More details and precise mathematical arguments are provided in Appendix B.

Lemma 4. Consider the JSQ system as presented in Section 2.1. Suppose the system satisfies the condition $\lambda_n = n\mu(1 - \epsilon_n)$, where $\alpha > 1$. Let

$$\gamma_n(\theta) := n\mu - \lambda_n e^{\epsilon_n \theta} \qquad \qquad \beta_n(\mathbf{q}; \theta) := \mu \sum_{i=1}^n \mathbb{1}_{\{q_i=0\}} e^{\theta \epsilon_n \sum_{i=1}^n q_i}.$$

Then, for any $\theta < \theta_n := -\frac{1}{\epsilon_n} \log (1 - \epsilon_n)$, such that $\theta n \epsilon_n < \theta_{\perp}$, we have

$$\mathbb{E}_{\pi_n} \left[e^{\theta \epsilon_n \sum_{i=1}^n q_i} \right] = \frac{1}{\gamma_n(\theta)} \mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)].$$
(5)

Lemma 4 provides the MGF of the scaled total queue length $\epsilon_n \overline{q}$ and is an intermediate step to prove the results in Theorems 1 and 2. The proof of Lemma 4 is provided in Appendix B. Technical note. It is important to note that the condition $\theta n \epsilon_n < \theta_{\perp}$ in Lemma 4 holds true for all $\theta < 0$, and when ϵ_n is small for $\theta \in (0, \theta_n)$. Consequently, Eq. (5) is valid for all n and ϵ_n if $\theta < 0$, but if $\theta \in (0, \theta_n)$, it can only be used for ϵ_n small enough. Essentially, we need to use the SSC result given in Proposition 3 to obtain the bounds in Theorems 1 and 2, and these are valid only under the condition $\theta n \epsilon_n < \theta_{\perp}$.

Proof of Theorem 1. Recall that we use the notation $\lambda_n = n\mu(1 - \epsilon_n)$ and according to the theorem statement, $\epsilon_n = n^{-\alpha}$, where $\alpha > 1$. Also, $\overline{q} = \sum_{i=1}^n q_i$. In order to prove Theorem 1, we use Lemma 4. We first prove the following claim.

Claim 1. For $n \ge 2$, we have

$$\mathbb{E}_{\pi_n}\left[\mathbb{1}_{\{q_i=0\}}\right] = \epsilon_n, \quad \forall i \in \{1, 2, \dots, n\}.$$
(6)

And, for any $\theta < \theta_n$ and ϵ_n small enough such that $2|\theta|n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \theta_{\perp}$, we have

$$\sum_{i=1}^{n} \mathbb{E}_{\pi_n} \Big[\mathbb{1}_{\{q_i=0\}} \Big| e^{\theta \epsilon_n \overline{q}} - 1 \Big| \Big] \le \frac{2e\kappa_{\perp}^2}{\theta_{\perp}} |\theta| n^2 \epsilon_n^2 \log\left(\frac{1}{\epsilon_n}\right),$$

where θ_{\perp} and κ_{\perp} are as defined in Proposition 3.

Proof of Claim 1 is provide in Appendix B. Now, according to the assumption, $\epsilon_n = n^{-\alpha}$, where $\alpha > 1$. Under the $\alpha > 1$ condition, the condition $2|\theta|n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \theta_{\perp}$ can be satisfied for n large enough, i.e., we need n large enough such that $2\alpha|\theta|n^{1-\alpha}\log n < \theta_{\perp}$, or $\frac{\log n}{n^{\alpha-1}} < \frac{\theta_{\perp}}{2\alpha|\theta|}$. From Claim 1, we have that for any $\theta < 1$, (as $\theta_n \ge 1$), we have,

$$\lim_{n \to \infty} \frac{1}{n\epsilon_n} \sum_{i=1}^n \mathbb{E}_{\pi_n} \left[\mathbb{1}_{\{q_i=0\}} \right] = 1$$

and

$$\lim_{n \to \infty} \frac{1}{n\mu\epsilon_n} \Big| \mathbb{E}_{\pi_n} [\beta_n(\mathbf{q}; \theta)] - \mu \sum_{i=1}^n \mathbb{E}_{\pi_n} [\mathbbm{1}_{\{q_i=0\}}] \Big| \le \lim_{n \to \infty} \frac{1}{n\mu\epsilon_n} \sum_{i=1}^n \mathbb{E}_{\pi_n} \Big[\mathbbm{1}_{\{q_i=0\}} \Big| e^{\theta\epsilon_n \overline{q}} - 1 \Big| \Big] = 0.$$

Thus, we have $\lim_{n\to\infty} \frac{1}{n\mu\epsilon_n} \mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)] = 1$. Further, for any θ , we have

$$\lim_{n \to \infty} \frac{1}{n\mu\epsilon_n} \gamma_n(\theta) = \frac{1}{n\mu\epsilon_n} \left(n\mu - \lambda_n e^{\epsilon_n \theta} \right) = \frac{1}{\epsilon_n} \left(1 - e^{\epsilon_n \theta} + \epsilon_n e^{\epsilon_n \theta} \right) = 1 - \theta.$$

Thus, by using Lemma 4, for any $\theta < 1$,

$$\lim_{n \to \infty} \mathbb{E}[e^{\theta \epsilon_n \overline{q}}] = \lim_{n \to \infty} \frac{1}{\gamma_n(\theta)} \mathbb{E}_{\pi_n}[\beta_n(\mathbf{q}; \theta)] = \frac{1}{1 - \theta}$$

This shows that the Moment Generating Function (MGF) of $\epsilon_n \overline{q}$ converges to that of an exponential random variable with mean 1. Now, by using [56, Theorem 25.10], we have that $\epsilon_n \overline{q} \stackrel{d}{\to} \Upsilon$, where Υ is an exponential random variable with mean 1.

Next, by using Proposition 3, for $\theta \in (0, \theta_{\perp})$ we have that

$$1 \leq \lim_{n \to \infty} \mathbb{E}_{\pi_{\epsilon}} \left[e^{n\epsilon_n \theta |q_{\perp_i}|} \right] \leq \lim_{n \to \infty} \left(\mathbb{E}_{\pi_{\epsilon}} \left[e^{\theta |q_{\perp_i}|} \right] \right)^{n\epsilon_n} \leq \lim_{n \to \infty} \kappa_{\perp}^{n\epsilon_n} = 1,$$

where the second inequality follows by using Jensen's Inequality. This implies that, for all $i \in \{1, 2, ..., n\}$, we have $n\epsilon_n q_{\perp_i} \stackrel{d}{\to} 0$. Now, by definition of \mathbf{q}_{\perp} , we have, $n\epsilon_n \mathbf{q} = \epsilon_n \overline{q} \mathbf{1} + n\epsilon_n \mathbf{q}_{\perp}$, where $\epsilon_n \overline{q} \stackrel{d}{\to} \Upsilon$ and $n\epsilon_n q_{\perp_i} \stackrel{d}{\to} 0$ as $n \to \infty$. Thus, we have $n\epsilon_n \mathbf{q} \stackrel{d}{\to} \Upsilon \mathbf{1}$. This completes the proof. Proof of Theorem 2. Note that for $\epsilon_n \leq 1/2$, we have $\theta_n \leq 2$. Thus, the condition $2|\theta|n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \theta_{\perp}$ is satisfied for any $\theta \in (0, \theta_n)$ whenever ϵ_n is small enough such that $4n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \theta_{\perp}$. Then, from Claim 1, for any $\theta \in (0, \theta_n)$, we have

$$\mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)] \le \mu n \epsilon_n (1 - \kappa_2 n \epsilon_n \log \epsilon_n),$$

where, as $\theta \leq \theta_n \leq 2$, we can take $\kappa_2 := \frac{4e\kappa_{\perp}^2}{\theta_{\perp}}$. Now, by using Markov's inequality, we have that for all $\theta \in (0, \theta_n)$,

$$\mathbb{P}(\epsilon_n \overline{q} > x) \leq e^{-\theta x} \mathbb{E}[e^{\theta \epsilon_n \sum_{i=1}^n q_i}] \\
\leq \mu n \epsilon_n (1 - \kappa_2 n \epsilon_n \log \epsilon_n) \times \frac{1}{\gamma_n(\theta)} e^{-\theta x} \\
= \mu n \epsilon_n (1 - \kappa_2 n \epsilon_n \log \epsilon_n) \times \frac{1}{n\mu - \lambda_n e^{\epsilon_n \theta}} e^{-\theta x}.$$
(7)

Next, we optimize the upper bound over the values of $\theta \in (0, \theta_n)$. By differentiation,

$$\frac{d}{d\theta} \left(\frac{1}{\gamma_n(\theta)} e^{-\theta x} \right) = \frac{1}{\gamma_n^2(\theta)} \left(-x\gamma_n(\theta) + \lambda_n \epsilon_n e^{\epsilon_n \theta} \right) e^{-\theta x}$$

Then, the derivative is equal to zero at

$$\theta = \theta_{x,n} := \frac{1}{\epsilon_n} \log \left(\frac{n\mu x}{\lambda_n (x + \epsilon_n)} \right) = \frac{1}{\epsilon_n} \log \left(\frac{x}{(1 - \epsilon_n)(x + \epsilon_n)} \right) < \theta_n,$$

where the last inequality is valid for any x > 0. Also, to ensure that $\theta_{x,n} > 0$, we need that

$$\frac{x}{(1-\epsilon_n)(x+\epsilon_n)} > 1 \implies x > 1-\epsilon_n.$$

Then, by substituting $\theta = \theta_{x,n}$, we have

$$\frac{1}{n\mu - \lambda_n e^{\epsilon_n \theta_{x,n} x}} e^{-\theta_{x,n} x} = \frac{x + \epsilon_n}{\mu n \epsilon_n} \exp\left(-\frac{x}{\epsilon_n} \log\left(\frac{x}{(1 - \epsilon_n)(x + \epsilon_n)}\right)\right)$$
$$= \frac{x + \epsilon_n}{\mu n \epsilon_n} \exp\left(-\frac{x}{\epsilon_n} \log\frac{1}{1 - \epsilon_n} + \frac{x}{\epsilon_n} \log\frac{x + \epsilon_n}{x}\right)$$
$$\leq \frac{e(x + \epsilon_n)}{\mu n \epsilon_n} \exp\left(-\frac{x}{\epsilon_n} \log\frac{1}{1 - \epsilon_n}\right).$$

Using the above result in Eq. (7) and $x \ge \epsilon_n$, we have

$$\mathbb{P}(\epsilon_n \sum_{i=1}^n q_i > x) \le 2ex(1 - \kappa_2 n\epsilon_n \log \epsilon_n) \exp\left(-\frac{x}{\epsilon_n} \log \frac{1}{1 - \epsilon_n}\right).$$

This completes the first part of the proof. For the second, i.e., the lower bound in Eq. (2), we couple the JSQ system with a Single Server Queue (SSQ) as follows, i.e., we consider a SSQ process $\{q_{ssq}(t)\}_{t\geq 0}$, where all the servers of the original JSQ system are pooled together to create a single server with service rate $n\mu$. Thus, $\{q_{ssq}(t)\}_{t\geq 0}$ is an M/M/1 queue with arrival rate λ_n and service rate $n\mu$. Then, the stationary distribution of $\{q_{ssq}(t)\}_{t\geq 0}$ is given by,

$$\mathbb{P}(q_{ssq} = i) = \left(1 - \frac{\lambda_n}{n\mu}\right) \left(\frac{\lambda_n}{n\mu}\right)^i = \epsilon_n (1 - \epsilon_n)^i$$

This gives us

$$\mathbb{P}\Big(\epsilon_n \sum_{i=1}^n q_i > x\Big) \ge \mathbb{P}(\epsilon_n q_{ssq} > x) = (1 - \epsilon_n)^{\left\lceil \frac{x}{\epsilon_n} \right\rceil} \ge (1 - \epsilon_n)^{\frac{x}{\epsilon_n} - 1} = \frac{1}{1 - \epsilon_n} \exp\Big(-\frac{x}{\epsilon_n} \log\frac{1}{1 - \epsilon_n}\Big).$$

This completes the proof.

3. Discrete time Single server queue

In this section, we present the bound on the tail probability for a SSQ. The aim is to show the affect of a general arrival and service distribution on the decay rate of the tail probability $\mathbb{P}(\epsilon q > x)$. The model is as follows.

3.1. Model for Single Server Queue

We consider a sequence of discrete-time SSQ indexed by the HT parameter ϵ , where q(t) denotes the number of customers in the system (queue length) at the beginning of time slot t. Customers arrive to the system as an i.i.d. process $\{a(t)\}_{t\geq 0}$, with $\mathbb{E}[a(t)] = \lambda_{\epsilon}$, $\operatorname{Var}(a(t)) = \sigma_{\epsilon,a}^2$, and $a(t) \leq A$ almost surely. Once the customers join the queue, the server decides to serve up to s(t) jobs waiting in the queues, with $\mathbb{E}[s(t)] = \mu_{\epsilon}$, $\operatorname{Var}(s(t)) = \sigma_{\epsilon,s}^2$, and $s(t) \leq A$ almost surely. Even though the server can process s(t) customers, it is possible that there are not enough customers in the system, and some of the service gets wasted. Hence, we call s(t) as the 'potential' service. Similar to the arrival process, the potential services are also independent and identically distributed across time slots. Moreover, we assume that these potential services are independent of the queue length vector. The queue evolution process is given by

$$q(t+1) = [q(t) + a(t) - s(t)]^{+} = q(t) + a(t) - s(t) + u(t),$$
(8)

where $[x]^+ := \max\{x, 0\}$ is used because the queue length cannot be negative. The term u(t) is the unused service and represents the difference between the potential and actual service. By definition, the unused service u(t) is positive only if q(t+1) = 0, which implies q(t+1)u(t) = 0 for all t. Also, the unused service cannot be higher than the potential, and so, $0 \le u(t) \le s(t) \le A$ almost surely.

It is well known that $\lambda_{\epsilon} < \mu_{\epsilon}$ implies stability, that is, the Markov chain $\{q(t)\}_{t\geq 0}$ is positive recurrent. As a result, the system reaches steady state. We use π_{ϵ} to denote the steady state distribution of the queue length process $\{q(t)\}_{t\geq 0}$. We drop the symbol t to denote that variables in steady state, i.e., q follows the steady state distribution π_{ϵ} , and q^+ denote the state that comes after q, i.e., $q^+ = [q+a-s]^+ = q+a-s+u$, where a and s follow the same distribution as a(t) and s(t), respectively.

3.2. Results for SSQ

In this section, we establish an upper bound on the tail of the queue length distribution for a system in non-asymptotic HT. We consider $\lambda_{\epsilon} = \mu_{\epsilon}(1 - \epsilon)$, so that as $\epsilon \to 0$, the system approaches HT conditions. In Theorem 5, we provide a bound on the probability $\mathbb{P}(\epsilon q > x)$ when the HT parameter is bounded away from zero.

Theorem 5. Consider the SSQ as defined in Section 3.1, with $\lambda_{\epsilon} = \mu_{\epsilon}(1-\epsilon)$. Let $\sigma_{\epsilon}^2 := \sigma_{\epsilon,a}^2 + \sigma_{\epsilon,s}^2$,

$$\theta_{\epsilon} := \frac{2\mu_{\epsilon}}{\sigma_{\epsilon}^2}, \qquad \qquad \kappa_1 := \frac{2\mu_{\epsilon}E_3}{3\sigma_{\epsilon}^4} + \epsilon \frac{9\mu_{\epsilon}^2 A^4}{\sigma_{\epsilon}^6},$$

where $E_3 := \max\{0, \mathbb{E}[(a-s)^3]\}$. Let $\mathbb{E}[e^{\epsilon\theta q(0)}] < \infty$ for all $\theta \le \theta_{\epsilon}$. Then, for all $x > \frac{\theta_{\epsilon}}{(1+\kappa_1\epsilon)}$, we have

$$\mathbb{P}(\epsilon q > x) \le e\theta_{\epsilon} x e^{-\theta_{\epsilon}(1-\kappa_1\epsilon)x}, \quad \forall \epsilon \in (0,1).$$
(9)

Further, as $x \to \infty$, we have

$$\lim_{x \to \infty} \frac{1}{x} \log \mathbb{P}(\epsilon q > x) \le -\theta_{\epsilon} (1 - \kappa_1 \epsilon).$$
(10)

The exponential decay of the tail of the queue-length distribution of an SSQ has been established by the tail bound presented in Eq. (9). This result is consistent with the well-known fact that the distribution of the scaled steady-state queue length ϵq converges to an exponential distribution as ϵ tends to zero. However, our result provides a more detailed insight by characterizing the rate of convergence of the tail probability to its corresponding HT value.

Note that, SSQ is a one-dimensional system, and so, there is no concept of SSC in SSQ. Further, in case of SSQ, the pre-exponent error is given by $e\theta_{\epsilon}x$. The reasoning for this is same as that in the case of JSQ system. Therefore, we only discuss the pre-limit tail for the discrete time SSQ.

Pre-limit tail: As mentioned before, pre-limit tail refers to the actual decay rate of the tail probability for the pre-limit system. Under general arrival and service distribution, as in the case of discrete time SSQ, we provide an upper bound on the pre-limit tail, which given by $\theta_{\epsilon}(1 - \kappa_1 \epsilon)$. The steady-state distribution of the pre-limit system need not follow an exponential distribution. As such, to obtain an exponential decay rate on the tail probability, we employ a second order approximation to approximate the distribution with an exponential. It is important to note that as ϵ approaches zero, the upper bound on the pre-limit tail converges to the correct HT value, i.e. $\lim_{\epsilon \to 0} \theta_{\epsilon}(1 - \kappa_1 \epsilon) = \frac{2\mu}{\sigma^2}$, where μ and σ are the limiting values of μ_{ϵ} and σ_{ϵ} as $\epsilon \to 0$.

Note that the constant κ_1 in the pre-limit tail depends on the third moment of (a - s), while the actual decay rate of the tail probability in HT depends on the corresponding first and the second moment. Using exponential Lyapunov functions, the HT distribution of a queueing system is generally obtained using a second-order approximation of the MGF of (a - s), which involves the first two moments. As such, the 'strength' of this second-order approximation depends on the third moment of (a - s), which is essentially captured by the dependency of the constant κ_1 on E_3 in Theorem 5.

3.3. Proof of Theorem 5

The first step to prove Theorem 5 is characterizing the steady-state distribution of ϵq in terms of its MGF. For this goal, we consider the Lyapunov function $V(x;\theta) := e^{\theta \epsilon x}$ and obtain the following result.

Lemma 6. Consider an SSQ as defined in Section 3.1. Suppose the initial state q(0) satisfies $\mathbb{E}[e^{\theta \epsilon q(0)}] < \infty$ for all $\theta < \theta_0$ and for all $\epsilon \in (0, 1)$. Then, for any $\theta < \theta_0$, we have

$$\mathbb{E}[V(q(t+1);\theta)] = (1 - \gamma_{\epsilon}(\theta))^{t+1} \mathbb{E}[V(q(0);\theta)] + \sum_{i=0}^{\iota} (1 - \gamma_{\epsilon}(\theta))^{i} \mathbb{E}[\beta_{\epsilon}(t-i;\theta)],$$
(11)

where $\gamma_{\epsilon}(\theta) := 1 - \mathbb{E}\left[e^{\epsilon\theta(a-s)}\right]$ and $\beta_{\epsilon}(t;\theta) := 1 - \mathbb{E}\left[e^{-\theta\epsilon u(t)} | q(t)\right]$. Let $\Theta_{\epsilon} := \{\theta \in \mathbb{R} : \theta < \theta_0, \ \mathbb{E}[e^{\theta\epsilon(a-s)}] < 1\}$. Then, for all $\theta \in \Theta_{\epsilon} \cup \{\theta : \theta \le 0\}$, we have

$$\mathbb{E}_{\pi_{\epsilon}}[V(q;\theta)] := \mathbb{E}_{\pi_{\epsilon}}\left[e^{\theta\epsilon q}\right] = \frac{1 - \mathbb{E}_{\pi_{\epsilon}}\left[e^{-\theta\epsilon u}\right]}{1 - \mathbb{E}\left[e^{\epsilon\theta(a-s)}\right]}.$$
(12)

The expression in Eq. (12) was first introduced in [6] for θ in an interval around zero. In this work, we have expanded the range of validity for Eq. (12) to a much larger interval given by Θ_{ϵ} . Although it may be difficult to fully characterize Θ_{ϵ} for a fixed $\epsilon > 0$, we can use Taylor's approximation of the exponential function to obtain a close enough subset. Accurately characterizing Θ_{ϵ} is crucial to obtaining the correct tail bounds. Our finite-time characterization of the queue-length distribution, as shown in Eq. (11), allows us to achieve this expanded range of validity.

The proof of Lemma 6 follows by showing that, for any $\theta \in \Theta_{\epsilon}$, the right-hand side (RHS) in Eq. (11) converges to a finite value. The details of the proof of Lemma 6 are presented in Appendix C.

Lemma 7. Suppose X is a non-negative random variable, and the MGF of X satisfies the inequality $\mathbb{E}[e^{\theta X}] \leq \frac{1}{1-\theta/\lambda}, \forall \theta \in (0,\lambda)$, where $\lambda > 0$. Then, for any $x > \frac{1}{\lambda}$, we have

$$\mathbb{P}(X > x) \le e\lambda x e^{-\lambda x}.$$

The proof of Lemma 7 is provided in Appendix A.

Proof of Theorem 5. By equating the drift of the queue length q(t) to zero in steady state, we prove the following claim.

Claim 2. For any $\theta \in \mathbb{R}$, $1 - \mathbb{E}_{\pi_{\epsilon}}[e^{-\theta \epsilon u}] \leq \epsilon^2 \theta \mu_{\epsilon}$.

Next, using Taylor's expansion of the exponential function we obtain the following claim.

Claim 3. For any $\theta \in \left(0, \frac{\theta_{\epsilon}}{(1+\kappa_1\epsilon)}\right), 1-\mathbb{E}\left[e^{\epsilon\theta(a-s)}\right] \ge \theta\epsilon^2\mu_{\epsilon}\left(1-\frac{\theta}{\theta_{\epsilon}}\left(1+\kappa_1\epsilon\right)\right).$

The proofs of Claims 2 and 3 are provided in Appendix C. Next, we use the Lemma 6. Note that, by the assumption $\mathbb{E}[e^{\theta \epsilon q(0)}] < \infty$ for $\theta < \theta_{\epsilon}$, we have that $\left(0, \frac{\theta_{\epsilon}}{(1+\kappa_1\epsilon)}\right) \subseteq \Theta_{\epsilon}$. Then, by Lemma 6 and the above mentioned claims, we have

$$\mathbb{E}\left[e^{\epsilon\theta q}\right] \leq \left(1 - \frac{\theta}{\theta_{\epsilon}} \left(1 + \kappa_{1}\epsilon\right)\right)^{-1}, \quad \forall \theta \in \left(0, \frac{\theta_{\epsilon}}{(1 + \kappa_{1}\epsilon)}\right).$$

Afterwards, the result in Theorem 5 follows simply by using Markov's inequality as shown in Lemma 7. \Box

4. M/M/n system

4.1. Model

An M/M/n system (also known as Erlang-C) is a multi-server queue with n servers, a single queue, and follows a continuous-time first-come-first-serve service discipline. Customer arrivals follow a Poisson process with rate λ_n , and each server has an exponentially distributed service time with constant service rate μ . The rate μ is independent of the system size; thus, the subscript n is omitted.

To represent the system's dynamics, we use $q_n(t)$ to denote the number of customers in the system at time t. We use $w_n(t) = [q_n(t) - n]^+$ to denote the number of customers waiting in the queue, and $r_n(t) = [n - q_n(t)]^+$ for the number of idle servers at time t.

The queue-length process $\{q_n(t)\}_{t\geq 0}$ is a CTMC, and it is well known that it is stable (positive recurrent) if $\lambda_n < n\mu$. To be consequent with the previous sections, we assume that $\lambda_n = n\mu(1 - \epsilon_n)$. Further, a stationary distribution π_n exists under this stability condition. Similarly to previous sections, we drop the index t to denote steady-state variables.

Similarly to the JSQ system, to study the Many-Server-HT asymptotics of this system, we define the load $\rho_n := \frac{\lambda_n}{n\mu} = 1 - \epsilon_n$ and consider $\epsilon_n = cn^{-\alpha}$, where c > 0 is a constant, and $\alpha > 0$ a parameter (as in the JSQ system). In this case, we are interested in $\alpha \in (0, 1]$.

As explained in the introduction, the steady-state dynamics of the system vary with the value of α . We study three regimes: (i) Sub-HW, where $\alpha \in \left(0, \frac{1}{2}\right)$, (ii) HW, where $\alpha = \frac{1}{2}$, and (iii) Super-HW where

$$\alpha \in \left(\frac{1}{2}, \infty\right).$$

4.2. Results on w_n and r_n

In this section, we provide separate tail bounds on the number of idle servers and the number of waiting customers because the scaling parameter is different. In Theorem 8 we present our results for the number of idle servers r_n .

Theorem 8. Consider the M/M/n queue as given in Section 4.1 with $\lambda_n = n\mu(1-\epsilon_n)$. Let $\tilde{r}_n := r_n - n\epsilon_n$, and $\eta_n := \frac{1}{\sqrt{n(1-\epsilon_n)}}$.

(a) Super-HW regime: Suppose $\epsilon_n = cn^{-\alpha}$ with $\alpha > \frac{1}{2}$. Then, if $n^{2\alpha-1} > 4c^2$, we have

$$\mathbb{P}(r_n > 0) \le 4e\pi c n^{-\alpha + \frac{1}{2}}, \qquad \qquad \lim_{n \to \infty} \mathbb{P}(r_n > 0) = 0, \qquad (13)$$

and there exist a constant κ_1 , independent of n, such that,

$$\mathbb{P}(\eta_n \tilde{r}_n > x) \le \frac{\kappa_1 c}{n^{\alpha - \frac{1}{2}}} e^{-\frac{1}{2}x^2}.$$

(b) HW regime: Suppose $\epsilon_n = cn^{-\frac{1}{2}}$. Then,

$$\lim_{n \to \infty} \mathbb{P}(r_n > 0) = \frac{\sqrt{2\pi}c \exp\left(\frac{c^2}{2}\right) \Phi(c)}{1 + \sqrt{2\pi}c \exp\left(\frac{c^2}{2}\right) \Phi(c)},\tag{14}$$

and there exist constant κ_2 , such that

$$\mathbb{P}(\eta_n \tilde{r}_n > x) \le \kappa_2 e^{-\frac{1}{2}x^2}$$

(c) Sub-HW regime: Suppose $\epsilon_n = cn^{-\alpha}$ with $\alpha \in (0, \frac{1}{2})$. Then, there exists constant κ_3 such that

$$\mathbb{P}(r_n > 0) \ge 1 - \frac{\kappa_3}{c} n^{\alpha - \frac{1}{2}} e^{-cn^{\frac{1}{2} - \alpha}}, \qquad \qquad \lim_{n \to \infty} \mathbb{P}(r_n > 0) = 1,$$

and we have

$$\mathbb{P}(\eta_n \tilde{r}_n > x) \le e^{-\frac{1}{2}x^2}, \qquad \qquad \mathbb{P}(\eta_n \tilde{r}_n < -x) \le e^{-\frac{1}{2}x^2 + 2e\epsilon_n x^2}$$

Theorem 8 provides the tail bounds on the number of idle servers in the M/M/n queue in each of the three regimes considered in the paper.

Remark 2. Note that we do not provide a LD result for r_n , as we did in the case of SSQ and JSQ systems, as the number of idle servers r_n is always bounded by n. As such, for the pre-limit system (i.e., when the system size n is finite), we have that

$$\mathbb{P}(\eta_n \tilde{r}_n > x) = 0, \ \forall x > \sqrt{n\rho_n}.$$

Next, we provide the results for the number of waiting customers w_n .

Corollary 9. Consider the M/M/n queue as given in Section 4.1 with $\lambda_n = n\mu(1 - \epsilon_n)$. Then, for any $\{\epsilon_n\}_{n\geq 0}$ such that $\epsilon_n \to 0$ as $n \to \infty$, and for any $\theta < 1$, we have

$$\lim_{n \to \infty} \mathbb{E}\left[e^{\theta \epsilon_n w_n} \middle| w_n > 0\right] = \frac{1}{1 - \theta}, \qquad [\epsilon_n w_n \middle| w_n > 0] \stackrel{d}{\to} \Upsilon,$$

where Υ is an exponential random variable with mean 1. Further, for any $x > \epsilon_n$, we have

$$\frac{1}{(1-\epsilon_n)^2}e^{-\theta_n x} \le \mathbb{P}(\epsilon_n w_n \ge x | w_n > 0) \le \frac{1}{1-\epsilon_n}e^{-\theta_n x},$$

where $\theta_n = \frac{1}{\epsilon_n} \log \frac{1}{1-\epsilon_n}$. As a consequence, we get the LD result

$$\lim_{x \to \infty} \frac{1}{x} \mathbb{P}(\epsilon_n w_n \ge x) = \lim_{x \to \infty} \frac{1}{x} \mathbb{P}(\epsilon_n w_n \ge x | w_n > 0) = \theta_n.$$
(15)

Corollary 9 provides the limiting distribution of the steady state number of waiting customers in the system. For any n and ϵ_n , the distribution of $[w_n|w_n > 0]$ can be exactly characterized. As will be depicted in Lemma 10, it turns out that $[w_n|w_n > 0]$ follows a geometric distribution with parameter ϵ_n . Such a result can also be derived by simply solving for the steady-state distribution of an M/M/n queue. Furthermore, as we can exactly characterize the distribution of $[w_n|w_n > 0]$, there is no need for using Markov's inequality to get a tail bound on the number of waiting customers. Hence, we do not obtain the pre-limit nor the pre-exponent error in the case of the number of waiting customers.

Remark 3. It is worth noting that although the conditional distribution $[w_n|w_n > 0]$ follows a geometric distribution, it is possible for the probability $\mathbb{P}(w_n > 0)$ to approach zero. This can be observed by examining the bounds on $\mathbb{P}(r_n > 0)$ provided in Theorem 8. Specifically, in the Sub-HW regime, we have

$$\mathbb{P}(w_n > 0) \le 1 - \mathbb{P}(r_n > 0) \le \frac{\kappa_3}{c} n^{\alpha - \frac{1}{2}} e^{-cn^{\frac{1}{2} - \alpha}}.$$

It is well-known that $\mathbb{P}(w_n > 0)$ decreases to zero in this regime. However, we additionally characterize the rate at which this convergence occurs.

Remark 4. The results presented in Theorem 8 and Corollary 9 only consider Many-Server-HT regimes. There, the HT parameter ϵ_n approaches zero as the system size n grows. One can also consider a conventional HT regime, where the system size n is constant and the HT parameter $\epsilon \to 0$ (using the notation $\lambda_n = n\mu(1-\epsilon)$) independent of n. In this case, the tail bound for the relevant quantities closely aligns with the tail-bound results obtained in the Super-HW regime. By employing similar calculations as, it can be shown that the number of waiting customers exhibits the same tail bounds as presented in Corollary 9 after replacing ϵ_n by ϵ . Furthermore, the probability $\mathbb{P}(r_n > 0)$ is on the order of $O(\epsilon)$. Moreover, in conventional HT, there is no need for a tail bound on r_n as $r_n \leq n$ and n is a constant.

4.2.1. Steady-state distribution of w_n and r_n

In this section, we provide the complete characterization of the steady state distribution of $\{q_n(t)\}_{t\geq 0}$. Specifically, we provide the MGF of the steady-state number of waiting customers (w_n) and idle servers (r_n) .

Lemma 10. Consider the M/M/n queue as given in Section 4.1 and suppose $\lambda_n = n\mu(1 - \epsilon_n)$. Define

$$G_n(t) := \exp\left(-n\epsilon_n t - n(1-\epsilon_n)(e^{-t} + t - 1)\right).$$

Then, we have the following results.

(a) For any $\theta \in \mathbb{R}$, we have

$$\mathbb{E}\left[e^{\theta r_n}\Big|r_n>0\right] = G_n^{-1}(\theta)\left(\int_{-\infty}^0 G_n dt\right)^{-1}\int_{-\infty}^\theta G_n dt.$$

(b) For any $\theta < \log \frac{1}{1-\epsilon_n}$, we have

$$\mathbb{E}\left[e^{\theta w_n} \middle| w_n > 0\right] = \frac{1}{1 - \frac{1}{\epsilon_n} \left(1 - e^{-\theta}\right)}$$

Further, we have $\mathbb{P}(q_n = n) = \left(\frac{1}{\epsilon_n} + n \int_{-\infty}^0 G_n(t) dt\right)^{-1}$, and

$$\mathbb{P}(w_n > 0) = \frac{1 - \epsilon_n}{\epsilon_n} \mathbb{P}(q_n = n), \qquad \qquad \mathbb{P}(r_n > 0) = n \mathbb{P}(q_n = n) \int_{-\infty}^0 G_n(t) dt.$$

Lemma 10 provides the steady state distribution of r_n and w_n for any value of n and ϵ_n . The results in Theorem 8 are derived by using the result in Lemma 10.a, Markov's inequality. Lemma 10.a indicates that the distribution of $[r_n|r_n > 0]$ closely resembles that of a truncated normal random variable. By replacing $G_n(t)$ with $\tilde{G}_n(t) := \exp\left(-n\eta_n\epsilon_n t - \frac{1}{2}t^2\right)$, we have that the function

$$\tilde{G}_n^{-1}(\theta) \left(\int_{-\infty}^0 \tilde{G}_n dt \right)^{-1} \int_{-\infty}^\theta \tilde{G}_n dt$$

defines the MGF of a truncated normal random variable, specifically $[Y_n|Y_n > 0]$, where Y_n follows a normal distribution with mean $n\eta_n\epsilon_n$ and variance 1.

Intuitively, for large values of n, we can approximate $G_n(\eta_n t)$ by $\tilde{G}_n(t)$ because $\frac{1}{\eta_n^2} \left(e^{-\eta_n t} + \eta_n t - 1\right) \approx \frac{t^2}{2}$, where $\eta_n = \frac{1}{\sqrt{n\rho_n}}$. Consequently, we observe that, for large n, the distribution of $[\eta_n r_n | r_n > 0]$ closely matches the distribution of $[Y_n | Y_n > 0]$, where Y_n is defined above. Consequently, for large values of n, $[\eta_n \tilde{r}_n | r_n > 0]$ closely matches the distribution of a truncated standard normal random variable. In the proof of Theorem 8, use this idea to establish tail bounds on r_n in each of the three regimes.

The result in Corollary 9 immediately follows from Lemma 10.b just be observing that the MGF of $[w_n|w_n > 0]$ matches with that of a geometric random variable with parameter ϵ_n . The mathematical details for results provided in this section (Section 4.2) is provided in Appendix D.

References

- [1] V. Gupta, N. Walton, Load balancing in the Nondegenerate Slowdown Regime, Operations Research 67 (1) (2019) 281–294.
- D. Hurtado-Lange, S. T. Maguluri, A load balancing system in the many-server heavy-traffic asymptotics, Queueing Systems 101 (3-4) (2022) 353–391.
- [3] B. Hajek, Hitting-time and occupation-time bounds implied by drift analysis with applications, Advances in Applied Probability (1982) 502-525.
- [4] A. Eryilmaz, R. Srikant, Asymptotically tight steady-state queue length bounds implied by drift conditions, Queueing Systems 72 (3-4) (2012) 311–359.
- [5] S. T. Maguluri, R. Srikant, Heavy traffic queue length behavior in a switch under the MaxWeight algorithm, Stochastic Systems 6 (1) (2016) 211–250.
 - URL http://dx.doi.org/10.1214/15-SSY193
- [6] D. Hurtado-Lange, S. T. Maguluri, Transform methods for heavy-traffic analysis, Stochastic Systems 10 (4) (2020) 275–309.
 [7] A. Braverman, J. Dai, J. Feng, Stein's method for steady-state diffusion approximations: An introduction through the Enlang A and Enlang C models. Stochastic Systems 6, (2) (2017) 201–266.
- Erlang-A and Erlang-C models, Stochastic Systems 6 (2) (2017) 301–366.
 [8] J. Kingman, On queues in heavy traffic, Journal of the Royal Statistical Society. Series B (Methodological) (1962) 383–392.
- [9] J. M. Harrison, Heavy traffic analysis of a system with parallel servers: asymptotic optimality of discrete-review policies, Annals of applied probability (1998) 822–848.
- [10] R. Williams, On dynamic scheduling of a parallel server system with complete resource pooling, Fields Institute Communications 28 (49-71) (2000) 5–1.
- [11] J. M. Harrison, A. Zeevi, Dynamic scheduling of a multiclass queue in the halfin-whitt heavy traffic regime, Operations Research 52 (2) (2004) 243-257.
- [12] D. Gamarnik, A. Zeevi, Validity of heavy traffic steady-state approximations in Generalized Jackson Networks, The Annals of Applied Probability (2006) 56–90.
- [13] A. Braverman, J. Dai, Stein's method for steady-state diffusion approximations of M/Ph/n+M systems, The Annals of Applied Probability 27 (1) (Feb. 2017). doi:10.1214/16-AAP1211. URL https://doi.org/10.1214/16-AAP1211
- [14] A. Braverman, J. Dai, Stein's method for steady-state diffusion approximations of M/Ph/n+ M systems, The Annals of Applied Probability 27 (1) (2017) 550–581.
- [15] L. Ying, Stein's method for mean field approximations in light and heavy traffic regimes, Proc. ACM Meas. Anal. Comput. Syst. 1 (1) (2017) 12:1-12:27. doi:10.1145/3084449. URL http://doi.acm.org/10.1145/3084449
- [16] R. Gaunt, N. Walton, Stein's method for the single server queue in heavy traffic, Statistics & Probability Letters 156 (2020) 108566.
- [17] A. Braverman, J. Dai, M. Miyazawa, Heavy traffic approximation for the stationary distribution of a Generalized Jackson Network: The BAR approach, Stochastic Systems 7 (1) (2017) 143–196.
- [18] A. Braverman, J. Dai, M. Miyazawa, The bar-approach for multiclass queueing networks with sbp service policies, arXiv preprint arXiv:2302.05791 (2023).
- [19] J. Dai, P. Glynn, Y. Xu, Asymptotic product-form steady-state for generalized jackson networks in multi-scale heavy traffic, arXiv preprint arXiv:2304.01499 (2023).
- [20] P. R. Jhunjhunwala, M. Zubeldia, S. T. Maguluri, Join-the-shortest queue with abandonment: Critically loaded and heavily overloaded regimes, arXiv preprint arXiv:2211.15050 (2022).
- [21] S. Vallender, Calculation of the wasserstein distance between probability distributions on the line, Theory of Probability & Its Applications 18 (4) (1974) 784–786.
- [22] D. Mukherjee, S. C. Borst, J. S. Van Leeuwaarden, P. A. Whiting, Universality of power-of-d load balancing in many-server systems, Stochastic Systems 8 (4) (2018) 265–292.
- [23] M. Mitzenmacher, Load balancing and density dependent jump Markov processes, in: focs, IEEE, 1996, p. 213.
- [24] M. Mitzenmacher, The power of two choices in randomized load balancing, IEEE Transactions on Parallel and Distributed Systems 12 (10) (2001) 1094–1104.

- [25] A. Stolyar, Pull-based load distribution among heterogeneous parallel servers: The case of multiple routers, Queueing Systems 85 (1-2) (2017) 31–65.
- [26] X. Liu, L. Ying, A simple steady-state analysis of load balancing algorithms in the sub-Halfin-Whitt regime, ACM SIGMETRICS Performance Evaluation Review 46 (2) (2019) 15–17.
- [27] S. M. Varma, F. Castro, S. T. Maguluri, Power-of-d choices load balancing in the sub-halfin whitt regime (2022). arXiv: 2208.07539.
- [28] S. Bhamidi, A. Budhiraja, M. Dewaskar, Near equilibrium fluctuations for supermarket models with growing choices, The Annals of Applied Probability 32 (3) (2022) 2083–2138.
- [29] S. Banerjee, D. Mukherjee, Join-the-shortest queue diffusion limit in Halfin–Whitt regime: Tail asymptotics and scaling of extrema, The Annals of Applied Probability 29 (2) (2019) 1262–1309.
- [30] S. Banerjee, D. Mukherjee, Join-the-shortest queue diffusion limit in Halfin–Whitt regime: Sensitivity on the heavy-traffic parameter, The Annals of Applied Probability 30 (1) (2020) 80–144.
- [31] A. Braverman, Steady-state analysis of the join-the-shortest-queue model in the Halfin–Whitt regime, Mathematics of Operations Research (2020).
- [32] P. Eschenfeldt, D. Gamarnik, Join the shortest queue with many servers. the heavy-traffic asymptotics, Mathematics of Operations Research 43 (3) (2018) 867–886.
- [33] S. Halfin, W. Whitt, Heavy-traffic limits for queues with many exponential servers, Operations research 29 (3) (1981) 567–588.
- [34] X. Liu, L. Ying, Universal scaling of distributed queues under load balancing in the super-halfin-whitt regime, IEEE/ACM Transactions on Networking 30 (1) (2021) 190–201.
- [35] Z. Zhao, S. Banerjee, D. Mukherjee, Many-server asymptotics for join-the-shortest queue in the super-halfin-whitt scaling window (2023). arXiv:2106.00121.
- [36] G. Foschini, J. Salz, A basic dynamic routing problem and diffusion, IEEE Transactions on Communications 26 (3) (1978) 320–327.
- [37] R. Williams, Diffusion approximations for open multiclass queueing networks: Sufficient conditions involving state space collapse, Queueing Systems Theory and Applications (1998) 27 – 88.
- [38] J. M. Harrison, M. J. López, Heavy traffic resource pooling in parallel-server systems, Queueing systems 33 (4) (1999) 339–368.
- [39] V. Gupta, N. Walton, Load balancing in the nondegenerate slowdown regime, Operations Research 67 (1) (2019) 281–294.
- [40] A. J. Ganesh, Big queues, Springer Science & Business Media, 2004.
- [41] P. W. Glynn, W. Whitt, Logarithmic asymptotics for steady-state tail probabilities in a single-server queue, Journal of Applied Probability 31 (A) (1994) 131–156.
- [42] A. Puhalskii, Large deviation analysis of the single server queue, Queueing Systems 21 (1995) 5–66.
- [43] N. G. Duffield, N. O'connell, Large deviations and overflow probabilities for the general single-server queue, with applications, in: Mathematical Proceedings of the Cambridge Philosophical Society, Vol. 118, Cambridge University Press, 1995, pp. 363–374.
- [44] P. Dupuis, R. S. Ellis, The large deviation principle for a general class of queueing systems. i, Transactions of the American Mathematical Society 347 (8) (1995) 2689–2751.
- [45] A. Ridder, A. Shwartz, Large deviations without principle: Join the shortest queue, Mathematical Methods of Operations Research 62 (2005) 467–483.
- [46] R. D. Foley, D. R. McDonald, Join the shortest queue: stability and exact asymptotics, Annals of Applied Probability (2001) 569–607.
- [47] A. A. Puhalskii, A. A. Vladimirov, A large deviation principle for join the shortest queue, Mathematics of Operations Research 32 (3) (2007) 700–710.
- [48] X. Lin, V. Venkataramanan, On the large-deviations optimality of scheduling policies minimizing the drift of a lyapunov function, in: 2009 47th Annual Allerton Conference on Communication, Control, and Computing (Allerton), IEEE, 2009, pp. 919–926.
- [49] V. Venkataramanan, X. Lin, On the queue-overflow probability of wireless systems: A new approach combining large deviations with lyapunov functions, IEEE transactions on information theory 59 (10) (2013) 6367–6392.
- [50] S. Banerjee, Y. Kanoria, P. Qian, Large deviations optimal scheduling of closed queueing networks, arXiv preprint arXiv:1803.04959 (2018).
- [51] M. Zubeldia, M. Mandjes, Large deviations for acyclic networks of queues with correlated gaussian inputs, Queueing Systems 98 (3-4) (2021) 333–371.
- [52] M. Mandjes, Large deviations for Gaussian queues: modeling communication networks, John Wiley & Sons, 2007.
- [53] R. Vershynin, High-dimensional probability: An introduction with applications in data science, Vol. 47, Cambridge university press, 2018.
- [54] Y. Peres, K. Talwar, U. Wieder, Graphical balanced allocations and the $(1 + \beta)$ -choice process, Random Structures & Algorithms 47 (4) (2015) 760–775.
- [55] N. Bansal, O. N. Feldheim, The power of two choices in graphical allocation, in: Proceedings of the 54th Annual ACM SIGACT Symposium on Theory of Computing, 2022, pp. 52–63.
- [56] P. Billingsley, Probability and Measure, 2nd Edition, John Wiley and Sons, 1986.

A. Essential Lemmas and their proofs

Proof of Lemma 7. The proof simply follows by using the Markov's inequality and then optimizing over the value of θ . For any $\theta \in (0, 1/\lambda)$, we have

$$\mathbb{P}(X > x) \le e^{-\theta x} \mathbb{E}[e^{\theta X}] \le \frac{1}{1 - \theta/\lambda} e^{-\theta x}.$$
(A.1)

As the above inequality is valid for any $\theta \in (0, \lambda)$, we can minimize the RHS in the above inequality over the range $(0, \lambda)$. Now, the derivative of the RHS equates to zero when $\theta = \theta^* := \lambda \left(1 - \frac{1}{\lambda x}\right)$, where $\theta^* \in (0, \lambda)$ whenever $x > 1/\lambda$. Thus, for any $x > 1/\lambda$, by substituting $\theta = \theta^*$ in Eq. (A.1), we get the result.

Lemma 11. Consider a Discrete Time Markov Chain $\{\mathbf{x}(t)\}_{t\geq 0}$, where $\mathbf{x}(t) \in \mathbb{R}^d$ for all $t \geq 0$. Let the Markov chain be positive recurrent, with stationary distribution π , and \mathbf{x}_{∞} be a random variable that follows the distribution π . Suppose $f : \mathbb{R}^d \times \mathbb{R} \to \mathbb{R}_+$ is a Lyapunov function such that following conditions hold.

C1. There exists $\theta_0 > 0$ such that $\forall \theta < \theta_0$, we have,

$$\mathbb{E}[f(\mathbf{x}(t),\theta)] < \infty, \quad \forall t \ge 0$$

C2. There exists $\theta_1 > 0$ such that for all $\theta < \theta_1$, the function f satisfies the negative drift condition,

$$\mathbb{E}[\Delta f(\mathbf{x}(t+1),\theta)|\mathbf{x}(t)] = -\gamma(\theta)f(\mathbf{x}(t),\theta) + \beta(\mathbf{x}(t),\theta),$$
(A.2)

where for all $\theta < \theta_1$, one of the following conditions is satisfied

1. There exists $\bar{\beta}(\theta)$ such that

$$\gamma(\theta) > 0, \qquad \qquad \mathbb{E}[\beta(\mathbf{x}(t), \theta)] < \bar{\beta}(\theta), \qquad \qquad \lim_{t \to \infty} \mathbb{E}[\beta(\mathbf{x}(t), \theta)] \quad exists$$

2. There exists a function $b(\theta) < \infty$ such that

$$\mathbb{E}[f(\mathbf{x}(t), \theta)] \le b(\theta), \ \forall t \ge 0.$$

Then, we have that for all $\theta < \min\{\theta_0, \theta_1\}$,

$$\mathbb{E}_{\pi}[f(\mathbf{x}_{\infty},\theta)] = \frac{1}{\gamma(\theta)} \mathbb{E}_{\pi}[\beta(\mathbf{x}_{\infty},\theta)] < \infty.$$

Proof of Lemma 11. For the complete part of the proof, we assume that $\theta < \min\{\theta_0, \theta_1\}$. From, we Condition C1, we have that $\mathbb{E}[f(\mathbf{x}(t), \theta)] < \infty, \forall t \ge 0$. Thus, in Eq. (A.2) in Condition C2, we can take expectation on both sides to get that,

$$\mathbb{E}[f(\mathbf{x}(t+1),\theta)] = (1 - \gamma(\theta))\mathbb{E}[f(\mathbf{x}(t),\theta)] + \mathbb{E}[\beta(\mathbf{x}(t),\theta)]$$

Depending on the value of $\gamma(\theta)$, we consider two cases.

• Case 1: $\gamma(\theta) \ge 1$. In this case, we have that $1 - \gamma(\theta) < 0$. Then, as $f(\mathbf{x}(t), \theta) \ge 0$, from the previous equation, we have

$$\mathbb{E}[f(\mathbf{x}(t+1),\theta)] \le \mathbb{E}[\beta(\mathbf{x}(t),\theta)] \le \bar{\beta}(\theta).$$

As $\mathbb{E}[f(\mathbf{x}(t), \theta)]$ is uniformly bounded by $\bar{\beta}$ for all $t \ge 0$,

$$\mathbb{E}[f(\mathbf{x}_{\infty}, \theta)] = \lim_{t \to \infty} \mathbb{E}[f(\mathbf{x}(t+1), \theta)] \le \bar{\beta}(\theta).$$

Thus, under steady state, we can rewrite Eq. (A.2) in Condition C2 as,

$$\mathbb{E}[f(\mathbf{x}_{\infty}, \theta)] = (1 - \gamma(\theta))\mathbb{E}[f(\mathbf{x}_{\infty}, \theta)] + \mathbb{E}[\beta(\mathbf{x}_{\infty}, \theta)].$$

This gives us the result. Note that the above result also holds when we directly have that $\mathbb{E}[f(\mathbf{x}(t), \theta)]$ is uniformly bounded, as mentioned in Condition C2 part (2).

• Case 2: $\gamma(\theta) \in (0,1)$. In this case, we can use Eq. (A.2) in Condition C2 recursively, to get that

$$\mathbb{E}[f(\mathbf{x}(t+1),\theta)] = (1-\gamma(\theta))^{t+1} \mathbb{E}[f(\mathbf{x}(0),\theta)] + \sum_{i=0}^{t} (1-\gamma(\theta))^{i} \mathbb{E}[\beta(\mathbf{x}(t-i),\theta)].$$
(A.3)

Now, as $\lim_{t\to\infty} \mathbb{E}[\beta(\mathbf{x}(t),\theta)]$ exists, and is equal to $\mathbb{E}[\beta(\mathbf{x}_{\infty},\theta)]$, we have that

$$\lim_{t \to \infty} \sum_{i=0}^{t} (1 - \gamma(\theta))^{i} \mathbb{E}[\beta(\mathbf{x}(t-i), \theta)] = \frac{1}{\gamma(\theta)} \mathbb{E}[\beta(\mathbf{x}_{\infty}, \theta)],$$
$$\lim_{t \to \infty} (1 - \gamma(\theta))^{t+1} \mathbb{E}[f(\mathbf{x}(0), \theta)] = 0.$$

Substituting the above in Eq. (A.3) gives us the result.

B. Proof of results in Section 2

Proof of Proposition 3. For any $i \in \{1, 2, ..., n\}$, we consider the following exponential Lyapunov functions

$$V_{\perp i}(\mathbf{x};\theta) = e^{\theta \mathbf{x}_{\perp i}} = e^{\theta(x_i - \hat{x})}, \qquad \qquad U_{\perp i}(\mathbf{x};\theta) = e^{\theta \mathbf{x}_{\perp i}} = e^{-\theta(x_i - \hat{x})},$$

where $\hat{x} = \frac{1}{n} \sum_{i=1}^{n} x_i$.

First, note that the total queue length at any time t is less than the total number of arrivals till time t. And as the arrivals follow a Poisson process, the total number of arrivals till time t is a Poisson random variable with parameter $\lambda_n t$. This gives us that

$$\mathbb{E}[V_{\perp i}(\mathbf{q}(t);\theta)] \le \exp\left(\lambda_n t \left(e^{\theta} - 1\right)\right) < \infty.$$

Thus, $V_{\perp i}(\cdot)$ satisfies the Condition C1 of Lemma 11. Similarly, $U_{\perp i}(\cdot)$ also satisfies the Condition C1 of Lemma 11.

Suppose G denotes the generator matrix of the underlying CTMC of the JSQ system. Then, the drift of $V_{\perp i}(\mathbf{q}(t);\theta)$, denoted by $\Delta V_{\perp i}(\mathbf{q}(t);\theta)$, is given by

$$\mathbb{E}\left[\Delta V_{\perp i}(\mathbf{q}(t);\theta)\big|\mathbf{q}(t)\right] := \sum_{\mathbf{q}'} G(\mathbf{q}',\mathbf{q}(t)) \left[V_{\perp i}(\mathbf{q}';\theta) - V_{\perp i}(\mathbf{q}(t);\theta)\right].$$

Using this, we have

$$\begin{split} \mathbb{E}\left[\Delta V_{\perp i}(\mathbf{q}(t);\theta) \left| \mathbf{q}(t) \right] \\ &= \lambda_n \left(e^{\theta q_{\perp i}(t) + \theta \left(1 - \frac{1}{n}\right)} \mathbb{1}_{\{i=i^*\}} + e^{\theta q_{\perp i}(t) - \frac{\theta}{n}} \mathbb{1}_{\{i\neq i^*\}} - e^{\theta q_{\perp i}(t)} \right) \\ &+ \sum_{j=1}^n \mu \left(e^{\theta q_{\perp i}(t) - \theta \left(1 - \frac{1}{n}\right)} \mathbb{1}_{\{j=i\}} + e^{\theta q_{\perp i}(t) + \frac{\theta}{n}} \mathbb{1}_{\{j\neq i\}} - e^{\theta q_{\perp i}(t)} \right) \mathbb{1}_{\{q_j > 0\}} \\ &= \lambda_n e^{\theta q_{\perp i}(t)} \left(e^{\theta \left(1 - \frac{1}{n}\right)} \mathbb{1}_{\{i=i^*\}} + e^{-\frac{\theta}{n}} \mathbb{1}_{\{i\neq i^*\}} - 1 \right) \\ &+ \mu e^{\theta q_{\perp i}(t)} \sum_{j=1}^n \left(e^{-\theta \left(1 - \frac{1}{n}\right)} \mathbb{1}_{\{j=i\}} + e^{\frac{\theta}{n}} \mathbb{1}_{\{j\neq i\}} - 1 \right) \mathbb{1}_{\{q_j > 0\}} \end{split}$$

Exponential Tail Bounds on Queues

$$= \lambda_n e^{\theta q_{\perp i}(t)} \left(\left(e^{\theta} - 1 \right) e^{-\frac{\theta}{n}} \mathbb{1}_{\{i=i^*\}} + e^{-\frac{\theta}{n}} - 1 \right) \\ + \mu e^{\theta q_{\perp i}(t)} \left(\left(e^{-\theta} - 1 \right) e^{\frac{\theta}{n}} \mathbb{1}_{\{q_i(t)>0\}} + \left(e^{\frac{\theta}{n}} - 1 \right) \sum_j \mathbb{1}_{\{q_j>0\}} \right).$$

where i^* is the index chosen by the JSQ dispatcher.

Next, we consider the function $V_{\perp}(\mathbf{x};\theta) = \sum_{i=1}^{n} V_{\perp i}(\mathbf{x};\theta)$. Then, the drift of $V_{\perp}(\mathbf{q}(t);\theta)$ is just the sum of the drifts of $V_{\perp i}(\mathbf{q}(t);\theta)$, that is,

$$\begin{split} \mathbb{E} \Big[\Delta V_{\perp}(\mathbf{q}(t);\theta) \big| \mathbf{q}(t) \Big] \\ &= \sum_{i=1}^{n} \mathbb{E} \Big[\Delta V_{\perp i}(\mathbf{q}(t);\theta) \big| \mathbf{q}(t) \Big] \\ &= \lambda_{n} \left(e^{\theta} - 1 \right) e^{-\frac{\theta}{n}} e^{\theta q_{\perp,\min}(t)} + \lambda_{n} \left(e^{-\frac{\theta}{n}} - 1 \right) \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} \\ &+ \mu \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} \left(\left(e^{-\theta} - 1 \right) e^{\frac{\theta}{n}} \mathbbm{1}_{\{q_{i}(t)>0\}} + \left(e^{\frac{\theta}{n}} - 1 \right) \sum_{j} \mathbbm{1}_{\{q_{j}>0\}} \right) \\ &= \lambda_{n} \left(e^{\theta} - 1 \right) e^{-\frac{\theta}{n}} e^{\theta q_{\perp,\min}(t)} + \lambda_{n} \left(e^{-\frac{\theta}{n}} - 1 \right) \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} \\ &+ \mu \left(e^{-\theta} - 1 \right) e^{\frac{\theta}{n}} \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} + n\mu \left(e^{\frac{\theta}{n}} - 1 \right) \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} \\ &- \mu \left(e^{-\theta} - 1 \right) e^{\frac{\theta}{n}} \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)} \mathbbm{1}_{\{q_{i}(t)=0\}} - \mu \left(e^{\frac{\theta}{n}} - 1 \right) \sum_{j=1}^{n} \mathbbm{1}_{\{q_{j}(t)=0\}} \sum_{i=1}^{n} e^{\theta q_{\perp i}(t)}. \end{split}$$

Next, by using $\theta > 0$, $q_{\perp,\min}(t) = \min_i \left\{ q_i(t) - \frac{1}{n} \sum_{j=1}^n q_j(t) \right\} < 0$, and $e^{\theta q_{\perp i}(t)} \mathbbm{1}_{\{q_i(t)=0\}} = e^{-\frac{\theta}{n} \sum_{j=1}^n q_j(t)} \mathbbm{1}_{\{q_i(t)=0\}} \le 1$, we have

$$\mathbb{E}\left[\Delta V_{\perp}(\mathbf{q}(t);\theta)\big|\mathbf{q}(t)\right] \leq \lambda_n \left(e^{\theta} - 1\right) e^{-\frac{\theta}{n}} + n\mu \left(1 - e^{-\theta}\right) e^{\frac{\theta}{n}} \\ + \left(\lambda_n \left(e^{-\frac{\theta}{n}} - 1\right) + \mu \left(e^{-\theta} - 1\right) \right) \\ + n\mu \left(e^{\frac{\theta}{n}} - 1\right) \sum_{i=1}^n e^{\theta q_{\perp i}(t)}$$

For any $\theta > 0$, we have that

$$-\frac{1}{n}e^{\theta} \leq \frac{e^{-\frac{\theta}{n}}-1}{1-e^{-\theta}} \leq -\frac{1}{n}, \qquad \qquad \frac{1}{n} \leq \frac{e^{\frac{\theta}{n}}-1}{1-e^{-\theta}} \leq \frac{1}{n}e^{\theta},$$

This gives us that, for any $\theta > 0$,

$$\frac{1}{1-e^{-\theta}} \mathbb{E}\left[\Delta V_{\perp}(\mathbf{q}(t);\theta) \big| \mathbf{q}(t)\right] \leq \lambda_n e^{\theta} + n\mu e^{\frac{\theta}{n}} + \left(-\frac{\lambda_n}{n} + \mu(e^{\theta}-1)\right) \sum_{i=1}^n e^{\theta q_{\perp i}(t)}$$
$$\leq 2n\mu e^{\theta} - \frac{\lambda_n}{n} \sum_{i=1}^n e^{\theta q_{\perp i}(t)} + \mu \theta e^{\theta} \sum_{i=1}^n e^{\theta q_{\perp i}(t)}$$
(B.1)

By using similar calculation as in Eq. (B.1), and defining $U_{\perp}(\mathbf{x}; \theta) := \sum_{i=1}^{n} U_{\perp i}(\mathbf{x}; \theta)$, we have that

$$\mathbb{E}\left[\Delta U_{\perp}(\mathbf{q}(t);\theta) | \mathbf{q}(t)\right]$$

$$\begin{split} &= \lambda_n \left(e^{-\theta} - 1 \right) e^{\frac{\theta}{n}} e^{-\theta q_{\perp,\min}(t)} + \lambda_n \left(e^{\frac{\theta}{n}} - 1 \right) \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} \\ &+ \mu \left(e^{\theta} - 1 \right) e^{-\frac{\theta}{n}} \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} + n\mu \left(e^{-\frac{\theta}{n}} - 1 \right) \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} \\ &- \mu \left(e^{\theta} - 1 \right) e^{-\frac{\theta}{n}} \sum_{j=1}^n e^{-\theta q_{\perp i}(t)} \mathbbm{1}_{\{q_i(t)=0\}} \\ &- \mu \left(e^{-\frac{\theta}{n}} - 1 \right) \sum_{j=1}^n \mathbbm{1}_{\{q_j(t)=0\}} \sum_{i=1}^n e^{-\theta q_{\perp i}(t)}. \end{split}$$

Thus, for any $\theta > 0$, by using $q_{\perp i}(t) = q_{\perp,\min}(t) = -\frac{1}{n} \sum_{j=1}^{n} q_i(t)$ whenever $q_i(t) = 0$, we have

$$\frac{1}{1-e^{-\theta}} \mathbb{E}\left[\Delta U_{\perp}(\mathbf{q}(t);\theta) | \mathbf{q}(t)\right] \\
\leq -\lambda_{n} e^{-\theta q_{\perp,\min}(t)} + \left(\frac{\lambda_{n}}{n} e^{\theta} + \mu(e^{\theta} - 1)\right) \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)} \\
- \mu e^{-\theta q_{\perp,\min}(t)} \sum_{j=1}^{n} \mathbb{1}_{\{q_{j}(t)=0\}} + \frac{\mu}{n} e^{\theta} \sum_{j=1}^{n} \mathbb{1}_{\{q_{j}(t)=0\}} \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)} \\
\stackrel{(a)}{\leq} -\lambda_{n} e^{-\theta q_{\perp,\min}(t)} + \left(\frac{\lambda_{n}}{n} e^{\theta} + 2\mu(e^{\theta} - 1)\right) \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)} \\
\stackrel{(b)}{\leq} -\lambda_{n} e^{-\theta q_{\perp,\min}(t)} + \frac{\lambda_{n}}{n} \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)} + 3\mu \theta e^{\theta} \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)},$$
(B.2)

where (a) follows by using $e^{-\theta q_{\perp,\min}(t)} \ge \sum_{i=1}^{n} e^{-\theta q_{\perp i}(t)}$ and $\sum_{j=1}^{n} \mathbbm{1}_{\{q_j(t)=0\}} \le n$; and (b) follows by using $\frac{\lambda_n}{n}e^{\theta} = \frac{\lambda_n}{n} + \frac{\lambda_n}{n}(e^{\theta} - 1) \le \frac{\lambda_n}{n} + \mu\theta e^{\theta}$. Before proceeding with the next step, we provide a bound on the following function.

$$D(\mathbf{x};\theta) := -e^{-\theta x_{\min}} + \frac{1}{n} \sum_{i=1}^{n} e^{-\theta x_i} - \frac{1}{n} \sum_{i=1}^{n} e^{\theta x_i},$$

where $\mathbf{x} \in \mathbb{R}^n$ such that $\sum_{i=1}^n x_i = 0$ and $x_{\min} = \min_i x_i$. Then, $x_{\min} \leq 0$. Suppose $\mathcal{I}^+ = \{i : x_i \geq 0\}$ and $\mathcal{I}^- = [n] \setminus \mathcal{I}^+$. Suppose $n_+ = |\mathcal{I}^+|$ and $n_- = |\mathcal{I}^-|$

1. Case 1: Consider the case

$$n_+ \ge n/4$$
, and $n_- \le 3n/4$

In this case, we have

$$\begin{split} D(\mathbf{x};\theta) &= -e^{-\theta x_{\min}} + \frac{1}{n} \sum_{i \in \mathcal{I}^+} e^{-\theta x_i} + \frac{1}{n} \sum_{i \in \mathcal{I}^-} e^{-\theta x_i} - \frac{1}{n} \sum_{i=1}^n e^{\theta x_i} \\ &\stackrel{(a)}{\leq} -e^{-\theta x_{\min}} + \frac{n_+}{n} + \frac{n_-}{n} e^{-\theta x_{\min}} - \frac{1}{n} \sum_{i=1}^n e^{\theta x_i} \\ &= \frac{n_+}{n} (1 - e^{-\theta x_{\min}}) - \frac{1}{n} \sum_{i=1}^n e^{\theta x_i} \\ &\stackrel{(b)}{\leq} \frac{1}{4} (1 - e^{-\theta x_{\min}}) - \frac{1}{n} \sum_{i=1}^n e^{\theta x_i} \\ &\leq \frac{1}{4} - \frac{1}{4n} \sum_{i=1}^n e^{-\theta x_i} - \frac{1}{n} \sum_{i=1}^n e^{\theta x_i}, \end{split}$$

where (a) follows by using $x_i \ge 0$ for all $i \in \mathcal{I}^+$; and (b) follows by using $x_{\min} \le 0$.

2. Case 2: Consider the case

$$\frac{1}{n}\sum_{i=1}^{n}e^{-\theta x_{i}} \le \frac{1}{2}e^{-\theta x_{\min}} + \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}}$$

In this case, we easily have that,

$$D(\mathbf{x};\theta) \le -\frac{1}{2}e^{-\theta x_{\min}} - \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}} \le -\frac{1}{2n}\sum_{i=1}^{n}e^{-\theta x_{i}} - \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}}$$

3. Case 3: Consider the case

$$\frac{1}{n}\sum_{i=1}^{n}e^{-\theta x_{i}} > \frac{1}{2}e^{-\theta x_{\min}} + \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}} \text{ and } n_{+} \le n/4.$$

In this case, suppose $\sum_{i \in \mathcal{I}^+} x_i = B$, then $\sum_{i \in \mathcal{I}^-} -x_i = B$ as $\sum_{i=1}^n x_i = 0$. Further, as $n_- \geq 3n/4$, suppose \mathcal{I}_1^- is the collection of indices of $\frac{n}{2}$ smallest elements in \mathcal{I}^- and $\mathcal{I}_2^- = \mathcal{I}^- \setminus \mathcal{I}_1^-$. Then, for any $i \in \mathcal{I}_1^-$ and $j \in \mathcal{I}_2^-$, we have $x_i \leq x_j$.

We can argue that for any $j \in \mathcal{I}_2^-$, we have $-x_j \leq \frac{2B}{n}$. This can be shown by contraction. Suppose $\exists j \in I_2^-$ such that $-x_j > \frac{2B}{n}$. Then, $-x_i > \frac{2B}{n}$ for all $i \in I_1^-$. As $|I_1^-| = \frac{n}{2}$, we have $\sum_{i \in I^-} -x_i \geq \sum_{i \in I_1^-} -x_i > B$. This creates a contradiction as $\sum_{i \in \mathcal{I}^-} -x_i = B$. Thus, $-x_j \leq \frac{2B}{n}$ for all $j \in \mathcal{I}_2^-$. This implies that,

$$\frac{1}{n}\sum_{i=1}^{n}e^{-\theta x_{i}} \leq \frac{1}{n}\sum_{i\in\mathcal{I}^{+}}e^{-\theta x_{i}} + \frac{1}{n}\sum_{i\in\mathcal{I}^{-}_{2}}e^{-\theta x_{i}} + \frac{1}{n}\sum_{i\in\mathcal{I}^{-}_{1}}e^{-\theta x_{i}} \\
\leq \frac{n_{+}}{n} + \frac{1}{n}\sum_{i\in\mathcal{I}^{-}_{2}}e^{\frac{2\theta B}{n}} + \frac{1}{2}e^{-\theta x_{\min}} \\
\leq \frac{1}{4} + \frac{1}{2}e^{\frac{2\theta B}{n}} + \frac{1}{2}e^{-\theta x_{\min}}.$$
(B.3)

Next, we have further two cases

- Case 3.1: $\theta B \leq \frac{n}{2}$. In this case, we have that $e^{\frac{2\theta B}{n}} \leq e$.

- Case 3.2: $\theta B \geq \frac{n}{2}$. In this case, by using the Jensen's inequality,

$$\frac{1}{n}\sum_{i=1}^n e^{\theta x_i} \geq \frac{1}{n}\sum_{i\in\mathcal{I}^+}e^{\theta x_i} \geq \frac{n_+}{n}e^{\frac{\theta B}{n_+}} \geq \frac{1}{4}e^{\frac{4\theta B}{n}},$$

where the last inequality holds as $\theta B \ge \frac{n}{2}$, and so, $\frac{n_+}{n} e^{\frac{\theta B}{n_+}}$ is minimized by taking $n_+ = \frac{n}{4}$. Thus, by using the assumption $\frac{1}{n} \sum_{i=1}^{n} e^{-\theta x_i} > \frac{1}{2} e^{-\theta x_{\min}} + \frac{1}{2n} \sum_{i=1}^{n} e^{\theta x_i}$ and Eq. (B.3), we have that

$$\frac{1}{8}e^{\frac{4\theta B}{n}} \leq \frac{1}{4} + \frac{1}{2}e^{\frac{2\theta B}{n}} \leq \frac{3}{4}e^{\frac{2\theta B}{n}}.$$

This gives us that $e^{\frac{2\theta B}{n}} \leq 6$.

Thus, in both the above cases, $e^{\frac{2\theta B}{n}} \leq 6$. This gives us

$$\frac{1}{n}\sum_{i=1}^{n}e^{-\theta x_{i}} \le \frac{1}{4} + 3 + \frac{1}{2}e^{-\theta x_{\min}} \le 4 + \frac{1}{2}e^{-\theta x_{\min}}$$

And thus,

$$D(\mathbf{x};\theta) \le 4 - \frac{1}{2}e^{-\theta x_{\min}} - \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}} \le 4 - \frac{1}{2n}\sum_{i=1}^{n}e^{-\theta x_{i}} - \frac{1}{2n}\sum_{i=1}^{n}e^{\theta x_{i}}$$

By combining the results from Case 1, Case 2 and Case 3, we get

$$D(\mathbf{x};\theta) \le 4 - \frac{1}{4n} \sum_{i=1}^{n} e^{-\theta x_i} - \frac{1}{2n} \sum_{i=1}^{n} e^{\theta x_i}$$
(B.4)

Now, from Eq. (B.1) and (B.2), for any $\theta > 0$, we have

$$\begin{split} \frac{1}{1-e^{-\theta}} \left(\mathbb{E}[\Delta V_{\perp}(\mathbf{q}(t);\theta)|\mathbf{q}(t)] + \mathbb{E}[\Delta U(\mathbf{q}(t);\theta)|\mathbf{q}(t)] \right) \\ &\leq 2n\mu e^{\theta} + \lambda_n D(\mathbf{q}(t);\theta) + \mu \theta e^{\theta} \sum_{i=1}^n e^{\theta q_{\perp i}(t)} + 3\mu \theta e^{\theta} \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} \\ &\stackrel{(a)}{\leq} 2n\mu e^{\theta} + 4\lambda_n + \left(\mu \theta e^{\theta} - \frac{\lambda_n}{2n}\right) \sum_{i=1}^n e^{\theta q_{\perp i}(t)} \\ &\quad + \left(3\mu \theta e^{\theta} - \frac{\lambda_n}{4n}\right) \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} \\ &\stackrel{(b)}{\leq} 8n\mu + \left(2\mu \theta - \frac{\lambda_n}{2n}\right) \sum_{i=1}^n e^{\theta q_{\perp i}(t)} + \left(6\mu \theta - \frac{\lambda_n}{4n}\right) \sum_{i=1}^n e^{-\theta q_{\perp i}(t)} \\ &\stackrel{(c)}{\leq} 8n\mu - \frac{\lambda_n}{8n} \sum_{i=1}^n e^{\theta q_{\perp i}(t)} - \frac{\lambda_n}{8n} \sum_{i=1}^n e^{-\theta q_{\perp i}(t)}, \end{split}$$

where (a) follows using Eq. (B.4); (b) follows by taking $\theta < 1/2$, and so $e^{\theta} < 2$; and (c) follows by using $\theta \leq \frac{\lambda_n}{48n\mu}$. Thus, the Lyapunov function $V_{\perp}(\cdot) + U_{\perp}(\cdot)$ satisfies the Condition C2 of Lemma 11. Thus, by using Lemma 11, the drift is zero in steady state, i.e.,

$$\mathbb{E}_{\pi_n}[\Delta V_{\perp}(\mathbf{q};\theta)] + \mathbb{E}_{\pi_n}[\Delta U(\mathbf{q};\theta)] = 0.$$

So, for any $n \ge 2$, by picking $\theta_{\perp} := \frac{1}{96} \le \frac{\lambda_n}{48n\mu}$, we have that for any $0 < \theta \le \theta_{\perp}$,

$$\mathbb{E}_{\pi_n}\Big[\sum_{i=1}^n e^{-\theta q_{\perp i}} + \sum_{i=1}^n e^{\theta q_{\perp i}}\Big] \le \frac{64n^2\mu}{\lambda_n} = \frac{64n}{1-\epsilon_n} \le 128n,$$

where the last inequality holds whenever $\epsilon_n \leq 1/2$. Now, as the servers are homogeneous, by using symmetry of the JSQ system, we have

$$\mathbb{E}_{\pi_n} \left[e^{\theta |q_{\perp i}|} \right] \le \mathbb{E}_{\pi_n} \left[e^{\theta q_{\perp i}} \right] + \mathbb{E}_{\pi_n} \left[e^{-\theta q_{\perp i}} \right] \le 128.$$

This completes the proof.

Proof of Lemma 4. Recall that we use the notation $\lambda_n = n\mu(1 - \epsilon_n)$ and $\overline{q} = \sum_{i=1}^n q_i$. Consider the exponential Lyapunov function

$$V(\mathbf{x};\theta) := \exp\left(\theta\epsilon_n \sum_{i=1}^n x_i\right).$$
(B.5)

Note that the total queue length at any time t is less than the total number of arrivals till time t. And as the arrivals follow a Poisson process, the total number of arrivals till time t is a Poisson random variable with parameter $\lambda_n t$. This gives us that

$$\mathbb{E}[V(\mathbf{q}(t);\theta)] \le \exp\left(\lambda_n t \left(e^{\theta \epsilon_n} - 1\right)\right) < \infty.$$

Thus, $V(\cdot)$ satisfies the Condition C1 of Lemma 11.

Next, the drift of the function $V(\cdot)$ is given by

$$\begin{split} \mathbb{E}[\Delta V(\mathbf{q}(t);\theta)|\mathbf{q}(t)] &= \left[\lambda_n \left(e^{\epsilon_n \theta} - 1\right) + \mu \left(e^{-\epsilon_n \theta} - 1\right) \sum_{i=1}^n \mathbbm{1}_{\{q_i(t)>0\}}\right] V(\mathbf{q}(t);\theta) \\ &= \left(1 - e^{-\epsilon_n \theta}\right) \left[\lambda_n e^{\epsilon_n \theta} - n\mu\right] V(\mathbf{q}(t);\theta) \\ &+ \mu \left(1 - e^{-\epsilon_n \theta}\right) \sum_{i=1}^n \mathbbm{1}_{\{q_i(t)=0\}} V(\mathbf{q}(t);\theta) \\ &= \left(1 - e^{-\epsilon_n \theta}\right) \left(-\gamma_n(\theta) V(\mathbf{q}(t);\theta) + \beta_n(\mathbf{q}(t);\theta)\right), \end{split}$$

where

$$\gamma_n(\theta) := n\mu - \lambda_n e^{\epsilon_n \theta}, \qquad \qquad \beta_n(\mathbf{q}(t); \theta) := \mu \sum_{i=1}^n \mathbb{1}_{\{q_i(t)=0\}} V(\mathbf{q}(t); \theta).$$

Then, we have that

$$\gamma_n(\theta) > 0 \text{ for any } \theta < \theta_n := \frac{1}{\epsilon_n} \log\left(\frac{n\mu}{\lambda_n}\right) = -\frac{1}{\epsilon_n} \log\left(1 - \epsilon_n\right).$$
 (B.6)

Note that as $\theta_n \ge 1$, $\gamma_n(\theta) > 0$ holds for any $\theta \le 1$.

Next, by using Eq. (B.5), we have

$$\mathbb{E}[\beta_n(\mathbf{q}(t);\theta)] \le n\mu \mathbb{E}[V(\mathbf{q}(t);\theta)] < \infty.$$
(B.7)

And, in steady state, for $\theta \in (0, \theta_n)$

$$\mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)] = \mu \sum_{i=1}^n \mathbb{E}_{\pi_n} \left[\mathbbm{1}_{\{q_i=0\}} e^{\theta \epsilon_n \overline{q}} \right]$$

$$\stackrel{(a)}{=} \mu \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} e^{\theta n \epsilon_{n}(q_{i}-q_{\perp i})} \Big]$$

$$= \mu \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} e^{-\theta n \epsilon_{n} q_{\perp i}} \Big]$$

$$\leq \mu \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[e^{\theta n \epsilon_{n} |q_{\perp i}|} \Big]$$

$$\stackrel{(b)}{=} \kappa_{\perp} n \mu,$$

$$(B.8)$$

where (a) follows by using $q_{\perp i} = q_i - \frac{1}{n}\overline{q}$ by definition; and (b) follows by using Proposition 3, whenever $\theta n \epsilon_n < \theta_{\perp}$ for all $\theta \in (0, \theta_n)$, where θ_{\perp} is as given in Proposition 3. Then, for any $\theta \in (0, \theta_n)$ and $\epsilon_n \leq \frac{\theta_{\perp}}{\theta n}$, by combining Eq. (B.6), (B.7) and (B.8), we have that $V(\cdot)$ satisfies the Condition C2 of Lemma 11. Thus, by using Lemma 11, we have that for any $\theta \in (0, \theta_n)$

$$\mathbb{E}_{\pi_n}[V(\mathbf{q};\theta)] = \frac{1}{\gamma_n(\theta)} \mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)].$$
(B.9)

Further, when $\theta < 0$, we easily have

$$\mathbb{E}_{\pi_n}[V(\mathbf{q};\theta)] = \mathbb{E}_{\pi_n}\left[e^{\theta\epsilon_n\overline{q}}\right] \le 1, \qquad \mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)] \le \mu \sum_{i=1}^n \mathbb{E}_{\pi_n}\left[\mathbbm{1}_{\{q_i=0\}}\right] \le n\mu.$$

Thus, as $\mathbb{E}_{\pi_n}[V(\mathbf{q};\theta)]$ and $\mathbb{E}_{\pi_n}[\beta_n(\mathbf{q};\theta)]$ are bounded, by Condition C2 of Lemma 11, we have that for any $\theta < 0$, the Eq. (B.9) is still satisfied. This completes the proof.

Proof of Claim 1. By using Lemma 4 and taking $\theta < 0$ and $\theta \to 0$, we have

$$\mu \mathbb{E}_{\pi_n} \left[\sum_{i=1}^n \mathbb{1}_{\{q_i=0\}} \right] = \lim_{\theta \to 0^-} \mathbb{E}_{\pi_n} [\beta_n(\mathbf{q}; \theta)]$$
$$= \lim_{\theta \to 0^-} \gamma_n(\theta) \mathbb{E}_{\pi_n} [V(\mathbf{q}; \theta)]$$
$$= n\mu - \lambda_n$$
$$= n\epsilon_n \mu.$$

Also, by symmetry, we have $\mathbb{E}_{\pi_n}[\mathbbm{1}_{\{q_i=0\}}] = \mathbb{E}_{\pi_n}[\mathbbm{1}_{\{q_j=0\}}]$ for all i, j. Thus, we get,

$$\mathbb{E}_{\pi_n} \big[\mathbb{1}_{\{q_i=0\}} \big] = \epsilon_n, \quad \forall i \in \{1, 2, \dots, n\}.$$

Further, we have

$$\begin{split} \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} \big| e^{\theta \epsilon_{n} \overline{q}} - 1 \big| \Big] &\stackrel{(a)}{=} \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} \big| e^{\theta n \epsilon_{n}(q_{i}-q_{\perp i})} - 1 \big| \Big] \\ &= \sum_{i=1}^{n} \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} \big| e^{-\theta n \epsilon_{n} q_{\perp i}} - 1 \big| \Big] \\ &\stackrel{(b)}{=} n \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} \big| e^{-\theta n \epsilon_{n} q_{\perp i}} - 1 \big| \Big] \\ &\stackrel{(c)}{\leq} n \mathbb{E}_{\pi_{n}} \Big[\mathbb{1}_{\{q_{i}=0\}} \big]^{\frac{1}{p}} \mathbb{E}_{\pi_{n}} \Big[\big| e^{-\theta n \epsilon_{n} q_{\perp i}} - 1 \big| \Big] \\ &\stackrel{(d)}{\leq} \theta n^{2} \epsilon_{n}^{1+\frac{1}{p}} \mathbb{E}_{\pi_{n}} \Big[|q_{\perp i}|^{r} e^{r|\theta|n\epsilon_{n}|q_{\perp i}|} \Big]^{\frac{1}{r}} \\ &\frac{27}{4} \end{split}$$

$$\stackrel{(e)}{\leq} \theta n^{2} \epsilon_{n}^{1+\frac{1}{p}} \mathbb{E}_{\pi_{n}} \left[|q_{\perp i}|^{2r} \right]^{\frac{1}{2r}} \mathbb{E}_{\pi_{n}} \left[e^{2r|\theta|n\epsilon_{n}|q_{\perp i}|} \right]^{\frac{1}{2r}}$$

$$\stackrel{(f)}{\leq} \theta n^{2} \epsilon_{n}^{1+\frac{1}{p}} \times \frac{2\kappa_{\perp}r}{\theta_{\perp}} \times \kappa_{\perp},$$
(B.10)

where (a) follows by using $q_{\perp i} = q_i - \overline{q}$; (b) follows by using the symmetry of the system as the servers are homogeneous; (c) and (e) follows by using Hölder's inequality by choosing r, p > 0 and $\frac{1}{r} + \frac{1}{p} = 1$; (d) follows as $|e^x - 1| \le |x|e^{|x|}$ for all $x \in \mathbb{R}$ and Eq. (6); and (f) follows by using Proposition 3, by choosing n large enough such that $2r|\theta|n\epsilon_n < \theta_{\perp}$.

Now, by choosing $r = -\log \epsilon_n$ and $p = \frac{\log \epsilon_n}{\log \epsilon_n + 1}$, we have

$$\sum_{i=1}^{n} \mathbb{E}_{\pi_n} \Big[\mathbb{1}_{\{q_i=0\}} \Big| e^{\theta \epsilon_n \overline{q}} - 1 \Big| \Big] \le \frac{2e\kappa_{\perp}^2}{\theta_{\perp}} \alpha \theta n^2 \epsilon_n^2 \log \Big(\frac{1}{\epsilon_n}\Big),$$

whenever $2|\theta| n\epsilon_n \log \Big(\frac{1}{\epsilon_n}\Big) < \theta_{\perp}.$

where the inequality holds whenever $2|\theta|n\epsilon_n \log\left(\frac{1}{\epsilon_n}\right) < \theta_{\perp}$.

C. Proof of results in Section 3

Proof of Lemma 6. First, we show that the MGF of q(t) exists for any $t \ge 0$ and for any $\theta < \theta_0$. Note that as $a(t) \le A$ almost surely, we have, $q(t) \le q(0) + At$, $\forall t \ge 0$. Thus, by assumption $V(q(0); \theta) < \infty$, we have

$$\mathbb{E}\left[e^{\theta \epsilon q(t)}\right] \le \mathbb{E}[V(q(0);\theta)]e^{\theta \epsilon At} < \infty, \ \forall t \ge 0.$$

This shows that $V(\cdot)$ satisfies the Condition C1 of Lemma 11. Further, the MGF of a(t), s(t) and u(t) also exists as they are bounded random variables, as given in Section 3.1. Next, the recursion in Eq. (8) gives us that, q(t+1) - u(t) = q(t) + a(t) - s(t). We write this in an exponential form and take expectation to get

$$\mathbb{E}\left[e^{\theta\epsilon(q(t+1)-u(t))}|q(t)\right] = \mathbb{E}\left[e^{\theta\epsilon(q(t)+a(t)-s(t))}|q(t)\right]$$
$$= \mathbb{E}\left[e^{\theta\epsilon(a-s)}\right]V(q(t);\theta)$$
$$= (1-\gamma_{\epsilon}(\theta))V(q(t);\theta),$$
(C.1)

where the second equality follows as the arrival and the potential service are identically distributed across time, and are independent of the queue length process. Using the relation $q(t+1)u(t) = 0, \forall t \ge 0$, we have

$$\mathbb{E}\left[e^{\theta\epsilon(q(t+1)-u(t))}\big|q(t)\right] = \mathbb{E}\left[V(q(t+1);\theta)|q(t)\right] + \mathbb{E}\left[e^{-\theta\epsilon u(t)}\big|q(t)\right] - 1$$
$$= \mathbb{E}\left[V(q(t+1);\theta)|q(t)\right] - \beta_{\epsilon}(t;\theta).$$

Substituting this in the Eq. (C.1), we arrive at

$$\mathbb{E}[V(q(t+1);\theta)|q(t)] = (1 - \gamma_{\epsilon}(\theta))V(q(t);\theta) + \beta_{\epsilon}(t;\theta).$$
(C.2)

By taking expectation on both sides, we get

$$\mathbb{E}[V(q(t+1);\theta)] = (1 - \gamma_{\epsilon}(\theta))\mathbb{E}[V(q(t);\theta)] + \mathbb{E}[\beta_{\epsilon}(t;\theta)].$$
(C.3)

Now the result in Eq. (11) follows by recursively using the Eq. (C.3).

Further, we can rewrite Eq. (C.2) as

$$\mathbb{E}[\Delta V(q(t);\theta)|q(t)] = -\gamma_{\epsilon}(\theta)V(q(t);\theta) + \beta_{\epsilon}(t;\theta)$$

As $u(t) \leq A$, for any $\theta \in \mathbb{R}$, $\beta_{\epsilon}(t;\theta) \leq e^{|\theta|\epsilon A}$. Further, $\gamma_{\epsilon}(\theta) > 0$ for any $\theta < \theta_{\epsilon}$. Thus, $V(\cdot)$ satisfies the Condition C2 of Lemma 11. Thus, from Lemma 11, we get that for all $\theta < \min\{\theta_0, \theta_{\epsilon}\}$,

$$\mathbb{E}[V(q;\theta)] = \lim_{t \to \infty} \mathbb{E}[V(q(t);\theta)] = \frac{1}{\gamma_{\epsilon}(\theta)} \lim_{t \to \infty} \mathbb{E}[\beta_{\epsilon}(t;\theta)] = \frac{1 - \mathbb{E}_{\pi_{\epsilon}}[e^{-\theta\epsilon u}]}{1 - \mathbb{E}[e^{\epsilon\theta(a-s)}]}.$$

This completes the proof.

Proof of Claim 2. By simply equating the drift of the queue length to zero in steady state, that is

$$\mathbb{E}_{\pi_{\epsilon}} \left[\mathbb{E}[q(t+1) - q(t)|q(t)] \right] = 0,$$

we have that

$$\mathbb{E}_{\pi_{\epsilon}}[u] = \mathbb{E}[s-a] = \epsilon \mu_{\epsilon}.$$

Since $e^y \ge 1 + y$ for all $y \in \mathbb{R}$, we have

$$\mathbb{I} - \mathbb{E}_{\pi_{\epsilon}}[e^{-\theta \epsilon u}] \le \theta \epsilon \mathbb{E}_{\pi_{\epsilon}}[u] = \epsilon^2 \theta \mu_{\epsilon}.$$

Proof of Claim 3. We have

$$\mathbb{E}\left[e^{\epsilon\theta(a-s)}\right] \le 1 + \theta\epsilon\mathbb{E}[a-s] + \frac{1}{2}\epsilon^2\theta^2\mathbb{E}\left[(a-s)^2\right] + \frac{1}{6}\epsilon^3\theta^3\mathbb{E}\left[(a-s)^3\right] + \epsilon^4\theta^4\mathbb{E}\left[(a-s)^4e^{\epsilon\theta(a-s)}\right]$$
(C.4)

$$\leq 1 - \theta \epsilon^2 \mu_{\epsilon} + \frac{1}{2} \epsilon^2 \theta^2 (\sigma_{\epsilon}^2 + \epsilon^2 \mu_{\epsilon}^2) + \frac{1}{6} \epsilon^3 \theta^3 E_3 + (\epsilon \theta A)^4 \mathbb{E}[e^{\epsilon \theta (a-s)}]$$
(C.5)

where the second inequality holds by the properties of the arrival and the service process and by using $E_3 := \max\{0, \mathbb{E}[(a-s)^3]\}$. This implies that

$$\mathbb{E}\big[e^{\epsilon\theta(a-s)}\big] \leq \frac{1}{1-(\epsilon\theta A)^4} \Big(1-\theta\epsilon^2\mu_\epsilon + \frac{1}{2}\epsilon^2\theta^2(\sigma_\epsilon^2+\epsilon^2\mu_\epsilon^2) + \frac{1}{6}\epsilon^3\theta^3E_3\Big).$$

Thus, to ensure $\mathbb{E}\left[e^{\epsilon\theta(a-s)}\right] < 1$, we only need

$$1 - \theta \epsilon^2 \mu_{\epsilon} + \frac{1}{2} \epsilon^2 \theta^2 (\sigma_{\epsilon}^2 + \epsilon^2 \mu_{\epsilon}^2) + \frac{1}{6} \epsilon^3 \theta^3 E_3 \le 1 - (\epsilon \theta A)^4.$$

This translates to the condition,

$$-\mu_{\epsilon} + \frac{1}{2}\theta(\sigma_{\epsilon}^2 + \epsilon^2 \mu_{\epsilon}^2) + \frac{1}{6}\epsilon\theta^2 E_3 \leq -\epsilon^2\theta^3 A^4.$$

By simple calculations, we have that the above inequality holds true whenever

$$\theta \in \left(0, \mu_{\epsilon} \left(\frac{1}{2}\sigma_{\epsilon}^2 + \epsilon \frac{\mu_{\epsilon}E_3}{3\sigma_{\epsilon}^2} + \epsilon^2 \mu_{\epsilon}^2 + \epsilon^2 \frac{4\mu_{\epsilon}^2 A^4}{\sigma_{\epsilon}^4}\right)^{-1}\right).$$

Thus, by choosing

$$\kappa_1 := \frac{2\mu_{\epsilon}E_3}{3\sigma_{\epsilon}^4} + \epsilon \frac{9\mu_{\epsilon}^2 A^4}{\sigma_{\epsilon}^6}, \qquad \qquad \tilde{\theta}_{\epsilon} := \frac{2\mu_{\epsilon}}{\sigma_{\epsilon}^2(1+\kappa_1\epsilon)}.$$

we get that

$$\mathbb{E}\left[e^{\epsilon\theta(a-s)}\right] < 1, \quad \forall \theta \in \left(0, \tilde{\theta}_{\epsilon}\right).$$
(C.6)

Further, from Eq. (C.4), for any $\theta \in (0, \tilde{\theta}_{\epsilon})$, we also have

$$\begin{split} 1 - \mathbb{E} \Big[e^{\epsilon \theta(a-s)} \Big] &\geq \theta \epsilon^2 \mu_{\epsilon} - \frac{1}{2} \epsilon^2 \theta^2 (\sigma_{\epsilon}^2 + \epsilon^2 \mu_{\epsilon}^2) - \frac{1}{6} \epsilon^3 \theta^3 E_3 - (\epsilon \theta A)^4 \mathbb{E} [e^{\epsilon \theta(a-s)}] \\ &\stackrel{(a)}{\geq} \theta \epsilon^2 \mu_{\epsilon} - \frac{1}{2} \epsilon^2 \theta^2 (\sigma_{\epsilon}^2 + \epsilon^2 \mu_{\epsilon}^2) - \frac{1}{6} \epsilon^3 \theta^3 E_3 - (\epsilon \theta A)^4 \\ &\stackrel{(b)}{\geq} \theta \epsilon^2 \mu_{\epsilon} \Big(1 - \theta \frac{\sigma_{\epsilon}^2}{2\mu_{\epsilon}} \big(1 + \kappa_1 \epsilon \big) \Big), \end{split}$$

where (a) follows by using Eq. (C.6); and (b) follows by using $\theta < \tilde{\theta}_{\epsilon} \leq \frac{2\mu_{\epsilon}}{\sigma_{\epsilon}^2}$.

D. Proof of results in Section 4

D.1. Proof of Lemma 10

Proof. Consider the function $V_1(x;\theta) = e^{\theta[x-n]^+}$. Then, from the definition of $q_n(t)$ and $w_n(t)$, $V_1(q_n(t);\theta) = e^{\theta w_n(t)}$, and similarly, $V_1(q_n;\theta) = e^{\theta w_n}$. Suppose $a_n(t)$ is the total number of arrivals till time t. Then, $a_n(t)$ follows Poission distribution with parameter $\lambda_n t$, and $w_n(t) \le q_n(t) \le a_n(t)$. This gives us that,

$$\mathbb{E}[V_1(q_n(t);\theta)] = \mathbb{E}[e^{\theta w_n(t)}] \le \mathbb{E}[e^{\theta a_n(t)}] = e^{\lambda_n t(e^{\theta} - 1)} < \infty$$

Thus, $V_1(\cdot)$ satisfies Condition C1 of Lemma 11.

Suppose G denotes the generator matrix of the underlying CTMC of M/M/n + M queue. When the number of jobs in the system is q, the drift of $V_1(q;\theta)$, denoted by $\Delta V_1(q;\theta)$ is defined as

$$\mathbb{E}[\Delta V_1(q(t);\theta)|q(t)=q] := \sum_{q'\in\mathbb{W}} G(q',q) \left(V_1(q';\theta) - V_1(q;\theta)\right).$$

Then,

$$\begin{split} \mathbb{E}[\Delta V_1(q(t);\theta)|q(t) &= q] \\ &= \lambda_n V_1(q+1;\theta) + \mu \min\{n,q\} V_1(q-1;\theta) - (\lambda_n + \mu \min\{n,q\}) V_1(q;\theta) \\ &= e^{\theta[q-n]^+} \left(\lambda_n \left(e^{\theta} - 1\right) \mathbb{1}_{\{q \ge n\}} + n\mu \left(e^{-\theta} - 1\right) \mathbb{1}_{\{q > n\}}\right) \\ &= \left(e^{\theta} - 1\right) \left[e^{\theta[q-n]^+} \left(\lambda_n - n\mu e^{-\theta}\right) \mathbb{1}_{\{q > n\}} + \lambda_n \mathbb{1}_{\{q = n\}}\right] \\ &= \left(e^{\theta} - 1\right) \left(\lambda_n - n\mu e^{-\theta}\right) e^{\theta[q-n]^+} \\ &- \left(e^{\theta} - 1\right) \left(\lambda_n \mathbb{1}_{\{q < n\}} - n\mu e^{-\theta} \mathbb{1}_{\{q \le n\}}\right). \end{split}$$

We define

$$\gamma_n(\theta) := - \left(\lambda_n - n\mu e^{-\theta}\right),$$

$$\beta_n(q;\theta) := \left(n\mu e^{-\theta} \mathbb{1}_{\{q \le n\}} - \lambda_n \mathbb{1}_{\{q < n\}}\right).$$

Then, we have

$$\mathbb{E}[\Delta V_1(q_n(t);\theta)|q_n(t)] = \left(e^{\theta} - 1\right) \left(-\gamma_n(\theta)V_1(q_n(t);\theta) + \beta_n(q_n(t);\theta)\right).$$

Note that for any

$$\theta < \theta_n := -\log\left(\frac{\lambda_n}{n\mu}\right) = \log\frac{1}{1-\epsilon_n},$$

we have $\gamma_n(\theta) > 0$. Further, $\beta_n(\cdot)$ is a bounded function as, $\beta_n(q_n(t);\theta) \leq n\mu (e^{\theta} - 1)$, and then, as $q_n(t)$ converges to q_n in distribution, we have

$$\mathbb{E}[\beta_n(q_n;\theta)] = \left(e^{\theta} - 1\right) \left(n\mu e^{-\theta} \mathbb{P}(q_n \le n) - \lambda_n \mathbb{P}(q_n < n)\right)$$

Thus, $V_1(\cdot)$ also satisfies Condition C2 of Lemma 11. Then, we get that for all $\theta < \theta_n$, we have

$$\mathbb{E}[e^{\theta w_n}] = \frac{n\mu e^{-\theta} \mathbb{P}(q_n \le n) - \lambda_n \mathbb{P}(q_n < n)}{n\mu e^{-\theta} - \lambda_n}.$$
 (D.1)

By putting $\theta = 0$ in the above equation, we get

$$n\mu\mathbb{P}(q_n \le n) - \lambda_n\mathbb{P}(q_n < n) = n\mu - \lambda_n = n\mu\epsilon_n.$$

This implies that,

$$-n\mu\epsilon_n \mathbb{P}(w_n > 0) + \lambda_n \mathbb{P}(q_n = n) = 0.$$
(D.2)

Further, $\mathbb{E}[e^{\theta w_n}] = \mathbb{E}[e^{\theta w_n} \mathbb{1}_{\{w_n > 0\}}] + \mathbb{P}(q_n \le n)$. Combining this with Eq. (D.1) and (D.2), we arrive at

$$\mathbb{E}\left[e^{\theta w_n}\mathbb{1}_{\{w_n>0\}}\right] = \frac{n\mu\epsilon_n\mathbb{P}(w_n>0)}{n\mu e^{-\theta} - \lambda_n}.$$

This implies that,

$$\mathbb{E}\left[e^{\theta w_n} \middle| w_n > 0\right] = \frac{n\mu\epsilon_n}{n\mu e^{-\theta} - \lambda_n} = \frac{1}{1 - \frac{1}{\epsilon_n} \left(1 - e^{-\theta}\right)}.$$

Next, we evaluate the MGF of r_n . Consider the function $V_2(x;\theta) = e^{\theta[n-x]^+}$. Then, from the definition of $q_n(t)$ and $r_n(t)$, $V_2(q_n(t);\theta) = e^{\theta r_n(t)}$, and similarly, $V_2(q_n;\theta) = e^{\theta r_n}$. The drift of $V_2(q;\theta)$, denoted by $\Delta V_2(q;\theta)$ is defined as

$$\mathbb{E}[\Delta V_2(q_n(t);\theta)|q_n(t)=q] := \sum_{q'\in\mathbb{W}} G(q',q) \left(V_2(q';\theta) - V_2(q;\theta)\right).$$

Then,

$$\begin{split} \mathbb{E}[\Delta V_2(q_n(t);\theta)|q_n(t) &= q] \\ &= \lambda_n V_2(q+1;\theta) + \mu \min\{n,q\} V_2(q-1;\theta) - (\lambda_n + \mu \min\{n,q\}) V_2(q;\theta) \\ &= e^{\theta[n-q]^+} \left(\lambda_n \left(e^{-\theta} - 1\right) \mathbb{1}_{\{q < n\}} + \mu \min\{n,q\} \left(e^{\theta} - 1\right) \mathbb{1}_{\{q \le n\}}\right) \\ &= e^{\theta[n-q]^+} \left(\lambda_n \left(e^{-\theta} - 1\right) + \mu(n - [n-q]^+) \left(e^{\theta} - 1\right)\right) \\ &+ \lambda_n \left(1 - e^{-\theta}\right) \mathbb{1}_{\{q \ge n\}} - n\mu \left(e^{\theta} - 1\right) \mathbb{1}_{\{q > n\}} \\ &= (1 - e^{-\theta}) \left(-\tilde{\gamma}_n(q;\theta) V_2(q;\theta) + \tilde{\beta}_n(q;\theta)\right), \end{split}$$

where

$$\tilde{\gamma}_n(q;\theta) := \left(\lambda_n - \mu(n - [n-q]^+)e^\theta\right),\\ \tilde{\beta}_n(q;\theta) := \left(\lambda_n \mathbb{1}_{\{q \ge n\}} - n\mu e^\theta \mathbb{1}_{\{q > n\}}\right).$$

Note that $V_2(q;\theta) \leq e^{\theta n}$. Thus, as $V_2(\cdot)$ is uniformly bounded, in steady-state, for any $\theta \in \mathbb{R}$, we have

$$\mathbb{E}[\Delta V_2(q_n;\theta)] = 0$$

and so,

$$\mathbb{E}[\tilde{\gamma}_n(q_n;\theta)V_2(q_n;\theta)] = \mathbb{E}[\tilde{\beta}_n(q_n;\theta)].$$

This gives us that for any $\theta \in \mathbb{R}$,

$$\left(\lambda_n e^{-\theta} - n\mu\right) \mathbb{E}\left[e^{\theta r_n}\right] + \mu \mathbb{E}\left[r_n e^{\theta r_n}\right] = \lambda_n e^{-\theta} \mathbb{P}(q_n \ge n) - n\mu \mathbb{P}(q_n > n).$$
(D.3)

Further, $\mathbb{E}[e^{\theta r_n}] = \mathbb{E}[e^{\theta r_n} \mathbbm{1}_{\{r_n > 0\}}] + \mathbb{P}(q_n \ge n)$, and

$$\mathbb{E}[r_n e^{\theta r_n}] = \mathbb{E}[r_n e^{\theta r_n} \mathbb{1}_{\{r_n > 0\}}] = \frac{d}{d\theta} \mathbb{E}[e^{\theta r_n} \mathbb{1}_{\{r_n > 0\}}].$$

Substituting this in Eq. (D.3), we have

$$(n(1-\epsilon_n)e^{-\theta}-n)\mathbb{E}[e^{\theta r_n}\mathbb{1}_{\{r_n>0\}}] + \frac{d}{d\theta}\mathbb{E}[e^{\theta r_n}\mathbb{1}_{\{r_n>0\}}] = n\mathbb{P}(q_n=n).$$

Solving the above differential equation gives us that

$$\mathbb{E}\left[e^{\theta r_n}\mathbb{1}_{\{r_n>0\}}\right] = n\mathbb{P}(q_n = n)\tilde{G}_n^{-1}(\theta)\int_{-\infty}^{\theta}G_n(t)dt$$

where

$$G_n(t) = \exp\left(-nt - (1 - \epsilon_n)(e^{-t} - 1)\right) = \exp\left(-n\epsilon_n t - (1 - \epsilon_n)(e^{-t} + t - 1)\right).$$

By substituting $\theta = 0$, we get

$$\mathbb{P}(r_n > 0) = n \mathbb{P}(q_n = n) \int_{-\infty}^0 G_n(t) dt.$$
 (D.4)

Thus, we finally get

$$\mathbb{E}\left[e^{\theta r_n}\mathbb{1}_{\{r_n>0\}}\right] = \mathbb{P}(r_n>0)\tilde{G}_n^{-1}(\theta)\left(\int_{-\infty}^0 G_n(t)dt\right)^{-1}\int_{-\infty}^\theta G_n(t)dt$$

This implies that, for any $\theta \in \mathbb{R}$,

$$\mathbb{E}\left[e^{\theta r_n}|r_n>0\right] = \tilde{G}_n^{-1}(\theta) \left(\int_{-\infty}^0 G_n(t)dt\right)^{-1} \int_{-\infty}^\theta G_n(t)dt$$

Next, we calculate the steady state values of $\mathbb{P}(r_n > 0)$, $\mathbb{P}(w_n > 0)$ and $\mathbb{P}(q_n = n)$. Note that

$$\mathbb{P}(r_n > 0) + \mathbb{P}(q_n = n) + \mathbb{P}(w_n > 0) = \mathbb{P}(q_n < n) + \mathbb{P}(q_n = n) + \mathbb{P}(q_n > n) = 1.$$

We use this to solve for $\mathbb{P}(q_n = n)$. By using Eq. (D.2) and (D.4) to get that

$$\left(\frac{\lambda_n}{n\mu\epsilon_n} + 1 + n\int_{-\infty}^0 G_n(t)dt\right)\mathbb{P}(q_n = n) = 1.$$

This gives us that

$$\mathbb{P}(q_n = n) = \left(\frac{1}{\epsilon_n} + n \int_{-\infty}^0 G_n(t) dt\right)^{-1}.$$

Now, the result follows using Eq. (D.2) and (D.4).

Proof of Theorem 8. First note that for any a, b and s_1, s_2 , we have

$$\int_{s_1}^{s_2} \exp\left(at - \frac{1}{2}bt^2\right)$$
$$= \sqrt{\frac{2\pi}{b}} \exp\left(\frac{a^2}{2b}\right) \left[\Phi\left(\sqrt{b}s_2 - \frac{a}{\sqrt{b}}\right) - \Phi\left(\sqrt{b}s_1 - \frac{a}{\sqrt{b}}\right)\right],\tag{D.5}$$

where $\Phi(\cdot)$ denotes the CDF of a standard normal random variable. First, we prove the result for the number of idle servers when the system is in Super-HW regime. In this regime, $\epsilon_n = cn^{-\alpha}$ where $\alpha > \frac{1}{2}$. For convenience, we use $\rho_n = 1 - \epsilon_n$ and $\zeta_n := \sqrt{\frac{n\epsilon_n^2}{\rho_n}}$. For system in Super-HW regime,

$$\zeta_n = \left(\frac{c^2 n^{1-2\alpha}}{1-cn^{-\alpha}}\right)^{\frac{1}{2}} \le 2cn^{\frac{1}{2}-\alpha},$$

where the last inequality follows by assuming $n \ge 4c^2$. Further, if $n^{\alpha-\frac{1}{2}} > 2c$, we have $\zeta_n \le 1$ Now,

$$\begin{split} \int_{-\infty}^{\theta} G_n(t) dt \\ &= \int_{-\infty}^{\theta} \exp\left(-n\epsilon_n t - n\rho_n(e^{-t} + t - 1)\right) dt \\ &= \int_{-\theta}^{\infty} \exp\left(n\epsilon_n t - n\rho_n(e^t - t - 1)\right) dt \\ &= \int_{-\theta}^{\infty} \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) \exp\left(-n\rho_n\left(e^t - \frac{1}{2}t^2 - t - 1\right)\right) dt \\ &\stackrel{(a)}{\leq} \int_{0}^{\infty} \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) dt \\ &\quad + \exp\left(-\frac{1}{6}n\rho_n \theta^3\right) \int_{-\theta}^{0} \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) dt \\ &\stackrel{(b)}{\leq} \sqrt{\frac{2\pi}{n\rho_n}} \exp\left(\frac{1}{2}\zeta_n^2\right) \Phi\left(\zeta_n\right) \\ &\quad + \sqrt{\frac{2\pi}{n\rho_n}} \exp\left(\frac{1}{2}\zeta_n^2 - \frac{1}{6}n\rho_n \theta^3\right) \left[\Phi\left(\sqrt{n\rho_n}\theta + \zeta_n\right) - \Phi\left(\zeta_n\right)\right], \end{split}$$

where (a) follows by using $e^t - \frac{1}{2}t^2 - t - 1 \ge \frac{1}{6}t^3$ for t < 0; and (b) follows by using Eq. (D.5). This also gives us that

$$n\epsilon_n \int_{-\infty}^0 G_n(t)dt \le \sqrt{2\pi}\zeta_n \exp\left(\frac{1}{2}\zeta_n^2\right) \Phi\left(\zeta_n\right) \le \sqrt{2\pi e}\zeta_n,\tag{D.6}$$

where the last inequality follows by using $\zeta_n \leq 1$ whenever $n^{\alpha-\frac{1}{2}} > 2c$. This implies that, in Super-HW regime,

$$\mathbb{P}(r_n > 0) \stackrel{(a)}{=} \frac{n\epsilon_n \int_{-\infty}^0 G_n(t)dt}{1 + n\epsilon_n \int_{-\infty}^0 G_n(t)dt} \le n\epsilon_n \int_{-\infty}^0 G_n(t)dt \le 4e\pi c n^{-\alpha + \frac{1}{2}},\tag{D.7}$$

where (a) follows by using Lemma 10.b. This proves the probability bound presented in Eq. (13). Next, we provide a bound on the tail probability. We also have,

$$\begin{aligned} \int_{-\infty}^{0} G_n(t) dt \\ &= \int_{-\infty}^{0} \exp\left(-n\epsilon_n t - n\rho_n(e^{-t} + t - 1)\right) dt \\ &= \int_{0}^{\infty} \exp\left(n\epsilon_n t - n\rho_n(e^t - t - 1)\right) dt \\ &\ge \exp\left(\zeta_n\right) \int_{1/\sqrt{n\rho_n}}^{\infty} \exp\left(-n\rho_n(e^t - t - 1)\right) dt \\ &\stackrel{(a)}{\ge} \exp\left(\zeta_n\right) \int_{1/\sqrt{n\rho_n}}^{\infty} \exp\left(-\frac{1}{2}n\rho_n t^2 - n\rho_n t^3 e^t\right) dt \\ &\ge \exp\left(\zeta_n\right) \int_{1/\sqrt{n\rho_n}}^{1} \exp\left(-\frac{1}{2}n\rho_n t^2 - en\rho_n t^2\right) dt \\ &\ge \exp\left(\zeta_n\right) \int_{1/\sqrt{n\rho_n}}^{1} \exp\left(-4n\rho_n t^2\right) dt \\ &\ge \exp\left(\zeta_n\right) \int_{1/\sqrt{n\rho_n}}^{1} \exp\left(-4n\rho_n t^2\right) dt \end{aligned}$$

where (a) follows by using $e^t - t - 1 \leq \frac{1}{2}t^2 + t^3e^t$ for all $t \geq 0$. This gives us that

$$\int_{-\infty}^{0} G_n(t)dt \ge \frac{1}{\tilde{\kappa}_1} \times \frac{1}{\sqrt{n\rho_n}} \exp\left(\zeta_n\right),\tag{D.8}$$

where $\tilde{\kappa}_1 := \left(\int_1^2 \exp\left(-4t^2\right) dt\right)^{-1}$. Further, we have that for any $\theta \in (0, \sqrt{n\rho_n})$,

$$\int_{0}^{\rho/\sqrt{n\rho_{n}}} G_{n}(t)dt = \int_{0}^{\theta/\sqrt{n\rho_{n}}} \exp\left(-n\epsilon_{n}t - n\rho_{n}(e^{-t} + t - 1)\right)dt \\
\leq \int_{0}^{\theta/\sqrt{n\rho_{n}}} \exp\left(-\frac{1}{2}n\rho_{n}t^{2} - n\rho_{n}\left(e^{-t} - \frac{1}{2}t^{2} + t - 1\right)\right)dt \\
\stackrel{(a)}{\leq} \int_{0}^{\theta/\sqrt{n\rho_{n}}} \exp\left(-\frac{1}{2}n\rho_{n}t^{2} + \frac{1}{6}n\rho_{n}t^{3}\right)dt \\
\stackrel{(b)}{\leq} \frac{1}{\sqrt{n\rho_{n}}} \int_{0}^{\theta} \exp\left(-\frac{1}{2}t^{2} + \frac{1}{6\sqrt{n\rho_{n}}}t^{3}\right)dt \\
\stackrel{(c)}{\leq} \frac{1}{\sqrt{n\rho_{n}}} \int_{0}^{\theta} \exp\left(-\frac{1}{3}t^{2}\right)dt \\
\leq \frac{1}{\sqrt{n\rho_{n}}} \int_{0}^{\infty} \exp\left(-\frac{1}{3}t^{2}\right)dt \\
\stackrel{(d)}{\leq} \frac{\sqrt{3\pi}}{2\sqrt{n\rho_{n}}}, \qquad (D.9)$$

where (a) follows by using $e^{-t} - \frac{1}{2}t^2 + t - 1 \ge -\frac{1}{6}t^3$; (b) follows by substituting $t \to \frac{1}{\sqrt{n\rho_n}}t$; (c) follows for $\theta < \sqrt{n\rho_n}$; and finally, (d) follows by using Eq. (D.5). Now, by using Lemma 10.a and substituting $\tilde{r}_n = r_n - n\epsilon_n$, for $\theta \in (0, \sqrt{n\rho_n})$, we have

$$\mathbb{E}\Big[\exp\Big(\frac{1}{\sqrt{n\rho_n}}\theta\tilde{r}_n\Big)\Big|r_n>0\Big] \\
= \exp\Big(n\rho_n(e^{-\frac{\theta}{\sqrt{n\rho_n}}}+\frac{\theta}{\sqrt{n\rho_n}}-1)\Big)\Big[1+\left(\int_{-\infty}^0 G_n(t)dt\right)^{-1}\int_0^{\frac{\theta}{\sqrt{n\rho_n}}}G_n(t)dt\Big] \\
\stackrel{(a)}{\leq} \exp\Big(\frac{\theta^2}{2}\Big)\Big[1+\tilde{\kappa}_1\frac{\sqrt{3\pi}}{2}\exp\big(-\zeta_n\big)\Big],$$
(D.10)

where (a) follows by using $e^{-x} + x - 1 \le \frac{x^2}{2}$ for any $x \ge 0$ and Eq. (D.8) and (D.9). Note that the result in Eq. (D.10) is valid for all three regimes.

Next, by using Markov's inequality, for any $\theta > 0$ and $x \ge 0$,

$$\mathbb{P}\Big(\frac{1}{\sqrt{n\rho_n}}r_n > x + \zeta_n \Big| r_n > 0\Big) \le \Big[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2}\Big] \exp\Big(-\theta x + \frac{\theta^2}{2}\Big).$$

As $r_n \leq n$, we only consider the range of x for which $x + \zeta_n \leq \frac{1}{\sqrt{n\rho_n}}n$, which gives $x < \sqrt{n\rho_n}$. Then, by substituting $\theta = x < \sqrt{n\rho_n}$ in the previous equation, we have

$$\mathbb{P}\left(\frac{1}{\sqrt{n\rho_n}}r_n > x + \zeta_n \left| r_n > 0\right) \le \left[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2}\right] e^{-\frac{x^2}{2}}.$$
(D.11)

Now, result in Theorem 8.a follows by using Eq. (D.7) and picking

$$\kappa_1 := 4e\pi \Big[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2} \Big].$$

This completes the proof of Theorem 8.a.

Next, we consider the HW regime. In this case, we have $\epsilon_n = cn^{-\frac{1}{2}}$, and

$$\zeta_n = \frac{c}{\sqrt{1 - cn^{-\frac{1}{2}}}}.$$

Further,

$$\begin{split} &\int_{-\infty}^{0} G_n(t)dt - \sqrt{\frac{2\pi}{n\rho_n}} \exp\left(\frac{1}{2}\zeta^2\right) \Phi\left(\zeta_n\right) \\ &= \int_{0}^{\infty} \exp\left(n\epsilon_n t - n\rho_n(e^t - t - 1)\right) dt - \int_{0}^{\infty} \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) dt \\ &= \int_{0}^{\infty} \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) \left(\exp\left(-n\rho_n\left(e^t - \frac{1}{2}t^2 - t - 1\right)\right) - 1\right) dt \\ &\stackrel{(a)}{\geq} -n\rho_n \int_{0}^{\infty} \left(e^t - \frac{1}{2}t^2 - t - 1\right) \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) dt \\ &\stackrel{(b)}{\geq} -n\rho_n \int_{0}^{\infty} t^3 e^t \exp\left(n\epsilon_n t - \frac{1}{2}n\rho_n t^2\right) dt \\ &\stackrel{(c)}{\geq} -\frac{1}{n\rho_n} \int_{0}^{\infty} t^3 \exp\left((1 + 2c)t - \frac{1}{2}t^2\right) dt \\ &\stackrel{(d)}{\geq} -\frac{\tilde{\kappa}_2}{n\rho_n}, \end{split}$$

where (a) follows by using $e^x - 1 \ge x$ for all $x \in \mathbb{R}$; (b) follows by using $e^t - \frac{1}{2}t^2 - t - 1 \le t^3e^t$ for all $t \ge 0$; (c) follows by substituting $t \to \frac{1}{\sqrt{n\rho_n}}t$ and using $\frac{1}{\sqrt{n\rho_n}} + \zeta_n \le (1+2c)$ when the system is in HW regime and $n \ge 4c^2$; and (d) follows by taking $\tilde{\kappa}_2 := \int_0^\infty t^3 \exp\left((1+2c)t - \frac{1}{2}t^2\right)dt$. This gives us that

$$n\epsilon_n \int_{-\infty}^0 G_n(t)dt \ge \sqrt{2\pi}\zeta_n \exp\left(\frac{1}{2}\zeta^2\right) \Phi\left(\zeta_n\right) - \tilde{\kappa}_2 \frac{\epsilon_n}{\rho_n}.$$

Combining the above inequality with Eq. (D.6), we have

$$-\tilde{\kappa}_2 \frac{\epsilon_n}{\rho_n} \le n\epsilon_n \int_{-\infty}^0 G_n(t)dt - \sqrt{2\pi}\zeta_n \exp\left(\frac{1}{2}\zeta^2\right) \Phi(\zeta_n) \le 0.$$

Thus, we have

$$\lim_{n \to \infty} n\epsilon_n \int_{-\infty}^0 G_n(t)dt = \lim_{n \to \infty} \sqrt{2\pi}\zeta_n \exp\left(\frac{1}{2}\zeta_n^2\right) \Phi\left(\zeta_n\right) = \sqrt{2\pi}c \exp\left(\frac{c^2}{2}\right) \Phi\left(c\right),$$

and so,

$$\lim_{n \to \infty} \mathbb{P}(r_n > 0) = \lim_{n \to \infty} \frac{n\epsilon_n \int_{-\infty}^0 G_n(t)dt}{1 + n\epsilon_n \int_{-\infty}^0 G_n(t)dt} = \frac{\sqrt{2\pi}c \exp\left(\frac{c^2}{2}\right) \Phi\left(c\right)}{1 + \sqrt{2\pi}c \exp\left(\frac{c^2}{2}\right) \Phi\left(c\right)}.$$

Next, by using Eq. (D.10) and Markov's inequality, we have

$$\mathbb{P}\left(\frac{1}{\sqrt{n\rho_n}}r_n > x + \zeta_n\right) \le \mathbb{P}\left(\frac{1}{\sqrt{n\rho_n}}r_n > x + \zeta_n \left| r_n > 0\right)$$
$$\le \left[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2}\right] e^{-\frac{x^2}{2}},$$

where the last inequality follows by same calculations as in Eq. (D.11). Now, result in Theorem 8.b follows by picking

$$\kappa_2 := \left[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2}\right].$$

This completes the proof of Theorem 8.b.

Finally, we consider the Sub-HW regime, i.e., $\epsilon_n = cn^{-\alpha}$ where $\alpha \in (0, \frac{1}{2})$. In this case,

$$\zeta_n = \left(\frac{c^2 n^{1-2\alpha}}{1-cn^{-\alpha}}\right)^{\frac{1}{2}} \ge cn^{\frac{1}{2}-\alpha}.$$

As such $\zeta \to \infty$ as $n \to \infty$. From Eq. (D.8), we have that

$$n\epsilon_n \int_{-\infty}^0 G_n(t)dt \ge \frac{1}{\tilde{\kappa}_1} \zeta_n \exp\left(\zeta_n\right).$$

Then,

$$1 - \mathbb{P}(r_n > 0) = \frac{1}{1 + n\epsilon_n \int_{-\infty}^0 G_n(t)dt} \le \tilde{\kappa}_1 \zeta_n^{-1} \exp\left(-\zeta_n\right) \le \frac{\tilde{\kappa}_1}{c} n^{\alpha - \frac{1}{2}} e^{-cn^{\frac{1}{2} - \alpha}}.$$

Thus, for $\alpha \in (0, \frac{1}{2})$, by choosing $\kappa_3 = \tilde{\kappa}_1$, we have

$$\mathbb{P}(r_n > 0) \ge 1 - \frac{\kappa_3}{c} n^{\alpha - \frac{1}{2}} e^{-cn^{\frac{1}{2} - \alpha}}, \qquad \qquad \lim_{n \to \infty} \mathbb{P}(r_n > 0) = 1.$$

Next, by using Eq. (D.10) and Markov's inequality, we have

$$\mathbb{P}\left(\frac{1}{\sqrt{n\rho_n}}\tilde{r}_n > x\right) \le \mathbb{P}\left(\frac{1}{\sqrt{n\rho_n}}r_n > x \middle| r_n > 0\right)$$
$$\le \left[1 + \tilde{\kappa}_1 \frac{\sqrt{3\pi}}{2} e^{-cn^{\frac{1}{2}-\alpha}}\right] e^{-\frac{x^2}{2}}.$$

Further, for $\theta > 0$, we have $\left(\int_{-\infty}^{0} G_n(t)dt\right)^{-1} \int_{\infty}^{-\theta} G_n(t)dt \le 1$. Then,

$$\mathbb{E}\Big[\exp\Big(-\frac{1}{\sqrt{n\rho_n}}\theta\tilde{r}_n\Big)\Big|r_n>0\Big] \le \exp\Big(n\rho_n\Big(e^{\frac{\theta}{\sqrt{n\rho_n}}}-\frac{\theta}{\sqrt{n\rho_n}}-1\Big)\Big)$$
$$\le \exp\Big(\frac{1}{2}\theta^2+\frac{\theta^3}{\sqrt{n\rho_n}}e^{\frac{\theta}{\sqrt{n\rho_n}}}\Big),$$

This gives us that

$$\mathbb{P}\Big(-\frac{1}{\sqrt{n\rho_n}}\tilde{r}_n > x\Big|r_n > 0\Big) \le e^{-\theta x} \exp\Big(\frac{1}{2}\theta^2 + \frac{\theta^3}{\sqrt{n\rho_n}}e^{\frac{\theta}{\sqrt{n\rho_n}}}\Big).$$

By substituting $\theta = x$ in the above equation, we get that for any $x \ge 0$,

$$\mathbb{P}\Big(-\frac{1}{\sqrt{n\rho_n}}\tilde{r}_n > x\Big|r_n > 0\Big) \le \exp\Big(-\frac{1}{2}x^2 + \frac{x^3}{\sqrt{n\rho_n}}e^{\frac{x}{\sqrt{n\rho_n}}}\Big).$$

Note that $-\tilde{r}_n \leq n\epsilon_n$ as $r_n \geq 0$. Hence, we consider the range of x such that $\sqrt{n\rho_n}x < n\epsilon_n$, or $x < \zeta_n$. For any $x \geq \zeta_n$, the above inequality holds trivially. Then, for $x < \zeta_n$, we have $\frac{x}{\sqrt{n\rho_n}} \leq \frac{\epsilon_n}{\rho_n} \leq 2\epsilon_n \leq 1$, assuming $n^{\alpha} > 2c$ (which holds when $n \geq 4c^2$). Then,

$$\mathbb{P}\Big(-\frac{1}{\sqrt{n\rho_n}}\tilde{r}_n > x\Big|r_n > 0\Big) \le \exp\Big(-\frac{1}{2}x^2 + 2e\epsilon_n x^2\Big).$$

This completes the proof.