Construction of topological representation of geometric patterns using Cantor self-similar set

Shousuke Ohmori^{1,2*)}, Yoshihiro Yamazaki^{3,4}, Tomoyuki Yamamoto^{3,4}, and Akihiko Kitada⁴

¹National Institute of Technology, Gunma College, Maebashi-shi, Gunma 371-8530, Japan ²Waseda Research Institute for Science and Engineering, Waseda University, Shinjuku, Tokyo 169-8555, Japan

³Faculty of Science and Engineering, Waseda University, Shinjuku-ku, Tokyo 169-8555, Japan
 ⁴Institute of Condensed-Matter Science, Comprehensive Resaerch Organization, Waseda University, Shinjuku-ku, Tokyo 169-8555, Japan

*corresponding author: 42261timemachine@ruri.waseda.jp

Abstract

Universal representation of geometric patterns of disordered matters is investigated with the aid of general topology. By utilizing the result obtained in the previous study (S. Ohmori, et.al., Phys. Scr. 94, 105213 (2019)) that any patterns can be represented by a specific topological space, a construction of topological representation of patterns using Cantor set is shown. The obtained topological representations are then demonstrated by the contractions that characterize the self-similarity of Cantor set. For some practical geometric patterns, e.g., network, dendritic, and clusterized patterns, their topological representations are focused on.

1 Introduction

A variety of geometric patterns found in solid and liquid states have been hugely studied from the viewpoint of the physics of disordered systems[1, 2]. The network (graphic) structures of polymers[3], the clusterized structure of molecular liquid[4, 5], and the dendrite found in the process of solidification[6] are practical and familiar examples. In the study of patterns of disordered matter, there is a historically significant issue of finding a mathematical description that can universally and uniformly express these geometric patterns, similar to the group theory in crystallography. To attempt this problem, several mathematical methods based on topological property have been developed. For instance, the persistent homology method that is based on the algebraic topology have been applied to the study of classification of geometric structures formed by amorphous materials[7, 8].

As the other approach, we have studied materials structure from the viewpoint of the most fundamental topology, namely, general topology[9, 10, 11, 12]. In our studies, the geometric patterns are discussed in the context of Continuum Theory, which is one of the field of general topology[13]. A topological space is called a continuum if it is a connected compact Hausdorff-space, and each geometric pattern is specified by using continua. For instance, a dendritic pattern is described as the topological dendrite that is a locally-connected continuum without simple closed curve. (For details of this description, see Sec. 2.) The continua corresponding to the geometric patterns can be expressed indirectly as the formation of a set of equivalence classes for a specific topological space with the 0-dim, perfect, compact properties. Here, the collection of subsets of a topological space relative to equivalence classes is called a decomposition space ¹. Then, each pattern is associated with a decomposition space, homeomorphically, and hence, the patterns can be discussed uniformly through the corresponding decomposition spaces.

Among our methodology, we recently proposed a mathematical model called Cantor cube model[12]. This model provides the practical form of the decomposition spaces of a Cantor cube $X = \{0, 1\}^{\Lambda}$ ² corresponding to some geometric patterns. For instance, focus on a finite graph Y_g composed of arcs E_1, \ldots, E_r ($r < \infty$). (For the definition and the sketch of a finite graph, see Sec. 2.) Then, the corresponding decomposition space \mathcal{D}_g of X can be represented as follows : for a node x connecting with arcs E_{t_1}, \ldots, E_{t_q} and for a point y in an arc E_i of Y_g ,

$$x \doteq \cup_{j=1}^{q} (X^{t_j} \cap S_x^{t_j}), \quad y \doteq X^i \cap S_y^i.$$

$$\tag{1}$$

Here, the right hand sides $\bigcup_{j=1}^{q} (X^{t_j} \cap S_x^{t_j})$ and $X^i \cap S_y^i$ are the points of \mathscr{D}_g and the sign " \doteq " shows identification of the point in Y_g with the corresponding point of \mathscr{D}_g homeomorphically. Intuitively, $\bigcup_{j=1}^{q} X^{t_j}$ in (1) represents that the point *x* possesses just the bonds E_{t_1}, \ldots, E_{t_q} emanated from *x*, and $S_x^{t_j}$ determines the position of *x* in E_{t_j} ³. By the topological representation (1), the network pattern Y_g can be completely copied to \mathscr{D}_g . In the previous study[12], such topological representations using Cantor cube are found for the other geometric patterns. However, it seems that these representations are too abstract to study geometric pattern even more. In particular, the relation (1) does not tell us the information on which property of Cantor cube is essential to represent the geometric pattern.

In the present article, we consider Cantor set instead of Cantor cube. Cantor set is a topological space homeomorphic to Cantor Middle-Third Set (abbreviated as CMTS hereafter)[14]. This space has been hugely studied from viewpoints of chaos and fractal science[15, 16], because it is well known as the most typical self-similar fractal set; its self-similarity can be characterized by equipping the contractions. Also, it is known that Cantor set is the practical form of Cantor cube ⁴. The purpose of this article is to find the essential structure of topological representations by handling Cantor set. Then, it is found that each representation is obtained as a composition of the contractions characterizing the self-similarity of Cantor set.

The current article is organized as follows. In the next section, we review the classification of some geometric patterns based on continua used in our study. In Sec. 3, we show the basic topological properties for the self-similarity of a Cantor set and we sketch the procedure of representing any pattern by a decomposition space of the Cantor set. The obtained result in previous section is applied to some continua with graphic, dendritic, or clusterized structures in Sec. 4. A conclusion is given in Sec. 5.

¹Let (A, τ) be a topological space with a topology τ . A partition \mathscr{D} of A is a set $\{D\}$ of nonempty subsets of A such that $D \cap D' = \emptyset$ for $D \neq D', D, D' \in \mathscr{D}$ and $\bigcup \mathscr{D}(= \bigcup_{D \in \mathscr{D}}) = A$. A decomposition space $(\mathscr{D}, \tau(\mathscr{D}))$ of (A, τ) is a topological space whose topology $\tau(\mathscr{D})$ on a partition \mathscr{D} of A is defined by $\tau(\mathscr{D}) = \{\mathscr{U} \subset \mathscr{D}; \bigcup \mathscr{U} \in \tau\}$. The space $(\mathscr{D}, \tau(\mathscr{D}))$ is a kind of quotient space of (A, τ) . See, for the detailed discussions, [13].

²A Cantor cube is a topological space $(X, \tau) = (\{0, 1\}^{\Lambda}, \tau_0^{\Lambda})$, Card $\Lambda \ge \aleph_0$ where $(\{0, 1\}^{\Lambda}, \tau_0^{\Lambda})$ is the Λ -product space of $(\{0, 1\}, \tau_0)$ for a set Λ and τ_0 is a discrete topology for $\{0, 1\}$. See for details [12].

³According to accurate definitions of the signs X and S in (1), see Sec. 3 in [12].

⁴Cantor cube $(\{0,1\}^{\Lambda}, \tau_0^{\Lambda})$ becomes a Cantor set when Card $\Lambda = \aleph_0$, e.g., $\Lambda = \mathbb{N}$, where \mathbb{N} is the set of natural numbers.

2 Classification of geometric patterns

In this section, classification of geometric patterns based on the concept of continua is reviewed. The patterns are basically classified into the four types of topological spaces; topological graph, topological tree, topological dendrite, and the direct sum of them. Note that graph, tree, dendrite are continua, whereas the direct sum is not a continuum but a disconnected space.

2.1 Topological Graph (Network Pattern)

We first focus on the geometric pattern of network (graphic) configuration. Figure 1 (a) shows an example of such pattern. This pattern is widely found in many disordered matters, e.g., polymers and amorphous. In Continuum Theory, this pattern can be completely associated with a topological space called a topological graph. A topological space is called a topological graph (a finite graph, or a graph) provided that it is a continuum that can be described as the union of finitely many arcs, any two of which are either disjoint or intersect in terms of their end points ⁵. Here, an arc is a topological space homeomorphic to the closed interval [0,1] and its end points are the points mapped homeomorphically into 0 and 1. Figure 1 (b) shows an arc in which e_1 and e_2 indicate its end points. In practice, each arc in a graph characterizes a bond in a network pattern. Note that any arc is itself a topological graph. The other basic example of a graph is a simple closed curve. This is the space homeomorphic to $S^1 := \{x = (x_1, x_2) \in \mathbb{R}^2; |x| = \sqrt{x_1^2 + x_2^2} = 1\}$ (Fig. 1 (c)).



Figure 1: Illustrations of (a) a topological graph (network pattern), (b) an arc with the end points e_1 and e_2 , (c) a simple closed curve.

2.2 Topological Tree (Branching Pattern)

The special pattern of the graph is a topological tree. A topological tree (a tree, or an acyclic graph) is a graph that contains no simple closed curve. Figure 2 (a) shows an example of a topological tree. This pattern corresponds to the branching pattern such as the dendritic pattern. The simplest example of a tree is an arc. Note that a tree that is not an arc can be composed of the family of simple *n*-od spaces, where a space is a simple *n*-od (n = 3, 4, ...) if it is a tree that has a unique point connecting with *n* number of arcs and the other points do not connect with l (> 2) number of arcs. Figure 2 (b) and (c) show the

$$rd(x,X) \le \beta,$$
 (2)

⁵Any graph can be also characterized by focusing on the vicinity of each point. Set a point x in a topological space (X, τ) and a cardinal number β . We say x is of less than or equal to β in X, written

if for any open set $U \in \tau$ containing *x*, there exists $V \in \tau$ such that $x \in V \subset U$ and $Card\partial V \leq \beta$. (∂V shows the boundary of *V*.) Then, we can prove the statement that a continuum *X* is a graph if and only if for any $x \in X$, $ord(x,X) \leq n$ for some $n \in \mathbb{N}$ and $ord(x,X) \leq 2$ for all but finitely many $x \in X$ [13]. In particular, the necessary and sufficient condition that a continuum *X* becomes an arc or a simple closed curve is $ord(x,X) \leq 2$ for any $x \in X$. These statements show that a topological graph can be discussed based on the abstract topological concept such as (2). In Continuum Theory, several general topological methods are used to investigate what shapes the topological graph takes.

simple 3-od and 4-od spaces, respectively. The simple *n*-od space is known as one of the key space to investigate a topological tree in Continuum Theory.



Figure 2: Illustrations of (a) a topological tree, (b) the simple 3-od, (c) the simple 4-od.

2.3 Topological Dendrite (Branching or Self-Similar Pattern)

As a similar pattern to a tree, a topological dendrite is well-known. This space is defined as a Peanocontinuum that contains no simple closed curve. Here, a Peano-continuum is a continuum with locallyconnected. The crucial difference between a topological tree and a topological dendrite is that a topological dendrite is no necessary to become a finite graph. In other words, a dendrite can contain the structure composed of infinitely many arcs and the self-similar structure. For instance, the fern structure that has a self-similarity can be characterized by a topological dendrite. Figure 3 shows an example of a topological dendrite. In this figure, the topological dendrite is composed of the infinitely many arcs of F_i with the end points p_i and l_i (i = 1, 2, ...) and their convergence point p_0 . Note that since any graph is a Peano-continuum, any tree becomes a dendrite. In the practical branching (dendritic) pattern found in nature, we can make use of either a topological tree or a topological dendrite for suiting the pattern.



Figure 3: Schematic explanation of a topological dendrite. F_i shows an arc with the end points p_i and l_i (i = 1, 2, ...).

2.4 Direct Sum (Clusterized Sturcture)

For describing the pattern with a clusterized structure, a topological space called the direct sum is suitable. For the given family of topological spaces $\{(X_i, \tau_i); i = 1, ..., s\}$ satisfying $X_i \cap X_j = \emptyset, i \neq j$, the direct sum $\left(\bigoplus_{i=1}^s X_i, \bigoplus_{i=1}^s \tau_i\right)$ of them is defined as the space $\bigoplus_{i=1}^s X_i = \bigcup_{i=1}^s X_i$ that equips the topology $\bigoplus_{i=1}^s \tau_i = \left\{U \subset \bigoplus_{i=1}^s X_i; U \cap X_i \in \tau_i, i = 1, ..., s\right\}$ [17]. The direct sum is always disconnected. By

corresponding each cluster to X_i , the direct sum provides the whole description of the set of the clusters as one topological space. For instance, the clusterized structure whose cluster possesses a network pattern is characterized by the the direct sum $\bigoplus_{i=1}^{s} X_i$ of the finite graphs X_i . Figure 4 (a) shows this case. By replacing a graphic structure with other continua such as a tree and a dendrite, their clusterized structure can be also constructed. Note that for the continua composed of just one point continuum $\{x_i\}$ (i = 1, ..., s) its direct sum becomes $\{x_1, ..., x_s\}$, which is a finite totally disconnected space (Fig. 4 (b)). Furthermore, we can consider a direct sum $X = \bigoplus_{i=1}^{n} X_i$ of the clusters X_i (i = 1, ..., n) where each X_i is also the direct sum of the other clusters, $X_i = \bigoplus_{k=1}^{s(i)} X_{i,k}$. Then, the direct sum X characterizes a hierarchic structure of the clusters.



Figure 4: Direct sums of (a) the three graphs C_1, C_2 and C_3 , and (b) the family of one-point continuum $\{x_i\}$.

It is noted that the property, the topological spaces appearing in the above classifications are every compact metric spaces, is crucial. Actually, in order to gain the topological representations of any pattern, we utilize the mathematical fact that any compact metric space is homeomorphic to a decomposition space of a 0-dim, perfect, compact Hausdorff-space. In the next section, a practical construction of such topological representation is shown based on a Cantor set.

3 Topological Self-similarity of Cantor set

In this section, we focus on the Cantor set and demonstrate that it has a topological self-similarity, which is characterized by a pair of contractions. The obtained self-similar property is used to construct a topological representation.

At first, we recall what topological self-similar is[18]. A topological space (A, τ) is topologically self-similar if it satisfies the following conditions; (i) (A, τ) is metrizable, namely, the topology τ is identical with a metric topology τ_d specified by a metric d on A and (ii) a set of contractions $\{f_j : (A, \tau_d) \rightarrow (A, \tau_d), j = 1, ..., m(2 \le m < \infty)\}$ that satisfies

$$\cup_{j=1}^{m} f_j(A) = A \tag{3}$$

is found. Here, a map $f : (A, \tau_d) \to (A, \tau_d)$ on a metric space (A, τ_d) is a contraction if there is a constant $0 \le \alpha < 1$ such that $d(f(x), f(y)) \le \alpha d(x, y)$ $(x, y \in A)$. The relation (3) shows the self-similar fractal property of *A* and hence the topological self-similarity is considered as a mathematical generalization of self-similar fractal sets. Indeed, several familiar self-similar sets such as CMTS, Sierpinski gasket, Koch curve, and Menger sponge are topological self-similar.

Next we set a Cantor set. Let $\{0,1\}^{\mathbb{N}}$ be the set of infinitely countable product of $\{0,1\}$ whose topology τ_d is a metric topology with the metric defined by $d(x,x') = \sum_{i=1}^{\infty} |x_i - x'_i|/2^i \ (x,x' \in \{0,1\}^{\mathbb{N}})$. Note that $(\{0,1\}^{\mathbb{N}}, \tau_d)$ is a 0-dim, perfect, compact metric space. And $(\{0,1\}^{\mathbb{N}}, \tau_d)$ is homeomorphic to CMTS= *C*. Indeed, set a function *h* from $\{0,1\}^{\mathbb{N}}$ onto *C* by

$$h: \{0,1\}^{\mathbb{N}} \to C, x \mapsto \sum_{i=1}^{\infty} \frac{2}{3^{i}} x_{i} \quad (x = (x_{1}, x_{2}, \dots) \in \{0,1\}^{\mathbb{N}}).$$
(4)

Then, it becomes a homeomorphism. Thus, $(\{0,1\}^{\mathbb{N}}, \tau_d)$ is a Cantor set. To show its self-similarity, we set

$$\begin{cases} F_0: \{0,1\}^{\mathbb{N}} \to \{0,1\}^{\mathbb{N}}, x = (x_1, x_2, \dots) \mapsto F_0(x) = (0, x_1, x_2, \dots), \\ F_1: \{0,1\}^{\mathbb{N}} \to \{0,1\}^{\mathbb{N}}, x = (x_1, x_2, \dots) \mapsto F_1(x) = (1, x_1, x_2, \dots). \end{cases}$$
(5)

It is confrimed that F_0 and F_1 are injective and the following relations hold : for each i = 0, 1,

$$d(F_i(x), F_i(x')) = \frac{1}{2} d(x, x') \quad (x, x' \in \{0, 1\}^{\mathbb{N}}),$$
(6)

and

$$F_0(\{0,1\}^{\mathbb{N}}) \cup F_1(\{0,1\}^{\mathbb{N}}) = \{0,1\}^{\mathbb{N}}.$$
(7)

The relation (6) shows that F_0 and F_1 are contractions on $\{0,1\}^{\mathbb{N}}$ in terms of the metric *d*. In addition, from (7), it is found that $\{0,1\}^{\mathbb{N}}$ has a topological self-similar structure.

For characterizing a geometric pattern *Y* based on the Cantor set $\{0,1\}^{\mathbb{N}}$, it is necessary to find a continuous map from $\{0,1\}^{\mathbb{N}}$ onto *Y*. The continuous map can be obtained, methodically, by decomposing $\{0,1\}^{\mathbb{N}}$ into its cones, where a subset forming $\{k_1\}_1 \times \cdots \times \{k_i\}_i \times \{0,1\}^{\mathbb{N}-\{1,\ldots,i\}} = \{x : \mathbb{N} \to \{0,1\}, x(l) = k_l \in \{0,1\}, l = 1,\ldots,i\}$ of $\{0,1\}^{\mathbb{N}}$ is called a cone. Note that the decomposition can be performed generally when the space is 0-dim perfect T₀-space. The cone is expressed by a finitely composition of the contractions F_0 and F_1 ,

$$\{k_1\}_1 \times \dots \times \{k_i\}_i \times \{0,1\}^{\mathbb{N} - \{1,\dots,i\}} = F_{k_1} \circ \dots \circ F_{k_i}(\{0,1\}^{\mathbb{N}}),$$
(8)

where $k_l \in \{0,1\}$ $(l = 1, ..., i < \infty)$. This representation (8) of the cone using the contractions F_0 and F_1 is called the "contraction-cone" of $\{0,1\}^{\mathbb{N}}$, hereafter. Note that each contraction-cone is also a Cantor set. By using the contraction-cones (8), $\{0,1\}^{\mathbb{N}}$ is decomposed as $\{0,1\}^{\mathbb{N}} = \bigoplus_{i=1}^{n} X_i$, where

$$\begin{cases} X_1 = F_0(\{0,1\}^{\mathbb{N}}), \\ X_i = F_1 \circ \cdots \circ F_1 \circ F_0(X) = F_1^{i-1} \circ F_0(\{0,1\}^{\mathbb{N}}) \ (i = 2, 3, \dots, n-1), \\ X_n = F_1^{n-1}(\{0,1\}^{\mathbb{N}}). \end{cases}$$
(9)

Each X_i is also decomposed by a set $\{X_{i_1}, \ldots, X_{i_{n_i}}\}$ of the contraction-cones where

$$\begin{cases} X_{i_1} = F_1^{i-1} \circ F_0 \circ F_0(\{0,1\}^{\mathbb{N}}) = F_1^{i-1} \circ F_0^2(\{0,1\}^{\mathbb{N}}), \\ X_{i_j} = F_1^{i-1} \circ F_0 \circ F_1^{j-1} \circ F_0^j(\{0,1\}^{\mathbb{N}}) \ (j = 2, 3, \dots, n_i - 1), \\ X_{i_{n_i}} = F_1^{i-1} \circ F_0 \circ F_1^{n_i-1}(\{0,1\}^{\mathbb{N}}). \end{cases}$$
(10)

Continuing the decomposing procedure provides a continuous map $f : \{0,1\}^{\mathbb{N}} \to Y[12]$. By using the continuous onto map f, the decomposition space $\mathscr{D}_f = \{f^{-1}(y); y \in Y\}$ with a decomposition topology

 $\tau(\mathscr{D}_f) = \{\mathscr{U} \subset \mathscr{D}_f; \bigcup \mathscr{U} \in \tau_d\} \text{ of } \{0,1\}^{\mathbb{N}} \text{ is obtained to be homeomorphic onto } Y. This decomposition space <math>\mathscr{D}_f$ becomes the topological representation of the geometric pattern Y.

As the simplest case, we see the topological representation of a closed interval [0, 1]. Note that this representation is immediately available to obtain the representation of the arc pattern because an arc is homeomorphic to [0, 1]. First the following relation that is a generalization of the relation (8) is satisfied : for $k_1, k_2, \dots \in \{0, 1\}$,

$$\{k_1\}_1 \times \{k_2\}_2 \times \dots \times \{0,1\}^{\mathbb{N} - \{1,2,\dots\}} = \lim F_{k_1} \circ F_{k_2} \circ \dots \circ F_{k_n}(\{0,1\}^{\mathbb{N}})$$

= $\{(k_1,k_2,\dots)\}$ (singleton), (11)

where lim stands for the limit on $\operatorname{Im}(\{0,1\}^{\mathbb{N}}) - \{\emptyset\}$, $\operatorname{Im}(\{0,1\}^{\mathbb{N}})$ is the family of closed sets of $\{0,1\}^{\mathbb{N}}$, and the first equation of the right hand side expresses the limit for the sequence $\{F_{k_1}(\{0,1\}^{\mathbb{N}}), F_{k_1} \circ F_{k_2}(\{0,1\}^{\mathbb{N}}), \dots\}$ in $\operatorname{Im}(\{0,1\}^{\mathbb{N}}) - \{\emptyset\}^6$. For the Cantor cube model $\{0,1\}^{\Lambda}$, the topological representation of [0,1] is obtained as follows; (i) when $y = \sum_{i=1}^{\infty} a_i/2^i \notin M$ for some $a_1, a_2, \dots \in \{0,1\}$, where $M \equiv \{l/2^n; n = 1, 2, \dots$ and $l = 1, \dots, 2^n - 1\}$,

$$f^{-1}(y) = \{a_1\}_{\lambda_1} \times \{a_2\}_{\lambda_2} \times \dots \times \{0,1\}^{\Lambda - \{\lambda_1, \lambda_2, \dots\}},$$
(12)

(ii) when $y = l/2^n \in M$,

$$f^{-1}(y) = \left[\{a_1\}_{\lambda_1} \times \{a_2\}_{\lambda_2} \times \dots \times \{a_{n-1}\}_{\lambda_{n-1}} \times \{0\}_{\lambda_n} \times \{1\}_{\lambda_{n+1}} \times \{1\}_{\lambda_{n+2}} \times \dots \times \{0,1\}^{\Lambda - \{\lambda_1,\lambda_2,\dots\}} \right] \\ \cup \left[\{a_1\}_{\lambda_1} \times \{a_2\}_{\lambda_2} \times \dots \times \{a_{n-1}\}_{\lambda_{n-1}} \times \{1\}_{\lambda_n} \times \{0\}_{\lambda_{n+1}} \times \{0\}_{\lambda_{n+2}} \times \dots \times \{0,1\}^{\Lambda - \{\lambda_1,\lambda_2,\dots\}} \right] (13)$$

for some a_1, \ldots, a_{n-1} , where $\lambda_1, \lambda_2, \cdots \in \Lambda$. Applying (11) to (12) and (13), the topological representation of [0, 1] is reconstructed as follows; (i) for $y = \sum_{i=1}^{\infty} a_i/2^i \notin M$

$$f^{-1}(\mathbf{y}) = \lim F_{a_1} \circ F_{a_2} \circ \dots \circ F_{a_n}(\{0,1\}^{\mathbb{N}}) = \{(a_1, a_2, \dots)\},\tag{14}$$

and (ii) for $y = l/2^n \in M$

$$f^{-1}(y) = \lim F_{a_1} \circ F_{a_2} \circ \dots \circ F_{a_{n-1}} \circ (F_1 \circ F_0^n(\{0,1\}^{\mathbb{N}}) \oplus F_0 \circ F_1^n(\{0,1\}^{\mathbb{N}})) = \{a^0, a^1\}$$
(15)

for some a_1, \ldots, a_{n-1} , where $a^0 = (a_1, \ldots, a_{n-1}, 0, 1, 1, \ldots)$ and $a^1 = (a_1, \ldots, a_{n-1}, 1, 0, 0, \ldots)$. Here, $f^{-1}(0) = \{e_0\}$ and $f^{-1}(1) = \{e_1\}$ where $e_0 = (0, 0, \ldots)$ and $e_1 = (1, 1, \ldots)$ are the elements of $\{0, 1\}^{\mathbb{N}}$.

4 Topological representations of geometric patterns

The method of topological representation obtained in the previous section is now applied to the topological spaces shown in Sec. 2.

⁶For a sequence $\{A_n\}_{n\in\mathbb{N}}$ of $2^{\{0,1\}^{\mathbb{N}}}$, $\lim A_n \equiv \liminf A_n = \limsup A_n$ where $\liminf A_n = \{x \in \{0,1\}^{\mathbb{N}}$; for each open set U of $\{0,1\}^{\mathbb{N}}$ containing $x, U \cap A_n \neq \emptyset$ for all but finitely many $n\}$ and $\limsup A_n = \{x \in \{0,1\}^{\mathbb{N}}$; for each open set U of $\{0,1\}^{\mathbb{N}}$ containing $x, U \cap A_n \neq \emptyset$ for infinitely many $n\}$. Note that $\lim(A_n \cup B_n) = \lim A_n \cup \lim B_n$. The relation (11) is obtained as follows; it is clear that $\{k_1\}_1 \times \{k_2\}_2 \times \cdots \times \{0,1\}^{\mathbb{N}-\{1,2,\cdots\}}$ c $\liminf F_{k_1} \circ F_{k_2} \circ \cdots \circ F_{k_n}(\{0,1\}^{\mathbb{N}})$. If $x \notin \{k_1\}_1 \times \{k_2\}_2 \times \cdots \times \{0,1\}^{\mathbb{N}-\{1,2,\cdots\}}$, $x_{i_0} \neq k_{i_0}$ for some $i_0 \in \mathbb{N}$. Letting an open set $u = \{x_{i_0}\}_{i_0} \times \{0,1\}^{\mathbb{N}-\{i_0\}}$ of $\{0,1\}^{\mathbb{N}}$ containing x, it follows that for any infinite countable index set I, there exists $l \in I$ with $l > i_0$ such that $u \cap F_{k_1} \circ \cdots \circ F_{k_l}(X) = \emptyset$. Therefore, $\limsup F_{k_1} \circ F_{k_2} \circ \cdots \circ F_{k_n}(\{0,1\}^{\mathbb{N}}) \subset \{k_1\}_1 \times \{k_2\}_2 \times \cdots \times \{0,1\}^{\mathbb{N}-\{1,2,\cdots\}}$. For the detailed discussion about the limits, see A. Illanes and S. B. Nadler Jr., *Hyperspaces* (Marcel Dekker, New York, 1999).

First of all, we consider an arc *E* with two end points e_1 and e_2 as shown in (b) of Fig. 1. Setting a homeomorphism *h* from *E* to [0,1], a point $a \in E$ is mapped into $h(a) \in [0,1]$. With the aid of the representations (14) and (15) of [0,1], we can obtain the topological representation of *E* based on the contractions F_0 and F_1 of the Cantor set $\{0,1\}^{\mathbb{N}}$, as follows; if $h(a) \notin M (\equiv \{l/2^n; l = 1, \dots, 2^n - 1, n =$ $1,2,\dots\}), h(a)$ satisfies $h(a) = \sum_{i=1}^{\infty} k_i/2^i$ for some $k_1, k_2, \dots \in \{0,1\}$ and then

$$a \doteq \lim F_{k_1} \circ F_{k_2} \circ \dots \circ F_{k_n}(\{0,1\}^{\mathbb{N}}) = \{(k_1,k_2,\dots)\}.$$
(16)

If $h(a) \in M$, then

$$a \doteq \lim F_{k_1} \circ F_{k_2} \circ \dots \circ F_{k_{m-1}} \circ (F_1 \circ F_0^m(\{0,1\}^{\mathbb{N}}) \oplus F_0 \circ F_1^m(\{0,1\}^{\mathbb{N}})) = \{k^0, k^1\},$$
(17)

for some *m* and some $k_1, \ldots, k_{m-1} \in \{0, 1\}$ where $k^0 = (k_1, \ldots, k_{m-1}, 0, 1, 1, \ldots)$ and $k^1 = (k_1, \ldots, k_{m-1}, 1, 0, 0, \ldots)$. Here, \doteq stands for the sign of identification of *a* with a corresponding point $f^{-1}(a)$ of the decomposition space. By introducing the sign L_a defined as,

$$L_a \equiv \begin{cases} (16), & h(a) \notin M, \\ (17), & h(a) \in M, \end{cases}$$

$$\tag{18}$$

the representations (16) and (17) can be simplified as

$$a \doteq L_a \quad (a \in E). \tag{19}$$

Note that assuming $h(e_1) = 0$ and $h(e_2) = 1$, the end points e_1 and e_2 form $e_1 \doteq \lim F_0^n(\{0,1\}^{\mathbb{N}})$ and $e_2 \doteq \lim F_1^n(\{0,1\}^{\mathbb{N}})$, respectively. The representations (16) and (17) provide that each geometric element *a* of the arc *E* is described as the limit point of the composition F_0 and F_1 .

Next, we focus on a finite graph. Set Y_g to be a finite graph composed of r numbers of arcs E_1, \ldots, E_r $(r < \infty)$. To associate with each of arcs E_i , we construct a partition $\{X^1, \ldots, X^r\}$ of $\{0, 1\}^{\mathbb{N}}$ composed of the following contraction-cones :

$$\begin{cases} X^{1} = F_{0}(\{0,1\}^{\mathbb{N}}), \\ X^{i} = F_{1} \circ \cdots \circ F_{1} \circ F_{0}(\{0,1\}^{\mathbb{N}}) = F_{1}^{i-1} \circ F_{0}(\{0,1\}^{\mathbb{N}}) \ (i = 2, 3, \dots, r-1), \\ X^{r} = F_{1}^{r-1}(\{0,1\}^{\mathbb{N}}). \end{cases}$$
(20)

For each arc E_i , the topological representation is obtained as the form (19). Considering this fact, the topological representation of Y_g is

$$y \doteq F_1^{i-1} \circ F_0(L_y^i) \tag{21}$$

for a point y in an arc E_i , where L_y^i is defined by (18) for the homeomorphism h_i from E_i onto [0,1] instead of h, and

$$x \doteq \bigcup_{j=1}^{q} F_{1}^{t_{j}-1} \circ F_{0}(L_{x}^{t_{j}})$$
(22)

for a node *x* connecting with arcs E_{t_1}, \dots, E_{t_q} . In (21) and (22), L_y^i and $L_x^{t_j}$ are the terms that represent the position of *y* and *x* in the arcs E_i and E_{t_j} , respectively. $F_1^{i-1} \circ F_0$ shows that *y* is contained only in E_i , whereas, $\bigcup_{j=1}^q F_1^{t_j-1} \circ F_0$ shows that *x* is the node connecting with the arcs E_{t_1}, \dots, E_{t_q} . Note that (21) and (22) reproduce the representations of (1) in Sec. 1, which is obtained by the Cantor cube model $\{0,1\}^{\Lambda}$. Indeed, it is found that for $z \in Y_g$, the factors S_z^l and X^l (l = 1, ..., r) in (1) correspond to L_z^l and $F_1^l \circ F_0$ and therefore S_z^l and X^l can be characterized by the composition of F_0 and F_1 . To obtain the topological representation for the Cantor set $\{0, 1\}^{\mathbb{N}}$ from Cantor cube model $\{0, 1\}^{\Lambda}$, therefore, the transformations

$$S_z^l \to L_z^l, \quad X^l \to F_1^l \circ F_0$$
 (23)

are applicable. Then, the topological representations of a tree or a topological dendrite for $\{0,1\}^{\mathbb{N}}$ are also obtained from them for $\{0,1\}^{\Lambda}$ shown in the previous study[12].

The clusterized structure of network configuration can be realized by the topological space of the direct sum of finite graphs. We focus on this structure and denote it by $Y_c = (\bigoplus_{i=1}^{s} C_i, \bigoplus_{i=1}^{s} \tau_i)$, where (C_i, τ_i) is a finite graph composed of $E_1^i, \ldots, E_{r(i)}^i$ $(r(i) < \infty)$. To the disjoint clusters C_1, \ldots, C_s , the contraction-cones J_1, \ldots, J_s can be assigned, where

$$\begin{cases} J_1 = F_0(\{0,1\}^{\mathbb{N}}), \\ J_i = F_1^{i-1} \circ F_0(\{0,1\}^{\mathbb{N}}) \ (i = 2, 3, \dots, s-1), \\ J_s = F_1^{r-1}(\{0,1\}^{\mathbb{N}}). \end{cases}$$
(24)

Each finite graph C_i has the topological representation of the forms (21) and (22). Therefore, we obtain the representation for the whole space Y_c as follows; assuming that $a \in Y_c$ belongs to a cluster C_{i_0} , then

$$a \doteq F_1^{i_0 - 1} \circ F_0 \circ \begin{cases} \cup_{j=1}^q F_1^{i_j - 1} \circ F_0(L_a^{i_j}), \\ F_1^{i-1} \circ F_0(L_a^{i}), \end{cases}$$
(25)

where *a* is located either in the node connecting with the arcs $E_{t_1}^{i_0}, \ldots, E_{t_q}^{i_0}$ ($t_q \le r(i_0)$) or in the arc $E_i^{i_0}$. In the right hand side of (25), the first composition of the contractions $F_1^{i_0-1} \circ F_0$ is the term that indicates a graph C_{i_0} to which *a* belongs, and the successive compositions gives the positioning of *a* in the graph C_{i_0} .

For the special case where each cluster is composed of just one point, $C_i = \{x_i\}$, shown in Fig. 4 (b), we have $\bigoplus_{i=1}^{s} C_i = \bigcup_{i=1}^{s} \{x_i\} = \{x_1, \dots, x_s\}$. Then, by using the contraction-cones J_i $(j = 1, \dots, s)$ of (24), the topological representation is obtained as

$$x_i \doteq J_i \ (i = 1, \dots, s). \tag{26}$$

As an application of the topological representation for the clusterized pattern, we now consider a polycrystal filled with an *n* number of single crystals whose single crystal is supposed to form a specific geometric pattern, e.g., dendritic pattern. Such a situation is often found in solidification[19]. Fig. 5 (a) shows the sketch of such polycrystal where the dendritic structure characterizes its each single crystal. The topological representation of this type has been discussed in our studies, so far[11, 12]. In particular, by identifying this polycrystal with a kind of clusterized structure whose cluster is a tree (Fig. 5 (b)), we derived the topological representation using Cantor cube model, as follows. Each single crystal $Y_d^i (i = 1, ..., n)$ is represented by $\mathscr{D}_{Y_d^i} = \left\{ y \doteq J_i \cap \bigcup_{j=1}^q (X^{t_j} \cap S_y^{t_j}); y \in Y_d^i \right\} \cup \left\{ b \doteq J_i \cap X^j \cap S_b^j; b \in Y_d^i \right\}$ and their polycrystal is represented as the union of $\mathscr{D}_{Y_d^i}$,

$$\mathscr{D} = \bigcup_{i=1}^{n} \mathscr{D}_{Y_{d}^{i}}.$$
(27)

Note that $\mathscr{D}_{Y_d^i}$ and $\mathscr{D}_{Y_d^j}$ are mutually disjoint for $i \neq j$. With the aid of the transformations (23) and the contraction-cones (24), $\mathscr{D}_{Y_d^i}$ is reproduced as

$$\mathscr{D}_{Y_d^i} = \left\{ y \doteq F_1^{i-1} \circ F_0 \circ \left(\cup_{j=1}^q F_1^{t_j-1} \circ F_0(L_y^{t_j}) \right); y \in Y_d^i \right\} \cup \left\{ b \doteq F_1^{i-1} \circ F_0 \circ F_1^{j-1} \circ F_0(L_b^j); b \in Y_d^i \right\},$$
(28)

for i = 1, ..., n. (28) provides the representation of the dendritic single crystal characterized by the contraction F_0 and F_1 of the Cantor set, and (27) is the relationship of the single crystals to a whole polycrystal composed of them.



Figure 5: Schematic explanations of (a) a polycrystal tiled by dendritic single crystals and (b) each dendritic single crystal that is identified with a tree cluster. $\mathscr{D}_{Y_d^i}$ is given as the relation (28) that represents a tree cluster where y is a node and b is a point in an arc.

As we have shown in this section, each geometric pattern can be represented topologically by the Cantor set and its contractions, independently of the detail properties of each matter. The construction of these representations in our method is based on the topological and fractal properties of the Cantor set. In fact, the Cantor set has the topological properties of a 0-dim, perfect, compact Hausdorff-space, which supply the partition composed of cones. In addition, the contractions characterizing the self-similarity of the Cantor set make up a contraction-cone that is the key for representing each pattern topologically. So far, mathematical methods using topological and fractal viewpoints, such as the scaling method, have been contributed to the development of the study of pattern formation physics[16, 20]. Therefore, we believe that analyzing the Cantor set by our current method may have the potential to provide universal description of the geometric patterns of disordered matter.

5 Conclusion

We have investigated mathematical description that expresses universally the geometric patterns found in disordered matters from the viewpoint of the general topology. These geometric patterns can be characterized by the concept of continua, and focusing on the practical patterns such as a network, branching, and clusterized structures, we have successfully classified them under the topological graph, tree (or topological dendrite), and the direct sum of continua. To such topological spaces, we have associated the topological representations obtained in the basis of a Cantor set $(\{0,1\}^{\mathbb{N}}, \tau_d)$ with the contractions F_0 and F_1 characterizing the self-similarity. It is found that in the topological representations, each element of the geometric patterns can be identified with the limit of the compositions of F_0 and F_1 .

acknowledgments

The authors are grateful to Prof. S. Matsutani at Kanazawa University for useful comments and encouragements. This work was supported by Institute of Mathematics for Industry, Joint Usage/Research Center in Kyushu University. (FY2022 Workshop II "Geometry and Algebra in Material Science III" (2022a003)).

References

- N. E. Cusack, *The Physics of Structurally Disordered Matter: An Introduction* (Adam Hilger, London, 1987).
- [2] G. R. Strobl, Condensed Matter Physics: Crystals, Liquids, Liquid Crystals, and Polymers (Springer Berlin, Heidelberg, 2004).
- [3] Y. Tezuka and H. Oike, J. Am. Chem. Soc., 123 (2001), 11570.
- [4] Y. Katayama, T. Mizutani, W. Utsumi, O. Shimomura, M. Yamakata, and K. Funakoshi, Nature (London), 403 (2000), 170.
- [5] T. Morishita, Phys. Rev. Lett., 87 (2001), 105701.
- [6] W. Kurz and D. J. Fisher, *Fundamentals of Solidification* (Trans Tech Publication Ltd, Switzerland, 1998).
- [7] A. Hirata, K. Matsue, and M. Chen, Structural Analysis of Metallic Glasses with Computational Homology (Springer Tokyo, Tokyo, 2016)
- [8] Y. Hiraoka, T. Nakamura, A. Hirata, E. G. Escolar, K. Matsue, and Y. Nishiura, PNAS, 113 (2016), 7035.
- [9] A. Kitada and Y. Ogasawara, Chaos, Solitons & Fractals, 24 (2005), 785.
- [10] A. Kitada, Y. Ogasawara, and T. Yamamoto, Chaos, Solitons & Fractals, 34 (2007), 1732.
- [11] A. Kitada, S. Ohmori, and T. Yamamoto, J. Phys. Soc. Jpn., 85 (2016), 045001.
- [12] S. Ohmori, Y. Yamazaki, T. Yamamoto, and A. Kitada, Phys. Scr., 94 (2019), 105213.
- [13] S. B. Nadler Jr., Continuum Theory (Marcel Dekker, New York, 1992).
- [14] R. W. Vallin, *The Elements of Cantor sets With Applications* (John Wiley & Sons, New Jersey, 2013).
- [15] R. L. Devaney, An introduction to Chaotic Dynamical Systems, 2ed edition (CRC Press, Boca Raton, 2003).
- [16] J. Feder, Fractals (Plenum Press, New York, 1988).
- [17] R. Engelking, General Topology (Heldermann Verlag, Berlin, 1989).

- [18] M. Hata, Japan J. Appl. Math., 2 (1985), 381.
- [19] D. A. Porter and K. E. Easterling, *Phase Transformations in Metals and Alloys, 2nd ed.* (Chap-man&Hall, London, 1992).
- [20] M. Cross and H. Greenside, Pattern Formation and Dynamics in Nonequilibrium Systems (Cambridge University Press, Cambridge, 2009).