# EXPLICIT FORMULAS FOR MATRICES ASSOCIATED TO LADDER, CIRCULAR LADDER, AND MÖBIUS LADDER GRAPHS

ALI AZIMI AND M. FARROKHI D. G.

Abstract. We give explicit formulas for resistance distance matrices and Moore-Penrose inverses of incidence and Laplacian matrices of ladder, circular ladder, and Möbius ladder graphs. As a result, we compute the Kirchhoff index of these graphs and give new combinatorial formulas for the number of their spanning trees.

### 1. INTRODUCTION

The incidence matrix  $Q(\Gamma)$  of an oriented graph  $\Gamma$  with vertex-set  $V(\Gamma)$  =  $\{v_1, \ldots, v_n\}$  and edge-set  $E(\Gamma) = \{e_1, \ldots, e_m\}$  is an  $n \times m$  matrix defined as follows: The rows and columns of Q are indexed by vertices and edges of Γ, respectively. The  $(i, j)$  entry of Q is 0 if the vertex  $v_i$  and the edge  $e_j$  are not incident, otherwise it is 1 or  $-1$  depending on whether  $e_j$  starts from or ends at  $v_i$ . The Laplacian matrix  $L(\Gamma)$  of  $\Gamma$ , that is actually equal to  $Q(\Gamma)Q'(\Gamma)$ , is an  $n \times n$  matrix whose rows and columns are indexed by vertices of Γ. The  $(i, j)$  entry of  $L(\Gamma)$  is equal to the degree of the vertex  $v_i$  if  $i = j$ , and it is -1 or 0 if the vertices  $v_i$  and  $v_j$  are adjacent or non-adjacent, respectively.

The Moore-Penrose inverse of an  $m \times n$  real matrix A, denoted by  $A^+$ , is the (unique)  $n \times m$  real matrix that satisfies the following equations

$$
AA^{+}A = A
$$
,  $A^{+}AA^{+} = A^{+}$ ,  $(AA^{+})' = AA^{+}$ ,  $(A^{+}A)' = A^{+}A$ 

(see  $[9, 10]$  $[9, 10]$ ). The Moore-Penrose inverse of the incidence matrix of a graph was first studied by Ijira in  $[17]$ . We refer the interested reader to  $[1, 2, 3]$  $[1, 2, 3]$  $[1, 2, 3]$  $[1, 2, 3]$  for previous studies on the Moore-Penrose inverses of matrices associated to various classes of graphs.

Incidence matrices and their inverses, such as the Moore-Penrose inverses, are important tools used in analysis of graphs. Let Γ be a simple connected graph. The resistance distance between two vertices  $v_i$  and  $v_j$  can be computed via the formula

$$
r(v_i, v_j) = L^+_{v_i, v_i} + L^+_{v_j, v_j} - L^+_{v_i, v_j} - L^+_{v_j, v_i},
$$

where  $L_{v_i,v_j}^+$  denotes the  $(i, j)$ -entry of the Moore-penrose inverse  $L^+$  of  $L = L(\Gamma)$ . The notion of resistance distance between vertices of a graph is introduced by Klein and Randic in 1993 [\[19\]](#page-15-6) as the effective resistance between the given vertices (computed by Ohm's law) when a battery is attached between them assuming that all edges are unit resistors. Beside many applications in physics, chemistry, and

<sup>2000</sup> *Mathematics Subject Classification.* Primary 05C20, 05C50, Secondary 15A09.

*Key words and phrases.* Moore-Penrose inverse, resistance distance, Kirchhoff index, ladder graph, circular ladder graph, Möbius ladder graph.

network sciences (see [\[13,](#page-15-7) [18,](#page-15-8) [22\]](#page-15-9)), resistance distances can also be studied from a purely mathematical point of view as they represent new metrics on graph as well as making connections to graph theoretical objects like the Moore-Penrose inverse of Laplacian matrices above, see  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$  $[5, 7, 8, 12, 14, 15, 16]$ . A strictly related parameter is the so-called Kirchhoff index of the graph, namely

$$
Kf(\Gamma) = \frac{1}{2} \sum_{i=1}^{n} \sum_{j=1}^{n} r(v_i, v_j).
$$

In this paper, we consider three familiar classes of graphs, namely ladder, circular ladder, and Möbius graphs, and study their resistance distances and the Moore-Penrose inverse of their Laplacian matrices. Recall that the ladder graph  $L_n$  on 2n vertices is composed of two paths of length  $n - 1$  with spokes/rungs connecting corresponding vertices. The circular ladder graph  $CL_n$  defines analogous to ladder graphs with paths replaced by cycles. Also, the Möbius ladder graph  $M_n$  obtains from the circular ladder graph by switching two non-spoke edges of a square.

### 2. Preliminaries

In this section, we introduce some recursive sequences arising from the study of ladder, circular ladder, and Möbius ladder graphs.

**Definition.** By a  $(p, q)$ -sequence  $\{x_n\}$  we mean a sequence satisfying  $x_n = px_{n-1}$  $qx_{n-2}$  for all n provided that  $x_{n-1}$  and  $x_{n-2}$  are defined.

- ${a_n}_{n=0}^{\infty}$  is the  $(4, 1)$ -sequence [A001835](https://oeis.org/A001835) with initial values  $(a_0, a_1) = (1, 1)$ .
- $\{s_n\}_{n=0}^{\infty}$  is the  $(4, 1)$ -sequence [A001353](https://oeis.org/A001353) with initial values  $(s_0, s_1) = (0, 1)$ .

Lemma [2.1](#page-1-0) presents a list of identities satisfied by these sequences and will be used frequently in the rest of the paper.

<span id="page-1-0"></span>**Lemma 2.1.** *The following identities hold among sequences*  $\{a_n\}$  *and*  $\{s_n\}$ *:* 

<span id="page-1-1"></span>
$$
(1) \t s_n = \sum_{i=1}^n a_i,
$$

<span id="page-1-2"></span>(2) 
$$
2s_n = a_{n+1} - a_n,
$$

<span id="page-1-3"></span>(3) 
$$
a_{n+1} - a_n = a_{k+1}a_{n-k+1} - a_ka_{n-k},
$$

<span id="page-1-4"></span>(4) 
$$
s_n = a_k s_{n-k} + a_{n-k+1} s_k,
$$

<span id="page-1-5"></span>(5) 
$$
(a_{n+1} + a_n)^2 = 3(a_{n+1} - a_n)^2 + 4,
$$

<span id="page-1-6"></span>(6) 
$$
\sum_{i+j=n+1}^{n} a_i a_j = \frac{n(a_{n+1} + a_n) + (a_{n+1} - a_n)}{6} = \frac{(n+1)s_n + na_n}{3}
$$

*for all*  $n > k > 0$ *.* 

*Proof.* The equations [\(1\)](#page-1-1) and [\(2\)](#page-1-2) follow by induction on n and the equation [\(3\)](#page-1-3) follows by induction on k. Let  $\lambda := 2 + \sqrt{3}$ ,  $\alpha_a := (3 - \sqrt{3})/6$ ,  $\beta_a := (3 + \sqrt{3})/6$ , and  $\alpha_s = -\beta_s = \sqrt{3}/6$ . Then  $a_n$  and  $s_n$  are given by the Binet formulas

$$
a_n = \alpha_a \lambda^n + \beta_b \lambda^{-n}
$$
 and  $s_n = \alpha_s \lambda^n + \beta_s \lambda^{-n}$ .

The equations [\(4\)](#page-1-4) and [\(5\)](#page-1-5) can be obtained simply from the Binets formulas. Now, we prove equation [\(6\)](#page-1-6). Let  $b_n := \sum_{i=1}^n a_i a_{n+1-i}$  for all  $n \ge 1$ . First observe that

<span id="page-2-1"></span>
$$
b_n = \sum_{i=1}^n a_i a_{n+1-i} = \sum_{i=1}^n (4a_i a_{n-i} - a_i a_{n-1-i}),
$$

which implies that

(7) 
$$
b_n - 4b_{n-1} + b_{n-2} = a_n - a_{n-1}
$$

On the other hand, by equation [\(3\)](#page-1-3),

$$
b_n - b_{n-2} = a_1 a_n + ((a_2 a_{n-1} - a_1 a_{n-2}) + \dots + (a_{n-1} a_2 - a_{n-2} a_1)) + a_n a_1
$$
  
=  $2a_n + (n-2)(a_n - a_{n-1})$ 

so that

(8) 
$$
b_n - b_{n-2} = na_n - (n-2)a_{n-1}.
$$

Summing up the equations  $(8)$  when n is replaced by  $3, \ldots, n$ , it follows that

(9) 
$$
b_n + b_{n-1} = na_n + s_{n-1}.
$$

Now, by solving the system of linear equations [\(7\)](#page-2-1), [\(8\)](#page-2-0), and [\(9\)](#page-2-2) for  $b_n$ ,  $b_{n-1}$ , and  $b_{n-2}$ , the result follows.

 $\Box$ 

## <span id="page-2-2"></span><span id="page-2-0"></span>3. ladder graph

Let  $L_n$  denote the ladder graph of order  $2n$ . Suppose vertices along the top and bottom of  $L_n$ , as depicted in Fig. [1,](#page-2-3) are labeled  $u_1^+, \ldots, u_n^+$  and  $u_1^-, \ldots, u_n^-$ , respectively. Also, suppose the edges of  $L_n$  are oriented as  $e_i^{\varepsilon} = (u_i^{\varepsilon}, u_{i+1}^{\varepsilon})$  for  $i = 1, \ldots, n - 1$  with  $\varepsilon = \pm 1$ , and  $f_i = (u_i^+, u_i^-)$  for  $i = 1, \ldots, n$ , where  $f_i$  are the spokes.



<span id="page-2-3"></span>FIGURE 1. Ladder graph  $L_n$ 

Let  $S(\Gamma)$  denote the set of all spanning trees of the underlying graph of  $\Gamma$  and  $s(\Gamma) := \#S(\Gamma)$ . The graph obtained from  $\Gamma$  by contracting edge e is denoted by  $\Gamma/e$ . It is known from [\[20,](#page-15-17) p.35] that  $s(L_n) = 4s(L_{n-1}) - s(L_{n-2})$  for all  $n \geq 2$ . Since,  $s(L_0) = 0$  and  $s(L_1) = 1$ , we observe that  $s(L_n) = s_n$  for all  $n \geq 0$ .

Let  $Q = [q_{u,e}]$  be the incidence matrix of  $L_n$  with the orientation given in Fig. [1](#page-2-3) whose rows and columns are indexed by ordered sets

$$
V(L_n) = \{u_1^+, \ldots, u_n^+, u_1^-, \ldots, u_n^-\}
$$

and

$$
E(L_n) = \{f_1, \ldots, f_n, e_1^+ \ldots, e_{n-1}^+, e_1^- \ldots, e_{n-1}^-\},\
$$

respectively. Also, let

(10) 
$$
H = \begin{pmatrix} B & -B \\ C & D \\ D & C \end{pmatrix},
$$

be the  $(3n-2) \times 2n$  matrix defined as follows: The rows and columns of H are indexed by ordered sets  $E(L_n)$  and  $V(L_n)$ , respectively. Let  $B = [b_{i,j}]$  be the  $n \times n$ symmetric matrix given by  $b_{i,j} = a_i a_{n-j+1}/2s_n$ , and  $C = [c_{i,j}]$  and  $D = [d_{i,j}]$  be the  $(n - 1) \times n$  matrices given by

$$
c_{i,j} = \begin{cases} \delta_{i,j} - \frac{i}{2n} - \frac{a_{n-j+1}s_i}{2s_n}, & i \leq j, \\ \frac{1}{2} - \frac{i}{2n} + \frac{a_j s_{n-i}}{2s_n}, & i > j, \end{cases}
$$

and

$$
d_{i,j} = \begin{cases} -\frac{i}{2n} + \frac{a_{n-j+1}s_i}{2s_n}, & i \leq j, \\ \frac{1}{2} - \frac{i}{2n} - \frac{a_j s_{n-i}}{2s_n}, & i > j, \end{cases}
$$

respectively. In the following series of lemmas, we provide the machinery to prove that  $H$  is the Moore-Penrose inverse of  $Q$ .

## <span id="page-3-2"></span>**Lemma 3.1.**  $H1 = 0$ .

*Proof.* It is obvious from the definition that  $(C+D)\mathbf{1} = 0$ . Thus

$$
H\mathbf{1} = \begin{bmatrix} (B-B)\mathbf{1} \\ (C+D)\mathbf{1} \\ (D+C)\mathbf{1} \end{bmatrix} = 0,
$$

<span id="page-3-0"></span>as required  $\Box$ 

# Lemma 3.2.  $QH = I - \frac{1}{2n}J$ . *Proof.* Let  $\delta'_{i,j} := 1 - \delta_{i,j}$ . For  $1 \leq i \leq j \leq n$ , we have

$$
(QH)_{u_i^+, u_j^+} = b_{i,j} - \delta'_{i,1} c_{i-1,j} + \delta'_{i,n} c_{i,j}
$$
  
= 
$$
\frac{a_i a_{n-j+1}}{2s_n} - \delta'_{i,1} \left( -\frac{i-1}{2n} - \frac{a_{n-j+1} s_{i-1}}{2s_n} \right) + \delta'_{i,n} \left( \delta_{i,j} - \frac{i}{2n} - \frac{a_{n-j+1} s_k}{2s_n} \right)
$$
  
= 
$$
\delta_{i,j} - \frac{1}{2n},
$$

by using equation [\(1\)](#page-1-1) when  $i = n$ . Also, an analogous argument yields

$$
(QH)_{u_j^+, u_i^+} = (QH)_{u_i^-, u_j^-} = (QH)_{u_j^-, u_i^-} = \delta_{i,j} - \frac{1}{2n}.
$$

Similarly, by discussing the cases where  $i = 1$ ,  $i = n$ , and  $i \neq 1, n$  and either  $i \leq j$ ,  $i = j + 1$ , and  $i > j + 1$ , we obtain

$$
(QH)_{u_i^+, u_j^-} = (QH)_{u_i^-, u_j^+} = -b_{i,j} - \delta_{1,i}' d_{i-1,j} + \delta_{i,n}' d_{i,j} = -\frac{1}{2n}
$$

<span id="page-3-1"></span>for all  $1 \le i, j \le n$ . The proof is complete.

Lemma 3.3. HQ *is a symmetric matrix.*

*Proof.* Let  $1 \leq i, j \leq n$ . Since B is a symmetric matrix,

$$
(HQ)_{f_i, f_j} = 2b_{i,j} = 2b_{j,i} = (HQ)_{f_j, f_i}.
$$

From the definition of  $H$ , we get

$$
(HQ)_{e_i^+, f_j} = -(HQ)_{e_i^-, f_j} = c_{i,j} - d_{i,j} = \begin{cases} \delta_{i,j} - \frac{a_{n-j+1}s_i}{s_n}, & i \leq j, \\ \frac{a_j s_{n-i}}{s_n}, & i > j, \end{cases}
$$

and

$$
(HQ)_{f_j, e_i^+} = -(HQ)_{f_j, e_i^-} = b_{j,i} - b_{j,i+1} = \begin{cases} \frac{a_i a_{n-j+1} - a_{i+1} a_{n-j+1}}{2s_n}, & i < j, \\ \frac{a_j a_{n-i+1} - a_j a_{n-i}}{2s_n}, & i \ge j. \end{cases}
$$

Utilizing equations  $(1)$ ,  $(2)$ , and  $(4)$  and a simple case-by-case analysis, it yields  $(HQ)_{e_i^+, f_j} = (HQ)_{f_j, e_i}$  and  $(HQ)_{e_i^-, f_j} = (HQ)_{f_j, e_i^-}$ . On the other hand,

$$
(HQ)_{e_i^+, e_j^+} = c_{i,j} - c_{i,j+1} = \begin{cases} \delta_{i,j} + \frac{(a_{n-j} - a_{n-j+1})s_i}{2s_n}, & i \leq j, \\ \frac{(a_j - a_{j+1})s_{n-i}}{2s_n}, & i > j+1, \\ -\frac{1}{2} + \frac{a_j s_{n-i} + a_{n-j} s_i}{2s_n}, & i = j+1, \end{cases}
$$

$$
(HQ)_{e_i^+, e_j^-} = d_{i,j} - d_{i,j+1} = \begin{cases} \frac{(a_{n-j+1} - a_{n-j})s_i}{2s_n}, & i \leq j, \\ \frac{1}{2} - \frac{a_j s_{n-i} + a_{n-j} s_i}{2s_n}, & i = j+1, \\ \frac{(a_{j+1} - a_j)s_{n-i}}{2s_n}, & i > j+1, \end{cases}
$$

$$
(HQ)_{e_j^-,e_i^+} = c_{j,i} - c_{j,i+1} = \begin{cases} \frac{(a_i - a_{i+1})s_{n-j}}{2s_n}, & j > i+1, \\ -\frac{1}{2} + \frac{a_i s_{n-j} + a_{n-i} s_j}{2s_n}, & j = i+1, \\ \frac{(a_{n-i} - a_{n-i+1})s_j}{2s_n}, & i \ge j \end{cases}
$$

and

$$
(HQ)_{e_i^-,e_j^-} = d_{i,j} - d_{i,j+1} = \begin{cases} \frac{(a_{n-j+1} - a_{n-j})s_i}{2s_n}, & i \leq j, \\ \frac{1}{2} - \frac{a_j s_{n-i} + a_{n-j} s_i}{2s_n}, & i = j+1, \\ \frac{(a_{j+1} - a_j)s_{n-i}}{2s_n}, & i > j+1, \end{cases}
$$

<span id="page-4-0"></span>from which the result follows.  $\Box$ 

**Theorem 3.4.** Let  $L_n$  be a ladder graph with incident matrix Q. Then  $Q^+ = H$ .

*Proof.* By Lemmas [3.2](#page-3-0) and [3.3,](#page-3-1) QH and HQ are symmetric. On the other hand,  $QHQ = Q$  for  $\mathbf{1}'Q = 0$ . Since  $HQH = H$  by Lemmas [3.1](#page-3-2) and [3.2,](#page-3-0) it follows that  $H = Q^+$ .

The following lemma will be used frequently in the reminder of this section.

<span id="page-4-1"></span>Lemma 3.5 ([\[4,](#page-15-18) Corollary 3.2]). *Let* Γ *be a connected graph and* P *be a path of length* d *between vertices*  $u, v \in V(\Gamma)$ *. If all edges in* P *have the same direction from* u *to* v *then,*

$$
r(u, v) = \sum_{e \in E(P)} (q_{e,u}^+ - q_{e,v}^+),
$$

*where*  $r(u, v)$  *is the resistance distance between vertices u and v*.

**Corollary 3.6.** *If*  $\mathcal{B} = 2s(L_n)B$ . Then  $\mathcal{B}$  has eigenvalue  $s(L_n)$  with associated *eigenvector* 1*. Also, the diagonal entries of* B *satisfy the equations*

$$
\mathcal{B}_{i,i}=s(L_n/f_i)
$$

*for*  $i = 1, ..., n$ *.* 

*Proof.* By [\[3,](#page-15-5) Lemma 2.1]

$$
\sum_{i=1}^{n} q_{f_i, u_j^+}^+ = 1 - \frac{n}{2n} = \frac{1}{2},
$$

On the other hand, by Theorem [3.4,](#page-4-0)

$$
\sum_{i=1}^{n} q_{f_i, u_j^+}^+ = \frac{a_i \sum_{i \le j} a_{n-j+1} + a_{n-i+1} \sum_{i > j} a_j}{2s(L_n)}
$$

so that

$$
s(L_n) = a_i \sum_{i \le j} a_{n-j+1} + a_{n-i+1} \sum_{i > j} a_j = \sum_{j=1}^n b_{i,j},
$$

that is  $\mathcal{B}1 = s(L_n)1$ . Now, from Lemma [3.5](#page-4-1) and Theorem [3.4,](#page-4-0) and the definition of the resistance distance between two vertices, we get

<span id="page-5-1"></span>
$$
\frac{s(L_n/f_i)}{s(L_n)} = r(u_i^+, u_i^-) = 2q_{f_i, u_i^+}^+ = \frac{a_i a_{n-i+1}}{s(L_n)}
$$

(see  $[6, p. 133]$ ). Therefore

(11) 
$$
s(L_n/f_i) = a_i a_{n-i+1} = B_{i,i},
$$

as required.  $\square$ 

In [\[11\]](#page-15-20) the author uses circus reduction to obtain the resistance distance and Kirchhoff index of ladder graphs. Here we use the Moore-Penrose inverses of Laplacian matrices of ladder graphs to drive new formulas for resistance distance and Kirchhoff index of ladder graphs.

<span id="page-5-0"></span>**Theorem 3.7.** Let  $R = [r(u, v)]$  be the resistance distance matrix of the ladder *graph* L<sup>n</sup> *of order* 2n*. Then*

$$
r(u_i^{\varepsilon}, u_j^{\varepsilon'}) = \frac{d}{2} - \varepsilon \varepsilon' \frac{a_i a_{n-j+1}}{2s_n} + \alpha(i, j),
$$

 $where d = d(u_i^+, u_j^+), \, \varepsilon, \varepsilon' = \pm 1, \, and \, \alpha(i, j) = (s(L_n/f_i) + s(L_n/f_j))/4s_n.$ 

*Proof.* Let  $d = j - i = d(u_i^+, u_j^+)$  and  $P : e_i^+, \ldots, e_{j-1}^+$  be the shortest path between  $u_i^+$  and  $u_j^+$ , where  $1 \leq i < j \leq n$ . Then

$$
\sum_{k=i}^{j-1} q^+_{e^+_k, u^+_i} = \sum_{k=i}^{j-1} c_{k,i} = c_{i,i} + \sum_{k=i+1}^{j-1} c_{k,i},
$$

Thus

$$
\sum_{k=i}^{j-1} q^+_{e^+_k, u^+_i} = \left(1 - \frac{i}{2n} - \frac{a_{n-i+1}s_i}{2s_n}\right) + \sum_{k=i+1}^{j-1} \left(\frac{1}{2} - \frac{k}{2n} + \frac{a_i s_{n-k}}{2s_n}\right).
$$

By equations  $(2)$  and  $(3)$ ,

$$
\sum_{k=i}^{j} q_{e_k^+,u_i^+}^+ = \frac{d+1}{2} - \frac{id}{2n} - \frac{d(d-1)}{4n} + \frac{a_i(a_{n-i} - a_{n-j+1}) - a_{n-i+1}(a_{i+1} - a_i)}{4s_n}
$$
  
=  $\frac{d+1}{2} - \frac{id}{2n} - \frac{d(d-1)}{4n} + \frac{a_n - a_{n+1} + a_i a_{n-i+1} - a_i a_{n-j+1}}{4s_n}$   
=  $\frac{d+1}{2} - \frac{id}{2n} - \frac{d(d-1)}{4n} + \frac{-2s_n + s(L_n/f_i) - a_i a_{n-j+1}}{4s_n}$   
=  $\frac{d}{2} - \frac{id}{2n} - \frac{d(d-1)}{4n} - \frac{a_i a_{n-j+1}}{4s_n} + \frac{s(L_n/f_i)}{4s_n}.$ 

The same argument for the vertex  $u_j^+$  yields

$$
\sum_{k=i}^{j-1} q_{e_k^+, u_j^+}^+ = \sum_{k=i}^{j-1} c_{k,j} = -\frac{id}{2n} - \frac{d(d-1)}{4n} + \frac{a_{n-j+1}(a_i - a_{d+i})}{4s_n}
$$

$$
= -\frac{id}{2n} - \frac{d(d-1)}{4n} + \frac{a_i a_{n-j+1}}{4s_n} - \frac{s(L_n/f_j)}{4s_n}.
$$

Therefore,

$$
r(u_i^+, u_j^+) = \frac{d}{2} - \frac{a_i a_{n-j+1}}{2s_n} + \frac{s(L_n/f_i) + s(L_n/f_j)}{4s_n}.
$$

From the block structure of H it follows that  $r(u_i^-, u_j^-) = r(u_i^+, u_j^+)$ . Finally, from the equality

$$
\sum_{k=i}^{j-1} q_{e_k^+, u_j^-}^+ = \sum_{k=i}^{j-1} d_{k,j} = \sum_{k=i}^{j-1} \left( -\frac{k}{2n} + \frac{a_{n-j+1} s_k}{2s_n} \right)
$$
  
= 
$$
-\frac{id}{2n} - \frac{d(d-1)}{4n} - \frac{a_i a_{n-j+1}}{4s_n} + \frac{s(L_n/f_j)}{4s_n}
$$

we obtain

$$
r(u_j^-, u_i^+) = r(u_i^+, u_j^-) = \sum_{k=i}^{j-1} q_{e_k^+, u_i^+}^+ + b_{j,i} - \sum_{k=i}^{j-1} q_{e_k^+, u_j^-}^+ + b_{j,j}
$$
  
= 
$$
\frac{d}{2} + \frac{a_i a_{n-j+1}}{2s_n} + \frac{s(L_n/f_i) + s(L_n/f_j)}{4s_n},
$$

as required.  $\hfill \square$ 

Utilizing Theorem [3.7,](#page-5-0) we can compute the Kirchhoff index of ladder graphs.

Corollary 3.8. For any  $n \geq 1$ ,

$$
Kf(L_n) = \frac{n^2}{3} \left( n + 1 + \frac{a_n}{s_n} \right).
$$

*Proof.* We have

$$
Kf(L_n) = \sum_{\{u,v\} \subseteq V(L_n)} r(u,v)
$$
  
= 
$$
\frac{4n \sum_{i=1}^n s(L_n/f_i)}{4s_n} + 2Kf(P_n)
$$
  
= 
$$
\frac{n \sum_{i=1}^n a_i a_{n+1-i}}{s_n} + \frac{n(n^2 - 1)}{3}
$$

from which the result follows by equation  $(6)$ .

## 4. Circular ladder graph

In this section, we compute the Moore-Penrose inverse of incidence matrices of circular ladder graphs. Accordingly, we obtain the resistance distance matrix and Kirchhoff index of circular ladders. In what follows,  $CL_n$  denotes the circular ladder graph of order  $2n$  and  $u_1^+, \ldots, u_n^+$  and  $u_1^-, \ldots, u_n^-$  stand for the vertices of the internal and external cycles of  $CL_n$ , respectively, as in Fig. [2.](#page-7-0) Also,  $u_{i+kn}^{\varepsilon}$ denotes the vertex  $u_i^{\varepsilon}$  for all  $1 \leq i \leq n$ , integer k, and  $\varepsilon = \pm 1$ .



<span id="page-7-0"></span>FIGURE 2. Circular ladder  $CL_n$ 

Let  $\Gamma$  be a planar graph embedded on a shaded cylinder as in Fig. [3.](#page-8-0) A Jordan path is a smooth simple path starting from outer border and ending at the inner border such that it meets every face and edge at most once and never paths through vertices (see [\[20\]](#page-15-17)). Removing edges crossed by a Jordan path  $\mathcal J$  results in a graph whose spanning trees depend only on  $\mathcal J$ . On the other hand, every spanning tree of Γ determines a Jordan path. Therefore, the set of all spanning trees of Γ can be partitioned into sets  $S_{\mathcal{J}}(\Gamma)$  where each set  $S_{\mathcal{J}}(\Gamma)$  corresponds to spanning trees associated to a Jordan path J. Let  $s_{\mathcal{J}}(\Gamma) := \#S_{\mathcal{J}}(\Gamma)$ .

<span id="page-7-1"></span>**Theorem 4.1.** Let  $CL_n$  be the circular ladder graph of order  $2n$ . If f is a spoke, *then*

$$
s(CL_n/f) = ns(L_n).
$$

*Proof.* Consider a natural embedding of circular ladder  $CL_n$  on a cylinder as in Fig. [3](#page-8-0) and assume without loss of generality that  $f = u_1^+ u_1^-$ . We have two (equivalent up to reflection) classes of Jordan paths:



<span id="page-8-0"></span>Figure 3. Jordan path

- (1) Jordan paths  $\mathcal{J}_{i,j}^1$  through edges  $u_i^+ u_{i+1}^+ u_{i+1}^- u_{i+1}^- \ldots, u_{j-1}^+ u_{j-1}^- u_j^-$  for  $1 \le i < j \le n+1$ ;
- (2) Jordan paths  $\mathcal{J}_{i,j}^2$  through edges  $u_j^+ u_{j-1}^+ u_{j-1}^+ u_{j-1}^- \ldots, u_{i+1}^+ u_{i+1}^- u_i^- u_{i+1}^$ for  $1 \le i < j \le n+1$ .

Note that the Jordan paths in parts (1) and (2) coincide when  $j = i + 1$ . Also  $s_{\mathcal{J}_{i,j}^1}(CL_n/f) = s_{\mathcal{J}_{i,j}^2}(CL_n/f)$ . Utilizing the above Jordan paths in conjunction with equation  $(11)$ , yields

$$
s(CL_n/f) = 2 \sum_{1 \le i < j \le n+1} s_{\mathcal{J}_{i,j}^1}(CL_n/f) - \sum_{i=1}^n s_{\mathcal{J}_{i,i+1}^1}(CL_n/f)
$$
\n
$$
= 2 \sum_{1 \le i < j \le n+1} s(L_{n+1+i-j}/f_i) - \sum_{i=1}^n s(L_n/f_i)
$$
\n
$$
= 2 \sum_{1 \le i < j \le n+1} a_i a_{n+2-j} - \sum_{i=1}^n a_i a_{n+1-i}
$$
\n
$$
= 2 \sum_{2 \le j \le n+1} s_{j-1} a_{n+2-j} - \sum_{i=1}^n a_i a_{n+1-i}
$$
\n
$$
= 2 \sum_{1 \le i \le n} a_i s_{n+1-i} - \sum_{i=1}^n a_i a_{n+1-i},
$$

where  $L_s/f_t$  denotes the contraction of the t's spoke of  $L_s$  from right or left. Let  $x_n := \sum_{i=1}^n a_i a_{n+1-i}$  and  $y_n := \sum_{i=1}^n a_i s_{n+1-i}$  for all  $n \ge 1$ . Clearly,  $y_n - y_{n-1} =$  $x_n$ . Also, from equation [\(4\)](#page-1-4), it follows that  $y_n + y_{n-1} = ns_n$ . Thus  $2y_n = ns_n + x_n$ , which implies that

<span id="page-8-1"></span>as required.  $\Box$ 

$$
s(CL_n/f) = 2y_n - x_n = ns_n,
$$

**Theorem 4.2.** Let  $CL_n$  be the circular ladder graph of order  $2n$ . Then

$$
s(CL_n) = \frac{1}{2}n(a_{n+1} + a_n - 2).
$$

*Proof.* Analogous to the proof of Theorem [4.1,](#page-7-1) we consider two families of (equivalent up to reflection) Jordan paths given by  $\mathcal{J}_{i,j}^1$  and  $\mathcal{J}_{i,j}^2$  as in Theorem [4.1](#page-7-1) where

i is behind j regarding clockwise rotations. From the rotational symmetry of  $CL_n$ and the equality of  $\mathcal{J}_{i,j}^1$  and  $\mathcal{J}_{i,j}^2$  for consecutive values of i and j, it follows that

$$
s(CL_n) = 2n \sum_{i=1}^{n} s(L_i) - ns(L_n) = \frac{1}{2}n(a_{n+1} - a_n - 2)
$$
  
by equation (2).

In what follows,  $e_i^+$ ,  $e_i^-$ , and  $f_i$  stand for directed edges  $(u_i^+, u_{i+1}^+)$ ,  $(u_i^-, u_{i+1}^-)$ , and  $(u_i^+, u_i^-)$  for  $i = 1, \ldots, n$ . Accordingly, we consider the orientation of  $CL_n$ whose directed edges are  $e_i^+, e_i^-,$  and  $f_i$  for  $i = 1, ..., n$ .

<span id="page-9-2"></span>**Theorem 4.3.** Let  $CL_n$  be the oriented circular ladder graph of order  $2n$  with *incidence matrix Q. The Moore-Penrose inverse*  $Q^+ = [q_{e,u}^+]$  *of*  $Q$  *is given by* 

$$
q_{e_{i+t}^{\varepsilon},u_i^{\varepsilon}}^+ = -\varepsilon \varepsilon' \frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \varepsilon \varepsilon' \frac{a_{t+1} + \varepsilon \varepsilon' 1}{4} - \frac{2t + 1}{4n},
$$
  

$$
\varepsilon q_{f_{i+t},u_i^{\varepsilon}}^+ = \frac{a_{t+1} + a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \frac{a_{t+1} - a_t}{4}
$$

*for all*  $1 \leq i \leq n$ ,  $t = 0, \ldots, n-1$ , and  $\varepsilon, \varepsilon' = \pm 1$ .

*Proof.* Let  $d = d(f_i, u_j)$ . From the symmetry of graph, it follows that  $q_t^+$ .  $^{+}_{f_{i},u^{-}_{i}} =$  $-q_{f_i}^+$  $t_{i}$ <sub>t<sub>i</sub>, $u_i$ <sup>+</sup></sub>. Hence

$$
2q_{f_i, u_i^+}^+ = r(u_i^+, u_i^-) = \frac{s(CL_n/f_i)}{s(CL_n)} = \frac{s_n}{a_{n+1} + a_n - 2}
$$

by Lemma [3.5](#page-4-1) and Theorems [4.1](#page-7-1) and [4.2.](#page-8-1)

Consider the edge cut-set  $\{e_j^+, f_{j+1}, e_{j+1}^+\}$ , where  $j = i+t$  with  $0 \le t < n$ . From [\[3,](#page-15-5) Lemma 2.1], it follows that

(12) 
$$
q_{e_{j+1}^+, u_i^+}^+ = \delta_{i+1,j} - \frac{1}{2n} - q_{f_{j+1}, u_i^+}^+ + q_{e_j^+, u_i^+}^+.
$$

Analogously, by considering the edge cut-set  $\{e_j^-,f_{j+1},e_{j+1}^-\}$ , we obtain

(13) 
$$
q_{e_{j+1},u_i^+}^+ = -\frac{1}{2n} + q_{f_{j+1},u_i^+}^+ + q_{e_j^-,u_i^+}^+.
$$

for all  $j \geq i$ . If we replace j by  $i - 1$ , and use the equalities

<span id="page-9-0"></span>
$$
q^+_{e^+_{i-1},u^+_i} = - q^+_{e^+_{i},u^+_i} \quad \text{and} \quad q^+_{e^-_{i-1},u^+_i} = - q^+_{e^-_{i},u^+_i}
$$

obtained from the symmetry of the graph, then we get the initial values

$$
q_{e_i^+, u_i^+}^+ = \frac{1}{2} \left( 1 - \frac{1}{2n} - q_{f_i, u_i^+}^+ \right) \quad \text{and} \quad q_{e_i^-, u_i^+}^+ = \frac{1}{2} \left( \frac{-1}{2n} + q_{f_i, u_i^+}^+ \right).
$$

On the other hand, the incidence vector of the cycle induced by  $\{u_j^+, u_{j+1}^+, u_j^-, u_{j+1}^-\}$ lies in the left null space of  $Q^+$ , from which we obtain

<span id="page-9-1"></span>
$$
q_{f_{j+1},u_i^+}^+ - q_{e_j^-,u_i^+}^+ - q_{f_j,u_i^+}^+ + q_{e_j^+,u_i^+}^+ = 0
$$

so that

(14) 
$$
q_{f_{j+1},u_i^+}^+ = q_{e_j^-,u_i^+}^+ + q_{f_j,u_i^+}^+ - q_{e_j^+,u_i^+}^+
$$

for all  $j \geq i$ . Let

$$
C := \begin{pmatrix} 2 & -1 & -1 \\ -1 & 2 & 1 \\ -1 & 1 & 1 \end{pmatrix} \text{ and } D := -\frac{1}{2n} \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.
$$

If  $X_j$  denotes the column matrix  $(q_{\rho}^+)$  $e^+_{j}, u^+_i q^+_{e^-_j}$  $e_{j}^{+}, u_{i}^{+}$   $q_{f_{j}}^{+}$  $(\sum_{j=1}^{j+1} a_j + i)^T$  for  $i \leq j < i+n$ , then from the equalities  $(13)$  and  $(14)$ , it follows that

$$
X_{j+1} = CX_j + D
$$

for all  $i \leq j < i + n$ . Using induction on  $t \geq 0$ , one can prove that

$$
C^{t} = \begin{pmatrix} \frac{a_{t+1}+1}{2} & -\frac{a_{t+1}-1}{2} & -s_t\\ -\frac{a_{t+1}-1}{2} & \frac{a_{t+1}+1}{2} & s_t\\ -s_t & s_t & a_t \end{pmatrix}
$$

and

$$
I + C + \dots + C^{t-1} = \begin{pmatrix} \frac{s_t + t}{2} & -\frac{s_t - t}{2} & -\frac{a_t - 1}{2} \\ -\frac{s_t - t}{2} & \frac{s_t + t}{2} & \frac{a_t - 1}{2} \\ -\frac{a_t - 1}{2} & \frac{a_t - 1}{2} & s_{t-1} + 1 \end{pmatrix}.
$$

Now, from the equality

$$
X_{i+t} = C^t X_i + (I + C + \dots + C^{t-1})D,
$$

it follows that

$$
q_{e_{i+t}^+,u_i^+}^+ = -\frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \frac{a_{t+1} + 1}{4} - \frac{2t + 1}{4n},
$$
  
\n
$$
q_{e_{i+t}^+,u_i^+}^+ = \frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \frac{a_{t+1} - 1}{4} - \frac{2t + 1}{4n},
$$
  
\n
$$
q_{f_{i+t},u_i^+}^+ = \frac{a_{t+1} + a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \frac{a_{t+1} - a_t}{4}
$$

for all  $t = 0, ..., n - 1$ . Finally, the result follows by applying the same argument or using the symmetry of graph for vertices  $u_1, ..., u_m$ . or using the symmetry of graph for vertices  $u_1^-, \ldots, u_n^ \overline{n}$ .

Corollary 4.4. *If*  $n = 2k + 1$ *, then* 

$$
s(CL_n) = ns(L_n) \left( \frac{s(L_{k+1}) + s(L_k)}{s(L_{k+1}) - s(L_k)} \right)
$$

*and if*  $n = 2k$ *, then* 

$$
s(CL_n) = ns(L_n) \left( \frac{s(L_{k+1}) + 2s(L_k) + s(L_{k-1})}{s(L_{k+1}) - s(L_{k-1})} \right).
$$

*Proof.* We know that  $s(CL_n/f_i)/s(CL_n) = r(u_i^+, u_i^-) = 2q_{f_i}^+$  $t_{i}$ <sub>t<sub>i</sub>, $u_i^+$ . Thus</sub>

$$
s(CL_n) = ns(L_n) \cdot \frac{a_{n+1} + a_n - 2}{a_{n+1} - a_n}.
$$

Using Binet's formulas, one can easily show that

$$
\frac{a_{n+1} + a_n - 2}{a_{n+1} - a_n} = \frac{s_{k+1} + s_k}{s_{k+1} - s_k} \quad \text{or} \quad \frac{s_{k+1} + 2s_k + s_{k-1}}{s_{k+1} - s_k}
$$

according to  $n = 2k + 1$  or  $n = 2k$ , respectively. The result follows.

Corollary 4.5. *Let*  $1 \le i \le n$ ,  $0 \le t < n$ , and  $\varepsilon, \varepsilon' = \pm 1$ *. Then* 

$$
r(u_i^{\varepsilon}, u_{i+t}^{\varepsilon'}) = -\varepsilon \varepsilon' \frac{a_{t+1} + a_t - \varepsilon \varepsilon' 2}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \varepsilon \varepsilon' \frac{a_{t+1} - a_t}{4} + \frac{t}{2} - \frac{t^2}{2n}.
$$

*Proof.* First observe that  $r(u_i^-, u_{i+t}^-) = r(u_i^+, u_{i+t}^+) = r(u_0^+, u_t^+)$  by Theorem [4.3](#page-9-2) and Lemma [3.5.](#page-4-1) We have

$$
\sum_{s=0}^{t-1} q^+_{e^+_s, u^+_0} = -\frac{a_{t+1} + a_t - 2}{8} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \frac{a_{t+1} - a_t}{8} + \frac{t}{4} - \frac{t^2}{4n}
$$

and

$$
\sum_{s=0}^{t-1} q^+_{e^+_{s}, u^-_0} = \frac{a_{t+1} + a_t - 2}{8} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \frac{a_{t+1} - a_t}{8} + \frac{t}{4} - \frac{t^2}{4n}.
$$

Also,

$$
\sum_{s=0}^{t-1} q^+_{e^+_{s}, u^+_{t}} = \sum_{s=0}^{t-1} q^+_{e^+_{n+s-t}, u^+_{0}} = -\sum_{s=0}^{t-1} q^+_{e^+_{s}, u^+_{0}}
$$

and

$$
\sum_{s=0}^{t-1} q^+_{e^+_s,u^-_t} = \sum_{s=0}^{t-1} q^+_{e^+_{n+s-t},u^-_0} = - \sum_{s=0}^{t-1} q^+_{e^+_s,u^-_0}
$$

by symmetry of the graph. Let  $1 \leq i \leq n$  and  $0 \leq t < n$ . Then, by Lemma [3.5,](#page-4-1)

$$
r(u_i^+, u_{i+t}^+) = r(u_0^+, u_t^+) = 2 \sum_{s=0}^{t-1} q_{e_s^+, u_0^+}^+
$$
  
=  $-\frac{a_{t+1} + a_t - 2}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \frac{a_{t+1} - a_t}{4} + \frac{t}{2} - \frac{t^2}{2n}$ 

and

$$
r(u_i^+, u_{i+t}^-) = r(u_0^+, u_t^-) = \sum_{s=0}^{t-1} q_{e_s^+, u_0^+}^+ - \sum_{s=0}^{t-1} q_{e_s^+, u_t^-}^+ + q^+(f_t, u_0^+) - q^+(f_t, u_t^-)
$$
  

$$
= \sum_{s=0}^{t-1} q_{e_s^+, u_0^+}^+ + \sum_{s=0}^{t-1} q_{e_s^+, u_0^-}^+ + q^+(f_t, u_0^+) + q^+(f_0, u_0^+)
$$
  

$$
= \frac{a_{t+1} + a_t + 2}{4} \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \frac{a_{t+1} - a_t}{4} + \frac{t}{2} - \frac{t^2}{2n},
$$

as required.  $\square$ 

**Corollary 4.6.** For every  $n \geq 3$ , we have

$$
Kf(CL_n) = n^2 \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} + \frac{1}{6}n(n^2 - 1)
$$

It is known from [\[19,](#page-15-6) Theorem F] that  $Kf(\Gamma) = n \text{tr}(L^+(\Gamma))$  for every graph  $\Gamma$  of order n, where  $L(\Gamma)$  denotes the Laplacian matrix of  $\Gamma$ . Also, from the definition, we know that

$$
r(u, v) = L_{u, u}^{+} + L_{v, v}^{+} - L_{u, v}^{+} - L_{v, u}^{+}
$$

for all  $u, v \in V(\Gamma)$ , where  $L^+$  denotes the Moore-Penrose inverse of  $L = L(\Gamma)$  and  $r(u, v)$  is the resistance distance between vertices u and v of Γ. Utilizing the fact that circular ladders are vertex transitive graphs, we obtain the following result immediately.

<span id="page-12-2"></span>**Corollary 4.7.** Let  $L := LCL_n$  be the Laplacian matrix of  $CL_n$  and  $L^+ = [l_{uv}^+]$ *be its Moore-Penrose inverse. Then*

$$
l_{u_i^{\varepsilon}, u_{i+t}^{\varepsilon'}} = \varepsilon \varepsilon' \frac{a_{t+1} + a_t}{8} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n - 2} - \varepsilon \varepsilon' \frac{a_{t+1} - a_t}{8} - \frac{t}{4} + \frac{t^2}{4n} + \frac{1}{24} \left( n - \frac{1}{n} \right)
$$

*for all*  $1 \leq i \leq n$ ,  $0 \leq t < n$ , and  $\varepsilon, \varepsilon' = \pm 1$ .

# 5. MÖBIUS LADDER GRAPH

Following the same techniques used in the previous section, we shall compute the Moore-Penrose inverse of incidence matrices of Möbius ladder graphs. Likewise, we obtain the resistance distance matrix and Kirchhoff index of Möbius ladders. In what follows,  $M_n$  denotes the Möbius ladder graph of order  $2n$  and  $u_1, \ldots, u_n$ and  $v_1, \ldots, v_n$  stand for the vertices of the external and internal cycles of  $M_n$ , respectively, as in Fig. [4\(](#page-12-0)a). Also, we may set  $u_{i+n} := v_i$  for all  $i = 1, ..., n$ , and that  $u_{i+2nk} := u_i$  for all  $1 \leq i \leq 2n$  and integers k. In what follows,  $f_i := u_i v_i$  $(1 \leq i \leq n)$  is the *i*-th spoke and  $e_i = u_i u_{i+1}$  is the *i*-th edge of the 2n-cycle  $u_1, \ldots, u_{2n}$   $(1 \leq i \leq 2n)$ , see the second drawing in Fig. [4\(](#page-12-0)b).



<span id="page-12-0"></span>FIGURE 4. Möbius ladder graph  $M_n$ 

<span id="page-12-1"></span>**Theorem 5.1.** Let  $M_n$  be the Möbius ladder graph of order  $2n$  and  $Q$  be its inci*dence matrix. Then the Moore-Penrose inverse of*  $Q = [q_{e,v}^+]$  *is given by* 

$$
q_{e_{i+t},u_i}^+ = -\frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} + \frac{a_{t+1} + 1}{4} - \frac{2t + 1}{4n},
$$
  

$$
q_{e_{n+i+t},u_i}^+ = \frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} - \frac{a_{t+1} - 1}{4} - \frac{2t + 1}{4n},
$$
  

$$
\varepsilon_{i+t} q_{f_{i+t},u_i}^+ = \frac{a_{t+1} + a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} - \frac{a_{t+1} - a_t}{4}
$$

*for all*  $t = 0, ..., n - 1$ *.* 

*Proof.* First observe that  $s(M_n/f_i) = s(CL_n/f_i)$  and that by [\[20,](#page-15-17) p. 41]

$$
s(M_n) = s(CL_n) + 2n = \frac{n}{2}(a_{n+1} + a_n + 2).
$$

Therefore, by Lemma [3.5](#page-4-1) and Theorem [4.1,](#page-7-1)

<span id="page-13-0"></span>
$$
q_{f_i, u_i}^+ = -q_{f_i, v_i}^+ = \frac{r(u_i, v_i)}{2} = \frac{s(M_n/f_i)}{2s(M_n)} = \frac{s_n}{a_{n+1} + a_n + 2}.
$$

Consider the edge cut-set  $\{e_j, f_{j+1}, e_{j+1}\}$ . From [\[3,](#page-15-5) Lemma 2.1], it follows that

(15) 
$$
q_{e_{j+1},u_i}^+ = \delta_{i,j+1} - \frac{1}{2n} + q_{e_j,u_i}^+ - \varepsilon_{j+1} q_{f_{j+1},u_i}^+,
$$

where  $\varepsilon_j = (-1)^{[(j-1)/n]}$ . Since  $q_{e_i,u_i}^+ = -q_{e_{i-1},u_i}^+$  by the symmetry of the graph, equation [\(15\)](#page-13-0) yields the initial values

$$
q_{e_i, u_i}^+ = \frac{1}{2} \left( 1 - \frac{1}{2n} - \varepsilon_i q_{f_i, u_i}^+ \right)
$$

for all  $i = 1, \ldots, 2n$ . Likewise, from  $f_{n+i} = f_i$  and  $q_{e_{n+i},u_i}^+ = -q_{e_{n+i-1},u_i}^+$  it follows that

$$
q_{e_{n+i},u_i}^{+} = \frac{1}{2} \left( -\frac{1}{2n} + \varepsilon_i q_{f_i,u_i}^{+} \right)
$$

for all  $i = 1, \ldots, 2n$ . Notice that the values of  $q_{e_{i+t},u_i}^+$  are independent of i by the symmetry of the graph, so we may restrict ourselves to i with  $1 \leq i \leq n$  and assume  $\varepsilon_i = 1$ .

On the other hand, the incidence vector of the cycle  $u_j, u_{j+1}, u_{n+j}, u_{n+j+1}$  lies in the left null space of  $Q^+$ , from which it follows that

<span id="page-13-1"></span>
$$
q_{e_j, u_i}^+ + \varepsilon_{j+1} q_{f_{j+1}, u_i}^+ - q_{e_{n+j}, u_i}^+ - \varepsilon_j q_{f_j, u_i}^+ = 0
$$

or

(16) 
$$
\varepsilon_{j+1} q_{j_{j+1},u_i}^+ = q_{e_{n+j},u_i}^+ - q_{e_j,u_i}^+ + \varepsilon_j q_{f_j,u_i}^+.
$$

Let

$$
C := \begin{pmatrix} 2 & -1 & -1 & 0 \\ -1 & 2 & 0 & -1 \\ -1 & 1 & 1 & 0 \\ 1 & -1 & 0 & 1 \end{pmatrix} \text{ and } D := -\frac{1}{2n} \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix}.
$$

If  $X_j$  denotes the column matrix  $(q_{e_j,u_i}^+ q_{e_{n+j},u_i}^+ \varepsilon_j q_{f_j,u_i}^+ \varepsilon_{n+j} q_{f_{n+j},u_i}^+)^T$  for  $i \leq j <$  $i + n$ , then from the equalities  $(15)$  and  $(16)$ , it follows that

$$
X_{j+1} = CX_j + D
$$

for all  $i \leq j < n + i - 1$ . Using induction on  $t \geq 0$ , one can prove that

$$
C^{t} = \begin{pmatrix} \frac{a_{t+1}+1}{2} & -\frac{a_{t+1}-1}{2} & -\frac{s_t+t}{2} & \frac{s_t-t}{2} \\ -\frac{a_{t+1}-1}{2} & \frac{a_{t+1}+1}{2} & \frac{s_t-t}{2} & -\frac{s_t+t}{2} \\ -s_t & s_t & \frac{a_t+1}{2} & -\frac{a_t-1}{2} \\ s_t & -s_t & -\frac{a_t-1}{2} & \frac{a_t+1}{2} \end{pmatrix}
$$

and

$$
I + C + \dots + C^{t-1} = \begin{pmatrix} \frac{s_t + t}{2} & -\frac{s_t - t}{2} & -\frac{a_t + t^2 - t - 1}{4} & \frac{a_t - t^2 + t - 1}{4} \\ -\frac{s_t - t}{2} & \frac{s_t + t}{2} & \frac{a_t - t^2 + t - 1}{4} & -\frac{a_t + t^2 - t - 1}{4} \\ -\frac{a_t - 1}{2} & \frac{a_t - 1}{2} & \frac{s_{t-1} + t + 1}{2} & -\frac{s_{t-1} - (t-1)}{2} \\ \frac{a_t - 1}{2} & -\frac{a_t - 1}{2} & -\frac{s_{t-1} - (t-1)}{2} & \frac{s_{t-1} + t + 1}{2} \end{pmatrix}.
$$

Now, from the equality

$$
X_{i+t} = C^t X_i + (I + C + \dots + C^{t-1})D,
$$

it follows that

$$
q_{e_{i+t},u_i}^+ = -\frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} + \frac{a_{t+1} + 1}{4} - \frac{2t + 1}{4n}
$$

$$
q_{e_{n+i+t},u_i}^+ = \frac{2a_{t+1} - a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} - \frac{a_{t+1} - 1}{4} - \frac{2t + 1}{4n},
$$

$$
\varepsilon_{i+t} q_{f_{i+t},u_i}^+ = \frac{a_{t+1} + a_t}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} - \frac{a_{t+1} - a_t}{4}
$$

for all  $t = 0, \ldots, n - 1$ , as required.

<span id="page-14-0"></span>**Corollary 5.2.** *Let*  $1 \leq i \leq 2n$  *and*  $0 \leq t \leq n$ *. Then* 

$$
r(u_i, u_{i+t}) = -\frac{a_{t+1} + a_t - 2}{4} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} + \frac{a_{t+1} - a_t}{4} + \frac{t}{2} - \frac{t^2}{2n}
$$

*Proof.* The result follows from Lemma [3.5](#page-4-1) and Theorem [5.1](#page-12-1) in conjunction with the fact that

$$
\sum_{j=0}^{t-1} q^+_{e_{i+j},u_{i+t}} = -\sum_{j=0}^{t-1} q^+_{e_{i+j},u_i}
$$

by the symmetry of the graph.

Theorem [5.2](#page-14-0) in conjunction with equation [\(5\)](#page-1-5) yields

**Corollary 5.3.** *For every*  $n \geq 3$ *, we have* 

$$
Kf(CL_n) = n^2 \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} + \frac{1}{6}n(n^2 - 1).
$$

Finally, the Moore-Penrose inverse of the Laplacian matrix of  $M_n$  can be obtained analogous to Corollary [4.7.](#page-12-2)

**Corollary 5.4.** Let  $L := L(M_n)$  be the Laplacian matrix of  $M_n$  and  $L^+ = [l_{uv}^+]$  be *its Moore-Penrose inverse. Then*

$$
l_{u_i, u_{i+t}}^+ = \frac{a_{t+1} + a_t}{8} \cdot \frac{a_{n+1} - a_n}{a_{n+1} + a_n + 2} - \frac{a_{t+1} - a_t}{8} - \frac{t}{4} + \frac{t^2}{4n} + \frac{1}{24} \left( n - \frac{1}{n} \right)
$$

*for all*  $1 \leq i \leq 2n$  *and*  $0 \leq t < n$ *.* 

$$
f_{\rm{max}}
$$

.

,

### **REFERENCES**

- <span id="page-15-3"></span>[1] A. Azimi and R. B. Bapat, Moore-Penrose inverse of the incidence matrix of a distance regular graph, *Linear Algebra Appl.* 551 (2018), 92–103.
- <span id="page-15-4"></span>[2] A. Azimi and R. B. Bapat, The Moore-Penrose inverse of the incidence matrix of complete multipartite and bi-block graphs, *Discrete Math.* 342(8) (2019), 2393–2401.
- <span id="page-15-5"></span>[3] A. Azimi, R. B. Bapat, and E. Estaji, Moore-Penrose inverse of incidence matrix of graphs with complete and cyclic blocks, *Discrete Math.* 342 (2019), 10–17.
- <span id="page-15-18"></span>[4] A. Azimi, R. B. Bapat, and M. Farrokhi D. G., Resistance distance of blow-ups of trees, *Discrete Appl. Math.* 320 (2022), 311–323.
- <span id="page-15-10"></span>[5] P. Ali, F. Atik, and R. B. Bapat, Identities for minors of the Laplacian, resistance and distance matrices of graphs with arbitrary weights, *Linear Multilinear Algebra* 68(2) (2020), 323–336.
- <span id="page-15-19"></span><span id="page-15-11"></span>[6] R. B. Bapat, Graphs and Matrices, Second Edition, Springer, 2014.
- [7] R. B. Bapat and S. Gupta, Resistance distance in wheels and fans, *Indian J. Pure Appl. Math.* 41 $(1)$   $(2010)$ , 1-13.
- <span id="page-15-12"></span>[8] N. L. Biggs, Potential theory on distance-regular graphs, *Combin. Probab. Comput.* 2(3) (1993), 243–255.
- <span id="page-15-0"></span>[9] A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory and Applications*, Second Edition, Springer, New York, 2003.
- <span id="page-15-1"></span>[10] S. L. Campbell and C. D. Meyer, *Generalized Inverses of Linear Transformations*, Pitman, London, 1979.
- <span id="page-15-20"></span>[11] Z. Cinkir, Effective resistances and Kirchhoff index of ladder graphs, *J. Math. Chem.* 54(4) (2016), 955–966.
- <span id="page-15-13"></span>[12] K. C. Das and Y. Yang, Eigenvalues of the resistance-distance matrix of complete multipartite graphs, *J. Inequal. Appl.* (2017), No. 296, 11 pp.
- <span id="page-15-7"></span>[13] E. Ernesto and M. Delio, Hubs-biased resistance distances on graphs and networks, *J. Math. Anal. Appl.* 507, no. 1, (2022), Paper No. 125728, 22 pp.
- <span id="page-15-14"></span>[14] X. Gao, Y. Luo, and W. Liu, Resistance distances and the Kirchhoff index in Cayley graphs, *Discrete Appl. Math.* 159(17) (2011), 2050–2057.
- <span id="page-15-15"></span>[15] S. V. Gervacio, Resistance distance in complete n-partite graphs, *Discrete Appl. Math.* 203 (2016), 53–61.
- <span id="page-15-16"></span>[16] Q. Huang, H. Chen, and Q. Deng, Resistance distances and the Kirchhoff index in double graphs, *J. Appl. Math. Comput.* 50 (2016), 1–14.
- <span id="page-15-2"></span>[17] Y. Ijiri, On the generalized inverse of an incidence matrix, *J. Soc. Indust. Appl. Math.* 13 (1965), 827–836.
- <span id="page-15-8"></span>[18] M. Kagan and B. Mata, A physics perspective on the resistance distance for graphs, *Math. Comput. Sci.* 13(1-2) (2019), 105–115.
- <span id="page-15-17"></span><span id="page-15-6"></span>[19] D. J. Klein and M. Randić, Resistance distance, *J. Math. Chem.* **12** (1993), 81–95.
- [20] M. Rubey, *Counting Spanning Trees*, Diplomarbeit, Universitat Wein, Mai, 2000.
- <span id="page-15-9"></span>[21] N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, <https://oeis.org>.
- [22] Y. Yi and S. Yue, Resistance-distance betweenness: Definition, properties, and application in complex linguistic networks, *Phys. A*, 610, (2023), Paper No. 128408.

Department of Mathematics and Applied Mathematics, Xiamen University Malaysia, 43900, Sepang, Selangor Darul Ehsan, Malaysia

#### *Email address*: ali.azimi61@gmail.com, ali.azimi@xmu.edu.my

Research Center for Basic Sciences and Modern Technologies (RBST), Institute for Advanced Studies in Basic Sciences (IASBS), Zanjan 45137-66731, Iran

*Email address*: m.farrokhi.d.g@gmail.com, farrokhi@iasbs.ac.ir