Tristan Rivière

Abstract : In any 5-dimensional closed Sasakian manifold we prove that any minmax operation on the area among Legendrian surfaces is achieved by a continuous conformal Legendrian map from a closed riemann surface S into N^5 equipped with an integer multiplicity bounded in L^{∞} . Moreover this map, equipped with this multiplicity, satisfies a weak version of the Hamiltonian Minimal Equation. We conjecture that any solution to this equation is a smooth branched Legendrian immersion away from isolated Schoen-Wolfson conical singularities with non zero Maslov class.

Math. Class. 53D12, 49Q05, 53A10, 58E12, 49Q10

I Introduction

I.1 Foreword

The simplest framework of our investigations in this work could be formulated as follows: study area variations among maps u from an oriented 2-manifold Σ into \mathbb{C}^2 under the Lagrangian constraint

$$u^*\omega = 0 , \qquad (I.1)$$

where ω is the standard symplectic form $\omega = dy_1 \wedge dy_2 + dy_3 \wedge dy_4$. More precisely we ask ourselves how to implement variational operations (minimization under various constraints, minmax) for the standard area Lagrangian

$$A(u) := \int_{\Sigma} |\partial_{x_1} u \wedge \partial_{x_2} u| \ dx_1 \wedge dx_2$$

among maps satisfying the pointwise constraint¹

$$du_1 \wedge du_2 + du_3 \wedge du_4 = 0$$

The Lagrangian A is notoriously known in calculus of variations to be delicate to work with (due mostly to the very large invariance group in the domain : the "gauge group" of diffeomorphisms of the surface Σ). In order to "break" this invariance and reduce the gauge group to a finite dimensional one, it is preferable instead to consider the Dirichlet Energy of the map u for a variable metric g of Σ

$$E(u,g) := \frac{1}{2} \int_{\Sigma} |du|_g^2 \, dvol_g \ge A(u) \;. \tag{I.2}$$

with equality if and only if u is conformal with respect to the metric g that is in isothermic or conformal coordinates such that $g = e^{2\lambda} [dx_1^2 + dx_2^2]$ there holds

$$|\partial_{x_1}u|^2 = |\partial_{x_2}u|^2$$
 and $\partial_{x_1}u \cdot \partial_{x_2}u = 0$, (I.3)

¹In case u is a graph $u(x_1, x_2) = (x_1, x_2, v(x_1, x_2))$ the constraint (I.1) is equivalent to the incompressibility condition

$$\det(\nabla v) = -1$$

where \cdot denotes the standard scalar product in \mathbb{C}^2 . Stationary critical points for fixed g have an holomorphic Hopf differential

$$H(u) := \partial_z u \cdot \partial_z u \, dz \otimes dz$$

while making g vary simultaneously in the Moduli space of constant Gauss curvature metrics is known to imply H(u) = 0 and equality holds in (I.2). Hence critical points of E(u, g) are critical points of the area.

Assuming u is a conformal critical point of the Dirichlet energy under the Lagrangian constraint, in local conformal coordinates, <u>formally</u> the Euler Lagrange Equation of our constrained variational problem is given by

$$\begin{cases} \operatorname{div}\left(\mathfrak{g}\,\nabla u\right) = 0\\ \operatorname{div}\left(\mathfrak{g}^{-1}\,\nabla\mathfrak{g}\right) = 0 \end{cases}, \tag{I.4}$$

where $\mathfrak{g} = e^{i\beta}$ is an S^1 valued map², which is the Lagrange multiplier associated to the Lagrangian pointwise constraint (I.1). The first equation is a structural equation shared by any conformal parametrization of any Lagrangian surface in \mathbb{C}^2 while the second one is the Euler Lagrange equation itself. It is saying that the Lagrange multiplier \mathfrak{g} is an S^1 harmonic map. These equations are obviously calling upon the use of classical elliptic theory for solving regularity or existence questions. The difficulty however is that, while the natural assumption is to consider $u \in W_{loc}^{1,2}(\Sigma, \mathbb{C}^2)$, there is a-priori no information on the function space to which \mathfrak{g} belongs to in order to start implementing this theory. Hence these questions (regarding existence and regularity) cannot be settled at the strictly "PDE level" whose formulation even is problematic. Indeed in order to give a distributional meaning to (I.4) one needs at least an assumption like \mathfrak{g} is in $W^{1,1}$ or $H^{1/2}$. Studying the regularity³ of solution to the system (I.4) assuming the conformality condition⁴ (I.3) is certainly an interesting elliptic PDE problem by itself but somehow artificial though since these assumptions are not given a-priori by any variational operation we are aiming at doing.

I.2 Hamiltonian and Lagrangian Stationary Immersions

A Kähler surface is a complex two dimensional manifold (M^2, J) with a compatible⁵ symplectic form ω . Lagrangian sub-manifolds are the ones on which the restriction of the symplectic form vanishes. In particular, Lagrangian sub-manifolds are the ones for which the complex structure realizes an isometry between the tangent and the normal spaces at every points.

The study of area variations under Lagrangian constraints has been initiated in a series of work by Yong-Geun Oh ([12], [13]). He proposed to minimize the area under the Lagrangian constraint within a Lagrangian homology class with the objective to obtain in this way very special representative of these classes. He called the sub-manifolds critical points of the area within Lagrangian sub-manifolds stationary Lagrangians. He introduced also a less restrictive class of surfaces⁶ which are the so called Hamiltonian stationary sub-manifolds. This later class of surfaces are critical points of the area for local deformations which preserve infinitesimally the Lagrangian condition. These deformations are generated by the Hamiltonian vector-fields which are of the form

$$J \nabla h$$
. (I.5)

²The function β plays the role of a Lagrange multiplier issued from the incompressibility constraint similarly as the role played by the "pressure" in the variational formulation of Euler equations.

 $^{^{3}}$ Observe that everywhere discontinuous solutions to the second equation of (I.4) have been constructed in [2].

⁴A weaker hypothesis would be to assume that the Hopf differential H(u) is holomorphic, which is equivalent for the critical point to be a stationary critical point that is critical for variation in the domain.

⁵A symplectic form is said to be compatible with a complex or with an almost complex structure J if $g(X, Y) := \omega(X, JY)$ defines a riemannian metric.

 $^{^{6}}$ The class of Hamiltonian stationary surfaces is in general richer and more interesting than the class of Lagrangian stationary surfaces according to the author of the present paper.

where h is an arbitrary smooth function. Such surfaces are also called *Hamiltonian minimal* or simply H-minimal surfaces. These pioneered works on the subject contain very interesting and now classical conjectures in the field about possible volume minimizing candidates under hamiltonian deformations within \mathbb{C}^2 or $\mathbb{C}P^2$.

Y.-G. Oh computed in particular the Euler Lagrange equations associated to this variational problem. In the special case of Kähler-Einstein manifolds (satisfying $Ric(g) = \lambda g$), it has been discovered by Dazor [6] that the mean curvature of a Lagrangian surface Σ is given by

$$\vec{H} := J \, \nabla^{\Sigma} \beta \; ,$$

where β is a locally well defined function on Σ and ∇^{Σ} is the gradient of β for the induced metric on Σ . In the particular case when the manifold is Ricci flat (which implies that M^2 is Calabi Yau), if Ω is a normalized global holomorphic section of the canonical bundle $K := \wedge^{2,0} M$ the function β is given by

$$e^{i\beta}\iota_{\Sigma}^*\Omega = dvol_q$$

where ι_{Σ} is the canonical embedding of Σ in M^2 . Hence β is a globally well defined function in $\mathbb{R}/2\pi\mathbb{Z}$ and is called *Lagrangian angle*. For general Kähler-Einstein manifolds β is still a well defined function in $\mathbb{R}/2\pi\mathbb{Z}$ in case the induced canonical connection of the restriction to Σ of the canonical bundle $\iota_{\Sigma}^* K$ of M^2 (which is flat) has no monodromy (see [24]).

In [12] and [13] it is proved that a sub-manifold is *hamiltonian stationary* (or *H*-minimal) if the contraction of the mean curvature vector with the Kähler form

$$\omega \bot \dot{H} = \alpha_{\vec{H}}$$

defines an <u>harmonic</u> one form of the sub-manifold Σ . This is equivalent to the fact that

$$\Delta^{\Sigma}\beta = 0$$

which is nothing but the second equation in (I.4).

In the particular case when β is a globally univalued function, then $\beta = \beta_0$ and the Euler-Lagrange equation is simply the minimal surface equation $\vec{H} = 0$. Such a sub-manifold is called *minimal lagrangian*. If the Kähler-Einstein manifolds is Ricci flat then any *minimal lagrangian* sub-manifold is calibrated by $e^{i\beta_0} \Omega$ and and realizes in this way an absolute minimizer⁷ in its homology class (see [10]). While it is a very interesting object of research with numerous applications in geometry (and in high energy physics⁸) in general there are no reasons why $d\beta$ should be equal to zero and the general hamiltonian stationary equation in Kähler-Einstein Manifold is the 3rd order equation⁹. It is interesting to observe that, while having $d\beta$ being harmonic we have a "finite dimensional perturbation" of $\beta = \beta_0$ but nevertheless Hminimal surfaces have very different features. For instance there exists closed H-minimal surfaces in \mathbb{C}^2 (the Clifford torus $S^1 \times S^1 \subset \mathbb{C} \times \mathbb{C}$ is an example of such a surface¹⁰) while the existence of closed minimal surfaces is prevented by the maximum principle.

⁷The local stability of arbitrary minimal lagrangian sub-manifolds generalizes to the situation of general Kähler-Einstein manifolds of non positive Ricci. This being said we will be interested in the other case in the present work that is Ric(g) > 0. ⁸Minimal Lagrangian submanifolds of Calabi Yau 3 and 4-folds appear in string theory under the name of "super-

symmetric cycle" and they play a key role in the Strominger Yau Zaslow theory of mirror symmetry. ⁹The equation is second order with respect to the multivalued Lagrangian angle function β but the expression of $d\beta$ involves 2 derivatives with respect to the Lagrangian immersion.

 $^{^{10}}$ One of the Oh conjecture says that this torus should be minimizing in its Hamiltonian isotopy class : The class of tori which can be obtained through hamiltonian deformations, that is, deformations which infinitesimally are given by (I.5) (see [12]).

I.3 The Schoen Wolfson existence and regularity result

In a breakthrough paper [20], Rick Schoen and Jon Wolfson performed the very first variational construction of minimizing stationary lagrangian surfaces in homology classes. They proved, in particular, the existence of an area minimizing Lagrangian surface in any 2 dimensional lagrangian homology class of a Kähler surface. The main issue is about proving the regularity of these minimizers. The two authors proved that these area minimizing lagrangian stationary surfaces are branched immersions with possible isolated "conical" singularities¹¹. At this singularities a_i the S^1 valued map \mathfrak{g} realizes some topological degree d_i and solves in conformal coordinates the equation

$$\operatorname{curl}\left[\mathfrak{g}^{-1}\nabla\mathfrak{g}\right] = \sum_{i\in I} d_i\,\delta_{a_i}$$

In fact these conical singularities, whose description is fully given in [20], are proved to exist even for minimizers inside some spherical Lagrangian homology classes (see [11] and [25]) in some Kähler-Einstein surfaces. This last fact is a surprise and is one of the reasons why the variational analysis of Lagrangian surfaces, even in the lowest co-dimension 2, is particularly involved.

The starting point for performing such an analysis is the search of a monotonicity formula. In classical minimal surface theory it says that the quantity of area of a closed minimal surface inside a ball of radius r divided by r^2 is increasing. At this point Schoen and Wolfson made the following crucial observation that such a monotonicity formula is not true in \mathbb{C}^2 for H-minimal surfaces in general (even minimizing !). They give an explicit simple counter-example in [19] (section 4). The existence of such a counter-example could be explained by the fact that the generator of dilation¹² in \mathbb{C}^2 is not hamiltonian :

$$y_1 \,\partial_{y_1} + y_2 \,\partial_{y_2} + y_3 \,\partial_{y_3} + y_4 \,\partial_{y_4} \neq i\nabla h$$

The major observation made by Schoen and Wolfson is that by adding the Legendrian coordinate (if it locally exists) this problem can be overcome. If u is Lagrangian in \mathbb{C}^2 there holds

$$d(u_1 \, du_2 - u_2 \, du_1 + u_3 \, du_4 - u_4 \, du_3) = 0$$

and one introduces the local Legendrian coordinate

$$d\varphi_u := u_1 \, du_2 - u_2 \, du_1 + u_3 \, du_4 - u_4 \, du_3 \; .$$

In \mathbb{R}^5 one introduce the form

$$\alpha := -d\varphi + y_1 \, dy_2 - y_2 \, dy_1 + y_3 \, dy_4 - y_4 \, dy_3$$

If u is Lagrangian the map $v := (\varphi_u, u_1, u_2, u_3, u_4)$ is cancelling the form α which is non integrable $\alpha \wedge d\alpha \wedge d\alpha \neq 0$. The map v is called *Legendrian*. Infinitesimal variations among Legendrian maps in \mathbb{R}^5 are given by Hamiltonian vector fields in the contact space (\mathbb{R}^5, α) of the form

$$\vec{X} := J_H \,\nabla^H h - 2 \,h \,\partial_\varphi \,\,. \tag{I.6}$$

where *H* is the 4-plane given by the kernel of α , ∇^H is the projection on *H* of the gradient in \mathbb{R}^5 with respect to the metric such that π_* , the differential of the canonical projection $\pi(\varphi, y_1, y_2, y_3, y_4) = (y_1, y_2, y_3, y_4)$ from *H* into \mathbb{C}^2 , realizes an isometry and the length of ∂_{φ} is 1. \mathbb{R}^5 equipped with this metric

¹¹The "conical" singularities are singularities with non flat *tangent cones* and around which β is not a univalent \mathbb{R} -valued map and is contributing to the total Maslov class. They are called nowadays "Schoen Wolfson cones".

 $^{1^{2}}$ The existence of monotonicity formula, which is a conservation law, is intimately linked to the existence of infinitesimal isotropy of the space (see [1]) and technically in general is obtained by considering the variation of the surface with respect to the generator of dilations "exploring" this isotropy.

defines the Heisenberg group \mathbb{H}^2 . Finally J_H is the complex structure on H such that it projects by π_* onto the standard complex structure of \mathbb{C}^2 . The authors in this work introduce the notion of H-minimal Legendrian for the immersion which are critical point of the area with respect to all infinitesimal variations generated by Hamiltonian vector fields in \mathbb{R}^5 of the form (I.6). One then observe

 $u = \pi \circ v$ is H-Minimal in $\mathbb{C}^2 \iff v$ is H-Minimal Legendrian in (\mathbb{R}^5, α) .

Testing $h = -\varphi$ produces the generator of dilations in \mathbb{C}^2 plus dilations in the 5th Legendrian direction but with a different scale.

$$-i\nabla^H \varphi + 2\varphi \,\partial_{\varphi} = y_1 \,\partial_{y_2} - y_2 \,\partial_{y_1} + y_3 \,\partial_{y_4} - y_4 \,\partial_{y_3} + 2\varphi \,\partial_{\varphi} \,.$$

This fact is an encouragement for looking for a monotonicity formula for H-Minimal Legendrian surfaces. However, as a matter of fact, it cannot be deduced from a simple insertion of this generator of the dilation in the stationarity condition. The derivation of a monotonicity formula for H-Minimal Legendrian immersions in (\mathbb{R}^5 , α) proposed in [20], which is maybe one of the main achievement of this work, is quite indirect and involved (going through the resolution of a wave equation, the use of Bessel functions...etc). Another drawback is that the formula is semi-explicit.

Recently the author in [18] obtained a new monotonicity formula by inserting in the Legendrian stationarity condition an explicit hamiltonian : $h := \chi(\mathfrak{r}) \arctan(2 \varphi/|y|^2)$, where \mathfrak{r} is the Folland-Korányi gauge given by $\mathfrak{r}^4 = |y|^4 + 2 \varphi^2$ and χ is some well chosen cut-off function. Because of its explicit nature the formula is more effective and is the starting point of the present work and the proof of the main result which was out of reach without it.

I.4 Main result

Let (N^5, g) be a 5 dimensional oriented Riemannian manifold and α a non degenerate 1-form on N^5 (i.e. (N^5, α) is contact) satisfying

$$\alpha \wedge d\alpha \wedge d\alpha > 0 \; ,$$

The triple (N^5, g, α) is called a Sasakian structure if the cone $(N^5 \times \mathbb{R}_+, k := dt^2 + t^2g)$ with the non degenerate symplectic form $\Omega := 2^{-1} d(t^2 \alpha)$ is Kähler. Let J be the compatible complex structure $(\Omega(\cdot, J \cdot) = g)$. The tangent vector field along N^5 given by $\vec{R} := J(t\partial_t)$ is unit on N^5 and is orthogonal to the "horizontal hyperplanes" given by $H = \text{Ker}(\alpha)$. It is called *Reeb Vector-field* of the distribution H. This distribution of plane is invariant under the action of J in $N^5 \times \mathbb{R}_+$ and we shall denote J_H its restriction on H. Such a structure is called Sasakian structure (see [22]). Classical examples are the Heisenberg group \mathbb{H}^2 , the unit sphere S^5 for the Reeb vector-field tangent to the fibers of the Hopf fibration into $\mathbb{C}P^2$ or the Stiefel manifold $V_2(\mathbb{R}^4) \simeq S^3 \times S^2$ of orthonormal 2-frames in \mathbb{R}^4 for the Reeb vector-field given by fibers of the tautological projection onto the Grassman manifold $G_2(\mathbb{R}^4)$ of 2-planes in \mathbb{R}^4 .

A Sasakian structure being given, we introduce the Sobolev space of Legendrian $W^{2,4}$ immersions of a closed oriented surface Σ in N^5

$$\mathfrak{E}^{2,4}_{\Sigma,Leg}(\Sigma,N^5):=\left\{\vec{\Lambda}\in W^{2,4}(\Sigma,N^5)\quad;\quad |d\vec{\Lambda}\,\dot{\wedge}\,d\vec{\Lambda}|>0 \ \text{ and } \ \vec{\Lambda}^*\alpha=0 \quad \text{ on } \Sigma\right\}\;.$$

We prove in section II that $\mathfrak{M} := \mathfrak{E}_{\Sigma,Leg}^{2,4}(\Sigma, N^5)$ has the structure of Banach manifold and possesses a compatible Finsler structure for which the associated *Palais distance* is complete.

An admissible family \mathcal{A} in \mathfrak{M} is a set of subsets of \mathfrak{M} which is invariant under isotopies in \mathfrak{M} . The min-max value or the "width" associated to such a family is the number given by

$$\beta := \inf_{A \in \mathcal{A}} \sup_{\vec{\Lambda} \in A} \int_{\Sigma} dvol_{\vec{\Lambda}} .$$

Our main result in the present work says that any such min-max is achieved by a continuous conformal Legendrian map from a closed riemann surface S into N^5 equipped with an integer multiplicity bounded in L^{∞} and satisfying a weak version of the H-minimal equation. Such a surface is called Legendrian Hamiltonian stationary parametrized integral varifold. Precisely we have:

Theorem I.1. Let \mathcal{A} be an admissible family for the space of $W^{2,4}$ Legendrian immersions of a closed oriented surface Σ into a closed Sasakian manifold (N^5, g, α) . Assume that the associated width is strictly positive

$$\beta := \inf_{A \in \mathcal{A}} \sup_{\vec{\Lambda} \in A} \int_{\Sigma} dvol_{\vec{\Lambda}} > 0 \; .$$

Then there exists a closed Riemann surface¹³ (S,h) with $genus(S) \leq genus(\Sigma)$ a map $\vec{\Lambda} \in C^0(S, N^5) \cap W^{1,2}(S, N^5)$ and $N \in L^{\infty}(S, \mathbb{N}^*)$ such that

i)

$$\vec{\Lambda}^* \alpha = 0$$

ii)

$$\partial \vec{\Lambda} \otimes \partial \vec{\Lambda} := \partial_z \vec{\Lambda} \cdot \partial_z \vec{\Lambda} \, dz \otimes dz = 0 \qquad a. \ e. \ on \ S$$

iii)

$$\int_{S} N \, dvol_{\vec{\Lambda}} = \frac{1}{2} \, \int_{S} N \, |\nabla \vec{\Lambda}|_{h}^{2} \, dvol_{h} = \beta$$

iv)

$$\forall h \in C^{3}(S) \quad , \quad \forall f \in C^{1}(S) \quad for \ a.e. \ \lambda \in \mathbb{R} \quad s.t. \quad \vec{\Lambda}(f^{-1}\{\lambda\}) \cap Supp(h) = \emptyset$$

$$\int_{f^{-1}((\lambda, +\infty))} N \,\nabla \vec{\Lambda} \cdot \nabla \left[\vec{X}_{h} \circ \vec{\Lambda}\right] \ dvol_{h} = 0 \ , \tag{I.7}$$

where \vec{X}_h is the Hamiltonian vector-field associated to h and given by

$$\vec{X}_h := J_H(\nabla^H h) - 2 h \vec{R} ,$$
 (I.8)

where ∇^H is the projection onto H of the gradient in (N^5, g) .

In [17] the corresponding result to theorem I.1 was proven without the pointwise Legendrian constraint. Solutions to (I.7) for arbitrary \vec{X} (non necessarily Hamiltonian) were called *parametrised stationary integral varifolds*. In an analogous way we shall be calling any triple $(S, \vec{\Lambda}, N)$ solving (I.7) for any Hamiltonian vector-field of the form (I.8) Legendrian parametrised Hamiltonian stationary integral varifolds.

We stress that the stationarity identity (I.7) holds for a.e. $\lambda \in \mathbb{R}$ foor which the image by $\vec{\Lambda}$ of the level set $f^{-1}\{\lambda\}$ avoids the support of the testing vector-field \vec{X} . This is the called "localisation hypothesis" (It has been first introduced in [16]). Combining the nullity of the variation of the area for perturbation \vec{X} in the target with the "localisation hypothesis" is a substitute to the Euler Lagrange equation (I.4), which has no satisfying weak formulation as we discussed in the foreword.

The strategy adopted to establish theorem I.1 is the penalisation approach introduced by the author in [17] under the name "viscosity method". This method has then been implemented for free boundary

 $^{^{13}}$ We equip the surface with a compatible metric.

surfaces in [14] and we will adopt some of the improvements of the method brought in this work. Precisely we study the smoothed minmax operation

$$\beta(\varepsilon) := \inf_{A \in \mathcal{A}} \sup_{\vec{\Lambda} \in A} E_{\varepsilon}(\vec{\Lambda}) ,$$

with

$$E_{\varepsilon}(\vec{\Lambda}) := \int_{\Sigma} dvol_{\vec{\Lambda}} + \varepsilon^4 \int_{\Sigma} (1 + |\vec{\mathbb{I}}_{\vec{\Lambda}}|^2)^2 \, dvol_{\vec{\Lambda}}$$

where $\vec{\mathbb{I}}_{\vec{\Lambda}}$ is the second fundamental form of $\vec{\Lambda}$. In section II We construct almost critical points $\vec{\Lambda}_k$ of E_{ε_k} for a sequence $\varepsilon_k \to 0$ such that

$$E_{\varepsilon_k}(\vec{\Lambda}_k) - \beta(\varepsilon_k) \longrightarrow 0 \quad \text{and} \quad \varepsilon_k^4 \int_{\Sigma} (1 + |\vec{\mathbb{I}}_{\vec{\Lambda}}|^2)^2 \, dvol_{\vec{\Lambda}_k} = o\left(\frac{1}{\log \varepsilon_k^{-1}}\right) \,.$$

The main difficulty to overcome in sections III and IV consists in passing to the limit for some well chosen subsequence of $\vec{\Lambda}_k$ in order to obtain the map $\vec{\Lambda}$ satisfying the conclusions of theorem I.1. While we broadly follow the same scheme as the one introduced in [17], the Legendrian constraint however introduces new serious challenges for the proof of theorem I.1.

In [15] it is proved that any *parametrised stationary varifolds* is a smooth branched minimal immersion equipped with a smooth multiplicity. We make the following conjecture.

Open Problem I.1. Prove that 2-dimensional Legendrian parametrised Hamiltonian stationary integral varifolds are branched Legendrian immersions and smooth away from isolated Schoen Wolfson conical singularities.

We first present the proof of theorem I.1 for $N^5 = V_2(\mathbb{R}^4)$ which has motivated the present works for reasons explained in [17] in relation with the Willmore conjecture. The case $N^5 = V_2(\mathbb{R}^4)$ has the advantage of clarity in the presentation due to the symmetric nature of the Stiefel space of orthonormal 2 frames in \mathbb{R}^4 . This being said the global symmetry of $V_2(\mathbb{R}^4)$ has no incidence at all on the analysis presented in sections II, IV and V. What is crucial is the common nature of the asymptotic of any arbitrary Sasakian manifold while dilating at any arbitrary point. It always converge to to the Heisenberg Group \mathbb{H}^2 . Hence the proof of theorem I.1 for general target is formally identical and requires some obvious adaptation of the computation of sections III, IV and V. This is explained in the last section VI.

II Notations and Preliminaries on $V_2(\mathbb{R}^4)$

II.1 Contact structure; complex structure on horizontal planes

We denote by $V_2(\mathbb{R}^4)$ the Stiefel Manifold of orthonormal 2 frames in \mathbb{R}^4 :

$$V_2(\mathbb{R}^4) := \left\{ (\vec{a}, \vec{b}) \in S^3 \times S^3 \quad ; \quad \vec{a} \cdot \vec{b} = 0 \right\}$$

We consider on $V_2(\mathbb{R}^4)$ the following 2-form

$$\alpha := \vec{a} \cdot d\vec{b} - \vec{b} \cdot d\vec{a} = \sum_{i=1}^4 a^i \, db^i - b^i \, da^i \quad .$$

where $\vec{a} = (a^1, \dots, a^4)$ and $\vec{b} = (b^1, \dots, b^4)$. The Stiefel manifold $V_2(\mathbb{R}^4)$ defines an S^1 -bundle over $G_2(\mathbb{R}^4) \simeq S^2_+ \times S^2_-$. The projection "Hopf map" Π which to (\vec{a}, \vec{b}) assigns $((\vec{a} \wedge \vec{b})^+, (\vec{a} \wedge \vec{b})^-)$ where $(\vec{a} \wedge \vec{b})^{\pm}$ are respectively the dual and anti-dual parts of $\sqrt{2} \vec{a} \wedge \vec{b}$ that is where

$$(\vec{a} \wedge \vec{b})^{\pm} = \frac{1}{\sqrt{2}} \left[\vec{a} \wedge \vec{b} \pm * (\vec{a} \wedge \vec{b}) \right]$$

and * is the Hodge operator on 2 vectors in \mathbb{R}^4 such that

$$\vec{e} \wedge \vec{f} \wedge * (\vec{e} \wedge \vec{f}) = |\vec{e} \wedge \vec{f}|^2 \ \vec{\varepsilon_1} \wedge \vec{\varepsilon_2} \wedge \vec{\varepsilon_3} \wedge \vec{\varepsilon_4} ,$$

where $(\vec{\varepsilon}_1 \cdots \vec{\varepsilon}_4)$ is the canonical basis of \mathbb{R}^4 .

So we have where we recall that Π is the tautological projection from $V_2(\mathbb{R}^4)$ into $G_2(\mathbb{R}^4) \simeq S^2_+ \times S^2_$ given by

$$\Pi : (\vec{a}, \vec{b}) \longrightarrow ((\vec{a} \wedge \vec{b})^+, (\vec{a} \wedge \vec{b})^-) = \left(\frac{\vec{a} \wedge \vec{b} + *\vec{a} \wedge \vec{b}}{\sqrt{2}}, \frac{\vec{a} \wedge \vec{b} - *\vec{a} \wedge \vec{b}}{\sqrt{2}}\right)$$

We shall denote π_+ and π_- the canonical projections

$$\pi_+$$
 : $S^2_+ \times S^2_- \longrightarrow S^2_+$ and π_- : $S^2_+ \times S^2_- \longrightarrow S^2_-$

Using the quaternionic representatives of \vec{a} and \vec{b} that we denote respectively

$$\mathbf{a} := a^1 + a^2 \mathbf{i} + a^3 \mathbf{j} + a^4 \mathbf{k}$$
 and $\mathbf{b} := b^1 + b^2 \mathbf{i} + b^3 \mathbf{j} + b^4 \mathbf{k}$

the projection Π identifies to the following map

$$\Pi : \vec{a} \wedge \vec{b} \in G_2(\mathbb{R}^4) \longleftrightarrow \left(\begin{pmatrix} <\mathbf{a}, \mathbf{ib} > \\ <\mathbf{a}, \mathbf{jb} > \\ <\mathbf{a}, \mathbf{kb} > \end{pmatrix}, \begin{pmatrix} <\mathbf{a}, \mathbf{bi} > \\ <\mathbf{a}, \mathbf{bj} > \\ <\mathbf{a}, \mathbf{bk} > \end{pmatrix} \right) = (\mathbf{ab}^*, \mathbf{b}^*\mathbf{a}) \in S^2_+ \times S^2_-$$

where $\langle \cdot, \cdot \rangle$ is the scalar product on $\mathbf{H} \simeq \mathbb{R}^4$. This diffeomorphism will naturally induce the complex structure of $\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}$ on $G_2(\mathbb{R}^4)$ as we will see where $\mathbb{C}P^1$ is referring to the reversed complex structure (i.e reversed orientation) with respect to the canonical one J_+ on S^2 : $J_-X = -J_+X$.

Let $\vec{\Lambda} = (\vec{a}, \vec{b})$ we have

$$T_{\vec{\Lambda}} V_2(\mathbb{R}^4) = \left\{ (\vec{V}, \vec{W}) \ ; \ \vec{V} \cdot \vec{a} = 0 \ , \ \vec{W} \cdot \vec{b} = 0 \ \text{ and } \ \vec{a} \cdot \vec{W} + \vec{V} \cdot \vec{b} = 0 \right\} \ .$$

Let $(\vec{c}, \vec{d}) \in V_2(\mathbb{R}^4)$ such that $*(\vec{a} \wedge \vec{b}) = \vec{c} \wedge \vec{d}$. We denote $\vec{G} = \vec{a} \wedge \vec{b}$ and note that

$$T_{\vec{G}}G_2(\mathbb{R}^4) = \left\{ \vec{a} \wedge \vec{W} + \vec{V} \wedge \vec{b} \quad \text{where} \quad \vec{V} \,, \, \vec{W} \in \operatorname{Span}(\vec{a}, \vec{b})^{\perp} = \operatorname{Span}(\vec{c}, \vec{d}) \right\} \,,$$

and

$$\Pi_* T_{\vec{\Lambda}} V_2(\mathbb{R}^4) = \left\{ ((\vec{a} \wedge \vec{W})^+ + (\vec{V} \wedge \vec{b})^+, (\vec{a} \wedge \vec{W})^- + (\vec{V} \wedge \vec{b})^-) \quad \text{where} \quad \vec{V}, \, \vec{W} \in \text{Span}(\vec{a}, \vec{b})^\perp \right\}$$
$$= TS_+^2 \oplus TS_-^2 \,.$$

We denote respectively $\vec{G}^+ := \pi_+(\Pi(\vec{\Lambda})) = (\vec{a} \wedge \vec{b})^+$ and $\vec{G}^- := \pi_-(\Pi(\vec{\Lambda})) = (\vec{a} \wedge \vec{b})^-$. Denote by π_\perp the projection onto the span of $\{\vec{c}, \vec{d}\}$. First we introduce We equip the unit spheres of unit self-dual (resp. anti-self-dual) 2-vectors of \mathbb{R}^4 with the following complex structure

$$J_{(\vec{a}\wedge\vec{b})^{\pm}}((\vec{a}\wedge\vec{c})^{\pm}) = \pm (\vec{a}\wedge\vec{d})^{\pm}$$

Then we equip the product $S^2_+ \times S^2_-$ with the product complex structure

$$J_{\vec{G}\,\pm}(\vec{a}\wedge\vec{c}\pm\vec{d}\wedge\vec{b})=\pm[\vec{a}\wedge\vec{d}\mp\vec{c}\wedge\vec{b}]$$

For $\vec{X} \in \text{Span}(\vec{c}, \vec{d})$ we have $(\vec{a} \wedge \vec{X})^{\pm} = \sqrt{2^{-1}} (\vec{a} \wedge \vec{X} \pm \vec{X}^{\perp} \wedge \vec{b})$ where $\vec{X}^{\perp} := \star (\vec{a} \wedge \vec{b} \wedge X)$ or in other words $\vec{a} \wedge \vec{b} \wedge \vec{X} \wedge \vec{X}^{\perp} = |X|^2 \star 1$. With this notation we have

$$\forall \vec{X} \in \operatorname{Span}(\vec{c}, \vec{d}) \qquad J_{\vec{G}^{\pm}}\left((\vec{a} \wedge \vec{X})^{\pm}\right) = \pm (\vec{a} \wedge \vec{X}^{\perp})^{\pm}$$

Let $X, Y \in T_{\vec{\Lambda}} V_2(\mathbb{R}^4)$, we have¹⁴

$$-\left((\vec{a}\wedge\vec{b})^{+}\right)^{*}\omega_{S^{2}}\left(X,Y\right)=\left(d_{X}(\vec{a}\wedge\vec{b})^{+},J_{\vec{G}^{+}}(d_{Y}(\vec{a}\wedge\vec{b})^{+})\right)$$

where $d_X(\vec{a} \wedge \vec{b})^+$ is denoting the push forward by Π of the tangent vector X at (\vec{a}, \vec{b}) . Let $P^+(\vec{a}, \vec{b}) = \vec{a}$ and $P^-(\vec{a}, \vec{b}) = \vec{b}$ the push forward of X by P^+ (resp. by P^-) will simply be denoted $d_X\vec{a}$ (resp. $d_X\vec{b}$)

$$\begin{aligned} d_X(\vec{a} \wedge \vec{b})^+ &= \frac{1}{\sqrt{2}} (d_X(\vec{a} \wedge \vec{b}) + *d_X(\vec{a} \wedge \vec{b})) = d_X \vec{a} \cdot \vec{c} \ (\vec{c} \wedge \vec{b})^+ + d_X \vec{a} \cdot \vec{d} \ (\vec{d} \wedge \vec{b})^+ \\ &+ d_X \vec{b} \cdot \vec{c} \ (\vec{a} \wedge \vec{c})^+ + d_X \vec{b} \cdot \vec{d} \ (\vec{a} \wedge \vec{d})^+ \\ &= [d_X \vec{a} \cdot \vec{c} - d_X \vec{b} \cdot \vec{d}] \ (\vec{c} \wedge \vec{b})^+ + [d_X \vec{a} \cdot \vec{d} + d_X \vec{b} \cdot \vec{c}] \ (\vec{d} \wedge \vec{b})^+ \end{aligned}$$

Hence

$$\begin{split} J_{\vec{G}^{+}}(d_{Y}(\vec{a}\wedge\vec{b})^{+}) &= [d_{Y}\vec{a}\cdot\vec{c} - d_{Y}\vec{b}\cdot\vec{d}] \ J_{\vec{G}^{+}}(\vec{c}\wedge\vec{b})^{+} + [d_{Y}\vec{a}\cdot\vec{d} + d_{Y}\vec{b}\cdot\vec{c}] \ J_{\vec{G}^{+}}(\vec{d}\wedge\vec{b})^{+} \\ &= [d_{Y}\vec{a}\cdot\vec{c} - d_{Y}\vec{b}\cdot\vec{d}] \ (\vec{d}\wedge\vec{b})^{+} - [d_{Y}\vec{a}\cdot\vec{d} + d_{Y}\vec{b}\cdot\vec{c}] \ (\vec{c}\wedge\vec{b})^{+} \ . \end{split}$$

We deduce

$$\begin{pmatrix} d_X(\vec{a} \wedge \vec{b})^+, J_{\vec{G}^+}(d_Y(\vec{a} \wedge \vec{b})^+) \end{pmatrix}$$

$$= \left(\begin{bmatrix} d_X \vec{a} \cdot \vec{c} - d_X \vec{b} \cdot \vec{d} \end{bmatrix} (\vec{c} \wedge \vec{b})^+ + \begin{bmatrix} d_X \vec{a} \cdot \vec{d} + d_X \vec{b} \cdot \vec{c} \end{bmatrix} (\vec{d} \wedge \vec{b})^+,$$

$$\begin{bmatrix} d_Y \vec{a} \cdot \vec{c} - d_Y \vec{b} \cdot \vec{d} \end{bmatrix} (\vec{d} \wedge \vec{b})^+ - \begin{bmatrix} d_Y \vec{a} \cdot \vec{d} + d_Y \vec{b} \cdot \vec{c} \end{bmatrix} (\vec{c} \wedge \vec{b})^+ \right)$$

$$= -\begin{bmatrix} d_X \vec{a} \cdot \vec{c} - d_X \vec{b} \cdot \vec{d} \end{bmatrix} \begin{bmatrix} d_Y \vec{a} \cdot \vec{d} + d_Y \vec{b} \cdot \vec{c} \end{bmatrix} + \begin{bmatrix} d_X \vec{a} \cdot \vec{d} + d_X \vec{b} \cdot \vec{c} \end{bmatrix} \begin{bmatrix} d_Y \vec{a} \cdot \vec{c} - d_Y \vec{b} \cdot \vec{d} \end{bmatrix}$$

$$= -d_X \vec{a} \cdot \vec{c} \, d_Y \vec{a} \cdot \vec{d} + d_X \vec{a} \cdot \vec{d} \, d_Y \vec{a} \cdot \vec{c} - d_X \vec{b} \cdot \vec{c} \, d_Y \vec{b} \cdot \vec{d} + d_X \vec{b} \cdot \vec{d} \, d_Y \vec{b} \cdot \vec{c} - d_X \vec{a} \cdot d_Y \vec{b} + d_Y \vec{a} \cdot d_X \vec{b}$$

$$= - \star \begin{bmatrix} \vec{a} \wedge \vec{b} \wedge d_X \vec{a} \wedge d_Y \vec{a} \end{bmatrix} - \star \begin{bmatrix} \vec{a} \wedge \vec{b} \wedge d_X \vec{b} \wedge d_Y \vec{b} \end{bmatrix} - d_X \vec{a} \cdot d_Y \vec{b} + d_Y \vec{a} \cdot d_X \vec{b}$$

¹⁴The - sign comes from the fact that the relation between the Kähler metric and its associated Kähler form is given by $g(u, v) = \omega(u, Jv)$ and hence $-g(u, Jv) = \omega(u, v)$.

We have

$$d\vec{a} \wedge d\vec{a} = 2 \left(d\vec{a} \cdot \vec{b} \right) \wedge \left(d\vec{a} \cdot \vec{c} \right) \quad \vec{b} \wedge \vec{c} + 2 \left(d\vec{a} \cdot \vec{b} \right) \wedge \left(d\vec{a} \cdot \vec{d} \right) \quad \vec{b} \wedge \vec{d} + 2 \left(d\vec{a} \cdot \vec{c} \right) \wedge \left(d\vec{a} \cdot \vec{d} \right) \quad \vec{c} \wedge \vec{d}$$

Hence

$$\frac{1}{2}\,\vec{a}\wedge\vec{b}\wedge d\vec{a}\wedge d\vec{a} = (d\vec{a}\cdot\vec{c})\wedge(d\vec{a}\cdot\vec{d})\,\,\vec{a}\wedge\vec{b}\wedge\vec{c}\wedge\vec{d}\;.$$

Similarly

$$\frac{1}{2}\vec{a}\wedge\vec{b}\wedge d\vec{b} \wedge d\vec{b} = (d\vec{b}\cdot\vec{c})\wedge (d\vec{b}\cdot\vec{d}) \vec{a}\wedge\vec{b}\wedge\vec{c}\wedge\vec{d}.$$

Denote

$$d\vec{c} \wedge d\vec{d} := \sum_{l=1}^{4} dc^{l} \wedge dd^{l} , \qquad (\text{II.1})$$

there holds

$$d\vec{c} \wedge d\vec{d} = (d\vec{c} \cdot \vec{a}) \wedge (\vec{a} \cdot d\vec{d}) + (d\vec{c} \cdot \vec{b}) \wedge (\vec{b} \cdot d\vec{d}) = (\vec{c} \cdot d\vec{a}) \wedge (d\vec{a} \cdot \vec{d}) + (\vec{c} \cdot d\vec{b}) \wedge (d\vec{b} \cdot \vec{d}) .$$

Hence

$$d\vec{c} \wedge d\vec{d} = \frac{1}{2} \star_{\wedge \mathbb{R}^4} \vec{a} \wedge \vec{b} \wedge \left[d\vec{a} \wedge d\vec{a} + d\vec{b} \wedge d\vec{b} \right]$$

Thus

$$-\left((\vec{a}\wedge\vec{b})^{+}\right)^{*}\omega_{S^{2}}^{+}\left(X,Y\right) = -d\vec{c}\wedge d\vec{d}(X,Y) - d\vec{a}\wedge d\vec{b}(X,Y)$$

Similarly one computes

$$-\left((\vec{a}\wedge\vec{b})^{-}\right)^{*}\omega_{S^{2}}^{-}(X,Y)=\left(d_{X}(\vec{a}\wedge\vec{b})^{-},J_{\vec{G}^{-}}(d_{Y}(\vec{a}\wedge\vec{b})^{-})\right)$$

 but

$$\begin{aligned} d_X(\vec{a} \wedge \vec{b})^- &= \frac{1}{\sqrt{2}} (d_X(\vec{a} \wedge \vec{b}) - *d_X(\vec{a} \wedge \vec{b})) = d_X \vec{a} \cdot \vec{c} \ (\vec{c} \wedge \vec{b})^- + d_X \vec{a} \cdot \vec{d} \ (\vec{d} \wedge \vec{b})^- \\ &+ d_X \vec{b} \cdot \vec{c} \ (\vec{a} \wedge \vec{c})^- + d_X \vec{b} \cdot \vec{d} \ (\vec{a} \wedge \vec{d})^- \\ &= [d_X \vec{a} \cdot \vec{c} + d_X \vec{b} \cdot \vec{d}] \ (\vec{c} \wedge \vec{b})^- + [d_X \vec{a} \cdot \vec{d} - d_X \vec{b} \cdot \vec{c}] \ (\vec{d} \wedge \vec{b})^- \end{aligned}$$

Hence

$$\begin{aligned} J_{\vec{G}^{-}}(d_{Y}(\vec{a}\wedge\vec{b})^{-}) &= [d_{Y}\vec{a}\cdot\vec{c} + d_{Y}\vec{b}\cdot\vec{d}] \ J_{\vec{G}^{-}}(\vec{c}\wedge\vec{b})^{-} + [d_{Y}\vec{a}\cdot\vec{d} - d_{Y}\vec{b}\cdot\vec{c}] \ J_{\vec{G}^{-}}(\vec{d}\wedge\vec{b})^{-} \\ &= -[d_{Y}\vec{a}\cdot\vec{c} + d_{Y}\vec{b}\cdot\vec{d}] \ (\vec{d}\wedge\vec{b})^{-} + [d_{Y}\vec{a}\cdot\vec{d} - d_{Y}\vec{b}\cdot\vec{c}] \ (\vec{c}\wedge\vec{b})^{-} \ . \end{aligned}$$

We deduce

$$\begin{pmatrix} d_X(\vec{a}\wedge\vec{b})^-, J_{\vec{G}^-}(d_Y(\vec{a}\wedge\vec{b})^-) \end{pmatrix} = [d_X\vec{a}\cdot\vec{c} + d_X\vec{b}\cdot\vec{d}] [d_Y\vec{a}\cdot\vec{d} - d_Y\vec{b}\cdot\vec{c}]$$

$$-[d_X\vec{a}\cdot\vec{d} - d_X\vec{b}\cdot\vec{c}] [d_Y\vec{a}\cdot\vec{c} + d_Y\vec{b}\cdot\vec{d}]$$

$$= d_X\vec{a}\cdot\vec{c} d_Y\vec{a}\cdot\vec{d} - d_X\vec{a}\cdot\vec{d}d_Y\vec{a}\cdot\vec{c} + d_X\vec{b}\cdot\vec{c} d_Y\vec{b}\cdot\vec{d} - d_X\vec{b}\cdot\vec{d}d_Y\vec{b}\cdot\vec{c} - d_X\vec{a}\cdot d_Y\vec{b} + d_Y\vec{a}\cdot d_X\vec{b}$$

$$= \star[\vec{a}\wedge\vec{b}\wedge d_X\vec{a}\wedge d_Y\vec{a}] + \star[\vec{a}\wedge\vec{b}\wedge d_X\vec{b}\wedge d_Y\vec{b}] - d_X\vec{a}\cdot d_Y\vec{b} + d_Y\vec{a}\cdot d_X\vec{b} .$$

Hence

$$-\left((\vec{a}\wedge\vec{b})^{-}\right)^{*}\omega_{S^{2}}^{-}(X,Y) = d\vec{c}\wedge d\vec{d}(X,Y) - d\vec{a}\wedge d\vec{b}(X,Y) \ .$$

We deduce the following lemma

Lemma II.1. [The contact Form] Let $(\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$ and $(\vec{c}, \vec{d}) \in V_2(\mathbb{R}^4)$ such that

$$*\vec{a}\wedge\vec{b}=\vec{c}\wedge\vec{b}$$

Then the following identities hold

$$\left((\vec{a}\wedge\vec{b})^{\pm}\right)^{*}\omega_{S^{2}}^{\pm}=\pm d\vec{c}\wedge d\vec{d}+d\vec{a}\wedge d\vec{b}$$

and

$$d\vec{c} \wedge d\vec{d} = \frac{1}{2} \star_{\wedge \mathbb{R}^4} \vec{a} \wedge \vec{b} \wedge \left[d\vec{a} \wedge d\vec{a} + d\vec{b} \wedge d\vec{b} \right]$$

 $In \ particular^{15}$

$$d\alpha = 2 \, d\vec{a} \wedge d\vec{b} = \left((\vec{a} \wedge \vec{b})^+ \right)^* \omega_{S^2} + \left((\vec{a} \wedge \vec{b})^- \right)^* \omega_{S^2} = \Pi^* \left(\pi_+^* \omega_{S^2}^+ + \pi_-^* \omega_{S^2}^- \right) = \Pi^* \omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}}$$

and

$$\alpha \wedge d\alpha \wedge d\alpha = 2 \alpha \wedge \left((\vec{a} \wedge \vec{b})^+ \right)^* \omega_{S^2}^+ \wedge \left((\vec{a} \wedge \vec{b})^- \right)^* \omega_{S^2}^- = 2 \alpha \wedge \Pi^* \pi_+^* \omega_{S^2}^+ \wedge \Pi^* \pi_-^* \omega_{S^2}^-$$

We introduce the Reeb Vector Field

$$\vec{R} := \frac{\partial}{\partial \theta} (\vec{a}^{\,\theta}, \vec{b}^{\,\theta})_{\theta=0} := \left. \frac{d}{d\theta} (\cos\theta \, \vec{a} + \sin\theta \, \vec{b} \, , \, -\sin\theta \, \vec{a} + \cos\theta \, \vec{b}) \right|_{\theta=0} = \left(\vec{b}, -\vec{a} \right)$$

then

$$\alpha \wedge d\alpha \wedge d\alpha \bigsqcup \vec{R} = -2 \Pi^* \left[\pi_+^* \omega_{S^2} \wedge \pi_-^* \omega_{S^2} \right] = -2 \Pi^* \Omega_{S_+^2 \times S_+^2}$$

where $\Omega_{S^2_+ \times S^2_-}$ is the volume form on $S^2_+ \times S^2_-$. Denote by $H_{(\vec{a},\vec{b})} := \operatorname{Ker}(\Pi_*)^{\perp} = \operatorname{Span}(\vec{R})^{\perp}$. We have

$$H_{(\vec{a},\vec{b})} = \left\{ (\vec{V},\vec{W}) \in \mathbb{R}^4 \times \mathbb{R}^4 \ ; \ \vec{V},\vec{W} \in \operatorname{Span}(\vec{a},\vec{b})^{\perp} \right\}$$

and

$$\Pi_*(\vec{V}, \vec{W}) = ((\vec{V} \land \vec{b})^+ + (\vec{a} \land \vec{W})^+, (\vec{V} \land \vec{b})^- + (\vec{a} \land \vec{W})^-)$$

Thus Π_* realises an isometry between $H_{(\vec{a},\vec{b})}$ and $T_{\Pi(\vec{a},\vec{b})}S^2_+ \times S^2_-$. On H we define the complex structure J_H such that

$$\forall \, \vec{X} \in H \quad \Pi_* J_H \vec{X} := J_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}} \Pi_* \vec{X} \; .$$

This gives in particular

$$\forall \vec{X} , \vec{Y} \in H \quad \vec{Y} \cdot J_H \vec{X} = \omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}} (\vec{X}, \vec{Y}) \;.$$

Finally we are proving a last preliminary result

Lemma II.2. [The transverse Complex structure] Under the previous notations, $(\vec{V}, \vec{W}) \in H_{(\vec{a}, \vec{b})}$ being given there holds

$$J_H(\vec{V},\vec{W}) = (-\vec{W},\vec{V}) \ .$$

Proof of Lemma II.2 It remains to prove the last assertion on J_H . Let $(\vec{V}, \vec{W}) \in H_{(\vec{a}, \vec{b})}$ we have

$$\Pi_* J_H(\vec{V}, \vec{W}) = J_{\vec{a} \wedge \vec{b}} \Pi_*(\vec{V}, \vec{W}) = J_{\vec{a} \wedge \vec{b}} ((\vec{V} \wedge \vec{b})^+ + (\vec{a} \wedge \vec{W})^+, (\vec{V} \wedge \vec{b})^- + (\vec{a} \wedge \vec{W})^-)$$
$$= ((\vec{V}^\perp \wedge \vec{b})^+ + (\vec{a} \wedge \vec{W}^\perp)^+, -(\vec{V}^\perp \wedge \vec{b})^- - (\vec{a} \wedge \vec{W}^\perp)^-)$$

¹⁵Observe that $\pi_{+}^{*}\omega_{S^{2}} + \pi_{-}^{*}\omega_{S^{2}} = \pi_{+}^{*}\omega_{\mathbb{C}P^{1}} - \pi_{-}^{*}\omega_{\mathbb{C}P^{1}}$.

We have respectively

$$\begin{split} (\vec{V}^{\perp} \wedge \vec{b})^{+} &= \frac{1}{2} \left[\vec{V}^{\perp} \wedge \vec{b} + *(\vec{V}^{\perp} \wedge \vec{b}) \right] = \frac{1}{2} \left[\vec{V}^{\perp} \wedge \vec{b} + \vec{a} \wedge \vec{V} \right] = (\vec{a} \wedge \vec{V})^{+} \\ (\vec{a} \wedge \vec{W}^{\perp})^{+} &= \frac{1}{2} \left[\vec{a} \wedge \vec{W}^{\perp} + *(\vec{a} \wedge \vec{W}^{\perp}) \right] = \frac{1}{2} \left[\vec{a} \wedge \vec{W}^{\perp} + \vec{b} \wedge \vec{W} \right] = (\vec{b} \wedge \vec{W})^{+} \\ -(\vec{V}^{\perp} \wedge \vec{b})^{-} &= -\frac{1}{2} \left[\vec{V}^{\perp} \wedge \vec{b} - *(\vec{V}^{\perp} \wedge \vec{b}) \right] = -\frac{1}{2} \left[\vec{V}^{\perp} \wedge \vec{b} - \vec{a} \wedge \vec{V} \right] = (\vec{a} \wedge \vec{V})^{-} \\ -(\vec{a} \wedge \vec{W}^{\perp})^{-} &= -\frac{1}{2} \left[\vec{a} \wedge \vec{W}^{\perp} - *(\vec{a} \wedge \vec{W}^{\perp}) \right] = -\frac{1}{2} \left[\vec{a} \wedge \vec{W}^{\perp} - \vec{b} \wedge \vec{W} \right] = (\vec{b} \wedge \vec{W})^{-} \end{split}$$

Thus

$$\Pi_* J_H(\vec{V}, \vec{W}) = \left((\vec{a} \wedge \vec{V})^+ + (\vec{b} \wedge \vec{W})^+, (\vec{a} \wedge \vec{V})^- + (\vec{b} \wedge \vec{W})^- \right) = \Pi_* (-\vec{W}, \vec{V})$$
(II.2)

II.2 The space of $W^{2,4}$ Legendrian immersions of a surface

Let Σ be a closed oriented 2-dimensional manifold. In [17] section II the author introduces the space of Sobolev Immersions

$$\mathfrak{E}_{\Sigma}^{2,4} := W_{imm}^{2,4}(\Sigma, V_2(\mathbb{R}^4)) := \left\{ \vec{\Lambda} \in W^{2,4}(\Sigma, V_2(\mathbb{R}^4)) \text{ s. t. } \operatorname{rank}(d\vec{\Lambda})(x) = 2 \quad \forall x \in \Sigma \right\}$$

which is proven to be a C^2 submanifold of the Banach vector space $W^{2,4}(\Sigma, \mathbb{R}^4 \times \mathbb{R}^4)$; moreover [17] introduces a Finsler Structure over $\mathfrak{E}_{\Sigma}^{2,4}$ for which the associated Palais distance defines a complete metric space structure on $\mathfrak{E}_{\Sigma,2}$. We now prove the following theorem.

Theorem II.1. The space of $W^{2,4}$ Legendrian immersions of a closed oriented surface Σ into $V_2(\mathbb{R}^4)$ (i.e. elements from $\mathfrak{E}_{\Sigma}^{2,4}$ satisfying $\vec{\Lambda}^* \alpha = 0$) realises a C^2 sub-manifold of $W_{imm}^{2,4}(\Sigma, V_2(\mathbb{R}^4))$.

Proof of theorem II.1. We consider the map

$$\mathfrak{H} : \vec{\Lambda} \in \mathfrak{E}^{2,4}_{\Sigma} \longrightarrow \vec{\Lambda}^* \alpha \in \Gamma_{W^{1,4}}(T^*\Sigma) .$$

We claim that \mathfrak{H} realizes a C^2 submersion from the Banach manifold $\mathfrak{E}_{\Sigma}^{2,4}$ onto the Banach vector space of $W^{1,4}$ one forms on Σ . Moreover there holds

$$\forall \ \vec{\Lambda} = (\vec{\Phi}, \vec{n}) \in \mathfrak{E}_{\Sigma}^{2,4} \quad \forall \ (\vec{V}, \vec{W}) \in W^{2,4}(\Sigma, \mathbb{R}^4 \times \mathbb{R}^4) \ ; \quad \vec{V} \cdot \vec{\Phi} = 0 \ , \ \vec{W} \cdot \vec{n} = 0 \ \text{ and } \ \vec{\Phi} \cdot \vec{W} + \vec{n} \cdot \vec{V} = 0$$
$$d\mathfrak{H}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W}) = \vec{V} \cdot d\vec{n} + \vec{\Phi} \cdot d\vec{W} - \vec{n} \cdot d\vec{V} - \vec{W} \cdot d\vec{\Phi} \ .$$

Indeed, first, \mathfrak{H} is the restriction to the C^2 sub-manifold $\mathfrak{E}_{\Sigma}^{2,4}$ of the Banach vector space $W^{2,4}(\Sigma, \mathbb{R}^4 \times \mathbb{R}^4)$ of the (obviously C^{∞}) bilinear form

$$B(\vec{a}, \vec{b}) := \vec{a} \cdot d\vec{b} - \vec{b} \cdot d\vec{a}$$

It remains to prove in a second step that \mathfrak{H} realises a submersion that is at every point $\vec{\Lambda} = (\vec{\Phi}, \vec{n}) \in \mathfrak{E}_{\Sigma}^{2,4}$ the linear map $d\mathfrak{H}_{\vec{\Lambda}}$ is surjective from $T_{\vec{\Lambda}}\mathfrak{E}_{\Sigma}^{2,4}$ onto $\Gamma_{W^{1,4}}(T^*\Sigma)$.

Instead of proving it directly, we introduce the map

$$\mathfrak{K} : \vec{\Lambda} \in \mathfrak{E}_{\Sigma}^{2,4} \longrightarrow \left(\vec{\Lambda}^* d\alpha \,, \, \Pi_0(\vec{\Lambda}^* \alpha) \,, \, \int_{\Gamma_1} \vec{\Lambda}^* \alpha, \cdots \,, \int_{\Gamma_{2\,g(\Sigma)}} \vec{\Lambda}^* \alpha \right) \,,$$

where $\Pi_0(\vec{\Lambda}^*\alpha)$ is the exact part in the Hodge decomposition of $\vec{\Lambda}^*\alpha$ with respect to the smooth reference metric g_0

$$\Pi_0(\vec{\Lambda}^*\alpha) := d^{*_{g_0}} \Delta_{g_0}^{-1} \left[\vec{\Lambda}^*\alpha \right]$$

and Γ_j is a fixed basis of closed loops generating $H_1(\Sigma, \mathbb{Z})$. We have that

$$\mathfrak{K} : \mathfrak{E}^{2,4}_{\Sigma} \longrightarrow W^{1,4}(\wedge^2 \Sigma) \times W^{2,4}(\Sigma) \times \mathbb{R}^{2g(\Sigma)} .$$

Since \mathfrak{H} is $C^1 \mathrm{we}$ deduce that \mathfrak{K} is also C^1 and one has

$$\forall \ \vec{\Lambda} = (\vec{\Phi}, \vec{n}) \in \mathfrak{E}_{\Sigma}^{2,4} \quad \forall \ (\vec{V}, \vec{W}) \in W^{2,4}(\Sigma, \mathbb{R}^4 \times \mathbb{R}^4) \ ; \quad \vec{V} \cdot \vec{\Phi} = 0 \ , \ \vec{W} \cdot \vec{n} = 0 \ \text{ and } \ \vec{\Phi} \cdot \vec{W} + \vec{n} \cdot \vec{V} = 0$$

$$d\mathfrak{K}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W}) = \left(2 \, d\vec{V} \wedge d\vec{n} + 2 \, d\vec{\Phi} \wedge d\vec{W}, \Pi_0(d\mathfrak{H}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W})), 2 \, \int_{\Gamma_1} \vec{V} \cdot d\vec{n} - \vec{W} \cdot d\vec{\Phi} \cdots 2 \, \int_{\Gamma_{2\,g(\Sigma)}} \vec{V} \cdot d\vec{n} - \vec{W} \cdot d\vec{\Phi} \right)$$

Let $(\Omega, \phi, s_1 \cdots s_{2g(\Sigma)}) \in W_0^{1,4}(\wedge^2 \Sigma) \times W^{2,4}(\Sigma) \times \mathbb{R}^{2g(\Sigma)}$ where $W_0^{1,4}(\wedge^2 \Sigma)$ is the space of $W^{2,4}$ -forms on Σ with integral 0. Assuming $\vec{\Lambda}^* \alpha = 0$, we are looking for $(\vec{V}, \vec{W}) \in W^{2,4}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4))$ such that

$$d\mathfrak{K}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W}) = (\Omega, \phi, s_1 \cdots s_{2g(\Sigma)}) \; .$$

For any $\vec{\Lambda} = (\vec{\Phi}, \vec{n}) \in W^{2,4}_{imm}(\Sigma, V_2(\mathbb{R}^4))$ we introduce the corresponding Gauss Map

$$\vec{T} := \frac{\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}}{|\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}|} \in W^{1,4}(\Sigma, G_2(TV_2(\mathbb{R}^4))) \hookrightarrow C^{0,1/2}(\Sigma, G_2(TV_2(\mathbb{R}^4)))$$

Claim : There exists a radius r > 0 depending only on the $W^{1,4}$ norm of \vec{T} such that on any ball¹⁶ $B_r(x_0)$ for x_0 in Σ the image of $B_r(x_0)$ by $\vec{\Lambda}$ is injective and described by a local graph in $V_2(\mathbb{R}^4)$: that is there exists a local chart $(y_1, y_2, y_3, y_4, y_5)$ in an open set \mathcal{O} containing $\vec{\Lambda}(x)$ such that

$$\bar{\Lambda}(B_r(x)) = \{(y_1, y_2, f(y_1, y_2), g(y_1, y_2), h(y_1, y_2)) \in y(\omega) \ ; \ (y_1, y_2) \in U\}$$

where U is an open set in \mathbb{R}^2 . Moreover $f, g, h \in W^{2,4}(U, \mathbb{R})$.

Proof of the claim : We can produce this chart by taking smooth coordinates (y_1, y_2) in the tangent plate $\vec{\Lambda}_* T_{x_0} \Sigma$ with $y_1(\vec{\Lambda}(x)) = 0$ and $y_2(\vec{\Lambda}(x)) = 0$ that we complete in order to have a local chart inn the neighborhood of $\vec{\Lambda}(x)$. Using the local inversion theorem we have that the map which assigns to any point $x \in B_r(x_0)$ the two first coordinates $\tilde{\Lambda}(x) := (\Lambda_1(x), \Lambda_2(x))$ of $\vec{\Lambda}(x)$ realizes a C^1 diffeomorphism between $B_r(x)$ and U, moreover since $\det(d\tilde{\Lambda}) \in C^0(B_r(x))$ and since $\det(d\tilde{\Lambda})(x_0) > 0$, for r > 0 small enough we have the existence of C > 1 such that

$$\forall x \in B_r(x_0) \quad C^{-1} \le \det(d\tilde{\Lambda})(x) \le C$$

Moreover using the chain rule in Sobolev space we have

$$\partial_{y_i y_j}^2 \tilde{\Lambda}^{-1} = \partial_{y_i} \left[(d\tilde{\Lambda})_j^{-1} \circ \tilde{\Lambda}^{-1}(y) \right] = \partial_{x_l} (d\tilde{\Lambda})_j^{-1} \ \partial_{y_j} (\tilde{\Lambda}^{-1})_l(y) \ .$$

Hence

$$\int_{B_r(x)} |\partial_{y_i y_j}^2 \tilde{\Lambda}^{-1}|^4 (\tilde{\Lambda}(x)) \, \det(d\tilde{\Lambda})(x) \, dx^2 \le C \, \int_{B_r(x)} |\nabla_x (d\tilde{\Lambda})^{-1}|^4 (x) \, \|d(\tilde{\Lambda})^{-1}\|_{\infty}^4 dx^2 < +\infty$$

¹⁶We have fixed a reference smooth metric g_0 on Σ with respect to which the spaces $W^{k,p}(\Sigma)$ are defined and we take the geodesic balls with respect to this metric.

We deduce using the coarea formula that $\tilde{\Lambda}^{-1} \in W^{2,4}(U, B_r(x))$ and we take respectively

$$f(y_1, y_2) := \Lambda_3 \circ \tilde{\Lambda}^{-1}(y_1, y_2) \ , \ g(y_1, y_2) := \Lambda_4 \circ \tilde{\Lambda}^{-1}(y_1, y_2) \ \text{ and } \ h(y_1, y_2) := \Lambda_5 \circ \tilde{\Lambda}^{-1}(y_1, y_2) \ .$$

Using Chain rules (successively the classical one of C^1 functions and then the one in Sobolev Spaces) we have for any m = 1, 2, 3 and i, j = 1,

$$\partial_{y_i y_j}^2 [\Lambda_m \circ \tilde{\Lambda}^{-1}] = \partial_{x_l x_k}^2 \Lambda_m \circ \tilde{\Lambda}^{-1} \ \partial_{y_i} \tilde{\Lambda}_l^{-1} \ \partial_{y_j} \tilde{\Lambda}_k^{-1} + \partial_{x_l} \Lambda_m \circ \tilde{\Lambda}^{-1} \ \partial_{y_i y_j} \tilde{\Lambda}_l^{-1}$$

and using again the co-area formula we have that f, g, h are all three in $W^{2,4}(U)$. This ends the proof of the claim.

We fix a finite covering of Σ by balls $(B_r(x_i))_{i=1\cdots Q}$ and we consider an adapted partition of unity made of C^{∞} non negative functions χ_i with support in $B_r(x_i)$

$$\sum_{i=1}^{Q} \chi_i \equiv 1$$

There exists $\beta \in W^{2,4}(\wedge^1 \Sigma)$ such that $d\beta = \Omega$. Let $(h_k^0)_{k=1\cdots 2 g(\Sigma)}$ be a basis of harmonic 1 forms for g_0 generating $H^1(\Sigma)$. These forms are known to b smooth. We modify possibly β by adding the linear combination of h_k^0 such that

$$\forall k = 1 \cdots 2 g(\Sigma) \qquad \int_{\Gamma_k} \beta = s_k .$$
 (II.3)

We write $\Omega_i := d(\chi_i \beta) \in W_0^{1,4}(\wedge^1 B_r(x_i))$ and

$$\Omega_i = \omega_i(x) \ dx_1 \wedge dx_2$$

Hence

$$(\tilde{\Lambda}^{-1})^*\Omega_i = \omega_i(\tilde{\Lambda}^{-1}(y)) \det(\nabla_y \tilde{\Lambda}^{-1}) dy_1 \wedge dy_2 = \tilde{\omega}_i(y) dy_1 \wedge dy_2$$

where we have $\tilde{\omega}_i \in W_0^{1,4}(U)$ (using again chain rule in Sobolev spaces as well as co-area formula). Let $\gamma_i = (\tilde{\Lambda}^{-1})^*(\chi_i \beta) \in W_0^{1,4}(\wedge^1 U)$. We have

$$(\tilde{\Lambda}^{-1})^* \Omega_i = d\gamma_i$$
.

In particular $\int_U (\tilde{\Lambda}^{-1})^* \Omega_i = 0$. Let $\xi_i \in W^{3,4}(U)$ solving

$$\begin{cases} \Delta \xi_i = \tilde{\omega}_i(y) & \text{in } \mathcal{D}'(U) \\ \frac{\partial \xi_i}{\partial \nu} = 0 & \text{on } \partial U . \end{cases}$$

We have in particular $d(*d\xi_i) = (\tilde{\Lambda}^{-1})^* \Omega_i$ and $*d\xi_i$ can continuously be extended by 0 outside U. Observe that on U we have the existence of ϕ_i such that

$$\chi_i \circ (\tilde{\Lambda}^{-1}) \ (\tilde{\Lambda}^{-1})^* \beta = *d\xi_i + d\phi_i$$

On $U \setminus \operatorname{supp}(\chi_i \circ (\tilde{\Lambda}^{-1}))$ both ξ_i and ϕ_i are harmonic. We take a C^{∞} regularisation $\overline{\phi}_i$ of ϕ_i such that $\overline{\phi}_i - \phi_i$ is equal to zero in an open neighbourhood of ∂U in U and we have that $\sigma_i := *d\xi_i + d\overline{\phi}_i$ is $W^{2,4}$ in U with support strictly inside U and satisfies

$$d\sigma_i = \Omega_i$$

Recall that on $V_2(\mathbb{R}^4)$ we have

$$\alpha \wedge d\alpha \wedge d\alpha > 0$$

We have seen that the restriction to the Horizontal plane H of $d\alpha$ is non degenerate and that

$$\operatorname{Ker}(d\alpha)(\vec{a},\vec{b}) = \operatorname{Span}\left\{(\vec{b},-\vec{a})\right\} = \operatorname{Span}\left\{\vec{R}\right\}$$

while in the opposite way

$$\operatorname{Ker}(\alpha)(\vec{a}, \vec{b}) = H_{(\vec{a}, \vec{b})}$$
 and $\alpha(\vec{R}) = -2$.

Hence for any $(\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$

$$\Xi_{\vec{a},\vec{b}} : \vec{X} = (\vec{V},\vec{W}) \longrightarrow d\alpha \, \square \, \vec{X} + \alpha(\vec{X}) \, \alpha$$

realises an isomorphism from $T_{(\vec{a},\vec{b})}V_2(\mathbb{R}^4)$ into $T^*_{(\vec{a},\vec{b})}V_2(\mathbb{R}^4)$. The explicit form of Ξ^{-1} is the following : Let $\gamma \in \Gamma(T^*V_2(\mathbb{R}^4))$ we take V

$$\vec{X} = \vec{X}^H + \vec{X}$$

where

$$\left\{ \begin{array}{ll} \vec{X}^V(\vec{a},\vec{b}) := 2^{-1} \, < \alpha, \gamma > \ \vec{R}(\vec{a},\vec{b}) \\ \\ \vec{X}^H(\vec{a},\vec{b}) \in H_{(\vec{a},\vec{b})} \quad \text{s. t.} \quad \forall \vec{Y} \in H_{(\vec{a},\vec{b})} \quad \vec{Y} \cdot \vec{X}^H := d\alpha(\vec{X}^H, J_H \vec{Y}) = \gamma(\vec{Y}) \end{array} \right.$$

where we have used that on H the restriction of the two form $d\alpha$ is non degenerate $d\alpha \wedge d\alpha > 0$ on H. The explicit forms above imply $\Xi_{(\vec{a},\vec{b})}$ as a map from $\Gamma(TV_2(\mathbb{R}^4))$ into $\Gamma(T^*V_2(\mathbb{R}^4))$ as well as $\Xi_{(\vec{a},\vec{b})}^{-1}$ are obviously smooth.

We denote

$$\vec{X}_i \circ y^{-1} = (\vec{V}_i \circ y^{-1}, \vec{W}_i \circ y^{-1}) = \Xi_{y^{-1}(y_1, y_2, y_3, y_4, y_5)}^{-1}(y^* \sigma_i)$$

It satisfies

$$y^*\sigma_i = 2\left[\vec{V}_i \cdot d\vec{b} - \vec{W}_i \cdot d\vec{a}\right] + \left(\vec{a} \cdot \vec{W}_i - \vec{b} \cdot \vec{V}_i\right)\left[\vec{a} \cdot d\vec{b} - \vec{b} \cdot d\vec{a}\right]$$

The restriction of \vec{X}_i to $y \circ \vec{\Lambda}(B_r(x_i))$ satisfies

$$\vec{X}_i \circ y^{-1} = (\vec{V}_i \circ y^{-1}, \vec{W}_i \circ y^{-1}) = \Xi_{y^{-1}(y_1, y_2, f(y_1, y_2), g(y_1, y_2), h(y_1, y_2))}^{-1}(y^* \sigma_i)$$

Since y and y^{-1} are smooth, since f, g, h are in $W^{2,4}(U)$ and since $\sigma_i \in W^{2,4}(\wedge^1 U)$, the chain rule for Sobolev maps combined with the area formula as above gives that the extensions by zero outside $B_r(x_i)$ of $(\vec{V}_i \circ \vec{\Lambda}, \vec{W}_i \circ \vec{\Lambda})$ belong to $W^{2,4}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4))$. By restricting this identity on $\vec{\Lambda}(B_r(x_i))$, assuming $\vec{\Lambda}$ is Legendrian we obtain

$$\Omega_i = d \left[\vec{\Lambda}^* y^* \sigma_i \right] = 2 \left[d [\vec{V}_i \circ \vec{\Lambda}] \dot{\wedge} d\vec{n} - d [\vec{W}_i \circ \vec{\Lambda}] \dot{\wedge} d\vec{\Phi} \right]$$

Summing over i gives

$$(\vec{V}, \vec{W}) = \sum_{i} (\vec{V}_i \circ \vec{\Lambda}, \vec{W}_i \circ \vec{\Lambda}) \text{ in } W^{2,4}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4))$$

and we have

$$\Omega = 2 \left[d\vec{V} \dot{\wedge} d\vec{n} - d\vec{W} \dot{\wedge} d\vec{\Phi} \right] \; . \label{eq:Gamma-state}$$

Observe that

$$\sigma_i = \chi_i \circ (\tilde{\Lambda}^{-1}) \ (\tilde{\Lambda}^{-1})^* \beta + d(\phi_i - \overline{\phi}_i)$$

where $\phi_i - \overline{\phi}_i$ is supported i the open set U_i . Hence each $(\phi_i - \overline{\phi}_i) \circ \tilde{\Lambda}$ extends by 0 outside $B_r(x_i)$ as a $W^{1,4}$ function on the whole Σ and we have

$$2\left[\vec{V}\cdot d\vec{n} - \vec{W}\cdot d\vec{\Phi}\right] = \beta + d\left[\sum_{i}\phi_{i} - \overline{\phi}_{i}\right]$$

Recall that

$$d\mathfrak{H}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W}) = 2 \left[\vec{V} \cdot d\vec{n} - \vec{W} \cdot d\vec{\Phi} \right] + d \left[\vec{\Phi} \cdot \vec{W} - \vec{n} \cdot \vec{V} \right]$$

Observe moreover that for any $f \in W^{2,4}(\Sigma)$ there holds

$$d\mathfrak{H}_{\vec{\Lambda}} \cdot (-f \, \vec{n}, f \, \vec{\Phi}) = 2 \, df$$
.

Consider then f such that

$$2\,df = d^{*_{g_0}} \Delta_{g_0}^{-1} \left[d\mathfrak{H}_{\vec{\Lambda}} \cdot (\vec{V}, \vec{W}) \right] - d\phi \in W^{1,4}(\wedge^1 \Sigma)$$

This gives

$$d^{*_{g_0}}\Delta_{g_0}^{-1}\left[d\mathfrak{H}_{\vec{\Lambda}}\cdot(\vec{V}+f\vec{n},\vec{W}-f\vec{\Phi})\right]=d\phi$$

where we have used the fact that

$$d^{*_{g_0}}\Delta_{g_0}^{-1}[df] = df$$

Observe moreover

$$2\left[d[\vec{V}+f\vec{n}]\dot{\wedge}d\vec{n}-d[\vec{W}-f\vec{\Phi}]\dot{\wedge}d\vec{\Phi}\right] = 2\left[d\vec{V}\dot{\wedge}d\vec{n}-d\vec{W}\dot{\wedge}d\vec{\Phi}\right] = \Omega$$

and

$$\forall k = 1 \cdots 2g(\Sigma) \qquad 2 \int_{\Gamma_k} (\vec{V} + f\vec{n}) \cdot d\vec{n} - (\vec{W} - f\vec{\Phi}) \cdot d\vec{\Phi} = s_k \; .$$

Hence we have proved that if $\vec{\Lambda}$ is a $W^{2,4}$ Legendrian immersion into $V_2(\mathbb{R}^4)$ we have

$$\forall (\Omega, \phi, s_1, \cdots, s_{2g(\Sigma)}) \in W^{1,4}(\wedge^2 \Sigma) \times W^{2,4}(\Sigma) \times \mathbb{R}^{2g(\Sigma)}$$

$$\exists \, (\vec{V},\vec{W}) \in \Gamma_{W^{2,4}}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4)) \qquad d\mathfrak{K}_{\vec{\Lambda}}(\vec{V},\vec{W}) = (\Omega,\phi,s_1,\cdots,s_{2\,g(\Sigma)}) \ .$$

This implies that \mathfrak{K} realises s submersion at every point $\vec{\Lambda} \in \mathfrak{E}_{\Sigma}^{2,4}$ satisfying $\vec{\Lambda}^* \alpha = 0$. We claim that at a point $\vec{\Lambda} \in \mathfrak{E}_{\Sigma}^{2,4}$ satisfying $\vec{\Lambda}^* \alpha = 0$ the Kernel of $d\mathfrak{K}_{\vec{\Lambda}}$ is closed in $\Gamma_{W^{2,4}}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4))$ and splits, that is there exists a closed supplement of Ker $d\mathfrak{K}_{\vec{\Lambda}}$. We have

$$(\vec{V}, \vec{W}) \in \operatorname{Ker} d\mathfrak{K}_{\vec{\Lambda}} \quad \Longleftrightarrow \quad \vec{V} \cdot d\vec{n} - \vec{W} \cdot d\vec{\Phi} = 0 \quad \text{on } \Sigma$$

The Splitting of Ker $d\mathfrak{K}_{\vec{\Lambda}}$ can be seen by taking the intersection of its orthogonal complement in $L^2(\Sigma, \mathbb{R}^8)$ with $\Gamma_{W^{2,4}}(\vec{\Lambda}^{-1}TV_2(\mathbb{R}^4))$

Observe that

$$\mathfrak{K}(\vec{\Lambda}) = 0 \qquad \Longleftrightarrow \qquad \vec{\Lambda}^* \alpha = 0$$

Hence we have proved that the space of $W^{2,4}$ Legendrian immersions of a surface Σ is a C^2 sub-manifold of $\mathfrak{E}_{\Sigma}^{2,4}$ and this concludes the proof of theorem II.1.

We equip the tangent bundle of $\mathfrak{E}_{\Sigma,Leq}^{2,4}$ with the Finsler structure introduced in [17] (Proposition II.2).

$$\forall \ \vec{w} \in T_{\vec{\Lambda}} \mathfrak{E}^{2,4}_{\Sigma,Leg} \qquad \|\vec{w}\|_{\vec{\Lambda}} := \left[\int_{\Sigma} \left[|\nabla^2 \vec{w}|^2 + |\nabla \vec{w}|^2 + |\vec{w}|^2 \right]^2 \ dvol_{g_{\vec{\Lambda}}} \right]^{1/4} + \||\nabla \vec{w}|_{g_{\vec{\Lambda}}}\|_{L^{\infty}(\Sigma)}$$

The restriction to $\mathfrak{E}_{\Sigma,Leg}^{2,4}$ of the same Finsler structure is inducing a Palais distance d_{Leg} which is clearly larger than the restriction to $\mathfrak{E}_{\Sigma,Leg}^{2,4}$ of the Palais distance d on $\mathfrak{E}_{\Sigma}^{2,4}$. Hence since this last one is complete on $\mathfrak{E}_{\Sigma}^{2,4}$ (see [17] proposition II.3) and since the membership to $\mathfrak{E}_{\Sigma,Leg}^{2,4}$ passes to the limit under distance control, $(\mathfrak{E}_{\Sigma,Leg}^{2,4}, d_{Leg})$ is also complete (i.e. Cauchy sequences for d_{Leg} are obviously Cauchy for d).

We recall the definition of homotopic admissible families or simply admissible family.

Definition II.1. Let \mathfrak{M} be a Banach manifold. A subset \mathcal{A} of the power set of \mathfrak{M} (i.e. A is a set of subsets of \mathfrak{M}) is called admissible family if it is stable under isotopy that is : forall $H \in C^0([0,1] \times \mathfrak{M}, \mathfrak{M})$ such that H_t is an homeomorphism of \mathfrak{M} for any $t \in [0,1]$ and for any $A \in \mathcal{A}$ we have $H_1(A) \in \mathcal{A}$.

We recall the proposition which was central in the viscosity method introduced in [17] and more precisely the version given in [14] (which permits to avoid the use of the Palais-Smale condition which should hold for our problem but whose proof might be a bit tedious in the present context).

Proposition II.1. Let E_{ε} be a family of C^1 functionals on a complete Finsler manifold \mathfrak{M} , with E_{ε} differentiable with respect to ε and $\varepsilon \to E_{\varepsilon}(x)$, $\varepsilon \to \frac{dE_{\varepsilon}(x)}{d\varepsilon}$ both increasing in ε for any $x \in \mathfrak{M}$. Assume also that

$$\|dE_{\varepsilon_i}(x_i) - dE_{\varepsilon}(x_i)\|_{T^*\mathfrak{M}} \longrightarrow 0 , \qquad (\text{II.4})$$

whenever $0 < \varepsilon \leq \partial_i$, $\varepsilon_i \to \varepsilon$ and $\limsup_{i \to +\infty} E_{\varepsilon}(x_i) < +\infty$.

Then for any admissible family A we define the corresponding minmax value (or width) for E_{ε}

$$\beta(\varepsilon) := \inf_{A \in \mathcal{A}} \sup_{x \in A} E_{\varepsilon}(x)$$

there exists a sequence $\varepsilon_k \to 0$ and $x_k \in \mathfrak{M}$ such that

$$E_{\varepsilon_k}(x_k) - \beta(\varepsilon_k) \to 0, \quad \|dE_{\varepsilon_k}(x_k)\|_{T^*\mathfrak{M}} < f(\varepsilon_k) , \quad \varepsilon_k \log \varepsilon_k^{-1} \left. \frac{dE_{\varepsilon}(x_k)}{d\varepsilon} \right|_{\varepsilon = \varepsilon_k} \to 0$$

where f is any function fixed in advance.

II.3 Local Geometry of Legendrian Immersions

II.3.1 Infinitesimal variations of Legendrian Immersions

Let $\vec{\Lambda} = (\vec{\Phi}, \vec{n})$ be a C^2 immersion of a closed oriented surface Σ into $V_2(\mathbb{R}^4)$. We shall restrict to Legendrian immersions which are the immersions satisfying

$$\forall x \in \Sigma \qquad \vec{\Lambda}_* T_x \Sigma \subset H_{\vec{\Lambda}(x)}$$

Let Ψ_t be the flow of a vector-field X in $V_2(\mathbb{R}^4)$ such that Ψ_t is preserving the kernel of α for every t (i.e. is a contacto-morphism). We have then for all t

$$\operatorname{Ker}(\Psi_t^*\alpha) = \operatorname{Ker}(\alpha)$$

Two non zero one forms have the same kernel if and only if they are parallel to each other. Hence, since $|\alpha| = \sqrt{2}$ at every point one has

$$\Psi_t^* \alpha = f(t) \alpha$$
 where $2 f(t) := < \alpha, \Psi_t^* \alpha > .$

Hence

$$\frac{d\Psi_t^*\alpha}{dt} = \mathcal{L}_X \alpha = \dot{f} \,\alpha$$

Thus

$$d(\alpha(X)) + d\alpha \, \mathbf{L} \, X = \dot{f} \, \alpha$$

We decompose $\vec{X} = \vec{X}^V + \vec{X}^H$ where \vec{X}^H is the orthogonal projection of \vec{X} onto H. This gives in particular

$$d(\alpha(\vec{X}^V)) + (\Pi^* \omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}}) \, \mathbf{\bot} \, \vec{X}^H = f \, \alpha$$

Denote $h := \alpha(\vec{X}^V)$ that is

$$\vec{X}^V = \frac{h}{2} \vec{R} \; ,$$

and let $\vec{Y} = \vec{Y}^V + \vec{Y}^H$ we have

$$d_{\vec{Y}}h + \vec{Y}^H \cdot J_H \vec{X}^H = \dot{f} \ \alpha(\vec{Y}^V) \ .$$

This gives

$$\vec{X}^H = J_H \nabla^H h$$

where $\nabla^{H}h$ is the horizontal projection of the gradient of h. To conclude, infinitesimal variations preserving the Legendrian constraint are given by vector fields of the form

$$\vec{X}_h := J_H \nabla^H h + \frac{h}{2} \vec{R} ,$$

where h is an arbitrary function (i.e. Hamiltonian).

The projection of a Legendrian immersion is satisfying $\vec{\Lambda}^* \alpha = 0$ hence

$$0 = d\vec{\Lambda}^* \alpha = \vec{\Lambda}^* \Pi^* \omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}} = (\Pi \circ \vec{\Lambda})^* \omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}}$$

hence $\vec{G} := \Pi \circ \vec{\Lambda}$ is Lagrangian into $\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}$. If $\vec{e_1}, \vec{e_2}$ is an orthonormal basis of $\vec{G}_* T\Sigma$ we have since the restriction of $\omega_{\mathbb{C}P^1 \times \overline{\mathbb{C}P^1}}$ to $\vec{G}_* T\Sigma = 0$ we have

$$\vec{e}_2 \cdot J\vec{e}_1 = -\vec{e}_1 \cdot J\vec{e}_2 = \omega(\vec{e}_1, \vec{e}_2) = 0$$

hence $J\vec{e}_1, J, \vec{e}_2$ realises an orthonormal basis of the normal bundle to the immersion \vec{G} . By uniqueness of the horizontal lifts and the fact that Π_* realises an isometry between $H_{(\vec{a},\vec{b})}$ and $T_{\Pi(\vec{a},\vec{b})} \mathbb{C}P^1 \times \mathbb{C}P^1$ the same holds for the horizontal lifts as well and $(J_H \vec{e}_1^H, J_H \vec{e}_2^H)$ realises an orthonormal basis of the normal plane to $\vec{\Lambda}_* T\Sigma$ in H.

Let $\vec{c} \in \text{Span}\{\vec{a}, \vec{b}\}^{\perp}$. We have

$$\Pi_*(\vec{c},0) = \frac{d}{dt} \left[\Pi \circ (\vec{a}+t\,\vec{c},\vec{b}) \right] = \frac{d}{dt} \left(((\vec{a}+t\vec{c})\wedge\vec{b})^+, ((\vec{a}+t\vec{c})\wedge\vec{b})^- \right) = \left((\vec{c}\wedge\vec{b})^+, (\vec{c}\wedge\vec{b})^- \right) \;.$$

Hence

$$\Pi_* J_H(\vec{c}, 0) = \left(J_{\vec{G}^+}((\vec{c} \wedge \vec{b})^+), J_{\vec{G}^-}((\vec{c} \wedge \vec{b})^-) \right) = \left((\vec{d} \wedge \vec{b})^+, -(\vec{d} \wedge \vec{b})^- \right) = \left((\vec{a} \wedge \vec{c})^+, (\vec{a} \wedge \vec{c})^- \right) = \Pi_*(0, \vec{c})$$

where

$$\vec{d} := \star \left[\vec{a} \wedge \vec{b} \wedge \vec{c}
ight] \; .$$

Since $J_{H}^{2} = -I_{H}$ $J_{H}(0, \vec{c}) = -(\vec{c}, 0)$

II.3.2 The second fundamental Form of a Legendrian immersion

We assume that $\vec{\Lambda} = (\vec{\Phi}, \vec{n})$ is conformal Legendrian from the unit disc D^2 into $V_2(\mathbb{R}^4)$. Hence the unit Gauss Map is a 3-vector given by

$$\vec{N} := \frac{e^{-2\lambda}}{\sqrt{2}} \vec{R}(\vec{\Lambda}) \wedge J_H \frac{\partial \vec{\Lambda}}{\partial x_1} \wedge J_H \frac{\partial \vec{\Lambda}}{\partial x_2} ,$$

where

$$e^{2\lambda} = \left| \frac{\partial \vec{\Lambda}}{\partial x_1} \right|^2 = \left| \frac{\partial \vec{\Lambda}}{\partial x_2} \right|^2$$

We see $V_2(\mathbb{R}^4)$ canonically isometrically embedded in $S^3 \times S^3 \hookrightarrow \mathbb{R}^4 \times \mathbb{R}^4$. The second fundamental form of the immersion $\vec{\Lambda}$ is given by

$$ec{\mathbb{I}}_{ij} \, dx_i \otimes dx_j := \pi_{ec{N}} \partial^2_{x_i x_j} ec{\Lambda} \, dx_i \otimes dx_j$$

where $\pi_{\vec{N}}$ is the orthogonal projection onto the normal space to $\vec{\Lambda}_* T\Sigma$ in $TV_2(\mathbb{R}^4)$. Assuming as just above that $\vec{\Lambda}$ is in local conformal coordinates, this normal space is generated by the orthonormal basis

$$\sqrt{2}^{-1} \vec{R}(\vec{\Lambda}) , e^{-\lambda} J_H \partial_{x_1} \vec{\Lambda} , e^{-\lambda} J_H \partial_{x_2} \vec{\Lambda} .$$

Hence we have in these conformal coordinates

$$\vec{\mathbb{I}}_{ij} = e^{-2\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \partial^2_{x_i x_j} \vec{\Lambda} J_H \partial_{x_1} \vec{\Lambda} + e^{-2\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \partial^2_{x_i x_j} \vec{\Lambda} J_H \partial_{x_2} \vec{\Lambda} + \frac{1}{2} \vec{R} \cdot \partial^2_{x_i x_j} \vec{\Lambda} \vec{R}$$

Observe that the Legendrian condition is implying

$$\partial_{x_k}\partial_{x_l}\vec{\Lambda}\cdot\vec{R} = \partial_{x_k}\partial_{x_l}\vec{\Phi}\cdot\vec{n} - \partial_{x_k}\partial_{x_l}\vec{n}\cdot\vec{\Phi} = \partial_{x_k}\left[\partial_{x_l}\vec{\Phi}\cdot\vec{n} - \partial_{x_l}\vec{n}\cdot\vec{\Phi}\right] = 0.$$
(II.5)

Hence

$$\vec{\mathbb{I}}_{ij} = e^{-2\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_i x_j}^2 \vec{\Lambda} J_H \partial_{x_1} \vec{\Lambda} + e^{-2\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_i x_j}^2 \vec{\Lambda} J_H \partial_{x_2} \vec{\Lambda}$$

In particular

$$|\vec{\mathbb{I}}_{ij}|^2 = \left| \pi_T \left[J_H \nabla_{x_i} \partial_{x_j} \vec{\Lambda} \right] \right|^2 = \left| \pi_{\vec{N}} \partial^2_{x_i x_j} \vec{\Lambda} \right|^2 ,$$

where π_T is the orthogonal projection onto $\vec{\Lambda}_* T\Sigma$, ∇ is the covariant derivative on $V_2(\mathbb{R}^4)$ and we denote $\nabla_{x_k} := \nabla_{\partial_{x_k} \vec{\Lambda}}$. We have in particular

$$\begin{split} |\vec{\mathbb{I}}|_{g}^{2} &= e^{-4\lambda} \sum_{i,j=1}^{2} \left| \pi_{\vec{N}} \partial_{x_{i}x_{j}}^{2} \vec{\Lambda} \right|^{2} .\\ \nabla_{x_{i}} \vec{N} &= \left| \frac{1}{\sqrt{2}} \nabla_{x_{i}} [\vec{R}(\vec{\Lambda})] \wedge J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{1}} \wedge J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{2}} \right.\\ &\left. + \frac{1}{\sqrt{2}} \vec{R}(\vec{\Lambda}) \wedge \nabla_{x_{i}} \left[J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{1}} \right] \wedge J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{2}} \right.\\ &\left. + \frac{1}{\sqrt{2}} \vec{R}(\vec{\Lambda}) \wedge J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{1}} \wedge \nabla_{x_{i}} \left[J_{H} e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_{2}} \right] \right. \end{split}$$

We have first for i, k = 1, 2

$$\nabla_{x_i}[\vec{R}(\vec{\Lambda})] \cdot \vec{R}(\vec{\Lambda}) = 0 \quad , \quad \nabla_{x_i}[\vec{R}(\vec{\Lambda})] \cdot \partial_{x_k}\vec{\Lambda} = \partial_{x_k}\vec{\Phi} \cdot \partial_{x_i}\vec{n} - \partial_{x_i}\vec{\Phi} \cdot \partial_{x_k}\vec{n} = 0 \; ,$$

hence

$$\nabla_{x_i}[\vec{R}(\vec{\Lambda})] \wedge J_H \, e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_1} \wedge \ J_H \, e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_2} = 0 \; .$$

We have also for i, k = 1, 2

$$\nabla_{x_i} \left[J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_k} \right] \cdot \partial_{x_l} \vec{\Lambda} = -J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_k} \cdot \partial_{x_l x_i}^2 \vec{\Lambda}$$

Thus

$$\nabla_{x_i}\vec{N} = -\sum_{j=1}^2 \frac{e^{-2\lambda}}{\sqrt{2}} J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_1} \cdot \partial_{x_j x_i}^2 \vec{\Lambda} \ \vec{R}(\vec{\Lambda}) \wedge \partial_{x_l}\vec{\Lambda} \wedge \ J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_2}$$
$$-\sum_{j=1}^2 \frac{e^{-2\lambda}}{\sqrt{2}} J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_2} \cdot \partial_{x_j x_i}^2 \vec{\Lambda} \ \vec{R}(\vec{\Lambda}) \wedge \ J_H e^{-\lambda} \frac{\partial \vec{\Lambda}}{\partial x_1} \wedge \partial_{x_l}\vec{\Lambda}$$

This is implying in particular

$$\sum_{i=1}^{2} |\nabla_{x_i} \vec{N}|^2 = e^{-2\lambda} \sum_{i,j=1}^{2} |\pi_{\vec{N}} \partial_{x_i x_j}^2 \vec{\Lambda}|^2 .$$

We recall that the mean curvature vector is given by

$$\vec{H}_{\vec{\Lambda}} := \frac{1}{2} \operatorname{Tr}_{g} \mathbb{I} = \frac{e^{-2\lambda}}{2} \pi_{\vec{N}} \left[\partial_{x_{1}^{2}}^{2} \vec{\Lambda} + \partial_{x_{2}^{2}}^{2} \vec{\Lambda} \right] \; .$$

Hence

$$2\vec{H}_{\vec{\Lambda}} = e^{-4\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \Delta \vec{\Lambda} J_H \partial_{x_1} \vec{\Lambda} + e^{-4\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \Delta \vec{\Lambda} J_H \partial_{x_2} \vec{\Lambda}$$

Let

$$\gamma := e^{-2\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \, dx_1 + e^{-2\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \, dx_2 \; .$$

We have

$$d\gamma = -\partial_{x_2} \left[e^{-2\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \right] + \partial_{x_1} \left[e^{-2\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \right] dx_1 \wedge dx_2$$
$$= \partial_{x_2} \left[e^{-2\lambda} \nabla (J_H \partial_{x_1} \vec{\Lambda}) \cdot \nabla \vec{\Lambda} \right] - \partial_{x_1} \left[e^{-2\lambda} \nabla (J_H \partial_{x_2} \vec{\Lambda}) \cdot \nabla \vec{\Lambda} \right] dx_1 \wedge dx_2$$

Observe that

$$\nabla (J_H \partial_{x_i} \vec{\Lambda}) \cdot \nabla \vec{\Lambda} = - \star \nabla \left(\vec{\Phi} \wedge \vec{n} \wedge \partial_{x_i} \vec{\Phi} \right) \wedge \nabla \vec{\Phi} - \star \nabla \left(\vec{\Phi} \wedge \vec{n} \wedge \partial_{x_i} \vec{n} \right) \wedge \nabla \vec{n}$$
$$- \star \left(\vec{\Phi} \wedge \vec{n} \wedge \partial_{x_i} \nabla \vec{\Phi} \right) \wedge \nabla \vec{\Phi} - \star \left(\vec{\Phi} \wedge \vec{n} \wedge \partial_{x_i} \nabla \vec{n} \right) \wedge \nabla \vec{n}$$
$$= J_H \nabla^H \partial_{x_i} \vec{\Lambda} \cdot \nabla \vec{\Lambda} = -\nabla \partial_{x_i} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda}$$

Hence

$$d\gamma = -\partial_{x_2} \left[e^{-2\lambda} \nabla \partial_{x_1} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} \right] + \partial_{x_1} \left[e^{-2\lambda} \nabla \partial_{x_2} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} \right] dx_1 \wedge dx_2$$

$$= \left[2 \,\partial_{x_2} \lambda \, e^{-2\lambda} \nabla \partial_{x_1} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} - 2 \,\partial_{x_1} \lambda \, e^{-2\lambda} \nabla \partial_{x_2} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} \right] dx_1 \wedge dx_2 \tag{II.6}$$

$$- e^{-2\lambda} \left[\nabla \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_2} (J_H \nabla \vec{\Lambda}) - \nabla \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1} (J_H \nabla \vec{\Lambda}) \right] dx_1 \wedge dx_2$$

We have

$$\begin{split} \nabla_{x_k} \partial_{x_l} \vec{\Lambda} &= e^{-2\lambda} \left[\partial_{x_k} \partial_{x_l} \vec{\Lambda} \cdot \partial_{x_k} \vec{\Lambda} \ \partial_{x_k} \vec{\Lambda} + \partial_{x_k} \partial_{x_l} \vec{\Lambda} \cdot \partial_{x_{k+1}} \vec{\Lambda} \ \partial_{x_{k+1}} \vec{\Lambda} \right] \\ &+ e^{-2\lambda} \left[\partial_{x_k} \partial_{x_l} \vec{\Lambda} \cdot J_H(\vec{\Lambda}) \ \partial_{x_k} \vec{\Lambda} \ J_H(\vec{\Lambda}) \ \partial_{x_k} \vec{\Lambda} + \partial_{x_k} \partial_{x_l} \vec{\Lambda} \cdot J_H(\vec{\Lambda}) \partial_{x_{k+1}} \vec{\Lambda} \ J_H(\vec{\Lambda}) \ \partial_{x_{k+1}} \vec{\Lambda} \right] \\ &+ 2^{-1} \ \partial_{x_k} \partial_{x_l} \vec{\Lambda} \cdot \vec{R} \ \vec{R} \end{split}$$

(II.5) gives

$$\left\{ \begin{array}{l} \nabla_{x_k} \partial_{x_k} \vec{\Lambda} = \partial_{x_k} \lambda \ \partial_{x_k} \vec{\Lambda} - \partial_{x_{k+1}} \lambda \ \partial_{x_{k+1}} \vec{\Lambda} + e^{-2\lambda} \left[\partial_{x_k} \partial_{x_k} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} \right] J_H \nabla \vec{\Lambda} \\ \nabla_{x_k} \partial_{x_{k+1}} \vec{\Lambda} = \partial_{x_{k+1}} \lambda \ \partial_{x_k} \vec{\Lambda} + \partial_{x_k} \lambda \ \partial_{x_{k+1}} \vec{\Lambda} + e^{-2\lambda} \left[\partial_{x_k} \partial_{x_{k+1}} \vec{\Lambda} \cdot J_H \nabla \vec{\Lambda} \right] J_H \nabla \vec{\Lambda} \end{array} \right.$$

which implies in particular

$$\nabla \partial_{x_1} \vec{\Lambda} = \nabla \lambda \ \partial_{x_1} \vec{\Lambda} + \nabla^{\perp} \lambda \ \partial_{x_2} \vec{\Lambda} + e^{-2\lambda} \ \nabla \partial_{x_1} \vec{\Lambda} \cdot J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \ J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda}$$
$$+ e^{-2\lambda} \ \nabla \partial_{x_1} \vec{\Lambda} \cdot J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \ J_H \partial_{x_2} \vec{\Lambda}$$

Observe also that

$$J_H \partial_{x_j} \vec{\Lambda} \cdot \partial_{x_k} \left(J_H \partial_{x_l} \vec{\Lambda} \right) = \partial_{x_j} \vec{\Lambda} \cdot \partial_{x_k x_l}^2 \vec{\Lambda}$$

We deduce

$$\nabla \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_2} (J_H \nabla \vec{\Lambda}) - \nabla \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1} (J_H \nabla \vec{\Lambda}) = 2 \nabla \partial_{x_1} \vec{\Lambda} \cdot J_H \nabla \partial_{x_2} \vec{\Lambda}$$

Combining the previous gives then

$$\nabla \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_2} (J_H \nabla \vec{\Lambda}) - \nabla \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1} (J_H \nabla \vec{\Lambda}) = 2 \nabla \lambda \ \partial_{x_1} \vec{\Lambda} \cdot J_H \nabla \partial_{x_2} \vec{\Lambda}$$

+2\sum \lambda \lambda_{x_2} \vec{\Lambda} \cdot J_H \nabla \lambda_{x_2} \vec{\Lambda} + 2 \end{v}^{-\lambda \lambda} \lambda_{x_1} \vec{\Lambda} \cdot \lambda_{x_2} \vec{\Lambda} + 2 \end{v}^{-2\lambda} \nabla \delta_{x_1} \vec{\Lambda} \cdot \delta_{x_2} \vec{\Lambda} + 2 \end{v}^{-2\lambda} \nabla \delta_{x_1} \vec{\Lambda} \delta \delta_{x_2} \vec{\Lambda} + 2 \end{v}^{-2\lambda} \nabla \delta_{x_1} \vec{\Lambda} \delta \delta_{x_2} \vec{\Lambda} \delta \delta \delta_{x_2} \vec{\Lambda} \delta \delta \delta_{x_2} \vec{\Lambda} \delta \delta_{x_2} \vec{\Lambda} \delta \delta \delta_{x_2} \vec{\Lambda} \delta \delta_{x_2} \vec{\Lambda} \delta \delta \delta \delta \delta \delta_{x_2} \vec{\Lambda} \delta \del

Observe that

$$e^{-2\lambda}\partial_{x_2}\vec{\Lambda}\cdot\nabla\partial_{x_2}\vec{\Lambda} = \nabla\lambda$$
 and $e^{-2\lambda}\partial_{x_1}\vec{\Lambda}\cdot\nabla\partial_{x_2}\vec{\Lambda} = -\nabla^{\perp}\lambda$

Hence

$$\nabla \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_2} (J_H(\vec{\Lambda}) \nabla \vec{\Lambda}) - \nabla \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1} (J_H(\vec{\Lambda}) \nabla \vec{\Lambda})$$

$$= 2 \nabla \lambda \left[-J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \cdot \nabla \partial_{x_2} \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \cdot \nabla^{\perp} \partial_{x_2} \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \cdot \nabla^{\perp} \partial_{x_1} \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \cdot \nabla \partial_{x_1} \vec{\Lambda} \right]$$

Observe that

$$J_{H}(\vec{\Lambda})\partial_{x_{2}}\vec{\Lambda}\cdot\nabla\partial_{x_{1}}\vec{\Lambda} = \nabla\left[J_{H}(\vec{\Lambda})\partial_{x_{2}}\vec{\Lambda}\cdot\partial_{x_{1}}\vec{\Lambda}\right] - \nabla(J_{H}(\vec{\Lambda})\partial_{x_{2}}\vec{\Lambda})\cdot\partial_{x_{1}}\vec{\Lambda} = J_{H}(\vec{\Lambda})\partial_{x_{1}}\vec{\Lambda}\cdot\nabla\partial_{x_{2}}\vec{\Lambda} .$$

Thus

$$\nabla \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_2} (J_H(\vec{\Lambda}) \nabla \vec{\Lambda}) - \nabla \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1} (J_H(\vec{\Lambda}) \nabla \vec{\Lambda})$$

$$= 2 \nabla \lambda \left[J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \cdot \nabla^{\perp} \partial_{x_2} \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \cdot \nabla^{\perp} \partial_{x_1} \vec{\Lambda} \right]$$

$$= -2 \partial_{x_1} \lambda \left[J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_2 x_2}^2 \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_1 x_2}^2 \vec{\Lambda} \right] + 2 \partial_{x_2} \lambda \left[J_H(\vec{\Lambda}) \partial_{x_2} \vec{\Lambda} \cdot \partial_{x_1 x_2}^2 \vec{\Lambda} + J_H(\vec{\Lambda}) \partial_{x_1} \vec{\Lambda} \cdot \partial_{x_1 x_1}^2 \vec{\Lambda} \right]$$

$$= -2 \partial_{x_1} \lambda \left[J_H(\vec{\Lambda}) \nabla \vec{\Lambda} \cdot \nabla \partial_{x_2} \vec{\Lambda} \right] + 2 \partial_{x_2} \lambda \left[J_H(\vec{\Lambda}) \nabla \vec{\Lambda} \cdot \nabla \partial_{x_1} \vec{\Lambda} \right] .$$

$$(II 7)$$

Combining (II.6) and (II.7) we obtain that γ is closed and we have established the following lemma which was already known since the work of Dazor [6].

Lemma II.3. Let $\vec{\Lambda}$ be a Legendrian immersion into $V_2(\mathbb{R}^4)$ then the second fundamental form is horizontal and

$$d\left(J_H(\vec{\Lambda})\vec{H}_{\vec{\Lambda}}\right)^{\sharp} = 0 , \qquad (II.8)$$

where for any vector $\vec{X} \in H$ we assign the one form \vec{X}^{\sharp} on H such that

$$\forall \vec{Y} \in H \qquad \vec{X}^{\sharp}(\vec{Y}) := \vec{X} \cdot \vec{Y} \; .$$

The co-homology class in Σ defined by $\left(J_H(\vec{\Lambda})\vec{H}_{\vec{\Lambda}}\right)^{\sharp}$ is invariant under the action of Hamiltonian isotopies (this is one of the main results in [13]).

Let $d\beta = 2^{-1} \gamma$ (β is locally defined). We have

$$\begin{cases} \partial_{x_1}\beta = 2^{-1} e^{-2\lambda} J_H \partial_{x_1} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \\ \partial_{x_2}\beta = 2^{-1} e^{-2\lambda} J_H \partial_{x_2} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \end{cases}$$

This gives

$$J_H\left(\partial_{x_1}\beta\,\partial_{x_1}\vec{\Lambda} + \partial_{x_2}\beta\,\partial_{x_2}\vec{\Lambda}\right) = 2^{-1}\,\pi_H\left[\Delta\vec{\Lambda}\right] \;.$$

We have

$$\nabla^{\Sigma}\beta = \langle d\beta \cdot d\vec{\Lambda} \rangle_g = e^{-2\lambda} \left(\partial_{x_1}\beta \, \partial_{x_1}\vec{\Lambda} + \partial_{x_2}\beta \, \partial_{x_2}\vec{\Lambda} \right) = -2^{-1} J_H \, \Delta_g \vec{\Lambda}$$

Hence

$$\int_{\Sigma} \langle d\vec{w}, d\vec{\Lambda} \rangle_g \ dvol_g = -\int_{\Sigma} \vec{w} \cdot \Delta_g \vec{\Lambda} \ dvol_g = 2 \int_{\Sigma} \vec{w} \cdot J_H \nabla^{\Sigma} \beta \ dvol_g$$

Assume \vec{w} is of the form $\vec{w} := J^H \nabla^H h + 2^{-1} h \vec{R}$ this then gives

$$\int_{\Sigma} \langle d\vec{w}, d\vec{\Lambda} \rangle_g \ dvol_g = 2 \int_{\Sigma} \langle dh, d\beta \rangle_g \ dvol_g \ . \tag{II.9}$$

II.3.3 The variations of the Gauss unit multi-vector of a Legendrian immersion

We consider a smooth perturbation of a conformal Legendrian immersion and we denote by $\vec{w} := d\vec{\Lambda}/dt(0)$. We see $\vec{\Lambda}_t$ as an immersion into $V_2(\mathbb{R}^4) \hookrightarrow S^3 \times S^3 \hookrightarrow \mathbb{R}^4 \times \mathbb{R}^4$. Hence $\vec{N} \in \wedge^3 \mathbb{R}^8$. We have

$$\frac{d\vec{R}(\vec{\Lambda}_t)}{dt} = \frac{d}{dt} \begin{pmatrix} \vec{n}_t \\ -\vec{\Phi}_t \end{pmatrix} = I \vec{w}$$

where

$$I := \left(\begin{array}{cc} 0 & 1\\ & \\ -1 & 0 \end{array}\right)$$

Let

$$\vec{T} := \frac{\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}}{|\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}|} \; .$$

The link between \vec{T} and \vec{N} is made as follows. We have that

$$\vec{\Lambda} \cdot J_H \partial_{x_k} \vec{\Lambda} = \begin{pmatrix} \vec{\Phi} \\ \vec{n} \end{pmatrix} \cdot J_H \partial_{x_k} \vec{\Lambda} = \star \vec{\Phi} \wedge \vec{n} \wedge \partial_{x_k} \vec{\Phi} \wedge \vec{\Phi} + \star \vec{\Phi} \wedge \vec{n} \wedge \partial_{x_k} \vec{n} \wedge \vec{n} = 0$$

we have similarly

$$\begin{pmatrix} \vec{\Phi} \\ -\vec{n} \end{pmatrix} \cdot J_H \partial_{x_k} \vec{\Lambda} = 0 \quad \begin{pmatrix} \vec{\Phi} \\ -\vec{n} \end{pmatrix} \cdot \vec{R}(\vec{\Lambda}) = 0 \quad \cdots$$

Hence

$$\sqrt{2}^{-1}\vec{\Lambda} = \sqrt{2}^{-1} \begin{pmatrix} \vec{\Phi} \\ \vec{n} \end{pmatrix}, \ \sqrt{2}^{-1}\vec{U} := \sqrt{2}^{-1} \begin{pmatrix} \vec{\Phi} \\ -\vec{n} \end{pmatrix}, \ \sqrt{2}^{-1}\vec{V} := \sqrt{2}^{-1} \begin{pmatrix} \vec{n} \\ \vec{\Phi} \end{pmatrix}, \ \sqrt{2}^{-1}\vec{R} := \sqrt{2}^{-1} \begin{pmatrix} \vec{n} \\ -\vec{\Phi} \end{pmatrix}$$

together with

$$e^{-\lambda}\partial_{x_1}\vec{\Lambda}$$
, $e^{-\lambda}\partial_{x_2}\vec{\Lambda}$, $e^{-\lambda}J_H\partial_{x_1}\vec{\Lambda}$ and $e^{-\lambda}J_H\partial_{x_1}\vec{\Lambda}$

realises an orthonormal basis of \mathbb{R}^8 . We deduce that

$$\vec{N} := \pm \sqrt{2}^{-3} \, \star \left[\vec{\Lambda} \wedge \vec{U} \wedge \vec{V} \wedge \vec{T} \right] \; .$$

In particular

$$|d\vec{N}|_g^2 \leq 2^{-3} \left[|d\vec{T}|_g^2 + 2 \right] \, .$$

We have

$$\begin{aligned} \frac{d}{dt} |\partial_{x_1} \vec{\Lambda}_t \wedge \partial_{x_2} \vec{\Lambda}_t|^2 &= +2 \left[\partial_{x_1} \vec{w} \wedge \partial_{x_2} \vec{\Lambda} \right] \cdot \left[\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda} \right] + 2 \left[\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{w} \right] \cdot \left[\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda} \right] \\ &= 2 e^{2\lambda} \left[\partial_{x_1} \vec{w} \cdot \partial_{x_1} \vec{\Lambda} + \partial_{x_2} \vec{w} \cdot \partial_{x_2} \vec{\Lambda} \right] = 2 e^{2\lambda} \nabla \vec{w} \cdot \nabla \vec{\Lambda} \end{aligned}$$

This is implying

$$\begin{split} \frac{d\vec{T}}{dt} &= e^{-2\lambda} \left[\partial_{x_1} \vec{w} \wedge \partial_{x_2} \vec{\Lambda} + \partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{w} \right] - \frac{1}{2} \left| \partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda} \right|^{-3} \left. \frac{d}{dt} |\partial_{x_1} \vec{\Lambda}_t \wedge \partial_{x_2} \vec{\Lambda}_t|^2 \left[\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda} \right] \\ &= e^{-2\lambda} \left[\partial_{x_1} \vec{w} \wedge \partial_{x_2} \vec{\Lambda} + \partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{w} \right] - e^{-2\lambda} \nabla \vec{w} \cdot \nabla \vec{\Lambda} \vec{T} \end{split}$$

Now we compute the derivative with respect to t of the metric of $\vec{\Lambda}_t$.

$$\frac{dg_{ij}}{dt} = \partial_{x_i} \vec{w} \cdot \partial_{x_j} \vec{\Lambda} + \partial_{x_i} \vec{\Lambda} \cdot \partial_{x_j} \vec{w} , \qquad (\text{II.10})$$

which gives

$$\frac{dg^{ij}}{dt} = -e^{-4\lambda} \left[\partial_{x_i} \vec{w} \cdot \partial_{x_j} \vec{\Lambda} + \partial_{x_i} \vec{\Lambda} \cdot \partial_{x_j} \vec{w} \right] \; .$$

Hence

$$\begin{split} &\frac{d}{dt} \left[|d\vec{T}|_{g}^{2} \right] = -e^{-4\lambda} \left[\partial_{x_{i}} \vec{w} \cdot \partial_{x_{j}} \vec{\Lambda} + \partial_{x_{i}} \vec{\Lambda} \cdot \partial_{x_{j}} \vec{w} \right] \ \partial_{x_{i}} \vec{T} \, \partial_{x_{j}} \vec{T} + 2 \, e^{-2\lambda} \, \nabla \vec{T} \cdot \nabla \frac{d\vec{T}}{dt} \\ &= -2 \, \left\langle d\vec{T} \, \dot{\otimes} \, d\vec{T}, d\vec{w} \, \dot{\otimes} \, d\vec{\Lambda} \right\rangle_{g} + 2 \, e^{-2\lambda} \, \nabla \vec{T} \cdot \nabla \left\langle d\vec{w} \wedge d\vec{\Lambda} \right\rangle_{g} - 2 \, e^{-2\lambda} \, \nabla \vec{T} \cdot \nabla \left[< d\vec{w} \cdot d\vec{\Lambda} >_{g} \vec{T} \right] \\ &= -2 \, \left\langle d\vec{T} \, \dot{\otimes} \, d\vec{T}, d\vec{w} \, \dot{\otimes} \, d\vec{\Lambda} \right\rangle_{g} + 2 \, d\vec{T} \cdot d \left\langle d\vec{w} \wedge d\vec{\Lambda} \right\rangle_{g} - 2 \, |d\vec{T}|_{g}^{2} \, < d\vec{w} \cdot d\vec{\Lambda} >_{g} \, . \end{split}$$

We deduce from (II.10) that

$$\frac{d}{dt} \left[dvol_g \right] = \frac{d}{dt} \left[\sqrt{g_{11} g_{22} - g_{12}^2} \right] dx_1 \wedge dx_2 = \langle d\vec{w} \cdot d\vec{\Lambda} \rangle_g \ dvol_g \ .$$

Hence combining the previous gives

$$\begin{aligned} \frac{d}{dt} \int_{\Sigma} (1+|d\vec{T}|_{g}^{2})^{2} \, dvol_{g} &= 4 \int_{\Sigma} (1+|d\vec{T}|_{g}^{2}) \left[-\left\langle d\vec{T} \, \dot{\otimes} \, d\vec{T}, d\vec{w} \, \dot{\otimes} \, d\vec{\Lambda} \right\rangle_{g} + \, d\vec{T} \cdot d\left\langle d\vec{w} \wedge d\vec{\Lambda} \right\rangle_{g} \right] \, dvol_{g} \\ &+ \int_{\Sigma} \left[(1+|d\vec{T}|_{g}^{2})^{2} - 4 \, |d\vec{T}|_{g}^{2} \, (1+|d\vec{T}|_{g}^{2}) \right] < d\vec{w} \cdot d\vec{\Lambda} >_{g} \, dvol_{g} \end{aligned}$$

II.3.4 Almost critical point of E_{ε} with entropy estimate.

We are considering Legendrian immersions of a closed oriented surface Σ . We denote simply by g the metric induced by the immersion. We shall be using the following definition

Definition II.2. We call "admissible sequence of almost critical point" of

$$E_{\varepsilon}(\vec{\Lambda}) := \int_{\Sigma} dvol_g + \varepsilon^4 \int_{\Sigma} (1 + |d\vec{T}|_g^2)^2 \, dvol_g$$

any sequence of elements $\vec{\Lambda}_k \in \mathfrak{E}^{2,4}_{\Sigma,Leg}$ and parameter $\varepsilon_k \to 0$ satisfying respectively

$$\limsup_{k \to +\infty} E_{\varepsilon_k}(\vec{\Lambda}_k) < +\infty , \qquad (\text{II.11})$$

moreover

$$\|dE_{\varepsilon_k}(\vec{\Lambda}_k)\|_{\vec{\Lambda}_k} \le e^{-\varepsilon_k^{-2}} , \qquad (\text{II.12})$$

and satisfying the entropy condition

$$\varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2)^2 \, dvol_{g_k} = o\left(\frac{1}{\log \varepsilon_k^{-1}}\right) \,. \tag{II.13}$$

Hence in particular for any subdomain Ω of Σ and any smooth function h in $V_2(\mathbb{R}^4)$ such that $\operatorname{supp}(h) \subset V_2(\mathbb{R}^4) \setminus \vec{\Lambda}_k(\partial \Omega)$ we have

$$e^{-\varepsilon_{k}^{-2}}O(\|w\|_{\vec{\Lambda}_{k}}) = \int_{\Omega} \langle d\vec{w} \cdot d\vec{\Lambda}_{k} \rangle_{g_{k}} dvol_{g_{k}}$$

$$+4\varepsilon_{k}^{4}\int_{\Omega}(1+|d\vec{T}_{k}|_{g_{k}}^{2})\left[-\left\langle d\vec{T}_{k}\dot{\otimes}d\vec{T}_{k}, d\vec{w}\dot{\otimes}d\vec{\Lambda}_{k}\right\rangle_{g} + d\vec{T}_{k}\cdot d\left\langle d\vec{w} \wedge d\vec{\Lambda}_{k}\right\rangle_{g_{k}}\right] dvol_{g_{k}} \qquad (\text{II.14})$$

$$+\varepsilon_{k}^{4}\int_{\Omega}\left[(1+|d\vec{T}_{k}|_{g_{k}}^{2})^{2} - 4|d\vec{T}_{k}|_{g_{k}}^{2}(1+|d\vec{T}_{k}|_{g_{k}}^{2})\right] \langle d\vec{w} \cdot d\vec{\Lambda}_{k} \rangle_{g_{k}} dvol_{g_{k}}$$

where \vec{T}_k and g_k are denoting respectively the tangent 2-vector to $\vec{\Lambda}_k$ and the induced metric by $\vec{\Lambda}_k$ moreover

$$\vec{w}_h := \vec{X}_h(\vec{\Lambda}_k) = \left[J_H \nabla^H h + \frac{h}{2} \vec{R} \right] \circ \vec{\Lambda}_k .$$

Observe that since $|d\vec{\Lambda}_k|_{g_k}^2 = 2$ at every point we have - interpreting simply \vec{N}_k as a section of $\vec{\Lambda}^{-1}T\mathbb{R}^8$ -

$$|d\vec{w}|_{g} \leq |\nabla\vec{X}_{h}| \quad \text{and} \quad |\nabla^{g}d\vec{w}|_{g} \leq C |\nabla^{2}\vec{X}_{h}| + C |\nabla\vec{X}_{h}| |d\vec{N}|_{g} \leq C \left[|\nabla^{2}\vec{X}_{h}| + |\nabla\vec{X}_{h}| [1 + |d\vec{T}|_{g}] \right]$$
(II.15)

Hence the entropy condition is implying

$$\lim_{k \to +\infty} 4 \varepsilon_k^4 \int_{\Omega} (1 + |d\vec{T}_k|_{g_k}^2) \left[- \left\langle d\vec{T}_k \dot{\otimes} d\vec{T}_k, d\vec{w} \dot{\otimes} d\vec{\Lambda}_k \right\rangle_g + d\vec{T}_k \cdot d \left\langle d\vec{w} \wedge d\vec{\Lambda}_k \right\rangle_g \right] dvol_g + \varepsilon_k^4 \int_{\Omega} \left[(1 + |d\vec{T}_k|_{g_k}^2)^2 - 4 |d\vec{T}_k|_{g_k}^2 (1 + |d\vec{T}_k|_{g_k}^2) \right] < d\vec{w} \cdot d\vec{\Lambda}_k >_{g_k} dvol_{g_k} = 0 .$$
(II.16)

Observe that

$$\partial_{x_i} \vec{T} \, \mathbf{L} \, \vec{R} = \partial_{x_i} \left[\frac{\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}}{|\partial_{x_1} \vec{\Lambda} \wedge \partial_{x_2} \vec{\Lambda}|} \right] \mathbf{L} \, \vec{R} = 0 \tag{II.17}$$

since $\partial^2 \vec{\Lambda}$ and $\partial \vec{\Lambda}$ are horizontal. Hence

$$\left| d\vec{T}_k \cdot d\left\langle d\vec{w} \wedge d\vec{\Lambda}_k \right\rangle_g \right| \le \left| d\vec{T}_k \right|_g^2 |\nabla^H \vec{X}_h| + \left| d\vec{T}_k \right|_g |\nabla^H \nabla \vec{X}_h| \tag{II.18}$$

where $\nabla^H \vec{Y} = \nabla \vec{Y} - 2^{-1} \nabla \vec{Y} \cdot \vec{R} \ \vec{R}$. We have in particular $\nabla^H \vec{R} = \nabla \vec{R}$ and then

$$\nabla^H \nabla \vec{X} = \nabla^H \nabla^H \vec{X} + 2^{-1} \nabla \vec{X} \cdot \vec{R} \ \nabla \vec{R}$$

We have at the point $\vec{p} = (\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$

$$\nabla \vec{R}(\vec{p}) = \nabla^{\mathbb{R}^8} \vec{R} - 2^{-1} \nabla^{\mathbb{R}^8} \vec{R} \cdot \vec{p} \, \vec{p} - 2^{-1} \nabla^{\mathbb{R}^8} \vec{R} \cdot \vec{U} \, \vec{U} - 2^{-1} \nabla^{\mathbb{R}^8} \vec{R} \cdot \vec{V} \, \vec{V}$$
$$= \begin{pmatrix} d\vec{b} \\ -d\vec{a} \end{pmatrix} - 2^{-1} \left[d\vec{b} \cdot \vec{a} - d\vec{a} \cdot \vec{b} \right] \begin{pmatrix} \vec{a} \\ \vec{b} \end{pmatrix} - 2^{-1} \left[d\vec{b} \cdot \vec{a} + d\vec{a} \cdot \vec{b} \right] \begin{pmatrix} \vec{a} \\ -\vec{b} \end{pmatrix} .$$

In particular if \vec{Z} is horizontal, we have

$$\nabla_{\vec{Z}}\vec{R} = \begin{pmatrix} d_{\vec{Z}}\vec{b} \\ -d_{\vec{Z}}\vec{a} \end{pmatrix} . \tag{II.19}$$

This permits to refine a bit (II.18) in order to deduce

$$\left| d\vec{T}_k \cdot d\left\langle d\vec{w} \wedge d\vec{\Lambda}_k \right\rangle_g \right| \le \left| d\vec{T}_k \right|_g^2 |\nabla^H \vec{X}_h| + \left| d\vec{T}_k \right|_g \left[|\nabla^H \nabla^H \vec{X}_h| + |\nabla \vec{X}_h \cdot \vec{R}| \right]$$
(II.20)

It remains to estimate $e^{-\varepsilon_k^{-2}}O(\|w\|_{\vec{\Lambda}_k})$. One has

$$\|w\|_{\vec{\Lambda}_k} = \left[\int_{\Sigma} \left[|\nabla^2 \vec{w}|_g^2 + |\nabla \vec{w}|_g^2 + |\vec{w}|^2\right]^2 \, dvol_g\right]^{1/4} + \||\nabla \vec{w}|_g\|_{L^{\infty}(\Sigma)}$$

Using (II.15) we obtain

$$\|w\|_{\vec{\Lambda}_{k}} \leq \left[\int_{\Sigma} \left[|\nabla^{2}\vec{X}_{h}|^{2} \circ \vec{\Lambda}_{k} + |\nabla\vec{X}_{h}|^{2} \circ \vec{\Lambda}_{k} |1 + |d\vec{T}_{k}||^{2} + |\vec{X}_{h}|^{2} \right]^{2} dvol_{g} \right]^{1/4} + \|\nabla X_{h}\|_{\infty} .$$
(II.21)

Combining (II.9), (II.14), (II.16) and (II.21)

Lemma II.4. For any choice of smooth function $h \in C^3(V_2(\mathbb{R}^4))$ and any choice of admissible sequence of almost critical points of E_{ε} there holds

$$\lim_{k \to +\infty} \int_{\Sigma} \langle dh, d\beta_k \rangle_{g_k} \, dvol_{g_k} = 0 \,. \tag{II.22}$$

Let $G_2^{Leg}(V_2(\mathbb{R}^4))$ be the Grassmann bundle over $V_2(\mathbb{R}^4)$ of 2-planes which are Legendrians. Denote by π the projection from $G_2^{Leg}(V_2(\mathbb{R}^4))$ onto $V_2(\mathbb{R}^4)$ associated to this bundle. A point in $G_2^{Leg}(V_2(\mathbb{R}^4))$ is denoted (\mathcal{P}, \vec{p}) where \mathcal{P} is a 2 dimensional subspace of $H_{\vec{p}}$ the horizontal 2-plane at \vec{p} . Considering now an admissible sequence of almost critical points of E_{ε} we shall denote \mathbf{v}_k the varifold given by

$$\forall \ \Xi \in C^0(G_2^{Leg}(V_2(\mathbb{R}^4))) \quad \mathbf{v}_k(\Xi) := \int_{\Sigma} \Xi((\vec{\Lambda}_k)_* T_x \Sigma, \vec{\Lambda}_k(x)) \ dvol_{g_k}$$

Modulo extraction of a subsequence the non negative measure \mathbf{v}_k is converging in Radon measure to a limiting measure \mathbf{v}_{∞} . Given a C^1 vector-field \vec{X} in $V_2(\mathbb{R}^4)$ and a 2 plane \mathcal{P} in a fiber of the tangent bundle $TV_2(\mathbb{R}^4)$ we define the divergence of \vec{X} along \mathcal{P} (see [21] page 42) to be the number

$$\operatorname{div}_{\mathcal{P}} \vec{X} := \sum_{j=1} \vec{e}_j \cdot \nabla_{\vec{e}_j} \vec{X}$$

where $(\vec{e}_j)_{j=1,2}$ is an arbitrary orthonormal basis of \mathcal{P} (one easily verify that the definition is independent of the choice of such a basis). If \vec{X} is C^1 the map

$$(\mathcal{P}, \vec{p}) \longrightarrow \operatorname{div}_{\mathcal{P}} \vec{X}$$

is in particular a continuous function on the Grassmann bundle $G_2(V_2(\mathbb{R}^4))$. Observe that

$$\int_{\Sigma} \langle d\vec{w} \cdot d\vec{\Lambda}_k \rangle_{g_k} \ dvol_{g_k} = \int_{(\mathcal{P}, \vec{p}) \in G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \operatorname{div}_{\mathcal{P}} \vec{X}_h \ d\mathbf{v}_k(\mathcal{P}, \vec{p})$$

Hence we come naturally to the following definition

Definition II.3. A Radon measure \mathbf{v} on the Grassmann bundle $G_2^{Leg}(V_2(\mathbb{R}^4))$ of Legendrian 2-planes in $TV_2(\mathbb{R}^4)$ is called Legendrian stationary or contact stationary if for any C^2 function h on $V_2(\mathbb{R}^4)$ there holds

$$\int_{(\mathcal{P},\vec{p})\in G_2^{Leg}(V_2(\mathbb{R}^4))} div_{\mathcal{P}} \vec{X}_h \ d\mathbf{v}(\mathcal{P},\vec{p}) = 0$$

where

$$\vec{X}_h := J_H \, \nabla^H h + \frac{h}{2} \, \vec{R} \; .$$

The weak limit \mathbf{v}_{∞} of \mathbf{v}_k is clearly legendrian stationary thanks to (II.14), (II.16) and the weak convergence in Radon measures. The goal of the sections below is to describe \mathbf{v}_{∞} in more details.

III The almost monotonicity formula at the limit

In this section $\vec{\Lambda}_k$ denotes an admissible sequence of almost critical points of E_{ε} .

III.1 The "truncated" almost monotonicity formula at the level k

In this subsection we shall omit to write the subscript k. We assume $\vec{\Lambda}$ is passing through the point $\vec{p} := (\vec{\varepsilon_1}, \vec{\varepsilon_2})$ where $\vec{\varepsilon_i}$ is the canonical basis of \mathbb{R}^4 . In the neighbourhood of \vec{p} we introduce the Legendrian coordinate

$$\varphi := \vec{a} \cdot \vec{\varepsilon_2} - \vec{b} \cdot \vec{\varepsilon_1} = a_2 - b_1 ,$$

$$\rho^2 := |(\vec{a}, \vec{b}) - \vec{p}|^2$$

In the neighbourhood of \vec{p} we write $\vec{a} = \vec{\varepsilon_1} + \vec{v}$ and $\vec{b} = \vec{\varepsilon_2} + \vec{w}$. Using $|\vec{a}|^2 = 1$, $|\vec{b}|^2 = 1$ and $\vec{a} \cdot \vec{b} = 0$ gives

$$\begin{cases} 2 v^{1} = -|\vec{v}|^{2} \\ 2 w_{2} = -|\vec{w}|^{2} \\ w^{1} + v^{2} = -v^{3} w^{3} - v^{4} w^{4} - v^{1} w^{1} - v^{2} w^{2} \end{cases}$$
(III.1)

This gives

$$\begin{cases} v^1 = O(\rho^2) \\ w^2 = O(\rho^2) \\ w^1(1 + O((v^2)^2)) + v^2(1 + O((w^1)^2)) = O(\rho^2) \\ v^2 - w^1 = \varphi \end{cases}$$

which finally implies

$$\left\{ \begin{array}{l} v^{1} = O(\rho^{2}) \\ w^{2} = O(\rho^{2}) \\ w^{1} + v^{2} = O(\rho^{2}) \\ v^{2} - w^{1} = \varphi \end{array} \right.$$

We deduce

$$\alpha = \vec{a} \cdot d\vec{b} - \vec{b} \cdot d\vec{a} = -d\varphi + v^1 dw^1 - w^1 dv^1 + v^2 dw^2 - w^2 dv^2 + v^3 dw^3 - w^3 dv^3 + v^4 dw^4 - w^4 dv^4$$
. (III.2)
We introduce the multivalued function β defined on Σ such that

We introduce the multivalued function β defined on Σ such that

$$-d\beta := (J_H(\vec{\Lambda})\vec{H}_{\vec{\Lambda}})^{\sharp}$$
.

With these notations, in conformal coordinates we have

$$J_H \vec{H}_{\vec{\Lambda}} = \frac{e^{-2\lambda}}{2} \sum_{k=1}^2 J_H \nabla_{x_k} \partial_{x_k} \vec{\Lambda} = \langle d\beta, d\vec{\Lambda} \rangle_g = -e^{-2\lambda} \sum_{k=1}^2 \partial_{x_k} \beta \, \partial_{x_k} \vec{\Lambda} \; .$$

We take the scalar product of this identity with $\vec{\Lambda} - \vec{p}$. This gives

$$\sum_{k=1}^{2} [\vec{\Lambda} - \vec{p}] \cdot J_H \, \nabla_{x_k} \partial_{x_k} \vec{\Lambda} = -\sum_{k=1}^{2} \partial_{x_k} \beta \, \partial_{x_k} |\vec{\Lambda} - \vec{p}|^2 \tag{III.3}$$

Observe that

$$\left[\vec{\Lambda}-\vec{p}\right] \cdot J_H \nabla_{x_k} \partial_{x_k} \vec{\Lambda} = \partial_{x_k^2}^2 \vec{\Phi} \cdot (\vec{n}-\vec{\varepsilon}_2) - \partial_{x_k^2}^2 \vec{n} \cdot (\vec{\Phi}-\vec{\varepsilon}_1) = -\partial_{x_k} \left[\partial_{x_k} \vec{\Phi} \cdot \vec{\varepsilon}_2 - \partial_{x_k} \vec{n} \cdot \vec{\varepsilon}_1\right] = -\partial_{x_k^2}^2 \varphi .$$

Hence

$$\Delta \varphi = \sum_{k=1}^{2} \partial_{x_k} \beta \, \partial_{x_k} |\vec{\Lambda} - \vec{p}|^2 \, . \tag{III.4}$$

Multiplying now by $J_H(\vec{\varepsilon}_2, -\vec{\varepsilon}_1)^H$ where $(\vec{\varepsilon}_2, -\vec{\varepsilon}_1)^H$ is the orthogonal projection of $(\vec{\varepsilon}_2, -\vec{\varepsilon}_1)$ onto H

$$\sum_{k=1}^{2} J(\vec{\varepsilon_{2}},-\vec{\varepsilon_{1}}) \cdot \nabla_{x_{k}} \partial_{x_{k}} \vec{\Lambda} = 2 \sum_{k=1}^{2} \partial_{x_{k}} \beta \left(\vec{\varepsilon_{2}},-\vec{\varepsilon_{1}}\right) \cdot \partial_{x_{k}} \vec{\Lambda} = 2 \sum_{k=1}^{2} \partial_{x_{k}} \beta \partial_{x_{k}} \varphi$$

Recall that $\nabla_{x_k} \partial_{x_k} \vec{\Lambda}$ is horizontal and that

$$\begin{split} &\sum_{k=1}^{2} \nabla_{x_{k}} \partial_{x_{k}} \vec{\Lambda} = \Delta \vec{\Lambda} - 2^{-1} \vec{\Lambda} \cdot \Delta \vec{\Lambda} \ \vec{\Lambda} - 2^{-1} \vec{U} \cdot \Delta \vec{\Lambda} \ \vec{U} - 2^{-1} \vec{V} \cdot \Delta \vec{\Lambda} \ \vec{V} \\ &= \Delta \vec{\Lambda} + 2^{-1} \left[|\nabla \vec{\Lambda}|^{2} \vec{\Lambda} + \nabla \vec{U} \cdot \nabla \vec{\Lambda} \ \vec{U} + \nabla \vec{V} \cdot \nabla \vec{\Lambda} \ \vec{V} \right] \end{split}$$

We have

$$2^{-1} J(\vec{\varepsilon}_{2}, -\vec{\varepsilon}_{1}) \cdot \left[|\nabla \vec{\Lambda}|^{2} \vec{\Lambda} + \nabla \vec{U} \cdot \nabla \vec{\Lambda} \vec{U} + \nabla \vec{V} \cdot \nabla \vec{\Lambda} \vec{V} \right]$$

= $2 e^{2\lambda} + (v^{1} + w^{2}) e^{2\lambda} + 2^{-1} \nabla \vec{U} \cdot \nabla \vec{\Lambda} (v^{1} - w^{2}) + 2^{-1} \nabla \vec{V} \cdot \nabla \vec{\Lambda} (w^{1} + v^{2}) = 2 e^{2\lambda} \left(1 + O(\rho^{2}) + O(\varphi) \right) .$

hence

$$\Delta[v^1 + w^2] + 2e^{2\lambda} \left(1 + O(\rho^2) + O(\varphi)\right) = 2\sum_{k=1}^2 \partial_{x_k}\beta \,\partial_{x_k}\varphi$$

We have

$$2(v^{1} + w^{2}) = -\sum_{i=1}^{4} (v^{i})^{2} + (w^{i})^{2} = -|\vec{\Lambda} - \vec{p}|^{2}$$

This gives finally

$$\begin{cases} 1 + O(\rho^2) + O(\varphi) - 4^{-1}\Delta_g \rho^2 = \langle d\beta, d\varphi \rangle_g \\ \Delta_g \varphi = \langle d\beta, d\rho^2 \rangle_g \end{cases}$$
(III.5)

We have

$$\begin{split} |\vec{\Lambda} - \vec{p}|^2 &= 2^{-1} \left| (\vec{\Lambda} - \vec{p}) \cdot \vec{\Lambda} \right|^2 + 2^{-1} \left| (\vec{\Lambda} - \vec{p}) \cdot \vec{R} (\vec{\Lambda}) \right|^2 + 2^{-1} \left| (\vec{\Lambda} - \vec{p}) \cdot \vec{U} (\vec{\Lambda}) \right|^2 + 2^{-1} \left| (\vec{\Lambda} - \vec{p}) \cdot \vec{V} (\vec{\Lambda}) \right|^2 \\ &+ e^{-2\lambda} \left| (\vec{\Lambda} - \vec{p}) \cdot \nabla \vec{\Lambda} \right|^2 + e^{-2\lambda} \left| (\vec{\Lambda} - \vec{p}) \cdot J_H \nabla \vec{\Lambda} \right|^2 \end{split}$$

We have respectively

$$|(\vec{\Lambda} - \vec{p}) \cdot \vec{\Lambda}|^2 = |v^1 + w^2|^2 = O(\rho^4) , \ |(\vec{\Lambda} - \vec{p}) \cdot \vec{R}(\vec{\Lambda})|^2 = |w^1 - v^2|^2 = \varphi^2 ,$$

and

$$|(\vec{\Lambda} - \vec{p}) \cdot \vec{U}(\vec{\Lambda})|^2 = |v^1 - w^2|^2 = O(\rho^4), \ |(\vec{\Lambda} - \vec{p}) \cdot \vec{V}(\vec{\Lambda})|^2 = |w^1 + v^2|^2 = O(\rho^4) + O(\varphi^2) \ .$$

Moreover

$$e^{-2\lambda} \left| (\vec{\Lambda} - \vec{p}) \cdot \nabla \vec{\Lambda} \right|^2 = 4^{-1} \left| d\rho^2 \right|_g^2 , \ e^{-2\lambda} \left| (\vec{\Lambda} - \vec{p}) \cdot J_H \nabla \vec{\Lambda} \right|^2 = |d(v^2 - w^1)|_g^2 = |d\varphi|_g^2$$

Combining the previous is implying

$$1 = |d\rho|_g^2 + \rho^{-2} |d\varphi|_g^2 + O(\rho^2) + \rho^{-2} O(\varphi^2) .$$
 (III.6)

Away from $\rho = 0$ we introduce

$$\sigma := \frac{2\,\varphi}{\rho^2} \; .$$

We now follow the main computations of [18]. First of all we have (see (III.10) in [18])

$$< d\sigma, d\beta >_g = \left\langle [2 \rho^{-2} d\varphi - 2 \varphi \rho^{-4} d\rho^2], d\beta \right\rangle_g$$

= $4\rho^{-2} \left(1 + O(\rho^2) + O(\varphi) + \rho^{-2} O(\varphi^2)\right) + 4^{-1} \rho^{-4} \Delta_g \mathfrak{r}$
= $4 \frac{\sqrt{1 + \sigma^2}}{\mathfrak{r}^2} \left(1 + O(\rho^2) + O(\varphi) + \rho^{-2} O(\varphi^2)\right) - 4^{-1} \frac{1 + \sigma^2}{\mathfrak{r}^4} \Delta_g \mathfrak{r}^4$

where \mathfrak{r} is the *Folland-Korányi gauge* given by $\mathfrak{r} := (\rho^4 + 4 \varphi^2)^{1/4}$. This gives in particular

$$\left\langle \frac{d\sigma}{1+\sigma^2}, d\beta \right\rangle_g = \frac{4}{\mathfrak{r}^2} \frac{1}{\sqrt{1+\sigma^2}} \left(1 + O(\rho^2) + O(\varphi) + \rho^{-2} O(\varphi^2) \right) - 4 \left| d\log \mathfrak{r} \right|_g^2 - \Delta_g \log \mathfrak{r} .$$
(III.7)

Observe that from (III.2)

$$\nabla^H \varphi = \sum_{i=1}^4 v^i \, \nabla^H w^i - w^i \, \nabla^H v^i \; .$$

We have in \mathbb{R}^8 at a point $\vec{Q} = (\vec{\varepsilon_1} + \vec{v}, \vec{\varepsilon_2} + \vec{w}) \in V_2(\mathbb{R}^8)$

$$\begin{aligned} \left| \sum_{i=1}^{4} v^{i} \nabla^{H} w^{i} - w^{i} \nabla^{H} v^{i} \right|^{2} &= \left| \sum_{i=1}^{4} v^{i} \nabla w^{i} - w^{i} \nabla v^{i} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \nabla w^{i} - w^{i} \nabla v^{i} \right) \cdot \vec{R} \right|^{2} \\ &- 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \nabla w^{i} - w^{i} \nabla v^{i} \right) \cdot \vec{U} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \nabla w^{i} - w^{i} \nabla v^{i} \right) \cdot \vec{V} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \nabla w^{i} - w^{i} \nabla v^{i} \right) \cdot \vec{Q} \right|^{2} \\ &= \rho^{2} - 2^{-1} |-v^{1} (1+v^{1}) - (w^{1})^{2} - (v^{2})^{2} - w^{2} (1+w^{2}) - (v^{3})^{2} - (v^{4})^{2} - (w^{4})^{2} |^{2} \end{aligned}$$

$$\begin{aligned} &-2^{-1} |v^2 + w^1 + 2 \sum_{i=1}^4 v^i w^i|^2 \\ &-2^{-1} |v^1 (1+v^1) + (v^2)^2 + (v^3)^2 + (v^4)^2 - (w^1)^2 - w^2 (1+w^2) - (w^3)^2 - (w^4)^2|^2 - 2^{-1} |v^2 - w^1|^2 \\ &= \rho^2 + O(\rho^4) + O(\varphi^2) = \rho^2 + O(\mathfrak{r}^4) \end{aligned}$$

We have also

$$\begin{split} |\rho \, \nabla^{H} \rho|^{2} &= \left| \sum_{i=1}^{4} v^{i} \, \nabla^{H} v^{i} + w^{i} \, \nabla^{H} w^{i} \right|^{2} = \left| \sum_{i=1}^{4} v^{i} \, \nabla v^{i} + w^{i} \, \nabla w^{i} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \, \nabla v^{i} + w^{i} \, \nabla w^{i} \right) \cdot \vec{R} \right|^{2} \\ &- 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \, \nabla v^{i} + w^{i} \, \nabla w^{i} \right) \cdot \vec{U} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \, \nabla v^{i} + w^{i} \, \nabla w^{i} \right) \cdot \vec{V} \right|^{2} - 2^{-1} \left| \left(\sum_{i=1}^{4} v^{i} \, \nabla v^{i} + w^{i} \, \nabla w^{i} \right) \cdot \vec{Q} \right|^{2} \\ &= \rho^{2} - 2^{-1} \left| v^{2} - w^{1} \right|^{2} - 2^{-1} \left| v^{1} - w^{2} + \sum_{i=1}^{4} (v^{i})^{2} - (w^{i})^{2} \right|^{2} - 2^{-1} \left| v^{2} + w^{1} + 2 \sum_{i=1}^{2} v^{i} \, w^{i} \right|^{2} - 2^{-1} \left| v^{1} + w^{2} + \rho^{2} \right|^{2} \\ &= \rho^{2} + O(\varphi^{2}) + O(\rho^{4}) = \rho^{2} + O(\mathfrak{r}^{4}) \; . \end{split}$$

and similarly we prove

$$\nabla^H \varphi \cdot \nabla^H \rho = O(\mathfrak{r}^4) \; .$$

Following again [18] (III.13) we obtain

$$|\nabla^{H}\mathfrak{r}|^{2} = \frac{\rho^{6}}{\mathfrak{r}^{6}}|\nabla^{H}\rho|^{2} + \frac{|\nabla^{H}\varphi^{2}|}{\mathfrak{r}^{6}} + 4\frac{\varphi\rho^{3}}{\mathfrak{r}^{6}}\nabla^{H}\varphi\cdot\nabla^{H}\rho = \frac{\rho^{2}}{\mathfrak{r}^{2}} + O(\mathfrak{r}^{2}) = \frac{1}{\sqrt{1+\sigma^{2}}} + O(\mathfrak{r}^{2}) .$$
(III.8)

We denote by $(\nabla^{\Sigma} \mathfrak{r})^{\perp} := \nabla^{H} \mathfrak{r} - \nabla^{\Sigma} \mathfrak{r}$ where $\nabla^{\Sigma} \mathfrak{r}$ is denoting the gradient of $\mathfrak{r} \circ \vec{\Lambda}$ that is the orthogonal projection of $\nabla^{H} \mathfrak{r}$ onto the tangent plane $\vec{\Lambda}_{*}T\Sigma$. Hence one has in particular

$$|d\log \mathfrak{r}|_g^2 = |\nabla^{\Sigma}\mathfrak{r}|^2 \quad \text{and} \quad |(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^2 = |\nabla^H\mathfrak{r}|^2 - |\nabla^{\Sigma}\mathfrak{r}|^2 = \frac{1}{\sqrt{1+\sigma^2}} + O(\mathfrak{r}^2) - |\nabla^{\Sigma}\mathfrak{r}|^2 \ .$$

Combining this fact with (III.7) and (III.8) and arguing as in [18] we finally obtain away from $\rho = 0$

$$< d \arctan \sigma, d\beta >_g + \Delta_g \log \mathfrak{r} = 4 \frac{|(\nabla^{\Sigma} \mathfrak{r})^{\perp}|^2}{\mathfrak{r}^2} + \frac{4}{\mathfrak{r}^2} \frac{1}{\sqrt{1 + \sigma^2}} \left(O(\mathfrak{r}^2) + \rho^{-2} O(\varphi^2) \right) + O(1)$$

Observe that

$$\frac{1}{\rho^2} \frac{1}{\sqrt{1+\sigma^2}} = \frac{1}{\sqrt{\rho^4 + 4 \varphi^2}} = \frac{1}{\mathfrak{r}^2} \ .$$

Hence

$$\frac{4}{\mathfrak{r}^2} \frac{1}{\sqrt{1+\sigma^2}} \, \left(O(\mathfrak{r})^2 + \rho^{-2} \, O(\varphi^2) \right) = O(1) \ ,$$

and we get away from $\rho = 0$

$$< d \arctan \sigma, d\beta >_g + \Delta_g \log \mathfrak{r} = 4 \frac{|(\nabla^{\Sigma} \mathfrak{r})^{\perp}|^2}{\mathfrak{r}^2} + O(1)$$
 (III.9)

As in [18] again we compute away from $\rho = 0$

$$\begin{aligned} \mathbf{r}^{3} &< d\mathbf{r}, d\beta >_{g} = \rho^{3} < d\rho, d\beta >_{g} + < d\varphi^{2}, d\beta >_{g} \\ &= 2^{-1} \rho^{2} \Delta_{g} \varphi + 2 \varphi \left[1 + O(\mathbf{r}^{2}) - 4^{-1} \Delta_{g} \rho^{2} \right] \\ &= 2\varphi \left[1 + O(\mathbf{r}^{2}) \right] + 2^{-1} \mathrm{div}^{\Sigma} (\rho^{2} \nabla^{\Sigma} \varphi - \varphi \nabla^{\Sigma} \rho^{2}) \\ &= \mathbf{r}^{2} \frac{\sigma}{\sqrt{1 + \sigma^{2}}} + O(\mathbf{r}^{4}) + 4^{-1} \mathrm{div}^{\Sigma} \left(\mathbf{r}^{4} \frac{\nabla^{\Sigma} \sigma}{1 + \sigma^{2}} \right) \end{aligned}$$
(III.10)

Let 0 < r < 1 arbitrary and let χ be a cut-off function such that

$$\chi(t) = \begin{cases} 1 & \text{for } t < 1 \\ 0 & \text{for } t > 2 \end{cases} , \quad \chi' \le 0 \quad \text{on } \mathbb{R}_+ .$$

Let $1 > r > \eta > 0$ and consider the hamiltonian function given on a neighbourhood of \vec{p} in $V_2(\mathbb{R}^4)$ by

$$h_{r,\eta} := [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \arctan \sigma$$

We claim that away from $\mathfrak{r} = 0$ the function $\arctan \sigma$ defines a C^l function for any $l \in \mathbb{N}$. Indeed for $\rho = 0$ and $\varphi > 0$ it extends continuously by $\pi/2$ while for $\rho = 0$ and $\varphi < 0$ it extends continuously by $-\pi/2$. We have moreover

$$\nabla \arctan \sigma = \frac{2}{1+4\frac{\varphi^2}{\rho^4}} \left[\frac{\nabla \varphi}{\rho^2} - 2\frac{\varphi}{\rho^3} \nabla \rho \right] = 2\frac{\rho^2}{\mathfrak{r}^4} \nabla \varphi - \frac{2}{\mathfrak{r}^4} \varphi \nabla \rho^2 .$$
(III.11)

Hence, away from $\mathfrak{r} = 0$, $\nabla \arctan \sigma$ extends continuously by 0 on $\rho = 0$. \mathfrak{r} is obviously smooth away from $\mathfrak{r} = 0$ as well as $\rho^2 \nabla \varphi - \varphi \nabla \rho^2$ which is proving the claim. We multiply (III.9) by $[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)]$ and (III.10) by $\mathfrak{r}^{-3} \arctan \sigma [r^{-1}\chi'(\mathfrak{r}/r) - \eta^{-1}\chi'(\mathfrak{r}/\eta)]$, this gives

$$< dh_{r,\eta}, d\beta >_g + \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)\right] \Delta_g \log \mathfrak{r} = 4 \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)\right] \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}^2} + O(1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)\right]$$
$$+ \left[r^{-1}\chi'(\mathfrak{r}/r) - \eta^{-1}\chi'(\mathfrak{r}/\eta)\right] \mathfrak{r}^{-1} \arctan \sigma \frac{\sigma}{\sqrt{1+\sigma^2}} + O(\mathfrak{r}) \left[r^{-1}\chi'(\mathfrak{r}/r) - \eta^{-1}\chi'(\mathfrak{r}/\eta)\right]$$
$$+ 4^{-1}\mathfrak{r}^{-3} \left[r^{-1}\chi'(\mathfrak{r}/r) - \eta^{-1}\chi'(\mathfrak{r}/\eta)\right] \arctan \sigma \operatorname{div}^{\Sigma} \left(\mathfrak{r}^4 \frac{\nabla^{\Sigma}\sigma}{1+\sigma^2}\right)$$

Integrating over Σ gives

$$\begin{split} &\int_{\Sigma} < dh_{r,\eta}, d\beta >_g \ dvol_g - \int_{\Sigma} \left[\frac{\mathfrak{r}}{r} \,\chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \,\chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \frac{|\nabla^{\Sigma}\mathfrak{r}|^2}{\mathfrak{r}^2} \ dvol_g \\ &+ \int_{\Sigma} \ O(1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] + \left[O(\mathfrak{r}/r) \,\chi'(\mathfrak{r}/r) - O(\mathfrak{r}/\eta) \chi'(\mathfrak{r}/\eta) \right] \ dvol_g \\ &+ 4^{-1} \,\int_{\Sigma} \nabla^{\Sigma} \left[\mathfrak{r}^{-3} \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \ \arctan \sigma \right] \,\mathfrak{r}^4 \frac{\nabla^{\Sigma} \sigma}{1 + \sigma^2} \ dvol_g \\ &- \int_{\Sigma} \left[\frac{\mathfrak{r}}{r} \,\chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \,\chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \,\frac{1}{\mathfrak{r}^2} \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^2}} \ dvol_g = 4 \,\int_{\Sigma} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{\left| (\nabla^{\Sigma} \mathfrak{r})^{\perp} \right|^2}{\mathfrak{r}^2} \ dvol_g \end{split}$$

From which we deduce the "truncated¹⁷" almost monotonicity formula satisfied by $\vec{\Lambda}_k$:

$$\begin{split} &\int_{\Sigma} \langle dh_{r,\eta}, d\beta \rangle_{g} \ dvol_{g} - \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^{2}}} \right] \ dvol_{g} \\ &+ \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{r}\right) + O\left(\frac{\mathfrak{r}}{r}\right) \chi'\left(\frac{\mathfrak{r}}{r}\right) \right] \ dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{r^{2}} \chi''\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} \ dvol_{g} \\ &- \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} \ dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1+\sigma^{2})^{2}} \ dvol_{g} \\ &= 4 \int_{\Sigma} [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} \ dvol_{g} - \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^{2}}} \right] \ dvol_{g} \\ &+ 4 \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1+\sigma^{2})^{2}} \ dvol_{g} - \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{1+\sigma^{2}} \ dvol_{g} \\ &+ \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{\eta^{2}} \chi''\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} \ dvol_{g} + \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{\eta}\right) + O\left(\frac{\mathfrak{r}}{\eta}\right) \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \ dvol_{g} \\ &\qquad (III.12) \end{split}$$

III.2 Passing to the limit $k \to +\infty$ in the "truncated" almost monotonicity formula (III.12).

We have seen (II.22) that

$$\lim_{k \to +\infty} \int_{\Sigma} \langle dh_{r,\eta}, d\beta_k \rangle_{g_k} \ dvol_{g_k} = 0$$

More precisely, from (II.14), (II.13), (II.15), (II.20) and from (II.9) we obtain

$$\begin{aligned} \left| \int_{\Sigma} \langle dh_{r,\eta}, d\beta_k \rangle_{g_k} \ dvol_{g_k} \right| &\leq C \ \varepsilon_k^4 \ \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2)^2 \ |\nabla^H \vec{X}_{h_{r,\eta}}| \circ \vec{\Lambda}_k \ dvol_{g_k} \\ &+ C \ \varepsilon_k^4 \ \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2) \ |d\vec{T}_k|_{g_k} \ \left[|\nabla^H \nabla^H \vec{X}_{h_{r,\eta}}| \circ \vec{\Lambda}_k + |\nabla \vec{X}_{h_{r,\eta}} \cdot \vec{R}| \circ \vec{\Lambda}_k \right] \ dvol_{g_k} \end{aligned}$$

Recall

$$\vec{X}_h = J_H \nabla^H h + \frac{h}{2} \vec{R} \; .$$

We have in particular

$$J_H \nabla^H \left[\chi(\mathfrak{r}/\eta) \arctan \sigma \right] = \frac{J_H \nabla^H \mathfrak{r}}{\eta} \chi'(\mathfrak{r}/\eta) \arctan \sigma + \chi(\mathfrak{r}/\eta) \frac{J_H \nabla^H \sigma}{1 + \sigma^2}$$

We recall that a subsequence has been extracted in such a way that the varifold \mathbf{v}_k defined by $\vec{\Lambda}_k$ is converging towards a limiting varifold \mathbf{v}_{∞} . We also denote by μ_k (resp. μ_{∞}) the so called *weight* of \mathbf{v}_k (resp. \mathbf{v}_{∞}). It is given by

$$\forall A \subset V_2(\mathbb{R}^4)$$
 Borel $\mu_k(A) := \mathbf{v}_k(\pi^{-1}(A))$

¹⁷ "Truncated" is referring to the fact that we are not making η go to zero at this stage and that we have to "wait" for k going to infinity first.

where π denotes the bundle projection of the Grassmann bundle $G_2^{Leg}(V_2(\mathbb{R}^4) \text{ over } V_2(\mathbb{R}^4))$. The Radon measure convergence in $G_2^{Leg}(V_2(\mathbb{R}^4))$ is implying the following passages to the limit for any smooth functions F and G on $V_2(\mathbb{R}^4)$: we have respectively

$$\lim_{k \to +\infty} \int_{\Sigma} F(\vec{\Lambda}_k(x)) \, dvol_{g_k} = \int_{V_2(\mathbb{R}^4)} F(\vec{p}) \, d\mu_{\infty}(\vec{p}) \, ,$$

we have also

$$\lim_{k \to +\infty} \int_{\Sigma} G(\vec{\Lambda}_k) |\nabla^{\Sigma}(F(\vec{\Lambda}_k(x)))|^2 \, dvol_{g_k} = \lim_{k \to +\infty} \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} G(\vec{p}) |\nabla^{\mathcal{P}}F(\vec{p})|^2 \, d\mathbf{v}_k(\mathcal{P},\vec{p})$$
$$= \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} G(\vec{p}) |\nabla^{\mathcal{P}}F(\vec{p})|^2 \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p})$$

where $\nabla^{\mathcal{P}} F$ is denoting the orthogonal projection of the gradient of F on the Legendrian 2-plane \mathcal{P} in $T_{\vec{P}}V_2(\mathbb{R}^4)$. Passing to the limit in (III.12) gives

$$\begin{split} &-\int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma(\vec{p}) \arctan \sigma(\vec{p})}{\sqrt{1 + \sigma^{2}(\vec{p})}} \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+\int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O(1) \left(\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \right) + O\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+\frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}^{2}(\vec{p})}{r^{2}} \chi''\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &-\frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+\frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &= 4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right)\right] \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{1}|^{2}}{\mathfrak{r}(\vec{p})^{2}} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &- \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^{2}}}(\vec{p}) \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ 4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})^{2}} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &- \frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \end{aligned}$$

We first deduce the following lemma

Lemma III.1. Assume

$$\liminf_{\eta \to 0} \frac{1}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p}) < +\infty , \qquad (\text{III.14})$$

then

$$\lim_{\eta \to 0} \int_{\eta < \mathfrak{r} < 1} \frac{|\nabla^{\mathcal{P}} \sigma|^2}{(1 + \sigma^2(\vec{p}))^2} \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) < +\infty \,. \tag{III.15}$$

Proof of lemma III.1. We have from (III.11) away from $\mathfrak{r} = 0$ and for any $(\mathcal{P}, \vec{p}) \in G_2^{Leg}(V_2(\mathbb{R}^4))$

$$\left|\frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^{2}(\vec{p})}\right| = \frac{1}{\mathfrak{r}^{4}(\vec{p})} \left|\rho^{2}(\vec{p})\nabla^{\mathcal{P}}\varphi - \varphi(\vec{p})\nabla^{\mathcal{P}}\rho^{2}\right| \le \frac{|\nabla^{\mathcal{P}}\varphi|}{\mathfrak{r}^{2}(\vec{p})} + \frac{|\nabla^{\mathcal{P}}\rho|}{\mathfrak{r}(\vec{p})} \le \frac{2}{\mathfrak{r}(\vec{p})} .$$
(III.16)

Observe that we have respectively

$$\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \ge 0$$
, (III.17)

moreover

$$-\chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \ge 0 , \qquad (\text{III.18})$$

and

$$\frac{\sigma \arctan \sigma}{\sqrt{1+\sigma^2}}(\vec{p}) \ge 0 . \tag{III.19}$$

Using (III.16) we have respectively

$$4 \left| \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right| \\ \leq \frac{C}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p})$$
(III.20)

moreover

$$\left| \frac{3}{4} \int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^2(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \right| \\
\leq \frac{C}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p}) \tag{III.21}$$

 $\quad \text{and} \quad$

$$\frac{1}{4} \int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}^2(\vec{p})}{\eta^2} \chi''\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\
\leq \frac{C}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p}) \tag{III.22}$$

Combining (III.13), (III.20), (III.21) and (III.22) we finally obtain

$$\begin{split} \limsup_{\eta \to 0} & \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} 4 \left[\chi \left(\frac{\mathfrak{r}(\vec{p})}{r} \right) - \chi \left(\frac{\mathfrak{r}(\vec{p})}{\eta} \right) \right] \frac{|(\nabla^{\mathcal{P}} \mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ & - \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \, \chi' \left(\frac{\mathfrak{r}(\vec{p})}{\eta} \right) \, \frac{1}{\mathfrak{r}^2(\vec{p})} \left[|\nabla^{\mathcal{P}} \mathfrak{r}|^2 + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^2}} (\vec{p}) \right] \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) < +\infty \, . \end{split}$$
(III.23)

We then deduce for any r < 1

$$\lim_{\eta \to 0} \int_{\mathfrak{r} > \eta} \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) < +\infty \,. \tag{III.24}$$

At $\mathfrak{r} = 0$ we have that ∇v_i and ∇w_i for i = 3, 4 are orthogonal to $(\vec{\varepsilon_1}, \vec{\varepsilon_2})$ and hence horizontal. We have moreover thanks to (II.2)

$$J_H \nabla^H v_i = J_H(\vec{\varepsilon_i}, 0) = (0, \vec{\varepsilon_i}) = \nabla^H w_i$$

Then, in a neighbourhood of $(\vec{\varepsilon}_1, \vec{\varepsilon}_2)$ at the point $(\vec{\varepsilon}_1 + \vec{v}, \vec{\varepsilon}_2 + \vec{w})$ we have

$$J_H(\nabla^H \varphi) = J_H\left(\sum_{i=1}^4 v_i \,\nabla^H w_i - w_i \,\nabla^H v_i\right) = -\rho \,\nabla^H \rho + O(\mathfrak{r}^2) \;.$$

This gives

$$\mathfrak{r}^{3} J_{H}(\nabla^{H}\mathfrak{r}) = \rho^{3} J_{H}(\nabla^{H}\rho) + 2 \varphi J_{H}(\nabla^{H}\varphi) = \rho^{2} \nabla^{H}\varphi - 2 \varphi \rho \nabla^{H}\rho + O(\mathfrak{r}^{4})$$

$$= -\rho^{4} \nabla^{H} \left(\frac{\varphi}{\rho^{2}}\right) + O(\mathfrak{r}^{4}) .$$
(III.25)

For every $\mathcal{P} \in G_2^{Leg}(V_2(\mathbb{R}^4))$ we have that $J_H(\mathcal{P})$ is also horizontal, orthogonal to \mathcal{P} and

$$H_{(\vec{a},\vec{b})} = \mathcal{P} \oplus J_H(\mathcal{P})$$

Hence we deduce from (III.25)

$$-\rho^4 \nabla^{\mathcal{P}} \left(\frac{\varphi}{\rho^2}\right) + O(\mathfrak{r}^4) = \mathfrak{r}^3 J_H((\nabla^{\mathcal{P}}\mathfrak{r})^{\perp})$$

This gives

$$\frac{(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}}{\mathfrak{r}} = \frac{1}{2}\frac{\rho^4}{\mathfrak{r}^4}J_H(\nabla^{\mathcal{P}}\sigma) + O(1) = \frac{1}{2}J_H\left(\frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2}\right) + O(1) , \qquad (\text{III.26})$$

from which we deduce

$$\int_{\mathfrak{r}>\eta} \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) = \frac{1}{4} \, \int_{\mathfrak{r}>\eta} \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \left|\frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2(\vec{p})}\right|^2 \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) + O(\mu_{\infty}(\{\mathfrak{r}<2r\}))$$

Combining this last identity with (III.24) we get (III.15) and lemma III.1 is proved.

We deduce the following lemma

Lemma III.2. Assume

$$\limsup_{\eta \to 0} \frac{1}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p}) < +\infty , \qquad (\text{III.27})$$

then for any function $\phi \in C_0^{\infty}(\mathbb{R}_+, \mathbb{R}_+)$ such that $supp(\phi) \subset \mathbb{R}_+ \setminus \{0\}$ and $\int_0^{+\infty} \phi(t) = 1$

$$\lim_{\eta \to 0} \frac{1}{\eta^2} \int_{V_2(\mathbb{R}^4)} \frac{\eta}{\mathfrak{r}(\vec{p})} \phi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{1 + \sigma \arctan\sigma}{\sqrt{1 + \sigma^2}} (\vec{p}) \ d\mu_{\infty} = \theta_0$$
(III.28)

exists, is finite and is independent of ϕ .

Remark III.1. Observe that

$$\frac{d}{ds} \left[\frac{1+s \arctan s}{\sqrt{1+s^2}} \right] = \frac{\arctan s}{(1+s^2)^{3/2}}$$

Hence the function $(1 + s \arctan s)/\sqrt{1 + s^2}$ is decreasing on \mathbb{R}_- and increasing on \mathbb{R}_+ and we deduce

$$\forall s \in \mathbb{R}$$
 $1 \le \frac{1+s \arctan s}{\sqrt{1+s^2}} \le \frac{\pi}{2}$.

Remark III.2. Observe that if \mathbf{v}_{∞} is the varifold associated to a smooth Legendrian immersion Σ we have $\sigma \to 0$ as $\mathfrak{r} \to 0$ (see [18] section III) hence

$$\begin{aligned} \theta_0 &= \lim_{\eta \to 0} \frac{1}{\eta^2} \int_{V_2(\mathbb{R}^4)} \frac{\eta}{\mathfrak{r}(\vec{p})} \phi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \ d\mu_\infty &= \lim_{\eta \to 0} \frac{1}{\eta^2} \int_{V_2(\mathbb{R}^4)} \frac{\eta}{\mathfrak{r}(\vec{p})} \phi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \left|\nabla^\Sigma \mathfrak{r}\right| \ d\mu_\infty \\ &= \lim_{\eta \to 0} \int_0^\infty \phi\left(\frac{s}{\eta}\right) \frac{\mathcal{H}^1(\{\mathfrak{r}=s\})}{s} \ \frac{ds}{\eta} = 2\pi \ Card(\{\mathfrak{r}^{-1}(\{0\})\}) \ \int_0^\infty \phi(t) \ dt = 2\pi \ Card(\{\mathfrak{r}^{-1}(\{0\})\}) \ . \end{aligned}$$

Proof of lemma III.2. Let d > 1 such that $\text{Supp}(\phi) \in [d^{-1}, d]$. We introduce $\chi(t) := \int_t^{+\infty} \phi(s) \, ds$ and we apply (III.13). Because of (III.15) we have respectively

$$\left| \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \,\chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \,\frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} \,d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right|$$
$$\leq C \,\int_{d^{-1}\eta < \mathfrak{r} < d\eta} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} \,d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \longrightarrow 0$$

moreover

$$\begin{aligned} \left| \int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right| \\ &\leq C \left[\eta^{-2} \int_{d^{-1}\eta < \mathfrak{r} < d\eta} d\mu_{\infty} \right]^{1/2} \left[\int_{d^{-1}\eta < \mathfrak{r} < d\eta} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right]^{1/2} \longrightarrow 0 \end{aligned}$$

and

$$\begin{aligned} \left| \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}^2(\vec{p})}{\eta^2} \chi''\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) & \arctan\sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right| \\ & \leq C \left[\eta^{-2} \int_{d^{-1}\eta < \mathfrak{r} < d\eta} d\mu_{\infty} \right]^{1/2} \left[\int_{d^{-1}\eta < \mathfrak{r} < d\eta} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right]^{1/2} \longrightarrow 0 \end{aligned}$$

We have also

$$\int_{G_2^{Leg}(V_2(\mathbb{R}^4))} O\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \ d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \le C \eta^2 \left[\eta^{-2} \int_{d^{-1}\eta < \mathfrak{r} < d\eta} d\mu_{\infty}\right] \longrightarrow 0$$

and

$$\begin{split} &\int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \left[O(1) \left(\chi \left(\frac{\mathfrak{r}(\vec{p})}{r} \right) - \chi \left(\frac{\mathfrak{r}(\vec{p})}{\eta} \right) \right) \right] \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &= \int_{G_2^{Leg}(V_2(\mathbb{R}^4)) \setminus \pi^{-1}\{\mathfrak{r}=0\}} O(1) \, \chi \left(\frac{\mathfrak{r}(\vec{p})}{r} \right) \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &- \sum_{j=0}^{\infty} \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} O(1) \left[\chi \left(\frac{\mathfrak{r}(\vec{p})}{2^{-j}\eta} \right) - \chi \left(\frac{\mathfrak{r}(\vec{p})}{2^{-j-1}\eta} \right) \right] \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \end{split}$$
We have

$$\begin{split} & \left| \sum_{j=0}^{\infty} \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} O(1) \left[\chi\left(\frac{\mathfrak{r}(\vec{p})}{2^{-j\eta}}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{2^{-j-1}\eta}\right) \right] \ d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right. \\ & \leq C \, \eta^2 \sum_{j=0}^{\infty} 2^{-2j} \, (2^{-j\eta})^{-2} \, \int_{d^{-1}2^{-j-1}\eta < \mathfrak{r} < d \, 2^{-j\eta}} \, d\mu_{\infty} \ \longrightarrow 0 \end{split}$$

Finally we observe that

$$4 \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \left[\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \right] \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p})$$

is a monotone function of η . Hence

$$\begin{split} &\lim_{\eta\to 0} 4 \, \int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \left[\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \right] \, \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &= 4 \, \int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4)) \setminus \pi^{-1}(\{0\})} \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \; . \end{split}$$

Combining all the previous we obtain that

$$-\int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \,\chi'\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \,\frac{1}{\mathfrak{r}^2(\vec{p})} \,\left[|\nabla^{\mathcal{P}}\mathfrak{r}|^2 + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^2}}(\vec{p})\right] \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p})$$

is converging as η converges to zero. Hence

$$\lim_{\eta \to 0} \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \phi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{1}{\mathfrak{r}^2(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^2 + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^2}}(\vec{p}) \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) = \theta_0(\phi)$$
(III.29)

exists and is finite. Recall that for any $\mathcal{P} \in G_2^{Leg}(V_2(\mathbb{R}^4))$

$$|\nabla^{H}\mathfrak{r}|^{2} = |\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + |(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^{2}$$

From (III.26) and using also (III.8) we have

$$\frac{|\nabla^{\mathcal{P}}\mathfrak{r}|^2}{\mathfrak{r}^2} = \frac{|\nabla^{\mathcal{H}}\mathfrak{r}|^2}{\mathfrak{r}^2} - \frac{1}{4}\frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1) = \frac{1}{\mathfrak{r}^2}\frac{1}{\sqrt{1+\sigma^2}} - \frac{1}{4}\frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1)$$

We deduce

$$\lim_{\eta \to 0} \int_{G_2^{Leg}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{\eta} \phi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \frac{1}{\mathfrak{r}^2(\vec{p})} \left[\frac{1+\sigma \arctan\sigma}{\sqrt{1+\sigma^2}}(\vec{p})\right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) = \theta_0(\phi)$$

By taking $h_{r,\eta} := [\chi(\mathfrak{r}/r) - \int_{\mathfrak{r}/\eta}^{+\infty} \phi(s) ds] \arctan \sigma$ where χ is fixed and ϕ is arbitrary satisfying the

assumption of the lemma we observe that

$$\begin{split} \theta_{0}(\phi) &= -4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))\backslash\pi^{-1}(\{0\})} \chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}(\vec{p})^{2}} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &- \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \, \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma(\vec{p}) \arctan \sigma(\vec{p})}{\sqrt{1 + \sigma^{2}(\vec{p})}} \right] \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))\backslash\pi^{-1}(\{0\})} \left[O(1) \left(\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \right) + O\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \right] \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}^{2}(\vec{p})}{r^{2}} \, \chi''\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \, \arctan \sigma(\vec{p}) \, \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &- \frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \, \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \, \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \, \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{|\nabla^{\mathcal{P}}\sigma|^{2}}{(1 + \sigma^{2}(\vec{p}))^{2}} \, d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \end{split}$$

Obviously $\theta_0(\phi)$ is independent of ϕ and this concludes the proof of lemma III.2.

We have the following lemma

Lemma III.3. There exists a universal constant C > 1 such that, for any 0 < 2s < r < 1, assuming we have

$$\limsup_{\eta \to 0} \frac{1}{\eta^2} \int_{\eta < \mathfrak{r} < 2\eta} d\mu_{\infty}(\vec{p}) < +\infty , \qquad (\text{III.30})$$

then

$$C^{-1}\left[\theta_0 + \int_{0 < \mathfrak{r} < s/2} \left| \frac{\nabla^{\mathcal{P}} \sigma}{1 + \sigma^2(\vec{p})} \right|^2 d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \right] \le \frac{1}{s^2} \int_{0 < \mathfrak{r} < s} d\mu_{\infty} \le C \frac{1}{r^2} \int_{\mathfrak{r} < r} d\mu_{\infty} .$$
(III.31)

We replace r by s in (III.13) and we make η converge to zero. Thanks to the previous lemma we obtain

$$\begin{split} &-\int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma(\vec{p}) \arctan \sigma(\vec{p})}{\sqrt{1 + \sigma^{2}(\vec{p})}} \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4})) \setminus \{\mathfrak{r}=0\}} \left[O(1) \chi\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) + O\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \right] d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}^{2}(\vec{p})}{s^{2}} \chi''\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &- \frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{|\nabla^{\mathcal{P}}\sigma|^{2}}{(1 + \sigma^{2}(\vec{p}))^{2}} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \\ &= 4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4})) \setminus \{\mathfrak{r}=0\}} \chi\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}(\vec{p})^{2}} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) + \theta_{0} \end{split}$$

Using thee fact that we have

$$|\nabla^{\mathcal{P}}\mathfrak{r}| \le 1$$
 and $\left|\frac{\nabla^{\mathcal{P}}\sigma}{1+\sigma^2(\vec{p})}\right| \le \frac{2}{\mathfrak{r}}$

we deduce

$$\theta_0 + 4 \int_{G_2^{Leg}(V_2(\mathbb{R}^4)) \setminus \{\mathfrak{r}=0\}} \chi\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}(\vec{p})^2} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \leq \frac{C}{s^2} \int_{s < \mathfrak{r} < 2s} d\mu_{\infty} + \int_{0 < \mathfrak{r} < 2s} d\mu_{\infty}$$

Using

$$\frac{|(\nabla^{\mathcal{P}}\mathfrak{r})^{\perp}|^2}{\mathfrak{r}^2} = \frac{1}{4} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1)$$

we deduce the first inequality in (III.31). For 2s < r we have

$$\begin{split} &-\int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma(\vec{p}) \arctan \sigma(\vec{p})}{\sqrt{1 + \sigma^{2}(\vec{p})}} \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O(1) \left(\chi\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) - \chi\left(\frac{\mathfrak{r}(\vec{p})}{\eta}\right) \right) + O\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}^{2}(\vec{p})}{r^{2}} \chi''\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &- \frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{1}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{r} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{r}\right) \frac{|\nabla^{\mathcal{P}}\sigma|^{2}}{(1 + \sigma^{2}(\vec{p}))^{2}} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &= 4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[\chi\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^{2}}(\vec{p})} \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &- \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{1}{\mathfrak{r}^{2}(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^{2} + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^{2}}(\vec{p})} \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ 4 \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{3}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \frac{4}{4} \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \arctan \sigma(\vec{p}) \frac{\nabla^{\mathcal{P}}\mathfrak{r}}{\mathfrak{r}(\vec{p})} \cdot \frac{\nabla^{\mathcal{P}}\sigma}{1 + \sigma^{2}(\vec{p})} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \\ &+ \int_{G_{2}^{Leg}(V_{2}(\mathbb{R}^{4}))} \left[O\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \det \sigma(\mathcal{P},\vec{p}) \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \end{aligned}$$

We have that $\chi' \leq 0$ on $\mathbb R$ and $\mathrm{supp}(\chi') \subset [1,2]$ hence we deduce

$$-\int_{G_2^{L^{eg}}(V_2(\mathbb{R}^4))} \frac{\mathfrak{r}(\vec{p})}{s} \chi'\left(\frac{\mathfrak{r}(\vec{p})}{s}\right) \frac{1}{\mathfrak{r}^2(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^2 + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^2}}(\vec{p}) \right] d\mathbf{v}_{\infty}(\mathcal{P},\vec{p})$$

$$\leq \frac{C}{r^2} \int_{r<\mathfrak{r}<2r} d\mu_{\infty} + \int_{0<\mathfrak{r}<2r} d\mu_{\infty} + C \left[\frac{1}{s^2} \int_{s<\mathfrak{r}<2s} d\mu_{\infty} \right]^{1/2} \left[\int_{s<\mathfrak{r}<2s} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P},\vec{p}) \right]^{1/2}$$

We choose χ such that

$$\chi' \le 0$$
 on \mathbb{R} and $-\chi' > \frac{1}{2}$ on $[5/4, 7/4]$.

Since

$$\frac{|\nabla^{\mathcal{P}}\mathfrak{r}|^2}{\mathfrak{r}^2} = \frac{|\nabla^H\mathfrak{r}|^2}{\mathfrak{r}^2} - \frac{1}{4}\frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1) = \frac{1}{\mathfrak{r}^2}\frac{1}{\sqrt{1+\sigma^2}} - \frac{1}{4}\frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1) \ .$$

Using remark III.1 we deduce

$$\frac{1}{\mathfrak{r}^2(\vec{p})} \left[|\nabla^{\mathcal{P}}\mathfrak{r}|^2 + \frac{\sigma \arctan \sigma}{\sqrt{1+\sigma^2}}(\vec{p}) \right] \ge \frac{1}{\mathfrak{r}^2} - \frac{1}{4} \frac{|\nabla^{\mathcal{P}}\sigma|^2}{(1+\sigma^2)^2} + O(1)$$

We deduce that for r<1

$$\frac{1}{s^2} \int_{5s/4 < \mathfrak{r} < 7s/4} d\mu_{\infty} \leq \frac{C}{r^2} \int_{0 < \mathfrak{r} < 2r} d\mu_{\infty} + C \left[\frac{1}{s^2} \int_{s < \mathfrak{r} < 2s} d\mu_{\infty} \right]^{1/2} \left[\int_{s < \mathfrak{r} < 2s} \frac{|\nabla^{\mathcal{P}} \sigma|^2}{(1 + \sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \right]^{1/2} \\ + C \int_{s < \mathfrak{r} < 2s} \frac{|\nabla^{\mathcal{P}} \sigma|^2}{(1 + \sigma^2(\vec{p}))^2} d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) .$$

Let

$$A := \sup_{2s < r} \frac{1}{s^2} \int_{5s/4 < \mathfrak{r} < 7s/4} d\mu_{\infty}$$

Using the first inequality in (III.31) we just proved, we deduce

$$A \le \frac{C}{r^2} \int_{0 < \mathfrak{r} < 2r} d\mu_{\infty} + C A^{1/2} \left[r^{-2} \int_{0 < \mathfrak{r} < 2r} d\mu_{\infty} \right]^{1/2}$$

Hence for any s < r we have

$$\frac{1}{s^2} \int_{5s/4 < \mathfrak{r} < 7s/4} d\mu_{\infty} \le \frac{C}{r^2} \int_{0 < \mathfrak{r} < 2r} d\mu_{\infty} .$$

This gives for any s < r

$$\frac{1}{s^2} \int_{0 < \mathfrak{r} < s} \ d\mu_{\infty} = \frac{1}{s^2} \sum_{j=0}^{\infty} \int_{5^{j+1}s/7^{j+1} < \mathfrak{r} < 5^j s/7^j} \ d\mu_{\infty} \le C \ \sum_{j=0}^{\infty} \left(\frac{5}{7}\right)^j \ \frac{1}{r^2} \ \int_{0 < \mathfrak{r} < 2r} \ d\mu_{\infty}$$

which implies the second inequality in (III.31). Lemma III.3 is proved.

III.3 The finiteness and a global bound of the upper 2-density of μ_{∞} at every point in $V_2(\mathbb{R}^4)$.

We denote by $\mathfrak{r}_{\vec{p}_0}$ the Folland Korányi gauge with respect to $\vec{p}_0 = (\vec{a}_0, \vec{b}_0) \in V_2(\mathbb{R}^4)$. That is, for any $\vec{p} = (\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$

$$\rho_{\vec{p}_0}^2(\vec{p}) = |\vec{p} - \vec{p}_0|^2 \quad , \quad \varphi_{\vec{p}_0}(\vec{p}) := \vec{a} \cdot \vec{b}_0 - \vec{a}_0 \cdot \vec{b} \quad \text{ and } \quad \mathfrak{r}_{\vec{p}_0}^4 := \rho_{\vec{p}_0}^4 + 4 \, \varphi_{\vec{p}_0}^2 \; .$$

For $\theta \in \mathbb{R}$ and $\vec{p} = (\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$ we denote

$$\vec{p}^{\,\theta} := (\cos\theta\,\vec{a} + \sin\theta\,\vec{b}, -\sin\theta\,\vec{a} + \cos\theta\,\vec{b}) \ .$$

We shall now prove the following lemma

Lemma III.4. For any $\vec{p}_0 \in V_2(\mathbb{R}^4)$ the following holds

$$\limsup_{r \to 0} \frac{1}{r^2} \int_{\mathfrak{r}_{\vec{p}_0} < r} d\mu_{\infty}(\vec{p}) < +\infty .$$
(III.34)

Hence there exists C > 1 universal such that for any $\vec{p}_0 \in V_2(\mathbb{R}^4)$ and any 0 < s < r

$$C^{-1} \left[\theta_0 + \int_{0 < \mathfrak{r}_{\vec{p}_0} < s/2} \left| \frac{\nabla^{\mathcal{P}} \sigma}{1 + \sigma^2(\vec{p})} \right|^2 \, d\mathbf{v}_{\infty}(\mathcal{P}, \vec{p}) \right] \le \frac{1}{s^2} \, \int_{\mathfrak{r}_{\vec{p}_0} < s} \, d\mu_{\infty} \le C \, \frac{1}{r^2} \, \int_{\mathfrak{r}_{\vec{p}_0} < r} \, d\mu_{\infty} \, . \tag{III.35}$$

Proof of lemma III.4. Let M > 0 and denote \vec{p} such that

$$E_{\infty} := \left\{ \vec{p} \in V_2(\mathbb{R}^4) ; \limsup_{r \to 0} \frac{1}{r^2} \int_{\mathfrak{r}_{\vec{p}} < r} d\mu_{\infty} = +\infty \right\} .$$

Let M > 0 and $\delta > 0$. For any $\vec{p} \in E_{\infty}$ we choose $\delta > r_{\vec{p}} > 0$ such that

$$\frac{1}{r_{\vec{p}}^2}\int_{\mathfrak{r}_{\vec{p}} < r_{\vec{p}}} d\mu_\infty > M \ .$$

Let

$$\mathfrak{M}_{\infty} := \Pi^* \mu_{\infty}$$
 .

where we recall that Π is the tautological projection from $V_2(\mathbb{R}^4)$ into $G_2(\mathbb{R}^4)$ such that $\Pi(\vec{a}, \vec{b}) = \vec{a} \wedge \vec{b}$. Let $\vec{G}_{\vec{p}} := \Pi(\vec{p})$. We have for any pair (\vec{a}, \vec{b}) and (\vec{c}, \vec{d}) in $V_2(\mathbb{R}^4)$

$$|\vec{a} \wedge \vec{b} - \vec{c} \wedge \vec{d}| = |(\vec{a} - \vec{c}) \wedge \vec{b} - \vec{c} \wedge (\vec{d} - \vec{b})| \le |\vec{a} - \vec{c}| + |\vec{d} - \vec{b}| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right| \le \sqrt{2} \left| (\vec{a}, \vec{b}) - (\vec{c}, \vec{d}) \right|$$

Let $\vec{G}_{\vec{p}} := \Pi(\vec{p})$, because of the previous

$$\Pi\left(\{\vec{q} \in V_2(\mathbb{R}^4) ; \ \mathfrak{r}_{\vec{p}}(\vec{q}) < r_{\vec{p}}\}\right) \subset B_{\sqrt{2}r_{\vec{p}}}(\vec{G}_{\vec{p}})$$

Hence we have

$$\frac{1}{r_{\vec{p}}^2}\int_{B_{\sqrt{2}r_{\vec{p}}}(\vec{G}_{\vec{p}})}d\mathfrak{M}_\infty > M$$

Obviously $B_{\sqrt{2}r_{\vec{p}}}(\vec{G}_{\vec{p}})$ realizes an open cover of $\Pi(E_{\infty})$. Extracting a Besicovitch covering $(B_{\sqrt{2}r_i}(\vec{G}_i))_{i\in I}$, we deduce that

$$M \sum_{i \in I} r_i^2 \le C \int_{G_2(\mathbb{R}^4)} d\mathfrak{M}_{\infty}$$

where C is a constant depending only on $G_2(\mathbb{R}^4)$ given by the Besicovitch covering theorem. Since M > 0and $\delta > 0$ are arbitrary, we deduce

$$\mathcal{H}^2(\Pi(E_\infty)) = 0$$

This implies in particular that E_{∞} has no interior. Recall that from lemma III.3, for any $\vec{p} \in V_2(\mathbb{R}^4) \setminus E_{\infty}$, we have

$$\forall r < 1 \qquad \frac{1}{r^2} \int_{0 < \mathfrak{r}_{\vec{p}} < r} d\mu_{\infty} \le C \, \mu_{\infty}(V_2(\mathbb{R}^4)) \, .$$

where C > 0. Moreover, since by assumption

١

$$\limsup_{r \to 0} \frac{1}{r^2} \int_{\mathfrak{r}_{\vec{p}} < r} d\mu_{\infty} < +\infty ,$$

we have

$$\forall \vec{p} \in V_2(\mathbb{R}^4) \setminus E_{\infty} \qquad \mu_{\infty}(\{\vec{p}\}) = 0 .$$

Hence

$$\forall \vec{p} \in V_2(\mathbb{R}^4) \setminus E_\infty \qquad \forall r < 1 \qquad \frac{1}{r^2} \int_{\mathfrak{r}_{\vec{p}} < r} d\mu_\infty \leq C \,\mu_\infty(V_2(\mathbb{R}^4)) \;.$$

Let now $\vec{p_0} = (\vec{a}_0, \vec{b}_0) \in E_{\infty}$ and $\vec{p}_k = (\vec{a}_k, \vec{b}_k) \to \vec{p}_{\infty}$ with $\vec{p}_k \in V_2(\mathbb{R}^4) \setminus E_{\infty}$. Let r > 0 and $k \in \mathbb{N}$ such that $\mathfrak{r}_{\vec{p}_0}(\vec{p}_k) < 2^{-1} C_0^{-1} r$ where $C_0 > 1$ is going to be fixed later on below. We have for any $\vec{p} \in V_2(\mathbb{R}^4)$

$$\mathfrak{r}_{\vec{p}_0}(\vec{p}) = \left[|\vec{p} - \vec{p}_0|^4 + 4\,\varphi_{\vec{p}_0}^2(\vec{p}) \right]^{1/4} = \left[|\vec{p} - \vec{p}_0|^4 + 4\,(\vec{a}\cdot\vec{b}_0 - \vec{a}_0\cdot\vec{b})^2 \right]^{1/4}$$

We write

$$\begin{split} \varphi_{\vec{p}_0}(\vec{p}) &= \vec{a} \cdot \vec{b}_0 - \vec{b} \cdot \vec{a}_0 = (\vec{a} - \vec{a}_0) \cdot \vec{b}_0 - (\vec{b} - \vec{b}_0) \cdot \vec{a}_0 \\ &= (\vec{a} - \vec{a}_0) \cdot (\vec{b}_0 - \vec{b}_k) - (\vec{b} - \vec{b}_0) \cdot (\vec{a}_0 - \vec{a}_k) + (\vec{a} - \vec{a}_0) \cdot \vec{b}_k - (\vec{b} - \vec{b}_0) \cdot \vec{a}_k \\ &= (\vec{a} - \vec{a}_0) \cdot (\vec{b}_0 - \vec{b}_k) - (\vec{b} - \vec{b}_0) \cdot (\vec{a}_0 - \vec{a}_k) + \varphi_{\vec{p}_k}(\vec{p}) + \varphi_{\vec{p}_0}(\vec{p}_k) \\ &\qquad |\varphi_{\vec{p}_0}(\vec{p})| \le \rho_{\vec{p}_0}(\vec{p}) \ \rho_{\vec{p}_0}(\vec{p}_k) + |\varphi_{\vec{p}_k}(\vec{p})| + |\varphi_{\vec{p}_0}(\vec{p}_k)| \\ &\le \rho_{\vec{p}_0}(\vec{p}_k) \ \rho_{\vec{p}_0}(\vec{p}_k) + \rho_{\vec{p}_k}(\vec{p}) \ \rho_{\vec{p}_0}(\vec{p}_k) + |\varphi_{\vec{p}_k}(\vec{p})| + |\varphi_{\vec{p}_0}(\vec{p}_k)| \end{split}$$

Hence

Hence

$$\begin{aligned} \mathbf{r}_{\vec{p}_{0}}(\vec{p}) &\leq \rho_{\vec{p}_{0}}(\vec{p}) + \sqrt{2} \left| \varphi_{\vec{p}_{0}}(\vec{p}) \right| \\ &\leq \rho_{\vec{p}_{0}}(\vec{p}_{k}) + \rho_{\vec{p}_{k}}(\vec{p}) + \sqrt{2} \rho_{\vec{p}_{0}}(\vec{p}_{k}) + \sqrt{2} \rho_{\vec{p}_{0}}(\vec{p}_{k}) + \sqrt{2} \left| \varphi_{\vec{p}_{0}}(\vec{p}_{k}) \right| + \sqrt{2} \left| \varphi_{\vec{p}_{0}}(\vec{p}_{k}) \right| \end{aligned}$$
(III.36)
$$&\leq C_{0} \left[\mathbf{r}_{\vec{p}_{k}}(\vec{p}) + \mathbf{r}_{\vec{p}_{0}}(\vec{p}_{k}) \right] \end{aligned}$$

where $C_0 > 1$ is universal. We choose \vec{p}_k such that $\mathfrak{r}_{\vec{p}_0}(\vec{p}_k) < 2^{-1} C_0^{-1} r$. Hence

$$\left\{ \vec{p} \in V_2(\mathbb{R}^4) \; ; \; \mathfrak{r}_{\vec{p}_0}(\vec{p}) < r \right\} \subset \left\{ \vec{p} \in V_2(\mathbb{R}^4) \; ; \; \mathfrak{r}_{\vec{p}_k}(\vec{p}) < 2^{-1} C_0^{-1} r \right\}$$

Since $\vec{p}_k \in V_2(\mathbb{R}^4) \setminus E_\infty$ we have

$$r^{-2} \int_{\mathfrak{r}_{\vec{p}_k}(\vec{p}) < 2^{-1} C_0^{-1} r} d\mu_{\infty} \le C \, \mu_{\infty}(V_2(\mathbb{R}^4))$$

where C is universal. Hence we deduce

$$r^{-2} \int_{\mathfrak{r}_{\vec{p}_0}(\vec{p}) < r} d\mu_\infty \le C \, \mu_\infty(V_2(\mathbb{R}^4)) \; .$$

This holds for any 0 < r < 1 which contradicts the fact that $\vec{p}_0 \in E_{\infty}$. Hence $E_{\infty} = \emptyset$ and lemma III.4 is proved.

III.4 Bounding the limiting 2-density of μ_{∞} from below

We take again

$$h_{r,\eta} := [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \arctan \sigma$$
.

we have

$$\nabla^H \arctan \sigma = \frac{\rho^2}{\mathfrak{r}^4} \, \nabla^H \varphi - \frac{1}{\mathfrak{r}^4} \, \varphi \, \nabla^H \rho^2 \, \, .$$

Hence we have

$$\begin{split} \vec{X}_{h_{r,\eta}} &:= J_H \nabla^H h_{r,\eta} + \frac{h_{r,\eta}}{2} \vec{R} \\ &= \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \left[\frac{\rho^2}{\mathfrak{r}^4} J_H \nabla^H \varphi - \frac{1}{\mathfrak{r}^4} \varphi J_H \nabla^H \rho^2 + \arctan \sigma \; \frac{\vec{R}}{2} \right] + \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \arctan \sigma \nabla^H \mathfrak{r} \; . \end{split}$$

Recall that

$$\nabla^H \varphi = \sum_{i=1}^4 v^i \, \nabla^H w^i - w^i \, \nabla^H v^i \; .$$

Hence

$$\begin{cases} |\vec{X}_{h_{r,\eta}}| \leq C \sum_{j=0}^{1} \mathfrak{r}^{-1+j} \left| r^{-j} \chi^{(j)}(\mathfrak{r}/r) - \eta^{-j} \chi^{(j)}(\mathfrak{r}/\eta) \right| \\ |\nabla^{H} \vec{X}_{h_{r,\eta}}| \leq C \sum_{j=0}^{2} \mathfrak{r}^{-2+j} \left| r^{-j} \chi^{(j)}(\mathfrak{r}/r) - \eta^{-j} \chi^{(j)}(\mathfrak{r}/\eta) \right| \\ |\nabla^{H} \nabla^{H} \vec{X}_{h_{r,\eta}}| \leq C \sum_{j=0}^{3} \mathfrak{r}^{-3+j} \left| r^{-j} \chi^{(j)}(\mathfrak{r}/r) - \eta^{-j} \chi^{(j)}(\mathfrak{r}/\eta) \right| \end{cases}$$

We have also

$$\nabla \vec{X}_h \cdot \vec{R} = \nabla \left[J_H \nabla^H h_{r,\eta} \right] \cdot \vec{R} + \nabla h_{r,\eta} = -J_H \nabla^H h_{r,\eta} \cdot \nabla^H \vec{R} + \nabla h_{r,\eta}$$

which gives

$$|\nabla \vec{X}_h \cdot \vec{R}| \le C \,\mathfrak{r}^{-1} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] + C \,\left| \left[r^{-1} \,\chi'(\mathfrak{r}/r) - \eta^{-1} \,\chi'(\mathfrak{r}/\eta) \right| \;.$$

Recall

$$\int_{\Sigma} \langle d\vec{w}_{r,\eta}, d\vec{\Lambda}_k \rangle_g \ dvol_g = 2 \int_{\Sigma} \langle dh_{r,\eta}, d\beta_k \rangle_g \ dvol_g \ . \tag{III.37}$$

 $\quad \text{and} \quad$

$$\begin{aligned} \int_{\Sigma} &< d\vec{w}_{r,\eta} \cdot d\vec{\Lambda}_k >_{g_k} dvol_{g_k} + e^{-\varepsilon_k^{-2}} O(\|w\|_{\vec{\Lambda}_k}) \\ &= -4 \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2) \left[-\left\langle d\vec{T}_k \dot{\otimes} d\vec{T}_k, d\vec{w} \dot{\otimes} d\vec{\Lambda}_k \right\rangle_g + d\vec{T}_k \cdot d\left\langle d\vec{w} \wedge d\vec{\Lambda}_k \right\rangle_{g_k} \right] dvol_{g_k} \quad (\text{III.38}) \\ &- \varepsilon_k^4 \int_{\Sigma} \left[(1 + |d\vec{T}_k|_{g_k}^2)^2 - 4 |d\vec{T}_k|_{g_k}^2 (1 + |d\vec{T}_k|_{g_k}^2) \right] < d\vec{w} \cdot d\vec{\Lambda}_k >_{g_k} dvol_{g_k} \end{aligned}$$

and we recall from (II.15) and from (II.20)

$$|d\vec{\Lambda}_k \otimes d\vec{w}_{r,\eta}|_{g_k} \le |\nabla^H \vec{X}_{h_{r,\eta}}| \tag{III.39}$$

and

$$\left| d\vec{T}_k \cdot d\left\langle d\vec{w}_{r,\eta} \wedge d\vec{\Lambda}_k \right\rangle_g \right| \le \left| d\vec{T}_k \right|_g^2 \left| \nabla^H \vec{X}_{h_{r,\eta}} \right| + \left| d\vec{T}_k \right|_g \left[\left| \nabla^H \nabla^H \vec{X}_{h_{r,\eta}} \right| + \left| \nabla \vec{X}_{h_{r,\eta}} \cdot \vec{R} \right| \right]$$
(III.40)

Hence

$$\begin{split} \left| \int_{\Sigma} &< dh_{r,\eta}, d\beta_k >_g \ dvol_g \right| \le C \, \varepsilon_k^4 \, \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2)^2 \ |\nabla^H \vec{X}_{h_{r,\eta}}| \ dvol_{g_k} \\ &+ C \, \varepsilon_k^4 \, \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2)^{3/2} \ [|\nabla^H \nabla^H \vec{X}_{h_{r,\eta}}| + |\nabla \vec{X}_{h_{r,\eta}} \cdot \vec{R}|] \ dvol_{g_k} + e^{-\varepsilon_k^{-2}} \, O(||w||_{\vec{\Lambda}_k}) \end{split}$$

This gives

$$\left| \int_{\Sigma} \langle dh_{r,\eta}, d\beta_k \rangle_g \ dvol_g \right| \leq C \,\varepsilon_k^4 \, \int_{\Sigma \cap \{\eta < \mathfrak{r} < 2r\}} \mathfrak{r}^{-2} \, (1 + |d\vec{T}_k|_{g_k}^2)^2 \ dvol_{g_k} + C \,\varepsilon_k^4 \, \int_{\Sigma \cap \{\eta < \mathfrak{r} < 2r\}} \mathfrak{r}^{-3} \, (1 + |d\vec{T}_k|_{g_k}^2)^{3/2} \ dvol_{g_k} + e^{-\varepsilon_k^{-2}} \, O(\|w\|_{\vec{\Lambda}_k})$$
(III.41)

We recall that the function

$$d(\vec{p},\vec{q}) := \mathfrak{r}_{\vec{p}}(\vec{q}) = \left[|\vec{p} - \vec{q}|^4 + 4 \, |\vec{a} \cdot \vec{d} - \vec{b} \cdot \vec{c}|^2 \right]^{1/4} \quad \text{where} \quad \vec{p} = (\vec{a},\vec{b}) \quad \text{and} \ \vec{q} = (\vec{c},\vec{d})$$

and we have seen in - see (III.36) - that it satisfies the quasi-distance condition

$$\mathfrak{r}_{\vec{p}}(\vec{q}) = \mathfrak{r}_{\vec{q}}(\vec{p}) \quad \text{and} \quad d(\vec{p}_1, \vec{p}) \le C_0 \left[d(\vec{p}_1, \vec{q}) + d(\vec{p}, \vec{q}) \right]$$
(III.42)

for some universal $C_0 > 1$. We are now proving the following lemma.

Lemma III.5. There exist two universal constants C > 0 and $c_0 > 0$ such that for any $\delta_k \to 0$ and for k large enough and for any point $\vec{p} \in \vec{\Lambda}_k(\Sigma)$ satisfying

$$\forall r < 1 \qquad \varepsilon_k^4 \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < r\}} (1 + |d\vec{T}_k|_{g_k}^2)^2 \ dvol_{g_k} \le \delta_k \ \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < 5\,C_0^2\,r\}} \ dvol_{g_k} \ , \tag{III.43}$$

we have for any 0 < r < 1/2

$$\int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < r\}} |\nabla^{\Sigma} \arctan \sigma_{k}|^{2} dvol_{g_{k}} \leq \frac{C}{r^{2}} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < r\}} dvol_{g_{k}}
+ C \left[\delta \log\left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4}\left(\frac{r}{\varepsilon}\right)\right] \sup_{s < r} \frac{1}{s^{2}} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_{g_{k}}
+ C e^{-\varepsilon_{k}^{-2}} \varepsilon_{k}^{-3},$$
(III.44)

and for any 0 < s < r < 1/2

$$\begin{bmatrix} c_0 - \left[\delta \log\left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4}\left(\frac{r}{\varepsilon}\right)\right]^{1/2} \end{bmatrix} \frac{1}{s^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_{g_k} \\ \leq \frac{C}{r^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < r\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \varepsilon_k^{-3} .$$
(III.45)

Proof of lemma III.5. We take $\vec{p} := (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$. Since the second fundamental form of $\vec{\Lambda}_k$ is in L^4 the weak Immersion is C^1 . Hence \vec{p} has finitely many pre-images by $\vec{\Lambda}_k$ and we denote their number by $N_{\vec{p}} \in \mathbb{N}$. We choose \vec{p} such that $N_{\vec{p}} \geq 1$. Let $x_0 \in \Sigma$ such that $\vec{\Lambda}_k(x_0) = \vec{p}$. We take local conformal coordinates around x_0 such that $x_i(x_0) = 0$ and we omit to write explicitly the subscript k. We can assume modulo rotation in the image and dilation of the coordinates that $\partial_{x_1}\vec{\Lambda}(0) = (\vec{\varepsilon}_3, 0)$. Since both $J_H(\partial_{x_1}\vec{\Lambda}(0)) = (0, \vec{\varepsilon}_3)$ and $J_H(\partial_{x_2}\vec{\Lambda}(0))$ are orthogonal to $\vec{\Lambda}_*T_{x_0}\Sigma$ we can assume modulo the action of rotations preserving H and J_H that $\partial_{x_2}\vec{\Lambda}(0) = (\vec{\varepsilon}_4, 0)$ and $J_H(\partial_{x_2}\vec{\Lambda}(0)) = (0, \vec{\varepsilon}_4)$. Hence, since $\vec{\Lambda}$ is C^1 we have in the neighbourhood of 0

$$\Lambda(x_1, x_2) = (\vec{\varepsilon}_1 + x_1 \, \vec{\varepsilon}_3 + x_2 \, \vec{\varepsilon}_4 + o(|x|), \vec{\varepsilon}_2 + o(|x|))$$

Recall that for $\vec{a} = \vec{\varepsilon}_1 + \vec{v}$ and $\vec{b} := \vec{\varepsilon}_2 + \vec{w}$ with $(\vec{a}, \vec{b}) \in V_2(\mathbb{R}^4)$ recall that $\varphi = v^2 - w^1$

$$\nabla^H \varphi(\vec{a}, \vec{b}) = \sum_{i=1}^4 v^i \, \nabla^H w^i - w^i \, \nabla^H v^i \, .$$

Recall for $\vec{p} = (\vec{a}, \vec{b}) = (\vec{\varepsilon}_1 + \vec{v}, \vec{\varepsilon}_2 + \vec{w})$

$$\begin{aligned} H_{\vec{p}} &= T_{\vec{p}} V_2(\mathbb{R}^4) \cap (\vec{R})^{\perp} \\ &= \left\{ (\vec{V}, \vec{W}) \in \mathbb{R}^8 \ ; \ V_1 + \vec{V} \cdot \vec{v} = 0 \ ; \ V_2 + \vec{V} \cdot \vec{w} = 0 \ ; \ W_1 + \vec{W} \cdot \vec{v} = 0 \ ; \ W_2 + \vec{W} \cdot \vec{w} = 0 \right\} \end{aligned}$$

we have respectively

 $\nabla^H v^i = \pi_H(\vec{\varepsilon_i}, 0)$ and $\nabla^H w^i = \pi_H(0, \vec{\varepsilon_i})$.

First of all we have at the point $\vec{p} := (\vec{a}, \vec{b})$

$$\nabla^{V_2(\mathbb{R}^4)} v^i = (\vec{\varepsilon}_i - a^i \, \vec{a} - b^i \, \vec{b}, 0) \quad \text{and} \quad \nabla^{V_2(\mathbb{R}^4)} w^i = (0, \vec{\varepsilon}_i - a^i \, \vec{a} - b^i \, \vec{b})$$

Hence

$$\nabla^{H} v^{i} = (\vec{\varepsilon_{i}} - a^{i} \vec{a} - b^{i} \vec{b}, 0) \quad \text{and} \quad \nabla^{H} w^{i} = (0, \vec{\varepsilon_{i}} - a^{i} \vec{a} - b^{i} \vec{b})$$
(III.46)

In particular at $\vec{p} = (\vec{a}, \vec{b}) = (\vec{\varepsilon}_1 + \vec{v}, \vec{\varepsilon}_2 + \vec{w})$, thanks to (III.1) there holds

$$\nabla^{H} v^{1} = (-\vec{v} - w^{1} \ (\vec{\varepsilon}_{2} + \vec{w}) - v^{1} \ (\vec{\varepsilon}_{1} + \vec{v}), 0)$$

$$\nabla^{H} v^{2} = (-\vec{w} - v^{2} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{2} \ (\vec{\varepsilon}_{2} + \vec{w}), 0)$$

$$\nabla^{H} v^{3} = (\vec{\varepsilon}_{3} - v^{3} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{3} \ (\vec{\varepsilon}_{2} + \vec{w}), 0)$$

$$\nabla^{H} v^{4} = (\vec{\varepsilon}_{4} - v^{4} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{4} \ (\vec{\varepsilon}_{2} + \vec{w}), 0) ,$$
(III.47)

and

$$\begin{cases} \nabla^{H}w^{1} = (0, -\vec{v} - w^{1} \ (\vec{\varepsilon}_{2} + \vec{w}) - v^{1} \ (\vec{\varepsilon}_{1} + \vec{v})) \\ \nabla^{H}w^{2} = (0, -\vec{w} - v^{2} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{2} \ (\vec{\varepsilon}_{2} + \vec{w})) \\ \nabla^{H}w^{3} = (0, \vec{\varepsilon}_{3} - v^{3} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{3} \ (\vec{\varepsilon}_{2} + \vec{w})) \\ \nabla^{H}w^{4} = (0, \vec{\varepsilon}_{4} - v^{4} \ (\vec{\varepsilon}_{1} + \vec{v}) - w^{4} \ (\vec{\varepsilon}_{2} + \vec{w})) . \end{cases}$$
(III.48)

Observe that for i = 1...4

$$J_H \nabla^H v^i = \nabla^H w^i . (III.49)$$

Thus in particular

$$\begin{aligned} \nabla^{H}v^{1} \circ \vec{\Lambda}(x) &= O(|x|) \\ \nabla^{H}v^{2} \circ \vec{\Lambda}(x) &= O(|x|) \\ \nabla^{H}v^{3} \circ \vec{\Lambda}(x) &= (\vec{\varepsilon_{3}}, 0) + O(|x|) \\ \nabla^{H}v^{4} \circ \vec{\Lambda}(x) &= (\vec{\varepsilon_{4}}, 0) + O(|x|) \end{aligned} \qquad \text{and} \qquad \begin{cases} \nabla^{H}w^{1} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{H}w^{2} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{H}w^{3} \circ \vec{\Lambda}(x) = (0, \vec{\varepsilon_{3}}) + O(|x|) \\ \nabla^{H}w^{4} \circ \vec{\Lambda}(x) &= (0, \vec{\varepsilon_{4}}) + O(|x|) \end{aligned} \qquad (III.50)$$

Thus

$$\begin{cases} \nabla^{\Sigma} v^{1} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{\Sigma} v^{2} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{\Sigma} v^{3} \circ \vec{\Lambda}(x) = (\vec{\varepsilon_{3}}, 0) + O(|x|) \\ \nabla^{H} v^{4} \circ \vec{\Lambda}(x) = (\vec{\varepsilon_{4}}, 0) + O(|x|) \end{cases} \quad \text{and} \quad \begin{cases} \nabla \Sigma w^{1} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{\Sigma} w^{2} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{\Sigma} w^{3} \circ \vec{\Lambda}(x) = O(|x|) \\ \nabla^{\Sigma} w^{4} \circ \vec{\Lambda}(x) = O(|x|) \end{cases}$$

Hence we deduce in particular that

$$w^3 \circ \vec{\Lambda} = O(|x|^2)$$
 and $w^4 \circ \vec{\Lambda} = O(|x|^2)$

Since

$$\nabla^H \varphi \circ \vec{\Lambda} = \sum_{i=3}^4 v^i \circ \vec{\Lambda} \, \nabla^H w^i \circ \vec{\Lambda} - w^i \circ \vec{\Lambda} \, \nabla^H v^i \circ \vec{\Lambda} + O(|x|^2) \; .$$

we deduce from the previous that

$$abla^{\Sigma} \varphi = O(|x|^2) \quad \text{and} \quad \varphi(x) = O(|x|^3) \ .$$

Since $\rho^2 = |x|^2 (1 + o(1))$ we deduce

$$|\nabla^{\Sigma} \arctan \sigma| = O(1)$$
 and $|\arctan \sigma| = O(|x|)$. (III.51)

Since $\vec{\Lambda}$ is a C^1 immersion we have

$$\lim_{s \to 0} \left\| \frac{\rho}{\mathfrak{r}} - 1 \right\|_{L^{\infty}(\vec{\Lambda}^{-1}B_{\eta}(\vec{p}))} = 0$$

and then, using again that $\vec{\Lambda}$ is a C^1 immersion we have

$$\lim_{\eta \to 0} \frac{1}{\eta^2} \int_{\mathfrak{r}_{\vec{p}} < \eta} dvol_{g_k} = \lim_{\eta \to 0} \frac{1}{\eta^2} \int_{\rho_{\vec{p}} < \eta} dvol_{g_k} = \pi N_{\vec{p}}$$
(III.52)

We recall the identity (III.12)

$$\begin{split} &\int_{\Sigma} < dh_{r,\eta}, d\beta >_{g} dvol_{g} - \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^{2}}} \right] dvol_{g} \\ &+ \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{r}\right) + O\left(\frac{\mathfrak{r}}{r}\right) \chi'\left(\frac{\mathfrak{r}}{r}\right) \right] dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{r^{2}} \chi''\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} \\ &- \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1+\sigma^{2})^{2}} dvol_{g} \\ &= 4 \int_{\Sigma} [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} dvol_{g} - \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^{2}}} \right] dvol_{g} \\ &+ 4 \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1+\sigma^{2})^{2}} dvol_{g} - \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} + \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{\eta}\right) + O\left(\frac{\mathfrak{r}}{\eta}\right) \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] dvol_{g} \\ &+ \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{\eta^{2}} \chi''\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} + \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{\eta}\right) + O\left(\frac{\mathfrak{r}}{\eta}\right) \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] dvol_{g} \\ &(\text{III.53}) \end{split}$$

Using (III.51) we have

$$\int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{1}{\mathfrak{r}^2} \left[|\nabla^{\Sigma}\mathfrak{r}|^2 + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^2}} \right] dvol_g$$
$$= \frac{1}{\eta^2} \int_{\mathfrak{r}<2\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{\eta}{\mathfrak{r}} |\nabla^{\Sigma}\mathfrak{r}|^2 dvol_g + O\left(\frac{1}{\eta} \int_{\mathfrak{r}<2\eta} dvol_g\right)$$

From (III.8), (III.26) and (III.51) we also have

$$1 + O(\mathfrak{r}^2) = |\nabla^H \mathfrak{r}|^2 = |\nabla^\Sigma \mathfrak{r}|^2 + |(\nabla^\Sigma \mathfrak{r})^\perp|^2 = |\nabla^\Sigma \mathfrak{r}|^2 + 4^{-1} \mathfrak{r}^2 |\nabla^\Sigma (\arctan \sigma)|^2 = |\nabla^\Sigma \mathfrak{r}|^2 + O(\mathfrak{r}^2)$$

Con ng the previous gives the

$$-\int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^{2}}} \right] dvol_{g}$$

$$= -\frac{1}{\eta} \int_{\mathfrak{r}<2\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{\eta}{\mathfrak{r}} \left| \nabla^{\Sigma}\left(\frac{\mathfrak{r}}{\eta}\right) \right| dvol_{g} + o(1) = -\int_{0}^{\infty} \chi'(s) \frac{\mathcal{H}^{1}(\{\mathfrak{r}_{\vec{p}} = \eta s\})}{\eta s} ds + o(1)$$
(III.54)
ence

Hence

$$\lim_{\eta \to 0} -\int_{\Sigma} \frac{\mathfrak{r}}{\eta} \,\chi'\left(\frac{\mathfrak{r}}{\eta}\right) \,\frac{1}{\mathfrak{r}^2} \left[|\nabla^{\Sigma}\mathfrak{r}|^2 + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^2}} \right] \,dvol_g = 2 \,\pi \,N_{\vec{p}} \tag{III.55}$$

Thus

$$\begin{split} \lim_{\eta \to 0} 4 \int_{\Sigma} [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} \, dvol_{g} - \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1 + \sigma^{2}}} \right] \, dvol_{g} \\ + 4 \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1 + \sigma^{2})^{2}} \, dvol_{g} - \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1 + \sigma^{2}} \, dvol_{g} \\ + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{\eta^{2}} \chi''\left(\frac{\mathfrak{r}}{\eta}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1 + \sigma^{2}} \, dvol_{g} + \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{\eta}\right) + O\left(\frac{\mathfrak{r}}{\eta}\right) \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \, dvol_{g} \\ = 4 \int_{\Sigma} \chi\left(\frac{\mathfrak{r}}{r}\right) \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} \, dvol_{g} + 2\pi N_{\vec{p}} \end{split}$$
(III.56)

We have also

$$\begin{split} &\int_{\Sigma} < dh_{r,\eta}, d\beta >_g \ dvol_g = \frac{1}{2} \int_{\Sigma} < d\vec{w}_{r,\eta}, d\vec{\Lambda} >_g \ dvol_g = \frac{1}{2} \int_{\Sigma} \vec{X}_{r,\eta} \circ \Lambda_k \cdot \Delta_g \vec{\Lambda} \ dvol_g \\ &= -\int_{\Sigma} \nabla^H \left[(\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)) \arctan \sigma \right] \cdot J_H \pi_{\vec{N}} \Delta_g \vec{\Lambda} dvol_g \\ &= -\int_{\Sigma} \nabla^{\Sigma} \left[(\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)) \arctan \sigma \right] \cdot J_H \pi_{\vec{N}} \vec{H}_{\vec{\Lambda}} \ dvol_g \end{split}$$

where we have used the fact that $\vec{R} \circ \Lambda \cdot \Delta \vec{\Lambda} = 0$ - see (II.5) - and the fact that J_H realizes an isomorphism between the horizontal projection of the normal bundle of the immersion and the tangent bundle to the immersion. We have then

$$\begin{split} \left| \int_{\Sigma} < dh_{r,\eta}, d\beta >_{g} dvol_{g} \right| &\leq C \int_{\eta < \mathfrak{r} < 2r} |\nabla^{\Sigma} \arctan \sigma| \sqrt{1 + |d\vec{T}|_{g}^{2}} dvol_{g} \\ &+ C \int_{\eta < \mathfrak{r} < 2\eta} |x|^{-1} |\arctan \sigma| \sqrt{1 + |d\vec{T}_{k}|_{g_{k}}^{2}} dvol_{g} + C \int_{r < \mathfrak{r} < 2r} |x|^{-1} |\arctan \sigma| \sqrt{1 + |d\vec{T}|_{g}^{2}} dvol_{g} \\ &\leq C \left[\int_{\eta < \mathfrak{r} < 2r} |\nabla^{\Sigma} \arctan \sigma|^{2} dvol_{g} \right]^{1/2} \left[\int_{\eta < \mathfrak{r} < 2r} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \right]^{1/4} \left[\int_{\eta < \mathfrak{r} < 2r} dvol_{g} \right]^{1/4} \\ &+ C \left[\int_{\eta < \mathfrak{r} < 2\eta} |x|^{-2} |\arctan \sigma|^{2} dvol_{g} \right]^{1/2} \left[\int_{\eta < \mathfrak{r} < 2\eta} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \right]^{1/4} \left[\int_{\eta < \mathfrak{r} < 2\eta} dvol_{g} \right]^{1/4} \\ &+ C \int_{r < \mathfrak{r} < 2r} |x|^{-1} |\arctan \sigma| \sqrt{1 + |d\vec{T}|_{g}^{2}} dvol_{g} \end{split}$$
(III.57)

Since $|\nabla^{\Sigma} \arctan \sigma| = O(1)$ and $|\arctan \sigma| = O(|x|)$ we deduce

$$\begin{split} & \limsup_{\eta \to 0} \left| \int_{\Sigma} \langle dh_{r,\eta}, d\beta \rangle_g \ dvol_g \right| \le C \ \int_{r < \mathfrak{r} < 2r} |x|^{-1} |\arctan\sigma| \ \sqrt{1 + |d\vec{T}|_g^2} \ dvol_g \\ &+ C \left[\int_{\mathfrak{r} < 2r} |\nabla^{\Sigma} \arctan\sigma|^2 \ dvol_g \right]^{1/2} \ \left[\int_{\mathfrak{r} < 2r} (1 + |d\vec{T}|_g^2)^2 \ dvol_g \right]^{1/4} \ \left[\int_{\mathfrak{r} < 2r} \ dvol_g \right]^{1/4} \end{split}$$
(III.58)

Combining (III.53), (III.56) and (III.58) gives

$$4 \int_{\Sigma} \chi\left(\frac{\mathfrak{r}}{r}\right) \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} dvol_{g} + 2\pi N_{\vec{p}} \leq -\int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{1}{\mathfrak{r}^{2}} \left[|\nabla^{\Sigma}\mathfrak{r}|^{2} + \frac{\sigma \arctan\sigma}{\sqrt{1+\sigma^{2}}} \right] dvol_{g} \\ + \int_{\Sigma} \left[O(1)\chi\left(\frac{\mathfrak{r}}{r}\right) + O\left(\frac{\mathfrak{r}}{r}\right)\chi'\left(\frac{\mathfrak{r}}{r}\right) \right] dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}^{2}}{r^{2}} \chi''\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} \\ - \frac{3}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \arctan\sigma \frac{\nabla^{\Sigma}\mathfrak{r}}{\mathfrak{r}} \cdot \frac{\nabla^{\Sigma}\sigma}{1+\sigma^{2}} dvol_{g} + \frac{1}{4} \int_{\Sigma} \frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) \frac{|\nabla^{\Sigma}\sigma|^{2}}{(1+\sigma^{2})^{2}} dvol_{g} \qquad (\text{III.59}) \\ + C \int_{r < \mathfrak{r} < r} |x|^{-1} |\arctan\sigma| \sqrt{1 + |d\vec{T}|_{g}^{2}} dvol_{g} \\ + C \left[\int_{\mathfrak{r} < 2r} |\nabla^{\Sigma}\arctan\sigma|^{2} dvol_{g}\right]^{1/2} \left[\int_{\mathfrak{r} < 2r} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g}\right]^{1/4} \left[\int_{\mathfrak{r} < 2r} dvol_{g}\right]^{1/4}$$

Using (III.26) we deduce for any $0<\alpha<1$

$$\begin{split} & 4 \int_{\mathfrak{r}_{\vec{p}} < r} |\nabla^{\Sigma} \arctan \sigma|^2 + O(1) \ dvol_g + 2 \pi N_{\vec{p}} \le C \ \mathfrak{r}^{-2} \ \int_{r < \mathfrak{r}_{\vec{p}} < 2r} \ dvol_g \\ & + C \ \mathfrak{r}^{-1} \ \int_{r < \mathfrak{r} < 2r} \sqrt{1 + |d\vec{T}|_g^2} \ dvol_g + C \ \alpha \ \int_{\mathfrak{r} < 2r} |\nabla^{\Sigma} \arctan \sigma|^2 \ dvol_g \\ & + C \ \alpha^{-1} \ \left[\int_{\mathfrak{r} < 2r} (1 + |d\vec{T}|_g^2)^2 \ dvol_g \right]^{1/2} \ \left[\int_{\mathfrak{r} < 2r} \ dvol_g \right]^{1/2} . \end{split}$$

This gives

$$\begin{split} &(4-C\alpha) \int_{\mathfrak{r}_{\vec{p}}< r} |\nabla^{\Sigma} \arctan \sigma|^{2} dvol_{g} + 2 \pi N_{\vec{p}} \leq C \mathfrak{r}^{-2} \int_{\mathfrak{r}_{\vec{p}}< 2 r} dvol_{g} \\ &+ C \mathfrak{r}^{-1} \int_{r < \mathfrak{r}_{\vec{p}}< 2 r} \sqrt{1 + |d\vec{T}|_{g}^{2}} dvol_{g} + C \alpha^{-2} \mathfrak{r}^{2} \int_{\mathfrak{r}_{\vec{p}}< 2 r} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \\ &\leq C \mathfrak{r}^{-2} \int_{\mathfrak{r}_{\vec{p}}< 2 r} dvol_{g} + C \mathfrak{r}^{-3/2} \left[\mathfrak{r}^{2} \int_{\mathfrak{r}_{\vec{p}}< 2 r} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \right]^{1/4} \left[\int_{\mathfrak{r}_{\vec{p}}< 2 r} dvol_{g} \right]^{3/4} \\ &+ C \alpha^{-2} \mathfrak{r}^{2} \int_{\mathfrak{r}_{\vec{p}}< 2 r} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \, . \end{split}$$

We choose $C\alpha \leq 2$ and we finally get using the assumption of the lemma

$$2 \int_{\mathfrak{r}_{\vec{p}} < r} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g + 2\pi N_{\vec{p}} \le C \left[1 + \delta \frac{r^4}{\varepsilon^4} \right] \frac{1}{\mathfrak{r}^2} \int_{\mathfrak{r}_{\vec{p}} < 10 C_0^2 r} dvol_g .$$

Observe that $|\nabla^\Sigma \arctan \sigma|^2 \leq C\,\mathfrak{r}^{-2}$ and this implies

$$\int_{r < \mathfrak{r}_{\vec{p}} < 10 C_0^2 r} |\nabla^{\Sigma} \arctan \sigma|^2 \, dvol_g \le \frac{C}{\mathfrak{r}^2} \, \int_{r < \mathfrak{r}_{\vec{p}} < 10 C_0^2 r} \, dvol_g \; .$$

Hence we get

$$\forall r < C_0^{-2} \varepsilon \,\delta^{-1/4} \qquad 2 \int_{\mathfrak{r}_{\vec{p}} < r} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g + 2 \pi N_{\vec{p}} \le C \,\frac{1}{\mathfrak{r}^2} \,\int_{\mathfrak{r}_{\vec{p}} < r} \,dvol_g \,. \tag{III.60}$$

Let $1 > r > \varepsilon \, \delta^{-1/4}$ and let $\varepsilon \le s \le \varepsilon \, \delta^{-1/4}$, from (III.58)

$$\begin{split} &\lim_{\eta \to 0} \sup \left| \int_{\Sigma} \langle dh_{s,\eta}, d\beta \rangle_g \ dvol_g \right| \le C \ \int_{s < \mathfrak{r} < 2\,s} |x|^{-1} |\arctan\sigma| \ \sqrt{1 + |d\vec{T}|_g^2} \ dvol_g \\ &+ C \left[\int_{\mathfrak{r} < 2\,s} |\nabla^{\Sigma} \arctan\sigma|^2 \ dvol_g \right]^{1/2} \left[\int_{\mathfrak{r} < 2\,s} (1 + |d\vec{T}|_g^2)^2 \ dvol_g \right]^{1/4} \left[\int_{\mathfrak{r} < 2\,s} \ dvol_g \right]^{1/4} \end{split}$$
(III.61)

We have obviously $h_{r,\eta} = h_{\varepsilon,\eta} + h_{r,\varepsilon}$. Combining (III.41) with (III.61) gives then

$$\begin{split} &\lim_{\eta \to 0} \left| \int_{\Sigma} < dh_{r,\eta}, d\beta_k >_g dvol_g \right| \le \limsup_{\eta \to 0} \left| \int_{\Sigma} < dh_{\varepsilon,\eta}, d\beta_k >_g dvol_g \right| \\ &+ \left| \int_{\Sigma} < dh_{\varepsilon,\eta}, d\beta_k >_g dvol_g \right| \le C \varepsilon^4 \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}^{-2} \left(1 + |d\vec{T}|_g^2 \right)^2 dvol_g \\ &+ C \varepsilon^4 \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}^{-3} \left(1 + |d\vec{T}_k|_g^2 \right)^{3/2} dvol_g + C \int_{\varepsilon < \mathfrak{r} < 2\varepsilon} |x|^{-1} |\arctan \sigma| \sqrt{1 + |d\vec{T}|_g^2} dvol_g \\ &+ C \left[\int_{\mathfrak{r} < 2\varepsilon} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g \right]^{1/2} \left[\int_{\mathfrak{r} < 2\varepsilon} (1 + |d\vec{T}|_g^2)^2 dvol_g \right]^{1/4} \left[\int_{\mathfrak{r} < 2\varepsilon} dvol_g \right]^{1/4} \\ &+ e^{-\varepsilon^{-2}} O(||\vec{w}_{\varepsilon,r}||_{\vec{\Lambda}_k}) \end{split}$$
(III.62)

where $\vec{w}_{\varepsilon,r} := \nabla \vec{X}_{h_{r,\varepsilon}}$. We first estimate using (II.21)

$$\|\vec{w}_{\varepsilon,r}\|_{\vec{\Lambda}_k}$$

$$\leq \left[\int_{\Sigma} \left[|\nabla^2 \vec{X}_{h_{r,\varepsilon}}|^2 \circ \vec{\Lambda}_k + |\nabla \vec{X}_{h_{r,\varepsilon}}|^2 \circ \vec{\Lambda}_k |1 + |d\vec{T}_k||^2 + |\vec{X}_{h_{r,\varepsilon}}|^2 \right]^2 dvol_g \right]^{1/4} + \|\nabla \vec{X}_{h_{r,\varepsilon}}\|_{\infty} .$$
(III.63)

This gives

$$\|\vec{w}_{\varepsilon,r}\|_{\vec{\Lambda}_k} \le O(\varepsilon^{-3}) . \tag{III.64}$$

We have using the hypothesis of the lemma

$$\varepsilon^{4} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r}_{\vec{p}} < 2r\}} \mathfrak{r}_{\vec{p}}^{-2} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} = \varepsilon^{4} \int_{\varepsilon/2}^{r} \frac{ds}{s^{2}} \int_{\mathfrak{r}=2s} \frac{(1 + |d\vec{T}|_{g}^{2})^{2}}{|\nabla^{\Sigma}\mathfrak{r}|} dl$$

$$= \varepsilon^{4} \frac{1}{4r^{2}} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r}_{\vec{p}} < 2r\}} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} + 2\varepsilon^{4} \int_{\varepsilon/2}^{r} \frac{ds}{s^{3}} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r}_{\vec{p}} < 2s\}} (1 + |d\vec{T}|_{g}^{2})^{2} dvol_{g} \quad (\text{III.65})$$

$$\leq C \,\delta \, \log\left(\frac{r}{\varepsilon}\right) \sup_{\varepsilon < s < 2C_{0}^{2}r} \frac{1}{s^{2}} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_{g}$$

we have moreover

$$\varepsilon^{4} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}_{\vec{p}}^{-3} \left(1 + |d\vec{T}_{k}|_{g}^{2}\right)^{3/2} dvol_{g}$$

$$\leq \varepsilon^{4} \left[\int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}_{\vec{p}}^{-6} dvol_{g} \right]^{1/4} \left[\int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}_{\vec{p}}^{-2} \left(1 + |d\vec{T}_{k}|_{g}^{2}\right)^{2} dvol_{g} \right]^{3/4}$$
(III.66)

We bound

$$\int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}_{\vec{p}}^{-6} \, dvol_g = \int_{\varepsilon/2}^r \frac{ds}{s^6} \int_{\mathfrak{r}=2s} \frac{1}{|\nabla^{\Sigma}\mathfrak{r}|} dl = (2r)^{-6} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} dvol_g + 6 \int_{\varepsilon/2}^r \frac{ds}{s^7} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r}_{\vec{p}} < 2s\}} dvol_g \le C \sup_{\varepsilon < s < 2r} \frac{1}{s^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_g \int_{\varepsilon/2}^r \frac{ds}{s^5}.$$
(III.67)

Combining (III.65), (III.66) and (III.67) gives

$$\varepsilon^{4} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < 2r\}} \mathfrak{r}_{\vec{p}}^{-3} \left(1 + |d\vec{T}_{k}|_{g}^{2}\right)^{3/2} dvol_{g}$$

$$\leq C \,\delta^{3/4} \,\log^{3/4}\left(\frac{r}{\varepsilon}\right) \,\sup_{\varepsilon < s < 2 \,C_{0}^{2} \,r} \,\frac{1}{s^{2}} \,\int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_{g} \,. \tag{III.68}$$

We have also

$$\int_{\varepsilon < \mathfrak{r} < 2\varepsilon} |x|^{-1} |\arctan\sigma| \sqrt{1 + |d\vec{T}|_g^2} \, dvol_g$$

$$\leq C \varepsilon^{-1} \left[\int_{\varepsilon < \mathfrak{r} < 2\varepsilon} \, dvol_g \right]^{3/4} \left[\int_{\varepsilon < \mathfrak{r} < 2\varepsilon} (1 + |d\vec{T}|_g^2)^2 \, dvol_g \right]^{1/4} \leq C \, \delta^{1/4} \, \varepsilon^{-2} \, \int_{\varepsilon < \mathfrak{r} < 2\varepsilon} \, dvol_g \,.$$
(III.69)

Using moreover (III.60) for $r = 2\varepsilon$ we also get

$$\left[\int_{\mathfrak{r}<2\varepsilon} |\nabla^{\Sigma} \arctan \sigma|^2 \, dvol_g \right]^{1/2} \left[\int_{\mathfrak{r}<2\varepsilon} (1+|d\vec{T}|_g^2)^2 \, dvol_g \right]^{1/4} \left[\int_{\mathfrak{r}<2\varepsilon} \, dvol_g \right]^{1/4}$$

$$\leq C \, \delta^{1/4} \, \varepsilon^{-2} \, \int_{\mathfrak{r}<10 \, C_0^2 \, \varepsilon} \, dvol_g$$
(III.70)

Combining (III.62)....(III.70) we obtain

$$\begin{split} & \limsup_{\eta \to 0} \left| \int_{\Sigma} \langle dh_{s,\eta}, d\beta \rangle_g \ dvol_g \right| \\ & \leq C \left[\delta \ \log\left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \ \log^{3/4}\left(\frac{r}{\varepsilon}\right) \right] \ \sup_{\varepsilon < s < 10 \ C_0^2 \ r} \ \frac{1}{s^2} \ \int_{\Sigma \cap \{\mathfrak{r}_{\overline{\rho}} < s\}} \ dvol_g + C \ e^{-\varepsilon^{-2}} \ \varepsilon^{-3} \end{split}$$
(III.71)

Combining (III.53), (III.56) and (III.71) we obtain

$$\int_{\mathfrak{r}_{\vec{p}}<2\,r} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g \leq C \, \frac{1}{r^2} \, \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}}<2\,r\}} dvol_g + C \left[\delta \, \log\left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \, \log^{3/4}\left(\frac{r}{\varepsilon}\right)\right] \, \sup_{s<10 \, C_0^2 \, r} \, \frac{1}{s^2} \, \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}}$$

where we have also used the obvious inequality

$$\int_{r < \mathfrak{r}_{\vec{p}} < 2r} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g \le \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < r, p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < r, p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < p < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < 2r\}} dvol_g = \frac{1}{r^2} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < 2r\}} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{r < 2r\}} dvol_g = \frac{1}{r^2} dvol_g = \frac{1}{r^2} \int_{\Sigma \cap \{$$

This implies (III.44).

Let $0 < s_0 < r$ such that

$$\frac{1}{s_0^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s_0\}} dvol_g := \sup_{s < 10 \ C_0^2 \ r} \frac{1}{s^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_g$$
(III.73)

Since χ has been chosen such that $\chi' \leq 0$ and $\chi'(t) < -1/2$ on [5/4, 7/4] (III.53) implies for any $\eta > \varepsilon$

$$\begin{aligned} &4 \int_{\Sigma} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] |\nabla^{\Sigma} \arctan \sigma|^{2} \, dvol_{g} + \eta^{-2} \int_{\Sigma \cap \{5\eta/4 < \mathfrak{r} < 7\eta/4\}} \, dvol_{g} \\ &\leq \frac{C}{r^{2}} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2\,r\}} \, dvol_{g} + C \left[\int_{\Sigma \cap \{\eta < \mathfrak{r} < 2\eta\}} |\nabla^{\Sigma} \arctan \sigma|^{2} \, dvol_{g} \right]^{1/2} \left[\eta^{-2} \int_{\Sigma \cap \{\eta < \mathfrak{r} < 2\eta\}} \, dvol_{g} \right]^{1/2} \\ &+ C \left| \int_{\Sigma} < dh_{r,\eta}, d\beta >_{g} \, dvol_{g} \right| \end{aligned}$$

Combining (III.41), (III.65) , (III.68) and (III.73) for $\eta \geq \varepsilon$ one obtains

$$\left| \int_{\Sigma} \langle dh_{r,\eta}, d\beta \rangle_g \ dvol_g \right| \leq C \left[\delta^{3/4} \ \log^{3/4} \left(\frac{r}{\varepsilon} \right) + \delta \ \log \left(\frac{r}{\varepsilon} \right) \right] \frac{1}{s_0^2} \int_{\Sigma \cap \{ \mathfrak{r}_{\vec{p}} < s_0 \}} dvol_g + C \ e^{-\varepsilon^{-2}} \ \varepsilon^{-3}$$

While for $\eta < \varepsilon$ combining (III.57), (III.60) and (III.72) gives

$$\begin{split} \left| \int_{\Sigma} < dh_{\varepsilon,\eta}, d\beta >_g dvol_g \right| \\ \leq C \left[\frac{1}{s_0^2} \int_{\Sigma \cap \{\mathfrak{r}_{\overline{p}} < s_0\}} dvol_g \right]^{1/2} \left[\varepsilon^{-2} \int_{\eta < \mathfrak{r} < 2\varepsilon} \varepsilon^4 \left(1 + |d\vec{T}|_g^2 \right)^2 dvol_g \right]^{1/4} \left[\varepsilon^{-2} \int_{\eta < \mathfrak{r} < 2\varepsilon} dvol_g \right]^{1/4} \\ + C \varepsilon^{-2} \int_{\varepsilon < \mathfrak{r} < 2\varepsilon} \sqrt{\varepsilon^2 \left(1 + |d\vec{T}|_g^2 \right)} dvol_g \leq C \, \delta^{1/4} \, \frac{1}{s_0^2} \int_{\Sigma \cap \{\mathfrak{r}_{\overline{p}} < s_0\}} dvol_g + C \, e^{-\varepsilon^{-2}} \, \varepsilon^{-3} \end{split}$$
(III.74)

We assume first that $s_0 < r$. For $\eta := (7/5)^{-j} s_0$ for j going from -1 to $+\infty$ we obtain, using again the fact that $h_{r,\eta} = h_{r,\varepsilon} + h_{\varepsilon,\eta}$,

$$(7/5)^{2j} s_0^{-2} \int_{\Sigma \cap \{5(7/5)^{-j} s_0/4 < \mathfrak{r} < 7(7/5)^{-j} s_0/4\}} dvol_g \leq \frac{C}{r^2} \int_{\Sigma \cap \{r < \mathfrak{r}_{\vec{p}} < 2r\}} dvol_g$$

$$+ C \left[\int_{\Sigma \cap \{\mathfrak{r} < 2r\}} |\nabla^{\Sigma} \arctan \sigma|^2 dvol_g \right]^{1/2} \left[s_0^{-2} \int_{\Sigma \cap \{\varepsilon < \mathfrak{r} < s_0\}} dvol_g \right]^{1/2}$$

$$+ C \left[\delta^{1/4} + \delta^{3/4} \log^{3/4} \left(\frac{r}{\varepsilon}\right) + \delta \log \left(\frac{r}{\varepsilon}\right) \right] \frac{1}{s_0^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s_0\}} dvol_g$$

$$+ C e^{-\varepsilon^{-2}} \varepsilon^{-3}$$

$$(III.75)$$

Multiplying by $(7/5)^{-2j}$ and summing from j = -1 up to $+\infty$, using the fact that $7(7/5)^{-j-1} = 5(7/5)^{-j}$ we obtain

$$\frac{1}{s_0^2} \int_{\Sigma \cap \{\mathbf{r} < s_0\}} dvol_g \leq C \left[\sum_{j=-1}^{+\infty} \left(\frac{5}{7}\right)^{2j} \right] \frac{1}{r^2} \int_{\Sigma \cap \{r < \mathbf{r}_{\vec{p}} < 2r\}} dvol_g$$

$$+ C \left[\sum_{j=-1}^{+\infty} \left(\frac{5}{7}\right)^{2j} \right] \left[\frac{1}{r^2} \int_{\Sigma \cap \{\mathbf{r}_{\vec{p}} < 2r\}} dvol_g \right]^{1/2} \left[\frac{1}{s_0^2} \int_{\Sigma \cap \{\varepsilon < \mathbf{r} < s_0\}} dvol_g \right]^{1/2}$$

$$+ C \left[\sum_{j=-1}^{+\infty} \left(\frac{5}{7}\right)^{2j} \right] \left[\delta \log \left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4} \left(\frac{r}{\varepsilon}\right) \right]^{1/2} \frac{1}{s_0^2} \int_{\Sigma \cap \{\mathbf{r} < s_0\}} dvol_g$$

$$+ C e^{-\varepsilon^{-2}} \varepsilon^{-3}$$
(III.76)

If $r < s_0 < 10 C_0^2 r$ we trivially write

$$\frac{1}{s_0^2} \int_{\Sigma \cap \{\mathfrak{r} < s_0\}} dvol_g \leq \frac{1}{r^2} \int_{\Sigma \cap \{\mathfrak{r} < 10 C_0^2 \ r\}} dvol_g$$

We then deduce (III.45) and this concludes the proof of lemma III.5. From the previous we shall deduce the following lemma

Lemma III.6. Under the previous notations there exists a universal constant $c_* > 0$ such that

for
$$\mu_{\infty}$$
 a. e. $\vec{p} \in V_2(\mathbb{R}^4)$, $\forall s > 0$ then $c_* < \frac{\mu_{\infty}(\{\mathfrak{r}_{\vec{p}} < s\})}{s^2} \le C \ \mu_{\infty}(V_2(\mathbb{R}^4))$. (III.77)

Proof of lemma III.6. Let $\vec{p} \in V_2(\mathbb{R}^4)$ and s > 0 such that $\mu_{\infty}(\{\mathfrak{r}_{\vec{p}} < s\}) \neq 0$. Let $\delta_k \to 0$ given by

$$\delta_k^2 := (\log \varepsilon_k^{-1})^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \tag{III.78}$$

Observe that with this choice we garantee

$$\lim_{k \to +\infty} \delta_k \log \varepsilon_k^{-1} = 0 \quad \text{and} \quad \lim_{k \to +\infty} \delta_k^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = 0 \,. \tag{III.79}$$

Let E_k be the set of $\vec{p} \in V_2(\mathbb{R}^4)$ such that there exists s > 0 with

$$\delta_k \, \mu_k(\{\mathfrak{r}_{\vec{p}} < 5 \, C_0^2 \, s\}) \le \varepsilon_k^4 \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}} < s\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k}$$

Observe that for $\vec{p} \neq \vec{q}$

$$\sup_{\vec{o}\in V_2(\mathbb{R}^4)} \frac{\mathfrak{r}_{\vec{p}}(\vec{q})}{\max\{\mathfrak{r}_{\vec{p}}(\vec{o}),\mathfrak{r}_{\vec{q}}(\vec{o})\}} \le 2C_0$$

We can then use the Vitali type Covering Lemma in Quasi-metric spaces given by Lemma 2.7 in [3] and cover E_k by at most countably many balls $(B^{\mathfrak{r}}_{5C_0^2 s_i}(\vec{p}_i))_{i\in I}$ for the quasi-distance \mathfrak{r} such that the balls $B^{\mathfrak{r}}_{5C_0^2 s_i}(\vec{p}_i)$ for $i \in I$ are disjoint to each other. Hence we have

$$\mu_k(E_k) \le \sum_{i \in I} \tilde{\mu}_k(\{\mathfrak{r}_{\vec{p}_i} < 5 C_0^2 s_i\}) \le \delta_k^{-1} \sum_{i \in I} \varepsilon_k^4 \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}_i} < s_i\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \le \delta_k^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = o(1) \; .$$

Assume that there exists k_0 for any $k \ge k_0$ as large as we want

$$\bar{\Lambda}_k(\Sigma) \cap B_s^{\mathfrak{r}}(\vec{p}) \subset E_k \cap B_s^{\mathfrak{r}}(\vec{p}) ,$$

then we would have

$$\mu_k\left(\{\vec{q} ; \, \mathfrak{r}_{\vec{p}}(\vec{q}) < s\}\right) \le \mu_k(E_k) \longrightarrow 0 \;,$$

and this would imply

$$\mu_{\infty}\left(\left\{\mathfrak{r}_{\vec{p}} < s\right\}\right) = 0 ,$$

which contradicts our assumption. Hence, we can find k as large as we want such that

$$\left(\vec{\Lambda}_k(\Sigma) \cap B_s^{\mathfrak{r}}(\vec{p})\right) \bigcap \left(V_2(\mathbb{R}^4) \setminus E_k\right) \neq \emptyset$$

For such a k we consider then $\vec{p}_k \in \vec{\Lambda}_k(\Sigma) \cap B_s^{\mathfrak{r}}(\vec{p})$ and $\vec{p}_k \notin E_k$. Applying lemma III.5 and more specifically (III.45) to this point, for the choice of δ_k we are making we obtain that for any $\eta < s$ there holds

$$[c_0 - o(1)] \frac{1}{\eta^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}_k} < \eta\}} dvol_{g_k}$$
$$\leq \frac{C}{s^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}_k} < s\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \ \varepsilon_k^{-3}$$

as explained in the first part of the proof of lemma III.5, we have

$$\lim_{\eta \to 0} \frac{1}{\eta^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < \eta\}} dvol_{g_k} = \pi N_{\vec{p}_k} \ge \pi$$

Using one more time the fact that \mathfrak{r} is a quasi distance we deduce the existence of three universal positive constants $c_0 > 0$, $c_1 > 1$ and C > 0 such that, for some subsequence k' we have

$$C \mu_{k'}(\{\mathfrak{r}_{\vec{p}} < c_1 s\}) \ge s^2 \pi [c_0 - o(1)]$$
.

Passing to the limit we have proved the following implication

$$\mu_{\infty}\left(\{\mathfrak{r}_{\vec{p}} < s\}\right) \neq 0 \quad \Longrightarrow \quad \mu_{\infty}\left(\{\mathfrak{r}_{\vec{p}} \le c_1 s\}\right) \ge s^2 \ \pi c_0$$

Assume there exists G, measurable, such that

$$\mu_{\infty}(G) > 0 \quad \text{ and } \quad \forall \vec{p} \in G \quad \exists \, s_{\vec{p}} > 0 \quad \text{ s. t. } \mu_{\infty}\left(\{\mathfrak{r}_{\vec{p}} \leq c_1 \, s_{\vec{p}}\}\right) < s_{\vec{p}}^2 \, \pi \, c_0$$

We then have

$$\forall \ \vec{p} \in G \quad \exists \ s_{\vec{p}} > 0 \quad \text{s. t.} \quad \mu_{\infty}\left(\{\mathfrak{r}_{\vec{p}} < s_{\vec{p}}\}\right) = 0$$

It implies that $\mu_{\infty}(G) = 0$ which is a contradiction and this concludes the proof of lemma III.6.

IV The limit of the area density of $\vec{\Lambda}_k$.

In this section $\vec{\Lambda}_k$ denotes an admissible sequence of almost critical points of E_{ε} . We assume first that the conformal class of the induced metric $g_k := \vec{\Lambda}_k^* g_{V_2(\mathbb{R}^4)}$ is constant (the general case will be treated in a subsequent section) and, thanks to the uniformization principle, we can then compose by a diffeomorphism such that there exists a fixed constant Gauss curvature h on Σ such that

$$g_k = \vec{\Lambda}_k^* g_{V_2(\mathbb{R}^4)} = e^{2\lambda_k} h .$$

We are interested in this section with the limiting behaviour of

$$\nu_k := \frac{1}{2} |d\vec{\Lambda}_k|_h^2 \, dvol_h = e^{2\lambda_k} \, dvol_h \; .$$

Again we denote \mathbf{v}_k to be the 2-varifold associated to the immersion $\vec{\Lambda}_k$

$$\forall \ \Xi \in C^0(G_2^{Leg}(V_2(\mathbb{R}^4))) \quad \mathbf{v}_k(\Xi) := \int_{\Sigma} \Xi((\vec{\Lambda}_k)_* T_x \Sigma, \vec{\Lambda}_k(x)) \ dvol_{g_k}$$

and by μ_k the Radon measure given by the integration along the fibers that is

$$\forall A \subset V_2(\mathbb{R}^4)$$
 Borel $\mu_k(A) := \mathbf{v}_k(\pi^{-1}(A))$.

which is $\mu_k := \pi_* \mathbf{v}_k$. We clearly have for any $\Psi \in C^0(V_2(\mathbb{R}^4))$

$$<\mu_k,\Psi>=\int_{\Sigma}\Psi(\vec{\Lambda}_k)\ dvol_{g_k}=<\nu_k,\Psi\circ\vec{\Lambda}_k>,$$

or in other words

$$(\Lambda_k)_*\nu_k = \mu_k$$

Modulo extraction of a subsequence we can assume that in the sense of Radon measure convergence we have the existence of ν_{∞} , μ_{∞} and \mathbf{v}_{∞} such that

$$\nu_k \rightharpoonup \nu_\infty$$
, $\mu_k \rightharpoonup \mu_\infty$ and $\mathbf{v}_k \rightharpoonup \mathbf{v}_\infty$.

Since the $W^{1,2}$ -norm of $\vec{\Lambda}_k$ is bounded on (Σ, h) we can also assume that

$$\vec{\Lambda}_k \rightharpoonup \vec{\Lambda}_\infty$$
 weakly in $W_h^{1,2}(\Sigma, V_2(\mathbb{R}^4))$.

Thanks to the theorem of Rellich Kondrachov we have that $\vec{\Lambda}_k$ converges strongly to $\vec{\Lambda}_{\infty}$ in $L^2(\Sigma)$ and for any smooth function f on $V_2(\mathbb{R}^4)$ (which is compact) we have that $f \circ \vec{\Lambda}_k$ is also strongly converging in L^2 towards $f \circ \vec{\Lambda}_{\infty}$. Hence in particular

$$\vec{\Lambda}_k^* \alpha \ \rightharpoonup \ \vec{\Lambda}_\infty^* \alpha \quad \text{ in } \mathcal{D}'(\Sigma)$$

Since $0 = \vec{\Lambda}_k^* \alpha$ we deduce that $\vec{\Lambda}_\infty^* \alpha = 0$ and $\vec{\Lambda}_\infty$ is then *weakly Legendrian*. There is however no reason a-priori for $\vec{\Lambda}_\infty$ to be weakly conformal from (Σ, h) into $V_2(\mathbb{R}^4)$. We now prove the following lemma which is maybe the most important building-block of the present work after the almost monotonicity formula.

Lemma IV.1. There exists a universal constant $c_Q > 0$ such that, for any $x_0 \in \Sigma$, $\rho > 0$ such that for some subsequence k'

$$\limsup_{k' \to +\infty} \int_{\partial B_{\rho}(x_0)} |d\vec{\Lambda}_{k'}|_h \ dl_h < +\infty , \qquad (\text{IV.1})$$

and two positive constants $c_1, c_2 > 0$ depending only on

0

$$\Theta := \limsup_{k' \to +\infty} \sup_{r < 1} \sup_{\vec{p} \in V_2(\mathbb{R}^4)} \frac{\mu_k(\{\mathfrak{r}_{\vec{p}} < r\})}{r^2} , \qquad (\text{IV.2})$$

such that if

$$< t := \limsup_{k' \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}) \le c_1$$
(IV.3)

then either

$$\nu_{\infty}(B_{\rho}(x_0)) > c_Q , \qquad (\text{IV.4})$$

or

$$\lim_{k' \to +\infty} \sup_{B_{\rho}(x_0)} |d\vec{\Lambda}_{k'}|_h^2 \, \mathbf{1}_{\left\{x \in B_{\rho}(x_0) \; ; \; \mathfrak{r}^*_{\vec{\Lambda}_{k'}(x)} > c_2 \, t\right\}} \, dvol_h = 0 \; , \tag{IV.5}$$

where

$$\mathfrak{r}_{\vec{p}}^* := \inf \left\{ \mathfrak{r}_{\vec{p}}(\vec{q}) \quad ; \quad \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0)) \right\} \;,$$

and $\mathbf{1}_{\left\{x \; ; \; \mathfrak{r}^*_{\vec{\Lambda}_{k'}(x)} > c_2 t\right\}}$ is the characteristic function of the points $x \in B_{\rho}(x_0)$ such that $\mathfrak{r}^*_{\vec{\Lambda}_{k'}(x)} > c_2 t$. In particular, if (IV.4) does not hold, we have

$$\nu_{\infty}(B_{\rho}(x_{0})) \leq \limsup_{k \to +\infty} \nu_{k}(B_{\rho}(x_{0})) = \limsup_{k \to +\infty} \left[\left(\left. \vec{\Lambda}_{k} \right|_{B_{\rho}(x_{0})} \right)_{*} \nu_{k} \right] (V_{2}(\mathbb{R}^{4}))$$

$$\leq \limsup_{k \to +\infty} \mu_{k} \left(\left\{ \vec{p} \ ; \ \mathfrak{r}_{\vec{p}}^{*} < c_{2} t \right\} \right) \leq C t^{2} \Theta .$$
(IV.6)

Proof of Lemma IV.1. We denote by k the subsequence given by the hypothesis of the lemma. Let $\vec{\gamma}(s) = (\vec{a}(s), \vec{b}(s))$ be a C^1 Legendrian path passing through $\vec{p}_0 := (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$ at the time s = 0. We denote $\vec{\gamma}(s) := (\vec{a}(s), \vec{b}(s)) = (\vec{\varepsilon}_1 + \vec{v}(s), \vec{\varepsilon}_2 + \vec{w}(s))$. Recall

$$\varphi_{\vec{p}_0}(\vec{\gamma}(s)) = v_2(s) - w_1(s)$$

Hence we have from (III.2)

$$\frac{d\varphi_{\vec{p}_0}}{ds} = \frac{d(v_2 - w_1)}{ds} = \sum_{i=1}^4 v_i(s) \frac{dw_i}{ds} - w_i(s) \frac{dv_i}{ds} \,.$$

We deduce for any $s_0 > 0$

$$\begin{aligned} |\varphi_{\vec{p}_{0}}|(s_{0}) &\leq \int_{0}^{s_{0}} \left| \frac{d\varphi_{\vec{p}_{0}}}{ds} \right| \, ds \leq \|\vec{v}\|_{L^{\infty}([0,t],\mathbb{R}^{4})} \, \int_{0}^{s_{0}} \left| \frac{d\vec{w}}{ds} \right| \, ds + \|\vec{w}\|_{L^{\infty}([0,s_{0}],\mathbb{R}^{4})} \, \int_{0}^{s_{0}} \left| \frac{d\vec{v}}{ds} \right| \, ds \\ &\leq 2 \int_{0}^{s_{0}} \left| \frac{d\vec{v}}{ds} \right| \, ds \int_{0}^{s_{0}} \left| \frac{d\vec{w}}{ds} \right| \, ds \leq \left[\int_{0}^{s_{0}} \left| \frac{d\vec{v}}{ds} \right| \, ds \right]^{2} + \left[\int_{0}^{s_{0}} \left| \frac{d\vec{w}}{ds} \right| \, ds \right]^{2} \leq 2 \left[\int_{0}^{s_{0}} \left| \frac{d\vec{\gamma}}{ds} \right| \, ds \right]^{2} \end{aligned}$$

While we proved it for notations conveniences for $\vec{p}_0 := (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$, it holds by homogeneity of $V_2(\mathbb{R}^4)$ for any choice of \vec{p}_0 . This gives for $x(s) := \rho(\cos s, \sin s)$ and any $s \in [0, 2\pi]$ and $\vec{p}_0 := \vec{\Lambda}_{k'}(x(0))$

$$\begin{aligned} \mathfrak{r}_{\vec{p}_0}\left(\vec{\Lambda}_{k'}(x(s))\right) &= \sqrt[4]{\left|\int_0^s \frac{d(\vec{\Lambda}_{k'} \circ x)(s)}{ds} \, ds\right|^4 + 4\,\varphi_{\vec{p}_0}^2\left(\vec{\Lambda}_{k'}(x(s))\right)} \\ &\leq \sqrt[4]{17} \, \int_0^s \left|\frac{d(\vec{\Lambda}_{k'} \circ x)(s)}{ds}\right| \, ds \end{aligned} \tag{IV.7}$$

We deduce from (IV.1)

$$t := \limsup_{k \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}) < +\infty .$$
(IV.8)

Hence

$$\sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{\infty}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,) \leq \limsup_{k \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,) = t \;. \tag{IV.9}$$

We pick $x_1 \in \partial B_{\rho}(x_0)$ such that $\vec{\Lambda}_k(x_1) \to \vec{\Lambda}_{\infty}(x_1) = \vec{p}_1$ (since $\vec{\Lambda}_k$ is weakly converging to $\vec{\Lambda}_{\infty}$ in $W^{1,2}(\Sigma)$ the trace on $\partial B_{\rho}(x_0)$ is converging weakly in $H^{1/2}$ and hence almost everywhere thanks to Rellich Kondrachov).

We recall - see (III.36) - that the function

$$d(\vec{p}, \vec{q}) := \mathfrak{r}_{\vec{p}}(\vec{q}) = \left[|\vec{p} - \vec{q}|^4 + 4 \, |\vec{a} \cdot \vec{d} - \vec{b} \cdot \vec{c}|^2 \right]^{1/4} \quad \text{where} \quad \vec{p} = (\vec{a}, \vec{b}) \quad \text{and} \ \vec{q} = (\vec{c}, \vec{d})$$

satisfies the quasi-distance condition

$$\mathfrak{r}_{\vec{p}}(\vec{q}) = \mathfrak{r}_{\vec{q}}(\vec{p}) \quad \text{and} \quad d(\vec{p}_1, \vec{p}) \le C_0 \left[d(\vec{p}_1, \vec{q}) + d(\vec{p}, \vec{q}) \right]$$
(IV.10)

for some universal $C_0 > 1$. In particular, for any $\vec{p} \in V_2(\mathbb{R}^4)$ and any $\vec{q} \in \vec{\Lambda}_k(\partial B_\rho(x_0))$ there holds

$$\mathfrak{r}_{\vec{p}_1}(\vec{p}) \leq C_0 \,\mathfrak{r}_{\vec{p}_1}(\vec{q}) + C_0 \,\mathfrak{r}_{\vec{p}}(\vec{q}) \;,$$

which implies for k large enough

$$\mathfrak{r}_{\vec{p}_1}(\vec{p}) > 8 \, C_0 \ t \quad \Longrightarrow \quad \inf_{\vec{q} \in \vec{\Lambda}_k(\partial B_\rho(x_0))} \mathfrak{r}_{\vec{p}}(\vec{q}) \geq 4 \ t \ .$$

We denote by $\tilde{\mu}_k := (\vec{\Lambda}_k | B_{\rho}(x_0))_* \nu_k$ and we denote by $\tilde{\mathbf{v}}_k$ the corresponding varifold and $\tilde{\mu}_{\infty}$ is the limit of $\tilde{\mu}_k$.

We introduce for the same function χ as in the previous section which is equal to 1 in [0, 1] equal to zero on $[2, +\infty)$ and satisfying $\chi' \leq 0$ and consider

$$\zeta(r) := 1 - \chi\left(\frac{r}{8 C_0 s}\right)$$

where s > t will be chosen later. The function ζ has been constructed in such a way that, for k' large enough, $\zeta \circ \mathfrak{r}_{\vec{p}_1}$ is supported away from an open neighbourhood of $\vec{\Lambda}_k(\partial B_\rho(x_0))$. More precisely

$$\zeta(\mathfrak{r}_{\vec{p}_1}(\vec{p})) \neq 0 \implies \inf_{\vec{q} \in \vec{\Lambda}_k(\partial B_\rho(x_0))} \mathfrak{r}_{\vec{p}}(\vec{q}) \ge 2 \ t \ . \tag{IV.11}$$

We shall now be considering the infinitesimal variation for E_{ε_k} at the restriction of $\vec{\Lambda}_k$ to $B_{\rho}(x_0)$ for the Hamiltonian given by

$$h_{r,\eta} := \zeta h_{r,\eta} = \zeta(\mathfrak{r}_{\vec{p}_1}) \left[\chi(\mathfrak{r}_{\vec{p}}/r) - \chi(\mathfrak{r}_{\vec{p}}/\eta) \right] \arctan \sigma_{\vec{p}}$$

for some $\vec{p}, r > 0$ and $\eta > 0$ to be fixed later such that

$$\begin{cases} \zeta \circ \mathfrak{r}_{\vec{p}_1}(\vec{q}) \equiv 1 & \text{for} \quad \mathfrak{r}_{\vec{p}}(\vec{q}) < 2\eta \\ \\ \zeta \circ \mathfrak{r}_{\vec{p}_1}(\vec{q}) \equiv 1 & \text{for} \quad \mathfrak{r}_{\vec{p}}(\vec{q}) > r . \end{cases}$$

We denote

$$\mathfrak{r}_{\vec{p}}^* := \inf \left\{ \mathfrak{r}_{\vec{p}}(\vec{q}) \; ; \; \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0)) \right\} \; .$$

Because of (IV.11) we have $\mathfrak{r}_{\vec{p}}^* > 2t$. We consider respectively

$$\vec{X}_{\tilde{h}_{r,\eta}} := J_H \nabla^H \tilde{h}_{r,\eta} + \frac{\tilde{h}_{r,\eta}}{2} \ \vec{R} \quad \text{ and } \quad \vec{w}_{\tilde{h}_{r,\eta}} := \vec{X}_{\tilde{h}_{r,\eta}} \circ \vec{\Lambda}_k \ .$$

To simplify the notations, in the rest of the proof of the lemma we write

$$\mathfrak{r} := \mathfrak{r}_{\vec{p}}$$
, $\sigma := \sigma_{\vec{p}}$, $\mathfrak{r}_1 := \mathfrak{r}_{\vec{p}_1}$, $\sigma_1 := \sigma_{\vec{p}_1}$ and $\mathfrak{r}^* := \mathfrak{r}_{\vec{p}}^*$

We multiply (III.9) by $\zeta(\mathfrak{r}_1) [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)]$, (III.10) by $\mathfrak{r}^{-3} \arctan \sigma \zeta(\mathfrak{r}_1) [r^{-1}\chi'(\mathfrak{r}/r) - \eta^{-1}\chi'(\mathfrak{r}/\eta)]$

and (III.10) for \mathfrak{r}_1 and σ_1 instead of \mathfrak{r} and σ by \mathfrak{r}_1^{-3} $(8 C_0 s)^{-1} \zeta'(\mathfrak{r}_1) [\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)] \arctan \sigma$ this gives

$$\begin{split} &< d\tilde{h}_{r,\eta}, d\beta >_g + \zeta(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \Delta_g \log \mathfrak{r} = 4 \, \zeta(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{|(\nabla^{\Sigma} \mathfrak{r})^{\perp}|^2}{\mathfrak{r}^2} \\ &+ O(1) \, \zeta(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] + O(\mathfrak{r}) \, \zeta(\mathfrak{r}_1) \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \\ &+ \zeta(\mathfrak{r}_1) \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \mathfrak{r}^{-1} \arctan \sigma \frac{\sigma}{\sqrt{1 + \sigma^2}} \\ &+ 4^{-1} \, \mathfrak{r}^{-3} \, \zeta(\mathfrak{r}_1) \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \arctan \sigma \operatorname{div}^{\Sigma} \left(\mathfrak{r}^4 \frac{\nabla^{\Sigma} \sigma}{1 + \sigma^2} \right) \\ &+ (8 \, C_0 \, s)^{-1} \, \zeta'(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \arctan \sigma \, O(\mathfrak{r}_1) \\ &+ (8 \, C_0 \, s)^{-1} \, \zeta'(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \arctan \sigma \, \mathfrak{r}_1^{-1} \, \frac{\sigma_1}{\sqrt{1 + \sigma_1^2}} \\ &+ 4^{-1} \left(8 \, C_0 \, s)^{-1} \, \mathfrak{r}_1^{-3} \, \zeta'(\mathfrak{r}_1) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \arctan \sigma \, \operatorname{div}^{\Sigma} \left(\mathfrak{r}_1^4 \frac{\nabla^{\Sigma} \sigma_1}{1 + \sigma_1^2} \right) \end{split}$$

Integrating over $B_{\rho}(x_0)$ gives

$$\begin{split} &\int_{B_{\rho}(x_{0})} < d\tilde{h}_{r,\eta}, d\beta >_{g} dvol_{g} - \int_{B_{\rho}(x_{0})} \zeta(\mathfrak{r}_{1}) \left[\frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \frac{|\nabla^{\Sigma}\mathfrak{r}|^{2}}{\mathfrak{r}^{2}} dvol_{g} \\ &+ \int_{B_{\rho}(x_{0})} O(1) \zeta(\mathfrak{r}_{1}) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] + \zeta(\mathfrak{r}_{1}) \left[O(\mathfrak{r}/r) \chi'(\mathfrak{r}/r) - O(\mathfrak{r}/\eta) \chi'(\mathfrak{r}/\eta) \right] dvol_{g} \\ &+ 4^{-1} \int_{B_{\rho}(x_{0})} \zeta(\mathfrak{r}_{1}) \nabla^{\Sigma} \left[\mathfrak{r}^{-3} \left[r^{-1} \chi'(\mathfrak{r}/r) - \eta^{-1} \chi'(\mathfrak{r}/\eta) \right] \arctan \sigma \right] \mathfrak{r}^{4} \frac{\nabla^{\Sigma} \sigma}{1 + \sigma^{2}} dvol_{g} \\ &- \int_{B_{\rho}(x_{0})} \zeta(\mathfrak{r}_{1}) \left[\frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \frac{1}{\mathfrak{r}^{2}} \frac{\sigma \arctan \sigma}{\sqrt{1 + \sigma^{2}}} dvol_{g} \\ &- 4 \int_{B_{\rho}(x_{0})} \zeta(\mathfrak{r}_{1}) \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{|(\nabla^{\Sigma}\mathfrak{r})^{\perp}|^{2}}{\mathfrak{r}^{2}} dvol_{g} \\ &= -4^{-1} \int_{B_{\rho}(x_{0})} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \left[\frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \arctan \sigma \frac{\nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\sigma}{1 + \sigma^{2}} dvol_{g} \\ &+ \int_{B_{\rho}(x_{0})} \frac{34}{8C_{0}s} - \frac{\mathfrak{r}_{1}\zeta''(\mathfrak{r}_{1})}{(8C_{0}s)^{2}} \right] \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{\nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\sigma_{1}}{1 + \sigma^{2}} dvol_{g} \\ &+ \int_{B_{\rho}(x_{0})} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \left[O(\mathfrak{r}_{1}) + \mathfrak{r}^{-1} \nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\mathfrak{r} + \arctan \sigma \mathfrak{r}_{1}^{-1} \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \right] dvol_{g} \end{aligned}$$

We have the following bounds (using (III.35) in lemma III.4), since $\zeta' \circ \mathfrak{r}_1$ is supported in the domain $8 C_0 s < \mathfrak{r}_1 < 16 C_0 s$

$$\begin{split} &\lim_{k \to +\infty} \left| \int_{B_{\rho}(x_0)} \frac{\zeta'(\mathfrak{r}_1)}{8 C_0 s} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \left[O(\mathfrak{r}_1) + \mathfrak{r}^{-1} \nabla^{\Sigma} \mathfrak{r}_1 \cdot \nabla^{\Sigma} \mathfrak{r} \right] dvol_g \right| \\ &\leq C \, s^{-1} \left(1 + \mathfrak{r}_*^{-1} \right) \tilde{\mu}_{\infty}(\mathfrak{r}_1 < 16 \, C_0 \, s) \leq C \, s \left(1 + \mathfrak{r}_*^{-1} \right) \mu_{\infty}(\mathfrak{r}_1^{-1}[0, 1]) \,. \end{split}$$
(IV.13)

We are choosing η and r such that $\zeta'(\mathfrak{r}_1) \chi'\left(\frac{\mathfrak{r}}{r}\right) \equiv 0$ and $\zeta'(\mathfrak{r}_1) \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \equiv 0$ hence

$$\left| 4^{-1} \int_{B_{\rho}(x_0)} \frac{\zeta'(\mathfrak{r}_1)}{8 C_0 s} \left[\frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \arctan \sigma \frac{\nabla^{\Sigma} \mathfrak{r}_1 \cdot \nabla^{\Sigma} \sigma}{1 + \sigma^2} \, dvol_g \right| = 0 \,. \tag{IV.14}$$

We have also

$$\begin{split} &\lim_{k \to +\infty} \left| \int_{B_{\rho}(x_{0})} \left[\frac{3}{4} \frac{\zeta'(\mathfrak{r}_{1})}{8 C_{0} s} - \frac{\mathfrak{r}_{1} \zeta''(\mathfrak{r}_{1})}{(8 C_{0} s)^{2}} \right] \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{\nabla^{\Sigma} \mathfrak{r}_{1} \cdot \nabla^{\Sigma} \sigma_{1}}{1 + \sigma_{1}^{2}} \, dvol_{g} \right| \\ &\leq C \left(C_{0} \, s \right)^{-1} \left[\tilde{\mu}_{\infty} (8 \, C_{0} \, s < \mathfrak{r}_{1} < 16 \, C_{0} \, s) \right]^{1/2} \left[\int_{8 \, C_{0} \, s < \mathfrak{r}_{1}(\vec{q}) < 16 \, C_{0} \, s} \frac{|\nabla^{\mathcal{P}} \sigma_{1}|^{2}}{(1 + \sigma_{1}^{2})^{2}} \, d\tilde{\mathbf{v}}_{\infty}(\mathcal{P}, \vec{q}) \right]^{1/2} \end{split}$$
(IV.15)

We recall from (III.11) that away from $\mathfrak{r} = 0$

$$\left|\nabla^{H} \arctan \sigma\right| = \left|2 \frac{\rho^{2}}{\mathfrak{r}^{4}} \nabla \varphi - \frac{2}{\mathfrak{r}^{4}} \varphi \nabla \rho^{2}\right| \le \frac{C}{\mathfrak{r}} .$$
(IV.16)

.

We have

$$\left\|\nabla^{H}\left[\left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)\right] \operatorname{arctan} \sigma\right]\right\|_{L^{\infty}(\mathfrak{r}_{1} < 16 C_{0} s)} = \left\|\nabla^{H} \operatorname{arctan} \sigma\right\|_{L^{\infty}(\mathfrak{r}_{1} < 16 C_{0} s)}$$

Hence we deduce from (IV.16)

$$\left\|\nabla^{H}\left[\left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta)\right] \operatorname{arctan} \sigma\right]\right\|_{L^{\infty}(\mathfrak{r}_{1} < 16 C_{0} s)} \leq \frac{C}{\mathfrak{r}^{*}} .$$
(IV.17)

Hence

$$\begin{aligned} \zeta'(\mathfrak{r}_1(\vec{q})) &\neq 0 \implies \\ &|[\chi(\mathfrak{r}(\vec{q})/r) - \chi(\mathfrak{r}(\vec{q}/\eta)] \arctan \sigma(\vec{q}) - [\chi(\mathfrak{r}(\vec{p}_1)/r) - \chi(\mathfrak{r}(\vec{p}_1)/\eta)] \arctan \sigma(\vec{p}_1)| \\ &= |\arctan \sigma(\vec{q}) - \arctan \sigma(\vec{p}_1)| \le C \ C_0 \ \frac{s}{\mathfrak{r}^*} \ . \end{aligned}$$
(IV.18)

We consider (III.10) with \mathfrak{r}_1 and σ_1 instead of \mathfrak{r} and σ

$$\mathfrak{r}_1^3 < d\mathfrak{r}_1, d\beta >= \mathfrak{r}_1^2 \frac{\sigma_1}{\sqrt{1+\sigma_1^2}} + O(\mathfrak{r}_1^4) + 4^{-1} \operatorname{div}^{\Sigma} \left(\mathfrak{r}_1^4 \frac{\nabla^{\Sigma} \sigma_1}{1+\sigma_1^2} \right)$$

and we multiply this identity by $\zeta'(\mathfrak{r}_1) \mathfrak{r}_1^{-3}$ and integrate over $B_{\rho}(x_0)$ this gives

$$\begin{split} \int_{B_{\rho}(x_{0})} &< d(\zeta \circ \mathfrak{r}_{1}), d\beta > \ dvol_{g} = \int_{B_{\rho}(x_{0})} \zeta'(\mathfrak{r}_{1}) \left[\mathfrak{r}_{1}^{-1} \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} + O(\mathfrak{r}_{1}) \right] \ dvol_{g} \\ &+ \frac{1}{4} \int_{B_{\rho}(x_{0})} \zeta'(\mathfrak{r}_{1}) \ \mathfrak{r}_{1}^{-3} \operatorname{div}^{\Sigma} \left(\mathfrak{r}_{1}^{4} \frac{\nabla^{\Sigma} \sigma_{1}}{1 + \sigma_{1}^{2}} \right) \ dvol_{g} \end{split}$$
(IV.19)

Since $\zeta \circ \mathfrak{r}_1$ is a smooth function supported away from $\vec{\Lambda}_k(\partial B_\rho(x_0))$ lemma II.4 is implying

$$\lim_{k \to 0} \int_{B_{\rho}(x_0)} \langle d(\zeta \circ \mathfrak{r}_1), d\beta \rangle \ dvol_g = 0$$
(IV.20)

Hence, after integrating by parts the last integral in (IV.19) one gets

$$\begin{split} & \limsup_{k \to 0} \left| \int_{B_{\rho}(x_{0})} \zeta'(\mathfrak{r}_{1}) \, \mathfrak{r}_{1}^{-1} \, \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \, dvol_{g} \right| \\ & \leq \left[\tilde{\mu}_{\infty} (8 \, C_{0} \, s < \mathfrak{r}_{1} < 16 \, C_{0} \, s) \right]^{1/2} \, \left[\int_{8 \, C_{0} \, s < \mathfrak{r}_{1}(\vec{q}) < 16 \, C_{0} \, s} \frac{|\nabla^{\mathcal{P}} \sigma_{1}|^{2}}{(1 + \sigma_{1}^{2})^{2}} \, d\tilde{\mathbf{v}}_{\infty}(\mathcal{P}, \vec{q}) \right]^{1/2} \end{split} \tag{IV.21}$$

Combining (IV.18) and (IV.21) gives

$$\begin{split} & \limsup_{k \to 0} \left| \int_{B_{\rho}(x_{0})} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \arctan \sigma \, \mathfrak{r}_{1}^{-1} \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \, dvol_{g} \right| \\ & \leq \frac{\left| \left[\chi(\mathfrak{r}(\vec{p}_{1})/r) - \chi(\mathfrak{r}(\vec{p}_{1})/\eta) \right] \, \arctan \sigma(\vec{p}_{1}) \right|}{8C_{0}s} \, \lim_{k \to 0} \left| \int_{B_{\rho}(x_{0})} \zeta'(\mathfrak{r}_{1}) \, \mathfrak{r}_{1}^{-1} \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \, dvol_{g} \right| \\ & + C \, C_{0} \, s \, (\mathfrak{r}^{*})^{-1} \, \limsup_{k \to 0} \left| \int_{B_{\rho}(x_{0})} \left| \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \right| \, \mathfrak{r}_{1}^{-1} \, \left| \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \right| \, dvol_{g} \right| \end{split}$$
(IV.22)
$$& \leq C \, (C_{0} \, s)^{-1} \, \left[\mu_{\infty}(C_{0} \, s < \mathfrak{r}_{1} < 2 \, C_{0} \, s) \right]^{1/2} \, \left[\int_{C_{0} \, s < \mathfrak{r}_{1}(\vec{q}) < 2 \, C_{0} \, s} \frac{|\nabla^{\mathcal{P}} \sigma_{1}|^{2}}{(1 + \sigma_{1}^{2})^{2}} \, d\tilde{\mathbf{v}}_{\infty}(\mathcal{P}, \vec{q}) \right]^{1/2} \\ & + C \, (C_{0} \, s)^{-1} \, (\mathfrak{r}^{*})^{-1} \, \tilde{\mu}_{\infty}(C_{0} \, s < \mathfrak{r}_{1} < 2 \, C_{0} \, s) \leq C \, C_{0} \, s \, (\mathfrak{r}^{*})^{-1} \, \mu_{\infty}(\{\mathfrak{r}_{1} < 1\}) \end{split}$$

Let $N \in \mathbb{N}$ such that $\mathfrak{r}^* > 2^{N+1}t$, because of lemma III.4 there exists $j_k \in \{1 \cdots N\}$ such that

$$\int_{C_0 2^{j_k} t < \mathfrak{r}_1(\vec{q}) < 2^{j_k+1} C_0 t} \frac{|\nabla^{\mathcal{P}} \sigma_1|^2}{(1+\sigma_1^2)^2} d\tilde{\mathbf{v}}_{\infty}(\mathcal{P}, \vec{q}) \le \frac{C}{N} \mu_{\infty}(\{\mathfrak{r}_1 < 1\})$$
(IV.23)

We take $s := 2^{j_k} t$ and we shall fix N later on. Combining (IV.13)...(IV.21) we obtain

$$\begin{split} &\lim_{k \to +\infty} \left| -4^{-1} \int_{B_{\rho}(x_{0})} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \left[\frac{\mathfrak{r}}{r} \chi'\left(\frac{\mathfrak{r}}{r}\right) - \frac{\mathfrak{r}}{\eta} \chi'\left(\frac{\mathfrak{r}}{\eta}\right) \right] \arctan \sigma \frac{\nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\sigma}{1 + \sigma^{2}} \, dvol_{g} \\ &+ \int_{B_{\rho}(x_{0})} \left[\frac{3}{4} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} - \frac{\mathfrak{r}_{1} \zeta''(\mathfrak{r}_{1})}{(8C_{0}s)^{2}} \right] \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \frac{\nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\sigma_{1}}{1 + \sigma_{1}^{2}} \, dvol_{g} \\ &+ \int_{B_{\rho}(x_{0})} \frac{\zeta'(\mathfrak{r}_{1})}{8C_{0}s} \left[\chi(\mathfrak{r}/r) - \chi(\mathfrak{r}/\eta) \right] \left[O(\mathfrak{r}_{1}) + \nabla^{\Sigma}\mathfrak{r}_{1} \cdot \nabla^{\Sigma}\mathfrak{r} + \arctan \sigma \, \mathfrak{r}_{1}^{-1} \frac{\sigma_{1}}{\sqrt{1 + \sigma_{1}^{2}}} \right] \, dvol_{g} \\ &\leq C \, 2^{N} t \, \mu_{\infty}(\{\mathfrak{r}_{1} < 1\}) + C \, C_{0} \, 2^{N} t \, (\mathfrak{r}^{*})^{-1} \, \mu_{\infty}(\{\mathfrak{r}_{1} < 1\}) + C \, \frac{1}{\sqrt{N}} \, \mu_{\infty}(\{\mathfrak{r}_{1} < 1\}) \, . \end{split}$$

Let $\delta_k \to 0$ given by

$$\delta_k^2 := (\log \varepsilon_k^{-1})^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \tag{IV.25}$$

Observe that with this choice we garantee

$$\lim_{k \to +\infty} \delta_k \log \varepsilon_k^{-1} = 0 \quad \text{and} \quad \lim_{k \to +\infty} \delta_k^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = 0 \,. \tag{IV.26}$$

Let F_k be the set of $\vec{p} \in V_2(\mathbb{R}^4)$ with $\mathfrak{r}_{\vec{p}_1}(\vec{p}) > N \, 2^{N+1} t$ such that there exists s > 0 with

$$\delta_k \, \tilde{\mu}_k(\{\mathfrak{r}_{\vec{p}} < 5 \, C_0^2 \, s\}) \le \varepsilon_k^4 \, \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}} < s\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k}$$

Observe that for $\vec{p} \neq \vec{q}$

$$\sup_{\vec{o} \in V_2(\mathbb{R}^4)} \frac{\mathfrak{r}_{\vec{p}}(\vec{q})}{\max\{\mathfrak{r}_{\vec{p}}(\vec{o}), \mathfrak{r}_{\vec{q}}(\vec{o})\}} \le 2 C_0 .$$

We can then use the Vitali type Covering Lemma in Quasi-metric spaces given by Lemma 2.7 in [3] and cover F_k by at most countably many balls $(B_{5C_0^2 s_i}^{\mathfrak{r}}(\vec{p_i}))_{i \in I}$ for the quasi-distance \mathfrak{r} such that the balls $B_{5C_0^2 s_i}^{\mathfrak{r}}(\vec{p_i})$ for $i \in I$ are disjoint to each other. Hence we have

$$\begin{split} \tilde{\mu}_k(F_k) &\leq \sum_{i \in I} \tilde{\mu}_k(\{\mathfrak{r}_{\vec{p}_i} < 5 C_0^2 \, s_i\}) \leq \delta_k^{-1} \, \sum_{i \in I} \varepsilon_k^4 \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}_i} < s_i\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \\ &\leq \delta_k^{-1} \, \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = o(1) \; . \end{split}$$

Assume that for every k and every \vec{p} satisfying $\mathfrak{r}_1(\vec{p}) > 2^N N t$ either $\vec{p} \in F_k$ or $\vec{p} \in V_2(\mathbb{R}^4) \setminus \vec{\Lambda}_k(B_\rho(x_0))$ then

$$\tilde{\mu}_k\left(\{\vec{p} ; \mathfrak{r}_1(\vec{p}) > 2^N N t\}\right) = \tilde{\mu}_k(F_k) \longrightarrow 0 ,$$

from which we deduce

$$\tilde{\mu}_{\infty}(\{\vec{p}\ ;\ \mathfrak{r}_{1}(\vec{p}) > 2^{N}t\}) = 0$$
. (IV.27)

Alternatively, assume $\vec{\Lambda}_k(B_{\rho}(x_0)) \cap \{\vec{p} ; \mathfrak{r}_1(\vec{p}) > 2^N N t\} \cap (V_2(\mathbb{R}^4) \setminus F_k) \neq \emptyset$, for \vec{p}_k in this intersection we can apply lemma III.5 for r = t and any s < t which gives

$$\begin{bmatrix} c_0 - \left[\delta \log\left(\frac{t}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4}\left(\frac{t}{\varepsilon}\right)\right]^{1/2} \end{bmatrix} \frac{1}{s^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}} < s\}} dvol_{g_k}$$

$$\leq \frac{C}{t^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}} < t\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \varepsilon_k^{-3} .$$
(IV.28)

Since $\vec{p}_k \in \vec{\Lambda}_k(B_t(x_0)) \setminus \vec{\Lambda}_k(\partial B_t(x_0))$ we have using (III.52)

$$\lim_{s \to 0} \frac{1}{s^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}_k} < s\}} dvol_{g_k} = \pi N_{\vec{p}_k} \ge \pi \; .$$

This gives

$$\pi \left[c_0 - \left[\delta \log\left(\frac{t}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4}\left(\frac{t}{\varepsilon}\right) \right]^{1/2} \right]$$

$$\leq \frac{C}{t^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}_k} < t\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \ \varepsilon_k^{-3} \ .$$
(IV.29)

We deduce

$$\pi c_0 \le \liminf_{k \to +\infty} \frac{C}{t^2} \int_{B_{\rho}(x_0) \cap \{ \mathfrak{r}_{\vec{p}_k} < t \}} dvol_{g_k} .$$
(IV.30)

Assume that for k large enough there exists $\vec{p}_k \in \vec{\Lambda}_k(B_\rho(x_0)) \cap \{\vec{p} ; \mathfrak{r}_1(\vec{p}) > 2^N t\} \cap (V_2(\mathbb{R}^4) \setminus F_k)$. If $r^*_{\vec{p}_k} > 1$ then, since in the domain $0 < r_{\vec{p}_k} < 1/2$ we have $\zeta \circ \mathfrak{r}_1 \equiv 1$ we can apply (III.45) as such and we get

$$\pi \left[c_0 - \left[\delta \log\left(\frac{r}{\varepsilon}\right) + \delta^{1/4} + \delta^{3/4} \log^{3/4}\left(\frac{r}{\varepsilon}\right) \right]^{1/2} \right]$$

$$\leq C_* \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}_k} < 1/2\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \ \varepsilon_k^{-3} \ .$$
(IV.31)

and passing to the limit we get

$$\nu_{\infty}(B_{\rho}(x_0)) \ge \pi \, \frac{c_0}{C_*} \tag{IV.32}$$

and by choosing $0 < c_Q < \frac{c_0}{C_*}$ we have proved the lemma in this case.

We now fix $N \in \mathbb{N}^*$ such that

$$C \frac{1}{\sqrt{N}} \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) \le \frac{\pi}{4} c_0 ,$$
 (IV.33)

and assume now $0 < 2^N t < r^*_{\vec{p}_k} < 1$. We extract a subsequence that we keep denoting \vec{p}_k such that $\vec{p}_k \to \vec{p}_\infty$. We apply (IV.24) for $\vec{p} = \vec{p}_\infty$ that we combine with the the proof of lemma III.5 but with an error term coming from $\partial \vec{\Lambda}_k(B_t(x_0))$ in the r.h.s. of (IV.12) which is controlled by (IV.24) we get

$$\begin{split} &\lim_{k \to +\infty} \sup_{\tau^2} \frac{C}{f^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}_{\infty}} < 2t\}} dvol_{g_k} - C \, 2^N t \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) - C \, C_0 \, 2^N t \, (\mathfrak{r}_{\vec{p}_{\infty}}^*)^{-1} \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) \\ &- C \, \frac{1}{\sqrt{N}} \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) \leq C_* \, \mu_{\infty}(\mathfrak{r} < 1) \; . \end{split}$$
(IV.34)

From (IV.30) we have

$$\liminf_{k \to +\infty} \frac{C}{t^2} \int_{B_{\rho}(x_0) \cap \{\mathfrak{r}_{\vec{p}_{\infty}} < 2t\}} dvol_{g_k} \ge \pi c_0 \tag{IV.35}$$

Combining (IV.34) and (IV.35) is giving finally

$$\pi c_0 - C \, 2^N t \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) - C \, C_0 \, 2^N t \, (\mathfrak{r}_{\vec{p}_{\infty}}^*)^{-1} \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) - C \, \frac{1}{\sqrt{N}} \, \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) \leq C_* \, \tilde{\mu}_{\infty}(V_2(\mathbb{R}^4) = C_* \nu_{\infty}(B_{\rho}(x_0)) \, .$$
(IV.36)

We restrict to 0 < t such that

$$C 2^N t \ \mu_{\infty}(\{\mathfrak{r}_1 < 1\}) \le \frac{\pi}{4} c_0$$
, (IV.37)

and we choose $\mathfrak{r}^*_{\overrightarrow{p}_\infty}$ large enough compare to t so that

$$C C_0 2^N t (\mathfrak{r}^*_{\vec{p}_\infty})^{-1} \mu_\infty(\{\mathfrak{r}_1 < 1\}) \le \frac{\pi}{4} c_0 .$$
 (IV.38)

Combining (IV.33) assuming $t := \limsup_{k' \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q})$ satisfies (IV.37), assume one can find a subsequence of points $\vec{p}_k \in \vec{\Lambda}_k(B_{\rho}(x_0)) \cap \{\vec{p}; \mathfrak{r}_1(\vec{p}) > 2^N t\} \cap (V_2(\mathbb{R}^4) \setminus F_k)$ converging to a point \vec{p}_{∞} such that (IV.38) is satisfied, then we obtain

$$\frac{\pi}{4} c_0 \le C_* \nu_\infty(B_\rho(x_0)) . \tag{IV.39}$$

This concludes the proof of lemma IV.1.

Lemma IV.2. The limiting measure ν_{∞} decomposes as follows

$$\nu_{\infty} = f \, dvol_h + \sum_{i=1}^{Q} c_i \, \delta_{x_i} , \qquad (\text{IV.40})$$

where $f \in L^1(\Sigma, \mathbb{R}_+)$ and $c_i \geq c_Q$ (where c_Q is the constant given in lemma IV.1). Moreover the weak $W^{1,2}$ -limit $\vec{\Lambda}_{\infty}$ of the sequence $\vec{\Lambda}_k$ is in $C^0_{loc}(\Sigma \setminus \{x_1 \cdots x_Q\}, V_2(\mathbb{R}^4))$). Finally for any open set U in Σ with $x_i \notin \overline{U}$ for any $i = 1 \cdots Q$ there holds

$$\lim_{k \to +\infty} (\vec{\Lambda}_k)_* (\nu_k \, \boldsymbol{\sqcup} \, U) = (\vec{\Lambda}_\infty)_* (\nu_\infty \, \boldsymbol{\sqcup} \, U) \ . \tag{IV.41}$$

Proof of lemma IV.2 Let $x \in \Sigma$ and let r > 0. Using Fubini combined with the mean value argument, there exists $\rho \in (r/2, r)$ and a subsequence that we keep denoting k such that

$$\int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_k|_h^2 \, dl_h \leq \frac{C}{r} \, \int_{B_r(x)} |d\vec{\Lambda}_k|_h^2 \, dvol_h \leq 2\frac{C}{r} \, \mu_k(V_2(\mathbb{R}^4)) \, .$$

and simultaneously

$$\int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_{\infty}|_{h}^{2} dl_{h} \leq \frac{C}{r} \int_{B_{r}(x)} |d\vec{\Lambda}_{\infty}|_{h}^{2} dvol_{h} \leq 2\frac{C}{r} \mu_{\infty}(V_{2}(\mathbb{R}^{4})) .$$

Hence the restriction of $\vec{\Lambda}_k$ to $\partial B_\rho(x)$ is sequentially weakly pre-compact in $W^{1,2}(\partial B_\rho(x), V_2(\mathbb{R}^4))$ as well as in $C^{0,1/2}(\partial B_\rho(x), V_2(\mathbb{R}^4))$. By Arzela-Ascoli we deduce that $\vec{\Lambda}_k$ is strongly pre-compact in $C^{0,\alpha}(\partial B_\rho(x), V_2(\mathbb{R}^4))$ for any $\alpha < 1/2$. Since $\vec{\Lambda}_k$ converges weakly towards $\vec{\Lambda}_\infty$ in $W^{1,2}(B_\rho(x), V_2(\mathbb{R}^4))$, by continuity of the trace operation into $W^{1/2,2}(\partial B_\rho, V_2(\mathbb{R}^4))$ we deduce

$$\forall \alpha < 1/2 \qquad \vec{\Lambda}_k \longrightarrow \vec{\Lambda}_\infty \quad \text{strongly in } C^{0,\alpha}(\partial B_\rho(x), V_2(\mathbb{R}^4)) \ . \tag{IV.42}$$

and moreover

$$\int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_{\infty}|_{h}^{2} dl_{h} \leq \liminf_{k \to +\infty} \int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_{k}|_{h}^{2} dl_{h} \leq \frac{2}{r} \liminf_{k \to +\infty} \int_{B_{r}(x)} |d\vec{\Lambda}_{k}|_{h}^{2} dvol_{h} \\
\leq \frac{4}{r} \nu_{\infty}(\overline{B_{r}(x)}) .$$
(IV.43)

Cauchy Schwartz inequality gives

$$\int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_k|_h \ dl_h \le C \rho^{1/2} \left[\int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_k|_h^2 \ dl_h \right]^{1/2} \le C \sqrt{\frac{\rho}{r}} \sqrt{\int_{B_r(x)} |d\vec{\Lambda}_k|_h^2 \ dvol_h} \ . \tag{IV.44}$$

Hence

$$\limsup_{k \to +\infty} \int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_k|_h \ dl_h < +\infty \ .$$

Let

$$t := \lim_{k \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{k'}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,) = \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_{\infty}(\partial B_{\rho}(x_0))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,)$$

Applying lemma IV.1 we deduce that

either
$$\nu_{\infty}(B_{\rho}(x)) > c_Q$$
 or $\nu_{\infty}(B_{\rho}(x)) \le C t^2 V_2(\mathbb{R}^4)$. (IV.45)

Assuming

$$\nu_{\infty}(\{x\}) > 0$$

the second alternative cannot happen because, taking r small enough we can make t as small as we want. Indeed

$$t \le C \int_{\partial B_{\rho}(x)} |d\vec{\Lambda}_{\infty}|_{h} dl_{h} \le C \sqrt{\frac{\rho}{r}} \sqrt{\int_{B_{r}(x)} |d\vec{\Lambda}_{\infty}|_{h}^{2} dvol_{h}} \longrightarrow 0 \quad \text{as } r \to 0 .$$
 (IV.46)

Hence we have proved the following quantization phenomenon

$$\nu_{\infty}(\{x\}) > 0 \implies \nu_{\infty}(\{x\}) > c_Q , \qquad (\text{IV.47})$$

and this can only happen at at most finitely many points. We denote by $x_1 \cdots x_Q$ the set of atoms (when it is not empty). Let K be a compact set containing no atom for ν_{∞} we claim that there exists C independent of K such that

$$\nu_{\infty}(K) \le C \int_{K} |d\vec{\Lambda}_{\infty}|_{h}^{2} \, dvol_{h} \, . \tag{IV.48}$$

This claim is obviously implying the decomposition (IV.40) of the measure ν_{∞} .

Proof of (IV.48). Since there are at most finitely many atoms, there exists $r_0 > 0$ such that $K \cap B_{r_0}(x_i) = \emptyset$ for any $i = 1 \cdots Q$. Let U be the open set given by

$$U := \Sigma \setminus \bigcup_{i=1}^{Q} \overline{B_{r_0}(x_i)}$$

and we choose V open, arbitrary, such that $K \subset V$ and $\overline{V} \subset U$. Let r > 0 such that $B_{5r}(y) \subset V$ for any $y \in K$ and

$$\sup_{y \in K} \nu_{\infty}(B_{5r}(y)) < c_Q$$

We take a maximal finite family of balls $(B_r(y_j))_{j \in J}$ for $y_j \in K$ and the distance between two centers $y_j \neq y'_j$ is at least r. Since it is a maximal family we have

$$K \subset \bigcup_{j \in J} B_r(y_j)$$
 and $\sum_{j \in J} \mathbf{1}_{B_{5r}(y_j)} \leq C$

where C > 0 is universal and $\mathbf{1}_{B_{5r}(y_j)}$ denotes the characteristic function of the ball $B_{5r}(y_j)$. For each $j \in J$ we chose $\rho_j \in [2r, 4r]$ and we extract a subsequence that we keep denoting k (recall that the cardinal of J is finite) such that $\forall j \in J$ there holds

$$\int_{\partial B_{\rho_j}(y_j)} |d\vec{\Lambda}_k|_h^2 \ dl_h \le \frac{C}{r} \ \int_{B_{4r}(y_j)} |d\vec{\Lambda}_k|_h^2 \ dvol_h \le 2\frac{C}{r} \ \mu_k(V_2(\mathbb{R}^4)) \ ,$$

and simultaneously

$$\int_{\partial B_{\rho_j}(y_j)} |d\vec{\Lambda}_{\infty}|_h^2 \ dl_h \le \frac{C}{r} \ \int_{B_{4r}(y_j)} |d\vec{\Lambda}_{\infty}|_h^2 \ dvol_h \le 2\frac{C}{r} \ \mu_{\infty}(V_2(\mathbb{R}^4))$$

Hence, as for the proof of (IV.42), we have

$$\vec{\Lambda}_k \longrightarrow \vec{\Lambda}_\infty$$
 in $C^0\left(\bigcup_{j \in J} \partial B_{\rho_j}(y_j), V_2(\mathbb{R}^4)\right)$

and because of (IV.46) we have

$$\begin{split} t_j &:= \lim_{k \to +\infty} \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_k(\partial B_{\rho_j}(y_j))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,) = \sup_{\{\vec{p}, \vec{q} \in \vec{\Lambda}_\infty(\partial B_{\rho_j}y_j))\}} \mathfrak{r}_{\vec{p}}(\vec{q}\,) \\ &\leq C \, \sqrt{\int_{B_{4r}(y_j)} |d\vec{\Lambda}_\infty|_h^2 \, dvol_h} \end{split}$$

Because of lemma IV.1, since for every $j \in J$ we have $\nu_{\infty}(B_{\rho_j}(y_j)) < c_Q$, using more precisely (IV.6) we deduce

$$\begin{split} \nu_{\infty}(K) &\leq \sum_{j \in J} \nu_{\infty}(B_{\rho_{j}}(y_{j})) \leq C \sum_{j \in J} t_{j}^{2} \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \leq C \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \sum_{j \in J} \int_{B_{4r}(y_{j})} |d\vec{\Lambda}_{\infty}|_{h}^{2} \ dvol_{h} \\ &\leq C \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \int_{\Sigma} \sum_{j \in J} \mathbf{1}_{B_{4r}(y_{j})} \ |d\vec{\Lambda}_{\infty}|_{h}^{2} \ dvol_{h} \leq \ C \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \int_{V} |d\vec{\Lambda}_{\infty}|_{h}^{2} \ dvol_{h} \end{split}$$

Since this holds for any open set V containing K we deduce (IV.48) and the claim is proved. We deduce in particular that

$$\int_{K} |d\vec{\Lambda}_{\infty}|_{h}^{2} \, dvol_{h} = 0 \quad \Longrightarrow \quad \nu_{\infty}(K) = 0 \quad .$$

and we deduce that away from the atoms the measure ν_{∞} is absolutely continuous with respect to $dvol_h$. Again from lemma IV.1 we have

$$\lim_{k \to +\infty} \sup_{B_{\rho_j}(y_j) \cap \mathfrak{r}^j_{\bar{\Lambda}_k(x)} > c_2 t_j} |d\vec{\Lambda}_k|_h^2 \, dvol_h = 0 , \qquad (\text{IV.49})$$

where

$$\mathfrak{r}^{j}_{\vec{p}} := \inf \left\{ \mathfrak{r}_{\vec{p}}(\vec{q}) \quad ; \quad \vec{q} \in \vec{\Lambda}_{k}(\partial B_{\rho_{j}}(y_{j})) \right\} \; .$$

Let $\chi_j(\vec{p}) := (\mathfrak{r}_{\vec{p}}^j - c_2 t_j)^+$ we have that the restriction of $\chi_j \circ \vec{\Lambda}_k$ to $B_{\rho_j}(y_j)$ is strongly converging to zero in $W^{1,2}$ and we deduce that for all $j \in J$

$$\forall \vec{p} \in \vec{\Lambda}_{\infty}(B_{\rho_j}(y_j)) \quad \text{and} \quad \forall \vec{q} \in \vec{\Lambda}_{\infty}(\partial B_{\rho_j}(y_j)) \quad \mathfrak{r}_{\vec{p}}(\vec{q}) < C\left(c_2 + 1\right) t_j$$

where C denotes the constant in the triangular inequality of the quasi distance \mathfrak{r} . Thus for x and y in K such that |x - y| < r we have that x and y belong to two balls $B_{2r}(y_j)$ which intersect each-other and using again the fact that \mathfrak{r} is a quasi-distance we deduce that

$$\mathfrak{r}_{\vec{\Lambda}_{\infty}(x)}(\vec{\Lambda}_{\infty}(y)) < C \max_{j \in J} \{t_j\} \leq C \max_{j \in J} \sqrt{\int_{B_{4r}(y_j)} |d\vec{\Lambda}_{\infty}|_h^2 \, dvol_h}$$

Let now $\varepsilon > 0$, we choose r > 0 such that

$$\max_{y \in K} \sqrt{\int_{B_{4r}(y)} |d\vec{\Lambda}_{\infty}|_h^2 \, dvol_h} < \varepsilon$$

we have that

$$\forall \, x,y \in K \quad |x-y| < r \implies \mathfrak{r}_{\vec{\Lambda}_{\infty}(x)}(\vec{\Lambda}_{\infty}(y)) < \varepsilon \; .$$

This implies that $\vec{\Lambda}_{\infty}$ is continuous on K.

Finally we have

$$\begin{split} &\int_{K} |\vec{\Lambda}_{k} - \vec{\Lambda}_{\infty}| \ d\nu_{k} \leq \sum_{j \in J} \int_{B_{\rho_{j}}(y_{j})} |\vec{\Lambda}_{k} - \vec{\Lambda}_{\infty}| \ d\nu_{k} \leq C \ \sum_{j \in J} t_{j} \ \int_{B_{\rho_{j}}(y_{j}) \cap \mathfrak{r}_{\vec{\Lambda}_{k}(x)}^{j} \leq c_{2} t_{j}} \ |d\vec{\Lambda}_{k}|_{h}^{2} \ dvol_{h} \\ &+ \sum_{j \in J} \int_{B_{\rho_{j}}(y_{j}) \cap \mathfrak{r}_{\vec{\Lambda}_{k}(x)}^{j} > c_{2} t_{j}} \ |d\vec{\Lambda}_{k}|_{h}^{2} \ dvol_{h} \end{split}$$

From (IV.49) we deduce

$$\begin{split} &\lim_{k \to +\infty} \sup_{K} |\vec{\Lambda}_{k} - \vec{\Lambda}_{\infty}| \ d\nu_{k} \leq C \ \max_{j \in J} t_{j} \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \\ &\leq C \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \max_{j \in J} \sqrt{\int_{B_{4r}(y_{j})} |d\vec{\Lambda}_{\infty}|_{h}^{2} \ dvol_{h}} \leq C \ \mu_{\infty}(V_{2}(\mathbb{R}^{4})) \ \max_{y \in K} \sqrt{\int_{B_{4r}(y)} |d\vec{\Lambda}_{\infty}|_{h}^{2} \ dvol_{h}} \end{split}$$

By making r go to zero implies

$$\lim_{k \to +\infty} \int_{K} |\vec{\Lambda}_{k} - \vec{\Lambda}_{\infty}| \ d\nu_{k} = 0 , \qquad (\text{IV.50})$$

For any $\Psi \in C^0(V_2(\mathbb{R}^4))$, because of (IV.50) and since $\vec{\Lambda}_{\infty}$ is continuous on K, the weak convergence in Radon measure is implying

$$\lim_{k \to +\infty} \int_{K} \Psi(\vec{\Lambda}_{k}) \ d\nu_{k} = \lim_{k \to +\infty} \int_{K} \Psi(\vec{\Lambda}_{\infty}) \ d\nu_{k} = \int_{K} \Psi(\vec{\Lambda}_{\infty}) \ d\nu_{\infty}$$
(IV.51)

This holds for any compact not containing the atoms of ν_∞ and this is closing the proof of lemma IV.2 . \Box

Lemma IV.3. There exists $N \in L^{\infty}(\Sigma, \mathbb{N}^*)$ such that

$$\nu_{\infty} = N \frac{|d\vec{\Lambda}_{\infty} \wedge d\vec{\Lambda}_{\infty}|_{h}}{2} \, dvol_{h} + \sum_{i=1}^{Q} c_{i} \, \delta_{x_{i}} \; .$$

Proof of lemma IV.3. We take local conformal coordinates (x_1, x_2) for h around a point which is 0 = (0, 0) in these coordinates and we can write in these coordinates

$$h = e^{2\lambda} \left[dx_1^2 + dx_2^2 \right]$$
.

We can assume again that $\vec{\Lambda}_{\infty}(0) = (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$. We assume that 0 is not one of the atoms of ν_{∞} , we assume moreover that 0 is a Lebesgue point for f in such a way that

$$\lim_{r \to 0} \frac{\nu_{\infty}(B_r(0))}{\pi r^2} = f(0) e^{2\lambda(0)} .$$
 (IV.52)

This is satisfied $\mathcal{H}^2 - a.e.$ on Σ . The goal in this lemma is to prove that there exists a non zero integer N such that

$$f(0) = N \left| \partial_{x_1} \vec{\Lambda}_{\infty}(0) \wedge \partial_{x_2} \vec{\Lambda}_{\infty}(0) \right| e^{-2\lambda(0)} .$$
 (IV.53)

where we recall the notation $\nu_{\infty} = f \, dvol_h + \sum_{i=1}^Q c_i \, \delta_{x_i}$. We can also assume that

$$\lim_{k \to +\infty} \vec{\Lambda}_k(0) = \vec{\Lambda}_\infty(0) . \tag{IV.54}$$

We can assume without loss of generality that 0 is a Lebesgue point for $d\vec{\Lambda}_{\infty}$ and that $\vec{\Lambda}_{\infty}$ is approximate differentiable at 0. This holds for \mathcal{H}^2 -a.e. points in Σ as well since $\vec{\Lambda}_{\infty} \in W^{1,2}$ (see [7] section 6.1). Hence we have

$$\lim_{r \to 0} \int_{B_r(0)} \frac{|d\vec{\Lambda}_{\infty}(x) - d\vec{\Lambda}_{\infty}(0)|^2}{r^2} \, dx^2 = 0 \; . \tag{IV.55}$$

as well as

$$\lim_{r \to 0} \int_{B_r(0)} \frac{\left| \vec{\Lambda}_{\infty}(x) - \vec{\Lambda}_{\infty}(0) - d\vec{\Lambda}_{\infty}(0) \cdot x \right|}{r^3} \, dx^2 = 0 \,, \tag{IV.56}$$

and

$$\lim_{r \to 0} \int_{B_r(0)} \frac{\left| \vec{\Lambda}_{\infty}(x) - \vec{\Lambda}_{\infty}(0) - d\vec{\Lambda}_{\infty}(0) \cdot x \right|^2}{r^4} \, dx^2 = 0 \; . \tag{IV.57}$$

Since 0 is chosen to be a point of approximate differentiability of $\vec{\Lambda}_{\infty}$ Hence in particular for any t > 0 there exists $r \in [t/2, t]$ and $\theta_r \in [0, 2\pi]$ such that

$$\lim_{r \to 0} \int_0^{2\pi} \left| d\vec{\Lambda}_{\infty}(r,\theta) - d\vec{\Lambda}_{\infty}(0) \right| \ d\theta = 0$$

and

$$\lim_{r \to 0} r^{-1} \int_0^{2\pi} \left| \vec{\Lambda}_{\infty}(r,\theta) - \vec{\Lambda}_{\infty}(0) - r \cos \theta \,\partial_{x_1} \vec{\Lambda}_{\infty}(0) - r \sin \theta \,\partial_{x_2} \vec{\Lambda}_{\infty}(0) \right| \, d\theta = 0$$

By the mean value theorem there exists $\theta_r \in [0, 2\pi]$ such that

$$\left|\vec{\Lambda}_{\infty}(r,\theta_r) - \vec{\Lambda}_{\infty}(0) - r\,\cos\theta_r\,\partial_{x_1}\vec{\Lambda}_{\infty}(0) - r\,\sin\theta_r\,\partial_{x_2}\vec{\Lambda}_{\infty}(0)\right| = o(r) \;.$$

We have

$$\begin{aligned} \forall \, \theta \in [0, 2\pi] \quad \vec{\Lambda}_{\infty}(r, \theta) - \vec{\Lambda}_{\infty}(r, \theta_{r}) &= r \, \int_{\theta_{r}}^{\theta} -\sin \, \phi \, \partial_{x_{1}} \vec{\Lambda}_{\infty}(r, \phi) + \cos \, \phi \, \partial_{x_{2}} \vec{\Lambda}_{\infty}(r, \phi) \, d\phi \\ &= r \, \int_{\theta_{r}}^{\theta} -\sin \, \phi \, \partial_{x_{1}} \vec{\Lambda}_{\infty}(0) + \cos \, \phi \, \partial_{x_{2}} \vec{\Lambda}_{\infty}(0) \, d\phi + o(r) \end{aligned}$$

Combining the previous we obtain

$$\left\|\vec{\Lambda}_{\infty}(r,\theta) - \vec{\Lambda}_{\infty}(0) - r\,\cos\theta\,\partial_{x_1}\vec{\Lambda}_{\infty}(0) - r\,\sin\theta\,\partial_{x_2}\vec{\Lambda}_{\infty}(0)\right\|_{L^{\infty}([0,2\pi])} = o(r) \,\,. \tag{IV.58}$$

As in the proof of the previous lemma we can also choose r such that there exists a subsequence (we keep denoting k) satisfying

$$\int_{\partial B_r(0)} |d\vec{\Lambda}_k|^2 \, dl \le r^{-1} \, \int_{B_{4r}(0) \setminus B_{r/4}(0)} |\nabla\vec{\Lambda}_k|^2 \, dx^2 \tag{IV.59}$$

The radii such that (IV.58) and (IV.59) hold are called good radii.

$$\|\vec{\Lambda}_{\infty}(y) - \vec{\Lambda}_{\infty}(x)\|_{L^{\infty}(\partial B_{r}(0)) \times \partial B_{r}(0))} \leq \pi |d\vec{\Lambda}_{\infty}(0)| \ r + o(r)$$

we deduce from (IV.27) that there exists a constant C > 0 independent of r such that

$$\tilde{\mu}_{\infty}\left(\left\{\vec{p}\;;\;\inf\left\{\mathfrak{r}_{\vec{q}}(\vec{p})\;;\vec{q}\in\partial\vec{\Lambda}_{\infty}(\partial B_{r}(0))\right\}>C|d\vec{\Lambda}_{\infty}(0)|\;r\right\}\right)=0\;.$$
(IV.60)

where we adopt again the notation

$$\widetilde{\mu}_{\infty} := \lim_{k \to +\infty} \left(\left. \overrightarrow{\Lambda}_k \right| B_r(0) \right)_* \nu_k .$$

Since 0 is a Lebesgue point for f we have

$$\begin{split} \nu_{\infty}(B_{4^{-4\,l}}(0)) &= \int_{B_{4^{-4\,l}}(0)} f(x) \ e^{2\lambda} \ dx^2 \\ &= \int_{B_{4^{-4\,l}}(0)} f(0) \ e^{2\lambda} \ dx^2 + o(4^{-4\,l}) \end{split}$$

(recall $\nu_{\infty}(\{0\}) = 0$) we can choose $t_j = 4^{-4j} \to 0$ and from the previous fact we have

$$\nu_{\infty} \left(B_{4t_j}(0) \right) = O(t_j^2) \tag{IV.61}$$

Finally Taking $\delta_k \to 0$ given by

$$\delta_k^2 := (\log \varepsilon_k^{-1})^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \tag{IV.62}$$

Recall that this choice is made in such a way that

$$\lim_{k \to +\infty} \delta_k \log \varepsilon_k^{-1} = 0 \quad \text{and} \quad \lim_{k \to +\infty} \delta_k^{-1} \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = 0 \,. \tag{IV.63}$$

Let F_k be the set of $\vec{p} \in V_2(\mathbb{R}^4)$ with such that there exists s > 0 with

$$\delta_k \, \mu_k(\{\mathfrak{r}_{\vec{p}} < 5 \, C_0^2 \, s\}) \le \varepsilon_k^4 \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}} < s\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k}$$

Observe that for $\vec{p} \neq \vec{q}$

$$\sup_{\vec{o}\in V_2(\mathbb{R}^4)} \frac{\mathfrak{r}_{\vec{p}}(\vec{q})}{\max\{\mathfrak{r}_{\vec{p}}(\vec{o}),\mathfrak{r}_{\vec{q}}(\vec{o})\}} \le 2C_0$$

We can then use the Vitali type Covering Lemma in Quasi-metric spaces given by Lemma 2.7 in [3] and cover F_k by at most countably many balls $(B^{\mathfrak{r}}_{5C_0^2 s_i}(\vec{p}_i))_{i \in I}$ for the quasi-distance \mathfrak{r} such that the balls $B^{\mathfrak{r}}_{5C_0^2 s_i}(\vec{p}_i)$ for $i \in I$ are disjoint to each other. Hence we have

$$\begin{split} \mu_k(F_k) &\leq \sum_{i \in I} \tilde{\mu}_k(\{\mathfrak{r}_{\vec{p}_i} < 5 \, C_0^2 \, s_i\}) \leq \delta_k^{-1} \, \sum_{i \in I} \varepsilon_k^4 \int_{B_\rho(x_0) \cap \{\mathfrak{r}_{\vec{p}_i} < s_i\}} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} \\ &\leq \delta_k^{-1} \, \varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|^2)^2 \, dvol_{g_k} = o(1) \; . \end{split}$$

We construct a sequence of "good radii" $r_j \in [t_j/2, t_j]$ a sub sequence $\phi_j(k)$ and an increasing sequence index k_j as follows. Assuming $(r_l)_{l < j}$ as well as $(\phi_l(k))_{l < j}$ and $(k_l)_{l < j}$ have been constructed, we choose r_j as explained above such that

$$\left\|\vec{\Lambda}_{\infty}(r_j,\theta) - \vec{\Lambda}_{\infty}(0) - r_j \cos\theta \,\partial_{x_1}\vec{\Lambda}_{\infty}(0) - r_j \sin\theta \,\partial_{x_2}\vec{\Lambda}_{\infty}(0)\right\|_{L^{\infty}([0,2\pi])} = o(r_j) \,. \tag{IV.64}$$

and such that there exists a subsequence of $\phi_{j-1}(k)$ that we denote $\phi_j(k)$ such that

$$\int_{\partial B_{r_j}(0)} |d\vec{\Lambda}_{\phi_j(k)}|^2 \, dl \le r_j^{-1} \, \int_{B_{4r_j}(0) \setminus B_{r_j/4}(0)} |\nabla\vec{\Lambda}_{\phi_j(k)}|^2 \, dx^2 \, . \tag{IV.65}$$

Combining (concl-222), (IV.54), (IV.61) with (IV.65), we can "wait enough" and pick k_j large enough so that

$$|\vec{\Lambda}_{\phi_j(k_j)}(0) - \vec{\Lambda}_{\infty}(0)| = o(r_j) , \qquad (IV.66)$$

 $\left|\nu_{\phi_j(k_j)}(B_{r_j}(0)) - \nu_{\infty}(B_{r_j}(0))\right| = o(r_j^2)$, (IV.67)

$$\int_{\partial B_{r_j}(0)} |d\vec{\Lambda}_{\phi_j(k_j)}|^2 \, dl \le r_j^{-1} \, O(t_j^2) = O(r_j) \,, \tag{IV.68}$$

$$\int_{B_{r_j}} |\vec{\Lambda}_{\phi_j(k_j)} - \vec{\Lambda}_{\infty}|^2 \ dx^2 = o(r_j^4) \ , \tag{IV.69}$$

$$\int_{B_{r_j}(0)} |\vec{\Lambda}_{\phi_j(k_j)} - \vec{\Lambda}_{\infty}| \ d\nu_{\phi_j(k_j)} = o(r_j^3) \ , \tag{IV.70}$$

$$\int_{B_{r_j}(0)} |\vec{\Lambda}_{\phi_j(k_j)} - \vec{\Lambda}_{\infty}|^2 \ d\nu_{\phi_j(k_j)} = o(r_j^4) \ , \tag{IV.71}$$

$$r_j^{-1}\sqrt{\varepsilon_{\phi_j(k_j)}} \longrightarrow 0$$
, (IV.72)

$$\frac{\varepsilon_{\phi_j(k_j)}^4}{r_j^2} \int_{B_{r_j}(0)} (1 + |d\vec{T}_{\phi_j(k_j)}|_{g_{\phi_j(k_j)}}^2)^2 \, dvol_{g_{\phi_j(k_j)}} = o\left(\frac{1}{\log \varepsilon_{\phi_j(k_j)}^{-1}}\right) \,, \tag{IV.73}$$

$$\mu_{\phi_j(k_j)} \left(F_{\phi_j(k_j)} \right) = o(r_j^2) .$$
 (IV.74)

Denoting $\tilde{\mathbf{v}}_{k,r}$ the varifold generated by the restriction to $B_r(0)$ of $\vec{\Lambda}_k$ we introduce the dilated varifold

 $\hat{\mathbf{v}}_{k,r} := (D_{r^{-1}})_* \tilde{\mathbf{v}}_{k,r}$

Where

$$D_{r^{-1}} : V_2(\mathbb{R}^4) \longrightarrow r^{-1}(V_2(\mathbb{R}^4) - \vec{\Lambda}_{\infty}(0))$$
$$(\vec{\varepsilon}_1 + \vec{v}, \vec{\varepsilon}_2 + \vec{w}) \longrightarrow (r^{-1}\vec{v}, r^{-1}\vec{w})$$

Let $\vec{v}_r := r^{-1}\vec{v}$, $\vec{w}_r := r^{-1}\vec{w}$ and introduce $\varphi_r := r^{-2}\varphi = r^{-2}(v^2 - w^1)$. The inverse of D_r is $(D_r)^{-1}$ such that

$$(D_{r^{-1}})^{-1}(\vec{v}_r, \vec{w}_r) := r(\vec{v}_r, \vec{w}_r) + (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$$

we have that

$$\left((D_{r^{-1}})^{-1} \right)^* \alpha = \left((D_{r^{-1}})^{-1} \right)^* \left[(\vec{\varepsilon_1} + \vec{v}) \cdot d\vec{w} - (\vec{\varepsilon_2} + \vec{w}) \cdot d\vec{v} \right]$$
$$= r \left(dw_r^1 - dv_r^2 \right) + r^2 \left[\vec{v_r} \cdot d\vec{w_r} - \vec{w_r} \cdot d\vec{v_r} \right] = r^2 \alpha_r$$

where

 $\alpha_r := -d\varphi_r + \vec{v}_r \cdot d\vec{w}_r - \vec{w}_r \cdot d\vec{v}_r$

Observe that $\hat{\mathbf{v}}_{k,r}$ is Legendrian for α_r . We denote moreover

$$\mathbf{r}_r := \left[\left(\left(|\vec{v}_r|^2 + |\vec{w}_r|^2 \right)^2 + 4 \, \varphi_r^2 \right)^{1/4} = r^{-1} \, \mathbf{r} \right]$$

For the above constructed subsequence $\phi_j(k_j)$ we introduce respectively

$$\hat{\Lambda}_{j}(y) := r_{j}^{-1} \left(\vec{\Lambda}_{\phi_{j}(k_{j})}(r_{j} y) - \vec{\Lambda}_{\infty}(0) \right) \quad , \quad \hat{\Lambda}_{\infty,j}(y) := r_{j}^{-1} \left(\vec{\Lambda}_{\infty}(r_{j} y) - \vec{\Lambda}_{\infty}(0) \right)$$

and

$$\hat{\nu}_j := \frac{1}{2} |\nabla_y \hat{\Lambda}_j|^2 \, dy^2 = \frac{1}{2 \, r_j^2} |\nabla_x \Lambda_{\phi_j(k_j)}|^2 \, dx^2 \quad , \quad \hat{\mathbf{v}}_j := \hat{\mathbf{v}}_{\phi_j(k_j), r_j} \quad \text{and} \quad \alpha_j := \alpha_{r_j}$$

We write

$$\vec{v}_j := \vec{v}_{r_j} = r_j^{-1} \vec{v} \quad , \quad \vec{w}_j := \vec{w}_{r_j} = r_j^{-1} \vec{w} \quad , \quad \varphi_j := \varphi_{r_j} := r_j^{-2} \varphi \quad \text{and} \quad \mathfrak{r}_j := r_j^{-1} \mathfrak{r} \quad .$$

We see α_j as a one form in \mathbb{R}^9 given by

$$\alpha_j := -d\varphi_j + \vec{v}_j \cdot d\vec{w}_j - \vec{w}_j \cdot d\vec{v}_j$$

We also denote

$$\hat{v}_j := \vec{v} \circ \hat{\Lambda}_j = r_j^{-1} \, \vec{v} \circ \vec{\Lambda}_j \quad \text{and} \quad \hat{w}_j := \vec{w} \circ \hat{\Lambda}_j = r_j^{-1} \, \vec{w} \circ \vec{\Lambda}_j$$

and naurally

$$\hat{v}_{l,j} := r_j^{-1} v_l \circ \vec{\Lambda}_j \quad , \quad \hat{w}_{l,j} := r_j^{-1} w_l \circ \vec{\Lambda}_j \quad , \quad \hat{\varphi}_j := r_j^{-2} \varphi \circ \vec{\Lambda}_j \quad \text{and} \quad \hat{\mathfrak{r}}_j := r_j^{-1} \mathfrak{r} \circ \vec{\Lambda}_j \ .$$

We have then in particular $\hat{\Lambda}_j = (\hat{v}_j, \hat{w}_j)$. Because of (IV.55), (IV.56), (IV.64), (IV.66)...(IV.73) we have respectively that

$$\int_{B_1(0)} |\nabla \hat{\Lambda}_{\infty,j}(y) - \nabla \vec{\Lambda}_{\infty}(0)|^2 \, dy^2 \longrightarrow 0 \quad , \quad \int_{B_1(0)} |\hat{\Lambda}_{\infty,j}(y) - \nabla \vec{\Lambda}_{\infty}(0) \cdot y| \, dy^2 \longrightarrow 0 \; . \tag{IV.75}$$

and

$$\left\|\hat{\Lambda}_{\infty,j}(1,\theta) - \cos\theta \,\partial_{x_1}\vec{\Lambda}_{\infty}(0) - \sin\theta \,\partial_{x_2}\vec{\Lambda}_{\infty}(0)\right\|_{L^{\infty}([0,2\pi])} = o(1) \tag{IV.76}$$

 $\quad \text{and} \quad$

$$\lim_{j \to +\infty} \int_{B_1(0)} |\hat{\Lambda}_j - \hat{\Lambda}_{j,\infty}|^2 \, dy^2 = 0 \tag{IV.77}$$

as well as

$$\lim_{j \to +\infty} \int_{B_1(0)} |\hat{\Lambda}_j - \hat{\Lambda}_{j,\infty}|^2 \, d\hat{\nu}_j = 0 \tag{IV.78}$$

,

.

、

Moreover the map $\hat{\Lambda}_j$ is a critical point of

$$\hat{E}_{j}(\hat{\Lambda}) := \int_{B_{1}(0)} dvol_{g_{\hat{\Lambda}}} + \hat{\varepsilon}_{j}^{4} \int_{B_{1}(0)} (r_{j}^{2} + |d\hat{T}|_{g_{\hat{\Lambda}}}^{2})^{2} dvol_{g_{\hat{\Lambda}}}$$

for perturbations within $r_j^{-1}(V_2(\mathbb{R}^4) - \vec{\Lambda}_{\infty}(0))$ preserving the Legendrian condition relative to α_j supported away from $\hat{\Lambda}_j(\partial B_1(0))$ and satisfying

$$\limsup_{j \to +\infty} \hat{E}_j(\hat{\Lambda}_j) < +\infty \quad , \quad \hat{\varepsilon}_j^4 \int_{B_1(0)} (r_j^2 + |d\hat{T}_j|_{g_{\hat{\Lambda}_j}}^2)^2 \, dvol_{g_{\hat{\Lambda}}} = o\left(\frac{1}{\log \hat{\varepsilon}_j^{-1}}\right) \tag{IV.79}$$

where

$$\hat{T}_j := \frac{\partial_{x_1} \hat{\Lambda}_j \wedge \partial_{x_2} \hat{\Lambda}_j}{|\partial_{x_1} \hat{\Lambda}_j \wedge \partial_{x_2} \hat{\Lambda}_j|} \quad \text{and} \quad \hat{\varepsilon}_j := \frac{\varepsilon_{\phi_j(k_j)}}{\sqrt{r_j}}$$

Observe that $r_j^{-1}(V_2(\mathbb{R}^4) - \vec{\Lambda}_{\infty}(0))$ equipped with α_j in the local chart given by $(\varphi_j, v_j^3, w_j^3, v_j^4, w_j^4)$ is converging in any C^l norm locally towards the Heisenberg group : $\mathbb{R}^5 := \{(\varphi_{\infty}, v_{\infty}^3, w_{\infty}^3, v_{\infty}^4, w_{\infty}^4)\}$ equipped with $\alpha_{\infty} := -d\varphi_{\infty} + v_{\infty}^3 dw_{\infty}^3 - w_{\infty}^3 dv_{\infty}^3 + v_{\infty}^4 dw_{\infty}^4 - w_{\infty}^4 dv_{\infty}^4$. We can extract a subsequence that we keep denoting j such that

 $\hat{\nu}_j \rightharpoonup \hat{\nu}_\infty$ weakly as Radon measure on $B_1(0)$ and $\hat{\mathbf{v}}_j \rightharpoonup \hat{\mathbf{v}}_\infty$ as varifold in $\mathcal{M}(G_2(\mathbb{R}^8))$

The previous lemma apply and we can then deduce that there exists \hat{f} supported in $B_1(0)$ and finitely many points y_j such that

$$\hat{\nu}_{\infty} \sqcup B_1(0) = \hat{f} \, dy^2 + \sum_{j=1}^Q c_j \, \delta_{y_j}$$

where the masses c_j are bounded from below by a universal constant $c_Q > 0$. By possibly having dilated a bit less in the image (i.e. instead of taking $\hat{\Lambda}_j := r_j^{-1}(\vec{\Lambda}_k(r_j y) - \vec{\Lambda}_\infty(0))$ one would have taken $\hat{\Lambda}_j := \beta r_j^{-1}(\vec{\Lambda}_k(r_j y) - \vec{\Lambda}_\infty(0))$ for some $\beta < 1$ we can assume that $\hat{\nu}_\infty(B_1(0)) < c_Q$ and assume that there is no mass concentration. Hence $\hat{\nu}_\infty$ is absolutely continuous with respect to Legesgue. Let $\hat{\Lambda}_{\infty,\infty}(y) = \nabla \vec{\Lambda}(0) \cdot y$. Because of (IV.57) and (IV.77) there holds respectively

$$\int_{B_1(0)} |\hat{\Lambda}_j(y) - \hat{\Lambda}_{\infty,\infty}(y)| \, dy^2 = o(1)$$

Hence we have

$$\hat{\Lambda}_j \rightarrow \hat{\Lambda}_{\infty,\infty}$$
 weakly in $W^{1,2}(B_1(0))$. (IV.80)

We can apply¹⁸ Lemma IV.2 and in particular (IV.51) for compactly supported Ψ to deduce that

$$\lim_{j \to +\infty} (\hat{\Lambda}_j)_* (\hat{\nu}_j \, \bigsqcup B_1(0)) = (\hat{\Lambda}_{\infty,\infty})_* (\hat{\nu}_\infty \, \bigsqcup B_1(0))$$
(IV.81)

This implies that $|\hat{\mathbf{v}}_{\infty}|$ is supported in the (possibly degenerated) ellipse $\mathcal{E} := \{\vec{p} := \nabla \vec{\Lambda}(0) \cdot y ; |y| < 1\}$. This ellipse is itself contained in the Legendrian hyperplane given by $\varphi_{\infty} = 0$.

For every $\vec{p} \in r_i^{-1}([V_2(\mathbb{R}^4) \setminus F_k] - \vec{\Lambda}_{\infty}(0))$ and any s > 0 we have from (III.45)

$$\begin{bmatrix} c_0 - \left[\delta_k \log\left(\frac{s\,r_j}{\varepsilon_k}\right) + \delta_k^{1/4} + \delta_k^{3/4} \log^{3/4}\left(\frac{s\,r_j}{\varepsilon_k}\right)\right]^{1/2} \end{bmatrix} \frac{1}{s^2} \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < s\,r_j\}} dvol_{g_k} \\ \leq C \int_{\Sigma \cap \{\mathfrak{r}_{\vec{p}} < 1\}} dvol_{g_k} + C \ e^{-\varepsilon_k^{-2}} \varepsilon_k^{-3} \leq C E_{\varepsilon_k} \ . \tag{IV.82}$$

where δ_k is given by (IV.62) (we have simply written k for $\phi_j(k_j)$) in the two lines above). Because of (IV.74) we deduce from (IV.82) that for $|\hat{\mathbf{v}}_{\infty}|$ a.e. \vec{p} and any s > 0 there holds

$$|\hat{\mathbf{v}}_{\infty}|(B_s^{\mathfrak{r}}(\vec{p})) \leq C s^2$$

Assume first that the rank of $\nabla \vec{\Lambda}_0$ is less or equal than one that is, the ellipse \mathcal{E} is degenerated and is either a segment or a point. We cover the support of $|\hat{\mathbf{v}}_{\infty}|$ by at most s^{-1} balls and we deduce

$$|\hat{\mathbf{v}}_{\infty}|(\mathbb{R}^5) \leq Cs$$

¹⁸Lemma IV.2 is proved in a compact ambiant space $V_2(\mathbb{R}^4)$, while we are now considering an "expanding space" converging to the Heisenberg group \mathbb{H}^2 . Nevertheless the proof of Lemma IV.2 is the same except that one has to consider compactly supported test functions exclusively.

This holds for any s we deduce

$$(\hat{\Lambda}_{\infty,\infty})_*(\hat{\nu}_{\infty} \sqcup B_1(0)) \le |\hat{\mathbf{v}}_{\infty}|(\mathbb{R}^5) = 0$$

Hence $\hat{\nu}_{\infty} \sqcup B_1(0) = 0$ in that case which implies f = 0 (thanks to (IV.52) and (IV.67)) as stated in the lemma.

We work now under the assumption that $\operatorname{rank}(\nabla \vec{\Lambda}(0)) = 2$ without loss of generality one can assume that

$$\mathcal{P}_{\infty} := \operatorname{Span}(\partial_{x_1} \vec{\Lambda}_{\infty}(0), \partial_{x_2} \vec{\Lambda}_{\infty}(0)) = \operatorname{Span}\left((\vec{\varepsilon}_3, 0), (\vec{\varepsilon}_4, 0)\right)$$

where we recall that we are using the following notations for the coordinates in \mathbb{R}^8 in the neighbourhood of $(\vec{\varepsilon_1}, \vec{\varepsilon_2})$:

$$\vec{p} := ((1+v^1)\ \vec{\varepsilon_1} + v^2\ \vec{\varepsilon_2} + v^3\ \vec{\varepsilon_3} + v^4\ \vec{\varepsilon_4}, w^1\ \vec{\varepsilon_1} + (1+w^2)\ \vec{\varepsilon_2} + w^3\vec{\varepsilon_3} + w^4\ \vec{\varepsilon_4})$$

We claim that

$$\hat{\mathbf{v}}_{\infty} = \delta_{\mathcal{P}_{\infty}} \otimes |\hat{\mathbf{v}}_{\infty}| .$$
 (IV.83)

This is equivalent to

$$\int_{B_1(0)} |\nabla \hat{v}_j^1|^2 + |\nabla \hat{w}_j^1|^2 + |\nabla \hat{v}_j^2|^2 + |\nabla \hat{w}_j^2|^2 + |\nabla \hat{w}_j^3|^2 + |\nabla \hat{w}_j^4|^2 + |\nabla \hat{\varphi}_j|^2 \, dy^2 = o(1) \,. \tag{IV.84}$$

We know from (IV.80) that $\hat{\Lambda}_j$ is weakly converging towards $\hat{\Lambda}_{\infty,\infty}$ in $W^{1,2}(B_1(0))$. Because of (IV.68) we have

$$\limsup_{j \to +\infty} \int_{\partial B_1(0)} |d\hat{\Lambda}_j|^2 \ dl < +\infty$$

Hence $\hat{\Lambda}_j$ is strongly pre-compact in $C^{0,\mu}(\partial B_1(0))$ for any $\mu < 1/2$

$$\lim_{j \to +\infty} \|\hat{\Lambda}_j - \hat{\Lambda}_{\infty,\infty}\|_{L^{\infty}(\partial B_1(0))} = 0 .$$
 (IV.85)

We have $\hat{\nu}_{\infty}(B_1(0)) < c_Q$. Let $t := \operatorname{diam}(\hat{\Lambda}_{\infty,\infty}(\partial B_1(0)))$. Thanks to (IV.5), there exists c_2

$$\lim_{j \to +\infty} \int_{B_1(0)} |\nabla \hat{\Lambda}_j|^2 \, \mathbf{1}_{\omega_j(t)} \, dy^2 = 0 \,, \qquad (\text{IV.86})$$

where $\mathbf{1}_{\omega_i(t)}$ is the characteristic function of

$$\omega_j(t) := \left\{ x \in B_1(0) \ ; \ \mathfrak{r}_j^*(\hat{\Lambda}_j(x)) > c_2 t \right\} \,,$$

and where

$$\mathfrak{r}_{j}^{*}(\hat{p}) := \inf \left\{ \mathfrak{r}_{j,\hat{p}}(\hat{q}) \quad ; \quad \hat{q} \in \hat{\Lambda}_{j}(\partial B_{1}(0)) \right\}$$

For any $\hat{p}, \hat{q} \in r_j^{-1}(V_2(\mathbb{R}^4) - \vec{\Lambda}_{\infty}(0))$ we are using the notation

$$\mathfrak{r}_{j,\hat{p}}(\hat{q}) := r_j^{-1} \mathfrak{r}_{\vec{p}_j}(\vec{q}_j) \text{ and } \vec{p}_j := r_j \,\hat{p} + \vec{\Lambda}_{\infty}(0) \text{ and } \vec{q}_j := r_j \,\hat{q} + \vec{\Lambda}_{\infty}(0) .$$

Recall that in a neighbourhood of $\vec{\Lambda}_j(0) = (\vec{\varepsilon}_1, \vec{\varepsilon}_2)$ there holds

$$\nabla^{H}v^{1} = O(\rho) \quad , \quad \nabla^{H}w^{2} = O(\rho) \quad \text{ and } \quad \nabla^{H}(w^{1} + v^{2}) = O(\rho)$$
This implies

$$\begin{aligned} \nabla \hat{v}_{j}^{1} &= |\nabla^{H} v^{1} \circ \vec{\Lambda}_{j} \cdot \nabla \hat{\Lambda}_{j}| = O(\rho \circ \vec{\Lambda}_{j}) |\nabla \hat{\Lambda}_{j}| \\ &|\nabla \hat{w}_{j}^{2}| = |\nabla^{H} w^{2} \circ \vec{\Lambda}_{j} \cdot \nabla \hat{\Lambda}_{j}| = O(\rho \circ \vec{\Lambda}_{j}) |\nabla \hat{\Lambda}_{j}| \\ &|\nabla (\hat{w}_{j}^{1} + \hat{v}_{j}^{2})| = |\nabla^{H} (w^{1} + v^{2}) \circ \vec{\Lambda}_{j} \cdot \nabla \hat{\Lambda}_{j}| = O(\rho \circ \vec{\Lambda}_{j}) |\nabla \hat{\Lambda}_{j}| \\ &\int |\nabla (\hat{w}_{j}^{1} + \hat{v}_{j}^{2})|^{2} + |\nabla (\hat{w}^{2})|^{2} + |\nabla (\hat{w}^{1} + \hat{w}^{2})|^{2} dv^{2} = o(1) \end{aligned}$$
(IV.87)

Hence

$$\int_{B_1(0)} |\nabla \hat{v}_j^1|^2 + |\nabla \hat{w}_j^2|^2 + |\nabla (\hat{w}_j^1 + \hat{v}_j^2)|^2 \, dy^2 = o(1) \,. \tag{IV.88}$$

We consider $\xi(z_1, z_3)$ to be an arbitrary smooth non negative cut-off function on \mathcal{P}_{∞} such that $\operatorname{Supp}(\xi) \subset \mathcal{E}_{\infty}$. We introduce the following Hamiltonian function

$$h_{j} := \chi(\mathfrak{r}_{j}^{*}/s) \,\xi(v_{j}^{3}, v_{j}^{4}) \, \left[v_{j}^{3} \,w_{j}^{3} + v_{j}^{4} \,w_{j}^{4} - \varphi_{j}\right]$$

$$= \chi(\mathfrak{r}^{*}/r_{j} \,s) \, r_{j}^{-2} \,\xi(r_{j}^{-1} \,v^{3}, r_{j}^{-1} \,v^{4}) \, \left[v^{3} \,w^{3} + v^{4} \,w^{4} - v^{2} + w^{1}\right] \,, \qquad (IV.89)$$

where χ is a cut-off function equal to one on [0, 1] and supported in $[2, +\infty)$. We choose $s > c_2 t$ large enough and fixed such that $\chi(\mathfrak{r}^*/s)$ is identically equal to 1 on \mathcal{E}_{∞} . Because of (IV.86) there holds for any smooth function f

$$\lim_{j \to +\infty} \int_{B_1(0)} |\nabla \hat{\Lambda}_j|^2 f(|\hat{\Lambda}_j|) \left[|\chi'(\mathfrak{r}^*/s)| + |\chi''(\mathfrak{r}^*/s)| \right] \circ \hat{\Lambda}_j \, dy^2 = 0 \,. \tag{IV.90}$$

For j large enough, because of (IV.85), $\hat{\Lambda}_j(\partial B_1(0)) \cap \text{Supp}(\xi) = \emptyset$. We have

$$\begin{split} \vec{X}_{h_j} &:= r_j^{-1} \, s^{-1} \, J_H \nabla^H \mathfrak{r}_* \, \chi'(\mathfrak{r}_j^*/s) \, \xi(v_j^3, v_j^4) \, \left[v_j^3 \, w_j^3 + v_j^4 \, w_j^4 - \varphi_j \right] \\ &+ \chi(\mathfrak{r}^*/r_j \, s) \, r_j^{-2} \, J_H \, \nabla^H \left[\xi(r_j^{-1} \, v^3, r_j^{-1} \, v^4) \, \left[v^3 \, w^3 + v^4 \, w^4 - v^2 + w^1 \right] \right] + \frac{h_j}{2} \, \left(\vec{\varepsilon}_2 + \vec{w}, -\vec{\varepsilon}_1 - \vec{v} \right) \, . \end{split}$$

which gives in particular

$$\|\nabla^{H}\vec{X}_{h_{j}}\|_{L^{\infty}(B_{Rr_{j}}(\vec{\varepsilon}_{1},\vec{\varepsilon}_{2}))} \leq C(Rr_{j})^{-2} \quad \text{and} \quad \|\nabla^{H}(\nabla^{H}\vec{X}_{h_{j}})\|_{L^{\infty}(B_{Rr_{j}}(\vec{\varepsilon}_{1},\vec{\varepsilon}_{2}))} \leq C(Rr_{j})^{-3} \quad (\text{IV.91})$$

Let

$$\hat{X}_{h_j}(\hat{p}) := r_j \, \vec{X}_{h_j}(r_j \, \hat{p} + \vec{\Lambda}_{\infty}(0))$$

Recall

$$\alpha_j := -d\varphi_j + \vec{v}_j \cdot d\vec{w}_j - \vec{w}_j \cdot d\vec{v}_j$$

and denote

$$H_j := \operatorname{Ker}(\alpha_j)$$
 and $J_{H_j} := (D_{r_j^{-1}})_* \circ J_H \circ (D_{r_j^{-1}})_*^{-1}$

We have in particular $\hat{X}_{h_j} \circ \hat{\Lambda}_j(y) = r_j \vec{X}_{h_j}(\vec{\Lambda}_j)(r_j y)$ and with the previously introduced notation there holds

$$\hat{X}_{h_j}(\hat{p}) = r_j J_H \nabla^H h_j (r_j \,\hat{p} + \vec{\Lambda}_\infty(0)) + r_j \,\frac{h_j}{2} \,\left(\vec{\varepsilon}_2 + \vec{w}(r_j \,\hat{p} + \vec{\Lambda}_\infty(0)), -\vec{\varepsilon}_1 - \vec{v}(r_j \,\hat{p} + \vec{\Lambda}_\infty(0))\right) \\ = J_{H_j} \nabla^{H_j} (r_j \,\hat{p} + \vec{\Lambda}_\infty(0)) + r_j \,\frac{\hat{h}_j}{2} \,\left(\vec{\varepsilon}_2 + \vec{w}(\vec{p}), -\vec{\varepsilon}_1 - \vec{v}(\vec{p})\right)$$

where we are using the induced metric from \mathbb{R}^8 on $r_j^{-1}(V_2(\mathbb{R}^4) - \vec{\Lambda}_{\infty}(0))$. $\hat{h}_j := h_j(r_j \hat{\Lambda}_j + \vec{\Lambda}_{\infty}(0))$. Observe that

$$\nabla^{H_j} \hat{X}_{h_j} = r_j^2 \, (\nabla^H \vec{X}_{h_j}) (r_j \, \hat{p} + \vec{\Lambda}_\infty(0)) \quad \text{and} \quad \nabla^{H_j} (\nabla^{H_j} \hat{X}_{h_j}) = r_j^3 \, \nabla^H (\nabla^H \vec{X}_{h_j}) (r_j \, \hat{p} + \vec{\Lambda}_\infty(0))$$

which gives thanks to (IV.91)

$$\|\nabla^{H_j} \hat{X}_{h_j}\|_{L^{\infty}(B_R(0))} + \|\nabla^{H_j} (\nabla^{H_j} \hat{X}_{h_j})\|_{L^{\infty}(B_R(0))} \le C$$
(IV.92)

Thanks to the entropy condition (IV.79) and the computations (II.16), the criticality of $\hat{\Lambda}_j$ of \hat{E}_j for the Hamiltonian deformations generated by \hat{X}_{h_j} is then giving

$$\int_{B_1(0)} \langle d\hat{\Lambda}_j, d(\hat{X}_{h_j} \circ \hat{\Lambda}_j) \rangle \ dy^2 = o(1) \ . \tag{IV.93}$$

Observe that

$$\left\langle d(\hat{v}_{j}, \hat{w}_{j}), d\left(\hat{h}_{j}\left(\vec{\varepsilon}_{2} + r_{j}\,\hat{w}_{j}, -\vec{\varepsilon}_{1} + r_{j}\,\hat{v}_{j}\right)\right) \right\rangle$$

$$= r_{j}\,\hat{h}_{j}\,\left\langle d(\hat{v}_{j}, \hat{w}_{j}), d(\hat{w}_{j}, -\hat{v}_{j}) \right\rangle - \left(d\hat{v}_{j}^{2} - d\hat{w}_{j}^{1} + r_{j}\,\hat{w}_{j} \cdot d\hat{v}_{j} - r_{j}\,\hat{v}_{j} \cdot d\hat{w}_{j}\right) \cdot d\hat{h}_{j}$$
(IV.94)

Recall $\varphi_j = r_j^{-2} \varphi = r_j^{-2} (v^2 - w^1) = r_j^{-1} (v_j^2 - w_j^1)$. Since $\hat{\Lambda}_j^* \alpha_j = 0$ there holds

$$r_j^{-1} d(\hat{v}_j^2 - \hat{w}_j^1) = -\hat{w}_j \cdot d\hat{v}_j + \hat{v}_j \cdot d\hat{w}_j .$$
 (IV.95)

Combining (IV.90), (IV.93), (IV.94) and (IV.95) we deduce

$$\int_{B_1(0)} \chi\left(\frac{\mathfrak{r}^*}{r_j s}\right) \frac{1}{r_j} \left\langle d\hat{\Lambda}_j, d\left(J_H \nabla^H \left[\xi(r_j^{-1} v^3, r_j^{-1} v^4) \left[v^3 w^3 + v^4 w^4 - \varphi\right]\right] \circ \vec{\Lambda}_j \right) \right\rangle \ dy^2 = o(1) \ .$$
(IV.96)

We have

From (IV.81), since $\hat{\Lambda}_{\infty,\infty}(B_1) \subset \text{Span}\{(\tilde{\varepsilon}_3,0),(\tilde{\varepsilon}_4,0)\}$, we deduce for q > 0

$$\int_{B_1(0)} \chi\left(\frac{\mathfrak{r}^*}{r_j s}\right) \left[|\hat{v}_j^1|^q + |\hat{v}_j^2|^q + |\hat{w}_j^1|^q + |\hat{w}_j^2|^q + |\hat{w}_j^3|^q + |\hat{w}_j^3|^q + |\hat{\psi}_j^4|^q + |\hat{\varphi}_j|^q \right] |d\hat{\Lambda}_j|^2 \, dy^2 = o(1) \,. \tag{IV.98}$$

Recall also from (III.47) and (III.48) that $\nabla^H w^i \circ (\hat{\Lambda}_j)$ are polynomials of \hat{v}_j^l and \hat{w}_j^l (of grade at most 3). Hence we deduce

$$\begin{split} &\int_{B_{1}(0)} \chi\left(\frac{\mathfrak{r}^{*}}{r_{j}\,s}\right) \,\frac{1}{r_{j}} \,\left\langle d\hat{\Lambda}_{j}, d\left(J_{H}\,\nabla^{H}\left[\xi(r_{j}^{-1}\,v^{3},r_{j}^{-1}\,v^{4})\right] \,\left[v^{3}\,w^{3}+v^{4}\,w^{4}-\varphi\right]\circ\vec{\Lambda}_{j}\right)\right\rangle \,dy^{2} \\ &= \int_{B_{1}(0)} \chi\left(\frac{\mathfrak{r}^{*}}{r_{j}\,s}\right) \,\left\langle d\hat{\Lambda}_{j}, \left[\nabla^{H}w^{3}(\vec{\Lambda}_{j})\,\partial_{v_{j}^{3}}\xi(\hat{v})+\nabla^{H}w^{4}(\vec{\Lambda}_{j})\,\partial_{v_{j}^{4}}\xi(\hat{v})\right] \,\left[\hat{v}_{j}^{3}\,d\hat{w}_{j}^{3}+\hat{v}_{j}^{4}\,d\hat{w}_{j}^{4}-d\hat{\varphi}_{j}\right]\right\rangle \,dy^{2} \\ &+ o(1) \;. \end{split}$$
(IV.99)

We write

$$\hat{v}_{j}^{3} d\hat{w}_{j}^{3} + \hat{v}_{j}^{4} d\hat{w}_{j}^{4} - d\hat{\varphi}_{j} = \hat{w}_{j}^{3} d\hat{v}_{j}^{3} + \hat{w}_{j}^{4} d\hat{v}_{j}^{4} - \hat{v}_{j}^{1} d\hat{w}_{j}^{1} + \hat{w}_{j}^{1} d\hat{v}_{j}^{1} - \hat{v}_{j}^{2} d\hat{w}_{j}^{2} + \hat{w}_{j}^{2} d\hat{v}_{j}^{2}$$

Substituting this expression in (IV.99) and using (IV.98) we obtain finally

$$\int_{B_{1}(0)} \chi\left(\frac{\mathbf{r}^{*}}{r_{j}s}\right) \frac{1}{r_{j}} \left\langle d\hat{\Lambda}_{j}, d\left(J_{H} \nabla^{H}\left[\xi(r_{j}^{-1}v^{3}, r_{j}^{-1}v^{4})\right] \left[v^{3}w^{3} + v^{4}w^{4} - \varphi\right] \circ \vec{\Lambda}_{j}\right) \right\rangle dy^{2}$$
(IV.100)
= $o(1)$.

Combining (IV.96) and (IV.100) is giving

$$\int_{B_1(0)} \chi\left(\frac{\mathbf{r}^*}{r_j s}\right) \frac{1}{r_j} \left\langle d\hat{\Lambda}_j, d\left(\left[\xi(r_j^{-1} v^3, r_j^{-1} v^4) J_H \nabla^H \left[v^3 w^3 + v^4 w^4 - \varphi\right]\right] \circ \vec{\Lambda}_j\right) \right\rangle \ dy^2 = o(1) .$$
(IV.101)

Recall from (III.47) and (III.48)

$$\nabla^{H}\varphi - \sum_{i=3}^{4} v^{i} \nabla^{H} w^{i} - w^{i} \nabla^{H} v^{i} = v^{1} \nabla^{H} w^{1} - w^{1} \nabla^{H} v^{1} + v^{2} \nabla w^{2} - w^{2} \nabla^{H} v^{2}$$

$$= (w^{1} \vec{v} + (w^{1})^{2} (\vec{\varepsilon}_{2} + \vec{w}) + w^{1} v^{1} (\vec{\varepsilon}_{1} + \vec{v}), -v^{1} \vec{v} - v^{1} w^{1} (\vec{\varepsilon}_{2} + \vec{w}) - (v^{1})^{2} (\vec{\varepsilon}_{1} + \vec{v})) \qquad (\text{IV.102})$$

$$+ (w^{2} \vec{w} + w^{2} v^{2} (\vec{\varepsilon}_{1} + \vec{v}) + (w^{2})^{2} (\vec{\varepsilon}_{2} + \vec{w}), -v^{2} \vec{w} - (v^{2})^{2} (\vec{\varepsilon}_{1} + \vec{v}) - v^{2} w^{2} (\vec{\varepsilon}_{2} + \vec{w})) \qquad (\text{IV.102})$$

Hence

$$\nabla^{H} \left[v^{3} w^{3} + v^{4} w^{4} - \varphi \right] - 2 w_{3} \nabla^{H} v^{3} - 2 w_{4} \nabla^{H} v^{4}$$

$$= (w^{1} \vec{v} + (w^{1})^{2} (\vec{\varepsilon}_{2} + \vec{w}) + w^{1} v^{1} (\vec{\varepsilon}_{1} + \vec{v}), -v^{1} \vec{v} - v^{1} w^{1} (\vec{\varepsilon}_{2} + \vec{w}) - (v^{1})^{2} (\vec{\varepsilon}_{1} + \vec{v})) \qquad (\text{IV.103})$$

$$+ (w^{2} \vec{w} + w^{2} v^{2} (\vec{\varepsilon}_{1} + \vec{v}) + (w^{2})^{2} (\vec{\varepsilon}_{2} + \vec{w}), -v^{2} \vec{w} - (v^{2})^{2} (\vec{\varepsilon}_{1} + \vec{v}) - v^{2} w^{2} (\vec{\varepsilon}_{2} + \vec{w}))$$

Hence

$$\int_{B_1(0)} \chi\left(\frac{\mathfrak{r}^*}{r_j s}\right) \frac{1}{r_j} \left\langle d\hat{\Lambda}_j, d\left(\left[\xi(r_j^{-1} v^3, r_j^{-1} v^4) \left[w^3 \nabla^H w^3 + w^4 \nabla^H w^4\right]\right] \circ \vec{\Lambda}_j\right) \right\rangle \ dy^2 = o(1) \ . \ (\text{IV.104})$$

Using one more time (IV.98) we finally obtain

$$\int_{B_1(0)} \chi\left(\frac{\mathfrak{r}^*}{r_j \, s}\right) \, \left[|d\hat{w}_j^3|^2 + |d\hat{w}_j^4|^2 \right] \, \xi(\hat{v}_j^3, \hat{v}_j^4) \, dy^2 = o(1) \, . \tag{IV.105}$$

Combining (IV.88) and (IV.105) is giving finally (IV.84) which itself is implying (IV.83).

Any infinitesimal variation ϕ_t such that $\operatorname{supp}((\phi_t)_*|\hat{\mathbf{v}}_{\infty}|) \subset \mathcal{E}_{\infty}$ is preserving the legendrian property of the varifold hence $\hat{\mathbf{v}}_{\infty}$ is stationary in a classical sense within the Legendrian two plane spanned by $\partial_{x_1}\vec{\Lambda}(0)$ and $\partial_{x_2}\vec{\Lambda}(0)$ relative to its boundary $\partial \mathcal{E}_{\infty}$. We deduce using the constancy theorem (see [21] chapter 8 theorem 41.1) that there exists a constant $\theta_0 > 0$ such that

$$|\hat{\mathbf{v}}_{\infty}| = \theta_0 \ d\mathcal{H}^2 \square \mathcal{E}$$
.

It remains to prove that m is a positive integer.

For any $0 < \beta < 1$ we denote

$$\mathcal{E}_{\beta} := \left\{ \vec{p} := \nabla \vec{\Lambda}_{\infty}(0) \cdot y \quad ; \quad |y| < \beta \right\}$$

Let $a(v_{\infty}^3, v_{\infty}^4) \in C_0^{\infty}(\mathcal{E}_{\beta}(0))$ and $b(v_{\infty}^3, v_{\infty}^4) \in C_0^{\infty}(\mathcal{E}_{\beta})$ arbitrary. Hence, because of (IV.85), for j large enough, there holds

$$a \circ \hat{\Lambda}_j \equiv 0$$
 and $b \circ \hat{\Lambda}_j \equiv 0$ on $\partial B_1(0)$. (IV.106)

We introduce now the following Hamiltonian

$$h_j(\vec{v}_j, \vec{w}_j) := \chi\left(\frac{\mathfrak{r}^*}{r_j \, s}\right) \left[a(v_j^3, v_j^4) \, w_j^3 + b(v_j^3, v_j^4) \, w_j^4\right] \; .$$

Arguing exactly as for the previous Hamiltonian gives then

$$\lim_{j \to 0} \int_{B_1(0)} \chi\left(\frac{\mathfrak{r}^*}{r_j s}\right) \nabla\left[a(\hat{v}_j^3, \hat{v}_j^4) \vec{\varepsilon}_3\right] \nabla\tilde{\Lambda}_j + \nabla\left[b(\hat{v}_j^3, \hat{v}_j^4) \vec{\varepsilon}_4\right] \nabla\tilde{\Lambda}_j \ dx^2 = 0 \tag{IV.107}$$

where we have introduced the notation $\tilde{\Lambda}_j := (\hat{v}_j^3, \hat{v}_j^4)$ for the projection on the two Legendrian plane \mathcal{L} generated by $(0, \vec{\varepsilon}_3, 0)$ and $(0, \vec{\varepsilon}_4, 0)$. Recall that $\hat{\Lambda}_j$ is conformal on $B_1(0)$. Denote

$$\hat{\Lambda}_{j}^{*}g_{\mathbb{R}^{8}} = e^{2\lambda_{j}} \left[dx_{1}^{2} + dx_{2}^{2} \right]$$

Because of (IV.84) we have in particular for l = 3, 4

$$\lim_{j \to +\infty} \int_{B_1(0)} |e^{2\hat{\lambda}_j} \left(\vec{\varepsilon}_l, 0\right) - \left(\vec{\varepsilon}_l, 0\right) \cdot \nabla \hat{\Lambda}_j \nabla \hat{\Lambda}_j | dx^2 = 0 ,$$

which implies for l = 3, 4

$$\lim_{j \to +\infty} \int_{B_1(0)} |e^{2\hat{\lambda}_j} - |\nabla \hat{v}_j^l|^2 | \, dx^2 = 0 \,. \tag{IV.108}$$

Observe that we have also

$$\lim_{j \to +\infty} \int_{B_1(0)} \left| e^{\hat{\lambda}_j} \left(\vec{\varepsilon}_l, 0 \right) - \left(\vec{\varepsilon}_l, 0 \right) \cdot \nabla \hat{\Lambda}_j \nabla \hat{\Lambda}_j \ e^{-\hat{\lambda}_j} \right|^2 \ dx^2 = 0$$

Using the fact that $(\vec{\varepsilon_3},0) \cdot (\vec{\varepsilon_4},0) = 0$ we deduce from the previous line

$$\lim_{j \to +\infty} \int_{B_1(0)} |\nabla \hat{v}_j^4 \cdot \nabla \hat{v}_j^3| \, dx^2 = 0 \, . \tag{IV.109}$$

We have

$$\begin{split} \partial_{x_1} \left[a(\hat{v}_j^3, \hat{v}_j^4) \right] \partial_{x_1} \hat{v}_j^3 + \partial_{x_1} \left[b(\hat{v}_j^3, \hat{v}_j^4) \right] \partial_{x_1} \hat{v}_j^4 + \partial_{x_2} \left[a(\hat{v}_j^3, \hat{v}_j^4) \right] \partial_{x_2} \hat{v}_j^3 + \partial_{x_2} \left[b(\hat{v}_j^3, \hat{v}_j^4) \right] \partial_{x_2} \hat{v}_j^4 \\ &= \partial_{\hat{v}_j^3} a \left(\partial_{x_1} \hat{v}_j^3 \right)^2 + \partial_{\hat{v}_j^4} a \partial_{x_1} \hat{v}_j^4 \partial_{x_1} \hat{v}_j^3 + \partial_{\hat{v}_j^3} a \left(\partial_{x_2} \hat{v}_j^3 \right)^2 + \partial_{\hat{v}_j^4} a \partial_{x_2} \hat{v}_j^4 \partial_{x_2} \hat{v}_j^3 \\ &+ \partial_{\hat{v}_j^3} b \partial_{x_1} \hat{v}_j^3 \partial_{x_1} \hat{v}_j^4 + \partial_{\hat{v}_j^4} b \left(\partial_{x_1} \hat{v}_j^4 \right)^2 + \partial_{\hat{v}_j^3} b \partial_{x_2} \hat{v}_j^3 \partial_{x_2} \hat{v}_j^4 + \partial_{\hat{v}_j^4} b \left(\partial_{x_2} \hat{v}_j^4 \right)^2 \\ &= e^{2\hat{\lambda}_j} \left[\partial_{\hat{v}_j^3} a + \partial_{\hat{v}_j^4} b \right] + \partial_{\hat{v}_j^3} a \left[|\nabla \hat{v}_j^3|^2 - e^{2\hat{\lambda}_j} \right] + \partial_{\hat{v}_j^4} b \left[|\nabla \hat{v}_j^4|^2 - e^{2\hat{\lambda}_j} \right] + \left[\partial_{\hat{v}_j^4} a + \partial_{\hat{v}_j^3} b \right] \nabla \hat{v}_j^4 \cdot \nabla \hat{v}_j^3 \ . \end{split}$$

Combining this identity with (IV.107), (IV.108) and (IV.109) gives then

$$\lim_{j \to +\infty} \int_{B_1(0)} \left[\partial_{\hat{v}_j^3} a + \partial_{\hat{v}_j^4} b \right] \circ \hat{\Lambda}_j \ |\partial_{x_1} \hat{\Lambda}_j \wedge \partial_{x_1} \hat{\Lambda}_j| \ dx_1 \wedge dx_2 = 0$$

The area formula gives then

$$\lim_{j \to +\infty} \int_{\mathcal{L}} \left[\partial_{y_1} a + \partial_{y_2} b \right](y) \ N_j(y) \ dy^2 = 0$$
 (IV.110)

where $N_j(y)$ is the number of (x_1, x_2) in $B_1(0)$ such that $(\hat{v}_3(x_1, x_2), \hat{v}_4(x_1, x_2)) = (y_1, y_2)$. At this stage we have used a uniform L^{∞} control of $(\nabla_y^l a, \nabla_y^l b)$ for $l \leq 3$ and we have proved

$$\lim_{j \to +\infty} \sup_{\sum_{l=0}^{3} \|\nabla_{y}^{l}(a,b)\|_{\infty} \leq 1} \int_{\mathcal{L}} \left[\partial_{y_{1}}a + \partial_{y_{2}}b \right](y) \ N_{j}(y) \ dy^{2} = 0$$
(IV.111)

Let's take now a small parameter $\tau_j \to 0$ and consider the convolution (a_j, b_j) of a and b respectively with $\chi_j(y_1, y_2) := \tau_j^{-1} \chi((y_1/\tau_j, y_2/\tau_j))$ where $\chi \in C_0^{\infty}(\mathbb{R}^2)$ with $\int_{\mathbb{R}^2} \chi(y) \, dy^2 = 1$. We have obviously

$$\|\nabla_{y}[(a,b) - (a_{j},b_{j})]\|_{L^{\infty}(\mathbb{R}^{2})} = o(1) \|\nabla_{y}(a,b)\|_{L^{\infty}(\mathbb{R}^{2})}$$

which implies in particular

$$\lim_{j \to +\infty} \int_{\mathcal{L}} \left[\partial_{y_1}(a - a_j) + \partial_{y_2}(b - b_j) \right](y) \ N_j(y) \ dy^2 = o(1) \ \|\nabla_y(a, b)\|_{L^{\infty}(\mathbb{R}^2)} \ .$$

We have also for $l \leq 3$

$$\|\nabla_{y}^{l}(a_{j}, b_{j})\|_{L^{\infty}(\mathbb{R}^{2})} \leq \tau_{j}^{-l+1} \|\nabla_{y}(a, b)\|_{L^{\infty}(\mathbb{R}^{2})}$$

We choose τ_j in such a way that τ_j^2 goes infinitesimally slower to zero than all the terms in the estimates above which go to zero. In particular we require

$$\tau_j^{-2} \hat{\varepsilon}_j^4 \int_{B_1(0)} (r_j^2 + |d\hat{T}_j|^2_{g_{\hat{\Lambda}_j}})^2 \, dvol_{g_{\hat{\Lambda}}} = o(1) \, \|\nabla_y(a,b)\|_{L^{\infty}(\mathbb{R}^2)} \, .$$

In this way we deduce foor any $\beta < 1$

$$\lim_{j \to +\infty} \sup_{\|\nabla_y(a,b)\|_{\infty} \le 1} \sup_{\text{Supp}(a,b) \subset \mathcal{E}_{\beta}} \int_{\mathcal{L}} \left[\partial_{y_1} a + \partial_{y_2} b \right](y) \ N_j(y) \ dy^2 = 0 \ . \tag{IV.112}$$

Using Lemma A.7 of [15] we obtain

$$\lim_{j \to +\infty} \left| N_j - |\mathcal{E}_{1/2}|^{-1} \int_{\mathcal{E}_{1/2}} N_j(y) \, dy^2 \right|_{L^{1,\infty}(\mathcal{E}_{1/2})} = 0 \tag{IV.113}$$

The area formula gives also

$$N_j(y) dy^2
ightarrow |\hat{\mathbf{v}}_{\infty}| = \theta_0 d\mathcal{H}^2 \square \mathcal{E} \text{ in } (C^0(\mathcal{E}_{1/2}))^*$$

This implies in particular

$$\lim_{j \to +\infty} |\mathcal{E}_{1/2}|^{-1} \int_{\mathcal{E}_{1/2}} N_j(y) \, dy^2 = \theta_0$$

and from (IV.113), we obtain

$$\lim_{j \to +\infty} \operatorname{dist} \left(|\mathcal{E}_{1/2}|^{-1} \int_{\mathcal{E}_{1/2}} N_j(y) \, dy^2, \mathbb{N} \right) = 0$$

hence

$$\theta_0 \in \mathbb{N}^*$$
.

Because of (IV.67) we have

$$\pi \,\theta_0 \,\left| \partial_{x_1} \vec{\Lambda}_\infty(0) \wedge \partial_{x_2} \vec{\Lambda}_\infty(0) \right| = \theta_0 \,\left| \mathcal{E} \right| = \lim_{j \to +\infty} \hat{\nu}_j(B_1(0)) = \lim_{r \to 0} \frac{\nu_\infty(B_r(x))}{r^2} = \pi \, f(0) \, e^{2\lambda(0)}$$

This implies (IV.53). Since $\mu_k = (\vec{\Lambda}_k)_* \nu_k \rightharpoonup (\vec{\Lambda}_\infty)_* \nu_\infty$, we have that $\mu_\infty = (\vec{\Lambda}_\infty)_* \nu_\infty$. Thus for $|d\vec{\Lambda}_\infty|_h^2 dvol_h$ almost every $x \in \Sigma$

$$\pi N_x \le \limsup_{t \to 0} \frac{\mu_{\infty}(B_t^{\mathfrak{r}}(\vec{\Lambda}_{\infty}(x)))}{t^2} \le C \,\mu_{\infty}(V_2(\mathbb{R}^4))$$

Hence N is in $L^{\infty}(\Sigma)$. This concludes the proof of lemma IV.3.

We now identify the limiting varifold \mathbf{v}_{∞} as being the varifold associated to $\vec{\Lambda}_{\infty}$ with the weight N. Precisely we have.

Lemma IV.4. We denote by $\mathbf{v}_{\infty}^{\eta}$ the limit of the varifold induced by $\vec{\Lambda}_k$ on $\Sigma \setminus \bigcup_{i=1}^Q B_{\eta}(x_i)$ where $\eta > 0$ is arbitrary

$$\forall \ \Xi \in C^0(G_2^{Leg}(V_2(\mathbb{R}^4))) \quad \mathbf{v}_{\infty}^{\eta}(\Xi) := \int_{\Sigma \setminus \cup_{i=1}^Q B_{\eta}(x_i)} N \ \Xi((\vec{\Lambda}_{\infty})_* T_x \Sigma, \vec{\Lambda}_{\infty}(x)) \ dvol_{g_{\infty}} \ .$$

Proof of lemma IV.4 For almost every $x_0 \in \Sigma$ one has

$$\lim_{r \to 0} \int_{B_r(x_0)} \frac{|d\vec{\Lambda}_{\infty}(x) - d\vec{\Lambda}_{\infty}(0)|^2}{r^2} \, dx^2 = 0 \,. \tag{IV.114}$$

as well as

$$\lim_{r \to 0} \int_{B_r(x_0)} \frac{\left| \vec{\Lambda}_{\infty}(x) - \vec{\Lambda}_{\infty}(0) - d\vec{\Lambda}_{\infty}(0) \cdot x \right|}{r^3} \ dx^2 = 0 \tag{IV.115}$$

We take such a point x_0 and we consider conformal coordinates for h such that $x_0 = (0,0)$. As in the proof of the previous lemma, we have a sequence of radii such that

$$\left\|\vec{\Lambda}_{\infty}(r_j,\theta) - \vec{\Lambda}_{\infty}(x_0) - r_j \cos\theta \,\partial_{x_1}\vec{\Lambda}_{\infty}(0) - r_j, \sin\theta \,\partial_{x_2}\vec{\Lambda}_{\infty}(0)\right\|_{L^{\infty}([0,2\pi])} = o(r_j) \,. \tag{IV.116}$$

Let $\delta > 0$. We claim first that for any $\Xi \in C^0(G_2^{Leg}(V_2(\mathbb{R}^4))))$, for j large enough

$$\limsup_{k \to +\infty} \int_{B_{r_j}(0)} \left| \Xi((\vec{\Lambda}_k)_* T_x \Sigma, \vec{\Lambda}_k(x)) - \Xi((\vec{\Lambda}_\infty)_* T_x \Sigma, \vec{\Lambda}_\infty(x)) \right| \, dvol_{g_k} \le \delta \, r_j^2$$

This claim is proved as in the proof of (5.10) in [14] and the lemma follows for the same reasons as the one given in the proof of proposition 5.7 of [14].

V Proof of theorem I.1 for $N^5 = V_2(\mathbb{R}^4)$.

Let \mathcal{A} be an admissible family in $\mathfrak{M} := \mathfrak{E}^{2,4}_{\Sigma,Leq}$ such that

$$\beta := \inf_{A \in \mathcal{A}} \sup_{\vec{\Lambda} \in A} \int_{\Sigma} dvol_{g_{\vec{\Lambda}}} > 0$$

We apply proposition II.1 for $f(t) := e^{-1/t^2}$ and we obtain a sequence $\varepsilon_k \to 0$ as well as $\vec{\Lambda}_k \in \mathfrak{E}^{2,4}_{\Sigma,Leg}$ satisfying

$$\limsup_{k \to +\infty} E_{\varepsilon_k}(\vec{\Lambda}_k) = \beta \tag{V.117}$$

moreover

$$|dE_{\varepsilon_k}(\vec{\Lambda}_k)||_{\vec{\Lambda}_k} \le e^{-\varepsilon_k^{-2}} , \qquad (V.118)$$

and satisfying the entropy condition

$$\varepsilon_k^4 \int_{\Sigma} (1 + |d\vec{T}_k|_{g_k}^2)^2 \, dvol_{g_k} = o\left(\frac{1}{\log \varepsilon_k^{-1}}\right) \,. \tag{V.119}$$

The section III and IV apply. At the concentration points x_i we proceed to a blow-up analysis exactly as in the proof of lemma III.6 in [17], using (IV.6) of lemma IV.1 in the same way as the global energy quantization lemma III.3 is used in [17], to prove that there is no area dissipation in the neck regions (see also 6.3 in [14]). Finally the case when the Riemann structure induced by $\vec{\Lambda}_k$ is degenerating is treated as in section IV of [17] or section 6 in [14]. This is ending the proof of theorem I.1.

VI The proof of theorem I.1 in the general case.

We recall the definition

Definition VI.4. A compact Riemannian manifold (N^5, g) is Sasakian if its metric cone $(C(S) := \mathbb{R}^*_+ \times N^5, \overline{g} = dt^2 + t^2 g)$ is Kähler.

As a consequence of the definition C(S) is equipped with an integrable complex structure J and a Kähler 2–form. On $\{1\} \times N^5 \simeq N^5$ we define the Reeb vector-field

$$\vec{R} := \sqrt{2} J(\partial_t)$$

as well as the 1-form

$$\alpha := \sqrt{2} J dt$$

so that $\alpha(\vec{R}) = 2$. In N^5 we define $H := \ker(\alpha)$ and J^H to be the restriction to H of J (bearing in mind that J realizes an isometry of H). Observe that, since \overline{g} is compatible with J the explicit expression of \overline{g} gives

$$\forall \vec{X} \in H \qquad 0 = \overline{g}(\partial_t, J\vec{X}) = -g(J\partial_t, \vec{X})$$

Hence $H \ker(\alpha)$ coincides with the orthogonal to the Reeb vector-field \vec{R} . The symplectic form Ω on $(C(S) := \mathbb{R}^*_+ \times N^5, \overline{g} = dt^2 + t^2 g)$ is given by

$$\Omega := dt \wedge J \, dt + \frac{t^2}{2\sqrt{2}} \, d\alpha = \frac{i}{2} \, \overline{\partial} \partial t^2$$

Since Ω is non degenerate we deduce that α is contact on N^5 that is

$$lpha \wedge dlpha \wedge dlpha
eq 0$$
 .

C

A 5-Sasakian manifold is then often seen as the collection of $(N^5, g, \alpha, \vec{R}, J^H)$ satisfying the above conditions. In [8] the existence of a "Sasakian potential" associated to every Sasakian structure has been discovered (see also a presentation in [22] section 1.2). At the neighbourhood of any point \vec{p} there exists on N^5 a local chart $(\psi, z_1 = x_1 + iy_1, z_2 = x_2 + iy_2)$ where $\psi(\vec{p}) = 0, z_j(\vec{p}) = 0$ and a real valued function h depending only on (x_1, y_1, x_2, y_2) such that

$$\begin{cases} \vec{R} = \sqrt{2} \,\partial_{\psi} \\ H^{0,1} = (H \otimes \mathbb{C})^{(1,0)} \quad \text{is generated by } \partial_{z_j} - \sqrt{2}^{-1} \,\alpha(\partial_{z_j}) \,\partial_{\psi} \\ J^H(\partial_{z_j} - \sqrt{2}^{-1} \,\alpha(\partial_{z_j}) \,\partial_{\psi}) = i \left(\partial_{z_j} - \sqrt{2}^{-1} \,\alpha(\partial_{z_j}) \,\partial_{\psi}\right) \\ \alpha = \sqrt{2} \,d\psi + i \,\sqrt{2} \left[\sum_{j=1}^2 \,\partial_{z_j} h \,dz_j - \partial_{\overline{z}_j} h \,d\overline{z}_j\right] \end{cases}$$
(VI.1)

Moreover h is a Kähler potential for the transversally Kähler foliation $\psi = cte$ and we can also assume¹⁹ (modulo an holomorphic change of coordinates $z \to w$ on the leafs) that

$$\frac{\partial^2 h}{\partial z_j \partial \overline{z}_k}(0) = \delta_{jk} \tag{VI.2}$$

so that the transverse Kähler metric on the leafs $\psi = cte$ satisfies

$$g^{T}(X,Y) = \frac{1}{2\sqrt{2}} d\alpha(X,JY) = g_{j\overline{k}}^{T} dz_{j} \otimes d\overline{z}_{k} \quad \text{and} \quad g_{j\overline{k}}^{T}(0) = \delta_{jk}$$

Now we introduce

$$\varphi := \psi + i \sum_{j=1}^{2} \partial_{z_j} h(0) \, z_j - i \sum_{j=1}^{2} \partial_{\overline{z}_j} h(0) \, \overline{z}_j + \frac{i}{2} \sum_{j,k=1}^{2} \partial_{z_j z_k}^2 h(0) \, z_j \, z_k - \frac{i}{2} \sum_{j,k=1}^{2} \partial_{\overline{z}_j \overline{z}_k}^2 h(0) \, \overline{z}_j \, \overline{z}_k \; .$$

We have

$$\sqrt{2}^{-1} \alpha = d\varphi + i \sum_{j=1}^{2} \left[\partial_{z_j} h - \partial_{z_j} h(0) - \sum_{k=1}^{2} \frac{\partial^2 h}{\partial z_j \partial z_k}(0) z_k \right] dz_j$$
$$-i \sum_{j=1}^{2} \left[\partial_{\overline{z}_j} h - \partial_{\overline{z}_j} h(0) - \sum_{k=1}^{2} \partial_{\overline{z}_j \overline{z}_k}^2 h(0) \overline{z}_k \right] d\overline{z}_j \qquad (VI.3)$$
$$= d\varphi + i \sum_{j=1}^{2} (\overline{z}_j + O(|z|^2)) dz_j - i \sum_{j=1}^{2} (z_j + O(|z|^2)) d\overline{z}_j$$

Hence by dilating at \vec{p} we are asymptotically converging to the Heisenberg group \mathbb{H}^2 as in the special case $V_2(\mathbb{R}^4)$ above and the proof of theorem I.1 given for $V_2(\mathbb{R}^4)$ in sections II...V can be adapted word by word with the same error terms (which were not taking into account the particular structure of $V_2(\mathbb{R}^4)$ but only the convergence to \mathbb{H}^2 in the blow-up).

Conflict of interest The author declares no conflict of interest.

¹⁹This change of coordinate is called *special foliated coordinates* in [23]

References

- Allard, William K. A characterization of the area integrand. Symposia Mathematica, Vol. XIV (Convegno di Geometria Simplettica e Fisica Matematica & Convegno di Teoria Geometrica dell'Integrazione e Varietà Minimali, INDAM, Rome, 1973), pp. 429-444. Academic Press, London, 1974.
- [2] Almeida, Luis The regularity problem for generalized harmonic maps into homogeneous spaces. Calc. Var. Partial Differential Equations 3 (1995), no. 2, 193-242.
- [3] Alvarado, Ryan; Mitrea, Marius Hardy spaces on Ahlfors-regular quasi metric spaces. A sharp theory. Lecture Notes in Mathematics, 2142. Springer, Cham, 2015.
- Boyer, Charles P.; Galicki, Krzysztof On Sasakian-Einstein geometry. Internat. J. Math. 11 (2000), no. 7, 873-909.
- [5] Boyer, Charles P.; Galicki, Krzysztof Sasakian geometry. Oxford Mathematical Monographs. Oxford University Press, Oxford, 2008
- [6] Dazord, Pierre Sur la géométrie des sous-fibrés et des feuilletages lagrangiens. (French) [On the geometry of sub-bundles and Lagrange foliations] Ann. Sci. École Norm. Sup. (4) 14 (1981), no. 4, 465-480 (1982).
- [7] Evans, Lawrence C.; Gariepy, Ronald F. Measure theory and fine properties of functions. Studies in Advanced Mathematics. CRC Press, Boca Raton, FL, 1992.
- [8] Godlinski, Michal; Kopczynski, Wojciech; Nurowski, Pawel Locally Sasakian manifolds. Classical Quantum Gravity 17(2000), no.18, 105-115
- [9] Guan, Pengfei; Zhang, Xi A geodesic equation in the space of Sasakian metrics. Geometry and analysis. No. 1, 303-318. Adv. Lect. Math. (ALM), 17 International Press, Somerville, MA, 2011
- [10] Harvey, Reese; Lawson, H. Blaine, Jr. Calibrated geometries. Acta Math. 148 (1982), 47-157.
- [11] Micallef, M.; Wolfson, J. Area minimizers in a K3 surface and holomorphicity. Geom. Funct. Anal.16(2006), no.2, 437-452
- [12] Oh, Yong-Geun Volume minimization of Lagrangian submanifolds under Hamiltonian deformations. Math. Z. 212 (1993), no. 2, 175-192.
- [13] Oh, Yong-Geun Mean curvature vector and symplectic topology of Lagrangian submanifolds in Einstein-Kähler manifolds. Math. Z. 216 (1994), no. 3, 471-482.
- [14] Pigati, Alessandro The viscosity method for min-max free boundary minimal surfaces. Arch. Ration. Mech. Anal. 244 (2022), no. 2, 391-441.
- [15] Pigati, Alessandro; Rivière, Tristan The regularity of parametrized integer stationary varifolds in two dimensions. Comm. Pure Appl. Math. 73 (2020), no. 9, 1981-2042
- [16] Rivière, Tristan The regularity of conformal target harmonic maps. Calc. Var. Partial Differential Equations 56 (2017), no.4, Paper No. 117
- [17] Rivière, Tristan A viscosity method in the min-max theory of minimal surfaces. Publ. Math. Inst. Hautes Études Sci. 126 (2017), 177-246.

- [18] Rivière, Tristan Almost Monotonicity Formula for H-minimal Legendrian Surfaces in the Heisenberg Group Comm. Pure Appl. Math. 77 (2024), no.3, 1940-1957
- [19] Schoen, Richard; Wolfson, Jon Minimizing volume among Lagrangian submanifolds. Differential equations: La Pietra 1996 (Florence), 181-199, Proc. Sympos. Pure Math., 65, Amer. Math. Soc., Providence, RI, 1999.
- [20] Schoen, Richard; Wolfson, Jon Minimizing area among Lagrangian surfaces: the mapping problem. J. Differential Geom. 58 (2001), no. 1, 1-86.
- [21] Simon, Leon Lectures on geometric measure theory. Proceedings of the Centre for Mathematical Analysis, Australian National University, 3. Canberra, 1983.
- [22] Sparks, James Sasaki-Einstein manifolds. Surveys in differential geometry. Volume XVI. Geometry of special holonomy and related topics, 265-324, Surv. Differ. Geom., 16, Int. Press, Somerville, MA, 2011
- [23] Vezzoni, Luigi; Zedda, Michela On the J-flow in Sasakian manifolds. Ann. Mat. Pura Appl. (4)195(2016), no.3, 757-774
- [24] Wolfson, Jon G. Minimal Lagrangian diffeomorphisms and the Monge-Ampère equation. J. Differential Geom. 46 (1997), no. 2, 335-373
- [25] Wolfson, Jon Lagrangian homology classes without regular minimizers. J. Differential Geom. 71 (2005), no. 2, 307-313.