Finite Population Survey Sampling: An Unapologetic Bayesian Perspective

Sudipto Banerjee^{1*}

^{1*}UCLA Department of Biostatistics; UCLA Department of Statistics and Data Science, University of California Los Angeles, 650 Charles E. Young Drive South, Los Angeles, 90095-1772, California, USA.

Corresponding author(s). E-mail(s): sudipto@ucla.edu;

Abstract

This article attempts to offer some perspectives on Bayesian inference for finite population quantities when the units in the population are assumed to exhibit complex dependencies. Beginning with an overview of Bayesian hierarchical models, including some that yield design-based Horvitz-Thompson estimators, the article proceeds to introduce dependence in finite populations and sets out inferential frameworks for ignorable and nonignorable responses. Multivariate dependencies using graphical models and spatial processes are discussed and some salient features of two recent analyses for spatial finite populations are presented.

Keywords: Bayesian inference, Finite population survey sampling, Graphical models, Hierarchical models, Spatial data

1 Introduction

Basu (1971) tells a delightful and impressionable tale of a circus statistician and an elephant to cogently illustrate, with characteristic humour, the potential pitfalls of the widely used design-based Horvitz-Thompson estimator (Narain, 1951; Horvitz and Thompson, 1952) in survey sampling. Little (2004), in a splendidly written review paper, quotes this tale and offers substantial insights on inference for finite populations in an illuminating broader discussion of model-based and design-based approaches in survey sampling. These two paradigms have received significant attention in survey sampling and I do not intend to undertake a comprehensive review here. Finite population survey sampling has an extensive literature that is beyond the scope of a single

article (see classic texts such as Cochran, 1977; Kish, 1965, emphasizing design-based approaches). Bayesian inference for finite population survey sampling has been discussed from diverse perspectives, for example, in Ericson (1969), Rao and Ghangurde (1972), Arora et al (1997), Ghosh and Meeden (1997), Little and Rubin (2002) Gelman (2007), Little (2004) and Gelman et al (2014), while an excellent appraisal of classical and Bayesian approaches is offered by Rao (2011). Di Zio et al (2023) offer a wonderfully insightful review of Debabrata Basu's legacy in the development of survey sampling, which I will not undertake here.

I will share some perspectives on Bayesian inference for finite population quantities with an emphasis on dependent finite populations. Little (2004) comments that advocating Bayesian inference for survey sampling is akin to "swimming upstream", given the aversion that many survey statisticians have to modelling assumptions, but notes the richness and flexibility available in models for improving design-based inference and for addressing survey weights. Advocates of design-based survey sampling, where inference is based on the randomisation distribution, have often criticised modelling assumptions such as the population units being "independent and identically distributed", which fails to account for the sampling design (see, e.g., Kish, 1995, Section 9). However, with computational resources available to statisticians today, building models for dependent finite population units yield rich and flexible classes of Bayesian models. A key point is that the usual model-based approaches (likelihood or Bayesian) often proceed from an assumption of "ignorability" that the analysis includes all variables that affect the probability of a unit being included in the sample. A different way to make this point is to say that the probability of obtaining a given sample does not depend upon the values of the population units. Under such assumptions, the likelihood, or the model of how the data was realised, drives the inference and the sampling design can be ignored (see, e.g., Rubin, 1976; Gelman, 2007; Gelman et al, 2014, for further discussions).

The Bayesian approach to statistical inference seeks to provide full uncertainty quantification of unknown quantities of interest conditional upon known (or seen) quantities. Denoting \mathcal{U} to be a collection of unknowns and \mathcal{K} a collection of "knowns", we wish to report the posterior distribution $[\mathcal{U} \mid \mathcal{K}] \propto [\mathcal{U}, \mathcal{K}]$. Probabilistic inference arrives at this distribution by specifying $[\mathcal{K} \mid \mathcal{U}]$ (the likelihood) and $[\mathcal{U}]$ (the prior). Treating Bayesian survey sampling from the perspective of missing data is appealing and conceptually simple. Here \mathcal{K} represents the sampled data and \mathcal{U} represents all unknowns including the population units not in the sample as well as parameters needed to specify a probability model for $[\mathcal{U}]$. The plethora of applications within this paradigm, especially in the context of complex surveys, is daunting due to sheer volume (see, e.g., Little, 1982; Little and Rubin, 2002; Tang et al, 2003; Gelman et al, 2014; Kalton, 2019, and references therein).

Following conventional notations, let us assume that $\mathcal{N} = \{1, 2, \ldots, N\}$ is a finite set of integers labelling the population units. Given a sample $\{i_1, i_2, \ldots, i_n\}$ of size n, which is a subset of \mathcal{N} , units in the population and sample are described by $\{(Y_t, Z_t) : t \in \mathcal{N}\}$, where Y_t denotes the value of population unit labelled t and Z_t is a binary variable indicating if unit t is in the sample or not. The binary vector $Z = (Z_1, Z_2, \ldots, Z_N)^{\mathrm{T}}$ completely encodes the sampled units in the population and,

analogously, $Y = (Y_1, Y_2, \ldots, Y_N)^{\mathsf{T}}$ denotes the values of the all units in the finite population. Model-based inference (likelihood or Bayesian) assigns a probability law to the finite collection (Y, Z), which defines the sampling design. It is sometimes convenient to imagine the exercise of sampling as a process over time where units enter the sample one by one. At time zero, then, a probability model for Y, p(Y), can be interpreted as a prior distribution on the unknown population units. The sampling design determines the joint distribution over Y and Z by specifying p(Z | Y). Bayesian inference proceeds from $p(Y | \mathcal{D}, Z) \propto p(Y)p(Z | Y)p(\mathcal{D} | Y, Z)$, where \mathcal{D} is the sampled data.

In the domain of dependent finite population models, there is a substantial literature on small area estimation for regionally aggregated data (see, e.g., Rao and Molina, 2015; Clayton and Kaldor, 1987; Datta and Ghosh, 1991; Ghosh and Rao, 1994; Arora and Lahiri, 1997; Ghosh et al, 1998), where we model dependencies across regions. If one considers "being adjacent to" or "being a neighbour of" as a relation over the set of regions, then one can build dependencies using a graph $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$, where vertices in \mathcal{V} correspond to the regions and the edges in \mathcal{E} are incident upon pairs of vertices that represent adjacent regions. Graphical models recognise conditional dependence among adjacent nodes, and conditional independence otherwise, given all other nodes.

Much of this article revisits familiar models in Bayesian finite population inference, but attempts to offer additional insights from a more contemporary perspective of modelling complex multivariate dependencies. Section 2 presents Bayesian inference for simple random sampling without replacement. Section 3 discusses Bayesian models that reproduce inference from Horvitz-Thompson estimators. Section 4 discusses Bayesian multistage sampling as devised in Scott and Smith (1969) and Malec and Sedransk (1985), but with a modest generalisation to modelling unknown variances. I offer derivations in a systematic manner using conjugate Bayesian updating with minimal use of matrix computations. Section 5 introduces models for dependent population units with an emphasis on spatial processes and multivariate dependencies. Ignorable and nonignorable responses are discussed in this context. Section 6 briefly discusses model choice and assessment, while Section 7 presents some key results from analysis of finite population surveys in ignorable and nonignorable contexts. Section 8 concludes the paper with a brief discussion.

2 SRSWOR and Bayesian hierarchical models

In simple random sampling without replacement (SRSWOR) from a population of N units, $p(Z \mid Y) = 1/{N \choose n}$ whenever $\sum_{i=1}^{N} Z_i = n$ and 0 otherwise. Since this distribution does not depend upon Y, we can write p(Y, Z) = p(Y)p(Z) and we seek the posterior distribution $p(Y \mid \mathcal{D}, Z) \propto p(Y)p(\mathcal{D} \mid Y, Z)$. Inference on the population is based solely on the prior distribution for Y, which induces the distribution $p(\mathcal{D} \mid Y, Z)$. If the probability model assumes that the units are exchangeable, then we can reorder the labels as necessary and denote \mathcal{D} by $y = (y_1, y_2, \ldots, y_n)^{\mathrm{T}}$ as the values of the n sampled units and $Y_u = (Y_{n+1}, \ldots, Y_N)^{\mathrm{T}}$ as the unknown values of units not sampled. As a familiar example, consider the simple hierarchical model for population units:

$$Y_i \stackrel{iid}{\sim} N(\mu, \sigma^2), \ i = 1, 2, \dots, N; \quad \mu \sim N(\theta, \sigma^2/n_0); \quad \sigma^2 \sim IG(a_0, b_0) ,$$
 (1)

where θ , n_0 , a_0 and b_0 are known scalars. The joint posterior factorises as

$$p(\mu, \sigma^2 \mid y) \propto \underbrace{IG(\sigma^2 \mid a_1, b_1)}_{p(\sigma^2 \mid y)} \times \underbrace{N\left(\mu \mid \frac{n_0 \theta + n\bar{y}}{n_0 + n}, \frac{\sigma^2}{n_0 + n}\right)}_{p(\mu \mid \sigma^2, y)},$$

where $\bar{y} = \sum_{i=1}^{n} y_i/n$ is the sample mean, $a_1 = a_0 + \frac{n}{2}$ and $b_1 = b_0 + \frac{f(\theta, \bar{y})}{2}$, $f(\theta, \bar{y}) = (n-1)s^2 + \frac{n_0n}{n_0+n}(\bar{y}-\theta)^2$ and $s^2 = \sum_{i=1}^{n} (y_i - \bar{y})^2/(n-1)$. Conditional on σ^2 , it is easy to derive the joint posterior distribution $p(Y_u, \mu \mid \sigma^2, y) = N(\mu \mid Mm, \sigma^2M) \times N(Y_u \mid 1_{N-n}\mu, \sigma^2 I_{N-n})$, where $m = n_0\theta + n\bar{y}$ and $M = 1/(n+n_0)$. The marginal posterior distribution $p(Y_u \mid \sigma^2, y)$ is easily obtained (without requiring explicit integration) by substituting $\mu = Mm + \omega$, where $\omega \sim N(0, \sigma^2M)$, as $N(Y_u \mid 1_{N-n}(n_0\theta + n\bar{y})/(n + n_0), \sigma^2(1_{N-n}1_{N-n}^T/(n+n_0) + I_{N-n}/n))$. These calculations are straightforward in Bayesian statistics and are straightforward to adapt to finite population models. The posterior distribution for any linear function of the population units, say $p(a^{\mathrm{T}}Y \mid \sigma^2, y)$, where $a = (a_1, \ldots, a_N)^{\mathrm{T}}$, is again Normal with mean

$$\sum_{i=1}^{n} a_i y_i + \sum_{i=n+1}^{N} a_i \mathbb{E}[Y_{u,i} \mid \sigma^2, y] = \sum_{i=1}^{n} \left(a_i + \frac{\sum_{i=n+1}^{N} a_i}{n_0 + n} \right) y_i + \frac{n_0 \sum_{i=n+1}^{N} a_i}{n_0 + n} \theta .$$
 (2)

The posterior covariance matrix of Y_u is the sum of $\mathbb{E}[\mathbb{V}(Y_u \mid \sigma^2, \mu, y)]$ and $\mathbb{V}(\mathbb{E}[Y_u \mid \sigma^2, \mu, y])$, which yields $\sigma^2(1_{N-n}1_{N-n}^{\mathrm{T}}/(n+n_0)+I_{N-n})$. Therefore,

$$\mathbb{V}\left(a^{\mathrm{T}}Y_{u} \mid \sigma^{2}, y\right) = \sigma^{2}\left(\frac{\left(\sum_{i=n+1}^{N} a_{i}\right)^{2}}{n+n_{0}} + \sum_{i=n+1}^{N} a_{i}^{2}\right).$$
(3)

For full inference when σ^2 is unknown, we first draw a sample of values of $\sigma^2 \sim IG(a_1, b_1)$ followed by one draw from a Normal distribution with mean and variance given by (2) and (3), respectively, for each drawn value of σ^2 .

For the population total, $a_i = 1$ for each i = 1, 2, ..., N so the mean and variance of $p(\sum_{i=1}^{N} Y_i \mid \sigma^2, y)$ is given by $\sum_{i=1}^{n} (n_0 + N)y_i/(n_0 + n)$ and the variance is $\sigma^2((N-n)(n_0+N)/(n_0+n))$. Letting f = n/N and $f_0 = n_0/N$, the posterior distribution of the population mean, $p(\bar{Y} \mid \sigma^2, y)$, can be expressed as

$$\left[\frac{\bar{Y} - \left\{f\bar{y} + (1-f)\frac{n_0\theta + n\bar{y}}{n_0 + n}\right\}}{\sigma\sqrt{\frac{(1-f)(1+f_0)}{n+n_0}}}\right|\sigma^2, y\right] \sim N(0, 1) .$$
(4)

If the prior distribution on μ becomes vague, i.e., $n_0 \to 0$, then the posterior mean and variance in (4) are \bar{y} and $(1-f)\sigma^2/n$, respectively. Further, as $n_0 \to 0$ and $a_0 \to -1/2$ (corresponds to a uniform prior over the real line for $(\mu, \log \sigma^2)$), the resulting posterior distribution $p(\sigma^2 \mid y) \to IG((n-1)/2, (n-1)s^2/2)$ and the marginal posterior

distribution for the population mean is given by $\left[\frac{\bar{Y}-\bar{y}}{s\sqrt{(1-f)/n}} \mid y\right] \sim T_{n-1}$, where T_{n-1} denotes the Student's *t* distribution with n-1 degrees of freedom. This, numerically, reproduces the inference from the classical theory of design-based estimators while retaining the appealing interpretation of direct probability statements about the population mean and without recourse to the specialised mathematical developments of Central Limit Theorems for SRSWOR sampling from finite populations.

3 Bayesian and Horvitz-Thompson estimates

The posterior distribution of the population total, $N\bar{Y}$, has mean $\sum_{i=1}^{n} y_i/\pi_i$, where $\pi_i = n/N$ is the inclusion probability of the *i*-th unit in the sample. This is precisely the design-based Horvitz-Thompson estimator (Narain, 1951; Horvitz and Thompson, 1952) of the population total. This, then, begs the question whether there exists a Bayesian model that will yield a posterior mean equalling the Horvitz-Thompson estimator. The answer is in the affirmative and it is worth taking a look at a simple model (Little, 2004),

$$Y_i = \pi_i \beta + \pi_i \epsilon_i; \quad \epsilon_i \stackrel{ind}{\sim} N(0,1), \ i = 1, 2, \dots, N; \quad p(\beta) \propto 1 ,$$
(5)

where each π_i denotes the probability of population unit *i* being included in the sample. For now, assume that π_i 's are known. Again, without loss of generality and appropriate relabelling of the population units, we assume that the labels $i = 1, 2, \ldots, n$ represent units that have been sampled. The posterior distribution $p(\beta \mid y)$ is Normal with mean

$$\mathbb{E}[\beta \mid y] = (\pi^{\mathrm{T}} \mathrm{diag}(1/\pi_i^2)\pi)^{-1} \pi^{\mathrm{T}} \mathrm{diag}(1/\pi_i^2)y = (1/n) \sum_{i=1}^n (y_i/\pi_i) + \sum_{$$

where $\pi = (\pi_1, \ldots, \pi_n)^{\mathrm{T}}$ and $\operatorname{diag}(1/\pi_i^2)$ is the $n \times n$ diagonal matrix with elements $1/\pi_i^2$. Therefore, the posterior mean of the population total is

$$\mathbb{E}\left[\sum_{i=1}^{N} Y_{i} \middle| y\right] = \sum_{i=1}^{n} y_{i} + \sum_{i=n+1}^{N} \mathbb{E}[Y_{i} \mid y] = \sum_{i=1}^{n} y_{i} + \sum_{i=n+1}^{N} \pi_{i} \mathbb{E}[\beta \mid y]$$
$$= \sum_{i=1}^{n} y_{i} + \left(\sum_{i=n+1}^{N} \pi_{i}\right) \mathbb{E}[\beta \mid y] = \sum_{i=1}^{n} y_{i} + \left(n - \sum_{i=1}^{n} \pi_{i}\right) \mathbb{E}[\beta \mid y] \quad (6)$$
$$= n\mathbb{E}[\beta \mid y] + \sum_{i=1}^{n} (y_{i} - \pi_{i}\mathbb{E}[\beta \mid y]) = \sum_{i=1}^{n} (y_{i}/\pi_{i}) + \sum_{i=1}^{n} (y_{i} - \hat{y}_{i}) ,$$

where we have used the fact that $\sum_{i=1}^{N} \pi_i = \sum_{i=1}^{N} \mathbb{E}[Z_i] = \mathbb{E}\left[\sum_{i=1}^{N} Z_i\right] = n$ in the fourth equality and $\hat{y}_i = \mathbb{E}[\pi_i\beta \mid y] = (\pi_i/n) \sum_{j=1}^{n} (y_j/\pi_j)$ in the last equality. The first term in the last expression is the Horvitz-Thompson estimator, while the second term is a sum of the differences between the observed and model-predicted

values of the sampled units. Letting n approach the population size, i.e., if the entire population is sampled, the second term vanishes and the posterior mean is the Horvitz-Thompson estimator. This is a model where the Bayesian estimate of the population total approximates the Horvitz-Thompson estimator.

To derive a model where the posterior expectation of the population total is *exactly* the Horvitz-Thompson estimator, we modify (5) as follows,

$$Y_i = \pi_i \beta + \pi_i \epsilon_i; \quad \epsilon_i \stackrel{ind}{\sim} N(0, \sigma_i^2), \ i = 1, 2, \dots, N; \quad p(\beta) \propto 1 ,$$
(7)

where the σ_i^2 's are assumed to be fixed constants. The posterior mean of β is

$$\mathbb{E}[\beta \mid y] = (\pi^{\mathrm{T}} \mathrm{diag}(1/(\sigma_i \pi_i)^2)\pi)^{-1} \pi^{\mathrm{T}} \mathrm{diag}(1/(\sigma_i \pi_i)^2)y = \sum_{i=1}^n w_i(y_i/\pi_i) ,$$

where $w_i = (1/\sigma_i^2) / (\sum_{i=1}^n 1/\sigma_i^2)$. Proceeding analogous to (6), we again have

$$\mathbb{E}\left[\left|\sum_{i=1}^{N} Y_{i}\right| y\right] = \sum_{i=1}^{n} y_{i} + \left(\sum_{i=n+1}^{N} \pi_{i}\right) \mathbb{E}[\beta \mid y] = \sum_{i=1}^{n} y_{i} + \left(\sum_{i=n+1}^{N} \pi_{i}\right) \left(\sum_{i=1}^{n} w_{i}(y_{i}/\pi_{i})\right)$$
$$= \sum_{i=1}^{n} \left(1 + \left(n - \sum_{i=1}^{n} \pi_{i}\right) w_{i}/\pi_{i}\right) y_{i} = \sum_{i=1}^{n} \tilde{w}_{i}y_{i},$$
(8)

where $\tilde{w}_i = 1 + (n - \sum_{i=1}^n \pi_i) w_i/\pi_i$. We will obtain the Horvitz-Thompson estimator in the last expression if $\tilde{w}_i = 1/\pi_i$. Letting $\sigma_i^2 = 1/(1 - \pi_i)$ yields $w_i = (1 - \pi_i)/(n - \sum_{i=1}^n \pi_i)$ so that $\tilde{w}_i = 1 + (1 - \pi_i)/\pi_i = 1/\pi_i$. Therefore, setting $\sigma_i^2 = 1/(1 - \pi_i)$ in (7) yields a Bayesian model where the posterior expectation of the population total exactly agrees with the Horvitz-Thompson estimator (Ghosh and Sinha, 1990). See Ghosh (2012) for generalisations of this result to exponential families and other families of priors.

4 Bayesian multistage sampling

Multistage sampling designs draw samples from the population in multiple stages. The population is assumed to be composed of primary (level 1) units from which a random sample is drawn using SRSWOR. Next, for each of these sampled primary units, a random sample of secondary (level 2) units is drawn and, if needed, this process is continued to further sample from the secondary units. Consider 2-stage sampling. Let i = 1, 2, ..., N be the labels for the primary population units and let $j = 1, 2, ..., M_i$ be the number of secondary units within the *i*-th primary unit. Consider a sampling design that draws *n* primary units using SRSWOR and, then, for each of these primary units draws a sample of m_i secondary unit units using SRSWOR. If $Y_{(i,j)}$ denotes the population unit referenced by secondary unit *j* within primary unit *i*, then the Horvitz-Thompson estimator for the population total is $\sum_{i,j=1}^{n,m_i} y_{(i,j)}/\pi_{i,j}$,

where $\pi_{i,j} = (n/N)(m_i/M_i) = P(Z_{(i,j)} = 1)$, where $Z_{(i,j)} = 1$ if the secondary unit j within primary unit i is sampled and $Z_{(i,j)} = 0$ otherwise.

4.1 Bayesian hierarchical model for 2-stage sampling

A Bayesian model for 2-stage sampling reproducing the Horvitz-Thompson estimator as its posterior mean is easily derived from (7) by writing $Y_{(i,j)} = \pi_{(i,j)}\beta + \pi_{(i,j)}\epsilon_{(i,j)}$, where $\epsilon_{(i,j)} \stackrel{ind}{\sim} N(0, 1/(1 - \pi_{(i,j)}))$. However, richer classes of models arise from Bayesian hierarchical models for 2-stage sampling (Scott and Smith, 1969), while Malec and Sedransk (1985) discuss Bayesian inference for more general multistage sampling. Here, we will take a closer look at the 2-stage model but in a more contemporary context of "BIG DATA" settings, where the CPU storage and memory allocations will not allow us to deal with data of size exceeding $O(\max_{1 \le i \le N} m_i)$. Bayesian analysis will, therefore, have to proceed sequentially as samples stream in using a "divide and conquer" mechanism.

Let us turn to a 2-stage sampling design in the context of a finite population of massive size, where resources allow us to sample only n out of N population units and data of size no more than $\max_{1 \le i \le n} m_i$ units for analysis. We consider the model

$$Y_{ij} \stackrel{ind}{\sim} N(\beta_i, \delta^2 / \gamma_i^2), \ j = 1, 2, \dots, M_i;$$

$$\beta_i \stackrel{ind}{\sim} N(\nu, \delta^2), \ i = 1, 2, \dots, N;$$

$$\nu \sim N(c_0, \delta^2 / n_0); \quad \delta^2 \sim IG(a_0, b_0) ,$$

(9)

where γ_i^2 , n_0 , c_0 , a_0 and b_0 are fixed scalars. Suppose, with legitimate relabelling of the population units that is allowed due to exchangeable random variables in (9), we sample i = 1, 2, ..., n primary units from the population and $j = 1, 2, ..., m_i$ secondary units from the *i*-th primary unit. Let $y_i = (y_{i1}, ..., y_{im_i})^T$ be the $m_i \times 1$ vector of sampled observations from the *i*-th primary unit in the population. The first equation in (9) yields $y_i = 1_{m_i}\beta_i + \epsilon_i$, where 1_{m_i} is the $m_i \times 1$ vector of ones, β_i is the scalar parameter corresponding to the mean of the *i*-th primary unit, and $\epsilon_i \stackrel{ind}{\sim} N\left(0, (\delta^2/\gamma_i^2)I_{m_i}\right)$ with I_{m_i} denoting the $m_i \times m_i$ identity matrix. Substituting the distribution of β_i in the second equation of (9) results in integrating out β_i from the model and we obtain $p(y_i \mid \nu, \delta^2)$ as the model $y_i = 1_{m_i}\nu + \omega_i$, where $\omega_i \stackrel{ind}{\sim} N\left(0, \delta^2 V_i\right)$, where $V_i = (1/\gamma_i^2)I_{m_i} + 1_{m_i}1_{m_i}^{\mathrm{T}}$. To optimise storage and computations we follow sequential Bayesian updating where the posterior distribution is updated using data from one primary unit only at each iteration. The resulting posterior distribution then acts as the prior distribution to be updated using data from the next primary unit.

4.2 Inference for $\{\nu, \delta^2\}$

To illustrate in the context of (9), we first derive the posterior distribution $p(\nu, \delta^2 \mid y_1) \propto IG(\delta^2 \mid a_0, b_0) \times N(\nu \mid c_0, \delta^2/n_0) \times N(y_1 \mid 1_{m_i}\nu, \delta^2 V_1)$, which, after standard

Bayesian Normal-Normal calculations, yields

$$p(\nu, \delta^2 \mid y_1) = \frac{IG(\delta^2 \mid a_1, b_1)}{p(\delta^2 \mid y_1)} \times \underbrace{\frac{N\left(\nu \mid C_1c_1, \delta^2C_1\right)}{p(\nu \mid \delta^2, y_1)}}_{p(\nu \mid \delta^2, y_1)}$$

where $c_1 = n_0 c_0 + 1_{m_1}^{T} V_1^{-1} y_1$, $C_1^{-1} = n_0 + 1_{m_1}^{T} V_1^{-1} 1_{m_1}$, $a_1 = a_0 + m_1/2$ and $b_1 = b_0 + q_1/2$ with $q_1 = n_0 c_0^2 + y_1^{T} V_1^{-1} y_1 - c_1^2 C_1$. Simplifications arise from applying the Sherman-Morrison matrix identity to obtain

$$V_1^{-1} = \gamma_1^2 I_{m_1} - \frac{\gamma_1^2 \mathbf{1}_{m_1} \mathbf{1}_{m_1}^{\mathrm{T}} \gamma_1^2}{1 + m_1 \gamma_1^2} = \gamma_1^2 \left(I_{m_1} - \frac{\gamma_1^2}{1 + m_1 \gamma_1^2} \mathbf{1}_{m_1} \mathbf{1}_{m_1}^{\mathrm{T}} \right).$$
(10)

Writing $\lambda_1 = m_1 \gamma_1^2 / (1 + m_1 \gamma_1^2)$, we obtain the following simplified expressions: $c_1 = n_0 c_0 + \lambda_1 \bar{y}_1$ and $C_1^{-1} = n_0 + \lambda_1$, which yields $\mathbb{E}[\nu \mid y_1] = n_0 / (n_0 + \lambda_1) c_0 + \lambda_1 / (n_0 + \lambda_1) \bar{y}_1$. Treating this as the base case, we apply induction to conclude for each $t = 1, 2, \ldots, n$

$$p(\nu, \delta^2 \mid y_1, \dots, y_t) = \frac{IG(\delta^2 \mid a_t, b_t)}{p(\delta^2 \mid y_1, \dots, y_t)} \times \frac{N\left(\nu \mid C_t c_t, \delta^2 C_t\right)}{p(\nu \mid \delta^2, y_1, \dots, y_t)},$$
(11)

where $c_t = c_{t-1} + 1_{m_t}^{\mathrm{T}} V_t^{-1} y_t$, $C_t^{-1} = C_{t-1}^{-1} + 1_{m_t}^{\mathrm{T}} V_t^{-1} 1_{m_t}$, $a_t = a_{t-1} + m_t/2$ and $b_t = b_{t-1} + q_t/2$ with $q_t = c_{t-1}^2/C_{t-1} + y_t^{\mathrm{T}} V_t^{-1} y_t - c_t^2 C_t$. Applying (10) to V_t^{-1} again yields $c_t = c_{t-1} + \lambda_t \bar{y}_t$ and $C_t^{-1} = C_{t-1} + \lambda_t$. In particular, for each $t = 1, \ldots, n$

$$\mathbb{E}[\nu \mid \delta^2, y_1, \dots, y_t] = \frac{n_0}{n_0 + \sum_{i=1}^t \lambda_i} c_0 + \sum_{i=1}^t \frac{\lambda_i}{n_0 + \sum_{i=1}^t \lambda_i} \bar{y}_i .$$
(12)

Since $\mathbb{E}[\nu \mid \delta^2, y_1, \dots, y_t]$ does not depend on δ , (12) is also the value of the marginal posterior mean $\mathbb{E}[\nu \mid y_1, \dots, y_t]$. Furthermore, $\mathbb{V}(\mathbb{E}[\nu \mid \delta^2, y_1, \dots, y_t]) = 0$ with respect to the density $p(\delta^2 \mid y_1, \dots, y_t)$ and, hence, the marginal posterior variance of ν is

$$\mathbb{V}(\nu \mid y_1, \dots, y_t) = \frac{b_t}{a_t - 1} C_t = \frac{b_0 + \sum_{i=1}^t q_i/2}{\left(a_0 + \sum_{i=1}^t m_i/2 - 1\right) \left(n_0 + \sum_{i=1}^t \lambda_i\right)} .$$
(13)

4.3 Inference for $\{\beta_1, \ldots, \beta_N\}$

Turning to inference for $\beta = (\beta_1, \beta_2, \dots, \beta_N)^{\mathrm{T}}$, note that

$$p(\beta_t \mid \nu, \delta^2, y_t) \propto N(\beta_t \mid \nu, \delta^2) \times N(y_t \mid 1_{m_t} \beta_t, \delta^2 / \gamma_t^2)$$

= $N(\beta_t \mid C_{\beta_t} c_{\beta_t}, \delta^2 C_{\beta_t}) = N(\beta_t \mid (1 - \lambda_t)\nu + \lambda_t \bar{y}_t, \delta^2 (1 - \lambda_t))$

where the last equality follows from $c_{\beta_t} = \nu + \gamma_1^2 m_t \bar{y}_t = \nu + (\lambda_t/(1-\lambda_t)) \bar{y}_t$ and $C_{\beta_1}^{-1} = 1 + m_t \gamma_t^2 = 1/(1-\lambda_t)$. The posterior distribution $p(\beta_1, \ldots, \beta_t \mid \nu, \delta^2, y_1, \ldots, y_t)$

for each $t = 1, \ldots, n$ is proportional to

$$\underbrace{\prod_{i=1}^{t-1} N(\beta_i \mid 1 - \lambda_i)\nu + \lambda_i \bar{y}_i, \delta^2(1 - \lambda_i))}_{p(\beta_1, \dots, \beta_{t-1} \mid \nu, \delta^2, y_1, \dots, y_{t-1})} \times \underbrace{N(\beta_t \mid \nu, \delta^2)}_{p(\beta_t \mid \nu, \delta^2)} \times \underbrace{N(y_t \mid \beta_t, \delta^2 \gamma_t^2)}_{p(y_t \mid \beta_t, \delta^2)} \\ \propto \underbrace{\prod_{i=1}^{t-1} N(\beta_i \mid 1 - \lambda_i)\nu + \lambda_i \bar{y}_i, \delta^2(1 - \lambda_i))}_{p(\beta_1, \dots, \beta_{t-1} \mid \nu, \delta^2, y_1, \dots, y_{t-1})} \times \underbrace{N(\beta_t \mid (1 - \lambda_t)\nu + \lambda_t \bar{y}_t, \delta^2(1 - \lambda_t))}_{p(\beta_t \mid \nu, \delta^2, y_t)}.$$
(14)

Therefore, $p(\beta_1, \ldots, \beta_t \mid \nu, \delta^2, y_1, \ldots, y_t) = \prod_{i=1}^t N(\beta_i \mid (1 - \lambda_i)\nu + \lambda_i \bar{y}_i, \delta^2(1 - \lambda_i))$ for each $t = 1, 2, \ldots, n$. For $t = n+1, \ldots, N$, the primary units have not been sampled and, hence, there is no data available to update the priors. Hence, we can write

$$p(\beta \mid \nu, \delta^2, y_1, \dots, y_n) = \prod_{i=1}^N N\left(\beta_i \mid (1 - \lambda_i)\nu + \lambda_i \bar{y}_i, \delta^2(1 - \lambda_i)\right) , \qquad (15)$$

where we define $m_i = 0$ and, hence, $\lambda_i = 0$, for $i = n + 1, \ldots, N$. The marginal posterior distribution $p(\beta \mid \delta^2, y_1, \ldots, y_n)$ is easily obtained by writing (15) as $\beta = (1_N - \lambda)\nu + \Lambda \bar{y} + \omega$, where $1_N - \lambda = (1 - \lambda_1, \ldots, 1 - \lambda_N)^{\mathrm{T}}$ is $N \times 1$, $\Lambda = \mathrm{diag}(\lambda_1, \ldots, \lambda_N)$ is the $N \times N$ diagonal matrix with elements $1 - \lambda_i$, $\bar{y} = (\bar{y}_1, \ldots, \bar{y}_n, 0, \ldots, 0)^{\mathrm{T}}$ is $N \times 1$ with the last N - n terms being arbitrarily set to 0 and $\omega \sim N(0, \delta^2(I_N - \Lambda))$. Therefore,

$$\beta \mid \delta^2, y_1, \dots, y_n \sim N\left((1_N - \lambda)\bar{y}_{\cdot} + \Lambda \bar{y}, \delta^2(I_N - \Lambda) + \eta^2(1_N - \lambda)(1_N - \lambda)^{\mathrm{T}}\right)\right), \quad (16)$$

where $\bar{y}_{\cdot} = \mathbb{E}[\nu \mid \delta^2, y_1, \dots, y_n]$ given in (12) and $\eta^2 = \delta^2 / (\sum_{i=1}^N \lambda_i)$.

We now have the necessary distributions to sample from the posterior distribution of the model parameters $\{\delta^2, \nu, \beta\}$. We draw a sample from $p(\nu, \delta^2 \mid y_1, \ldots, y_n)$ as described earlier and draw a value of β from (15) for each drawn value of $\{\delta^2, \nu\}$. The resulting samples of β are draws from $p(\beta \mid y_1, \ldots, y_n)$. Note that this is computationally more efficient than drawing samples of β from (16) for each drawn δ^2 since drawing from (16) will require expensive matrix computations involving the $N \times N$ covariance matrix, where N may exceed our data processing capabilities.

4.4 Inference for all unobserved population units

We now turn to inference for the unobserved population units. Let $Y_i = (Y_{i,m_i+1},\ldots,Y_{i,M_i})^{\mathrm{T}}$ for each $i = 1, 2, \ldots, N$ and let $Y_u = (Y_1^{\mathrm{T}},\ldots,Y_N^{\mathrm{T}})^{\mathrm{T}}$ be the vector of all unobserved population units, where we are using the definition $m_i = 0$ for $i = n + 1, \ldots, N$. The random variable Y_u is independent of the observed data given $\{\beta, \delta^2\}$ and its full conditional distribution is

$$p(Y_u \mid \beta, \delta^2) = \prod_{i=1}^{N} N\left(Y_i \mid 1_{M_i - m_i}\beta_i, (\delta^2 / \gamma_i^2) I_{M_i - m_i}\right) .$$
(17)

Drawing samples from the posterior distribution $p(Y_{ij} | y_1, \ldots, y_n) = \int p(Y_{ij} | \beta_i, \delta^2) \times p(\beta_i, \delta^2 | y_1, \ldots, y_n)$ for any unobserved population unit (i, j) now proceeds by drawing one value of $Y_{ij} \sim N(\beta_i, \delta^2/\gamma_i^2)$ for each value of the posterior samples of β_i and δ^2 . Again, no matrices are involved in this sampling step.

Posterior inference for finite population quantities such as $\Omega = \sum_{i=1}^{N} \sum_{j=1}^{M_i} a_{ij} Y_{ij}$ is straightforward. We split up the sum as $\sum_{i=1}^{n} \sum_{j=1}^{m_i} a_{ij} y_{ij} + \sum_{i=1}^{N} \sum_{j=m_i+1}^{M_i} a_{ij} Y_{ij}$ and obtain posterior samples of this quantity by simply substituting the sampled values of Y_{ij} in the second sum. Explicit algebraic expressions for the posterior mean and variance are not required for computations. Nevertheless, we have all the ingredients to obtain these moments. Let $a_i = \sum_{j=m_i+1}^{M_i} a_{ij}$ and recall that $\lambda_i = 0$ for $i = n+1, \ldots, n$. Since $\mathbb{E}[Y_{ij} \mid y_1, \ldots, y_n] = \mathbb{E}[\beta_i \mid y_1, \ldots, y_n] = (1 - \lambda_i)\overline{y}_i + \lambda_i \overline{y}_i$, the finite population estimate of Ω in the limit as $n_0 \to 0$ in (12) (given in Scott and Smith, 1969) is obtained as

$$\mathbb{E}\left[\Omega \mid y_{1}, \dots, y_{n}\right] = \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} a_{ij}y_{ij} + \sum_{i=1}^{N} \sum_{j=m_{i}+1}^{M_{i}} a_{ij}\mathbb{E}\left[Y_{ij} \mid y_{1}, \dots, y_{n}\right]$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} a_{ij}y_{ij} + \sum_{i=1}^{N} a_{i}\left\{(1-\lambda_{i})\bar{y}_{.} + \lambda_{i}\bar{y}_{i}\right\}$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} a_{ij}y_{ij} + \sum_{i=1}^{N} a_{i}\lambda_{i}\bar{y}_{i} + \left(\sum_{i=1}^{N} a_{i}(1-\lambda_{i})\right)\bar{y}_{.}$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} a_{ij}y_{ij} + \sum_{i=1}^{n} \frac{a_{i}\lambda_{i}}{m_{i}} \sum_{j=1}^{m_{i}} y_{ij} + \frac{\sum_{i=1}^{N} a_{i}(1-\lambda_{i})}{\sum_{i=1}^{N} \lambda_{i}} \sum_{i=1}^{n} \frac{\lambda_{i}}{m_{i}} \sum_{j=1}^{m_{i}} y_{ij}$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} \left\{a_{ij} + a_{i}\frac{\lambda_{i}}{m_{i}} + \frac{\sum_{i=1}^{N} a_{i}(1-\lambda_{i})}{\sum_{i=1}^{N} \lambda_{i}} \frac{\lambda_{i}}{m_{i}}\right\} y_{ij}$$

$$= \sum_{i=1}^{n} \sum_{j=1}^{m_{i}} \left\{a_{ij} + (a_{i} + k)\frac{\lambda_{i}}{m_{i}}\right\} y_{ij}, \text{ where } k = \frac{\sum_{i=1}^{N} a_{i}(1-\lambda_{i})}{\sum_{i=1}^{N} \lambda_{i}}.$$
(18)

The posterior estimate of the population total is obtained from (18) by setting $a_{ij} = 1$, which implies $a_i = M_i - m_i$. The relationship between the Bayesian estimate and several design-based estimators is explored in Scott and Smith (1969). The Horvitz-Thompson estimator does not emerge naturally from the posterior mean. Nevertheless, some simplifications are possible. If we further assume that m_i and γ_i^2 vary in a manner that $\lambda_i = \lambda$ for each primary unit, then (18) yields

$$\sum_{i=1}^{n} \sum_{j=1}^{m_i} \left\{ 1 - \lambda + \lambda \frac{M_i}{m_i} + (1 - \lambda) \frac{\bar{a}}{m_i} \right\} y_{ij} = \lambda \sum_{i=1}^{n} M_i \bar{y}_i + (1 - \lambda) \sum_{i=1}^{n} \left\{ 1 + \frac{\bar{a}}{m_i} \right\} m_i \bar{y}_i ,$$

where $\bar{a} = \sum_{i=1}^{N} a_i/N$. In the limit as $\lambda \to 1$, the above expression equals $\sum_{i=1}^{N} M_i \bar{y}_i$. Hence, a simple modification of (9) with $Y_{ij} \stackrel{ind}{\sim} N(x_i \beta_i, \delta^2/\gamma_i^2)$ and each $x_i = n/N$

for i = 1, 2, ..., N will yield a posterior mean $(N/n) \sum_{i=1}^{N} M_i \bar{y}_i$ in the limit as $\lambda \to 1$, which is precisely the Horvitz-Thompson estimator.

4.5 Conjugate Bayesian regression in "BIG DATA" settings

It is worth remarking, more generally, that conjugate Bayesian linear regression analysis renders itself as an extremely viable option for massive data sets. If the size of the data exceeds available memory capabilities and cannot be accessed in its entirety due to memory allocations, we split up the data into n mutually exclusive blocks, each of size m_i , so that $\sum_{i=1}^{n} m_i$ is the total number of observations. This partitioning of the data is performed in a manner so that our memory resources are capable of analysing the individual data blocks. For example, consider the model

$$y_i \stackrel{ind}{\sim} N(X_i\beta, \sigma^2 V_i), \ i = 1, \dots, n; \quad \beta \mid \sigma^2 \sim N(C_0 c_0, \sigma^2 C_0); \quad \sigma^2 \sim IG(a_0, b_0),$$
(19)

where each y_i is $m_i \times 1$, X_i is an $m_i \times p$ matrix consisting of the values of known explanatory variables, V_i is an $m_i \times m_i$ positive definite covariance matrix, β is the $p \times 1$ vector of slopes that is assigned a *p*-variate multivariate Gaussian prior specified using a $p \times p$ matrix C_0 and $p \times 1$ vector c_0 and σ^2 is the residual variance. Sequential Bayesian updating uses the simple observation that $p(\beta, \sigma^2 \mid y_1, \ldots, y_t) \propto p(\beta, \sigma^2 \mid y_1, \ldots, y_{t-1}) \times p(y_t \mid \beta, \sigma^2)$. This yields the joint posterior for each $t = 1, 2, \ldots, n$ as

$$p(\beta, \sigma^2 \mid y_1, \dots, y_t) = \frac{IG(\sigma^2 \mid a_t, b_t)}{p(\sigma^2 \mid y_1, \dots, y_t)} \times \frac{N\left(\beta \mid C_t c_t, \sigma^2 C_t\right)}{p(\beta \mid \sigma^2, y_1, \dots, y_t)} , \qquad (20)$$

where $a_t = a_{t-1} + m_t/2$, $b_t = b_{t-1} + q_t/2$, $q_t = y_t^{\mathrm{T}} V_t^{-1} y_t + c_{t-1}^{\mathrm{T}} C_{t-1} c_{t-1} - c_t^{\mathrm{T}} C_t c_t$, $c_t = c_{t-1} + X_t^{\mathrm{T}} V_t^{-1} y_t$ and $C_t^{-1} = C_{t-1}^{-1} + X_t^{\mathrm{T}} V_t^{-1} X_t$. Since each step now involves only storage and computation in the order of at most $O(m^3)$ flops, where $m = \max_{1 \le t \le n} m_t$, we obtain our desired posterior distribution at t = n by applying (19) to one block of data in each iteration and overwriting these objects as we move to the next iteration. Posterior inference for an unobserved population unit Y_{ij} that belongs to block i will only access information from that block. For each sampled values β and σ^2 , we will draw one value $Y_{ij} \sim N(X_i\beta, \sigma^2 V_i)$, which results in a draw from $p(Y_{ij} \mid y_1, \ldots, n)$.

4.6 Extensions

Extending the above models to three or higher stages is straightforward, although the algebra becomes more tedious. In practice, we can conveniently build hierarchical multistage models in programming environments such as BUGS (Lunn et al, 2009), rstan (Stan Development Team, 2024) and nimble (de Valpine et al, 2017, 2023). As we discussed in the preceding section, the conditional independence across primary units accrue significant computational benefits when dealing with large, or even massive, data sets since we can exploit the sequential nature of Bayesian updating in (20). Section 5 turns to multivariate models to infer on populations with spatially dependent units.

5 Dependent population models

A key advantage of the model-based approach to survey sampling is its ability to incorporate information, either known or posited, about the population units. Subsequent inference about the finite population quantities will follow from the probability distribution of unobserved quantities while accounting for the dependence structure in the population. Here, it will be instructive to recall the modelling framework promulgated by Rubin (1976); also see Gelman et al (2014). As in Section 1, let $Y = (y_1, \ldots, y_N)^{\mathrm{T}}$ be the units in the population and let $Z = (Z_1, \ldots, Z_N)^{\mathrm{T}}$ be the sampling inclusion indicators. A probability model for the entire finite population is given by $p(Y, Z \mid \theta, \phi) = p(Y \mid \theta) \times p(Z \mid Y, \phi)$, where θ and ϕ are unknown parameters specifying the distribution of Y and the conditional distribution of Z given Y, respectively. We sample from

$$p(Y_u \mid y, Z) \propto \int p(Y_u \mid y, Z, \theta, \phi) \times p(\theta, \phi \mid y, Z) \, d\theta \, d\phi \,, \tag{21}$$

where $Y = \{Y_u, y\}$ with y denoting observed samples and Y_u unknown values in population units excluded in the sampling design. In order to sample from (21), we note that $p(Y_u, \theta, \phi \mid y, Z) = p(Y_u \mid y, Z, \theta, \phi) \times p(\theta, \phi \mid y, Z)$. Therefore, we first sample values of θ and ϕ from their joint posterior distribution $p(\theta, \phi \mid y, Z)$ and, subsequently, we draw one value of Y_u from $p(Y_u \mid y, Z, \theta, \phi)$ using each sampled value of $\{\theta, \phi\}$.

5.1 Ignorable sampling designs

A crucial observation is that the conditional distribution $p(Z | Y, \phi)$ models the sampling design and is accounted for in the finite population inference. For example, in the models considered in Sections 2–4 $p(Z | y, \phi) = p(Z)$ does not depend on the sampled values and there are no unknown design parameters ϕ . Only the sizes of the population and sample are used, which means that information about the design does not transmit to inference on the unknown population values and, hence, these designs are referred to as *ignorable*. As seen in the aforementioned sections, purely modelbased approaches can reproduce the Horvitz-Thompson estimator without requiring any information on the sampling design.

Inference for finite population units from an ignorable design is, therefore, essentially a model-based prediction or Bayesian imputation problem. Linear regression models with dependent population units modify (19) and formulated as

$$\begin{bmatrix} y\\ Y_u \end{bmatrix} = \begin{bmatrix} X_s\\ X_u \end{bmatrix} \beta + \begin{bmatrix} \epsilon_s\\ \epsilon_u \end{bmatrix}; \quad \begin{bmatrix} \epsilon_s\\ \epsilon_u \end{bmatrix} \sim N\left(\begin{bmatrix} 0\\ 0 \end{bmatrix}, \begin{bmatrix} V_s(\theta) & V_{su}(\theta)\\ V_{us}(\theta) & V_u(\theta) \end{bmatrix}\right);$$

$$\beta = A\mu + \eta; \quad \eta \sim N(0, V_\beta); \quad \theta \sim p(\theta),$$
(22)

where X_s and X_u are fixed design matrices corresponding to sampled and non-sampled units, respectively, $\begin{bmatrix} V_s(\theta) & V_{su}(\theta) \\ V_{us}(\theta) & V_u(\theta) \end{bmatrix}$ is a positive definite covariance matrix partitioned accordingly, and A and μ are assumed fixed. Inference proceeds from the following

steps: (i) Draw samples from the posterior distribution $p(\beta, \theta \mid y)$; (ii) Draw samples from $p(Y_u \mid y)$ so that $\mathbb{E}_{\beta,\theta\mid y} \left[N\left(Y_u \mid X_u\beta + V_{us}V_s^{-1}(y - X_s\beta), V_u - V_{us}V_s^{-1}V_{su} \right) \right]$. The posterior samples Y_u and the observations y are then substituted into desired functions of the population units to obtain the posterior samples of the desired functions. The posterior estimator obtained from (22) is consistent in the sense that $\mathbb{E}[Y_u \mid y, \beta, \theta] = X_u\beta + V_{us}V_s^{-1}(y - X_s\beta)$ equals y if the entire finite population is sampled whereupon $X_u = X_s$ and $V_{us} = V_s$ and the expression becomes free of model parameters $\{\beta, \theta\}$. Furthermore, the conditional variance $\mathbb{V}(Y_u \mid y, \beta, \theta)$ equals 0 when the finite population is sampled implying that the posterior estimator equals the finite population quantity with no uncertainty. In spatial contexts, which we discuss below, this reveals an interpolation property for finite populations using (22).

5.2 Spatial structured populations

A contemporary problem in survey sampling concerns estimation in spatially oriented finite populations, where each unit emerges as a (partial) realisation of a spatial process. In this domain, there is a substantial literature on small area estimation for regionally aggregated data (see, e.g., Rao and Molina, 2015; Clayton and Kaldor, 1987; Datta and Ghosh, 1991; Ghosh and Rao, 1994; Arora and Lahiri, 1997; Ghosh et al, 1998), where interest lies in modelling dependencies across regions such as counties or districts within states, states within a country, or census-tracts or postal codes within a state or city. The population units correspond to regions are assumed to be correlated through a Markov Random Field positing that each region is conditionally dependent only on its neighbours, given all other regions. Conditional Auto-Regression (CAR) models are popular in small area estimation to model such dependence; see the aforementioned literature.

Modelling considerations differ from CAR models when we consider quantities that, at least conceptually, exist in continuum over the entire domain, although they are recorded at a finite number of point-referenced locations. Now, dependence among the population units is induced from a spatial process, which assigns a probability law to an uncountable subset within a *d*-dimensional Euclidean domain. In general, spatial process modelling (Banerjee et al 2014; Cressie and Wikle 2011; and Ripley 2004) follows the generic paradigm

$$[data \mid process] \times [process \mid parameters] \times [parameters], \qquad (23)$$

which accommodates complex dependencies and multiple sources of variation. The literature on finite population sampling in spatial process settings appears to be considerably sparser than small area estimation. Here, Ver Hoef (2002) discuss connections between geostatistical models and classical design-based sampling and develop methods for executing model-based block "kriging". Cicchitelli and Montanari (2012) present a spline regression model-assisted, design-based estimator of the mean for use on a random sample from both finite and infinite spatial populations, while Bruno et al (2013) use linear spatial interpolation to create a design-based predictor of values at unobserved locations.

For ignorable designs, the framework in (22) is often sufficient. Specifically in spatial contexts, the elements of the covariance matrix in (22) are the values of a positive definite covariance function. Several choices for the correlation function are available (see, e.g. Banerjee et al, 2014). The isotropic Matérn function, which models spatial covariance as a function of distance between spatial locations, while also introducing a parameter to control smoothness of process realisations, is defined as $C(d_{ab}) = \sigma^2 + \tau^2$ if $d_{ab} = 0$ and $C(d_{ab}) = \tau^2 \frac{2^{1-\eta}}{\Gamma(\eta)} \left(\sqrt{2\eta} d_{ab} \phi\right)^{\eta} K_{\eta} \left(\sqrt{2\eta} d_{ab} \phi\right)$ if $d_{ab} > 0$, where $K_{\eta}(\cdot)$ is the modified Bessel function and d_{ab} is the distance between two locations ℓ_a and ℓ_b . The process parameters $\theta = \{\sigma^2, \tau^2, \phi, \eta\}$ capture variation due to measurement error or micro-resolution variation, variation attributable to spatial structure, the rate of decline in spatial association, and smoothness of the process, respectively. The exponential covariance function $C(d_{ab}) = \tau^2 \exp(-\phi d_{ab})$ is a special case of Matérn when $\eta = 1/2$.

5.3 Non-ignorable spatial designs

A more general framework for spatial survey sampling that accommodates nonignorable sampling designs is built from the approach surrounding (21). We introduce some further notations. Formally, define a spatial domain $\mathcal{L} \subseteq \mathbb{R}^2$ and let $\mathcal{L}_{FP} = \{\ell_1, \ldots, \ell_N\}$ denote a finite collection of N spatial locations. This set represents the units in our finite population and is made up of locations that are sampled, denoted by \mathcal{L}_s , and those that are not, denoted by \mathcal{L}_u ; therefore, $\mathcal{L}_s \cup \mathcal{L}_u = \mathcal{L}_{FP}$. In addition, certain spatial applications will require a "structural zero" subset, denoted by $\mathcal{L}_0 \subset \mathcal{L} \setminus \mathcal{L}_{FP}$, which represents all locations where sampling is precluded due to geographic considerations. For example, points situated over lakes and rivers may be precluded from the finite population of interest when one is interested in household economic surveys. Likewise, heavily urbanised metropolitan areas are precluded from studies concerning biomass interpolation from a finite population of trees in a forest.

It is of interest to learn about the process in (23) from the sampled observations. A spatial process endows a probability law over the uncountable collection of random variables $\{(Y(\ell), Z(\ell)) : \ell \in \mathcal{L}\}$ and, hence, induces one for the finite population $\{(Y(\ell), Z(\ell)) : \ell \in \mathcal{L}_{FP}\}$. We consider three underlying stochastic processes generating the data: (i) a possibly vector-valued latent spatial process, $\{w(\ell) : \ell \in \mathcal{L}\}$, which models the variable(s) of interest at ℓ ; (ii) a sampling indicator process, $\{Z(\ell) : \ell \in \mathcal{L}\}$, which indicates whether a location is sampled or not; and possibly (iii) a structural zero indicator process (partially observed), $\{u(\ell) : \ell \in \mathcal{L}\}$, which indicates whether $\ell \in \mathcal{L}_{FP}$ or not. A joint hierarchical model, conceptually, is constructed as

$$[Y(\cdot) \mid w(\cdot), u(\cdot)] \times [Z(\cdot) \mid Y(\cdot), u(\cdot)] \times [w(\cdot) \mid u(\cdot)] \times [u(\cdot)] .$$

$$(24)$$

We turn to specifying probability models for each of these components in the spirit of Finley et al (2011), who predicted continuous forest variables at new locations accounting for uncertainty in whether the new locations were forested or not.

5.3.1 Modelling $[Y(\cdot) \mid w(\cdot), u(\cdot)]$

Let $u(\ell) = 1$ if $\ell \in \mathcal{L}_{FP}$ and $u(\ell) = 0$ if $\ell \notin \mathcal{L}_{FP}$. The first component in (24) is modelled as a customary spatial (or spatial-temporal) regression model for point-referenced data if $u(\ell) = 1$ and is distributed as the Dirac measure, δ_0 , with point mass at 0 if $u(\ell) = 0$. Thus,

$$Y(\ell) = u(\ell) \{ x(\ell)^{\mathrm{T}} \beta + \tilde{x}(\ell)^{\mathrm{T}} w(\ell) + \epsilon(\ell) \} + (1 - u(\ell)) \delta_0 ; \quad \epsilon(\ell) \stackrel{iid}{\sim} N(0, \sigma^2)$$

where $x(\ell)$ is a vector of explanatory (perhaps including design) variables at ℓ with regression coefficients β , $\tilde{x}(\ell)$ is a subset of $x(\ell)$ whose regression coefficients, $w(\ell)$, vary over space, that is, the impact of the predictors on the population units vary over space and $w(\ell)$ is either a multivariate Gaussian process with a valid cross-covariance matrix or a vector of independent univariate processes (see, e.g., Banerjee et al, 2014) or using matrix-variate normal distributions (Zhang and Banerjee, 2022). The white noise process $\epsilon(\ell)$ captures micro-scale variation existing at fine scales (smaller than the minimum distances among locations in the finite population) that is attributed to measurement errors in surveys.

5.3.2 Modelling $[Z(\cdot) \mid Y(\cdot), \omega(\cdot), u(\cdot)]$

While the sampling indicator process $Z(\cdot)$ "disappears" from the likelihood or the posterior for ignorable designs, we can model non-ignorable designs using such a process. For example, we consider the model

$$Z(\ell) \sim u(\ell) \text{Ber}(\pi(\ell)) + (1 - u(\ell)) \delta_0; \quad \text{logit}(\pi(\ell)) = \beta_{0Z} + \beta_{1Z} Y(\ell) + x_Z^{\mathrm{T}} \beta_{[2:p]Z},$$

where, for an arbitrary location ℓ , the indicator process is distributed as a Bernoulli distribution with probability $\pi(\ell)$ if $u(\ell) = 1$, that is, if $\ell \in \mathcal{L}_{FP}$, and is set equal to zero if $u(\ell) = 0$, that is, if $\ell \notin \mathcal{L}_{FP}$. We introduce external predictors, x_Z , along with the finite population units $Y(\ell)$ to inform the sampling indicator process. In fact, one can conceive of introducing another spatial process

$$\operatorname{logit}(\pi(\ell)) = \beta_{0Z} + \beta_{1Z}Y(\ell) + x_Z^{\mathrm{T}}\beta_Z + \omega(\ell) ; \quad \omega(\ell) \sim GP(0, C_{\theta_{\omega}}(\cdot)) ,$$

where $\omega(\ell)$ is a zero centred Gaussian process with a specified covariance kernel $C_{\theta_{\omega}}(\cdot)$.

5.3.3 Modelling $[w(\cdot) \mid u(\cdot)] = [w(\cdot)]$ and $[u(\cdot)]$

The spatial process $w(\ell)$ is assumed to be independent of the population indicator $u(\ell)$ and is modelled as a multivariate Gaussian process. Let $w(\ell) = (w_1(\ell), \ldots, w_q(\ell))^{\mathrm{T}}$ be a $q \times 1$ vector, where $w_j(\ell)$ is a real-valued stochastic process over \mathcal{L} . The process is customarily specified by its mean $\mathbb{E}[w_j(\ell)] = \mu_j(\ell)$, which we assume to be zero, and the collection of stationary covariance functions $C_{ij}(h) = \operatorname{Cov}(w_i(\ell), w_j(\ell+h))$, for $i, j = 1, 2, \ldots, q$. The $q \times q$ matrix $C(h) = (C_{ij}(h))$ is the cross-covariance matrix and must satisfy (i) $C(h)^{\mathrm{T}} = C(-h)$ (note that C(h) itself need not be symmetric), and (ii) $\sum_{i,j=1}^{n} a_i^{\mathrm{T}} C_{ij}(h_{ij}) a_j \geq 0$ for any set of vectors a_1, \ldots, a_q in \mathbb{R}^q , where $h_{ij} =$



Fig. 1: Left: A directed graph (Bayesian network) with arrows pointing from parents to children. Right: An undirected conditional independence graph where the absence of edges between two nodes represent conditional independence given all other nodes.

 $\ell_i - \ell_j$, to ensure a valid multivariate spatial process for $w(\ell)$. A variety of methods are available for constructing multivariate spatial processes (Banerjee et al, 2014; Gelfand and Banerjee, 2010; Genton and Kleiber, 2015; Zhang and Banerjee, 2022).

The quantity $u(\ell)$ is a population indicator process modelled by

$$u(\ell) \sim \operatorname{Ber}(\pi(\ell)); \quad \operatorname{logit}(\pi(\ell)) = x_u(\ell)^{\top} \beta_u + v(\ell); \quad v(\ell) \sim GP(0, C_{\theta_v}(\cdot)),$$

where $v(\ell)$ is a GP with covariance function $C_{\theta_v}(\cdot)$. Here, the model for $\pi(\ell) = P(u(\ell) = 1)$ is a logistic regression accommodating explanatory variables that inform about a location being a part of the finite population or not.

5.4 Graphical models for finite populations

The ability to endow probability laws over the finite population units makes the Bayesian framework substantially richer than design-based or classical model-based frameworks. For example, we introduce complex multivariate dependencies among the finite population units using graphical models. Such models are based upon joint probability distributions over directed acyclic graphs (DAGs or Bayesian networks) and undirected graphs (left and right panels in Figure 1, respectively), where each node represents the value(s) of our variable(s) of interest from a population unit. An edge between two nodes in an undirected graph indicates that the random variables from the two nodes are conditionally dependence given the remaining variables. A DAG, due to its acyclic property, always has at least one source node (or a node that does not have an edge incident on it, hence has no parents) and one can always construct a sequence of ordered nodes (topological order) from its structure. Using this, the DAG defines a valid joint probability distribution as a product of the probability densities of the variable(s) from a node conditional on its parents.

Graphical models are especially significant in contemporary multivariate data analysis (Cox and Wermuth, 1996, 1993). Let $Y_i = (Y_{i1}, \ldots, Y_{iN})^{\mathsf{T}}$ be an $N \times 1$ random vector with elements Y_{ij} denoting the values of a variable of interest indexed by $i = 1, 2, \ldots, q$ over population units $j = 1, 2, \ldots, N$. Let $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ be a posited undirected graph, where \mathcal{V} is the set of q vertices representing the variables and \mathcal{E} is the set of all edges between pairs of variables that are assumed to be conditionally dependent with each other given all other variables. We specify the following sequence of full conditional distributions for modelling Y_i 's given all other Y_j 's for $j \in \{1, 2, \ldots, q\} \setminus \{i\}$:

$$Y_i \mid Y_{(-i)} \sim N\left(\sum_{j=1}^N A_{ij}Y_j, \Gamma_i\right), \quad i = 1, 2, \dots, q,$$
 (25)

where $Y_{-(i)} = \{Y_1, Y_2, \ldots, Y_q\} \setminus \{Y_i\}$, i.e., the collection of Y_j 's for $j = 1, 2, \ldots, q$ but excluding Y_i, A_{ij} 's are fixed $N \times N$ matrices, $A_{ii} = O$ (the matrix of zeroes), and Γ_i 's are fixed positive definite matrices. Since each variable can exhibit its own dependence structure, the Γ_i varies by variable. Brook's Lemma provides a straightforward method for deriving the joint density from (25) using the identity:

$$p(Y_1, Y_2, \dots, Y_q) = \prod_{i=1}^q \frac{p(Y_i \mid \tilde{Y}_1, \dots, \tilde{Y}_{i-1}, Y_{i+1}, \dots, Y_q)}{p(\tilde{Y}_i \mid \tilde{Y}_1, \dots, \tilde{Y}_{i-1}, Y_{i+1}, \dots, Y_q))} \times p(\tilde{Y}_1, \tilde{Y}_2, \dots, \tilde{Y}_q) , \quad (26)$$

where $\tilde{Y} = (\tilde{Y}_1^T, \tilde{Y}_2^T, \dots, \tilde{Y}_q^T)^T$ is any fixed point in the support of p(Y) and we assume that the joint density $p(\cdot) > 0$ over its entire support. The proof is a straightforward verification proceeding from the last element of the right hand side (i.e., the joint density on the right hand side). Thus, we begin with the observation that

$$\frac{p(Y_q \mid \tilde{Y}_1, \dots, \tilde{Y}_{q-1})}{p(\tilde{Y}_q \mid \tilde{Y}_1, \dots, \tilde{Y}_{q-1})} \times p(\tilde{Y}_1, \tilde{Y}_2, \dots, \tilde{Y}_q) = p(\tilde{Y}_1, \tilde{Y}_2, \dots, \tilde{Y}_{q-1}, Y_q) \ .$$

Proceeding as above will continue to replace \tilde{Y}_i with Y_i in the joint density on the right hand side of (26) for each i = q - 1, q - 2, ..., 1 and we eventually arrive at $p(Y_1, Y_2, ..., Y_q)$. Applying (26) to the full conditional distributions in (25) with $\tilde{Y}_i = 0$ for each i = 1, ..., q yields the joint density

$$p(Y) \propto \exp\left\{-\frac{1}{2}\left(\sum_{i=1}^{q} Y_i^{\mathrm{T}} \Gamma_i^{-1} Y_i - \sum_{i=1}^{q} \sum_{j \neq i}^{q} Y_i^{\mathrm{T}} \Gamma_i^{-1} A_{ij} Y_j\right)\right\} \propto \exp\left(-\frac{1}{2} Y^{\mathrm{T}} Q Y\right),$$
(27)

where $Y = (Y_1^{\mathrm{T}}, \ldots, Y_q^{\mathrm{T}})^{\mathrm{T}}$ is $Nq \times 1$ and $Q = M^{-1}(I - A)$ is $Nq \times Nq$, $M = \bigoplus \Gamma_i$ is block-diagonal with (i, i)-th block Γ_i for $i = 1, \ldots, q$ and $A = (A_{ij})$ is the $Nq \times Nq$ block matrix with A_{ij} as the (i, j)-th block. For (27) to be a valid density, Q needs to be symmetric and positive definite so that $Y \sim N(0, Q^{-1})$. In addition, Q must conform to the conditional independence relationships in \mathcal{G} . To be precise, if two distinct nodes

i and *j* in the graph do not have an edge, then $Y_i \perp Y_j \mid Y_{-(i,j)}$ or, equivalently, the $N \times N$ block of the precision $Q_{ij} = -\Gamma_i^{-1} A_{ij} = O$. Since $A_{ii} = O$ in (25), the (*i*, *i*)-th block of Q is Γ_i^{-1} . We devise one such construction below.

Given the inter-variable graph $\mathcal{G} = \{\mathcal{V}, \mathcal{E}\}$ let $\Lambda = (\lambda_{ij}) = D - \rho W$ be the $q \times q$ graph Laplacian, where $W = (w_{ij})$ is the adjacency matrix with nonzero w_{ij} only if there is an edge between i and j, $D = (d_{ii})$ is a $q \times q$ diagonal matrix with the sum of each row of W along the diagonal, i.e., $d_{ii} = \sum_{j=1}^{q} w_{ij}$, and ρ is a scalar parameter that ensures positive-definiteness of Λ as long as $\rho \in (1/\zeta_{min}, \zeta_{max})$, where ζ_{min} and ζ_{max} are the minimum and maximum eigenvalues of $D^{-1/2}WD^{-1/2}$, respectively. Let R_i be any nonsingular $N \times N$ factor (e.g., a triangular Cholesky) of the positive definite precision matrix of Y_i , i.e., $(\operatorname{var}(Y_i))^{-1} = C_{ii}^{-1} = R_i^{\mathsf{T}}R_i$ for each $i = 1, 2, \ldots, q$. We set $\Gamma_i^{-1} = \lambda_{ii}R_i^{\mathsf{T}}R_i$ and $A_{ij} = -(\lambda_{ij}/\lambda_{ii})R_i^{-1}R_j$ in (25). Then $Q_{ij} = \lambda_{ij}R_i^{\mathsf{T}}R_j$ and $Q = \tilde{R}^{\mathsf{T}}(\Lambda \otimes I)\tilde{R}$, where $\tilde{R} = \bigoplus_{i=1}^{q}R_i$. Since each R_i is nonsingular and Λ is positive definite, it follows that Q is positive definite and, hence, $\det(Q) = \prod_{i=1}^{q} (\det(R_i))^2 (\det(\Lambda))^n$. The above construction yields proper densities in (27) that will conform to conditional independence among the variables represented by the nodes of a posited undirected graph and can also be adapted for analysing high-dimensional multivariate data, where at least one of or both N and q are large. The special case where each $R_i = R$ for $i = 1, 2, \ldots, q$ yields the separable model $Q = \Lambda \otimes (R^{\mathsf{T}}R)$.

If, for example, the population units are spatially structured, as is often the case in small area estimation, then spatial dependencies are introduced in the R_i 's. The CAR model, which is a popular choice in small area estimation, uses the Laplacian of a spatial graph, say $\mathcal{G}_s = \{\mathcal{V}_s, \mathcal{E}_s\}$, for modelling spatial concentrations. The vertices in \mathcal{V}_s are the population units and the edges in \mathcal{E}_s between two units indicate they are spatial neighbours. The suffix s is used to distinguish this graph from the intervariable graph in the previous paragraph. Each R_i satisfies $R_i^{T}R_i = D_s - \rho_{s,i}W_s$, where D_s is $N \times N$ diagonal with the number of neighbours for each region as its diagonal elements and W_s is the $N \times N$ spatial adjacency matrix. The parameter $\rho_{s,i}$ depends on the variable index i because each variable is allowed to have its own spatial correlation parameter. Since $D_s^{-1/2}WD_s^{-1/2}$ is symmetric, there is an orthonormal basis of its eigenvectors $\{u_1, u_2, \ldots, u_N\}$ in \mathbb{R}^N consisting of its eigenvectors, and we can write $I_N - \rho_{s,i}D_s^{-1/2}WD_s^{-1/2} = \sum_{j=1}^N (1 - \zeta_{s,j}\rho_{s,i})u_ju_j^{\mathrm{T}}$, where $\zeta_{s,j}$ is the eigenvalue corre-sponding to u_j . Since \mathcal{G}_s is fixed from the map of population units, the eigenvalues and eigenvectors are computed only once and fixed. Each spatial concentration parameter $\rho_{s,i}$ is restricted to be in an interval so that $1 - \zeta_{s,j}\rho_{s,i} > 0$ for each $j = 1, 2, \dots, N$. If $R_i = \sum_{j=1}^N \sqrt{(1-\zeta_{s,j}\rho_{s,i})} u_j u_j^{\mathrm{T}} / \sqrt{d_{s,j}}$, where $d_{s,j}$ is the *j*-th diagonal element of D_s , then R_i^2 is the symmetric square root of the spatial concentration matrix and depends only on one unknown parameter, $\rho_{s,i}$, which is estimated from its posterior.

6 Model assessments and comparisons

The preceding discussions demonstrates the richness of Bayesian hierarchical modelling for estimating finite population quantities. This richness comes at an expense of justifying the choice of the model for the population, which requires using Bayesian model selection metrics. For example, consider the two models (5) and (7). Both these

models emulate (one approximate and the other exact) the Horvitz-Thompson estimator, but differ in their variance structure. A design-based perspective favouring the Horvitz-Thompson estimator focuses primarily on its unbiased nature and on estimating its variance. A model-based perspective, including a fully Bayesian perspective, entails deciding between these models based upon a realised sample.

To elucidate further, assume that we wish to compare two models within the framework of (22) but with possibly different specifications for the design matrix and the dependence structure. A key quantity in Bayesian model selection is the point-wise predictive distribution (see, e.g., Gelman et al, 2014) defined as

$$p(Y_{rep,i} \mid y) = \mathbb{E}_{\beta,\theta \mid y} \left[p(Y_{rep,i} \mid \beta, \theta) \right] = \int p(Y_{rep,i} \mid \beta, \theta) \times p(\theta, \beta \mid y) \, d\beta \, d\theta \qquad (28)$$

for each unit *i* in the population. We sample one value of $Y_{rep,i} \sim p(Y_{rep,i} | \beta, \theta)$ for each posterior sample of β and θ . An ideal measure of a model's performance is its outof-sample predictive performance for new data produced from the true data-generating process. This, unfortunately, is impracticable (except in synthetic experiments that simulate the population) because we do not know the true data generating process. Therefore, the predictive distribution (28) encodes the distribution of alternative data that would be replicated from the fitted model.

Gelfand and Ghosh (1998) uses the predictive distribution to define

$$D = \sum_{i \in \mathcal{S}} (y_i - \mathbb{E}[Y_{rep,i} \mid y])^2 + \sum_{i \in \mathcal{S}} \mathbb{V}(Y_{rep,i} \mid y)$$
(29)

as a model choice metric, while Gneiting and Raftery (2007) prefer

$$GRS = -\sum_{i \in \mathcal{S}} \frac{(y_i - \mathbb{E}[Y_{rep,i} \mid y])^2}{\mathbb{V}(Y_{rep,i} \mid y)} - \sum_{i \in \mathcal{S}} \log \mathbb{V}(Y_{rep,i} \mid y) , \qquad (30)$$

as a proper scoring rule, where S is the set of sampled units. Lower values of D and higher values of GRS indicate better model fit. A third, more recently developed, metric is the Watanabe-Akaike Information Criterion (see, e.g., Watanabe, 2013; Vehtari et al, 2017), which computes the logarithm of (28) evaluated at each data point and adjusts with a penalty term as

$$WAIC = -2\sum_{i\in\mathcal{S}} \log \mathbb{E}_{\beta,\theta|y}[p(y_i \mid \beta, \theta)] + 2\sum_{i\in\mathcal{S}} \mathbb{V}_{\beta,\theta|y}\left(\log p(y_i \mid \beta, \theta)\right) .$$
(31)

The WAIC computed as above is in the same scale as the deviance with smaller values indicating superior performance.

Each of the metrics (29)—(31) are computed without requiring any information on a true data generating model. The quantities (29) and (30) are computed using samples of $Y_{rep,i}$ from (28) to calculate $\mathbb{E}[Y_{rep,i} \mid y]$ and $\mathbb{V}(Y_{rep,i} \mid y)$. The remaining information used comes from the realised sample of values from the finite population. We compute the quantities in (31) using $\sum_{l=1}^{L} p(y_i \mid \beta^{(l)}, \theta^{(l)})/L$ and $\operatorname{var}_{l=1,...,L}(\log p(y_i \mid$

 $\beta^{(l)}, \theta^{(l)}))$ to estimate $\mathbb{E}_{\beta,\theta|y}[p(y_i \mid \beta, \theta)]$ and $\mathbb{V}_{\beta,\theta|y}(\log p(y_i \mid \beta, \theta))$, respectively, where $\operatorname{var}_{l=1,\ldots,L}(a^{(l)}) = \sum_{l=1}^{L} \left(a^{(l)} - \left(\sum_{l=1}^{L} a^{(l)}/L\right)\right)^2/(L-1)$ and $\beta^{(l)}, \theta^{(l)}$ are posterior samples $l = 1, \ldots, L$. Vehtari et al. (Vehtari et al, 2017) provide further details on *WAIC*, including methods to compute standard errors of *WAIC* (used by Chan-Golston et al, 2020, in finite population inference) and its connections to predictive leave-one-out (LOO) comparisons.

7 Examples

I present some salient features of an analysis from two examples. The first is a study of ground-water nitrate levels in the population of California Central Valley wells situated in the Tulare Lake Basin (TLB), where health officials seek estimates of mean nitrate levels. The data we consider here has been extracted from the California Ambient Spatio-Temporal Information on Nitrate in Groundwater (CASTING) Database, which is described in Harter et al (2017) and Boyle et al (2012). The spatial locations of these wells are available. Health implications of high nitrate levels in wells are dire. At high levels, infants and pregnant women are more susceptible to nitrate poisoning, which makes it more difficult for oxygen to be distributed to body and can be fatal to infants less than six months old.

Our finite population consists of 6,117 unique wells among 63 zip codes. We follow a two-stage cluster sampling design described in Chan-Golston et al (2020), where 21 zip codes were randomly chosen in the first stage and 50%–90% of the wells in each zip code were randomly sampled in the second stage. The nitrate levels in the sample wells were measured in milligrams per litre (mg/L). The following models were used to analyse the sampled data: (i) the two-stage hierarchical cluster model described in Section 4; (ii) a spatial random field model over the entire region that ignores the structure of the cluster sampling design (so a single set of spatial process parameters for the entire region); (iii) a two-stage model with an additive global spatial process; and (iv) a regional spatial process model where each cluster has its own mean parameter and spatial process. All of these models can, in fact, be subsumed into (22) with appropriately constructed design and covariance matrices.

Table 1 presents the posterior mean and 95% credible intervals for the finite population means obtained from each model. There are some discrepancies in these estimates among the different models. Hence, it will be appropriate to The WAIC scores for each of the models are also presented. Some key features are worth pointing out. Notably, models that either ignore the two-stage structure of the sampled data (models ii and iv) or that ignore the spatial dependence (model i) appear to perform considerably worse (in terms of WAIC) than the model that accommodates both the sampling structure and the spatial dependence (model iii). Our recommendation is to report the analysis from model (iii). Also, while the credible intervals around the finite population mean from different models largely overlap, the point estimates from (i) and (iv) are lower than (ii) or (iii). This is, perhaps, attributed to the former two models not having a single spatial random field over the entire domain. Finally, the spatial clustering introduced in the latter two models result in shorter credible intervals.

Table 1: Results of California Nitrate Data Analysis

Model	Finite Population Mean in mg/L (95% CI)	WAIC
(i) Two-Stage	$26.6\ (19.2,\ 35.1)$	4701.6
(ii) Spatial	$32.6\ (27.2,\ 38.7)$	4858.4
(iii) Two-Stage + Spatial	31.2(24.2, 37.3)	2697.0
(iv) Regional Spatial	26.2(18.7, 34.4)	4536.6

Unlike the above analysis, which assumed an ignorable design, I now present some key aspects of a second analysis from a study to estimate the percentage of annual reported income spent on fruits and vegetables (PIFV) in two low-income communities in Los Angeles, California (see Chan-Golston et al, 2022, and references therein for a detailed description of the study). This study follows the framework described in Section 5.3. There are N = 635 locations in \mathcal{L}_{FP} and within each location we have a number of households comprising the population units. The total number of population units is 2015. We randomly sampled n = 555 locations and then sampled households within each of these locations to obtain a sample of 1294 population units. These units reported amounts spent on fruits and vegetables on weekly, bi-weekly, or monthly scale. These values were multiplied by 52, 26, and 12, respectively, to reflect the annual amount spent on fruits and vegetables in a household. Reported yearly incomes in the population units were also collected.

We consider a simpler version of the model presented in Section 5.3 and its subsections with the process $u(\cdot) = 1$. We assume that the PIFV of the *j*-th household within location ℓ_i is modelled as $Y_j(\ell_i) = x_j(\ell_i)^{\mathrm{T}}\beta + w(\ell_i) + \epsilon(\ell_i)$, $x_j(\ell_i)$ is a vector of fixed explanatory variables, $w(\ell_i)$ is a spatial random effect and $\epsilon(\ell_i)$ is white noise representing measurement errors and micro-scale variation. This is the model for $[Y(\cdot) \mid w(\cdot)]$ in Section 5.3 with $u(\ell) = 1$. Based upon this, we consider four models: (i) a simple linear regression model that sets $w(\ell) = 0$ for all locations (yields an ignorable response); (ii) a model with nonignorable response that models $[Z(\cdot) \mid Y(\cdot), \omega(\cdot)]$ in addition to $[Y(\cdot) \mid w(\cdot)]$, but with no spatial effect (hence, $w(\cdot) = 0$ and $\omega(\cdot) = 0$); (iii) spatial random effects in $[Y(\cdot) \mid w(\cdot)]$, but no spatial effects in $[Z(\cdot) \mid Y(\cdot)]$; and (iv) fully spatial model with nonignorable responses with spatial effects $w(\cdot)$ and $\omega(\cdot)$ as described in Section 5.3.

Table 2 presents the posterior estimates of the finite population PIFV (first row) from the four models described above. The performance of these models was evaluated using D and GRS defined in (29) and (30), respectively. Based upon these metrics, model (iii) is marginally preferred to model (iv), which suggests a lack of spatial dependence in the sampling inclusion model. Both the spatial models, (iii) and (iv), considerably outperform the non-spatial models including (ii) which still accommodates the nonignorable response. Model (i), which assumes ignorability, performs the worse. The credible interval from the ignorable model seems very tight, while those from the other models are wider because of the propagation of uncertainty from the additional parameters being estimated. The PIFV from model (iii) is recommended for reporting given the model's superior performance.

Table 2: Results of regression models predicting percentage of income spent on fruits and vegetables (log-scale). The finite population mean income is presented in units of \$ 10,000.

	Model 1	Model 2	Model 3	Model 4
PIFV $\%$	$17\% \ (16\%, \ 19\%)$	26% (22%, 31%)	35% (28%, 43%)	26% (22%, 32%)
D	3527.8	3692.2	3701.2	3482
GRS	-1841.2	-1903.6	-1863	-1767.8

8 Discussion

The article presents Bayesian inference for finite populations using survey data with an emphasis on settings where the finite population units are posited to exhibit complex dependence structures. Spatial process models for the finite population units are devised in ignorable and nonignorable response settings to bring out the richness and flexibility of Bayesian inference adapted to finite population survey sampling. Bayesian inference for finite populations in stochastic process settings has received relatively limited attention and can form a part of future extensions to this paper. Of particular interest to spatial statisticians will be the connections between models for nonignorable spatial responses and the preferential sampling framework described by Diggle et al (2010). Modelling massive spatially oriented survey data is gaining attention in environmental sciences, ecology and forestry (see, e.g., Finley et al, in press, for a case study in forestry that compares Bayesian spatial models with design-based estimates for biomass and demonstrates the inferential benefits of the former). New methods for analysing massive spatially referenced survey data can adapt, perhaps even simplify, spatial meta-kriging approaches (Guhaniyogi and Banerjee, 2018) to finite population inference for ignorable and non-ignorable settings. Finally, attention is needed in the development and dissemination of software products for implementing Bayesian models for spatially oriented survey data.

Acknowledgments. The author wishes to thank the editors and an anonymous referee for valuable feedback. The author is especially grateful to Professors Roderick J. Little and Trivellore Raghunathan from the University of Michigan, Ann Arbor, U.S.A., and Professor J.N.K. Rao from Carleton University, Ottawa, Canada, for insightful discussions on inference for finite populations. The work of the author has been supported, in part, by the National Science Foundation (NSF) from grants NSF/DMS 1916349 and NSF/DMS 2113778, by the National Institute of Environmental Health Sciences (NIEHS) from grants R01ES030210 and R01ES027027 and by the National Institute of General Medical Science from grant R01GM148761.

Disclosures and declarations

• Funding: The work of the author has been supported, in part, by the National Science Foundation (NSF) from grants NSF/DMS 1916349 and NSF/DMS 2113778, by the National Institute of Environmental Health Sciences (NIEHS) from grants R01ES030210 and R01ES027027 and by the National Institute of General Medical Science from grant R01GM148761.

• Conflict of interest: The author declares that there are no financial or nonfinancial conflicts of interest.

References

- Arora V, Lahiri P (1997) On the superiority of the bayesian method over the blup in small area estimation problems. Statistica Sinica 7(4):1053–1063. URL http:// www.jstor.org/stable/24306172
- Arora V, Lahiri P, Mukherjee K (1997) Empirical bayes estimation of finite population means from complex surveys. Journal of the American Statistical Association 92(440):1555–1562
- Banerjee S, Carlin BP, Gelfand AE (2014) Hierarchical Modeling and Analysis for Spatial Data, 2nd edn. Chapman & Hall/CRC, Boca Raton, FL
- Basu D (1971) An essay on the logical foundations of survey sampling, part 1. Holt, Rinehart and Winston, Toronto, p $203{-}242$
- Boyle D, King A, Kourakos G, et al (2012) Groundwater Nitrate Occurrence, Technical Report 4. Tech. rep., Center for Watershed Sciences, University of California, Davis, Davis, CA
- Bruno F, Cocchi D, Vagheggini A (2013) Finite population properties of individual predictors based on spatial pattern. Environmental and Ecological Statistics 20(3):467–494
- Chan-Golston AM, Banerjee S, Handcock MS (2020) Bayesian inference for finite populations under spatial process settings. Environmetrics 31(3):e2606.https://doi.org/https://doi.org/10.1002/env.2606, URL https://onlinelibrary.wiley.com/doi/abs/10.1002/env.2606, e2606 env.2606, https://onlinelibrary.wiley.com/doi/pdf/10.1002/env.2606
- Chan-Golston AM, Banerjee S, Belin TR, et al (2022) Bayesian finitepopulation inference with spatially correlated measurements. Japanese Journal of Statistics and Data Science 5:407–430. https://doi.org/https://doi. org/10.1007/s42081-022-00178-8, URL https://link.springer.com/article/10.1007/ s42081-022-00178-8#citeas
- Cicchitelli G, Montanari GE (2012) Model-assisted estimation of a spatial population mean. International Statistical Review 80(1):111–126
- Clayton D, Kaldor J (1987) Empirical Bayes estimates of age-standardized relative risks for use in disease mapping. Biometrics 43(3):671–681

Cochran WG (1977) Sampling Techniques, 3rd edn. John Wiley & Sons, Hoboken, NJ

- Cox D, Wermuth N (1996) Multivariate Dependencies. Chapman & Hall/CRC, Boca Raton, FL
- Cox DR, Wermuth N (1993) Linear Dependencies Represented by Chain Graphs. Statistical Science 8(3):204 – 218. https://doi.org/10.1214/ss/1177010887, URL https://doi.org/10.1214/ss/1177010887
- Cressie N, Wikle CK (2011) Statistics for Spatio-Temporal Data. John Wiley & Sons, Hoboken, NJ
- Datta GS, Ghosh M (1991) Bayesian prediction in linear models: Applications to small area estimation. The Annals of Statistics pp 1748–1770
- de Valpine P, Turek D, Paciorek C, et al (2017) Programming with models: writing statistical algorithms for general model structures with NIMBLE. Journal of Computational and Graphical Statistics 26:403–413. https://doi.org/10.1080/10618600. 2016.1172487
- de Valpine P, Paciorek C, Turek D, et al (2023) NIMBLE: MCMC, Particle Filtering, and Programmable Hierarchical Modeling. https://doi.org/10.5281/zenodo. 1211190, URL https://cran.r-project.org/package=nimble, R package version 1.0.1
- Di Zio M, Liseo B, Ranalli MG (2023) Bayesian ideas in survey sampling: The legacy of Basu. Sankhya A https://doi.org/10.1007/s13171-023-00327-5, URL https://doi.org/10.1007/s13171-023-00327-5
- Diggle PJ, Menezes R, Su TL (2010) Geostatistical inference under preferential sampling. Journal of the Royal Statistical Society: Series C 59(2):191–232
- Ericson WA (1969) Subjective Bayesian models in sampling finite populations. Journal of the Royal Statistical Society, Series B 31(2):195–233
- Finley AO, Banerjee S, MacFarlane DW (2011) A hierarchical model for quantifying forest variables over large heterogeneous landscapes with uncertain forest areas. Journal of the American Statistical Association 106(493):31–48. https://doi.org/10. 1198/jasa.2011.ap09653, URL https://doi.org/10.1198/jasa.2011.ap09653, pMID: 26139950, https://doi.org/10.1198/jasa.2011.ap09653
- Finley AO, Andersen HE, Babcock C, et al (in press) Models to support forest inventory and small area estimation using sparsely sampled lidar: A case study involving g-liht lidar in tanana, alaska. Journal of Agricultural, Biological, and Environmental Statistics
- Gelfand AE, Banerjee S (2010) Multivariate spatial process models. In: Gelfand A, Diggle P, Fuentes M, et al (eds) Handbook of Spatial Statistics. CRC Press, Boca Raton, FL, p 495–516

- Gelfand AE, Ghosh SK (1998) Model choice: a minimum posterior predictive loss approach. Biometrika 85(1):1–11
- Gelman A (2007) Struggles with Survey Weighting and Regression Modeling. Statistical Science 22(2):153–164
- Gelman A, Carlin JB, Stern HS, et al (2014) Bayesian Data Analysis, 3rd edn. Chapman & Hall/CRC, Boca Raton, FL
- Genton MG, Kleiber W (2015) Cross-covariance functions for multivariate geostatistics. Statistical Science pp 147–163
- Ghosh M (2012) Finite population sampling: A model-design synthesis. Statistics in Transition new series 13(2):235–242
- Ghosh M, Meeden G (1997) Bayesian Methods for Finite Population Sampling. Chapman & Hall, London
- Ghosh M, Rao JNK (1994) Small Area Estimation: An Appraisal. Statistical Science 9(1):55–93
- Ghosh M, Sinha BK (1990) On the consistency between model-and design-based estimators in survey sampling. Communications in Statistics - Theory and Methods 19(2):689–702. https://doi.org/10.1080/03610929008830226, URL https://doi.org/ 10.1080/03610929008830226, https://doi.org/10.1080/03610929008830226
- Ghosh M, Natarajan K, Stroud TWF, et al (1998) Generalized linear models for smallarea estimation. Journal of the American Statistical Association 93(441):273–282
- Gneiting T, Raftery AE (2007) Strictly proper scoring rules, prediction, and estimation. Journal of the American Statistical Association 102(447):359–378
- Guhaniyogi R, Banerjee S (2018) Meta-kriging: Scalable bayesian modeling and inference for massive spatial datasets. Technometrics 60(4):430–444
- Harter T, Dzurella K, Kourakos G, et al (2017) Nitrogen Fertilizer Loading to Groundwater in the Central Valley, Final Report to the Fertilizer Research Education Program Projects 11-0301 and 15-0454. Tech. rep., California Department of Food and Agriculture and University of California Davis, Davis, CA
- Horvitz DG, Thompson DJ (1952) A Generalization of Sampling Without Replacement From a Finite Universe. Journal of the American Statistical Association 47(260):663–685
- Kalton G (2019) Developments in survey research over the past 60 years: A personal perspective. International Statistical Review 87(S1):S10–S30. https://doi.org/ https://doi.org/10.1111/insr.12287, URL https://onlinelibrary.wiley.com/doi/abs/ 10.1111/insr.12287, https://onlinelibrary.wiley.com/doi/pdf/10.1111/insr.12287

- Kish L (1965) Survey Sampling. John Wiley & Sons, Inc., Hoboken, New Jersey
- Kish L (1995) The hundred years' wars of survey sampling. Statistics in Transition 2:813–830
- Little R, Rubin D (2002) Statistical Analysis with Missing Data. John Wiley & Sons, Inc., Hoboken, New Jersey
- Little RJ (1982) Models for nonresponse in sample surveys. Journal of the American statistical Association 77(378):237–250
- Little RJ (2004) To Model or Not To Model? Competing Modes of Inference for Finite Population Sampling. Journal of the American Statistical Association 99(466):546– 556
- Lunn D, Spiegelhalter D, Thomas A, et al (2009) The bugs project: Evolution, critique and future directions. Statistics in Medicine 28(25):3049–3067. https://doi.org/ https://doi.org/10.1002/sim.3680, URL https://onlinelibrary.wiley.com/doi/abs/ 10.1002/sim.3680, https://onlinelibrary.wiley.com/doi/pdf/10.1002/sim.3680
- Malec D, Sedransk J (1985) Bayesian Inference for Finite Population Parameters in Multistage Cluster Sampling. Journal of the American Statistical Association 80(392):897–902
- Narain R (1951) On sampling without replacement with varying probabilities. Journal of the Indian Society of Agricultural Statistics 3:169–175
- Rao JNK (2011) Impact of Frequentist and Bayesian Methods on Survey Sampling Practice: A Selective Appraisal. Statistical Science 26(2):240 – 256. https://doi.org/ 10.1214/10-STS346, URL https://doi.org/10.1214/10-STS346
- Rao JNK. Ghangurde PD(1972)Bayesian optimization insampling finite populations. Journal of the American Statistical Association https://doi.org/10.1080/01621459.1972.10482406, URL 67(338):439-443.https://www.tandfonline.com/doi/abs/10.1080/01621459.1972.10482406, https://www.tandfonline.com/doi/pdf/10.1080/01621459.1972.10482406
- Rao JNK, Molina I (2015) Small Area Estimation, 2nd edn. John Wiley & Sons, Hoboken, NJ
- Ripley BD (2004) Spatial Statistics. John Wiley & Sons, Hoboken, NJ
- Rubin DB (1976) Inference and missing data. Biometrika 63(3):581–592
- Scott A, Smith TMF (1969) Estimation in Multi-Stage Surveys. Journal of the American Statistical Association 64(327):830–840

- Stan Development Team (2024) RStan: the R interface to Stan. URL https://mc-stan. org/, r package version 2.32.5
- Tang G, Little RJ, Raghunathan TE (2003) Analysis of multivariate missing data with nonignorable nonresponse. Biometrika 90(4):747–764
- Vehtari A, Gelman A, Gabry J (2017) Practical Bayesian model evaluation using leaveone-out cross-validation and WAIC. Statistics and Computing 27(5):1413–1432
- Ver Hoef J (2002) Sampling and geostatistics for spatial data. Écoscience 9(2):152-161
- Watanabe S (2013) A widely applicable bayesian information criterion. Journal of Machine Learning Research 14(27):867–897. URL http://jmlr.org/papers/v14/watanabe13a.html
- Zhang L, Banerjee S (2022) Spatial factor modeling: A bayesian matrix-normal approach for misaligned data. Biometrics 78(2):560–573. https://doi.org/https://doi.org/10.1111/biom.13452, URL https://onlinelibrary.wiley.com/doi/abs/10. 1111/biom.13452, https://onlinelibrary.wiley.com/doi/pdf/10.1111/biom.13452