A bounded partition approach to identifying one fake coin and its type

Takehiro Tokuda^a, Yoshimichi Watanabe^{b,*}

^a Emeritus Professor, Tokyo Institute of Technology, 2-12-1, Ookayama, Meguro, 152-8552, Tokyo, Japan
^b University of Yamanashi, 4-3-11, Takeda, Kofu, 400-8511, Japan

Abstract

Fake coin problems using balance scales to identify one fake coin and its type among n coins (n > 2) were solved by Dyson in 1946. Dyson gave adaptive solutions with the minimum number of weighings where later weighings may be dependent on results of past weighings. In 2003 Born et al. gave non-adaptive solutions where all weighings are predetermined. Both solutions require the computation of a Dyson set, which is a list of placement of each coin for each weighing. The computation of a Dyson set requires substantial amount of time when n gets larger.

We present a bounded partition approach to the fake coin problems without computing any Dyson set. Our approach uses bounded partition of coins recursively until the problem size may be small enough to use at most one weighing.

Keywords: fake coin problem, bounded partition, problem size, recursive algorithm, ternary search

1. Introduction

We have $n \operatorname{coins} (n > 2)$ and they are identical in appearance. Exactly one of $n \operatorname{coins}$ is a fake coin which is heavier or lighter than genuine coins. All genuine coins have the same weight. The fake coin problem of $n \operatorname{coins}$ is to identify one fake coin and its type, heavy or light, using the minimum number of weighings on balance scales.

In 1946, Dyson showed that if $6 \le 2n \le 3^k - 3$ $(k \ge 2)$, then we can solve the fake coin problems of n coins using at most k weighings, and if $2n > 3^k - 3$, then we cannot solve the problems of n coins using at most k weighings. Dyson gave adaptive solutions to the fake coin problems where later weighings may be dependent on results of past weighings [1]. In 2003, Born et al. gave non-adaptive solutions to the fake coin problems where all weighings are predetermined [2].

^{*}Corresponding author

E-mail addresses: tokuda@cs.titech.ac.jp (T. Tokuda), nabe@yamanashi.ac.jp (Y. Watanabe).

Both solutions use a Dyson set, which is a list of placement of each coin for each weighing. The computation of a Dyson set requires substantial amount of time when n gets larger.

In this note, we present a bounded partition approach for solving the fake coin problems without computing any Dyson set. We directly solve the fake con problems based on the size of a problem. Our approach uses bounded partition of coins recursively until the problem size may be small enough to use at most one weighing. Historically Dyson used the size of a problem for deriving the necessary condition of solutions for the fake coin problems.

2. Basic Idea

We consider three versions of fake coin problems P[n], P[(m)], and P[(x, y)], which are closely related to solve the original problem. Notice that each weighing has the same number of coins on the left side and the right side of balance scales.

- P[n] Original fake coin problem of n coins.
- P[(m)] Fake coin problem of m coins where one genuine coin outside of given m coins can be used to place the same number of coins on both sides of balance scales, when the number of the left side coins and the number of the right side coins differ by 1.
- P[(x, y)] Fake coin problem of x heavy coin candidates and y light coin candidates. A heavy coin candidate is a genuine or heavy coin, and a light coin candidate is a genuine or light coin. Also one genuine coin outside of given x + y coins can be used to place the same number of coins on both sides, when the number of the left side coins and the number of the right side coins differ by 1.

Notice that original problem P[1] has no solution, because we cannot identify the type of the given fake coin, and P[2] has no solution, because we cannot tell whether a heavier coin or a lighter coin is fake. Solutions of original problems P[3] and P[4] require 2 weighings and 3 weighings respectively. Also P[1] cannot be considered as a subproblem of P[3] in any sense. But if we were allowed to use one genuine coin other than given coins, then P[(1)] has a solution where we compare the given coin with one genuine coin once, and P[(2)] has a solution where we compare one of given coins with one genuine coin at most twice. Both P[(3)] and P[(4)] can be solved using 2 weighings. Naturally P[(1)] is a subproblem of P[(3)] as we observe later.

Relations of three versions of fake coin problems are as follows. For P[n], if we have a partition (x, y, m) where n = x + y + m, x coins on the left side, y coins on the right side, and m coins on the outside, then the result of one weighing makes P[n] into one of the following three subproblems as shown in the following description of a basic method for solving P[n].

P[n]:

 $n \rightarrow \text{partition}(x, y, m);$

If the left side goes down, P[(x, y)]; $\{m \text{ coins on the outside are genuine}\}$

If the right side goes down, P[(y, x)]; {*m* coins on the outside are genuine} If balanced, P[(m)]; {*x* + *y* coins on both sides are genuine}

We respectively refer to P[(x, y)], P[(y, x)], and P[(m)] as the left subproblem, the right subproblem, and the balanced subproblem of P[n]. Notice that one genuine coin is available after the first weighing except when m = 0. As we observe later, when the number of the outside coins is 0 in the first weighing, one genuine coin is available after the second weighing.

For P[(x, y)], if we have a partition $((x_1, y_1), (x_2, y_2), (x_3, y_3))$, where $x = x_1 + x_2 + x_3$, $y = y_1 + y_2 + y_3$, x_1 heavy coin candidates and y_1 light coin candidates on the left side, x_2 heavy coin candidates and y_2 light coin candidates on the right side, and x_3 heavy coin candidates and y_3 light coin candidates on the outside, then the result of one weighing makes P[(x, y)] into one of the following three subproblems as shown in the following description of a basic method for solving P[(x, y)].

P[(x,y)]:

 $(x, y) \rightarrow \text{partition}((x_1, y_1), (x_2, y_2), (x_3, y_3));$

If the left side goes down, $P[(x_1, y_2)]$; $\{x_2 + y_1 + x_3 + y_3 \text{ coins are genuine}\}$ If the right side goes down, $P[(x_2, y_1)]$; $\{x_1 + y_2 + x_3 + y_3 \text{ coins are genuine}\}$ If balanced, $P[(x_3, y_3)]$; $\{x_1 + y_1 + x_2 + y_2 \text{ coins are genuine}\}$

We respectively refer to $P[(x_1, y_2)]$, $P[(x_2, y_1)]$, and $P[(x_3, y_3)]$ as the left subproblem, the right subproblem, and the balanced subproblem of P[(x, y)].

For P[(m)], if we have a partition (x, y, t) where m = x + y + t, then the result of one weighing makes P[(m)] into one of the following three subproblems as shown in the following description of a basic method for solving P[(m)].

P[(m)]:

 $m \rightarrow \text{partition}(x, y, t);$

If the left side goes down, P[(x, y)]; {t coins on the outside are genuine} If the right side goes down, P[(y, x)]; {t coins on the outside are genuine} If balanced, P[(t)]; {x + y coins on both sides are genuine}

We respectively refer to P[(x, y)], P[(y, x)], and P[(t)] as the left subproblem, the right subproblem, and the balanced subproblem of P[(m)].

For selecting an appropriate partition, we use the size of a problem. The size of a problem P, denoted by Size[P], is the sum of the number of heavy coin candidates and the number of light coin candidates of P. Initially any one of n coins can be a fake coin. Hence all n coins are heavy coin candidates and all n coins are light coin candidates. Therefore Size[P[n]] = 2n. Values of the size of problems P[n], P[(x, y)], and P[(m)] are as follows.

Size [P[n]] = n + n = 2n.

Size [P[(x,y)] = x + y.

Size [P[(m)]] = m + m = 2m.

After each weighing, we have a new total number of heavy coin candidates and light coin candidates.

We use a bounded partition of coins recursively. Partition must be made to ensure that subproblems of smaller size can be solved using fewer weighings than the given problem. Notice that problems P[(x, y)] and P[(m)] of size ≤ 3 can be solved using at most one weighing as described in Induction Basis.

A partition of a problem P with bound c is a partition where P1 is the left subproblem of P, P2 is the right subproblem of P, P3 is the balanced subproblem of P, Size[P1] $\leq c$, Size[P2] $\leq c$, and Size[P3] $\leq c$.

Using appropriate partition bounds, our method has the following Properties 1.1, 1.2, and 1.3. For convenience we assume problems P[(0)] and P[(0,0)] are solvable without doing anything.

Property 1.1 (P[n]). If $6 \le 2n \le 3^k - 3$ for $k \ge 2$, then P[n] is solvable with at most k weighings.

Property 1.2 (P[(m)]). If $2m \le 3^k - 1 < 3^k$ for $k \ge 1$, then P[(m)] is solvable with at most k weighings.

Property 1.3 (P[(x,y)]). If $x + y \le 3^k$ for $k \ge 1$, then P[(x,y)] is solvable with at most k weighings.

A partition of a fake coin problem P into three subproblems is well-behaved, if the following properties hold.

- (1) Size[P] $\leq 3^k$ for $k \geq 2$ implies that Size[left subproblem of P] $\leq 3^{k-1}$, Size[right subproblem of P] $\leq 3^{k-1}$, and Size[balanced subproblem of P] $\leq 3^{k-1}$.
- (2) Subproblems of size ≤ 3 can be solved with at most one weighing.

To show Properties 1.1, 1.2, and 1.3, it suffices to prove that our method has the following Properties 2.1, 2.2, 2.3, and 2.4 using 3^{k-1} as the partition bound of the three subproblems for a given problem of size S such that $3^{k-1} < S \leq 3^k$ in P[(m)] and P[(x, y)], and for a given problem of size S such that $3^{k-1} < S + 3 \leq 3^k$ in P[n]. Proofs of Properties 2.1, 2.2, and 2.3. are given in the Appendix. A proof of Property 2.4 is given in Induction Basis.

Property 2.1 (Size[P[n]]). If $6 \leq \text{Size}[P[n]] \leq 3^k - 3$ for $k \geq 2$, then Size[left subproblem of $P[n] \geq 3^{k-1}$, Size[right subproblem of $P[n] \geq 3^{k-1}$, and Size[balanced subproblem of $P[n] \geq 3^{k-1}$.

Property 2.2 (Size[P[(m)]]). If Size[P[(m)]] $\leq 3^k$ for $k \geq 2$, then Size[left subproblem of P[(m)]] $\leq 3^{k-1}$, Size[right subproblem of P[(m)]] $\leq 3^{k-1}$, and Size[balanced subproblem of P[(m)]] $\leq 3^{k-1}$.

Property 2.3 (Size[P[(x, y)]]). If Size[P[(x, y)]] $\leq 3^k$ for $k \geq 2$, then Size[left subproblem of P[(x, y)]] $\leq 3^{k-1}$, Size[right subproblem of P[(x, y)]] $\leq 3^{k-1}$, and Size[balanced subproblem of P[(x, y)]] $\leq 3^{k-1}$.

Property 2.4 (Problems of size ≤ 3). Problems P[(m)] and P[(x, y)] of size ≤ 3 can be solved with at most one weighing.

The maximum number of coins of P[(m)] solvable with at most k weighings is m such that $2m = 3^k - 1$ for $k \ge 1$. The maximum number of coins of P[(x, y)]solvable with at most k weighings is 3^k , when $x \ne y$ and $x + y = 3^k$, or $3^k - 1$, when x = y and $x + y = 3^k - 1$, for $k \ge 1$.

3. Related Work

A Dyson set is a set of vectors where each element is -1 or 0 or +1, the sum of each dimension part is 0, and all vectors and their (-1) multiplied vectors are distinct. For example, { (-1, +1), (0, -1), (+1, 0) } is a Dyson set for three-coin problem P[3].

3.1. Dyson method

Dyson method uses the following Dyson set for the fake coin problem of 4 coins. Here -1, 0, and +1 respectively stand for the left side, the outside, and the right side for the first and the second weighing. Dyson method uses the Dyson set for P[3] and the fourth coin d is kept outside for the first and the second weighing.

 $\begin{array}{rrr} a: & (-1,+1) \\ b: & (0,-1) \\ c: & (+1,0) \\ d: & (0,0) \end{array}$

A vector v of the Dyson set and its (-1) multiplied vector -v respectively show the results of all weighings when the coin is heavy and when the coin is light, where -1, 0, and +1 respectively stand for the left side going down, balanced, and the right side going down.

Based on this Dyson set, the first and the second weighings are as follows. The third weighing is performed when the first and the second weighings are all balanced.

1. a versus c

2. *b* versus *a* (if all balanced)

3. d versus a

3.2. Born method

Born method uses the following Dyson set for 4 coins.

- a: (-1, -1, -1)
- b: (+1,0,0)
- c: (0, +1, 0)
- d: (0, 0, +1)

Based on this Dyson set, three weighings are as follows. All weighings are predetermined.

- 1. a versus b
- 2. a versus c
- 3. a versus d

3.3. Our Method

We recursively produce subproblems for a given problem P[4] until we have subproblems of size ≤ 3 . We select 9 as the partition bound for P[4].

Bound computation for 4 $3^{\lceil \log_3(2 \times 4 + 3) \rceil - 1} = 3^2 = 9$

 $P[4]: 4 \rightarrow (2,2,0)$ with bound 9

 $P[(2,2)]: (2,2) \to ((1,1),(1,1),(0,0))$ with bound 3

P[(0)], P[(1,1)], and P[(0,0)] are of size ≤ 3 , which can be solved using at most one weighing.

Based on this solution, when the left side always goes down for example, three weightings are as follows,

- 1. a, b versus c, d with outside none
- 2. a, c versus b, d with outside none
- 3. a versus c (genuine coin) with outside d

3.4. Problem of 12 coins

Both Dyson method and Born method use the following Dyson set for P[12]. $\{(-1, -1, 0), (0, 0, +1), (+1, +1, -1), (-1, 0, -1), (0, +1, 0), (+1, -1, +1), (-1, 0, -1), (-1, 0,$

(0, -1, -1), (+1, 0, 0), (-1, +1, +1), (-1, 0, +1), (0, +1, -1), (+1, -1, 0)

Hence the resulting three weighings are as follows.

- 1. a, d, i, j versus c, f, h, l with outside b, e, g, k
- 2. a, f, g, l versus c, e, i, k with outside b, d, h, j
- $3. \quad c,d,g,k \text{ versus } b,f,i,j \text{ with outside } a,e,h,l$

Our method produces the following subproblems using partitions with bounds. Bound computation for 12 $3^{\lceil \log_3(2 \times 12+3) \rceil - 1} = 3^2 = 9$

 $P[12]: 12 \to (4, 4, 4)$ with bound 9

 $P[(4,4)]: (4,4) \to ((2,1), (2,1), (0,2))$ with bound 3

P[(2,1)] and P[(0,2)] are of size ≤ 3 , which can be solved with one weighing P[(4)]: $(4) \rightarrow (2,1,1)$ with bound 3

P[(2,1)], P[(1,2)], and P[(1)] are of size ≤ 3 , which can be solved with one weighing.

Based on this solution, when the left side always goes down for example, three weightings are as follows,

- 1: a, b, c, d versus e, f, g, h with outside i, j, k, l
- 2: a, b, e versus c, d, f with outside g, h
- 3: a versus b with outside f

4. Induction Basis

We give solutions to problems P[(x, y)] and P[(m)] of size ≤ 3 . As shown in Table 1, we can confirm that if $x + y \leq 3$, then P[(x, y)] is solvable with at most one weighing. As shown in Table 2, we can confirm that if $2m \leq 3$, then P[(m)] is solvable with at most one weighing. Notice that cases P[(0)] and P[(0, 0)] are not selected in actual weighings.

		Table 1: $P[(x, y)]$ of size ≤ 3
Problem	Partition	Description
(0,0)	_	We are done.
(1, 0)	_	The coin is heavy.
(0,1)	_	The coin is light.
(2,0)	(1, 0, 1)	One coin on the left side, another coin on the outside, and one genuine coin on the right side. If the left side goes down, then it is heavy. If balanced, then the outside coin is heavy.
(1, 1)	(1, 0, 1)	One heavy coin candidate on the left side, one light coin candi- date on the outside, and one genuine coin on the right side. If the left side goes down, then it is heavy. If balanced, then the outside coin is light.
(0,2)	(1, 0, 1)	One coin on the left side, another coin on the outside, and one genuine coin on the right side. If the left side goes up, then it is light. If balanced, then the outside coin is light.
(3,0)	(1, 1, 1)	One coin on the left side and another clin on the right side. If one of sides goes down, then the down side coin is heavy. If balanced, then the outside coin is heavy.
(2,1)	(1, 1, 1)	One heavy coin candidate on the left side and another heavy coin candidate on the right side. If one of sides goes down, then the down side coin is heavy. If balanced, then the outside coin is light.
(1,2)	(1, 1, 1)	One light coin candidate on the left side and another light coin candidate on the right side. If one of sides goes up, then the up side coin is light. If balanced, then the outside coin is heavy.
(0,3)	(1, 1, 1)	One coin on the left side and another coin on the right side. If one of sides goes up, then the up side coin is light. If balanced, then the outside coin is light.

Table 2: P[(m)] of size ≤ 3

		Table 2. $\Gamma[(m)]$ of size ≤ 5
Problem	Partition	Description
(0)	_	We are done.
(1)	(1, 0, 0)	One coin on the left side and one genuine coin on the right side.
		If the left side goes down, then it is heavy. If the left side goes
		up, then it is light.

5. Bounded Partition Method

We describe our bounded partition method in pseudo-code. For producing partitions, we use Standard Partition for P[n], Pair Partition for P[(x, y)], and Special Partition for P[(m)]. To implement P[n], P[(x, y)], and P[(m)], we respectively use procedures F[c, n], G[c, (x, y)], and H[c, m] with partition bound c. Partition bound c is computed from given n in procedure FC[n]. We use partition bound 3^{k-1} for the smallest integer k such that $2n \leq 3^k - 3$ for given n.

In Standard Partition, the number of the left side coins and the number of the right side coins are always same. This number is greater than or equal to the number of the outside coins except when n = 1.

 $\begin{array}{l} \text{StandardPartition}[n]:\\ \text{If }n=0 \text{ then }(0,0,0);\\ \text{If }n=1 \text{ then }(0,0,1);\\ \text{If }n=2 \text{ then }(1,1,0);\\ \text{If }(n\geq 3) \text{ and }(\text{Mod}[n,3]=0) \text{ then }\left(\frac{n}{3},\frac{n}{3},\frac{n}{3}\right);\\ \left\{(j,j,j) \text{ if }n=3j\right\}\\ \text{If }(n\geq 3) \text{ and }(\text{Mod}[n,3]=1) \text{ then }\left(\frac{n+2}{3},\frac{n+2}{3},\frac{n-4}{3}\right);\\ \left\{(j+1,j+1,j-1) \text{ if }n=3j+1\right\}\\ \text{If }(n\geq 3) \text{ and }(\text{Mod}[n,3]=2) \text{ then }\left(\frac{n+1}{3},\frac{n+1}{3},\frac{n-2}{3}\right);\\ \left\{(j+1,j+1,j) \text{ if }n=3j+2\right\}\end{array}$

In Special Partition, the number of the left side coins and the number of the right side coins are not same for m = 3j + 1 $(j \ge 0)$ case or m = 2 case. For these weighing cases, we add one genuine coin on the right side so that we have the same number of coins on both sides.

$$\begin{split} &\text{SpecialPartition}[m]:\\ &\text{If } m=0 \text{ then } (0,0,0);\\ &\text{If } m=1 \text{ then } (1,0,0);\\ &\text{If } m=2 \text{ then } (1,0,1);\\ &\text{If } (m\geq 3) \text{ and } (\text{Mod}[m,3]=0) \text{ then } \left(\frac{m}{3},\frac{m}{3},\frac{m}{3}\right);\\ &\left\{(j,j,j) \text{ if } m=3j\right\}\\ &\text{If } (m\geq 3) \text{ and } (\text{Mod}[m,3]=1) \text{ then } \left(\frac{m+2}{3},\frac{m-1}{3},\frac{m-1}{3}\right);\\ &\left\{(j+1,j,j) \text{ if } m=3j+1\right\}\\ &\text{If } (m\geq 3) \text{ and } (\text{Mod}[m,3]=2) \text{ then } \left(\frac{m+1}{3},\frac{m+1}{3},\frac{m-2}{3}\right);\\ &\left\{(j+1,j+1,j) \text{ if } m=3j+2\right\} \end{split}$$

Pair Partition produces $((x_1, y_1), (x_2, y_2), (x_3, y_3))$ such that $x_1 = x_2, y_1 = y_2, x_1 + y_1 \leq c$ and $x_2 + y_2 \leq c$ for (x, y) such that x + y > 3. When $x_1 + y_1 > c$ or $x_2 + y_2 > c$, overflow coins are added to y_3 .

PairPartition[c, (x, y)]: If $x + y \leq 3$ then ((x, y), (0, 0), (0, 0)) else ($(x_1, x_2, x_3) \leftarrow$ StandardPartition[x];

```
\begin{array}{l} (y_1, y_2, y_3) \leftarrow \text{StandardPartition}[y];\\ \text{If } x_1 + y_1 > c \text{ then } (y_1, y_3) \leftarrow (c - x_1, y_3 + y_1 + x_1 - c);\\ \text{If } x_2 + y_2 > c \text{ then } (y_2, y_3) \leftarrow (c - x_2, y_3 + y_2 + x_2 - c);\\ ((x_1, y_1), (x_2, y_2), (x_3, y_3));\\ ); \end{array}
```

```
 \begin{split} & \operatorname{FC}[n]: \\ & c \leftarrow 3^{\lceil \log_3(2n+3)\rceil -1}; \, \operatorname{F}[c,n]; \end{split}
```

As for availability of the use of one genuine coin, notice that if $n \ge 3$ and $n \ne 4$, then we can use one genuine coin after the first weighing by Standard Partition. If n = 4, then we can use one genuine coin after the second weighing by Pair Partition.

$$\begin{split} & \mathbf{F}[c,n]:\\ & (x,y,m) \leftarrow \mathrm{StandardPartition}[n];\\ & \mathrm{Print}[c,(x,y,m)];\\ & \mathrm{If \ the \ left \ side \ goes \ down \ and \ x+y>3 \ then \ \mathbf{G}[\frac{c}{3},(x,y)];\\ & \mathrm{If \ the \ right \ side \ goes \ down \ and \ y+x>3 \ then \ \mathbf{G}[\frac{c}{3},(y,x)];\\ & \mathrm{If \ balanced \ and \ }2m>3 \ then \ \mathbf{H}[\frac{c}{3},m]; \end{split}$$

 $\begin{array}{l} {\rm G}[c,(x,y)]:\\ ((x_1,y_1),(x_2,y_2),(x_3,y_3)) \leftarrow {\rm PairPartition}[c,(x,y)];\\ {\rm Print}[c,((x_1,y_1),(x_2,y_2),(x_3,y_3))];\\ {\rm If the left side goes down and } x_1+y_2>3 \ {\rm then} \ {\rm G}[\frac{c}{3},(x_1,y_2)];\\ {\rm If the right side goes down and } x_2+y_1>3 \ {\rm then} \ {\rm G}[\frac{c}{3},(x_2,y_1)];\\ {\rm If balanced and } x_3+y_3>3 \ {\rm then} \ {\rm G}[\frac{c}{3},(x_3,y_3)]; \end{array}$

$$\begin{split} & \mathrm{H}[c,m]:\\ & (x,y,t) \leftarrow \mathrm{SpecialPartition}[m];\\ & \mathrm{Print}[c,(x,y,t)];\\ & \{\mathrm{If}\;y=x-1\;\mathrm{then}\;\mathrm{we}\;\mathrm{add}\;\mathrm{one}\;\mathrm{genuine}\;\mathrm{coin}\;\mathrm{on}\;\mathrm{the}\;\mathrm{right}\;\mathrm{side}\;\mathrm{for}\;\mathrm{weighing}\;\\ & \mathrm{purpose}\;\mathrm{only}\}\\ & \mathrm{If}\;\mathrm{the}\;\mathrm{left}\;\mathrm{side}\;\mathrm{goes}\;\mathrm{down}\;\mathrm{and}\;x+y>3\;\mathrm{then}\;\mathrm{G}[\frac{c}{3},(x,y)];\\ & \mathrm{If}\;\mathrm{the}\;\mathrm{right}\;\mathrm{side}\;\mathrm{goes}\;\mathrm{down}\;\mathrm{and}\;y+x>3\;\mathrm{then}\;\mathrm{G}[\frac{c}{3},(y,x)];\\ & \mathrm{If}\;\mathrm{balanced}\;\mathrm{and}\;2t>3\;\mathrm{then}\;\mathrm{H}[\frac{c}{3},t]; \end{split}$$

Notice that Special Partition and Pair Partition are well-behaved partitions. But Standard Partition is not well-behaved. If we use Standard Partition for P[(4)], then Standard Partition of 4 is (2, 2, 0). It is not acceptable with bound 3, where $2 \times 4 \leq 9$ and 2 + 2 > 3. Special Partition of 4 is (2, 1, 1) and it is acceptable with bound 3, because $2 \times 4 \leq 9$, $2 + 1 = 1 + 2 \leq 3$ and $2 \times 1 \leq 3$.

Example (Problem of 39 coins). Subproblems of 4 weighings for P[39] are as follows.

Bound computation for 39 $3^{\lceil \log_3(2 \times 39 + 3) \rceil - 1} = 3^3 = 27$

 $\begin{array}{l} P[39]: \ 39 \to (13, 13, 13) \ \text{with bound } 27 \\ P[(13, 13)]: \ (13, 13) \to ((5, 4), (5, 4), (3, 5)) \ \text{with bound } 9 \\ P[(5, 4)]: \ (5, 4) \to ((2, 1), (2, 1), (1, 2)) \ \text{with bound } 3 \\ P[(3, 5)]: \ (3, 5) \to ((1, 2), (1, 2), (1, 1)) \ \text{with bound } 3 \\ P[(13)]: \ (13) \to (5, 4, 4) \ \text{with bound } 9 \\ P[(5, 4)]: \ (5, 4) \to ((2, 1), (2, 1), (1, 2)) \ \text{with bound } 3 \\ P[(4, 5)]: \ (4, 5) \to ((2, 1), (2, 1), (0, 3)) \ \text{with bound } 3 \\ P[(4)]: \ (4) \to (2, 1, 1) \ \text{with bound } 3 \\ P[(2, 1)], \ P[(1, 2)], \ P[(1, 1)], \ P[(0, 3)], \ \text{and } P[(1)] \ \text{are of size} \leq 3, \ \text{which can be solved with one weighing.} \end{array}$

These procedures, F[c, n], G[c, (x, y)], and H[c, m], replace one problem by its subproblem recursively until the size of a subproblem ≤ 3 in search mode. If we delete component conditions, "the left side goes down and", "the right side goes down and", and "balanced and", from if-conditions of these procedures, then we can enumerate all subproblems until the size of subproblems ≤ 3 .

6. Concluding Remarks

We have shown our bounded partition approach to the fake coin problems. With the help of one genuine coin available after the first or the second weighing, we can naturally have the three subproblems of a given problem. Our method is essentially a direct ternary search method for finding one fake coin based on the size of subproblems solvable with fewer weighings than the given problem.

References

- F. J. Dyson, The problem of the pennies, The Mathematical Gazette 30 (291) (1946) 231–234.
- [2] A. Born, C. A. J. Hurkens, G. J. Woeginger, How to detect a counterfeit coin: Adaptive versus non-adaptive solutions, Information Processing Letters 86 (3) (2003) 137–141.

Appendix

Property 2.1 (F[c, n]: $(x, y, m) \leftarrow \text{StandardPartition}[n]$). If $6 \le 2n \le 3^k - 3$ for $k \ge 2$, then $x + y \le 3^{k-1}$, $y + x \le 3^{k-1}$, and $2m \le 3^{k-1}$.

Proof of Property 2.1.

Case 1. n = 3j and (x, y, m) = (j, j, j) for $j \ge 1$ If $2n = 6j \le 3^k - 3$, then $2j \le 3^{k-1} - 1 \le 3^{k-1}$. Hence $x + y = y + x \le 3^{k-1}$ and $2m \le 3^{k-1}$.

Case 2. n = 3j + 1 and (x, y, m) = (j + 1, j + 1, j - 1) for $j \ge 1$ If $2n = 6j + 2 \le 3^k - 3$, then $6j + 6 \le 3^k + 1$. Hence 2m = 2j - 2, x + y = y + x = 2j + 2, and $2j - 2 < 2j + 2 \le 3^{k-1} + \frac{1}{3}$. Therefore $2j + 2 \le 3^{k-1}$, because both 2j + 2 and 3^{k-1} are integers. Notice that if we change the required precondition from $2n \le 3^k - 3$ to $2n \le 3^k$, then $2j + 2 \le 3^{k-1} + \frac{4}{3}$ and x + y can be larger than 3^{k-1} .

Case 3. n = 3j + 2 and (x, y, m) = (j + 1, j + 1, j) for $j \ge 1$ If $2n = 6j + 4 \le 3^k - 3$, then $6j + 6 \le 3^k - 1$. Hence 2m = 2j, x + y = y + x = 2j + 2, and $2j < 2j + 2 \le 3^{k-1} - \frac{1}{3} \le 3^{k-1}$.

Property 2.2 (H[c, m]: $(x, y, t) \leftarrow \text{SpecialPartition}[m]$). If $2m \leq 3^k$ for $k \geq 2$, then $x + y \leq 3^{k-1}, y + x \leq 3^{k-1}$, and $2t \leq 3^{k-1}$.

Proof of Property 2.2.

When m = 2 and $(2) \rightarrow (1, 0, 1)$, this property holds. Cases where $m \ge 3$ are as follows.

Case 1. m = 3j and (x, y, t) = (j, j, j) for $j \ge 1$ If $2m = 6j \le 3^k$, then $2j \le 3^{k-1}$. Hence $x + y = y + x = 2j \le 3^{k-1}$ and $2t = 2j \le 3^{k-1}$. Case 2. m = 3j + 1 and (x, y, t) = (j + 1, j, j) for $j \ge 1$ If $2m = 6j + 2 \le 3^k$, then $6j + 3 \le 3^k + 1$. Hence 2t = 2j, x + y = y + x = 2j + 1, and $2j < 2j + 1 \le 3^{k-1} + \frac{1}{3}$. Therefore $2j + 1 \le 3^{k-1}$, because both 2j + 1 and 3^{k-1} are integers. Case 3. m = 3j + 2 and (x, y, t) = (j + 1, j + 1, j) for $j \ge 1$ If $2m = 6j + 4 \le 3^k$, then $6j + 6 \le 3^k + 2$. Hence 2t = 2j, x + y = y + x = 2j + 2, $2j < 2j + 2 \le 3^{k-1} + \frac{2}{3}$.

Therefore $2j + 2 \leq 3^{k-1}$, because both 2j + 2 and 3^{k-1} are integers.

Property 2.3 (G[c, (x, y)]: $((x_1, z_1), (x_2, z_2), (x_3, z_3)) \leftarrow \text{PairPartition}[c, (x, y)])$. $(x_1, x_2, x_3) \leftarrow \text{StandardPartition}[x]; (y_1, y_2, y_3) \leftarrow \text{StandardPartition}[y]; \text{ and}$ Pair Partition with bound c produces $((x_1, z_1), (x_2, z_2), (x_3, z_3))$. If $x + y \leq 3^k$ for $k \geq 2$, then $c = 3^{k-1}, x_1 + z_2 \leq 3^{k-1}, x_2 + z_1 \leq 3^{k-1}$, and $x_3 + z_3 \leq 3^{k-1}$.

PROOF OF PROPERTY 2.3.

General Case: When neither x nor y is 1.

Because $x_1 = x_2$, $y_1 = y_2$, and $z_1 = z_2$, it suffices to show that if $x + y \le 3^k$ for $k \ge 2$, then $x_1 + z_1 \le 3^{k-1}$, $x_2 + z_2 \le 3^{k-1}$, and $x_3 + z_3 \le 3^{k-1}$. Notice that $c = 3^{k-1}$, $x + y \le 3c = 3^k$, $x = x_1 + x_2 + x_3$, and $y = y_1 + y_2 + y_3$. Case 1. If $x_1 + y_1 \le c$ and $x_2 + y_2 \le c$, then $(z_1, z_2, z_3) = (y_1, y_2, y_3)$. Hence $x_3 \le x_1 = x_2$, $y_3 \le y_1 = y_2$ and $x_3 + y_3 \le x_1 + y_1 \le c$. Case 2. If $x_1 + y_1 > c$ and $x_2 + y_2 > c$, then $(x_1 + z_1, x_2 + z_2, x_3 + z_3) = (c, c, x_3 + y_3 + (x_1 + y_1 - c) + (x_2 + y_2 - c))$. If $x + y \le 3c$, then $x + y = x_1 + x_2 + x_3 + z_1 + z_2 + z_3 \le 3c$. Hence $x_3 + z_3 \le 3c - (x_1 + z_1 + x_2 + z_2) = 3c - 2c = c$.

Exceptional Case: When x or y is 1.

$$\begin{split} & \mathbf{P}[(x,y)] \text{ can be } \mathbf{P}[(1,1)] \text{ or } \mathbf{P}[(x,1)] \text{ or } \mathbf{P}[(1,y)] \text{ for } x \neq 1 \text{ and } y \neq 1. \text{ It suffices to show that three cases of } \mathbf{P}[(x,1)], \text{ cases of } x+1=3^k, x+1=3^k-1, \\ & \text{and } x+1=3^k-2 \text{ for } k \geq 2, \text{ produce the size of subproblems } \leq 3^{k-1}. \\ & \text{Case 1.} \\ & \text{ If } x=3^k-1=3(3^{k-1}-1)+2, \text{ then } (x,1) \rightarrow ((3^{k-1}-1+1,0),(3^{k$$

Hence the size of all subproblems $\leq 3^{k-1}$.