A NOTE ON GENERALIZED REPUNIT NUMERICAL SEMIGROUPS

FEIHU LIU¹, GUOCE XIN^{2,*}, SUTING YE³ AND JINGJING YIN⁴

ABSTRACT. Let $A = (a_1, a_2, ..., a_n)$ be relative prime positive integers with $a_i \geq 2$. The Frobenius number F(A) is the largest integer not belonging to the numerical semigroup $\langle A \rangle$ generated by A. The genus g(A) is the number of positive integer elements that are not in $\langle A \rangle$. The Frobenius problem is to find F(A) and g(A) for a given sequence A. In this note, we study the Frobenius problem of $A = \left(a, ba + d, b^2a + \frac{b^2-1}{b-1}d, ..., b^ka + \frac{b^k-1}{b-1}d\right)$ and obtain formulas for F(A) and g(A) when $a \geq k-1$. Our formulas simplifies further for some special cases, such as repunit, Mersenne and Thabit numerical semigroups. The idea is similar to that in [[13],arXiv:2306.03459].

Mathematic subject classification: Primary 11D07; Secondary 11A67, 11B75, 20M14. Keywords: Numerical semigroup; Apéry set; Frobenius number; Genus; Pseudo-Frobenius number.

1. INTRODUCTION

In [13], we studied the Frobenius problem for the numerical semigroup $\langle A \rangle = \langle a, 2a + d, 2^2a + 3d, ..., 2^ka + (2^k - 1)d \rangle$. We show in this note that the same idea applies to the Frobenius problem for the numerical semigroup $\langle A \rangle = \langle a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d \rangle$. Let b = 2, we can obtain the results in [13]. The material in this note is better merged with [13], but that paper has been submitted.

Throughout this paper, \mathbb{Z} , \mathbb{N} and \mathbb{P} denote the set of all integers, non-negative integers and positive integers, respectively. We need to introduce some basic definitions in [25].

A subset S is a submonoid of N if $S \subseteq \mathbb{N}$, $0 \in S$ and S is closed under the addition in N. If $\mathbb{N} \setminus S$ is finite, then we say that S is a numerical semigroup. For a sequence (or set) $A = (a_1, a_2, ..., a_n)$ with $a_i \in \mathbb{P}$, we denoted by $\langle A \rangle$ the submonoid of N generated by A, that is,

$$\langle A \rangle = \left\{ \sum_{i=1}^{n} x_i a_i \mid n \in \mathbb{P}, x_i \in \mathbb{N}, 1 \le i \le n \right\}$$

If gcd(A) = 1, then $\langle A \rangle$ is a numerical semigroup (see [25]). If $\langle A \rangle$ is a numerical semigroup, then we say that A is a system of generators of $\langle A \rangle$. If no proper subsequence (or subset) of A generates $\langle A \rangle$, then we say that A is a minimal system of generators of $\langle A \rangle$, and the number of A is the embedding dimension of $\langle A \rangle$, denoted by e(A). Moreover, the minimal system of generators is finite and unique (see [25]).

If gcd(A) = 1, then we have the following definitions:

Date: June 19, 2023.

^{*} This work was partially supported by NSFC(12071311).

- (1) The Frobenius number F(A): The greatest integer not belonging to $\langle A \rangle$.
- (2) The genus (or Sylvester number) g(A): The cardinality of $\mathbb{N}\setminus\langle A\rangle$.
- (3) The pseudo-Frobenius numbers u: If $u \in \mathbb{Z} \setminus \langle A \rangle$ and $u + s \in \langle A \rangle$ for all $s \in \langle A \rangle \setminus \{0\}$.
- (4) The set PF(A): The set of pseudo-Frobenius numbers of $\langle A \rangle$ (see [21]).
- (5) The type t(A) of $\langle A \rangle$: The cardinality of PF(A).

For more knowledge about numerical semigroup, see [2, 25].

The Frobenius number F(A) has been widely studied. For $A = (a_1, a_2)$, Sylvester [29] obtained $F(A) = a_1a_2 - a_1 - a_2$ in 1882. For $n \ge 3$, F. Curtis [6] proved that the F(A) can not be given by closed formulas of a certain type. However, many special cases have been studied such as arithmetic progression in [3, 18, 26], geometric sequences in [16], triangular and tetrahedral sequences in [19]. For more special sequences, see [11, 12, 17, 20, 30, 31]. Many special numerical semigroups are also considered, such as Fibonacci in [15], Mersenne in [24], repunit in [23], squares and cubes in [10], Thabit in [22] and other numerical semigroups in [8, 27, 28].

The motivation of this paper comes from the repunit numerical semigroup: $\langle \{\frac{b^{n+i}-1}{b-1} \mid i \in \mathbb{N}\} \rangle$. Essentially, its minimal system of generators is $S(b,n) = \left(\frac{b^{n}-1}{b-1}, \frac{b^{n+1}-1}{b-1}, \dots, \frac{b^{2n-1}-1}{b-1}\right)$ ([23]), that is the embedding dimension e(S(b,n)) = n. In this paper, we study the following more general case

$$A = (a, Ha + dB) = \left(a, ba + d, b^{2}a + \frac{b^{2} - 1}{b - 1}d, \dots, b^{k}a + \frac{b^{k} - 1}{b - 1}d\right)$$

where $a, d, b, k \in \mathbb{P}, b \geq 2$ and gcd(a, d) = 1. Here we have the embedding dimension $e(A) \leq k+1$. We can observe that

- (1) If $a = 2^n 1$, b = 2, d = 1, k = n 1, then $\langle A \rangle$ is the Mersenne numerical semigroup S(n) in [24].
- (2) If $a = 3 \cdot 2^n 1$, b = 2, d = 1, k = n + 1, then $\langle A \rangle$ is the Thabit numerical semigroup T(n) in [22].
- (3) If $a = (2^m 1) \cdot 2^n 1$, b = 2, d = 1, k = n + m 1, then $\langle A \rangle$ is a class of numerical semigroups in [7].
- (4) If $a = (2^m + 1) \cdot 2^n (2^m 1)$, b = 2, $d = 2^m 1$, $k \in \{n + 1, n + m 1, n + m\}$, then $\langle A \rangle$ is a class of numerical semigroups in [28].
- (5) If $a = m(2^k 1) + 2^{k-1} 1$, b = 2, $k \ge 3$, then $\langle A \rangle$ is a class of numerical semigroups in [13].
- (6) If $a = \frac{b^n 1}{b 1}$, $b \ge 2$, d = 1, k = n 1, then $\langle A \rangle$ is the repunit numerical semigroups S(b, n) in [23].
- (7) If $a = b^{n+1} + \frac{b^n 1}{b-1}$, $b \ge 2$, d = 1, k = n+1, then $\langle A \rangle$ is a class of numerical semigroups in [8].
- (8) If $a = (b+1)b^n 1$, $b \ge 2$, d = b 1, k = n + 1, then $\langle A \rangle$ is the Thabit numerical semigroup $T_{b,1}(n)$ of the first kind base b in [27].

The main purpose of this paper is to solve the Frobenius problem of $A = A = (a, Ha + dB) = \left(a, ba + d, b^2a + \frac{b^2-1}{b-1}d, \dots, b^ka + \frac{b^k-1}{b-1}d\right)$. Using similar discussions as in paper [13], we

can obtain formulas of F(A) and g(A) in Theorem 3.5. Specifically, if k = n - 1 and $a = \frac{b^n - 1}{b - 1}$, then we can further obtain

- (1) The Frobenius number $F(A) = (b^n + d 1) \cdot \frac{b^n 1}{b 1} d$. (2) The genus $g(A) = \frac{(b^n b)(b^n + d 1)}{2(b 1)} + \frac{b^n(n 1)}{2}$. (3) The set of Pseudo-Frobenius number $PF(A) = \{F(A), F(A) d, ..., F(A) (n 2)d\}$.
- (4) The type t(A) = n 1.

The paper is organized as follows. In Section 2, we provide some necessary lemmas and related results. In Section 3, we establish Theorem 3.5, which gives formulas of Frobenius number F(A) and genus g(A) where $\left(a, ba+d, b^2a+\frac{b^2-1}{b-1}d, \dots, b^ka+\frac{b^k-1}{b-1}d\right)$ and $a \ge k-1$. Section 4 focus on application of Theorem 3.5 for special a and k. It is related to repunit numerical semigroup. Section 6 is a concluding remark.

2. Preliminary

It is convenient to use the short hand notation $A := (a, B) = (a, b_1, b_2, ..., b_k)$. Let $gcd(A) = 1, a, b_i \in \mathbb{P}$. The set

$$\langle A \rangle = \left\{ ax + \sum_{i=1}^{k} b_i x_i \mid x, x_i \in \mathbb{N} \right\}$$

is a numerical semigroup. Let $w \in \langle A \rangle \setminus \{0\}$. The Apéry set of w in $\langle A \rangle$ is $Ape(A, w) = \{s \in A\}$ $\langle A \rangle \mid s - w \notin \langle A \rangle$. In [25], we can obtain

$$Ape(A, w) = \{N_0, N_1, N_2, ..., N_{w-1}\},\$$

where $N_r := \min\{a_0 \mid a_0 \equiv r \mod w, a_0 \in \langle A \rangle\}, 0 \leq r \leq w - 1$. We usually take w := a.

A. Brauer and J. E. Shockley [4], E. S. Selmer [26] gave the following results respectively.

Lemma 2.1 ([4], [26]). Suppose $A := (a, B) = (a, b_1, b_2, ..., b_k)$. The Apéry set of a in $\langle A \rangle$ is $Ape(A, a) = \{N_0, N_1, N_2, ..., N_{a-1}\}$. Then the Frobenius number and genus of A are respectively:

$$F(A) = F(a, B) = \max_{r \in \{0, 1, \dots, a-1\}} N_r - a,$$
$$g(A) = g(a, B) = \frac{1}{a} \sum_{r=1}^{a-1} N_r - \frac{a-1}{2}.$$

Now we define the following order relation in \mathbb{Z} : $a \preceq_{\langle A \rangle} b$ if $b - a \in \langle A \rangle$. It is proved that this relation $\preceq_{\langle A \rangle}$ is an order relation in [25].

Lemma 2.2 (Proposition 2.20, [25]). Let $\langle A \rangle$ be a numerical semigroup. Then

$$PF(A) = \left\{ w - a \mid w \in \max_{\preceq_{\langle A \rangle}} Ape(A, a) \right\},\$$

where $Ape(A, a) = \{N_0, N_1, ..., N_{a-1}\}.$

Note that $N_0 = 0$ for all A. By the definition of N_r , we can easily obtain the following result.

Proposition 2.3. Let $A = (a, b_1, ..., b_k)$, gcd(a, d) = 1, $d \in \mathbb{P}$. Then we have

$$\{N_0, N_1, N_2, \dots, N_{a-1}\} = \{N_{d \cdot 0}, N_{d \cdot 1}, N_{d \cdot 2}, \dots, N_{d \cdot (a-1)}\}.$$
(1)

Lemma 2.4 ([13]). Let $A = (a, h_1a + db_1, ..., h_ka + db_k)$, $k, h, d \in \mathbb{P}$ and gcd(A) = 1, $m \in \mathbb{N}$, gcd(a, d) = 1. For a given $0 \le r \le a - 1$, we have

$$N_{dr} = \min\left\{O_B^H(ma+r) \cdot a + (ma+r)d \mid m \in \mathbb{N}\right\},\tag{2}$$

where

$$O_B^H(M) := \min\left\{\sum_{i=1}^k h_i x_i \mid \sum_{i=1}^k b_i x_i = M, \ x_i, M \in \mathbb{N}, 1 \le i \le k\right\}.$$

By Lemma 2.4, we can define an intermediate function with respect to $m \in \mathbb{N}$, namely:

$$N_{dr}(m) := O_B^H(ma+r) \cdot a + (ma+r)d.$$

We also need the following definition. For a given positive integers sequence $B = (b_1, b_2, ..., b_k)$, $1 = b_1 < b_2 < \cdots < b_k$ and $M \in \mathbb{N}$, let

$$opt_B(M) := \min\left\{\sum_{i=1}^k x_i \mid \sum_{i=1}^k b_i x_i = M, \ M, x_i \in \mathbb{N}, 1 \le i \le k\right\}.$$
 (3)

The problem $opt_B(M)$ is called the change-making problem [1]. A strategy is called greedy, that is to use as many of the maximum as possible, than as many of the next one as possible, and so on. We denote $grd_B(M)$ the number of elements used in B by the greedy strategy. Then we have $opt_B(M) \leq grd_B(M)$. If the greedy solution is always optimal, i.e., $opt_B(M) = grd_B(M)$ for all M > 0, then we call the sequence B orderly; Otherwise, we call sequence B non-orderly.

Lemma 2.5 (One-Point Theorem, [5, 9, 14]). Suppose $B' = (1, b_1, ..., b_k)$ is orderly and $b_{k+1} > b_k$. Let $s = \lceil b_{k+1}/b_k \rceil$. Then the sequence $B = (1, b_1, ..., b_k, b_{k+1})$ is orderly if and only if $opt_B(sb_k) = grd_B(sb_k)$.

3. The Frobenius Problem for
$$A = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right)$$

Now we consider the case $A = (a, Ha + dB) = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right), b \ge 2$, i.e., $H = (b, b^2, ..., b^k)$ and $B = \left(1, \frac{b^2 - 1}{b - 1}, ..., \frac{b^k - 1}{b - 1}\right)$. We first consider

$$opt_B(M) = \min\left\{\sum_{i=1}^k x_i \mid \sum_{i=1}^k \frac{b^i - 1}{b - 1} x_i = M, \ M, x_i \in \mathbb{N}, 1 \le i \le k\right\}.$$

By the *One-Point Theorem*, we can obtain the following result.

Lemma 3.1. Let $k \in \mathbb{P}$ and $b \geq 2$. Then the sequence $B = \left(1, \frac{b^2-1}{b-1}, \dots, \frac{b^k-1}{b-1}\right)$ is orderly, i.e., $opt_B(M) = grd_B(M)$ for all $M \in \mathbb{P}$.

Proof. By induction, we first know that the sequence (1) and $\left(1, \frac{b^2-1}{b-1}\right)$ are orderly sequences. Suppose $\left(1, \frac{b^2-1}{b-1}, \dots, \frac{b^{k-1}-1}{b-1}\right)$ is orderly. We have $s = \left\lceil \frac{b^k-1}{b^{k-1}-1} \right\rceil = \left\lceil \frac{b(b^{k-1}-1)+b-1}{b^{k-1}-1} \right\rceil = b+1$. By $(b+1) \cdot \frac{b^{k-1}-1}{b-1} = \frac{b^k-1}{b-1} + b \cdot \frac{b^{k-2}-1}{b-1}$, we have $opt_B(s(\frac{b^{k-1}-1}{b-1})) = grd_B(s(\frac{b^{k-1}-1}{b-1})) = b+1$. By Lemma 2.5, the sequence B is orderly. This completes the proof.

For a given $M \in \mathbb{P}$ and $B = \left(1, \frac{b^2-1}{b-1}, \dots, \frac{b^k-1}{b-1}\right)$, by $b \cdot \frac{b^i-1}{b-1} + 1 = \frac{b^{i+1}-1}{b-1}$ and Lemma 3.1, we have the following properties for x_i in $opt_B(M)$.

(1) $x_k = \left\lfloor \frac{(b-1)M}{b^k - 1} \right\rfloor$. (2) $x_i \in \{0, 1, ..., b\}$ for every $1 \le i \le k - 1$. (3) if $2 \le i \le k - 1$ and $x_i = b$, then $x_1 = \cdots = x_{i-1} = 0$.

If a solution $X = (x_1, x_2, ..., x_k)$ of $opt_B(M)$ satisfies the above conditions (1), (2) and (3), we call X the greedy presentation of M. We define R(M) to be the set of all greedy presentations of 0, 1, 2, ..., M. We define a colexicographic order on R(M) as follows:

$$\begin{aligned} &(x_1', x_2', ..., x_k') \preceq (x_1, x_2, ..., x_k) \\ &\Longleftrightarrow x_i' = x_i \text{ for any } i > 0 \text{ or} \\ &x_j' < x_j, x_i' = x_i \text{ for a certain } j > 0 \text{ and any } i > j. \end{aligned}$$

Obviously, the order relation \leq is a total order on R(M).

Remark 3.2. Lemma 3.1 essentially provides the construction process of $X = (x_1, x_2, ..., x_k)$ in R(M).

Now we consider the $O_B^H(M)$ for $B = \left(1, \frac{b^2-1}{b-1}, \dots, \frac{b^k-1}{b-1}\right)$, $H = (b, b^2, \dots, b^k)$ and any $M \in \mathbb{P}$. We have

$$O_B^H(M) = \min\left\{\sum_{i=1}^k b^i x_i \mid \sum_{i=1}^k \frac{b^i - 1}{b - 1} x_i = M, \ x_i \in \mathbb{N}, 1 \le i \le k\right\}$$
$$= \min\left\{(b - 1)M + \sum_{i=1}^k x_i \mid \sum_{i=1}^k \frac{b^i - 1}{b - 1} x_i = M, \ x_i \in \mathbb{N}, 1 \le i \le k\right\}.$$

Lemma 3.3. Let $A = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right)$, $a, d, b, k \in \mathbb{P}$, $b \geq 2$ and gcd(a, d) = 1. Suppose $a \geq k - 1$. Then $N_{dr}(m)$ is increasing with respect to $m \in \mathbb{N}$. More precisely, if $X = (x_1, x_2, ..., x_k)$ is a greedy presentation of r, then

$$N_{dr} = \left(\sum_{i=1}^{k} x_i\right) a + r((b-1)a + d) = \left(\sum_{i=1}^{k} b^i x_i\right) a + rd.$$
 (4)

Proof. Given a $m \in \mathbb{N}$. Assuming $X' = (x'_1, ..., x'_k)$ and $X = (x_1, ..., x_k)$ are greedy presentations of (m+1)a + r and ma + r, respectively. We can easily get $x'_k \ge x_k$. Furthermore

$$N_{dr}(m) = \left((b-1)(ma+r) + \sum_{i=1}^{k} x_i \right) a + (ma+r)d.$$

Therefore we have

$$N_{dr}(m+1) - N_{dr}(m) = (b-1)a^2 + ad + \left(\sum_{i=1}^k x'_i - \sum_{i=1}^k x_i\right)a$$

$$\geq \left((b-1)a + d + \sum_{i=1}^{k-1} x'_i - \sum_{i=1}^{k-1} x_i \right) a$$

$$\geq ((b-1)a + d - b - (k-2)(b-1))a$$

$$= ((b-1)(a-k+1) + d - 1)a \geq 0.$$

Then $N_{dr}(m)$ is increasing with respect to $m \in \mathbb{N}$. Moreover $N_{dr} = N_{dr}(0)$. This completes the proof.

In Equation (4), suppose $X = (x_1, x_2, ..., x_k)$ is a greedy presentation of r, we call $w(r) = \sum_{i=1}^{k} b^i x_i$ the *weight* of r. Now we have the following result.

Lemma 3.4. Suppose $X' = (x'_1, ..., x'_k) \in R(M)$ and $X = (x_1, ..., x_k) \in R(M)$ are greedy presentations of r_1 and r_2 , respectively. If $(x'_1, ..., x'_k) \preceq (x_1, ..., x_k)$, then we have $w(r_1) \leq w(r_2)$.

Proof. For $X = (x_1, x_2, ..., x_k) \in R(M)$, it is easy to obtain that the maximum of $\sum_{i=1}^{k-1} b^i x_i$ is b^k . At this point, the possible values for $(x_1, ..., x_{k-1})$ are

(b, b-1, b-1, ..., b-1), (0, b, b-1, ..., b-1), ..., (0, 0, 0, ..., b).

We consider the following three cases.

If $x'_i = x_i$ for any $1 \le i \le k$, then $w(r_1) = w(r_2)$. If $x'_k < x_k$, then $x_k \ge x'_k + 1$ and $\frac{k}{k-1}$

$$w(r_1) = \sum_{i=1}^{n} b^i x'_i = \sum_{i=1}^{n-1} b^i x'_i + b^k x'_k \le b^k + b^k x'_k = b^k (1 + x'_k) \le b^k x_k \le w(r_2).$$

If there is a certain $1 \le u \le k-1$, so that $x'_u < x_u$ and $x'_{u+1} = x_{u+1}, ..., x'_k = x_k$. Similarly we have

$$w(r_1) = \sum_{i=1}^k b^i x'_i = \sum_{i=1}^u b^i x'_i + \sum_{j=u+1}^k b^j x'_j$$

$$\leq b^u (1+x'_u) + \sum_{j=u+1}^k b^j x'_j = b^u (1+x'_u) + \sum_{j=u+1}^k b^j x_j$$

$$\leq b^u x_u + \sum_{j=u+1}^k b^j x_j \leq w(r_2).$$

This completes the proof.

Now we can obtain the main result of this section.

Theorem 3.5. Let $A = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right)$, $a, d, b, k \in \mathbb{P}$, gcd(a, d) = 1, $b \ge 2$ and $a \ge k - 1$. We have

$$F(A) = \left((b-1)a - b + d + \left(\sum_{i=1}^{k} x_i\right)_{a-1} \right) a - d,$$

$$g(A) = \sum_{r=1}^{a-1} \left(\sum_{i=1}^{k} x_i\right)_r + \frac{(a-1)((b-1)a+d-1)}{2},$$

where $(\sum_{i=1}^{k} x_i)_r$ is the sum of elements in the greedy presentation of r. Obviously $x_k = \left|\frac{(b-1)r}{b^k-1}\right|$.

Proof. By Lemma 3.4 and Equation (4), we have

$$\max N_{dr} = N_{d(a-1)} = \left(\sum_{i=1}^{k} x_i\right)_{a-1} \cdot a + (a-1)((b-1)a+d).$$

Further from Lemma 2.1, we can obtain the Frobenius number formula F(A). From Equation (4) and Lemma 2.1 again, we can obtain the genus g(A). This completes the proof.

4. A GENERALIZATION OF REPUNIT NUMERICAL SEMIGROUP

In this section we mainly focus on the generalization of repunit numerical semigroup. For $A = \left(a, ba + d, b^2a + \frac{b^2-1}{b-1}d, \dots, b^ka + \frac{b^k-1}{b-1}d\right)$, suppose k := n-1, $a := \frac{b^n-1}{b-1}$, $b \ge 2$. Therefore we have

$$A = \left(\frac{b^n - 1}{b - 1}, b \cdot \frac{b^n - 1}{b - 1} + d, b^2 \cdot \frac{b^n - 1}{b - 1} + \frac{b^2 - 1}{b - 1}d, \dots, b^{n - 1} \cdot \frac{b^n - 1}{b - 1} + \frac{b^{n - 1} - 1}{b - 1}d\right).$$

We consider the numerical semigroup $\langle A \rangle$ generated by A.

By
$$a - 1 = \frac{b^n - 1}{b - 1} - 1 = b \cdot \frac{b^{n - 1} - 1}{b - 1}$$
, we have

$$R(a - 1) = \left\{ (x_1, x_2, ..., x_{n - 1}) \mid 0 \le x_i \le b \text{ for } 1 \le i \le n - 1; \\ \text{if } x_i = b, \text{ then } x_j = 0 \text{ for } j \le i - 1 \right\}.$$
(5)

Theorem 4.1. For $A = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right)$, suppose k := n - 1, $a := \frac{b^n - 1}{b - 1}$, $b, n \ge 2$ and gcd(a, d) = 1. We have

$$F(A) = (b^n + d - 1) \cdot \frac{b^n - 1}{b - 1} - d,$$

$$g(A) = \frac{(b^n - b)(b^n + d - 1)}{2(b - 1)} + \frac{b^n(n - 1)}{2}$$

Proof. By Theorem 3.5 and $a - 1 = b \cdot \frac{b^{n-1}-1}{b-1}$, we have

$$F(A) = \left((b-1) \cdot \frac{b^n - 1}{b-1} - b + d + b \right) \cdot \frac{b^n - 1}{b-1} - d$$
$$= (b^n + d - 1) \cdot \frac{b^n - 1}{b-1} - d.$$

Now we consider the genus g(A). Based on the composition of R(a-1) mentioned above, we have

$$\begin{split} \sum_{r=1}^{a-1} \left(\sum_{i=1}^{n-1} x_i \right)_r &= (n-1)b^{n-2} \cdot \sum_{i=0}^{b-1} i + \sum_{j=1}^{n-1} \left(b \cdot b^{n-1-j} + (n-1-j) \cdot b^{n-2-j} \cdot \sum_{i=0}^{b-1} i \right) \\ &= \frac{(n-1)b^{n-1}(b-1)}{2} + \sum_{j=1}^{n-1} \left(b^{n-j} + \frac{b^{n-1-j}(b-1)(n-j-1)}{2} \right) \\ &= \frac{b^n - b}{2(b-1)} + \frac{b^n(n-1)}{2}. \end{split}$$

By Theorem 3.5, we have

$$g(A) = \frac{(b^n - b)(b^n + d - 1)}{2(b - 1)} + \frac{b^n(n - 1)}{2}.$$

This completes the proof.

Let d = 1 in Theorem 4.1, the $\langle A \rangle$ becomes the repunit numerical semigroup ([23]). Clearly, we have the following corollary.

Corollary 4.2 (Theorem 20, Theorem 25, [23]). Let $n, b \ge 2$, d = 1 in Theorem 4.1. Then $S(b, n) = \langle A \rangle$ is the repunit numerical semigroup in [23]. Furthermore, we have

$$F(S(b,n)) = \frac{b^n - 1}{b - 1}b^n - 1,$$

$$g(S(b,n)) = \frac{b^n}{2}\left(\frac{b^n - 1}{b - 1} + n - 1\right).$$

Theorem 4.3. For $A = \left(a, ba + d, b^2a + \frac{b^2 - 1}{b - 1}d, ..., b^ka + \frac{b^k - 1}{b - 1}d\right)$, suppose k := n - 1, $a := \frac{b^n - 1}{b - 1}$, $b, n \ge 2$ and gcd(a, d) = 1. Then t(A) = n - 1. Furthermore

$$PF(A) = \{F(A), F(A) - d, \dots, F(A) - (n-2)d\}.$$

Proof. Let $X = (x_1, x_2, ..., x_{n-1})$ be a greedy presentation of $r, 0 \le r \le a-1$. By Equation (4), we have

$$N_{dr} = \left(\sum_{i=1}^{n-1} b^{i} x_{i}\right) a + \sum_{i=1}^{n-1} \frac{b^{i} - 1}{b - 1} x_{i} d$$
$$= \left(\sum_{i=1}^{n-1} b^{i} x_{i}\right) a + \sum_{i=1}^{n-1} \frac{b^{i} x_{i}}{b - 1} d - \sum_{i=1}^{n-1} \frac{x_{i}}{b - 1} d.$$

For a given $0 \le r \le a-1$, we have $r = s \cdot \frac{b^{n-1}-1}{b-1} + r'$, $0 \le s \le b$ and $0 \le r' \le \frac{b^{n-1}-b}{b-1}$. Therefore, under the order relation $\preceq_{\langle A \rangle}$, the greedy presentations of the maximal elements in R(a-1) are contained in the following elements:

$$(b,b-1,b-1,...,b-1,b-1),\;(0,b,b-1,...,b-1,b-1),\;(0,0,b,...,b-1,b-1),\;...,\;(0,0,0,...,0,b).$$

For the above greedy presentations, we always have

$$\sum_{i=1}^{n-1} b^i x_i = b^n.$$

For the above maximal elements, let $X' = (x'_1, ..., x'_{n-1}) \in R(a-1)$ and $X = (x_1, ..., x_{n-1}) \in R(a-1)$ be greedy presentations of r_1 and r_2 , respectively. We need to consider the following case

$$N_{dr_1} - N_{dr_2} = \sum_{i=1}^{n-1} \frac{x_i}{b-1} d - \sum_{i=1}^{n-1} \frac{x'_i}{b-1} d = td, \quad t = 1, 2, ..., n-2$$

Now we need to determine whether td belongs to $\langle A \rangle$. If $td \in \langle A \rangle$, there would exist $(y_0, y_1, ..., y_{n-1})$ such that

$$td = y_0 a + y_1 (ba + d) + \dots + y_{n-1} \left(b^{n-1} a + \frac{b^{n-1} - 1}{b - 1} d \right).$$

These $y_0, ..., y_{n-1} \in \mathbb{N}$ are not all 0. Then we have

$$(n-2)d \ge \left(t - y_1 - y_2\frac{b^2 - 1}{b - 1} - \dots - y_{n-1}\frac{b^{n-1} - 1}{b - 1}\right)d = (y_0 + by_1 + \dots + b^{n-1}y_{n-1})a > 0.$$

By gcd(a,d) = 1, we have $a | \left(t - y_1 - y_2 \frac{b^2 - 1}{b - 1} - \dots - y_{n-1} \frac{b^{n-1} - 1}{b - 1} \right)$. However $a > n - 2 \ge t$, we have $t = y_1 + y_2 \frac{b^2 - 1}{b - 1} + \dots + y_{n-1} \frac{b^{n-1} - 1}{b - 1}$. So $(y_0 + by_1 + \dots + b^{n-1}y_{n-1})a = 0$, a contradiction.

Therefore we have

$$\max_{\leq \langle A \rangle} Ape(A, a) = \{ N_{dr} \mid r = a - \{1, 2, ..., n - 1\} \}.$$

By Lemma 2.2, we have

$$PF(A) = \{N_{dr} - a \mid r = a - \{1, 2, ..., n - 1\}\}$$

= $\left\{ (b^n - 1)a + \frac{b^n d}{b - 1} - \left\{ \frac{bd}{b - 1}, \frac{(2b - 1)d}{b - 1}, \frac{(3b - 2)d}{b - 1}, ..., \frac{((n - 1)b - (n - 2))d}{b - 1} \right\} \right\}$
= $\{F(A), F(A) - d, ..., F(A) - (n - 2)d\}.$

This completes the proof.

Similarly, we can obtain the following corollary.

Corollary 4.4 (Theorem 23, [23]). Let $n, b \ge 2$, d = 1 in Theorem 4.1. Then $S(b, n) = \langle A \rangle$ is the repunit numerical semigroup in [23]. We have t(S(b, n)) = n - 1 and

$$PF(S(b,n)) = \{F(S(n)), F(S(n)) - 1, \dots, F(S(n)) - (n-2)\}.$$

5. Concluding Remark

Our model $A = \left(a, ba + d, b^2a + \frac{b^2-1}{b-1}d, \dots, b^ka + \frac{b^k-1}{b-1}d\right)$ can be used to explain the following eight numerical semigroups in the literature.

1. If
$$a = 2^n - 1$$
, $b = 2$, $d = 1$, $k = n - 1$ and $n \ge 2$, then

$$A = (2^n - 1, 2^{n+1} - 1, 2^{n+2} - 1..., 2^{2n-1} - 1),$$

and the $\langle A \rangle$ is the Mersenne numerical semigroups S(n) in [24].

2. If $a = 3 \cdot 2^n - 1$, b = 2, d = 1, k = n + 1 and $n \ge 1$, then

$$A = (3 \cdot 2^{n} - 1, 3 \cdot 2^{n+1} - 1, 3 \cdot 2^{n+2} - 1, ..., 3 \cdot 2^{2n+1} - 1),$$

and the $\langle A \rangle$ is the Thabit numerical semigroups T(n) in [22].

If
$$a = (2^m - 1) \cdot 2^n - 1$$
, $b = 2$, $d = 1$, $k = n + m - 1$, $n \ge 1$ and $2 \le m \le 2^n$, then

$$A = ((2^{m} - 1) \cdot 2^{n} - 1, (2^{m} - 1) \cdot 2^{n+1} - 1, ..., (2^{m} - 1) \cdot 2^{2n+m-1} - 1),$$

and the $\langle A \rangle$ is a class of numerical semigroups S(m, n) in [7].

4. Let $m, n \in \mathbb{N}$ and $m \geq 2$ and

$$\delta = \begin{cases} 1 & \text{if } n = 0; \\ m & \text{if } n \neq 0, m \le n; \\ m - 1 & \text{if } n \neq 0, m > n. \end{cases}$$

If $a = (2^m + 1) \cdot 2^n - (2^m - 1)$, b = 2, $d = 2^m - 1$ and $k = n + \delta$, then the $\langle A \rangle$ is a class of numerical semigroups GT(n,m) in [28].

5. If $a = m(2^k - 1) + 2^{k-1} - 1$, b = 2, $m \ge 1$, $k \ge 3$, then the $\langle A \rangle$ is a class of numerical semigroups in [13].

6. If
$$a = \frac{b^n - 1}{b - 1}$$
, $b \ge 2$, $d = 1$, $k = n - 1$ and $n \ge 2$, then

$$A = \left(\frac{b^n - 1}{b - 1}, \frac{b^{n+1} - 1}{b - 1}, \frac{b^{n+2} - 1}{b - 1}, \dots, \frac{b^{2n-1} - 1}{b - 1}\right),$$

and the $\langle A \rangle$ is the repunit numerical semigroups S(b, n) in [23].

7. If
$$a = b^{n+1} + \frac{b^n - 1}{b - 1}$$
, $b \ge 2$, $d = 1$, $k = n + 1$, then

$$A = \left(b^{n+1} + \frac{b^n - 1}{b - 1}, b^{n+2} + \frac{b^{n+1} - 1}{b - 1}, \dots, b^{2n+2} + \frac{b^{2n+1} - 1}{b - 1}\right),$$

and the $\langle A \rangle$ is a class of numerical semigroups S(b, n) in [8].

8. If
$$a = (b+1)b^n - 1$$
, $b \ge 2$, $d = b - 1$, $k = n + 1$, then

$$A = ((b+1)b^n - 1, (b+1)b^{n+1} - 1, ..., (b+1)b^{2n+1} - 1),$$

and the $\langle A \rangle$ is the Thabit numerical semigroup $T_{b,1}(n)$ of the first kind base b in [27].

Acknowledgements: This work was partially supported by the National Natural Science Foundation of China [12071311].

References

- A. Adamaszek, M. Adamaszek, Combinatorics of the change-making problem, European. J. Comb. 31 (2010), 47–63.
- [2] A. Assi, M. D'Anna, P. A. García-Sánchez, Numerical Semigroups and Applications, Second Edition. Vol.3, RSMS Springer, Cham (2020).
- [3] A. Brauer, On a problem of partitions, Amer. J. Math. 64 (1942), 299–312.
- [4] A. Brauer, J. E. Shockley, On a problem of Frobenius, J. Reine Angew. Math. 211 (1962), 215– 220.
- [5] L. J. Cowen, R. Cowen, A. Steinberg, Totally greedy coin sets and greedy obstructions, Electron. J. Comb. 15 (2008), #R90.

3.

- [6] F. Curtis, On formulas for the Frobenius number of a numerical semigroup, Math. Scand. 67 (1990), 190–192.
- [7] Z. Gu, X. Tang, The Frobenius probelm for a class of numerical semigroups, Int. J. Number Theory. 13 (2017), 1335–1347.
- [8] Z. Gu, On the numerical semigroup generated by $\{b^{n+1+i} + \frac{b^{n+i}-1}{b-1} \mid i \in \mathbb{N}\}$, Discrete Math. Appl. 30(4) (2020), 257–264.
- T. C. Hu, M. L. Lenard, Optimality of a Heuristic solution for a class of Knapsack problems, Operations Research. 24(1) (1976), 193–196.
- [10] M. Lepilov, J. O'Rourke, I. Swanson, Frobenius numbers of numerical semigroups generated by three consecutive squares or cubes, Semigroup Forum. 91 (2015), 238–259.
- [11] F. Liu, G. Xin, On Frobenius formulas of power sequences, arXiv:2210.02722, (2022).
- [12] F. Liu, G. Xin, A combinatorial approach to Frobenius numbers of some special sequences (Complete Version), arXiv:2303.07149. (2023).
- [13] F. Liu, G. Xin, A generalization of Mersenne and Thabit numerical semigroups, arXiv:2306.03459. (2023).
- [14] M. J. Magazine, G. L. Nemhauser, L. E. Trotter Jr, When the greedy solution solves a class of Knapsack problems, Operations Research. 23(2) (1975), 207–217.
- [15] J. M. Marín, J. L. Ramírez Alfonsín, M. P. Revuelta, On the Frobenius number of Fibonacci numerical semigroups, Integers. 7 (2007), A14.
- [16] D. C. Ong, V. Ponomarenko, The Frobenius number of geometric sequences, Integers. 8 (2008), A33.
- [17] J. L. Ramírez Alfonsín, The Diophantine Frobenius Problem, Oxford Lecture Series in Mathematics and Its Applications, vol. 30, Oxford University Press. (2005).
- [18] J. B. Roberts, Note on linear forms, Proc. Amer. Math. Soc. 7 (1956), 465–469.
- [19] A. M. Robles-Pérez, J. C. Rosales, The Frobenius number for sequences of triangular and tetrahedral numbers, J. Number Theory. 186 (2018), 473–492.
- [20] O. J. Rödseth, On a linear diophantine problem of Frobenius II, J. Reine Angew Math. 307/308 (1979), 431–440.
- [21] J. C. Rosales, M. B. Branco, Numerical semigroups that can be expressed as an intersection of symmetric numerical semigroups, J. Pure Appl. Algebra. 171 (2002), 303–314.
- [22] J. C. Rosales, M. B. Branco, D. Torrão, The Frobenius problem for Thabit numerical semigroups, J. Number Theory. 155 (2015), 85–99.
- [23] J. C. Rosales, M. B. Branco, D. Torrão, The Frobenius problem for repunit numerical semigroups, Ramanujan J. 40 (2016), 323–334.
- [24] J. C. Rosales, M. B. Branco, D. Torrão, The Frobenius problem for Mersenne numerical semigroups, Math. Z. 286 (2017), 1–9.
- [25] J. C. Rosales, P. A. García-Sánchez, Numerical Semigroups, Developments in Mathematics. Vol.20, Springer, New York (2009).
- [26] E. S. Selmer, On the linear Diophantine problem of Frobenius, J. Reine Angew. Math. 293/294 (1977), 1–17.
- [27] Kyunghwan Song, The Frobenius problem for numerical semigroups generated by the Thabit numbers of the first, second kind base b and the Cunningham numbers, Bull. Korean Math. Soc. 57 (2020), 623–647.
- [28] Kyunghwan Song, The Frobenius problem for extended Thabit numerical semigroups, Integers. 21 (2021), A17.
- [29] J. J. Sylvester, On sub-invariants, i.e., semi-invariants to binary quanties of an unlimited order, Amer. J. Math. 5 (1882), 119–136.
- [30] A. Tripathi, On the Frobenius problem for $\{a, ha + d, ha + bd, ha + b^2d, ..., ha + b^kd\}$, J. Number Theory. 162 (2016), 212–223.

[31] A. Tripathi, Formulate for the Frobenius number in three variables, J. Number Theory. 170 (2017), 368–389.

 $^{1,2,3,4}{\rm School}$ of Mathematical Sciences, Capital Normal University, Beijing 100048, PR China

Email address: ¹liufeihu7476@163.com & ²guoce_xin@163.com & ³yesuting0203@163.com

& ⁴yinjingj@163.com