ON THE HILBERT DEPTH OF CERTAIN MONOMIAL IDEALS AND APPLICATIONS

SILVIU BĂLĂNESCU¹ AND MIRCEA CIMPOEAŞ²

ABSTRACT. We study the Stanley depth and the Hilbert depth for I and S/I, where $I \subset S = K[x_1, \ldots, x_n]$ is the intersection of monomial prime ideals with disjoint sets of variables. As an application, we obtain bounds for the Stanley depth of $I_{n,m}^t$ and $J_{n,m}^t$, where $I_{n,m}$ is the *m*-path ideal of the path graph of length n and $J_{n,m}$ is the the *m*-path ideal of the path graph of length n and $J_{n,m}$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the path graph of length n and $J_{n,m}$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the the *m*-path ideal of the stanley depth of $I_{n,m}^t$ is the stanley depth

INTRODUCTION

Let K be a field and $S = K[x_1, \ldots, x_n]$ the polynomial ring over K. Let M be a \mathbb{Z}^n graded S-module. A Stanley decomposition of M is a direct sum $\mathcal{D} : M = \bigoplus_{i=1}^r m_i K[Z_i]$ as a \mathbb{Z}^n -graded K-vector space, where $m_i \in M$ is homogeneous with respect to \mathbb{Z}^n -grading, $Z_i \subset \{x_1, \ldots, x_n\}$ such that $m_i K[Z_i] = \{um_i : u \in K[Z_i]\} \subset M$ is a free $K[Z_i]$ -submodule of M. We define sdepth(\mathcal{D}) = $\min_{i=1,\ldots,r} |Z_i|$ and sdepth(M) = $\max\{\text{sdepth}(\mathcal{D}) | \mathcal{D} \text{ is a}$ Stanley decomposition of $M\}$. The number sdepth(M) is called the Stanley depth of M.

Herzog, Vladoiu and Zheng show in [9] that sdepth(M) can be computed in a finite number of steps if M = I/J, where $J \subset I \subset S$ are monomial ideals. In [1], J. Apel restated a conjecture firstly given by Stanley in [13], namely that $sdepth(M) \ge depth(M)$ for any \mathbb{Z}^n -graded S-module M. This conjecture proves to be false, in general, for M = S/I and M = J/I, where $0 \ne I \subset J \subset S$ are monomial ideals, see [8]. For a friendly introduction in the thematic of Stanley depth, we refer the reader [10].

Let M be a finitely generated graded S-module. The Hilbert depth of M, denoted by hdepth(M), is the maximal depth of a finitely generated graded S-module N with the same Hilbert series as M. In [6] we introduced a new method to compute the Hilbert depth of a quotient J/I of two squarefree monomial ideals $I \subset J \subset S$; see Section 1.

In Section 2 we consider the edge ideal of a complete bipartite graph, that is

$$I := (x_1, \dots, x_n) \cap (x_{n+1}, \dots, x_{n+m}) \subset S := K[x_1, \dots, x_{n+m}],$$

and we study the Stanley depth and the Hilbert depth of I and S/I.

¹Silviu Bălănescu, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, E-mail: silviu.balanescu@stud.fsa.upb.ro

²Mircea Cimpoeaş, University Politehnica of Bucharest, Faculty of Applied Sciences, Bucharest, 060042, Romania and Simion Stoilow Institute of Mathematics, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania, E-mail: mircea.cimpoeas@upb.ro, mircea.cimpoeas@imar.ro

²⁰²⁰ Mathematics Subject Classification. 05A18, 06A07, 13C15, 13P10, 13F20.

Key words and phrases. Stanley depth, Hilbert depth, depth, monomial ideal, path ideal.

Assume $m \leq n$. In Proposition 2.2 we show that

$$m \ge \operatorname{sdepth}(S/I) \ge \min\{m, \left\lceil \frac{n}{2} \right\rceil\}$$

Also, in Theorem 2.6 we prove that

hdepth
$$(S/I) \leq \left\lfloor n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}} \right\rfloor.$$

In particular, we note that hdepth(S/I) < m if and only if $n \leq 2m - 2$. In Theorem 2.9 we prove that

$$\mathrm{hdepth}(I) = \mathrm{sdepth}(I) = \left\lceil \frac{m}{2} \right\rceil + \left\lceil \frac{n}{2} \right\rceil$$

if n and m are not both even. Also, we prove that if n = 2s and m = 2t then

$$t + s \leq \text{sdepth}(I) \leq \text{hdepth}(S/I) = t + s + 1.$$

In particular, we have $hdepth(I) = \lfloor \frac{n+m+2}{2} \rfloor$ for any $n \ge m \ge 1$. In Section 3 we consider a generalization of the ideal from the previous section, namely

$$I := I_{n_1,\dots,n_r} := (x_1,\dots,x_{n_1}) \cap (x_{n_1+1},\dots,x_{n_1+n_2}) \cap \dots \cap (x_{n_1+\dots+n_{r-1}+1},\dots,x_N) \subset S,$$

where $N = n_1 + \cdots + n_r$ and $S = K[x_1, \ldots, x_N]$. In Theorem 3.3 we prove that

$$\left\lfloor \frac{N+r}{2} \right\rfloor \ge \text{hdepth}(I) \ge \text{sdepth}(I) \ge \left\lceil \frac{n_1}{2} \right\rceil + \dots + \left\lceil \frac{n_r}{2} \right\rceil$$

Also, we conjecture that

$$\mathrm{hdepth}(I) = \left\lfloor \frac{N+r}{2} \right\rfloor.$$

This formula holds for r = 2 and if $r \ge 3$ and at most one of the numbers n_1, \ldots, n_r is even. In Proposition 3.8 we characterize hdepth(S/I) and hdepth(I) in combinatorial terms. In Proposition 3.10 we show that

hdepth
$$(S/I) \le \min\{d \ge r : \binom{N-d+r-1}{r} < n_1 n_2 \cdots n_r\} - 1.$$

Proposition 3.13 yields us to conjecture that

hdepth
$$(S/I) \approx N - \left\lceil \sqrt[r]{r!n_1n_2\cdots n_r} \right\rceil$$
.

Let $n > m \ge 2$ and $t \ge 1$ be some integers. In Section 4 we apply the results from Section 3 in order to obtain sharper bounds for the Stanley depth of $I_{n,m}^t$ and $J_{n,m}^t$, where

$$I_{n,m} = (x_1 x_2 \cdots x_m, x_2 x_3 \cdots x_{m+1}, \dots, x_{n-m+1} \cdots x_n) \subset S := K[x_1, \dots, x_n],$$

is the m-path ideal associated to path graph of length n and

$$J_{n,m} = I_{n,m} + (x_{n-m+2} \cdots x_n x_1, \dots, x_n x_1 \cdots x_{m-1}) \subset S$$

is the *m*-path ideal associated to the cycle graph of length n.

In Theorem 4.3 we show that

sdepth
$$(I_{n,m}^t) \le \min\{n - \left\lceil \frac{t_0}{2} \right\rceil, n - \left\lfloor \frac{n - t_0 + 1}{m + 1} \right\rfloor + 1\},\$$

where $t_0 = \min\{t, n - m\}$. In Theorem 4.5 we show that

$$\operatorname{sdepth}(J_{n,m}^t) \le \left\lfloor \frac{n+d}{2} \right\rfloor - 1,$$

for any $t \ge n-1$, where $d = \gcd(n, m)$.

1. Preliminaries

First, we fix some notations and we recall the main result of [6].

We denote $[n] := \{1, 2, ..., n\}$ and $S := K[x_1, ..., x_n]$.

For a subset $C \subset [n]$, we denote $x_C := \prod_{j \in C} x_j \in S$.

For two subsets $C \subset D \subset [n]$, we denote $[C, D] := \{A \subset [n] : C \subset A \subset D\}$, and we call it the *interval* bounded by C and D.

Let $I \subset J \subset S$ be two square free monomial ideals. We let:

$$\mathbf{P}_{J/I} := \{ C \subset [n] : x_C \in J \setminus I \} \subset 2^{[n]}.$$

A partition of $P_{J/I}$ is a decomposition:

$$\mathcal{P}: \mathbf{P}_{J/I} = \bigcup_{i=1}^{r} [C_i, D_i],$$

into disjoint intervals.

If \mathcal{P} is a partition of $P_{J/I}$, we let $\operatorname{sdepth}(\mathcal{P}) := \min_{i=1}^r |D_i|$. The Stanley depth of $P_{J/I}$ is

$$sdepth(P_{J/I}) := \max\{sdepth(\mathcal{P}) : \mathcal{P} \text{ is a partition of } P_{J/I}\}.$$

Herzog, Vlădoiu and Zheng proved in [9] that:

$$\operatorname{sdepth}(J/I) = \operatorname{sdepth}(\mathcal{P}_{J/I}).$$

Let $P := P_{J/I}$, where $I \subset J \subset S$ are square-free monomial ideals. For any $0 \le k \le n$, we denote:

$$\mathbf{P}_k := \{A \in \mathbf{P} \ : \ |A| = k\} \text{ and } \alpha_k(J/I) = \alpha_k(\mathbf{P}) = |\mathbf{P}_k|$$

For all $0 \le d \le n$ and $0 \le k \le d$, we consider the integers

$$\beta_k^d(J/I) := \sum_{j=0}^k (-1)^{k-j} \binom{d-j}{k-j} \alpha_j(J/I).$$
(1.1)

From (1.1) we can easily deduce that

$$\alpha_k(J/I) = \sum_{j=0}^k \binom{d-j}{k-j} \beta_k^d(J/I), \text{ for all } 0 \le k \le d.$$

$$(1.2)$$

Also, we have that

$$\beta_k^d(J/I) = \alpha_k(J/I) - \binom{d}{k} \beta_0^d(J/I) - \binom{d-1}{k-1} \beta_1^d(J/I) - \dots - \binom{d-k+1}{1} \beta_{k-1}^d(J/I).$$
(1.3)

Theorem 1.1. ([6, Theorem 2.4]) With the above notations, the Hilbert depth of J/I is

$$hdepth(J/I) := \max\{d : \beta_k^d(J/I) \ge 0 \text{ for all } 0 \le k \le d\}.$$

As a basic property of the Hilbert depth, we state the following:

Proposition 1.2. Let $I \subset J \subset S$ be two square-free monomial ideals. Then sdepth $(J/I) \leq hdepth(J/I)$.

2. Edge ideal of a complete bipartite graph

Let n and m be two positive integers. We let $S = K[x_1, x_2, ..., x_{n+m}]$ and we consider the square free monomial ideal:

 $I := (x_1, \ldots, x_n) \cap (x_{n+1}, \ldots, x_{n+m}) \subset S.$

Our aim is to study the Stanley depth and the Hilbert depth of I and S/I.

As usual, given a positive integer k, we denote $[k] := \{1, 2, \dots, k\}$.

Remark 2.1. Let $K_{n,m} = (V, E)$ be the complete bipartite graph, that is $V = V' \cup V''$, where $V' = \{1, \ldots, n\}, V'' = \{n + 1, \ldots, n + m\}$ and $E = \{\{i, j\} : i \in [n], j - n \in [m]\}$. Note that $I = (x_i x_{n+j} : i \in [n], j \in [m])$ is the edge ideal of $K_{n,m}$.

Also, we mention that depth(S/I) = 1, which can be easily checked.

Proposition 2.2. Let $n \ge m \ge 1$ be two integers. Then:

- (1) $m \ge \operatorname{sdepth}(S/I) \ge \min\{m, \lceil \frac{n}{2} \rceil\}.$
- (2) $m + \lceil \frac{n}{2} \rceil \ge \text{sdepth}(I) \ge \lceil \frac{n}{2} \rceil + \lceil \frac{m}{2} \rceil.$
- (3) If $n \ge 2m 1$ then sdepth(S/I) = m.

Proof. (1) Since $I = I'S \cap I''S$, where $I' = (x_1, \ldots, x_n) \subset S' = K[x_1, \ldots, x_n]$ and $I'' = (x_{n+1}, \ldots, x_{n+m}) \subset S'' = K[x_{n+1}, \ldots, x_{n+m}]$, from [3, Theorem 1.3(2)] it follows that

$$\operatorname{sdepth}(S/I'S) \ge \operatorname{sdepth}(S/I) \ge \min\{\operatorname{sdepth}(S/I'S), \operatorname{sdepth}_{S''}(S''/I'') + \operatorname{sdepth}_{S'}(I')\}.$$

As $S/I'S \cong S''$, we have that sdepth(S/I'S) = m.

Also, $S''/I'' \cong K$, so sdepth_{S''}(S''/I'') = 0.

Finally, sdepth_{S'}(I') = $\lceil \frac{n}{2} \rceil$, see [2, Theorem 2.2].

(2) Since $(I : x_{n+1}) = I'S$, from [12, Proposition 1.3] (see arXiv version), [2, Theorem 2.2] and [9, Lemma 3.6] we have

$$\operatorname{sdepth}(I) \leq \operatorname{sdepth}(I:x_{n+1}) = \operatorname{sdepth}(I'S) = m + \operatorname{sdepth}_{S'}(I') = m + \left\lceil \frac{n}{2} \right\rceil.$$

The other inequality follows from [11, Lemma 1.1] and [2, Theorem 2.2].

(3) If $n \ge 2m - 1$ then $\left\lceil \frac{n}{2} \right\rceil \ge m$, hence the result follows from (1).

Lemma 2.3. Let $n \ge m \ge 1$ be two integers and N := n + m. We have that

(1)
$$\alpha_k(I) = \begin{cases} 0, & 0 \le k \le 1\\ \sum_{j=1}^{k-1} {n \choose j} {m \choose k-j}, & 2 \le k \le N \end{cases}$$

(2) $\alpha_k(I) = {N \choose k} - {n \choose k} - {m \choose k} + \delta_{k0}, \text{ for all } 0 \le k \le N.$
(3) $\alpha_k(S/I) = {n \choose k} + {m \choose k} - \delta_{k0}, \text{ for all } 0 \le k \le N.$

Proof. (1) Since I is generated in degree 2, we have $\alpha_0(I) = \alpha_1(I) = 1$. Any squarefree monomial $u \in I$ with $\deg(u) = k \geq 2$ can be written as $u = v \cdot w$, where $v \in S' = K[x_1, \ldots, x_n]$ and $w \in S'' = K[x_{n+1}, \ldots, x_N]$ are squarefree monomials. Assume $\deg(v) = j$ with $1 \leq j \leq k - 1$. Then $\deg(w) = k - j$. Since there are $\binom{n}{j}$ squarefree monomials of degree j in S' and $\binom{m}{k-j}$ squarefree monomials of degree k - j in S'', we easily get the required conclusion.

(2) For $k \leq 1$ the identity can be easily checked. Assume $k \geq 2$. From (1) and the well known combinatorial formula

$$\sum_{j=0}^{k} \binom{n}{j} \binom{m}{k-j} = \binom{n+m}{k} = \binom{N}{k},$$

we get the required conclusion.

(3) It follows immediately from (2).

Lemma 2.4. For any integers $0 \le k \le d$ and $n \ge 0$ we have that

$$\sum_{j=0}^{k} (-1)^{k-j} \binom{d-j}{k-j} \binom{n}{j} = (-1)^k \binom{d-n}{k} = \binom{n-d+k-1}{k}.$$

Proof. Using the identity $\binom{x}{k} = \binom{-x+k-1}{k}$ and the Chu–Vandermonde summation, we get

$$\sum_{j=0}^{k} (-1)^{k-j} \binom{d-j}{k-j} \binom{n}{j} = \sum_{j=0}^{k} \binom{-d+k-1}{k-j} \binom{n}{j} = \binom{n-d+k-1}{k},$$

as required.

Lemma 2.5. Let $n \ge m \ge 1$ and $0 \le k \le d \le N := n + m$ some integers. We have that

(1)
$$\beta_k^d(S/I) = \binom{n-d+k-1}{k} + \binom{m-d+k-1}{k} + (-1)^{k+1} \binom{d}{k},$$

(2) $\beta_k^d(I) = \binom{N-d+k-1}{k} - \binom{n-d+k-1}{k} - \binom{m-d+k-1}{k} + (-1)^k \binom{d}{k}.$

Proof. (1) From (1.1), Lemma 2.3(3) and Lemma 2.4 we have that

$$\beta_k^d(S/I) = \sum_{j=1}^k (-1)^{k-j} \binom{d-j}{k-j} \binom{n}{j} + \sum_{j=1}^k (-1)^{k-j} \binom{d-j}{k-j} \binom{m}{j} - (-1)^k \binom{d}{k} = \binom{n-d+k-1}{k} + \binom{m-d+k-1}{k} + (-1)^{k+1} \binom{d}{k},$$

as required. (2) The proof is similar, using (1.1), Lemma 2.3(2) and Lemma 2.4.

Note that, if $n \ge 2m - 1$ then, according to Proposition 2.2(3) and Proposition 1.2 we have $hdepth(S/I) \ge sdepth(S/I) = m$. Also, $sdepth(S/I) \le m$, for any $n \ge m$.

Hence, it is interesting to consider the case $n \leq 2m - 2$, in order to find a better upper bound for sdepth(S/I).

Theorem 2.6. Let $n \ge m \ge 1$ be two integers. Then

$$\operatorname{sdepth}(S/I) \le \operatorname{hdepth}(S/I) \le \left[n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}}\right].$$

In particular, if $n \leq 2m - 2$ then hdepth(S/I) < m.

Proof. The first inequality follows from Proposition 1.2. We consider the quadratic function

$$\varphi(t) = \frac{1}{2}t(t-1) - (n+m)t + \frac{1}{2}n(n+1) + \frac{1}{2}m(m+1).$$

Note that, according to Lemma 2.5(1), $\beta_2^d(S/I) = \varphi(d)$.

The roots of $\varphi(t) = 0$ are $t_{1,2} = n + m + \frac{1}{2} \pm \sqrt{2mn + \frac{1}{4}}$ and therefore

$$\varphi(t) < 0$$
 if and only if $t \in \left(n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}}, n + m + \frac{1}{2} + \sqrt{2mn + \frac{1}{4}}\right)$.

From the fact that $\beta_2^d(S/I) = \varphi(d)$ and the above, it follows that

$$\beta_2^d(S/I) < 0 \text{ for } d \ge \left\lfloor n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}} \right\rfloor + 1$$

From Theorem 1.1, we get hdepth $(S/I) \leq \lfloor n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}} \rfloor$. In order to prove the last part, we consider the function

$$\psi(x) = x + m + \frac{1}{2} - \sqrt{2mx + \frac{1}{4}}, \ x \in [m, \infty)$$

Since $\frac{d\psi}{dx}(x) > 0$, $m \le n \le 2m - 2$ and $\psi(2m - 1) = m$, it follows that

$$\left\lfloor 2m + \frac{1}{2} - \sqrt{2m^2 + \frac{1}{4}} \right\rfloor \le \lfloor \psi(n) \rfloor = \left\lfloor n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}} \right\rfloor < \psi(2m - 1) = m,$$
 required.

as re

Theorem 2.7. Let $n \ge m \ge 2$ be two integers. Then

 $\operatorname{hdepth}(S/I) = \max\{d \le n + m : \binom{d-n}{2\ell} + \binom{d-m}{2\ell} \ge \binom{d}{2\ell} \text{ for all } 1 \le \ell \le \lfloor \frac{d}{2} \rfloor\}.$ Moreover, $\operatorname{hdepth}(S/I) < m \text{ if } n \le 2m - 2. \text{ Also, } m \le \operatorname{hdepth}(S/I) \le n - m + 1 \text{ if } n \ge 2m - 1.$

Proof. Let $q := \max\{d \le n + m : \binom{d-n}{2\ell} + \binom{d-m}{2\ell} \ge \binom{d}{2\ell}$ for all $1 \le \ell \le \lfloor \frac{d}{2} \rfloor\}$. From Lemma 2.5(1) and the identity $\binom{x}{k} = \binom{-x+k-1}{k}$ it follows that

$$\beta_{2\ell}^d(S/I) = \binom{d-n}{2\ell} + \binom{d-m}{2\ell} - \binom{d}{2\ell}.$$

Hence hdepth $(S/I) \leq q$.

On the other hand, from the proof of Theorem 2.6 and (2.1), it follows that

$$q \leq \left[n + m + \frac{1}{2} - \sqrt{2mn + \frac{1}{4}} \right].$$
 (2.2)

We consider two cases:

(i) $n \leq 2m - 2$. From Theorem 2.6, it follows that q < m. From Lemma 2.5(2) and $0 \leq k \leq q$ with k odd, we have

$$\beta_k^q(S/I) = \binom{n-q+k-1}{k} + \binom{m-q+k-1}{k} + \binom{q}{k} \ge \binom{n-m+k}{k} + 1 + \binom{q}{k} > 0.$$

Since by the definition of a we have $\beta_k^q(S/I) \ge 0$ for all $0 \le k \le q$ with k even we

Since, by the definition of q, we have $\beta_k^q(S/I) \ge 0$ for all $0 \le k \le q$ with k even, we conclude that $\mathrm{hdepth}(S/I) \ge q$. Hence $\mathrm{hdepth}(S/I) = q < m$, as required.

(ii) $n \ge 2m - 1$. First, note that

 $m = \operatorname{sdepth}(S/I) \le \operatorname{hdepth}(S/I) \le q.$

From (2.2) and the above it follows that

$$m \le q \le \left\lfloor n + m + \frac{1}{2} - \sqrt{2m(2m-1) + \frac{1}{4}} \right\rfloor = \left\lfloor n + m + \frac{1}{2} - (2m - \frac{1}{2}) \right\rfloor = n - m + 1.$$

From Lemma 2.5(2) and $0 \le k \le q$ with k odd, we have

$$\beta_k^q(S/I) = \binom{n-q+k-1}{k} + \binom{q-m}{k} + \binom{q}{k} \ge \binom{m}{k} + 0 + \binom{m}{k} \ge 0.$$

Using the same argument as in the case (i), it follows that hdepth(S/I) = q, as required.

Thus, the proof is complete.

Lemma 2.8. Let $n \ge m \ge 1$ be two integers. Then

$$\operatorname{hdepth}(I) \le \left\lfloor \frac{n+m+2}{2} \right\rfloor.$$

(2.1)

Proof. If n + m = 2, that is n = m = 1, then there is nothing to prove. Assume $n + m \ge 3$. From Lemma 2.5(2) and straightforward computations, it follows that

$$\beta_3^d(I) = \frac{nm(n+m-2d+2)}{2} < 0,$$

if and only if $d > \frac{n+m+2}{2}$. Hence, we get the required result.

Theorem 2.9. Let $n \ge m \ge 1$ be two integers.

(1) If n and m are not both even then we have that:

$$\operatorname{sdepth}(I) = \operatorname{hdepth}(I) = \left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{m}{2} \right\rceil.$$

(2) If n = 2t and m = 2s then we have that:

$$t + s \leq \operatorname{sdepth}(I) \leq \operatorname{hdepth}(I) \leq t + s + 1.$$

In both cases, we have $hdepth(I) = \lfloor \frac{n+m+2}{2} \rfloor$.

Proof. (1) According to Proposition 2.2(2), we have that

$$\operatorname{sdepth}(I) \ge \left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{m}{2} \right\rceil.$$
 (2.3)

On the other hand, according to Proposition 1.2 and Lemma 2.8, we have that

$$\operatorname{sdepth}(I) \le \operatorname{hdepth}(I) \le \left\lfloor \frac{n+m}{2} \right\rfloor + 1.$$
 (2.4)

Note that, if n and m are not both even, then

$$\left\lceil \frac{n}{2} \right\rceil + \left\lceil \frac{m}{2} \right\rceil = \left\lfloor \frac{n+m}{2} \right\rfloor + 1.$$
(2.5)

Hence, (1) follows from (2.3), (2.4) and (2.5).

(2) From (2.3) and (2.4) we have that

$$t + s \le \operatorname{sdepth}(I) \le \operatorname{hdepth}(I) \le t + s + 1.$$

On the other hand, for $0 \le k \le t + s + 1$, from Lemma 2.5(2) we have

$$\beta_k^{t+s+1}(I) = \binom{t+s-2+k}{k} - \binom{t-s-2+k}{k} - \binom{s-t-2+k}{k} + (-1)^k \binom{t+s+1}{k}.$$
(2.6)

By direct computations, from (2.6) it follows that

$$\beta_0^{t+s+1}(I) = 0, \ \beta_1^{t+s+1}(I) = 0, \ \beta_2^{t+s+1}(I) = 4st \text{ and } \beta_3^{t+s+1}(I) = 0.$$

Also, by straightforward computations, we get

$$\beta_4^{t+s+1}(I) = \beta_5^{t+s+1}(I) = \frac{ts(2s^2 + 2t^2 - 1)}{3} > 0$$

Now, assume $6 \le k \le t + s + 1$. Without any loss of generality, we assume that t = s + a, where a is a nonnegative integer. In order to prove that $\beta := \beta_k^{t+s+1}(I) \ge 0$, we consider the following cases:

(i) k is even.
(i.1)
$$a = 0$$
. From (2.6) and the fact that $s \ge 1$ it follows that

$$\beta = \binom{2s-2+k}{k} - \binom{k-2}{k} - \binom{k-2}{k} + \binom{2s+1}{k} \ge \binom{k}{k} + \binom{2s+1}{k} > 0.$$
(i.2) $a = 1$. From (2.6) and the fact that $s \ge 1$ it follows that

$$\beta = \binom{2s-1+k}{k} - \binom{k-1}{k} - \binom{k-3}{k} + \binom{2s+2}{k} \binom{k+1}{k} + \binom{2s+2}{k} > 0.$$
(i.3) $a \ge 2$ and $k \ge a+2$. From (2.6) we get

$$\beta = \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} + \binom{2s+a+1}{k} > \binom{2s+a+1}{k} \ge 0.$$
(i.4) $a \ge 5$ and $k \le a+1$. From (2.6), using $\binom{-k}{k} = (-1)^k \binom{x+k-1}{k}$, we get

$$\beta = \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} - \binom{a+1}{k} + \binom{2s+a+1}{k} = \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} + \binom{2s+a+1}{k} - \binom{a+1}{k} = \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} + \binom{2s+a+1}{k} - \binom{a+1}{k} > 0.$$
(ii) k is odd.
(iii) k is odd.
(iii.1) $a = 0$. From (2.6) and the fact that $k \ge 6$ it follows that

$$\beta = \binom{2s-2+k}{k} - \binom{k-2}{k} - \binom{k-2}{k} - \binom{2s+1}{k} \ge \binom{2s+4}{k} - \binom{2s+1}{k} > 0.$$
(ii.2) $a = 0$. From (2.6) and the fact that $k \ge 6$ it follows that

$$(2s-1+k) - \binom{k-1}{k} - \binom{k-2}{k} - \binom{2s+2}{k} - \binom{2s+4}{k} - \binom{2s+2}{k} - \binom{2s+2}{k} - \binom{2s+2}{k} = \binom{2s+4}{k} - \binom{2s+2}{k} - \binom{2s+2}{k}$$

$$\begin{split} \beta &= \binom{2s-1+k}{k} - \binom{k-1}{k} - \binom{k-3}{k} - \binom{2s+2}{k} \ge \binom{2s+5}{k} - \binom{2s+2}{k} > 0. \\ \text{(ii.3)} \ a \ge 2 \ \text{and} \ k \ge a+2. \ \text{From (2.6)} \ \text{and the fact that} \ k \ge 6 \ \text{and} \ s \ge 1 \ \text{we get} \\ \beta &= \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} - \binom{2s+a+1}{k} > \binom{2s+a+k-3}{k-1} - \\ -\binom{a+k-2}{k} \ge \binom{a+k-1}{k-1} - \binom{a+k-2}{k} = \frac{ak+k^2-k-a^2+a}{a(a-1)} \binom{a+k-2}{k} > 0. \\ \text{(ii.4)} \ a \ge 5 \ \text{and} \ k \le a+1. \ \text{From (2.6)}, \ \text{using} \ \binom{-x}{k} = (-1)^k \binom{x+k-1}{k}, \ \text{we get} \\ \beta &= \binom{2s+a+k-2}{k} - \binom{a+k-2}{k} - \binom{a+k-2}{k} + \binom{a+1}{k} - \binom{2s+a+1}{k} = \\ &= \binom{2s+a+k-2}{k} - \binom{2s+a+1}{k} - \binom{2s+a+1}{k} - \binom{2s+a+1}{k} = \\ &= \binom{2s+a+k-2}{k} - \binom{2s+a+1}{k} - \binom{2s+a+1}{k} - \binom{a+k-2}{k} - \binom{a+1}{k} = \\ &= \binom{2s+a+k-2}{k} - \binom{2s+a+1}{k} - \binom{2s+a+1}{k} - \binom{a+1}{k} = \\ &= \binom{2s+a+k-3}{k} \binom{\ell}{k-1} - \sum_{\ell=a+1}^{a+k-3} \binom{\ell}{k-1} > 0. \end{split}$$

Hence, hdepth(I) = s + t + 1 and the proof is complete.

3. A GENERALIZATION

Let n_1, n_2, \ldots, n_r be some positive integers, $N = n_1 + \cdots + n_r$ and $S := K[x_1, \ldots, x_N]$. We consider the ideal

 $I := I_{n_1, n_2, \dots, n_r} := (x_1, \dots, x_{n_1}) \cap (x_{n_1+1}, \dots, x_{n_1+n_2}) \cap \dots \cap (x_{n_1+\dots+n_{r-1}+1}, \dots, x_N) \subset S,$ which generalize the ideal I from the previous section.

Lemma 3.1. With the above notations, we have that

$$\alpha_k(I) = \begin{cases} 0, & k \le r-1 \\ \sum_{\substack{\ell_1, \ell_2, \dots, \ell_r \ge 1 \\ \ell_1 + \dots + \ell_r = k}} {\binom{n_1}{\ell_1} \binom{n_2}{\ell_2} \cdots \binom{n_r}{\ell_r}, & k \ge r \end{cases}.$$

Proof. Since I is generated by square free monomials of degree k, it is clear that $\alpha_k(I) = 0$ for $k \leq r-1$. Assume $k \geq r$ and let $u \in I$ a square free monomial of degree k. It follows that $u = u_1 \cdots u_r$, where

$$1 \neq u_j \in I_j := (x_{n_1 + \dots + n_{j-1} + 1}, \dots, x_{n_1 + \dots + n_j})$$
 for all $1 \leq j \leq r$.

Let $\ell_j = \deg(u_j) \ge 1$. Since there are $\binom{n_j}{\ell_j}$ squarefree monomials of degree ℓ_j in I_j , we get the required conclusion.

Lemma 3.2. With the above notations, we have that

(1) $\alpha_k(I) = 0$ for $k \le r - 1$. (1) $\alpha_r(I) = n_1 n_2 \cdots n_r.$ (2) $\alpha_r(I) = n_1 n_2 \cdots n_r.$ (3) $\alpha_{r+1}(I) = n_1 n_2 \cdots n_r \cdot \frac{n_1 + \cdots + n_r - r}{2}.$ (4) $\alpha_k(I) = \binom{N}{k} \text{ for } k \ge N - \min_{i=1} n_i + 1.$

Proof. (1), (2) and (3) follow immediately from Lemma 3.1. In order to prove (4), it is enough to observe that any squarefree monomial $u \in S$ of degree $k \geq N - \min_{i=1}^{n} n_i + 1$, \square belongs to I.

Theorem 3.3. With the above notations, we have that:

$$\left\lfloor \frac{N+r}{2} \right\rfloor \ge \operatorname{hdepth}(I) \ge \operatorname{sdepth}(I) \ge \left\lceil \frac{n_1}{2} \right\rceil + \dots + \left\lceil \frac{n_r}{2} \right\rceil.$$

Proof. In order to prove the first inequality, let $d > \frac{n_1 + \dots + n_r + r}{2}$ be an integer. From Theorem 1.1 and Lemma 3.2(1,2,3) it follows that

$$\beta_k^d(I) = 0 \text{ for } 0 \le k \le r - 1, \ \beta_r^d(I) = \alpha_r(I) = n_1 n_2 \cdots n_r \text{ and}$$

$$\beta_{r+1}^d(I) = \alpha_{r+1}(I) - (d-r)\alpha_r(I) = n_1 n_2 \cdots n_r \left(\frac{n_1 + \dots + n_r + r}{2} - d\right) < 0.$$

whether the height of the second second

Hence

The inequality $hdepth(I) \ge sdepth(I)$ follows from Proposition 1.2, and the last inequality follows from [3, Corollary 1.9(1)] and [2, Theorem 2.2].

Based on our computer experiments, we propose the following conjecture:

Conjecture 3.4. With the above notations, we have

$$\mathrm{hdepth}(I) = \left\lfloor \frac{N+r}{2} \right\rfloor.$$

Note that, according to Theorem 2.9, Conjecture 3.4 holds for r = 2. Also, according to Theorem 3.3, Conjecture 3.4 is true when at most one of the numbers n_1, \ldots, n_r is even.

Proposition 3.5. With the above notations, we have that

$$N - \min_{i=1}^{r} n_i \ge \operatorname{hdepth}(S/I) \ge \operatorname{sdepth}(S/I) \ge \left\lceil \frac{n_1}{2} \right\rceil + \dots + \left\lceil \frac{n_r}{2} \right\rceil - \min_{i=1}^{r} \left\lceil \frac{n_i}{2} \right\rceil$$

Proof. From Lemma 3.2(4) it follows that

$$\alpha_k(S/I) = 0 \text{ for all } k \ge N - \max_{i=1}^r n_i + 1.$$

Hence, from [6, Lemma 1.3], we get the first inequality.

The second inequality follows from [3, Corollary 1.9] and [2, Theorem 2.2].

Note that, Proposition 3.5 reproves the fact that $hdepth(S/I) \ge depth(S/I) = r - 1$.

Lemma 3.6. For any $0 \le k \le N$, we have that:

(1) $\alpha_k(I) = \sum_{J \subseteq [n]} (-1)^{|J|} {\binom{N - \sum_{i \in J} n_i}{k}}.$ (2) $\alpha_k(I) = \sum_{\emptyset \neq J \subseteq [n]} (-1)^{|J|+1} {\binom{N - \sum_{i \in J} n_i}{k}}.$

Proof. (1) For all $1 \le i \le r$ we let:

$$A_i = \{(\ell_1, \dots, \ell_d) : \ell_1 + \dots + \ell_r = k, \ell_i = 0 \text{ and } \ell_j \ge 0 \text{ for } j \ne i\}.$$

Also, we consider the set:

$$A = \{ (\ell_1, \dots, \ell_d) : \ell_1 + \dots + \ell_r = k \text{ and } \ell_i \ge 0 \text{ for all } 1 \le i \le r \}$$

For any nonempty subset $J \subset [n]$, we let $A_J = \bigcup_{i \in J} A_i$. Also, we denote $A_{\emptyset} = A$. From Lemma 3.1 it follows that

$$\alpha_k(I) = \sum_{(\ell_1,\dots,\ell_r)\in A_{\emptyset}\setminus(\bigcup_{i=1}^r A_i)} \binom{n_1}{\ell_1} \cdots \binom{n_r}{\ell_r}.$$
(3.1)

Note that this equality holds also for k < r as both terms are zero in this case. It is well known that

$$\sum_{(\ell_1,\dots,\ell_r)\in A} \binom{n_1}{\ell_1}\cdots\binom{n_r}{\ell_r} = \binom{n_1+\cdots+n_r}{k} = \binom{N}{k}.$$

11

Similarly, if $J \subset [n]$ then

$$\sum_{(\ell_1,\dots,\ell_r)\in\bigcap_{i\in J}A_i} \binom{n_1}{\ell_1}\cdots\binom{n_r}{\ell_r} = \binom{N-\sum_{i\in J}n_i}{k}.$$
(3.2)

From (3.1) and (3.2), using the inclusion exclusion principle, we get the required conclusion. (2) Follows from (1) and the fact that $\alpha(S/I) = {N \choose k} - \alpha(I)$.

Note that Lemma 3.6 generalizes Lemma 2.3(2,3). From Lemma 3.6 and Lemma 2.4 we get the following generalization of Lemma 2.5:

Lemma 3.7. For any $0 \le k \le d \le N$, we have that:

(1)
$$\beta_k^d(I) = \sum_{J \subseteq [n]} (-1)^{|J|} {\binom{N - \sum_{i \in J} n_i - d + k - 1}{k}}.$$

(2) $\beta_k^d(S/I) = \sum_{\emptyset \neq J \subseteq [n]} (-1)^{|J| + 1} {\binom{N - \sum_{i \in J} n_i - d + k - 1}{k}}.$

Proposition 3.8. With the above notations, we have that:

$$(1) \text{ hdepth}(I) = \max\{d : \left\lceil \frac{n_1}{2} \right\rceil + \dots + \left\lceil \frac{n_r}{2} \right\rceil \le d \le \left\lfloor \frac{N+r}{2} \right\rfloor \text{ and} \\ \sum_{J \subseteq [n]} (-1)^{|J|} \binom{N - \sum_{i \in J} n_i - d + k - 1}{k} \ge 0 \text{ for all } r \le k \le d\}.$$

$$(2) \text{ hdepth}(S/I) = \max\{d : \left\lceil \frac{n_1}{2} \right\rceil + \dots + \left\lceil \frac{n_r}{2} \right\rceil - \min_{i=1}^r \left\lceil \frac{n_i}{2} \right\rceil \le d \le N - \min_{i=1}^r n_i \text{ and} \\ \sum_{\emptyset \ne J \subseteq [n]} (-1)^{|J|+1} \binom{N - \sum_{i \in J} n_i - d + k - 1}{k} \ge 0 \text{ for all } r \le k \le d\}.$$

Proof. (1) It follows from Theorem 3.3 and Lemma 3.7(1).

(1) It follows from Proposition 3.5 and Lemma 3.7(2).

Lemma 3.9. Let $d \ge r$. We have that

$$\beta_r^d(S/I) = \binom{N-d+r-1}{r} - n_1 n_2 \cdots n_r.$$

Proof. From Lemma 3.2 it follows that

$$\alpha_k(S/I) = \binom{N}{k}$$
 for $k \le r-1$, $\alpha_r(S/I) = \binom{N}{r} - n_1 n_2 \cdots n_r$.

Hence, the required result follows from (1.1) and Lemma 2.4.

Proposition 3.10. With the above notations, we have that:

hdepth
$$(S/I) \le \min\{d \ge r : \binom{N-d+r-1}{r} < n_1 n_2 \cdots n_r\} - 1.$$

Proof. First of all, note that, according to (1.3), we have

$$\beta_0^N(S/I) = 1$$
 and $\beta_k^N(S/I) = 0$ for all $1 \le k \le r - 1$

Moreover, according to Lemma 3.9, (1.1) and (1.3), we have

$$\beta_r^N(S/I) = \binom{N - N + r - 1}{r} - n_1 n_2 \cdots n_r = -n_1 n_2 \cdots n_r < 0.$$

Therefore, we have that

$$q := \min\{d \ge r : \binom{N-d+r-1}{r} < n_1 n_2 \cdots n_r\}$$

is well defined and $q \leq N$. Now, it is enough to notice that, from above, $\beta_r^q(S/I) < 0$ and thus hdepth $(S/I) \leq q - 1$, as required.

Lemma 3.11. We have that
$$\binom{N-d+r-1}{r} \ge n_1 n_2 \cdots n_r \text{ for all } d \le N - \left\lceil \sqrt[r]{r!n_1 n_2 \cdots n_r} \right\rceil.$$

Proof. We assume that $d = \lfloor aN \rfloor$, where $a \in (0, 1)$. Then

$$\binom{N-d+r-1}{r} = \frac{(N-d+r-1)(N-d+r-2)\cdots(N-d)}{r!} \ge \frac{(N-aN+r-1)(N-aN+r-2)\cdots(N-aN)}{r!} \ge \frac{N^r(1-a)^r}{r!}.$$
 (3.3)

On the other hand

$$\frac{N^r(1-a)^r}{r!} \ge n_1 n_2 \cdots n_r \Leftrightarrow (1-a)^r \ge \frac{r! n_1 n_2 \cdots n_r}{r! N^k} \Leftrightarrow a \le 1 - \frac{\sqrt[r]{r! n_1 n_2 \cdots n_r}}{N}$$
(3.4)

The conclusion follows from (3.3) and (3.4).

Remark 3.12. Note that, from the inequality of means, we have

$$\sqrt[r]{r!n_1n_2\cdots n_r} \le \frac{N\sqrt[r]{r!}}{r},$$

with equality for $n_1 = n_2 = \cdots = n_r = n = \frac{N}{r}$. Therefore, from Lemma 3.11, we have that

$$\binom{N-d+r-1}{r} \ge n^r \text{ for all } d \le N \left(1 - \sqrt[r]{\frac{r!}{r^r}}\right).$$

Proposition 3.13. With the above notations, we have that

$$\beta_r^d(S/I) \ge 0 \text{ for all } d \le N - \left\lceil \sqrt[r]{r!n_1n_2\cdots n_r} \right\rceil$$

Proof. It follows from Lemma 3.9 and Lemma 3.11.

Proposition 3.13 allows us to conjecture that

hdepth
$$(S/I) \approx N - \left\lceil \sqrt[r]{r!n_1n_2\cdots n_r} \right\rceil$$

4. Applications

The *m*-path ideal of a path graph. Let $n \ge m \ge 1$ be two integers and

 $I_{n,m} = (x_1 x_2 \cdots x_m, \ x_2 x_3 \cdots x_{m+1}, \ \dots, x_{n-m+1} \cdots x_n) \subset S = K[x_1, \dots, x_n],$

be the *m*-path ideal associated to the path graph of length *n*. Let $t \ge 1$. We define:

$$\varphi(n,m,t) := \begin{cases} n-t+2 - \left\lfloor \frac{n-t+2}{m+1} \right\rfloor - \left\lceil \frac{n-t+2}{m+1} \right\rceil, & t \le n+1-m \\ m-1, & t > n+1-m \end{cases}$$

According to [4, Theorem 2.6] we have that

 $\operatorname{sdepth}(S/I_{n,m}^t) \ge \operatorname{depth}(S/I_{n,m}^t) = \varphi(n, m, t).$

Assume that $t \leq n - m$ and let $S_{t+m} := K[x_1, \ldots, x_{m+t}]$. We consider the ideal

 $U_{m,t} = (x_{i_1} \cdots x_{i_m} : i_j \equiv j \pmod{m}, 1 \le j \le m) \subset S_{t+m}.$

By Euclidean division, we write t + m = am + b, where $1 \le b \le m$. According to the proof of [4, Lemma 2.4], we have that

$$U_{m,t} = V_{m,1,a+1} \cap \cdots \cap V_{m,b,a+1} \cap V_{m,b+1,a} \cap \cdots \cap V_{m,m,a}, \text{ where } V_{m,j,k} = (x_j, x_{j+m}, \dots, x_{j+(k-1)m}).$$
(4.1)

Proposition 4.1. With the above notations, we have that:

$$sdepth(U_{m,t}) \le hdepth(U_{m,t}) \le m + \left\lfloor \frac{t}{2} \right\rfloor.$$

Proof. According to Theorem 3.3, we have that

hdepth
$$(U_{m,t}) \le \left\lfloor \frac{m+t+m}{2} \right\rfloor = m + \left\lfloor \frac{t}{2} \right\rfloor$$

Now, apply Proposition 1.2.

We recall the following well known results:

Lemma 4.2. Let $I \subset S$ be a monomial ideal and $u \in S$ a monomial which do no belong to I. Then

- (1) $\operatorname{sdepth}(S/(I:u)) \ge \operatorname{sdepth}(S/I)$. ([3, Proposition 2.7(2)])
- (2) $\operatorname{sdepth}(I:u) \ge \operatorname{sdepth}(I)$. ([12, Proposition 1.3] arXiv version)

Theorem 4.3. Let
$$n \ge m \ge 1$$
 and $t \ge 1$. Let $t_0 := \min\{t, n-m\}$. We have that

$$\operatorname{sdepth}(I_{n,m}^t) \le \min\{n - \left\lceil \frac{t_0}{2} \right\rceil, n - \left\lfloor \frac{n - t_0 + 1}{m + 1} \right\rfloor + 1\}.$$

Proof. If $t \ge n - m + 1$, then $t_0 = n - m$ and, according to [4, Lemma 2.1], we have

$$I_{n,m}^{t_0} = I_{n,m}^t : (x_{n-m+1} \cdots x_n)^{t-t_0}$$

Therefore, from Lemma 4.2(2) it follows that

$$\operatorname{sdepth}(I_{n,m}^{t}) \leq \operatorname{sdepth}(I_{n,m}^{t_0})$$

Hence, we can assume that $t \leq n - m$ and $t_0 = t$.

By Euclidean division, we write n - t + 1 = q(m + 1) + r, where $0 \le r \le m$. According to [4, Lemma 2.5], there exists a monomial $w \in S$ such that:

$$(I_{n,m}^t : w) = \begin{cases} U_{m,t} + P_{m,t,q}, & r < m\\ U_{m,t} + P_{m,t,q} + (x_{n-m+1} \cdots x_n), & r = m \end{cases},$$
(4.2)

where $P_{m,t,q} \subset K[x_{t+m+1}, \ldots, x_{t+q(m+1)-1}]$ is a prime monomial ideal of height 2(q-1). If r < m then, from (4.2) and [3, Theorem 1.3] it follows that

 $\operatorname{sdepth}(I_{n,m}^t: w) \leq \min\{\operatorname{sdepth}(U_{m,t}S), \operatorname{sdepth}(P_{m,t,q}S)\}.$

Using Proposition 4.1, [9, Lemma 3.6] and [2, Theorem 2.2] we deduce that

$$\operatorname{sdepth}(I_{n,m}^t:w) \le \min\{n - \left\lceil \frac{t}{2} \right\rceil, n - q + 1\}$$

In the case r = m, we obtain the same inequality. Therefore, the required conclusion follows from Lemma 4.2(2) and the fact that $q = \lfloor \frac{n-t+1}{m+1} \rfloor$.

The *m*-path ideal of a cycle graph. Let $n > m \ge 2$ be two integer and

$$J_{n,m} = I_{n,m} + (x_{n-m+2} \cdots x_n x_1, \dots, x_n x_1 \cdots x_{m-1}) \subset S = K[x_1, \dots, x_n],$$

the m-path ideal associated to the cycle graph of length n.

Let $d := \operatorname{gcd}(n, m)$. We consider the ideal

 $U'_{n,d} = (x_1, x_{d+1}, \cdots, x_{d(r-1)+1}) \cap (x_2, x_{d+2}, \cdots, x_{d(r-1)+2}) \cap \cdots \cap (x_d, x_{2d}, \dots, x_{rd}), \quad (4.3)$ where $r := \frac{n}{d}$. Note that $U'_{n,1} = \mathbf{m} = (x_1, \dots, x_n).$

Proposition 4.4. With the above notations, we have that:

$$sdepth(U'_{n,d}) \le hdepth(U'_{n,d}) \le \left\lfloor \frac{n+d}{2} \right\rfloor.$$

Proof. According to Theorem 3.3, we have that

$$\mathrm{hdepth}(U'_{n,d}) \le \left\lfloor \frac{n+d}{2} \right\rfloor$$

Now, apply Proposition 1.2.

Let $t_0 := t_0(n, m)$ be the maximal integer such that $t_0 \le n-1$ and there exists a positive integer α such that

$$mt_0 = \alpha n + d.$$

Let $t \ge t_0$ be an integer.

Theorem 4.5. Let $n > m \ge 2$ and $t \ge t_0$. We have that

$$\operatorname{sdepth}(J_{n,m}^t) \le \left\lfloor \frac{n+d}{2} \right\rfloor.$$

15

Proof. According to [5, Lemma 2.2], there exists a monomial $w_t \in S$ such that

$$J_{n,m}^t:w_t)=U_{n,d}'.$$

The conclusion follows from Lemma 4.2(2) and Proposition 4.4.

Aknowledgments. We gratefully acknowledge the use of the computer algebra system Cocoa ([7]) for our experiments.

The second author, Mircea Cimpoeaş, was supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS - UEFISCDI, project number PN-III-P1-1.1-TE-2021-1633, within PNCDI III.

References

- J. Apel, On a conjecture of R. P. Stanley; Part II Quotients Modulo Monomial Ideals, J. of Alg. Comb. 17 (2003), 57-74.
- [2] C. Biro, D. M. Howard, M. T. Keller, W. T. Trotter, S. J. Young, Interval partitions and Stanley depth, Journal of Combinatorial Theory Series A 117(4) (2010), 475-482.
- [3] M. Cimpoeaş, Several inequalities regarding Stanley depth, Romanian Journal of Math. and Computer Science 2(1) (2012), 28-40.
- [4] S. Bălănescu, M. Cimpoeaş, Depth and Stanley depth of powers of the path ideal of a path graph, (2023), arXiv:2303.01132v1
- [5] S. Bălănescu, M. Cimpoeaş, Depth and Stanley depth of powers of the path ideal of a cycle graph, (2024), arXiv:2303.15032v3
- [6] S. Bălănescu, M. Cimpoeaş, C. Krattenthaller, On the Hilbert depth of monomial ideals, (2024), arXiv:2306.09450v4
- [7] CoCoATeam, CoCoA: a system for doing Computations in Commutative Algebra, Available at http://cocoa.dima.unige.it
- [8] A. M. Duval, B. Goeckneker, C. J. Klivans, J. L. Martine, A non-partitionable Cohen-Macaulay simplicial complex, Advances in Mathematics 299 (2016), 381-395.
- [9] J. Herzog, M. Vladoiu, X. Zheng, How to compute the Stanley depth of a monomial ideal, Journal of Algebra 322(9), (2009), 3151-3169.
- [10] J. Herzog, A survey on Stanley depth, In Monomial Ideals, Computations and Applications, Springer, (2013), 3-45.
- [11] A. Popescu, Special Stanley decompositions, Bull. Math. Soc. Sci. Math. Roumanie 53(101) No. 4, 2010, 363-372.
- [12] D. Popescu, An inequality between depth and Stanley depth, Bull. Math. Soc. Sci. Math. Roumanie 52 (100) (2009), 377-382, arXiv:0905.4597v2.
- [13] R. P. Stanley, Linear Diophantine equations and local cohomology, Invent. Math. 68, (1982), 175-193.