Extremal results on the Mostar index of trees with fixed parameters

Fazal Hayat^{*}, Shou-Jun Xu[†]

School of Mathematics and Statistics, Gansu Center for Applied Mathematics, Lanzhou University, Lanzhou 730000, P.R. China

Abstract

For a graph G, the Mostar index of G is the sum of $|n_u(e) - n_v(e)|$ over all edges e = uv of G, where $n_u(e)$ denotes the number of vertices of G that have a smaller distance in G to u than to v, and analogously for $n_v(e)$. We determine all the graphs that maximize and minimize the Mostar index respectively over all trees in terms of some fixed parameters like the number of odd vertices, the number of vertices of degree two, and the number of pendent paths of fixed length.

Keywords: Mostar index, tree, odd vertex, vertex of degree two, pendent path.

2010 Mathematics Subject Classification: 05C12; 05C90

1 Introduction

Let G be a simple connected graph with vertex set V(G) and edge set E(G). The distance between u and v in G is the least length of the path connecting u and v and is denoted by $d_G(u, v)$.

A molecular graph is a simple graph such that its vertices correspond to the atoms and the edges to the bonds of a molecule. A topological index of G is a real number related to

^{*}E-mail: fhayatmaths@gmail.com

[†]Corresponding author E-mail: shjxu@lzu.edu.cn

G. They are widely used for characterizing molecular graphs, establishing relationships between the structure and properties of molecules, predicting the biological activity of chemical compounds, and making their chemical applications.

Doslić et al. [8] introduced a bond-additive structural invariant as a quantitative refinement of the distance non-balancedness and also a measure of peripherality in graphs, named the Mostar index. For graph G, the Mostar index is defined as

$$Mo(G) = \sum_{e=uv \in E(G)} \psi(uv),$$

where $\psi(uv) = |n_u - n_v|$, and n_u denotes the number of vertices of G closer to u than to v and n_v is the number of vertices closer to v than to u.

Finding extremal (maximum and minimum) graphs either for the whole class of graphs of certain fixed order or within some sub-classes of these graphs is an important line of research and has received much attention. For example, Doslić et al. [8] determined all the graphs that maximize and minimize the Mostar index among all trees and unicyclic graphs. Tepeh [18] obtained the graphs having the smallest and largest Mostar index over all bicyclic graphs. Hayat and Zhou [13] determined the *n*-vertex cacti with the largest Mostar index, and obtained a sharp upper bound for the Mostar index among cacti of fixed order and cycles, and characterized the extremal graphs. In [14], Hayat and Zhou identified all the graphs that minimize and/or maximize the Mostar index among all trees with fixed parameters like the maximum degree, the diameter, and the number of pendent vertices. Hayat and Xu [12] investigated the lower bound on the Mostar index in tricyclic graphs, and characterized the graphs that attain the bound. Deng and Li [4] obtained those trees with a fixed degree sequence having the largest Mostar index. In [5], Deng and Li studied the extremal problem for the Mostar index among trees with a given number of segments sequence. In [6], Deng and Li determined the greatest Mostar index over all chemical trees of fixed order, and the least Mostar index among trees with fixed order and diameter. Liu and Deng [16] obtained the maximum Mostar index among unicyclic graphs with a given diameter. More related work about the Mostar index can be found in [1, 2, 3, 7, 9, 15, 17, 19, 20] and the references cited therein.

To have a full understanding of the relationship between the Mostar index and the structural properties of the graphs. In this paper, we further consider the Mostar index among trees in terms of some other fixed parameters, and more precisely, we identify all the graphs that maximize and minimize the Mostar index among *n*-vertex trees with fixed parameters like the number of odd vertices, the number of vertices of degree two, and the number of pendent paths of fixed length.

2 Preliminaries

For $v \in V(G)$, $N_G(v)$ denotes the set of vertices that are adjacent to v in G. The degree of $v \in V(G)$, denoted by $d_G(v)$, is the cardinality of $N_G(v)$. Suppose e is an edge and x, yare two non-adjacent vertices of G. Then G - e is the subgraph of G obtained by deleting the edge e and G + xy is a graph obtained from G by adding an edge connecting x, y. The graph formed from G by deleting a vertex $v \in V(G)$ (and its incident edges) is denoted by G - v. The diameter of G is the largest distance between any two vertices of G. A path $P = u_0 u_1 \cdots u_r$ with $r \ge 1$ in a graph G is called a pendent path of length r at u_r , if $d_G(u_0) = 1, d_G(u_r) \ge 2$, and if $r \ge 2, d_G(u_i) = 2$ for $1 \le i \le r-1$. Particularly, a pendent path of length one at v is just a pendent edge at v. A pendent vertex is a vertex of degree one. A vertex is called an odd (resp. even) vertex if its degree is odd (resp. even). A vertex in a tree with a degree of at least three is known as a branch vertex. A vertex is called an internal vertex if its degree is of at least 2. A tree whose all internal vertices have degree of at least three is called a series-reduced tree or a homeomorphically irreducible tree [10, 11]. By S_n and P_n we denote the star and path on n vertices, respectively. A caterpillar is a tree for which there exists some fixed path with every vertex either on the path or adjacent to a vertex on the path (the path is called the spine of the caterpillar). Note that S_n and P_n are caterpillars.

Theorem 2.1. [8] Among all trees of order n, S_n (resp. P_n) is the unique tree with maximum (resp. minimum) Mostar index.

Let G be a graph with a cut edge e = uv, and let G/e be the graph obtained from G by contracting the edge e into a new vertex w such that it is adjacent to each vertex in $N_G(u) \cup N_G(v) \setminus \{u, v\}$ and then attaching a pendent edge at w.

Lemma 2.1. [8] Let G be a graph with a cut edge e. If e is not a pendent edge, then Mo(G/e) > Mo(G).

Let G be a nontrivial graph with $u \in V(G)$. For two nonnegative integers ℓ and m, let $G_{u;\ell,m}$ be the graph obtained from G by attaching two pendent paths of length ℓ and m, respectively at u. In particular, $G_{u;0,0} = G$ and $G_{u;\ell,0}$ is obtained from G by attaching a pendent path of length ℓ .

Lemma 2.2. [8] Let G be a nontrivial graph with $u \in V(G)$. If $\ell \ge m \ge 1$, then $Mo(G_{u;\ell,m}) > Mo(G_{u;\ell+1,m-1})$.

Lemma 2.3. [13] Let G be a graph of order n with a cut edge e = uv. Then $\psi_G(e) \leq (n-2)$ with equality if and only if e = uv is a pendent edge.

Lemma 2.4. [14] Let T be a tree with $x, y \in V(T)$. Suppose that x_1, \ldots, x_s and y_1, \ldots, y_t are pendent vertices adjacent to x and y, respectively. Let x' and y' be respectively the neighbors of x and y in the path connecting x and y. Let $T' = T - \{xx_i : i = 1, \ldots, s\} + \{x'x_i : i = 1, \ldots, s\}$, and $T'' = T - \{yy_i : i = 1, \ldots, t\} + \{y'y_i : i = 1, \ldots, t\}$. Then $Mo(T) < \max\{Mo(T'), Mo(T'')\}.$

For $3 \leq r \leq n-2$, let $S_{n,r}$ be the tree consisting of r pendent paths of almost equal lengths i.e., t pendent paths of length $\lfloor \frac{n-1}{r} \rfloor + 1$ and r-t pendent paths of length $\lfloor \frac{n-1}{r} \rfloor$ at a common vertex with $t = n - 1 - r \lfloor \frac{n-1}{r} \rfloor$. Particularly, $S_{n,n-1} = S_n$, and $S_{n,2} = P_n$.

Theorem 2.2. [14] Let T be a tree of order n with r pendent vertices, where $3 \le r \le n-2$, with maximum Mostar index. Then $T \cong S_{n,r}$.

Corollary 2.1. If $2 \le r \le n-2$, then $Mo(S_{n,r}) < Mo(S_{n,r+1})$.

Proof. Let u be a vertex in $S_{n,r}$ with degree r and uv be an edge in the longest path of $S_{n,r}$, where if r = 2, u is any vertex of degree two and uv is in the longer sub-path with a terminal vertex u. By Lemma 2.1, $Mo(S_{n,r}/uv) > Mo(S_{n,r})$. Note that $S_{n,r}/uv$ is a tree of order n with r + 1 pendent vertices. By Theorem 2.2, $Mo(S_{n,r}/uv) \le Mo(S_{n,r+1})$. It follows that $Mo(S_{n,r}) < Mo(S_{n,r+1})$.

Let $T_z(d_{i_1}, d_{i_2}, ..., d_{i_z})$ be the caterpillar of order n such that $v_1 v_2 \cdots v_z$ is the spine of $T_z(d_{i_1}, d_{i_2}, ..., d_{i_z})$, where the degree of v_j is d_{i_j} with j = 1, 2, ..., z.

Lemma 2.5. [4] Let T be a tree with degree sequence $(d_1, d_2, ..., d_z)$, where $d_1 \ge d_2 \ge \cdots \ge d_z \ge 2$, $z \ge 2$ such that T minimizes the Mostar index. Then $T \cong T_z(d_{i_1}, d_{i_2}, ..., d_{i_z})$, where $d_{i_1} \ge d_{i_2} \ge \ldots \ge d_{i_s}$ and $d_{i_{s+1}} \le d_{i_{s+2}} \le \ldots \le d_{i_z}$ for some $s \in \{1, 2, \ldots, z-1\}$.

Lemma 2.6. Let T be a tree with a longest path $P_r = v_0v_1 \cdots v_r$. Let $\{v_{i-1}, v_{i+1}, u_1, \ldots, u_t\}$, where $t \ge 1$ be the neighbours of a vertex $v_i \in P_r$ $(1 \le i \le r-1)$. Suppose that n_0 (resp. n_r) be the order of the components of T - E(P) containing v_0 (resp. v_r). Let $T' = T - \{v_iu_j : 1 \le j \le c\} + \{v_0u_j : 1 \le j \le c\}$, where $1 \le c \le t$. If $n_r \ge n_0 + d_T(v_i, v_0)$, then Mo(T') < Mo(T).

Proof. Note that $n_r \ge n_0 + d_T(v_i, v_0)$, we have $n_0 + d_T(v_i, v_0) \le \frac{n}{2}$. Let $s = d_T(v_i, v_0)$ and let $z_0 \dots z_s$ be the path from v_i to v_0 , where $z_0 = v_i$ and $z_s = v_0$. Let n'_c be the number of vertices of the components of $T - v_i$ containing one of u_1, \dots, u_c . For $k = 0, \dots, s$, we have $n_0 + s - k < \min\{n_0 + n'_c + s - k, n - (n_0 + n'_c + s - k)\} \le \frac{n}{2}$, implying that $\psi_T(z_{k-1}z_k) = |(n_0 + s - k) - (n - (n_0 + s - k))| > |n_0 + n'_c + s - k - (n - (n_0 + n'_c + s - k))| =$ $\psi_{T'}(z_{k-1}z_k)$. Therefore, $Mo(T) - Mo(T') = \sum_{k=1}^s \psi_T(z_{k-1}z_k) - \sum_{k=1}^s \psi_{T'}(z_{k-1}z_k) > 0$, i.e., Mo(T) > Mo(T'). □

For $a, b \ge 0$ with $2(a + b) \le n - 1$, let $C_{n,a,b}$ be the tree obtained from the path $P_{n-a-b} = v_1 v_2 \dots v_{n-a-b}$ by attaching a pendent vertex u_i to v_i for each i with $2 \le i \le a+1$ and each i with $n - a - 2b \le i \le n - a - b - 1$. In particular, $C_{n,0,0} = P_n$.

Lemma 2.7. Let a and b be positive integers with $a \ge b+2$ and $2(a+b) \le n-3$. Then, $Mo(C_{n,a-1,b+1}) < Mo(C_{n,a,b}).$

Proof. Let $P_{1,n-a-b}$ be the path from v_1 to v_{n-a-b} . Let $T = C_{n,a,b}$, and $T' = T - v_{a+1}u_{a+1} + v_{n-a-2b-1}u_{a+1}$. Then, $T' \cong C_{n,a-1,b+1}$. From T and T', one has $\psi_T(e) = \psi_{T'}(e)$ for each $e \in T \setminus E(P_{a+1,n-a-2b-2})$. If $2a \leq \frac{n}{2}$, then $2b + 2 < 2a \leq \frac{n}{2}$, and otherwise, $2b+2 < n-2a < \frac{n}{2}$. Then Mo(T) - Mo(T') = |2b+2-(n-2b-2)| - |2a-(n-2a)| > 0. Thus Mo(T') < Mo(T), implying that $Mo(C_{n,a-1,b+1}) < Mo(C_{n,a,b})$. □

3 Mostar index and the number of odd vertices

For integers n and k with $1 \le k \le \lfloor \frac{n}{2} \rfloor$, let $\mathcal{O}(n,k)$ be the set of trees of order n with 2k odd vertices.

Theorem 3.1. Let T be in $\mathcal{O}(n,k)$, where $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$ with maximum Mostar index. Then $T \cong S_{n,2k}$, where if $k = \frac{n}{2}$, then $S_{n,n} = S_n$. *Proof.* The case k = 1 is trivial as in this case, $T \cong S_{n,2} = P_n$.

If $k = \lfloor \frac{n}{2} \rfloor$, then $S_{n,2\lfloor \frac{n}{2} \rfloor} = S_n \in \mathcal{O}(n,k)$, and so the result follows by Theorem 2.1.

Suppose that $2 \le k < \lfloor \frac{n}{2} \rfloor$, i.e., $4 \le 2k \le n-2$. Let r be the number of pendent vertices of T, then $r \le 2k$. By Theorem 2.2 and Corollary 2.1, $Mo(T) \le Mo(S_{n,r}) \le Mo(S_{n,2k})$ with equalities if and only if $T \cong S_{n,r}$ and r = 2k, i.e., $T \cong S_{n,2k}$.

Theorem 3.2. Let T be in $\mathcal{O}(n,k)$, where $1 \leq k \leq \lfloor \frac{n}{2} \rfloor$ with minimum Mostar index. Then $T \cong C_{n,\lceil \frac{k-1}{2} \rceil, \lfloor \frac{k-1}{2} \rfloor}$.

Proof. Let $T \in \mathcal{O}(n,k)$ such that Mo(T) is as small as possible. Suppose that $(d_1, d_2, ..., d_z)$ is the degree sequence of T, where $d_1 \ge d_2 \ge ... \ge d_z \ge 2$, $z \ge 2$. Then by Lemma 2.5, $T \cong T_z(d_{i_1}, d_{i_2}, ..., d_{i_z})$, where $d_{i_1} \ge d_{i_2} \ge ..., \ge d_{i_s}$ and $d_{i_{s+1}} \le d_{i_{s+2}} \le ..., \le d_{i_z}$ for some $s \in \{1, 2, ..., z-1\}$.

The case k = 1 is trivial. Suppose that $k \ge 2$. Since T is a caterpillar, let $P = v_0v_1 \cdots v_rv_{r+1}$ be the spine of T. We claim that the maximum odd degree of T is exactly 3. Otherwise, there exist a vertex v_i $(1 \le i \le r)$ such that $d_T(v_i) = 2t + 1$, where $t \ge 2$. Let u_1, \ldots, u_{2t-1} be the neighbors of v_i outside P in T. Let T' be the tree obtained from Tby deleting the pendent vertex u_{2t-1} and the edges $v_iu_1, v_iu_2, \ldots, v_iu_{2t-2}$, splitting v_i into two adjacent vertices v'_i and v''_i , joining $v_iu_1, \ldots, v_iu_{2t-3}$ to v'_i and u_{2t-2} to v''_i . Obviously, $T' \in \mathcal{O}(n,k)$. We denote by $e_1, e_2, \ldots e_{r-1}$ (resp. $f_1, f_2, \ldots f_r$) the consecutive edges in the spine $P = v_1 \cdots v_i \cdots v_r$ of T (resp. $P = v_1 \cdots v'_i v''_i \cdots v_r$ of T'). Note that the number of pendent edges in T is |E(T)| - (r-1) = n - r, and the number of pendent edges in T' is |E(T')| - r = n - 1 - r.

From T and T', one has $\psi_T(e_j) = \psi_{T'}(f_j)$ for any $j \in \{1, 2, ..., i-1\}$, and $\psi_T(e_\ell) = \psi_{T'}(f_{\ell+1})$ for any $\ell \in \{i, i+1, ..., r-1\}$. Then $Mo(T) - Mo(T') = (n-2)(n-r) - ((n-2)(n-r-1) + \psi_{T'}(f_i)) = (n-2) - \psi_{T'}(f_i) > 0$. By Lemma 2.3, we have Mo(T') < Mo(T), a contradiction. Thus, the maximum odd degree of T is exactly 3.

If $k = \frac{n}{2}$ i.e., 2k = n, then by Proposition 2.5, $T \cong C_{n, \lceil \frac{k-1}{2} \rceil, \lceil \frac{k-1}{2} \rceil}$.

Suppose that $k < \frac{n}{2}$. Then there is at least one even vertex in T. We claim that all the even vertices have a degree exactly 2. Otherwise, for some v_i $(1 \le i \le r)$, we have $d_T(v_i) = 2t$ such that $N_T(v_i) = \{v_{i-1}, v_{i+1}, u_1, \ldots, u_{2t-2}\}$, where $t \ge 2$. Suppose that n_1 (resp. n_2) be the order of the components of $T - v_i$ containing v_0 (resp. v_{r+1}). Assume that $n_1 \le n_2$. Let $T'' = T - \{v_i u_3 + v_0 u_3\}$, where v_0 is the pendent vertex in T. Note that in T'' the degree of v_i is odd and the degree of v_0 is even, implying that $T'' \in \mathcal{O}(n, k)$. By Lemma 2.6, Mo(T'') < Mo(T), a contradiction. Thus, all the even vertices in T have a degree exactly 2.

Consequently, T is a tree with exactly n - 2k vertices of a degree 2. Note that T is a caterpillar, therefore the number of pendent vertices in T is k + 2. Let t be the numbers of vertices of a degree 3 in T. The relation $\sum_{v \in V(G)} d_T(v) = 2|E(T)| = 2(n-1)$ gives that k + 2 + 2(n-2k) + 3t = 2n-2, so t = k-1. Therefore, the degree sequence of T is $(\underbrace{3, \ldots, 3, 2, \ldots, 2, 1, \ldots, 1}_{k-1})$. By Lemma 2.5, $T \cong C_{n,a,b}$, for some a, b with a + b = k - 1. By Lemma 2.7, $T \cong C_{n, \lceil \frac{k-1}{2} \rceil, \lfloor \frac{k-1}{2} \rfloor}$.

Let T be a tree of order n with k branch vertices. Let s be the number of pendent vertices in T. Then $s + 2(n - s - k) + 3k \le 2(n - 1)$, i.e., $s \ge k + 2$. This implies that $2k + 2 \le k + s \le n$, and thus, $k \le \frac{n}{2} - 1$

Corollary 3.1. Let T be a tree of order n with k branch vertices, where $0 \le k \le \frac{n}{2} - 1$ with minimum Mostar index. Then $T \cong C_{n, \lceil \frac{k}{2} \rceil, \lceil \frac{k}{2} \rceil}$.

Proof. Let T be a tree of order n with k branch vertices such that Mo(T) is as small as possible. If k = 0, then $T \cong P_n$ and it is trivial.

Suppose that $k \ge 1$. Let (d_1, d_2, \ldots, d_z) be the degree sequence of T, where $d_1 \ge d_2 \ge \cdots \ge d_z \ge 2, z \ge 2$. Then by Lemma 2.5, $T \cong T_z(d_{i_1}, d_{i_2}, \ldots, d_{i_z})$, where $d_{i_1} \ge d_{i_2} \ge, \ldots, \ge d_{i_s}$ and $d_{i_{s+1}} \le d_{i_{s+2}} \le, \ldots, \le d_{i_z}$ for some $s \in \{1, 2, \ldots, z - 1\}$.

Since T is a caterpillar, let $P = v_0 v_1 \dots v_r$ be the spine of T. Let Δ be the maximum degree of T. We claim that $\Delta = 3$. Suppose that $\Delta \ge 4$, then for some $v_i \in V(T)$, where $(1 \le i \le r - 1)$ such that $N_T(v_i) = \{v_{i-1}, v_{i+1}, u_1, \dots, u_{\Delta-2}\}$. Suppose that n_1 (resp. n_2) be the order of the components of $T - v_i$ containing v_0 (resp. v_r). Without loss of generality, we assume that $n_1 \le n_2$. Let $T' = T - \{v_i u_1 + v_0 u_1\}$, where v_0 is a pendent vertex of T. Obviously, T' is a tree of order n with k branch vertices. By Lemma 2.6, Mo(T') < Mo(T), a contradiction. Thus, $\Delta = 3$. Let s be the number of pendent vertices in T. As s + 2(n - s - k) + 3k = 2(n - 1), we have s = k + 2, and thus T is a tree of order n with 2k + 2 odd vertices. By Theorem 3.2, we have $T \cong C_{n, \lceil \frac{k}{2} \rceil, \lceil \frac{k}{2} \rceil}$.

Let \mathcal{O}_{2n} be the set of all trees with 2n vertices, which are all odd. By Theorem 2.1 and Theorem 3.2, we have the following result.

Theorem 3.3. Let T be in \mathcal{O}_{2n} , with maximum (resp. minimum) Mostar index. Then $T \cong S_{2n}$ (resp. $T \cong C_{2n,0,n-1}$).

4 Mostar index and the number of vertices of degree two

For integers n and t with $0 \le t \le n-2$, let $\mathcal{E}(n,t)$ be the set of trees of order n with t vertices of degree two. Note that $\mathcal{E}(n, n-2) = \{P_n\}$ and $\mathcal{E}(n, n-3) = \emptyset$. So we only consider the trees in $\mathcal{E}(n,t)$ with $0 \le t \le n-4$.

Theorem 4.1. Let T be in $\mathcal{E}(n,t)$, where $0 \le t \le n-4$ with maximum Mostar index. Then $T \cong S_{n,n-t-1}$.

Proof. Let $T \in \mathcal{E}(n, t)$ such that Mo(T) is as large as possible.

If t = 0, then $S_{n,n-1} = S_n \in \mathcal{E}(n,t)$, so the result follows by Theorem 2.1.

Suppose that $t \ge 1$. We claim that T contains exactly one vertex having degree at least 3. Otherwise, T contains two vertices, say u, v having degree at least 3. Suppose that u_1, \ldots, u_x and v_1, \ldots, v_y be the pendent vertices adjacent to u and v, respectively. Let u' (resp. v') be the neighbor of u (resp. v) in the path connecting u and v. Let $T' = T - \{uu_i : i = 1, \ldots, x\} + \{u'u_i : i = 1, \ldots, x\}$, and $T'' = T - \{vv_i : i = 1, \ldots, y\} + \{v'v_i : i = 1, \ldots, y\}$. Then by Lemma 2.4, $Mo(T) < \max\{Mo(T'), Mo(T'')\}$, a contradiction. Thus, T contains exactly one vertex having degree at least 3. Let z be the number of pendent vertices of T, then $z \le n - t - 1$. By Theorem 2.2 and Corollary 2.1, we have $Mo(T) \le Mo(S_{n,z}) \le Mo(S_{n,n-t-1})$ with equalities if and only if $T \cong S_{n,z}$ and z = n - t - 1, i.e., $T \cong S_{n,n-t-1}$.

For $a, b \ge 0$ with $2(a + b) \le n - 5$, let $F_{n,a,b}$ be the tree obtained from the path $P_{n-a-b-2} = v_1v_2\cdots v_{n-a-b-2}$ by attaching a pendent vertex u_1 to v_2 , and then attaching pendent vertex u_i to v_i for each i with $2 \le i \le a+2$ and each i with $n-a-2b-2 \le i \le n-a-b-3$.

Lemma 4.1. For $a \ge b \ge 1$ and 2(a+b) < n-5, we have $Mo(F_{n,a-1,b+1}) < Mo(F_{n,a,b})$.

Proof. Let $P_{1,n-a-b-2}$ be the path from v_1 to $v_{n-a-b-2}$. Let $T = F_{n,a,b}$, and $T' = T - v_{a+2}u_{a+2} + v_{n-a-2b-3}u_{a+2}$. Then $T' \cong F_{n,a-1,b+1}$. From T and T', one has $\psi_T(e) = \psi_{T'}(e)$ for each $e \in T \setminus E(P_{a+2,n-a-2b-4})$. If $2a+3 \leq \frac{n}{2}$, then $2b+2 < 2a+3 \leq \frac{n}{2}$, and otherwise, $2b+2 < n-2a-3 < \frac{n}{2}$. Then Mo(T) - Mo(T') = |2b+2 - (n-2b-2)| - |2a+3 - (n-2a-3)| > 0. Thus Mo(T') < Mo(T), implying that $Mo(F_{n,a-1,b+1}) < Mo(F_{n,a,b})$. □

Theorem 4.2. Let T be in $\mathcal{E}(n,t)$, where $0 \le t \le n-4$ with minimum Mostar index.

(i) If n-t=5, then $T \cong F_{n,0,0}$. If n-t is odd and at least 7, then $T \cong F_{n,\lceil \frac{n-t-5}{4}\rceil-1,\lfloor \frac{n-t-5}{4}\rfloor+1}$.

(ii) If
$$n-t$$
 is even, then $T \cong C_{n,\lceil \frac{n-t}{4} - \frac{1}{2} \rceil, \lfloor \frac{n-t}{4} - \frac{1}{2} \rfloor}$.

Proof. Let T be in $\mathcal{E}(n,t)$ such that Mo(T) is as small as possible. Suppose that $(d_1, d_2, ..., d_z)$ be the degree sequence of T, where $d_1 \ge d_2 \ge ... \ge d_z \ge 2, z \ge 2$. By lemma 2.5, $T \cong T_z(d_{i_1}, d_{i_2}, ..., d_{i_z})$, where $d_{i_1} \ge d_{i_2} \ge ... \ge d_{i_s}$ and $d_{i_{s+1}} \le d_{i_{s+2}} \le ..., \le d_{i_z}$ for some $s \in \{1, 2, ..., z - 1\}$.

Since T is a caterpillar, let $P = v_0 v_1 \cdots v_r$ be the spine of T. Let Δ be the maximum degree of T. We claim that $\Delta \leq 4$. Suppose that $\Delta \geq 5$, then for some $v_i \in V(T)$, where $(1 \leq i \leq r-1)$ such that $N_T(v_i) = \{v_{i-1}, v_{i+1}, u_1, \ldots, u_{\Delta-2}\}$. Suppose that n_1 (resp. n_2) be the order of the components of $T - v_i$ containing v_0 (resp. v_r). Assume that $n_1 \leq n_2$. Let $T' = T - \{v_i u_j : 2 \leq j \leq \Delta - 2\} + \{v_0 u_j : 2 \leq j \leq \Delta - 2\}$, where v_0 is a pendent vertex of T. Obviously, $T' \in \mathcal{E}(n, t)$. By Lemma 2.6, Mo(T') < Mo(T), a contradiction. Thus, $\Delta \leq 4$.

Further, we claim that T has at most one vertex of degree 4. Suppose that T contains at least two vertices of degree 4, say x, y. Let n_x (resp. n_y) be the order of the component of T - E(P) containing x (resp. y). Assume that $n_x \ge n_y + d_T(x, y)$. Then $n_y + d_T(x, y) \le \frac{n}{2}$. Let u be the neighbor of x outside P in T. Let T'' = T - xu + yu. Obviously, $T'' \in \mathcal{E}(n, t)$. By Lemma 2.6, Mo(T') < Mo(T), a contradiction. Thus, T contains at most one vertex of degree 4.

Let d_i be the number of vertices of degree i in T, where $1 \leq i \leq 4$. Note that $d_1 + d_2 + d_3 + d_4 = n$ and $d_1 + 2d_2 + 3d_3 + 4d_4 = 2(n-1)$. Since $d_2 = t$ and $d_4 \leq 1$. Case 1. n - t is odd.

From the above relation, we have $d_1 = \frac{n-t+3}{2}$, $d_3 = \frac{n-t-5}{2}$, and $d_4 = 1$. If n-t = 5, then the degree sequence of T is $(4, 2, ..., 2, \underbrace{1, ..., 1}_{n-t-1})$. By Lmma 2.5, $T \cong F_{n,0,0}$. If $n-t \ge 7$, then the degree sequence of T is $(4, \underbrace{3, ..., 3}_{n-t-5}, \underbrace{2, ..., 2}_{t}, \underbrace{1, ..., 1}_{n-t+3})$. By Lemma 2.5, $T \cong F_{n,a,b}$, for some a, b with $a + b = \frac{n-t-5}{2}$. By Lemma 4.1, $T \cong F_{n,\lceil \frac{n-t-5}{4}\rceil - 1,\lfloor \frac{n-t-5}{4}\rfloor + 1}$. Case 2. n-t is even.

From the above relation, we have $d_1 = \frac{n-t+2}{2}$, $d_3 = \frac{n-t-2}{2}$, and $d_4 = 0$. Since $d_4 = 0$,

then the maximum degree of T is 3, and the degree sequence of T is $(\underbrace{3, ..., 3}_{\frac{n-t-2}{2}}, \underbrace{2, ..., 2}_{t}, \underbrace{1, ..., 1}_{\frac{n-t+2}{2}})$ by Proposition 2.5, $T \cong C_{n,a,b}$, for some a, b with $a + b = \frac{n-t-2}{2}$. By Lemma 2.7, $T \cong C_{n,\lceil \frac{n-t}{4} - \frac{1}{2}\rceil, \lfloor \frac{n-t}{4} - \frac{1}{2}\rfloor}$.

As an immediate consequence of Theorem 4.2, we obtain the following result.

Corollary 4.1. Let T be a series-reduced tree of order $n \ge 5$ with minimum Mostar index. Then $T \cong F_{n,\lceil \frac{n-5}{4}\rceil-1,\lfloor \frac{n-5}{4}\rfloor+1}$ (resp. $T \cong C_{n,\lceil \frac{n-2}{4}\rceil,\lfloor \frac{n-2}{4}\rfloor}$) for odd n (resp. even n).

5 Mostar index and the number of pendent paths of fixed length

For integers n, r and k, let $S_{n,k}^r$ be the tree consisting of k pendent paths of length r and n-kr-1 pendent paths of length one at a common vertex, where k = 1 and $2 \le r \le n-3$, or $k \ge 2$, $r \ge 2$ and $kr \le n-2$.

Theorem 5.1. Let T be a tree of order n with k pendent paths of length r, where k = 1and $2 \le r \le n-3$, or $k \ge 2$, $r \ge 2$ and $kr \le n-2$, with maximum Mostar index. Then $T \cong S_{n,k}^r$.

Proof. Let T be a tree of order n with k pendent paths of length r such that Mo(T) is as large as possible.

Let X be the set of vertices at which there is a pendent path of length r in T. Obviously, the order of X is at least one. We claim that X contains exactly one vertex. Suppose that $|V(X)| \ge 2$, and $v_1, v_2 \in V(X)$ such that $d_T(v_1, v_2)$ is as small as possible. Let P be the path connecting v_1 and v_2 . If $d_T(v_1, v_2) > 1$, then each internal vertex of P is of degree two. Let n_1 (resp. n_2) be the order of the component of T - E(P) containing v_1 (resp. v_2). Assume that $n_1 \ge n_2$. Obviously, $n_2 \le \frac{n}{2}$. Let u be the neighbor of v_2 in P and w be one other neighbor of v_2 . Let $n_w = n_w(v_2w|T)$. Let $T' = T - v_2w + uw$. Obviously, T' is a tree of order n with k pendent paths of length r. Note that $\psi_T(e) = \psi_{T'}(e)$ for $e \in E(T) \setminus \{uv_2, v_2w\} = E(T') \setminus \{uv_2, uw\}$ and $\psi_T(v_2w) = \psi_{T'}(uw)$. Then

$$Mo(T) - Mo(T') = \psi_T(uv_2) - \psi_{T'}(uv_2) = |n_2 - (n - n_2)| - |(n_2 - n_w) - n - (n_2 - n_w)| < 0,$$

implying that Mo(T) < Mo(T'), a contradiction. Thus, T consists of k pendent paths of length r at a common vertex. Let $X = \{z\}$

Note that if k = 1 (resp. $k \ge 2$) then there are two (resp. one) neighbours of z not lying on the pendent path of length r. We claim that all neighbours of z not lying on the pendent path of length r are pendent. Let s be any such neighbours of z. Suppose that $d_T(s) \ge 2$. Let T/zs be the tree obtained from T by contracting the edge e = zs into a new vertex, say u (such that it is adjacent to each vertex in $N_T(z) \cup N_T(s) \setminus \{z, s\}$ and then attaching a pendent edge at u. Obviously, T/zs is a tree of order n with k pendent paths of length r. By Lemma 2.1, Mo(T) < Mo(T/zs), a contradiction. Thus, $d_T(s) = 1$. That is, $T \cong S_{n,k}^r$.

For positive integers n, r, a and b with $a \ge b$ and $(a + b)r \le n - 2$, let $A_{n,a,b}^r$ be the *n*-vertex tree obtained from the path $P_{n-a-b} = v_1v_2 \dots v_{n-a-b}$ by attaching a and bpendent paths of length r at v_1 and v_{n-a-b} , respectively.

Lemma 5.1. Suppose that $a \ge b + 2$ and (a + b)r < n - 1. Then $Mo(A^r_{n,a-1,b+1}) < Mo(A^r_{n,a,b})$.

Proof. Let $P_{1,n-a-b}$ be the path from v_1 to v_{n-a-b} . Let $T = A_{n,a,b}^r$. Let V_1 be all the vertices of a pendent paths at v_1 except v_1 and $u_1u_2\cdots u_rv_1$ be one pendent path at v_1 . Let V_2 be all the vertices of b pendent paths at v_{n-a-b} except v_{n-a-b} and $w_1w_2\cdots w_rv_{n-a-b}$ be one pendent path at v_{n-a-b} . Let $T' = T - v_1u_r + v_{n-a-b}u_r$. Clearly, $T' \cong A_{n,a-1,b+1}^r$. From T and T', one has $\psi_T(e) = \psi_{T'}(e)$ for each $e \in T \setminus E(P_{1,n-a-b})$.

If $ar \leq \frac{n}{2}$, then $(b+1)r < ar \leq \frac{n}{2}$, and otherwise, $(b+1)r < n - ar < \frac{n}{2}$. Then $Mo(A^r_{n,a,b}) - Mo(A^r_{n,a-1,b+1}) = |(b+1)r - (n - br - r)| - |ar - (n - ar)| > 0$. Thus, $Mo(A^r_{n,a,b}) > Mo(A^r_{n,a-1,b+1})$.

Theorem 5.2. Let T be a tree of order n with k pendent paths of length r with minimum Mostar index.

- (i) If k = 1 and $2 \le r \le n 3$, then $T \cong A_{n,1,2}^1$.
- (ii) If k = 2 and $1 \le r \le n 2$, then $T \cong P_n$.
- (iii) If $k \geq 3$ and $1 \leq r \leq \frac{n-2}{k}$, then $T \cong A^r_{n, \lceil \frac{k}{2} \rceil, \lceil \frac{k}{2} \rceil}$.

Proof. Let T be a tree of order n with k pendent paths of length r such that Mo(T) is as small as possible.

If k = 1 and $2 \le r \le n-3$, then $T \cong P_n$. As P_n is *n*-vertex tree with two pendent paths of length *r* for each r = 2, ..., n-2 and $A_{n,1,2}^1$ is *n*-vertex tree with one pendent path of length *r* for each r = 2, ..., n-3 it follows that $T \cong A_{n,1,2}^1$. This proves Item (i).

If k = 2 and $1 \le r \le n - 2$, then the result follows by Theorem 2.1. As P_n is the unique tree with minimum Mostar index among all *n*-vertex trees. This proves Item (ii).

Suppose that $k \ge 3$ and $1 \le r \le \frac{n-2}{k}$. Let X be the set of vertices at which there is a pendent path of length r in T. Clearly, the order of X is at least one. We claim that X contains exactly two vertices, we need to show that $|X| \le 2$ and $|X| \ge 2$.

First, we show that $|X| \leq 2$. Suppose that $|X| \geq 3$. We may choose two vertices, say u, v such that $d_T(u, v)$ is as large as possible. Let P be the path connecting u and v. Let n_u (resp. n_v) be the order of the component of T - E(P) containing u (resp. v). We may choose vertices w and c in P such that both $d_T(u, w)$ and $d_T(c, v)$ is as small as possible. Assume that $n_u + d_T(u, w) \geq n_v + d_T(c, v)$. Then $n_v + d_T(c, v) \leq \frac{n}{2}$. Let $x_1, \ldots x_p$ be the neighbors of c outside P in T, where $p = d_T(c) - 2$. Let $T' = T - \{cx_i : i = 1, \ldots, p\} + \{vx_i : i = 1, \ldots, p\}$. Obviously, T' is a tree of order n with k pendent paths of length r. By Lemma 2.6, Mo(T) > Mo(T'), a contradiction. Thus, $|X| \leq 2$.

Secondly, we show that $|X| \ge 2$. If r = 1, and k < n - 1, we have $|X| \ge 2$. Suppose that $r \ge 2$ and |X| = 1. Then k pendent paths of length r of T are k branches of T at a common vertex, say v which we denote by Q_1, \ldots, Q_k . Let $T_v = T - \bigcup_{i=1}^k (V(Q_i) \setminus \{v\})$. Then T_v is a nontrivial tree as rk < n - 1. Moreover, T_v is not pendent path of length greater than or equal to r at v, as otherwise, T must have k + 1 pendent paths of length r, which is impossible.

Let $X = V(Q_1) \setminus \{v\}$. Suppose that there is a pendent path, say B_1 , at v in T, which is also a path in T_v . Then the length ℓ of B_ℓ is less than r. Let $X' = V(B_1) \setminus \{v\}$ and G = T - X - X'. Then $T \cong G_{v,r,l}$. It is easy to see that $G_{v,r+l,0}$ is a tree of order n with k pendent paths of length r. By Lemma 2.2, we have $Mo(T) = Mo(G_{v,r,\ell}) >$ $Mo(G_{v,r+1,\ell-1}) > \cdots > Mo(G_{v,r+\ell,0})$, a contradiction. Thus, there is no pendent path at v in T, which is also a path in T_v . There must be a vertex different from v with degree at least 3 in T_v . Choose such a vertex u so that the distance to v is as small as possible. Let Q be the branch of T_v at v containing u. Note that there are two pendent paths at u in T_v . Let s_1 and s_2 be their lengths with $s_1 \leq s_2$. Then $T \cong H_{u,s_1,s_2}$, where H is a subtree of T. By repeating this process, we may obtain a tree T^* from T such that the branch Qof T_v at v is changed into a pendent path B_2 at v with length s = |V(Q| - 1). By Lemma 2.2, we have $Mo(T) > Mo(H_{v,s_1+s_2,0}) > \cdots > Mo(T^*)$. Note that Q_1, \ldots, Q_k have still k pendent paths of length r in T^* . Let $Y = V(B_2) \setminus \{v\}$. Let $H' = T^* - X - Y$. Then $T^* \cong H'_{v,r,s}$. Clearly, $H'_{v,r+s,0}$ ia a tree of order n with k pendent paths of length r. By Lemma 2.2, we have $Mo(T^*) = Mo(H'_{v,r,s}) > \cdots > Mo(H'_{v,r+s,0})$, a contradiction. Thus, $|X| \geq 2$. Hence, |X| = 2, as claimed.

Now we show that $T \cong A_{n,a,b}^r$ for some a and b with $a \ge b \ge 1$ and a+b=k. If r=1, it is trivial. Suppose that $r \ge 2$. Assume that $X = \{s_1, s_2\}$. Let P be the path connecting s_1 and s_2 . Suppose that there is a vertex of degree at least three different from s_1 and s_2 , say w on P. Let n_1 (resp. n_2) be the order of the component of T - w containing s_1 (resp. s_2). Assume that $n_1 \ge n_2 + d_T(w, s_2)$. Let x_1, \ldots, x_p be the neighbors of w outside P in T. Let $T^* = T - \{wx_i : i = 1, \ldots, p\} + \{s_2x_i : i = 1, \ldots, p\}$. Obviously, T^* is a tree of order n with k pendent paths of length r. By Lemma 2.6, $Mo(T) > Mo(T^*)$, a contradiction. Thus, all internal vertices of P (if any exist) is of degree 2.

Denote by a (resp. b) the number of pendent paths of length r at s_1 (resp. s_2). Obviously, a + b = k, $d_T(s_1) \ge a + 1$ and $d_T(s_2) \ge b + 1$. If $d_T(s_1) \ge a + 2$, then there is a branch, say Q at s_1 not containing s_2 , which is not a pendent path of length r, but, as above by Lemma 2.2, Q can not be a pendent path and it can not have a vertex different from s_1 of degree at least 3. So $d_T(s_1) = a + 1$. Similarly, $d_T(s_2) = b + 1$. Thus, $T \cong A_{n,a,b}^r$ for some a and b with $a, b \ge 1$ and a + b = k. Assume that $a \ge b$. If $a \ge b + 2$, then by Lemma 5.1, $Mo(A_{n,a-1,b+1}^r) < Mo(A_{n,a,b}^r)$, a contradiction. Therefore, a - b = 0, 1, implying that $T \cong A_{n,\lceil \frac{k}{2} \rceil, \lfloor \frac{k}{2} \rfloor}^r$. The case r = 1 has been proved in [14]. This completes the proof.

6 Concluding remarks

In this paper, we identified all the graphs having the largest and smallest Mostar index respectively among n-vertex trees with fixed parameters like the number of odd vertices, the number of vertices of degree two and the number of pendent paths of fixed length. We also identified the unique graph that maximizes and minimizes the Mostar index respectively among trees with all odd vertices. Moreover, we identified the unique graph that minimizes the Mostar index over all trees of order n with a fixed number of branch vertices and among all series-reduced trees, respectively. For future work, it will be interesting to determine all the graphs that maximize and minimize the Mostar index over all series-reduced trees of order n with fixed parameters like the diameter, the maximum degree, and the number of pendent vertices.

Acknowledgement: This work was partially supported by the National Natural Science Foundation of China (Grant Nos. 12071194, 11571155).

References

- M. Arockiaraj, J. Clement and N. Tratnik, Mostar indices of carbon nanostructures and circumscribed donut benzenoid systems. Int. J. Quantum Chem. 119 (2019) e26043.
- [2] A. Ali, T. Došlić, Mostar index: results and perspectives. Appl. Math. Comput. 404 (2021) 19. 126245.
- [3] Y. Alizadeh, K. Xu, S. Klavžar, On the Mostar index of trees and product graphs. Filomat 35 (2021) 4637–4643.
- [4] K. Deng, S. Li, On the extremal values for the Mostar index of trees with given degree sequence. Appl. Math. Comput. 390 (2021) 125598.
- [5] K. Deng, S. Li, On the extremal Mostar indices of trees with a given segment sequence. Bull. Malays. Math. Sci. Soc. 45 (2021) 593–612.
- [6] K. Deng, S. Li, Chemical trees with extremal Mostar index. MATCH Commun. Math. Comput. Chem. 85 (2021) 161–180.
- [7] K. Deng, S. Li, Extremal catacondensed benzenoid with respect to the Mostar index. J. Math. Chem. 58 (2020) 1437–1465.
- [8] T. Došlić, I. Martinjak, R. Skrekovski, S. Tipurić Spužević, I. Zubac, Mostar index. J. Math. Chem. 56 (2018) 2995–3013.

- [9] F. Gao, K. Xu, T. Došlić, On the difference between Mostar index and irregularity of graphs. Bull. Malays. Math. Sci. Soc. 44 (2021) 905–926.
- [10] J. Haslegrave, Extremal results on average subtree density of series-reduced trees. J. Combin. Theory Ser. B. 107 (2014) 26–41.
- [11] F. Harary, G. Prins, The number of homeomorphically irreducible trees, and other species. Acta Math. 101 (1959) 141–162.
- [12] F. Hayat, S. J. Xu, A lower bound on the Mostar index of tricyclic graphs. Filomat (Accepted).
- [13] F. Hayat, B. Zhou, On cacti with large Mostar index. Filomat 33 (2019) 4865–4873.
- [14] F. Hayat, B. Zhou, On Mostar index of trees with parameters. Filomat 33 (2019) 6453–6458.
- [15] S. Huang, S. Li, M. Zhang, On the extremal Mostar indices of hexagonal Chains. MATCH Commun. Math. Comput. Chem. 84 (2020) 249–271.
- [16] G. Liu, K. Deng, The maximum Mostar indices of unicyclic graphs with given diameter. Appl. Math. Comput. 439 (2023) 127636.
- [17] S. Miklavič, J. Pardey, D. Rautenbach, F. Werner, Maximizing the Mostar index for bipartite graphs and split graphs. Discrete Optim. 48 (2023) 100768.
- [18] A. Tepeh, Extremal bicyclic graphs with respect to Mostar index. Appl. Math. Comput. 355 (2019) 319–324.
- [19] Q. Xiao, M. Zeng, Z. Tang, The hexagonal chains with the first three maximal Mostar indices. Discrete Appl. Math. 288 (2020) 180–191.
- [20] Q. Xiao, M. Zeng, Z. Tang, H. Deng, H. Hua, Hexagonal chains with first three minimal Mostar indices. MATCH Commun. Math. Comput. Chem. 85 (2021) 47–61.