

# Composition of nested embeddings with an application to outlier removal

Shuchi Chawla<sup>1</sup> and Kristin Sheridan<sup>1</sup>

<sup>1</sup>Department of of Computer Science, University of Texas, Austin, TX

## Abstract

We study the design of embeddings into Euclidean space with outliers. Given a metric space  $(X, d)$  and an integer  $k$ , the goal is to embed all but  $k$  points in  $X$  (called the "outliers") into  $\ell_2$  with the smallest possible distortion  $c$ . Finding the optimal distortion  $c$  for a given outlier set size  $k$ , or alternately the smallest  $k$  for a given target distortion  $c$  are both NP-hard problems. In fact, it is UGC-hard to approximate  $k$  to within a factor smaller than 2 even when the metric sans outliers is isometrically embeddable into  $\ell_2$ . We consider bi-criteria approximations. Our main result is a polynomial time algorithm that approximates the outlier set size to within an  $O(\log^4 k)$  factor and the distortion to within a constant factor.

The main technical component in our result is an approach for constructing a composition of two given embeddings from subsets of  $X$  into  $\ell_2$  which inherits the distortions of each to within small multiplicative factors. Specifically, given a low  $c_S$  distortion embedding from  $S \subset X$  into  $\ell_2$  and a high(er)  $c_X$  distortion embedding from the entire set  $X$  into  $\ell_2$ , we construct a single embedding that achieves the same distortion  $c_S$  over pairs of points in  $S$  and an expansion of at most  $O(\log k) \cdot c_X$  over the remaining pairs of points, where  $k = |X \setminus S|$ . Our composition theorem extends to embeddings into arbitrary  $\ell_p$  metrics for  $p \geq 1$ , and may be of independent interest. While unions of embeddings over disjoint sets have been studied previously, to our knowledge, this is the first work to consider compositions of *nested* embeddings.

# 1 Introduction

Low distortion metric embeddings are an important algorithmic tool with a myriad of applications. The goal is to transform a dataset that lies in an unwieldy metric space into one lying in a nicer space, enabling clean and fast algorithms. Embeddings play an important role in the design of approximation algorithms, in finding good low dimensional representations for data in machine learning and data science, in data visualization, in the design of fast algorithms, and more. (See, e.g., [20].)

One of the most enduring concepts from the vast literature on embeddings is that of distortion which is defined to be the maximum ratio over all pairs of points in the metric by which the distance between the points is expanded or contracted by the embedding. Since this is a worst case notion, it can be particularly sensitive to errors or noise or even intentional corruption of the underlying data. Indeed adversarial or random data corruption is a frequent problem in data science contexts.

Motivated by these applications, Sidiropoulos et al. [29] introduced the notion of embeddings with outliers. Formally, given a metric space  $(X, d_X)$  of finite size and a target space  $(Y, d_Y)$  our goal is to find a low-distortion embedding into  $Y$  of all but a few points in  $X$ ; these points are called the outliers. Specifically, we say that the space  $(X, d_X)$  has a  $(k, c)$ -outlier embedding into space  $(Y, d_Y)$  if there exists an outlier set  $K \subset X$  of size at most  $k$  and a map  $\alpha$  from  $X \setminus K$  to  $Y$  with distortion at most  $c$ . [29] showed that for many host spaces  $(Y, d_Y)$  of interest, computing the optimal outlier set size  $k$  for a given target distortion  $c$ , or vice versa the optimal distortion  $c$  for a given target outlier set size  $k$ , is NP-hard.

In this work, we study the design of approximately optimal outlier embeddings into the Euclidean metric. Given a metric  $(X, d_X)$  that admits a  $(k, c)$ -outlier embedding into  $\ell_2$ , we provide a polynomial time algorithm that constructs an  $(O(k \text{ polylog } k), O(c))$ -outlier embedding into  $\ell_2$ . In other words, our algorithm removes  $O(k \text{ polylog } k)$  outliers and returns an embedding of the remaining metric into  $\ell_2$  that has distortion only a small constant factor worse than the desired one. In fact, our algorithm allows for a tradeoff between the outlier set size and the distortion obtained – allowing us to obtain a distortion of  $(1 + \epsilon)c$  for any  $\epsilon > 0$  at the cost of blowing up the outlier set size by an additional multiplicative factor of at most  $1/\epsilon$ .

To our knowledge, the only approximations to multiplicative distortion for outlier embeddings known prior to our work considered the special case of embedding unweighted graphs into a line. For this setting, Chubarian and Sidiropoulos [11] developed an algorithm that constructs an  $(O(c^\delta k \log^{5/2}(n)), O(c^{13}))$ -outlier embedding when the input metric is an unweighted graph metric. Here  $n$  is the size of the given metric  $X$ . While our result is incomparable to theirs (as it does not limit the dimension of the embedding), we emphasize that our approximation factors do not depend on the size of the metric  $X$  and has a vastly improved dependence on the distortion  $c$ , as we discuss below.

**Composition of nested embeddings.** The main technical tool in our work is what we call a composition of nested embeddings, which may be of independent interest. Consider a metric space  $(X, d_X)$  and a subspace  $S \subset X$ . Let  $\alpha_X : X \rightarrow Y$  and  $\alpha_S : S \rightarrow Y$  be two embeddings from  $X$  and  $S$  into  $Y$  with distortions  $c_X$  and  $c_S$  respectively. We call these *nested embeddings*. In general we would expect that the distortion of  $\alpha_S$  is smaller than that of  $\alpha_X$ . In fact, the distortion of  $\alpha_X$  even restricted to just the subset  $S$  may be larger than  $c_S$ . In that case, we ask: is it possible to obtain a new embedding from the entire set  $X$  into  $Y$  such that the distortion of this embedding over just the “inner” set  $S$  is the same as before –  $c_S$  – while the distortion over all of  $X$  is comparable to  $c_X$ ? In particular, we want a combination of the two embeddings that inherits their respective distortions, with small multiplicative worsening, over the corresponding sets of points.

We show that for embeddings into the  $\ell_p$  metric for  $p \geq 1$ , this is indeed achievable with a small caveat. In particular, we construct a new embedding based on  $\alpha_S$  and  $\alpha_X$  that achieves a distortion of  $c_S$  over the set  $S$ , while at the same time expanding distances over the remaining pairs of points in  $X$  by a factor of at most  $c_X$  times  $O(\log(|X \setminus S|))$ . The caveat is that some of the latter distances may shrink. We call such a composition a *weak nested composition* (see Definition 2.4).<sup>1</sup>

---

<sup>1</sup>When the host metric is  $\ell_1$ , we also achieve expansion, obtaining a net distortion of at most  $O(\log(|X \setminus S|))c_X$  over all pairs of points.

This concept of composition is similar in some ways to unions of embeddings studied previously in [23] and [26]. The goal in these works is to combine two embeddings over *disjoint* subsets  $A$  and  $B$  of a metric space into a single embedding over their union such that the distortion over the union is comparable to the distortions  $c_A$  and  $c_B$  of the given embeddings. When the host metric is  $\ell_2$  or  $\ell_\infty$ , these works construct unions whose distortion is some constant factor times the *product*,  $c_A \cdot c_B$ , of the distortions of the given embeddings. Additionally, both the sets inherit the same distortion bound – if, for example,  $c_A \ll c_B$ , it is not guaranteed that points in  $A$  will retain similarly lower distortion in the composition. Extending these results to  $\ell_p$  metrics for  $p \notin \{2, \infty\}$  is open. By contrast, our composition guarantees *the same* distortion  $c_S$  as the given embedding on the “inner” set of points  $S$ , and a bound on the expansion over the remaining points that is linear in  $c_X$  (although the latter does not imply a bound the distortion over  $X$  as the composed embedding may contract some pairs of points). Moreover, our approach works for all  $\ell_p$  metrics for  $p \geq 1$ . Nesting provides us with crucial structure to obtain these bounds.

Our composition theorem provides an even stronger guarantee when all small enough subsets of  $X$  embed into  $\ell_p$  with low distortion. In particular, suppose that every subset of  $X$  of size  $k$  embeds into  $\ell_p$  with distortion  $\zeta_k$ , and suppose that in addition we are given an embedding  $\alpha_S$  from some  $S \subset X$  into  $\ell_p$ . Then we obtain a composition where the distortion of all pairs of points in  $S$  is at most  $c_S$  and distances over all remaining pairs of points in  $X$  expand by at most  $O(\zeta_k \log k)$  for  $k := |X \setminus S| + 1$ .

This sort of guarantee is reminiscent of local versus global guarantees for metric embeddings. In particular, Arora et al. [3] ask: suppose that *every* subset of metric space  $X$  of size at most  $k$  admits a low distortion embedding into  $\ell_1$ , does  $X$  also admit a low distortion embedding into  $\ell_1$ ? Charikar et al. [9] show that this is indeed possible but that the distortion blows up by a factor of  $\Theta(\log n/k)$ . Our setting is slightly different in that we not only need every small set of size  $k$  to be embeddable with low distortion, but we also need a good embedding for *one* set of size  $n - k + 1$ .

**A tradeoff between outlier set size and distortion.** Armed with a weak nested composition into  $\ell_2$  space, we develop an SDP-rounding algorithm for obtaining bicriteria approximations for outlier embeddings into  $\ell_2$ . For a metric  $(X, d_X)$  that admits a  $(k, c)$ -outlier embedding into  $\ell_2$ , and any given target distortion  $c' = \gamma c$ ,  $\gamma \geq 1$ , our algorithm constructs a  $(k', c')$ -outlier embedding with at most  $k' = O(\frac{\beta^2}{\gamma^2 - 1} \frac{\zeta^2}{c^2} \cdot k)$  outliers. Here  $\beta$  is the multiplicative factor incurred in the distortion of the nested composition we construct, and  $\zeta$  is the worst-case distortion of embedding any size  $k$  subset of  $X$  into  $\ell_2$ . Both  $\beta$  and  $\zeta$  are  $O(\log k)$ , implying a polylogarithmic approximation for the number of outliers. Note further that our approximation factor depends inversely on  $\zeta/c$  – as the distortion of the non-outliers becomes closer and closer to the overall worst case distortion for  $X$ , it becomes easier to approximate the outlier set.<sup>2</sup> Finally, we achieve a tradeoff between the outlier set size  $k'$  and the target distortion  $c'$ . Setting  $c' = (1 + \epsilon)c$  for a small  $\epsilon > 0$ , for example, increases the outlier set size by an  $O(1/\epsilon)$  factor.

**Hardness of approximation.** Finally, we note that Sidiropoulos *et al.* [29] showed that it is NP-hard to determine the size of the smallest outlier set such that the remaining metric is isometrically embeddable into  $\ell_2^d$  for any fixed dimension  $d > 1$ . Designing an outlier embedding into  $\ell_2$  with arbitrary dimension is potentially an easier problem. Since we do not limit the dimension of the outlier embeddings we construct, we revisit and strengthen [29]’s result along these lines. We show that NP-hardness continues to hold even for embeddings into  $\ell_2$  without a specified dimension bound, and also when the given metric is the shortest path metric of an unweighted undirected graph. Our construction of a hard instance is arguably simpler than that of Sidiropoulos *et al.*, and is readily seen to extend to  $\ell_p$  metrics for  $p > 1$ . We present a separate, more involved, construction for  $p = 1$ . As with Sidiropoulos *et al.*’s results we show that, under the unique games conjecture, it is also hard to obtain a  $2 - \epsilon$  approximation for the minimum outlier set size, for any  $\epsilon > 0$ .

## Further related work

**Approximations for distortion.** Much of the work on low distortion embeddings focuses on providing uniform bounds for embedding any given finite metric into a structured space. Indyk and Matousek [20] give an excellent

<sup>2</sup>In contrast, [11]’s approximation factor for  $k$  for embeddings into a line has a polynomial dependence on  $c$ .

overview of many of these results. Bourgain’s Theorem [7, 22] shows that all finite metric spaces of size  $n$  have an  $O(\log n)$ -distortion embedding into  $\ell_p$ -space for  $p \geq 1$  and that such a randomized version of such an embedding can be computed quickly. A derandomized such embedding can be computed in polynomial time with  $\Theta(n^2)$  dimensions.

More closely related to our work, a number of papers study the objective of approximating the instance-specific minimum distortion for embedding into various host metrics. This includes, e.g., embeddings into constant dimensional Euclidean space [5, 13, 17, 24, 28], the line [24, 18, 4, 25], trees [10, 6], and ultrametrics [2]. These works tend to use combinatorial arguments as they focus on low dimensional embeddings, whereas ours relies on an SDP formulation of the problem.

**Outlier embeddings.** The notion of outlier embeddings was first introduced by Sidiropoulos *et al.* in [29]. In that paper, they showed that it is NP-hard to find the size of a minimum outlier set for embedding a metric into ultrametrics, tree metrics, or  $\ell_2^d$  for constant  $d$ . Under the Unique Games Conjecture, it is also NP-hard to approximate these values to a factor better than 2. On the algorithmic side, they gave polynomial time algorithms to 3, 4, or 2-approximate minimum outlier set size for isometric embeddings into ultrametrics, tree metrics, or  $\ell_2^d$  for fixed constant  $d$ , respectively. The algorithm for  $\ell_2^d$  embeddings is exponential in  $d$ , so  $d$  cannot grow with the size of the input while remaining efficient.

Sidiropoulos *et al.* also gave bi-criteria approximations for  $\ell_\infty$  (i.e. additive) distortion. In particular, they give a polynomial time algorithm to find embeddings with at most  $2k$  outliers and  $O(\sqrt{\delta})$   $\ell_\infty$ -distortion when there exists an embedding of the metric with at most  $k$  outliers that has  $\ell_\infty$ -distortion at most  $\delta$ . The algorithm is polynomial in  $k$ ,  $\delta$ , and  $n$  but exponential in  $d$  which is taken to be a constant.

Chubarian *et al.* [11] expanded on the results of Sidiropoulos *et al.* by giving the first bicriteria approximation for minimum outlier sets with distortion. In particular, they showed that given an unweighted graph metric and tuple  $(k, c)$ , there is a polynomial time algorithm that either correctly decides that there does not exist an embedding of the metric into the real line with at most  $k$  outliers and at most  $c$  distortion, or outputs an  $(O(c^6 k \log^{5/2} n), O(c^{13}))$ -outlier embedding into the real line.

**Embeddings with slack.** A different notion of distortion that is robust to noise in the data was introduced by Abraham *et al.* [1]. In this *embeddings with slack* model, a budget of slack is applied to pairs of vertices in the metric space (as opposed to individual vertices, as in our model). An embedding of a metric space  $(X, d_X)$  into another space  $(Y, d_Y)$  has distortion  $c$  with  $\epsilon$ -slack if all but an  $\epsilon$  fraction of the distances are distorted by at most  $c$ . Abraham *et al.* [1] showed that there exists a polynomial time algorithm that finds an  $O(\log \frac{1}{\epsilon})^{1/p}$ -distortion embedding with  $\epsilon$  slack for embeddings into  $\ell_p$  for  $p \geq 1, \epsilon > 0$ . Chan *et al.* [8] showed that there is a polynomial time algorithm for embedding a metric  $(V, d)$  of  $n$  points into a spanner graph of at most  $O(n)$  edges with  $\epsilon$ -slack and  $O(\log \frac{1}{\epsilon})$  distortion. Lammersen *et al.* [21] extended results in this topic to the streaming setting by giving an algorithm using poly-logarithmic space that computes embeddings with slack into finite metrics. From an algorithmic viewpoint, defining outliers in terms of edges versus nodes leads to very different optimization problems. Furthermore, an important difference between our work and these previous works on embeddings with slack is that we are interested in instance-specific approximations, whereas these latter works aim to find uniform bounds on distortion with slack that hold for all input metric spaces.

## 2 Definitions and main results

We begin by defining terms used in this paper and discussing our main results.

### Outlier embeddings and distortion

**Definition 2.1.** A metric space is a pair  $(X, d_X)$  such that  $X$  is a set of elements we call points or nodes and  $d_X : X \times X \rightarrow \mathbb{R}_{\geq 0}$  is a function that has the following properties:

1. For all  $x, y \in X$ ,  $d_X(x, y) = 0$  if and only if  $x = y$
2. For all  $x, y \in X$ ,  $d_X(x, y) = d_X(y, x)$
3. For all  $x, y, z \in X$ ,  $d_X(x, z) \leq d_X(x, y) + d_X(y, z)$

In this paper we will focus on expanding embeddings from a given finite metric into  $\ell_p$ .

**Definition 2.2.** An expanding embedding  $\alpha : X \rightarrow Y$  of a metric space  $(X, d_X)$  into another metric space  $(Y, d_Y)$  has distortion  $c \geq 1$  if for all  $u, v \in X$ :

$$d_X(u, v) \leq d_Y(\alpha(u), \alpha(v)) \leq c \cdot d_X(u, v)$$

Following [11] and [29], we consider so-called *outlier embeddings* that embed all but a small set of outliers from the given metric into the host space.

**Definition 2.3.** An embedding  $\alpha : X \rightarrow Y$  of a metric space  $(X, d_X)$  into another metric space  $(Y, d_Y)$  is a  $(k, c)$ -outlier embedding if there exists  $K \subseteq X$  such that  $|K| \leq k$  and  $\alpha|_{X \setminus K}$  (the restriction of  $\alpha$  to the domain  $X \setminus K$ ) is an embedding of  $(X \setminus K, d|_{X \setminus K})$  with distortion at most  $c$ .

## Nested compositions

A main component of our approach is to compose two different embeddings that map subsets of  $X$  into  $Y$ , where the composition preserves the distortions on the respective subsets to within constant factors. In particular, we assume that the entire set  $X$  can be embed into  $Y$  with some distortion  $c$ , but that we also have access to a much better embedding of a subset of  $X$ , call it  $S$ , into  $Y$  that has distortion  $c' < c$ . We want to combine these two embeddings into a single mapping of  $X$  into  $Y$  that has overall distortion at most  $g(c)$  for some small-growing function  $g$ , but that preserves the smaller distortion  $c'$  over pairs of points in  $S$ . We call such a construction a *composition of nested embeddings*.

**Definition 2.4** (Composition of nested embeddings). Let  $(X, d_X)$  and  $(Y, d_Y)$  be two metric spaces and  $g : [0, \infty) \rightarrow [1, \infty)$ . A  $g$ -nested composition is an algorithm that, given a set  $S \subseteq X$  with  $k := |X \setminus S|$ , and two expanding embeddings,  $\alpha_S : S \rightarrow Y$  with distortion  $c_S$  and  $\alpha_X : X \rightarrow Y$  with distortion  $c_X \geq c_S$ , returns an embedding  $\alpha : X \rightarrow Y$  such that,

$$\text{for all } u, v \in S, \quad d_X(u, v) \leq d_Y(\alpha(u), \alpha(v)) \leq c_S \cdot d_X(u, v), \quad (1)$$

$$\text{and, for all } u, v \in X, \quad d_X(u, v) \leq d_Y(\alpha(u), \alpha(v)) \leq g(k)c_X \cdot d_X(u, v). \quad (2)$$

We say that the nested composition is weak if the embedding  $\alpha$  restricted to  $X \setminus S$  is not an expanding embedding. That is, the inequality  $d_X(u, v) \leq d_Y(\alpha(u), \alpha(v))$  does not hold for all  $u, v \in X$ .

For nested compositions where the host space is the  $\ell_p$  metric, we will consider randomized embeddings. For a randomized embedding  $\alpha$  into  $\ell_p$  space, the distortion is measured relative to the expectation of the  $p$ th power of the  $\ell_p$  norm. In particular, we will require the following version of the constraint (2) above:

$$\mathbb{E} [ \|\alpha(x) - \alpha(y)\|_p^p ]^{1/p} \leq g(k)c_X \cdot d_X(u, v).$$

With this definition, it is easy to see that the existence of a randomized (weak) nested embedding into  $\ell_p$  implies the existence of a deterministic (weak) nested embedding into  $\ell_p$  with the same parameters. In particular, a scaled concatenation of the embeddings in the support of the claimed distribution provides such an embedding.

## Main results

**Nested compositions.** Our main technical result shows that we can efficiently construct randomized  $O(\log k)$ -weak nested compositions when the host space  $Y$  is an  $\ell_p$  metric. For  $p = 2$ , the case of interest for us, we can even efficiently construct a *deterministic*  $O(\log k)$ -weak nested composition into  $\ell_2$  space by solving an appropriate semi-definite program. Finally, for  $p = 1$ , our construction ensures expansion and satisfies both inequalities in constraint (2). We leave open the question of constructing (strong) nested compositions satisfying expansion for  $p > 1$ .

**Theorem 2.5.** *Let  $(X, d_X)$  be any finite metric and  $p \geq 1$ . There exists a  $382H_k$ -weak nested composition from  $X$  into the  $\ell_p$  metric, where  $H_k$  is the  $k$ th Harmonic number. Further, for  $p = 1$ , there exists a  $382H_k$ -nested composition from  $X$  into  $\ell_1$ .*

For metric spaces  $(X, d_X)$  where the distortion of the embedding into  $\ell_p$  depends on the size of the subset being embedded, we can in fact obtain a stronger guarantee – the distortion of the composition depends only on the size of the outlier set  $k = |X \setminus S|$  and not on the size of the entire space  $X$ . In particular, we can replace the quantity  $c_X$  in constraint (2) by the worst case distortion from embedding any subset of size  $k + 1$  into the host metric.

**Theorem 2.6.** *Let  $(X, d_X)$  be any finite metric and  $p \geq 1$ . Suppose that for  $k \in \mathbb{Z}^+$  every subset of  $X$  of size  $k$  can be embedded into  $\ell_p$  with distortion  $\zeta_k$ . Then, there exists a weak nested composition that given a subset  $S \subseteq X$  with  $|X \setminus S| \leq k$ , and an expanding embedding  $\alpha_S$  from  $S$  into  $\ell_p$  with distortion  $c_S$ , returns an embedding  $\alpha : X \rightarrow \ell_p$  such that,*

$$\begin{aligned} \text{for all } u, v \in S, & \quad d_X(u, v) \leq \|\alpha(x) - \alpha(y)\|_p \leq c_S \cdot d_X(u, v), \\ \text{and, for all } u, v \in X, & \quad \|\alpha(x) - \alpha(y)\|_p \leq 382H_{k+1}\zeta_k \cdot d_X(u, v). \end{aligned}$$

**Outlier embeddings.** With these results in hand, we obtain the following bicriteria approximations for outlier embeddings from finite metrics into  $\ell_2$ .

**Theorem 2.7.** *Let  $(X, d_X)$  be a metric space that admits a  $(0, \zeta)$ -outlier embedding as well as a  $(k, c)$ -outlier embedding into  $\ell_2$ . Then, given a (randomized)  $g$ -weak nested composition from  $X$  into  $\ell_2$ , there exists a polynomial time algorithm  $\mathcal{A}$  that, for any  $\gamma > 1$ , finds a subset  $K \subseteq X$  and an embedding  $\alpha : X \setminus K \rightarrow \ell_2$  such that  $\alpha$  has distortion at most  $\gamma c$ , and*

$$|K| \leq 2 \frac{g^2(k)\zeta^2/c^2 + \gamma^2}{\gamma^2 - 1} k$$

Choosing  $\gamma = 1 + \epsilon$  for  $\epsilon \in (0, 1]$ , in particular, provides an  $\left(O\left(\frac{g^2(k)\zeta^2}{\epsilon^2}k\right), (1 + \epsilon)c\right)$ -outlier embedding from  $X$  into  $\ell_2$ .

It is worth observing that the above theorem only requires the *existence* of a *weak* nested composition into  $\ell_2$ . In particular, the nested composition does not have to be efficiently computable. The outlier embedding algorithm simply uses the parameters of such a nested embedding in formulaating a semi-definite program with a certain value, as we discuss in Section 3.

Putting Theorems 2.5 and 2.7 together implies that we can find an  $(O(k \log^2 k \log^2 n/c^2), O(c))$ -outlier embedding of  $X$  into  $\ell_2$  space, where we have applied Bourgain’s result that all metrics have an  $O(\log n)$  embedding into  $\ell_2$  space [7]. Using Theorem 2.6 instead of Theorem 2.5, and using Bourgain’s embedding on subsets of size  $k$ , gives us a stronger approximation guarantee:

**Corollary 2.8.** *Let  $(X, d_X)$  be a metric space that admits a  $(k, c)$ -outlier embedding into  $\ell_2$ . Then there exists a polynomial time algorithm  $\mathcal{A}$  that, given any  $\epsilon > 0$ , finds a subset  $K \subseteq X$  and an embedding  $\alpha : X \setminus K \rightarrow \ell_2$  such that  $\alpha$  has distortion at most  $(1 + \epsilon)c$ , and*

$$|K| \leq O\left(\frac{\log^4 k}{\epsilon c^2}\right) \cdot k.$$

**Hardness of approximation.** Finally, we provide a strengthening of Sidiropoulos *et al.* [29]’s hardness result for outlier embeddings, showing that it is NP-hard to determine the size of the smallest outlier set such that the remaining metric is isometrically embeddable into  $\ell_2$  even when the dimension of the embedding is unrestricted. As with Sidiropoulos *et al.*’s results, under the unique games conjecture, it is also hard to obtain a  $2 - \epsilon$  approximation for the minimum outlier set size, for any  $\epsilon > 0$ . Furthermore, our construction achieves two other properties that [29]’s doesn’t: (1) Our hardness results apply also to shortest path metrics over unweighted undirected graphs. (2) We show that the hardness result holds for embedding into the  $\ell_p$  metric for any  $p \geq 1$ .

**Theorem 2.9.** *Let  $(X, d)$  be the distance metric for an unweighted undirected graph  $G = (V, E)$ . then, given  $(X, d, k)$  it is NP-hard to decide if there exists a subset  $K \subseteq X$  with  $|K| = k$  such that  $(X \setminus K, d|_{X \setminus K})$  is isometrically embeddable into  $\ell_p$  for any finite integer  $p \geq 1$ .*

*Under the unique games conjecture, it is NP-hard to find a  $2 - \epsilon$  approximation for the minimum such  $k$ , for any  $\epsilon > 0$ .*

### 3 SDP relaxation and approximation

In this section we will prove Theorem 2.7. We begin with a semi-definite programming formulation for constructing an outlier embedding into  $\ell_2$ . In the absence of outliers, the optimal embedding of any finite metric into  $\ell_2$  can be found using an SDP. In particular, for a given such metric  $(X, d)$ , let  $\vec{v}_x$  for  $x \in X$  denote the mapping of  $x$  into  $\ell_2$ . Then, the constraint  $d^2(x, y) \leq \|\vec{v}_x - \vec{v}_y\|^2 \leq c^2 \cdot d^2(x, y)$  ensures that the distance between points  $x$  and  $y$  is distorted by a factor of at most  $c$ . The challenge is to incorporate outliers into this formulation.

Consider the finite metric space  $(X, d)$ , and suppose that there exists a  $(k, c)$ -outlier embedding from  $(X, d)$  into Euclidean space for some integer  $k > 0$  and real number  $c \geq 1$ . We use the vector  $\vec{v}_x$  for  $x \in X$  to denote the mapping of  $x$  into  $\ell_2$ , and  $\delta_x \in [0, 1]$  as an indicator for whether  $x$  is an outlier. We then construct the following SDP:

$$\begin{aligned} \min_{\delta, \vec{v}} \quad & \sum_{x \in X} \delta_x && \text{(Outlier SDP)} \\ \text{s.t.} \quad & \forall x, y \in X : && (1 - \delta_x - \delta_y) \cdot d^2(x, y) \leq \|\vec{v}_x - \vec{v}_y\|^2 \leq (c^2 + (\delta_x + \delta_y)f(k)) \cdot d^2(x, y), \\ & \forall x \in X : && \delta_x \in [0, 1]. \end{aligned} \quad (3)$$

Here  $f(k)$  is a function to be determined. We claim that for an appropriate choice of  $f$ , this SDP is a relaxation for the problem of minimizing the outlier set size such that all non-outlier elements in  $X$  can be embed into  $\ell_2$  with distortion  $c$ . In particular, given a  $(k, c)$ -outlier embedding from  $(X, d)$  into  $\ell_2$ , we can find a feasible solution for the SDP with value at most  $k$ . It will be sufficient to set  $f(k) := (g(k)\zeta)^2$  where  $\zeta$  is the distortion of an outlier-free embedding of  $X$  into  $\ell_2$  and  $g$  is the function corresponding to the weak nested composition guaranteed in the statement of Theorem 2.7.

**Lemma 3.1.** *Let  $(X, d)$  be a finite metric space that admits a  $(0, \zeta)$ -outlier embedding as well as a  $(k, c)$ -outlier embedding into  $\ell_2$  for  $c \leq \zeta$ . If there exists a  $g$ -weak nested composition from  $(X, d)$  into  $\ell_2$ , then (Outlier SDP) with  $f(k) := (g(k)\zeta)^2$  has a feasible solution with value equal to  $k$ .*

*Proof.* Let  $K \subseteq X$  be a subset of size at most  $k$  such that  $X \setminus K$  is embeddable into  $\ell_2$  with distortion at most  $c$ . By definition of  $k$ , such a set must exist. Let  $\alpha_S : X \setminus K \rightarrow \ell_2$  be an expanding embedding of  $X \setminus K$  into  $\ell_2$  with distortion at most  $c$ ; and  $\alpha_X : X \rightarrow \ell_2$  be an expanding embedding with distortion at most  $\zeta$ . Then, by Definition 2.4, there exists an embedding  $\alpha : X \rightarrow \ell_2$  such that for all  $x, y \notin K$ , we have  $d(x, y) \leq \|\alpha(x) - \alpha(y)\|_2 \leq c \cdot d(x, y)$  and for all  $x, y \in X$ , we have  $\|\alpha(x) - \alpha(y)\|_2 \leq g(k)\zeta \cdot d(x, y)$ .

We will now construct a feasible solution for (Outlier SDP). Set  $\delta_x$  to 1 if  $x \in K$  and 0 otherwise. Set  $\vec{v}_x$  to  $\alpha(x)$ . Clearly  $\sum_{x \in X} \delta_x = k$ . We show that this setting of the variables satisfies the given constraints.

Consider a constraint corresponding to  $x, y \in X \setminus K$ . Then we have that  $(1 - \delta_x - \delta_y) \cdot d(x, y)^2 = d(x, y)^2 \leq \|\alpha(x) - \alpha(y)\|_2^2 \leq c^2 \cdot d(x, y)^2 = (c^2 + (\delta_x + \delta_y)f(k)) \cdot d(x, y)^2$  by the facts we have already asserted about  $\alpha$ .

Next, consider a constraint corresponding to  $x \in X, y \in K$ . In this case,  $\delta_y = 1$  and  $(1 - \delta_x - \delta_y) \leq 0$ , so the first inequality in (3) is satisfied. On the other hand, by the definition of  $f$  and  $\alpha$ ,  $\|\alpha(x) - \alpha(y)\|_2^2 \leq f(k) \cdot (d(x, y))^2 \leq (c^2 + (\delta_x + \delta_y)f(k)) \cdot (d(x, y))^2$ , which gives us the second inequality. Thus, the solution  $(\delta, \vec{v})$  is a feasible solution with value  $k$ .  $\square$

Observe that  $f$  is a function of  $k$ , and so in order to set up and solve the SDP, we require knowing the value of the parameter  $k$ . We can get around this by setting  $k = 1, 2, \dots$ , and so on until we find the smallest value of  $k$  for which the SDP with parameter  $f(k)$  has a feasible solution of value at most  $k$ . Rounding this solution then gives the desired theorem.

We are now ready to prove Theorem 2.7.

*Proof of Theorem 2.7.* Suppose that  $(X, d)$  admits a distortion  $\zeta$  embedding into  $\ell_2$  as well as a  $(k, c)$ -outlier embedding into  $\ell_2$ . By Lemma 3.1, there exists a solution to the SDP Outlier SDP with value at most  $k$ , which we can find efficiently by solving the SDP. Let  $\{(v_x, \delta_x)\}_{x \in X}$  denote such a solution. Let  $\Delta$  be a parameter to be defined. Define  $\alpha(x) \mapsto \frac{1}{\sqrt{1-2\Delta}}v_x$ , and  $K \mapsto \{x : \delta_x \geq \Delta\}$ . We claim that for an appropriate choice of  $\Delta$ , the solution  $(K, \alpha)$  satisfies the requirements of the theorem.

In particular, we note that  $|K| \leq (\sum_{x \in X} \delta_x) / \Delta \leq k / \Delta$ . To bound the distortion of  $\alpha$  restricted to  $X \setminus K$  by  $\gamma c$ , let us consider some pair of points  $x, y \in X \setminus K$ , and recall that we have  $\delta_x, \delta_y < \Delta$ . Then, substituting  $v_x = \sqrt{1-2\Delta}\alpha(x)$  in (3) gives us:

$$(1 - 2\Delta) \cdot d^2(x, y) \leq (1 - 2\Delta)\|\alpha(x) - \alpha(y)\|_2^2 \leq (c^2 + 2\Delta f(k)) \cdot d(x, y)^2$$

The first inequality implies expansion. The second provides an upper bound on the distortion of:

$$\frac{c^2 + 2\Delta f(k)}{1 - 2\Delta}$$

Setting this quantity equal to  $\gamma^2 c^2$  and solving for  $\Delta$  gives us  $\Delta = \frac{c^2(\gamma^2 - 1)}{2f(k) + 2c^2\gamma^2}$  and  $|K| \leq \frac{2f(k) + 2c^2\gamma^2}{c^2(\gamma^2 - 1)}k$ .  $\square$

Corollary 2.8 follows similarly from Theorem 2.6 by setting the function  $f$  in the SDP constraint (3) to  $f(k) := (382H_{k+1} \cdot \zeta_k)^2$  — the square of the upper bound on distortion of the nested composition in the statement of Theorem 2.6.

## 4 Composition of nested embeddings

In this section, we will present an algorithm for composing nested embeddings into  $\ell_p$  spaces, thereby proving Theorems 2.5 and 2.6. Our algorithm, Algorithm 1, is formally specified below.

Let  $c_X$  and  $c_S$  denote the distortion of  $\alpha_S$  and  $\alpha_X$  respectively. We now show that if  $\alpha$  is the output of Algorithm 1 on input  $((X, d), S, p, \alpha_S, \alpha_X, \tau)$  with  $\tau = 2$ , the distortion between elements in  $S$  is at most  $c_S$  and all other distortion is at most  $(\frac{155}{2} \cdot H_k \cdot c_S + (\frac{225}{2} \cdot H_k + 1) \cdot c_X)$  in expectation. Our argument is broken into two parts: Lemma 4.1 shows that the embedding  $\alpha$  has low contraction; Lemma 4.2 bounds the amount by which every distance expands.

We state the lemmas first and then prove them in the following subsections. Throughout these arguments we assume that  $\tau = 2$ , although it is possible to obtain slightly better distortion bounds by choosing a value for  $\tau$  carefully. We present general versions of the lemmas, exhibiting the dependence of the bounds on  $\tau$  in Appendix A. Theorem 2.5 follows in a straightforward manner from these lemmas; Section 4.3 contains a proof of this theorem as well as a proof of the strengthened composition result — Theorem 2.6.



---

**Algorithm 1** Algorithm for finding a nested embedding

---

**Input:** Metric space  $(X, d)$ , subset  $S \subseteq X$ , finite integer  $p \geq 1$ , expanding embedding  $\alpha_S : X - S \rightarrow \ell_p$ , expanding embedding  $\alpha_X : X \rightarrow \ell_p$ , and real number  $\tau$

**Output:** A randomized expanding embedding  $\alpha : X \rightarrow \ell_p$  such that for all  $x, y \in S$ ,  $\|\alpha(x) - \alpha(y)\|_2 \leq f(c_S) \cdot d(x, y)$  and for all  $x, y \in X$ ,  $E[\|\alpha(x) - \alpha(y)\|_2] \leq g(c_S, c_X) \cdot d(x, y)$ .

- 1:  $K \leftarrow X \setminus S$ .
- 2: Define a function  $\gamma : K \rightarrow S$  such that  $\gamma(u) \in \arg \min_{v \in S} d(u, v)$ .  $\triangleright$  ie  $\gamma(u)$  is one of  $u$ 's closest neighbors in  $S$
- 3: Select  $b$  uniformly at random from the range  $[2, \tau + 2]$
- 4: Select a uniformly random permutation  $\pi : X \setminus S \rightarrow [k]$  of the vertices in  $X \setminus S$
- 5:  $K' \leftarrow K, i \leftarrow 1$
- 6: **while**  $K' \neq \emptyset$  **do**
- 7:      $u_i \leftarrow \pi^{-1}(i)$
- 8:      $K_i \leftarrow \{v \in K' \mid d(v, u_i) \leq b \cdot d(v, \gamma(v))\}$   $\triangleright$  Call  $u_i$  the “center” of  $K_i$
- 9:     Define an embedding  $\alpha_i : X \rightarrow \ell_p$  such that

$$\alpha_i(v) = \begin{cases} \alpha_X(v) & v \in K_i \\ \alpha_X(\gamma(u_i)) & v \notin K_i \end{cases}$$

- 10:      $K' \leftarrow K' \setminus K_i, t \leftarrow i, i \leftarrow i + 1$
- 11: Define an embedding  $\alpha' : X \rightarrow \ell_p$  such that

$$\alpha'(v) = \begin{cases} \alpha_S(v) & \text{if } v \in S \\ \alpha_S(\gamma(u_i)) & \text{if } v \in K_i \text{ and with } u_i \text{ being the center of } K_i \end{cases}$$

- 12: Define an embedding  $\alpha : X \rightarrow \ell_p$  such that  $\alpha(v) \mapsto (\alpha'(v) | \alpha_1(v) | \dots | \alpha_t(v))$   $\triangleright$  here  $|$  denotes concatenation
  - 13: **Output**  $\alpha$
-

**Lemma 4.1.** Let  $\alpha \leftarrow \text{Algorithm 1}((X, d), S, p, \alpha_S, \alpha_X, \tau)$  with  $\tau = 2$ . Then for all  $x, y \in X$ , we have  $3^{-1+\frac{1}{p}} \cdot \|\alpha(x) - \alpha(y)\|_p \geq d(x, y)$ . Furthermore, for  $x, y \in S$ , we have  $\|\alpha(x) - \alpha(y)\|_p \geq d(x, y)$ .

**Lemma 4.2.** Let  $\alpha \leftarrow \text{Algorithm 1}((X, d), S, p, \alpha_S, \alpha_X, \tau)$  with  $\tau = 2$ . Then we have the following bounds on the expansion for each pair  $x, y \in X$ :

- (a) If  $x, y \in S$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq c_S \cdot d(x, y)$ .
- (b) If  $x, y \in K_i$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq c_X \cdot d(x, y)$ .
- (c) If  $x \in S, y \in X \setminus S$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq [7 \cdot c_S + 9 \cdot c_X] \cdot d(x, y)$ .
- (d) If  $x, y \in X \setminus S$  and  $d(x, \gamma(x)) \leq 2 \cdot d(x, y)$  for  $\gamma$  as defined in line 2 of the algorithm, then  $\|\alpha(x) - \alpha(y)\|_p \leq [31 \cdot c_S + 45 \cdot c_X] \cdot d(x, y)$ .
- (e) If  $x, y \in X \setminus S$  and  $d(x, \gamma(x)), d(y, \gamma(y)) > 2 \cdot d(x, y)$  for  $\gamma$  as defined in line 2 of the algorithm, then  $E_\alpha[\|\alpha(x) - \alpha(y)\|_p] \leq (\frac{155}{2} \cdot H_k \cdot c_S + (\frac{225}{2} \cdot H_k + 1) \cdot c_X) \cdot d(x, y)$ .

## 4.1 Proof of contraction bounds

*Proof of Lemma 4.1.* Let  $\gamma$  be as defined in line 2 of the algorithm and let the  $u_i$  and  $K_i$  be as defined in lines 6 and 7 of the algorithm. We divide into cases.

1. If  $x, y \in S$ , then  $\|\alpha(x) - \alpha(y)\|_p = \|\alpha'(x) - \alpha'(y)\|_p \geq d(x, y)$  because  $\alpha_S$  is expanding
2. If  $x, y \in K_i$  for some  $K_i$  defined in line 7 of the algorithm or if  $x \in K_i, y = \gamma(u_i)$ , their embeddings differ only on the coordinates associated with  $\alpha_i$ , so we have

$$\begin{aligned} \|\alpha(x) - \alpha(y)\|_p &= \|\alpha_i(x) - \alpha_i(y)\|_p \\ &= \|\alpha_X(x) - \alpha_X(y)\|_p \\ &\geq d(x, y), \end{aligned}$$

where the second line is by definition of  $\alpha_i$  and the last line is because  $\alpha_X$  is expanding.

3. If  $x \in S, y \notin S$ , then let  $y \in K_i$  for some  $i$  as defined in line 7 of the algorithm and let  $u_i$  be the center for this  $i$  defined in line 6. This implies that  $\alpha(x)$  and  $\alpha(y)$  differ only on coordinates associated with  $\alpha'$  and  $\alpha_i$ .

$$\begin{aligned} \|\alpha(x) - \alpha(y)\|_p &= (\|\alpha'(x) - \alpha'(y)\|_p^p + \|\alpha_i(x) - \alpha_i(y)\|_p^p)^{1/p} \\ &\geq (\|\alpha_S(x) - \alpha_S(\gamma(u_i))\|_p^p + \|\alpha_X(\gamma(u_i)) - \alpha_X(y)\|_p^p)^{1/p} \\ &\geq (d(x, \gamma(u_i))^p + d(\gamma(u_i), y)^p)^{1/p} \\ &\geq \frac{2^{1/p}}{2} (d(x, \gamma(u_i)) + d(\gamma(u_i), y)) \\ &\geq \frac{1}{2^{1-1/p}} d(x, y), \end{aligned}$$

where the second line is by definition of  $\alpha'$  and  $\alpha_i$ , the third is by cases 1 and 2 of this lemma, the fourth is by the power mean inequality, and the fifth is by the triangle inequality.

4. If  $x \in K_i, y \in K_j$  for some  $K_i, K_j$  defined in line 7 of the algorithm. Let  $u_i, u_j$  be the centers of these sets as defined in line 6 of the algorithm. Note that  $\alpha(x)$  and  $\alpha(y)$  differ only on the indices associated with  $\alpha', \alpha_i$ , and

membership of $x$ and $y$	restrictions on $d(x, y)$	upper bound on expected distortion
$x, y \in S$	none	$c_S$ (a)
$x, y \in K_i$	none	$c_X$ (b)
$x \in S, y \in X - S$	none	$[7 \cdot c_S + 9 \cdot c_X]$ (c)
$x, y \in X - S$	$d(x, \gamma(x)) \leq 2 \cdot d(x, y)$	$[31 \cdot c_S + 45 \cdot c_X]$ (d)
$x, y \in X - S$	$d(x, \gamma(x)), d(y, \gamma(y)) > 2 \cdot d(x, y)$	$(\frac{155}{2} \cdot H_k \cdot c_S + (\frac{225}{2} \cdot H_k + 1) \cdot c_X)$ (e)

Table 1: Summary of the bounds in Lemma 4.2. Let  $\alpha \leftarrow \text{Algorithm 1}((X, d), S, p, \alpha_S, \alpha_X, \tau)$  where  $\alpha_S : S \rightarrow \ell_p$  is an expanding embedding of distortion at most  $c_S$  and  $\alpha_X : X \rightarrow \ell_p$  is an expanding embedding of distortion at most  $c_X$ . Then the third column of the table gives an upper bound on the the expected value of  $\|\alpha(x) - \alpha(y)\|_p$  where  $x$  and  $y$  meet the criteria of the first two columns. Here  $\gamma$  and  $K_i$  are as defined in lines 2 and 6-7 of the algorithm.

$\alpha_j$ . Thus we get

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p^p &= (\|\alpha'(x) - \alpha'(y)\|_p^p + \|\alpha_i(x) - \alpha_i(y)\|_p^p + \|\alpha_j(x) - \alpha_j(y)\|_p^p)^{1/p} \\
&\geq (\|\alpha_S(\gamma(u_i)) - \alpha_S(\gamma(u_j))\|_p^p + \|\alpha_X(\gamma(u_i)) - \alpha_X(x)\|_p^p + \|\alpha_X(\gamma(u_j)) - \alpha_X(y)\|_p^p)^{1/p} \\
&\geq (d(\gamma(u_i), \gamma(u_j))^p + d(\gamma(u_i), x)^p + d(\gamma(u_j), y)^p)^{1/p} \\
&\geq \frac{3^{1/p}}{3} (d(\gamma(u_i), \gamma(u_j)) + d(\gamma(u_i), x) + d(\gamma(u_j), y)) \\
&\geq 3^{-1+\frac{1}{p}} \cdot d(x, y),
\end{aligned}$$

where the second line is by definition of the three embeddings, the third line is by cases 1 and 2 of this lemma, the fourth is by the power mean inequality, and the fifth is by the triangle inequality.  $\square$

## 4.2 Proofs of expansion bounds

The bounds in Lemma 4.2 are summarized in Table 1. We will prove each statement separately.

*Proof of Lemma 4.2 (a).*  $u, v \notin K_i$  for all  $i$ , so  $\alpha_i(u) = \alpha_i(v)$  for all  $i$ . Thus,

$$\begin{aligned}
\|\alpha(u) - \alpha(v)\|_p &= \|\alpha'(u) - \alpha'(v)\|_p \\
&= \|\alpha_S(u) - \alpha_S(v)\|_p \\
&\leq c_S \cdot d(u, v),
\end{aligned}$$

where the last equality is by definition of  $\alpha_S$ .  $\square$

*Proof of Lemma 4.2 (b).* Since  $x, y \in K_i$  for a fixed  $i$ , we have that  $\alpha(x), \alpha(y)$  differ only on coordinates associated with  $\alpha_i$ . Thus, we get

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p &= \|\alpha_i(x) - \alpha_i(y)\|_p \\
&= \|\alpha_X(x) - \alpha_X(y)\|_p \\
&\leq c_X \cdot d(x, y),
\end{aligned}$$

where we have used the fact that  $\alpha_X(u) = \alpha_i(u)$  for  $x \in K_i$ .  $\square$

Now we consider the distortion between outliers and non-outliers. To prove these bounds, we will need to define extra points in the  $\ell_p$  space that  $X$  is mapped to. Consider the points  $u_i$  defined in Step (7) of the algorithm. We will now define a point  $\beta(u_i)$  for each such  $u_i \in K$ . If  $u_i \in K_i$  we set  $\beta(u_i) = \alpha(u_i)$ . Otherwise if  $u_i \in K_j$  for  $j < i$ , we set all of the coordinates of  $\beta(u_i)$  to be the same as  $\alpha(u_i)$ , except for the coordinates associated with  $\alpha_j(u_i)$ , that




Figure 1: Left: Visualization of nodes referenced in Lemma 4.2 (a); Right: Visualization of nodes referenced in Lemma 4.2 (b)

are replaced with those of  $\alpha_X(\gamma(u_j))$ ; and the coordinates associated with  $\alpha_i(u_i)$ , that are replaced with those for  $\alpha_X(u_i)$ . In other words,  $\beta(u_i)$  is the point that  $u_i$  would be mapped to by  $\alpha$  if it had so happened that  $u_i$  belonged to  $K_i$ .

Now we are ready to prove Lemma 4.2 (c). We break this proof up into two parts: Lemmas 4.3 and 4.4.

**Lemma 4.3.** *Let  $\alpha \leftarrow \text{Algorithm 1}((X, d), S, p, \alpha_S, \alpha_X, \tau)$  with  $\tau = 2$ . Consider  $x \in S$ ;  $u_i$  as defined in Step (7) of the algorithm for some  $i$ ; and  $\beta(u_i)$  as defined above. If  $x = \gamma(u_i)$  where  $\gamma : X \setminus S \rightarrow S$  is as defined in Step (2) of the algorithm, then  $\|\alpha(x) - \beta(u_i)\|_p \leq c_X \cdot d(x, u_i)$ . For all other  $x$ ,  $\|\alpha(x) - \beta(u_i)\|_p \leq (c_S + c_X) \cdot d(x, u_i)$ .*

*Proof.* First, note that the point  $\beta(u_i)$  is exactly the point that would have been assigned to  $\alpha(u_i)$  if it was not placed in some  $K_j$  for  $j < i$ . Additionally, this means that  $\beta(u_i)$  and  $\alpha(x)$  are the same on the indices associated with  $\alpha_j$  for all  $j \neq i$  (namely they are assigned  $\alpha_X(\gamma(u_j))$ ).

Now we have  $\alpha_j(x) = \alpha_j(u_i)$  for all  $j \neq i$ . Thus, we have

$$\begin{aligned} \|\alpha(x) - \beta(u_i)\|_p &= \|\alpha'(x)|\alpha_i(x) - \alpha'(u_i)|\alpha_X(u_i)\|_p \\ &\leq \|\alpha'(x) - \alpha'(u_i)\|_p + \|\alpha_i(x) - \alpha_X(u_i)\|_p. \end{aligned}$$

We have two cases to consider.

1. If  $x = \gamma(u_i)$ , then we have  $\alpha'(u_i) = \alpha'(x)$  by definition, so we have

$$\begin{aligned} \|\alpha(x) - \beta(u_i)\|_p &\leq \|\alpha_i(x) - \alpha_X(u_i)\|_p \\ &= \|\alpha_X(x) - \alpha_X(u_i)\|_p \\ &\leq c_X \cdot d(x, u_i), \end{aligned}$$

where the last line is by the fact that  $\alpha_X$  is an embedding with at most  $c_X$  distortion.

2. If  $x \neq \gamma(u_i)$ , we have  $d(\gamma(u_i), u_i) \leq d(x, u_i)$  by definition of  $\gamma(u_i)$ . Then we use the triangle inequality as follows.

$$\begin{aligned} \|\alpha(x) - \beta(u_i)\|_p &\leq \|\alpha(x) - \alpha(\gamma(u_i))\|_p + \|\alpha(\gamma(u_i)) - \beta(u_i)\|_p \\ &\leq c_S \cdot d(x, u_i) + c_X \cdot d(\gamma(u_i), u_i) \\ &\leq c_S \cdot d(x, u_i) + c_X \cdot d(x, u_i) \\ &= (c_S + c_X)d(x, u_i) \end{aligned}$$

where the third inequality comes from Lemma 4.2 (a) and the first case of this lemma.




Figure 2: Visualization of nodes referenced in Lemmas 4.3 and 4.4 when  $u_i \in K_i$ .

□

Next we want to consider the distortion between more general members of a set  $K_i$  and members of  $X \setminus K_i$ . For the coming proofs, to show that  $x$  and  $y$  are not too distorted, we will generally focus on showing that the distance between  $x$  and  $y$  is at least a constant factor larger than the distance from  $x$  or  $y$  to some other point whose distance to  $x$  or  $y$  is already known not to be too distorted.

**Lemma 4.4.** *Let  $\alpha \leftarrow \text{Algorithm 1}((X, d), S, p, \alpha_S, \alpha_X, \tau)$  with  $\tau = 2$ . Let  $K_i$  be as defined in Step (8) of the algorithm for some  $i$ . Consider  $x \in S$  and  $y \in K_i$  for some  $i$ . If  $x = \gamma(y)$  where  $\gamma$  is as defined in line 2 of the algorithm, then we have  $\|\alpha(x) - \alpha(y)\|_p \leq [5 \cdot c_S + 9 \cdot c_X] \cdot d(x, y)$ . For all other  $x$ ,  $\|\alpha(x) - \alpha(y)\|_p \leq [7 \cdot c_S + 9 \cdot c_X] \cdot d(x, y)$ .*

*Proof.* Let  $\gamma : X \setminus S \rightarrow S$  be as defined in Step (2) of the algorithm,  $u_i$  be as defined in Step (7) for  $i$  consistent with the lemma statement, and  $\beta(u_i)$  be as defined above. We divide analysis into 2 cases.

1. If  $x = \gamma(y)$ , the lemma essentially comes from the triangle inequality and  $y$ 's relative closeness to  $u_i$  that it must have if it is assigned to  $K_i$ .

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p &= \\
\|\alpha(\gamma(y)) - \alpha(y)\|_p &\leq \|\alpha(x) - \beta(u_i)\|_p + \|\alpha(y) - \beta(u_i)\|_p \\
&\leq [(c_S + c_X) \cdot d(x = \gamma(y), u_i) + c_X d(y, u_i)] \\
&\leq [(c_S + c_X) \cdot [d(\gamma(y), y) + d(y, u_i)] + c_X \cdot d(y, u_i)] \\
&= (c_S + c_X) \cdot d(\gamma(y), y) + (c_S + 2c_X) \cdot d(y, u_i) \\
&\leq (c_S + c_X) \cdot d(\gamma(y), y) + 4(c_S + 2c_X) \cdot d(y, \gamma(y)) \\
&= [5c_S + 9c_X] \cdot d(y, \gamma(y) = x).
\end{aligned}$$

Here the first inequality is by the fact that  $\alpha(x)$  and  $\alpha(y)$  may only differ on the indices associated with  $\alpha'$  and the indices associated with  $\alpha_i$ . The second inequality is by Lemma 4.3 and Lemma 4.2 (b), the third is by the triangle inequality, and the fourth line is by rearranging terms. The fifth line is by the fact that  $d(y, u_i) \leq b \cdot d(y, \gamma(y))$  since  $y$  is in  $K_i$  and  $u_i$  is the center, and the fact that  $b \leq 4$  no matter the result of the random choice in the algorithm.




Figure 3: Left: Visualization of nodes referenced in Lemma 4.2 (d); Right: Visualization of nodes referenced in Lemma 4.2 (e)

2. If  $x \neq \gamma(y)$ , we have  $d(\gamma(y), y) \leq d(x, y)$  by definition of  $\gamma(y)$ . Thus, we get

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p &\leq \|\alpha(x) - \alpha(\gamma(y))\|_p + \|\alpha(\gamma(y)) - \alpha(y)\|_p \\
&\leq [c_S \cdot d(x, \gamma(y)) + [5c_S + 9c_X] \cdot d(\gamma(y), y)] \\
&\leq [c_S \cdot [d(x, y) + d(y, \gamma(y))] + [5c_S + 9c_X] \cdot d(\gamma(y), y)] \\
&\leq [7c_S + 9c_X] \cdot d(x, y),
\end{aligned}$$

where the second inequality is by Lemma 4.2 a and by the previous case of this lemma, the third inequality is by the triangle inequality, and the last inequality is because  $\gamma(y)$  is a closest node in  $S$  to  $y$ . □

Finally, we consider comparing two the distance of two nodes  $x, y \in X \setminus S$ . First we consider the case that at least one of the nodes has a relatively short distance to  $S$  compared to the distance to the other node.

*Proof of Lemma 4.2 (d).* Let  $\gamma$  be as defined in line 2 of the algorithm. We get

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p &\leq \|\alpha(x) - \alpha(\gamma(x))\|_p + \|\alpha(\gamma(x)) - \alpha(y)\|_p \\
&\leq [5c_S + 9c_X] \cdot d(x, \gamma(x)) + [7c_S + 9c_X] \cdot d(\gamma(x), y) \\
&\leq [12c_S + 18c_X] \cdot d(x, \gamma(x)) + [7c_S + 9c_X] \cdot d(x, y) \\
&\leq [31c_S + 45c_X] \cdot d(x, y),
\end{aligned}$$

where the first inequality is by the triangle inequality, the second is by Lemma 4.3, the third is by the triangle inequality applied to  $d(\gamma(x), y)$  and rearranging terms, and the fourth is by the fact that  $d(x, \gamma(x)) \leq 2 \cdot d(x, y)$  and rearranging terms. □

Now we consider the final case, where we must consider expected distance.

*Proof of Lemma 4.2 (e).* Let  $\gamma$  be as defined in line (2) of the algorithm. We will say that  $x$  and  $y$  are “split” if  $x \in K_i, y \in K_j$  for  $i \neq j$  and  $K_i, K_j$  as defined in line (8) of the algorithm. Let  $u_i, u_j$  be as defined in line (7) of the algorithm for the same choice of  $i, j$  respectively.

- First consider the worst-case distortion when  $x$  and  $y$  are *not* split. Then  $x, y \in K_i$  for some  $i$ . We have  $\|\alpha(x) - \alpha(y)\|_p \leq c_X \cdot d(x, y)$  by Lemma 4.2 b, no matter the choice of  $b$ .

- Now consider the worst-case distortion when  $x$  and  $y$  are split such that  $x$  is placed in  $K_i$  and  $y$  is placed in  $K_j$  for  $i < j$ . We have

$$\begin{aligned}
\|\alpha(x) - \alpha(y)\|_p &\leq \|\alpha(x) - \alpha(\gamma(x))\|_p + \|\alpha(\gamma(x)) - \alpha(y)\|_p \\
&\leq [5c_S + 9c_X] \cdot d(x, \gamma(x)) + [7c_S + 9c_X] \cdot d(\gamma(x), y) \\
&\leq [5c_S + 9c_X] \cdot d(x, \gamma(x)) + [7c_S + 9c_X] \cdot [d(x, y) + d(x, \gamma(x))] \\
&\leq [5c_S + 9c_X] \cdot d(x, \gamma(x)) + [7c_S + 9c_X] \cdot \frac{3}{2}d(x, \gamma(x)) \\
&= \left[\frac{31}{2}c_S + \frac{45}{2}c_X\right] \cdot d(x, \gamma(x)),
\end{aligned}$$

where the first inequality is by the triangle inequality, the second is by Lemma 4.4, the third inequality is by the triangle inequality on  $d(\gamma(x), y)$ , the fourth inequality is by the fact that  $d(x, y) < \frac{1}{2}d(x, \gamma(x))$  by the condition of this lemma, and the final equality is by rearranging terms.

Now consider the probability of  $x$  and  $y$  being split. First, let us fix some node  $u$  chosen in some iteration of Step (7) of the algorithm and let  $K_u$  be the cluster formed by this vertex. Suppose that the placement of  $x$  and  $y$  is undetermined prior to this point of time. We will bound the probability that  $x$  and  $y$  are split by  $u$ , that is, exactly one of these vertices ends up in the cluster  $K_u$ . Without loss of generality, assume that  $\frac{d(x, u)}{d(x, \gamma(x))} \leq \frac{d(y, u)}{d(y, \gamma(y))}$ . This implies that if  $x, y$  are split, then  $x \in K_u, y \notin K_u$  and we have

$$\frac{d(x, u)}{d(x, \gamma(x))} \leq b \leq \frac{d(y, u)}{d(y, \gamma(y))}.$$

This implies  $b$  must fall in a range of width

$$\begin{aligned}
W &\leq \frac{d(y, u)}{d(y, \gamma(y))} - \frac{d(x, u)}{d(x, \gamma(x))} \\
&\leq \frac{d(x, u) + d(x, y)}{d(x, \gamma(y)) - d(x, y)} - \frac{d(x, u)}{d(x, \gamma(x))} \\
&\leq \frac{d(x, u) + d(x, y)}{d(x, \gamma(x)) - d(x, y)} - \frac{d(x, u)}{d(x, \gamma(x))} \\
&= d(x, y) \cdot \frac{d(x, \gamma(x)) + d(x, u)}{d(x, \gamma(x)) \cdot (d(x, \gamma(x)) - d(x, y))} \\
&\leq d(x, y) \cdot \frac{5}{(d(x, \gamma(x)) - d(x, y))} \\
&\leq 10 \cdot \frac{d(x, y)}{d(x, \gamma(x))},
\end{aligned}$$

where the second inequality is by applying the triangle inequality twice, the third inequality is by the fact that  $\gamma(x)$  is a closest node in  $S$  to  $x$ , and the fourth is by cross-multiplying. The fifth line is by the fact that  $x \in K_u$ , which implies  $d(x, u) \leq \max$  possible value of  $b \cdot d(x, \gamma(x))$ . Finally, the sixth follows from the fact that  $d(x, y) < \frac{1}{2} \cdot d(x, \gamma(x))$ , a condition of the lemma.

Next we will bound the overall probability that  $x$  and  $y$  are split by *some* node  $u$ . For a vertex  $u$ , define  $\beta_u = \min \left\{ \frac{d(x, u)}{d(x, \gamma(x))}, \frac{d(y, u)}{d(y, \gamma(y))} \right\}$ . This is the smallest value of  $b$  at which the cluster  $K_u$  formed by  $u$  contains either  $x$  or  $y$ . Consider ordering vertices  $u$  in  $K$  in increasing order of  $\beta_u$ , and let  $\text{index} : K \rightarrow [k]$  denote this ordering. We say that a node  $u$  "decides" the pair  $(x, y)$  if at least one of  $x$  and  $y$  is in  $K_u$ .  $u$  decides  $(x, y)$  iff  $b \geq \beta_u$ . This implies that if  $u$  and  $u'$  satisfy  $\text{index}(u) < \text{index}(u')$  and  $u$  appears before  $u'$  in the ordering  $\pi$ , then at the time we consider  $u'$  in Step (7), either it is the case that  $b \geq \beta_u$  and  $(x, y)$  has already been decided by  $u$ , or it is the case that  $b < \beta_u < \beta_{u'}$ ,

in which case  $u'$  cannot decide  $(x, y)$ . Therefore, in either case,  $u'$  does not decide  $(x, y)$ , and consequently does not split them.

In other words, in order for a vertex  $u$  to be able to split  $(x, y)$ , it must be the case that among the  $\text{index}(u)$  vertices before  $u$  in the  $\text{index}$  ordering (and including  $u$  itself),  $u$  is the first vertex to appear in the ordering  $\pi$ . Let us call this latter event  $E_u$ , and observe that this event is independent of the choice of  $b$  — it only depends on the choice of the permutations  $\pi$  and  $\text{index}$ . We also note that  $\Pr_\pi[E_u] = 1/\text{index}(u)$ .

We can now write down the probability that  $x$  and  $y$  are split as follows, using the fact that from our discussion above,  $\Pr_b[u \text{ splits } (x, y) | \neg E_u] = 0$  for all  $u$ .

$$\begin{aligned} \Pr_{\pi, b}[x, y \text{ are split}] &= \sum_{u \in K} \Pr_b[u \text{ splits } (x, y) | E_u] \cdot \Pr_\pi[E_u] + \Pr_b[u \text{ splits } (x, y) | \neg E_u] \cdot \Pr_\pi[\neg E_u] \\ &\leq \sum_{u \in K} \frac{W}{\tau} \cdot \frac{1}{\text{index}(u)} \\ &= \sum_{u \in K} \frac{10}{2} \frac{d(x, y)}{d(x_u, \gamma(x_u))} \cdot \frac{1}{\text{index}(u)}. \end{aligned}$$

Here we used  $x_u$  to denote the node in  $(x, y)$  that is closer to  $u$ , and substitute expressions from above for  $W$ ,  $\tau$ , and  $\Pr_\pi[E_u]$ .

Finally, let us consider the expected distance between  $\alpha(x)$  and  $\alpha(y)$ . Note that by our earlier analysis, the distance between  $x$  and  $y$  when they are split by  $u$  is at most  $[\frac{31}{2}c_S + \frac{45}{2}c_X] \cdot d(x_u, \gamma(x_u))$ . Thus, we can compute the expected distance between  $x$  and  $y$  as follows:

$$\begin{aligned} E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p] &\leq c_X \cdot d(x, y) \cdot \Pr[(x, y) \text{ are not split}] \\ &\quad + \sum_{u \in K} \Pr[x, y \text{ are split by } u | E_u] \cdot \Pr_\pi[E_u] \cdot \left(\frac{31}{2}c_S + \frac{45}{2}c_X\right) \cdot d(x_u, \gamma(x_u)) \\ &\leq c_X \cdot d(x, y) + \sum_{u \in K | \eta(u) \leq \tau+2} \frac{5}{\text{index}(u)} \cdot \frac{d(x, y)}{d(x_u, \gamma(x_u))} \cdot \left(\frac{31}{2}c_S + \frac{45}{2}c_X\right) \cdot d(x_u, \gamma(x_u)) \\ &\leq c_X \cdot d(x, y) + 5 \cdot \left(\frac{31}{2}c_S + \frac{45}{2}c_X\right) \cdot d(x, y) \cdot \sum_{i=1}^k \frac{1}{i} \\ &= \left(\frac{155}{2} \cdot H_k \cdot c_S + \left(\frac{225}{2} \cdot H_k + 1\right) \cdot c_X\right) \cdot d(x, y), \end{aligned}$$

where  $H_k$  is the  $k$ th Harmonic number. □

### 4.3 Nested compositions

In this section we prove Theorems 2.5 and 2.6.

*Proof of Theorem 2.5.* We have shown that if we pick  $b$  at random from the range  $[2, \tau + 2]$  with  $\tau = 2$ , we get for all  $x, y \in X \setminus K$ ,  $\|\alpha(x) - \alpha(y)\|_p = d(x, y)$  and for all  $x, y \in X$ ,  $E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_2] \leq \left(\frac{155}{2} \cdot H_k \cdot c_S + \left(\frac{225}{2} \cdot H_k + 1\right) \cdot c_X\right) \cdot d(x, y)$ . Let  $\chi : [2, \tau + 2] \rightarrow \{f : X \rightarrow \mathbb{R}^t\}$  be a random variable dependent on  $\pi$  and  $b$  whose value is the embedding  $\alpha$  defined by this choice of  $b$ . Note that there are at most  $2^{2^{|X|}}$  values in the support of  $\chi$ , as  $\alpha$  is completely defined by the choice of the  $K_i$  and there are at most  $2^{|X|}$  ways to select a  $K_i$ , with at most  $|X|$  total values of  $i$ . This is finite, so we can concatenate the entire set and obtain an embedding of finite size. Let  $q_\alpha := \Pr_b[\chi = \alpha]$  and let  $\zeta \circ \alpha$  be the



portion of  $\alpha$  that follows  $\alpha'$  (i.e.  $\zeta \circ \alpha(v)$  is  $\alpha_1(v) \cdots \alpha_t(v)$ ) and note that  $\alpha'$  is independent of the choice of  $\pi$  and  $b$ . Consider the following embedding:

$$\alpha^*(x) := \alpha'(x) \parallel_{\alpha} q_{\alpha} \cdot (\zeta \circ \alpha)(x),$$

where  $\parallel_{\alpha}$  refers to the concatenation of the value over all choices of  $\alpha$  and  $q_{\alpha} \cdot (\zeta \circ \alpha)(x)$  is the vector  $(q_{\alpha} \cdot \alpha_1(x), q_{\alpha} \cdot \alpha_2(x), \dots, q_{\alpha} \cdot \alpha_t(x))$  where  $\cdot$  denotes element-wise multiplication.

Note that:

$$\begin{aligned} \|\alpha^*(x) - \alpha^*(y)\|_p &= \left( \|\alpha'(x) - \alpha'(y)\|_p^p + \sum_{\alpha} \|q_{\alpha} \cdot \zeta \circ \alpha(x) - q_{\alpha} \cdot \zeta \circ \alpha(y)\|_p^p \right)^{1/p} \\ &\leq \|\alpha'(x) - \alpha'(y)\|_p + \sum_{\alpha} \|q_{\alpha} \cdot \zeta \circ \alpha(x) - q_{\alpha} \cdot \zeta \circ \alpha(y)\|_p \\ &\leq E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p] + \sum_{\alpha} q_{\alpha} \cdot \|\zeta \circ \alpha(x) - \zeta \circ \alpha(y)\|_p \\ &\leq E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p] + \sum_{\alpha} q_{\alpha} \cdot \|\alpha(x) - \alpha(y)\|_p \\ &\leq 2E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p], \end{aligned}$$

where the last line is by how we defined  $q_{\alpha}$  and the fact we are summing over all possible choices of  $\alpha$  (which we have already shown comes from a finite size set). For all  $x, y \in X$ ,  $E[\|\alpha(x) - \alpha(y)\|_2] \leq (\frac{155}{2} \cdot H_k \cdot c_S + (\frac{225}{2} \cdot H_k + 1) \cdot c_X) \cdot d(x, y)$ . The second line is by the power mean inequality and the third is by the fact that  $\|\alpha'(x) - \alpha'(y)\|_p$  is a lower bound on  $E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p]$  due to the fact  $\alpha'(x)$  is a sub-vector of  $\alpha(x)$  no matter the choice of  $b$ , as well as facts about distance between scaled vectors. The fourth line is by the fact that  $\zeta \circ \alpha(x)$  is a sub-vector of  $\alpha(x)$ , and the fifth is by definition of  $E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p]$ .

In finding this embedding, we did blow the dimension up by an exponential factor, but any  $\ell_p$  metric on  $n$  points is isometrically embeddable in at most  $n$  dimensions, so we have still proven existence.

Additionally, note that for arbitrary  $x, y \in X$ , contraction is arbitrary, but for  $x, y \in X \setminus K$ ,  $\alpha^*(x), \alpha^*(y)$  differ only on the indices associated with  $\alpha'(x), \alpha'(y)$ , which are the same as  $\alpha_S(x), \alpha_S(y)$ , and thus for these pairs, the distance is non-contracting and has expansion at most  $c_S$ . Also note that for  $p = 1$ , we get the following simpler analysis (using the fact that  $\sum_{\alpha} q_{\alpha} = 1$ ).

$$\begin{aligned} \|\alpha^*(x) - \alpha^*(y)\|_1 &= \left( \|\alpha'(x) - \alpha'(y)\|_1 + \sum_{\alpha} \|q_{\alpha} \cdot \zeta \circ \alpha(x) - q_{\alpha} \cdot \zeta \circ \alpha(y)\|_1 \right)^1 \\ &= \sum_{\alpha} q_{\alpha} \|\alpha'(x) - \alpha'(y)\|_1 + \sum_{\alpha} q_{\alpha} \cdot \|\zeta \circ \alpha(x) - \zeta \circ \alpha(y)\|_1 \\ &= \sum_{\alpha} q_{\alpha} \cdot \|\alpha(x) - \alpha(y)\|_p \\ &= E_{\pi, b}[\|\alpha(x) - \alpha(y)\|_p] \end{aligned}$$

Additionally, we get that contraction from Lemma 4.1 is at most  $3^{1-1/p}$ , which is 1 for  $p = 1$ .  $\square$

*Proof of Theorem 2.6.* Notice that Algorithm 1 can replace its use of  $\alpha_X$  in line 8 with any embedding of  $K_i \cup \{\gamma(u_i)\}$ , and if we have an upper bound on such an embedding's distortion, we can replace  $c_X$  in all of the theorems and lemmas in this section with that bound.  $\square$

## 5 Conclusion

In this paper, we give a bi-criteria approximation algorithm that given a constant  $c$  and metric  $(X, d)$  finds an  $(O(k \log^4 k), O(c))$ -outlier embedding into  $\ell_2$  if the metric has a  $(k, c)$ -outlier embedding into  $\ell_2$ . In doing so, we give an algorithm that given a metric space  $(X, d)$ , a  $c_S$ -distortion embedding of a subset  $S \subseteq X$  into  $\ell_p$  and a  $c_X$ -distortion embedding of  $X$  into  $\ell_p$ , finds a single (composition) embedding of  $X$  into  $\ell_p$  such that distortion between points in  $S$  is at most  $c_S$  and expansion between all pairs of points is at most  $O(c_S + H_k \cdot c_X)$ . We also leave several open questions on this topic. Among them, we ask:

- Is there a polynomial time algorithm that given constant  $c$ , finds an  $(O(k), O(c))$ -embedding into  $\ell_2$ ?
- What bicriteria approximations can be obtained for outlier embeddings into  $\ell_p$  for other values of  $p$ ?
- Do there exist nested compositions that do not incur the contraction presented in this paper (i.e. non-weak nested embeddings)?
- Can the parameters for our nested embedding algorithm be improved?
- Can our hardness of approximation result be extended to non-isometric outlier embeddings?

## References

- [1] I. Abraham, Y. Bartal, J. Kleinberg, T.-H.H. Chan, O. Neiman, Kedar Dhamdhere, A. Slivkins, and Anupam Gupta. Metric embeddings with relaxed guarantees. In *46th Annual IEEE Symposium on Foundations of Computer Science (FOCS'05)*, pages 83–100, 2005.
- [2] Noga Alon, Mihai Badoiu, Erik D. Demaine, Martin Farach-Colton, Mohammad Taghi Hajiaghayi, and Anastasios Sidiropoulos. Ordinal embeddings of minimum relaxation: General properties, trees, and ultrametrics. *ACM Trans. Algorithms*, 4(4):46:1–46:21, 2008.
- [3] Sanjeev Arora, László Lovász, Ilan Newman, Yuval Rabani, Yuri Rabinovich, and Santosh Vempala. Local versus global properties of metric spaces. *SIAM Journal on Computing*, 41(1):250–271, 2012.
- [4] Mihai Badoiu, Julia Chuzhoy, Piotr Indyk, and Anastasios Sidiropoulos. Low-distortion embeddings of general metrics into the line. In Harold N. Gabow and Ronald Fagin, editors, *Proceedings of the 37th Annual ACM Symposium on Theory of Computing, Baltimore, MD, USA, May 22-24, 2005*, pages 225–233. ACM, 2005.
- [5] Mihai Badoiu, Julia Chuzhoy, Piotr Indyk, and Anastasios Sidiropoulos. Embedding ultrametrics into low-dimensional spaces. In Nina Amenta and Otfried Cheong, editors, *Proceedings of the 22nd ACM Symposium on Computational Geometry, Sedona, Arizona, USA, June 5-7, 2006*, pages 187–196. ACM, 2006.
- [6] Mihai Badoiu, Piotr Indyk, and Anastasios Sidiropoulos. Approximation algorithms for embedding general metrics into trees. In Nikhil Bansal, Kirk Pruhs, and Clifford Stein, editors, *Proceedings of the Eighteenth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2007, New Orleans, Louisiana, USA, January 7-9, 2007*, pages 512–521. SIAM, 2007.
- [7] Jean Bourgain. On lipschitz embedding of finite metric spaces in hilbert space. *Israel Journal of Mathematics*, 52:46–52, 1985.
- [8] T. H. Hubert Chan, Michael Dinitz, and Anupam Gupta. Spanners with slack. In Yossi Azar and Thomas Erlebach, editors, *Algorithms – ESA 2006*, pages 196–207, Berlin, Heidelberg, 2006. Springer Berlin Heidelberg.
- [9] Moses Charikar, Konstantin Makarychev, and Yury Makarychev. Local global tradeoffs in metric embeddings. *SIAM Journal on Computing*, 39(6):2487–2512, 2010.
- [10] Victor Chepoi, Feodor F. Dragan, Ilan Newman, Yuri Rabinovich, and Yann Vaxès. Constant approximation algorithms for embedding graph metrics into trees and outerplanar graphs. *Discret. Comput. Geom.*, 47(1):187–214, 2012.

- [11] Karine Chubarian and Anastasios Sidiropoulos. Computing bi-lipschitz outlier embeddings into the line. In *International Workshop and International Workshop on Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, 2020.
- [12] James A. Clarkson. Uniformly convex spaces. *Transactions of the American Mathematical Society*, 40:396–414, 1936.
- [13] Mark de Berg, Krzysztof Onak, and Anastasios Sidiropoulos. Fat polygonal partitions with applications to visualization and embeddings. *J. Comput. Geom.*, 4:212–239, 2010.
- [14] M. Deza and S. Shpectorov. Recognition of the  $\ell_1$ -graphs with complexity  $O(nm)$ , or football in a hypercube. *European Journal of Combinatorics*, 17(2-3):279–289, 1996.
- [15] Michel Deza and Monique Laurent. The cut cone and  $\ell_1$  metrics. In R. L. Graham, B. Korte, Bonn L. Lovasz, A. Wigderson, and G.M. Ziegler, editors, *Geometry of Cuts and Metrics*, chapter 4, pages 37–52. Springer, 1997.
- [16] D. Ž. Djoković. Distance-preserving subgraphs of hypercubes. *Journal of Combinatorial Theory, Series B*, 14(3):263–267, 1973.
- [17] Jeff Edmonds, Anastasios Sidiropoulos, and Anastasios Zouzias. Inapproximability for planar embedding problems. In Moses Charikar, editor, *Proceedings of the Twenty-First Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2010, Austin, Texas, USA, January 17-19, 2010*, pages 222–235. SIAM, 2010.
- [18] Michael R. Fellows, Fedor V. Fomin, Daniel Lokshtanov, Elena Losievskaja, Frances A. Rosamond, and Saket Saurabh. Distortion is fixed parameter tractable. *ACM Trans. Comput. Theory*, 5(4):16:1–16:20, 2013.
- [19] Ronald L. Graham and Peter M. Winkler. On isometric embeddings of graphs. *Transactions of the American Mathematical Society*, 288(2):527–536, 1985.
- [20] Piotr Indyk and Jirí Matousek. Low-distortion embeddings of finite metric spaces. In Jacob E. Goodman and Joseph O’Rourke, editors, *Handbook of Discrete and Computational Geometry, Second Edition*, pages 177–196. Chapman and Hall/CRC, 2004.
- [21] Christiane Lammersen, Anastasios Sidiropoulos, and Christian Sohler. Streaming embeddings with slack. In Frank Dehne, Marina Gavrilova, Jörg-Rüdiger Sack, and Csaba D. Tóth, editors, *Algorithms and Data Structures*, pages 483–494, Berlin, Heidelberg, 2009. Springer Berlin Heidelberg.
- [22] Nathan Linial, Eran London, and Yuri Rabinovich. The geometry of graphs and some of its algorithmic applications. In *35th Annual Symposium on Foundations of Computer Science, Santa Fe, New Mexico, USA, 20-22 November 1994*, pages 577–591. IEEE Computer Society, 1994.
- [23] Konstantin Makarychev and Yury Makarychev. A union of euclidean metric spaces is euclidean. *Discrete Analysis*, 14, 02 2016.
- [24] Jirí Matousek and Anastasios Sidiropoulos. Inapproximability for metric embeddings into  $\mathbb{R}^d$ . In *49th Annual IEEE Symposium on Foundations of Computer Science, FOCS 2008, October 25-28, 2008, Philadelphia, PA, USA*, pages 405–413. IEEE Computer Society, 2008.
- [25] Amir Nayyeri and Benjamin Raichel. Reality distortion: Exact and approximate algorithms for embedding into the line. In Venkatesan Guruswami, editor, *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015, Berkeley, CA, USA, 17-20 October, 2015*, pages 729–747. IEEE Computer Society, 2015.
- [26] Mikhail I. Ostrovskii and Beata Randrianantoanina. On  $\ell_1$ -embeddability of unions of  $\ell_1$ -embeddable metric spaces and of twisted unions of hypercubes. *Analysis and Geometry in Metric Spaces*, 10(1):313–329, 2022.
- [27] S. V. Shpectorov. On scale embeddings of graphs into hypercubes. *Eur. J. Comb.*, 14(2):117–130, March 1993.

- [28] Anastasios Sidiropoulos, Mihai Badoiu, Kedar Dhamdhere, Anupam Gupta, Piotr Indyk, Yuri Rabinovich, Harald Räcke, and R. Ravi. Approximation algorithms for low-distortion embeddings into low-dimensional spaces. *SIAM J. Discret. Math.*, 33(1):454–473, 2019.
- [29] Anastasios Sidiropoulos, Dingkang Wang, and Yusu Wang. Metric embeddings with outliers. In *ACM-SIAM Symposium on Discrete Algorithms*, 2015.

## A Full constants for Section 4

Here we give a version of Lemma 4.2 more generally in terms of  $\tau$  and other flexible parameters. The proof would be identical to that given in Section 4.

**Lemma A.1.** *Let  $\alpha \leftarrow$  Algorithm 1( $(X, d), S, p, \alpha_S, \alpha_X, \tau$ ). Then we have the following bounds on the expansion for each pair  $x, y \in X$ :*

- (a) *If  $x, y \in S$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq c_S \cdot d(x, y)$ .*
- (b) *If  $x, y \in K_i$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq c_X \cdot d(x, y)$ .*
- (c) *If  $x \in S, y \in X \setminus S$ , then  $\|\alpha(x) - \alpha(y)\|_p \leq [(\tau + 5)c_S + (2\tau + 5)c_X] \cdot d(x, y)$ .*
- (d) *If  $x, y \in X \setminus S$  and  $d(x, \gamma(x)) \leq \kappa \cdot d(x, y)$  for positive  $\kappa$  and  $\gamma$  as defined in line 2 of the algorithm, then*

$$\|\alpha(x) - \alpha(y)\|_p \leq [((2\tau + 8)\kappa + \tau + 5) \cdot c_S + ((4\tau + 10)\kappa + 2\tau + 5) \cdot c_X] \cdot d(x, y),$$

where  $b$  is as chosen in line  $x$  of the algorithm.

- (e) *If  $x, y \in X \setminus S$  and  $d(x, \gamma(x)), d(y, \gamma(y)) > \kappa \cdot d(x, y)$  for  $\gamma$  as defined in line 2 of the algorithm and real number  $\kappa > 1$ , then*

$$\begin{aligned} & E_\alpha[\|\alpha(x) - \alpha(y)\|_p] \\ & \leq \left[ \left[ \frac{\tau + 3}{(\kappa - 1)\tau} \cdot H_k \cdot (\kappa(\tau + 3) + (\kappa + 1)(\tau + 5)) \right] c_S + \left[ 1 + \frac{\tau + 3}{(\kappa - 1)\tau} \cdot H_k \cdot (2\tau + 5)(2\kappa + 1) \right] c_X \right] \cdot d(x, y). \end{aligned}$$

## B Hardness of finding outlier sets

Sidiropoulos *et al.* [29] showed that for any  $d \geq 2$ , it is NP-hard to determine the size of the smallest outlier set for a finite metric  $(X, d)$  such that the metric without the outlier set is isometrically embeddable into  $\ell_2$ . Additionally, because they reduce from Vertex Cover, they show that under the Unique Games Conjecture it is NP-hard to approximate the size of such a set to a factor better than  $2 - \epsilon$ .

In this appendix, we will give an alternate proof of a similar conclusion, but we extend their result to show that it holds even if the input metric is an unweighted graph metric. We note that unlike the Sidiropoulos *et al.* proof, our proof does not apply for arbitrary choice of dimension  $d$ .

First, we claim the following Lemma B.1 which we will prove later. Note that it is NP-hard to decide if a general metric is isometrically  $\ell_1$ -embeddable and thus it is hard to decide if the minimum outlier set for such an embedding has size 0 or size larger than 0, implying hardness of any approximation for this value. However,  $\ell_1$ -embeddability for can be decided in polynomial time for unweighted graph metrics [27, 14, 15], and we show that even with this restriction on the input, it is hard to determine minimum outlier set size.

**Lemma B.1.** *Let  $(X, d)$  be the distance metric for an unweighted graph  $G = (V, E)$ . then, given  $(X, d, k)$  it is NP-hard to decide if there exists a subset  $K \subseteq X$  with  $|K| = k$  such that  $(X \setminus K, d|_{X \setminus K})$  is isometrically embeddable into  $\ell_1$ , even when the input metric is an unweighted graph metric.*

*Under the unique games conjecture, it is NP-hard to find a  $2 - \epsilon$  approximation for the minimum such  $k$ , for any  $\epsilon > 0$ .*




Figure 4: Example of a subgraph formed by nodes  $\{u_1 = x, u_2 = y, v_1 = w, v_2 = z\}$  as defined in the proof of Theorem 2.9.

We can now use this result to prove Theorem 2.9.

*Proof of Theorem 2.9.* We appeal to Lemma B.1 to show that the theorem holds for  $p = 1$  and here we show that it holds for  $1 < p < \infty$ .

Consider a graph  $G = (V, E)$ . Given  $G$ , we give a polynomial time construction of an unweighted graph metric  $(V', d_{G'})$  such that the size of the minimum outlier set for embedding the metric into  $\ell_p$  is the same as the size of the minimum vertex cover on  $G$ . In this proof, minimum outlier set refers to the minimum outlier set for distortion 1 and we consider embeddings into  $\ell_p$  for finite integer  $p > 1$ .

**Construction:** We will construct an unweighted graph  $G' = (V', E')$  and let  $d_{G'}$  be the distance metric on this graph. In particular, let  $V' = \{u_1 | u \in V\} \cup \{u_2 | u \in V\}$ . Add edges between every pair of nodes in the graph, but omit edges  $u_2 v_2$  for  $uv \in E$ .

**Correctness:** Let  $K$  be a minimum outlier set on  $(V', d_{G'})$  and let  $\hat{V}$  be a minimum vertex cover on  $G$ . We claim  $|K| = |\hat{V}|$ .

$|K| \leq |\hat{V}|$  : We construct an outlier set of size at most  $|\hat{V}|$ . Define  $K' := \{u_2 | u \in \hat{V}\}$ . We claim that  $(V' \setminus K', d|_{V' \setminus K'})$  is the equidistant metric with distance 1 between all points, which is always embeddable in  $\ell_p$  for any  $p$ . Assume this is not the case. Note that all distances in  $G'$  are 1 or 2, so there exists a pair of nodes with distance 2 between them. The only such pairs are of the form  $\{u_2, v_2\}$  for  $uv \in E$ . However,  $u_2, v_2 \in V' \setminus K'$  implies that  $\hat{V}$  does not cover edge  $uv$  so it is not a vertex cover and we reach a contradiction. Thus since  $K$  is minimum, we get  $|K| \leq |K'| \leq |\hat{V}|$ .

$|\hat{V}| \leq |K|$  : We construct a vertex cover of size at most  $|K|$ . Define  $\hat{V}' := \{u | u_1 \in K \text{ or } u_2 \in K\}$ . We see  $|\hat{V}'| \leq |K|$  and we claim it is a vertex cover. Assume otherwise. Then there exists an edge  $uv$  that is not covered by  $\hat{V}'$  and there is a subgraph of the form in Figure 4 in the induced subgraph of  $G'$  on  $V' \setminus K'$ ,  $G'[V' \setminus K']$ . Note that distances in this subgraph are exactly distances in the entire graph by our construction. Thus we need only show that this subgraph is not isometrically embeddable in  $\ell_p$ .

Note that  $\ell_p$  is a strictly convex space for  $1 < p < \infty$  [12]. Thus, for  $a, b \in \mathbb{R}^t$  for any fixed  $t$ , if  $\|a\|_p = \|b\|_p = 1$  and  $a \neq b$ , then  $\|\frac{a+b}{2}\|_p < 1$ , which implies  $\|a+b\|_p < 2^{1/p}$ . Let  $w, x, y, z$  be four points in  $\ell_p$  such that the distance between all pairs of points is 1, except between  $y$  and  $z$  which are a distance 2 apart. Then let  $a = y - x$  and  $b = x - z$ . We have  $\|y - x\|_p = \|x - z\|_p = 1$ , so  $\|y - z\|_p < 2^{1/p}$  unless  $y - x = x - z$ . Since  $2^{1/p} < 2$  for  $p > 1$ , we get that  $y - x = x - z \implies x = \frac{y+z}{2}$ . However, equivalent analysis on  $w$  implies  $w = \frac{y+z}{2}$ . This means that  $x$  and  $w$  are the same point and their distance is 0, not 1. Thus, this set of four points cannot exist in  $\ell_p$  for  $1 < p < \infty$  and the subgraph in Figure 4 is not isometrically embeddable into this space.  $\square$

*Proof of Lemma B.1.* As in the previous lemma, we reduce the vertex cover problem to this problem. Consider a graph  $G = (V, E)$ . We give a polynomial time construction of an unweighted graph metric  $(V', d_{G'})$  such that that size of the minimum outlier set for embedding the metric into  $\ell_1$  is the same as the size of the minimum vertex cover on  $G$ . In this proof, minimum outlier set refers to the minimum outlier set for distortion 1 and we consider embeddings into  $\ell_1$ .

**Construction:** We will construct an unweighted graph  $G' = (V', E')$  and let  $d_{G'}$  be the distance metric on this graph. In particular, let  $V' = \{x_i, y_i, z_i, w_i | i \in V\}$ . Add edges between every pair of nodes in the graph, but omit edges  $x_i y_i$  for all  $i$ . Additionally, omit edges  $x_i x_j$  if  $ij \in E$ .

**Correctness:** Let  $K$  be a minimum outlier set on  $(V', d_{G'})$  and let  $\hat{V}$  be a minimum vertex cover on  $G$ . We claim  $|K| = |\hat{V}|$ .

$|K| \leq |\hat{V}|$  : Define  $K' := \{x_i | i \in \hat{V}\}$ . We claim that  $(V' \setminus K', d_{G'}|_{V' \setminus K'})$  is  $\ell_1$  embeddable. In particular, [27, 14], and [15] show that if an unweighted graph is such that each node has at most one other node it does not have an edge to (i.e. if the graph is a subgraph of a cocktail party graph), then the graph is  $\ell_1$ -embeddable. Note that  $d_{G'}$  restricted to the nodes in  $V' \setminus K'$  is in fact the distance metric on the induced subgraph of  $G'$  on those same nodes,  $G'[V' \setminus K']$ . This is because we only removed some of the  $x_i$ , which cannot affect distances between any remaining pairs.

Thus, we are left with showing that  $G'[V' \setminus K']$  is such that each node has at most one other node to which it does not have an edge. Assume otherwise. Then there exists  $a, b, c \in V' - K'$  such that  $a$  does not have an edge to  $b$  or to  $c$ . The only nodes in  $G'$  that are missing an edge to more than one other node in  $G'$  are some of the  $x_i$ , so  $a$  must be  $x_i$  for some  $i$ . Additionally, the nodes that  $x_i$  does not have an edge to are  $y_i$  and all  $x_j$  such that  $ij \in E$ . Thus, at least one of  $b$  and  $c$  must be  $x_j$  for some  $j$  such that  $ij \in E$ . However, this implies  $x_i, x_j \notin K' \implies i, j \notin \hat{V}$ , so  $\hat{V}$  is not a vertex cover and we reach a contradiction. Thus  $K'$  is an outlier set and we get  $|K| \leq |K'| = |\hat{V}|$

$|\hat{V}| \leq |K|$  : Define  $\hat{V}' := \{i | x_i, y_i, z_i, \text{ or } w_i \in K\}$ . We have  $|\hat{V}'| \leq |K|$ , so if  $\hat{V}'$  is a valid vertex cover then we obtain the desired bound. Assume  $\hat{V}'$  is not a vertex cover. Then there exists  $u_i u_j \in E$  such that  $\{x_i, y_i, z_i, w_i, x_j, y_j, z_j, w_j\} \subseteq V' - K$ . These nodes form the subgraph pictured in Figure 5, and  $d_{G'}$  on this subset of nodes has the same value as the graph metric on this subgraph. Thus, we need only show that the subgraph in Figure 5, which we call  $G''$ , is not  $\ell_1$ -embeddable.

By [15], an unweighted graph is  $\ell_1$ -embeddable if and only if there exists an integer  $t \in \mathbb{Z}^+$  such that the same graph with all edge weights set to  $t$  is hypercube embeddable. Deza and Shpectorov [27, 14] show that if an unweighted graph is  $\ell_1$ -embeddable and it is not “reducible” as defined by [19], then it must be an isometric subgraph of a cocktail party graph or a half-cube (a type of graph that is hypercube embeddable at scale 2). In Lemma B.2 we use Graham and Winkler’s [19] techniques to show that  $G''$  is not reducible.

Additionally,  $x_i$  lacks neighbors  $y_i$  and  $x_j$ , so this is not a subgraph of a cocktail party graph, which is a graph in which each node is a neighbor of all but one node. This leaves us with showing that  $G''$  is not a subgraph of a half-cube, which we prove by showing that it is not hypercube embeddable at scale 2 in Lemma B.3. Thus, we conclude  $G''$  cannot be a subgraph of  $G'[V' \setminus K]$  and  $\hat{V}'$  must be a vertex cover. □

**Lemma B.2.** *The graph appearing in Figure 5 is not reducible, where reducible is as defined by [19]*

*Proof.* Graham and Winkler show that a graph is not reducible if all of its edges are in the same equivalence class of the equivalence relation  $\hat{\theta}$ .  $\hat{\theta}$  is defined to be the transitive closure of  $\theta$ , which is defined on the edges of a graph as follows:

For a graph  $G'' = (V'', E'')$ , edges  $ab, cd \in E''$  are related by  $\theta$  if and only if

$$[d_{G''}(a, c) - d_{G''}(a, d)] - [d_{G''}(b, c) - d_{G''}(b, d)] \neq 0.$$




Figure 5: Example of a subgraph formed by nodes  $\{x_i, y_i, z_i, w_i, x_j, y_j, z_j, w_j\}$  as defined in the proof of Lemma B.1, where  $ij \in E$ .

We will show that all edges of  $G''$  are in the same equivalence class of  $\hat{\theta}$ .

First, notice that in the big/main clique of  $y_i, y_j, z_i, z_j, w_i, w_j$  in  $G''$ , all edges must be related by  $\hat{\theta}$ . Take two adjacent edges  $ab, bc$ . Since all distances in the clique are 1, we get  $[d_{G''}(a, b) - d_{G''}(a, c)] - [d_{G''}(b, b) - d_{G''}(b, c)] = 1 \neq 0$ . Thus all adjacent edges in the clique are related by  $\theta$  and thus all edges in the clique are related by  $\hat{\theta}$ . This just leaves us to consider the edges to  $x_i$  and  $x_j$ . We see that  $z_i x_i \theta z_i w_i$  because again we have adjacent edges with distances between all three vertices being 1. The same analysis goes for all the other edges to  $x_i$  and to  $x_j$  since they are part of forming a smaller clique with  $x_i$  or  $x_j$  and two members of the main clique. This means they are all in the same equivalence class as the edges in the main clique and thus all edges are in the same equivalence class, meaning it is irreducible.  $\square$

**Lemma B.3.** *The graph appearing in Figure 5 is not hypercube embeddable, and the same graph with all edge weights scaled to 2 is also not hypercube embeddable.*

*Proof.* The graph is not bipartite, so it is not hypercube embeddable at scale 1 [16].

To begin the second part of this part of the proof, we first consider an alternative view of a hypercube embedding. In particular, if we have a hypercube embedding in dimension  $t$  for a metric space on  $n$  nodes, we can write an  $n \times t$  matrix in which each row of the matrix is the binary string associated with a particular node. The number of of columns where two rows differ is then the distance between the corresponding nodes in the graph. Notice that this matrix defines a hypercube embedding and a hypercube embedding defines this matrix. Additionally, we can remove any columns in which all rows have the same value without affecting the “distance” between rows. Additionally, if we swap the order of the columns or if we pick a column of this matrix and flip all bits in that column, it has no effect on the distance between the rows. Thus, we can always assume that the top row of the matrix for a hypercube embedding is made up of all 0s, as such a choice of embedding must exist if some hypercube embedding does.

Let’s begin constructing a hypercube embedding for the graph  $G''$ . In particular, we define a matrix in which the top row corresponds to the node  $x_i \in V(G'')$  and the row consists of all 0s (which we’ve already argued is a fine assumption). Then we know that all rows except those corresponding to  $y_i$  and  $x_j$  must have exactly two 1s in their rows, and  $y_i$  and  $x_j$  have exactly four 1s in their rows (in order to make their distance from  $x_i$  correct). We will assume that all the columns in the matrix we construct have at least two distinct values in each column, as we have argued such an embedding must exist if any embedding exists since we can delete columns where all values are equal. Thus, we can assume that  $M$  has at most 18 columns since there are at most eighteen 1s in the entire matrix. This leads us to the following partial embedding where most values are unassigned so far.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$x_i$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_i$																		
$y_i$																		
$x_j$																		
$w_i$																		
$w_j$																		
$z_j$																		
$y_j$																		

We will next assume that the two 1s in  $z_i$ 's row are in the first two columns. (As we mentioned, we can always rearrange the columns of a hypercube embedding matrix to make one that looks this way since those columns at this point are indistinguishable.) This gives us the following matrix:

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$x_i$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_i$	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$y_i$																		
$x_j$																		
$w_i$																		
$w_j$																		
$z_j$																		
$y_j$																		

Next, consider node  $y_i$  that is a distance 2 closer to  $z_i$  than to  $x_i$  (in the scaled graph, as it is a distance 1 closer in the original graph). Notice that in all columns after the first two columns, if  $y_i$  doesn't match  $x_i$ , then it also doesn't match  $z_i$ . Thus, the only place where it can have these two differences with  $x_i$  that it does not have those differences with  $z_i$  is the first two columns. Thus,  $y_i$  must have 1s in the first two columns. The same goes for  $x_j$  because it is also a distance 2 closer to  $z_i$  than to  $x_i$ . For all the other nodes, they are equidistant from  $x_i$  and  $z_i$ . Thus, because they must have the same number of differences with  $x_i$ 's and  $z_i$ 's rows after the first two columns, they must also have the same number of differences in the first two columns. This means they either have 10 or 01 in the first two columns. Because we don't know which of these to put in those columns, for now we don't fill those values in yet. Because  $y_i$  and  $x_j$  are a distance 4 from  $x_i$ , they must have four 1s total. We will assume that the first four 1s for  $y_i$  are in the first four columns, because we can rearrange any matrix in which the other two 1s are in later columns such that they are in the second two columns. This leaves us with the following matrix constructed so far.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$x_i$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_i$	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$y_i$	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$x_j$	1	1																
$w_i$																		
$w_j$																		
$z_j$																		
$y_j$																		

Now we can consider the distance between  $x_j$  and  $y_i$ . They are a distance 2 apart in the scaled graph, and we only have two more 1s that we can place in  $x_j$ 's row. If we put both of these 1s in positions 3 and 4, then there are no differences between  $x_j$  and  $y_i$ , and if we put none of these 1s in positions 3 or 4, the two have a distance 4 apart. Thus, we must have exactly a single 1 in columns 3 and 4. We will assume that this is in column 3, as at the moment these columns are indistinguishable so if a good embedding has the opposite assignment, we can swap the columns to get



something consistent with this embedding. We will also assume that the last 1 is in column 5 since the columns after 4 are indistinguishable at this point.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$x_i$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_i$	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$y_i$	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$x_j$	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
$w_i$																		
$w_j$																		
$z_j$																		
$y_j$																		

Now we can consider labeling  $w_i, w_j, z_j$ . We already established that due to the fact they are equidistant from  $x_i$  and  $z_i$ , they must all have either 10 or 01 in the first two columns. Then they have a single 1 left for the rest of the columns since they are a distance 2 from  $x_i$ . We notice that if we put this second 1 in column 4, then the distance between the node and  $x_j$  is 4 (one difference in the first two columns and one each in each of columns 3 through 5), when it should be 2. Analogously, if the 1 is in column 5, the distance to  $y_i$  is 4 instead of 2. If the 1 is in column 6 or larger, the distance to  $y_i$  and  $x_j$  is 4 instead of 2. Thus, this leaves us with putting the 1 in column 3 for all three of these nodes. Everything after this must be 0s since we used up our only other 1 in the first two columns. This gives us the following partial embedding.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18
$x_i$	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_i$	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$y_i$	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$x_j$	1	1	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
$w_i$			1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$w_j$			1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$z_j$			1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
$y_j$																		

Now we notice that  $w_i, w_j, z_j$  have no differences after column 3. Thus, all of their differences must be in the first two columns. This means that we must come up with three length 2 binary strings that are all Hamming distance 2 from each other, which is impossible. Because all of the decisions we made in constructing this embedding were necessary, this partial construction is required which means it's impossible to construct a hypercube embedding for this graph at scale 2.

There are also other ways to verify that this graph is not hypercube embeddable at scale 2 or that it is not a halved cube, including running the algorithm of Deza and Shpectorov [14] for doing so. We presented the proof this way because their algorithm is more complex and generalized than what is needed for our specific purposes.  $\square$