

ON HARADA'S IDENTITY AND SOME OTHER CONSEQUENCES OF BURNSIDE'S VANISHING PROPERTY

SEBASTIAN BURCIU

ABSTRACT. In this short note, we prove some consequences of Burnside's vanishing property [BP23]. It is known that Harada's identity concerning the product of all conjugacy classes of a finite group is a consequence of Burnside's vanishing property of characters. We prove a similar formula for any weakly-integral fusion category. In the second part, we prove some structural results concerning nilpotent and modular fusion categories. As an application we show that any fusion category of dimension $p^2q^2r^2d$ with $p < q < r$ prime numbers and d a square-free integer is weakly-group theoretical.

CONTENTS

1. Introduction	1
2. Preliminaries	3
3. Proof of Theorem 1.2	5
4. Proof of Theorem 1.3 and some other applications	6
References	7

1. INTRODUCTION

A classical vanishing result of Burnside states that any irreducible non-linear character of a finite group vanishes on at least one element of the group.

Based on this vanishing result, in [Har08, Theorem 1], K. Harada proved that the product of all conjugacy classes of a finite group form a coset with respect to the commutator subgroup G' .

Theorem 1.1. (*Harada*, [Har08]) *For any finite group G , the following identity holds:*

$$\left(\prod_{j=0}^m \frac{C_j}{|C^j|} \right)^2 = \frac{1}{|G'|} \sum_{C^j \subseteq G'} C_j$$

Date: June 21, 2023.

2020 *Mathematics Subject Classification.* 18M20; 16T30.

The first author is supported by a grant of the Ministry of Research, Innovation and Digitization, CNCS/CCCDI - UEFISCDI, project number PN-III-P4-ID-PCE-2020-0878, within PNCDI III.

where G' is the commutator subgroup of G .

In the above theorem $\{\mathcal{C}^j\}_{j=0}^m$ are all the conjugacy classes of G and $|\mathcal{C}^j|$ is the size of the conjugacy class. The elements $C_j := \sum_{g \in \mathcal{C}^j} g$ are the class sums associated to \mathcal{C}^j .

One of the goals of this short note is to present a similar formula for any weakly-integral fusion category. For weakly-integral fusion categories Burnside's vanishing result was extended in [Bur23]. Some other consequences of Brauer's vanishing result at the level of fusion categories are also presented.

Let \mathcal{C} be a weakly-integral fusion category, \mathcal{C}^j the set of all its conjugacy classes and C_j their associated class sums. Our first main result is the following:

Theorem 1.2. *For any weakly-integral fusion category, with the above notations, one has*

$$\left(\prod_{j=0}^m \frac{C_j}{\text{FPdim}(\mathcal{C}^j)} \right)^2 = \frac{\text{FPdim}(\mathcal{C}_{\text{pt}})}{\text{FPdim}(\mathcal{C})} \left(\sum_{\mu_j \in \mathcal{J}_{\mathcal{C}_{\text{pt}}}} C_j \right)$$

For the definition of conjugacy classes \mathcal{C}^j and their associated class sums C_j , see Section 2. Moreover, the support of the fusion subcategory \mathcal{C}_{pt} is defined in [Bur20, Subsection 4.2].

In [OY23, Theorem 4.2] it was shown that any weakly group-theoretical modular tensor category whose dimension is nd with $(n, d) = 1$ is in fact a Deligne product of $\mathcal{C}(\mathbb{Z}_q, d) \boxtimes \mathcal{D}$ of two other modular tensor categories.

Suppose that \mathcal{C} is a weakly-integral modular category. As usually we suppose that \mathcal{C} is of type $(d_1, n_1; d_2, n_2; \dots, d_r, n_r)$ with $d_1=1$. Our second main result of this note is the following:

Theorem 1.3. *Suppose that \mathcal{C} is a weakly-integral modular tensor category with $\text{FPdim}(\mathcal{C}) = nd$ with $(n, d) = 1$ and d a square-free integer.*

Then, with the above notations one has $d|n_i$ for all $i \geq 1$.

As an application we prove the following:

Theorem 1.4. *Let \mathcal{C} be an integral modular fusion category of dimension $p^2q^2r^2d$ with d a square-free number and $p < q < r$ three prime numbers. Then any such fusion category is weakly group-theoretical and can be written as a Deligne product $\mathcal{A} \boxtimes \mathcal{B}$ with $\mathcal{A} \simeq \mathcal{C}(\mathbb{Z}_q, d)$ and \mathcal{B} a weakly group theoretical fusion category of dimension $p^2q^2r^2$.*

Note that the case $d = 1$ of the above result was treated in [CP22, Prop. 3.8.]. Proposition 4.5 proves another consequence of Burnside vanishing result for nilpotent fusion categories.

Shortly, this note is organized as follows: In Section 2 we remind the basics on pivotal fusion categories that are needed through the paper. In Section 3 we prove Harada's identity for weakly-integral fusion categories. In Section 4 we prove Theorem 1.3 and Proposition 4.5

2. PRELIMINARIES

Throughout this paper we work over the ground field \mathbb{C} . For the basic theory of fusion categories, we refer the reader to the monograph [EGNO15].

Recall that a pivotal structure of a rigid monoidal category \mathcal{C} is an isomorphism $j : \text{id}_{\mathcal{C}} \rightarrow ()^{**}$ of monoidal functors. A pivotal monoidal category is a rigid monoidal category endowed with a pivotal structure.

For any fusion category \mathcal{C} , the monoidal center (or the Drinfeld center) of \mathcal{C} is a braided fusion category $\mathcal{Z}(\mathcal{C})$ endowed with a forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$, see [EGNO15].

The forgetful functor F also admits a right adjoint functor $R : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ and $Z := FR : \mathcal{C} \rightarrow \mathcal{C}$ is a Hopf comonad defined as an end. Indeed, following [Shi17, Section 2.6] one has that

$$(2.1) \quad Z(V) \simeq \int_{X \in \mathcal{C}} X \otimes V \otimes X^*$$

It is known that $A := Z(\mathbf{1})$ has the structure of central commutative algebra in $\mathcal{Z}(\mathcal{C})$.

The vector space $\text{CE}(\mathcal{C}) := \text{Hom}_{\mathcal{C}}(\mathbf{1}, A)$ is called *the set of central elements*. There is a canonical bijection:

$$(2.2) \quad \text{CE}(\mathcal{C}) \xrightarrow{\psi} \text{End}(\text{id}_{\mathcal{C}}), \quad \psi(a)_X = \rho_X(a \otimes \text{id}_X), \quad X = \mathbf{1} \otimes X \xrightarrow{a \otimes \text{id}_X} X \otimes Z(\mathbf{1}) \xrightarrow{\rho_X} X$$

The vector space $\text{CF}(\mathcal{C}) := \text{Hom}_{\mathcal{C}}(A_{\mathcal{C}}, \mathbf{1})$ is called the *space of class functions* of \mathcal{C} . To any object $X \in \text{Irr}(\mathcal{C})$ Shimizu has attached an element $\chi_X \in \text{CF}(\mathcal{C})$ called *the character of X* .

Recall $R : \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ is a right adjoint to the forgetful functor $F : \mathcal{Z}(\mathcal{C}) \rightarrow \mathcal{C}$. As explained in [Shi17, Theorem 3.8] this adjunction gives an isomorphism of \mathbb{C} -algebras

$$(2.3) \quad \text{CF}(\mathcal{C}) \xrightarrow{\cong} \text{End}_{\mathcal{Z}(\mathcal{C})}(R(\mathbf{1})), \quad \chi \mapsto Z(\chi) \circ \delta_{\mathbf{1}}.$$

where $\delta_{\mathbf{1}}$ is the comultiplication of the comonad Z associated at $\mathbf{1}$.

Let $\lambda \in \text{CF}(\mathcal{C})$ be a non-zero cointegral of \mathcal{C} . The *Fourier transform* of \mathcal{C} associated to λ is the linear map

$$(2.4) \quad \mathcal{F}_{\lambda} : \text{CE}(\mathcal{C}) \rightarrow \text{CF}(\mathcal{C}) \quad \text{given by} \quad a \mapsto \lambda \leftarrow \mathcal{S}(a)$$

where $\mathcal{S} : \text{CE}(\mathcal{C}) \rightarrow \text{CE}(\mathcal{C})$ is the above antipodal operator. The Fourier transform is a bijective \mathbb{k} -linear map whose inverse is given in [Shi17, Equation 5.17]. Since the monoidal center $\mathcal{Z}(\mathcal{C})$ is also a fusion category we can write $R(\mathbf{1}) = \bigoplus_{i=0}^m \mathcal{C}_i$ as a direct sum of simple objects in $\mathcal{Z}(\mathcal{C})$. Thus $\mathcal{C}^0, \dots, \mathcal{C}^m$ are the conjugacy classes of \mathcal{C} . Since the unit object $\mathbf{1}_{\mathcal{Z}(\mathcal{C})}$ is always a subobject of $R(\mathbf{1})$, we can assume $\mathcal{C}^0 = \mathbf{1}_{\mathcal{Z}(\mathcal{C})}$.

Let $\tilde{F}_0, \tilde{F}_1, \dots, \tilde{F}_m \in \text{End}_{\mathcal{Z}(\mathcal{C})}(R(\mathbf{1}))$ be the canonical projections on each of these conjugacy classes. Let also F_0, F_1, \dots, F_m be also the corresponding primitive idempotents of $\text{CF}(\mathcal{C})$ under the canonical adjunction isomorphism $\text{CF}(\mathcal{C}) \simeq \text{End}_{\mathcal{Z}(\mathcal{C})}(R(\mathbf{1}))$ from Equation (2.2).

We define $C_i := \mathcal{F}_\lambda^{-1}(F_i) \in \text{CE}(\mathcal{C})$ to be the *conjugacy class sums* corresponding to the conjugacy class \mathcal{C}^i where $\lambda \in \text{CF}(\mathcal{C})$ is a cointegral such that $\langle \lambda, u \rangle = 1$.

We denote by $\mu_j : \text{CF}(\mathcal{C}) \rightarrow \mathbb{C}$ the algebra homomorphisms corresponding to the central primitive idempotent F_j .

We adapt the following from [BP23, Definition 1.4].

Definition 2.1. *A fusion category \mathcal{C} has Burnside's vanishing property (or shortly, is Burnside) if for all $X_i \in \text{Irr}(\mathcal{C})$, the following two assertions are equivalent:*

- (1) *There is some $\mu_j \in \widehat{\mathcal{B}}$ such that $\mu_j(\chi_{X_i}) = 0$,*
- (2) *X_i is not an invertible object.*

By [Bur23, Theorem 2] any weakly integral fusion category is Burnside.

2.1. Definition of the support $\mathcal{J}_{\mathcal{D}}$. Let \mathcal{C} be a fusion subcategory with a commutative Grothendieck ring. For any fusion subcategory $\mathcal{D} \subseteq \mathcal{C}$, recall the definition of its support form [Bur20, Subsection 4.2]. It is well known that there is an inclusion of \mathbb{C} -algebras $\iota : \text{CF}(\mathcal{D}) \hookrightarrow \text{CF}(\mathcal{C})$. Thus there is a subset $\mathcal{J}_{\mathcal{D}} \subseteq \{0, \dots, m\}$ such that

$$(2.5) \quad \lambda_{\mathcal{D}} = \sum_{j \in \mathcal{J}_{\mathcal{D}}} F_j$$

since $\lambda_{\mathcal{D}}$ is an idempotent element inside $\text{CF}(\mathcal{C})$. The set $\mathcal{J}_{\mathcal{D}}$ is called the *support* of the fusion subcategory \mathcal{D} .

2.2. The dual hypergroup of $K(\mathcal{C})$. Let \mathcal{C} be a fusion category with a commutative Grothendieck group $K(\mathcal{C})$. On the dual space $\widehat{K(\mathcal{C})} := \text{Hom}_{\mathbb{C}}(K(\mathcal{C}), \mathbb{C})$ one defines a multiplicative structure as in [Bur21], such as, for all $f, g \in K(\mathcal{C})$ one has:

$$(2.6) \quad [f \star g]\left(\frac{\chi_i}{d_i}\right) := f\left(\frac{\chi_i}{d_i}\right)g\left(\frac{\chi_i}{d_i}\right), \text{ for all } \chi_i \in \text{Irr}(\mathcal{C}).$$

Then $\widehat{K(\mathcal{C})}$ is not anymore a fusion ring but an abelian normalized hypergroup, see [BP23, Definition 1.1].

Let (H, \mathcal{B}) be an abelian normalized hypergroup. By the commutativity assumption on $K(\mathcal{C})$, the set of algebra morphisms $\widehat{\text{Irr}(\mathcal{C})} := \{\mu_j\}_{j=0}^m$ forms a basis for the dual $\widehat{K(\mathcal{C})}$.

2.3. The isomorphism α . Note that the dual hypergroup $\widehat{K(\mathcal{C})}$ is denoted by $\widehat{\text{CF}(\mathcal{C})}$ in [Bur21]. Theorem 3.4 from the same paper implies that for any pseudo-unitary fusion category there is a canonical isomorphism of hypergroups

$$(2.7) \quad \alpha : \widehat{\text{CF}(\mathcal{C})} \rightarrow \text{CE}(\mathcal{C}), \mu_j \mapsto \frac{C_j}{\dim(\mathcal{C}^j)}.$$

3. PROOF OF THEOREM 1.2

Proof. Since \mathcal{C} is weakly-integral, it is a Burnside category, by [Bur23, Theorem 2]. Then [BP23, Corollary 4.5] implies that

$$(3.1) \quad \left(\prod_{j=0}^m \mu_j \right)^2 = \sum_{[X_i] \in \text{Irr}(\mathcal{C}_{\text{pt}})} \widehat{E}_i.$$

where \widehat{E}_i is the primitive central idempotent of $\widehat{K(\mathcal{C})}$ associated to $\chi_i := \chi_{X_i}$. Note that $\alpha(\widehat{E}_i) = E_i$, the primitive idempotent of $\text{CE}(\mathcal{C})$ associated to the irreducible character E_i , see [Shi17, Subsection 6.2]. Applying the isomorphism α to Equation (3.1) one obtains the following identity in $\text{CE}(\mathcal{C})$.

$$\left(\prod_{j=0}^m \frac{C_j}{\text{FPdim}(\mathcal{C}^j)} \right)^2 = \sum_{[X_i] \in \text{Irr}(\mathcal{C}_{\text{pt}})} E_i.$$

On the other hand one has by [Bur20, Eq. (4.13) and Eq. (4.15)] one has

$$\frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{\text{pt}})} \left(\sum_{[X_i] \in \text{Irr}(\mathcal{C}_{\text{pt}})} E_i \right) = \sum_{j \in \mathcal{J}_{\mathcal{C}_{\text{pt}}}} C_j$$

and this implies that

$$\left(\prod_{j=0}^m \frac{C_j}{\text{FPdim}(\mathcal{C}^j)} \right)^2 = \frac{\dim(\mathcal{C}_{\text{pt}})}{\dim(\mathcal{C})} \left(\sum_{j \in \mathcal{J}_{\mathcal{C}_{\text{pt}}}} C_j \right).$$

The proof of the theorem is finished. □

Corollary 3.1. *With the above notations, if the structure constants are integers then one has that*

$$(3.2) \quad \frac{\dim(\mathcal{C})}{\dim(\mathcal{C}_{\text{pt}})} \mid \prod_{j=0}^m \text{FPdim}(\mathcal{C}^j)$$

Corollary 3.2. *If \mathcal{C} is Burnside and modular with integral structure constants then*

$$(3.3) \quad \text{FPdim}(\mathcal{C}_{\text{ad}}) = \frac{\text{FPdim}(\mathcal{C})}{\text{FPdim}(\mathcal{C}_{\text{pt}})} \mid \prod_{j=0}^m \text{FPdim}(d_j)^2$$

Proof. It follows that $\text{FPdim}(\mathcal{C}^j) = d_j^2$ since \mathcal{C} is modular. □

For a semisimple Hopf algebra H recall the notion of commutator left coideal subalgebra H' , see [Bur12]. In this case Harada's identity can be written as:

Corollary 3.3. *For a semisimple Hopf algebra one has*

$$\left(\prod_{j=0}^m \frac{C_j}{\text{FPdim}(\mathcal{C}^j)}\right)^2 = \Lambda_{H'} = \frac{1}{\text{dim}(H')} \left(\sum_{\{j \mid \mathcal{C}^j \subseteq H'\}} C_j \right)$$

4. PROOF OF THEOREM 1.3 AND SOME OTHER APPLICATIONS

Suppose that \mathcal{C} is a weakly-integral modular category. As usually we suppose that \mathcal{C} is of type $(d_1, n_1; d_2, n_2; \dots, d_r, n_r)$ with $d_1=1$. We will use the notation $\text{FPdim}(\mathcal{C}) = \text{FPdim}(\mathcal{C})_1 \text{FPdim}(\mathcal{C})_{\geq 2}$ where $\text{FPdim}(\mathcal{C})_1$ is the free square part of $\text{FPdim}(\mathcal{C})$. Thus $\text{FPdim}(\mathcal{C})_1$ is the unique square free divisor of $\text{FPdim}(\mathcal{C})$ such that $\text{g.c.d}(\text{FPdim}(\mathcal{C})_1, \text{FPdim}(\mathcal{C})_{\geq 2}) = 1$. Note that $\text{FPdim}(\mathcal{C})_1 \mid n_1$ by [BP23, Corollary 1.17].

Let $\text{Irr}(\mathcal{C})$ denote the set of isomorphism classes of simple object of \mathcal{C} . For any subset $\mathcal{A} \subseteq \text{Irr}(\mathcal{C})$ we denote $\text{FPdim}(\mathcal{A}) := \sum_{X \in \mathcal{A}} \text{FPdim}(X)^2$.

Remark 4.1. *if \mathcal{C} is an integral modular category then for any $d_i > 1$ one clearly has $\text{gcd}(d_i, \text{FPdim}(\mathcal{C})_1) = 1$ since $d_i^2 \mid \text{FPdim}(\mathcal{C})$ by [ENO11, Theorem 2.11].*

4.1. Proof of Theorem 1.3. Note that Theorem 1.3 can be reformulated as

Theorem 4.2. *In any modular category \mathcal{C} one has $\text{FPdim}(\mathcal{C})_1 \mid n_i$ for all $i \geq 1$.*

Proof. Recall that for any $a > 0$ by $\text{Irr}_a(\mathcal{C})$ is denoted the set of isomorphism classes of simple objects of \mathcal{C} of dimension a .

Let $G(\mathcal{C})$ be the set of invertible objects of \mathcal{C} . Consider the action of $G(\mathcal{C})$ on $\text{Irr}(\mathcal{C})$ and note that each set $\text{Irr}_{d_i}(\mathcal{C})$ is a union of orbits of this action. By [ENO05, Remark 8.17] one has that $|G(\mathcal{C})| \mid \text{FPdim}(\text{Irr}_{d_i}(\mathcal{C})) = n_i d_i^2$.

Thus $\text{FPdim}(\mathcal{C})_1 \mid n_i d_i^2$ and since $(d_i, \text{FPdim}(\mathcal{C})_1) = 1$ for all i with $d_i > 1$ it follows that $\text{FPdim}(\mathcal{C})_1 \mid n_i$. \square

Proposition 4.3. *The first divisibility $d \mid n_1$ is true if $(d, \text{FPdim}(V)^2) = 1$ for any simple V as in [OY23].*

Proof. By [BP23, Theorem 1.15] one has that $\mathcal{V}(\mathcal{C}) = \mathcal{V}(\mathcal{C}_{\text{pt}}) \cup \mathcal{V}(d_i^2)$. On the other hand note that if $p \mid d$ since $(p, d_i^2) = 1$ it follows that $p \mid \text{FPdim}(\mathcal{C}_{\text{pt}})$. \square

Remark 4.4. *Note that by [Nat14, Thm 7. 4] any modular fusion category \mathcal{C} with $\text{FPdim}(\mathcal{C}) = p^a q^b d$ with d a square free number is WGT. Thus by [OY23][Theorem 4.2] these fusion categories decompose as a Deligne product as above.*

4.2. Proof of Theorem 1.4. Our proof goes as follows. By [OY23, Theorem 4.2], it suffices to show that \mathcal{C} is weakly group-theoretical. If $d \neq 1$ then by Theorem 1.3 it follows that $d \mid n_i$ for all $i > 1$. Thus $n_i = dn'_i$ and Frobenius-Perron dimension of \mathcal{C} can be written as

$$(pqr)^2 = \sum_{i=0}^r n'_i d_i^2.$$

Then one can do something similar as in the proof of from [ENO11, Lemma 9.3] to deduce that \mathcal{C} has a symmetric fusion subcategory. If the subcategory \mathcal{C}_{pt} is degenerate, then its Mueger center is a symmetric fusion subcategory. Otherwise $\mathcal{C} \simeq \mathcal{C}_{pt} \boxtimes \mathcal{C}_{ad}$ and one can repeat the argument for \mathcal{C}_{ad} .

4.3. Nilpotent fusion categories. Note that by Theorem 1.11 and Theorem 1.12 of [BP23] all nilpotent fusion categories with $K(\mathcal{C})$ commutative are both Burnside and dual Burnside. Moreover they are weakly-integral as iterated extensions of Vec. We also prove the following result for nilpotent fusion categories:

Proposition 4.5. *If \mathcal{C} is a non-pointed integral nilpotent fusion category with $K(\mathcal{C})$ commutative then*

- (1) $\text{FPdim}(\mathcal{C}_{ad})$ has no square free parts.
- (2) Moreover $\text{FPdim}(\mathcal{C})_1 \mid |U(\mathcal{C})|$ where $U(\mathcal{C})$ is the universal grading group of the fusion category \mathcal{C} .

Proof. By [BP23, Theorem 1.15] one has $\mathcal{V}(\mathcal{C}_{ad}) = \cup_{i=0}^m \mathcal{V}(d_i)$. For any prime p it follows that if $p \mid \text{FPdim}(\mathcal{C}_{ad})$ then $p \mid d_i$. By [GN08, Corollary 5.3] then $p^2 \mid d_i^2 \mid \text{FPdim}(\mathcal{C}_{ad})$ which shows that $\text{FPdim}(\mathcal{C}_{ad})$ has no square free part. On the other hand, since $\text{FPdim}(\mathcal{C}) = |U(\mathcal{C})| \text{FPdim}(\mathcal{C}_{ad})$ it follows that $\text{FPdim}(\mathcal{C})_1 \mid |U(\mathcal{C})|$. \square

REFERENCES

- [BP23] S. Burciu and S. Palcoux. Burnside type results for fusion rings. *arXiv:2302.07604*, 2023.
- [Bur12] S. Burciu. Normal coideal subalgebras of semisimple Hopf algebras. *Journal of Physics: Conference Series*, page 012004, 2012.
- [Bur20] S. Burciu. Conjugacy classes and centralizers for pivotal fusion categories. *Monatshefte für Mathematik*, 193(2):13–46, 2020.
- [Bur21] S. Burciu. Structure constants for premodular categories. *Bull. Lond. Math. Soc.*, 53(3):777–791, 2021.
- [Bur23] S. Burciu. On the Galois symmetries for the character table of an integral fusion category. *J. Algebra Appl.*, 22(1):Paper No. 2350026, 26, 2023.
- [CP22] A. Czenky and J. Plavnik. On odd-dimensional modular tensor categories. *Algebra Number Theory*, 16(8):1919–1939, 2022.
- [DMNO13] A. Davydov, M. Müger, D. Nikshych, and V. Ostrik. The Witt group of non-degenerate braided fusion categories. *Journal für die reine und angewandte Mathematik.*, 677:135–177, 2013.

- [EGNO15] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik. *Tensor categories*, volume 205. Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2015.
- [ENO05] P. Etingof, D. Nikshych, and V. Ostrik. On fusion categories. *Ann. of Math. (2)*, 162(2):581–642, 2005.
- [ENO11] P. Etingof, D. Nikshych, and V. Ostrik. Weakly group-theoretical and solvable fusion categories. *Adv. Math.*, 226(1):176–205, 2011.
- [GN08] S. Gelaki and D. Nikshych. Nilpotent fusion categories. *Adv. in Math.*, 217(3):1053–1071, 2008.
- [Har79] D. K. Harrison. Double coset and orbit spaces. *Pacific J. of Math.*, 80(2):451–491, 1979.
- [Har08] K. Harada. On a theorem of Brauer and Wielandt. *Proceedings of the American Mathematical Society*, 136(11):3825–3829, 2008.
- [LPR22] Z. Liu, S. Palcoux, and Y. Ren. Classification of Grothendieck rings of complex fusion categories of multiplicity one up to rank six. *Lett Math Phys*, 112(3):Paper No. 54, 2022.
- [Nat14] S. Natale. On weakly group-theoretical non-degenerate braided fusion categories. *J. Noncommut. Geom.*, 8:1043–1060, 2014.
- [OY23] V. Ostrik and Z. Yu. On the minimal extension and structure of braided weakly group-theoretical fusion categories. *Adv. Math.*, 419:108961, 2023.
- [Shi17] K. Shimizu. The monoidal center and the character algebra. *J. Pure Appl. Alg.*, 221(9):2338–2371, 2017.

INST. OF MATH. “SIMION STOILOW” OF THE ROMANIAN ACADEMY P.O. BOX 1-764,
RO-014700, BUCHAREST, ROMANIA

Email address: `sebastian.burciu@imar.ro`