# Group irregularity strength of disconnected graphs

Sylwia Cichacz, Barbara Krupińska<sup>∗</sup> AGH University of Krakow, Poland

June 22, 2023

#### Abstract

We investigate the group irregular strength  $(s_g(G))$  of graphs, i.e the smallest value of s such that for any Abelian group Γ of order s exists a function  $g: E(G) \to \Gamma$  such that sums of edge labels at every vertex is distinct. We give results for bound and exact values of  $(s_q(G))$  for graphs without small stars as components.

### 1 Introduction

One of the first facts we can observe in any simple graph  $G$  is that there are at least two vertices with equal amounts of adjacent edges (of the same degree). However, if we consider edge labeling  $f: E(G) \to \{1...s\}$  and calculate the weight of each vertex v as the sum of labels on the adjacent edges, then it is possible that every vertex could have a different weight. A labeling f that produces a unique weighted degree for every vertex  $v \in V(G)$  is called *irregular*. The maximum label s is called the *strength* of the labeling. *Irregularity strength of*  $G$ , denoted by  $s(G)$ , is the minimum value of s allowing some irregular labeling.

Chartrand  $[4]$  was the first to introduce the problem of finding  $s(G)$  and many authors investigated this problem further. In general it is known that  $s(G) \leq n-1$  for any graph G of order n containing no isolated edges and at most one isolated vertex, with the exception of the graph  $K_3$ , as proved by Aigner and Triesch [\[16\]](#page-6-0) and Nierhoff [\[5\]](#page-5-1). Though the family of stars witnesses the tightness of this upper bound, this result can be improved for graphs with large enough minimum degree  $\delta$ . The best general upper bound was proven by Kalkowski [\[6\]](#page-5-2) and states that  $s(G) \leq 6n/\delta + 6$ . This result was improved recently by Przybyło and Wei, who proved that for any  $\varepsilon \in (0, 0.25)$  there exist absolute constants  $c_1, c_2$  such that for all graphs G on n vertices with minimum degree  $\delta \geq 1$  and without isolated edges,  $s(G) \leq n/\delta(1 + c_1/\delta^{\varepsilon}) + c_2$  [\[17\]](#page-6-1).

Assume  $\Gamma$  is an Abelian group of order  $m \geq n$  with the operation denoted by  $+$  and the identity element by 0. To denote  $a + a + \ldots + a$ , where element a appears k times, we will use ka, we will write  $-a$  to denote the inverse of a, and instead of  $a + (-b)$  we will use  $a - b$ . Furthermore the notation  $\sum$  $\sum_{a \in S} a$  will be used as a shortened version of  $a_1 + a_2 + a_3 + \ldots$ , where  $a_1, a_2, a_3, \ldots$  are elements of the S set.

In this paper we will focus on edge labeling  $f : E(G) \to \Gamma$  which implies the weighted degrees defined as the sum in Γ:

$$
w(v) = \sum_{u \in N(v)} f(uv).
$$

The labeling f is called  $\Gamma$ -irregular if for all  $v, u \in V(G)$   $w(v) \neq w(u)$ . Furthermore the *group irregularity strength* of G, which we denote as  $s_q(G)$ , is the smallest integer s such that for every Abelian group  $\Gamma$  of order s there exists a  $\Gamma$ -irregular labeling f of G. If we consider the Abelian group to be only a cyclic group  $Z_k$ , then we have a *modular edge-graceful labeling* which was introduced by Jones in [\[10,](#page-5-3) [11,](#page-5-4) [12\]](#page-5-5). He described *modular edge-gracefulness of graphs* as the smallest integer

<sup>∗</sup>This work was partially supported by the Faculty of Applied Mathematics AGH UST statutory tasks within subsidy of Ministry of Science and Higher Education.

 $k(G) = k \geq n$  for which there exists an edge labeling  $f: E(G) \to \mathbb{Z}_k$  such that the induced vertex labeling  $g: V(G) \to \mathbb{Z}_k$  defined as

$$
g(u) = \sum_{v \in N(u)} f(uv) \bmod k
$$

is one-to-one.

We can easily deduct the correlation between  $k(G)$  and  $s_q(G)$ . Namely, for every graph G with no component of the order less than 3, there is  $k(G) \leq s_q(G)$ , [\[1\]](#page-5-6).

In a paper [\[2\]](#page-5-7) Anholcer, Cichacz and Milanic̆ have shown the following theorem that describes the value of  $s_q(G)$  for all connected graphs G of order  $n \geq 3$ .

<span id="page-1-0"></span>**Theorem 1.1** ([\[2\]](#page-5-7)). Let G be an arbitrary connected graph of order  $n \geq 3$ . Then

$$
s_g(G) = \begin{cases} n+2 & \text{if } G \cong K_{1,3^{2q+1}-2} \text{ for some integer } q \geq 1 \\ n+1 & \text{if } n \equiv 2 (mod 4) \land G \ncong K_{1,3^{2q+1}-2} \text{ for any integer } q \geq 1 \\ n & \text{otherwise} \end{cases}
$$

It is easy to see that to distinguish all  $n$  vertices in an arbitrary graph of that order we need at least n different element of Γ. Although the following lemma shows that an Abelian group of order  $n$ is not always enough to have a  $\Gamma$ -irregular labeling of  $G$ .

<span id="page-1-3"></span>**Lemma 1.2** ([\[1\]](#page-5-6)). *Let* G *be a graph of order* n. If  $n \equiv 2 \pmod{4}$ , then there is no Γ*-irregular labeling* G *for any Abelian group* Γ *of order* n*.*

So far the best upper bound for any graph was given by Anholcer, Cichacz, and Przybyło in [\[3\]](#page-5-8)

<span id="page-1-2"></span>Corollary 1.3 ([\[3\]](#page-5-8)). *Let* G *be an arbitrary graph of order* n *having no component of the order less than 3. Then*  $s_q(G) \leq 2n$ .

The following theorem [\[1\]](#page-5-6) by Anholcer and Cichacz determines the value of  $k(G)$  for an arbitrary graph G without any components of the order less than 3 or isomorphic to a star of even order.

<span id="page-1-1"></span>Theorem 1.4 ([\[1\]](#page-5-6)). *Let* G *be a graph of order* n *having neither a component of the order less than 3 nor a*  $K_{1,2u+1}$  *component for any integer*  $u \geq 1$ *. Then* 

$$
k(G) = n \quad \text{if} \quad n \equiv 1 \pmod{2}
$$
  
\n
$$
k(G) = n + 1 \quad \text{if} \quad n \equiv 2 \pmod{4}
$$
  
\n
$$
k(G) \le n + 1 \quad \text{if} \quad n \equiv 0 \pmod{4}.
$$

In the literature is also considered irregular labeling of digraphs. Cichacz and Tuza showed that if n is large enough with respect to an arbitrarily fixed  $\varepsilon > 0$  then  $\overline{G}$  has a Γ-irregular labeling for any Γ such that  $|\Gamma| > (1 + \varepsilon)n$  [\[14\]](#page-5-9). This bound was recently improved for digraphs with no weakly connected components of cardinality less than 4. Namely, it was shown that they have a Γ-irregular labeling for every Γ such that  $|\Gamma| \geq n+6$  [\[15\]](#page-5-10). Therefore on one hand we could expect that there exists a constant K such that any graph of order  $n$  without components of the order less than 3 has a Γ-irregular labeling for any group Γ of order at least  $n+K$ . On the other hand, we see in Theorem [1.1](#page-1-0) that some even stars can cause problems.

In this paper we will generalize Theorem [1.4](#page-1-1) for all Abelian groups, hence we show that such a constant exists (and is equal to 1) if a graph  $G$  does not have even stars as components. Moreover, we will improve the bound given in Corollary [1.3.](#page-1-2) We will show that if Γ has order at least  $n+K(q_0)$ , where  $q_0$  is the number of components being even stars in G, then G has a Γ-irregular labeling. However, this bound is not better than those given in Corollary [1.3](#page-1-2) for G being a union of  $K_{1,3}$ , it is much better for graphs such that the number of even stars is small compared to the order of G. In particular, we proved, that if G is a graph of order  $n$  having neither a component of the order less than 3 nor a component being a star  $K_{1,1+2u}$  for  $u \in \{1,2\}$ , then  $s(G) \leq n+3$ .

## 2 Proof of the main result

In the proof of the main result of this paper, we will use the construction of partition of  $\Gamma \setminus \{0\}$  into triplets and pairs, which was introduced in [\[13\]](#page-5-11). We call triplets  $T_i = \{a_i, b_i, c_i\}$  and  $T_j = \{a_j, b_j, c_j\}$ *complementary* if  $-a_i = a_j, -b_i = b_j$  and  $-c_i = c_j$ . We will denote the  $T_j$  triplet as  $-T_i$ . Let  $\Gamma$ be an Abelian group of odd order  $n = 6m + s$  ( $s = 1, 3$  or 5). A partition of  $\Gamma \setminus \{0\}$  into m pairs of complementary triplets  $T_1, -T_1, \ldots, T_m, -T_m$  and  $\frac{s-1}{2}$  sets  $\{d_j, -d_j\}$ , where  $\sum_{a \in T_i}$  $a = 0$ , is called a

<span id="page-2-0"></span>*Skolem partition* of  $\{0\}$ .

Theorem 2.1 ([\[13\]](#page-5-11)). *Every Abelian group* Γ *of odd order admits a Skolem partition.*

For any two given vertices  $v_1$  and  $v_2$  from the same connected component of a graph G exist walks from v<sup>1</sup> to v2. If a walk has an even number of vertices then we are going to call it an *even walk*. Otherwise, it is called an *odd walk*. In this proof, we will consider the shortest odd or even walk.

While labeling edges of a graph G we will start with 0 on all of them. Next, we will choose  $v_1$  and  $v_2$  and modify all of the edges of a chosen walk from  $v_1$  to  $v_2$  by adding some element of an Abelian group Γ. Roughly speaking we will be adding some element  $a \in \Gamma$  to all the labels of the edges on an odd position on the walk, starting from  $v_1$ . However, to the rest of the labels of edges on the walk we will add  $-a$ . This modification of labels we will denote with  $\phi_e(v_1, v_2) = a$  if we are considering the shortest even walk and  $\phi_o(v_1, v_2) = a$  otherwise. It is easy to see that  $\phi_e(v_1, v_2) = a$  adds a to a weighted degree of both  $v_1$  and  $v_2$ , while  $\phi_o(v_1, v_2) = a$  increases the weighted degree of  $v_1$  by a and decreases the weighted degree of  $v_2$  by a. We should also note that in both cases the operation does not modify the weighted degree of any other vertices of the walk.

We will need the following lemma:

<span id="page-2-1"></span>Lemma 2.2. *Let* G *be a graph of order* n *having neither a component of the order less than 3 nor a*  $K_{1,2u+1}$ ,  $u \geq 1$ *. Let*  $\Gamma$  *be an Abelian group of order*  $t \geq n$ *. Let*  $q_1$  *be the number of bipartite components of* G *with both color classes odd and* q<sup>2</sup> *be the number of components of odd order. Let* S *be a subsets of elements of*  $\Gamma$  *such that*  $S$  *contains*  $2m = 2q_1 + 2\left\lfloor \frac{q_2}{2} \right\rfloor$  *complementary triples*  $T_1, T_2, \ldots, T_{2m-1}, T_{2m}$  $\text{such that } T_{2i} = -T_{2i-1} \text{ for } i = 1, 2, \ldots, m \text{ and } k = \left[\frac{n-2m}{2}\right] \text{ zero-sum pairs } D_1, D_2, \ldots, D_k, \text{ and the}$ *identity element* {0}*.*

*Then there exists a*  $\Gamma$ *-irregular labeling*  $f: E(G) \to \Gamma$  *such that*  $w(v) \in S$  *for any*  $v \in V(G)$ *.* 

**Proof:** This proof is based on the proof of Lemma 2.5 [\[1\]](#page-5-6). Let us denote the elements of a triplet  $T_i$  by  $x_i, y_i$  and  $z_i$  for  $i = 1, 2, \ldots, 2m$ . If  $q_2$  is odd, then let  $T_{2m+1} = \{0, d_k, -d_k\}$ . If a component is a bipartite graph with both color classes even then we divide the vertices of every class into pairs  $(v_i^1, v_i^2)$ . Next we put

$$
\phi_o(v_i^1, v_i^2) = d_i
$$

for every such pair. Therefore  $w(v_i^1) = d_i$  and  $w(v_i^2) = -d_i$ . If a component is not bipartite but is even, then we label it analogically. When a component is a bipartite graph with both color classes odd, then we can be sure that there are at least 3 vertices in both classes. Let us choose three of them from each class and denote them with  $a_i, b_i$  and  $c_{i+1}$  in one of the classes and in the second with  $a_{i+1}, b_{i+1}$ and  $c_i$ . Than we can put

$$
\phi_e(a_i, c_i) = y_i,
$$
  
\n
$$
\phi_e(b_i, c_i) = z_i,
$$
  
\n
$$
\phi_e(a_{i+1}, c_{i+1}) = y_{i+1},
$$
  
\n
$$
\phi_e(b_{i+1}, c_{i+1}) = z_{i+1}.
$$

 $\sqrt{2}$ 

Thus  $w(a_{2j-1}) = y_{2j-1}$ ,  $w(b_{2j-1}) = z_{2j-1}$ ,  $w(c_{2j-1}) = -(y_{2j-1} + z_{2j-1}) = -x_{2j-1}$ ,  $w(a_{2j}) = y_{2j} =$  $-y_{2j-1}, w(b_{2j}) = z_{2j} = -z_{2j-1}$  and  $w(c_{2j+1}) = x_{2j-1}$  for  $j = 1, 2, ..., q_1$ . The rest of the vertices we label in the same way as a bipartite component of both classes even. If a component is of odd order we choose three vertices  $a_i, b_i$  and  $c_i$ , when it is a bipartite component then  $c_i$  we choose from a color class of odd order, and  $a_i, b_i$  from the even class. Next we put

$$
\phi_e(a_i, c_i) = y_i,
$$

$$
\phi_e(b_i, c_i) = z_i.
$$

The rest of the vertices we label analogically to earlier cases.

It is easy to see that the weights of vertices are pairwise distinct and  $w(v) \in S$  for any  $v \in V(G)$ .

Note that now we easily obtain the following results:

Corollary 2.3. *Let* G *be a graph of order* n *having neither a component of the order less than 3 nor*  $a K_{1,2u+1}$ ,  $u \geq 1$ . Then for every odd integer  $t \geq n$  there exists a  $\Gamma$ -irregular labeling, where  $\Gamma$  is an *Abelian group of order* t*.*

**Proof:** Let  $q_1$  as the number of bipartite components of G with both color classes odd and  $q_2$  be the number of components of odd order. Let  $2m = 2q_1 + 2 \left\lfloor \frac{q_2}{2} \right\rfloor$  and  $k = \left\lfloor \frac{n-2m}{2} \right\rfloor$ .

There exists a Skolem partition of Γ by Theorem [2.1.](#page-2-0) It is easy to see that if a Skolem partition of Γ exists, then there is a partition of  $\Gamma \setminus \{0\}$  into complementary triples  $T_1, T_2, \ldots, T_{2m-1}, T_{2m}$  such that  $T_{2i} = -T_{2i-1}$  for  $i = 1, 2, ..., m$  and  $k' = \lfloor \frac{t-2m}{2} \rfloor \ge k$  zero-sum pairs  $D_1, D_2, ..., D_{k'}$ . Now we apply Lemma [2.2](#page-2-1) and we are done.

The above corollary and Lemma [1.2](#page-1-3) imply the following bound analogous to those from Theorem [1.4](#page-1-1) for cyclic groups.

<span id="page-3-0"></span>Theorem 2.4. *Let* G *be a graph of order* n *having neither a component of the order less than 3 nor a*  $K_{1,2u+1}$  *component for any integer*  $u \geq 1$ *. Then* 

$$
s_g(G) = n \quad \text{if} \quad n \equiv 1 \pmod{2}
$$
  
\n
$$
s_g(G) = n+1 \quad \text{if} \quad n \equiv 2 \pmod{4}
$$
  
\n
$$
s_g(G) \le n+1 \quad \text{if} \quad n \equiv 0 \pmod{4}.
$$

For graphs with star components, we will need now the following.

<span id="page-3-1"></span>**Lemma 2.5.** Let  $G$  be a graph of order n having no component of the order less than 3. Let  $q_0$  be the *number of components being stars of even order,*  $q_1$  *be the number of bipartite components of* G *with both color classes odd and*  $q_2$  *be the number of components of odd order. Let*  $q_4 = \left[\frac{n-6\left\lceil \frac{q_0}{2}\right\rceil - 6q_1 - 6\left\lfloor \frac{q_2}{2}\right\rfloor - 1}{2}\right]$ 2  $\overline{1}$ 

*if*  $q_0$  *and n are both odd and*  $q_4 = \left[\frac{n-3q_0-6q_1-6\frac{q_2}{2}}{2}\right]$ 2 *otherwise.* Let  $\varepsilon(q_0) = 3$  *for*  $q_0$  *odd and n even,* and  $\varepsilon(q_0) = 0$  *otherwise.* Let  $\Gamma$  *be an Abelian group of odd order*  $t \geq n + K(q_0, q_4)$ *, where* 

 $K(q_0, q_4) = \max \{\varepsilon(q_0), 5q_0 - 2q_4 + \varepsilon(q_0) - 1\}.$ 

*Then there exists a*  $\Gamma$ *-irregular labeling*  $f: E(G) \to \Gamma$ *.* 

**Proof:** Let  $m = 3q_1 + 3\left\lfloor \frac{q_2}{2} \right\rfloor$ ,  $m' = \left\lceil \frac{q_0}{2} \right\rceil$ . Let  $k' = \left\lfloor \frac{t-2m-6m'-1}{2} \right\rfloor$  if  $q_0$  and n are both odd,  $k' = \left\lfloor \frac{t-2m-6m'}{2} \right\rfloor \geq q_4$  otherwise. Observe that  $k' \geq 5m'-2$  for  $q_0$  even and  $k' \geq 5m'-4$  otherwise.

A Skolem partition of Γ by Theorem [2.1](#page-2-0) exists. It is easy to see that if a Skolem partition of Γ exists, then there is a partition of  $\Gamma \setminus \{0\}$  into complementary triples  $T_1, T_2, \ldots, T_{2m-1}, T_{2m}, T_{2m+1}, \ldots$  $T_{2m+2m'-1}, T_{2m+2m'}$ , (i.e.  $T_{2i} = -T_{2i-1}$  for  $i = 1, 2, ..., m + m'$ ) and zero-sum pairs  $D_1, D_2, ..., D_{k'}$ . Let us denote the elements of a triplet  $T_i$  by  $x_i, y_i$  and  $z_i$  for  $i = 1, 2, ..., 2m + 2m'$ , and the elements of a pair  $D_j$  by  $d_j$ ,  $-d_j$  for  $j = 1, 2, \ldots, k'$ .

Let  $K_{1,1+2u_1}, K_{1,1+2u_2},...,K_{1,1+2u_{q_0}}$  be stars of even order such that  $c_i$  is the central vertex and  $v_i^1, v_i^2, \ldots, v_i^{1+2u_i}$  are the leaves in  $K_{1,1+2u_i}$ . Let

$$
\phi_e(c_i, v_i^1) = x_{2m+i},
$$
  

$$
\phi_e(c_i, v_i^2) = y_{2m+i},
$$

The vertices  $v_i^1$  and  $v_i^2$  get the permanent weights  $w(v_i^1) = x_{2m+i}$ ,  $w(v_i^2) = y_{2m+i}$  resp., whereas we will change the weight of vertices  $c_i$  for  $i = 1, 2, \ldots, q_0$ .

Suppose first that  $q_0$  or n is even. Let

$$
S_1 = \bigcup_{i=1}^{k'} \{d_i, -d_i\} \cup \bigcup_{i=0}^{m'-1} \{z_{2m+i}, -z_{2m+i}\},\
$$

$$
W_1 = \bigcup_{i=0}^{m'-1} \{x_{2m+2i+1}, -x_{2m+2i+1}, y_{2m+2i+1}, -y_{2m+2i+1}\}.
$$

Note that  $w(v_i^1), w(v_i^2) \in W_1$ . During the process of labeling in each step the set  $S_i$  will be the set with available labels and the set  $W_i$  is the set of occupied labels (i.e. the permanent weights). In the  $(2i + 1)$ th step for for  $i = 0, 1, \ldots, m' - 1$  we will pick an element  $g \in S_{2i+1}$  in such a way that  $g - z_{2m+i+1} \notin W_{2i+1} \cup \{0, -g\}.$  Note that  $|W_{2i+1}| = 4m' + 4i$  and  $|S_{2i+1}| = 2k' + 2m' - 4i.$  Since Γ has an odd order there exists exactly one element  $x ∈ Γ$  such that  $x - z_{2m+i+1} = -x$ . Moreover because  $k' > 5m' - 3$ , there is  $|W_{2i+1}| + 2 < |S_{2i+1}|$  and we can always choose such element  $g \in S_{2i+1}$ for  $i = 0, 1, \ldots, m' - 1$ . Then we put  $\phi_e(c_i, v_i^2) = g$  and  $\phi_e(c_{i+1}, v_{i+1}^2) = -g$  and we change the set of available labels

$$
S_{2(i+1)+1} = S_{2i+1} \setminus \{g, -g, g - z_{2m+i+1}, -g + z_{2m+i+1}\}
$$

and the set of permanent weights

$$
W_{2(i+1)+1} = W_{2i+1} \cup \{g, -g, g - z_{2m+i+1}, -g + z_{2m+i+1}\}.
$$

We got the permanent weights  $w(c_{i+1}) = g - z_{2m+i+1}$ ,  $w(v_{i+1}^3) = g$  and  $w(c_{i+2}) = -g + z_{2m+i+1}$ ,  $w(v_{i+2}^3) = -g$  for  $i = 0, 1, ..., m' - 1$ . One can easily see that  $w(c_{i+2}), w(v_{i+2}^3) \notin W_{2i+1} \cup \{0, g\}$ .

Observe that the set  $S_{2m'-1}$  contains at least  $t - 2m - 8m'$  elements which form zero-sum pairs. The remaining elements of stars  $K_{1,1+2u_i}$  we divide into pairs  $(v_i^{2j}, v_i^{2j+1})$  for  $j = 2, 3, ..., u_i$ . Next we put

$$
\phi_o(v_i^{2j}, v_i^{2j+1}) = g \in S_{2m'-1}
$$

for every such pair. Therefore  $w(v_i^1) = g$  and  $w(v_i^2) = -g$  from the set  $S_{2m'-1}$ . Finally, we are left with the set that has at least  $n-2m-2q_0-\sum_{i=1}^{q_0} 2u_i$  elements that form zero-sum pairs, moreover, we have 2m complementary triplets  $T_1, T_2, \ldots, T_{2m-1}, T_{2m}$ . Thus for the remaining components of the graph G we apply Lemma [2.2](#page-2-1) and we are done.

Suppose now that  $q_0$  is odd. For a star  $K_{1,1+2u_1}$  put  $\phi_e(c_1, v_1^1) = -x_{2m+1}, \phi_e(c_1, v_1^2) = y_{2m+1}$ , and  $\phi_e(c_1, v_1^3) = 0$ . Let

$$
S_3 = \bigcup_{i=1}^{k'} \{d_i, -d_i\} \cup \bigcup_{i=1}^{m'-1} \{z_{2m+i}, -z_{2m+i}\},
$$
  

$$
W_3 = \bigcup_{i=0}^{m'-1} \{x_{2m+2i+1}, -x_{2m+2i+1}, y_{2m+2i+1}, -y_{2m+2i+1}\} \cup \{z_{2m+1}, -z_{2m+1}\}.
$$

If additionally, n is odd, then  $q_2$  is odd. Let C be a component of the odd order. As in the proof of Lemma  $2.2$ , we can label edges of  $C$  in such a way that there exist three vertices with weights  $x_{2m+1}, y_{2m+1}$  and  $-z_{2m+1}$ , whereas all the remaining vertices in C we can divide into pairs that each pair is connected by an even walk. Since now  $4m' + 4i = |W_{2i+1}| + 2 < |S_{2i+1}| = 2k' + 2m' - 4i + 2$ for  $i = 0, 1, \ldots, m' - 1$ , thus we proceed now analogously as above and we are done.

From the above lemma, we obtain immediately the following:

Theorem 2.6. *Let* G *be a graph of order* n *having neither a component of the order less than 3 nor a*  $K_{1,1+2u}$  *component for*  $u \in \{1,2\}$ *. Then* 

$$
s_g(G) = n \t if \t n \equiv 1 \pmod{2}
$$
  
\n
$$
s_g(G) \le n+3 \t if \t n \equiv 0 \pmod{2}.
$$

**Proof:** If there is no  $K_{1,1+2u}$  for  $u \geq 1$ , then we are done by Theorem [2.4.](#page-3-0) Therefore we can assume that  $q_0 > 0$  and apply Lemma [2.5.](#page-3-1) Since  $u \geq 2$ , there is  $n \geq 8q_0 + 6q_1 + 3q_2$ , what implies that  $2q_4 \ge 5q_0 - 1$ , thus  $K(q_0, q_4) = \varepsilon(q_0)$ . If n is odd, then  $\varepsilon(q_0) = 0$  and  $\varepsilon(q_0) \le 3$  for n even.

Note that for some families of graphs containing small stars as components we can obtain better a bound than those in Theorem [1.3.](#page-1-2) Let  $tG$  denote the union of t disjoint copies of  $G$ , then if  $G = tK_{1,3} \cup lC_4$  $G = tK_{1,3} \cup lC_4$  $G = tK_{1,3} \cup lC_4$  and  $l \geq t$  we have  $s(G) \leq 4(t+l)+3$  by Lemma [2.5.](#page-3-1) Moreover in Figure 1 we show that  $s_q(2K_{1,3}) = 8$ . Therefore at the end of this section, we state the following conjecture.

Conjecture 2.7. *There exists a constant* K *such that any graph* G *of order* n *with no connected components of order less than* 3 *has a*  $\Gamma$ -*irregular labeling for every*  $\Gamma$  *such that*  $|\Gamma| \geq n + K$ *.* 



<span id="page-5-12"></span>Figure 1: Irregular labelings of  $2K_{1,3}$  in groups  $\mathbb{Z}_8$ ,  $\mathbb{Z}_2 \times \mathbb{Z}_4$  and  $\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_2$ , respectively.

# <span id="page-5-6"></span>References

- [1] Anholcer, M. & Cichacz, S. Group irregular labelings of disconnected graphs. *Contributions To Discrete Mathematics*. 12, 158-166 (2017)
- <span id="page-5-7"></span>[2] Anholcer, M., Cichacz, S. & Milanič, M. Group irregular strenght of connected graphs. *Journal Of Combinatorial Optimization*. 30 pp. 1-17 (2013)
- <span id="page-5-8"></span>[3] Anholcer, M., Cichacz, S. & Przybyło, J. Linear bound on nowhere-zero group irregularity strenght and nowhere-zero group sum chromatic number of graphs. *Applied Mathematics And Computation*. 343 pp. 149-155 (2019)
- <span id="page-5-0"></span>[4] Chartrand, G., Jacobson, M., Lehel, J., Oellermann, O., S.Ruiz & Saba, F. Irregular networks. *Congr. Numer.*. 64 pp. 187-192 (2017)
- <span id="page-5-1"></span>[5] Nierhoff, T. A tight bound on the irregularity strength of graphs. *SIAM J. Discr. Math*. 13, 313-323 (2000)
- <span id="page-5-2"></span>[6] Kalkowski, M., Karoński, M. & Pfender, F. A new upper bound for the irregularity strength of graphs. *SIAM J. Discr. Math*. 25 pp. 1319-1321 (2011)
- [7] Majerski, P. & Przybyło, J. On the irregularity strength of dense graphs. *SIAM J. Discr. Math*. 28, 197-205 (2014)
- [8] Lo, S. On edge-graceful labelings of graphs. *Cong. Numer.*. 50 pp. 231-241 (1985)
- [9] Jones, R. Modular and graceful edge colorings of graphs. *Ph.D. Thesis, Western Michigan University*. (2011)
- <span id="page-5-4"></span><span id="page-5-3"></span>[10] Jones, R. Modular and Graceful Edge Colorings of Graphs. *Ph.D. Thesis*. (2011)
- [11] Fuije-Okamoto, F., Jones, R., Kolasinski, K. & Zhang, P. On modular edge-graceful graphs. *Graphs Combin.*. 29 pp. 901-912 (2012)
- <span id="page-5-5"></span>[12] Jones, R. & Zhamg, P. Nowhere-zero modular edge-graceful graphs. *Discuss. Math. Graph Theory*. 32 pp. 487-505 (2012)
- <span id="page-5-11"></span>[13] Tannenbaum, P. Partition of abelian groups into sets with zero sums. *Congressus Numerantium*. 33 pp. 341-348 (1981)
- <span id="page-5-9"></span>[14] Cichacz, S. & Tuza, Z. Realization of digraphs in Abelian groups and its consequences. *J. Graph Theory*. pp. 1-15 (2021)
- <span id="page-5-10"></span>[15] Cichacz, S. S. Cichacz, Irregular labeling on Abelian groups of digraphs. *Preprint ArXiv:2303.08712 [math.CO]* . (2023)
- <span id="page-6-0"></span>[16] Aigner, M. & E., T. Irregular assignments of trees and forests. *SIAM J. Discrete Math.* . 3(4) pp. 439-449 (1990)
- <span id="page-6-1"></span>[17] Przybyło, J. & Wei, F. Irregular labeling on Abelian groups of digraphs. *Preprint ArXiv:2109.04317v2 [math.CO], To Appear In Combinatorica* . (2021)
- [18] Bača, M., Muthugurupackiam, K., Kathiresan, K. & Ramya, S. Modular irregularity strength of graphs. *Electron. J. Graph Theory Appl.*. 8 pp. 435-443 (2020)