

# On Two problems of Defective Choosability of Graphs

Jie Ma <sup>\*</sup>      Rongxing Xu <sup>†</sup>      Xuding Zhu <sup>‡</sup>

June 26, 2023

## Abstract

Given positive integers  $p \geq k$ , and a non-negative integer  $d$ , we say a graph  $G$  is  $(k, d, p)$ -choosable if for every list assignment  $L$  with  $|L(v)| \geq k$  for each  $v \in V(G)$  and  $|\bigcup_{v \in V(G)} L(v)| \leq p$ , there exists an  $L$ -coloring of  $G$  such that each monochromatic subgraph has maximum degree at most  $d$ . In particular,  $(k, 0, k)$ -choosable means  $k$ -colorable,  $(k, 0, +\infty)$ -choosable means  $k$ -choosable and  $(k, d, +\infty)$ -choosable means  $d$ -defective  $k$ -choosable. This paper proves that there are 1-defective 3-choosable planar graphs that are not 4-choosable, and for any positive integers  $\ell \geq k \geq 3$ , and non-negative integer  $d$ , there are  $(k, d, \ell)$ -choosable graphs that are not  $(k, d, \ell + 1)$ -choosable. These results answer questions asked by Wang and Xu [SIAM J. Discrete Math. 27, 4(2013), 2020-2037], and Kang [J. Graph Theory 73, 3(2013), 342-353], respectively. Our construction of  $(k, d, \ell)$ -choosable but not  $(k, d, \ell + 1)$ -choosable graphs generalizes the construction of Král' and Sgall in [J. Graph Theory 49, 3(2005), 177-186] for the case  $d = 0$ .

## 1 Introduction

A coloring of a graph  $G$  is a mapping  $\phi$  which assigns to each vertex  $v$  a color. The *defect* of a vertex  $v$ , denoted by  $\lambda_G(v, \phi)$ , is the number of neighbors of  $v$  which have the same color as  $v$ . A coloring  $\phi$  is  $d$ -defective if  $\lambda_G(v, \phi) \leq d$  for each vertex  $v \in V(G)$ . A 0-defective coloring is also called a proper coloring.

Assume  $G$  is a graph and  $f : V(G) \rightarrow \mathbb{N}^+$  is a mapping. An  $f$ -list assignment of  $G$  is a list assignment  $L$  of  $G$  which assigns to each vertex  $v$  a set  $L(v)$  of  $f(v)$  colors. Given a

---

<sup>\*</sup>School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, China. E-mail: jiema@ustc.edu.cn. Supported in part by the National Key R and D Program of China 2020YFA0713100, National Natural Science Foundation of China grants 12125106, Innovation Program for Quantum Science and Technology 2021ZD0302904, and Anhui Initiative in Quantum Information Technologies grant AHY150200.

<sup>†</sup>School of Mathematical Sciences, University of Science and Technology of China, Hefei, Anhui, 230026, China and School of Mathematical Sciences, Zhejiang Normal University, Jinhua, Zhejiang, 321000, China. E-mail: xurongxing@ustc.edu.cn. Supported by Anhui Initiative in Quantum Information Technologies grant AHY150200. Supported also by NSFC grant 11871439.

<sup>‡</sup>School of Mathematical Sciences, Zhejiang Normal University, Jinhua, Zhejiang, 321000, China. E-mail: xdzhu@zjnu.edu.cn. Grant Number: NSFC 11971438, U20A2068.

list assignment  $L$  and nonnegative integer  $d$ , a  $d$ -defective  $L$ -coloring  $\phi$  of  $G$  is a  $d$ -defective coloring  $\phi$  of  $G$  such that  $\phi(v) \in L(v)$  for each vertex  $v$ . We say  $G$  is  $d$ -defective  $f$ -choosable if  $G$  has a  $d$ -defective  $L$ -coloring for any  $f$ -list assignment  $L$ .

Given a mapping  $f : V(G) \rightarrow \mathbb{N}^+$  and two nonnegative integers  $d, p$ , we say  $G$  is  $(f, d, p)$ -choosable if for any  $f$ -list assignment  $L$  with  $|\bigcup_{v \in V(G)} L(v)| \leq p$ , there exists a  $d$ -defective  $L$ -coloring of  $G$ . In particular,  $(k, 0, k)$ -choosable is called  $k$ -colorable,  $(k, 0, +\infty)$ -choosable is called  $k$ -choosable,  $(k, d, k)$ -choosable is called  $d$ -defective  $k$ -colorable,  $(k, d, +\infty)$ -choosable is called  $d$ -defective  $k$ -choosable.

Defective coloring of planar graphs was first studied by Cowen, Cowen and Woodall [1]. They proved that every outerplanar graph is 2-defective 2-colorable and that every planar graph is 2-defective 3-colorable. These results were strengthened to defective list coloring by Eaton and Hull [2] and Škrekovski [12] independently. They proved that every planar graph is 2-defective 3-choosable and every outerplanar graph is 2-defective 2-choosable.

These results motivated some problems on the relation among defective colorability, defective choosability and choosability of planar graphs. It is known that there are 4-choosable planar graphs that are not 1-defective 3-colorable [13] (hence not 1-defective 3-choosable). Wang and Xu [13] asked the following.

**Question 1** *Is every 1-defective 3-choosable graph 4-choosable?*

This paper gives a negative answer to this question in a stronger form as following.

**Theorem 2** *There are 1-defective 3-choosable planar graphs that are 4-choosable.*

By definition, for any  $d \geq 0$ , if  $G$  is  $(k, d, +\infty)$ -choosable, then it is  $(k, d, p)$ -choosable for any  $p \geq k$ . Král' and Sgall [7] showed that for each  $\ell \geq k \geq 3$ , there exists a  $(k, 0, \ell)$ -choosable graph which is not  $(k, 0, \ell + 1)$ -choosable. Kang [6] asked the following question.


**Question 3** *Given positive integers  $k, d$ , does there exist an integer  $\ell_{k,d}$  such that every  $(k, d, \ell_{k,d})$ -choosable graph is  $(k, d, +\infty)$ -choosable?*

Our second result answers this question in negative for  $k \geq 3$ , which generalizes the construction of Král' and Sgall [7] to the cases of  $d \geq 0$ .

**Theorem 4** *For any integers  $d \geq 0$  and  $\ell \geq k \geq 3$ , there exists a  $(k, d, \ell)$ -choosable graph which is not  $(k, d, \ell + 1)$ -choosable.*

## 2 Proof of Theorem 2

The gadget graph  $T$  depicted in Figure 1 was constructed by Gutner [4] (see Fig. 1) and used in the construction of many counterexamples for several topics related to list coloring of planar graphs [9, 10, 11, 14].



**Fig. 1.** The gadget graph  $T$  in [4].

For a positive integer  $k$ , let  $T(k)$  be the graph obtained from the disjoint union of  $k$  copies of  $T$  by identifying all the copies of top vertex, denoted still by  $u$ , and identifying all the copies of the bottom vertex, denoted still by  $v$ . It is known [4, 11] that for any  $k \geq 16$ ,  $T(k)$  is not 4-choosable.

To prove Theorem 2, it suffices to show that if  $k \leq 26$ , then  $T(k)$  is 1-defective 3-choosable.

We first construct some 1-defective 3-colorings for some special list assignments of  $T$ . We assume  $T$  are labeled as in Fig.1.

**Lemma 5** *Let  $H = T - \{u, v\}$ , and  $f(w) = 2$  for  $w \in V(H) \setminus \{x, y, z\}$  and  $f(x), f(y), f(z) \geq 1$ . If one of  $f(x), f(y), f(z) = 2$ , then  $H$  is 1-defective  $f$ -choosable.*

**Proof.** Assume  $f(y) = f(z) = 1$  and  $f(x) = 2$  and  $L$  is an  $f$ -list assignment of  $H$ . Let  $\phi(y), \phi(z)$  be the unique color in  $L(y), L(z)$ , respectively. Let  $\phi(u_2) \in L(u_2) - \{\phi(y)\}$ ,  $\phi(v_2) \in L(v_2) - \{\phi(z)\}$ ,  $\phi(u_1) \in L(u_1) - \{\phi(z)\}$ ,  $\phi(v_1) \in L(v_1) - \{\phi(z)\}$  and  $\phi(x) \in L(x) - \{\phi(u_1)\}$ . It is straightforward to verify that  $\phi$  is a 1-defective  $L$ -coloring of  $H$ .

Assume  $f(z) = 2$  and  $f(x) = f(y) = 1$  and  $L$  is an  $f$ -list assignment of  $H$ . Let  $\phi(x), \phi(y)$  be the unique color in  $L(x), L(y)$ , respectively. Let  $\phi(u_1) \in L(u_1) - \{\phi(x)\}$ ,  $\phi(z) \in L(z) - \{\phi(u_1)\}$ ,  $\phi(v_1) \in L(v_1) - \{\phi(z)\}$ ,  $\phi(u_2) \in L(u_2) - \{\phi(z)\}$  and  $\phi(v_2) \in L(v_2) - \{\phi(y)\}$ . Again it is straightforward to verify that  $\phi$  is a 1-defective  $L$ -coloring of  $H$ . ■

**Corollary 6** *Let  $L$  be a list assignment of  $T$  with  $L(u) = \{\alpha\}$ ,  $L(v) = \{\beta\}$  and  $|L(w)| \geq 3$  for  $w \in V(T) \setminus \{u, v\}$ . If  $\alpha = \beta$ , or  $\alpha \neq \beta$  and  $\{\alpha, \beta\} \not\subseteq L(x) \cap L(y) \cap L(z)$ , or  $\alpha \neq \beta$  and  $L(x) \cap L(y) \cap L(z) - \{\alpha, \beta\} \neq \emptyset$ , then  $T$  has a 1-defective  $L$ -coloring  $\phi$  such that  $\lambda_T(u, \phi) = \lambda_T(v, \phi) = 0$ .*

**Proof.** Let  $L'$  be the list assignment of  $H = T - \{u, v\}$  defined as  $L'(w) = L(w) - \{\alpha, \beta\}$  if  $w \in \{x, y, z\}$ , and  $L'(w) = L(w) - \{\alpha\}$  for  $w \in \{u_1, u_2\}$  and  $L'(w) = L(w) - \{\beta\}$  for  $w \in \{v_1, v_2\}$ . If  $\alpha = \beta$  or  $\{\alpha, \beta\} \not\subseteq L(x) \cap L(y) \cap L(z)$ , then it follows from Lemma 5 that  $H$  has a 1-defective  $L'$ -coloring  $\phi$ . Extend  $\phi$  to  $T$  by letting  $\phi(u) = \alpha$  and  $\phi(v) = \beta$ , then  $\phi$  is a 1-defective  $L$ -coloring  $\phi$  such that  $\lambda_T(u, \phi) = \lambda_T(v, \phi) = 0$ .

If  $\alpha \neq \beta$  and  $L(x) \cap L(y) \cap L(z) - \{\alpha, \beta\} \neq \emptyset$ , say  $c \in L(x) \cap L(y) \cap L(z) - \{\alpha, \beta\}$ , then let  $\phi(x) = \phi(y) = \phi(z) = c$ ,  $\phi(w) \in L(w) - \{\alpha, c\}$  for  $w \in \{u_1, u_2\}$  and  $\phi(w) \in L(w) - \{\beta, c\}$  for  $w \in \{v_1, v_2\}$ . It is straightforward to verify that  $\phi$  is a 1-defective  $L$ -coloring  $\phi$  such that  $\lambda_T(u, \phi) = \lambda_T(v, \phi) = 0$ . ■

**Lemma 7** *Let  $L$  be a list assignment of  $T$  with  $L(u) = \{\alpha\}$ ,  $L(v) = \{\beta\}$  and  $|L(w)| \geq 3$  for  $w \in V(T) \setminus \{u, v\}$ . Then  $T$  has a 1-defective  $L$ -coloring  $\phi$  such that  $\lambda_T(u, \phi) = 0$ , and a 1-defective  $L$ -coloring  $\phi$  such that  $\lambda_T(v, \phi) = 0$ .*

**Proof.** By Corollary 6, it suffices to consider the case that  $\alpha \neq \beta$  and  $\{\alpha, \beta\} \subseteq L(x) \cap L(y) \cap L(z)$ . Let  $\phi(u) = \alpha$ ,  $\phi(v) = \beta$ ,  $\phi(x), \phi(z)$  be the unique color in  $L(x) - \{\alpha, \beta\}, L(z) - \{\alpha, \beta\}$ , respectively. Let  $\phi(u_1) \in L(u_1) - \{\alpha, \phi(x)\}, \phi(v_1) \in L(v_1) - \{\beta, \phi(z)\}, \phi(u_2) \in L(u_2) - \{\alpha, \phi(z)\}, \phi(v_2) \in L(v_2) - \{\beta, \phi(z)\}$ . If  $\phi(u_2) \neq \phi(v_2)$ , then let  $\phi(y) \in L(y) - \{\phi(u_2), \phi(v_2)\}$ . Otherwise, let  $\phi(y)$  be any color in  $\{\alpha, \beta\}$ . It is easy to verify that  $\phi$  is a 1-defective coloring of  $T$ . In most cases,  $\lambda_T(u, \phi) = \lambda_T(v, \phi) = 0$ , except that in the last case, if  $\phi(y) = \alpha$ , then  $\lambda_T(v, \phi) = 0$ , if  $\phi(y) = \beta$ , then  $\lambda_T(u, \phi) = 0$ . ■

Now we are ready to prove Theorem 2.

**Proof of Theorem 2.** Let  $L$  be any 3-list assignment of  $T(k)$ , we shall give a 1-defective 3-coloring  $\phi$  of  $T(k)$ . Without loss of generality, we assume that  $|L(w)| = 3$  for each vertex  $w$ . If  $L(u) \cap L(v) \neq \emptyset$ , say  $\alpha \in L(u) \cap L(v)$ , then let  $\phi(u) = \phi(v) = \alpha$ . By Corollary 6,  $\phi$  can be extended to a 1-defective coloring of  $T(k)$  so that  $\lambda_T(u, \phi) = \lambda_T(v, \phi) = 0$ .

Assume  $L(u) \cap L(v) = \emptyset$ . For each pair of colors  $(\alpha, \beta)$  with  $\alpha \in L(u), \beta \in L(v)$ , a copy of  $T$  is called  $(\alpha, \beta)$ -tight if  $\{\alpha, \beta\} = L(x) \cap L(y) \cap L(z)$  (here  $x, y, z$  refer to the corresponding vertices in that copy of  $T$ ). Note that if a copy of  $T$  is  $(\alpha, \beta)$ -tight, then it is not  $(\alpha', \beta')$ -tight for any  $(\alpha', \beta') \neq (\alpha, \beta)$ . So by pigeonhole principle, there is a pair  $(\alpha, \beta)$  with  $\alpha \in L(u), \beta \in L(v)$  such that there are at most two copies of  $T$ , say  $T^1$  and  $T^2$ , that are  $(\alpha, \beta)$ -tight. Let  $\phi(u) = \alpha, \phi(v) = \beta$ . By Corollary 6, for each other copy  $T'$  of  $T$ ,  $\phi$  can be extended to a 1-defective  $L$ -coloring of  $T'$  so that  $\lambda_{T'}(u, \phi) = \lambda_{T'}(v, \phi) = 0$ . By Lemma 7,  $\phi$  can be extended to a 1-defective  $L$ -coloring of  $T^1$  so that  $\lambda_{T^1}(u, \phi) = 0$ , and  $\phi$  can be extended to a 1-defective  $L$ -coloring of  $T^2$  so that  $\lambda_{T^2}(v, \phi) = 0$ . Thus  $\phi$  can be extended to a 1-defective  $L$ -coloring of  $T(k)$ . ■

### 3 Proof of Theorem 4

Let  $\ell \geq k \geq 3$  and  $d \geq 0$  be any fixed integers. In this section we prove Theorem 4 by constructing graphs that are  $(k, d, \ell)$ -choosable but not  $(k, d, \ell + 1)$ -choosable.

Lovász [8] proved that every graph  $G$  with  $\Delta(G) + 1 \leq k(d + 1)$  is  $d$ -defective  $k$ -colorable. This was generalized to list-version by Hendrey and Wood (see Corollary 3.2 in [5]).

**Lemma 8 ([5])** *Every graph  $G$  with  $\Delta(G) + 1 \leq k(d + 1)$  is  $d$ -defective  $k$ -choosable.*

Lemma 8 implies that every complete graph with  $k(d+1)$  vertices is  $d$ -defective  $k$ -choosable. In this paper, we need the following slightly stronger statement.

**Lemma 9** *Suppose  $G$  is a complete graph of order  $k(d+1)$ , and  $v$  is any vertex in  $G$ . Assume  $L$  is a list assignment with  $|L(u)| \geq k$  for any  $u \in V(G) \setminus \{v\}$  and  $|L(v)| = 1$ . Then  $G$  has a  $d$ -defective  $L$ -coloring.*

**Proof.** Assume  $L(v) = \{c\}$ . Let  $V_c = \{u \in V(G) : c \in L(u)\}$ . If  $|V_c| \geq d$ , then we choose arbitrary  $d$ -subset  $D \subset V_c$ , and color all the vertices in  $D$  and  $v$  with color  $c$ , also we delete the color  $c$  from  $V_c - D$ . Note that  $G - v - D$  is a complete graph of order  $(k - 1)(d + 1)$ ,

and each vertex has at least  $(k - 1)$  colors available. By Lemma 8, we can color  $G - v - D$  without using  $c$  and with defect  $d$  by colors from the lists. So assume that  $|V_c| \leq d - 1$ . Then we color all the vertices in  $V_c$  with color  $c$ . Note that  $G' = G - V_c$  still satisfies that  $\Delta(G') + 1 \leq k(d + 1)$  and each vertex has  $k$  colors available. Then we can extend the coloring to whole  $G$  by Lemma 8. ■


The following lemma follows from Lemma 3.3 in [5].

**Lemma 10** ([5]) *Let  $L$  be a  $k$ -list assignment of a graph  $G$ . Let  $A, B$  be a partition of  $V(G)$ , where  $G[A]$  is  $d$ -defective  $L$ -colorable. If for every vertex  $v \in B$ ,*

$$(d + 1)|N_G(v) \cap A| + \deg_B(v) + 1 \leq (d + 1)k,$$

*then  $G$  is  $d$ -defective  $L$ -colorable.*

Given a graph  $G$  and an non-negative integer  $d$ , we denote by  $G * d$  the graph obtained from the disjoint union of  $G$  and  $|V(G)|$  copies of the complete graph  $K_d$ , denoted as  $\{B_v : v \in V(G)\}$ , by identifying  $v$  with one vertex of  $B_v$ . Fig. 2 shows the graph  $C_9 * 4$ .



**Fig. 2.**  $C_9 * 4$

**Lemma 11**  $H = C_{2k+1} * (2d + 2)$  is not  $d$ -defective 2-colorable.

**Proof.** Let  $v_1, v_2, \dots, v_{2k+1}$  be the vertices of  $C_{2k+1}$  in order, and  $B_1, B_2, \dots, B_{2k+1}$  be the corresponding vertex-set of the  $2k + 1$  copies of  $K_{2d+2}$ . Assume there is a  $d$ -defective 2-coloring  $\phi$  of  $H$ . Since  $B_i$  is a clique of order  $2(d + 1)$ , each of the two colors is used  $d + 1$  times by  $\phi$  in  $B_i$ . Therefore,  $\phi(v_i) \neq \phi(v_j)$  whenever  $v_i v_j \in E(C_{2k+1})$ . This is a contradiction, as  $C_{2k+1}$  is not 2-colorable. ■

**Lemma 12** *Let  $H = C_{2k+1} * (2d + 2)$ , with  $v_1, v_2, \dots, v_{2k+1}$  be the vertices of  $C_{2k+1}$  in this cyclic order. Let  $B_1, B_2, \dots, B_{2k+1}$  be the corresponding vertex-set of the  $2k + 1$  copies of  $K_{2d+2}$ . Assume  $L$  is a list assignment of  $H$  with  $|L(u)| \geq 2$  for  $u \in \bigcup_{i=1}^{2k} B_i$  and  $|L(u)| \geq 3$  for  $u \in B_{2k+1}$ . Then for any color  $c$ ,  $H$  has a  $d$ -defective  $L$ -coloring  $\phi$  such that at most one vertex in  $B_{2k+1}$  is colored with  $c$  in  $\phi$ .*

**Proof.** As  $|L(v_{2k+1})| \geq 3$ , there is a proper  $L$ -coloring  $\phi$  of  $C_{2k+1}$ . We then extend  $\phi$  from each  $v_i$  to  $B_i$  by Lemma 9 so that color  $c$  is not used on other vertices of  $B_{2k+1}$ . This is possible since each vertex in  $B_{2k+1}$  still has at least two colors available. ■

**Lemma 13** *Assume  $k \geq 3$ ,  $t \geq 2$ ,  $d \geq 0$  and  $\ell = k - 2 + t$ . There exists a graph  $H(t, d, k) = (V, E)$  with a precolored independent set  $T = \{u_1, u_2, \dots, u_t\}$  for which the following hold:*

- *Assume the precoloring  $\phi$  of  $T$  uses  $t$  distinct colors in  $[\ell + 1]$ . Then there is a  $k$ -list assignment  $L$  of  $H(t, d, k)$  with  $L(v) \subseteq [\ell + 1]$  for each vertex  $v$  such that  $\phi$  cannot be extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, k)$  with  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$  for each  $u_i \in T$ . On the other hand, if  $d \geq 1$ , then for any  $k$ -list assignment  $L$  of  $H(t, d, k) - T$ ,  $\phi$  can be extended to a  $d$ -defective  $L$ -coloring  $\psi$  of  $H(t, d, k)$  such that  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$  for  $i = 1, 2, \dots, t - 1$  and  $\lambda_{H(t, d, k)}(u_t, \psi) \leq 1$ .*
- *Assume the precoloring  $\phi$  of  $T$  uses with at most  $t - 1$  colors. Then for any  $k$ -list assignment  $L$  of  $H(t, d, k) - T$ ,  $\phi$  can be extended to a  $d$ -defective  $L$ -coloring  $\psi$  of  $H(t, d, k)$  such that  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$  for each  $u_i \in T$ .*

**Proof.** We consider three cases according to the values of  $\ell$  and  $k$ .

**Case 1.**  $t \leq k - 1$  (hence  $3 \leq k \leq \ell \leq 2k - 3$ ).

Let  $H(t, d, k)$  be the join of a complete graph  $K_{(d+1)(k-t)+1}$  and an independent set  $T = \{u_1, u_2, \dots, u_t\}$ . Assume  $\phi$  is a precoloring of  $T$ .

Assume  $\phi$  uses  $t$  distinct colors in  $[\ell + 1]$ . Let  $S$  be  $k$  colors in  $[\ell + 1]$  that contains  $\phi(T)$ , and let  $L(v) = S$  for all  $v \in V(H(t, d, k) - T)$ .

We then show that  $\phi$  cannot be extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, k)$  with  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$ . Assume to the contrary that  $\phi$  is extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, k)$  with  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$ . Then vertices in the complete graph  $K_{(d+1)(k-t)+1}$  are colored by  $k - t$  colors. So one color class contains at least  $d + 2$  vertices, which is a contradiction. On the other hand, if  $d \geq 1$ , then for any  $k$ -list assignment  $L$  of  $H(t, d, k) - T$ , let  $w$  be an arbitrary vertex in  $V(H(t, d, k) - T)$ ,  $L'(v) = L(v) - \phi(T)$  for  $v \in V(H(t, d, k) - T) - \{w\}$ , and  $L'(w) = L(w) - \phi(T - \{u_t\})$ . We have  $|L'(v)| \geq k - t$  for any  $v \in V(K_{(d+1)(k-t)+1}) - \{w\}$ , and  $|L'(w)| \geq k - t + 1$ . By Lemma 8,  $K_{(d+1)(k-t)+1}$  has a  $d$ -defective  $L'$ -coloring, which is an extension of  $\phi$  to a  $d$ -defective  $L$ -coloring of  $H(t, d, k)$  with  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$  for each  $u_i \in T - \{u_t\}$ , and  $\lambda_{H(t, d, k)}(u_t, \psi) \leq 1$ .


If  $\phi$  use at most  $t - 1$  colors, then for any  $k$ -list assignment  $L$  of  $V(H(t, d, k)) - T$ , let  $L'(v) = L(v) - \phi(T)$  for  $v \in V(H(t, d, k)) - T$ . We have  $|L'(v)| \geq k - t + 1$  for any  $v \in V(K_{(d+1)(k-t)+1})$ . By Lemma 8,  $K_{(d+1)(k-t)+1}$  has a  $d$ -defective  $L'$ -coloring, which is an extension of  $\phi$  to a  $d$ -defective  $L$ -coloring of  $H(t, d, k)$  with  $\lambda_{H(t, d, k)}(u_i, \psi) = 0$  for each  $u_i \in T$ .

**Case 2.**  $t \geq k = 3$  (hence  $\ell \geq 2k - 2 = 4$ ).

Let  $T = \{u_1, u_2, \dots, u_t\}$  be an independent set. Let  $s \in \{\binom{t}{2}, \binom{t}{2} + 1\}$  be an odd integer, and let  $\mathcal{X} = \{X_a : 1 \leq a \leq s\}$  be a set disjoint copies of  $K_{d+1}$ . Let  $\pi : [s] \rightarrow \{(i, j) : 1 \leq i < j \leq t\}$  be a surjective map, and for each  $a \in [s]$  with  $\pi(a) = (i, j)$ , connect each vertex of  $X_a$  to  $u_i$  and  $u_j$ . Add the graph  $H = C_s * 2(d + 1)$ , with  $\{B_a : 1 \leq a \leq s\}$  be the corresponding vertex-set of the  $s$  copies of  $K_{2d+2}$ . For each  $a \in [s]$ , choose one vertex  $x_a \in X_a$  and connect  $x_a$  to all the vertices in  $B_a$ . This completes the construction of  $H(t, d, 3)$ . Fig. 3 is an example with  $t = 4$  and  $d = 3$ .

Assume  $\phi$  is a precoloring of  $T$ .

Assume  $\phi$  uses  $t$  distinct colors in  $[\ell + 1]$ . Let  $L$  be a 3-list assignment of  $H(t, d, 3) - T$  defined as follows: For  $x \in X_a$  with  $a \in [s]$  and  $\pi(a) = (i, j)$ , let  $L(x) = \{\phi(u_i), \phi(u_j), \beta\}$  and



**Fig. 3.**  $H(4, 3, 3)$ ; The green vertices are  $x_i$ s and red vertices are  $v_i$ s in the copy of  $C_7 * 8$ . We used the pair  $(u_3, u_4)$  twice as  $\binom{4}{2}$  is even.

$L(y) = \{\phi(u_1), \phi(u_2), \beta\}$  for  $y \in V(H)$ , where  $\beta \in [\ell+1] \setminus \phi(T)$  is a fixed color. We shall show that  $\phi$  cannot be extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, 3)$  with  $\lambda_{H(t, d, 3)}(u_i, \psi) = 0$  for  $i = 1, 2, \dots, t$ . Otherwise, assume  $\phi$  is extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, 3)$  with  $\lambda_{H(t, d, 3)}(u_i, \psi) = 0$ . Then vertices in  $X_a$  are all colored by  $\beta$  for each  $a \in [s]$ , and hence vertices in  $H$  are colored with  $\phi(u_1)$  and  $\phi(u_2)$ . This is in contrary to Lemma 11.

On the other hand, for any  $k$ -list assignment  $L$  of  $H(t, d, 3) - T$ , let  $L'$  be the list assignment of  $H$  obtained from  $L$  by deleting all the colors used in  $\phi$  on  $\{x_1, x_2, \dots, x_{s-1}\}$ . It is clear that

$$|L'(w)| \geq \begin{cases} 2, & \text{if } w \in \bigcup_{i=1}^{s-1} B_i, \\ 3, & \text{if } w \in B_s. \end{cases}$$

Therefore, by Lemma 12, we can extend the  $\phi$  to a  $d$ -defective coloring of  $H$ , such that at most one vertex in  $B_s$  uses  $\phi(x_s)$ , which yields a  $d$ -defective  $L$ -coloring as desired.

Assume  $\phi$  uses at most  $t - 1$  distinct colors. Then at least two vertices  $u_i, u_j$  in  $T$  received the same color. Without loss of generality, assume  $\pi(s) = (i, j)$ . Then we extend  $\phi$  to  $V(H(t, d, 3)) \setminus V(H)$  by assigning  $\phi(w) \in L(w) \setminus \phi(N_{H(t, d, 3)}(w)) \cap T$  for each vertex  $w \in \bigcup_{i=1}^s V(X_i)$  such that  $w$  has defect at most  $d$  if  $w \neq x_s$  and at most  $d - 1$  if  $w = x_s$ , this is possible as each  $X_i$  has exactly  $d + 1$  vertices and  $x_s$  has two colors available.

Finally, we need to extend  $\phi$  to  $H$ . Let  $L'$  be the list assignment of  $H$  defined as

$$L'(v) = \begin{cases} L(v) - \{\phi(x_i) : i = 1, 2, \dots, s\}, & \text{if } v \in V(H) \setminus B_s, \\ L(v) - \{\phi(x_i) : i = 1, 2, \dots, s - 1\}, & \text{if } v \in B_s. \end{cases}$$

Then  $|L'(u)| \geq 2$  for  $u \in V(H) \setminus B_s$ , and  $|L'(u)| \geq 3$  if  $u \in B_s$ . Since  $d \geq 1$ , then by Lemma 12, we can extend  $\phi$  to a  $d$ -defective  $L'$ -coloring of  $H$  such that at most one vertex in  $B_s$  is colored with  $\phi(x_s)$ , which yields a desired coloring.

**Case 3.**  $t \geq k > 3$  (hence  $\ell \geq 2k - 2 \geq 6$ ).

Let  $T = \{u_1, u_2, \dots, u_t\}$  be an independent set. Let  $s \in \{\binom{t}{k-2}, \binom{t}{k-2} + 1\}$  be an odd integer. Denote by  $\binom{[t]}{k-2}$  the set of  $(k - 2)$ -subsets of  $[t] = \{1, 2, \dots, t\}$ . Let  $H$  be a copy of  $C_s * (2d + 2)$ ,  $B_1, B_2, \dots, B_s$  be the corresponding vertex-set of the  $s$  copies of  $K_{2d+2}$ . Let



$\pi : [s] \rightarrow \binom{[t]}{k-2}$  be a surjective map. For  $a \in [s]$ , connect each vertex in  $B_a$  to  $\{u_i : i \in \pi(a)\}$ . This completes the construction of  $H(t, d, k)$ .

Suppose  $\phi$  is a precoloring of  $T$  with  $t$  distinct colors in  $[\ell + 1]$ . For each  $B_i$ , let  $T_i \subset T$  be the neighbors of vertices of  $B_i$ . Let  $L$  be a  $k$ -list assignment for vertices in  $H$ , such that for any  $w \in B_i$ ,  $L(w) = \bigcup_{v \in T_i} \{\phi(v)\} \cup \{\beta_1, \beta_2\}$ , where  $\beta_1, \beta_2 \in [\ell + 1] \setminus \phi(T)$  (note that  $\ell + 1 - t \geq 2$ ). If  $\phi$  can be extended to a  $d$ -defective coloring  $\psi$  of  $H(t, d, k)$  with  $\lambda(u_i, \psi) = 0$ , then vertices in  $H$  can be only colored with two colors  $\{\beta_1, \beta_2\}$ . But this is not possible by Lemma 11.

On the other hand, if  $d \geq 1$ , for any  $k$ -list assignment  $L$  of  $H(t, d, 3) - T$ , let  $L'$  be the list assignment of  $H$  obtained from  $L$  as follows (without loss of generality, we assume that  $t \in \pi(s)$ ):

$$L'(w) = \begin{cases} L(w) \setminus \phi(T), & \text{if } w \in \bigcup_{i=1}^{s-1} B_i, \\ L(w) \setminus \phi(T - u_t), & \text{if } w \in B_s. \end{cases}$$

It is clear that  $|L'(w)| \geq 2$  if  $w \in \bigcup_{i=1}^{s-1} B_i$  and  $|L'(w)| \geq 3$  if  $w \in B_s$ . Therefore, by Lemma 12, we can extend the  $\phi$  to a  $d$ -defective coloring of  $H$ , such that at most one vertex in  $B_s$  uses  $\phi(x_s)$ , which yields a  $d$ -defective  $L$ -coloring as desired.

Assume that  $\phi$  is a precoloring of  $T$  with  $t - 1$  distinct colors, and  $L$  is a  $k$ -list assignment of  $H$ . It follows that at least two vertices in  $T$  received same color, say  $u_1$  and  $u_2$ . Without loss of generality, assume that  $\{1, 2\} \subset \pi(s)$ . Let  $L'$  be the list assignment of  $H$  obtained from  $L$  by deleting all the colors used in  $\phi$  on  $T$ . It is clear that  $|L(w)| \geq 2$  and moreover,  $|L(w)| \geq 3$  if  $w \in B_s$ . So we can extend the precoloring to  $H$  by Lemma 12 as desired. ■

Now we are ready to prove Theorem 4.

**Proof of Theorem 4.** Assume  $\ell \geq k \geq 3$  and  $d \geq 0$ . We shall construct a graph which is  $(k, d, \ell)$ -choosable but not  $(k, d, \ell + 1)$ -choosable.

Let  $t = \ell + 2 - k \geq 2$ . Let  $G_1 = K_{k(d+1)}$ , and  $z \in V(G_1)$ . Take  $k$  disjoint copies of  $K_{(k-1)(d+1)}$ , which are denoted as  $C_1, C_2, \dots, C_k$ . Connect all the vertices of  $C_1 \cup \dots \cup C_k$  to the vertex  $z$ . The resulting graph is  $G_2$ .

Let  $q = \binom{(k-1)(d+1)}{k-1}$ . For  $i = 1, 2, \dots, k$ , let  $X_{i,1}, X_{i,2}, \dots, X_{i,q}$  be the family of all the  $(k-1)$ -subsets of  $V(C_i)$ . For  $1 \leq i \leq k, 1 \leq j \leq q$ , take  $t$  copies of disjoint complete graphs  $K_{d+1}$ , which are denoted by  $D_{i,j,1}, D_{i,j,2}, \dots, D_{i,j,t}$ . For  $s = 1, 2, \dots, t$ , connect all the vertices in  $D_{i,j,s}$  to all vertices in  $X_{i,j}$ . The resulting graph is  $G_3$ .

For  $1 \leq i \leq k, 1 \leq j \leq q$ , take a  $t$ -set  $T_{i,j} = \{u_{i,j,1}, u_{i,j,2}, \dots, u_{i,j,t}\}$  with  $u_{i,j,s} \in V(D_{i,j,s})$ , and build a copy  $H_{i,j}(t, d, k)$  of  $H(t, d, k)$  described in Lemma 13 with  $T = T_{i,j}$ . This completes the construction of the graph  $G$ . See Fig. 4 for illustration.


We first show that  $G$  is not  $(k, d, \ell + 1)$ -choosable. We shall construct a  $k$ -list assignment  $L$  of  $G$ , with  $L(v) \subseteq [\ell + 1]$  for each vertex  $v$ , such that  $G$  is not  $d$ -defective  $L$ -colorable.

For each vertex  $v$  in the subgraph  $G_2$ , let  $L(v) = \{1, 2, \dots, k\}$ . For  $i = 1, 2, \dots, k, 1 \leq j \leq q, 1 \leq s \leq t$ , let  $\theta_{i,j} : [t] \rightarrow \{i, k + 1, k + 2, \dots, \ell + 1\}$  be a bijection. For  $v \in D_{i,j,s}$ , let  $L(v) = ([k] \setminus \{i\}) \cup \{\theta_{i,j}(s)\}$ .

For  $1 \leq i \leq k$  and  $1 \leq j \leq q$ , for  $v \in V(H_{i,j}(t, d, k)) - T_{i,j}$ , let  $L(v) = L_{i,j}(v)$ , where  $L_{i,j}$  is a list assignment defined as follows:

Let  $\phi_{i,j}$  be the coloring of  $T_{i,j}$  defined as  $\phi_{i,j}(u_{i,j,s}) = \theta_{i,j}(s)$ . By Lemma 13, there is a  $k$ -list assignment  $L_{i,j}$  of  $H_{i,j}(t, d, k)$  with  $L_{i,j}(v) \subseteq [\ell + 1]$  such that  $\phi_{i,j}$  cannot be extended to a  $d$ -defective coloring  $\psi_{i,j}$  of  $H_{i,j}(t, d, k)$  with  $\lambda_{H_{i,j}(t, d, k)}(u_{i,j,s}, \psi_{i,j}) = 0$  for each  $u_{i,j,s} \in T_{i,j}$ .





**Fig. 4.** Illustration of the main construction

Now we prove that  $G$  is not  $d$ -defective  $L$ -colorable. Assume to the contrary that  $G$  has an  $L$ -coloring  $\psi$  with defect  $d$ . Assume  $\pi(z) = r$ . Since  $G_1$  is a complete graph with  $k(d+1)$  vertices, there must be  $d$  neighbors of  $z$  in  $G_1$  colored with  $r$ . Then the  $(k-1)(d+1)$  vertices of  $C_r$  are colored with  $\{1, 2, \dots, k\} \setminus \{r\}$ . Hence each of the  $k-1$  colors is used  $d+1$  times in  $\psi$ . Assume  $X_{r,p}$  is a  $(k-1)$ -subset of  $C_r$  that are colored by distinct colors from  $\{1, 2, \dots, k\} \setminus \{r\}$ . Note that each vertex  $x \in X_{r,p}$  has  $d$  neighbors in  $C_r$  that are colored the same color as  $x$ . As all vertices in  $D_{r,p,1}, D_{r,p,2}, \dots, D_{r,p,t}$  are adjacent to  $X_{r,p}$ . For each  $1 \leq s \leq t$ , we have  $\psi(v) = \theta_{r,p}(s)$  for each vertex  $v \in V(D_{r,p,s})$ . In particular,  $\psi(u_{r,p,s}) = \theta_{r,p}(s)$  and  $u_{r,p,s}$  has  $d$  neighbors in  $D_{r,p,s}$  that are colored the same color as  $u_{r,p,s}$ . This implies that  $\lambda_{H_{r,p}(t,d,k)}(u_{r,p,s}, \psi) = 0$  for  $s = 1, 2, \dots, t$ . This is a contradiction, as  $\psi(u_{r,p,s}) = \theta_{r,p}(s) = \phi_{r,p}(u_{r,p,s})$ , and by Lemma 13,  $\phi_{r,p}$  cannot be extended to a  $d$ -defective  $L$ -coloring of  $H_{r,p}(t, d, k)$  with  $\lambda_{H_{r,p}(t,d,k)}(u_{r,p,s}, \psi) = 0$  for  $s = 1, 2, \dots, t$ .

Lastly, we show that  $G$  is  $(k, d, \ell)$ -choosable. Assume  $L$  is any  $k$ -list assignment of  $G$  with  $L(v) \subseteq [\ell]$ . We shall show that  $G$  is  $d$ -defective  $L$ -colorable.

By Lemma 8, there is a  $d$ -defective  $L$ -coloring  $\phi$  of  $G_1$ . By Lemma 10,  $\phi$  can be extended to a  $d$ -defective  $L$ -coloring of  $G_2$  (by setting  $A = V(G_1)$  and  $B = V(G_2) \setminus V(G_1)$ ). Similarly, by Lemma 10, we can extend  $\phi$  to a  $d$ -defective  $L$ -coloring of  $G_3$ . Note that each vertex in  $D_{i,j,s}$  has  $(k-1)$  neighbors in  $C_i$  and  $d$  neighbors in  $D_{i,j,s}$ , so it also satisfies the inequality in Lemma 10 by setting  $A = V(G_2)$  and  $B = V(G_3) \setminus V(G_2)$ .

It remains to show that we can extend  $\phi$  to  $H_{i,j}(t, d, k)$  for  $1 \leq i \leq k, 1 \leq j \leq q$ .

First assume that for any  $u, v \in X_{i,j}$ ,  $\phi(u) \neq \phi(v)$  and for any  $x \in X_{i,j}$ ,  $\lambda_{G_3}(x, \phi) = d$ . In this case, there are only  $\ell - (k-1) = t - 1$  colors available for vertices in  $\bigcup_{s=1}^t V(D_{i,j,s})$ . So at least two of vertices in  $T_{i,j}$  received the same color. By Lemma 13, we can extend  $\phi$  to  $H_{i,j}(t, d, k)$ . In particular, if  $d = 0$ , then  $\phi$  is a 0-defective coloring, i.e., a proper coloring of  $G_3$ , implying that for any  $u, v \in X_{i,j}$ ,  $\phi(u) \neq \phi(v)$  and for any  $x \in X_{i,j}$ ,  $\lambda_{G_3}(x, \phi) = 0 = d$ . Hence  $\phi$  can be extended to  $H_{i,j}(t, d, k)$ .

Next assume that  $\phi(u) = \phi(v)$  for some  $u, v \in X_{i,j}$  or  $\lambda_{G_3}(x, \phi) \leq d - 1$  for some  $x \in X_{i,j}$ . In either case, it is enough to modify  $\phi$  on  $D_{i,j,t}$  (if necessary) such that  $\lambda_{G_3}(u_{i,j,t}, \phi) \leq d - 1$ . So it suffices to show that we can extend  $\phi$  from  $T$  to  $H_{i,j}(t, d, k)$  such that  $\lambda_{H_{i,j}(t,d,k)}(u_{i,j,s}) = 0$  for  $s = 1, 2, \dots, t - 1$  and  $\lambda_{H_{i,j}(t,d,k)}(u_{i,j,t}) \leq 1$ . By Lemma 13, this is possible. The proof of Theorem 4 now is completed. ■

## 4 Open problems

For  $d \geq 0$ , a graph is  $d$ -defective 1-choosable if and only if  $\Delta(G) \leq d$ , and which is equivalent to be  $d$ -defective 1-colorable. So Question 3 has a positive answer for  $k = 1$  and  $\ell_{1,d} = 1$ . For  $k = 2$  and  $d = 0$ , Král' and Sgall [7] showed that every  $(2, 0, 4)$ -choosable graph is  $(2, 0, +\infty)$ -choosable. The proof relies on a characterization of 2-choosable graphs by [3]. For  $d \geq 1$ , no characterization of  $d$ -defective 2-choosable graphs is known, and Question 3 remains open for  $k = 2$  and  $d \geq 1$ .

The following problem asked by Král' and Sgall [7] also remains open.

**Problem 14** *Is it true that for each  $k \geq 3$ , there exists a number  $\ell$  such that each  $(k, 0, \ell)$ -choosable graph is  $(k + 1)$ -choosable? If so, what is the least number  $\ell$  with this property?*

## References

- [1] L. J. Cowen, R. H. Cowen, and D. R. Woodall. Defective colorings of graphs in surfaces: partitions into subgraphs of bounded valency. *J. Graph Theory*, 10(2):187–195, 1986.
- [2] N. Eaton and T. Hull. Defective list colorings of planar graphs. *Bull. Inst. Combin. Appl.*, 25:79–87, 1999.
- [3] P. Erdős, A. L. Rubin, and H. Taylor. Choosability in graphs. In *Proceedings of the West Coast Conference on Combinatorics, Graph Theory and Computing (Humboldt State Univ., Arcata, Calif., 1979)*, Congress. Numer., XXVI, pages 125–157. Utilitas Math., Winnipeg, Man., 1980.
- [4] S. Gutner. The complexity of planar graph choosability. *Discrete Math.*, 159(1-3):119–130, 1996.
- [5] K. Hendrey and D. R. Wood. Defective and clustered choosability of sparse graphs. *Combin. Probab. Comput.*, 28(5):791–810, 2019.
- [6] R. J. Kang. Improper choosability and Property B. *J. Graph Theory*, 73(3):342–353, 2013.
- [7] D. Král' and J. Sgall. Coloring graphs from lists with bounded size of their union. *J. Graph Theory*, 49(3):177–186, 2005.
- [8] L. Lovász. On decomposition of graphs. *Studia Sci. Math. Hungar.*, 1:237–238, 1966.

- [9] E. Smith-Roberge. On the choosability with separation of planar graphs and its correspondence colouring analogue. *arXiv:2203.13348*, 2022.
- [10] M. Voigt. On list colourings and choosability of graphs. *Habilitationsschrift, Tu Ilmenau*, 1998.
- [11] M. Voigt and B. Wirth. On 3-colorable non-4-choosable planar graphs. *J. Graph Theory*, 24(3):233–235, 1997.
- [12] R. Škrekovski. List improper colourings of planar graphs. *Combin. Probab. Comput.*, 8(3):293–299, 1999.
- [13] Y. Wang and L. Xu. Improper choosability of planar graphs without 4-cycles. *SIAM J. Discrete Math.*, 27(4):2029–2037, 2013.
- [14] R. Xu and X. Zhu. The strong fractional choice number and the strong fractional paint number of graphs. *SIAM J. Discrete Math.*, 36(4):2585–2601, 2022.