Condition numbers for the Moore-Penrose inverse and the least squares problem involving rank-structured matrices

Sk. Safique Ahmad^a and Pinki Khatun^a

^aDepartment of Mathematics, Indian Institute of Technology Indore, Khandwa Road, Indore, 453552, Madhya Pradesh, India

ARTICLE HISTORY

Compiled August 6, 2024

Abstract

Perturbation theory plays a crucial role in sensitivity analysis, which is extensively used to assess the robustness of numerical techniques. To quantify the relative sensitivity of any problem, it becomes essential to investigate structured condition numbers (CNs) via componentwise perturbation theory. This paper addresses and analyzes structured mixed condition number (MCN) and componentwise condition number (CCN) for the Moore-Penrose (M-P) inverse and the minimum norm least squares (MNLS) solution involving rank-structured matrices, which include the Cauchy-Vandermonde (CV) matrices and $\{1, 1\}$ -quasiseparable (QS) matrices. A general framework has been developed to compute the upper bounds for MCN and CCN of rank deficient parameterized matrices. This framework leads to faster computation of upper bounds of structured CNs for CV and $\{1, 1\}$ -QS matrices. Furthermore, comparisons of obtained upper bounds are investigated theoretically and experimentally. In addition, the structured effective CNs for the M-P inverse and the MNLS solution of $\{1, 1\}$ -QS matrices are presented. Numerical tests reveal the reliability of the proposed upper bounds as well as demonstrate that the structured effective CNs are computationally less expensive and can be substantially smaller compared to the unstructured CNs.

KEYWORDS

Rank-structured matrices; Condition number; Moore-Penrose inverse; Minimum norm least squares solution; Cauchy-Vandermonde matrices; Quasiseparable matrices

AMS subject classification. 15A09; 15A12; 65F20; 65F35

1. Introduction

Condition numbers (CNs) are one of the most important tools in matrix computation since they are widely employed to investigate the stability and robustness of numerical algorithms; see [17]. It measures how sensitive, in the worst-case scenario, a problem is to a slight change in input data and is crucial in determining whether a numerical solution makes sense. On the other hand, the backward error is used to find a nearly perturbed problem with minimal magnitude perturbations so that the calculated solution becomes an actual solution to the perturbed problem. One can estimate the forward error of an approximate solution by combining the backward error with the CN (see [2] for more on CNs).

Most likely, for the first time, Rice [26] introduced the classical theory of CNs. In accordance with [26], the normwise CN, which has been extensively considered in the literature, quantifies the input and output data errors using the norms. The normwise CN has the drawback of ignoring the input and output data structures when the data is poorly scaled or sparse. Consequently, even minor relative normwise perturbations can have a disproportionate impact on entries that are small or zero, thereby potentially compromising data sparsity. To address

CONTACT Sk. Safique Ahmad Email: safique@iiti.ac.in; Pinki Khatun Email: pinki996.pk@gmail.com

this challenge, mixed condition number (MCN) and componentwise condition number (CCN) [27,28] emerge as a viable solution. MCN measures the input perturbations componentwise and the output error using norms, whereas CCN measures both the error in output data and the input perturbations componentwisely.

The Moore-Penrose (M-P) inverse holds a pivotal position in matrix computation, offering a generalization of the standard inverse for rectangular or rank deficient matrices $M \in \mathbb{R}^{m \times n}$. It is a unique matrix $Y \in \mathbb{R}^{n \times m}$ that satisfies the following equations: [29]

$$MYM = M, YMY = Y, (MY)^{\top} = MY, (YM)^{\top} = YM.$$

Typically, it is denoted by $Y = M^{\dagger}$. The M-P inverse finds its practical significance in solving the linear least squares (LS) problem, where the goal is to minimize the expression:

$$\min_{z \in \mathbb{R}^n} \|Mz - b\|_2,\tag{1}$$

where $M \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. Here, $\|\cdot\|_2$ denotes the spectral norm. If M is of full column rank, the LS problem has a unique solution given by $\mathbf{x} = M^{\dagger}b$. However, when M is rank deficient, the LS problem has infinitely many solutions, and the set of all solutions forms a closed and convex set. Consequently, this set uniquely yields the minimum norm LS (MNLS) solution \mathbf{x} , defined as $\mathbf{x} = M^{\dagger}b$. The M-P inverse and the LS problem have various applications in digital image restoration and reconstruction [4,5], Gauss–Markov model [25], and so on. The literature on CNs for the M-P inverse [10] and LS problems [37] is quite rich. The normwise CN for the M-P inverse and the LS problem is investigated in [10,15,22], while MCN and CCN are considered in [7,8]. For structured matrices, structured condition numbers for the M-P inverse and the LS problem have been investigated in [7,38], which involves the preservation of the inherent matrix structure within the perturbation matrices.

In the past few years, many fast algorithms have been developed for various problems involving rank-structured matrices, such as computing eigenvalues and singular values [32,39], solving linear systems [31] and LS problems [18], and computing the M-P inverse [3]. Quasiseparable (QS) [14], Cauchy [18], and Cauchy-Vandermonde (CV) [18] matrices are popular examples of rank-structured matrices that arise in many applications such as in boundary value problem [16,20], acoustic and electromagnetic scattering theory [6], interpolation problems [24], rational models of regression and E-optimal design [19], and so on.

One of the striking properties of the rank-structured matrices is that they can be parameterized by $\mathcal{O}(m+n)$ parameters rather than mn entries. Based on this property, many fast algorithms with lower computational costs have been developed [30–32]. Plenty of works involving rank-structured matrices have been done in recent years to investigate the structured CNs for eigenvalue problems [9,12], the solution of a linear system having a single as well as multiple right-hand sides [13,23], the Sylvester matrix equation [11], and so on, by considering perturbations on the parameters. Based on the above discussions, it is more sensible to investigate structured CNs by addressing perturbations on the parameters rather than directly on the matrix entries and to identify which set of parameters will be more suited for the development of fast algorithms. Thus, the forgoing discussion motivates us to consider perturbations on parameters instead of directly on entries in this paper.

In [37], authors have presented general parameterized QS representation and Givens-vector (GV) representation for the rectangular $m \times n$ ($m \ge n$) {1,1}-QS matrices (a special case of QS matrices), which are natural extensions of the square matrix case discussed in [12,13,30]. Then, the authors studied the structured MCN for the LS problems when the coefficient matrix is a full column rank $m \times n$ {1,1}-QS matrices. Explicit expressions are derived with respect to both QS and GV representations. Additionally, comparison results between obtained structured MCN are presented. Numerical experiments in [37] showed that the structured MCNs can be much smaller than the unstructured ones. However, the above investigations do not address the rank

deficient case. Furthermore, the MCN and CCN for the M-P inverse of rank deficient rankstructured matrices still need to be explored in the literature. Nevertheless, when dealing with rank deficient matrices, a prominent challenge in analyzing the CNs arises from the fact that even slight changes to the matrix can yield enormous variations in the computed M-P inverse. In light of this, normwise CNs for rank deficient unstructured matrices have been considered in [34,36], and for structured matrices in [38] under the assumptions: $\mathcal{R}(\Delta M) \subset \mathcal{R}(M)$ and $\mathcal{R}(\Delta M^{\top}) \subset \mathcal{R}(M^{\top})$, on the perturbation matrix ΔM in M, where $\mathcal{R}(M)$ denotes the range of M. Whereas, in [35], upper bounds are investigated for CCN for unstructured matrices under the above assumptions.

This paper's central aim is to study the structured MCN and CCN for the M-P inverse and the LS problem when dealing with rank deficient rank-structured matrices. This investigation adheres to the rank-preserving constraint, denoted as $\operatorname{rank}(M + \Delta M) = \operatorname{rank}(M)$, which encompasses a broader class of perturbation matrices than those constrained by $\mathcal{R}(\Delta M) \subset \mathcal{R}(M)$ and $\mathcal{R}(\Delta M^{\top}) \subset \mathcal{R}(M^{\top})$. This perspective expands the horizons of our study and offers valuable insights into structured CNs for this class of matrices.

The following highlights the main contributions of this paper:

- The MCN and CCN for two problems, the M-P inverse and the MNLS solution of the LS problem, involving rank deficient CV and $\{1, 1\}$ -QS matrices are considered under the broader rank condition, i.e., rank $(M + \Delta M) = \operatorname{rank}(M)$.
- By considering matrix entries to be differentiable functions of a set of real parameters, we develop a general framework to compute the upper bounds of the MCN and CCN of the M-P inverse and LS problem for rank deficient parameterized matrices. In addition, exact expressions in the full column rank case of the MCN and CCN are also obtained.
- For the CV and {1,1}-QS matrices, compact upper bounds are obtained for structured MCN and CCN. Two important parameter representations for {1,1}-QS matrices are considered: the QS representation and GV representation.
- For {1,1}-QS matrices, structured effective CNs are proposed and shown that they can reliably estimate the actual conditioning of these matrices. Numerical experiments are reported to demonstrate that structured CNs are significantly smaller compared to unstructured CNs and align consistently with the theoretical results.

The remaining part of this paper is structured as follows. Section 2 provides a few notations and preliminary results. In Section 3, for the M-P inverse and the MNLS solution, we develop expressions of upper bounds for MCN and CCN for a general class of parameterized matrices. These frameworks are utilized in Sections 4 and 5 to derive the bounds for structured MCN and CCN for CV and $\{1, 1\}$ -QS matrices. Further, Section 5 studies comparison results between different structured and unstructured CNs. In Section 6, numerical experiments are performed to illustrate our findings. Section 7 ends with conclusions and a line of future research.

2. Notation and preliminaries

The following notations are adopted throughout this work. $\mathbb{R}^{m \times n}$ denotes the collection of all $m \times n$ matrices and \mathbb{R}^m denotes the space of all column vectors of dimension m, where m, n are positive integers. To indicate the M-P inverse and transpose of any matrix $M \in \mathbb{R}^{m \times n}$, M^{\dagger} and M^{\top} are used, respectively. The *i*-th column of the identity matrix I_m of order m is denoted as e_i^m . For $M \in \mathbb{R}^{m \times n}$, set $\mathbf{E}_M := I_m - MM^{\dagger}$ and $\mathbf{F}_M := I_n - M^{\dagger}M$. Denote **0** as the zero matrix with conformal dimension. The Hadamard product of $M = [m_{ij}] \in \mathbb{R}^{m \times n}$ and $N = [n_{ij}] \in \mathbb{R}^{m \times n}$ is defined by $M \odot N := [m_{ij}n_{ij}] \in \mathbb{R}^{m \times n}$. For $x = [x_i] \in \mathbb{R}^n$ and $M \in \mathbb{R}^{m \times n}$, the infinity and max norm are defined by $\|x\|_{\infty} := \max_i |x_i|$ and $\|M\|_{\max} := \max_{i,j} |m_{ij}|$, respectively. We denote $E_{ij}^{mn} = e_i^m (e_j^n)^{\top}$ as the matrix with *ij*-element is 1 and zero elsewhere. The notation i = 1 : n indicates that *i* takes the values $1, 2, \ldots, n$. For any $M \in \mathbb{R}^{m \times n}$, let $|M| = [|m_{ij}|]$. For matrices $M, N \in \mathbb{R}^{m \times n}$, $|M| \le |N|$ means $|m_{ij}| \le |n_{ij}|$ for i = 1 : m and j = 1 : n. We

define M/N as $(M/N)_{ij} = n_{ij}^{\ddagger} m_{ij}$, where for any $a \in \mathbb{R}$, $a^{\ddagger} = \frac{1}{a}$ when $a \neq 0$, otherwise $a^{\ddagger} = 1$. $\Theta_x = \operatorname{diag}(x) \in \mathbb{R}^{n \times n}$ is the diagonal matrix, where $x \in \mathbb{R}^n$. For any $a \in \mathbb{R}$, $\operatorname{sign}(a) := \frac{a}{|a|}$ for $a \neq 0$ and sign(a) := 0 for a = 0, and sign $(M) := [sign(m_{ij})]$. The *i*-th row and the *j*-th column of M, are represented as M(i, :) and M(:, j), respectively.

Next, we discuss some important properties of the M-P inverse, which will be crucial for our main finding results. The following lemma states that for a full column rank matrices M, its M-P inverse is a continuous function of its data entries.

Lemma 2.1. [33] Let $M \in \mathbb{R}^{m \times n}$ with full column rank and $\{E_i\}$ be a collection of real $m \times n$ matrices satisfying $\lim_{j\to 0} E_j = 0$. Then, $(M + E_j)$ has full column rank when j is small enough and $\lim_{j \to 0} (M + E_j)^{\dagger} = M^{\dagger}.$

However, M does not share the above property when it is singular or rank deficient. Small perturbation ΔM on M can produce the computed M-P inverse far from the actual one. To tackle this situation, perturbation theory for the M-P inverse has been studied in certain specific constraints. Next, we recall the definition of 'acute' perturbation.

Definition 2.2. [29] An acute perturbation $\tilde{M} = M + \Delta M \in \mathbb{R}^{m \times n}$ of $M \in \mathbb{R}^{m \times n}$ is a perturbed matrix for which $\|MM^{\dagger} - \tilde{M}\tilde{M}^{\dagger}\|_{2} < 1$ and $\|M^{\dagger}M - \tilde{M}^{\dagger}\tilde{M}\|_{2} < 1$, where $\|\cdot\|_{2}$ is the spectral norm.

Proposition 2.3 provides an if and only if condition for the continuity of M^{\dagger} of any matrix $M \in \mathbb{R}^{m \times n}$.

Proposition 2.3. Let $M \in \mathbb{R}^{m \times n}$. Consider the set $S^1(M) = \{\Delta M \in \mathbb{R}^{m \times n} : \|M^{\dagger}\|_2 \|\Delta M\|_2 < \|\Delta M\|_2 < \|M\|_2$ 1]. Then $\lim_{\Delta M \to \mathbf{0}} (M + \Delta M)^{\dagger} = M^{\dagger}$ if and only if $\operatorname{rank}(M + \Delta M) = \operatorname{rank}(M)$, where $\Delta M \in \mathbb{N}^{M \times M}$. $\mathcal{S}^1(M)$.

Proof. For $\Delta M \in \mathcal{S}^1(M)$, we have $\|M^{\dagger}\|_2 \|\Delta M\|_2 < 1$. Then $M + \Delta M$ is an acute perturbation of M if and only if $\operatorname{rank}(M + \Delta M) = \operatorname{rank}(M)$ [21, Lemma 1]. Since on the set of acute perturbations of M, its M-P inverse M^{\dagger} is a continuous function about M [29, Page 140]. Therefore, it follows that M^{\dagger} is continuous on the set $S^{1}(M)$ if and only if rank $(M + \Delta M) =$ $\operatorname{rank}(M)$. Hence, the proof is completed.

Remark 1. When *M* has full column rank (or row rank), from Lemma 2.1, the rank condition in Proposition 2.3 holds trivially.

3. MCN and CCN for general parameterized matrices

In this part, initially, we define structured MCN and CCN for the M-P inverse and the unique MNLS solution for a general class of parameterized matrices. Suppose that each entry of $M \in$ $\mathbb{R}^{m \times n}$ is a differentiable function of a set of real parameters $\Psi = [\psi_1, \psi_2, \dots, \psi_p]^\top \in \mathbb{R}^p$ and write the matrix as $M(\Psi)$. We employ this notation for the rest of the paper. Due to the fact that a number of important classes of matrices can be parameterized by a collection of parameters, it is reasonable to consider perturbations on the parameters rather than directly on their entries. Let $\Delta \Psi \in \mathbb{R}^p$ be the perturbation on the parameters set $\Psi \in \mathbb{R}^p$, we consider the admissible perturbation in the matrix $M(\Psi)$ as $M(\Psi + \Delta \Psi) - M(\Psi) = \Delta M(\Psi)$. For maintaining the continuity property for $M^{\dagger}(\Psi)$, according to the Proposition 2.3, we restrict the perturbation on Ψ to the following set

$$\mathcal{S}(\Psi) := \left\{ \Delta \Psi \in \mathbb{R}^p : \operatorname{rank}(M(\Psi)) = \operatorname{rank}(M(\Psi + \Delta \Psi)) = r, \, \|M^{\dagger}(\Psi)\|_2 \|\Delta M(\Psi)\|_2 < 1 \right\}.$$
(2)

Next, we provide few examples to show that $\mathcal{S}(\Psi)$ is nonempty.

Example 3.1. In this example, we consider a symmetric matrix $M(\Psi) = [m_{ij}] \in \mathbb{R}^{3 \times 3}$, where $\Psi = [m_{11} = 1, m_{21} = 2, m_{31} = 3, m_{22} = 4, m_{32} = 6, m_{33} = 1]^{\top} \in \mathbb{R}^6$. Then

$$M(\Psi) = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \\ 3 & 6 & 1 \end{bmatrix}.$$

If we take $\Delta \Psi = [\Delta m_{11} = 0, \Delta m_{21} = 0, \Delta m_{31} = \mu, \Delta m_{22} = 0, \Delta m_{32} = 2\mu, \Delta m_{33} = 0]^{\top} \in \mathbb{R}^6$, where $\mu \neq -8/3$. Then, we get

$$M(\Psi + \Delta \Psi) = \begin{bmatrix} 1 & 2 & 3 + \mu \\ 2 & 4 & 6 + 2\mu \\ 3 + \mu & 6 + 2\mu & 1 \end{bmatrix}.$$

Here, $\|M^{\dagger}(\Psi)\|_{2} = 0.25$ and $\|\Delta M^{\dagger}(\Psi)\|_{2} = \sqrt{3}|\mu|$. Clearly, rank $(M(\Psi)) = \operatorname{rank}(M(\Psi + \Delta \Psi))$ and $\|M^{\dagger}(\Psi)\|_{2}\|\Delta M(\Psi)\|_{2} < 1$, whenever $|\mu| < 2.3094$. Therefore, the set $\mathcal{S}(\Psi)$ includes all perturbations $\Delta \Psi$ such that $|\mu| < 2.3094$.

Example 3.2. Consider the parameter set $\Psi = [\{2,4\},\{1\},\{-3,1\},\{5,2,6\},\{1,3\},\{2\},\{3,1\}]^{\top} \in \mathbb{R}^{13}$ of a $\{1,1\}$ -QS matrix given as in (27) and using the formula provided in Definition 5.1, we have

$$M(\Psi) = \begin{bmatrix} 5 & 3 & 2 \\ -6 & 2 & 3 \\ -12 & 4 & 6 \end{bmatrix}.$$
 (3)

Taking $\Delta \Psi = [\{0,0\},\{0\},\{0,0\},\{\mu,0,0\},\{\mu,0\},\{0\},\{0,0\}]^{\top} \in \mathbb{R}^{13}$, we get

$$M(\Psi + \Delta \Psi) = \begin{bmatrix} 5 + \mu & 3 + 3\mu & 2 + 2\mu \\ -6 & 2 & 3 \\ -12 & 4 & 6 \end{bmatrix}.$$

Here, $\|M^{\dagger}(\Psi)\|_{2} = 0.1812$ and $\|\Delta M^{\dagger}(\Psi)\|_{2} = \sqrt{14}|\mu|$. Clearly, $\operatorname{rank}(M(\Psi)) = \operatorname{rank}(M(\Psi + \Delta \Psi))$ and $\|M^{\dagger}(\Psi)\|_{2}\|\Delta M(\Psi)\|_{2} < 1$, whenever $|\mu| < 1.4747$. Thus, $\mathcal{S}(\Psi)$ contains all perturbations $\Delta \Psi$ such that $|\mu| < 1.4747$.

Example 3.3. Consider the parameter set $\Psi = [\{1, 2, 3\}, \{4, 5\}]^{\top} \in \mathbb{R}^5$ of a CV matrix and using the formula provided in Definition 4.1, we have

$$M(\Psi) = \begin{bmatrix} -\frac{1}{3} & -\frac{1}{4} & 1 & 1\\ -\frac{1}{2} & -\frac{1}{3} & 1 & 2\\ -1 & -\frac{1}{2} & 1 & 6 \end{bmatrix}.$$

Considering $\Delta \Psi = [\{0, 10^{-5}, 0\}, \{0, 0\}]^{\top} \in \mathbb{R}^5$, we have

$$M(\Psi + \Delta \Psi) = \begin{bmatrix} -\frac{1}{3} & -\frac{1}{4} & 1 & 1\\ -\frac{1}{2-10^{-5}} & -\frac{1}{3-10^{-5}} & 1 & 2+10^{-5}\\ -1 & -\frac{1}{2} & 1 & 6 \end{bmatrix}.$$

Here, rank $(M(\Psi))$ = rank $(M(\Psi+\Delta\Psi))$ = 3 and we obtain $||M^{\dagger}(\Psi)||_2$ = 7.5690 and $||\Delta M(\Psi)||_2$ = 1.0367 × 10⁻⁵. Consequently, $||M^{\dagger}(\Psi)||_2 ||\Delta M(\Psi)||_2$ = 7.8472 × 10⁻⁵ < 1, which shows that $\Delta \Psi \in \mathcal{S}(\Psi)$.

3.1. M-P inverse of general parameterized matrices

We introduce structured MCN and CCN in Definition 3.4 for M-P inverse of a general class of parameterized matrices. We provide general expressions for the upper bounds of these CNs in Theorem 3.6. Also, we present exact formulae for these CNs in Theorem 3.9 for full column rank matrices.

Definition 3.4. Let $M(\Psi) \in \mathbb{R}^{m \times n}$, rank $(M(\Psi)) = r \le \min\{m, n\}$. Then, we define structured MCN and CCN for $M^{\dagger}(\Psi)$ as follows:

$$\begin{aligned} \mathscr{M}^{\dagger}(M(\Psi)) &:= \lim_{\varepsilon \to 0} \sup \left\{ \frac{\|M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi)\|_{\max}}{\varepsilon \|M^{\dagger}\|_{\max}} : \|\Delta \Psi/\Psi\|_{\infty} \leq \varepsilon, \Delta \Psi \in \mathcal{S}(\Psi) \right\}, \\ \mathscr{C}^{\dagger}(M(\Psi)) &:= \lim_{\varepsilon \to 0} \sup \left\{ \frac{1}{\varepsilon} \left\| \frac{M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi)}{M^{\dagger}} \right\|_{\max} : \|\Delta \Psi/\Psi\|_{\infty} \leq \varepsilon, \Delta \Psi \in \mathcal{S}(\Psi) \right\}. \end{aligned}$$

To formulate the general expressions for the upper bounds of the CNs outlined in Definition 3.4, we present the following perturbation expression for $M^{\dagger}(\Psi)$.

Lemma 3.5. Let $M(\Psi) \in \mathbb{R}^{m \times n}$ and $\operatorname{rank}(M(\Psi)) = r$. Suppose $\Delta \Psi \in \mathcal{S}(\Psi)$ is the perturbation on the parameter set Ψ . Then

$$M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi) = \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} + M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} + \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \Delta \psi_{k} + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}).$$

Proof. Given that the elements of $M(\Psi)$ are differentiable functions of $\Psi = [\psi_1, \psi_2, ..., \psi_p]^\top$, using matrix differentiation, for an infinitesimal change in $M(\Psi)$, we get

$$\Delta M(\Psi) = M(\Psi + \Delta \Psi) - M(\Psi) = \sum_{k=1}^{p} \frac{\partial M(\Psi)}{\partial \psi_k} \Delta \psi_k + \mathcal{O}(\|\Delta \Psi\|_{\infty}^2), \tag{4}$$

where $\Delta \Psi = [\Delta \psi_1, \dots, \Delta \psi_p]^{\top}$. Since $\Delta \Psi \in \mathcal{S}(\Psi)$, using the perturbation expression for the M-P inverse [29], we obtain

$$\Delta M^{\dagger}(\Psi) = M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi) = -M^{\dagger} \Delta M(\Psi) M^{\dagger} + M^{\dagger} M^{\dagger^{\top}} (\Delta M(\Psi))^{\top} \mathbf{E}_{M} + \mathbf{F}_{M} (\Delta M(\Psi))^{\top} M^{\dagger^{\top}} M^{\dagger} + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}).$$
(5)

Putting (4) in (5), we get

$$\begin{split} \Delta M^{\dagger}(\Psi) &= -M^{\dagger} \Big(\sum_{k=1}^{p} \frac{\partial M(\Psi)}{\partial \psi_{k}} \Delta \psi_{k} \Big) M^{\dagger} + M^{\dagger} M^{\dagger^{\top}} \Big(\sum_{k=1}^{p} \frac{\partial M(\Psi)}{\partial \psi_{k}} \Delta \psi_{k} \Big)^{\top} \mathbf{E}_{M} \\ &+ \mathbf{F}_{M} \Big(\sum_{k=1}^{p} \frac{\partial M(\Psi)}{\partial \psi_{k}} \Delta \psi_{k} \Big)^{\top} M^{\dagger^{\top}} M^{\dagger} + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}). \end{split}$$

Hence, the desired expression is obtained.

In Theorem 3.6, for $M^{\dagger}(\Psi)$, we derive general expressions for upper bounds of the proposed CNs when rank $(M(\Psi)) = r$.

Theorem 3.6. For $M(\Psi) \in \mathbb{R}^{m \times n}$ with $\operatorname{rank}(M(\Psi)) = r$, we have

$$\mathscr{M}^{\dagger}(M(\Psi)) \leq \frac{\|\mathcal{X}_{\Psi}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} := \tilde{\mathscr{M}}^{\dagger}(M(\Psi)) \quad and \quad \mathscr{C}^{\dagger}(M(\Psi)) \leq \left\|\frac{\mathcal{X}_{\Psi}^{\dagger}}{M^{\dagger}}\right\|_{\max} := \tilde{\mathscr{C}}^{\dagger}(M(\Psi)),$$

where

$$\mathcal{X}_{\Psi}^{\dagger} = \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right| + \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} M^{\dagger} \right| \right) |\psi_{k}|.$$

Proof. From Lemma 3.5 and using the properties of absolute values, we have

$$\begin{split} \left| M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi) \right| &\leq \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right| \\ &+ \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \right| \right) |\Delta \psi_{k}| + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}). \end{split}$$

Now, by Definition 3.4, $\|\Delta \Psi/\Psi\|_{\infty} \leq \varepsilon$ implies that $|\Delta \psi_k| \leq \varepsilon |\psi_k|$ for all $k = 1, 2, \ldots, p$, and using the properties of the max norm, we find that

$$\begin{split} \|M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi)\|_{\max} &\leq \varepsilon \left\| \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right| \right. \\ &+ \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} M^{\dagger} \right| \right) |\psi_{k}| \right\|_{\max} + \mathcal{O}(\varepsilon^{2}). \end{split}$$

Then, if we take $\varepsilon \to 0$, and from Definition 3.4, we get the desired result of the first claim. In a similar manner, we obtain the second part of the claim.

In the next corollary, we obtain bounds for the CNs for M^{\dagger} in the unstructured case.

Corollary 3.7. Suppose $M \in \mathbb{R}^{m \times n}$ with $\operatorname{rank}(M) = r$. Then

$$\widetilde{\mathscr{M}}^{\dagger}(M) = \frac{1}{\|M^{\dagger}\|_{\max}} \left\| |M^{\dagger}| |M| |M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}| |M^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M}| |M^{\top}| |M^{\dagger^{\top}}M^{\dagger}| \right\|_{\max}, \\
\widetilde{\mathscr{C}}^{\dagger}(M) = \left\| \frac{1}{M^{\dagger}} \left(|M^{\dagger}| |M| |M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}| |M^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M}| |M^{\top}| |M^{\dagger^{\top}}M^{\dagger}| \right) \right\|_{\max}.$$

Proof. For any $M = [m_{ij}] \in \mathbb{R}^{m \times n}$ and any two column vectors a and b, we have

$$\frac{\partial M}{\partial m_{ij}} = e_i^m (e_j^n)^\top$$
 and (6)

$$|ab^{\top}| = |a||b^{\top}|. \tag{7}$$

By considering the parameters as the entries of M itself, i.e., $\Psi = [\{m_{ij}\}_{i,j=1}^{m,n}]^{\top} \in \mathbb{R}^{mn}$, and using (6), the sum expression in Theorem 3.6 can be written as:

$$\sum_{i=1}^{m} \sum_{j=1}^{n} \left(\left| M^{\dagger} \frac{\partial M}{\partial m_{ij}} M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger \top} \left(\frac{\partial M}{\partial m_{ij}} \right)^{\top} \mathbf{E}_{M} \right| + \left| \mathbf{F}_{M} \left(\frac{\partial M}{\partial m_{ij}} \right)^{\top} M^{\dagger \top} M^{\dagger} \right| \right) |m_{ij}|$$
$$= \sum_{i=1}^{m} \sum_{j=1}^{n} \left(\left| M^{\dagger} e_{i}^{m}(e_{j}^{n})^{\top} M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger \top} e_{j}^{n}(e_{i}^{m})^{\top} \mathbf{E}_{M} \right| + \left| \mathbf{F}_{M} e_{j}^{n}(e_{i}^{m})^{\top} M^{\dagger \top} M^{\dagger} \right| \right) |m_{ij}|.$$
(8)

Again, using (7), we can write (8) as

$$\sum_{i=1}^{m} \sum_{j=1}^{n} \left(|M^{\dagger}(:,i)| |m_{ij}| |M^{\dagger}(j,:)| + |M^{\dagger}M^{\dagger^{\top}}(:,j)| |m_{ij}| |\mathbf{E}_{M}(i,:)| + |\mathbf{F}_{M}(:,j)| |m_{ij}| |M^{\dagger^{\top}}M^{\dagger}(i,:)| \right)$$

$$= |M^{\dagger}| |M| |M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}| |M^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M}| |M^{\top}| |M^{\dagger^{\top}}M^{\dagger}|.$$
(9)

The desired upper bounds will be obtained by substituting (9) in Theorem 3.6.

Next, we estimate the bounds for CNs under the constraints $\mathcal{R}(\Delta M(\Psi)) \subseteq \mathcal{R}(M(\Psi))$ and $\mathcal{R}(\Delta M^{\top}(\Psi)) \subseteq \mathcal{R}(M^{\top}(\Psi))$.

Proposition 3.8. Let $M(\Psi) \in \mathbb{R}^{m \times n}$ be such that $\operatorname{rank}(M(\Psi)) = r$. Suppose that $\Delta \Psi \in \mathbb{R}^p$ is the perturbation on the parameter set Ψ satisfying the conditions, $\|M^{\dagger}\|_2 \|\Delta M(\Psi)\|_2 < 1$, $\mathcal{R}(\Delta M(\Psi)) \subseteq \mathcal{R}(M(\Psi))$ and $\mathcal{R}(\Delta M^{\top}(\Psi)) \subseteq \mathcal{R}(M^{\top}(\Psi))$. Then

$$M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi) = -\sum_{k=1}^{p} M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} \Delta \psi_{k} + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}).$$

Furthermore,

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi)) = \frac{\|\sum_{k=1}^{p} |M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger}| |\Delta \psi_{k}| \|_{\max}}{\|M^{\dagger}\|_{\max}},$$
$$\tilde{\mathscr{C}}^{\dagger}(M(\Psi)) = \left\| \frac{1}{M^{\dagger}} \sum_{k=1}^{p} |M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger}| |\Delta \psi_{k}| \right\|_{\max}.$$

Proof. If $M(\Psi), \Delta M(\Psi) \in \mathbb{R}^{m \times n}$ satisfies the assumptions $\mathcal{R}(\Delta M(\Psi)) \subseteq \mathcal{R}(M(\Psi))$ and $\mathcal{R}(\Delta M^{\top}(\Psi)) \subseteq \mathcal{R}(M^{\top}(\Psi))$. Then

$$M(\Psi)M^{\dagger}(\Psi)\Delta M(\Psi) = \Delta M(\Psi), \quad M^{\top}(\Psi)M^{\dagger}(\Psi)^{\top}\Delta M^{\top}(\Psi) = \Delta M^{\top}(\Psi).$$
(10)

In addition, if $||M^{\dagger}(\Psi)||_2 ||\Delta M(\Psi)||_2 < 1$ holds, it is shown in [1] that

$$M^{\dagger}(\Psi + \Delta \Psi) = (I_n + M^{\dagger}(\Psi)\Delta M(\Psi))^{-1}M^{\dagger}(\Psi).$$
(11)

Now, (10) implies $\Delta M^{\top}(\Psi) \mathbf{E}_M = \mathbf{0}$ and $\mathbf{F}_M \Delta M^{\top}(\Psi) = \mathbf{0}$. Again, (11) implies that rank $(M(\Psi + \Delta \Psi)) = \operatorname{rank}(M(\Psi))$. Therefore, $\Delta \Psi \in \mathcal{S}(\Psi)$. Hence, from Lemma 3.5, we get the desired expression.

The proof of the second part is a direct consequence of the derived perturbation expansion and the proof technique employed in Theorem 3.6. $\hfill\blacksquare$

Remark 2. Using Proposition 3.8, and in an analogous approach to Corollary 3.7, we can recover the bounds for unstructured CNs obtained in [35].

For the matrices with full column rank, the next theorem provides exact expressions of CNs for $M^{\dagger}(\Psi)$, introduced in Definition 3.4.

Theorem 3.9. For $M(\Psi) \in \mathbb{R}^{m \times n}$ with full column rank, we have

$$\mathscr{M}^{\dagger}(M(\Psi)) = \frac{\|\hat{\mathcal{X}}_{\Psi}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad and \quad \mathscr{C}^{\dagger}(M(\Psi)) = \left\|\frac{\hat{\mathcal{X}}_{\Psi}^{\dagger}}{M^{\dagger}}\right\|_{\max},$$

where

$$\hat{\mathcal{X}}_{\Psi}^{\dagger} = \sum_{k=1}^{p} \left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} - (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right| |\psi_{k}|.$$

Proof. Since $M(\Psi)$ is of full column rank matrix, we have $\mathbf{F}_M = \mathbf{0}$. Now, applying Remark 1 on Lemma 3.5, we get following perturbation expression for $M^{\dagger}(\Psi)$

$$\Delta M^{\dagger}(\Psi) = M^{\dagger}(\Psi + \Delta \Psi) - M^{\dagger}(\Psi) = \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} + (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right) \Delta \psi_{k} + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2}).$$
(12)

By employing a similar proof technique as in Theorem 3.6, and considering the given condition $\|\Delta\Psi/\Psi\|_{\infty} \leq \varepsilon$, we can establish the following bound:

$$\mathscr{M}^{\dagger}(M(\Psi)) \leq \frac{\left\|\sum_{k=1}^{p} \left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} - (M^{\top} M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} \right\| \psi_{k} \right\|_{\max}}{\|M^{\dagger}\|_{\max}}.$$
 (13)

On the other hand, from Lemma 2.1 and Remark 1, it follows that we can choose arbitrary perturbation $\Delta \Psi \in \mathbb{R}^p$ on the parameter set Ψ . Choose

$$\Delta \psi_k = -\varepsilon \operatorname{sign}(M_k)_{lq} \operatorname{sign}(\psi_k) \psi_k, \qquad (14)$$

where $M_k = M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_k} M^{\dagger} - (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_k}\right)^{\top} \mathbf{E}_M$, for $k = 1 : p, (M_k)_{lq}$ denotes the lq-th entry of the matrix M_k , and the indices l and q are such that

$$\left\|\sum_{k=1}^{p} |M_{k}| |\psi_{k}|\right\|_{\max} = \left(\sum_{k=1}^{p} |M_{k}| |\psi_{k}|\right)_{lq}.$$

The upper bound in (13) is obtained by inserting the values of (14) in the perturbation expression (12) and from the Definition 3.4. Therefore, the proof of the first claim is concluded. The second part of the claim can be obtained in a similar approach.

3.2. MNLS solution for general parameterized coefficient matrices

Let us consider the LS problem (1) for the parameterized matrix $M(\Psi) \in \mathbb{R}^{m \times n}$

$$\min_{z \in \mathbb{R}^n} \|M(\Psi)z - b\|_2 \tag{15}$$

with rank $(M(\Psi)) = r$ and $b \in \mathbb{R}^m$. When $M(\Psi)$ is rank deficient, the unique MNLS solution is provided by $\mathbf{x} = M(\Psi)^{\dagger}b$. Moreover, in this situation, \mathbf{x} is not even a continuous function of the data, and small changes in $M(\Psi)$ can produce large changes to \mathbf{x} . This happens as a consequence of the behavior of the M-P inverse for any rank deficient matrix. Thus, according to Proposition 2.3, we consider the perturbation $\Delta \Psi \in \mathcal{S}(\Psi)$ for the parameters, and then the perturbed problem

$$\min_{z \in \mathbb{R}^n} \|M(\Psi + \Delta \Psi)z - (b + \Delta b)\|_2$$

has the MNLS solution $\tilde{\mathbf{x}} = M(\Psi + \Delta \Psi)^{\dagger}(b + \Delta b)$. Consider $\Delta \mathbf{x} = \tilde{\mathbf{x}} - \mathbf{x}$.

In Definition 3.10, for the MNLS solution \mathbf{x} , we introduce its structured MCN and CCN.

Definition 3.10. Let $M(\Psi) \in \mathbb{R}^{m \times n}$ with $\operatorname{rank}(M(\Psi)) = r$ and $b \in \mathbb{R}^m$. Then, we define structured MCN and CCN of **x** as follows:

$$\mathcal{M}^{\dagger}(M(\Psi), b) := \lim_{\varepsilon \to 0} \sup \left\{ \frac{\|\Delta \mathbf{x}\|_{\infty}}{\varepsilon \|\mathbf{x}\|_{\infty}} : \|\Delta \Psi/\Psi\|_{\infty} \le \varepsilon, \|\Delta b/b\|_{\infty} \le \varepsilon, \ \Delta \Psi \in \mathcal{S}(\Psi), \ \Delta b \in \mathbb{R}^{m} \right\},$$
$$\mathcal{C}^{\dagger}(M(\Psi), b) := \lim_{\varepsilon \to 0} \sup \left\{ \frac{1}{\varepsilon} \left\| \frac{\Delta \mathbf{x}}{\mathbf{x}} \right\|_{\infty} : \|\Delta \Psi/\Psi\|_{\infty} \le \varepsilon, \|\Delta b/b\|_{\infty} \le \varepsilon, \ \Delta \Psi \in \mathcal{S}(\Psi), \ \Delta b \in \mathbb{R}^{m} \right\}$$

Our main objective of this section is to find general expressions of bounds for the CNs introduced in Definition 3.10, and the following lemma provides the perturbation expansion for the MNLS solution.

Lemma 3.11. Let $M(\Psi) \in \mathbb{R}^{m \times n}$ with $\operatorname{rank}(M(\Psi)) = r$ and $b \in \mathbb{R}^m$. Suppose $\Delta \Psi \in \mathcal{S}(\Psi)$ and $\Delta b \in \mathbb{R}^m$, and set $\mathbf{r} := b - M(\Psi)\mathbf{x}$. Then

$$\Delta \mathbf{x} = \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} + M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} + \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} \mathbf{x} \right) \Delta \psi_{k}$$
$$+ \sum_{i=1}^{m} M^{\dagger} e_{i}^{m} \Delta b_{i} + \mathcal{O}(\|[\Delta \Psi, \Delta b]\|_{\infty}^{2}).$$

Proof. Since $\Delta \mathbf{x} = M^{\dagger}(\Psi + \Delta \Psi)(b + \Delta b) - M^{\dagger}(\Psi)b$ and $\Delta \Psi \in \mathcal{S}(\Psi)$, using Lemma 3.5, we get

$$\Delta \mathbf{x} = \left(\sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} M^{\dagger} + M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{E}_{M} + \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \right) \Delta \psi_{k} + M^{\dagger} (\Psi) \right) (b + \Delta b) - M^{\dagger} (\Psi) b + \mathcal{O}(\|\Delta \Psi\|_{\infty}^{2})$$
(16)
$$= \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} + M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} + \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger^{\top}} \mathbf{x} \right) \Delta \psi_{k} + M^{\dagger} (\Psi) \Delta b + \mathcal{O}(\|[\Delta \Psi, \Delta b]\|_{\infty}^{2}).$$
(17)

On the other hand, for the perturbation Δb in b, we can write $\Delta b = \sum_{i=1}^{m} e_i^m \Delta b_i$. Therefore, from (17), we get

$$\Delta \mathbf{x} = \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} + M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} + \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} \mathbf{x} \right) \Delta \psi_{k}$$
$$+ M^{\dagger} \sum_{i=1}^{m} e_{i}^{m} \Delta b_{i} + \mathcal{O}(\|[\Delta \Psi, \Delta b]\|_{\infty}^{2}),$$

and hence, the desired result is obtained.

In Theorem 3.12, we provide general expressions for the upper bounds of $\mathscr{M}^{\dagger}(M(\Psi), b)$ and $\mathscr{C}^{\dagger}(M(\Psi), b)$.

Theorem 3.12. Let $M(\Psi) \in \mathbb{R}^{m \times n}$ be a matrix having $\operatorname{rank}(M(\Psi)) = r$ and $b \in \mathbb{R}^m$. Then,

$$\begin{aligned} \mathscr{M}^{\dagger}(M(\Psi), b) &\leq \quad \frac{\|\mathcal{X}_{\Psi}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} := \widetilde{\mathscr{M}}^{\dagger}(M(\Psi), b), \\ \mathscr{C}^{\dagger}(M(\Psi), b) &\leq \quad \left\|\Theta_{\mathbf{x}^{\dagger}}\mathcal{X}_{\Psi}^{ls}\right\|_{\infty} := \widetilde{\mathscr{C}}^{\dagger}(M(\Psi), b) \end{aligned}$$

where

$$\mathcal{X}_{\Psi}^{ls} = \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} \right| + \left| M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} \right| + \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} \mathbf{x} \right| \right) |\psi_{k}| + |M^{\dagger}||b|$$

and $\Theta_{\mathbf{x}^{\ddagger}} = \operatorname{diag}(\mathbf{x}^{\ddagger}).$

Proof. From Lemma 3.11 and utilizing the properties of absolute values, we obtain

$$|\Delta \mathbf{x}| \leq \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} \right| + \left| M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{T} \mathbf{r} \right| + \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger \top} \mathbf{x} \right| \right) |\Delta \psi_{k}| + \sum_{i=1}^{m} |M^{\dagger}| |\Delta b_{i}| + \mathcal{O}(\|[\Delta \Psi, \Delta b]_{\infty}\|^{2}).$$
(18)

Now, by Definition 3.10, $\|\Delta \Psi/\Psi\|_{\infty} \leq \varepsilon$ and $\|\Delta b/b\|_{\infty} \leq \varepsilon$ implies that for k = 1 : p, $|\Delta \psi_k| \leq \varepsilon |\psi_k|$, and for i = 1 : m, $|\Delta b_i| \leq \varepsilon |b_i|$, respectively. Taking infinity norm in (18), we deduced that

$$\|\Delta \mathbf{x}\|_{\infty} \leq \varepsilon \left\| \sum_{k=1}^{p} \left(\left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} \right| + \left| M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{T} \mathbf{r} + \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} M^{\dagger^{\top}} \mathbf{x} \right| \right) |\psi_{k}| + |M^{\dagger}||b| \right\|_{\infty} + \mathcal{O}(\varepsilon^{2}).$$
(19)

Then, if we take $\varepsilon \to 0$ in (19) and from Definition 3.10, we attain the desired result of the first assertion. The second assertion follows in a similar manner, as we can express $\left\|\frac{\Delta \mathbf{x}}{\mathbf{x}}\right\|_{\infty} = \|\Theta_{\mathbf{x}^{\dagger}} \Delta \mathbf{x}\|_{\infty}$.

Next, we discuss the bounds of the CNs for the MNLS solution of the LS problem (15) corresponding to unstructured matrices.

Corollary 3.13. For $M \in \mathbb{R}^{m \times n}$ having rank(M) = r and $b \in \mathbb{R}^m$, we have

$$\begin{split} \tilde{\mathscr{M}}^{\dagger}(M,b) &:= \frac{\left\| |M^{\dagger}||M||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||M^{\top}||\mathbf{r}| + |\mathbf{F}_{M}||M^{\top}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||b| \right\|_{\infty}}{\|\mathbf{x}\|_{\infty}}, \\ \tilde{\mathscr{C}}^{\dagger}(M,b) &:= \left\| \Theta_{\mathbf{x}^{\dagger}} \left(|M^{\dagger}||M||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||M^{\top}||\mathbf{r}| + |\mathbf{F}_{M}||M^{\top}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||b| \right) \right\|_{\infty} \end{split}$$

Proof. The proof follows in an analogous way to the Corollary 3.7 by considering $\Psi = [\{m_{ij}\}_{i,j=1}^{m,n}]^{\top} \in \mathbb{R}^{mn}$ in Theorem 3.12 and using (6) and (7).

The next theorem offers explicit formulae of structured CNs for the unique LS solution $\mathbf{x} = M^{\dagger}(\Psi)b$ for full column rank matrices.

Theorem 3.14. For $M(\Psi) \in \mathbb{R}^{m \times n}$ having full column rank and $b \in \mathbb{R}^m$, we get

$$\mathscr{M}^{\dagger}(M(\Psi), b) = \frac{\|\hat{\mathcal{X}}_{\Psi}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad and \quad \mathscr{C}^{\dagger}(M(\Psi), b) = \left\|\Theta_{\mathbf{x}^{\dagger}}\hat{\mathcal{X}}_{\Psi}^{ls}\right\|_{\infty},$$

where

$$\hat{\mathcal{X}}_{\Psi}^{ls} = \sum_{k=1}^{p} \left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} - (M^{\top} M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} \right| |\psi_{k}| + |M^{\dagger}| |b|.$$
(20)

Proof. In Lemma 3.11, using the fact that for any full column rank matrix $\mathbf{F}_M = \mathbf{0}$, we have

$$\Delta \mathbf{x} = \sum_{k=1}^{p} \left(-M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} + M^{\dagger} M^{\dagger \top} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} \right) \Delta \psi_{k} + \sum_{i=1}^{m} M^{\dagger} e_{i}^{m} \Delta b_{i} + \mathcal{O}(\|[\Delta \psi, \Delta b]\|_{\infty}^{2}).$$
(21)

Now, by applying the proof method used in Theorem 3.9 and considering the given conditions $\|\Delta\Psi/\Psi\|_{\infty} \leq \varepsilon$ and $\|\Delta b/b\|_{\infty} \leq \varepsilon$, we obtain

$$\mathscr{M}^{\dagger}(M(\Psi),b) \leq \frac{\left\|\sum_{k=1}^{p} \left| M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_{k}} \mathbf{x} - (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_{k}} \right)^{\top} \mathbf{r} \right| |\psi_{k}| + |M^{\dagger}| |b| \right\|_{\infty}}{\|\mathbf{x}\|_{\infty}}.$$
 (22)

From Lemma 2.1 and Remark 1, we can choose the perturbation $\Delta \Psi$ on the parameters set Ψ arbitrarily from \mathbb{R}^p . Consider the following perturbations

$$\Delta b = \varepsilon \, \Theta_{M^{\dagger}} \Theta_b \, b,$$

where $\Theta_{M^{\dagger}}$ and Θ_b are the diagonal matrices having diagonal entries $\Theta_{M^{\dagger}}(i, i) = \text{sign}(M^{\dagger}(l, i))$, and $\Theta_b(i, i) = \text{sign}(b_i)$ for i = 1 : m, respectively, and

$$\Delta \psi_k = -\varepsilon \operatorname{sign}(M_{\mathbf{x},k})_l \operatorname{sign}(\psi_k) \,\psi_k,$$

where $M_{\mathbf{x},k} := M^{\dagger} \frac{\partial M(\Psi)}{\partial \psi_k} \mathbf{x} - (M^{\top} M)^{-1} \left(\frac{\partial M(\Psi)}{\partial \psi_k}\right)^{\top} \mathbf{r}$ and l is the index so that

$$\left\|\sum_{k=1}^{p} |M_{\mathbf{x},k}| |\psi_{k}| + |M^{\dagger}| |b|\right\|_{\infty} = \left(\sum_{k=1}^{p} |M_{\mathbf{x},k}| |\psi_{k}| + |M^{\dagger}| |b|\right)_{l}.$$

The upper bound in (22) will be attained by substituting these perturbations in (21) and from Definition 3.10, and hence the desired expression is attained. Analogously, we can obtain the expression for the CCN.

Remark 3. The formula for the MCN $\mathscr{M}^{\dagger}(M(\phi), b)$ in Theorem 3.14 is also presented in [37]. However, our approach to proving the theorem differs slightly. Interested readers may also refer to the proof method in [37].

4. Cauchy-Vandermonde (CV) matrices

In this section, we start by reviewing the definition of CV matrices. Subsequently, we provide the derivative expressions for the matrix with respect to its parameter set. These expressions play a pivotal role in the derivation of computationally feasible upper bounds for the structured CNs of the M-P inverse and the solution of the LS problem for a rank deficient CV matrix given in Theorems 4.3 and 4.5, respectively. Explicit formulations for these CNs are also provided in the Theorems 4.4 and 4.6, respectively, when the matrix has full column rank. **Definition 4.1.** [18] A matrix $M \in \mathbb{R}^{m \times n}$ is classified as a CV matrix if it satisfies the following conditions: for $c = [c_1, c_2, \ldots, c_m]^\top \in \mathbb{R}^m$ and $d = [d_1, d_2, \ldots, d_l]^\top \in \mathbb{R}^l$, where $c_i \neq d_j$ for i = 1 : m and j = 1 : l, with $0 \le l \le n$, the matrix M can be represented as follows:

$$M = \begin{bmatrix} \frac{1}{c_1 - d_1} & \frac{1}{c_1 - d_2} & \cdots & \frac{1}{c_1 - d_l} & 1 & c_1 & c_1^2 & \cdots & c_1^{n-l-1} \\ \frac{1}{c_2 - d_1} & \frac{1}{c_2 - d_2} & \cdots & \frac{1}{c_2 - d_l} & 1 & c_2 & c_2^2 & \cdots & c_2^{n-l-1} \\ \vdots & \vdots \\ \frac{1}{c_m - d_1} & \frac{1}{c_m - d_2} & \cdots & \frac{1}{c_m - d_l} & 1 & c_m & c_m^2 & \cdots & c_m^{n-l-1} \end{bmatrix}.$$
(23)

M becomes the Vandermonde matrix when l = 0, and the Cauchy matrix when l = n.

For a CV matrix of the form (23), $\Psi_{\mathbb{CV}} := [\{c_i\}_{i=1}^m, \{d_i\}_{i=1}^l]^\top \in \mathbb{R}^{m+l}$ represents its parameter set. We use the notation $M(\Psi_{\mathbb{CV}})$ to refer a CV matrix parameterized by $\Psi_{\mathbb{CV}}$.

For the M-P inverse and the MNLS solution involving the CV matrix, our objective is to estimate the structured CNs. Lemma 4.2 accomplishes our claim. Before that, we will construct the following matrices. For any positive integers p, q and any vector $y = [y_1, y_2, \ldots, y_p]^{\top} \in \mathbb{R}^p$, define the matrices

$$\mathcal{Q}_{y,i}^{pq} := \begin{bmatrix} \mathbf{1}, \dots, \mathbf{1}, y, \mathbf{1}, \dots, \mathbf{1} \end{bmatrix} \in \mathbb{R}^{p \times q},$$

for i = 1 : q, with the *i*-th column is y and $\mathbf{1} \in \mathbb{R}^p$ have all entries equal to 1. Also, set

$$\mathcal{M}_1 := \left| -M(\Psi_{\mathbb{C}\mathbb{V}})(:, 1:l) \quad \mathbf{0} \quad M(\Psi_{\mathbb{C}\mathbb{V}})(:, l+2:n) \right| \in \mathbb{R}^{m \times n},\tag{24}$$

$$\mathcal{M}_2 := \begin{bmatrix} M(\Psi_{\mathbb{CV}})(:, 1:l) & \mathbf{0} \end{bmatrix} \in \mathbb{R}^{m \times n}, \tag{25}$$

where ${\bf 0}$ is the zero matrix with conformal dimensional.

The following lemma presents the derivative expressions of a CV matrix for the parameters in $\Psi_{\mathbb{CV}}$.

Lemma 4.2. Suppose $M(\Psi_{\mathbb{CV}}) \in \mathbb{R}^{m \times n}$ having rank r, represented by a set of real parameter $\Psi_{\mathbb{CV}} = \left[\{c_i\}_{i=1}^m, \{d_i\}_{i=1}^l\right]^\top \in \mathbb{R}^{m+l}$, where $c_i \neq d_j$, i = 1 : m and j = 1 : l. Then, each entry of $M(\Psi_{\mathbb{CV}})$ is a differentiable function of $\Psi_{\mathbb{CV}}$, and

(1)
$$\frac{\partial M(\Psi_{\mathbb{C}\mathbb{V}})}{\partial c_i} = e_i^m (\mathcal{M}_1 \odot (\mathcal{Q}_{\mathbf{c}'_i,i}^{nm})^\top)(i,:) \text{ for } i = 1:m,$$

(2)
$$\frac{\partial M(\Psi_{\mathbb{C}\mathbb{V}})}{\partial d_j} = (\mathcal{M}_2 \odot \mathcal{Q}_{\mathbf{d}'_j,j}^{mn})(:,j)(e_j^n)^\top \text{ for } j = 1:l,$$

where

$$\mathbf{c}'_{i} := \left[\frac{1}{c_{i} - d_{1}}, \frac{1}{c_{i} - d_{2}}, \dots, \frac{1}{c_{i} - d_{l}}, 1, \frac{1}{c_{i}}, \frac{2}{c_{i}}, \dots, \frac{(n - l - 1)}{c_{i}}\right]^{\top} \in \mathbb{R}^{n}$$
$$\mathbf{d}'_{j} := \left[\frac{1}{c_{1} - d_{j}}, \frac{1}{c_{2} - d_{j}}, \dots, \frac{1}{c_{m} - d_{j}}\right]^{\top} \in \mathbb{R}^{m},$$

for i = 1 : m and j = 1 : l.

Proof. By observing that, when $c_i \neq d_j$, where i = 1 : m and j = 1 : l, partial derivatives

corresponding to the parameters $\{c_i\}_{i=1}^m$ will be

$$\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial c_i} = \begin{bmatrix} 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ \frac{-1}{(c_i - d_1)^2} & \cdots & \frac{-1}{(c_i - d_l)^2} & 0 & 1 & 2c_i & \cdots & (n - l - 1)c_i^{n - l - 2} \\ 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 0 & 0 & 0 & 0 & \cdots & 0 \end{bmatrix}.$$

Now, on the right-hand side of the above, using Hadamard product with the matrix $(\mathcal{Q}_{\mathbf{c}'_i,i}^{nm})^{\top}$, we get

$$\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial c_i} = \left(e_i^m \left[-M(\Psi_{\mathbb{CV}})(i,1:l) \quad 0 \quad M(\Psi_{\mathbb{CV}})(i,l+2:n) \right] \right) \odot (\mathcal{Q}_{\mathbf{c}_i',i}^{nm})^\top \\ = \left(e_i^m \mathcal{M}_1(i,:) \right) \odot (\mathcal{Q}_{\mathbf{c}_i',i}^{nm})^\top = e_i^m (\mathcal{M}_1 \odot (\mathcal{Q}_{\mathbf{c}_i',i}^{nm})^\top)(i,:).$$

Hence, proof of the first statement follows.

In a similar argument, we can prove the second part of the statement.

For the structured CNs, computationally feasible upper bounds are provided in the following theorem for $M^{\dagger}(\Psi_{\mathbb{CV}})$ addressed in Definition 3.4.

Theorem 4.3. Suppose $M(\Psi_{\mathbb{CV}}) \in \mathbb{R}^{m \times n}$ with $\operatorname{rank}(M(\Psi_{\mathbb{CV}})) = r$. Then

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \frac{\|\mathcal{X}_{\mathbb{C}\mathbb{V}}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \left\|\frac{\mathcal{X}_{\mathbb{C}\mathbb{V}}^{\dagger}}{M^{\dagger}}\right\|_{\max},$$

where

$$\begin{aligned} \mathcal{X}_{\mathbb{CV}}^{\dagger} = & |M^{\dagger}||\Theta_{c}||(\mathcal{M}_{1}\odot\mathcal{Q})M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}(\mathcal{M}_{1}\odot\mathcal{Q})^{\top}||\Theta_{c}||\mathbf{E}_{M}| \\ & + |\mathbf{F}_{M}(\mathcal{M}_{1}\odot\mathcal{Q})^{\top}||\Theta_{c}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}(\mathcal{M}_{2}\odot\mathcal{M}_{2})||\Theta_{d'}||M^{\dagger}| \\ & + |M^{\dagger}M^{\dagger^{\top}}||\Theta_{d'}||(\mathcal{M}_{2}\odot\mathcal{M}_{2})^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}||\Theta_{d'}||(\mathcal{M}_{2}\odot\mathcal{M}_{2})^{\top}M^{\dagger^{\top}}M^{\dagger}|, \end{aligned}$$

 $d' = [d_1, \ldots, d_l, 0, \ldots, 0]^\top \in \mathbb{R}^n, \ \mathcal{Q} = [\mathbf{c}_1'^\top, \mathbf{c}_2'^\top, \ldots, \mathbf{c}_m'^\top]^\top \in \mathbb{R}^{m \times n} \ and \ \mathcal{M}_1, \mathcal{M}_2 \ as \ defined \ in$ (24) and (25), respectively.

Proof. For deriving the desired expressions for $\tilde{\mathcal{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$ and $\tilde{\mathcal{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$, we calculate the contribution of each parameter subset to the expressions outlined in Theorem 3.6. For the parameters $\{c_i\}_{i=1}^l$, using (1) of Lemma 4.2 we get:

$$\mathcal{E}_{c} := \sum_{i=1}^{m} \left(\left| M^{\dagger} \frac{\partial M(\Psi_{\mathbb{C}\mathbb{V}})}{\partial c_{i}} M^{\dagger} \right| + \left| M^{\dagger^{\top}} M^{\dagger} \left(\frac{\partial M(\Psi_{\mathbb{C}\mathbb{V}})}{\partial c_{i}} \right)^{\top} \mathbf{E}_{M} \right| \right. \\ \left. + \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{C}\mathbb{V}})}{\partial c_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \right| \right) |c_{i}| \\ = \sum_{i=1}^{m} \left(\left| M^{\dagger} e_{i}^{m} ((\mathcal{Q}_{\mathbf{c}_{i}',i}^{nm})^{\top} \odot \mathcal{M}_{1})(i,:) M^{\dagger} \right| + \left| M^{\dagger} M^{\dagger^{\top}} \left(e_{i}^{m} ((\mathcal{Q}_{\mathbf{c}_{i}',i}^{nm})^{\top} \odot \mathcal{M}_{1})(i,:) \right)^{\top} \mathbf{E}_{M} \right| \\ \left. + \left| \mathbf{F}_{M} \left(e_{i}^{m} ((\mathcal{Q}_{\mathbf{c}_{i}',i}^{nm})^{\top} \odot \mathcal{M}_{1})(i,:) \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \right| \right) |c_{i}|.$$

$$(26)$$

Using (7) in (26), we get

$$\begin{aligned} \mathcal{E}_{c} &= \sum_{i=1}^{m} \left(|M^{\dagger}(:,i)||c_{i}||((\mathcal{Q}_{\mathbf{c}_{i}^{\prime},i}^{nm})^{\top} \odot \mathcal{M}_{1})(i,:)M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}(\mathcal{Q}_{\mathbf{c}_{i}^{\prime},i}^{nm})^{\top} \odot \mathcal{M}_{1})^{\top})^{\top}(i,:)||c_{i}||\mathbf{E}_{M}(i,:)| \right. \\ &+ |\mathbf{F}_{M}(\mathcal{Q}_{\mathbf{c}_{i}^{\prime},i}^{nm})^{\top} \odot \mathcal{M}_{1})^{\top}(i,:)||c_{i}||M^{\dagger^{\top}}M^{\dagger}(i,:)| \right) \\ &= |M^{\dagger}||\Theta_{c}||(\mathcal{M}_{1} \odot \mathcal{Q})M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}(\mathcal{M}_{1} \odot \mathcal{Q})^{\top}||\Theta_{c}||\mathbf{E}_{M}| + |\mathbf{F}_{M}(\mathcal{M}_{1} \odot \mathcal{Q})^{\top}||\Theta_{c}||M^{\dagger^{\top}}M^{\dagger}|. \end{aligned}$$

Analogously, for the parameters $\{d_i\}_{i=1}^l$, using (2) of Lemma 4.2, we have

$$\begin{aligned} \mathcal{E}_{d} &:= \sum_{i=1}^{l} \left(\left| M^{\dagger} \frac{\partial M(\Psi_{\mathbb{CV}})}{\partial d_{i}} M^{\dagger} \right| + \left| M^{\dagger^{\top}} M^{\dagger} \left(\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial d_{i}} \right)^{\top} \mathbf{E}_{M} \right| \\ &+ \left| \mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial d_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} \right| \right) |d_{i}| \\ &= |M^{\dagger} (\mathcal{M}_{2} \odot \mathcal{M}_{2})||\Theta_{d'}||M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}}||\Theta_{d'}||(\mathcal{M}_{2} \odot \mathcal{M}_{2})^{\top} \mathbf{E}_{M}| \\ &+ |\mathbf{F}_{M}||\Theta_{d'}||(\mathcal{M}_{2} \odot \mathcal{M}_{2})^{\top} M^{\dagger^{\top}} M^{\dagger}|. \end{aligned}$$

Applying Theorem 3.6 yields

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \frac{\|\mathcal{E}_d + \mathcal{E}_c\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad \text{and} \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \left\|\frac{\mathcal{E}_d + \mathcal{E}_c}{M^{\dagger}}\right\|_{\max}.$$

Hence, the proof is completed.

We now employ Theorem 3.9 to deduce the explicit formulations for structured CNs of $M^{\dagger}(\Psi_{\mathbb{CV}})$ by considering $M(\Psi)$ has full column rank, which is presented next.

Theorem 4.4. Suppose $M(\Psi_{\mathbb{CV}}) \in \mathbb{R}^{m \times n}$ has full column rank. Then,

$$\mathscr{M}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \frac{\|\hat{\mathcal{X}}_{\mathbb{C}\mathbb{V}}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad and \quad \mathscr{C}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \left\|\frac{\hat{\mathcal{X}}_{\mathbb{C}\mathbb{V}}^{\dagger}}{M^{\dagger}}\right\|_{\max},$$

where

$$\hat{\mathcal{X}}_{\mathbb{CV}}^{\dagger} = \sum_{i=1}^{m} \left| M^{\dagger} E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}) M^{\dagger} - (M^{\top} M)^{-1} (E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}))^{\top} \mathbf{E}_{M} \right| |c_{i}|$$
$$+ \sum_{i=1}^{l} \left| M^{\dagger} (\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn} M^{\dagger} - (M^{\top} M)^{-1} ((\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn})^{\top} \mathbf{E}_{M} \right| |d_{i}|,$$

 $E_{ij}^{mn} = e_i^m (e_j^n)^\top$ and \mathcal{Q} is as defined in Theorem 4.3.

Proof. To employ the expressions given in Theorem 3.9, we need to compute the contribution for each subset of parameters in an analogous method to the proof of Theorem 4.3. For the

parameters $\{c_i\}_{i=1}^m$, using (1) of Lemma 4.2, we have

$$\begin{aligned} \mathcal{E}'_{c} &:= \sum_{i=1}^{m} \left| M^{\dagger} \frac{\partial M(\Psi_{\mathbb{CV}})}{\partial c_{i}} M^{\dagger} - (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial c_{i}} \right)^{\top} \mathbf{E}_{M} \right| |c_{i}| \\ &= \sum_{i=1}^{m} \left| M^{\dagger} \left(e_{i}^{m} (\mathcal{M}_{1} \odot (\mathcal{Q}_{\mathbf{c}_{i}',i}^{nm})^{\top})(i,:) \right) M^{\dagger} - (M^{\top}M)^{-1} \left(e_{i}^{m} (\mathcal{M}_{1} \odot (\mathcal{Q}_{\mathbf{c}_{i}',i}^{nm})^{\top})(i,:) \right)^{\top} \mathbf{E}_{M} \right| |c_{i}| \\ &= \sum_{i=1}^{m} \left| M^{\dagger} E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}) M^{\dagger} - (M^{\top}M)^{-1} (E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}))^{\top} \mathbf{E}_{M} \right| |c_{i}|. \end{aligned}$$

Similarly, for the parameters $\{d_i\}_{i=1}^l$, using (2) of Lemma 4.2, we get

$$\mathcal{E}'_{d} := \sum_{i=1}^{l} \left| M^{\dagger} \frac{\partial M(\Psi_{\mathbb{CV}})}{\partial d_{i}} M^{\dagger} - (M^{\top}M)^{-1} \left(\frac{\partial M(\Psi_{\mathbb{CV}})}{\partial d_{i}} \right)^{\top} \mathbf{E}_{M} \right| |d_{i}|$$
$$= \sum_{i=1}^{l} \left| M^{\dagger} (\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn} M^{\dagger} - (M^{\top}M)^{-1} ((\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn})^{\top} \mathbf{E}_{M} \right| |d_{i}|.$$

From Theorem 3.9, we have

$$\mathscr{M}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \frac{\|\mathscr{E}'_c + \mathscr{E}'_d\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad \text{and} \quad \mathscr{C}^{\dagger}(M(\Psi_{\mathbb{C}\mathbb{V}})) = \left\|\frac{\mathscr{E}'_c + \mathscr{E}'_d}{M^{\dagger}}\right\|_{\max},$$

and hence, our proof is completed.

Remark 4. If we consider l = 0 or l = n in the preceding results, we can calculate the structured CNs for the M-P inverse Vandermonde and Cauchy matrices, respectively.

Next, we consider the LS problem (15) corresponding to a rank deficient CV matrix. Using the expressions given in Theorem 3.12, we deduce upper bounds for structured CNs of \mathbf{x} , presented next.

Theorem 4.5. Suppose $M(\Psi_{\mathbb{CV}}) \in \mathbb{R}^{m \times n}$ with rank r and $b \in \mathbb{R}^m$. Set $\mathbf{r} := b - M(\Psi_{\mathbb{CV}})\mathbf{x}$. Then,

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}), b) = \frac{\|\mathcal{X}_{\mathbb{CV}}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}, b)) = \left\|\Theta_{\mathbf{x}^{\dagger}}\mathcal{X}_{\mathbb{CV}}^{ls}\right\|_{\infty},$$

where

$$\begin{aligned} \mathcal{X}_{\mathbb{C}\mathbb{V}}^{ls} = & |M^{\dagger}||b| + |M^{\dagger}||\Theta_{c}||(\mathcal{M}_{1}\odot\mathcal{Q})\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}(\mathcal{M}_{1}\odot\mathcal{Q})^{\top}||\Theta_{c}||\mathbf{r}| \\ & + |\mathbf{F}_{M}(\mathcal{M}_{1}\odot\mathcal{Q})^{\top}||\Theta_{c}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}(\mathcal{M}_{2}\odot\mathcal{M}_{2})||\Theta_{d'}||\mathbf{x}| \\ & + |M^{\dagger}M^{\dagger^{\top}}||\Theta_{d'}||\mathcal{M}_{2}\odot\mathcal{M}_{2})^{\top}\mathbf{r}| + |\mathbf{F}_{M}||\Theta_{d'}||(\mathcal{M}_{2}\odot\mathcal{M}_{2})^{\top}M^{\dagger^{\top}}\mathbf{x}|. \end{aligned}$$

Proof. By evaluating in an analogous method to the proof of Theorem 4.3, for each subset of parameters using the expressions given in Theorem 3.12, the proof is followed. Hence, we omit it here.

The structured CNs to the LS problem (15) corresponding to a full column rank CV matrix are stated next.

Theorem 4.6. Let $M(\Psi_{\mathbb{CV}}) \in \mathbb{R}^{m \times n}$ and $b \in \mathbb{R}^m$. Set $\mathbf{r} := b - M\mathbf{x}$. Then,

$$\mathscr{M}^{\dagger}(M(\Psi_{\mathbb{CV}}), b) = \frac{\|\hat{\mathcal{X}}_{\mathbb{CV}}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad and \quad \mathscr{C}^{\dagger}(M(\Psi_{\mathbb{CV}}), b) = \left\|\Theta_{\mathbf{x}^{\ddagger}}\hat{\mathcal{X}}_{\mathbb{CV}}^{ls}\right\|_{\infty},$$

where

$$\hat{\mathcal{X}}_{\mathbb{CV}}^{ls} = \sum_{i=1}^{m} \left| M^{\dagger} E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}) \mathbf{x} - (M^{\top} M)^{-1} (E_{ii}^{mm} (\mathcal{M}_{1} \odot \mathcal{Q}))^{\top} \mathbf{r} \right| |c_{i}|$$
$$+ \sum_{i=1}^{l} \left| M^{\dagger} (\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn} \mathbf{x} - (M^{\top} M)^{-1} ((\mathcal{M}_{2} \odot \mathcal{M}_{2}) E_{jj}^{nn})^{\top} \mathbf{r} + |M^{\dagger}| |b|$$

Proof. Since the proof follows in an analogous method to the proof of Theorem 4.4, by finding the contribution of each parameter set in the expressions of Theorem 3.14. Hence, we omit the proof.

5. Quasiseparable (QS) matrices

The outset of this section begins with a quick introduction to QS matrices, which is a specific type of rank-structured matrices. Specifically, CNs are investigated for two important representations known as QS representation [12,14] and GV representation [30]. Upper bounds for the CNs of the M-P inverse and the MNLS solution are obtained corresponding to QS representation in Subsection 5.1 and for the GV representation in Subsection 5.2. The relationship between different CNs is also investigated in Subsection 5.3.

For the first time, QS matrices were investigated in [14]. In this work, we focus solely on considering $\{1,1\}$ -QS matrices, which is a special case of QS matrices. Let M be in $\mathbb{R}^{n \times n}$. If every submatrix of M completely contained in the strictly lower triangular (resp. upper triangular) part is of rank at most 1 (resp. 1), and there is at least one of these submatrices has rank equal to 1 (resp. 1), then M is called a $\{1,1\}$ -QS matrix. Equivalently, we can write: $\max_{i=1}^{n} \operatorname{rank}(M(i+1:n,1:i)) = 1$ and $\max_{i=1}^{n} \operatorname{rank}(M(1:i,i+1:n)) = 1$.

5.1. CNs corresponding to QS representation

In [12,14], the notion of QS representation was proposed for $\{1,1\}$ -QS matrices. In this subsection, we first recall this representation and then discuss the structured MCN and CCN.

Definition 5.1. [12,14] A matrix $M \in \mathbb{R}^{n \times n}$ is classified to be a $\{1, 1\}$ -QS matrix if it can be parameterized by the following set of 7n - 8 real parameters

$$\Psi_{\mathbb{QS}} = \left[\{\mathbf{a}_{\mathbf{i}}\}_{i=2}^{n}, \{\mathbf{e}_{\mathbf{i}}\}_{i=2}^{n-1}, \{\mathbf{b}_{\mathbf{i}}\}_{i=1}^{n-1}, \{\mathbf{d}_{\mathbf{i}}\}_{i=1}^{n}, \{\mathbf{f}_{i}\}_{i=1}^{n-1}, \{\mathbf{g}_{i}\}_{i=2}^{n-1}, \{\mathbf{h}_{\mathbf{i}}\}_{i=2}^{n} \right]^{\top} \in \mathbb{R}^{7n-8},$$
(27)

as follows,

$$M = \begin{bmatrix} \mathbf{d}_1 & \mathbf{f}_1 \mathbf{h}_2 & \mathbf{f}_1 \mathbf{g}_2 \mathbf{h}_3 & \cdots & \mathbf{f}_1 \mathbf{g}_2 \cdots \mathbf{g}_{n-1} \mathbf{h}_n \\ \mathbf{a}_2 \mathbf{b}_1 & \mathbf{d}_2 & \mathbf{f}_2 \mathbf{h}_3 & \cdots & \mathbf{f}_2 \mathbf{g}_3 \cdots \mathbf{g}_{n-1} \mathbf{h}_n \\ \mathbf{a}_3 \mathbf{e}_2 \mathbf{b}_1 & \mathbf{a}_3 \mathbf{b}_2 & \mathbf{d}_3 & \cdots & \mathbf{f}_3 \mathbf{g}_4 \cdots \mathbf{g}_{n-1} \mathbf{h}_n \\ \mathbf{a}_4 \mathbf{e}_3 \mathbf{e}_2 \mathbf{b}_1 & \mathbf{a}_4 \mathbf{e}_3 \mathbf{b}_2 & \mathbf{a}_4 \mathbf{b}_3 & \cdots & \mathbf{f}_4 \mathbf{g}_5 \cdots \mathbf{g}_{n-1} \mathbf{h}_n \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \mathbf{a}_n \mathbf{e}_{n-1} \dots \mathbf{e}_2 \mathbf{b}_1 & \mathbf{a}_n \mathbf{e}_{n-1} \dots \mathbf{e}_3 \mathbf{b}_2 & \mathbf{a}_n \mathbf{e}_{n-1} \dots \mathbf{e}_4 \mathbf{b}_3 & \cdots & \mathbf{d}_n \end{bmatrix}.$$

The set of real parameters $\Psi_{\mathbb{QS}}$ as in (5.1) is called *QS representation* of *M*. We use the notation $M(\Psi_{\mathbb{QS}})$ to refer a $\{1,1\}$ -QS matrix parameterized by the set $\Psi_{\mathbb{QS}}$. For the rest part, we set

$$M(\Psi_{\mathbb{OS}}) := \mathcal{L}_M + \mathcal{D}_M + \mathcal{U}_M,\tag{28}$$

where L_M and U_M denote the strictly lower and upper triangular part of $M(\Psi_{\mathbb{QS}})$, respectively, and D_M denotes the diagonal part of $M(\Psi_{\mathbb{QS}})$.

In Lemma 5.2, we recall the derivative expressions of a $\{1,1\}$ -QS matrix $M(\Psi_{\mathbb{QS}})$ for the parameters in $\Psi_{\mathbb{QS}}$ provided in Definition 5.1, which are useful to obtain the desired upper bounds for CNs. These results are discussed in [12].

Lemma 5.2. [12] Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ be a $\{1, 1\}$ -QS matrix. Then $M(\Psi_{\mathbb{QS}})$ has entries that are differentiable functions of $\Psi_{\mathbb{QS}}$ defined as in (27) and

$$\begin{array}{l} (1) \ \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{d}_{i}} = e_{i}^{n}(e_{i}^{n})^{\top}, \ for \ i = 1:n. \\ (2) \ \mathbf{a}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{a}_{i}} = e_{i}^{n} \mathcal{L}_{M}(i,:), \ for \ i = 2:n. \\ (3) \ \mathbf{e}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{e}_{i}} = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ M(\Psi_{\mathbb{QS}})(i+1:n,1:i-1) & \mathbf{0} \end{bmatrix} \coloneqq \mathcal{F}_{i}, \ for \ i = 2:n-1. \\ (4) \ \mathbf{b}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{b}_{i}} = \mathcal{L}_{M}(:,i)(e_{i}^{n})^{\top}, \ for \ i = 1:n-1. \\ (5) \ \mathbf{g}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{g}_{i}} = \begin{bmatrix} \mathbf{0} & M(\Psi_{\mathbb{QS}})(1:i-1,i+1:n) \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \coloneqq \mathcal{G}_{i}, \ for \ i = 2:n-1. \\ (6) \ \mathbf{f}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{f}_{i}} = e_{i}^{n} \mathcal{U}_{M}(i,:), \ for \ i = 1:n-1. \\ (7) \ \mathbf{h}_{i} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{h}_{i}} = \mathcal{U}_{M}(i,:)(e_{i}^{n})^{\top}, \ for \ i = 2:n. \end{array}$$

Next, we use the derivative expressions given in Lemma 5.2 and Theorem 3.6 to compute the bounds of the structured CNs for $M^{\dagger}(\Psi_{\mathbb{QS}})$.

Theorem 5.3. For $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ with $\operatorname{rank}(M(\Psi_{\mathbb{QS}})) = r$, we have

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}})) = \frac{\|\mathcal{X}_{\mathbb{QS}}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}})) = \left\|\frac{\mathcal{X}_{\mathbb{QS}}^{\dagger}}{M^{\dagger}}\right\|_{\max},$$

where

$$\begin{split} \mathcal{X}_{\mathbb{QS}}^{\dagger} &:= |M^{\dagger}||\mathbf{D}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}||\mathbf{L}_{M}M^{\dagger}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}\mathbf{L}_{M}^{T}||\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathbf{L}_{M}^{\top}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}\mathbf{L}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{\top}\mathbf{E}_{M}| \\ &+ |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}||\mathbf{U}_{M}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}M^{\dagger}| \\ &+ |M^{\dagger}\mathbf{U}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{U}_{M}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{U}_{M}^{\top}M^{\dagger^{\top}}M^{\dagger}| \\ &+ \sum_{i=2}^{n-1} \left(|M^{\dagger}\mathcal{F}_{i}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{F}_{i}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathcal{F}_{i}^{\top}M^{\dagger^{\top}}M^{\dagger}| \right) \\ &+ \sum_{j=2}^{n-1} \left(|M^{\dagger}\mathcal{G}_{j}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{G}_{j}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathcal{G}_{j}^{\top}M^{\dagger^{\top}}M^{\dagger}| \right), \end{split}$$

 \mathcal{F}_i and \mathcal{G}_i are defined as in Lemma 5.2.

Proof. The proof of the above assertions involves determining the contribution of each subset of parameters to the expressions provided in Theorem 3.6. Using (1) of Lemma 5.2 for the

parameters $\{\mathbf{d}_i\}_{i=1}^n$, we have

$$\mathcal{E}_{\mathbf{d}} := \sum_{i=1}^{n} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{d}_{i}} M^{\dagger}| |\mathbf{d}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{d}_{i}} \right)^{\top} \mathbf{E}_{M}| |\mathbf{d}_{i}| + |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{d}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger}| |\mathbf{d}_{i}| \right)$$
$$= \sum_{i=1}^{n} \left(|M^{\dagger} e_{i}^{n}(e_{i}^{n})^{\top} M^{\dagger}| |\mathbf{d}_{i}| + |M^{\dagger} M^{\dagger^{\top}} e_{i}^{n}(e_{i}^{n})^{\top} \mathbf{E}_{M}| |\mathbf{d}_{i}| + |\mathbf{F}_{M} e_{i}^{n}(e_{i}^{n})^{\top} M^{\dagger^{\top}} M^{\dagger}| |\mathbf{d}_{i}| \right).$$
(29)

Using (7) in (29), we deduce

$$\mathcal{E}_{\mathbf{d}} = \sum_{i=1}^{n} \left(|M^{\dagger}(:,i)| |\mathbf{d}_{i}| |M^{\dagger}(i,:)| + |M^{\dagger}M^{\dagger^{\top}}(:,i)| |\mathbf{d}_{i}| |\mathbf{E}_{M}(i,:)| + |\mathbf{F}_{M}(:,i)| |\mathbf{d}_{i}| |M^{\dagger^{\top}}M^{\dagger}(i,:)| \right) \\ = |M^{\dagger}| |\mathbf{D}_{M}| |M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}| |\mathbf{D}_{M}| |\mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{D}_{M}| |M^{\dagger^{\top}}M^{\dagger}|.$$

Similarly, for $\{\mathbf{a}_i\}_{i=2}^n$ and using (2) of Lemma 5.2, we get

$$\begin{aligned} \mathcal{E}_{\mathbf{a}} &:= \sum_{i=2}^{n} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{Q}\mathbb{S}})}{\partial \mathbf{a}_{i}} M^{\dagger}||\mathbf{a}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{Q}\mathbb{S}})}{\partial \mathbf{a}_{i}}\right)^{\top} \mathbf{E}_{M}||\mathbf{a}_{i}| \\ &+ |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{Q}\mathbb{S}})}{\partial \mathbf{a}_{i}}\right)^{\top} M^{\dagger^{\top}} M^{\dagger}||\mathbf{a}_{i}| \right) \\ &= \sum_{i=2}^{n} \left(|M^{\dagger} e_{i}^{n} \mathbf{L}_{M}(i,:) M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{L}_{M}^{\top}(:,i)(e_{i}^{n})^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathbf{L}_{M}^{\top}(:,i)(e_{i}^{n})^{\top} M^{\dagger^{\top}} M^{\dagger}| \right) \\ &= |M^{\dagger}||\mathbf{L}_{M} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{L}_{M}^{\top}||\mathbf{E}_{M}| + |\mathbf{F}_{M} \mathbf{L}_{M}^{\top}||M^{\dagger^{\top}} M^{\dagger}|. \end{aligned}$$

For the parameters $\{\mathbf{b}_i\}_{i=1}^{n-1}$ and using (4) of Lemma 5.2, we deduce

$$\begin{aligned} \mathcal{E}_{\mathbf{b}} &:= \sum_{i=2}^{n} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{b}_{i}} M^{\dagger}| |\mathbf{b}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{b}_{i}} \right)^{\top} \mathbf{E}_{M} ||\mathbf{b}_{i}| \right. \\ &+ |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{b}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} ||\mathbf{b}_{i}| \right) \\ &= |M^{\dagger} \mathbf{L}_{M} ||M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} ||\mathbf{L}_{M}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{L}_{M}^{\top} M^{\dagger^{\top}} M^{\dagger}|. \end{aligned}$$

For the parameters $\{\mathbf{e}_i\}_{i=2}^{n-1}$, using (3) of Lemma 5.2, we have

$$\begin{aligned} \mathcal{E}_{\mathbf{e}} &:= \sum_{i=2}^{n-1} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{e}_{i}} M^{\dagger}| |\mathbf{e}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{e}_{i}} \right)^{\top} \mathbf{E}_{M}| |\mathbf{e}_{i}| \right) \\ &+ |\mathbf{F} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{e}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger}| |\mathbf{e}_{i}| \right) \\ &= \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{F}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{F}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{F}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right). \end{aligned}$$

In a similar approach, for the parameters $\{\mathbf{f}_i\}_{i=1}^{n-1}$, $\{\mathbf{g}_i\}_{i=2}^{n-1}$ and $\{\mathbf{h}_i\}_{i=2}^{n-1}$, we get

$$\mathcal{E}_{\mathbf{f}} := \sum_{i=1}^{n-1} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{f}_{i}} M^{\dagger}| |\mathbf{f}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{f}_{i}} \right)^{\top} \mathbf{E}_{M} ||\mathbf{f}_{i}| + |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{f}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} ||\mathbf{f}_{i}| \right)$$
$$= |M^{\dagger}| |\mathbf{U}_{M} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{U}_{M}^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M} \mathbf{U}_{M}^{\top}| |M^{\dagger^{\top}} M^{\dagger}|.$$

$$\begin{aligned} \mathcal{E}_{\mathbf{h}} &:= \sum_{i=2}^{n} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{h}_{i}} M^{\dagger}| |\mathbf{h}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{h}_{i}} \right)^{\top} \mathbf{E}_{M} ||\mathbf{h}_{i}| \\ &+ |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{h}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} ||\mathbf{h}_{i}| \right) \\ &= |M^{\dagger} \mathbf{U}_{M}| |M^{\dagger}| + |M^{\dagger}| |M^{\dagger^{\top}} \mathbf{U}_{M}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{U}_{M}^{\top} M^{\dagger^{\top}} M^{\dagger}|. \end{aligned}$$

$$\begin{split} \mathcal{E}_{\mathbf{g}} &:= \sum_{i=2}^{n-1} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{g}_{i}} M^{\dagger}||\mathbf{g}_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{g}_{i}} \right)^{\top} \mathbf{E}_{M}||\mathbf{g}_{i}| \\ &+ |\mathbf{F}_{M} \left(\frac{\partial M(\Psi_{\mathbb{QS}})}{\partial \mathbf{g}_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger}||\mathbf{g}_{i}| \right) \\ &= \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{G}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{G}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{G}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right). \end{split}$$

Now, it is straightforward from Theorem 3.6 that

$$\mathcal{X}_{\mathbb{QS}}^{\dagger} = \mathcal{E}_{\mathbf{d}} + \mathcal{E}_{\mathbf{a}} + \mathcal{E}_{\mathbf{b}} + \mathcal{E}_{\mathbf{e}} + \mathcal{E}_{\mathbf{f}} + \mathcal{E}_{\mathbf{g}} + \mathcal{E}_{\mathbf{h}},$$

and hence, the proof is completed.

Remark 5. When $M(\Psi_{\mathbb{QS}})$ is a $n \times n$ nonsingular $\{1, 1\}$ -QS matrix, we have $\mathbf{E}_M = \mathbf{0}$. Then, using the expressions in Theorem 3.9, and an analogous way to Theorem 5.3, exact formulae of the structured CNs can be deduced for the inversion of $M(\Psi_{\mathbb{QS}})$.

For any $\{1,1\}$ -QS matrix $M(\Psi_{\mathbb{QS}})$, it is worth noting that there exist infinitely many QS representations, as indicated by Dopico and Pomés [13]. The next result demonstrates that $\tilde{\mathcal{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathcal{C}}^{\dagger}((\Psi_{\mathbb{QS}}))$ are independent of the QS representation used.

Proposition 5.4. For any two representations $\Psi_{\mathbb{QS}}$ and $\Psi'_{\mathbb{QS}}$ of a $\{1,1\}$ -QS matrix $M \in \mathbb{R}^{m \times n}$, we get

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}})) = \tilde{\mathscr{M}}^{\dagger}(M(\Psi'_{\mathbb{QS}})) \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}})) = \tilde{\mathscr{C}}^{\dagger}(M(\Psi'_{\mathbb{QS}})).$$

Proof. Observe that the formulae given by the Theorem 5.3 only depend on the entries of $M, M^{\dagger}, \mathbf{E}_{M}$ and \mathbf{F}_{M} , but not on the specific selection of the parameter set; hence, the proof follows.

We provide upper bounds on the structured CNs of the MNLS solution of the LS problem (15) corresponding to $\{1, 1\}$ -QS matrices in the next theorem.

Theorem 5.5. Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ be such that $\operatorname{rank}(M(\Psi_{\mathbb{QS}})) = r$ and $b \in \mathbb{R}^n$. Set $\mathbf{r} :=$ $b - M(\Psi_{\mathbb{OS}})\mathbf{x}$. Then

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}), b) = \frac{\|\mathscr{X}_{\mathbb{QS}}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}), b) = \left\|\frac{\mathscr{X}_{\mathbb{QS}}^{ls}}{\mathbf{x}}\right\|_{\infty},$$

where

$$\begin{split} \mathcal{X}_{\mathbb{QS}}^{ls} &:= |M^{\dagger}||b| + |M^{\dagger}||\mathbf{D}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||\mathbf{L}_{M}\mathbf{x}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}\mathbf{L}_{M}^{T}||\mathbf{r}| + |\mathbf{F}_{M}\mathbf{L}_{M}^{\top}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}\mathbf{L}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{\top}\mathbf{r}| \\ &+ |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}|M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||\mathbf{U}_{M}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{r}| + |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}\mathbf{x}| \\ &+ |M^{\dagger}\mathbf{U}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{U}_{M}^{\top}\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{U}_{M}^{\top}M^{\dagger^{\top}}\mathbf{x}| \\ &+ \sum_{i=2}^{n-1} \left(|M^{\dagger}\mathcal{F}_{i}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{F}_{i}^{\top}\mathbf{r}| + |\mathbf{F}_{M}\mathcal{F}_{i}^{\top}M^{\dagger^{\top}}\mathbf{x}| \right) \\ &+ \sum_{j=2}^{n-1} \left(|M^{\dagger}\mathcal{G}_{j}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{G}_{j}^{\top}\mathbf{r}| + |\mathbf{F}_{M}\mathcal{G}_{j}^{\top}M^{\dagger^{\top}}\mathbf{x}| \right). \end{split}$$

Proof. The statement can be easily verified using a similar justification to that of the proof of the Theorem 5.3 and using the Theorem 3.12. Hence, we omit the proof.

Remark 6. By considering $M(\Psi_{\mathbb{QS}})$ nonsingular, for $A(\Psi_{\mathbb{QS}})\mathbf{x} = b$, using the expression in Theorem 3.14 and $\mathbf{r} = 0$, we can obtain following expression for the structured MCN of \mathbf{x} :

$$\mathcal{M}^{\dagger}(M(\Psi_{\mathbb{QS}}), b) = \frac{1}{\|\mathbf{x}\|_{\infty}} \left\| |M^{-1}||b| + |M^{-1}||\mathbf{D}_{M}||\mathbf{x}| + |M^{-1}||\mathbf{L}_{M}\mathbf{x}| + |M^{-1}\mathbf{L}_{M}||\mathbf{x}| + |M^{-1}||\mathbf{U}_{M}\mathbf{x}| + |M^{-1}\mathbf{U}_{M}||\mathbf{x}| + \sum_{i=2}^{n-1} |M^{-1}\mathcal{F}_{i}\mathbf{x}| + \sum_{j=2}^{n-1} |M^{-1}\mathcal{G}_{j}\mathbf{x}| \right\|_{\max}.$$

This result is the same as obtained by Dopico and Pomés [13].

5.2. CNs corresponding to GV representation

The GV representation, proposed initially in [30], is another essential representation for $\{1, 1\}$ -QS matrices. This representation is used to enhance the stability of fast algorithms. In this subsection, we first review the GV representation together with its minor variant called GV representation through tangent. Then, discuss the results for structured CNs corresponding to this representation.

Definition 5.6. [30] Any $M \in \mathbb{R}^{n \times n}$ is classified to be a $\{1, 1\}$ -QS matrix if it can be represented by the parameter set

$$\Psi_{\mathbb{QS}}^{\mathcal{GV}} = \left[\{p_i, q_i\}_{i=2}^{n-1}, \{u_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n, \{v_i\}_{i=1}^{n-1}, \{r_i, s_i\}_{i=2}^{n-1} \right]^\top \in \mathbb{R}^{7n-10},$$
(30)

satisfying the following properties,

- $\begin{array}{ll} (1) \ \{p_i,q_i\} \text{ is a cosine-sine pair with } p_i^2 + q_i^2 = 1, \text{ for every } i \in \{2:n-1\}, \\ (2) \ \{u_i\}_{i=1}^{n-1}, \ \{\mathbf{d}_i\}_{i=1}^n, \text{ and } \{v_i\}_{i=1}^{n-1} \text{ are independent parameters,} \\ (3) \ \{r_i,s_i\} \text{ is a cosine-sine pair with } r_i^2 + s_i^2 = 1, \text{ for every } i \in \{2:n-1\}, \end{array}$

as follows:

$$M = \begin{bmatrix} \mathbf{d}_1 & v_1 r_2 & v_1 s_2 r_3 & \dots & v_1 s_2 \dots s_{n-2} r_{n-1} & v_1 s_2 \dots s_{n-1} \\ p_2 u_1 & \mathbf{d}_2 & v_2 r_3 & \dots & v_2 s_3 \dots s_{n-2} r_{n-1} & v_2 s_3 \dots s_{n-1} \\ p_3 q_2 u_1 & p_3 u_2 & \mathbf{d}_3 & \dots & v_3 s_4 \dots s_{n-2} r_{n-1} & v_3 s_4 \dots s_{n-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ p_{n-1} q_{n-2} \dots q_2 u_1 & p_{n-1} q_{n-2} \dots q_3 u_2 & p_{n-1} q_{n-2} \dots q_4 u_3 & \dots & \mathbf{d}_{n-1} & v_{n-1} \\ q_{n-1} \dots q_2 u_1 & q_{n-1} \dots q_3 u_2 & q_{n-1} \dots q_4 u_3 & \dots & u_{n-1} & \mathbf{d}_n \end{bmatrix}.$$

Note that the $\Psi_{\mathbb{QS}}^{\mathcal{GV}}$ is a special case of $\Psi_{\mathbb{QS}}$ by considering $\{\mathbf{a}_i, \mathbf{e}_i\}_{i=2}^{n-1} = \{p_i, q_i\}_{i=2}^{n-1}, \{\mathbf{b}_i\}_{i=1}^{n-1} = \{u_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n = \{\mathbf{d}_i\}_{i=1}^n, \{\mathbf{f}_i\}_{i=1}^{n-1} = \{v_i\}_{i=1}^{n-1}, \{\mathbf{g}_i, \mathbf{h}_i\}_{i=2}^{n-1} = \{s_i, r_i\}_{i=2}^{n-1}, \text{ and } \mathbf{a}_n = \mathbf{h}_n = 1$ with additional conditions on the parameters. Since the parameters p_i and q_i are dependent, arbitrary perturbation to $\Psi_{\mathbb{QS}}^{\mathcal{GV}}$ will destroy the cosine-sine pairs and the same is true for r_i and s_i . Thus, it will be more sensible to restrict the perturbation that preserves the cosine-sine pair. Consequently, Dopico and Pomés in [12] introduced a new representation called GV representation through tangent using their tangents.

Definition 5.7. For the GV representation $\Psi_{\mathbb{QS}}^{\mathcal{GV}}$ as in (30), the GV representation through tangent is defined as

$$\Psi_{\mathcal{GV}} = [\{t_i\}_{i=2}^{n-1}, \{u_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n, \{v_i\}_{i=1}^{n-1}, \{w_i\}_{i=2}^{n-1}]^\top \in \mathbb{R}^{5n-6},$$
(31)

where $p_i = \frac{1}{\sqrt{1+t_i^2}}$, $q_i = \frac{t_i}{\sqrt{1+t_i^2}}$ and $r_i = \frac{1}{\sqrt{1+w_i^2}}$, $s_i = \frac{w_i}{\sqrt{1+w_i^2}}$, for i = 2: n-1.

We employ the notation $M(\Psi_{\mathcal{GV}})$ to refer a $\{1,1\}$ -QS matrix parameterized by the set $\Psi_{\mathcal{GV}}$. The derivative expressions corresponding to the parameters in the representation $\Psi_{\mathcal{GV}}$ are revisited in the next lemma:

Lemma 5.8. [12] Let $M(\Psi_{\mathcal{GV}}) \in \mathbb{R}^{n \times n}$ having rank $(M(\Psi_{\mathcal{GV}})) = r$. Then each entry of $M(\Psi_{\mathcal{GV}})$ is differentiable functions of the elements in $\Psi_{\mathcal{GV}}$, and

$$(1) \ t_i \frac{\partial M(\Psi_{\mathcal{GV}})}{\partial t_i} = \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ -q_i^2 M(\Psi_{\mathcal{GV}})(i, 1:i-1) & \mathbf{0} \\ p_i^2 M(\Psi_{\mathcal{GV}})(i+1:n, 1:i-1) & \mathbf{0} \end{bmatrix} := \mathcal{K}_i, \ for \ i = 2:n-1.$$

$$(2) \ w_i \frac{\partial M(\Psi_{\mathcal{GV}})}{\partial w_i} = \begin{bmatrix} \mathbf{0} & -s_i^2 M(\Psi_{\mathcal{GV}})(1:i-1,i) & r_i^2 M(\Psi_{\mathcal{GV}})(1:i-1,i+1:n) \\ \mathbf{0} & \mathbf{0} & \mathbf{0} \end{bmatrix} := \mathcal{L}_i, \ for \ i = 2:n-1.$$

Note: Partial derivative expressions corresponding to the parameters $\{u_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n$ and $\{v_i\}_{i=1}^{n-1}$ are same as the expression for the parameters $\{\mathbf{b}_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n$ and $\{\mathbf{f}_i\}_{i=1}^{n-1}$, respectively, given in the Lemma 5.2.

In Theorem 5.9, we discuss computationally feasible upper bounds for the structured CNs introduced in Definition 3.4 corresponding to the representation Ψ_{GV} .

Theorem 5.9. Let $M(\Psi_{\mathcal{GV}}) \in \mathbb{R}^{n \times n}$ with $\operatorname{rank}(M(\Psi_{\mathcal{GV}})) = r$, then

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{G}\mathcal{V}})) = \frac{\|\mathscr{X}_{\mathcal{G}\mathcal{V}}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{G}\mathcal{V}})) = \left\|\frac{\mathscr{X}_{\mathcal{G}\mathcal{V}}^{\dagger}}{M^{\dagger}}\right\|_{\max},$$

where

$$\begin{aligned} \mathcal{X}_{\mathcal{GV}}^{\dagger} &:= |M^{\dagger}||\mathbf{D}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}\mathbf{L}_{M}||M^{\dagger}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}||\mathbf{U}_{M}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{E}_{M}| \\ &+ |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}M^{\dagger}| + \sum_{i=2}^{n-1} \left(|M^{\dagger}\mathcal{K}_{i}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{K}_{i}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathcal{K}_{i}^{\top}M^{\dagger^{\top}}M^{\dagger}| \right) \\ &+ \sum_{j=2}^{n-1} \left(|M^{\dagger}\mathcal{L}_{j}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{L}_{j}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathcal{L}_{j}^{\top}M^{\dagger^{\top}}M^{\dagger}| \right). \end{aligned}$$

Here, \mathcal{K}_i and \mathcal{L}_i are defined as in Lemma 5.8.

Proof. To prove the claim, we need to find the contribution of each subset of parameters in $\Psi_{\mathcal{GV}}$ to the expressions given in Theorem 3.6. Since the derivative expressions for the parameters $\{u_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n$ and $\{v_i\}_{i=1}^{n-1}$ are same as the expressions for the parameters $\{\mathbf{b}_i\}_{i=1}^{n-1}, \{\mathbf{d}_i\}_{i=1}^n$ and $\{\mathbf{f}_i\}_{i=1}^{n-1}$, respectively, given in the Lemma 5.2, contribution for these parameters in the expressions given in Theorem 3.6 are same as $\mathcal{E}_{\mathbf{b}}$, $\mathcal{E}_{\mathbf{d}}$ and $\mathcal{E}_{\mathbf{f}}$, respectively. Therefore, we set $\mathcal{E}_u := \mathcal{E}_{\mathbf{b}}$ and $\mathcal{E}_v := \mathcal{E}_{\mathbf{f}}$. For the parameters $\{t_i\}_{i=2}^{n-1}$, we have

$$\begin{aligned} \mathcal{E}_t &:= \sum_{i=2}^{n-1} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial t_i} M^{\dagger}| |t_i| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial t_i} \right)^{\top} \mathbf{E}_M ||t_i| + |\mathbf{F}_M \left(\frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial t_i} \right)^{\top} M^{\dagger^{\top}} M^{\dagger} ||t_i| \right) \\ &= \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{K}_i M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{K}_i^{\top} \mathbf{E}_M || + |\mathbf{F}_M \mathcal{K}_i^{\top} M^{\dagger^{\top}} M^{\dagger}| \right). \end{aligned}$$

For the parameters $\{w_i\}_{i=2}^{n-1}$, we have

$$\mathcal{E}_{w} := \sum_{i=2}^{n-1} \left(|M^{\dagger} \frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial w_{i}} M^{\dagger}| |w_{i}| + |M^{\dagger} M^{\dagger^{\top}} \left(\frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial w_{i}} \right)^{\top} \mathbf{E}_{M}| |w_{i}| + |\mathbf{F} \left(\frac{\partial M(\Psi_{\mathcal{G}\mathcal{V}})}{\partial w_{i}} \right)^{\top} M^{\dagger^{\top}} M^{\dagger}| |w_{i}| \right)$$
$$= \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{L}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{L}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{L}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right).$$

Now, from Theorem 3.6, we have $\mathcal{X}_{\mathcal{GV}}^{\dagger} = \mathcal{E}_{\mathbf{d}} + \mathcal{E}_{v} + \mathcal{E}_{t} + \mathcal{E}_{w}$, and hence, the proof follows.

The bounds of the structured CNs for x with respect to the representation Ψ_{GV} are given in the next theorem.

Theorem 5.10. Let $M(\Psi_{\mathcal{GV}}) \in \mathbb{R}^{n \times n}$ with $\operatorname{rank}(M(\Psi_{\mathcal{GV}})) = r$. Set $\mathbf{r} := b - M(\Psi_{\mathcal{GV}})\mathbf{x}$, then

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{G}\mathcal{V}}), b) = \frac{\|\mathcal{X}_{\mathcal{G}\mathcal{V}}^{ls}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad and \quad \tilde{\mathscr{C}}(M(\Psi_{\mathcal{G}\mathcal{V}}), b) = \left\|\frac{\mathcal{X}_{\mathcal{G}\mathcal{V}}^{ls}}{\mathbf{x}}\right\|_{\infty},$$

where

$$\begin{aligned} \mathcal{X}_{\mathcal{GV}}^{ls} &:= |M^{\dagger}||\mathbf{D}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}\mathbf{L}_{M}||\mathbf{x}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{\top}\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||\mathbf{U}_{M}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{r}| \\ &+ |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}\mathbf{x}| + \sum_{i=2}^{n} \left(|M^{\dagger}\mathcal{K}_{i}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{K}_{i}^{\top}\mathbf{r}| + |\mathbf{F}_{M}\mathcal{K}_{i}^{\top}M^{\dagger^{\top}}\mathbf{x}| \right) \\ &+ \sum_{i=2}^{n} \left(|M^{\dagger}\mathcal{L}_{i}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathcal{L}_{i}^{\top}\mathbf{r}| + |\mathbf{F}_{M}\mathcal{L}_{i}^{\top}M^{\dagger^{\top}}\mathbf{x}| \right) + |M^{\dagger}||b|. \end{aligned}$$

Here, \mathcal{K}_i and \mathcal{L}_i are defined as in Lemma 5.8.

Proof. The proof follows by using the similar proof technique of Theorem 5.9. Hence, we omit it.

5.3. Comparisons between different CNs for $\{1,1\}$ -QS matrices

We compare structured and unstructured CNs for the M-P inverse in Proposition 5.11 and the MNLS solution in Proposition 5.12 for $\{1,1\}$ -QS matrix. For unstructured CNs, we use the expressions given in Corollary 3.7 and Corollary 3.13. Relationships between the CNs for representations Ψ_{QS} and Ψ_{GV} are also investigated in Propositions 5.13 and 5.14.

The next result describes that structured CNs for the parameter set $\Psi_{\mathbb{QS}}$ are smaller than unstructured ones for the M-P inverse up to an order of n.

Proposition 5.11. Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ be such that $\operatorname{rank}(M(\Psi_{\mathbb{QS}})) = r$, then we get the following relations

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{OS}})) \leq n \; \tilde{\mathscr{M}}^{\dagger}(M) \; and \; \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{OS}})) \leq n \; \tilde{\mathscr{C}}^{\dagger}(M).$$

Proof. Using the properties of absolute values and Theorem 5.3, we have

$$\begin{split} \mathcal{X}_{\mathbb{QS}}^{\dagger} &\leq |M^{\dagger}||\mathbf{D}_{M}||M^{\dagger}|+|M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{E}_{M}|+|\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}M^{\dagger}| \\ &+ 2|M^{\dagger}||\mathbf{L}_{M}||M^{\dagger}|+2|M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{T}||\mathbf{E}_{M}|+2|\mathbf{F}_{M}||\mathbf{L}_{M}^{T}||M^{\dagger^{\top}}M^{\dagger}| \\ &+ 2|M^{\dagger}||\mathbf{U}_{M}||M^{\dagger}|+2|M^{\dagger}M^{\dagger^{\top}}||\mathbf{U}_{M}^{T}||\mathbf{E}_{M}|+2|\mathbf{F}_{M}||\mathbf{U}_{M}^{T}||M^{\dagger^{\top}}M^{\dagger}| \\ &+ \sum_{i=2}^{n-1} \left(|M^{\dagger}||\mathcal{F}_{i}||M^{\dagger}|+|M^{\dagger}M^{\dagger^{\top}}||\mathcal{F}_{i}^{\top}||\mathbf{E}_{M}|+|\mathbf{F}_{M}||\mathcal{F}_{i}^{\top}||M^{\dagger^{\top}}M^{\dagger}|\right) \\ &+ \sum_{i=2}^{n-1} \left(|M^{\dagger}||\mathcal{G}_{i}||M^{\dagger}|+|M^{\dagger}M^{\dagger^{\top}}||\mathcal{G}_{i}^{\top}||\mathbf{E}_{M}|+|\mathbf{F}_{M}||\mathcal{G}_{i}^{\top}||M^{\dagger^{\top}}M^{\dagger}|\right). \end{split}$$

Using $|\mathcal{F}_i| \leq |\mathcal{L}_M|$ and $|\mathcal{G}_i| \leq |\mathcal{U}_M|$, we get

$$\mathcal{X}_{\mathbb{QS}}^{\dagger} \leq n \left(|M^{\dagger}| |M| |M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}| |M^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M}| |M^{\top}| |M^{\dagger^{\top}}M^{\dagger}| \right).$$

Therefore, the desired relations can be obtained from Theorem 5.3 and Corollary 3.7.

A similar type of result also holds for the LS problem, which is given next. We remove the proof since it is analogous to Proposition 5.11.

Proposition 5.12. Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ be as in Proposition 5.11 and $b \in \mathbb{R}^m$. Then, we get the following relations

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b) \leq n \ \tilde{\mathscr{M}}^{\dagger}(M,b) \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b) \leq \ n \ \tilde{\mathscr{C}}^{\dagger}(M,b).$$

Next result discuss about the relationship between the $\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ with $\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$ and $\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ with $\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$.

Proposition 5.13. For the representations $\Psi_{\mathbb{QS}}$ and $\Psi_{\mathcal{GV}}$ of a $\{1,1\}$ -QS matrix $M \in \mathbb{R}^{n \times n}$ with rank(M) = r, following holds:

$$\widetilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}})) \leq \widetilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}})) \quad and \quad \widetilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}})) \leq \widetilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}})).$$

Proof. The proof will be followed by observing that

$$\begin{aligned} \mathcal{K}_{i} &= \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ -q_{i}^{2}M(i,1:i-1) & \mathbf{0} \\ p_{i}^{2}M(i+1:n,1:i-1) & \mathbf{0} \end{bmatrix} = -q_{i}^{2} \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ M(i,1:i-1) & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} + p_{i}^{2} \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \\ M(i+1:n,1:i-1) & \mathbf{0} \end{bmatrix} \\ &= -q_{i}^{2} e_{i}^{m} \mathcal{L}_{M}(i,:) + p_{i}^{2} \mathcal{F}_{i}. \end{aligned}$$

Now, using the properties $|p_i|^2 \le 1$ and $|q_i|^2 \le 1$, and (7), we obtain

$$\sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{K}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{K}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{K}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right) \leq |M^{\dagger}| |\mathbf{L}_{M} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{L}_{M}^{\top}| |\mathbf{E}_{M}| + |\mathbf{F}_{M} \mathbf{L}_{M}^{\top}| |M^{\dagger^{\top}} M^{\dagger}| + \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{F}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{F}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{F}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right).$$

Similarly, we can write $\mathcal{L}_i = -s_i^2 U_M(:,i)(e_i^n)^\top + r_i^2 \mathcal{G}_i$. Therefore, using $|s_i|^2 \leq 1$ and $|r_i|^2 \leq 1$, and (7), we get

$$\sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{L}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{L}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{L}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right) \leq |M^{\dagger} \mathbf{U}_{M}| |M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}}| |\mathbf{U}_{M}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{U}_{M}^{\top} M^{\dagger^{\top}} M^{\dagger}| + \sum_{i=2}^{n-1} \left(|M^{\dagger} \mathcal{G}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{G}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{G}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right).$$

Hence, we get the desired relations from the above two inequalities, expressions from the Theorems 5.3 and 5.9. $\hfill\blacksquare$

Proposition 5.14 provides the relationship between CNs for LS problem (15) for any $\{1, 1\}$ -QS matrix corresponding to the parameter sets $\Psi_{\mathbb{QS}}$ and $\Psi_{\mathcal{GV}}$.

Proposition 5.14. For the representations $\Psi_{\mathbb{QS}}$ and $\Psi_{\mathcal{GV}}$ of a $\{1,1\}$ -QS matrix $M \in \mathbb{R}^{n \times n}$ having rank r and $b \in \mathbb{R}^n$, following holds:

$$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}), b) \leq \tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}), b) \quad and \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}), b) \leq \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}), b).$$

5.4. The structured effective CNs

The expressions in Theorems 5.3 and 5.5 can be computationally very expensive for large matrices due to the involvement of two sums. The effective CN for $\{1, 1\}$ -QS matrices was initially

considered by Dopico and Pomés in [12,13] for eigenvalue problem and linear system to reduce the computation complexity. In a similar fashion to avoid such problems, we propose in Definition 5.15, structured effective CNs $\tilde{\mathcal{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathcal{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$, which have similar contribution as $\tilde{\mathcal{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathcal{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$, respectively.

Definition 5.15. Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ having $\operatorname{rank}(M(\Psi_{\mathbb{QS}})) = r$. Then for $M^{\dagger}(\Psi_{\mathbb{QS}})$, we define structured effective MCN and CCN as

$$\widetilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) := \frac{\|\mathscr{X}_{f,\mathbb{QS}}^{\dagger}\|_{\max}}{\|M^{\dagger}\|_{\max}} \quad \text{and} \quad \widetilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) := \left\| \frac{\mathscr{X}_{f,\mathbb{QS}}^{\dagger}}{M^{\dagger}} \right\|_{\max}$$

where

$$\begin{split} \mathcal{X}_{f,\mathbb{QS}}^{\dagger} &:= |M^{\dagger}||\mathbf{D}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}||\mathbf{L}_{M}M^{\dagger}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}\mathbf{L}_{M}^{T}||\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathbf{L}_{M}^{\top}||M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}\mathbf{L}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{\top}\mathbf{E}_{M}| \\ &+ |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}M^{\dagger^{\top}}M^{\dagger}| + |M^{\dagger}||\mathbf{U}_{M}M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{E}_{M}| + |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}M^{\dagger}| \\ &+ |M^{\dagger}\mathbf{U}_{M}||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{U}_{M}^{\top}\mathbf{E}_{M}| + |\mathbf{F}_{M}||\mathbf{U}_{M}^{\top}M^{\dagger^{\top}}M^{\dagger}|. \end{split}$$

The following theorem demonstrates that the contribution of the sum terms in the expression of $\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ are negligible and reliably estimated up to a multiple of n.

Theorem 5.16. Under the same hypothesis as in Definition 5.15, following relations holds

$$\begin{split} \tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) &\leq \tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}})) \leq (n-1)\,\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})), \\ \tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) &\leq \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}})) \leq (n-1)\,\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})). \end{split}$$

Proof. The inequalities on the left side are trivial. For inequalities on the right side, we set

$$\begin{bmatrix} \mathbf{0} & \mathbf{0} \\ M(\Psi_{\mathbb{QS}})(i+1:n,1:i-1) & \mathbf{0} \end{bmatrix} = \begin{bmatrix} \mathbf{0} \\ \mathbf{L}_M(i+1:n,:) \end{bmatrix} + \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & -\mathbf{L}_M(i+1:n,i:n) \end{bmatrix}.$$

By using the above, we get

$$\left(|M^{\dagger} \mathcal{F}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{F}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{F}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right) \leq |M^{\dagger}| |\mathbf{L}_{M} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{L}_{M}^{T}| |\mathbf{E}_{M}|$$
$$+ |\mathbf{F}_{M} \mathbf{L}_{M}^{\top}| |M^{\dagger^{\top}} M^{\dagger}| + |M^{\dagger} \mathbf{L}_{M}| |M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}}| |\mathbf{L}_{M}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{L}_{M}^{\top}| M^{\dagger^{\top}} M^{\dagger}|.$$

Again,

$$\begin{bmatrix} \mathbf{0} & M(\Psi_{\mathbb{QS}})(1:i-1,i+1:n) \\ \mathbf{0} & \mathbf{0} \end{bmatrix} = \begin{bmatrix} \mathbf{0} & \mathbf{U}_M(:,i+1:n) \end{bmatrix} + \begin{bmatrix} \mathbf{0} & \mathbf{0} \\ \mathbf{0} & -\mathbf{U}_M(i:n,i+1:n) \end{bmatrix}.$$

By using the above, we get

$$\left(|M^{\dagger} \mathcal{G}_{i} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathcal{G}_{i}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M} \mathcal{G}_{i}^{\top} M^{\dagger^{\top}} M^{\dagger}| \right) \leq |M^{\dagger}| |\mathbf{U}_{M} M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}} \mathbf{U}_{M}^{\top}| |\mathbf{E}_{M}|$$
$$+ |\mathbf{F}_{M} \mathbf{U}_{M}^{\top}| |M^{\dagger^{\top}} M^{\dagger}| + |M^{\dagger} \mathbf{U}_{M}| |M^{\dagger}| + |M^{\dagger} M^{\dagger^{\top}}| |\mathbf{U}_{M}^{\top} \mathbf{E}_{M}| + |\mathbf{F}_{M}| |\mathbf{U}_{M}^{\top} M^{\dagger^{\top}} M^{\dagger}|$$

Hence, the desired result is straightforward from Definition 5.15.

Next, we compare the relation among effective CNs to the unstructured CNs given in Corollary 3.7.

Proposition 5.17. Under the same hypothesis as in Definition 5.15, following relations holds

$$\widetilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) \leq 2 \ \widetilde{\mathscr{M}}^{\dagger}(M) \quad and \quad \widetilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}})) \leq 2 \ \widetilde{\mathscr{C}}^{\dagger}(M).$$

Proof. By using properties of absolutes values and expression of $\mathcal{X}_{f,\mathbb{QS}}^{\dagger}$ given the Definition 5.15, we have

$$\mathcal{X}_{f,\mathbb{QS}}^{\dagger} \leq 2 \left(|M^{\dagger}||M||M^{\dagger}| + |M^{\dagger}M^{\dagger^{\top}}||M^{\top}||\mathbf{E}_{M}| + |\mathbf{F}_{M}||M^{\top}||M^{\dagger^{\top}}M^{\dagger}| \right).$$

Hence, the desired results attained from Definition 5.15.

Next, similarly to the Definition 5.15, we define structured effective CNs for the MNLS solution.

Definition 5.18. Let $M(\Psi_{\mathbb{QS}}) \in \mathbb{R}^{n \times n}$ having rank $(M(\Psi_{\mathbb{QS}})) = r$ and $b \in \mathbb{R}^m$. Then, for the MNLS solution **x**, we define structured effective MCN and CCN as follows:

$$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b) \quad := \frac{\|\mathscr{X}_{f,\mathbb{QS}}^{l_{s}}\|_{\infty}}{\|\mathbf{x}\|_{\infty}} \quad \text{and} \quad \tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b) := \left\|\frac{\mathscr{X}_{f,\mathbb{QS}}^{l_{s}}}{\mathbf{x}}\right\|_{\infty},$$

where

$$\begin{aligned} \mathcal{X}_{f,\mathbb{QS}}^{ls} : &= |M^{\dagger}||\mathbf{D}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{D}_{M}||\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{D}_{M}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||\mathbf{L}_{M}\mathbf{x}| \\ &+ |M^{\dagger}M^{\dagger^{\top}}\mathbf{L}_{M}^{T}||\mathbf{r}| + |\mathbf{F}_{M}\mathbf{L}_{M}^{\top}||M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}\mathbf{L}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{L}_{M}^{T}\mathbf{r}| \\ &+ |\mathbf{F}_{M}||\mathbf{L}_{M}^{\top}M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||\mathbf{U}_{M}\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}\mathbf{U}_{M}^{\top}||\mathbf{r}| + |\mathbf{F}_{M}\mathbf{U}_{M}^{\top}||M^{\dagger^{\top}}\mathbf{x}| \\ &+ |M^{\dagger}\mathbf{U}_{M}||\mathbf{x}| + |M^{\dagger}M^{\dagger^{\top}}||\mathbf{U}_{M}^{\top}\mathbf{r}| + |\mathbf{F}_{M}||\mathbf{U}_{M}^{\top}M^{\dagger^{\top}}\mathbf{x}| + |M^{\dagger}||b|. \end{aligned}$$

Similar results also hold for the MNLS solution as discussed in Theorem 5.16 and Proposition 5.17.

6. Numerical experiments

We reported a few numerical experiments in this section to illustrate the theoretical findings covered in Sections 4 and 5, respectively. For all numerical computations, we have used MATLAB R2022b, with unit roundoff errors 10^{-16} . Also, for structured and unstructured CNs, we give a comparison between their upper bounds. For unstructured CNs, we use the upper bounds for the M-P inverse and the MNLS solution given in Corollary 3.7 and 3.13, respectively. These examples reveal that structured CNs are significantly less than unstructured ones.

Example 6.1. Let $\Psi_{\mathbb{CV}} = [c = [1, 1, \frac{1}{2}, \frac{-1}{30}, \frac{1}{40}], d = [12, -0.75 \times 10^7, 25 \times 10^3\}]^{\top} \in \mathbb{R}^8$ be the parameter set of a 5 × 6 CV matrix M. For the M-P inverse, we compute the bounds for the structured MCN and CCN using Theorem 4.3. For the MNLS solution, we generate a random vector $b \in \mathbb{R}^5$ in MATLAB by the command *randn* and use Theorem 4.5 to compute the structured CNs. The computed results for the bounds of structured and unstructured CNs are listed in Table 1. We observed that the bounds for the structured CNs are of order one, whereas for the unstructured case, they are significantly large.

Example 6.2. [39] Let $M \in \mathbb{R}^{12 \times 20}$ be a CV matrix as in Definition 5.1 with the parameter set $\Psi_{\mathbb{CV}} = [\{c_i\}_{i=1}^{12}, \{d_i\}_{i=1}^{8}]^\top \in \mathbb{R}^{20}$, where

$$\begin{cases} c_i = \frac{i}{20}, & i = 1:12, \\ d_j = \frac{j+4}{50}, & j = 1:8. \end{cases}$$
(32)

Table 1.: Comparison between upper bounds of structured and unstructured CNs for $M^{\dagger}(\Psi_{\mathbb{CV}})$ and $M^{\dagger}(\Psi_{\mathbb{CV}}, b)$ for Example 6.1.

$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{C}^{\dagger}}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{M}^{\dagger}}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}),b)$	$\tilde{\mathscr{C}}^{\dagger}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}),b)$
1.9309e + 04	8.4149	2.6103e+0 6	63.7873	3.9137e+0 4	12.3655	2.1903e+0 6	12.3655

Table 2.: Comparison between upper bounds of structured and unstructured CNs for $M^{\dagger}(\Psi_{\mathbb{CV}})$ and $M^{\dagger}(\Psi_{\mathbb{CV}}, b)$ for Example 6.2.

$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{C}^{\dagger}}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}),b)$	$\tilde{\mathscr{C}}^{\dagger}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}),b)$
1.1568e + 05	8.5419e+01	5.3826e + 07	3.3542e + 04	1.8008e + 05	8.9389e + 01	3.6180e + 05	1.7959e + 02

For the MNLS solution, we generate a random vector $b \in \mathbb{R}^{12}$ in MATLAB by the command *randn*. The computed results for the bounds of structured and unstructured CNs are listed in Table 2. We observed that the bounds for the structured CNs are less than an order of 3 or 4 compared to the unstructured case.

Example 6.3. Let $M \in \mathbb{R}^{m \times n}$ be a CV matrix as in Definition 5.1 with the parameter set $\Psi_{\mathbb{CV}} = [c = \{c_i\}_{i=1}^m, d = \{d_i\}_{i=1}^l]^\top \in \mathbb{R}^{m+l}$, where c = 0.5 randn(m, 1) and d = randn(l, 1), generate by the command randn in MATLAB. For the MNLS solution, we take $b \in \mathbb{R}^m$. We choose n = 20 and l = 5. The computed results for the bounds of structured and unstructured

Table 3.: Comparison between upper bounds of structured and unstructured CNs for $M^{\dagger}(\Psi_{\mathbb{CV}})$ and $M^{\dagger}(\Psi_{\mathbb{CV}}, b)$ for Example 6.3.

m	$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{C}}^{\dagger}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}))$	$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{CV}}),b)$	$\tilde{\mathscr{C}^{\dagger}}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{CV}}), b)$
100	5.2481e + 04	8.5293e + 01	9.0595e + 07	2.8669e + 05	4.4045e + 04	1.5229e + 02	5.5927e + 04	4.3270e + 02
125	3.4639e + 04	6.4288e + 01	6.1046e + 07	2.5378e + 05	7.5900e + 04	2.3810e + 02	1.1277e + 06	5.4225e + 03
150	1.9732e + 04	5.7075e + 01	2.9498e + 07	1.0762e + 05	1.0708e + 04	6.2211e + 01	3.0915e + 04	2.8760e + 02
175	8.6870e + 04	8.6711e + 01	3.1972e + 07	1.1023e + 05	6.6658e + 04	2.1889e + 02	5.5834e + 05	7.4617e + 03
200	1.4574e + 04	8.7096e + 01	8.9156e + 07	3.0088e + 05	8.5138e + 03	7.8091e + 01	3.6041e + 05	4.3147e + 03

CNs are listed in Table 3 for different values of m ranging from 100 to 200 with step size 25. We observed that the structured CNs have significantly tighter bounds, almost 2 or 3 order smaller than those of the unstructured case for large matrices.

Example 6.4. We consider a random $\{1, 1\}$ -QS matrix M. For that, we first fixed n = 5 and use command *randn* to generate the vectors:

 $\mathbf{a} \in \mathbb{R}^{n-1}, \ \mathbf{b} \in \mathbb{R}^{n-1}, \ \mathbf{e} \in \mathbb{R}^{n-2}, \ \mathbf{d} \in \mathbb{R}^n, \ \mathbf{f} \in \mathbb{R}^{n-1}, \ \mathbf{g} \in \mathbb{R}^{n-2} \text{ and } \mathbf{h} \in \mathbb{R}^{n-1}.$

Using these vectors, we computed the $\{1, 1\}$ -QS matrix defined as (27). In Table 4, we see that for $M^{\dagger}(\Psi_{\mathbb{QS}})$ the bounds for structured CNs corresponding to $\Psi_{\mathbb{QS}}$ and structured effective CNs are substantially smaller than the unstructured ones. This validates the results of Proposition 5.11, Theorem 5.16 and Proposition 5.17. Next, for the MNLS solution, we generate a random

Table 4.: Comparison between upper bounds of unstructured, structured and effective CNs for the M-P inverse of $\{1, 1\}$ -QS matrices for Example 6.4.

$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}^{\dagger}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$
817.7856	3	2	5.3877e + 11	2.4189e + 05	1.5682e + 05

vector $b \in \mathbb{R}^n$ by the command *randn*. Table 5 illustrates that the results obtained for the computed bounds for structured CNs, effective CNs for MNLS solution are much smaller than those for unstructured ones and consistent with Proposition 5.12.

Table 5.: Comparison between upper bounds of unstructured, structured and effective CNs for the MNLS solution for $\{1, 1\}$ -QS matrices for Example 6.4.

$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}^{\dagger}}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$
1.7838e + 03	4.0024	3.0016	2.3669e + 03	8.8136	6.5184

Example 6.5. We consider several $\{1, 1\}$ -QS matrices of different orders. In fact, we choose n = 5, n = 7 and n = 10. We generate the random vectors $\mathbf{a}, \mathbf{b}, \mathbf{e}, \mathbf{d}, \mathbf{f}, \mathbf{g}$, and \mathbf{h} by the command randn in MATLAB as in the Example 6.4. After generating these vectors, we take following scaling

$$\mathbf{a} = a * 10^k, \ \mathbf{e} = e * 10^k, \ \mathbf{h} = h * 10^k,$$

where $k \in \{-1, -2, 0, 1, 2, 3\}$ to get unbalanced lower and upper right corner. In Table 6, we compare $\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ with $\tilde{\mathscr{M}}^{\dagger}(M)$ and $\tilde{\mathscr{C}}^{\dagger}(M)$, respectively, for the M-P inverse with different values of n. For the MNLS solution, we generate a random

Table 6.: Comparison between the upper bounds of unstructured, structured and effective CNs for the M-P inverse and the MNLS solution of $\{1, 1\}$ -QS matrices for Example 6.5.

n	$\tilde{\mathscr{M}^{\dagger}}(M)$	$\tilde{\mathscr{M}}^{\dagger}_{f}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}^{\dagger}(M)$	$\widetilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}}^{\dagger}(M,b)$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$
5	$1.4330e{+}04$	2	$2.1690e{+}12$	910	9.333e+04	3.0030	$1.1910e{+}05$	4.5451
7	$5.9139e{+}03$	3.0246	9.2340e + 08	74.2974	$6.5311e{+}04$	4.6655	2.0255e+04	6.7386
10	7.3293e + 04	2.0101	$1.5858e{+10}$	94.0499	$6.3988e{+}04$	1.0127	$2.5381e{+}05$	11.1271

vector $b \in \mathbb{R}^n$ for each choice of n. The computed bounds for the CNs are listed in Table 6. These results demonstrate the reliability of proposed CNs.

Example 6.6. In this example, we compare structured MCN and CCN with respect to QS representation and GV representation through tangent, structured effective CNs with their unstructured ones for M-P inverse and MNLS solution. On account of these, we generate the random vectors $t \in \mathbb{R}^{n-2}, u \in \mathbb{R}^{n-1}, v \in \mathbb{R}^{n-1}, w \in \mathbb{R}^{n-1}$ in MATLAB by the com and random vectors $t \in \mathbb{R}^{n-2}$.

for the parameter set $\Psi_{\mathcal{GV}}$. We set $\mathbf{d} = \mathbf{0} \in \mathbb{R}^n$ and rescale the vector v as v(1) = 0 and $v(n-1) = 10^2$. Then, we compute the $\{1,1\}$ -QS matrix M as in Definition 5.6. For different values of n, we generate 100 rank deficient $\{1,1\}$ -QS matrices. We use the formulae provided in Theorem 5.9 to compute the upper bounds $\tilde{\mathcal{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$ and $\tilde{\mathcal{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$. Again, we use the formulae for $\tilde{\mathcal{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathcal{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ presented as in Definition 5.15, and for $\tilde{\mathcal{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ and $\tilde{\mathcal{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$ presented as in Theorem 5.3.

We computed the above values for 100 randomly generated $\{1, 1\}$ -QS matrices for n = 30, 40, 50 and 60. In Table 7, average values of each upper bound of the CNs for the M-P inverse of these $\{1, 1\}$ -QS matrices are listed.

Table 7.: Comparison between upper bounds of unstructured, structured and effective CNs for M-P inverse of $\{1, 1\}$ -QS matrices for Example 6.6.

mean	n	$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{M}}^{\dagger}_{f}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}^{\dagger}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$
	30	1.0667e + 02	7.4873e + 01	1.2388e + 02	8.9215e + 01	2.3780e + 05	1.2730e + 04	2.2102e + 04	1.5815e + 04
	40	1.1429e + 02	7.4757e + 01	1.2356e + 02	8.9427e + 01	6.1267e + 08	1.4452e + 05	2.3753e + 05	1.7285e + 05
	50	1.6711e + 02	1.0323e + 02	1.7182e + 02	1.2296e + 02	6.9215e + 06	1.6133e + 05	3.2430e + 05	2.1980e + 05
	60	3.2135e + 02	1.8140e + 02	2.9452e + 02	2.1359e + 02	2.7856e + 07	3.0500e + 05	5.2242e + 05	3.9791e + 05

Table 8.: Comparison between upper bounds of unstructured, structured and effective CNs for the MNLS solution for $\{1, 1\}$ -QS matrices for Example 6.6.

mean	n	$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}),b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}^{\dagger}}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}),b)$	$\tilde{\mathscr{C}^{\dagger}}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$
	30	1.1278e + 02	7.8851e + 01	2.6113e + 02	9.4414e + 01	4.6667e + 03	1.7445e + 03	2.2102e + 04	2.1646e + 03
	40	1.1990e + 02	7.7823e + 01	2.4226e + 02	9.3005e + 01	7.3841e + 03	4.1774e + 03	2.3753e + 05	5.2069e + 03
	50	1.6512e + 02	1.0080e + 02	3.2184e + 02	1.2072e + 02	9.2143e + 03	4.2351e + 03	3.2430e + 05	5.3584e + 03
	60	3.3149e + 02	1.8715e + 02	7.6179e + 02	2.2120e + 02	7.0839e + 04	5.9482e + 04	5.2242e + 05	7.2054e + 04

Next, we generate a random vector $b \in \mathbb{R}^n$, for each value of n, and upper bounds for the structured CNs with respect to QS representation and GV representation through tangent, structured effective CNs and unstructured CNs for the MNLS solution are listed in Table 8.

The computed results in Tables 7 and 8 shows the consistency of Propositions 5.11–5.17 and Theorem 5.16. We can observe that the upper bounds of CNs for GV representation through tangent give more reliable bounds compared to the other CNs.

Example 6.7. In this example, we consider $\{1, 1\}$ -QS matrices of different orders. To construct $\{1, 1\}$ -QS matrices using the formula given in Definition 5.6, we generate the random vectors $t \in \mathbb{R}^{n-2}, u \in \mathbb{R}^{n-1}, v \in \mathbb{R}^{n-1}, w \in \mathbb{R}^{n-1}$ as in Example 6.6. Moreover, we generate $\mathbf{d} = randn(n) \in \mathbb{R}^n$ and set $\mathbf{d}(1) = 0$. Further, we rescale v by setting v(1) = 0 and v(n-1) = 1. For MNLS solution, we choose $b = randn(n) \in \mathbb{R}^n$. For different values of n, the computed upper bounds for structured CNs with respect to QS representation and GV representation through tangent, structured effective CNs and unstructured CNs for M-P inverse and MNLS solution are reported in Tables 9 and 10. These results confirm that our proposed upper bounds are reliable for large matrices, and structured ones are much sharper and smaller than unstructured ones.

Table 9.: Comparison between upper bounds of unstructured, structured and effective CNs for M-P inverse of $\{1, 1\}$ -QS matrices for Example 6.7.

n	$\tilde{\mathscr{M}}^{\dagger}(M)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}^{\dagger}(M)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}))$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}))$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}))$
100	4.3964e + 02	3.6260e + 02	5.3677e + 02	3.9861e + 02	4.5243e + 07	2.4837e + 04	3.6640e + 04	2.7217e + 04
150	4.1692e + 01	3.6846e + 01	5.2059e + 01	4.3613e + 01	1.0494e + 09	7.5769e + 06	1.1606e + 07	9.2937e + 06
200	3.3461e + 02	1.9504e + 02	2.9550e + 02	2.4237e + 02	1.5109e + 07	1.0192e + 05	1.9086e + 06	1.4250e + 05
250	2.1300e + 02	1.3904e + 02	2.1856e + 02	1.5019e + 02	4.8391e + 08	2.4495e + 06	4.1373e + 06	3.1253e + 06
300	2.0939e + 02	9.8454e + 01	1.6871e + 02	1.1458e + 02	1.1827e + 09	1.6759e + 06	1.7875e + 07	1.7776e + 06

Table 10.: Comparison between upper bounds of unstructured, structured and effective CNs for the MNLS solution for $\{1, 1\}$ -QS matrices for Example 6.7.

n	$\tilde{\mathscr{M}}^{\dagger}(M,b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathcal{GV}}),b)$	$\tilde{\mathscr{M}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{M}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}^{\dagger}}(M,b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathcal{GV}}),b)$	$\tilde{\mathscr{C}}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$	$\tilde{\mathscr{C}}_{f}^{\dagger}(M(\Psi_{\mathbb{QS}}),b)$
100	4.3739e + 02	3.6376e + 02	2.0884e + 02	4.0061e + 02	6.1525e + 03	4.8648e + 02	2.5191e + 03	5.4354e + 03
150	4.7556e + 01	4.3981e + 01	7.0553e + 01	5.1641e + 01	5.3904e + 03	3.3664e + 02	1.2982e + 03	3.8831e + 02
200	3.4402e + 02	2.0275e + 02	1.0998e + 02	2.5110e + 02	1.4342e + 04	5.3603e + 03	1.1052e + 04	6.5164e + 03
250	1.0986e + 02	1.0489e + 02	1.8630e + 02	1.1328e + 02	8.4676e + 04	5.1797e + 03	1.0935e + 04	6.0154e + 03
300	2.1839e + 02	1.0345e + 02	2.6652e + 02	1.2097e + 02	3.4116e + 03	1.4340e + 03	6.5971e + 03	1.7712e + 03

7. Conclusion

For the M-P inverse and the MNLS solution, we investigated structured MCN and CCN corresponding to a class of parameterized matrices, with each entry as a differentiable function of some real parameters. This framework has been used to derive the upper bounds of structured CNs for CV and {1,1}-QS matrices. QS representation and the GV representation through tangent are considered for {1,1}-QS matrices to investigate their structured CNs. It is proved that upper bounds for the structured CNs for GV representation through tangent are always smaller than the QS representation. Numerical examples demonstrate that the proposed structured effective CNs are significantly smaller in most cases. Our methodology can be applied to investigate CNs involving other rank-structured matrices. It would be interesting to generalize our findings to the weighted M-P inverse and weighted LS problems, as well as linear LS problems with multiple right-hand sides. Following that, research progress on the aforementioned topics will be reported elsewhere.

Acknowledgments

We gratefully acknowledge the financial assistance (File no. 09/1022(0098)/2020-EMR-I) provided as a fellowship to Pinki Khatun by the Council of Scientific & Industrial Research (CSIR), New Delhi, India. The authors are grateful to anonymous reviewers for their constructive comments and suggestions, which have contributed to enhancing the quality of this paper. We extend our sincere gratitude to Prof. Huaian Diao for providing us with the thesis [37].

Declarations

Conflict of interest The authors declare no competing interests.

References

- [1] A. Ben-Israel. On error bounds for generalized inverses. SIAM J. Numer. Anal., 3:585–592, 1966.
- [2] P. Bürgisser and F. Cucker. Condition: The Geometry of Numerical Algorithms, volume 349. Springer Science & Business Media, 2013.
- [3] S. Chandrasekaran, P. Dewilde, M. Gu, T. Pals, and A. J. van der Veen. Fast stable solver for sequentially semi-separable linear systems of equations. In *International Conference on High-Performance Computing*, pages 545–554, Springer, 2002.
- [4] S. Chountasis, V. N. Katsikis, and D. Pappas. Applications of the Moore-Penrose inverse in digital image restoration. *Math. Probl. Eng.*, page 12, 2009.
- [5] S. Chountasis, V. N. Katsikis, and D. Pappas. Digital image reconstruction in the spectral domain utilizing the Moore-Penrose inverse. *Math. Probl. Eng.*, page 14, 2010.
- [6] D. Colton and R. Kress. Inverse Acoustic and Electromagnetic Scattering Theory, volume 93 of Appl. Math. Sci. Springer-Verlag, Berlin, Second edition, 1998.
- [7] F. Cucker and H. Diao. Mixed and componentwise condition numbers for rectangular structured matrices. *Calcolo*, 44(2):89–115, 2007.
- [8] F. Cucker, H. Diao, and Y. Wei. On mixed and componentwise condition numbers for Moore-Penrose inverse and linear least squares problems. *Math. Comp.*, 76(258):947–963, 2007.
- [9] H. Diao and Q. Meng. Structured generalized eigenvalue condition numbers for parameterized quasiseparable matrices. *BIT*, 59(3):695–720, 2019.
- [10] H. Diao and Y. Wei. On Frobenius normwise condition numbers for Moore-Penrose inverse and linear least-squares problems. *Numer. Linear Algebra Appl.*, 14(8):603–610, 2007.
- [11] H. Diao, L. Li, and Q. Meng. Structured condition numbers for sylvester matrix equation with parameterized quasiseparable matrices. Comm. Anal. Comp., 1(3):183–213, 2023.
- [12] F. M. Dopico and K. Pomés. Structured eigenvalue condition numbers for parameterized quasiseparable matrices. *Numer. Math.*, 134(3):473–512, 2016.
- [13] F. M. Dopico and K. Pomés. Structured condition numbers for linear systems with parameterized quasiseparable coefficient matrices. *Numer. Algorithms*, 73(4):1131–1158, 2016.
- [14] Y. Eidelman and I. Gohberg. On a new class of structured matrices. Integral Equations Operator Theory, 34(3):293–324, 1999.
- [15] S. Gratton. On the condition number of linear least squares problems in a weighted Frobenius norm. BIT, 36(3):523–530, 1996.
- [16] L. Greengard and V. Rokhlin. On the numerical solution of two-point boundary value problems. Comm. Pure Appl. Math., 44(4):419–452, 1991.
- [17] N. J. Higham. Accuracy and Stability of Numerical Algorithms. SIAM, Philadelphia, Second edition, 2002.
- [18] R. Huang. Accurate solutions of weighted least squares problems associated with rank-structured matrices. Appl. Numer. Math., 146:416–435, 2019.
- [19] L. A. Imhof and W. J. Studden. E-optimal designs for rational models. Ann. Statist., 29(3):763–783, 2001.
- [20] J. Y. Lee and L. Greengard. A fast adaptive numerical method for stiff two-point boundary value problems. SIAM J. Sci. Comput., 18(2):403–429, 1997.
- [21] Z. Li, Q. Xu, and Y. Wei. A note on stable perturbations of Moore-Penrose inverses. Numer. Linear Algebra Appl., 20(1):18–26, 2013.
- [22] A. N. Malyshev. A unified theory of conditioning for linear least squares and Tikhonov regularization solutions. SIAM J. Matrix Anal. Appl., 24(4):1186–1196, 2003.
- [23] Q. Meng, H. Diao, and Q. Yu. Structured condition number for multiple right-hand side linear systems with parameterized quasiseparable coefficient matrix. J. Comput. Appl. Math., 368:112527, 20, 2020.

- [24] G. Mühlbach. Interpolation by Cauchy-Vandermonde systems and applications. J. Comput. Appl. Math., 122(1-2):203-222, 2000.
- [25] C. R. Rao and S. K. Mitra. Generalized Inverse of Matrices and its Applications. John Wiley & Sons, Inc., New York-London-Sydney, 1971.
- [26] J. R. Rice. A theory of condition. SIAM J. Numer. Anal., 3:287-310, 1966.
- [27] J. Rohn. New condition numbers for matrices and linear systems. Computing, 41(1-2):167–169, 1989.
- [28] R. D. Skeel. Scaling for numerical stability in Gaussian elimination. J. Assoc. Comput. Mach., 26 (3):494–526, 1979.
- [29] G. W. Stewart and J. G. Sun. Matrix Perturbation Theory. Academic Press, New York, 1990.
- [30] R. Vandebril, M. Van Barel, and N. Mastronardi. A note on the representation and definition of semiseparable matrices. *Numer. Linear Algebra Appl.*, 12(8):839–858, 2005.
- [31] R. Vandebril, M. Van Barel, and N. Mastronardi. *Matrix Computations and Semiseparable Matrices*, volume 1. Johns Hopkins University Press, Baltimore, MD, 2008.
- [32] R. Vandebril, M. Van Barel, and N. Mastronardi. *Matrix Computations and Semiseparable Matrices*, volume 2. Johns Hopkins University Press, Baltimore, MD, 2008.
- [33] G. Wang, Y. Wei, and S. Qiao. Generalized Inverses: Theory and Computations. Springer, Science Press, Beijing, Second edition, 2018.
- [34] Y. Wei and D. Wang. Condition numbers and perturbation of the weighted Moore-Penrose inverse and weighted linear least squares problem. Appl. Math. Comput., 145(1):45–58, 2003.
- [35] Y. Wei, W. Xu, S. Qiao, and H. Diao. Componentwise condition numbers for generalized matrix inversion and linear least squares. *Numer. Math. J. Chinese Univ.*, 14(3):277–286, 2005.
- [36] Y. Wei, H. Diao, and S. Qiao. Condition number for weighted linear least squares problem. J. Comput. Math., 25(5):561–572, 2007.
- [37] X. Wu. Structured condition numbers for least squares problems with parameterized quasiseparable coefficient matrices. Northeast Normal University, Master thesis, under the supervision of H. Diao, 2020.
- [38] W. Xu, Y. Wei, and S. Qiao. Condition numbers for structured least squares problems. *BIT*, 46(1): 203–225, 2006.
- [39] Z. Yang. Accurate computations of eigenvalues of quasi-Cauchy-Vandermonde matrices. Linear Algebra Appl., 622:268–293, 2021.